

Tutorial of fragment split by MolAICal

Qifeng Bai

Email: molaical@yeah.net

Homepage: <https://molaical.github.io>

School of Basic Medical Sciences

Lanzhou University

Lanzhou, Gansu 730000, P. R. China

1. Introduction

Sometime, to study the fragment properties of ligands, the ligands need to be split into smaller pieces. Here, MolAICal provides a way to split ligands into small fragments according to rotatable bonds. In this tutorial, the method of fragment split is introduced based on the protocol of MolAICal (<https://doi.org/10.1093/bib/bbaa161>).

2. Materials

2.1. Software requirement

1) MolAICal: <https://molaical.github.io>

2.2. Example files

1) All the necessary tutorial files are downloaded from:


<https://github.com/MolAICal/tutorials/tree/master/013-fragmentSplit>

3. Procedure

Change directory to “013-fragmentSplit”:

```
#> cd 013-fragmentSplit
```

Preparing the input file named “list.dat” that contains the files’ name of ligands:



1. Split ligands into small fragments without adding or deleting hydrogen

```
#> molaical.exe -tool fragSplit -i list.dat -o initial -oa results
```

Or

```
#> molaical.exe -tool fragSplit -i list.dat -o initial -oa results -w off
```

2. Split ligands into small fragments with adding hydrogen

```
#> molaical.exe -tool fragSplit -i list.dat -o initial -oa results -w add
```

3. Split ligands into small fragments with deleting hydrogen

```
#> molaical.exe -tool fragSplit -i list.dat -o initial -oa results -w del
```

Note: The folder “initial” includes split fragments without adding or deleting hydrogen. The folder “results” contains fragments that have been dealt with command parameter “-w”. The parameter “-w” indicates whether MolAICal deletes or adds hydrogens on the receptor.

If you want to merge the split fragments or ligands into one file, the command can be used as below:

```
#> molaical.exe -tool merge -m mol2 -i results -o all_frag.mol2
```