

Cc1cc(N)cc(C(=O)O[C@H]2[C@@H](OC)[C@H](OC)[C@H](OC)[C@H]2Oc3cc4c(cc3)c(=O)oc5cc(C)c(N)nc45)c1

The chemical structure shows a central benzene ring with three substituents: a furanone ring at the top, a phenol group at the bottom, and a chiral auxiliary at the right. The furanone ring is a five-membered ring with an oxygen atom and a carbonyl group. The phenol group is a benzene ring with a hydroxyl group. The chiral auxiliary is a complex group containing a furanone ring, a chiral center, and a hydroxyl group. The structure is drawn with black lines for the main framework, red lines for the auxiliary, and blue lines for the furanone rings. Stereochemistry is indicated with wedges and dashes.

CC1=CNC=C1C(=O)O[C@H]2[C@@H](OC(=O)c3cc(C)cc3)[C@H](OC(=O)c4cc5c(c3)oc(=O)c5c4O)[C@@H](OC(=O)c6ccc(N)cc6)[C@H]2O

The chemical structure shows a complex molecule. On the left, a 2-methylpyridine ring is connected via an ester linkage to a furanose sugar. The sugar is substituted with a methoxy group and a quaternary carbon. This sugar is linked via an ether bond to a large polycyclic aromatic system. This system includes a benzofuran moiety, a central benzene ring with a hydroxyl group, and a benzimidazole moiety with an amino group.

Cc1cc[nH]c1C(=O)O[C@H]2[C@@H](OC)[C@H](OC[C@H]3[C@@H](OC)[C@H](OC)[C@H]3OC(=O)c4cc(OC)c5cc(O)c(OC)c5c4)[C@H]2OCCc1cc[nH]c1C(=O)O[C@H]2[C@@H](OC)[C@H](OC)[C@@H](OC)[C@H](OC)[C@H]2Oc3cc(=O)c4c(c3)oc(=O)c4Oc5cc(=O)c6c(c5)oc(=O)c6NS(=O)(=O)Cc7cc(=O)c8c(c7)oc(=O)c8NCc1cc[nH]c1C(=O)O[C@H]2[C@@H](OC)[C@H](O)[C@@H](COc3c4c(c5cc(OC)c6c5c(=O)oc7c6cc(NC8CCCC8)cc7c)cc34)O2

Chemical structure of compound 10, showing a complex molecule with a pyridine ring, a sugar moiety, a naphthalene ring, and a carboxylic acid group.

Chemical structure of compound 10, showing a complex molecule with a pyridine ring, a sugar moiety, a naphthalene ring, and a carboxylic acid group.

The chemical structure of compound 10 is a complex molecule. It features a pyridine ring on the left, connected via an ester linkage to a furanose sugar. The sugar is further connected to a coumarin core, which is linked to a pyrazole ring on the right. The pyrazole ring has a methyl group and a carbonyl group. The coumarin core has a hydroxyl group and a carbonyl group. The pyridine ring has a methyl group. The furanose sugar has a methoxy group and a hydroxyl group. The pyrazole ring has a methyl group and a carbonyl group. The coumarin core has a hydroxyl group and a carbonyl group. The pyridine ring has a methyl group.

The chemical structure shows a complex molecule with several fused and linked rings. On the left, a pyridine ring is connected via a carbonyl group to a furanose sugar. The sugar is linked to a coumarin derivative, which is further connected to a pyrazole ring. The structure includes various functional groups such as hydroxyl groups, carbonyl groups, and a pyridine nitrogen atom.

The chemical structure is a complex molecule consisting of several interconnected rings and functional groups. On the left, a 4-methylpyridine ring is connected via its nitrogen atom to a carbonyl group, which is further linked to a furanose sugar ring. The sugar ring has a hydroxyl group and a methoxy group. This sugar is connected to a coumarin derivative, which has a hydroxyl group and a carbonyl group. The coumarin is linked to a biphenyl system, which has an amino group and a phenyl ring attached to it.

The chemical structure of compound 10 is a complex molecule. It features a pyridine ring on the left, connected via an ester linkage to a furanose sugar. The sugar is further connected to a naphthalene system, which is linked to a quinoxaline moiety on the right. The quinoxaline has a positive charge on the nitrogen atom and an amino group at the 8-position. The naphthalene system has a hydroxyl group at the 1-position and a methoxy group at the 2-position. The pyridine ring has a methyl group at the 4-position.

Combinatorial   RL   RL Docking