



筑波大学

University of Tsukuba

Linear models for Qualitative Responses

Marco Antonio Florenzano Mollinetti¹

¹**University of Tsukuba, Systems Optimization Laboratory**

mollinetti@syou.cs.tsukuba.ac.jp




Before we Begin

- Go to the github repo:
 - <https://github.com/Mollinetti/Experiment-Design-R>
 - Download the script for this class! (in the 'scripts' folder, class_7.r!)
- Run the snippet at the beginning to load/install the required libraries



Agenda

- Introduction
- Classification of Quantitative responses
- Logistic Regression
- Linear Determinant Analysis (LDA)
- Quadratic Determinant Analysis (QDA)

- 
- Introduction
 - Classification of Quantitative responses
 - Logistic Regression
 - Linear Determinant Analysis (LDA)
 - Quadratic Determinant Analysis (QDA)



Introduction

- Classification problems are **frequent**:
 1. Someone has a set of symptoms that can be attributed to three conditions. Which of the three does the person have?
 2. On a DNA sequence data of several patients with and without a disease, which DNA mutations are disease causing and which are not?




Introduction


- We will now fit a linear model to classify **categorical variables**
- For a 2 level qualitative response:
 - Logistic Regression*
- For more than two levels:
 - LDA
 - QDA

Introduction

- Why we can't use linear regression anymore?



Answer: Negative Prediction values!

- 
- Introduction
 - Classification of Quantitative responses
 - Logistic Regression
 - Linear Determinant Analysis (LDA)
 - Quadratic Determinant Analysis (QDA)

Classification of Quantitative Responses

- Remember, the response now has **many levels**
- For such, we can do a similar approach to the dummy variables
- **Define your contrasts beforehand**

$$Y = \begin{cases} 1 & \text{if response 1} \\ 2 & \text{if response 2} \\ 3 & \text{if response 3} \end{cases} \quad \text{or} \quad Y = \begin{cases} 01 & \text{if response 1} \\ 10 & \text{if response 2} \\ 00 & \text{if response 3} \end{cases}$$

Classification of Quantitative Responses

- How does a classifier outputs its predictions?
- Suppose we want to classify **apples, oranges and pears**
- We know by default that each variable is modeled in this specific way:

$$\text{apple} = \begin{bmatrix} 0 \\ 1 \end{bmatrix} \quad \text{orange} = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \quad \text{pear} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$


Classification of Quantitative Responses

- Values outputted by the classifier close to the labels corresponds to each level:

$$\begin{bmatrix} 0.0002 \\ 0.9986 \end{bmatrix} \approx \begin{bmatrix} 0 \\ 1 \end{bmatrix} = \text{apple}$$

$$\begin{bmatrix} 0.9780 \\ 0.0642 \end{bmatrix} \approx \begin{bmatrix} 1 \\ 0 \end{bmatrix} = \text{orange}$$

$$\begin{bmatrix} 0.9343 \\ 0.9846 \end{bmatrix} \approx \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \text{pear}$$

- 
- Introduction
 - Classification of Quantitative responses
 - **Logistic Regression**
 - Linear Determinant Analysis (LDA)
 - Quadratic Determinant Analysis (QDA)

Logistic Regression

- Logistic regression models that the **probability that the response belongs to a category**
- $p(X) = \Pr(Y = 1|X)$
- Best suited for two classes:
 - $p(X) < 0.5$ so Y belong to category 1
 - $p(X) \geq 0.5$ so Y belong to category 2

Logistic Regression

- If we follow the linear model (with additive and linear relations):

$$p(X) = \beta_0 + \beta_1 X_1 + \cdots \beta_p X_p + \epsilon$$

- Then $p(X)$ will have negative values, unacceptable for binary classification


Logistic Regression

- $p(X)$ has to be a model in the $[0,1]$ interval
- This can be done by the sigmoid function

$$p(X) = \frac{e^{\beta_0 + \beta_1 X_1 + \dots + \beta_p}}{1 + e^{\beta_0 + \beta_1 X_1 + \dots + \beta_p}}$$

Logistic Regression

- The **sigmoid function** has the following shape:



Logistic Regression

- The model is now fitted using the **maximum likelihood estimator**
- The **log-odds** or **logit** captures the linear relation of the model:

$$\log \left(\frac{p(X)}{1 - p(X)} \right) = e^{\beta_0 + \beta_1 X_1 + \dots + \beta_p X_p}$$

- In a logistic regression, **increasing X changes the log odds of the coefficients**



Logistic Regression

- Load the “BreastCancer” dataset from the mlbench library
- 33 columns (we’ll use 11)
- 1 qualitative variable (response)
 - 2 levels {benign, malignant}
- 10 quantitative variables (predictors)



Logistic Regression

- We will do first the regression with every quantitative variable, then we will adjust the model
- Verification of the assumptions are also required
- Goodness of fit:
 - Null Deviance (only the intercept)
 - Residual Deviance (all coefficients)
 - BIC/ AIC

Logistic Regression


- In R, the glm is called for logistic regression

```
#logistic regression with all variables but the id (we use glm for that)  
reg.fit <- glm(diagnosis~., data = BreastCancer, family = binomial)
```



Logistic Regression

- In R, the glm is called for logistic regression



Logistic Regression

- Calling summary to the model gives the following info:

```
Deviance Residuals:
    Min       1Q   Median       3Q      Max
-1.95590  -0.14839  -0.03943   0.00429   2.91690

Coefficients:
              Estimate Std. Error z value Pr(>|z|)
(Intercept)   -7.35952    12.85259  -0.573   0.5669
radius_mean   -2.04930     3.71588  -0.551   0.5813
texture_mean    0.38473     0.06454   5.961 2.5e-09 ***
perimeter_mean -0.07151     0.50516  -0.142   0.8874
area_mean      0.03980     0.01674   2.377   0.0174 *
smoothness_mean 76.43227    31.95492   2.392   0.0168 *
compactness_mean -1.46242    20.34249  -0.072   0.9427
concavity_mean  8.46870     8.12003   1.043   0.2970
concave_points_mean 66.82176    28.52910   2.342   0.0192 *
symmetry_mean  16.27824    10.63059   1.531   0.1257
fractal_dimension_mean -68.33703    85.55666  -0.799   0.4244
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

(Dispersion parameter for binomial family taken to be 1)

    Null deviance: 751.44  on 568  degrees of freedom
Residual deviance: 146.13  on 558  degrees of freedom
AIC: 168.13

Number of Fisher Scoring iterations: 9
```



Logistic Regression

- Calling summary to the model gives the following info:

```
Deviance Residuals:
    Min       1Q   Median       3Q      Max
-1.95590 -0.14839 -0.03943  0.00429  2.91690

Coefficients:
              Estimate Std. Error z value Pr(>|z|)
(Intercept)   -7.35952    12.85259  -0.573   0.5669
radius_mean    -2.04930     3.71588  -0.551   0.5813
texture_mean     0.38473     0.06454   5.961 2.5e-09 ***
perimeter_mean  -0.07151     0.50516  -0.142   0.8874
area_mean        0.03980     0.01674   2.377   0.0174 *
smoothness_mean  76.43227    31.95492   2.392   0.0168 *
compactness_mean -1.46242    20.34249  -0.072   0.9427
concavity_mean    8.46870     8.12003   1.043   0.2970
concave_points_mean 66.82176    28.52910   2.342   0.0192 *
symmetry_mean    16.27824    10.63059   1.531   0.1257
fractal_dimension_mean -68.33703    85.55666  -0.799   0.4244
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

(Dispersion parameter for binomial family taken to be 1)

    Null deviance: 751.44  on 568  degrees of freedom
Residual deviance: 146.13  on 558  degrees of freedom
AIC: 168.13
```

Z statistic:
Coefficient/standard error

Z statistic associated p-
value

Goodness of Fit for
Logistic regression

Number of Fisher Scoring iterations: 9



Logistic Regression

- We now predict the values
- Get the estimated value of each observation, then associate it to the class that it belongs

```
> glm.probs = predict(reg.fit, type = "response")
```

```
> glm.probs[1:10]
```

1	2	3	4	5	6	7	8	9	10
0.9999694	0.9999894	0.9999999	0.9822760	0.9999987	0.6692722	0.9995617	0.7736633	0.9923907	0.9647326



Logistic Regression

```
#convert the probabilities of glm.probs into proper classes "Malignant" or "Benign"  
#create a vector of 1250 "benign"  
glm.pred = rep("B",nrow(BreastCancer))  
#Fill with "Malignant" whatever probabilities above 0.5 (our chosen threshold)  
glm.pred[glm.probs > 0.5] = "M"  
glm.pred[1:10]
```

```
· glm.pred[1:10]  
[1] "M" "M" "M" "M" "M" "M" "M" "M" "M" "M"
```



Logistic Regression

- Calculate a **confusion matrix** to check for **false positive** and **false negatives**

```
> table(glm.pred, diagnosis)
      diagnosis
glm.pred  B    M
      B 347  19
      M  10 193
```

- Check the mean of the predictions for **accuracy**

```
> mean(glm.pred == diagnosis)
[1] 0.9490334
```

ALWAYS REMEMBER TO CHECK BOTH





Logistic Regression

- Logistic regression **does not** require:
 - ☐ Normality
 - ☐ Homoscedascity
 - ☐ Linearity




Logistic Regression

- However, Logistic regression adhere to:
 - Linearity of independent variables and log odds
 - Influential values
 - Collinearity
 - Large sample size

Logistic Regression

- Linearity of log odds
- inspecting the scatter plot between each predictor and the logit values






logit

Logistic Regression

Influential Values

- Plot of the studentized residuals against leverage



Logistic Regression

Collinearity


- Check the correlation
- Compute the VIF

```
> vif(reg.fit) # variance inflation factors
```

radius_mean	texture_mean
899.5200	1.8064
perimeter_mean	area_mean
698.9800	129.5600
smoothness_mean	compactness_mean
4.3729	15.2810
concavity_mean	concave_points_mean
5.2595	5.8564
symmetry_mean	fractal_dimension_mean
1.8395	9.7877

```
> sqrt(vif(reg.fit)) > 2 # problem? cutoff is 5 or 10
```

radius_mean	texture_mean
TRUE	FALSE
perimeter_mean	area_mean
TRUE	TRUE
smoothness_mean	compactness_mean
TRUE	TRUE
concavity_mean	concave_points_mean
TRUE	TRUE
symmetry_mean	fractal_dimension_mean
FALSE	TRUE

- 
- Introduction
 - Classification of Quantitative responses
 - Logistic Regression
 - Linear Determinant Analysis (LDA)
 - Quadratic Determinant Analysis (QDA)



Linear Determinant Analysis (LDA)

- Logistic Regression is not recommended for more than 2 classes
- Does not work well when classes are well separated
- LDA is more stable for smaller and normal X
- Assumes that probabilities are distributed from a gaussian distribution

Linear Determinant Analysis (LDA)

- Let Y be able to take on distinct K values of classes
- LDA models the probability of X belonging to a Y as an approximation of the Bayes' Theorem

$$\Pr(Y = K | X = x) = \frac{\pi_k f_k(x)}{\sum_{l=1}^K \pi_l f_l(x)}$$


Linear Determinant Analysis (LDA)

- Let Y be able to take on distinct K values of classes
- LDA models the probability of X belonging to a Y as an approximation of the Bayes' Theorem

$$\Pr(Y = K | X = x) = \frac{\pi_k f_k(x)}{\sum_{l=1}^K \pi_l f_l(x)}$$

Prior probability

Posterior probability
(Density function)



Linear Determinant Analysis (LDA)

- For a multivariate gaussian distribution the density function is:

$$f(x) = \frac{1}{(2\pi)^{p/2} |\Sigma|^{1/2}} \exp \left(-\frac{1}{2} (x - \mu)^T \Sigma^{-1} (x - \mu) \right).$$

Diagram illustrating the components of the multivariate Gaussian density function:

- Variance**: Points to the term $(2\pi)^{p/2}$.
- Covariance matrix**: Points to the term $|\Sigma|^{1/2}$.
- Mean**: Points to the term μ .

Linear Determinant Analysis (LDA)

- Values of parameters μ_1, \dots, μ_K and π_1, \dots, π_K must be estimated for each class
- Each value is calculated as:

$$\hat{\mu}_K = \frac{1}{n_K} \sum_{i:y_i=K} x_i$$
$$\hat{\sigma}^2_K = \frac{1}{n - K} \sum_{k=1}^K \sum_{i:y_i=k} x_i$$

Linear Determinant Analysis (LDA)

- The estimated values are plugged into the following formula and classified as the one that maximizes for given class
- The covariance matrix is the same for each class

$$\delta_k(x) = x^T \Sigma^{-1} \mu_k - \frac{1}{2} \mu_k^T \Sigma^{-1} \mu_k + \log \pi_k$$

Linear Determinant Analysis (LDA)


- In R, LDA is part of the MASS library

```
lda.fit <- lda(diagnosis~., data = BreastCancer)
```



Linear Determinant Analysis (LDA)

- In R, LDA is part of the MASS library



Linear Determinant Analysis (LDA)

■ Calling the object:

```
> lda.fit
```

```
Call:
```

```
lda(diagnosis ~ ., data = BreastCancer)
```

```
Prior probabilities of groups:
```

```
      B      M  
0.6274165 0.3725835
```

```
Group means:
```

	radius_mean	texture_mean	perimeter_mean	area_mean	smoothness_mean	compactness_mean	concavity_mean	concave_points_mean	symmetry_mean
B	12.14652	17.91476	78.07541	462.7902	0.09247765	0.08008462	0.04605762	0.02571741	0.174186
M	17.46283	21.60491	115.36538	978.3764	0.10289849	0.14518778	0.16077472	0.08799000	0.192909

	fractal_dimension_mean
B	0.06286739
M	0.06268009

```
Coefficients of linear discriminants:
```

	LD1
radius_mean	2.173832578
texture_mean	0.097479319
perimeter_mean	-0.243883158
area_mean	-0.004235635
smoothness_mean	8.610211091
compactness_mean	0.431476344
concavity_mean	3.592356858
concave_points_mean	28.529778564
symmetry_mean	4.489073661
fractal_dimension_mean	-0.529214778



Logistic Regression

- Calculate yet again a **confusion matrix** to check for **false positives** and **false negatives**

```
> table(lda.class, diagnosis)
      diagnosis
lda.class  B    M
B      351   29
M         6  183
```






Linear Determinant Analysis (LDA)

- Class-specific accuracy is **more important** than general accuracy in **some cases**
- Possible way to improve: **change the threshold values**
- However, how can we decide the best threshold value?

Linear Determinant Analysis (LDA)

- Plot of the ROC curve
- Plot all possible thresholds and measure the AUC (area under curve)
- A curve that hugs the top left is desired





Linear Determinant Analysis (LDA)

- Assumptions:

- ☐ Equality of Variance-Covariance
- ☐ Normality




Linear Determinant Analysis (LDA)

- Equality of Variance-Covariance:
 - ☐ Plot of the boxplots
 - ☐ Plot of the covariances homogeneity
 - ☐ Box M-test


Linear Determinant Analysis (LDA)

- Plot the boxplots of variances between classes




Linear Determinant Analysis (LDA)

■ Plot of the covariances homogeneity



Linear Determinant Analysis (LDA)

■ Box M-test



```
> boxM(BreastCancer[,-1], diagnosis)
```

Box's M-test for Homogeneity of Covariance Matrices

data: BreastCancer[, -1]

Chi-Sq (approx.) = 1738.8, df = 55, p-value < 2.2e-16






Linear Determinant Analysis (LDA)

- Normality:
 - QQ-plots
 - Shapiro-wilk test

Linear Determinant Analysis (LDA)

- QQ-plots: plotting the predicted value against the predictor



Linear Determinant Analysis (LDA)

- For predictors that are **clearly not normal**, run the **shapiro-wilk test against predicted values**

```
> #shapiro-wilk test on dubious columns  
> shapiro.test(pred.m$radius_mean)
```

Shapiro-Wilk normality test


```
data:  pred.m$radius_mean  
W = 0.97766, p-value = 0.001895
```

Transform
the data if
not normal

```
> #shapiro-wilk on transformed data  
> shapiro.test(sqrt(pred.m$radius_mean))
```

Shapiro-Wilk normality test

```
data:  sqrt(pred.m$radius_mean)  
W = 0.9884, p-value = 0.08347
```


- 
- Introduction
 - Classification of Quantitative responses
 - Logistic Regression
 - Linear Determinant Analysis (LDA)
 - Quadratic Determinant Analysis (QDA)



Quadratic Determinant Analysis

- QDA still assumes that probabilities are drawn from a multivariate gaussian distribution
- However, each class now has its own covariance matrix
- More flexible to fit curves

Quadratic Determinant Analysis



Quadratic Determinant Analysis

- Estimated values are plugged into the following function:

$$\delta_k(x) = -\frac{1}{2}x^T \Sigma_k^{-1}x + x^T \Sigma_k^{-1}\mu_k - \frac{1}{2}\mu_k^T \Sigma_k^{-1}\mu_k - \frac{1}{2}\log|\Sigma_k| + \log \pi_k$$

Quadratic Determinant Analysis


- In R, QDA is also part of the MASS library

```
#QUADRATIC DISCRIMINANT ANALYSIS  
qda.fit = qda(diagnosis~., data = BreastCancer)
```



Quadratic Determinant Analysis

- In R, QDA is also part of the MASS library



Quadratic Determinant Analysis

- **Predictions** and **confusion matrix** must also be estimated

```
> qda.class = predict(qda.fit, BreastCancer)$class
> table(qda.class, diagnosis)
```

	diagnosis	
qda.class	B	M
B	346	26
M	11	186





Quadratic Determinant Analysis

- Assumptions of QDA are the same as LDA
 - Equality of Variance-Covariance
 - Normality



Next Episode

- Now that we understand how to fit lines, we will go back to the Multiple Analysis of Variances (MANOVA)



筑波大学

University of Tsukuba

Multiple Linear Regression Model

Marco Antonio Florenzano Mollinetti¹

¹**University of Tsukuba, Systems Optimization Laboratory**

mollinetti@syoun.cs.tsukuba.ac.jp