


LSAP communities near mapped landfills

This report explains the source data, executed workflow and interpretation of lsap communities near mapped landfills. It documents the existing ArcGIS Pro output, including its coverage limits and retained ecclesiastical context. (OpenStreetMap contributors, 2026; OpenStreetMap Wiki contributors, n.d.).

Map P10 | Documentation date 6 October 2026 | Units Kilometres | Inventory retrieved 6 October 2026

LSAP communities near mapped landfills

Nearest mapped landfill footprint - geodesic distance in km



Distance to polygon edge; zero if inside. Site presence and proximity do not establish contamination. Member precision is unverified.

© OpenStreetMap contributors - ODbL - copied LSAP foundations

LSAP POLLUTION ATLAS / COPIED ECCLESIASTICAL AND MEMBER FOUNDATIONS / P10

Figure P10 World overview exported from the completed ArcGIS Pro layout.

Distance to polygon edge; zero if inside. Site presence and proximity do not establish contamination. Member precision is unverified.

All analyses use copies of 2,874 dioceses and 2,739 LSAP member points. Copies of 122 episcopal conference features and 194 country features preserve ecclesiastical context. Stable DIO_ID links outputs to the copied boundaries. Original foundational data were not edited (LSAP Pollution Atlas, 2026, foundation_manifest.json).

Analysis workflow

The following numbered sequence describes the executed ArcPy and Python analyses as a typed ModelBuilder style workflow. Custom raster, station aggregation and geometry operations are script steps, not a claim that an ArcGIS ModelBuilder model file was run. Source data: (OpenStreetMap contributors, 2026; OpenStreetMap Wiki contributors, n.d.).

1. Copy foundational members into Member_Landfill_Proximity.
2. Generate geodesic distance to nearest landfill polygon edge; members inside a footprint receive zero distance.
3. Generate all near matches within 20 km; summarize distinct sites within 1, 5 and 20 km and join to every member.
4. Classify nearest distances at 1, 5, 20 and 50 km; nearby members are darker. Retain local footprint detail and diocesan/conference context.
5. Create popups with nearest site ID, distance and buffer counts; preserve source inventory and all member attributes.

Output fields and files

Field or band	Meaning
NearSite_ID and NearSite_km	Nearest mapped site and distance to footprint edge
Sites1km Sites5km Sites20km	Distinct site counts at three distance thresholds
ProximityNote	Interpretation and positional precision note

Analysis dataset: PollutionAnalysis.gdb/Member_Landfill_Proximity. Styles/P10.lyrx and Styles/P10.mapx preserve layer and map definitions. Map_Exports/P10_World_overview.png preserves the layout preview. The editable project is LSAP_Pollution.aprx.

Interpretation and data handling

Site presence and proximity do not establish pollution or exposure. LSAP coordinates have unverified precision. Waste Atlas remains a documented supplement gap.

The working inventory contains 51,748 unique sites after removing 2,529 duplicate records. Of these, 50,996 receive a primary diocese assignment and 752 remain unassigned. Raw regional JSON, OSM_Landfill_Source and Landfills_Working remain available. Site control status and pollution severity are unknown. The European mirror is older than most other regions; retrieval date does not imply a uniform data vintage (LSAP Pollution Atlas, 2026, waste_QA.json).

Cartographic classes are descriptive display intervals, not regulatory or ecological quality grades. The pollution ramp is #E1E2DB, #B7B8A9, #898A7C, #74746A and #5D5E54. Lower oxygen and shorter proximity distances use the dark end. Foundation member dots use teal #247C80; proximity maps color member points by distance.

References

OpenStreetMap contributors. (2026). Landuse landfill polygon inventory [Data set]. Seven regional public ArcGIS mirrors, retrieved 6 October 2026. © OpenStreetMap contributors. Open Database License. [Source link](#)

OpenStreetMap Wiki contributors. (n.d.). Tag landuse landfill. Tag definition. Accessed 6 October 2026. [Source link](#)

Esri. (n.d.). Generate Near Table. ArcGIS Pro documentation. Reference for the geodesic proximity tools used in this project. Accessed 6 October 2026. [Source link](#)

Esri. (n.d.). Calculate Geometry Attributes. ArcGIS Pro documentation. Reference for geodesic area measurement. Accessed 6 October 2026. [Source link](#)

OpenStreetMap Landuse Areas for Africa. (n.d.). Landuse feature service layer 0 [Public regional mirror]. Queried for landuse=landfill on 6 October 2026. [Source link](#)

OpenStreetMap Landuse Areas for Asia. (n.d.). Landuse feature service layer 0 [Public regional mirror]. Queried for landuse=landfill on 6 October 2026. [Source link](#)

OpenStreetMap Landuse Areas for Australia and Oceania. (n.d.). Landuse feature service layer 0 [Public regional mirror]. Queried for landuse=landfill on 6 October 2026. [Source link](#)

OpenStreetMap Landuse Areas for Central America. (n.d.). Landuse feature service layer 0 [Public regional mirror]. Queried for landuse=landfill on 6 October 2026. [Source link](#)

OpenStreetMap Landuse Areas for Europe. (n.d.). Landuse feature service layer 0 [Public regional mirror]. Queried for landuse=landfill on 6 October 2026. [Source link](#)

OpenStreetMap Landuse Areas for North America. (n.d.). Landuse feature service layer 0 [Public regional mirror]. Queried for landuse=landfill on 6 October 2026. [Source link](#)

OpenStreetMap Landuse Areas for South America. (n.d.). Landuse feature service layer 0 [Public regional mirror]. Queried for landuse=landfill on 6 October 2026. [Source link](#)

LSAP Pollution Atlas. (2026). Internal source copies, executed analysis scripts and validation records [Project records]. LSAP_Pollution.aprx; FoundationCopies.gdb; PollutionAnalysis.gdb; foundation_manifest.json; raster_audit.json; freshwater_aggregate_QA.json; waste_QA.json; cleanup_audit.json; project_QA.json. Project root: Atlas Analyses/Pollution. Internal provenance is distinct from published source citations.