

LOGIC OF BRANCHING TIME

WAYS OF INDETERMINISM

Attila Molnár
Eötvös Loránd University



April 13, 2016

Trees

oooo

Ockhamist

ooo

Peircean

oooo

Leibnizian

oooooooo

Thomason

ooo


Łukasiewicz?

oooo

Tree of Time


INDETERMINIST FRAMES

Consider the tree on the right. Let p represent the sentence “There is a sea battle”. Suppose that w is today, and w_1 and w_2 are the two possible tomorrows. We have that $w \models \mathbf{F}p \wedge \mathbf{F}\neg p$, therefore,



INDETERMINIST FRAMES


Consider the tree on the right. Let p represent the sentence “There is a sea battle”. Suppose that w is today, and w_1 and w_2 are the two possible tomorrows. We have that $w \models \mathbf{F}p \wedge \mathbf{F}\neg p$, therefore,



- Since $\mathbf{F}p$ means “It **will** be true (tomorrow) that p ”, in w it is true that
 “It will be true (tomorrow) that there is sea battle and
 It will be true (tomorrow) that there is no sea battle”.

INDETERMINIST FRAMES

Consider the tree on the right. Let p represent the sentence “There is a sea battle”. Suppose that w is today, and w_1 and w_2 are the two possible tomorrows. We have that $w \models \mathbf{F}p \wedge \mathbf{F}\neg p$, therefore,




- Since $\mathbf{F}p$ means “It **will** be true (tomorrow) that p ”, in w it is true that

“It will be true (tomorrow) that there is sea battle and
It will be true (tomorrow) that there is no sea battle”.

- Since $\mathbf{F}p$ means “**In the future (tomorrow), it would be possible** that p ”, in w it is true that

“In the future (tomorrow), it would be possible that there is a sea battle and
In the future (tomorrow), it would be possible that there is no sea battle”.

INDETERMINIST FRAMES




Consider the tree on the right. Let p represent the sentence “There is a sea battle”. Suppose that w is today, and w_1 and w_2 are the two possible tomorrows. We have that $w \models \mathbf{F}p \wedge \mathbf{F}\neg p$, therefore,

- Since $\mathbf{F}p$ means “It **will** be true (tomorrow) that p ”, in w it is true that
 “It will be true (tomorrow) that there is sea battle and
 It will be true (tomorrow) that there is no see battle”.
- Since $\mathbf{F}p$ means “**In the future (tomorrow), it would be possible** that p ”, in w it is true that
 “In the future (tomorrow), it would be possible that there is a sea battle and
 In the future (tomorrow), it would be possible that there is no see battle”.

So trees are appropriate drawings but somehow not “will” is the appropriate word for \mathbf{F} . But then what is the meaning of “will” here?.


HISTORIES

Consider the trees as a **bundle** of linear frames, that are called **histories** in that context.




HISTORIES

Consider the trees as a **bundle** of linear frames, that are called **histories** in that context.




HISTORIES

Consider the trees as a **bundle** of linear frames, that are called **histories** in that context.




HISTORIES

Consider the trees as a **bundle** of linear frames, that are called **histories** in that context.




HISTORIES

Consider the trees as a **bundle** of linear frames, that are called **histories** in that context.



HISTORIES

Consider the trees as a **bundle** of linear frames, that are called **histories** in that context.



HISTORIES

Let $\mathfrak{F} = (W, <)$ be a tree.

DEFINITION: A **history** h is a maximally linear subset of W , i.e.,

- linear: $(\forall w, v \in h) \ w < v \vee w = v \vee w > v$.
- there is no proper linear extension of it:

$$(\forall h' \supseteq h) [h' \text{ is linear} \rightarrow h' \subseteq h.]$$

$h \overset{w}{\sim} h'$ will mean that histories h and h' share the same past until w . Since we are working with trees, this can be formalized simply by

$$h \overset{w}{\sim} h' \stackrel{\text{def}}{\iff} w \in h \cap h'$$

The set of all histories of a frame will be denoted by $H(\mathfrak{F})$:

$$H(\mathfrak{F}) \stackrel{\text{def}}{=} \{h \subseteq W : h \text{ is maximally linear}\}$$

HISTORIES

Let $\mathfrak{F} = (W, <)$ be a tree.

DEFINITION: A **history** h is a maximally linear subset of W , i.e.,

- linear: $(\forall w, v \in h) \ w < v \vee w = v \vee w > v$.
- there is no proper linear extension of it:

$$(\forall h' \supseteq h) [h' \text{ is linear} \rightarrow h' \subseteq h.]$$

$h \stackrel{w}{\sim} h'$ will mean that histories h and h' share the same past until w . Since we are working with trees, this can be formalized simply by

$$h \stackrel{w}{\sim} h' \stackrel{\text{def}}{\iff} w \in h \cap h'$$

The set of all histories of a frame will be denoted by $H(\mathfrak{F})$:

$$H(\mathfrak{F}) \stackrel{\text{def}}{=} \{h \subseteq W : h \text{ is maximally linear}\}$$

show that $\stackrel{w}{\sim}$ is an equivalence relation.

INDETERMINIST INTERPRETATIONS OF THE TENSE “WILL”.

Read $\mathbf{F}\varphi$ as “it will be the case that φ ”.

Is it plausible that

$$(\mathbf{F}\varphi \wedge \mathbf{F}\psi) \rightarrow [\mathbf{F}(\varphi \wedge \mathbf{F}\psi) \vee \mathbf{F}(\varphi \wedge \psi) \vee \mathbf{F}(\mathbf{F}\varphi \wedge \psi)] \quad ?$$

If φ will be true and ψ will be true, then at least one of the followings is true:

- φ will be true, and after that ψ will true.
- ψ will be true, and after that φ will true.
- φ and ψ will be true at the same time.

INDETERMINIST INTERPRETATIONS OF THE TENSE “WILL”.

Read $\mathbf{F}\varphi$ as “it will be the case that φ ”.

Is it plausible that

$$(\mathbf{F}\varphi \wedge \mathbf{F}\psi) \rightarrow [\mathbf{F}(\varphi \wedge \mathbf{F}\psi) \vee \mathbf{F}(\varphi \wedge \psi) \vee \mathbf{F}(\mathbf{F}\varphi \wedge \psi)] \quad ?$$

If φ will be true and ψ will be true, then at least one of the followings is true:

- φ will be true, and after that ψ will true.
- ψ will be true, and after that φ will true.
- φ and ψ will be true at the same time.

Yes: Ockhamist future

INDETERMINIST INTERPRETATIONS OF THE TENSE “WILL”.

Read $\mathbf{F}\varphi$ as “it will be the case that φ ”.

Is it plausible that

$$(\mathbf{F}\varphi \wedge \mathbf{F}\psi) \rightarrow [\mathbf{F}(\varphi \wedge \mathbf{F}\psi) \vee \mathbf{F}(\varphi \wedge \psi) \vee \mathbf{F}(\mathbf{F}\varphi \wedge \psi)] \quad ?$$

If φ will be true and ψ will be true, then at least one of the followings is true:

- φ will be true, and after that ψ will true.
- ψ will be true, and after that φ will true.
- φ and ψ will be true at the same time.

Yes: Ockhamist future

No: Peircean future

All the other options are variations of these two.

Trees
oooo

Ockhamist
ooo

Peircean
oooo

Leibnizian
oooooooo

Thomason
ooo

Łukasiewicz?
oooo

Ockhamist future

Ockhamist FUTURE

“Ockhamism” [...] holds that it is meaningless to ask about the truth value of “a will happen” at w without further specifications: the problem is correctly expressed only if, in addition to w , one of its possible futures is specified. “Will happen” has to be understood as “will happen in the specified future of w ”.

ZANARDO 1996 (about PRIOR 1967)

So the history is always a tacit parameter

OCKHAMIST FUTURE

“Ockhamism” [...] holds that it is meaningless to ask about the truth value of “a will happen” at w without further specifications: the problem is correctly expressed only if, in addition to w , one of its possible futures is specified. “Will happen” has to be understood as “will happen in the specified future of w ”.

ZANARDO 1996 (about PRIOR 1967)

So the history is always a tacit parameter

Indeterminism comes into the picture when we change the “specified possible future” (history) with an operator \Diamond .

Ockhamist FUTURE

“Ockhamism” [...] holds that it is meaningless to ask about the truth value of “a will happen” at w without further specifications: the problem is correctly expressed only if, in addition to w , one of its possible futures is specified. “Will happen” has to be understood as “will happen in the specified future of w ”.

ZANARDO 1996 (about PRIOR 1967)


So the history is always a tacit parameter

Indeterminism comes into the picture when we change the “specified possible future” (history) with an operator \Diamond .

Let $\mathfrak{M} = (W, <, V)$ be a tree model.

| | | |
|--|--|--|
| $\mathfrak{M}, h, w \models^{\circ} p$ | $\stackrel{\text{def}}{\Leftrightarrow}$ | $w \in V(p)$ |
| $\mathfrak{M}, h, w \models^{\circ} \neg \varphi$ | $\stackrel{\text{def}}{\Leftrightarrow}$ | it is not true that $\mathfrak{M}, h, w \models^{\circ} \varphi$ |
| $\mathfrak{M}, h, w \models^{\circ} \varphi \wedge \psi$ | $\stackrel{\text{def}}{\Leftrightarrow}$ | $\mathfrak{M}, h, w \models^{\circ} \varphi$ and $\mathfrak{M}, h, w \models^{\circ} \psi$ |
| $\mathfrak{M}, h, w \models^{\circ} \mathbf{P}\varphi$ | $\stackrel{\text{def}}{\Leftrightarrow}$ | $\exists v (v < w \wedge \mathfrak{M}, h, v \models^{\circ} \varphi)$ |
| $\mathfrak{M}, h, w \models^{\circ} \mathbf{F}\varphi$ | $\stackrel{\text{def}}{\Leftrightarrow}$ | $(\exists v \in \textcolor{red}{h}) (w < v \wedge \mathfrak{M}, h, v \models^{\circ} \varphi)$ |
| $\mathfrak{M}, h, w \models^{\circ} \Diamond \varphi$ | $\stackrel{\text{def}}{\Leftrightarrow}$ | $(\exists h' \overset{w}{\sim} h) \mathfrak{M}, h', w \models^{\circ} \varphi$ |

TRAINING



Ockhamist “WILL”

$$(\mathbf{F}\varphi \wedge \mathbf{F}\psi) \rightarrow [\mathbf{F}(\varphi \wedge \mathbf{F}\psi) \vee \mathbf{F}(\varphi \wedge \psi) \vee \mathbf{F}(\mathbf{F}\varphi \wedge \psi)] \quad ?$$

is valid because outside of φ and ψ there are no \Diamond -s, so once the meaning of φ and ψ is given, the meaning of the formula above is evaluated on a given history, which is a linear order of moments.

Trees
oooo

Ockhamist
ooo

Peircean
oooo

Leibnizian
oooooooo

Thomason
ooo

Łukasiewicz?
oooo

Peircean future

PEIRCEAN FUTURE

From the Peircean point of view, [...] “ φ will happen” is short for “ φ will happen, no matter what possible future of w is considered”, which is true just in case every possible future of w contains a moment at which φ is true.

ZANARDO 1996 (about PRIOR 1967)

So even if we use the histories to give the semantics of **F**, we do not relativize the truth of formulas to certain histories. Truth of a temporal statement is history-independent.

A set of worlds X **bars** w iff every history containing w goes through X :




$$X \text{ bars } w \stackrel{\text{def}}{\iff} (\forall h \in H(\mathfrak{F}))(w \in h \rightarrow h \cap X \neq \emptyset)$$


Let $\mathfrak{M} = (W, <, V)$ be a tree model.

$$\begin{array}{ll} \mathfrak{M}, w \models p & \stackrel{\text{def}}{\iff} w \in V(p) \\ \mathfrak{M}, w \models \neg\varphi & \stackrel{\text{def}}{\iff} \text{it is not true that } \mathfrak{M}, w \models \varphi \\ \mathfrak{M}, w \models \varphi \wedge \psi & \stackrel{\text{def}}{\iff} \mathfrak{M}, w \models \varphi \text{ and } \mathfrak{M}, w \models \psi \\ \mathfrak{M}, w \models \mathbf{P}\varphi & \stackrel{\text{def}}{\iff} \exists v (v < w \wedge \mathfrak{M}, v \models \varphi) \\ \mathfrak{M}, w \models \mathbf{F}\varphi & \stackrel{\text{def}}{\iff} \llbracket \varphi \rrbracket_{\mathbf{P}}^{\mathfrak{M}} \text{ bars } w \end{array}$$

where $\llbracket \varphi \rrbracket_{\mathbf{P}}^{\mathfrak{M}} \stackrel{\text{def}}{=} \{w : \mathfrak{M}, w \models \varphi\}$, i.e., the set of those worlds in which φ is true.




TRAINING



PEIRCEAN “WILL”

$$(\mathbf{F}\varphi \wedge \mathbf{F}\psi) \rightarrow [\mathbf{F}(\varphi \wedge \mathbf{F}\psi) \vee \mathbf{F}(\varphi \wedge \psi) \vee \mathbf{F}(\mathbf{F}\varphi \wedge \psi)] \quad ?$$


is invalid; let \mathfrak{M} be the following “twin lines”-model:



PEIRCEAN “WILL”

$$(\mathbf{F}\varphi \wedge \mathbf{F}\psi) \rightarrow [\mathbf{F}(\varphi \wedge \mathbf{F}\psi) \vee \mathbf{F}(\varphi \wedge \psi) \vee \mathbf{F}(\mathbf{F}\varphi \wedge \psi)] \quad ?$$

is invalid; let \mathfrak{M} be the following “twin lines”-model:




$V(p)$ and $V(q)$ bars w , so $\mathfrak{M}, w \models^{\text{P}} \mathbf{F}p \wedge \mathbf{F}q$

PEIRCEAN “WILL”

$$(\mathbf{F}\varphi \wedge \mathbf{F}\psi) \rightarrow [\mathbf{F}(\varphi \wedge \mathbf{F}\psi) \vee \mathbf{F}(\varphi \wedge \psi) \vee \mathbf{F}(\mathbf{F}\varphi \wedge \psi)] \quad ?$$

is invalid; let \mathfrak{M} be the following “twin lines”-model:




$V(p)$ and $V(q)$ bars w , so $\mathfrak{M}, w \models^{\text{P}} \mathbf{F}p \wedge \mathbf{F}q$

$V(p) \cap V(q) = \emptyset$, and \emptyset bars nothing, so $\mathfrak{M}, w \not\models^{\text{P}} \mathbf{F}(p \wedge q)$

PEIRCEAN “WILL”

$$(\mathbf{F}\varphi \wedge \mathbf{F}\psi) \rightarrow [\mathbf{F}(\varphi \wedge \mathbf{F}\psi) \vee \mathbf{F}(\varphi \wedge \psi) \vee \mathbf{F}(\mathbf{F}\varphi \wedge \psi)] \quad ?$$

is invalid; let \mathfrak{M} be the following “twin lines”-model:



$V(p)$ and $V(q)$ bars w , so $\mathfrak{M}, w \models^P \mathbf{F}p \wedge \mathbf{F}q$


$V(p) \cap V(q) = \emptyset$, and \emptyset bars nothing, so $\mathfrak{M}, w \not\models^P \mathbf{F}(p \wedge q)$

$\llbracket \mathbf{F}p \rrbracket_P^{\mathfrak{M}} = \{w\} \cup \{v_i : i \in \mathbb{N}\}$ $\llbracket \mathbf{F}q \rrbracket_P^{\mathfrak{M}} = \{w\} \cup \{u_i : i \in \mathbb{N}\}$

PEIRCEAN “WILL”

$$(\mathbf{F}\varphi \wedge \mathbf{F}\psi) \rightarrow [\mathbf{F}(\varphi \wedge \mathbf{F}\psi) \vee \mathbf{F}(\varphi \wedge \psi) \vee \mathbf{F}(\mathbf{F}\varphi \wedge \psi)] \quad ?$$

is invalid; let \mathfrak{M} be the following “twin lines”-model:



$V(p)$ and $V(q)$ bars w , so $\mathfrak{M}, w \models^p \mathbf{F}p \wedge \mathbf{F}q$

$V(p) \cap V(q) = \emptyset$, and \emptyset bars nothing, so $\mathfrak{M}, w \not\models^p \mathbf{F}(p \wedge q)$


$\llbracket \mathbf{F}p \rrbracket_P^{\mathfrak{M}} = \{w\} \cup \{v_i : i \in \mathbb{N}\}$ $\llbracket \mathbf{F}q \rrbracket_P^{\mathfrak{M}} = \{w\} \cup \{u_i : i \in \mathbb{N}\}$

$\llbracket \mathbf{F}p \rrbracket_P^{\mathfrak{M}} \cap V(q) = \{v_1\}$ $\llbracket \mathbf{F}q \rrbracket_P^{\mathfrak{M}} \cap V(p) = \{u_1\}$

PEIRCEAN “WILL”

$$(\mathbf{F}\varphi \wedge \mathbf{F}\psi) \rightarrow [\mathbf{F}(\varphi \wedge \mathbf{F}\psi) \vee \mathbf{F}(\varphi \wedge \psi) \vee \mathbf{F}(\mathbf{F}\varphi \wedge \psi)] \quad ?$$

is invalid; let \mathfrak{M} be the following “twin lines”-model:



$V(p)$ and $V(q)$ bars w , so $\mathfrak{M}, w \models^p \mathbf{F}p \wedge \mathbf{F}q$

$V(p) \cap V(q) = \emptyset$, and \emptyset bars nothing, so $\mathfrak{M}, w \not\models^p \mathbf{F}(p \wedge q)$

$\llbracket \mathbf{F}p \rrbracket_P^{\mathfrak{M}} = \{w\} \cup \{v_i : i \in \mathbb{N}\}$ $\llbracket \mathbf{F}q \rrbracket_P^{\mathfrak{M}} = \{w\} \cup \{u_i : i \in \mathbb{N}\}$

$\llbracket \mathbf{F}p \rrbracket_P^{\mathfrak{M}} \cap V(q) = \{v_1\}$ $\llbracket \mathbf{F}q \rrbracket_P^{\mathfrak{M}} \cap V(p) = \{u_1\}$

But neither of these bars w , so $\mathfrak{M}, w \not\models^p \mathbf{F}(\mathbf{F}p \wedge q)$ and $\mathfrak{M}, w \not\models^p \mathbf{F}(p \wedge \mathbf{F}q)$

“ALWAYS GOING TO BE ...”

Let p represent the sentence “The 3rd World War is on.”

How should we formalize the statement “There won’t be a 3rd World War.”?

“ALWAYS GOING TO BE ...”

Let p represent the sentence “The 3rd World War is on.”

How should we formalize the statement “There won’t be a 3rd World War.”?

- $\neg \mathbf{F}p$

“ALWAYS GOING TO BE ...”

Let p represent the sentence “The 3rd World War is on.”

How should we formalize the statement “There won’t be a 3rd World War.”?

- $\neg \mathbf{F}p$
- $\mathbf{F}\neg p$

“ALWAYS GOING TO BE ...”

Let p represent the sentence “The 3rd World War is on.”

How should we formalize the statement “There won’t be a 3rd World War.”?

- $\neg Fp$
 - $F\neg p$
-
- $\neg Fp$ says that $V(p)$ does not bar ‘us’. So there is an ‘escape’ history in which the 3rd World War won’t break out

“ALWAYS GOING TO BE ...”

Let p represent the sentence “The 3rd World War is on.”

How should we formalize the statement “There won’t be a 3rd World War.”?

- $\neg \mathbf{F}p$
 - $\mathbf{F}\neg p$
-
- $\neg \mathbf{F}p$ says that $V(p)$ does not bar ‘us’. So there is an ‘escape’ history in which the 3rd World War won’t break out
 - $\mathbf{F}\neg p$ says that $W - V(p)$ bar ‘us’. So no matter what happens, there will be moments in the future when there is no 3rd World War.

“ALWAYS GOING TO BE ...”

Let p represent the sentence “The 3rd World War is on.”

How should we formalize the statement “There won’t be a 3rd World War.”?

- $\neg Fp$
- $F\neg p$

-
- $\neg Fp$ says that $V(p)$ does not bar ‘us’. So there is an ‘escape’ history in which the 3rd World War won’t break out
 - $F\neg p$ says that $W - V(p)$ bar ‘us’. So no matter what happens, there will be moments in the future when there is no 3rd World War.

None of the above is correct, because the first is speaking about **only some** ‘escape’-history, and the second talks about **only some** moments in the future in which there is no 3rd World war. The reason of course is that we interpret F with a $\forall\exists$ -way, and no matter how we negate it, the mixed nature of it will survive.

If we want to formalize the ‘won’t-s, and “always going to be”-s, we need the old history-independent strong future operator for that purpose:

$$\mathfrak{M}, w \models^P \mathbf{G}\varphi \stackrel{\text{def}}{\iff} \forall v (w < v \wedge \mathfrak{M}, v \models^P \varphi)$$

Trees
oooo

Ockhamist
ooo

Peircean
oooo

Leibnizian
oooooooo

Thomason
ooo


Łukasiewicz?
oooo

Leibnizian

PARALLEL HISTORIES

KAMP FRAMES INSTEAD OF TREES


Consider the tree on the right. Let p represent the sentence “There is a sea battle”. Suppose that w and w' are today, and w_1 and w_2 are the two possible tomorrows.



PARALLEL HISTORIES

KAMP FRAMES INSTEAD OF TREES


Consider the tree on the right. Let p represent the sentence “There is a sea battle”. Suppose that w and w' are today, and w_1 and w_2 are the two possible tomorrows.




PARALLEL HISTORIES

KAMP FRAMES INSTEAD OF TREES


Consider the tree on the right. Let p represent the sentence “There is a sea battle”. Suppose that w and w' are today, and w_1 and w_2 are the two possible tomorrows.



Standard (tree) frame
($W, <$)



Kamp-frame
($W, <, \equiv$)



KAMP-FRAMES

A Kamp-frame is a triplet $(W, <, \equiv)$ where

- $<$ is irreflexive, transitive and non-branching:

- $w \not< w$
- $(w < v \wedge v < u) \rightarrow w < u$
- $(w < v \wedge w < u) \rightarrow (v < u \vee v = u \vee v > u)$
- $(w > v \wedge w > u) \rightarrow (v < u \vee v = u \vee v > u)$

- \equiv is reflexive, transitive and symmetric:

- $w \equiv w$
- $(w \equiv v \wedge v \equiv u) \rightarrow w \equiv u$
- $w \equiv v \rightarrow v \equiv w$

- $x \equiv y \rightarrow x \not< y$

- $(w \equiv v \wedge w' < w) \rightarrow (\exists v' < v) w' \equiv v'$

- $(\forall w, v)(\exists w' < w)(\exists v' < v) w \equiv v$

- $(\forall w, v)(w \equiv v \wedge w \neq v)(\exists w' > w)(\forall v' > v) w' \neq v'$

“sharing the same past”

class common root

maximality of histories

class irreflexivity

“sharing the same past”

class common root

maximality of histories

KAMP-FRAMES

A Kamp-frame is a triplet $(W, <, \equiv)$ where

- $<$ is irreflexive, transitive and non-branching:

- $w \not< w$
- $(w < v \wedge v < u) \rightarrow w < u$
- $(w < v \wedge w < u) \rightarrow (v < u \vee v = u \vee v > u)$
- $(w > v \wedge w > u) \rightarrow (v < u \vee v = u \vee v > u)$

- \equiv is reflexive, transitive and symmetric:

- $w \equiv w$
- $(w \equiv v \wedge v \equiv u) \rightarrow w \equiv u$
- $w \equiv v \rightarrow v \equiv w$

- $x \equiv y \rightarrow x \not< y$




- $(w \equiv v \wedge w' < w) \rightarrow (\exists v' < v) w' \equiv v'$

- $(\forall w, v)(\exists w' < w)(\exists v' < v) w \equiv v$

- $(\forall w, v)(w \equiv v \wedge w \neq v)(\exists w' > w)(\forall v' > v) w' \neq v'$

Show that

- class irreflexivity implies irreflexivity
- maximality of histories implies class irreflexivity



class irreflexivity

"sharing the same past"

class common root

maximality of histories

KAMP-MODELS

Let $\mathfrak{K} = (W, <, \equiv)$ be a Kamp-frame. A Kamp-valuation is a $V : \text{At} \rightarrow \wp W$ for which the following additional property holds:

$$w \in V(p) \Rightarrow (\forall v \equiv w) v \in V(p) \quad \text{for all } p \in \text{At}$$

a Kamp-frame $\mathfrak{K} = (W, <, \equiv)$ together with such a valuation V is a Kamp-model $\mathfrak{M}_K = (\mathfrak{K}, V)$.

$$\begin{array}{ll}
 \mathfrak{M}_K, w \models^{\text{K}} p & \stackrel{\text{def}}{\iff} w \in V(p) \\
 \mathfrak{M}_K, w \models^{\text{K}} \neg \varphi & \stackrel{\text{def}}{\iff} \text{it is not true that } \mathfrak{M}_K, w \models^{\text{K}} \varphi \\
 \mathfrak{M}_K, w \models^{\text{K}} \varphi \wedge \psi & \stackrel{\text{def}}{\iff} \mathfrak{M}_K, w \models^{\text{K}} \varphi \text{ and } \mathfrak{M}_K, w \models^{\text{K}} \psi \\
 \mathfrak{M}_K, w \models^{\text{K}} \mathbf{P}\varphi & \stackrel{\text{def}}{\iff} (\exists v < w) \mathfrak{M}_K, v \models^{\text{K}} \varphi \\
 \mathfrak{M}_K, w \models^{\text{K}} \mathbf{F}\varphi & \stackrel{\text{def}}{\iff} (\exists v > w) \mathfrak{M}_K, v \models^{\text{K}} \varphi \\
 \mathfrak{M}_K, w \models^{\text{K}} \Diamond \varphi & \stackrel{\text{def}}{\iff} (\exists v \equiv w) \mathfrak{M}_K, v \models^{\text{K}} \varphi
 \end{array}$$

KAMP-FRAMES

THEOREM: Every \models^{K} -valid formula is \models^{O} -valid.

KAMP-FRAMES

THEOREM: Every \models^{K} -valid formula is \models^{O} -valid.

PROPOSITION: There are \models^{O} -valid formulas that are not \models^{K} -valid.

KAMP-FRAMES

THEOREM: Every \models^K -valid formula is \models^O -valid.

PROPOSITION: There are \models^O -valid formulas that are not \models^K -valid.

This was expected: \Diamond in \models^K quantifies over worlds,
but \models^O quantify over **sets of** worlds.

\Diamond is interpreted with a 1st order quantification in \models^K
 \Diamond is interpreted with a 2nd order quantification in \models^O

KAMP-FRAMES

THEOREM: Every \models^K -valid formula is \models^O -valid.

PROPOSITION: There are \models^O -valid formulas that are not \models^K -valid.

This was expected: \Diamond in \models^K quantifies over worlds,
but \models^O quantify over **sets of** worlds.

\Diamond is interpreted with a 1st order quantification in \models^K
 \Diamond is interpreted with a 2nd order quantification in \models^O

It is not impossible, however, to replace a 2nd order quantification with a 1st order one, but to do so, you have to be able to name all the sets with an individuum uniquely.

KAMP-FRAMES

THEOREM: Every \models^K -valid formula is \models^O -valid.

PROPOSITION: There are \models^O -valid formulas that are not \models^K -valid.

This was expected: \Diamond in \models^K quantifies over worlds,
but \models^O quantify over **sets of** worlds.

\Diamond is interpreted with a 1st order quantification in \models^K
 \Diamond is interpreted with a 2nd order quantification in \models^O

It is not impossible, however, to replace a 2nd order quantification with a 1st order one, but to do so, you have to be able to name all the sets with an individuum uniquely.

Show that every history **in a finite tree** is uniquely identifiable with a world.

COUNTEREXAMPLE: INFINITE BINARY TREES

$W \stackrel{\text{def}}{=} \{w : w \text{ is a route to a point}\}$
 $= \{\langle w_1, \dots, w_n \rangle : n \in \omega, (\forall i \leq n) w_i \in \{U, R\}\}$

$w \sqsubseteq v \stackrel{\text{def}}{\iff} v$ is a continuation of w , i.e.,
 iff $(\forall i \leq n) w_i = v_i$ where n is the length of w .

Note that histories correspond to infinite routes!

Also note we can not name the histories by worlds (as was the case in the finite cases)! There are (infinitely) many infinite continuations of finite routes.




A set of histories $B \subseteq H(\mathfrak{F})$ is called a **bundle** iff

$$\bigcup B = W,$$

that is, for every $w \in W$ there is a history $h \in B$ s.t. $w \in h$.

We can find a proper bundle, which is in fact can be named by worlds:

$$\{h \in H(\mathfrak{F}) : \exists w (\forall v > w) v = \langle w, U, \dots, U \rangle\}$$



BUNDLED TREES

DEFINITION: A bundled tree is a triplet $\mathfrak{F}_B = (W, <, B)$ where $(W, <)$ is a tree and $B \subseteq H(W, <)$ is a bundle. A bundled model is a quadruple (\mathfrak{F}_B, V) where \mathfrak{F}_B is a bundled frame and $V : \text{At} \rightarrow \wp W$ is a valuation.

| | | |
|--|--|--|
| $\mathfrak{M}, h, w \models^B p$ | $\stackrel{\text{def}}{\Leftrightarrow}$ | $w \in V(p)$ |
| $\mathfrak{M}, h, w \models^B \neg \varphi$ | $\stackrel{\text{def}}{\Leftrightarrow}$ | it is not true that $\mathfrak{M}, h, w \models^B \varphi$ |
| $\mathfrak{M}, h, w \models^B \varphi \wedge \psi$ | $\stackrel{\text{def}}{\Leftrightarrow}$ | $\mathfrak{M}, h, w \models^B \varphi$ and $\mathfrak{M}, h, w \models^B \psi$ |
| $\mathfrak{M}, h, w \models^B \mathbf{P}\varphi$ | $\stackrel{\text{def}}{\Leftrightarrow}$ | $\exists v (v < w \wedge \mathfrak{M}, h, v \models^B \varphi)$ |
| $\mathfrak{M}, h, w \models^B \mathbf{F}\varphi$ | $\stackrel{\text{def}}{\Leftrightarrow}$ | $(\exists v \in h) (w < v \wedge \mathfrak{M}, h, v \models^B \varphi)$ |
| $\mathfrak{M}, h, w \models^B \Diamond \varphi$ | $\stackrel{\text{def}}{\Leftrightarrow}$ | $(\exists h' \sim^w h) (h \in B \wedge \mathfrak{M}, h', w \models^B \varphi)$ |

PROPOSITION: \models^K -validity corresponds to \models^B -validity.

QUOTE: “Belnap et al. have argued that it is implausible to assume that there could be some property which could »justify treating some maximal chains as real possibilities and others as not« (Belnap et al. 2001, p. 205)”

Hirokazu Nishimura (1979) – Stanford Enc.

DEFENSE OF BUNDLED TREES

Suppose we have discrete models (we are thinking in days instead of moments). Are these two sentences contradictory?

- “Inevitably, if today there is life on earth, then either this is the last day (of life on earth), or the last day will come.”
- “At any possible day on which there is life on earth, it is possible that there will be life on earth the following day.”

Hirokazu Nishimura (1979)– Stanford Enc.

Let p be “there is life on earth”. Let h be some history (it does not matter actually)

- $\mathfrak{M}, h, w \models \Box(p \rightarrow \underline{\mathbf{F}}\mathbf{G}\neg p)$
- For all $v \in W$: $\mathfrak{M}, h, v \models p \rightarrow \Diamond \mathbf{F}p$

DEFENSE OF BUNDLED TREES

finite zigzags: $\langle R, U, R, U, \dots, R, U \rangle$

infinite columns: $\langle U, U, U, U, \dots \rangle$

W : finite routes

R : continuation

$$B = \left\{ w \oplus v : \begin{array}{l} w \text{ fin. route,} \\ v \text{ inf. column} \end{array} \right\}$$

$V(p)$: the finite zigzags

life on Earth \leftrightarrow zigzagging


$\Box(p \rightarrow \underline{\mathbf{F}}\mathbf{G}\neg p) \leftrightarrow$ in every history,
sooner or later life (zigzags) will permanently end

$p \rightarrow \Diamond \mathbf{F}p \leftrightarrow$ one more day of life
(zigzag) is always possible

The Ockhamist will find the infinite
zigzag, which refutes

$$(p \rightarrow \underline{\mathbf{F}}\mathbf{G}\neg p)$$

But the bundled treehugger won't.



Trees
oooo

Ockhamist
ooo

Peircean
oooo

Leibnizian
oooooooo


Thomason
ooo

Łukasiewicz?
oooo

Thomason


TRUTH VALUE GAPS FOR THE FUTURE CONTINGENTS

Consider the tree on the right. Let p represent the sentence "There is a sea battle". Suppose that w is today, and w_1 and w_2 are the two possible tomorrows.



TRUTH VALUE GAPS FOR THE FUTURE CONTINGENTS


Consider the tree on the right. Let p represent the sentence “There is a sea battle”. Suppose that w is today, and w_1 and w_2 are the two possible tomorrows.



- $\mathbf{F}p$ is true iff $V(p)$ bars the moment of utterance. (standard Peircean)

TRUTH VALUE GAPS FOR THE FUTURE CONTINGENTS


Consider the tree on the right. Let p represent the sentence “There is a sea battle”. Suppose that w is today, and w_1 and w_2 are the two possible tomorrows.



- $\mathbf{F}p$ is true iff $V(p)$ bars the moment of utterance. (standard Peircean)
- $\mathbf{F}p$ is false iff $V(\neg p)$ bars the moment of utterance.

TRUTH VALUE GAPS FOR THE FUTURE CONTINGENTS

Consider the tree on the right. Let p represent the sentence “There is a sea battle”. Suppose that w is today, and w_1 and w_2 are the two possible tomorrows.




- $\mathbf{F}p$ is true iff $V(p)$ bars the moment of utterance. (standard Peircean)
- $\mathbf{F}p$ is false iff $V(\neg p)$ bars the moment of utterance.

(Not Peircean at all!!! Falsehood of $\mathbf{F}p$ correspond to the Peircean truth of $\mathbf{F}\neg p$, not to $\neg\mathbf{F}p$!! So from a Peircean point of view, falsehood is some kind of ‘inner negation’.)

TRUTH VALUE GAPS FOR THE FUTURE CONTINGENTS

Consider the tree on the right. Let p represent the sentence “There is a sea battle”. Suppose that w is today, and w_1 and w_2 are the two possible tomorrows.




- $\mathbf{F}p$ is true iff $V(p)$ bars the moment of utterance. (standard Peircean)
- $\mathbf{F}p$ is false iff $V(\neg p)$ bars the moment of utterance.

(Not Peircean at all!!! Falsehood of $\mathbf{F}p$ correspond to the Peircean truth of $\mathbf{F}\neg p$, not to $\neg\mathbf{F}p$!! So from a Peircean point of view, falsehood is some kind of ‘inner negation’.)

Partial semantics: What if neither of them bar the moment of utterance?

TRUTH VALUE GAPS FOR THE FUTURE CONTINGENTS

Consider the tree on the right. Let p represent the sentence “There is a sea battle”. Suppose that w is today, and w_1 and w_2 are the two possible tomorrows.



- $\mathbf{F}p$ is true iff $V(p)$ bars the moment of utterance. (standard Peircean)
- $\mathbf{F}p$ is false iff $V(\neg p)$ bars the moment of utterance.


(Not Peircean at all!!! Falsehood of $\mathbf{F}p$ correspond to the Peircean truth of $\mathbf{F}\neg p$, not to $\neg\mathbf{F}p$!! So from a Peircean point of view, falsehood is some kind of ‘inner negation’.)

Partial semantics: What if neither of them bar the moment of utterance?

Then $\mathbf{F}p$ is **undefined** in the moment of utterance.

TRUTH VALUE GAPS FOR THE FUTURE CONTINGENTS

Consider the tree on the right. Let p represent the sentence “There is a sea battle”. Suppose that w is today, and w_1 and w_2 are the two possible tomorrows.



- $\mathbf{F}p$ is true iff $V(p)$ bars the moment of utterance. (standard Peircean)
- $\mathbf{F}p$ is false iff $V(\neg p)$ bars the moment of utterance.

(Not Peircean at all!!! Falsehood of $\mathbf{F}p$ correspond to the Peircean truth of $\mathbf{F}\neg p$, not to $\neg\mathbf{F}p$!! So from a Peircean point of view, falsehood is some kind of ‘inner negation’.)

Partial semantics: What if neither of them bar the moment of utterance?

Then $\mathbf{F}p$ is **undefined** in the moment of utterance. Because these statements’ truth values are not **settled** yet. Tomorrow they will be settled, but now they are not.

Imre Ruzsa, the founder of this department, is a (the?) champion of partial modal semantics.

SUPERVALUATION

We will consider the word **undefined** as a 3rd truth value.

Ruzsa was a hardcore classical logician as well: He did not acknowledge any formal system as a logic if it did not fulfilled the Law of (Non-)Contradiction and the Law of Excluded Middle. But how did he give an account about these 3-valued thing as a logic? These classical laws are fulfilled – on those formulas where the interpretation is defined.

SUPERVALUATION

We will consider the word **undefined** as a 3rd truth value.

NOTATIONAL PROBLEM: we can not represent 3 value with our \models sign, so we switch to the intension-notation:

$$\mathfrak{M}, w \models \varphi \iff \llbracket \varphi \rrbracket_w^{\mathfrak{M}} = \text{true}$$

CHEAT: Instead of starting everything from the beginning, we can define truth via Ockhamist truth!

... Thomasonian truth **supervenes** on Ockhamist truth. . .

$$\llbracket \varphi \rrbracket_w^{\mathfrak{M}} \stackrel{\text{def}}{=} \begin{cases} \text{T} & \text{if } (\forall h \ni w) \mathfrak{M}, h, w \models^{\circ} \varphi \\ \text{F} & \text{if } (\forall h \ni w) \mathfrak{M}, h, w \models^{\circ} \neg \varphi \\ \text{U} & \text{otherwise} \end{cases}$$

Ruzsa was a hardcore classical logician as well: He did not acknowledge any formal system as a logic if it did not fulfilled the Law of (Non-)Contradiction and the Law of Excluded Middle. But how did he give an account about these 3-valued thing as a logic? These classical laws are fulfilled – on those formulas where the interpretation is defined.

COMPOSITIONALITY

Our cheat had a cost: the compositionality.

DEFINITION: A meaning function $\llbracket \cdot \rrbracket_w$ is compositional iff the meaning $\llbracket f(\varphi, \psi, \dots) \rrbracket_w$ of a complex expression $f(\varphi, \psi, \dots)$ is determined by the meanings $\llbracket \varphi \rrbracket_w, \llbracket \psi \rrbracket_w, \dots$ of the constituents φ, ψ, \dots .

That is a very general definition which is common
in the Andr  ka–N  meti–Madar  sz–Sain school

COMPOSITIONALITY

Our cheat had a cost: the compositionality.

DEFINITION: A meaning function $\llbracket \cdot \rrbracket_w$ is compositional iff the meaning $\llbracket f(\varphi, \psi, \dots) \rrbracket_w$ of a complex expression $f(\varphi, \psi, \dots)$ is determined by the meanings $\llbracket \varphi \rrbracket_w, \llbracket \psi \rrbracket_w, \dots$ of the constituents φ, ψ, \dots

That is a very general definition which is common in the Andr  ka–N  meti–Madar  sz–Sain school


Remember to the sea battle model:

$$\llbracket \mathbf{F}p \rrbracket_w = \mathbf{U}, \quad \llbracket \mathbf{F}\neg p \rrbracket_w = \mathbf{U}, \quad \llbracket \mathbf{F}p \vee \mathbf{F}\neg p \rrbracket_w = \mathbf{T}$$

But

$$\llbracket \mathbf{F}p \rrbracket_w = \mathbf{U}, \quad \llbracket \mathbf{F}p \rrbracket_w = \mathbf{U}, \quad \llbracket \mathbf{F}p \vee \mathbf{F}p \rrbracket_w = \mathbf{U}$$

In non-compositional logics you have to “look into” the formula



Trees
oooo

Ockhamist
ooo

Peircean
oooo

Leibnizian
oooooooo

Thomason
ooo

Łukasiewicz?
oooo

Łukasiewicz?

MODELLING THE UNDEFINED – KLEENE'S LOGIC(S)

ATOMS AND NEGATION

Every atom is either true or false, given by the model's valuation V .

| | | |
|-----------|---|---------------|
| | | $\neg\varphi$ |
| φ | F | T |
| | U | U |
| | T | F |

MODELLING THE UNDEFINED – KLEENE'S LOGIC(S)

CONJUNCTION AND DISJUNCTION

| $\varphi \wedge \psi$ | | ψ | | |
|-----------------------|---|--------|---|---|
| | | F | U | T |
| φ | F | F | ? | F |
| | U | ? | U | U |
| | T | F | U | T |

| $\varphi \vee \psi$ | | ψ | | |
|---------------------|---|--------|---|---|
| | | F | U | T |
| φ | F | F | U | T |
| | U | U | U | ? |
| | T | T | ? | T |

MODELLING THE UNDEFINED – KLEENE'S LOGIC(S)

CONJUNCTION AND DISJUNCTION

| $\varphi \wedge \psi$ | | ψ | | |
|-----------------------|---|--------|---|---|
| | | F | U | T |
| φ | F | F | ? | F |
| | U | ? | U | U |
| | T | F | U | T |

| $\varphi \vee \psi$ | | ψ | | |
|---------------------|---|--------|---|---|
| | | F | U | T |
| φ | F | F | U | T |
| | U | U | U | ? |
| | T | T | ? | T |

WEAK CONJUNCTIVES

| $\varphi \wedge \psi$ | | ψ | | |
|-----------------------|---|--------|---|---|
| | | F | U | T |
| φ | F | F | U | F |
| | U | U | U | U |
| | T | F | U | T |

| $\varphi \vee \psi$ | | ψ | | |
|---------------------|---|--------|---|---|
| | | F | U | T |
| φ | F | F | U | T |
| | U | U | U | U |
| | T | T | U | T |

MODELLING THE UNDEFINED – KLEENE'S LOGIC(S)

CONJUNCTION AND DISJUNCTION

| $\varphi \wedge \psi$ | | ψ | | |
|-----------------------|---|--------|---|---|
| | | F | U | T |
| φ | F | F | ? | F |
| | U | ? | U | U |
| | T | F | U | T |

| $\varphi \vee \psi$ | | ψ | | |
|---------------------|---|--------|---|---|
| | | F | U | T |
| φ | F | F | U | T |
| | U | U | U | ? |
| | T | T | ? | T |

WEAK CONJUNCTIVES

| $\varphi \wedge \psi$ | | ψ | | |
|-----------------------|---|--------|---|---|
| | | F | U | T |
| φ | F | F | U | F |
| | U | U | U | U |
| | T | F | U | T |

| $\varphi \vee \psi$ | | ψ | | |
|---------------------|---|--------|---|---|
| | | F | U | T |
| φ | F | F | U | T |
| | U | U | U | U |
| | T | T | U | T |

STRONG CONJUNCTIVES

| $\varphi \wedge \psi$ | | ψ | | |
|-----------------------|---|--------|---|---|
| | | F | U | T |
| φ | F | F | F | F |
| | U | F | U | U |
| | T | F | U | T |

| $\varphi \vee \psi$ | | ψ | | |
|---------------------|---|--------|---|---|
| | | F | U | T |
| φ | F | F | U | T |
| | U | U | U | T |
| | T | T | T | T |

MODELLING THE UNDEFINED – KLEENE'S LOGIC(S)

VARIATIONS FOR IMPLICATION

| weak $\neg\varphi \vee \psi$ | | ψ | | |
|---------------------------------|---|--------|---|---|
| | | F | U | T |
| φ | F | T | U | T |
| | U | U | U | U |
| | T | F | U | T |

| strong $\neg\varphi \vee \psi$ | | ψ | | |
|-----------------------------------|---|--------|---|---|
| | | F | U | T |
| φ | F | T | T | T |
| | U | U | U | T |
| | T | F | U | T |


| Łukasiewicz $\varphi \rightarrow \psi$ | | ψ | | |
|---|---|--------|---|---|
| | | F | U | T |
| φ | F | T | T | T |
| | U | U | T | T |
| | T | F | U | T |

ABOUT FUTURE

$$\llbracket \mathbf{F}\varphi \rrbracket_w^{\mathfrak{M}} \stackrel{\text{def}}{=} \begin{cases} \text{T} & \text{if } (\forall h \ni w)(\exists v > w) \llbracket \varphi \rrbracket_w^{\mathfrak{M}} = \text{T} \\ \text{F} & \text{if } (\forall h \ni w)(\exists v > w) \llbracket \varphi \rrbracket_w^{\mathfrak{M}} = \text{F} \\ \text{U} & \text{otherwise} \end{cases}$$

ABOUT FUTURE

$$\llbracket \mathbf{F}\varphi \rrbracket_w^{\mathfrak{M}} \stackrel{\text{def}}{=} \begin{cases} \text{T} & \text{if } (\forall h \ni w)(\exists v > w) \llbracket \varphi \rrbracket_w^{\mathfrak{M}} = \text{T} \\ \text{F} & \text{if } (\forall h \ni w)(\exists v > w) \llbracket \varphi \rrbracket_w^{\mathfrak{M}} = \text{F} \\ \text{U} & \text{otherwise} \end{cases}$$



Now go back to the sea battle

$$\llbracket \mathbf{F}p \rrbracket_w = \text{U}, \quad \llbracket \mathbf{F}\neg p \rrbracket_w = \text{U}, \quad \llbracket \mathbf{F}p \vee \mathbf{F}\neg p \rrbracket_w = \text{U}$$