

# Metropolis-Hastings

# Metropolis-Hastings

Sometimes Gibbs samples are too correlated

# Metropolis-Hastings

Sometimes Gibbs samples are too correlated

Apply rejection sampling to Markov Chains

# Metropolis-Hastings

For  $k = 1, 2, \dots$

- Sample  $x'$  from a wrong  $Q(x^k \rightarrow x')$

# Metropolis-Hastings

For  $k = 1, 2, \dots$

- Sample  $x'$  from a **wrong**  $Q(x^k \rightarrow x')$
- Accept proposal  $x'$  with probability  $A(x^k \rightarrow x')$

# Metropolis-Hastings

For  $k = 1, 2, \dots$

- Sample  $x'$  from a **wrong**  $Q(x^k \rightarrow x')$
- Accept proposal  $x'$  with probability  $A(x^k \rightarrow x')$
- Otherwise stay at  $x^k$

$$x^{k+1} = x^k$$

# Metropolis-Hastings

For  $k = 1, 2, \dots$

- Sample  $x'$  from a **wrong**  $Q(x^k \rightarrow x')$
- Accept proposal  $x'$  with probability  $A(x^k \rightarrow x')$
- Otherwise stay at  $x^k$

$$x^{k+1} = x^k$$

$$T(x \rightarrow x') = Q(x \rightarrow x')A(x \rightarrow x') \quad \text{for all } x \neq x'$$

# Metropolis-Hastings

For  $k = 1, 2, \dots$

- Sample  $x'$  from a **wrong**  $Q(x^k \rightarrow x')$
- Accept proposal  $x'$  with probability  $A(x^k \rightarrow x')$
- Otherwise stay at  $x^k$

$$x^{k+1} = x^k$$

$$T(x \rightarrow x') = Q(x \rightarrow x')A(x \rightarrow x') \quad \text{for all } x \neq x'$$

$$T(x \rightarrow x) = Q(x \rightarrow x)A(x \rightarrow x) + \sum_{x' \neq x} Q(x \rightarrow x')(1 - A(x \rightarrow x'))$$



# Metropolis-Hastings

For  $k = 1, 2, \dots$

- Sample  $x'$  from a **wrong**  $Q(x^k \rightarrow x')$
- Accept proposal  $x'$  with probability  $A(x^k \rightarrow x')$
- Otherwise stay at  $x^k$

$$x^{k+1} = x^k$$

$$T(x \rightarrow x') = Q(x \rightarrow x')A(x \rightarrow x') \quad \text{for all } x \neq x'$$

$$\begin{aligned} T(x \rightarrow x) &= Q(x \rightarrow x)A(x \rightarrow x) \\ &\quad + \sum_{x' \neq x} Q(x \rightarrow x')(1 - A(x \rightarrow x')) \end{aligned}$$

**How to choose A:**  $\pi(x') = \sum_x \pi(x)T(x \rightarrow x')$

# Detailed Balance

$$\pi(x') = \sum_x \pi(x) T(x \rightarrow x')$$

# Detailed Balance

**If**  $\pi(x)T(x \rightarrow x') = \pi(x')T(x' \rightarrow x)$

**Then**  $\pi(x') = \sum_x \pi(x)T(x \rightarrow x')$

# Detailed Balance

**If**  $\pi(x)T(x \rightarrow x') = \pi(x')T(x' \rightarrow x)$

**Then**  $\pi(x') = \sum_x \pi(x)T(x \rightarrow x')$

**Proof**  $\sum_x \pi(x)T(x \rightarrow x') = \sum_x \pi(x')T(x' \rightarrow x)$

# Detailed Balance

**If**  $\pi(x)T(x \rightarrow x') = \pi(x')T(x' \rightarrow x)$

**Then**  $\pi(x') = \sum_x \pi(x)T(x \rightarrow x')$

**Proof** 
$$\begin{aligned} \sum_x \pi(x)T(x \rightarrow x') &= \sum_x \pi(x')T(x' \rightarrow x) \\ &= \pi(x') \sum_x T(x' \rightarrow x) \end{aligned}$$

# Detailed Balance

**If**  $\pi(x)T(x \rightarrow x') = \pi(x')T(x' \rightarrow x)$

**Then**  $\pi(x') = \sum_x \pi(x)T(x \rightarrow x')$

**Proof** 
$$\begin{aligned} \sum_x \pi(x)T(x \rightarrow x') &= \sum_x \pi(x')T(x' \rightarrow x) \\ &= \pi(x') \underbrace{\sum_x T(x' \rightarrow x)}_1 \end{aligned}$$

# Detailed Balance

**If**  $\pi(x)T(x \rightarrow x') = \pi(x')T(x' \rightarrow x)$

**Then**  $\pi(x') = \sum_x \pi(x)T(x \rightarrow x')$

**Proof** 
$$\begin{aligned} \sum_x \pi(x)T(x \rightarrow x') &= \sum_x \pi(x')T(x' \rightarrow x) \\ &= \pi(x') \sum_x T(x' \rightarrow x) \\ &= \pi(x') \end{aligned}$$

# Metropolis-Hastings

For  $k = 1, 2, \dots$

- Sample  $x'$  from a **wrong**  $Q(x^k \rightarrow x')$
- Accept proposal  $x'$  with probability  $A(x^k \rightarrow x')$
- Otherwise stay at  $x^k$

$$x^{k+1} = x^k$$

$$T(x \rightarrow x') = Q(x \rightarrow x')A(x \rightarrow x') \quad \text{for all } x \neq x'$$

$$T(x' \rightarrow x') = Q(x' \rightarrow x')$$

**How to choose A:**

$$\pi(x') = \sum_x \pi(x)T(x \rightarrow x')$$



# Metropolis-Hastings

For  $k = 1, 2, \dots$

- Sample  $x'$  from a **wrong**  $Q(x^k \rightarrow x')$
- Accept proposal  $x'$  with probability  $A(x^k \rightarrow x')$
- Otherwise stay at  $x^k$

$$x^{k+1} = x^k$$

$$T(x \rightarrow x') = Q(x \rightarrow x')A(x \rightarrow x') \quad \text{for all } x \neq x'$$

$$T(x' \rightarrow x) = Q(x' \rightarrow x)$$

**How to choose A:**

$$\pi(x)T(x \rightarrow x') = \pi(x')T(x' \rightarrow x)$$