
a Calibration allocations
Four endpoints
+ random site
+ combination-directed
+ response-directed
Gold: stimulation site; pink: readouts.
b 40 networks, one noise draw
c 80 networks, three noise draws
Held-out response MSE
10^{-5}
10^{-7}
10^{-9}
10^{-11}
Four ends
+ random
+ comb.
+ response
d Fifth-site comparisons in both cohorts
Primary
Secondary
Adjusted, 3 tests
40 networks
Change, % [95% CI]
80 networks
Change, % [95% CI]
Profile width², combination vs random
-27.15 [-40.09, -15.18]
not measured
Combination MSE, combination vs random
-31.31 [-55.93, +0.83]
-31.07 [-42.35, -19.81]
Combination MSE, response vs random
-29.19 [-50.89, -4.38]
-17.73 [-28.39, -5.90]
Response MSE, combination vs random
-8.95 [-41.49, +33.68]
+13.50 [-37.19, +89.23]
Response MSE, response vs random
-35.90 [-57.62, -2.47]
+0.24 [-41.36, +74.08]
Response MSE, random fifth vs four ends
-18.26 [-49.02, +9.42]
-30.25 [-56.29, -5.98]
Response MSE, response fifth vs four ends
not tested
-30.08 [-52.31, -16.17]
−50%
0
+100%
Change in mean