

Submitted to: *Environ. Res. Lett.*

# The Effects of Weather on Maize Yields: New Evidence from Kenya

Monika Novackova, Pedram Rowhani, Martin Todd and  
Dominic Kniveton

Department of Geography, University of Sussex, Falmer, UK

E-mail: [monika.novac@gmail.com](mailto:monika.novac@gmail.com)

November 2018

**Abstract.** ...Applying the linear mixed effects models, we found that...

.. to be written later..

## 1. Introduction

### Paragraph 1

- Extreme weather causes disasters → early warning systems have been developed

### Paragraph 2

- What weather forecasts (measures) have been used in EWS? *ref. literature*
  - \* Mostly seasonal precip. totals and temperature averages

### Paragraph 3

- Identify difference between hazard and disaster
  - \* Not every hazard turns into disaster
  - \* For a hazard to become a disaster it needs to have **impact**
  - \* Here, we identify the key metrics which have impact on yield

### Paragraph 4

- Crop yield versus climate forecasting

## Paragraph 5

- Aim of the paper

## 2. Methods

...a case study looking at Kenya...

Our dataset consists of an yearly panel of 47 counties of Kenya describing the period of 1981-2017.

### 2.1. Data

- Source of the climate data: ( n.d.a) (BOKU) and ( n.d.b)
- Source of the yield data (Kenya MoA)

### 2.2. Statistical approach

- *We used commonly used measures of weather/drought (Only mention the significant weather measures/variables)*
- Describe the temporal aggregation of the weather variables, seasons

Kenya consists of 47 counties with semi-autonomous county governments (Barasa, Manyara, Molyneux & Tsofa 2017). As a result of the high degree of county-level autonomy, the policies and regulations often differ across the counties, hence the effects of weather on crop yield are likely to be different across the counties. Therefore, following the standard methodology, we estimated a battery of linear mixed effects models (also known as mixed models) commonly used to analyse longitudinal data (Bates, Pinheiro, Pinheiro & Bates 2000). Mixed models are suitable for analysis of panel data as they

account for the panel structure of the dataset. These types of models include both fixed effects parameters and random effects. Fixed effects are analogous to parameters in a classical linear regression model and value of each effect is assumed to be fixed over all counties (Bates n.d.). On the other hand, random effects are unobserved random variables. There are at least three benefits of treating a set of parameters as a random sample from some distribution. *(i)* Extrapolation of inference to a wider population *(ii)* improved accounting for system uncertainty and *(iii)* efficiency of estimation

Formally, a linear mixed model can be described by the distribution of two vectors of random variables: the response  $\mathcal{Y}$  and the vector of random effects  $\mathcal{B}$ . The distribution of  $\mathcal{B}$  is multivariate normal and the conditional distribution of  $\mathcal{Y}$  given  $\mathcal{B} = \mathbf{b}$  is multivariate normal of a form

$$(\mathcal{Y}|\mathcal{B} = \mathbf{b}) \sim N(\mathbf{X}\beta + \mathbf{Zb}, \sigma^2\mathbf{I}), \quad (1)$$

where  $\mathbf{X}$  is an  $n \times p$  model matrix of fixed effects,  $\beta$  is a  $p$ -dimensional fixed-effects parameter,  $\mathbf{Z}$  is an  $n \times q$  model matrix for the  $q$ -dimensional vector of random-effects variable  $\mathcal{B}$  evaluated at  $\mathbf{b}$  and  $\sigma$  a scale factor. The distribution of  $\mathcal{B}$  can be written as:

$$\mathcal{B} \sim N(0, \Sigma), \quad (2)$$

where  $\Sigma$  is a  $q \times q$  positive semi-definite variance-covariance matrix.

- Add a description of residual and other diagnostic tests (AIC, VIF, autocorrelation)
- Describe the procedure that I have applied to find the preferred combination of fixed effects and random effects
- Describe the procedure that I have applied to find the preferred way of modelling the correlation structure in errors (ARMA errors)

### 3. Results

- Tables of estimates of the preferred specification which includes the significant weather variables (see the preliminary results in Table 1)
  - For all counties
  - For Arid and semi-arid (ASAL) counties
  - For non-ASAL counties


**Table 1.** *Mixed effects model: Log of maize yield and weather, ARMA(1,1) errors*

	<i>All counties</i>		<i>ASAL</i>		<i>non-ASAL</i>	
<b>Fixed effects:</b>	<i>exp</i> ( $\beta$ )	<b>F-value<sup>a</sup></b>	<i>exp</i> ( $\beta$ )	<b>F-value<sup>a</sup></b>	<i>exp</i> ( $\beta$ )	<b>F-value<sup>a</sup></b>
Intercept	1.296***	19.916	1.276*	5.230	1.410**	10.061
Prec. total	1.081*	5.402	1.006	0.022	1.278***	19.386
Prec. total sq.	0.973*	4.289	1.004	0.051	0.880***	23.747
Prec. c. of var.	0.924 <sup>•</sup>	3.277	0.969	0.246	0.909	2.231
Dry spell -length	0.935*	4.810	0.833**	6.969	0.988	0.163
Dry spells $\geq$ 4 d.	0.939*	4.826	0.855**	8.065	0.989	0.096
Temp. - average	0.819***	12.127	0.808*	5.376	0.878	1.580
Temp. std. dev.	1.043 <sup>•</sup>	3.125	1.039	0.558	1.059 *	5.640
<b>Random effects:</b>						
Intercept						
<i>Number of observations:</i>	1300		698		602	

*Notes:* Standard errors in brackets; <sup>•</sup>  $p < 0.1$ ; \*  $p < 0.05$ ; \*\*  $p < 0.01$ ; \*\*\*  $p < 0.001$

<sup>a</sup> Marginal (type III) sum of squares. The F-statistics correspond to the sum of squares attributable to each fixed effect.

- Verbal description and interpretation of the results. Discussing goodness of fit using various criteria such as AIC or alternatives to  $R^2$ .



**Figure 1.** Marginal effects of seasonal rainfall and temperature

- If we get the yield data for the period from 2015 onwards: Out of sample predictions and comparison with the real data

University of Natural Resources and Life Sciences Vienna. Accessed in October 2018.

**URL:** *<http://ivfl-info.boku.ac.at/index.php/homepage?view=featured>*

Berkeley Earth. Accessed in October 2018.

**URL:** *<http://berkeleyearth.org/data/>*

Barasa, E., Manyara, A., Molyneux, S. & Tsofa, B. (2017).

Bates, D. M. (n.d.). lme4: Mixed-effects modeling with R.

Bates, J., Pinheiro, J., Pinheiro, J. & Bates, D. (2000). *Mixed-Effects Models in S and S-PLUS*, Statistics and Computing, Springer New York.