

1 The Big Picture

1.1 Why Study Statistics?

Communication, Quality Control, Decision Making, Big Data, ...

Statistics =

1.2 Descriptive and Inferential Statistics

Descriptive Statistics – quantifies characteristics of the distribution of data.

- Measures of central tendency
- Measures of variation
- Correlation, regression

Inferential Statistics – makes inferences about the population based on sample data.

- Estimation
- Hypothesis testing
- Statistical comparison

1.3 Under the hood of Statistics: Probability Distributions

- Binomial distribution
- Poisson distribution
- Normal distribution
- t-distribution
- Chi square distribution

1.4 Terminology: Population vs. Sample

Population: The complete collection of all elements (measurements, scores, etc.) to be studied.

Sample: A sub-collection of elements drawn from a population.
(simple random sampling, stratified sampling, etc.)

2 Distributions

2.1 Frequency Distribution: Histograms

The *frequency* of a particular observation is the number of times the observation occurs in the data

Consider the following data of cosmic radiation counts under various depths of snow.¹


Depth $x(\text{cm})$	Frequency (radiation count)	Relative Frequency	Mark (Label)
$0 \leq x \leq 3$	15	$\frac{15}{51} = 0.294 = 29.4\%$	1.5
$3 < x \leq 6$	12	$\frac{12}{51} = 0.235 = 23.5\%$	4.5
$6 < x \leq 9$	10	$\frac{10}{51} = 0.196 = 19.6\%$	7.5
$9 < x \leq 12$	4	$\frac{4}{51} = 0.078 = 7.8\%$	10.5
$12 < x \leq 15$	7	$\frac{7}{51} = 0.137 = 13.7\%$	13.5
$15 < x \leq 18$	3	$\frac{3}{51} = 0.059 = 5.9\%$	16.5

- The range of measurements (radiation counts) is divided into intervals called **classes**.
- The size (width) of each class is called the **class width**.
- **Class boundaries** can be set up so that there is no overlap between adjacent classes.
- **Frequency** is the number of elements belong to a given class.
- **Relative frequency** is frequency / total number of elements in the sample.

Exercise 1: What would happen if class width is too small or too large?

¹“Deep snow measurements suggested using cosmic radiation” by Bissell and Burson, *Water Resources Research*, 1974, vol 10, no.6, p.1243.

Histogram: Graphical representation of frequency distribution



Exercise 2: List a few examples of frequency distribution.

The most common distribution:

3 Measures of Central Tendency

Mean, Median, Mode, Mid-range

3.1 Mean (Arithmetic average)

For a *sample* with n elements $\{x_1, x_2, x_3, \dots, x_n\}$, the mean is defined to be

$$\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i \quad (\text{Sample mean})$$

- The mean of the whole *population* is denoted by $\mu = \frac{1}{N} \sum_{i=1}^N x_i$.
- \bar{x} is sensitive to extreme values in the data set.

Exercise 3: Calculate the mean of the GPA's of 3 students at BCIT. Their GPA's are $x_1 = 80, x_2 = 70, x_3 = 60$

3.2 Median

The *median* is the "middle" of a sorted list of numbers. If the number of elements n is odd, then the median is defined to be

$$\tilde{x} = x_m, \quad \text{where } m = \text{ceil}\left(\frac{n}{2}\right)$$

If n is even

$$\tilde{x} = \frac{x_{\frac{n}{2}+1} + x_{\frac{n}{2}}}{2}$$

The median is less affected by extreme values than the mean.

Exercise 4: Consider the following data set:

$$\{y_1 = 22, y_2 = 24, y_3 = 24, y_4 = 30, y_5 = 45, y_6 = 99\}$$

1. Find the mean and the median of $\{y_1, y_2, y_3, y_4, y_5\}$

2. Find the mean and the median of $\{y_1, y_2, y_3, y_4, y_5, y_6\}$

3.3 Mode

The value that has the highest frequency or "peak" in the frequency distribution.

- A distribution with 2 peaks is called **bimodal distribution**. **Unimodal** for one peak, **multimodal** for many peaks.
- The class that has the greatest frequency is called the **modal class**.

Exercise 5: What is the mode of the data set given in §2.1?

3.4 Mid-range

The middle of the range of the data values. That is,

$$(\text{mid-range}) = \frac{(\text{minimum}) + (\text{maximum})}{2}$$

Exercise 6: Find all the measures of central tendency for the data set $\{2, 2, 2, 20, 34, 45, 210\}$.

4 Measures of Variation (dispersion)

We often characterize the frequency distribution of given data using a representative value (e.g., mean, median) and a measure of variation, which is the "width" of the distribution. We will discuss two measures of variation; the range and the standard deviation.

4.1 Range

The range is the difference between the maximum value and the minimum value in the data set.

$$\text{Range} = \text{Maximum} - \text{Minimum}$$

Exercise 7: The measured values of some quantity were $\{2.13, 2.22, 2.09, 2.11\}$. Find the range.

Exercise 8: Find the range of the following two distributions. Is the range a good measure of variation in this case?

4.2 Standard deviation


The most commonly used measure of variation is the standard deviation. It is the *root mean square* (RMS) of $([\text{data value}] - [\text{the mean}])$. That is,

$$\sigma = \sqrt{\frac{\sum_{i=1}^N (x_i - \mu)^2}{N}} \quad \dots \quad \begin{array}{l} \text{for a population} \\ \text{[population standard deviation]} \end{array}$$

$$s = \sqrt{\frac{\sum_{i=1}^n (x_i - \bar{x})^2}{n - 1}} \quad \dots \quad \begin{array}{l} \text{for a sample} \\ \text{[sample standard deviation]} \end{array}$$

Note that in the standard deviation of a sample the sum is divided by $n - 1$, instead of n , in order to compensate the tendency that s underestimate σ when the sample size is small.

Exercise 9: Compare the standard deviations of A and B in the figure above.



Exercise 10: Calculate the standard deviation of the sample $\{1, 2, 2, 3\}$.

4.3 Variance

s^2 and σ^2 are called the variance.