
covid | horizon 2 | all locations | evaluation: Jun 2025 to the end of season 2025/26 | log1p(counts) updates | feedback delay 0 week(s)


All 53 locations | Jun 2025 to the end of season 2025/26

Small: single locations; large: means; axes: 1st to 99th percentile




All 53 locations | Up to peak (weeks -7 to 0)

Small: single locations; large: means; axes: 1st to 99th percentile




All 53 locations | After peak (weeks 1 to 8)

Small: single locations; large: means; axes: 1st to 99th percentile




United States | Jun 2025 to the end of season 2025/26

66 weeks




United States | Up to peak (weeks -7 to 0)

21 weeks




United States | After peak (weeks 1 to 8)

16 weeks




California | Jun 2025 to the end of season 2025/26

66 weeks




California | Up to peak (weeks -7 to 0)

18 weeks




California | After peak (weeks 1 to 8)

8 weeks




Texas | Jun 2025 to the end of season 2025/26

66 weeks




Texas | Up to peak (weeks -7 to 0)

19 weeks




Texas | After peak (weeks 1 to 8)

10 weeks




New York | Jun 2025 to the end of season 2025/26

66 weeks




New York | Up to peak (weeks -7 to 0)

13 weeks




New York | After peak (weeks 1 to 8)

8 weeks




Pennsylvania | Jun 2025 to the end of season 2025/26

66 weeks




Pennsylvania | Up to peak (weeks -7 to 0)

21 weeks




Pennsylvania | After peak (weeks 1 to 8)

15 weeks




Illinois | Jun 2025 to the end of season 2025/26

66 weeks




Illinois | Up to peak (weeks -7 to 0)

12 weeks



Illinois | After peak (weeks 1 to 8)

14 weeks



Single trackers
- Original
- Fixed MultiQT (eta=0.001*S0)
- Log-slope rate (min m=0.002)
- Target oracle (hindsight)

Rate-grid experts
- Rate grid: equal
- Rate grid: learned, no sharing (H=4)
- Rate grid: learned, no sharing, common weights (H=4)
- Rate grid: learned + sharing (H=2)
- Rate grid: learned + sharing, common weights (H=2)
- Rate grid: sharing + outer (outer m=0.001)
- Rate grid: sharing, common weights + outer (outer m=0.001)

Weekly starts: immediate admission
- First start only (m=0.05)
- Weekly starts: equal (m=0.05)
- Weekly starts: learned (m=0.05, lambda=0.25)
- Starts x rate grid: learned (lambda=0.25)
- Starts equal + outer (outer m=0.001)
- Starts learned + outer (outer m=0.001)
- Starts x rate grid + outer (outer m=0.001)

Diverse experts
- Diverse experts: equal (slow m=0.001)
- Diverse experts: learned + sharing (slow m=0.001, H=8)
- Diverse experts: learned + sharing, common weights (slow m=0.004)
- Diverse experts: soft switching (strength=1)

Studentized median
- Studentized median (RMS of 5, normal start, eta=0.003)
- Studentized median (base 80% width, normal start, eta=0.003)
- Median, constant sigma (RMS of 5, normal start, eta=0.1)

Studentized full base
- Studentized full base (base 80% width, eta=0.003)
- Studentized full base (level scales, eta=0.003)
- Studentized full base (level RMS of 5, eta=0.003)


Calibration error (CE): mean absolute coverage residual after averaging weeks within each quantile level.
Open circle: original forecasts. Filled circle: weights per quantile level; triangle: one weight vector shared by all levels; square / diamond: the same + outer MultiQT.

Studentized median methods: sigma_t = RMS of the last 5 released median residuals, or the base's 80% interval width / 2.563 (a normal sd).
With residuals, the original base is issued in the first 7 weeks, until sigma_t exists; these weeks are scored as the base.
Studentized full base: offsets start at the base; level scales = normal sd of the base's central interval at each level, or the RMS of each level's last 5 released residuals (weighted isotonic projection).

covid | horizon 2 | all locations | evaluation: Jun 2025 to the end of season 2025/26 | log1p(counts) updates | feedback delay 0 week(s)


All 53 locations | Jun 2025 to the end of season 2025/26

Small: single locations; large: means; axes: 1st to 99th percentile




All 53 locations | Up to peak (weeks -7 to 0)

Small: single locations; large: means; axes: 1st to 99th percentile




All 53 locations | After peak (weeks 1 to 8)

Small: single locations; large: means; axes: 1st to 99th percentile




United States | Jun 2025 to the end of season 2025/26

66 weeks




United States | Up to peak (weeks -7 to 0)

21 weeks




United States | After peak (weeks 1 to 8)

16 weeks




California | Jun 2025 to the end of season 2025/26

66 weeks




California | Up to peak (weeks -7 to 0)

18 weeks




California | After peak (weeks 1 to 8)

8 weeks




Texas | Jun 2025 to the end of season 2025/26

66 weeks




Texas | Up to peak (weeks -7 to 0)

19 weeks




Texas | After peak (weeks 1 to 8)

10 weeks




New York | Jun 2025 to the end of season 2025/26

66 weeks




New York | Up to peak (weeks -7 to 0)

13 weeks




New York | After peak (weeks 1 to 8)

8 weeks




Pennsylvania | Jun 2025 to the end of season 2025/26

66 weeks




Pennsylvania | Up to peak (weeks -7 to 0)

21 weeks




Pennsylvania | After peak (weeks 1 to 8)

15 weeks




Illinois | Jun 2025 to the end of season 2025/26

66 weeks




Illinois | Up to peak (weeks -7 to 0)

12 weeks



Illinois | After peak (weeks 1 to 8)

14 weeks



Single trackers
- Original
- Fixed MultiQT (eta=0.001*S0)
- Log-slope rate (min m=0.002)
- Target oracle (hindsight)

Rate-grid experts
- Rate grid: equal
- Rate grid: learned, no sharing (H=4)
- Rate grid: learned, no sharing, common weights (H=4)
- Rate grid: learned + sharing (H=2)
- Rate grid: learned + sharing, common weights (H=2)
- Rate grid: sharing + outer (outer m=0.001)
- Rate grid: sharing, common weights + outer (outer m=0.001)

Weekly starts: immediate admission
- First start only (m=0.05)
- Weekly starts: equal (m=0.05)
- Weekly starts: learned (m=0.05, lambda=0.25)
- Starts x rate grid: learned (lambda=0.25)
- Starts equal + outer (outer m=0.001)
- Starts learned + outer (outer m=0.001)
- Starts x rate grid + outer (outer m=0.001)

Diverse experts
- Diverse experts: equal (slow m=0.001)
- Diverse experts: learned + sharing (slow m=0.001, H=8)
- Diverse experts: learned + sharing, common weights (slow m=0.004)
- Diverse experts: soft switching (strength=1)

Studentized median
- Studentized median (RMS of 5, normal start, eta=0.003)
- Studentized median (base 80% width, normal start, eta=0.003)
- Median, constant sigma (RMS of 5, normal start, eta=0.1)

Studentized full base
- Studentized full base (base 80% width, eta=0.003)
- Studentized full base (level scales, eta=0.003)
- Studentized full base (level RMS of 5, eta=0.003)

Calibration error (CE): mean absolute coverage residual after averaging weeks within each quantile level.
Open circle: original forecasts. Filled circle: weights per quantile level; triangle: one weight vector shared by all levels; square / diamond: the same + outer MultiQT.

Studentized median methods: sigma_t = RMS of the last 5 released median residuals, or the base's 80% interval width / 2.563 (a normal sd).
With residuals, the original base is issued in the first 7 weeks, until sigma_t exists; these weeks are scored as the base.
Studentized full base: offsets start at the base; level scales = normal sd of the base's central interval at each level, or the RMS of each level's last 5 released residuals (weighted isotonic projection).