

covid | horizon 2 | all locations | full history | raw-count updates | feedback delay 2 week(s)


All 53 locations | Full observed history
Small: single locations; large: means; axes to the 99th percentile




All 53 locations | Up to peak (weeks -7 to 0)
Small: single locations; large: means; axes to the 99th percentile




All 53 locations | After peak (weeks 1 to 8)
Small: single locations; large: means; axes to the 99th percentile




United States | Full observed history
95 weeks




United States | Up to peak (weeks -7 to 0)
8 weeks




United States | After peak (weeks 1 to 8)
8 weeks




California | Full observed history
95 weeks




California | Up to peak (weeks -7 to 0)
8 weeks




California | After peak (weeks 1 to 8)
8 weeks




Texas | Full observed history
95 weeks




Texas | Up to peak (weeks -7 to 0)
8 weeks




Texas | After peak (weeks 1 to 8)
8 weeks




New York | Full observed history
95 weeks




New York | Up to peak (weeks -7 to 0)
8 weeks




New York | After peak (weeks 1 to 8)
7 weeks




Pennsylvania | Full observed history
95 weeks




Pennsylvania | Up to peak (weeks -7 to 0)
8 weeks




Pennsylvania | After peak (weeks 1 to 8)
7 weeks




Illinois | Full observed history
95 weeks



Illinois | Up to peak (weeks -7 to 0)
8 weeks



Illinois | After peak (weeks 1 to 8)
8 weeks



Single trackers
- Original
- Fixed MultiQT (eta=0.05*S0)
- Count-slope rate
- Target oracle (hindsight)

Rate-grid experts
- Rate grid: equal
- Rate grid: learned, no sharing
- Rate grid: learned, no sharing, common weights
- Rate grid: learned + sharing (H=8)
- Rate grid: learned + sharing, common weights
- Rate grid: sharing + outer (eta=0.005)
- Rate grid: sharing, common weights + outer

Weekly starts: immediate admission
- Weekly starts: equal (m=0.05)
- Weekly starts: learned (m=0.05)
- Starts x rate grid: learned
- Starts equal + outer
- Starts learned + outer
- Starts x rate grid + outer

Diverse experts
- Diverse experts: equal
- Diverse experts: learned + sharing (H=8)
- Diverse experts: learned + sharing, common weights

Studentized median
- Studentized median (RMS of 5, normal start, eta=0.1)
- Studentized median (base 80% width, normal start, eta=0.1)
- Median, constant sigma (RMS of 5, normal start, eta=0.1)

Studentized full base
- Studentized full base (base 80% width, eta=0.1)
- Studentized full base (level scales, eta=0.1)
- Studentized full base (level RMS of 5, eta=0.1)

Calibration error (CE): mean absolute coverage residual after averaging weeks within each quantile level.
Open circle: original forecasts. Filled circle: weights per quantile level; triangle: one weight vector shared by all levels; square / diamond: the same + outer MultiQT.
Not drawn (identical forecasts): First start only (m=0.05) = Fixed MultiQT (eta=0.05*S0)
Studentized median methods: sigma_t = RMS of the last 5 released median residuals, or the base's 80% interval width / 2.563 (a normal sd).
With residuals, the original base is issued in the first 9 weeks, until sigma_t exists; these weeks are scored as the base.
Studentized full base: offsets start at the base; level scales = normal sd of the base's central interval at each level, or the RMS of each level's last 5 released residuals (weighted isotonic projection).