

---

---

# *The $p$ -adic Method*

DRAFT

No Cover Image

Use `\coverimage{filename}` to add an image

阿巴阿巴!

---

---

# Contents

<b>1</b>	<b>Adelic topology</b>	<b>1</b>
1.1	Basic adelic subset . . . . .	1
1.2	General adelic subset . . . . .	1
1.3	Adelic topology . . . . .	1
<b>2</b>	<b>Interpolation</b>	<b>1</b>
2.1	Interpolation of analytic maps . . . . .	1
2.2	Interpolation on an analytic open subset of morphisms . . . . .	2
<b>3</b>	<b>Applications</b>	<b>2</b>
3.1	Existence of non-preperiodic points . . . . .	2
3.2	DML conjecture for étale morphisms . . . . .	2
3.3	DML conjecture for adelic general points . . . . .	2
	<b>References</b>	<b>2</b>

## 1 Adelic topology

The main reference of this section is [Xie25].

Let  $\mathbf{k}$  be a number field, and let  $M_{\mathbf{k}}$  be the set of places of  $\mathbf{k}$ .

### 1.1 Basic adelic subset

### 1.2 General adelic subset

### 1.3 Adelic topology

## 2 Interpolation

### 2.1 Interpolation of analytic maps

The main reference of this section is [Ame11; BGT10; Poo14]. We first state the main theorem of this section.

**Theorem 2.1** (ref.[Poo14, Theorem 1] cf.[BGT10, Theorem 3.3]). Let  $\mathbf{k}$  be a complete non-archimedean field of characteristic 0 with  $|p|_{\mathbf{k}} = 1/p$ . Set  $r_p = p^{-1/(p-1)}$ . Let  $E = E(0, 1) = \{x \in \mathbf{k}^d \mid \|x\| \leq 1\}$  be the closed unit ball in  $\mathbf{k}^d$ . Here the norm on  $\mathbf{k}^d$  or  $\mathbf{k}^{\circ}\{\underline{T}\}^d$  is the supremum norm, i.e.,  $|x| = \max_{1 \leq i \leq d} |x_i|$ .

Suppose that  $\Phi = (\Phi_1, \dots, \Phi_d) \in \mathbf{k}^{\circ}\{\underline{T}\}^d$  satisfies  $\|\Phi - \text{id}_E\| \leq r_p$ . Then there exists a function

$F \in \mathbf{k}\{\underline{T}, S\}^d$  such that for each  $n \in \mathbb{Z}_{\geq 0}$  and each  $x \in E$ ,

$$F(x, n) = \Phi^n(x).$$

| *Proof.* Yang: To be added. □

Yang: If  $f$  is invertible, can we see that  $g$  is unique?

## 2.2 Interpolation on an analytic open subset of morphisms

The main reference of this section is [Xie25, Section 3.2]. We first state the main theorem of this section.

**Theorem 2.2** (ref.[Xie25, Proposition 3.24]). Let  $\mathbf{k}$  be a finitely generated field over  $\mathbb{Q}$ ,  $X$  a projective variety defined over  $\mathbf{k}$ , and  $f : X \dashrightarrow X$  a dominant rational self-map defined over  $\mathbf{k}$ .

There exists an iteration  $g = f^m$  of  $f$ , an embedding  $\mathbf{k} \hookrightarrow \mathbb{C}_p$  for some prime  $p \geq 3$ , an analytic open subset  $U \subseteq X(\mathbb{C}_p)$  and an analytic map  $\Phi : \mathbb{C}_p^\circ \times U \rightarrow U$  such that

- (1)  $U \cong (\mathbb{C}_p^\circ)^d$  analytically, where  $d = \dim X$ ;
- (2)  $g$  is well-defined on  $U$ ,  $U$  is invariant under  $g$  and  $\|g|_U - \text{id}_U\| < 1/p$ ;
- (3)  $\Phi(n, x) = g^n(x)$  for each  $n \in \mathbb{Z}_{\geq 0}$  and each  $x \in U$ ;

**Lemma 2.3.** Yang: There is a good model for  $(X, f)$ .

**Example 2.4.** Let  $X = E \times E$  with  $E$  an elliptic curve without complex multiplication defined over a number field  $\mathbf{k}$ , and let  $f : X \rightarrow X$  be the endomorphism defined by  $(a, b) \mapsto (a + b, b)$ . Yang: To be continued.

## 3 Applications

### 3.1 Existence of non-preperiodic points

**Theorem 3.1** (ref.[Ame11, Corollary 9]). Let  $\mathbf{k}$  be an algebraically closed field of characteristic 0,  $X$  a projective variety defined over  $\mathbf{k}$ , and  $f : X \dashrightarrow X$  a dominant rational self-map defined over  $\mathbf{k}$ . Then there exists a point  $x \in X(\mathbf{k})$  such that the forward orbit  $\mathcal{O}_f(x) = \{f^n(x) : n \geq 0\}$  is well-defined and infinite.

### 3.2 DML conjecture for étale morphisms

### 3.3 DML conjecture for adelic general points

## References

- [Ame11] E Amerik. “Existence of non-preperiodic algebraic points for a rational self-map of infinite order”. In: *Mathematical Research Letters* 18.2 (2011), pp. 251–256 (cit. on pp. 1, 2).

- 
- [BGT10] Jason P Bell, Dragos Ghioca, and Thomas J Tucker. “The dynamical Mordell-Lang problem for étale maps”. In: *American journal of mathematics* 132.6 (2010), pp. 1655–1675 (cit. on p. [1](#)).
- [Poo14] Bjorn Poonen. “ $p$ -adic interpolation of iterates”. In: *Bulletin of the London Mathematical Society* 46.3 (2014), pp. 525–527 (cit. on p. [1](#)).
- [Xie25] Junyi Xie. “The existence of Zariski dense orbits for endomorphisms of projective surfaces”. In: *Journal of the American Mathematical Society* 38.1 (2025), pp. 1–62 (cit. on pp. [1](#), [2](#)).

DRAFT