

Affinoid algebras

1 The first properties

Definition 1. Let \mathbf{k} be a non-archimedean field. A banach \mathbf{k} -algebra A is called a *affinoid \mathbf{k} -algebra* if there exists an admissible surjective homomorphism

$$\varphi : \mathbf{k}\{\underline{T}/\underline{r}\} \twoheadrightarrow A$$

for some $\underline{r} = (r_1, \dots, r_n) \in \mathbb{R}_{>0}^n$.

If one can choose $r_1 = \dots = r_n = 1$, then we say that A is a *strict affinoid \mathbf{k} -algebra*.

Example 2. Yang: To be added.

Example 3. Let A be an affinoid \mathbf{k} -algebra and $f_i, g_i \in A$. Then the normed localization

$$A\{(f_i/g_i)/r_i\}_{i=1}^n = A \otimes_{\mathbf{k}\{\underline{T}\}} \mathbf{k}\{\underline{T}/\underline{r}, \underline{S}\}/(g_i T_i - f_i)_{i=1}^n$$

is again an affinoid \mathbf{k} -algebra. Yang: To be added.

Proposition 4. Let A be an affinoid \mathbf{k} -algebra. Then A is noetherian, and every ideal of A is closed.

Proof. Yang: To be completed. □

Proposition 5. Let A be an affinoid \mathbf{k} -algebra. Then there exists a constant $C > 0$ and $N > 0$ such that for all $f \in A$ and $n \geq N$, we have

$$\|f^n\| \leq C\rho(f)^n.$$

In particular, $\text{Qnil}(A) = \text{nil}(A)$.

Furthermore, if A is reduced, we have

$$\|f\| \leq C\rho(f)$$

for all $f \in A$.

Proof. Yang: To be completed. □

Proposition 6. Let A be an affinoid \mathbf{k} -algebra. If and only if $\rho(f) \in \sqrt{|\mathbf{k}|} \cup \{0\}$ for all $f \in A$, then A is strict. Yang: To be complete.

Proof. Yang: To be completed. □

Definition 7. Let \mathbf{k} be a non-archimedean field. We define the *ring of restricted Laurent series* over \mathbf{k} as

$$\mathbf{K}_r = \mathbf{L}_{\mathbf{k},r} = \left\{ \sum_{n \in \mathbb{Z}} a_n T^n : a_n \in \mathbf{k}, \lim_{|n| \rightarrow \infty} |a_n| r^n = 0 \right\}$$

equipped with the norm

$$\|f\| = \sup_{n \in \mathbb{Z}} |a_n| r^n.$$

Yang: Is \mathbf{K}_r always a field? Yang: Do we have $\mathbf{L}_{\mathbf{k},r} = \text{Frac}(\mathbf{k}\{T/r\})$?

Proposition 8. Let \mathbf{k} be a non-archimedean field. If $r \notin \sqrt{|\mathbf{k}^\times|}$, then \mathbf{K}_r is a complete non-archimedean field with non-trivial absolute value extending that of \mathbf{k} .

Yang: Tensor with \mathbf{K}_r .

Theorem 9. Let A be a strict affinoid \mathbf{k} -algebra. Then there exists a finite injective admissible homomorphism

$$\varphi : \mathbf{k}\{T\} \hookrightarrow A$$

for some $d \in \mathbb{N}$ and $r_1, \dots, r_d \in \mathbb{R}_{>0}$. Yang: To be checked.

2 Tate algebras and Weierstrass division

Definition 10. Let R be a non-archimedean banach ring and $r \in \mathbb{R}_{>0}$. A restricted power series $f = \sum_{\alpha \in \mathbb{N}^n} a_\alpha T^\alpha \in R\{\underline{T}/r\}$ is said to be *distinguished in the variable T_n of degree d* if

- $a_\alpha \in R$ is a unit for $\alpha = (0, \dots, 0, d)$;
- $\|a_\alpha\| r^\alpha < \|a_{(0, \dots, 0, d)}\| r_n^d$ for all $\alpha_n < d$.

Yang: To be revised.

Proposition 11. Let R be a non-archimedean banach ring. An element $f = \sum_{\alpha \in \mathbb{N}^n} a_\alpha T^\alpha \in R\{\underline{T}/r\}$ is invertible if and only if a_0 is invertible in R and $\|a_0\| > \|a_\alpha\| r^\alpha$ for all $\alpha \neq 0$.

Proof. Multiplying by a_0^{-1} , we can reduce to the case $a_0 = 1$. Let $g = \sum_{\alpha \in \mathbb{N}^n} b_\alpha T^\alpha$ be the inverse of f in $R[[\underline{T}]]$. Then we have

$$f \cdot g = \sum_{\alpha \in \mathbb{N}^n} a_\alpha T^\alpha \cdot \sum_{\beta \in \mathbb{N}^n} b_\beta T^\beta = \sum_{\gamma \in \mathbb{N}^n} \left(\sum_{\alpha+\beta=\gamma} a_\alpha b_\beta \right) T^\gamma = 1.$$

That is, for every $\gamma \neq 0 \in \mathbb{N}^n$,

$$b_\gamma = - \sum_{\substack{\alpha+\beta=\gamma \\ \alpha \neq 0}} a_\alpha b_\beta.$$

Let $A = \|f - 1\| < 1$. We show that for every $m \in \mathbb{N}$, there exists $C_m > 0$ such that for all $\alpha \in \mathbb{N}^n$ with $|\alpha| \geq C_m$, we have $\|b_\alpha\| r^\alpha \leq A^m$. For $m = 0$, note that $b_0 = 1$. By induction on γ with respect to the total order \leq_{total} , we have

$$\|b_\gamma\| r^\gamma \leq \max_{\substack{\alpha+\beta=\gamma \\ \alpha \neq 0}} \|a_\alpha\| r^\alpha \cdot \|b_\beta\| r^\beta \leq A \max_{\beta <_{\text{total}} \gamma} \|b_\beta\| r^\beta \leq 1.$$

Suppose that the claim holds for m . There exists $D_{m+1} \in \mathbb{N}$ such that for all $\alpha \in \mathbb{N}^n$ with $|\alpha| \geq D_{m+1}$, we have $\|a_\alpha\| r^\alpha \leq A^{m+1}$. Set $C_{m+1} = C_m + D_{m+1} + 1$. For any $\gamma \in \mathbb{N}^n$ with $|\gamma| \geq C_{m+1}$,

we have

$$\|b_\gamma\|r^\gamma \leq \max_{\substack{\alpha+\beta=\gamma \\ \alpha \neq 0}} \|a_\alpha\|r^\alpha \cdot \|b_\beta\|r^\beta \leq \max\{A^{m+1}, A \cdot A^m\} = A^{m+1}$$

since either $|\alpha| \geq D_{m+1}$ or $|\beta| \geq C_m$. Thus by induction, we have $\|b_\alpha\|r^\alpha \rightarrow 0$ as $|\alpha| \rightarrow +\infty$. It follows that $g \in R\{\underline{T}/r\}$. \square

Theorem 12 (Weierstrass preparation theorem). Let \mathbf{k} be a complete non-archimedean field. Let $f \in \mathbf{k}\{\underline{T}/r\}$ be a restricted power series that is distinguished in the variable T_n of degree d , i.e.,

$$f = \sum_{\alpha \in \mathbb{N}^{n-1}} a_\alpha T^\alpha + \sum_{\alpha_n \geq 1} a_\alpha T^\alpha$$

with $a_{(0,\dots,0,d)}$ being a unit in $\mathbf{k}\{\underline{T}/r\}$ and $\|a_\alpha\|r^\alpha < \|a_{(0,\dots,0,d)}\|r_n^d$ for all $\alpha_n < d$. Then there exists a unique monic polynomial $P \in \mathbf{k}\{\underline{T}/r\}[T_n]$ of degree d in T_n and a unique unit $U \in \mathbf{k}\{\underline{T}/r\}$ such that

$$f = P \cdot U.$$

Yang: To be checked.

Theorem 13 (Weierstrass division theorem). Let \mathbf{k} be a complete non-archimedean field. Let $f \in \mathbf{k}\{\underline{T}/r\}$ be a restricted power series that is distinguished in the variable T_n of degree d . Then for every $g \in \mathbf{k}\{\underline{T}/r\}$, there exists a unique $Q \in \mathbf{k}\{\underline{T}/r\}$ and a unique polynomial $R \in \mathbf{k}\{\underline{T}/r\}[T_n]$ of degree less than d in T_n such that

$$g = Q \cdot f + R.$$

Yang: To be checked.

Proposition 14. Let \mathbf{k} be a complete non-archimedean field and $r = (r_1, \dots, r_n) \in \mathbb{R}_+^n$. Then

$$\text{Spec } \mathbf{k}\{\underline{T}/r\} = \{\},$$

where

Appendix