

---

---

# *Non-archimedean Analysis*

DRAFT

No Cover Image  
Use \coverimage{filename} to add an image

# Contents

|          |   |          |
|----------|---|----------|
| <b>1</b> | <b>Valuation fields</b>   | <b>1</b> |
| 1.1      | Absolute values and completion . . . . .                        | 1        |
| 1.2      | Non-archimedean fields . . . . .                                | 3        |
| <b>2</b> | <b>Ultra-metric spaces</b>                                      | <b>4</b> |
| <b>3</b> | <b>Residue fields and reductions</b>                            | <b>5</b> |
| 3.1      | Recover non-archimedean complete fields algebraically . . . . . | 5        |
| 3.2      | Hensel's Lemma . . . . .  | 7        |
| 3.3      | Newton polygons . . . . .                                       | 8        |
| <b>4</b> | <b>Finite field extensions</b>                                  | <b>8</b> |
| 4.1      | Finite-dimensional vector space . . . . .                       | 8        |
| 4.2      | Finite field extensions . . . . .                               | 8        |
| <b>5</b> | <b>Analytic functions</b>                                       | <b>9</b> |
| 5.1      | Failure of continuous and differentiable functions . . . . .    | 9        |
| 5.2      | Power series . . . . .  | 9        |
| 5.3      | Analytic functions . . . . .                                    | 9        |
| <b>6</b> | <b>Example: p-adic fields</b>                                   | <b>9</b> |
| 6.1      | p-adic fields . . . . .   | 9        |
| 6.2      | Completion . . . . .  | 10       |

## 1 Valuation fields

### 1.1 Absolute values and completion

**Definition 1.1.** Let  $\mathbf{k}$  be a field. An *absolute value* on  $\mathbf{k}$  is a function  $\|\cdot\| : \mathbf{k} \rightarrow \mathbb{R}_{\geq 0}$  satisfying the following properties for all  $x, y \in \mathbf{k}$ :

- (a)  $\|x\| = 0$  if and only if  $x = 0$ ;
- (b)  $\|xy\| = \|x\| \cdot \|y\|$ ;
- (c)  $\|x + y\| \leq \|x\| + \|y\|$ .

A field  $\mathbf{k}$  equipped with an absolute value  $\|\cdot\|$  is called a *valuation field*.

**Remark 1.2.** Let  $\mathbf{k}$  be a field. Recall that a *valuation* on  $\mathbf{k}$  is a function  $v : \mathbf{k}^\times \rightarrow \mathbb{R}$  such that

- $\forall x, y \in \mathbf{k}^\times, v(xy) = v(x) + v(y)$ ;

- $\forall x, y \in \mathbf{k}^\times, v(x + y) \geq \min\{v(x), v(y)\}$ .

We can extend  $v$  to the whole field  $\mathbf{k}$  by defining  $v(0) = +\infty$ . Fix a real number  $\varepsilon \in (0, 1)$ . Then  $v$  induces an absolute value  $|\cdot|_v : \mathbf{k} \rightarrow \mathbb{R}_+$  defined by  $|x|_v = \varepsilon^{v(x)}$  for each  $x \in \mathbf{k}$ .

In some literature, the valuation  $v$  is called an *additive valuation* and the induced absolute value  $|\cdot|_v$  is called a *multiplicative valuation*. In this note, the term *valuation* always refers to the additive valuation.

**Example 1.3.** Let  $\mathbf{k}$  be a field. The *trivial absolute value* on  $\mathbf{k}$  is defined as

$$\|x\| := \begin{cases} 0, & x = 0; \\ 1, & x \neq 0. \end{cases}$$

**Definition 1.4.** The *(multiplicative) valuation group* of a valuation field  $(\mathbf{k}, \|\cdot\|)$  is defined as the subgroup of  $\mathbb{R}_{>0}$  given by

$$|\mathbf{k}^\times| := \{\|x\| : x \in \mathbf{k}^\times\}.$$

**Definition 1.5.** Let  $\mathbf{k}$  be a field. Two absolute values  $\|\cdot\|_1$  and  $\|\cdot\|_2$  on  $\mathbf{k}$  are said to be *equivalent* if there exists a real number  $c \in (0, 1)$  such that

$$\|x\|_1 = \|x\|_2^c, \quad \forall x \in \mathbf{k}.$$

Note that equivalent absolute values induce the same topology on the field  $\mathbf{k}$ . Moreover, the following lemma shows that the converse is also true.

**Lemma 1.6.** Let  $\mathbf{k}$  be a field and  $\|\cdot\|_1, \|\cdot\|_2$  be two absolute values on  $\mathbf{k}$ . Then the following statements are equivalent:

- (a)  $\|\cdot\|_1$  and  $\|\cdot\|_2$  are equivalent;
- (b)  $\|\cdot\|_1$  and  $\|\cdot\|_2$  induce the same topology on  $\mathbf{k}$ ;
- (c) The unit disks  $D_1 = \{x \in \mathbf{k} : \|x\|_1 < 1\}$  and  $D_2 = \{x \in \mathbf{k} : \|x\|_2 < 1\}$  are the same.

*Proof.* The implications (a)  $\Rightarrow$  (b) is obvious. Now we prove (b)  $\Rightarrow$  (c). For any  $x \in D_1$ , we have  $x^n \rightarrow 0$  as  $n \rightarrow \infty$  under the absolute value  $\|\cdot\|_1$  and thus under  $\|\cdot\|_2$ . Therefore,  $\|x\|_2^n \rightarrow 0$  as  $n \rightarrow \infty$ , which implies that  $\|x\|_2 < 1$ , i.e.,  $x \in D_2$ . Similarly, we can prove that  $D_2 \subseteq D_1$ .

Finally, we prove (c)  $\Rightarrow$  (a). If  $\|\cdot\|_1$  is trivial, then  $D_1 = \{0\}$  and thus  $\|\cdot\|_2$  is also trivial. In this case, they are equivalent. Suppose that both  $\|\cdot\|_1$  and  $\|\cdot\|_2$  are non-trivial. Pick any  $x, y \notin D_1 = D_2$ . Then there exist real numbers  $\alpha, \beta > 0$  such that  $\|x\|_1 = \|x\|_2^\alpha$  and  $\|y\|_1 = \|y\|_2^\beta$ . Suppose the contrary that  $\alpha \neq \beta$ . Consider the domain  $\Omega \subseteq \mathbb{Z}^2$  defined by

$$\begin{cases} n \log \|x\|_2 < m \log \|y\|_2; \\ n\alpha \log \|x\|_2 > m\beta \log \|y\|_2. \end{cases}$$

Since  $\alpha \neq \beta$ , the two lines defined by the equalities are not parallel. Thus  $\Omega$  is non-empty. Pick  $(n, m) \in \Omega$  and set  $z := x^n y^{-m}$ . Then we have  $\|z\|_2 < 1$  and  $\|z\|_1 > 1$ , a contradiction.  $\square$

**Definition 1.7.** Let  $(\mathbf{k}, \|\cdot\|)$  be a valuation field. We say that  $\mathbf{k}$  is *complete* if the metric  $d(x, y) := \|x - y\|$  makes  $\mathbf{k}$  a complete metric space.

**Lemma 1.8.** Let  $(\mathbf{k}, \|\cdot\|)$  be a valuation field and  $(\widehat{\mathbf{k}}, \|\cdot\|)$  its completion as a metric space. Then the operations of addition and multiplication on  $\mathbf{k}$  can be extended to  $\widehat{\mathbf{k}}$  uniquely, making  $(\widehat{\mathbf{k}}, \|\cdot\|)$  a complete valuation field containing  $\mathbf{k}$  as a dense subfield.

*Proof.* Simple analysis. □

**Example 1.9.** Let  $|\cdot|_\infty$  be the usual absolute value on the field  $\mathbb{Q}$  of rational numbers. Then  $(\mathbb{Q}, |\cdot|_\infty)$  is a valuation field. Its completion is the field  $\mathbb{R}$  of real numbers equipped with the usual absolute value.

**Example 1.10.** Let  $p$  be a prime number. For any non-zero rational number  $x \in \mathbb{Q}$ , we can write it as  $x = p^n \frac{a}{b}$ , where  $n \in \mathbb{Z}$  and  $a, b \in \mathbb{Z}$  are integers not divisible by  $p$ . The *p-adic absolute value* on  $\mathbb{Q}$  is defined as

$$|x|_p := \begin{cases} 0, & x = 0; \\ p^{-n}, & x = p^n \frac{a}{b} \text{ as above.} \end{cases}$$

Then  $(\mathbb{Q}, |\cdot|_p)$  is a valuation field. Its completion is the field

$$\mathbb{Q}_p = \left\{ \sum_{n=k}^{+\infty} a_n p^n : k \in \mathbb{Z}, a_n \in \{0, 1, \dots, p-1\} \right\}$$

of *p*-adic numbers equipped with the *p*-adic absolute value; see [Yang: to be added.](#)

Unlike the real number field  $\mathbb{R}$ , even a valuation field is complete, we can not expect the theorem of nested intervals to hold.

**Definition 1.11.** A valuation field  $(\mathbf{k}, \|\cdot\|)$  is called *spherically complete* if every decreasing sequence of closed balls in  $\mathbf{k}$  has a non-empty intersection.

**Example 1.12.** The field  $\mathbb{C}_p$  of *p*-adic complex numbers is not spherically complete, see [Yang: to be added.](#)

## 1.2 Non-archimedean fields

**Definition 1.13.** Let  $(\mathbf{k}, \|\cdot\|)$  be a valuation field. We say that  $\mathbf{k}$  is *non-archimedean* if its absolute value  $\|\cdot\|$  satisfies the *strong triangle inequality*:

$$\|x + y\| \leq \max\{\|x\|, \|y\|\}, \quad \forall x, y \in \mathbf{k}.$$

Otherwise, we say that  $\mathbf{k}$  is *archimedean*.

Let  $\mathbf{k}$  be a non-archimedean field. Then easily see that  $\{x \in \mathbf{k} : \|x\| \leq 1\}$  is a subring of  $\mathbf{k}$ . Moreover, it is a local ring whose maximal ideal is  $\{x \in \mathbf{k} : \|x\| < 1\}$ .

**Definition 1.14.** Let  $\mathbf{k}$  be a non-archimedean field. The *ring of integers* of  $\mathbf{k}$  is defined as

$$\mathbf{k}^\circ := \{x \in \mathbf{k} : \|x\| \leq 1\}.$$

Its maximal ideal is

$$\mathbf{k}^{\circ\circ} := \{x \in \mathbf{k} : \|x\| < 1\}.$$

The *residue field* of  $\mathbf{k}$  is defined as

$$\kappa_{\mathbf{k}} := \tilde{\mathbf{k}} := \mathbf{k}^{\circ}/\mathbf{k}^{\circ\circ}.$$

**Definition 1.15.** Let  $\mathbf{k}$  be a non-archimedean field. The *residue absolute value* on the residue field  $\kappa_{\mathbf{k}}$  is defined as

$$|x| := \inf_{y \in \varphi^{-1}(x)} \|y\|, \quad \forall x \in \kappa_{\mathbf{k}},$$

where  $\varphi : \mathbf{k}^{\circ} \rightarrow \kappa_{\mathbf{k}}$  is the canonical projection.

**Proposition 1.16.** Let  $\mathbf{k}$  be a non-archimedean field. Then the residue absolute value on the residue field  $\kappa_{\mathbf{k}}$  is trivial.

*Proof.* For any  $x \in \kappa_{\mathbf{k}}$ , if  $x = 0$ , then by definition  $|x| = 0$ . If  $x \neq 0$ , then  $\forall y \in \varphi^{-1}(x)$ , we have  $y \in \mathbf{k}^{\circ} \setminus \mathbf{k}^{\circ\circ}$ , i.e.,  $\|y\| = 1$ . Thus by definition  $|x| = 1$ .  $\square$

## 2 Ultra-metric spaces

We will use  $B(x, r)$  (resp.  $E(x, r)$ ) to denote the open ball (resp. closed ball) with center  $x$  and radius  $r$ .

**Definition 2.1.** A metric space  $(X, d)$  is called an *ultra-metric space* if its metric  $d$  satisfies the *strong triangle inequality*:

$$d(x, z) \leq \max\{d(x, y), d(y, z)\}, \quad \forall x, y, z \in X.$$

If  $(\mathbf{k}, \|\cdot\|)$  is a non-archimedean field, then the metric  $d(x, y) := \|x - y\|$  on  $\mathbf{k}$  makes  $(\mathbf{k}, d)$  an ultra-metric space.

**Proposition 2.2.** Let  $(X, d)$  be an ultra-metric space. Then for any  $x, y, z \in X$ , at least two of the three distances  $d(x, y), d(y, z), d(z, x)$  are equal. And the third distance is less than or equal to the common value of the other two.

*Proof.* Suppose that  $d(x, y) \geq d(y, z)$ . By the strong triangle inequality, we have

$$d(x, z) \leq \max\{d(x, y), d(y, z)\} = d(x, y).$$

On the other hand, by the strong triangle inequality again, we have

$$d(x, y) \leq \max\{d(x, z), d(z, y)\} = \max\{d(x, z), d(y, z)\} \leq d(x, y).$$

This shows that  $d(x, y) = \max\{d(x, z), d(y, z)\}$ . Thus either  $d(x, z) = d(x, y) \geq d(y, z)$  or  $d(y, z) = d(x, y) \geq d(x, z)$ .  $\square$

**Proposition 2.3.** Let  $(X, d)$  be an ultra-metric space. Let  $D_i$  be (open or closed) ball in  $X$  for  $i = 1, 2$ . If  $D_1 \cap D_2 \neq \emptyset$ , then either  $D_1 \subseteq D_2$  or  $D_2 \subseteq D_1$ .

*Proof.* Suppose that  $D_i$  has center  $x_i$  and radius  $r_i$  for  $i = 1, 2$ . Let  $y \in D_1 \cap D_2$ . We have

$$d(x_1, x_2) \leq \max\{d(x_1, y), d(y, x_2)\}.$$

Without loss of generality, we may assume that  $d(x_1, x_2) \leq d(x_1, y)$ . It follows that  $x_2 \in D_1$  since  $d(x_1, y) < r_1$  (or  $\leq r_1$ ).

If there exists  $z \in D_2 \setminus D_1$ , we claim that  $D_1 \subseteq D_2$ . We have  $d(x_1, z) > d(x_1, x_2)$ . Then by [Proposition 2.2](#),

$$r_1 \leq d(x_1, z) = d(x_2, z) \leq r_2.$$

In particular, if  $D_2$  is an open ball, then we have strict inequality  $r_1 < r_2$ . For any  $w \in D_1$ , we have

$$d(x_2, w) \leq \max\{d(x_2, x_1), d(x_1, w)\} \leq r_1 \leq r_2.$$

Thus  $w \in D_2$  whatever  $D_2$  is open or closed, and it shows that  $D_1 \subseteq D_2$ .  $\square$

**Proposition 2.4.** Let  $(X, d)$  be an ultra-metric space. Then both  $B(x, r)$  and  $E(x, r)$  are closed and open subsets of  $X$  for any  $x \in X$  and  $r > 0$ .

*Proof.* We show that the sphere  $S(x, r) := \{y \in X \mid d(x, y) = r\}$  is open in  $X$ . Note that if  $y \in S(x, r)$ , then for any  $r' < r$ , we have  $B(y, r') \cap E(x, r) \neq \emptyset$  and  $x \in E(x, r) \setminus B(y, r')$ . Thus by [Proposition 2.3](#), we have  $B(y, r') \subseteq E(x, r)$ . If  $B(y, r') \cap B(x, r) \neq \emptyset$ , then by [Proposition 2.3](#) again, we have  $B(y, r') \subseteq B(x, r)$ . However,  $y \in B(y, r') \setminus B(x, r)$ , a contradiction. Thus  $B(y, r') \subseteq E(x, r) \setminus B(x, r) = S(x, r)$ . It yields that  $S(x, r) = \bigcup_{y \in S(x, r)} B(y, r/2)$  is open in  $X$ .

Since  $E(x, r) = B(x, r) \cup S(x, r)$  and  $B(x, r) = E(x, r) \setminus S(x, r)$ , both  $B(x, r)$  and  $E(x, r)$  are open and closed in  $X$ .  $\square$

**Corollary 2.5.** Let  $(X, d)$  be an ultra-metric space. Then  $X$  is totally disconnected, i.e., the only connected subsets of  $X$  are the set with at most one point.

*Proof.* Suppose that  $S \subset X$  has at least two distinct points  $x, y \in S$ . Let  $r := d(x, y) > 0$ . Consider the open ball  $B(x, r/2)$ . By [Proposition 2.4](#),  $B(x, r/2)$  is both open and closed in  $X$ . Thus  $B(x, r/2) \cap S$  is both open and closed in  $S$ , however, it is non-empty and not equal to  $S$  since it contains  $x$  but not  $y$ . This shows that  $S$  is disconnected.  $\square$

## 3 Residue fields and reductions

### 3.1 Recover non-archimedean complete fields algebraically

In this subsection, let  $\mathbf{k}$  be a non-archimedean field. Set  $I_{r, <} := B(0, r)$  and  $I_{r, \leq} := E(0, r)$  for each  $r \in [0, 1]$ .

**Proposition 3.1.** The sets  $I_{r, <}$  and  $I_{r, \leq}$  are ideals of the ring of integers  $\mathbf{k}^\circ$ . Conversely, any ideal of  $\mathbf{k}^\circ$  is of the form  $I_{r, <}$  or  $I_{r, \leq}$  for some  $r \in (0, 1)$ .

*Proof.* Let  $I$  be an ideal of  $\mathbf{k}^\circ$ . Set  $r = \sup\{|a| : a \in I\}$  (resp.  $r = \max\{|a| : a \in I\}$  when the maximum exists). Then, by definition, we have  $I \subset I_{r, <}$  (resp.  $I \subset I_{r, \leq}$ ). For every  $x \in \mathbf{k}^\circ$  with

$|x| < r$  (resp.  $|x| \leq r$ ), there exists  $a \in I$  such that  $|x| \leq |a|$ . Thus,  $|x/a| \leq 1$  and so  $x/a \in \mathbf{k}^\circ$ . Since  $I$  is an ideal, we have  $x = (x/a)a \in I$ . Therefore,  $I_{r,<} \subset I$  (resp.  $I_{r,\leq} \subset I$ ).  $\square$

**Proposition 3.2.** Let  $I_r$  be either  $I_{r,<}$  or  $I_{r,\leq}$  for each  $r \in (0, 1)$ . Suppose  $\{r_n \in (0, 1)\}_{n \in \mathbb{N}}$  is a decreasing sequence converging to 0. Then the completion  $\widehat{\mathbf{k}}$  of  $\mathbf{k}$  is isomorphic to the projective limit

$$\widehat{\mathbf{k}}^\circ \cong \varprojlim_{n \in \mathbb{N}} \mathbf{k}^\circ / I_{r_n}.$$

*Proof.* For every  $x \in \widehat{\mathbf{k}}^\circ$ , there exists a cauchy sequence  $\{x_m\}_{m \in \mathbb{N}}$  in  $\mathbf{k}^\circ$  converging to  $x$ . Since  $\{r_n\}_{n \in \mathbb{N}}$  converges to 0, for each  $n \in \mathbb{N}$ , there exists  $M_n \in \mathbb{N}$  such that for all  $m, m' \geq M_n$ , we have  $|x_m - x_{m'}| < r_n$ . Thus, the sequence  $\{x_m + I_{r_n}\}_{m \in \mathbb{N}}$  is eventually constant in  $\mathbf{k}^\circ / I_{r_n}$ . Define a map

$$\Phi : \widehat{\mathbf{k}}^\circ \rightarrow \varprojlim_{n \in \mathbb{N}} \mathbf{k}^\circ / I_{r_n}, \quad x \mapsto \left( \lim_{m \rightarrow \infty} x_m + I_{r_n} \right)_{n \in \mathbb{N}}.$$

It is straightforward to verify that  $\Phi$  is a well-defined ring homomorphism.

Conversely, for every  $(a_n + I_{r_n})_{n \in \mathbb{N}} \in \varprojlim_{n \in \mathbb{N}} \mathbf{k}^\circ / I_{r_n}$ , we can choose a representative  $a_n \in \mathbf{k}^\circ$  for each  $n$ . We claim that the sequence  $\{a_n\}_{n \in \mathbb{N}}$  is a cauchy sequence in  $\mathbf{k}^\circ$ . Indeed, for every  $\varepsilon > 0$ , there exists  $N \in \mathbb{N}$  such that  $r_N < \varepsilon$ . For all  $m, n \geq N$ , since  $a_n + I_{r_n}$  maps to  $a_m + I_{r_m}$  under the natural projection, we have  $|a_n - a_m| < r_N < \varepsilon$ . Thus,  $\{a_n\}_{n \in \mathbb{N}}$  converges to some  $x \in \widehat{\mathbf{k}}^\circ$ . Easily see that the limit  $x$  is independent of the choice of representatives  $\{a_n\}_{n \in \mathbb{N}}$ . This gives a map

$$\Psi : \varprojlim_{n \in \mathbb{N}} \mathbf{k}^\circ / I_{r_n} \rightarrow \widehat{\mathbf{k}}^\circ, \quad (a_n + I_{r_n})_{n \in \mathbb{N}} \mapsto \lim_{n \rightarrow \infty} a_n.$$

Direct verification shows that  $\Psi = \Phi^{-1}$ .  $\square$

**Proposition 3.3.** Let  $\mathbf{k}$  be a non-archimedean field. Then  $\mathbf{k}^\circ$  is totally bounded iff  $\mathbf{k}^\circ / I_{r,<}$  and  $\mathbf{k}^\circ / I_{r,\leq}$  are finite for each  $r \in [0, 1]$ . Moreover, if  $\mathbf{k}$  is complete, then it is locally compact iff  $\mathbf{k}^\circ / I_r$  is finite for each  $r \in (0, 1)$ .

**Slogan** “Locally compact  $\iff$  pro-finite.”

*Proof.* We just prove the case for  $I_r = I_{r,<}$ . The case for  $I_r = I_{r,\leq}$  is similar.

Suppose that  $\mathbf{k}^\circ / I_r$  is finite for each  $r \in [0, 1]$ . Then for every  $\varepsilon > 0$ , there exists  $r \in (0, 1)$  such that  $r < \varepsilon$  and  $\mathbf{k}^\circ / I_r$  is finite. Let  $\{a_1 + I_r, \dots, a_n + I_r\}$  be the complete set of representatives of  $\mathbf{k}^\circ / I_r$ . Then the balls  $B(a_i, r)$  for  $i = 1, \dots, n$  cover  $\mathbf{k}^\circ$ .

Conversely, suppose that  $\mathbf{k}^\circ / I_r$  is infinite for some  $r \in [0, 1]$ . Then there exists an infinite set  $\{a_n\}$  with  $|a_n| \in [r, 1]$  such that their images in  $\mathbf{k}^\circ / I_r$  are distinct. In particular, for every  $m \neq n$ , we have  $|a_n - a_m| \geq r$ . Any subsequence of  $\{a_n\}$  is not cauchy. Thus,  $\mathbf{k}^\circ$  is not totally bounded.  $\square$

**Proposition 3.4.** The ring  $\mathbf{k}^\circ$  is noetherian iff  $\mathbf{k}$  is a discrete valuation field.

*Proof.* Note that  $|\mathbf{k}^\times| \subset \mathbb{R}_{>0}$  is a multiplicative subgroup. If  $\mathbf{k}$  is not a discrete valuation field, then  $|\mathbf{k}^\times|$  is dense in  $\mathbb{R}_{>0}$ . In particular, there exists a strictly ascending sequence  $r_n \in |\mathbf{k}^\times| \cap (0, 1)$ . Then the ideals  $I_{r_n,\leq}$  form a strictly ascending chain of ideals in  $\mathbf{k}^\circ$ .

The converse is standard since now  $\mathbf{k}^\circ$  is a discrete valuation ring.  $\square$

**Proposition 3.5.** Let  $\mathbf{k}$  be a complete non-archimedean field. Then  $\mathbf{k}$  is locally compact iff  $\mathbf{k}$  is a discrete valuation field and its residue field  $\mathcal{K}_{\mathbf{k}}$  is finite.

*Proof.* The necessity follows from Proposition 3.3. For the sufficiency, suppose that  $\mathbf{k}$  is a discrete valuation field whose residue field  $\mathcal{K}_{\mathbf{k}}$  is finite. Let  $\pi \in \mathbf{k}^\circ$  be a uniformizer. We only need to show that  $\mathbf{k}^\circ/\pi^n \mathbf{k}^\circ$  is finite for each  $n \in \mathbb{N}$ . Note that there is an isomorphism

$$\pi^{n-1} \mathbf{k}^\circ / \pi^n \mathbf{k}^\circ \cong \mathcal{K}_{\mathbf{k}}, \quad x + \pi^n \mathbf{k}^\circ \mapsto \overline{x/\pi^{n-1}}.$$

Thus, by induction on  $n$ , we conclude that  $\mathbf{k}^\circ/\pi^n \mathbf{k}^\circ$  is finite.  $\square$

## 3.2 Hensel's Lemma

**Theorem 3.6** (Hensel's lemma). Let  $\mathbf{k}$  be a complete non-archimedean field and  $F(T) \in \mathbf{k}^\circ[T]$  a monic polynomial. Suppose that the reduction  $f(T) \in \mathcal{K}_{\mathbf{k}}[T]$  of  $F(T)$  factors as

$$f(T) = g(T)h(T),$$

where  $g(T), h(T) \in \mathcal{K}_{\mathbf{k}}[T]$  are monic polynomials that are coprime in  $\mathcal{K}_{\mathbf{k}}[T]$ . Then there exist monic polynomials  $G(T), H(T) \in \mathbf{k}^\circ[T]$  such that

$$F(T) = G(T)H(T),$$

and the reductions of  $G(T), H(T)$  in  $\mathcal{K}_{\mathbf{k}}[T]$  are  $g(T), h(T)$  respectively.

*Proof.* Since  $\gcd(g, h) = 1$  in  $\mathcal{K}_{\mathbf{k}}[T]$ , there exist polynomials  $u(T), v(T) \in \mathcal{K}_{\mathbf{k}}[T]$  such that  $ug + vh = 1$  and  $\deg u < \deg h, \deg v < \deg g$ . Choose lifts  $G_0(T), H_0(T), U(T), V(T) \in \mathbf{k}^\circ[T]$  of  $g(T), h(T), u(T), v(T)$  respectively preserving their degrees such that  $G_0$  and  $H_0$  are monic. Then there exist  $r < 1$  such that

$$U(T)G_0(T) + V(T)H_0(T) \equiv 1 \pmod{I_r}, \quad F(T) - G_0(T)H_0(T) \equiv 0 \pmod{I_r},$$

where  $I_r = \{a \in \mathbf{k}^\circ : |a| < r\}$ .

We will construct a sequence of monic polynomials  $\{G_n(T)\}_{n \in \mathbb{N}}$  and  $\{H_n(T)\}_{n \in \mathbb{N}}$  in  $\mathbf{k}^\circ[T]$  such that for each  $n \in \mathbb{N}$ ,

$$G_n(T) \equiv G_{n-1}(T) \pmod{I_{r^n}}, \quad H_n(T) \equiv H_{n-1}(T) \pmod{I_{r^n}},$$

and

$$F(T) - G_n(T)H_n(T) \equiv 0 \pmod{I_{r^{n+1}}}.$$

If we have such sequences, then their coefficients converge in the complete ring  $\mathbf{k}^\circ$ . Let  $G(T)$  and  $H(T)$  be the limits of  $\{G_n(T)\}$  and  $\{H_n(T)\}$  respectively. Then we have  $F(T) = G(T)H(T)$  and the reductions of  $G(T), H(T)$  in  $\mathcal{K}_{\mathbf{k}}[T]$  are  $g(T), h(T)$  respectively.

The case  $n = 0$  is done by the above construction. Now suppose that we have constructed  $G_n(T)$  and  $H_n(T)$  for some  $n \geq 0$ . Since  $G_n - G_0 \equiv 0 \pmod{I_r}$  and  $H_n - H_0 \equiv 0 \pmod{I_r}$ , we have

$$UG_n + VH_n = UG_0 + VH_0 + U(G_n - G_0) + V(H_n - H_0) \equiv 1 \pmod{I_r}.$$



Set  $\Delta_n(T) = F(T) - G_n(T)H_n(T) \in I_{r^{n+1}}[T]$  and  $\epsilon_n = U\Delta_n, \delta_n = V\Delta_n \in I_{r^{n+1}}[T]$ . Then we have

$$\begin{aligned} (G_n + \epsilon_n)(H_n + \delta_n) - F_n &= G_n H_n + G_n \delta_n + H_n \epsilon_n + \epsilon_n \delta_n - F_n \\ &= (UG_n + VH_n - 1)\Delta_n + \epsilon_n \delta_n \in I_{r^{n+2}}[T]. \end{aligned}$$

Thus, we can set

$$G_{n+1}(T) = G_n(T) + \epsilon_n(T), \quad H_{n+1}(T) = H_n(T) + \delta_n(T).$$

□

**Corollary 3.7.** Let  $\mathbf{k}$  be a complete non-archimedean field and  $F(T) \in \mathbf{k}^\circ[T]$  a monic polynomial. Suppose that the reduction  $f(T) \in \mathbf{k}_{\mathbf{k}}[T]$  of  $F(T)$  has a simple root  $a \in \mathbf{k}_{\mathbf{k}}$ . Then there exists a root  $\alpha \in \mathbf{k}^\circ$  of  $F(T)$  whose reduction is  $a$ .

*Proof.* Since  $a$  is a simple root of  $f(T)$ , we have the factorization  $f(T) = (T - a)h(T)$  for some monic polynomial  $h(T) \in \mathbf{k}_{\mathbf{k}}[T]$  with  $h(a) \neq 0$ . Then the result follows from [Theorem 3.6](#). □

### 3.3 Newton polygons

Yang: To be filled.

## 4 Finite field extensions

### 4.1 Finite-dimensional vector space

**Proposition 4.1.** Let  $V$  be a finite-dimensional vector space over a complete non-archimedean field  $\mathbf{k}$ . Then all norms on  $V$  are equivalent. Yang: To be checked.

### 4.2 Finite field extensions

**Proposition 4.2.** Let  $\mathbf{k}$  be a complete non-archimedean field and  $\mathbf{l}$  a finite extension of  $\mathbf{k}$ . Then the absolute value on  $\mathbf{l}$  is uniquely determined by the absolute value on  $\mathbf{k}$ . Yang: To be checked.

**Proposition 4.3.** Let  $\mathbf{k}$  be an algebraically closed non-archimedean field. Then its completion  $\hat{\mathbf{k}}$  is also algebraically closed. Yang: To be checked.

## 5 Analytic functions

### 5.1 Failure of continuous and differentiable functions

**Definition 5.1.** Let  $(\mathbf{k}, \|\cdot\|)$  be a non-archimedean field and  $U \subset \mathbf{k}$  be an open subset. A function  $f : U \rightarrow \mathbf{k}$  is said to be *differentiable* at a point  $a \in U$  if the limit

$$f'(a) := \lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a}$$

exists in  $\mathbf{k}$ . If  $f$  is differentiable at every point in  $U$ , we say that  $f$  is differentiable on  $U$ . **Yang: to be revised.**

**Proposition 5.2.** Let  $(\mathbf{k}, \|\cdot\|)$  be a non-archimedean field. Then there exists a continuous function  $f : \mathbf{k} \rightarrow \mathbf{k}$  such that for any  $x, y \in \mathbf{k}$  with  $x \neq y$ , we have

$$\frac{f(x) - f(y)}{x - y} = 0.$$

### 5.2 Power series

**Proposition 5.3.** Let  $(\mathbf{k}, \|\cdot\|)$  be a complete non-archimedean field and  $\sum_{n=0}^{+\infty} a_n$  be a series in  $\mathbf{k}$ . Then the series  $\sum_{n=0}^{+\infty} a_n$  converges if and only if  $\lim_{n \rightarrow +\infty} a_n = 0$ . **Yang: To be checked.**

**Definition 5.4.** Let  $(\mathbf{k}, \|\cdot\|)$  be a complete non-archimedean field.

### 5.3 Analytic functions

As in the case of real analysis, we can define analytic functions over non-archimedean fields using power series.

## 6 Example: $p$ -adic fields

### 6.1 $p$ -adic fields

**Construction 6.1.** Let  $K$  be a number field and  $\mathfrak{p}$  be a prime ideal of the ring of integers  $\mathcal{O}_K$  of  $K$ . Considering the localization  $(\mathcal{O}_K)_{\mathfrak{p}}$  of  $\mathcal{O}_K$  at  $\mathfrak{p}$ , which is a discrete valuation ring, denote by  $v_{\mathfrak{p}} : K^{\times} \rightarrow \mathbb{Z}$  the corresponding discrete valuation. The  $p$ -adic absolute value on  $K$  associated to  $\mathfrak{p}$  is defined as

$$|x|_{\mathfrak{p}} := N(\mathfrak{p})^{-v_{\mathfrak{p}}(x)}, \quad \forall x \in K,$$

where  $N(\mathfrak{p}) := \#(\mathcal{O}_K/\mathfrak{p})$  is the norm of  $\mathfrak{p}$ .

The completion of  $K$  with respect to the  $p$ -adic absolute value  $|\cdot|_{\mathfrak{p}}$  is denoted by  $K_{\mathfrak{p}}$ , called the  $p$ -adic field.

One can just focus on the case  $K = \mathbb{Q}$  and  $\mathfrak{p} = (p)$  for a prime number  $p$ .

**Example 6.2.** Let  $p$  be a prime number. For every  $r \in \mathbb{Q}$ , we can write  $r$  as  $r = p^n \frac{a}{b}$ , where  $n \in \mathbb{Z}$  and  $a, b \in \mathbb{Z}$  are integers not divisible by  $p$ . The  $p$ -adic absolute value on  $\mathbb{Q}$  is defined as

$$|r|_p := p^{-n}.$$

The  $p$ -adic field  $\mathbb{Q}_p$  can be described concretely as follows:

$$\mathbb{Q}_p = \left\{ \sum_{i=n}^{+\infty} a_i p^i \mid n \in \mathbb{Z}, a_i \in \{0, 1, \dots, p-1\} \right\}.$$

For  $x = \sum_{i=n}^{+\infty} a_i p^i \in \mathbb{Q}_p$  with  $a_n \neq 0$ , its  $p$ -adic absolute value is given by  $|x|_p = p^{-n}$ . The operations of addition and multiplication on  $\mathbb{Q}_p$  are defined similarly as those on decimal expansions.

**Proposition 6.3.** The multiplicative group  $\mathbb{Q}_p^\times$  of the  $p$ -adic field  $\mathbb{Q}_p$  admits the following decomposition:

$$\mathbb{Q}_p^\times \cong p^\mathbb{Z} \times \mathbb{Z}_p^\times,$$

where  $p^\mathbb{Z} := \{p^n \mid n \in \mathbb{Z}\}$  and  $\mathbb{Z}_p^\times := \{x \in \mathbb{Q}_p \mid |x|_p = 1\}$  is the group of units of the ring of  $p$ -adic integers  $\mathbb{Z}_p$ . **Yang: To be checked.**

**Yang:** What is the relation between the finite extension of  $\mathbb{Q}_p$  and  $K_p$ ?

## 6.2 Completion

**Proposition 6.4.** The algebraic closure  $\overline{\mathbb{Q}_p}$  of  $\mathbb{Q}_p$  is not complete with respect to the extension of the  $p$ -adic absolute value  $|\cdot|_p$ .

**Construction 6.5.** Let  $p$  be a prime number. The field  $\mathbb{C}_p$  of  $p$ -adic complex numbers is defined as the completion of the algebraic closure of  $\mathbb{Q}_p$  with respect to the unique extension of the  $p$ -adic absolute value  $|\cdot|_p$  on  $\mathbb{Q}_p$ . The field  $\mathbb{C}_p$  is algebraically closed and complete with respect to  $|\cdot|_p$ .

**Yang: To be completed.**

**Proposition 6.6.** The field  $\mathbb{C}_p$  of  $p$ -adic complex numbers is not spherically complete.

**Construction 6.7.** Let  $p$  be a prime number. **Yang: We construct the spherically complete  $p$ -adic field  $\Omega_p$ .** **Yang: To be completed.**