
Non-archimedean Analysis

DRAFT

No Cover Image

Use `\coverimage{filename}` to add an image

Contents

1	Valuation fields	1
1.1	Absolute values and completion	1
1.2	Non-archimedean fields	3
2	Ultra-metric spaces	4
3	Algebraic structures of non-archimedean fields	5
3.1	Recover non-archimedean complete fields algebraically	5
3.2	Hensel's Lemma	7
3.3	Newton polygons	8
4	Finite field extensions	8
4.1	Finite-dimensional vector space	8
4.2	Finite field extensions	9
5	Analytic functions	9
5.1	Failure of continuous and differentiable functions	9
5.2	Power series	10
5.3	Analytic functions and maps	10
6	Example: p-adic fields	11
6.1	p-adic fields	11
6.2	Completion	11

1 Valuation fields

1.1 Absolute values and completion

Definition 1.1. Let \mathbf{k} be a field. An *absolute value* on \mathbf{k} is a function $\|\cdot\| : \mathbf{k} \rightarrow \mathbb{R}_{\geq 0}$ satisfying the following properties for all $x, y \in \mathbf{k}$:

- (a) $\|x\| = 0$ if and only if $x = 0$;
- (b) $\|xy\| = \|x\| \cdot \|y\|$;
- (c) $\|x + y\| \leq \|x\| + \|y\|$.

A field \mathbf{k} equipped with an absolute value $\|\cdot\|$ is called a *valuation field*.

Remark 1.2. Let \mathbf{k} be a field. Recall that a *valuation* on \mathbf{k} is a function $v : \mathbf{k}^\times \rightarrow \mathbb{R}$ such that

- $\forall x, y \in \mathbf{k}^\times, v(xy) = v(x) + v(y)$;

- $\forall x, y \in \mathbf{k}^\times, v(x + y) \geq \min\{v(x), v(y)\}$.

We can extend v to the whole field \mathbf{k} by defining $v(0) = +\infty$. Fix a real number $\varepsilon \in (0, 1)$. Then v induces an absolute value $|\cdot|_v : \mathbf{k} \rightarrow \mathbb{R}_+$ defined by $|x|_v = \varepsilon^{v(x)}$ for each $x \in \mathbf{k}$.

In some literature, the valuation v is called an *additive valuation* and the induced absolute value $|\cdot|_v$ is called a *multiplicative valuation*. In this note, the term *valuation* always refers to the additive valuation.

Example 1.3. Let \mathbf{k} be a field. The *trivial absolute value* on \mathbf{k} is defined as

$$\|x\| := \begin{cases} 0, & x = 0; \\ 1, & x \neq 0. \end{cases}$$

Definition 1.4. The (*multiplicative*) *valuation group* of a valuation field $(\mathbf{k}, \|\cdot\|)$ is defined as the subgroup of $\mathbb{R}_{>0}$ given by

$$|\mathbf{k}^\times| := \{\|x\| : x \in \mathbf{k}^\times\}.$$

Definition 1.5. Let \mathbf{k} be a field. Two absolute values $\|\cdot\|_1$ and $\|\cdot\|_2$ on \mathbf{k} are said to be *equivalent* if there exists a real number $c \in (0, 1)$ such that

$$\|x\|_1 = \|x\|_2^c, \quad \forall x \in \mathbf{k}.$$

Note that equivalent absolute values induce the same topology on the field \mathbf{k} . Moreover, the following lemma shows that the converse is also true.

Lemma 1.6. Let \mathbf{k} be a field and $\|\cdot\|_1, \|\cdot\|_2$ be two absolute values on \mathbf{k} . Then the following statements are equivalent:

- (a) $\|\cdot\|_1$ and $\|\cdot\|_2$ are equivalent;
- (b) $\|\cdot\|_1$ and $\|\cdot\|_2$ induce the same topology on \mathbf{k} ;
- (c) The unit disks $D_1 = \{x \in \mathbf{k} : \|x\|_1 < 1\}$ and $D_2 = \{x \in \mathbf{k} : \|x\|_2 < 1\}$ are the same.

Proof. The implications (a) \Rightarrow (b) is obvious. Now we prove (b) \Rightarrow (c). For any $x \in D_1$, we have $x^n \rightarrow 0$ as $n \rightarrow \infty$ under the absolute value $\|\cdot\|_1$ and thus under $\|\cdot\|_2$. Therefore, $\|x\|_2^n \rightarrow 0$ as $n \rightarrow \infty$, which implies that $\|x\|_2 < 1$, i.e., $x \in D_2$. Similarly, we can prove that $D_2 \subseteq D_1$.

Finally, we prove (c) \Rightarrow (a). If $\|\cdot\|_1$ is trivial, then $D_1 = \{0\}$ and thus $\|\cdot\|_2$ is also trivial. In this case, they are equivalent. Suppose that both $\|\cdot\|_1$ and $\|\cdot\|_2$ are non-trivial. Pick any $x, y \notin D_1 = D_2$. Then there exist real numbers $\alpha, \beta > 0$ such that $\|x\|_1 = \|x\|_2^\alpha$ and $\|y\|_1 = \|y\|_2^\beta$. Suppose the contrary that $\alpha \neq \beta$. Consider the domain $\Omega \subseteq \mathbb{Z}^2$ defined by

$$\begin{cases} n \log \|x\|_2 < m \log \|y\|_2; \\ n\alpha \log \|x\|_2 > m\beta \log \|y\|_2. \end{cases}$$

Since $\alpha \neq \beta$, the two lines defined by the equalities are not parallel. Thus Ω is non-empty. Pick $(n, m) \in \Omega$ and set $z := x^n y^{-m}$. Then we have $\|z\|_2 < 1$ and $\|z\|_1 > 1$, a contradiction. \square

Definition 1.7. Let $(\mathbf{k}, \|\cdot\|)$ be a valuation field. We say that \mathbf{k} is *complete* if the metric $d(x, y) := \|x - y\|$ makes \mathbf{k} a complete metric space.

Lemma 1.8. Let $(\mathbf{k}, \|\cdot\|)$ be a valuation field and $(\widehat{\mathbf{k}}, \|\cdot\|)$ its completion as a metric space. Then the operations of addition and multiplication on \mathbf{k} can be extended to $\widehat{\mathbf{k}}$ uniquely, making $(\widehat{\mathbf{k}}, \|\cdot\|)$ a complete valuation field containing \mathbf{k} as a dense subfield.

Proof. Simple analysis. □

Example 1.9. Let $|\cdot|_\infty$ be the usual absolute value on the field \mathbb{Q} of rational numbers. Then $(\mathbb{Q}, |\cdot|_\infty)$ is a valuation field. Its completion is the field \mathbb{R} of real numbers equipped with the usual absolute value.

Example 1.10. Let p be a prime number. For any non-zero rational number $x \in \mathbb{Q}$, we can write it as $x = p^n \frac{a}{b}$, where $n \in \mathbb{Z}$ and $a, b \in \mathbb{Z}$ are integers not divisible by p . The *p-adic absolute value* on \mathbb{Q} is defined as

$$|x|_p := \begin{cases} 0, & x = 0; \\ p^{-n}, & x = p^n \frac{a}{b} \text{ as above.} \end{cases}$$

Then $(\mathbb{Q}, |\cdot|_p)$ is a valuation field. Its completion is the field

$$\mathbb{Q}_p = \left\{ \sum_{n=k}^{+\infty} a_n p^n : k \in \mathbb{Z}, a_n \in \{0, 1, \dots, p-1\} \right\}$$

of *p*-adic numbers equipped with the *p*-adic absolute value; see [Yang: to be added.](#)

Unlike the real number field \mathbb{R} , even a valuation field is complete, we can not expect the theorem of nested intervals to hold.

Definition 1.11. A valuation field $(\mathbf{k}, \|\cdot\|)$ is called *spherically complete* if every decreasing sequence of closed balls in \mathbf{k} has a non-empty intersection.

Example 1.12. The field \mathbb{C}_p of *p*-adic complex numbers is not spherically complete, see [Yang: to be added.](#)

1.2 Non-archimedean fields

Definition 1.13. Let $(\mathbf{k}, \|\cdot\|)$ be a valuation field. We say that \mathbf{k} is *non-archimedean* if its absolute value $\|\cdot\|$ satisfies the *strong triangle inequality*:

$$\|x + y\| \leq \max\{\|x\|, \|y\|\}, \quad \forall x, y \in \mathbf{k}.$$

Otherwise, we say that \mathbf{k} is *archimedean*.

Let \mathbf{k} be a non-archimedean field. Then easily see that $\{x \in \mathbf{k} : \|x\| \leq 1\}$ is a subring of \mathbf{k} . Moreover, it is a local ring whose maximal ideal is $\{x \in \mathbf{k} : \|x\| < 1\}$.

Definition 1.14. Let \mathbf{k} be a non-archimedean field. The *ring of integers* of \mathbf{k} is defined as

$$\mathbf{k}^\circ := \{x \in \mathbf{k} : \|x\| \leq 1\}.$$

Its maximal ideal is

$$\mathbf{k}^{\circ\circ} := \{x \in \mathbf{k} : \|x\| < 1\}.$$

The *residue field* of \mathbf{k} is defined as

$$\kappa_{\mathbf{k}} := \tilde{\mathbf{k}} := \mathbf{k}^{\circ}/\mathbf{k}^{\circ\circ}.$$

Definition 1.15. Let \mathbf{k} be a non-archimedean field. The *residue absolute value* on the residue field $\kappa_{\mathbf{k}}$ is defined as

$$|x| := \inf_{y \in \varphi^{-1}(x)} \|y\|, \quad \forall x \in \kappa_{\mathbf{k}},$$

where $\varphi : \mathbf{k}^{\circ} \rightarrow \kappa_{\mathbf{k}}$ is the canonical projection.

Proposition 1.16. Let \mathbf{k} be a non-archimedean field. Then the residue absolute value on the residue field $\kappa_{\mathbf{k}}$ is trivial.

Proof. For any $x \in \kappa_{\mathbf{k}}$, if $x = 0$, then by definition $|x| = 0$. If $x \neq 0$, then $\forall y \in \varphi^{-1}(x)$, we have $y \in \mathbf{k}^{\circ} \setminus \mathbf{k}^{\circ\circ}$, i.e., $\|y\| = 1$. Thus by definition $|x| = 1$. \square

2 Ultra-metric spaces

We will use $B(x, r)$ (resp. $E(x, r)$) to denote the open ball (resp. closed ball) with center x and radius r .

Definition 2.1. A metric space (X, d) is called an *ultra-metric space* if its metric d satisfies the *strong triangle inequality*:

$$d(x, z) \leq \max\{d(x, y), d(y, z)\}, \quad \forall x, y, z \in X.$$

If $(\mathbf{k}, \|\cdot\|)$ is a non-archimedean field, then the metric $d(x, y) := \|x - y\|$ on \mathbf{k} makes (\mathbf{k}, d) an ultra-metric space.

Proposition 2.2. Let (X, d) be an ultra-metric space. Then for any $x, y, z \in X$, at least two of the three distances $d(x, y), d(y, z), d(z, x)$ are equal. And the third distance is less than or equal to the common value of the other two.

Proof. Suppose that $d(x, y) \geq d(y, z)$. By the strong triangle inequality, we have

$$d(x, z) \leq \max\{d(x, y), d(y, z)\} = d(x, y).$$

On the other hand, by the strong triangle inequality again, we have

$$d(x, y) \leq \max\{d(x, z), d(z, y)\} = \max\{d(x, z), d(y, z)\} \leq d(x, y).$$

This shows that $d(x, y) = \max\{d(x, z), d(y, z)\}$. Thus either $d(x, z) = d(x, y) \geq d(y, z)$ or $d(y, z) = d(x, y) \geq d(x, z)$. \square

Proposition 2.3. Let (X, d) be an ultra-metric space. Let D_i be (open or closed) ball in X for $i = 1, 2$. If $D_1 \cap D_2 \neq \emptyset$, then either $D_1 \subseteq D_2$ or $D_2 \subseteq D_1$.

Proof. Suppose that D_i has center x_i and radius r_i for $i = 1, 2$. Let $y \in D_1 \cap D_2$. We have

$$d(x_1, x_2) \leq \max\{d(x_1, y), d(y, x_2)\}.$$

Without loss of generality, we may assume that $d(x_1, x_2) \leq d(x_1, y)$. It follows that $x_2 \in D_1$ since $d(x_1, y) < r_1$ (or $\leq r_1$).

If there exists $z \in D_2 \setminus D_1$, we claim that $D_1 \subseteq D_2$. We have $d(x_1, z) > d(x_1, x_2)$. Then by [Proposition 2.2](#),

$$r_1 \leq d(x_1, z) = d(x_2, z) \leq r_2.$$

In particular, if D_2 is an open ball, then we have strict inequality $r_1 < r_2$. For any $w \in D_1$, we have

$$d(x_2, w) \leq \max\{d(x_2, x_1), d(x_1, w)\} \leq r_1 \leq r_2.$$

Thus $w \in D_2$ whatever D_2 is open or closed, and it shows that $D_1 \subseteq D_2$. \square

Proposition 2.4. Let (X, d) be an ultra-metric space. Then both $B(x, r)$ and $E(x, r)$ are closed and open subsets of X for any $x \in X$ and $r > 0$.

Proof. We show that the sphere $S(x, r) := \{y \in X \mid d(x, y) = r\}$ is open in X . Note that if $y \in S(x, r)$, then for any $r' < r$, we have $B(y, r') \cap E(x, r) \neq \emptyset$ and $x \in E(x, r) \setminus B(y, r')$. Thus by [Proposition 2.3](#), we have $B(y, r') \subseteq E(x, r)$. If $B(y, r') \cap B(x, r) \neq \emptyset$, then by [Proposition 2.3](#) again, we have $B(y, r') \subseteq B(x, r)$. However, $y \in B(y, r') \setminus B(x, r)$, a contradiction. Thus $B(y, r') \subseteq E(x, r) \setminus B(x, r) = S(x, r)$. It yields that $S(x, r) = \bigcup_{y \in S(x, r)} B(y, r/2)$ is open in X .

Since $E(x, r) = B(x, r) \cup S(x, r)$ and $B(x, r) = E(x, r) \setminus S(x, r)$, both $B(x, r)$ and $E(x, r)$ are open and closed in X . \square

Corollary 2.5. Let (X, d) be an ultra-metric space. Then X is totally disconnected, i.e., the only connected subsets of X are the set with at most one point.

Proof. Suppose that $S \subset X$ has at least two distinct points $x, y \in S$. Let $r := d(x, y) > 0$. Consider the open ball $B(x, r/2)$. By [Proposition 2.4](#), $B(x, r/2)$ is both open and closed in X . Thus $B(x, r/2) \cap S$ is both open and closed in S , however, it is non-empty and not equal to S since it contains x but not y . This shows that S is disconnected. \square

3 Algebraic structures of non-archimedean fields

3.1 Recover non-archimedean complete fields algebraically

In this subsection, let \mathbf{k} be a non-archimedean field. Set $I_{r, <} := B(0, r)$ and $I_{r, \leq} := E(0, r)$ for each $r \in [0, 1]$.

Proposition 3.1. The sets $I_{r, <}$ and $I_{r, \leq}$ are ideals of the ring of integers \mathbf{k}° . Conversely, any ideal of \mathbf{k}° is of the form $I_{r, <}$ or $I_{r, \leq}$ for some $r \in (0, 1)$.

Proof. Let I be an ideal of \mathbf{k}° . Set $r = \sup\{|a| : a \in I\}$ (resp. $r = \max\{|a| : a \in I\}$ when the maximum exists). Then, by definition, we have $I \subset I_{r, <}$ (resp. $I \subset I_{r, \leq}$). For every $x \in \mathbf{k}^\circ$ with

$|x| < r$ (resp. $|x| \leq r$), there exists $a \in I$ such that $|x| \leq |a|$. Thus, $|x/a| \leq 1$ and so $x/a \in \mathbf{k}^\circ$. Since I is an ideal, we have $x = (x/a)a \in I$. Therefore, $I_{r,<} \subset I$ (resp. $I_{r,\leq} \subset I$). \square

Proposition 3.2. Let I_r be either $I_{r,<}$ or $I_{r,\leq}$ for each $r \in (0, 1)$. Suppose $\{r_n \in (0, 1)\}_{n \in \mathbb{N}}$ is a decreasing sequence converging to 0. Then the completion $\hat{\mathbf{k}}$ of \mathbf{k} is isomorphic to the projective limit

$$\hat{\mathbf{k}}^\circ \cong \varprojlim_{n \in \mathbb{N}} \mathbf{k}^\circ / I_{r_n}.$$

Proof. For every $x \in \hat{\mathbf{k}}^\circ$, there exists a cauchy sequence $\{x_m\}_{m \in \mathbb{N}}$ in \mathbf{k}° converging to x . Since $\{r_n\}_{n \in \mathbb{N}}$ converges to 0, for each $n \in \mathbb{N}$, there exists $M_n \in \mathbb{N}$ such that for all $m, m' \geq M_n$, we have $|x_m - x_{m'}| < r_n$. Thus, the sequence $\{x_m + I_{r_n}\}_{m \in \mathbb{N}}$ is eventually constant in $\mathbf{k}^\circ / I_{r_n}$. Define a map

$$\Phi : \hat{\mathbf{k}}^\circ \rightarrow \varprojlim_{n \in \mathbb{N}} \mathbf{k}^\circ / I_{r_n}, \quad x \mapsto \left(\lim_{m \rightarrow \infty} x_m + I_{r_n} \right)_{n \in \mathbb{N}}.$$

It is straightforward to verify that Φ is a well-defined ring homomorphism.

Conversely, for every $(a_n + I_{r_n})_{n \in \mathbb{N}} \in \varprojlim_{n \in \mathbb{N}} \mathbf{k}^\circ / I_{r_n}$, we can choose a representative $a_n \in \mathbf{k}^\circ$ for each n . We claim that the sequence $\{a_n\}_{n \in \mathbb{N}}$ is a cauchy sequence in \mathbf{k}° . Indeed, for every $\varepsilon > 0$, there exists $N \in \mathbb{N}$ such that $r_N < \varepsilon$. For all $m, n \geq N$, since $a_n + I_{r_n}$ maps to $a_m + I_{r_m}$ under the natural projection, we have $|a_n - a_m| < r_N < \varepsilon$. Thus, $\{a_n\}_{n \in \mathbb{N}}$ converges to some $x \in \hat{\mathbf{k}}^\circ$. Easily see that the limit x is independent of the choice of representatives $\{a_n\}_{n \in \mathbb{N}}$. This gives a map

$$\Psi : \varprojlim_{n \in \mathbb{N}} \mathbf{k}^\circ / I_{r_n} \rightarrow \hat{\mathbf{k}}^\circ, \quad (a_n + I_{r_n})_{n \in \mathbb{N}} \mapsto \lim_{n \rightarrow \infty} a_n.$$

Direct verification shows that $\Psi = \Phi^{-1}$. \square

Proposition 3.3. Let \mathbf{k} be a non-archimedean field. Then \mathbf{k}° is totally bounded iff $\mathbf{k}^\circ / I_{r,<}$ and $\mathbf{k}^\circ / I_{r,\leq}$ are finite for each $r \in [0, 1]$. Moreover, if \mathbf{k} is complete, then it is locally compact iff \mathbf{k}° / I_r is finite for each $r \in (0, 1)$.

Slogan “Locally compact \iff pro-finite.”

Proof. We just prove the case for $I_r = I_{r,<}$. The case for $I_r = I_{r,\leq}$ is similar.

Suppose that \mathbf{k}° / I_r is finite for each $r \in [0, 1]$. Then for every $\varepsilon > 0$, there exists $r \in (0, 1)$ such that $r < \varepsilon$ and \mathbf{k}° / I_r is finite. Let $\{a_1 + I_r, \dots, a_n + I_r\}$ be the complete set of representatives of \mathbf{k}° / I_r . Then the balls $B(a_i, r)$ for $i = 1, \dots, n$ cover \mathbf{k}° .

Conversely, suppose that \mathbf{k}° / I_r is infinite for some $r \in [0, 1]$. Then there exists an infinite set $\{a_n\}$ with $|a_n| \in [r, 1]$ such that their images in \mathbf{k}° / I_r are distinct. In particular, for every $m \neq n$, we have $|a_n - a_m| \geq r$. Any subsequence of $\{a_n\}$ is not cauchy. Thus, \mathbf{k}° is not totally bounded. \square

Proposition 3.4. The ring \mathbf{k}° is noetherian iff \mathbf{k} is a discrete valuation field.

Proof. Note that $|\mathbf{k}^\times| \subset \mathbb{R}_{>0}$ is a multiplicative subgroup. If \mathbf{k} is not a discrete valuation field, then $|\mathbf{k}^\times|$ is dense in $\mathbb{R}_{>0}$. In particular, there exists a strictly ascending sequence $r_n \in |\mathbf{k}^\times| \cap (0, 1)$. Then the ideals $I_{r_n,\leq}$ form a strictly ascending chain of ideals in \mathbf{k}° .

The converse is standard since now \mathbf{k}° is a discrete valuation ring. \square

Proposition 3.5. Let \mathbf{k} be a complete non-archimedean field. Then \mathbf{k} is locally compact iff \mathbf{k} is a discrete valuation field and its residue field $\mathcal{K}_{\mathbf{k}}$ is finite.

Proof. The necessity follows from Proposition 3.3. For the sufficiency, suppose that \mathbf{k} is a discrete valuation field whose residue field $\mathcal{K}_{\mathbf{k}}$ is finite. Let $\pi \in \mathbf{k}^\circ$ be a uniformizer. We only need to show that $\mathbf{k}^\circ/\pi^n \mathbf{k}^\circ$ is finite for each $n \in \mathbb{N}$. Note that there is an isomorphism

$$\pi^{n-1} \mathbf{k}^\circ / \pi^n \mathbf{k}^\circ \cong \mathcal{K}_{\mathbf{k}}, \quad x + \pi^n \mathbf{k}^\circ \mapsto \overline{x/\pi^{n-1}}.$$

Thus, by induction on n , we conclude that $\mathbf{k}^\circ/\pi^n \mathbf{k}^\circ$ is finite. \square

3.2 Hensel's Lemma

Theorem 3.6 (Hensel's lemma). Let \mathbf{k} be a complete non-archimedean field and $F(T) \in \mathbf{k}^\circ[T]$ a monic polynomial. Suppose that the reduction $f(T) \in \mathcal{K}_{\mathbf{k}}[T]$ of $F(T)$ factors as

$$f(T) = g(T)h(T),$$

where $g(T), h(T) \in \mathcal{K}_{\mathbf{k}}[T]$ are monic polynomials that are coprime in $\mathcal{K}_{\mathbf{k}}[T]$. Then there exist monic polynomials $G(T), H(T) \in \mathbf{k}^\circ[T]$ such that

$$F(T) = G(T)H(T),$$

and the reductions of $G(T), H(T)$ in $\mathcal{K}_{\mathbf{k}}[T]$ are $g(T), h(T)$ respectively.

Proof. Since $\gcd(g, h) = 1$ in $\mathcal{K}_{\mathbf{k}}[T]$, there exist polynomials $u(T), v(T) \in \mathcal{K}_{\mathbf{k}}[T]$ such that $ug + vh = 1$ and $\deg u < \deg h, \deg v < \deg g$. Choose lifts $G_0(T), H_0(T), U(T), V(T) \in \mathbf{k}^\circ[T]$ of $g(T), h(T), u(T), v(T)$ respectively preserving their degrees such that G_0 and H_0 are monic. Then there exist $r < 1$ such that

$$U(T)G_0(T) + V(T)H_0(T) \equiv 1 \pmod{I_r}, \quad F(T) - G_0(T)H_0(T) \equiv 0 \pmod{I_r},$$

where $I_r = \{a \in \mathbf{k}^\circ : |a| < r\}$.

We will construct a sequence of monic polynomials $\{G_n(T)\}_{n \in \mathbb{N}}$ and $\{H_n(T)\}_{n \in \mathbb{N}}$ in $\mathbf{k}^\circ[T]$ such that for each $n \in \mathbb{N}$,

$$G_n(T) \equiv G_{n-1}(T) \pmod{I_{r^n}}, \quad H_n(T) \equiv H_{n-1}(T) \pmod{I_{r^n}},$$

and

$$F(T) - G_n(T)H_n(T) \equiv 0 \pmod{I_{r^{n+1}}}.$$

If we have such sequences, then their coefficients converge in the complete ring \mathbf{k}° . Let $G(T)$ and $H(T)$ be the limits of $\{G_n(T)\}$ and $\{H_n(T)\}$ respectively. Then we have $F(T) = G(T)H(T)$ and the reductions of $G(T), H(T)$ in $\mathcal{K}_{\mathbf{k}}[T]$ are $g(T), h(T)$ respectively.

The case $n = 0$ is done by the above construction. Now suppose that we have constructed $G_n(T)$ and $H_n(T)$ for some $n \geq 0$. Since $G_n - G_0 \equiv 0 \pmod{I_r}$ and $H_n - H_0 \equiv 0 \pmod{I_r}$, we have

$$UG_n + VH_n = UG_0 + VH_0 + U(G_n - G_0) + V(H_n - H_0) \equiv 1 \pmod{I_r}.$$

Set $\Delta_n(T) = F(T) - G_n(T)H_n(T) \in I_{r^{n+1}}[T]$ and $\epsilon_n = U\Delta_n, \delta_n = V\Delta_n \in I_{r^{n+1}}[T]$. Then we have

$$\begin{aligned} (G_n + \epsilon_n)(H_n + \delta_n) - F_n &= G_n H_n + G_n \delta_n + H_n \epsilon_n + \epsilon_n \delta_n - F_n \\ &= (UG_n + VH_n - 1)\Delta_n + \epsilon_n \delta_n \in I_{r^{n+2}}[T]. \end{aligned}$$

Thus, we can set

$$G_{n+1}(T) = G_n(T) + \epsilon_n(T), \quad H_{n+1}(T) = H_n(T) + \delta_n(T).$$

□

Corollary 3.7. Let \mathbf{k} be a complete non-archimedean field and $F(T) \in \mathbf{k}^\circ[T]$ a monic polynomial. Suppose that the reduction $f(T) \in \mathbf{k}_{\mathbf{k}}[T]$ of $F(T)$ has a simple root $a \in \mathbf{k}_{\mathbf{k}}$. Then there exists a root $\alpha \in \mathbf{k}^\circ$ of $F(T)$ whose reduction is a .

Proof. Since a is a simple root of $f(T)$, we have the factorization $f(T) = (T - a)h(T)$ for some monic polynomial $h(T) \in \mathbf{k}_{\mathbf{k}}[T]$ with $h(a) \neq 0$. Then the result follows from [Theorem 3.6](#). □

3.3 Newton polygons

Yang: To be filled.

4 Finite field extensions

4.1 Finite-dimensional vector space

Definition 4.1. Let \mathbf{k} be a valuation field and V a vector space over \mathbf{k} . A *norm* on V is a function $\|\cdot\| : V \rightarrow \mathbb{R}_{\geq 0}$ satisfying the following properties for all $x, y \in V$ and $a \in \mathbf{k}$:

- (a) $\|x\| = 0$ if and only if $x = 0$;
- (b) $\|ax\| = |a| \cdot \|x\|$;
- (c) $\|x + y\| \leq \|x\| + \|y\|$.

Definition 4.2. Let \mathbf{k} be a valuation field and V a vector space over \mathbf{k} . Two norms $\|\cdot\|_1$ and $\|\cdot\|_2$ on V are said to be *equivalent* if there exist positive constants $C_1, C_2 > 0$ such that for all $x \in V$,

$$C_1 \|x\|_1 \leq \|x\|_2 \leq C_2 \|x\|_1.$$

Lemma 4.3. Let \mathbf{k} be a valuation field and V a vector space over \mathbf{k} . Two norms $\|\cdot\|_1$ and $\|\cdot\|_2$ on V are equivalent if and only if they induce the same topology on V .

Proof. Yang: To be added. □

Proposition 4.4. Let V be a finite-dimensional vector space over a complete non-archimedean field \mathbf{k} . Then all norms on V are equivalent. Yang: To be checked.

Proof. Yang: To be added. □

Proposition 4.5. Let V be a finite-dimensional vector space over a complete valuation field \mathbf{k} . Then V is complete with respect to any norm on V .

4.2 Finite field extensions

Construction 4.6. Let \mathbf{k} be a valuation field and \mathbf{l} a finite extension of \mathbf{k} with degree $n = [\mathbf{l} : \mathbf{k}]$. For any $a \in \mathbf{l}$, define

$$|a|_{\mathbf{l}} := |N_{\mathbf{l}/\mathbf{k}}(a)|_{\mathbf{k}}^{1/n},$$

where $N_{\mathbf{l}/\mathbf{k}}(a)$ is the norm of a from \mathbf{l} to \mathbf{k} . By Lemma 4.7, $|\cdot|_{\mathbf{l}}$ is an absolute value on \mathbf{l} extending the absolute value $|\cdot|_{\mathbf{k}}$ on \mathbf{k} . Yang: To be checked.

Lemma 4.7. Let \mathbf{k} be a valuation field and \mathbf{l} a finite extension of \mathbf{k} . Then the function $|\cdot|_{\mathbf{l}}$ defined in Construction 4.6 is an absolute value on \mathbf{l} extending the absolute value on \mathbf{k} . Moreover, if \mathbf{k} is non-archimedean, then so is \mathbf{l} .

Proof. Yang: To be added. □

Proposition 4.8. Let \mathbf{k} be a complete non-archimedean field and \mathbf{l} a finite extension of \mathbf{k} . Then the absolute value on \mathbf{l} is uniquely determined by the absolute value on \mathbf{k} . Yang: To be checked.

Proof. Yang: To be added. □

Remark 4.9. Yang: I want to discuss some compatibility of extension and completion.

Proposition 4.10. Let \mathbf{k} be an algebraically closed non-archimedean field. Then its completion $\hat{\mathbf{k}}$ is also algebraically closed. Yang: To be checked.

Proof. Yang: To be added. □

5 Analytic functions

5.1 Failure of continuous and differentiable functions

Definition 5.1. Let $(\mathbf{k}, \|\cdot\|)$ be a non-archimedean field and $U \subset \mathbf{k}$ be an open subset. A function $f : U \rightarrow \mathbf{k}$ is said to be *differentiable* at a point $a \in U$ if the limit

$$f'(a) := \lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a}$$

exists in \mathbf{k} . If f is differentiable at every point in U , we say that f is differentiable on U . Yang: to be revised.

Proposition 5.2. Let $(\mathbf{k}, \|\cdot\|)$ be a non-archimedean field. Then there exists a continuous function $f : \mathbf{k} \rightarrow \mathbf{k}$ such that for any $x, y \in \mathbf{k}$ with $x \neq y$, we have

$$\frac{f(x) - f(y)}{x - y} = 0.$$

5.2 Power series

Lemma 5.3. Let $(\mathbf{k}, \|\cdot\|)$ be a non-archimedean field and $\sum_{n=0}^{+\infty} a_n$ be a series in \mathbf{k} . Then the series $\sum_{n=0}^{+\infty} a_n$ converges if and only if $\lim_{n \rightarrow +\infty} a_n = 0$. **Yang: To be checked.**

Proposition 5.4. Let $(\mathbf{k}, \|\cdot\|)$ be a non-archimedean field and $\sum_{n=0}^{+\infty} a_n(x-c)^n$ be a power series in \mathbf{k} . Then there exists a radius of convergence $R \in \mathbb{R}_{\geq 0} \cup \{+\infty\}$ such that the series converges for all $x \in \mathbf{k}$ with $\|x-c\| < R$ and diverges for all $x \in \mathbf{k}$ with $\|x-c\| > R$. **Yang: To be revised.**

Proposition 5.5. Let $(\mathbf{k}, \|\cdot\|)$ be a complete non-archimedean field and $\sum_{n=0}^{+\infty} a_n$ be a series in \mathbf{k} . Then the series $\sum_{n=0}^{+\infty} a_n$ converges if and only if $\lim_{n \rightarrow +\infty} a_n = 0$. **Yang: To be checked.**

Definition 5.6. Let $(\mathbf{k}, \|\cdot\|)$ be a complete non-archimedean field.

Proposition 5.7. Let $(\mathbf{k}, \|\cdot\|)$ be a complete non-archimedean field. Then the norm on the Tate algebra $\mathbf{k}\langle x_1, \dots, x_n \rangle$ coincides with the supremum norm on the closed unit polydisc in \mathbf{k}^n . **Yang: To be checked.**

5.3 Analytic functions and maps

As in the case of real analysis, we can define analytic functions over non-archimedean fields using power series.

Definition 5.8. Let $(\mathbf{k}, \|\cdot\|)$ be a complete non-archimedean field and $U \subset \mathbf{k}$ be an open subset. A function $f : U \rightarrow \mathbf{k}$ is said to be *analytic* at a point $c \in U$ if there exists a power series $\sum_{n=0}^{+\infty} a_n(x-c)^n$ that converges to $f(x)$ for all x in some neighborhood of c . If f is analytic at every point in U , we say that f is analytic on U . **Yang: to be revised.**

Theorem 5.9. Let $(\mathbf{k}, \|\cdot\|)$ be a complete non-archimedean field and $U \subset \mathbf{k}$ be an open subset. If $f : U \rightarrow \mathbf{k}$ is an analytic function, then f is locally Lipschitz continuous on U . **Yang: To be checked.**

Theorem 5.10. Let $(\mathbf{k}, \|\cdot\|)$ be a complete non-archimedean field and $U \subset \mathbf{k}$ be an open subset. If $f : U \rightarrow \mathbf{k}$ is an analytic function, then f satisfies the maximum modulus principle, i.e., if there exists a point $x_0 \in U$ such that $\|f(x_0)\| \geq \|f(x)\|$ for all $x \in U$, then f is constant on U . **Yang: To be checked.**

6 Example: p -adic fields

6.1 p -adic fields

Construction 6.1. Let K be a number field and \mathfrak{p} be a prime ideal of the ring of integers \mathcal{O}_K of K . Considering the localization $(\mathcal{O}_K)_{\mathfrak{p}}$ of \mathcal{O}_K at \mathfrak{p} , which is a discrete valuation ring, denote by $v_{\mathfrak{p}} : K^{\times} \rightarrow \mathbb{Z}$ the corresponding discrete valuation. The p -adic absolute value on K associated to \mathfrak{p} is defined as

$$|x|_{\mathfrak{p}} := N(\mathfrak{p})^{-v_{\mathfrak{p}}(x)}, \quad \forall x \in K,$$

where $N(\mathfrak{p}) := \#(\mathcal{O}_K/\mathfrak{p})$ is the norm of \mathfrak{p} .

The completion of K with respect to the p -adic absolute value $|\cdot|_{\mathfrak{p}}$ is denoted by $K_{\mathfrak{p}}$, called the p -adic field.

One can just focus on the case $K = \mathbb{Q}$ and $\mathfrak{p} = (p)$ for a prime number p .

Example 6.2. Let p be a prime number. For every $r \in \mathbb{Q}$, we can write r as $r = p^n \frac{a}{b}$, where $n \in \mathbb{Z}$ and $a, b \in \mathbb{Z}$ are integers not divisible by p . The p -adic absolute value on \mathbb{Q} is defined as

$$|r|_p := p^{-n}.$$

The p -adic field \mathbb{Q}_p can be described concretely as follows:

$$\mathbb{Q}_p = \left\{ \sum_{i=n}^{+\infty} a_i p^i \mid n \in \mathbb{Z}, a_i \in \{0, 1, \dots, p-1\} \right\}.$$

For $x = \sum_{i=n}^{+\infty} a_i p^i \in \mathbb{Q}_p$ with $a_n \neq 0$, its p -adic absolute value is given by $|x|_p = p^{-n}$. The operations of addition and multiplication on \mathbb{Q}_p are defined similarly as those on decimal expansions.

Proposition 6.3. The multiplicative group \mathbb{Q}_p^{\times} of the p -adic field \mathbb{Q}_p admits the following decomposition:

$$\mathbb{Q}_p^{\times} \cong p^{\mathbb{Z}} \times \mathbb{Z}_p^{\times},$$

where $p^{\mathbb{Z}} := \{p^n \mid n \in \mathbb{Z}\}$ and $\mathbb{Z}_p^{\times} := \{x \in \mathbb{Q}_p \mid |x|_p = 1\}$ is the group of units of the ring of p -adic integers \mathbb{Z}_p . Yang: To be checked.

Yang: What is the relation between the finite extension of \mathbb{Q}_p and K_p ?

6.2 Completion

Proposition 6.4. The algebraic closure $\overline{\mathbb{Q}_p}$ of \mathbb{Q}_p is not complete with respect to the extension of the p -adic absolute value $|\cdot|_p$.

Construction 6.5. Let p be a prime number. The field \mathbb{C}_p of p -adic complex numbers is defined as the completion of the algebraic closure of \mathbb{Q}_p with respect to the unique extension of the p -adic absolute value $|\cdot|_p$ on \mathbb{Q}_p . The field \mathbb{C}_p is algebraically closed and complete with respect to $|\cdot|_p$.

Yang: To be completed.

Proposition 6.6. The field \mathbb{C}_p of p -adic complex numbers is not spherically complete.

Construction 6.7. Let p be a prime number. *Yang: We construct the spherically complete p -adic field Ω_p . Yang: To be completed.*

DRAFT
