

Spectrum

1 Definition

Definition 1. Let R be a Banach ring. The *spectrum* $\mathcal{M}(R)$ of R is defined as the set of all multiplicative semi-norms on R that are bounded with respect to the given norm on R . For every point $x \in \mathcal{M}(R)$, we denote the corresponding multiplicative semi-norm by $|\cdot|_x$. We equip $\mathcal{M}(R)$ with the weakest topology such that for each $f \in R$, the evaluation map $\mathcal{M}(R) \rightarrow \mathbb{R}_{\geq 0}$, defined by $x \mapsto |f|_x =: f(x)$, is continuous.

Definition 2. Let $\varphi : R \rightarrow S$ be a bounded ring homomorphism of Banach rings. The *pullback* map $\mathcal{M}(\varphi) : \mathcal{M}(S) \rightarrow \mathcal{M}(R)$ is defined by $\mathcal{M}(\varphi)(x) = x \circ \varphi : f \mapsto |\varphi(f)|_x$ for each $x \in \mathcal{M}(S)$.

Definition 3. Let R be a Banach ring. A *character* of R is a bounded ring homomorphism $\chi : R \rightarrow K$, where K is a completed field. Two characters $\chi_1 : R \rightarrow K_1$ and $\chi_2 : R \rightarrow K_2$ are said to be *equivalent* if there exists a commutative diagram of bounded ring homomorphisms

$$\begin{array}{ccccc} & & R & & \\ & \swarrow \chi_1 & \downarrow & \searrow \chi_2 & \\ K_1 & \longleftarrow & K & \longrightarrow & K_2 \end{array}$$

for some completed field K .

Proposition 4. Let R be a Banach ring. The spectrum $\mathcal{M}(R)$ is in bijection with the equivalence classes of characters of R .

Proof. Yang: To be completed □

Proposition 5. Let R be a Banach ring. For each $x \in \mathcal{M}(R)$, let \wp_x be the kernel of the multiplicative semi-norm $|\cdot|_x$. Then \wp_x is a closed prime ideal of R , and $x \mapsto \wp_x$ defines a continuous map from $\mathcal{M}(R)$ to $\text{Spec}(R)$ equipped with the Zariski topology.

Proof. Yang: To be completed □

Definition 6. Let R be a Banach ring. For each $x \in \mathcal{M}(R)$, the *completed residue field* at the point x is defined as the completion of the residue field $\kappa(x) = \text{Frac}(R/\wp_x)$ with respect to the multiplicative norm induced by the semi-norm $|\cdot|_x$, denoted by $\mathcal{H}(x)$.

Definition 7. Let R be a Banach ring. The *Gel'fand transform* of R is the bounded ring homomorphism

$$\Gamma : R \rightarrow \prod_{x \in \mathcal{M}(R)} \mathcal{H}(x), \quad f \mapsto (f(x))_{x \in \mathcal{M}(R)},$$

where the norm on the product $\prod_{x \in \mathcal{M}(R)} \mathcal{H}(x)$ is given by the supremum norm.

Proposition 8. The Gel'fand transform $\Gamma : R \rightarrow \prod_{x \in \mathcal{M}(R)} \mathcal{H}(x)$ of a Banach ring R factors through the uniformization R^u of R , and the induced map $R^u \rightarrow \prod_{x \in \mathcal{M}(R)} \mathcal{H}(x)$ is an isometric embedding.

Yang: To be checked.

Theorem 9. Let R be a Banach ring. The spectrum $\mathcal{M}(R)$ is a nonempty compact Hausdorff space.

Proof. Yang: To be continued. \square

Lemma 10. Let $\{K_i\}_{i \in I}$ be a family of completed fields. Consider the Banach ring $R = \prod_{i \in I} K_i$ equipped with the product norm. The spectrum $\mathcal{M}(R)$ is homeomorphic to the Stone-Čech compactification of the discrete space I .

Remark 11. The Stone-Čech compactification of a discrete space is the largest compact Hausdorff space in which the original space can be densely embedded. Yang: To be checked.

Proposition 12. Let K/k be a Galois extension of complete fields, and let R be a Banach k -algebra. The Galois group $\text{Gal}(K/k)$ acts on the spectrum $\mathcal{M}(R \hat{\otimes}_k K)$ via

$$g \cdot x : f \mapsto |(1 \otimes g^{-1})(f)|_x$$

for each $g \in \text{Gal}(K/k)$, $x \in \mathcal{M}(R \hat{\otimes}_k K)$ and $f \in R \hat{\otimes}_k K$. Moreover, the natural map $\mathcal{M}(R \hat{\otimes}_k K) \rightarrow \mathcal{M}(R)$ induces a homeomorphism

$$\mathcal{M}(R \hat{\otimes}_k K) / \text{Gal}(K/k) \xrightarrow{\sim} \mathcal{M}(R).$$

Yang: To be checked.

2 Examples

Example 13. Let $(\mathbf{k}, |\cdot|)$ be a complete valuation field. The spectrum $\mathcal{M}(\mathbf{k})$ consists of a single point corresponding to the given absolute value $|\cdot|$ on \mathbf{k} . Yang: To be checked.

Example 14. Consider the Banach ring $(\mathbb{Z}, \|\cdot\|)$ with $\|\cdot\| = |\cdot|_\infty$ is the usual absolute value norm on \mathbb{Z} . Let $|\cdot|_p$ denote the p -adic norm for each prime number p , i.e., $|n|_p = p^{-v_p(n)}$ for each $n \in \mathbb{Z}$, where $v_p(n)$ is the p -adic valuation of n . The spectrum

$$\mathcal{M}(\mathbb{Z}) = \{|\cdot|_\infty^\varepsilon : \varepsilon \in (0, 1]\} \cup \{|\cdot|_p^\alpha : p \text{ is prime}, \alpha \in (0, \infty)\} \cup \{|\cdot|_0\},$$

where $|a|_p^\infty := \lim_{\alpha \rightarrow \infty} |a|_p^\alpha$ for each $a \in \mathbb{Z}$ and $|\cdot|_0$ is the trivial norm on \mathbb{Z} . Yang: To be checked.

Spectrum of Tate algebra in one variable Let \mathbf{k} be a complete non-archimedean field, and let $A = \mathbf{k}\{T/r\}$. We list some types of points in the spectrum $\mathcal{M}(A)$.

For each $a \in \mathbf{k}$ with $|a| \leq r$, we have the *type I* point x_a corresponding to the evaluation at a , i.e., $|f|_{x_a} := |f(a)|$ for each $f \in A$. For each closed disk $E = E(a, s) := \{b \in \mathbf{k} : |b - a| \leq s\}$ with center $a \in \mathbf{k}$ and radius $s \leq r$, we have the point $x_{a,s}$ corresponding to the multiplicative semi-norm defined by

$$|f|_{x_E} := \sup_{b \in E(a,s)} |f(b)|$$

for each $f \in A$. If $s \in |\mathbf{k}^\times|$, then the point x_E is called a *type II* point; otherwise, it is called a *type III* point.

Let $\{E^{(s)}\}_s$ be a family of closed disks in \mathbf{k} such that $E^{(s)}$ is of radius s , $E^{(s_1)} \subsetneq E^{(s_2)}$ for any $s_1 < s_2$ and $\bigcap_s E^{(s)} = \emptyset$. Then we have the point $x_{\{E^{(s)}\}}$ corresponding to the multiplicative semi-norm defined by

$$|f|_{x_{\{E^{(s)}\}}} := \inf_s |f|_{x_{E^{(s)}}}$$

for each $f \in A$. Such a point is called a *type IV* point.

Yang: To be completed.

Proposition 15. Let \mathbf{k} be a complete non-archimedean field, and let $r > 0$ be a positive real number. Consider the Tate algebra $\mathbf{k}\{r^{-1}T\}$ equipped with the Gauss norm. The points in the spectrum $\mathcal{M}(\mathbf{k}\{r^{-1}T\})$ can be classified into four types as described above. Yang: To be checked

Proof. Yang: To be completed. □

Proposition 16. Let \mathbf{k} be a complete non-archimedean field, and let $r > 0$ be a positive real number. Consider the Tate algebra $\mathbf{k}\{r^{-1}T\}$ equipped with the Gauss norm. The completed residue fields of the four types of points in the spectrum $\mathcal{M}(\mathbf{k}\{r^{-1}T\})$ are described as follows:

- For a type I point x_a with $a \in \mathbf{k}$ and $|a| \leq r$, the completed residue field $\mathcal{H}(x_a)$ is isomorphic to \mathbf{k} .
- For a type II point $x_{a,s}$ with $a \in \mathbf{k}$ and $s \in |\mathbf{k}^\times|$, the completed residue field $\mathcal{H}(x_{a,s})$ is isomorphic to the field of Laurent series over the residue field $\mathcal{K}_{\mathbf{k}}$, i.e., $\mathcal{K}_{\mathbf{k}}((t))$.
- For a type III point $x_{a,s}$ with $a \in \mathbf{k}$ and $s \notin |\mathbf{k}^\times|$, the completed residue field $\mathcal{H}(x_{a,s})$ is isomorphic to a transcendental extension of $\mathcal{K}_{\mathbf{k}}$ of degree one.
- For a type IV point $x_{\{E^{(s)}\}}$, the completed residue field $\mathcal{H}(x_{\{E^{(s)}\}})$ is isomorphic to a transcendental extension of $\mathcal{K}_{\mathbf{k}}$ of infinite degree.

Yang: To be checked.

Example 17. The completed residue field $\mathcal{H}(x_a)$ for a type I point x_a with $a \in \mathbf{k}$ and $|a| \leq r$ is isomorphic to \mathbf{k} . Yang: To be complete.