

Valuation fields

1 Absolute values and completion

Definition 1. Let \mathbf{k} be a field. An *absolute value* on \mathbf{k} is a function $\|\cdot\| : \mathbf{k} \rightarrow \mathbb{R}_{\geq 0}$ satisfying the following properties for all $x, y \in \mathbf{k}$:

- (a) $\|x\| = 0$ if and only if $x = 0$;
- (b) $\|xy\| = \|x\| \cdot \|y\|$;
- (c) $\|x + y\| \leq \|x\| + \|y\|$.

A field \mathbf{k} equipped with an absolute value $\|\cdot\|$ is called a *valuation field*.

Remark 2. Let \mathbf{k} be a field. Recall that a *valuation* on \mathbf{k} is a function $v : \mathbf{k}^\times \rightarrow \mathbb{R}$ such that

- $\forall x, y \in \mathbf{k}^\times, v(xy) = v(x) + v(y)$;
- $\forall x, y \in \mathbf{k}^\times, v(x + y) \geq \min\{v(x), v(y)\}$.

We can extend v to the whole field \mathbf{k} by defining $v(0) = +\infty$. Fix a real number $\varepsilon \in (0, 1)$. Then v induces an absolute value $|\cdot|_v : \mathbf{k} \rightarrow \mathbb{R}_+$ defined by $|x|_v = \varepsilon^{v(x)}$ for each $x \in \mathbf{k}$.

In some literature, the valuation v is called an *additive valuation* and the induced absolute value $|\cdot|_v$ is called a *multiplicative valuation*. In this note, the term *valuation* always refers to the additive valuation.

Example 3. Let \mathbf{k} be a field. The *trivial absolute value* on \mathbf{k} is defined as

$$\|x\| := \begin{cases} 0, & x = 0; \\ 1, & x \neq 0. \end{cases}$$

Definition 4. The *valuation group* of a valuation field $(\mathbf{k}, \|\cdot\|)$ is defined as the subgroup of $\mathbb{R}_{>0}$ given by

$$|\mathbf{k}^\times| := \{\|x\| : x \in \mathbf{k}^\times\}.$$

Yang: To be revised.

Definition 5. Let \mathbf{k} be a field. Two absolute values $\|\cdot\|_1$ and $\|\cdot\|_2$ on \mathbf{k} are said to be *equivalent* if there exists a real number $c \in (0, 1)$ such that

$$\|x\|_1 = \|x\|_2^c, \quad \forall x \in \mathbf{k}.$$

Note that equivalent absolute values induce the same topology on the field \mathbf{k} . Moreover, the following lemma shows that the converse is also true.

Lemma 6. Let \mathbf{k} be a field and $\|\cdot\|_1, \|\cdot\|_2$ be two absolute values on \mathbf{k} . Then the following statements are equivalent:

- (a) $\|\cdot\|_1$ and $\|\cdot\|_2$ are equivalent;
- (b) $\|\cdot\|_1$ and $\|\cdot\|_2$ induce the same topology on \mathbf{k} ;
- (c) The unit disks $D_1 = \{x \in \mathbf{k} : \|x\|_1 < 1\}$ and $D_2 = \{x \in \mathbf{k} : \|x\|_2 < 1\}$ are the same.

Proof. The implications (a) \Rightarrow (b) is obvious. Now we prove (b) \Rightarrow (c). For any $x \in D_1$, we have $x^n \rightarrow 0$ as $n \rightarrow \infty$ under the absolute value $\|\cdot\|_1$ and thus under $\|\cdot\|_2$. Therefore, $\|x\|_2^n \rightarrow 0$ as $n \rightarrow \infty$, which implies that $\|x\|_2 < 1$, i.e., $x \in D_2$. Similarly, we can prove that $D_2 \subseteq D_1$.

Finally, we prove (c) \Rightarrow (a). If $\|\cdot\|_1$ is trivial, then $D_1 = \{0\}$ and thus $\|\cdot\|_2$ is also trivial. In this case, they are equivalent. Suppose that both $\|\cdot\|_1$ and $\|\cdot\|_2$ are non-trivial. Pick any $x, y \notin D_1 = D_2$. Then there exist real numbers $\alpha, \beta > 0$ such that $\|x\|_1 = \|x\|_2^\alpha$ and $\|y\|_1 = \|y\|_2^\beta$. Suppose the contrary that $\alpha \neq \beta$. Consider the domain $\Omega \subseteq \mathbb{Z}^2$ defined by

$$\begin{cases} n \log \|x\|_2 < m \log \|y\|_2; \\ n\alpha \log \|x\|_2 > m\beta \log \|y\|_2. \end{cases}$$

Since $\alpha \neq \beta$, the two lines defined by the equalities are not parallel. Thus Ω is non-empty. Pick $(n, m) \in \Omega$ and set $z := x^n y^{-m}$. Then we have $\|z\|_2 < 1$ and $\|z\|_1 > 1$, a contradiction. \square

Definition 7. Let $(\mathbf{k}, \|\cdot\|)$ be a valuation field. We say that \mathbf{k} is *complete* if the metric $d(x, y) := \|x - y\|$ makes \mathbf{k} a complete metric space.

Lemma 8. Let $(\mathbf{k}, \|\cdot\|)$ be a valuation field and $(\widehat{\mathbf{k}}, \|\cdot\|)$ its completion as a metric space. Then the operations of addition and multiplication on \mathbf{k} can be extended to $\widehat{\mathbf{k}}$ uniquely, making $(\widehat{\mathbf{k}}, \|\cdot\|)$ a complete valuation field containing \mathbf{k} as a dense subfield.

Proof. Simple analysis. \square

Example 9. Let $|\cdot|_\infty$ be the usual absolute value on the field \mathbb{Q} of rational numbers. Then $(\mathbb{Q}, |\cdot|_\infty)$ is a valuation field. Its completion is the field \mathbb{R} of real numbers equipped with the usual absolute value.

Example 10. Let p be a prime number. For any non-zero rational number $x \in \mathbb{Q}$, we can write it as $x = p^n \frac{a}{b}$, where $n \in \mathbb{Z}$ and $a, b \in \mathbb{Z}$ are integers not divisible by p . The *p-adic absolute value* on \mathbb{Q} is defined as

$$|x|_p := \begin{cases} 0, & x = 0; \\ p^{-n}, & x = p^n \frac{a}{b} \text{ as above.} \end{cases}$$

Then $(\mathbb{Q}, |\cdot|_p)$ is a valuation field. Its completion is the field

$$\mathbb{Q}_p = \left\{ \sum_{n=k}^{+\infty} a_n p^n : k \in \mathbb{Z}, a_n \in \{0, 1, \dots, p-1\} \right\}$$

of *p-adic numbers* equipped with the *p-adic absolute value*.

Unlike the real number field \mathbb{R} , even a valuation field is complete, we can not expect the theorem of nested intervals to hold.

Definition 11. A valuation field $(\mathbf{k}, \|\cdot\|)$ is called *spherically complete* if every decreasing sequence of closed balls in \mathbf{k} has a non-empty intersection.

Example 12. The field \mathbb{C}_p of p -adic complex numbers is not spherically complete, see [Yang: to be added](#).

2 Non-archimedean fields

Definition 13. Let $(\mathbf{k}, \|\cdot\|)$ be a valuation field. We say that \mathbf{k} is *non-archimedean* if its absolute value $\|\cdot\|$ satisfies the *strong triangle inequality*:

$$\|x + y\| \leq \max\{\|x\|, \|y\|\}, \quad \forall x, y \in \mathbf{k}.$$

Otherwise, we say that \mathbf{k} is *archimedean*.

Let \mathbf{k} be a non-archimedean field. Then easily see that $\{x \in \mathbf{k} : \|x\| \leq 1\}$ is a subring of \mathbf{k} . Moreover, it is a local ring whose maximal ideal is $\{x \in \mathbf{k} : \|x\| < 1\}$.

Definition 14. Let \mathbf{k} be a non-archimedean field. The *ring of integers* of \mathbf{k} is defined as

$$\mathbf{k}^\circ := \{x \in \mathbf{k} : \|x\| \leq 1\}.$$

Its maximal ideal is

$$\mathbf{k}^{\circ\circ} := \{x \in \mathbf{k} : \|x\| < 1\}.$$

The *residue field* of \mathbf{k} is defined as

$$\mathcal{K}_{\mathbf{k}} := \tilde{\mathbf{k}} := \mathbf{k}^\circ / \mathbf{k}^{\circ\circ}.$$

Definition 15. Let \mathbf{k} be a non-archimedean field. The *residue absolute value* on the residue field $\mathcal{K}_{\mathbf{k}}$ is defined as

$$|x| := \inf_{y \in \varphi^{-1}(x)} \|y\|, \quad \forall x \in \mathcal{K}_{\mathbf{k}},$$

where $\varphi : \mathbf{k}^\circ \rightarrow \mathcal{K}_{\mathbf{k}}$ is the canonical projection.

Proposition 16. Let \mathbf{k} be a non-archimedean field. Then the residue absolute value on the residue field $\mathcal{K}_{\mathbf{k}}$ is trivial.

Proof. For any $x \in \mathcal{K}_{\mathbf{k}}$, if $x = 0$, then by definition $|x| = 0$. If $x \neq 0$, then $\forall y \in \varphi^{-1}(x)$, we have $y \in \mathbf{k}^\circ \setminus \mathbf{k}^{\circ\circ}$, i.e., $\|y\| = 1$. Thus by definition $|x| = 1$. \square