

Abelian Varieties



“如果是勇者辛美尔，他一定会这么做的！”

Contents

1	The First Properties of Abelian Varieties	1
1.1	Preliminaries	1
1.2	Definition and examples of Abelian Varieties	1
1.3	Complex abelian varieties	3
2	Picard Groups of Abelian Varieties	3
2.1	Pullback along group operations	3
2.2	Projectivity	5
2.3	Isogenies and finite subgroups	5
2.4	Dual abelian varieties	5

1 The First Properties of Abelian Varieties

1.1 Preliminaries

Proposition 1.1. Let $f : X \rightarrow Y$ be a morphism of varieties over a field \mathbf{k} . Then the function $y \mapsto \dim f^{-1}(y)$ is upper semicontinuous, i.e., for every integer m , the set $\{y \in Y : \dim f^{-1}(y) \geq m\}$ is closed in Y . **Yang: To be check.**

1.2 Definition and examples of Abelian Varieties

Theorem 1.2 (Rigidity Lemma). Let $\pi_i : X \rightarrow Y_i$ be proper morphisms of varieties over a field \mathbf{k} for $i = 1, 2$. Suppose that π_1 is a fibration and π_2 contracts $\pi_1^{-1}(y_0)$. Then there exists a rational map $\varphi : Y_1 \dashrightarrow Y_2$ such that $\pi_2 \circ \varphi = \pi_1$ and φ is well-defined near $Y_1 \setminus \{y_0\}$.

Definition 1.3. Let S be a scheme. An *abelian scheme over S* is a group object in the category \mathbf{Sch}_S such that the structure morphism is proper, smooth and a fibration. If $S = \operatorname{Spec} \mathbf{k}$ for some field \mathbf{k} , then it is called an *abelian variety over \mathbf{k}* .

Definition 1.4. Let \mathbf{k} be a field. An *abelian variety over \mathbf{k}* is a proper variety A over \mathbf{k} together with morphisms *identity* $e : \operatorname{Spec} \mathbf{k} \rightarrow A$, *multiplication* $m : A \times A \rightarrow A$ and *inversion* $i : A \rightarrow A$ such that the following diagrams commute:

(a) (Associativity)

$$\begin{array}{ccccc}
 & A \times A \times A & & & \\
 \text{id}_A \times m \swarrow & & m \times \text{id}_A \searrow & & \\
 A \times A & & & & A \times A \\
 & m \searrow & & m \swarrow & \\
 & A & & &
 \end{array}
 ;$$

(b) (Identity)

$$\begin{array}{ccccc}
 A \times \text{Spec } \mathbf{k} & \xrightarrow{\text{id}_A \times e} & A \times A & \xleftarrow{e \times \text{id}_A} & \text{Spec } \mathbf{k} \times A \\
 & \searrow \cong & \downarrow m & \swarrow \cong & \\
 & & A & &
 \end{array}
 ;$$

(c) (Inversion)

$$\begin{array}{ccccc}
 & A & & & \\
 \text{id}_A \times i \swarrow & \downarrow & i \times \text{id}_A \searrow & & \\
 A \times A & & \text{Spec } \mathbf{k} & & A \times A \\
 & m \searrow & \downarrow e & m \swarrow & \\
 & A & & &
 \end{array}
 .$$

Yang: Can we just say that $A(\mathbf{k})$ is a group with e, m, i satisfying the axioms?

Example 1.5. Let E be an elliptic curve over a field \mathbf{k} . Then E is an abelian variety of dimension 1.

Example 1.6.

Example 1.7.

In the following, we will always assume that A is an abelian variety over a field \mathbf{k} of dimension d .

Temporarily, we will use the notation e_A, m_A, i_A to denote the identity section, multiplication morphism and inversion morphism of an abelian variety A . The left translation by $a \in A(\mathbf{k})$ is defined as

$$l_a : A \xrightarrow{\cong} \text{Spec } \mathbf{k} \times A \xrightarrow{a \times \text{id}_A} A \times A \xrightarrow{m_A} A.$$

Similar definition applies to the right translation r_a .

Proposition 1.8. Let A be an abelian variety. Then A is smooth.

Proof. Note that there is an open subset $U \subset A$ which is smooth. Then apply the left translation morphism l_a . □

Proposition 1.9. Let A be an abelian variety. Then the cotangent bundle Ω_A is trivial, i.e., $\Omega_A \cong \mathcal{O}_A^{\oplus d}$ where $d = \dim A$.

Proof. Consider Ω_A as a geometric vector bundle of rank d . Then the conclusion follows from the fact that the left translation morphism l_a induces a morphism of varieties $\Omega_A \rightarrow \Omega_A$ for every $a \in A(\mathbf{k})$.

Yang: But how to show it is a morphism of varieties? Yang: To be completed. □

Lemma 1.10. Let X, Y, Z be proper varieties over a field \mathbf{k} and $g : X \times Y \rightarrow Z$ a morphism over \mathbf{k} . Suppose that g contracts $X \times y_0$ for some point $y_0 \in X(\mathbf{k})$. Then there exists a unique morphism $f : Y \rightarrow Z$ such that $g = f \circ p_Y$, where $p_Y : X \times Y \rightarrow Y$ is the projection to the second factor.

Proof. Yang: To be completed. □

Theorem 1.11. Let A and B be abelian varieties. Then any morphism $f : A \rightarrow B$ with $f(e_A) = e_B$ is a group homomorphism.

Proof. Let $f : A \times A \rightarrow B$ be given by $A \times A \xrightarrow{f \times f} B \times B \xrightarrow{m_B} B$. Then f contracts $A \times e_A$ to e_B .

Yang: To be completed. □

Proposition 1.12. Let A be an abelian variety. Then $A(\mathbf{k})$ is an abelian group.

Proof. Note that a group is abelian if and only if the inversion map is a homomorphism of groups. Then the conclusion follows from Theorem 1.11. □

From now on, we will use the notation $0, +, [-1]_A, t_a$ to denote the identity section, addition morphism, inversion morphism and translation by a of an abelian variety A . For every $n \in \mathbb{Z}_{>0}$, the homomorphism of multiplication by n is defined as

$$[n]_A : A \xrightarrow{\Delta} A \times A \xrightarrow{[n-1]_A \times \text{id}_A} A \times A \xrightarrow{+} A,$$

where Δ is the diagonal morphism.

Proposition 1.13. Let A be an abelian variety over \mathbb{k} and n a positive integer. Then the multiplication by n morphism $[n]_A : A \rightarrow A$ is finite surjective and étale.

Proof. Yang: To be completed. □

1.3 Complex abelian varieties

Theorem 1.14. Let A be a complex abelian variety. Then A is a complex torus, i.e., there exists a lattice $\Lambda \subset \mathbb{C}^d$ such that $A \cong \mathbb{C}^d / \Lambda$. Conversely, let $A = \mathbb{C}^n / \Lambda$ be a complex torus for some lattice Λ . Then A is a complex abelian variety if and only if Λ Yang: To be completed.

2 Picard Groups of Abelian Varieties

2.1 Pullback along group operations

Theorem 2.1 (Seesaw Theorem). Let A be an abelian variety over \mathbb{k} .

Theorem 2.2 (Theorem of the cube). Let X, Y, Z be completed varieties over \mathbb{k} and \mathcal{L} a line bundle on $X \times Y \times Z$. Suppose that there exist $x \in X(\mathbb{k}), y \in Y(\mathbb{k}), z \in Z(\mathbb{k})$ such that the restriction $\mathcal{L}|_{\{x\} \times Y \times Z}, \mathcal{L}|_{X \times \{y\} \times Z}$ and $\mathcal{L}|_{X \times Y \times \{z\}}$ are trivial. Then \mathcal{L} is trivial.

Proof. Yang: To be completed. □

Remark 2.3. If we assume the existence of the Picard scheme, then the theorem of the cube can be deduced from the Rigidity Lemma. Yang: To be completed.

Proposition 2.4. Let A be an abelian variety over \mathbb{k} , $f, g, h : X \rightarrow A$ morphisms from a variety X to A and \mathcal{L} a line bundle on A . Then

$$(f + g + h)^*\mathcal{L} \cong (f + g)^*\mathcal{L} \otimes (f + h)^*\mathcal{L} \otimes (g + h)^*\mathcal{L} \otimes f^*\mathcal{L}^{-1} \otimes g^*\mathcal{L}^{-1} \otimes h^*\mathcal{L}^{-1}.$$

Proof. Yang: To be completed. □

Proposition 2.5. Let A be an abelian variety over \mathbb{k} , $n \in \mathbb{Z}$ and \mathcal{L} a line bundle on A . Then we have

$$[n]_A^*\mathcal{L} \cong \mathcal{L}^{\otimes \frac{1}{2}(n^2+n)} \otimes [-1]_A^*\mathcal{L}^{\otimes \frac{1}{2}(n^2-n)}.$$

Proof. Yang: To be completed. □

Theorem 2.6 (Theorem of the square). Let A be an abelian variety over \mathbb{k} , $x, y \in A(\mathbb{k})$ two points and \mathcal{L} a line bundle on A . Then

$$t_{x+y}^*\mathcal{L} \otimes \mathcal{L} \cong t_x^*\mathcal{L} \otimes t_y^*\mathcal{L}.$$

Remark 2.7. We can define a map

$$\Phi_{\mathcal{L}} : A(\mathbb{k}) \rightarrow \text{Pic}(A), \quad x \mapsto t_x^*\mathcal{L} \otimes \mathcal{L}^{-1}.$$

Then theorem of the square implies that $\Phi_{\mathcal{L}}$ is a homomorphism of groups. When we vary \mathcal{L} , the map

$$\Phi_{\square} : \text{Pic}(A) \rightarrow \text{Hom}_{\mathbf{Grp}}(A(\mathbb{k}), \text{Pic}(A)), \quad \mathcal{L} \mapsto \Phi_{\mathcal{L}}$$

is a group homomorphism. For any $x \in A(\mathbb{k})$, we have

$$\Phi_{t_x^*\mathcal{L}} = \Phi_{\mathcal{L}}.$$

In the other words,

$$\Phi_{\mathcal{L}}(x) \in \text{Ker } \Phi_{\square}, \quad \forall \mathcal{L} \in \text{Pic}(A), x \in A(\mathbb{k}).$$

Yang: To be completed.

If we assume the scheme structure on $\text{Pic}(A)$, then $\Phi_{\mathcal{L}}$ is a morphism of scheme and factors through $\text{Pic}^0(A)$. Let $K(\mathcal{L}) := \text{Ker } \Phi_{\mathcal{L}}$, then $K(\mathcal{L})$ is a subgroup scheme of A . We give another description of $K(\mathcal{L})$. From this point, we can recover the dual abelian variety $A^\vee = \text{Pic}^0(A)$ as the quotient $A/K(\mathcal{L})$.

Yang: To be completed.

2.2 Projectivity

Proposition 2.8. Let A be an abelian variety over \mathbb{k} and D an effective divisor on A . Then $|2D|$ is base point free.

Theorem 2.9. Let A be an abelian variety over \mathbb{k} and D an effective divisor on A . TFAE:

- (a) the stabilizer $\text{Stab}(D)$ of D is finite;
- (b) the morphism $\Phi_{|2D|}$ induced by the complete linear system $|2D|$ is finite;
- (c) D is ample;
- (d) $K(\mathcal{O}_A(D))$ is finite.

Theorem 2.10. Let A be an abelian variety over \mathbf{k} . Then A is projective.

Proof. Yang: To be completed. □

2.3 Isogenies and finite subgroups

Theorem 2.11. Let A be an abelian variety of dimension d over \mathbb{k} . Then the subgroup $A[n]$ of n torsion points is finite and we have

- (a) if n is coprime to $\text{char}(\mathbf{k})$, then $A[n] \cong (\mathbb{Z}/n\mathbb{Z})^{2d}$;
- (b) if $n = p^k$ for $p = \text{char}(\mathbf{k}) > 0$

Proof. Yang: To be completed. □

Theorem 2.12. Let A be an abelian variety over \mathbb{k} . There is a bijection between the isogenies from A over \mathbb{k} and the finite subgroup schemes of A .

2.4 Dual abelian varieties

Theorem 2.13. Let A be an abelian variety over \mathbf{k} . Then $\text{Pic}^0(A)$ has a natural structure of an abelian variety, called the *dual abelian variety* of A , denoted by A^\vee .

Proposition 2.14. There exists a unique line bundle \mathcal{P} on $A \times A^\vee$ such that for every $y = \mathcal{L} \in A^\vee = \text{Pic}^0(A)$, we have $\mathcal{P}|_{A \times \{y\}} \cong \mathcal{L}$.