
Schemes and Varieties



“宇宙战警行为规范第四十条，遇见违反社会公德的行为，发现违法犯罪行为，要见义勇为，勇于斗争，善于斗争。第十四条，不说谎，不骗人，不弄虚作假，知错就改，诚实守信，言行一致，答应别人的事要做到。……榴莲，不用跟任何人道歉，你跟这儿的人不一样，你是来让世界变好的。”

Contents

0	Locally Ringed Space	2
0.1	Sheaves	2
0.2	Locally Ringed Space	4
0.3	Manifolds as locally ringed spaces	5
0.4	Vector bundles and \mathcal{O}_X -modules	5
1	The First Properties of Schemes	6
1.1	Schemes	6
1.2	Fiber product	7
1.3	Noetherian and finite type	9
1.4	Integral, reduced and irreducible	9
1.5	Dimension	10
1.6	Separated and proper	10
2	Category of sheaves of modules	11
2.1	Sheaves of modules, quasi-coherent and coherent sheaves	11
2.2	As abelian categories	11
2.3	Relevant functors	12
2.4	Locally free sheaves and vector bundles	12
2.5	Cohomological theory	12
3	Normal, Cohen-Macaulay, and regular schemes	12
3.1	Tangent spaces	12
4	Line Bundles and Divisors	12
4.1	Cartier Divisors	12
4.2	Line Bundles and Picard Group	12
4.3	Weil Divisors and Reflexive Sheaves	13
5	Line bundles induce morphisms	13
5.1	Ample and basepoint free line bundles	13
5.2	Linear systems	15
5.3	Asymptotic behavior	15
6	Differentials and duality	16
7	Flat, smooth and étale morphisms	16
7.1	Flat family and Hilbert polynomial	16
7.2	Base change and semicontinuity	16
7.3	Smooth morphisms	17
7.4	Étale morphisms	17

8	Relative objects	17
8.1	Relative schemes	17
8.2	Blowing up	18
8.3	Relative ampleness and relative morphisms	18
9	Finite morphisms and fibrations	18

0 Locally Ringed Space

0.1 Sheaves

Definition 0.1. Let X be a topological space. A *presheaf* of sets (resp. abelian groups, rings, etc.) on X is a contravariant functor $\mathcal{F} : \mathbf{Open}(X) \rightarrow \mathbf{Set}$ (resp. \mathbf{Ab} , \mathbf{Ring} , etc.), where $\mathbf{Open}(X)$ is the category of open subsets of X with inclusions as morphisms.

A presheaf \mathcal{F} is a *sheaf* if sections can be glued uniquely. More precisely, for every open covering $\{U_i\}_{i \in I}$ of an open set $U \subset X$ and every family of sections $s_i \in \mathcal{F}(U_i)$ such that $s_i|_{U_i \cap U_j} = s_j|_{U_i \cap U_j}$ for all $i, j \in I$, there exists a unique section $s \in \mathcal{F}(U)$ such that $s|_{U_i} = s_i$ for all $i \in I$.

For two open sets $V \subset U \subset X$, the morphism $\mathcal{F}(U) \rightarrow \mathcal{F}(V)$, often denoted by res_V^U , is called the *restriction map*.

Example 0.2. Let X be a real (resp. complex) manifold. The assignment $U \mapsto \mathcal{C}^\infty(U, \mathbb{R})$ (resp. $U \mapsto \{\text{holomorphic functions on } U\}$) defines a sheaf of rings on X .

Example 0.3. Let X be a non-connected topological space. The assignment

$$U \mapsto \{\text{constant functions on } U\}$$

defines a presheaf \mathcal{C} of rings on X but not a sheaf.

For a concrete example, let $X = (0, 1) \cup (2, 3)$ with the subspace topology from \mathbb{R} . Consider the open covering $\{(0, 1), (2, 3)\}$ of X . The sections $s_1 = 1 \in \mathcal{C}((0, 1))$ and $s_2 = 2 \in \mathcal{C}((2, 3))$ agree on the intersection (which is empty), but there is no global section $s \in \mathcal{C}(X)$ such that $s|_{(0, 1)} = s_1$ and $s|_{(2, 3)} = s_2$.

Definition 0.4. Let X be a topological space and \mathcal{F}, \mathcal{G} be presheaves on X with values in the same category (e.g., \mathbf{Set} , \mathbf{Ab} , \mathbf{Ring} , etc.). A *morphism of presheaves* $\varphi : \mathcal{F} \rightarrow \mathcal{G}$ is a natural transformation between the functors \mathcal{F} and \mathcal{G} . In other words, for every open set $U \subset X$, there is a morphism $\varphi(U) : \mathcal{F}(U) \rightarrow \mathcal{G}(U)$ such that for every inclusion of open sets $V \subset U$, the following diagram commutes:

$$\begin{array}{ccc} \mathcal{F}(U) & \xrightarrow{\varphi(U)} & \mathcal{G}(U) \\ \text{res}_V^U \downarrow & & \downarrow \text{res}_V^U \\ \mathcal{F}(V) & \xrightarrow{\varphi(V)} & \mathcal{G}(V). \end{array}$$

If \mathcal{F} and \mathcal{G} are sheaves, then φ is called a *morphism of sheaves*.

Definition 0.5. Let X be a topological space and \mathcal{F} a presheaf on X with values in a category \mathbf{C} . For a point $x \in X$, the *stalk* of \mathcal{F} at x , denoted by \mathcal{F}_x , is defined as the colimit

$$\mathcal{F}_x := \varinjlim_{U \ni x} \mathcal{F}(U),$$

where the colimit is taken over all open neighborhoods U of x . An element of \mathcal{F}_x is called a *germ* of sections of \mathcal{F} at x .

$$\mathcal{F}_x = \{(U, s) : U \in \mathbf{Open}(X), U \ni x, s \in \mathcal{F}(U)\} / \sim,$$

Definition 0.6. Let X be a topological space and \mathcal{F} a presheaf on X with values in **Set** (resp. **Ab**, **Ring**, etc.). A *sheafification* of \mathcal{F} is a sheaf \mathcal{F}^+ on X together with a morphism of presheaves $\eta : \mathcal{F} \rightarrow \mathcal{F}^+$ such that for every sheaf \mathcal{G} on X and every morphism of presheaves $\varphi : \mathcal{F} \rightarrow \mathcal{G}$, there exists a unique morphism of sheaves $\varphi^+ : \mathcal{F}^+ \rightarrow \mathcal{G}$ such that $\varphi = \varphi^+ \circ \eta$. In other words, the following diagram commutes:

$$\begin{array}{ccc} \mathcal{F} & \xrightarrow{\eta} & \mathcal{F}^\dagger \\ & \searrow \varphi & \downarrow \varphi^\dagger \\ & & \mathcal{G}. \end{array}$$

Yang: The concrete describe of sheafification.

Proposition 0.8. Let X be a topological space and $\varphi : \mathcal{F} \rightarrow \mathcal{G}$ be a homomorphism of sheaves of abelian groups on X . Then φ is injective if and only if for every open set $U \subset X$, the map $\varphi(U) : \mathcal{F}(U) \rightarrow \mathcal{G}(U)$ is injective. Yang: To be checked.

Remark 0.9. The surjectivity on stalks cannot imply the surjectivity on sections. A counterexample is given by the exponential map $\exp : \mathcal{O}_{\mathbb{C}} \rightarrow \mathcal{O}_{\mathbb{C}}^*$ defined by $\exp(f) = e^f$, where $\mathcal{O}_{\mathbb{C}}$ is the sheaf of holomorphic functions on \mathbb{C} and $\mathcal{O}_{\mathbb{C}}^*$ is the sheaf of non-vanishing holomorphic functions on \mathbb{C} . The induced map on stalks $\exp_z : \mathcal{O}_{\mathbb{C},z} \rightarrow \mathcal{O}_{\mathbb{C},z}^*$ is surjective for every $z \in \mathbb{C}$ by the existence of logarithm locally. However, the map on global sections $\exp(\mathbb{C}) : \mathcal{O}_{\mathbb{C}}(\mathbb{C}) \rightarrow \mathcal{O}_{\mathbb{C}}^*(\mathbb{C})$ is not surjective since there is no entire function f such that $e^{f(z)} = z$ for all $z \in \mathbb{C}^*$. **Yang: To be continued.**

Proposition 0.10. Let X be a topological space and $\varphi : \mathcal{F} \rightarrow \mathcal{G}$ be a homomorphism of sheaves of abelian groups on X . Then φ is an isomorphism if and only if it is injective and surjective.

Yang: Now we consider sheaves with values in an abelian category.

Definition 0.11. Let X be a topological space and $\varphi : \mathcal{F} \rightarrow \mathcal{G}$ be a homomorphism of sheaves of abelian groups on X . The *kernel* of φ , denoted by $\ker \varphi$, is the sheaf defined by

$$(\ker \varphi)(U) := \ker(\varphi(U) : \mathcal{F}(U) \rightarrow \mathcal{G}(U))$$

for every open set $U \subset X$.

The *cokernel* of φ , denoted by $\operatorname{coker} \varphi$, is the sheafification of the presheaf defined by

$$(\operatorname{coker} \varphi)_{\text{pre}}(U) := \operatorname{coker}(\varphi(U) : \mathcal{F}(U) \rightarrow \mathcal{G}(U))$$

for every open set $U \subset X$. Yang: To be continued.

Theorem 0.12. Let X be a topological space and \mathbf{C} be an abelian category (e.g., \mathbf{Ab}). Then the category of sheaves on X with values in \mathbf{C} is an abelian category.

Proof. Yang: To be continued. □

Yang: To be checked and continuous.

0.2 Locally Ringed Space

Definition 0.13. Let $f : X \rightarrow Y$ be a continuous map between topological spaces. The *push-forward* functor $f_* : \mathbf{Sh}(X, \mathbf{C}) \rightarrow \mathbf{Sh}(Y, \mathbf{C})$ is defined by

$$(f_* \mathcal{F})(V) := \mathcal{F}(f^{-1}(V))$$

for every open set $V \subset Y$ and sheaf $\mathcal{F} \in \mathbf{Sh}(X, \mathbf{C})$.

Definition 0.14. A *locally ringed space* is a pair (X, \mathcal{O}_X) where X is a topological space and \mathcal{O}_X is a sheaf of rings on X such that for every $x \in X$, the stalk $\mathcal{O}_{X,x}$ is a local ring.

A *morphism of locally ringed spaces* $f : (X, \mathcal{O}_X) \rightarrow (Y, \mathcal{O}_Y)$ consists of a continuous map $f : X \rightarrow Y$ and a morphism of sheaves of rings $f^\# : \mathcal{O}_Y \rightarrow f_* \mathcal{O}_X$ such that for every $x \in X$, the induced map on stalks $f_x^\# : \mathcal{O}_{Y,f(x)} \rightarrow \mathcal{O}_{X,x}$ is a local homomorphism, i.e., it maps the maximal ideal of $\mathcal{O}_{Y,f(x)}$ to the maximal ideal of $\mathcal{O}_{X,x}$.

Example 0.15. Let p be a prime number. Then the inclusion $\mathbb{Z}_{(p)} \rightarrow \mathbb{Q}$ is a homomorphism of local rings but not a local homomorphism. Here $\mathbb{Z}_{(p)}$ is the localization of \mathbb{Z} at the prime ideal (p) .

Example 0.16 (Glue morphisms). Let $f : (X, \mathcal{O}_X) \rightarrow (Y, \mathcal{O}_Y)$ be a morphism of locally ringed spaces. If $U \subset X$ and $V \subset Y$ are open subsets such that $f(U) \subset V$, then the restriction $f|_U : (U, \mathcal{O}_X|_U) \rightarrow (V, \mathcal{O}_Y|_V)$ is a morphism of locally ringed spaces. Conversely, if $\{U_i\}_{i \in I}$ is an open covering of X and for each $i \in I$, we have a morphism $f_i : (U_i, \mathcal{O}_X|_{U_i}) \rightarrow (Y, \mathcal{O}_Y)$ such that $f_i|_{U_i \cap U_j} = f_j|_{U_i \cap U_j}$ for all

$i, j \in I$, then there exists a unique morphism $f : (X, \mathcal{O}_X) \rightarrow (Y, \mathcal{O}_Y)$ such that $f|_{U_i} = f_i$ for all $i \in I$.

Example 0.17 (Glue locally ringed space). We construct a locally ringed space by gluing open subspaces. Let (X_i, \mathcal{O}_{X_i}) be locally ringed spaces for $i \in I$ and $(U_{ij}, \mathcal{O}_{X_i}|_{U_{ij}})$ be open subspaces for $i, j \in I$. Suppose we have isomorphisms $\varphi_{ij} : (U_{ij}, \mathcal{O}_{X_i}|_{U_{ij}}) \rightarrow (U_{ji}, \mathcal{O}_{X_j}|_{U_{ji}})$ such that

- (a) $\varphi_{ii} = \text{id}_{X_i}$ for all $i \in I$;
- (b) $\varphi_{ij}(U_{ij} \cap U_{ik}) = U_{ji} \cap U_{jk}$ for all $i, j \in I$;
- (c) $\varphi_{jk} \circ \varphi_{ij} = \varphi_{ik}$ on $U_{ij} \cap U_{ik}$ for all $i, j, k \in I$.

Then there exists a locally ringed space (X, \mathcal{O}_X) and open immersions $\psi_i : (X_i, \mathcal{O}_{X_i}) \rightarrow (X, \mathcal{O}_X)$ uniquely up to isomorphism such that

- (a) $\varphi_i(U_{ij}) = \psi_i(X_i) \cap \psi_j(X_j)$ for all $i, j \in I$;
- (b) the following diagram

$$\begin{array}{ccccc} (U_{ij}, \mathcal{O}_{X_i}|_{U_{ij}}) & \hookrightarrow & (X_i, \mathcal{O}_{X_i}) & \xrightarrow{\psi_i} & (X, \mathcal{O}_X) \\ \varphi_{ij} \downarrow & & & & \downarrow = \\ (U_{ji}, \mathcal{O}_{X_j}|_{U_{ji}}) & \hookrightarrow & (X_j, \mathcal{O}_{X_j}) & \xrightarrow{\psi_j} & (X, \mathcal{O}_X) \end{array}$$

commutes for all $i, j \in I$;

- (c) $X = \bigcup_{i \in I} \psi_i(X_i)$.

Such (X, \mathcal{O}_X) is called *the locally ringed space obtained by gluing the (X_i, \mathcal{O}_{X_i}) along the φ_{ij}* .

First φ_{ij} induces an equivalence relation \sim on the disjoint union $\coprod_{i \in I} X_i$. By taking the quotient space, we can glue the underlying topological spaces to get a topological space X . The structure sheaf \mathcal{O}_X is given by

$$\mathcal{O}_X(V) := \left\{ (s_i)_{i \in I} \in \prod_{i \in I} \mathcal{O}_{X_i}(\psi_i^{-1}(V)) \mid s_i|_{U_{ij}} = \varphi_{ij}^\#(s_j|_{U_{ji}}) \text{ for all } i, j \in I \right\}.$$

Easy to check that (X, \mathcal{O}_X) is a locally ringed space and satisfies the required properties. If there is another locally ringed space $(X', \mathcal{O}_{X'})$ with ψ'_i satisfying the same properties, then by gluing $\psi'_i \circ \psi_i^{-1}$ we get an isomorphism $(X, \mathcal{O}_X) \rightarrow (X', \mathcal{O}_{X'})$.

0.3 Manifolds as locally ringed spaces

0.4 Vector bundles and \mathcal{O}_X -modules

Let (X, \mathcal{O}_X) be a manifold (real or complex) and (\mathcal{E}, π, X) a vector bundle over X .

Yang: It can regard as a sheaf on X .

Definition 0.18. Let (X, \mathcal{O}_X) be a ringed space. A *sheaf of \mathcal{O}_X -modules* is a sheaf \mathcal{F} of abelian groups on X such that for every open set $U \subseteq X$, $\mathcal{F}(U)$ is an $\mathcal{O}_X(U)$ -module, and for every inclusion of open

sets $V \subseteq U$, the restriction map $\text{res}_{UV} : \mathcal{F}(U) \rightarrow \mathcal{F}(V)$ is $\mathcal{O}_X(U)$ -linear, where the $\mathcal{O}_X(U)$ -module structure on $\mathcal{F}(V)$ is induced by the restriction map $\text{res}_{UV} : \mathcal{O}_X(U) \rightarrow \mathcal{O}_X(V)$.

A *morphism of \mathcal{O}_X -modules* is a morphism of sheaves of abelian groups $\varphi : \mathcal{F} \rightarrow \mathcal{G}$ such that for every open set $U \subseteq X$, the map $\varphi(U) : \mathcal{F}(U) \rightarrow \mathcal{G}(U)$ is $\mathcal{O}_X(U)$ -linear. Yang: To be checked...

Yang: We will try to construct a sequence of subcategories of $\mathbf{Mod}_{\mathcal{O}_X}$.

Definition 0.19. A sheaf of \mathcal{O}_X -modules \mathcal{F} is said to be *finitely generated* if for every open set $U \subseteq X$, the $\mathcal{O}_X(U)$ -module $\mathcal{F}(U)$ is finitely generated. Yang: To be continued.

Definition 0.20. A sheaf of \mathcal{O}_X -modules \mathcal{F} is said to be *locally free of rank r* if for every point $x \in X$, there exists an open neighborhood U of x such that $\mathcal{F}|_U$ is isomorphic to \mathcal{O}_U^r , where \mathcal{O}_U^r is the direct sum of r copies of \mathcal{O}_U . Yang: To be continued.

Definition 0.21. A sheaf of \mathcal{O}_X -modules \mathcal{F} is said to be *quasi-coherent* if for every point $x \in X$, there exists an open neighborhood U of x such that $\mathcal{F}|_U$ is isomorphic to the cokernel of a morphism of free \mathcal{O}_U -modules, i.e., there exists an exact sequence of sheaves of \mathcal{O}_U -modules

$$\mathcal{O}_U^{(I)} \rightarrow \mathcal{O}_U^{(J)} \rightarrow \mathcal{F}|_U \rightarrow 0,$$

where I, J are (possibly infinite) index sets. Yang: To be checked...

Definition 0.22. A sheaf of \mathcal{O}_X -modules \mathcal{F} is said to be *coherent* if it is finitely generated, and for every open set $U \subseteq X$ and every morphism of sheaves of \mathcal{O}_U -modules $\varphi : \mathcal{O}_U^n \rightarrow \mathcal{F}|_U$, the kernel of φ is finitely generated. Yang: To be checked...

1 The First Properties of Schemes

If you learn the following content for the first time, it is recommended to skip all the proofs in this section and focus on the examples, remarks and the statements of propositions and theorems.

1.1 Schemes

Let R be a ring. Recall that the *spectrum* of R , denoted by $\text{Spec } R$, is the set of all prime ideals of R equipped with the Zariski topology, where the closed sets are of the form $V(I) = \{\mathfrak{p} \in \text{Spec } R : I \subset \mathfrak{p}\}$ for some ideal $I \subset R$.

For each $f \in R$, let $D(f) = \{\mathfrak{p} \in \text{Spec } R : f \notin \mathfrak{p}\}$. Such $D(f)$ is open in $\text{Spec } R$ and called a *principal open set*.

Proposition 1.1. Let R be a ring. The collection of principal open sets $\{D(f) : f \in R\}$ forms a basis for the Zariski topology on $\text{Spec } R$.

Proof. Yang: To be continued □

Define a sheaf of rings on $\text{Spec } R$ by

$$\mathcal{O}_{\text{Spec } R}(D(f)) = R[1/f].$$

Then $(\mathrm{Spec} R, \mathcal{O}_{\mathrm{Spec} R})$ is a locally ringed space.

Definition 1.2. An *affine scheme* is a locally ringed space isomorphic to $(\mathrm{Spec} R, \mathcal{O}_{\mathrm{Spec} R})$ for some ring R . A *scheme* is a locally ringed space (X, \mathcal{O}_X) which admits an open cover $\{U_i\}_{i \in I}$ such that $(U_i, \mathcal{O}_X|_{U_i})$ is an affine scheme for each $i \in I$.

A *morphism of schemes* is a morphism of locally ringed spaces.

These data form a category, denoted by **Sch**. If we fix a base scheme S , then an S -*scheme* is a scheme X together with a morphism $X \rightarrow S$. The category of S -schemes is denoted by **Sch**/ S or **Sch** $_S$.

Theorem 1.3. The functor $\mathrm{Spec} : \mathbf{Ring}^{\mathrm{op}} \rightarrow \mathbf{Sch}$ is fully faithful and induces an equivalence of categories between the category of rings and the category of affine schemes. Yang: To be continued

Definition 1.4. A morphism of schemes $f : X \rightarrow Y$ is an *open immersion* (resp. *closed immersion*) if f induces an isomorphism of X onto an open (resp. closed) subscheme of Y . An *immersion* is a morphism which factors as a closed immersion followed by an open immersion. Yang: To be continued

Example 1.5. Let R be a graded ring. The *projective scheme* $\mathrm{Proj} R$ is defined as the scheme associated to the sheaf of rings

$$\mathcal{O}_{\mathrm{Proj} R} = \bigoplus_{d \geq 0} R_d.$$

It can be covered by open affine subschemes of the form $\mathrm{Spec} R_f$ for homogeneous elements $f \in R$.

Yang: To be checked.

Example 1.6 (Glue open subschemes). The construction in Example 0.17 allows us to glue open subschemes to get a scheme. More precisely, let (X_i, \mathcal{O}_{X_i}) be schemes for $i \in I$ and $(U_{ij}, \mathcal{O}_{X_i}|_{U_{ij}})$ be open subschemes for $i, j \in I$. Suppose we have isomorphisms $\varphi_{ij} : (U_{ij}, \mathcal{O}_{X_i}|_{U_{ij}}) \rightarrow (U_{ji}, \mathcal{O}_{X_j}|_{U_{ji}})$ satisfying the cocycle condition as in Example 0.17. Then the locally ringed space (X, \mathcal{O}_X) obtained by gluing the (X_i, \mathcal{O}_{X_i}) along the φ_{ij} is a scheme.

Definition 1.7. Let $f : X \rightarrow Y$ be a morphism of schemes. The *scheme theoretic image* of f is the smallest closed subscheme Z of Y such that f factors through Z . More precisely, if $Y = \mathrm{Spec} A$ is affine, then the scheme theoretic image of f is $\mathrm{Spec}(A/\ker(f^\#))$, where $f^\# : A \rightarrow \Gamma(X, \mathcal{O}_X)$ is the induced map on global sections. In general, we can cover Y by affine open subsets and glue the scheme theoretic images on each affine open subset to get the scheme theoretic image of f . Yang: To be checked.

1.2 Fiber product

Definition 1.8. Let \mathcal{C} be a category and $X, Y, S \in \mathrm{Obj}(\mathcal{C})$ with morphisms $f : X \rightarrow S$ and $g : Y \rightarrow S$. A *fiber product* of X and Y over S is an object $Z \in \mathrm{Obj}(\mathcal{C})$ together with morphisms $p : Z \rightarrow X$ and

$q : Z \rightarrow Y$ such that the following diagram commutes:

$$\begin{array}{ccc} Z & \xrightarrow{q} & Y \\ p \downarrow & & \downarrow g \\ X & \xrightarrow{f} & S \end{array}$$

and satisfies the universal property that for any object $W \in \text{Obj}(\mathcal{C})$ with morphisms $u : W \rightarrow X$ and $v : W \rightarrow Y$ such that $f \circ u = g \circ v$, there exists a unique morphism $h : W \rightarrow Z$ such that $p \circ h = u$ and $q \circ h = v$.

If a fiber product exists, it is unique up to a unique isomorphism. We denote the fiber product by $X \times_S Y$. **Yang: To be checked.**

Example 1.9. In the category of sets, the fiber product $X \times_S Y$ is given by

$$X \times_S Y = \{(x, y) \in X \times Y : f(x) = g(y)\},$$

with the projections $p : X \times_S Y \rightarrow X$ and $q : X \times_S Y \rightarrow Y$ being the restrictions of the natural projections. **Yang: To be checked.**

Remark 1.10. If one reverses the arrows in [Definition 1.8](#), one gets the notion of *fiber coproduct*. It is also called the *pushout* or *amalgamated sum* in some literature. We denote the fiber coproduct of X and Y over S by $X \amalg_S Y$. Note that in the category of rings, the fiber coproduct $A \amalg_R B$ of R -algebras A and B over R is given by the tensor product $A \otimes_R B$. Dually, one can expect that fiber products of affine schemes correspond to tensor products of rings.

Theorem 1.11. The category of schemes admits fiber products. More precisely, given morphisms of schemes $f : X \rightarrow S$ and $g : Y \rightarrow S$, there exists a scheme Z together with morphisms $p : Z \rightarrow X$ and $q : Z \rightarrow Y$ such that the diagram

$$\begin{array}{ccc} Z & \xrightarrow{q} & Y \\ p \downarrow & & \downarrow g \\ X & \xrightarrow{f} & S \end{array}$$

commutes and satisfies the universal property of the fiber product. We denote this scheme by $X \times_S Y$.

Yang: To be continued

Definition 1.12. Let $f : X \rightarrow Y$ be a morphism of schemes and $y \in Y$ a point. The *scheme theoretic fiber* of f over y is the fiber product $X_y = X \times_Y \text{Spec } \kappa(y)$, where $\kappa(y)$ is the residue field of the local ring $\mathcal{O}_{Y,y}$. **Yang: To be checked.**

Definition 1.13. Let X be a scheme and $Z_1, Z_2 \subset X$ be closed subschemes defined by quasi-coherent sheaves of ideals $\mathcal{I}_1, \mathcal{I}_2 \subset \mathcal{O}_X$, respectively. The *scheme theoretic intersection* of Z_1 and Z_2 is the closed subscheme $Z_1 \cap Z_2$ defined by the quasi-coherent sheaf of ideals $\mathcal{I}_1 + \mathcal{I}_2$. **Yang: To be checked.**

1.5 Dimension

Definition 1.24. The *Krull dimension* of a topological space X , denoted by $\dim X$, is the supremum of the lengths n of chains of distinct irreducible closed subsets

$$Z_0 \subsetneq Z_1 \subsetneq \cdots \subsetneq Z_n$$

in X . If no such finite supremum exists, we say that X has infinite dimension. **Yang: To be checked.**

1.6 Separated and proper

Definition 1.25. A morphism of schemes $f : X \rightarrow Y$ is *separated* if the diagonal morphism $\Delta_f : X \rightarrow X \times_Y X$ is a closed immersion. A scheme X is *separated* if the structure morphism $X \rightarrow \operatorname{Spec} \mathbb{Z}$ is separated. **Yang: To be checked.**

Proposition 1.26. Any affine scheme is separated. More generally, any morphism between affine schemes is separated. **Yang: To be checked.**

Proposition 1.27. Let $f : X \rightarrow Y$ be a morphism of schemes. Then f is separated if and only if for any scheme T and any pair of morphisms $g_1, g_2 : T \rightarrow X$ such that $f \circ g_1 = f \circ g_2$, the equalizer of g_1 and g_2 is a closed subscheme of T . **Yang: To be checked.**

Proposition 1.28. A scheme X is separated if and only if for any pair of affine open subschemes $U, V \subset X$, the intersection $U \cap V$ is also an affine open subscheme. **Yang: To be checked.**

Proposition 1.29. The composition of separated morphisms is separated. Moreover, separatedness is stable under base change, i.e., if $f : X \rightarrow Y$ is a separated morphism and $Y' \rightarrow Y$ is any morphism, then the base change $X \times_Y Y' \rightarrow Y'$ is also separated. **Yang: To be checked.**

Proposition 1.30. A morphism of schemes $f : X \rightarrow Y$ is separated if and only if for every commutative diagram

$$\begin{array}{ccc} \operatorname{Spec} K & \xrightarrow{\quad} & X \\ \downarrow & \nearrow \text{dashed} & \downarrow f \\ \operatorname{Spec} R & \xrightarrow{\quad} & Y \end{array}$$

where R is a valuation ring with field of fractions K , there exists at most one morphism $\operatorname{Spec} R \rightarrow X$ making the entire diagram commute. **Yang: To be checked.**

Definition 1.31. A morphism of schemes $f : X \rightarrow Y$ is *universally closed* if for any morphism $Y' \rightarrow Y$, the base change $X \times_Y Y' \rightarrow Y'$ is a closed map. **Yang: To be checked.**

Definition 1.32. A morphism of schemes $f : X \rightarrow Y$ is *proper* if it is of finite type, separated, and universally closed (i.e., for any morphism $Y' \rightarrow Y$, the base change $X \times_Y Y' \rightarrow Y'$ is a closed map). A scheme X is *proper* if the structure morphism $X \rightarrow \operatorname{Spec} \mathbb{Z}$ is proper. **Yang: To be checked.**

Theorem 1.33. Any projective morphism is proper. In particular, any projective scheme is proper.

Yang: To be checked.

Proposition 1.34. The composition of proper morphisms is proper. Moreover, properness is stable under base change, i.e., if $f : X \rightarrow Y$ is a proper morphism and $Y' \rightarrow Y$ is any morphism, then the base change $X \times_Y Y' \rightarrow Y'$ is also proper. Yang: To be checked.

Proposition 1.35. A morphism of schemes $f : X \rightarrow Y$ is proper if and only if for every commutative diagram

$$\begin{array}{ccc} \operatorname{Spec} K & \longrightarrow & X \\ \downarrow & \nearrow & \downarrow f \\ \operatorname{Spec} R & \longrightarrow & Y \end{array}$$

where R is a valuation ring with field of fractions K , there exists a unique morphism $\operatorname{Spec} R \rightarrow X$ making the entire diagram commute. Yang: To be checked.

2 Category of sheaves of modules

2.1 Sheaves of modules, quasi-coherent and coherent sheaves

Definition 2.1. Let (X, \mathcal{O}_X) be a ringed space. An \mathcal{O}_X -module \mathcal{F} is called *quasi-coherent* if for every point $x \in X$, there exists an open neighborhood U of x such that $\mathcal{F}|_U$ is isomorphic to the cokernel of a morphism of free \mathcal{O}_U -modules, i.e., there exists an exact sequence of sheaves of \mathcal{O}_U -modules

$$\mathcal{O}_U^{(I)} \rightarrow \mathcal{O}_U^{(J)} \rightarrow \mathcal{F}|_U \rightarrow 0,$$

where I, J are (possibly infinite) index sets.

Definition 2.2. Let (X, \mathcal{O}_X) be a ringed space. An \mathcal{O}_X -module \mathcal{F} is called *finitely generated* if for every point $x \in X$, there exists an open neighborhood U of x such that there exists a surjective morphism of sheaves of \mathcal{O}_U -modules

$$\mathcal{O}_U^n \rightarrow \mathcal{F}|_U \rightarrow 0.$$

Definition 2.3. Let (X, \mathcal{O}_X) be a ringed space. An \mathcal{O}_X -module \mathcal{F} is called *coherent* if it is finitely generated, and for every open set $U \subseteq X$ and every morphism of sheaves of \mathcal{O}_U -modules $\varphi : \mathcal{O}_U^n \rightarrow \mathcal{F}|_U$, the kernel of φ is finitely generated.

2.2 As abelian categories

Theorem 2.4. The categories of sheaves of abelian groups, quasi-coherent sheaves, and coherent sheaves on a ringed space (X, \mathcal{O}_X) are all abelian categories. Yang: To be checked.

Theorem 2.5. Let (X, \mathcal{O}_X) be a ringed space. The category of sheaves of \mathcal{O}_X -modules has enough injectives. **Yang: To be checked.**

2.3 Relevant functors

2.4 Locally free sheaves and vector bundles

2.5 Cohomological theory

3 Normal, Cohen-Macaulay, and regular schemes

3.1 Tangent spaces

There are many description of the tangent space of a scheme at a point. Here we give one of them.

Let X be a scheme over a field \mathbf{k} , and let $x \in X(\mathbf{k})$.

Proposition 3.1. Let $\mathrm{Spec} \mathbf{k}[\epsilon]/(\epsilon^2)$ be the spectrum of the ring of dual numbers over \mathbf{k} with point $*$: $\mathrm{Spec} \mathbf{k} \rightarrow \mathrm{Spec} \mathbf{k}[\epsilon]/(\epsilon^2)$. The tangent space $T_x X$ is naturally isomorphic to the set of morphisms $\mathrm{Spec} \mathbf{k}[\epsilon]/(\epsilon^2) \rightarrow X$ that send $*$ to x , i.e.

$$T_x X \cong \{f : \mathrm{Spec} \mathbf{k}[\epsilon]/(\epsilon^2) \rightarrow X \mid f(*) = x\}.$$

Proof. **Yang: To be filled.** □

4 Line Bundles and Divisors

4.1 Cartier Divisors

4.2 Line Bundles and Picard Group

Definition 4.1. Let X be a scheme. The *Picard group* of X is defined to be $\mathrm{Pic}(X) = H^1(X, \mathcal{O}_X^*)$. The group operation is given by the tensor product of line bundles.

Definition 4.2. Let X be a scheme over a field \mathbf{k} and $\mathcal{L}, \mathcal{L}'$ two line bundles on X . We say that \mathcal{L} and \mathcal{L}' are *algebraically equivalent* if there exists a **Yang: non-singular** variety T over \mathbf{k} , two points $t_0, t_1 \in T(\mathbf{k})$ and a line bundle \mathcal{M} on $X \times T$ such that

$$\mathcal{M}|_{X \times \{t_0\}} \cong \mathcal{L}, \quad \mathcal{M}|_{X \times \{t_1\}} \cong \mathcal{L}'.$$

We denote it by $\mathcal{L} \sim_{\mathrm{alg}} \mathcal{L}'$. **Yang: To be checked.**

Example 5.4. Let $X = \mathbb{P}_A^m \times_A \mathbb{P}_A^n$ with A a ring and $\mathcal{L} = \pi_1^* \mathcal{O}_{\mathbb{P}^m}(1) \otimes \pi_2^* \mathcal{O}_{\mathbb{P}^n}(1)$, where π_1 and π_2 are the projections. Let T_0, \dots, T_m and S_0, \dots, S_n be the standard coordinates on \mathbb{P}^m and \mathbb{P}^n respectively. Then $\Gamma(X, \mathcal{L})$ is generated by the global sections $T_i S_j = \pi_1^* T_i \otimes \pi_2^* S_j$ for $0 \leq i \leq m$ and $0 \leq j \leq n$.

They induce a morphism $f : X \rightarrow \mathbb{P}_A^{(m+1)(n+1)-1}$. If $A = \mathbf{k}$ is a field, on \mathbf{k} -point level, it is given by

$$([x_0 : \cdots : x_m], [y_0 : \cdots : y_n]) \mapsto [\dots : x_i y_j : \dots],$$

where the coordinates on the right-hand side are indexed by all (i, j) with $0 \leq i \leq m$ and $0 \leq j \leq n$. This is called the *Segre embedding* of $\mathbb{P}^m \times \mathbb{P}^n$ into $\mathbb{P}^{(m+1)(n+1)-1}$.

Definition 5.5. A line bundle \mathcal{L} on a scheme X is *globally generated* if $\Gamma(X, \mathcal{L})$ generates \mathcal{L} , i.e., the natural map $\Gamma(X, \mathcal{L}) \otimes_{\mathcal{O}_X} \mathcal{O}_X \rightarrow \mathcal{L}$ is surjective. **Yang: To be continued.**

Example 5.6. Let

Example 5.7.

Definition 5.8. Let \mathcal{L} be a line bundle on a scheme X . **Yang: To be continued.**

Definition 5.9. A line bundle \mathcal{L} on a scheme X is *ample* if for every coherent sheaf \mathcal{F} on X , there exists $n_0 > 0$ such that for all $n \geq n_0$, $\mathcal{F} \otimes \mathcal{L}^{\otimes n}$ is globally generated. **Yang: To be continued.**

Definition 5.10. A line bundle \mathcal{L} on a scheme X is *very ample* if there exists a closed embedding $i : X \rightarrow \mathbb{P}_A^n$ such that $\mathcal{L} \cong i^* \mathcal{O}(1)$. **Yang: To be continued.**

Theorem 5.11. Let X be a scheme of finite type over a noetherian ring A and \mathcal{L} a line bundle on X . Then the following are equivalent:

- (a) \mathcal{L} is ample;
- (b) for some $n > 0$, $\mathcal{L}^{\otimes n}$ is very ample;
- (c) for all $n \gg 0$, $\mathcal{L}^{\otimes n}$ is very ample.

Yang: To be continued.

Proposition 5.12. Let X be a scheme of finite type over a noetherian ring A and \mathcal{L}, \mathcal{M} line bundles on X . Then we have the following:

- (a) if \mathcal{L} is ample and \mathcal{M} is globally generated, then $\mathcal{L} \otimes \mathcal{M}$ is ample;
- (b) if \mathcal{L} is very ample and \mathcal{M} is globally generated, then $\mathcal{L} \otimes \mathcal{M}$ is very ample;
- (c) if both \mathcal{L} and \mathcal{M} are ample, then so is $\mathcal{L} \otimes \mathcal{M}$;
- (d) if both \mathcal{L} and \mathcal{M} are globally generated, then so $\mathcal{L} \otimes \mathcal{M}$;
- (e) if \mathcal{L} is ample and \mathcal{M} is arbitrary, then for some $n > 0$, $\mathcal{L}^{\otimes n} \otimes \mathcal{M}$ is ample;

Yang: To be continued.

Proof. **Yang: To be continued.**

□

Proposition 5.13. Let X be a scheme of finite type over a noetherian ring A and \mathcal{L} a line bundle on X . Then \mathcal{L} is very ample if and only if the following two conditions hold:

- (a) (separate points) for any two distinct points $x, y \in X$, there exists $s!!!!!!!!!!!!$

Definition 5.18. A line bundle \mathcal{L} on a scheme X is *semiample* if for some $n > 0$, $\mathcal{L}^{\otimes n}$ is base-point free. Yang: To be continued.

Theorem 5.19. Let X be a scheme over a ring A and \mathcal{L} a semiample line bundle on X . Then there exists a morphism $f : X \rightarrow Y$ over A such that $\mathcal{L} \cong f^*\mathcal{O}_Y(1)$ for some very ample line bundle $\mathcal{O}_Y(1)$ on Y . Moreover, $Y = \text{Proj } R(X, \mathcal{L})$ and f is induced by the natural map $R(X, \mathcal{L}) \rightarrow \Gamma(X, \mathcal{L}^{\otimes n})$. Yang: To be continued.

Definition 5.20. A line bundle \mathcal{L} on a scheme X is *big* if the section ring $R(X, \mathcal{L})$ has maximal growth, i.e., there exists $C > 0$ such that

$$\dim \Gamma(X, \mathcal{L}^{\otimes n}) \geq Cn^{\dim X}$$

for all sufficiently large n . Yang: To be continued.

Example 5.21. Let $X = \mathbb{F}_2$ be the second Hirzebruch surface, i.e., the projective bundle $\mathbb{P}(\mathcal{O}_{\mathbb{P}^1} \oplus \mathcal{O}_{\mathbb{P}^1}(2))$ over \mathbb{P}^1 . Let $\pi : X \rightarrow \mathbb{P}^1$ be the projection and E the unique section of π with self-intersection -2 . Yang: To be continued.

6 Differentials and duality

7 Flat, smooth and étale morphisms

7.1 Flat family and Hilbert polynomial

Theorem 7.1. Let $X \subseteq \mathbb{P}_T^n$ be a closed subscheme, where T is a noetherian scheme. Then the following are equivalent:

- (a) $X \rightarrow T$ is flat.
- (b) The Hilbert polynomial of the fiber $X_t \subseteq \mathbb{P}_{k(t)}^n$ is independent of the choice of $t \in T$.

Yang: To be checked.

7.2 Base change and semicontinuity

Theorem 7.2 (Base change of flat morphisms). Let $f : X \rightarrow Y$ be a flat morphism of schemes. Then for any morphism $Y' \rightarrow Y$, the base change morphism $f' : X \times_Y Y' \rightarrow Y'$ is also flat.

Theorem 7.3 (Semicontinuity theorem). Let $f : X \rightarrow Y$ be a proper morphism of noetherian schemes, and let \mathcal{F} be a coherent sheaf on X which is flat over Y . Then for each integer $i \geq 0$, the function

$$h^i : Y \rightarrow \mathbb{Z}, \quad y \mapsto \dim_{k(y)} H^i(X_y, \mathcal{F}_y)$$

Yang: To be continued...

8.2 Blowing up

Definition 8.5. Let X be a scheme and $\mathcal{I} \subset \mathcal{O}_X$ be a quasi-coherent sheaf of ideals. The blowing up of X along \mathcal{I} , denoted by $\mathrm{Bl}_{\mathcal{I}} X$, is defined to be the relative Proj of the Rees algebra of \mathcal{I} :

$$\mathrm{Bl}_{\mathcal{I}} X = \mathrm{Proj}_X \bigoplus_{n=0}^{\infty} \mathcal{I}^n.$$

The projection morphism $\pi : \mathrm{Bl}_{\mathcal{I}} X \rightarrow X$ is projective and for any open subset $U \subset X$, we have $\pi^{-1}(U) \cong \mathrm{Bl}_{\mathcal{I}(U)} U$. The exceptional divisor of the blowing up is defined to be the closed subscheme $E = \pi^{-1}(V(\mathcal{I}))$ of $\mathrm{Bl}_{\mathcal{I}} X$. **Yang: To be continued...**

8.3 Relative ampleness and relative morphisms

9 Finite morphisms and fibrations