

# Definition and First Properties of Schemes

## 1 Locally Ringed Space

**Definition 1.** Let  $X$  be a topological space. A *presheaf* of sets (resp. abelian groups, rings, etc.) on  $X$  is a contravariant functor  $\mathcal{F} : \mathbf{Open}(X) \rightarrow \mathbf{Set}$  (resp.  $\mathbf{Ab}$ ,  $\mathbf{Ring}$ , etc.), where  $\mathbf{Open}(X)$  is the category of open subsets of  $X$  with inclusions as morphisms.

A presheaf  $\mathcal{F}$  is a *sheaf* if sections can be glued uniquely. More precisely, for every open covering  $\{U_i\}_{i \in I}$  of an open set  $U \subset X$  and every family of sections  $s_i \in \mathcal{F}(U_i)$  such that  $s_i|_{U_i \cap U_j} = s_j|_{U_i \cap U_j}$  for all  $i, j \in I$ , there exists a unique section  $s \in \mathcal{F}(U)$  such that  $s|_{U_i} = s_i$  for all  $i \in I$ .

**Example 2.** Let  $X$  be a real (resp. complex) manifold. The assignment  $U \mapsto \mathcal{C}^\infty(U, \mathbb{R})$  (resp.  $U \mapsto \{\text{holomorphic functions on } U\}$ ) defines a sheaf of rings on  $X$ .

**Example 3.** Let  $X$  be a non-connected topological space. The assignment

$$U \mapsto \{\text{constant functions on } U\}$$

defines a presheaf  $\mathcal{C}$  of rings on  $X$  but not a sheaf.

For a concrete example, let  $X = (0, 1) \cup (2, 3)$  with the subspace topology from  $\mathbb{R}$ . Consider the open covering  $\{(0, 1), (2, 3)\}$  of  $X$ . The sections  $s_1 = 1 \in \mathcal{C}((0, 1))$  and  $s_2 = 2 \in \mathcal{C}((2, 3))$  agree on the intersection (which is empty), but there is no global section  $s \in \mathcal{C}(X)$  such that  $s|_{(0, 1)} = s_1$  and  $s|_{(2, 3)} = s_2$ .

**Definition 4.** A *locally ringed space* is a pair  $(X, \mathcal{O}_X)$  where  $X$  is a topological space and  $\mathcal{O}_X$  is a sheaf of rings on  $X$  such that for every  $x \in X$ , the stalk  $\mathcal{O}_{X, x}$  is a local ring.

A *morphism of locally ringed spaces*  $f : (X, \mathcal{O}_X) \rightarrow (Y, \mathcal{O}_Y)$  consists of a continuous map  $f : X \rightarrow Y$  and a morphism of sheaves of rings  $f^\# : \mathcal{O}_Y \rightarrow f_* \mathcal{O}_X$  such that for every  $x \in X$ , the induced map on stalks  $f_x^\# : \mathcal{O}_{Y, f(x)} \rightarrow \mathcal{O}_{X, x}$  is a local homomorphism, i.e., it maps the maximal ideal of  $\mathcal{O}_{Y, f(x)}$  to the maximal ideal of  $\mathcal{O}_{X, x}$ .

**Example 5.** Let  $p$  be a prime number. Then the inclusion  $\mathbb{Z}_{(p)} \rightarrow \mathbb{Q}$  is a homomorphism of local rings but not a local homomorphism. Here  $\mathbb{Z}_{(p)}$  is the localization of  $\mathbb{Z}$  at the prime ideal  $(p)$ .

**Example 6** (Glue morphisms). Let  $f : (X, \mathcal{O}_X) \rightarrow (Y, \mathcal{O}_Y)$  be a morphism of locally ringed spaces. If  $U \subset X$  and  $V \subset Y$  are open subsets such that  $f(U) \subset V$ , then the restriction  $f|_U : (U, \mathcal{O}_X|_U) \rightarrow (V, \mathcal{O}_Y|_V)$  is a morphism of locally ringed spaces. Conversely, if  $\{U_i\}_{i \in I}$  is an open covering of  $X$  and for each  $i \in I$ , we have a morphism  $f_i : (U_i, \mathcal{O}_X|_{U_i}) \rightarrow (Y, \mathcal{O}_Y)$  such that  $f_i|_{U_i \cap U_j} = f_j|_{U_i \cap U_j}$  for all  $i, j \in I$ , then there exists a unique morphism  $f : (X, \mathcal{O}_X) \rightarrow (Y, \mathcal{O}_Y)$  such that  $f|_{U_i} = f_i$  for all  $i \in I$ .

**Example 7** (Glue locally ringed space). We construct a locally ringed space by gluing open subspaces. Let  $(X_i, \mathcal{O}_{X_i})$  be locally ringed spaces for  $i \in I$  and  $(U_{ij}, \mathcal{O}_{X_i}|_{U_{ij}})$  be open subspaces for  $i, j \in I$ . Suppose we have isomorphisms  $\varphi_{ij} : (U_{ij}, \mathcal{O}_{X_i}|_{U_{ij}}) \rightarrow (U_{ji}, \mathcal{O}_{X_j}|_{U_{ji}})$  such that

- (a)  $\varphi_{ii} = \text{id}_{X_i}$  for all  $i \in I$ ;

- (b)  $\varphi_{ij}(U_{ij} \cap U_{ik}) = U_{ji} \cap U_{jk}$  for all  $i, j \in I$ ;
- (c)  $\varphi_{jk} \circ \varphi_{ij} = \varphi_{ik}$  on  $U_{ij} \cap U_{ik}$  for all  $i, j, k \in I$ .

Then there exists a locally ringed space  $(X, \mathcal{O}_X)$  and open immersions  $\psi_i : (X_i, \mathcal{O}_{X_i}) \rightarrow (X, \mathcal{O}_X)$  uniquely up to isomorphism such that

- (a)  $\varphi_i(U_{ij}) = \psi_i(X_i) \cap \psi_j(X_j)$  for all  $i, j \in I$ ;
- (b) the following diagram

$$\begin{array}{ccccc} (U_{ij}, \mathcal{O}_{X_i}|_{U_{ij}}) & \hookrightarrow & (X_i, \mathcal{O}_{X_i}) & \xrightarrow{\psi_i} & (X, \mathcal{O}_X) \\ \varphi_{ij} \downarrow & & & & \downarrow = \\ (U_{ji}, \mathcal{O}_{X_j}|_{U_{ji}}) & \hookrightarrow & (X_j, \mathcal{O}_{X_j}) & \xrightarrow{\psi_j} & (X, \mathcal{O}_X) \end{array}$$

commutes for all  $i, j \in I$ ;

- (c)  $X = \bigcup_{i \in I} \psi_i(X_i)$ .

Such  $(X, \mathcal{O}_X)$  is called *the locally ringed space obtained by gluing the  $(X_i, \mathcal{O}_{X_i})$  along the  $\varphi_{ij}$* .

First  $\varphi_{ij}$  induces an equivalence relation  $\sim$  on the disjoint union  $\coprod_{i \in I} X_i$ . By taking the quotient space, we can glue the underlying topological spaces to get a topological space  $X$ . The structure sheaf  $\mathcal{O}_X$  is given by

$$\mathcal{O}_X(V) := \left\{ (s_i)_{i \in I} \in \prod_{i \in I} \mathcal{O}_{X_i}(\psi_i^{-1}(V)) \mid s_i|_{U_{ij}} = \varphi_{ij}^\#(s_j|_{U_{ji}}) \text{ for all } i, j \in I \right\}.$$

Easy to check that  $(X, \mathcal{O}_X)$  is a locally ringed space and satisfies the required properties. If there is another locally ringed space  $(X', \mathcal{O}_{X'})$  with  $\psi'_i$  satisfying the same properties, then by gluing  $\psi'_i \circ \psi_i^{-1}$  we get an isomorphism  $(X, \mathcal{O}_X) \rightarrow (X', \mathcal{O}_{X'})$ .

## 2 Schemes

Let  $R$  be a ring. Recall that the *spectrum* of  $R$ , denoted by  $\text{Spec } R$ , is the set of all prime ideals of  $R$  equipped with the Zariski topology, where the closed sets are of the form  $V(I) = \{\mathfrak{p} \in \text{Spec } R : I \subset \mathfrak{p}\}$  for some ideal  $I \subset R$ .

For each  $f \in R$ , let  $D(f) = \{\mathfrak{p} \in \text{Spec } R : f \notin \mathfrak{p}\}$ . Such  $D(f)$  is open in  $\text{Spec } R$  and called a *principal open set*.

**Proposition 8.** Let  $R$  be a ring. The collection of principal open sets  $\{D(f) : f \in R\}$  forms a basis for the Zariski topology on  $\text{Spec } R$ .

*Proof.* Yang: To be continued □

Define a sheaf of rings on  $\text{Spec } R$  by

$$\mathcal{O}_{\text{Spec } R}(D(f)) = R[1/f].$$

Then  $(\text{Spec } R, \mathcal{O}_{\text{Spec } R})$  is a locally ringed space.

**Definition 9.** An *affine scheme* is a locally ringed space isomorphic to  $(\operatorname{Spec} R, \mathcal{O}_{\operatorname{Spec} R})$  for some ring  $R$ . A *scheme* is a locally ringed space  $(X, \mathcal{O}_X)$  which admits an open cover  $\{U_i\}_{i \in I}$  such that  $(U_i, \mathcal{O}_X|_{U_i})$  is an affine scheme for each  $i \in I$ .

A *morphism of schemes* is a morphism of locally ringed spaces.

These data form a category, denoted by **Sch**. If we fix a base scheme  $S$ , then an  $S$ -*scheme* is a scheme  $X$  together with a morphism  $X \rightarrow S$ . The category of  $S$ -schemes is denoted by **Sch**/ $S$  or **Sch** $_S$ .

**Theorem 10.** The functor  $\operatorname{Spec} : \mathbf{Ring}^{\text{op}} \rightarrow \mathbf{Sch}$  is fully faithful and induces an equivalence of categories between the category of rings and the category of affine schemes. **Yang:** To be continued

**Definition 11.** A morphism of schemes  $f : X \rightarrow Y$  is an *open immersion* (resp. *closed immersion*) if  $f$  induces an isomorphism of  $X$  onto an open (resp. closed) subscheme of  $Y$ . An *immersion* is a morphism which factors as a closed immersion followed by an open immersion. **Yang:** To be continued

**Definition 12.** Let  $f : X \rightarrow Y$  be a morphism of schemes. The *scheme theoretic image* of  $f$  is the smallest closed subscheme  $Z$  of  $Y$  such that  $f$  factors through  $Z$ . More precisely, if  $Y = \operatorname{Spec} A$  is affine, then the scheme theoretic image of  $f$  is  $\operatorname{Spec}(A/\ker(f^\#))$ , where  $f^\# : A \rightarrow \Gamma(X, \mathcal{O}_X)$  is the induced map on global sections. In general, we can cover  $Y$  by affine open subsets and glue the scheme theoretic images on each affine open subset to get the scheme theoretic image of  $f$ . **Yang:** To be continued

**Example 13** (Glue open subschemes). The construction in [Example 7](#) allows us to glue open subschemes to get a scheme. More precisely, let  $(X_i, \mathcal{O}_{X_i})$  be schemes for  $i \in I$  and  $(U_{ij}, \mathcal{O}_{X_i}|_{U_{ij}})$  be open subschemes for  $i, j \in I$ . Suppose we have isomorphisms  $\varphi_{ij} : (U_{ij}, \mathcal{O}_{X_i}|_{U_{ij}}) \rightarrow (U_{ji}, \mathcal{O}_{X_j}|_{U_{ji}})$  satisfying the cocycle condition as in [Example 7](#). Then the locally ringed space  $(X, \mathcal{O}_X)$  obtained by gluing the  $(X_i, \mathcal{O}_{X_i})$  along the  $\varphi_{ij}$  is a scheme.

### 3 Integral, reduced and irreducible

### 4 Fiber product

### 5 Dimension

### 6 Noetherian and finite type

### 7 Separated and proper