


阿巴阿巴阿巴阿巴阿巴阿巴阿巴阿巴阿巴阿巴
巴阿巴阿巴阿巴阿巴阿巴阿巴阿巴阿巴阿巴
阿巴阿巴!

Elementary Results

1 Nakayama's Lemma

Theorem 1 (Nakayama's Lemma). Let A be a ring and \mathfrak{M} be its Jacobi radical. Suppose M is a finitely generated A -module. If $\mathfrak{a}M = M$ for $\mathfrak{a} \subset \mathfrak{M}$, then $M = 0$.

Proof. Suppose M is generated by x_1, \dots, x_n . Since $M = \mathfrak{a}M$, formally we have $(x_1, \dots, x_n)^T = \Phi(x_1, \dots, x_n)^T$ for $\Phi \in M_n(\mathfrak{a})$. Then $(\Phi - \text{id})(x_1, \dots, x_n)^T = 0$. Note that $\det(\Phi - \text{id}) = 1 + a$ for $a \in \mathfrak{a} \subset \mathfrak{M}$. Then $\Phi - \text{id}$ is invertible and then $M = 0$. \square

Proposition 2 (Geometric form of Nakayama's Lemma). Let $X = \text{Spec } A$ be an affine scheme, $x \in X$ a closed point and \mathcal{F} a coherent sheaf on X . If $a_1, \dots, a_k \in \mathcal{F}(X)$ generate $\mathcal{F}|_x = \mathcal{F} \otimes \kappa(x)$, then there is an open subset $U \subset X$ such that $a_i|_U$ generate $\mathcal{F}(U)$.

Proof. Yang: To be completed. \square

Corollary 3.

Proof. Yang: To be completed. \square

2 Nullstellensatz

Theorem 4 (Noether's Normalization Lemma). Let A be a k -algebra of finite type. Then there is an injection $k[T_1, \dots, T_d] \hookrightarrow A$ such that A is finite over $k[T_1, \dots, T_d]$.

Remark 5. Here A does not need to be integral. For example,

Theorem 6 (Hilbert's Nullstellensatz). Let A be a