Class field theory by Junyi Xie

DRAFT

阿巴阿巴!

Contents

1 March 02: to be completed **1**

2 March 06 **1**

2.1 Frobenius element (continue) 1

2.2 The Chebotarev density theorem 3

3 March 13: Introduction to Class Field Theory **5**

3.1 Local CFT: Lubin-Tate theory 5

1 March 02: to be completed

2 March 06

2.1 Frobenius element (continue)

Proposition 2.1. We have a tower of fields $M/L/K$. Suppose that M/K is Galois. Let $D \in \mathcal{O}_M$ a prime ideal and $\mathfrak{P} = D \cap \mathcal{O}_L$ and $\mathfrak{p} = D \cap \mathcal{O}_K$. Assume that L/K is Galois, then $\mathrm{Frob}_{(D,M/K)}|_L = \mathrm{Frob}_{(\mathfrak{P},L/K)}$.

Corollary 2.2. Let L/K be a Galois extension and $\mathfrak{p}$ a prime of K unramified in L. Then $\mathfrak{p}$ is split completely in L if and only if $\mathrm{Frob}_{(\mathfrak{P},L/K)} = 1$ for one (or equivalently, any) prime $\mathfrak{P}$ of L above $\mathfrak{p}$.

Proposition 2.3. Let L_1, L_2 be Galois extensions of K and $M = L_1L_2$. Then M/K is Galois and there is a natural embedding $j : \mathrm{Gal}(M/K) \hookrightarrow \mathrm{Gal}(L_1/K) \times \mathrm{Gal}(L_2/K)$. Under the embedding j, we have $j(\mathrm{Frob}_{(D,M/K)}) = (\mathrm{Frob}_{(\mathfrak{P},L_1/K)}, \mathrm{Frob}_{(\mathfrak{P},L_2/K)})$.

Hence in the situation of Proposition 2.3, $\mathfrak{p}$ is split completely in M if and only if $\mathfrak{p}$ is split completely in both L_1 and L_2.

Example 2.4. Let $K = \mathbb{Q}[\xi_n]$ with $\xi_n = e^{2\pi i/n}$ and p an odd prime. If $p \mid n$, then p is ramified in K. The Frobenius element $\mathrm{Frob}_{(p,K/\mathbb{Q})}$ is not defined. Suppose that $p \nmid n$. Then $\sigma = \mathrm{Frob}_{(p,K/\mathbb{Q})}$ is the unique element in $\mathrm{Gal}(K/\mathbb{Q}) \cong (\mathbb{Z}/n\mathbb{Z})^\times$ such that $\sigma(\alpha) = \alpha^p \pmod{\mathfrak{p}}$ for all $\alpha \in \mathcal{O}_K$ and all prime $\mathfrak{p}$ of K above p. We claim that σ is given by $\sigma(\xi_n) = \xi_n^p$. Fix a prime $\mathfrak{p}$ of K above p. We have

$$\sigma\left(\sum a_i\xi_n^i\right) = \sum a_i\sigma(\xi_n)^i = \sum a_i\xi_n^{pi} \equiv \sum a_i^p\xi_n^{pi} \equiv \left(\sum a_i\xi_n^i\right)^p \pmod{\mathfrak{p}}.$$

Note that $\sigma = \mathrm{Frob}_{(p,K/\mathbb{Q})}$ has order f where f is the smallest positive integer such that $p^f \equiv 1 \pmod{n}$.

Date: March 14, 2026, **Author**: Tianle Yang, My Homepage

DRAFT

Example 2.5. Let $K = \mathbb{Q}[\sqrt{d}]$ where d is a square-free integer. Let p be a prime unramified in K. Identify $\mathrm{Gal}(K/\mathbb{Q})$ with $\{\pm 1\}$ where -1 acts on K by sending $\sqrt{d}$ to $-\sqrt{d}$. Then by Corollary 2.2 we have

$$\mathrm{Frob}_{(p,K/\mathbb{Q})} = \begin{cases} 1 & \text{if } d \text{ is a quadratic residue modulo } p, \\ -1 & \text{if } d \text{ is not a quadratic residue modulo } p. \end{cases}$$

The quadratic reciprocity law by Frobenius elements Let $K = \mathbb{Q}[\xi]$ with $\xi = \exp(2\pi i/p)$ where p is an odd prime. Then $\mathrm{Gal}(K/\mathbb{Q}) \cong (\mathbb{Z}/p\mathbb{Z})^\times$ is cyclic of order $p-1$. There is a unique subgroup of $\mathrm{Gal}(K/\mathbb{Q})$ of index 2, consisting of the squares in $(\mathbb{Z}/p\mathbb{Z})^\times$, which corresponds to a unique quadratic extension F of $\mathbb{Q}$ contained in K.

We check the ramification in K and F. The only prime ramified in K is p. Hence the only prime ramified in F is p. If $p \equiv 1 \pmod 4$, then p is the only prime ramified in $\mathbb{Q}(\sqrt{p})$. And $\mathbb{Q}(\sqrt{p})$ is the unique quadratic extension of $\mathbb{Q}$ ramified only at p. Hence $F = \mathbb{Q}(\sqrt{p})$. Similarly, if $p \equiv 3 \pmod 4$, then $F = \mathbb{Q}(\sqrt{-p})$. Denote by $d = (-1)^{\frac{p-1}{2}} p$, then $F = \mathbb{Q}(\sqrt{d})$.

Let q be an odd prime different from p. Then q is unramified in F. Consider the Frobenius element $\mathrm{Frob}_{(q,K/\mathbb{Q})} : \xi \mapsto \xi^q$. We have

$$\begin{aligned} & q \text{ is a quadratic residue modulo } p \\ \iff & \mathrm{Frob}_{(q,K/\mathbb{Q})} \in \mathrm{Gal}(K/\mathbb{Q}) \text{is a square} \\ \iff & \mathrm{Frob}_{(q,F/\mathbb{Q})} = \mathrm{Frob}_{(q,K/\mathbb{Q})}|_F \in \mathrm{Gal}(F/\mathbb{Q}) \text{is the identity} \\ \iff & \text{the prime } q \text{ is split completely in } F \\ \iff & d \text{ is a quadratic residue modulo } q \end{aligned}$$

Hence

$$\left(\frac{d}{q}\right) = \left(\frac{(-1)^{\frac{p-1}{2}} p}{q}\right) = \left(\frac{-1}{q}\right)^{\frac{p-1}{2}} \left(\frac{p}{q}\right) = \left(\frac{p}{q}\right).$$

This gives a proof of the part of the quadratic reciprocity law for odd primes.

Theorem 2.6 (Quadratic reciprocity law)**.** Let p and q be two distinct odd primes. Then we have the following relations.

(a) $\left(\frac{p}{q}\right)\left(\frac{q}{p}\right) = (-1)^{\frac{p-1}{2}\cdot\frac{q-1}{2}}$. In other words,

$$\left(\frac{p}{q}\right) = \begin{cases} \left(\frac{q}{p}\right) & \text{if } p \equiv 1 \text{ or } q \equiv 1 \pmod 4, \\ -\left(\frac{q}{p}\right) & \text{if } p \equiv q \equiv 3 \pmod 4. \end{cases}$$

(b) $\left(\frac{-1}{p}\right) = (-1)^{\frac{p-1}{2}}$.

(c) $\left(\frac{2}{p}\right) = (-1)^{\frac{p^2-1}{8}}$.

We finish the proof of quadratic reciprocity law here.

DRAFT

Proof. The part (b) follows from the Fermat's theorem on sums of two squares. Then the part (a) follows from the discussion above.

For the part (c), let ξ be a primitive 8-th root of unity in $\overline{\mathbb{F}_p}$. Set $a = \xi + \xi^{-1}$. From $\xi^4 = -1$ we deduce that

$$X^4 + 1 = (X^2 - \xi^2)(X^2 - \xi^{-2}) \in \overline{\mathbb{F}_p}[X].$$

This implies that $\xi^2 + \xi^{-2} = 0$ and hence $a^2 = 2$. Note that the equality

$$\xi^p + \xi^{-p} = \xi + \xi^{-1}$$

holds if and only if $p \equiv \pm 1 \pmod 8$. Since $\xi^p + \xi^{-p} = a^p$, we have

$$a^p = a \iff p \equiv \pm 1 \pmod 8 \iff a \in \mathbb{F}_p.$$

We are done. □

2.2 The Chebotarev density theorem

Definition 2.7. Let K be a number field and $P = \operatorname{mSpec} \mathcal{O}_K$ the set of finite places of K. For every $S \subset P$, we say that S has *natrual density* δ if

$$\lim_{N\to\infty} \frac{\#\{\mathfrak{p} \in S : \operatorname{Nm}(\mathfrak{p}) \le N\}}{\#\{\mathfrak{p} \in P : \operatorname{Nm}(\mathfrak{p}) \le N\}} = \delta.$$

Theorem 2.8 (Chebotarev density theorem)**.** Let L/K be a Galois extension of number fields with Galois group $G = \operatorname{Gal}(L/K)$. Let C be a conjugacy class in G. Then the set of primes $\mathfrak{p}$ of K such that $\operatorname{Frob}_{(\mathfrak{p},L/K)} = C$ has natural density $\frac{|C|}{|G|}$.

Proof of Theorem 2.8 will be given later.

Corollary 2.9. Let L/K be a Galois extension of number fields. The set of primes of K that split completely in L has natural density $\frac{1}{[L:K]}$.

Remark 2.10. There exists a bound for the error in terms of discriminant of the polynomial, but it is large. Given a polynomial $f \in \mathbb{Z}[X]$, there exists a bound B such that if a given cycles type does not appear as the Frobenius element for any prime $p \le B$, then it does not appear as the Frobenius element for any prime p.

Here, by "cycle type" we mean the cycle type of the Frobenius element as a permutation of the roots of $f \pmod p$. It is equivalent to the factorization type of $f \pmod p$.

Example 2.11. Let $K = \mathbb{Q}[\xi_n]$, $\operatorname{Gal}(K/\mathbb{Q}) \cong (\mathbb{Z}/n\mathbb{Z})^\times$. We know that $\operatorname{Frob}_{(p,K/\mathbb{Q})}$ corresponds to $p \in (\mathbb{Z}/n\mathbb{Z})^\times$ for any prime p unramified in K. The Chebotarev density theorem implies that the set of primes p such that $p \equiv a \pmod n$ are equi-distributed of density $\frac{1}{\varphi(n)}$ for each $a \in (\mathbb{Z}/n\mathbb{Z})^\times$. This implies the Dirichlet's theorem on arithmetic progressions.

Application Let L/K be a Galois extension of number fields with Galois group G. Denote by

$$\operatorname{Spl}(L/K) = \{\mathfrak{p} \in \operatorname{mSpec} O_K : \mathfrak{p} \text{ is split completely in } L\}$$

the set of primes of K that split completely in L.

Theorem 2.12. If L/K and M/K is Galois extensions of number fields. Then $L \subset M$ if and only if $\mathrm{Spl}(M/K) \subset \mathrm{Spl}(L/K)$.

Proof. ($\Rightarrow$) obvious.

($\Leftarrow$) We have

$$\mathrm{Spl}(LM/K) = \mathrm{Spl}(L/K) \cap \mathrm{Spl}(M/K) = \mathrm{Spl}(M/K)$$

by Proposition 2.3. By the Chebotarev density theorem, we know that

$$\frac{1}{[LM : K]} = \text{ density of } \mathrm{Spl}(LM/K) = \text{ density of } \mathrm{Spl}(M/K) = \frac{1}{[M : K]}.$$

Hence $LM = M$ and then $L \subset M$. □

Corollary 2.13. Same as Theorem 2.12. Then

$$L = M \iff \mathrm{Spl}(L/K) = \mathrm{Spl}(M/K).$$

Slogan *We can use the information on base field (e.g. the splitting of primes) to determine the extension field.*

Theorem 2.14. For any number field K, integer N and finite set S of primes of K, there are only finite many fields extensions L/K of degree N unramified outside S.

Remark 2.15. Given a compact Riemann surface X and $P_1, \dots, P_n$ points on X, there are only finite many étale cover of $X \setminus \{P_1, \dots, P_n\}$ of degree N. This is since $\pi_1(X \setminus \{P_1, \dots, P_n\})$ is finitely generated.

For a finitely generated group G and integer N, there are only finite many subgroups of G of index N. This is since the subgroups of G of index N corresponds a homomorphism $G \to S_N$ by $G \curvearrowright G/H$ and there are only finite many homomorphisms $G \to S_N$.

Proof. Recall for all prime v and integer N, there are only finite many extensions of K_v of degree dividing N. This since every extension of K_v can be decomposed into an unramified extension and a totally ramified extension. For the unramified extension, there are only finite many of them since they are determined by the residue field extension. For the totally ramified extension, they are determined by the Eisenstein polynomials. And there exists M depending only on N such that for two Eisenstein polynomials f, g of degree N with coefficients in $\mathcal{O}_{K_v}$,

$$f \equiv g \pmod{\mathfrak{m}_v^M} \implies f \text{ and } g \text{ define the same extension of } K_v.$$

Hence the assertion follows from the finiteness of $\mathcal{O}_{K_v}/\mathfrak{m}_v^M$.

Next $\mathrm{disc}(L/K) = \prod_{w|v} \mathrm{disc}(L_w/K_v) \in \mathcal{O}_{K_v}$. Let L be an extension of K of degree N unramified outside S. Then $\mathrm{disc}(L/K)$ is bounded. The result follows from the Hermite's theorem below (Theorem 2.16) □

Theorem 2.16 (Hermite)**.** Let K be a number field. There are only finite many extensions of K with given discriminant.

DRAFT

Proof. For any extension L/K, we have $\mathrm{disc}(L/K) = \mathrm{Nm}_{K/\mathbb{Q}}(\mathrm{disc}(L/K)) \cdot \mathrm{disc}_K^{[L:K]}$. Hence we can assume that $K = \mathbb{Q}$ and we want to show that there are only finite many extensions L of $\mathbb{Q}$ with given discriminant d. By Minkowski's theorem, we have

$$|d| > \left(\frac{\pi}{4}\right)^{2n} \frac{n^n}{n!}.$$

Hence we may fix $n = [L : \mathbb{Q}]$. Set $D = |d|$.

Let $\sigma_1, \dots, \sigma_r \in M_{L,\infty}$ be the infinite places of L with completion $\mathbb{R}$ and $\sigma_{r+1}, \dots, \sigma_{r+s}$ be the infinite places of L with completion $\mathbb{C}$. Consider $\Sigma : L \to \mathbb{R}^r \times \mathbb{C}^s$ given by $x \mapsto (\sigma_1(x), \dots, \sigma_{r+s}(x))$. Define

$$X := \begin{cases} \{(x_1, \dots, x_{r+s}) : |x_1| < \sqrt{D+1}, \quad |x_i| < 1, i = 2, \dots, r+s\} & \text{if } r \neq 0, \\ \{(x_1, \dots, x_s) : |\operatorname{Re} x_1| < 1, |\operatorname{Im} x_1| < \sqrt{D+1}, \quad |x_i| < 1, i = 2, \dots, s\} & \text{if } r = 0. \end{cases}$$

We can compute the volume of X:

$$\mathrm{vol}(X) = \begin{cases} 2^r \pi^s \sqrt{D+1} & \text{if } r \neq 0, \\ 2\pi^{s-1}\sqrt{D+1} & \text{if } r = 0. \end{cases}$$

In either case, we have $\mathrm{vol}(X) > 2^r\sqrt{D}$. By the Minkowski's theorem on the lattice $\mathcal{O}_L$ and the convex body X, there exists a nonzero element $\alpha \in \mathcal{O}_L$ such that $\Sigma(\alpha) \in X$. All conjugates of α are absolutely bounded by a constant depending only on D. Hence there are only finite many possibilities of such $\alpha \in \mathcal{O}_{\overline{\mathbb{Q}}}$.

By the construction of X, if $\sigma(\alpha) = \alpha$ for some $\sigma \in \mathrm{Gal}(L/\mathbb{Q})$, then σ must be the identity on L. Hence $L = \mathbb{Q}(\alpha)$. □

3 March 13: Introduction to Class Field Theory

Reference: Milne's notes on class field theory

Slogan *CFT is to describe all abelian Galois extensions of a local or global field using objects associated to the field itself.*

For non-abelian extensions $\rightsquigarrow$ Langlands program. In this course, we will only talk about the abelian case, i.e. CFT.

3.1 Local CFT: Lubin-Tate theory

The goal of local CFT is to classify the abelian extensions of a local field. Also it describes the local components of the global Artin map.

Recall a *local field* is a field locally compact with respect to a non-trivial absolute value. It has a classification as following:

(a) K/bbQ_p finite extensions for some p;

(b) $K/bbF_p((t))$ finite extensions for some p;

DRAFT

(c) K/bbR or K/bbC.

When K is NA, we denote by π the uniformier of K. For all $a \in u\pi^m$ with $u \in \mathcal{O}_K^\times$, $m \in \mathbb{Z}$, denote by $\mathrm{ord}_K(a) = m$. We have an exact sequence

$$0 \to U_K \to K^\times \xrightarrow{\mathrm{ord}_K} \mathbb{Z} \to 0$$

where U_K is the unit group of K. The residue field k of K has q elements and $\mathrm{char}\, k = p$. The normalized absolute value on K is $|a| := q^{-\mathrm{ord}_K(a)}$. Let K^{al} denote the separeble algebraic closure of K.

By "extension of K", we always means subfield of K^{al} containing K. Both ord_K, $|\cdot|$ have unique extension to K^{al}.

Statement of the main theorem **Fact**: for two abelian extension L/K and M/K, LM/K is still abelian. Hence we can define

$$K^{\mathrm{ab}} := \bigcup_{L/K \text{abelian}} L.$$

It is still an abelian extension of K. We have

$$\mathrm{Gal}(K^{\mathrm{ab}}/K) = \mathrm{Gal}(K^{\mathrm{al}}/K)/[\mathrm{Gal}(K^{\mathrm{al}}/K), \mathrm{Gal}(K^{\mathrm{al}}/K)],$$

i.e., $\mathrm{Gal}(K^{\mathrm{ab}}/K)$ is the maximal abelian quotient of $\mathrm{Gal}(K^{\mathrm{al}}/K)$. Somehow we get the homological group $H^1(X)$ of a curve by consider the abelization of the fundamental group $\pi^1(X)$ of a topological space.

Let L/K be a finite unramified extension of K. Then L/K is Galois and $\mathrm{Gal}(L/K) \cong \mathrm{Gal}(l/k)$. Hence $\mathrm{Gal}(L/K) =< \sigma >$ for σ the Frobenius element. We also denote it by $\mathrm{Frob}_{L/K}$. The Frobenius is determined by

$$\mathrm{Frob}_{L/K}(\alpha) \equiv \alpha^q \mod \mathfrak{m}, \quad \forall \alpha \in \mathcal{O}_K.$$

Theorem 3.1 (Local reciprocity law)**.** Let K be a NA local field. There exists a unique homomorphism

$$\phi_K : K^\times \to \mathrm{Gal}(K^{\mathrm{ab}}/K)$$

with the following properties.

(a) For all prime element π of K and for all finite unramified extension L/K, $\phi_K(\pi)$ acts on L as $\mathrm{Frob}_{L/K}$.

(b) For all finite abelian extension L/K, $\mathrm{Nm}_{L/K}(L^\times)$ is contained in the kernel of $a \mapsto \phi_K(a)|_L$. And ϕ_K induce an isomorphism $\phi_{L/K} : K^\times/\mathrm{Nm}_{L/K}(L^\times) \to \mathrm{Gal}(L/K)$. In particular, $(K^\times : \mathrm{Nm}_{L/K}(L^\times)) = [L : K]$.

Notation 3.2. Write $\mathrm{Nm}(L^\times) := \mathrm{Nm}_{L/K}(L^\times)$.

We call $\phi_K, \phi_{L/K}$ the *local Artin maps* for K and L/K; or *local reciprocity maps*, denoted by rec_K and $\mathrm{rec}_{L/K}$; and also called the *norm residue map* or *symbol*, denoted by $a \mapsto (a, L/K)$.

Remark 3.3 (Something incorrect)**.** Compare $a \in K^\times$ with $f \in \mathbb{C}((t))^\times$. A non-zero element in $\mathbb{C}((t))^\times$ corresponds to a "formal" map from the infinity small neighborhood of 0 to itself. Topologically, it somehow can be viewed as a "loop" and hence gives an element in the fundamental group. Further more, it gives an element in the covering group of the universal covering. This can

DRAFT

be regarded as a geometry counterpart of the local Artin map.

The counterpart of the uniformier is the element $t \in \mathbb{C}((t))$. And the action of t on the universal covering is the monodromy around 0. Somehow it is the "Frobenius" element in the geometric setting.

Consider the covering tower $\tilde{X} \to Y \to X$. If a loop in X can be lifted to a loop in Y, then its action on the covering group of $Y \to X$ is trivial. This is the geometric counterpart of the condition (b) in Theorem 3.1.

The (b) says that $\forall$ finite abelian extension L of K, we have

$$\begin{tikzcd} K^\times \arrow[r, "\phi_K"] \arrow[d] & \mathrm{Gal}(K^{\mathrm{ab}}/K) \arrow[d] \\ K^\times/\mathrm{Nm}(L^\times) \arrow[r, "\cong", "\phi_{L/K}"'] & \mathrm{Gal}(L/K) \end{tikzcd}$$

Definition 3.4. The subgroups of $K^\times$ of the form $\mathrm{Nm}(L^\times)$ for some finite abelian extension L/K are called *norm groups*.

Corollary 3.5. Let K be a NA local field. Assume there exists a homomorphism $\phi : K^\times \to \mathrm{Gal}(K^{\mathrm{ab}}/K)$ satisfying condition (a) and (b) in Theorem 3.1. Then

(a) the map $L \mapsto \mathrm{Nm}(L^\times)$ is a bijection from the set of finite extension of K and the set of norm groups of $K^\times$;

(b) $L \subset L' \iff \mathrm{Nm}(L^\times) \supset \mathrm{Nm}(L'^\times)$;

(c) $\mathrm{Nm}((LL')^\times) = \mathrm{Nm}(L^\times) \cap \mathrm{Nm}(L'^\times)$;

(d) $\mathrm{Nm}((L \cap L')^\times) = \mathrm{Nm}(L^\times) \cdot \mathrm{Nm}(L'^\times)$;

(e) if a subgroup of $K^\times$ containing a norm group, the itself is a norm group.

Proof. Recall $\mathrm{Nm}_{L'/K} = \mathrm{Nm}_{L/K} \circ \mathrm{Nm}_{L'/K}$. Hence $L \subset L'$ implies $\mathrm{Nm}(L^\times) \subset \mathrm{Nm}(L'^\times)$. This gives one direction of (b). And we get $\mathrm{Nm}((LL')^\times) \subset \mathrm{Nm}(L^\times) \cap \mathrm{Nm}(L'^\times)$. Conversely, if $a \in \mathrm{Nm}(L^\times) \cap \mathrm{Nm}(L'^\times)$, then $\phi_{L/K}(a) = 1 = \phi_{L'/K}(a)$. But $\phi_{LL'/K}(a)|_L = \phi_{L/K}(a) = 1$ and similarly $\phi_{LL'/K}(a)|_{L'} = 1$. Hence $\phi_{LL'/K}(a) = 1$ and $a \in \mathrm{Nm}((LL')^\times)$. We get (c).

When $\mathrm{Nm}(L^\times) \supset \mathrm{Nm}(L'^\times)$, (c) implies $\mathrm{Nm}((LL')^\times) = \mathrm{Nm}(L'^\times)$. As index of a norm group is equal to the corresponding extension degree, we get $LL' = L'$ and hence $L \subset L'$. This proves (b).

Now prove (a). The definition gives surjective of $L \mapsto \mathrm{Nm}(L^\times)$. The (b) gives the injectivity.

Now prove (e). Let $N = \mathrm{Nm}(L^\times)$ be a norm group. Let $I \supset N$ a subgroup $K^\times$. Denote by M the fixed field of $\phi_{L/K}(I) \subset \mathrm{Gal}(L/K)$. Then the local Artin map $\phi_{L/K}$ maps I/N isomorphically onto $\mathrm{Gal}(L/M)$. Consider the commutative diagram

$$\begin{tikzcd} K^\times \arrow[r, "\phi_{L/K}"] \arrow[d, "="'] & \mathrm{Gal}(L/K) \arrow[d, "\theta"] \\ K^\times \arrow[r, "\phi_{M/K}"'] & \mathrm{Gal}(M/K), \end{tikzcd}$$

we get $\mathrm{Nm}(M^\times) = \ker \phi_{M/K} = \ker \theta \circ \phi_{L/K} = I$. Hence I is a norm group.

Finally, we prove (d). The map $L \mapsto \mathrm{Nm}(L^\times)$ is an order inverting bijection. Note that $L \cap L'$

is the largest extension of K which is contained in L and L' while $\mathrm{Nm}(L^\times)\cdot\mathrm{Nm}(L'^\times)$ is the smallest group containing $\mathrm{Nm}(L^\times)$ and $\mathrm{Nm}(L'^\times)$. By (e), $\mathrm{Nm}(L^\times)\cdot\mathrm{Nm}(L'^\times)$ is a norm group. This enforces $\mathrm{Nm}((L\cap L')^\times)=\mathrm{Nm}(L^\times)\cdot\mathrm{Nm}(L'^\times)$. □

Lemma 3.6. Let L be a extension of K. If $\mathrm{Nm}(L^\times)$ is of finite index of K, then it is open.

Proof. The group U_L is compact, then $\mathrm{Nm}(U_L)$ is closed in $K^\times$. By looking at orders, one has $\mathrm{Nm}(L^\times)\cap U_K=\mathrm{Nm}(U_L)$. so $U_K/\mathrm{Nm}(U_L)\hookrightarrow K^\times/\mathrm{Nm}(L^\times)$. Hence $\mathrm{Nm}(U_L)$ is closed of finite index in U_K. Then $\mathrm{Nm}(U_L)$ is open in U_K since the complement is finite union of closed subset. Note U_K is open in $K^\times$, so is $\mathrm{Nm}(U_L)$. As $\mathrm{Nm}(L^\times)$ contains an open subset $\mathrm{Nm}(U_L)$, itself is open by group structure. □

Theorem 3.7 (Local existence theorem)**.** The norm groups in $K^\times$ are exactly the open subgroups of finite index.

Corollary 3.8. The Corollary 3.5 gives a bijection between the set of finite abelian extensions of K and the set of open subgroups of finite index in $K^\times$.

Remark 3.9. The Corollary 3.8 also holds for archimedean local field. The abelian extension of $\mathbb{R}$ are $\mathbb{R}$ and $\mathbb{C}$. The norm subgroups of $\mathbb{R}^\times$ are $\mathbb{R}^\times$ and $\mathbb{R}_{>0}$. Let H be an open subgroup of $\mathbb{R}^\times$ of finite index of $\mathbb{R}$. Then $H\supset\mathbb{R}^{\times m}=\{a^m|a\in\mathbb{R}^\times\}$. But $\mathbb{R}^{\times m}=\mathbb{R}^\times$ or $\mathbb{R}_{>0}$. The unique isomorphism $\mathbb{R}^\times/\mathbb{R}_{>0}\cong\mathrm{Gal}(\mathbb{C}/\mathbb{R})$ gives the local Artin map for $\mathbb{R}$.

Remark 3.10. When $\mathrm{char}\,K=0$, any subgroup of finite index is open.

Indeed, let $H\subset K^\times$ be a finite index subgroup. Then $K^{\times m}\subset H$ for some m. Note that $a\in K^{\times m}$ if and only if $f(X)=X^m-a$ has a solution $u\in K$. Then $f'(u)=m\cdot u^{m-1}\neq 0$. Hence the map defined by f is locally invertible at u by Implicit Function Theorem. After perturbing a a little bit, f still has a solution in K. So $K^{\times m}$ is open and so is H.

DRAFT