

Algebraic Dynamics and Dynamical Itaka Theory

Tianle Yang

base on the joint work with Sheng Meng and Long Wang

ICCM 2025, January 7, 2026



SCHOOL OF
MATHEMATICAL SCIENCES
EAST CHINA NORMAL UNIVERSITY

Kawaguchi-Silverman Conjecture

Work over $\overline{\mathbb{Q}}$. X : smooth projective variety, $f : X \rightarrow X$: surjective endomorphism. H : ample divisor on X . $h : X(\overline{\mathbb{Q}}) \rightarrow \mathbb{R}_{\geq 1}$: a height function associated to H .

Conjecture: Kawaguchi-Silverman Conjecture = KSC

If the orbit $O_f(x) := \{f^n(x) \mid n \geq 0\}$ is Zariski dense in X , then

$$\alpha_f(x) = \delta_f.$$

here,

$$\alpha_f(x) := \lim_{n \rightarrow \infty} h(f^n(x))^{1/n},$$

arithmetic invariant at x ,


$$\delta_f := \lim_{n \rightarrow \infty} ((f^n)^*H \cdot H^{\dim X - 1})^{1/n},$$

geometric invariant of f .

Three orbit conjecture




An example




- $X = \mathbb{P}^2$,
 $f : [x : y : z] \mapsto [x^2 - y^2 : xy : z^2]$,
 $x = [1 : 2 : 1]$.

An example



- $X = \mathbb{P}^2$,
 $f : [x : y : z] \mapsto [x^2 - y^2 : xy : z^2]$,
 $x = [1 : 2 : 1]$.
- $f^*H \sim 2H \Rightarrow \delta_f = 2$.

An example




- $X = \mathbb{P}^2$,
 $f : [x : y : z] \mapsto [x^2 - y^2 : xy : z^2]$,
 $x = [1 : 2 : 1]$.

- $f^*H \sim 2H \Rightarrow \delta_f = 2$.

n	$h(f^n(x))$	
0	$\log 2$	≈ 0.7
1	$\log 3$	≈ 1.1
2	$\log 6$	≈ 1.8
3	$\log 30$	≈ 3.4
4	$\log 779$	≈ 6.7
5	$\log 558441$	≈ 13.2

An example



- $X = \mathbb{P}^2$,
 $f : [x : y : z] \mapsto [x^2 - y^2 : xy : z^2]$,
 $x = [1 : 2 : 1]$.

- $f^*H \sim 2H \Rightarrow \delta_f = 2$.

n	$h(f^n(x))$	
0	$\log 2$	≈ 0.7
1	$\log 3$	≈ 1.1
2	$\log 6$	≈ 1.8
3	$\log 30$	≈ 3.4
4	$\log 779$	≈ 6.7
5	$\log 558441$	≈ 13.2

- It is expected that $\alpha_f(x) = 2$.

Dynamical Itaka Theory



Settings



Main results



Strategies and Techniques



Further problem



Thank You!

