

Algebraic Dynamics and Dynamical Itaka Theory

Tianle Yang

based on the joint work with Sheng Meng and Long Wang

ICCM 2025, January 7, 2026



SCHOOL OF
MATHEMATICAL SCIENCES
EAST CHINA NORMAL UNIVERSITY

Kawaguchi-Silverman Conjecture

Work over $\overline{\mathbb{Q}}$. X : smooth projective variety, $f : X \rightarrow X$: surjective endomorphism. H : ample divisor on X . $h : X(\overline{\mathbb{Q}}) \rightarrow \mathbb{R}_{\geq 1}$: a height function associated to H .

Conjecture: Kawaguchi-Silverman Conjecture = KSC

If the orbit $O_f(x) := \{f^n(x) \mid n \geq 0\}$ is Zariski dense in X , then

$$\alpha_f(x) = \delta_f.$$

here,


$$\alpha_f(x) := \lim_{n \rightarrow \infty} h(f^n(x))^{1/n},$$

arithmetic invariant at x ,

$$\delta_f := \lim_{n \rightarrow \infty} ((f^n)^* H \cdot H^{\dim X - 1})^{1/n},$$


geometric invariant of f .

An example




- $X = \mathbb{P}^2$,
 $f : [x : y : z] \mapsto [x^2 - y^2 : xy : z^2]$,
 $x = [1 : 2 : 1]$.

An example



- $X = \mathbb{P}^2$,
 $f : [x : y : z] \mapsto [x^2 - y^2 : xy : z^2]$,
 $x = [1 : 2 : 1]$.
- $f^*H \sim 2H \Rightarrow (f^n)^*H \sim 2^n H \Rightarrow$
 $\delta_f = 2$.


An example



- $X = \mathbb{P}^2$,
 $f : [x : y : z] \mapsto [x^2 - y^2 : xy : z^2]$,
 $x = [1 : 2 : 1]$.
- $f^*H \sim 2H \Rightarrow (f^n)^*H \sim 2^n H \Rightarrow$
 $\delta_f = 2$.

n	$h(f^n(x))$	
0	$\log 2$	≈ 0.7
1	$\log 3$	≈ 1.1
2	$\log 6$	≈ 1.8
3	$\log 30$	≈ 3.4
4	$\log 779$	≈ 6.7
5	$\log 558441$	≈ 13.2

An example




- $X = \mathbb{P}^2$,
 $f : [x : y : z] \mapsto [x^2 - y^2 : xy : z^2]$,
 $x = [1 : 2 : 1]$.
- $f^*H \sim 2H \Rightarrow (f^n)^*H \sim 2^n H \Rightarrow$
 $\delta_f = 2$.

n	$h(f^n(x))$	
0	$\log 2$	≈ 0.7
1	$\log 3$	≈ 1.1
2	$\log 6$	≈ 1.8
3	$\log 30$	≈ 3.4
4	$\log 779$	≈ 6.7
5	$\log 558441$	≈ 13.2

- It is expected that $\alpha_f(x) = 2$.


Three orbit conjectures



The KSC relates the growth rate of heights along orbits to dynamical degrees.



Three orbit conjectures



The KSC relates the growth rate of heights along orbits to dynamical degrees.

The DML predicts if


$$\#O_f(x) \cap V = \infty,$$

then

$$O_g(y) \subseteq V$$

for some $y = f^r(x)$, $g = f^s$.

Three orbit conjectures



The KSC relates the growth rate of heights along orbits to dynamical degrees.

The DML predicts if

$$\#O_f(x) \cap V = \infty,$$

then

$$O_g(y) \subseteq V$$

for some $y = f^r(x)$, $g = f^s$.

The ZDO states that either


$$\exists x \text{ with } \overline{O_f(x)} = X,$$

or


$$\exists f^n \odot X \dashrightarrow Y \odot \text{id}_Y.$$

Three orbit conjectures

Main known cases:



Three orbit conjectures



Main known cases:


Endomorphisms of projective surfaces [Matsuzawa-Sano-Shibata];

Birational map of projective surfaces [Wang, to be checked];

Int-amplified endomorphisms [Meng-Zhong];

Non-isomorphic surjective endomorphism on threefolds [Meng-Zhang];

Three orbit conjectures




Main known cases:

DML in the étale case [Bell-Ghioca-Tucker];

DML for endomorphisms of \mathbb{A}^2 [Xie];

Three orbit conjectures



Main known cases:

DML in the étale case [Bell-Ghioca-Tucker];

DML for endomorphisms of \mathbb{A}^2 [Xie];

The ZDO for endomorphisms of surfaces [Xie, Jia-Xie-Zhang];

The ZDO for automorphisms of threefolds with positive entropy [Matsuzawa-Xie]

Dynamical Iitaka Theory

Coarse classification of
varieties via Kodaira
dimension $\kappa(X, K_X)$:

$\kappa(X, K_X)$	Typical geometry
$-\infty$	uniruled
0	Calabi-Yau
$0 < \kappa(X) < \dim X$	fibrations type
$\dim X$	general type

Dynamical Itaka Theory

Coarse classification of varieties via Kodaira dimension $\kappa(X, K_X)$:

$\kappa(X, K_X)$	Typical geometry
$-\infty$	uniruled
0	Calabi-Yau
$0 < \kappa(X) < \dim X$	fibrations type
$\dim X$	general type


Dynamical analogue: classify (X, f) via $\kappa_f(X, R_f)$.

$\kappa_f(X, R_f)$	Typical dynamics
0	f is log-étale
$0 < \kappa_f(X, R_f) < \dim X$	fibrations type
$\dim X$	R_{fs} is big for $s \geq 0$

Main results

Settings:

- $\pi : (X, f) \rightarrow (Y, g)$ smooth Fano fibration with X, Y smooth projective and $\rho(X) = \rho(Y) + 1$;
- Y is an abelian variety or of picard number one;



Main results

Settings:

- $\pi : (X, f) \rightarrow (Y, g)$ smooth Fano fibration with X, Y smooth projective and $\rho(X) = \rho(Y) + 1$;
- Y is an abelian variety or of picard number one;

Theorem: [Meng-Wang-Y]

KSC holds for (X, f) under the above settings.

Generalize the following known results:

•

Main results

Settings:

- $\pi : (X, f) \rightarrow (Y, g)$ smooth Fano fibration with X, Y smooth projective and $\rho(X) = \rho(Y) + 1$;
- Y is an abelian variety or of picard number one;

Theorem: [Meng-Wang-Y]

KSC holds for (X, f) under the above settings.

Sketch of idea: If $\delta_f = \delta_g$, then

KSC for abelian varieties
or polarized endomorphisms \Rightarrow KSC for $(Y, g) \Rightarrow$ KSC for (X, f) .

If $\delta_f > \delta_g$, under some extra conditions, we have another fibration by dynamical litaka theory.

Main results

Settings:

- $\pi : (X, f) \rightarrow (Y, g)$ smooth Fano fibration with X, Y smooth projective and $\rho(X) = \rho(Y) + 1$;
- Y is an abelian variety or of picard number one;
- f has a Zariski dense orbit;
- $\delta_f > \delta_g$;
- if Y is of picard number one, then $\delta_g = 1$.

Theorem: [Meng-Wang-Y]

There is equivariant fibrations

$$f \circlearrowleft X \overset{\varphi_{f,R_f}}{\dashrightarrow} Y \circlearrowleft g$$

with $\dim Y > 0$ and g is q -polarized.

Main results

Theorem: [Meng-Zhang]

Suppose that $f^*R_f \equiv qR_f$ for some $q > 1$. Then there exists an f -equivariant fibration (dynamical Iitaka fibration associated to R_f)

$$f \circlearrowleft X \overset{\varphi_{f,R_f}}{\dashrightarrow} Y \circlearrowleft g$$

with $\dim Y = \kappa_f(X, R_f)$ and g is q -polarized.

With help of this result, we only need to show that under our settings,

- $\kappa_f(X, R_f) \geq 1$;
- $f^*R_f \equiv \delta_f R_f$.

Then

KSC for polarized endomorphisms \implies KSC for $(Y, g) \implies$ KSC for (X, f) .

Main results

Theorem: [Meng-Zhang]

Suppose that $f^*R_f \equiv qR_f$ for some $q > 1$. Then there exists an f -equivariant fibration (dynamical Iitaka fibration associated to R_f)

$$f \circlearrowleft X \overset{\varphi_{f,R_f}}{\dashrightarrow} Y \circlearrowleft g$$

with $\dim Y = \kappa_f(X, R_f)$ and g is q -polarized.

With help of this result, we only need to show that under our settings,

- $\kappa_f(X, R_f) \geq 1$; \iff criterion for toric bundles
- $f^*R_f \equiv \delta_f R_f$.

Then

KSC for polarized endomorphisms \implies KSC for $(Y, g) \implies$ KSC for (X, f) .

Main results

Theorem: [Meng-Zhang]

Suppose that $f^*R_f \equiv qR_f$ for some $q > 1$. Then there exists an f -equivariant fibration (dynamical Iitaka fibration associated to R_f)

$$f \circlearrowleft X \overset{\varphi_{f,R_f}}{\dashrightarrow} Y \circlearrowleft g$$

with $\dim Y = \kappa_f(X, R_f)$ and g is q -polarized.

With help of this result, we only need to show that under our settings,

- $\kappa_f(X, R_f) \geq 1$; \iff criterion for toric bundles
- $f^*R_f \equiv \delta_f R_f$. \iff decomposition of cones + no rational curve on Y

Then

KSC for polarized endomorphisms \implies KSC for $(Y, g) \implies$ KSC for (X, f) .

Technique: decomposition of cones

Theorem: [Meng-Wang-Y]

$\pi : (X, f) \rightarrow (Y, g)$ flat;
 X, Y smooth projective;
 $\delta_f > \delta_g; \rho(X) = \rho(Y) + 1.$

\Rightarrow

$\exists D$ nef with $f^*D \equiv \delta_f D$ s.t.
 $\text{Psef}^1(X) = \pi^* \text{Psef}^1(Y) \oplus \mathbb{R}_{\geq 0} D,$
 $\text{Nef}^1(X) = \pi^* \text{Nef}^1(Y) \oplus \mathbb{R}_{\geq 0} D.$

Technique: decomposition of cones

Theorem: [Meng-Wang-Y]


$\pi : (X, f) \rightarrow (Y, g)$ flat;
 X, Y smooth projective;
 $\delta_f > \delta_g; \rho(X) = \rho(Y) + 1.$

\Rightarrow

$\exists D$ nef with $f^*D \equiv \delta_f D$ s.t.
 $\text{Psef}^1(X) = \pi^* \text{Psef}^1(Y) \oplus \mathbb{R}_{\geq 0} D,$
 $\text{Nef}^1(X) = \pi^* \text{Nef}^1(Y) \oplus \mathbb{R}_{\geq 0} D.$

Cones of a general fibration

$\pi : X \rightarrow Y:$



Technique: decomposition of cones

Theorem: [Meng-Wang-Y]


$\pi : (X, f) \rightarrow (Y, g)$ flat;
 X, Y smooth projective;
 $\delta_f > \delta_g; \rho(X) = \rho(Y) + 1$.

\Rightarrow


$\exists D$ nef with $f^*D \equiv \delta_f D$ s.t.
 $\text{Psef}^1(X) = \pi^* \text{Psef}^1(Y) \oplus \mathbb{R}_{\geq 0} D$,
 $\text{Nef}^1(X) = \pi^* \text{Nef}^1(Y) \oplus \mathbb{R}_{\geq 0} D$.

Cones of a general fibration

$\pi : X \rightarrow Y$:



Cones of fibration with dynamical restrictions:




Technique: criterion for toric bundles

Theorem: [Meng-Wang-Y]

Let $\pi : (X, f) \rightarrow (Y, g)$ be a Fano fibration with X smooth. Suppose that \exists reduced divisor D on X with $f^{-1}(D) = D, f^*D \sim qD$ for some $q > 1$ and $K_X + D \equiv_{\pi} 0$. Then $\pi : X \rightarrow Y$ is a toric bundle.

\mathbb{P}^2 as a toric variety with toric boundary $D = D_1 + D_2 + D_3$:



Further questions

Here are two further questions we are interested in:

- generalize the case $\rho(Y) = 1$ (hence $\rho(X) = 2$) to any X with picard number 2;
- study the decomposition of cones with the decomposition of K_X and R_f and use them to study the case R_f is big.

Thank You!

