

Algebraic Dynamics and Dynamical Itaka Theory

Tianle Yang

based on the joint work with Sheng Meng and Long Wang

Undergraduate Forum, ICCM 2025, January 7, 2026



SCHOOL OF
MATHEMATICAL SCIENCES
EAST CHINA NORMAL UNIVERSITY

Kawaguchi-Silverman Conjecture

$\overline{\mathbb{Q}}$. X smooth projective variety; $f : X \rightarrow X$ surjective endomorphism.



Kawaguchi-Silverman Conjecture

$\overline{\mathbb{Q}}$. X smooth projective variety; $f : X \rightarrow X$ surjective endomorphism.

Conjecture: Kawaguchi-Silverman Conjecture = KSC

If the orbit $O_f(x) := \{f^n(x) \mid n \geq 0\}$ is Zariski dense in X , then

$$\alpha_f(x) = \delta_f.$$

Kawaguchi-Silverman Conjecture

$\overline{\mathbb{Q}}$. X smooth projective variety; $f : X \rightarrow X$ surjective endomorphism.

Conjecture: Kawaguchi-Silverman Conjecture = KSC

If the orbit $O_f(x) := \{f^n(x) \mid n \geq 0\}$ is Zariski dense in X , then
$$\alpha_f(x) = \delta_f.$$

Let H be an ample divisor and $h \geq 1$ a height function associated to H .

Kawaguchi-Silverman Conjecture

$\overline{\mathbb{Q}}$. X smooth projective variety; $f : X \rightarrow X$ surjective endomorphism.

Conjecture: Kawaguchi-Silverman Conjecture = KSC

If the orbit $O_f(x) := \{f^n(x) \mid n \geq 0\}$ is Zariski dense in X , then
$$\alpha_f(x) = \delta_f.$$

Let H be an ample divisor and $h \geq 1$ a height function associated to H .

$$\bullet \quad \alpha_f(x) := \lim_{n \rightarrow \infty} h(f^n(x))^{\frac{1}{n}} \quad \text{arithmetic, local invariant at } x;$$

Kawaguchi-Silverman Conjecture

$\overline{\mathbb{Q}}$. X smooth projective variety; $f : X \rightarrow X$ surjective endomorphism.


Conjecture: Kawaguchi-Silverman Conjecture = KSC

If the orbit $O_f(x) := \{f^n(x) \mid n \geq 0\}$ is Zariski dense in X , then
$$\alpha_f(x) = \delta_f.$$

Let H be an ample divisor and $h \geq 1$ a height function associated to H .


- $\alpha_f(x) := \lim_{n \rightarrow \infty} h(f^n(x))^{\frac{1}{n}}$ arithmetic, local invariant at x ;
- $\delta_f := \lim_{n \rightarrow \infty} ((f^n)^*H \cdot H^{\dim X - 1})^{\frac{1}{n}}$ geometric, global invariant of f .

An example




- $X = \mathbb{P}^2$,
 $f : [x : y : z] \mapsto [x^2 - y^2 : xy : z^2]$,
 $x = [1 : 2 : 1]$.

An example



- $X = \mathbb{P}^2$,
 $f : [x : y : z] \mapsto [x^2 - y^2 : xy : z^2]$,
 $x = [1 : 2 : 1]$.
- $f^*H \sim 2H \Rightarrow (f^n)^*H \sim 2^n H \Rightarrow$
 $\delta_f = 2$.


An example



- $X = \mathbb{P}^2$,
 $f : [x : y : z] \mapsto [x^2 - y^2 : xy : z^2]$,
 $x = [1 : 2 : 1]$.
- $f^*H \sim 2H \Rightarrow (f^n)^*H \sim 2^n H \Rightarrow$
 $\delta_f = 2$.

| n | $h(f^n(x))$ | |
|-----|---------------|----------------|
| 0 | $\log 2$ | ≈ 0.7 |
| 1 | $\log 3$ | ≈ 1.1 |
| 2 | $\log 6$ | ≈ 1.8 |
| 3 | $\log 30$ | ≈ 3.4 |
| 4 | $\log 779$ | ≈ 6.7 |
| 5 | $\log 558441$ | ≈ 13.2 |

An example




- $X = \mathbb{P}^2$,
 $f : [x : y : z] \mapsto [x^2 - y^2 : xy : z^2]$,
 $x = [1 : 2 : 1]$.

- $f^*H \sim 2H \Rightarrow (f^n)^*H \sim 2^n H \Rightarrow$
 $\delta_f = 2$.

| n | $h(f^n(x))$ | |
|-----|---------------|----------------|
| 0 | $\log 2$ | ≈ 0.7 |
| 1 | $\log 3$ | ≈ 1.1 |
| 2 | $\log 6$ | ≈ 1.8 |
| 3 | $\log 30$ | ≈ 3.4 |
| 4 | $\log 779$ | ≈ 6.7 |
| 5 | $\log 558441$ | ≈ 13.2 |

- It is expected that $\alpha_f(x) = 2$.


Three orbit conjectures



The KSC relates the growth rate of heights along orbits to dynamical degrees.



Three orbit conjectures



The KSC relates the growth rate of heights along orbits to dynamical degrees.


The DML predicts if

$$\#O_f(x) \cap V = \infty \quad V \text{ a subvariety,}$$

then

$$O_{fr}(f^s(x)) \subseteq V \quad \text{for some } r, s.$$

Three orbit conjectures



The KSC relates the growth rate of heights along orbits to dynamical degrees.

The DML predicts if

$$\#O_f(x) \cap V = \infty \quad V \text{ a subvariety,}$$

then

$$O_{fr}(f^s(x)) \subseteq V \quad \text{for some } r, s.$$

The ZDO states that either


$$\exists x \text{ with } \overline{O_f(x)} = X,$$

or


$$\exists f^n \circ X \dashrightarrow Y \circ \text{id}_Y.$$

Three orbit conjectures

Main known cases:



Three orbit conjectures



Main known cases:

Smooth projective surfaces [Matsuzawa-Sano-Shibata];

Quasi-projective surfaces (assume DML) [Wang];

Smooth projective 3folds with $\deg f > 1$ [Meng-Zhang];

Abelian varieties [Kawaguchi-Silverman];


Hyperkähler manifolds [Lesieutre-Santriano];

Mori dream spaces [Matsuzawa];

Polarized endomorphisms [Kawaguchi-Silverman];

Int-amplified endomorphisms [Meng-Zhong].

Three orbit conjectures



Main known cases:

Étale case

Endomorphisms of \mathbb{A}^2


Projective surfaces in char $p > 0$

[Bell-Ghioca-Tucker];

[Xie];

[Xie-Yang].

Three orbit conjectures



Main known cases:

Étale case

Endomorphisms of \mathbb{A}^2

Projective surfaces in char $p > 0$

[Bell-Ghioca-Tucker];

[Xie];

[Xie-Yang].

Over \mathbb{C} (uncountable field)

[Amerik-Campana];

Existence of infinity orbits

[Amerik];

Projective surfaces

[Xie, Jia-Xie-Zhang];

Automorphisms of threefolds with positive entropy

[Matsuzawa-Xie].

Dynamical Itaka Theory

Coarse classification of
varieties via Kodaira
dimension $\kappa(X, K_X)$:

| $\kappa(X, K_X)$ | Typical geometry |
|--------------------------|------------------|
| $-\infty$ | uniruled |
| 0 | Calabi-Yau |
| $0 < \kappa(X) < \dim X$ | fibrations type |
| $\dim X$ | general type |

Dynamical Itaka Theory

Coarse classification of
varieties via Kodaira
dimension $\kappa(X, K_X)$:

| $\kappa(X, K_X)$ | Typical geometry |
|--------------------------|------------------|
| $-\infty$ | uniruled |
| 0 | Calabi-Yau |
| $0 < \kappa(X) < \dim X$ | fibrations type |
| $\dim X$ | general type |


Dynamical analogue: classify
 (X, f) via $\kappa_f(X, R_f)$.

| $\kappa_f(X, R_f)$ | Typical dynamics |
|---------------------------------|---------------------------------|
| 0 | f is log-étale |
| $0 < \kappa_f(X, R_f) < \dim X$ | fibrations type |
| $\dim X$ | R_{f^s} is big for $s \geq 0$ |

Main results

Settings:

- $\pi : (X, f) \rightarrow (Y, g)$ smooth Fano fibration with X, Y smooth projective and $\rho(X) = \rho(Y) + 1$, for example, \mathbb{P}^n -bundles over Y ;
- Y is an abelian variety or of picard number one;



Main results

Settings:

- $\pi : (X, f) \rightarrow (Y, g)$ smooth Fano fibration with X, Y smooth projective and $\rho(X) = \rho(Y) + 1$, for example, \mathbb{P}^n -bundles over Y ;
- Y is an abelian variety or of picard number one;

Theorem: [Meng-Wang-Y]

KSC holds for (X, f) under the above settings.

Generalize the following known results:

- [Li-Matsuzawa 2021, Theorem 4.1], projective bundles on smooth Fano varieties of Picard number one;
- [Nasserden-Zotine 2021, Theorem 1.4], projective bundles on elliptic curves.

Main results

Settings:

- $\pi : (X, f) \rightarrow (Y, g)$ smooth Fano fibration with X, Y smooth projective and $\rho(X) = \rho(Y) + 1$, for example, \mathbb{P}^n -bundles over Y ;
- Y is an abelian variety or of picard number one;

Theorem: [Meng-Wang-Y]

KSC holds for (X, f) under the above settings.

Sketch of idea: If $\delta_f = \delta_g$, then

KSC for abelian varieties
or polarized endomorphisms \Rightarrow KSC for $(Y, g) \Rightarrow$ KSC for (X, f) .

Main results

Settings:

- $\pi : (X, f) \rightarrow (Y, g)$ smooth Fano fibration with X, Y smooth projective and $\rho(X) = \rho(Y) + 1$, for example, \mathbb{P}^n -bundles over Y ;
- Y is an abelian variety or of picard number one;

Theorem: [Meng-Wang-Y]

KSC holds for (X, f) under the above settings.

Sketch of idea: If $\delta_f = \delta_g$, then

KSC for abelian varieties
or polarized endomorphisms \Rightarrow KSC for $(Y, g) \Rightarrow$ KSC for (X, f) .

If $\delta_f > \delta_g$, under some extra conditions, we have another fibration by dynamical litaka theory.

Main results

Settings:

- $\pi : (X, f) \rightarrow (Y, g)$ smooth Fano fibration with X, Y smooth projective and $\rho(X) = \rho(Y) + 1$, for example, \mathbb{P}^n -bundles over Y ;
- Y is an abelian variety or of picard number one;
- f has a Zariski dense orbit;
- $\delta_f > \delta_g$;
- if Y is of picard number one, then $\delta_g = 1$.

Theorem: [Meng-Wang-Y]

There is equivariant fibrations

$$f \circ X \xrightarrow{\varphi_{f,R_f}} Y \circ g$$

with $\dim Y > 0$ and g is q -polarized.

Main results

Theorem: [Meng-Zhang]

Suppose that $f^*R_f \equiv qR_f$ for some $q > 1$. Then there exists an f -equivariant fibration (dynamical Iitaka fibration associated to R_f)

$$f \circ X \xrightarrow{\varphi_{f,R_f}} Y \circ g$$

with $\dim Y = \kappa_f(X, R_f)$ and g is q -polarized.

Main results

Theorem: [Meng-Zhang]

Suppose that $f^*R_f \equiv qR_f$ for some $q > 1$. Then there exists an f -equivariant fibration (dynamical Iitaka fibration associated to R_f)

$$f \circ X \xrightarrow{\varphi_{f,R_f}} Y \circ g$$

with $\dim Y = \kappa_f(X, R_f)$ and g is q -polarized.

With help of this result, we only need to show that under our settings,

- $\kappa_f(X, R_f) \geq 1$;
- $f^*R_f \equiv \delta_f R_f$.

Main results

Theorem: [Meng-Zhang]

Suppose that $f^*R_f \equiv qR_f$ for some $q > 1$. Then there exists an f -equivariant fibration (dynamical Iitaka fibration associated to R_f)

$$f \circlearrowleft X \overset{\varphi_{f,R_f}}{\dashrightarrow} Y \circlearrowleft g$$

with $\dim Y = \kappa_f(X, R_f)$ and g is q -polarized.

With help of this result, we only need to show that under our settings,

- $\kappa_f(X, R_f) \geq 1$;
- $f^*R_f \equiv \delta_f R_f$.

Then

KSC for polarized endomorphisms \implies KSC for $(Y, g) \implies$ KSC for (X, f) .

Main results

Theorem: [Meng-Zhang]

Suppose that $f^*R_f \equiv qR_f$ for some $q > 1$. Then there exists an f -equivariant fibration (dynamical Iitaka fibration associated to R_f)

$$f \circ X \xrightarrow{\varphi_{f,R_f}} Y \circ g$$

with $\dim Y = \kappa_f(X, R_f)$ and g is q -polarized.

With help of this result, we only need to show that under our settings,

- $\kappa_f(X, R_f) \geq 1$; \Leftarrow criterion for toric bundles
- $f^*R_f \equiv \delta_f R_f$.

Then

KSC for polarized endomorphisms \implies KSC for $(Y, g) \implies$ KSC for (X, f) .

Main results

Theorem: [Meng-Zhang]

Suppose that $f^*R_f \equiv qR_f$ for some $q > 1$. Then there exists an f -equivariant fibration (dynamical Iitaka fibration associated to R_f)

$$f \circ X \xrightarrow{\varphi_{f,R_f}} Y \circ g$$

with $\dim Y = \kappa_f(X, R_f)$ and g is q -polarized.

With help of this result, we only need to show that under our settings,

- $\kappa_f(X, R_f) \geq 1$; \Leftarrow criterion for toric bundles
- $f^*R_f \equiv \delta_f R_f$. \Leftarrow decomposition of cones + no rational curve on Y

Then

KSC for polarized endomorphisms \implies KSC for $(Y, g) \implies$ KSC for (X, f) .

Technique: decomposition of cones

Theorem: [Meng-Wang-Y]

$\pi : (X, f) \rightarrow (Y, g)$ flat;
 X, Y smooth projective;
 $\delta_f > \delta_g; \rho(X) = \rho(Y) + 1.$

\Rightarrow

$\exists D$ nef with $f^*D \equiv \delta_f D$ s.t.
 $\text{Psef}^1(X) = \pi^* \text{Psef}^1(Y) \oplus \mathbb{R}_{\geq 0} D,$
 $\text{Nef}^1(X) = \pi^* \text{Nef}^1(Y) \oplus \mathbb{R}_{\geq 0} D.$

Technique: decomposition of cones

Theorem: [Meng-Wang-Y]


$\pi : (X, f) \rightarrow (Y, g)$ flat;
 X, Y smooth projective;
 $\delta_f > \delta_g; \rho(X) = \rho(Y) + 1.$

\Rightarrow

$\exists D$ nef with $f^*D \equiv \delta_f D$ s.t.
 $\text{Psef}^1(X) = \pi^* \text{Psef}^1(Y) \oplus \mathbb{R}_{\geq 0} D,$
 $\text{Nef}^1(X) = \pi^* \text{Nef}^1(Y) \oplus \mathbb{R}_{\geq 0} D.$

Cones of a general fibration

$\pi : X \rightarrow Y:$



Technique: decomposition of cones

Theorem: [Meng-Wang-Y]


$\pi : (X, f) \rightarrow (Y, g)$ flat;
 X, Y smooth projective;
 $\delta_f > \delta_g; \rho(X) = \rho(Y) + 1$.

\Rightarrow


$\exists D$ nef with $f^*D \equiv \delta_f D$ s.t.
 $\text{Psef}^1(X) = \pi^* \text{Psef}^1(Y) \oplus \mathbb{R}_{\geq 0} D$,
 $\text{Nef}^1(X) = \pi^* \text{Nef}^1(Y) \oplus \mathbb{R}_{\geq 0} D$.

Cones of a general fibration

$\pi : X \rightarrow Y$:



Cones of fibration with dynamical restrictions:



Technique: criterion for toric bundles

Theorem: [Meng-Wang-Y]

Let $\pi : (X, f) \rightarrow (Y, g)$ be a Fano fibration with X smooth. Suppose that \exists reduced divisor D on X with $f^{-1}(D) = D, f^*D \sim qD$ for some $q > 1$ and $K_X + D \equiv_{\pi} 0$. Then $\pi : X \rightarrow Y$ is a toric bundle.

Technique: criterion for toric bundles

Theorem: [Meng-Wang-Y]

Let $\pi : (X, f) \rightarrow (Y, g)$ be a Fano fibration with X smooth. Suppose that \exists reduced divisor D on X with $f^{-1}(D) = D, f^*D \sim qD$ for some $q > 1$ and $K_X + D \equiv_{\pi} 0$. Then $\pi : X \rightarrow Y$ is a toric bundle.

This theorem generalizes the following absolute case ($Y = \{\text{pt}\}$) criterion:

- [Hwang-Nakayama] $\rho(X) = 1$ case;
- [Meng-Zhang] polarized case;
- [Meng-Zhong] int-amplified case.

Technique: criterion for toric bundles


Theorem: [Meng-Wang-Y]

Let $\pi : (X, f) \rightarrow (Y, g)$ be a Fano fibration with X smooth. Suppose that \exists reduced divisor D on X with $f^{-1}(D) = D, f^*D \sim qD$ for some $q > 1$ and $K_X + D \equiv_{\pi} 0$. Then $\pi : X \rightarrow Y$ is a toric bundle.

This theorem generalizes the following absolute case ($Y = \{\text{pt}\}$) criterion:

- [Hwang-Nakayama] $\rho(X) = 1$ case;
- [Meng-Zhang] polarized case;
- [Meng-Zhong] int-amplified case.

The right sides is an example in the absolute case.



$$f : \mathbb{P}^2 \rightarrow \mathbb{P}^2,$$


$$[X : Y : Z] \mapsto [X^q : Y^q : Z^q];$$

$$D = \{XYZ = 0\}.$$

Further questions

Here are two further questions we are interested in:

- weaken the conditions on Y : can we just assume Y is normal projective?
-



Thank You!

