

Algebraic Dynamics and Dynamical Itaka Theory

Tianle Yang

base on the joint work with Sheng Meng and Long Wang

ICCM 2025, January 7, 2026



SCHOOL OF
MATHEMATICAL SCIENCES
EAST CHINA NORMAL UNIVERSITY

Kawaguchi-Silverman Conjecture

Work over $\overline{\mathbb{Q}}$. X : smooth projective variety, $f : X \rightarrow X$: surjective endomorphism. H : ample divisor on X . $h : X(\overline{\mathbb{Q}}) \rightarrow \mathbb{R}_{\geq 1}$: a height function associated to H .

Conjecture: Kawaguchi-Silverman Conjecture = KSC

If the orbit $O_f(x) := \{f^n(x) \mid n \geq 0\}$ is Zariski dense in X , then

$$\alpha_f(x) = \delta_f.$$

here,

$$\alpha_f(x) := \lim_{n \rightarrow \infty} h(f^n(x))^{1/n},$$

arithmetic invariant at x ,

$$\delta_f := \lim_{n \rightarrow \infty} ((f^n)^* H \cdot H^{\dim X - 1})^{1/n},$$

geometric invariant of f .

An example

Let $X = \mathbb{P}^2$ and $f : [x : y : z] \mapsto [x^d : y^d : z^d]$ with $d \geq 2$.

$$[1 : 2 : 3] \xrightarrow{f} [1 : 2^d : 3^d] \xrightarrow{f} [1 : 2^{d^2} : 3^{d^2}] \xrightarrow{f} \dots$$

Three orbit conjecture



Dynamical Itaka Theory



Settings



Main results



Strategies and Techniques



Thank You!

