

Formulaire d'Avis Médical à un Expert en IA

Champ	Valeur
ID du patient	-
Âge	-
Taille (cm)	-
Poids (kg)	-
Sexe	-
Notes médicales	-

Symptômes observés:

Anxiété

Traitements administrés:

Aucun traitement sélectionné

Maladie suspectée:

-

Description:

A common condition characterized by pain irritation or scratchiness in the throat often caused by viral infections such as the common cold or flu but can also result from bacterial infections like strep throat allergens dry air or irritants such as smoke symptoms may include difficulty swallowing redness or swelling of the throat hoarseness and sometimes fever or swollen lymph nodes if untreated bacterial sore throats can lead to complications like rheumatic fever or kidney inflammation while viral cases are typically self-limiting treatment depends on the cause and includes rest hydration pain relievers like acetaminophen or ibuprofen and antibiotics for confirmed bacterial infections prevention involves good hygiene practices avoiding irritants and staying hydrated

RÉSULTATS DE PRÉDICTION IA

Prédiction basée sur les symptômes:

Prédictions IA:

Modèle	Diagnostic
Description	Otitis Media in Children
Symptômes	UTI_in_Children

Prédiction basée sur la description:

Prédictions IA:

Modèle	Diagnostic
Description	Sore Throat
Symptômes	Influenza in Pregnancy

Recommandations d'antibiotiques:

Médicament	Dosage	Durée	Ligne	Âge	Sexe	Poids	Notes
Phenoxymethylp enicillin	666mg every 6 hours OR 500mg every 6 hours	5 days	1st choice	Adult	All	N/A	Avoid in penicillin allergy. Recommend to take 30 ...
Amoxicillin	500mg every 8 hours	5 days	2nd choice	Adult	All	N/A	Avoid in penicillin allergy.
Cefalexin	500mg every 8 hours	5 days	2nd choice	Adult	All	N/A	Cephalosporins should not be used in severe penici...
Clarithromycin	500mg every 12 hours	5 days	2nd choice	Adult	All	N/A	Use with caution in pregnancy. Suitable only in 2n...
Phenoxymethylp enicillin	62.5mg every 6 hours	5 days	1st choice	1-11 months	N/A	N/A	Avoid in penicillin allergy. Take 30 mins before f...
Phenoxymethylp enicillin	125mg every 6 hours	5 days	1st choice	1-5 years	N/A	N/A	Avoid in penicillin allergy. Take 30 mins before f...
Phenoxymethylp enicillin	250mg (Kopen) or 333mg (Calvepen) every 6 hours	5 days	1st choice	6-11 years	N/A	N/A	Avoid in penicillin allergy. Take 30 mins before f...
Phenoxymethylp enicillin	500mg (Kopen) or 666mg (Calvepen) every 6 hours	5 days	1st choice	"e 1 2 years and Adults	N/A	N/A	Avoid in penicillin allergy. Take 30 mins before f...
Amoxicillin	See Amoxicillin dosing for children	5 days	2nd choice	N/A	N/A	N/A	Avoid in penicillin allergy.
Cefalexin	See Cefalexin dosing for children	5 days	2nd choice	N/A	N/A	N/A	Cephalosporins contraindicated in severe penicilli...
Clarithromycin	See Clarithromycin dosing for children	5 days	2nd choice	N/A	N/A	N/A	Check macrolide warnings and drug interactions.
Trimethoprim	3mg/kg every 12 hours (max 4mg/kg)	N/A	N/A	6 weeks-5 months	N/A	5kg (6 w weeks)-7k g (5	N/A

Médicament	Dosage	Durée	Ligne	Âge	Sexe	Poids	Notes
						months)	
Trimethoprim	25-50mg every 12 hours (max 4mg/kg)	N/A	N/A	6 months-3 years	N/A	8kg (6 months)-14kg (3 years)	N/A
Trimethoprim	50-75mg every 12 hours (max 4mg/kg)	N/A	N/A	4-7 years	N/A	16kg (4 years)-23kg (7 years)	N/A
Trimethoprim	75-125mg every 12 hours (max 4mg/kg)	N/A	N/A	8-12 years	N/A	26kg (8 years)-39kg (12 years)	N/A
Trimethoprim	200mg every 12 hours	N/A	N/A	>12 years and Adults	N/A	N/A	N/A