

# Formulaire d'Avis Médical à un Expert en IA

Champ	Valeur
ID du patient	-
Âge	-
Taille (cm)	-
Poids (kg)	-
Sexe	-
Notes médicales	-

## Symptômes observés:

Anxiété

## Traitements administrés:

Aucun traitement sélectionné

## Maladie suspectée:

-

### Description:

A common condition characterized by pain irritation or scratchiness in the throat often caused by viral infections such as the common cold or flu but can also result from bacterial infections like strep throat allergens dry air or irritants such as smoke symptoms may include difficulty swallowing redness or swelling of the throat hoarseness and sometimes fever or swollen lymph nodes if untreated bacterial sore throats can lead to complications like rheumatic fever or kidney inflammation while viral cases are typically self-limiting treatment depends on the cause and includes rest hydration pain relievers like acetaminophen or ibuprofen and antibiotics for confirmed bacterial infections prevention involves good hygiene practices avoiding irritants and staying hydrated

## RÉSULTATS DE PRÉDICTION IA

## Prédiction basée sur les symptômes:

predictions: [object Object]

antibiotics:

precautions:

## Prédiction basée sur la description:

```
{"predictions":{"description_model":"Sore Throat","symptoms_model":"Influenza in Pregnancy"},"antibiotics":[{"drug":"Phenoxymethylpenicillin","dose":"666mg every 6 hours OR 500mg every 6 hours","duration":"5 days","notes":"Avoid in penicillin allergy. Recommend to take 30 minutes before a meal or 2 hours after food.","line of treatment":"1st choice","sexe":"All","age":"Adult","number of choice":"1","weight":null},{"drug":"Amoxicillin","dose":"500mg every 8 hours","duration":"5 days","notes":"Avoid in penicillin allergy.","line of treatment":"2nd choice","sexe":"All","age":"Adult","number of choice":"2","weight":null},{"drug":"Cefalexin","dose":"500mg every 8 hours","duration":"5 days","notes":"Cephalosporins should not be used in severe penicillin allergy.","line of treatment":"2nd choice","sexe":"All","age":"Adult","number of choice":"2","weight":null},{"drug":"Clarithromycin","dose":"500mg every 12 hours","duration":"5 days","notes":"Use with caution in pregnancy. Suitable only in 2nd and 3rd trimester. See macrolide warning and drug interactions. Alternative: Azithromycin 500mg stat then 250mg every 24 hours from Day 2 to Day 5.","line of treatment":"2nd choice","sexe":"All","age":"Adult","number of choice":"2","weight":null},{"drug":"Phenoxymethylpenicillin","dose":"62.5mg every 6 hours","duration":"5 days","notes":"Avoid in penicillin allergy. Take 30 mins before food or 2 hours after.","line of treatment":"1st choice","sexe":null,"age":"1-11 months","number of choice":"1st choice","weight":null},{"drug":"Phenoxymethylpenicillin","dose":"125mg every 6 hours","duration":"5 days","notes":"Avoid in penicillin allergy. Take 30 mins before food or 2 hours after.","line of treatment":"1st choice","sexe":null,"age":"1-5 years","number of choice":"1st choice","weight":null},{"drug":"Phenoxymethylpenicillin","dose":"250mg (Kopen) or 333mg (Calvepen) every 6 hours","duration":"5 days","notes":"Avoid in penicillin allergy. Take 30 mins before food or 2 hours after.","line
```