









SI Fig. S10. CA-inclusive CSP Confusion Matrix Histograms. Calculating CSPs with CAs included in the formula does drastically not change the percentage of allosteric CSPs ~72%. CSPs are calculated using formula described in **Eqn. 2**.



SI Fig. S11. Change in F1 scores when considering CA-inclusive CSPs. This plot shows that over the targets in the CSP DB, the average change in F1 score is negligible when comparing CSPs calculated from N/H shifts vs CSPs calculated with N/H/Ca shifts. CA-inclusive CSPs are calculated following **Eqn. 2**.




SI Fig. S12. CSP DB Confusion Matrix Histograms by closest interchain N-N distance.



SI Fig. S13. CSP DB Confusion Matrix Histograms by closest interchain atom-atom distance.

F1 score distributions for 1D single-atom CSPs (n = 126 targets)



SI Fig. S14. F1 Scores for 1D CSPs. This plot shows there are no statistical differences between distributions of calculated F1 scores from 1D CSPs H/N/CA atom types. CSPs are calculated as root-squared differences after applying an ideal holo offset for a 3D H/N/CA offset grid (see **Methods** for details). Statistical analysis is performed using paired Wilcoxon tests with Holm correction.

Structure Attributes

Select an **attribute** for searching (e.g., *Organism*, *Experimental Method*, *Resolution*). Add more conditions to narrow results. Examples: [Homo sapiens](#), [BRCA1](#), [Insulin](#).

AND

Number of Distinct Protein Entities

2

=

2

NOT

Experimental Method

2

is

SOLUTION NMR

NOT

Number of Distinct Molecular Entities

2

=

2

NOT

Oligomeric State

2

is

Hetero 2-mer

NOT


Attribute

Group


☐ Include CSM

Search


SI Fig. S15. PDB Advanced Search Settings. For the subset of targets manually scraped from the PDB, we restrict our analysis to these settings: entries resolved by Solution NMR, with two and only two distinct protein chains. At the time of curation, this advanced search resulted in 720 hits on the RCSB PDB. From this list of 720, entries were included in the CSP DB if there were N and H chemical shifts deposited in the BMRB for an identical sequence which was evaluated in both apo and holo experiments.



SI Fig. S16. Ideal offsets across the whole dataset for N/H. This heatmap shows the offsets used to align the apo/holo HSQC plots. The majority of targets require small adjustments within 0.025 ppm on the Hydrogen axis and 0.245 ppm on the Nitrogen axis.



SI Fig. S17. CSP significance threshold Histogram. This histogram summarizes the thresholds for determining whether a CSP for a given receptor is significant. For a small number of targets, the calculated significance threshold is significantly higher than one would expect, we attribute this to differences in experimental conditions between the apo and holo NMR experiments which aberrantly perturbs all observed shifts. We prune from the CSPdb any targets with a calculated significance threshold > 0.6 ppm. This includes the following targets: 2LMC, 1OO9, 2L12, 2L00, 2N01, 2MP0, 8VOI, 8U2Y, 6OSW.



SI Fig. S18. F1 vs MCC score comparison across CSP DB targets. This plot shows the correlation between calculated F1 and MCC metrics. MCC is a more reliable metric for larger systems with many True Negatives (small CSPs outside of the binding site) (7).

References

1. N. Juranic, *et al.*, Calmodulin Wraps around Its Binding Domain in the Plasma Membrane Ca²⁺ Pump Anchored by a Novel 18-1 Motif. *J. Biol. Chem.* **285**, 4015–4024 (2010).
2. A. Mondal, *et al.*, Structure Determination of Challenging Protein–Peptide Complexes Combining NMR Chemical Shift Data and Molecular Dynamics Simulations. *J. Chem. Inf. Model.* **63**, 2058–2072 (2023).
3. G. Alvarez, *et al.*, Loop Plasticity Drives Paralog-Specific Recognition in BET ET Domains. *J. Chem. Inf. Model.* **66**, 4685–4695 (2026).
4. S. Gianni, *et al.*, Demonstration of Long-Range Interactions in a PDZ Domain by NMR, Kinetics, and Protein Engineering. *Structure* **14**, 1801–1809 (2006).
5. R. Agrata, D. Komander, Ubiquitin—A structural perspective. *Mol. Cell* **85**, 323–346 (2025).
6. A. Hacısuleyman, B. Erman, Entropy Transfer between Residue Pairs and Allostery in Proteins: Quantifying Allosteric Communication in Ubiquitin. *PLOS Comput. Biol.* **13**, e1005319 (2017).
7. D. Chicco, M. J. Warrens, G. Jurman, The Matthews Correlation Coefficient (MCC) is More Informative Than Cohen's Kappa and Brier Score in Binary Classification Assessment. *IEEE Access* **9**, 78368–78381 (2021).