

# MONOIDAL CATEGORIES

## 1. INTRODUCTION

In this notes we study monoidal categories. After introducing monoidal categories and their basic properties in the first section we discuss monoidal functors and as an excursion (we do not use it in the rest of our notes) we present useful result about monoidal adjoint pairs (Theorem 3.3). Third section contains rigorous and formal proofs of both Mac Lane's coherence and strictness theorems. Next we discuss notion of dual pairs and rigid objects. We also introduce symmetric monoidal categories and symmetric monoidal functors in the closing section.

Throughout the notes our foundations is Tarski-Grothendieck set theory described in [Mon19, Introduction]. We indicate each usage of Tarski's axiom for the reader's convenience.

We explain now some conventions concerning notation that we use in this notes. There are two ways of denoting values of functions. In *prefix notation* a function symbol  $f$  precedes its arguments  $x_1, x_2, \dots, x_n$  and the expression is  $f(x_1, x_2, \dots, x_n)$  (parentheses are standard part of the prefix notation). On the other hand *infix notation* is used when a symbol  $f$  of a function is placed between each pair of arguments  $x_1, x_2, \dots, x_n$  and the expression is  $x_1 f x_2 f \dots f x_n$ . For real life example note that  $x_1 + x_2 + \dots + x_n$  is written in infix notation. Infix notation can be also used in the case of functors. For example let  $\otimes : \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$  be a functor and let  $\mathcal{C}$  be a category. Then using infix notation we can write the value of  $\otimes$  on objects  $X, Y$  of  $\mathcal{C}$  as  $X \otimes Y$ . We can also consider the composition  $\otimes \cdot \langle \otimes, 1_{\mathcal{C}} \rangle : \mathcal{C} \times \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$  and we can write its value on objects  $X, Y$  and  $Z$  of  $\mathcal{C}$  as  $(X \otimes Y) \otimes Z$ .

## 2. MONOIDAL CATEGORIES

**Definition 2.1.** Let  $\mathcal{C}$  be a category. Suppose that  $\otimes : \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$  is a functor (we use infix notation for values of this functor),  $I$  is an object of  $\mathcal{C}$ ,  $\alpha_{X,Y,Z} : X \otimes (Y \otimes Z) \rightarrow (X \otimes Y) \otimes Z$  is an isomorphism natural in objects  $X, Y, Z$  of  $\mathcal{C}$  and  $l_X : I \otimes X \rightarrow X$ ,  $r_X : X \otimes I \rightarrow X$  are isomorphisms natural in object  $X$  of  $\mathcal{C}$ . Assume that *Mac Lane's pentagon*

$$\begin{array}{ccc}
 X \otimes (Y \otimes (Z \otimes T)) & \xrightarrow{1_X \otimes \alpha_{Y,Z,T}} & X \otimes ((Y \otimes Z) \otimes T) \\
 \downarrow \alpha_{X,Y,Z \otimes T} & & \downarrow \alpha_{X,Y \otimes Z, T} \\
 (X \otimes Y) \otimes (Z \otimes T) & & (X \otimes (Y \otimes Z)) \otimes T \\
 \searrow \alpha_{X \otimes Y, Z, T} & & \swarrow \alpha_{X, Y \otimes Z, 1_T} \\
 & ((X \otimes Y) \otimes Z) \otimes T &
 \end{array}$$

is commutative for any objects  $X, Y, Z, T$  in  $\mathcal{C}$  and that *unit triangle*

$$\begin{array}{ccc}
 X \otimes (I \otimes Y) & \xrightarrow{\alpha_{X,I,Y}} & (X \otimes I) \otimes Y \\
 \downarrow 1_X \otimes l_Y & & \downarrow r_X \otimes 1_Y \\
 & X \otimes Y &
 \end{array}$$

is commutative for any objects  $X, Y$  in  $\mathcal{C}$ . Then  $(\otimes, I, \alpha, l, r)$  is called a *monoidal structure on  $\mathcal{C}$*  and  $(\mathcal{C}, \otimes, I, \alpha, r, l)$  is called a *monoidal category*. If  $\alpha, l, r$  are identities, then we say that  $(\mathcal{C}, \otimes, I, \alpha, r, l)$  is a *strict monoidal category*.

**Proposition 2.2.** *Let  $(\mathcal{C}, \otimes, I, \alpha, l, r)$  be a monoidal category. Then triangles*

$$\begin{array}{ccc} I \otimes (X \otimes Y) & \xrightarrow{\alpha_{I,X,Y}} & (I \otimes X) \otimes Y \\ & \searrow l_{X \otimes Y} \quad \swarrow l_X \otimes 1_Y & \\ & X \otimes Y & \end{array} \quad \begin{array}{ccc} X \otimes (Y \otimes I) & \xrightarrow{\alpha_{X,Y,I}} & (X \otimes Y) \otimes I \\ & \searrow 1_X \otimes r_Y \quad \swarrow r_{X \otimes Y} & \\ & X \otimes Y & \end{array}$$

are commutative for every pair  $X, Y$  of objects of  $\mathcal{C}$ .

*Proof.* We prove that the first triangle commutes (commutativity of the second can be proved by the similar method). Pick objects  $X, Y$  and consider the following diagram.

$$\begin{array}{ccccc} (I \otimes I) \otimes (X \otimes Y) & \xleftarrow{\alpha_{I,I,X \otimes Y}} & I \otimes (I \otimes (X \otimes Y)) & \xrightarrow{1_I \otimes \alpha_{I,X,Y}} & I \otimes ((I \otimes X) \otimes Y) \\ & \searrow r_I \otimes 1_{X \otimes Y} & \downarrow 1_I \otimes l_{X \otimes Y} & \swarrow 1_I \otimes (l_X \otimes 1_Y) & \\ & & I \otimes (X \otimes Y) & & \\ & \searrow \alpha_{I \otimes I, X, Y} & \downarrow \alpha_{I, X, Y} & \swarrow \alpha_{I, I \otimes X, Y} & \\ & & (I \otimes X) \otimes Y & & \\ & \swarrow (r_I \otimes 1_X) \otimes 1_Y & & \searrow (1_I \otimes l_X) \otimes 1_Y & \\ ((I \otimes I) \otimes X) \otimes Y & \xleftarrow{\alpha_{I, I, X} \otimes 1_Y} & (I \otimes (I \otimes X)) \otimes Y & & \end{array}$$

First note that all morphism in the diagram are isomorphisms. The outer pentagon in the diagram commutes, since it is an instance of the Mac Lane's pentagon. Moreover, the two triangles denoted by  $\circlearrowleft$  commute, since one is an instance of the unit triangle and the other is an image of an instance of the unit triangle under the functor  $(-) \otimes Y$ . Finally, the two squares denoted by  $\circlearrowleft$  are commutative according to the naturality of  $\alpha$ . This implies that the triangle denoted by  $\circlearrowright$  is commutative. This triangle is precisely the image under the functor  $I \otimes (-)$  of the first triangle in the statement. Since this  $I \otimes (-)$  is an equivalence of categories, it follows that the first triangle in the statement is commutative.  $\square$

Let  $\mathcal{C}$  be a category. By abuse of language we say that  $\mathcal{C}$  is a monoidal category when we have a certain monoidal structure on  $\mathcal{C}$  in mind. Also when we deal with two (or more) monoidal categories  $\mathcal{C}$  and  $\mathcal{D}$  we often use the same symbols to denote their monoidal structures. In these cases (as usual in categorical considerations) context helps us to avoid possible confusion.

## 3. MONOIDAL FUNCTORS, TRANSFORMATIONS AND ADJUNCTIONS

**Definition 3.1.** Let  $\mathcal{C}$  and  $\mathcal{D}$  be monoidal categories. Suppose that  $F : \mathcal{C} \rightarrow \mathcal{D}$  is a functor,  $\tau_{X,Y} : F(X) \otimes F(Y) \rightarrow F(X \otimes Y)$  is an morphism natural in pairs objects  $X, Y$  of  $\mathcal{C}$  and  $\phi : I \rightarrow F(I)$  is a morphism in  $\mathcal{D}$ . Assume that the diagram

$$\begin{array}{ccc}
 F(X) \otimes (F(Y) \otimes F(Z)) & \xrightarrow{\alpha_{F(X),F(Y),F(Z)}} & (F(X) \otimes F(Y)) \otimes F(Z) \\
 \downarrow 1_{F(X)} \otimes \tau_{Y,Z} & & \downarrow \tau_{X,Y} \otimes 1_{F(Z)} \\
 F(X) \otimes F(Y \otimes Z) & & F(X \otimes Y) \otimes F(Z) \\
 \downarrow \tau_{X,Y \otimes Z} & & \downarrow \tau_{X \otimes Y, Z} \\
 F(X \otimes (Y \otimes Z)) & \xrightarrow{F(\alpha_{X,Y,Z})} & F((X \otimes Y) \otimes Z)
 \end{array}$$

is commutative for any objects  $X, Y, Z$  in  $\mathcal{C}$  and diagrams

$$\begin{array}{ccc}
 I \otimes F(X) & & F(X) \otimes I \\
 \downarrow \phi \otimes 1_{F(X)} & \searrow l_{F(X)} & \downarrow 1_{F(X)} \otimes \phi \\
 F(I) \otimes F(X) & & F(X) \otimes F(I) \\
 \downarrow \tau_{I,X} & \nearrow F(l_X) & \downarrow \tau_{X,I} \\
 F(I \otimes X) & & F(X \otimes I) \\
 & \nearrow F(l_X) & \nearrow F(r_X) \\
 & F(X) & F(X)
 \end{array}$$

are commutative for every object  $X$  in  $\mathcal{C}$ . Then a triple  $(F, \tau, \phi)$  is called a *weak monoidal functor*. Moreover, if  $\tau$  is a natural isomorphism and  $\phi$  is an isomorphism, then we say that  $(F, \tau, \phi)$  is a *strong monoidal functor*. Finally, we say that  $(F, \tau, \phi)$  is a *strict monoidal functor* if  $\tau$  and  $\phi$  are identities.

If  $\mathcal{C}$  and  $\mathcal{D}$  are monoidal categories and  $(F, \tau, \phi)$  is a (weak, strong, strict) monoidal functor, where  $F : \mathcal{C} \rightarrow \mathcal{D}$  is the underlying functor, then by the usual abuse of language we say that  $F : \mathcal{C} \rightarrow \mathcal{D}$  is a (weak, strong, strict) monoidal functor.

**Definition 3.2.** Let  $\mathcal{C}$  and  $\mathcal{D}$  be monoidal categories. Suppose that  $(F, \tau^F, \phi^F)$  and  $(G, \tau^G, \phi^G)$  are (weak, strong, strict) monoidal functors. Let  $\sigma : F \rightarrow G$  be a natural transformation. Assume that the square

$$\begin{array}{ccc}
 F(X) \otimes F(Y) & \xrightarrow{\sigma_X \otimes \sigma_Y} & G(X) \otimes G(Y) \\
 \downarrow \tau_{X,Y}^F & & \downarrow \tau_{X,Y}^G \\
 F(X \otimes Y) & \xrightarrow{\sigma_{X \otimes Y}} & G(X \otimes Y)
 \end{array}$$

is commutative for every pair  $X, Y$  of objects in  $\mathcal{C}$  and also that the triangle

$$\begin{array}{ccc}
 F(I) & \xrightarrow{\sigma_I} & G(I) \\
 \nwarrow \phi^F & & \nearrow \phi^G \\
 & I &
 \end{array}$$

commutes. Then  $\sigma$  is a *natural transformation of monoidal functors* or a *monoidal transformation*.

The next theorem is not used in the remaining part of this notes. Therefore, we left some commutative diagram manipulations in its proof as exercises for the reader.

**Theorem 3.3.** *Let  $\mathcal{C}, \mathcal{D}$  be monoidal categories and let  $F : \mathcal{C} \rightarrow \mathcal{D}$  be a strong monoidal functor. Suppose that  $G : \mathcal{D} \rightarrow \mathcal{C}$  is a right adjoint functor to  $F$ . Assume that  $\eta : 1_{\mathcal{C}} \rightarrow GF$  and  $\xi : FG \rightarrow 1_{\mathcal{D}}$  are unit and counit of this adjunction, respectively. Then there exists a unique structure of a weak monoidal functor on  $G$  such that  $\eta$  and  $\xi$  are monoidal transformations.*

*Proof.* Denote the strong monoidal structure on  $F$  by  $(\tau^F, \phi^F)$ . Fix a pair  $X, Y$  of objects of  $\mathcal{D}$ . According to the universal property of  $\xi_{X \otimes Y}$  there exists a unique morphism  $\tau_{X,Y}^G : G(X) \otimes G(Y) \rightarrow G(X \otimes Y)$  that makes the following diagram commutative.

$$\begin{array}{ccc} F(G(X \otimes Y)) & \xleftarrow{F(\tau_{X,Y}^G)} & F(G(X) \otimes G(Y)) \\ \xi_{X \otimes Y} \downarrow & & \downarrow (\tau_{G(X), G(Y)}^F)^{-1} \\ X \otimes Y & \xleftarrow{\xi_X \otimes \xi_Y} & F(G(X)) \otimes F(G(Y)) \end{array}$$

We left to the reader verification that  $\tau_{X,Y}^G$  is natural both in  $X$  and  $Y$ . Next according to the universal property of  $\xi_I$  there exists a unique morphism  $\phi^G : I \rightarrow G(I)$  such that the following triangle is commutative.

$$\begin{array}{ccc} F(G(I)) & \xleftarrow{F(\phi^G)} & F(I) \\ \xi_I \downarrow & \nearrow (\phi^F)^{-1} & \\ I & & \end{array}$$

Now we verify that  $(G, \tau^G, \phi^G)$  is a weak monoidal functor. For this we need to check that for any objects  $X, Y, Z$  in  $\mathcal{D}$  the following diagram

$$\begin{array}{ccc} G(X) \otimes (G(Y) \otimes G(Z)) & \xrightarrow{\alpha_{G(X), G(Y), G(Z)}} & (G(X) \otimes G(Y)) \otimes G(Z) \\ \downarrow 1_{G(X)} \otimes \tau_{Y,Z}^G & & \downarrow \tau_{X,Y}^G \otimes 1_{G(Z)} \\ G(X) \otimes G(Y \otimes Z) & & G(X \otimes Y) \otimes G(Z) \\ \downarrow \tau_{X,Y \otimes Z}^G & & \downarrow \tau_{X \otimes Y, Z}^G \\ G(X \otimes (Y \otimes Z)) & \xrightarrow{G(\alpha_{X,Y,Z})} & G((X \otimes Y) \otimes Z) \end{array}$$

is commutative. Now definition of  $\tau^G$  implies (details are left to the reader) that

$$\xi_{(X \otimes Y) \otimes Z} \cdot F\left(G(\alpha_{X,Y,Z}) \cdot \tau_{X,Y \otimes Z}^G \cdot (1_{G(X)} \otimes \tau_{Y,Z}^G)\right) = \xi_{(X \otimes Y) \otimes Z} \cdot F\left(\tau_{X \otimes Y, Z}^G \cdot (\tau_{X,Y}^G \otimes 1_{G(Z)}) \cdot \alpha_{G(X), G(Y), G(Z)}\right)$$

By universal property of  $\xi_{(X \otimes Y) \otimes Z}$ , we deduce that the hexagonal diagram above is commutative. Using exactly the same trick we show that triangles

$$\begin{array}{ccc}
I \otimes G(X) & & G(X) \otimes I \\
\downarrow \phi^G \otimes 1_{G(X)} & \searrow l_{G(X)} & \downarrow 1_{G(X)} \otimes \phi^G \\
G(I) \otimes G(X) & & G(X) \otimes G(I) \\
\downarrow \tau_{I,X}^G & \nearrow G(l_X) & \downarrow \tau_{X,I}^G \\
G(I \otimes X) & & G(X \otimes I)
\end{array}$$

are commutative for every object  $X$  in  $\mathcal{D}$  (we apply  $F$  and use the universal property of  $\xi_X$ ). This shows that  $(G, \tau^G, \phi^G)$  is a weak monoidal functor. Note also that  $\tau^G$  and  $\phi^G$  are uniquely defined in such a way that  $\xi$  automatically is a monoidal transformation. This proves that  $(\tau^G, \phi^G)$  is the unique structure of a weak monoidal functor on  $G$  that makes  $\xi$  a monoidal transformation. It remains to verify that  $\eta$  is a monoidal transformation, when  $G$  is considered as a monoidal functor with respect to  $(\tau^G, \phi^G)$ . This also can be done by the same trick (apply  $F$  and use universality of  $\xi$ ). For instance we write the complete proof of the fact that  $\eta_I = G(\phi^F) \cdot \phi^G$ . First we apply  $F$  to the right hand side and then we compose it with  $\xi_{F(I)}$ . We obtain

$$\xi_{F(I)} \cdot F(G(\phi^F)) \cdot F(\phi^G) = \phi^F \cdot \xi_I \cdot F(\phi^G) = \phi^F \cdot (\phi^F)^{-1} = 1_{F(I)} = \xi_{F(I)} \cdot F(\eta_I)$$

Thus by universal property of  $\xi_{F(I)}$ , we deduce that  $\eta_I = G(\phi^F) \cdot \phi^G$ .  $\square$

#### 4. COHERENCE AND STRICTNESS FOR MONOIDAL CATEGORIES

The idea of coherence originated in algebraic topology. We refer the reader to interesting and enlightening article [Mac63] for history and explanation of this important concept. Let  $(\mathcal{C}, \otimes, I, \alpha, l, r)$  be a monoidal category. Coherence theorem states that appropriate diagrams involving  $\alpha, l, r$  and identities commute. To make this precise one needs to put considerable effort in constructing these diagrams in a formal way. This is our task in this section.

A magma consists of a set  $S$  equipped with a binary operation (we use infix notation for it)

$$\square : S \times S \rightarrow S$$

and a distinguished element  $e \in S$ . Morphism of magmas is a map of sets preserving binary operations and distinguished elements. The category of magmas is denoted by **Mgm**. According to [BDR94, Corollary 3.7.8] the forgetful functor  $|-| : \mathbf{Mgm} \rightarrow \mathbf{Set}$  admits a left adjoint. This means that for every set there exists a free magma generated by this set.

Now let  $S$  be any set and  $\mathbf{M}_S$  be a free magma generated by this set with operation  $\square$  and distinguished element  $e$ . We define a magma  $\mathbf{A}_S$  and a directed graph

$$\mathbf{A}_S \xrightleftharpoons[t]{s} \mathbf{M}_S$$

in which  $s, t$  are morphisms of magmas. The magma  $\mathbf{A}_S$  is a free magma generated by the set of symbols:

$$1_v \text{ for } v \in \mathbf{M}_S \setminus \{e\}, l_v, l_v^{-1}, r_v, r_v^{-1} \text{ for } v \in \mathbf{M}_S \text{ and } \alpha_{v,w,u}, \alpha_{v,w,u}^{-1} \text{ for } v, w, u \in \mathbf{M}_S.$$

By abuse of language we denote the binary operation of  $\mathbf{A}_S$  by  $\square$ . Its distinguished element is denoted by  $1_e$ . Now it remains to define morphisms  $s, t$ . For this we define

$$s(1_v) = v = t(1_v), s(l_v) = t(l_v^{-1}) = e \square v, t(l_v) = s(l_v^{-1}) = v, s(r_v) = t(r_v^{-1}) = v \square e, t(r_v) = s(r_v^{-1}) = v$$

$$s(\alpha_{v,w,u}) = t(\alpha_{v,w,u}^{-1}) = v \square (w \square u), t(\alpha_{v,w,u}) = s(\alpha_{v,w,u}^{-1}) = (v \square w) \square u$$

for every  $v, w, u \in \mathbf{M}_S$  and we extend these maps of sets to morphisms of magmas according to the fact that  $\mathbf{A}_S$  is free. We also define an automorphism  $i : \mathbf{A}_S \rightarrow \mathbf{A}_S$  by

$$i(1_v) = 1_v, i(l_v) = l_v^{-1}, i(l_v^{-1}) = l_v, i(r_v) = r_v^{-1}, i(r_v^{-1}) = r_v, i(\alpha_{v,w,u}) = \alpha_{v,w,u}^{-1}, i(\alpha_{v,w,u}^{-1}) = \alpha_{v,w,u}$$

for every  $v, w, u \in \mathbf{M}_S$  and we extend this map of sets to a magma endomorphism according to the fact that  $\mathbf{A}_S$  is free. We have  $i^2 = 1_{\mathbf{A}_S}$  and hence  $i$  is an automorphism. Moreover, we have  $s \cdot i = t$  and  $t \cdot i = s$ . Now we construct a groupoid  $\mathbf{Syn}_S$ . Objects of  $\mathbf{Syn}_S$  are elements of  $\mathbf{M}_S$ . Morphisms of  $\mathbf{Syn}_S$  are paths in the directed graph defined above modulo relation that asserts that edges  $1_v$  for  $v \in \mathbf{M}_S$  in the graph are identity morphisms in  $\mathbf{Syn}_S$  and for every edge  $\eta$  in the graph its inverse in  $\mathbf{Syn}_S$  is  $i(\eta)$ . Next  $\square$  define a functor  $\square : \mathbf{Syn}_S \times \mathbf{Syn}_S \rightarrow \mathbf{Syn}_S$  and we have distinguished object  $e$  in  $\mathbf{Syn}_S$ .

**Proposition 4.1.** *Let  $S$  be a set and let the graph*

$$\mathbf{A}_S \begin{array}{c} \xrightarrow{s} \\ \xrightarrow{t} \end{array} \mathbf{M}_S$$

*the automorphism  $i : \mathbf{A}_S \rightarrow \mathbf{A}_S$  and the groupoid  $\mathbf{Syn}_S$  be as defined above. Suppose that  $\mathcal{C}$  is a monoidal category. Then every function  $f$  that assigns to element of  $S$  an object of  $\mathcal{C}$  can be uniquely extended to a functor  $F_f : \mathbf{Syn}_S \rightarrow \mathcal{C}$  such that*

$$F_f(e) = I, F_f(v \square w) = F_f(v) \otimes F_f(w), F_f(l_v) = l_{F_f(v)}, F_f(r_v) = r_{F_f(v)}, F_f(\alpha_{v,w,u}) = \alpha_{F_f(v), F_f(w), F_f(u)}$$

*for any  $v, w, u \in \mathbf{M}_S$ .*

*Proof.* First using [Mon19, Introduction] we may enlarge our base universe so that  $\mathcal{C}$  is a small category. This does not affect construction of  $\mathbf{Syn}_S$  so without loss of generality assume that  $\mathcal{C}$  is small category. Note that  $\otimes$  and  $I$  give rise to a magma structure on the **set** of objects of  $\mathcal{C}$ . This implies that  $f$  can be uniquely extended to a morphism  $F_f : \mathbf{M}_S \rightarrow \text{ob}(\mathcal{C})$  of magmas. This is uniquely defined so that  $F_f(e) = I$  and  $F_f(v \square w) = F_f(v) \otimes F_f(w)$  for every  $v, w \in \mathbf{M}_S$ . We assign

$$F_f(1_v) = 1_{F_f(v)}$$

for  $v \in \mathbf{M}_S \setminus \{e\}$  and

$$F_f(l_v) = l_{F_f(v)}, F_f(l_v^{-1}) = l_v^{-1}, F_f(r_v) = r_{F_f(v)}, F_f(r_v^{-1}) = r_v^{-1}$$

$$F_f(\alpha_{v,w,u}) = \alpha_{F_f(v), F_f(w), F_f(u)}, F_f(\alpha_{v,w,u}^{-1}) = \alpha_{F_f(v), F_f(w), F_f(u)}^{-1}$$

for any  $v, w, u \in \mathbf{M}_S$ . One can also view the **set** of morphisms of  $\mathcal{C}$  as a magma with respect to binary operation  $\otimes$  and  $1_I$ . This implies that  $F_f$  can be extended to a morphism  $F_f : \mathbf{A}_S \rightarrow \text{Mor}(\mathcal{C})$  of magmas. Now  $F_f$  is a morphism of directed graphs

$$\mathbf{A}_S \begin{array}{c} \xrightarrow{s} \\ \xrightarrow{t} \end{array} \mathbf{M}_S$$

and

$$\text{Mor}(\mathcal{C}) \begin{array}{c} \xrightarrow{\text{dom}} \\ \xrightarrow{\text{cod}} \end{array} \text{ob}(\mathcal{C})$$

Moreover,  $F_f(\eta)^{-1}$  is identical to  $F_f(i(\eta))$  for every  $\eta$  in  $\mathbf{A}_S$ . Since morphisms of  $\mathbf{Syn}_S$  are paths in the first graph modulo the relation that asserts that  $1_v$  for  $v \in \mathbf{M}_S$  are identities and for every edge  $\eta$  in the graph its inverse in  $\mathbf{Syn}_S$  is  $i(\eta)$ , we deduce that  $F_f$  can be uniquely extended to a functor  $F_f : \mathbf{Syn}_S \rightarrow \mathcal{C}$  having all properties expressed in the statement.  $\square$

Let  $\mathcal{C}$  be a monoidal category and  $S$  be a subset of the class of its objects. We denote by  $F_S : \mathbf{Syn}_S \rightarrow \mathcal{C}$  the unique functor corresponding to the inclusion of  $S$  into the class of objects in  $\mathcal{C}$  by means of Proposition 4.1.

**Theorem 4.2** (Mac Lane's coherence result). *Let  $\mathcal{C}$  be a monoidal category and  $S$  be a subset of the class its objects. Then the functor  $F_S : \mathbf{Syn}_S \rightarrow \mathcal{C}$  sends any two parallel arrows in  $\mathbf{Syn}_S$  to the same arrow in  $\mathcal{C}$ .*

*Proof.* Suppose that  $\mathcal{D}$  is a monoidal category and suppose that a triple  $(F : \mathcal{C} \rightarrow \mathcal{D}, \tau, \phi)$  is a monoidal functor. Let  $f$  be a function given by the restriction of the functor  $F$  to a set  $S$ . Then  $f$  maps  $S$  into a class of objects of  $\mathcal{D}$ . There exists a unique functor  $F_f : \mathbf{Syn}_S \rightarrow \mathcal{D}$  that extends  $f$  and satisfies properties described in Proposition 4.1. Next for every  $v \in \mathbf{M}_S$  we define an isomorphism  $\sigma_v : F(F_S(v)) \rightarrow F_f(v)$ . This is done by induction. We define  $\sigma_e = \phi$  and  $\sigma_s = 1_{F(s)}$  for every  $s \in S$ . Next if  $\sigma_v$  and  $\sigma_w$  are defined for some  $v, w \in \mathbf{M}_S$ , then we define

$$\sigma_{v \square w} = (\sigma_v \otimes \sigma_w) \cdot \tau_{F_S(v), F_S(w)}$$

Now we prove that for any  $v, w \in \mathbf{M}_S$  and morphism  $\eta : v \rightarrow w$  in  $\mathbf{Syn}_S$  the square

$$(*) \quad \begin{array}{ccc} F(F_S(v)) & \xrightarrow{F(F_S(\eta))} & F(F_S(w)) \\ \sigma_v \downarrow & & \downarrow \sigma_w \\ F_f(v) & \xrightarrow{F_f(\eta)} & F_f(w) \end{array}$$

is commutative. Since each morphism in  $\mathbf{Syn}_S$  can be decomposed into arrows in  $\mathbf{A}_S$ , we derive that it suffices to check commutativity of  $(*)$  for an arrow in  $\mathbf{A}_S$ . Now the proof goes by induction. If  $\eta$  is  $1_v$  for some  $v \in \mathbf{M}_S$  then the commutativity of  $(*)$  boils down to the fact that  $\sigma_v = \sigma_v$ . Next assume that  $\eta = l_v$  for some  $v \in \mathbf{M}_S$ , then we have a commutative diagram

$$\begin{array}{ccc} F(F_S(e \square v)) & \xrightarrow{F(F_S(l_v))} & F(F_S(v)) \\ \downarrow = & & \downarrow = \\ F(I \otimes F_S(v)) & \xrightarrow{F(l_{F_S(v)})} & F(F_S(v)) \\ \downarrow (\phi \otimes 1_{F(F_S(v))}) \cdot \tau_{I, F_S(v)} & & \downarrow 1_{F(F_S(v))} \\ I \otimes F(F_S(v)) & \xrightarrow{l_{F_S(v)}} & F(F_S(v)) \\ \downarrow 1_I \otimes \sigma_v & & \downarrow \sigma_v \\ I \otimes F_f(v) & \xrightarrow{l_{F_f(v)}} & F_f(v) \\ \downarrow = & & \downarrow = \\ F_f(e \square v) & \xrightarrow{F_f(l_v)} & F_f(v) \end{array}$$

Indeed, the commutativity of the top square follows by definition of  $F_S$ , the second square from the top commutes as  $F$  is monoidal, the second square from the bottom commutes, since  $l_X : I \otimes X \rightarrow X$  is natural and finally the bottom square is commutative according to definition of  $F_f$ . Now the outer square in the diagram is an instance of  $(*)$  for  $\eta = l_v$ . This also gives the commutativity of  $(*)$  for  $\eta = l_v^{-1}$ . Similarly one can prove the commutativity of  $(*)$  for  $\eta = r_v$  and  $\eta = r_v^{-1}$ . Now suppose that  $\eta = \alpha_{v,w,u}$  for some  $v, w, u \in \mathbf{M}_S$ . We have a commutative diagram

$$\begin{array}{ccc}
F(F_S(v \square (w \square u))) & \xrightarrow{F(F_S(\alpha_{v,w,u}))} & F(F_S((v \square w) \square u)) \\
\downarrow = & & \downarrow = \\
F(F_S(v) \otimes (F_S(w) \otimes F_S(u))) & \xrightarrow{F(\alpha_{F_S(v), F_S(w), F_S(u)})} & F((F_S(v) \otimes F_S(w)) \otimes F_S(u)) \\
\downarrow \tau_{F_S(v), F_S(w) \otimes F_S(u)} & & \downarrow \tau_{F_S(v) \otimes F_S(w), F_S(u)} \\
F(F_S(v)) \otimes F(F_S(w) \otimes F_S(u)) & & F(F_S(v) \otimes F_S(w)) \otimes F(F_S(u)) \\
\downarrow 1_{F(F_S(v))} \otimes \tau_{F_S(w), F_S(u)} & & \downarrow \tau_{F_S(v), F_S(w)} \otimes 1_{F(F_S(u))} \\
F(F_S(v)) \otimes (F(F_S(w)) \otimes F(F_S(u))) & \xrightarrow{\alpha_{F(F_S(v)), F(F_S(w)), F(F_S(u))}} & (F(F_S(v)) \otimes F(F_S(w))) \otimes F(F_S(u)) \\
\downarrow \sigma_v \otimes (\sigma_w \otimes \sigma_u) & & \downarrow (\sigma_v \otimes \sigma_w) \otimes \sigma_u \\
F_f(v) \otimes (F_f(w) \otimes F_f(u)) & \xrightarrow{\alpha_{F_f(v), F_f(w), F_f(u)}} & (F_f(v) \otimes F_f(w)) \otimes F_f(u) \\
\downarrow = & & \downarrow = \\
F_f(v \square (w \square u)) & \xrightarrow{F_f(\alpha_{v,w,u})} & F_f((v \square w) \square u)
\end{array}$$

Indeed, the first square from the top commutes by definition of  $F_S$ , the second from the top commutes according to the fact that  $F$  is monoidal, the second square from the bottom is commutative, since  $\alpha_{X,Y,Z} : X \otimes (Y \otimes Z) \rightarrow (X \otimes Y) \otimes Z$  is natural and finally the bottom square is commutative by definition of  $F_f$ . Now the outer square is an instance of (\*) for  $\eta = \alpha_{v,w,u}$ . This also gives the commutativity of (\*) for  $\eta = \alpha_{v,w,u}^{-1}$ . Thus we know that (\*) is commutative for  $\eta$  in the generating set of  $\mathbf{A}_S$ . It remains to check that if  $\eta = \beta \square \gamma$  and instances of (\*) commute both for  $\beta$  and  $\gamma$ , then the instance of (\*) for  $\eta$  is commutative. Suppose that  $\beta : v \rightarrow u$ ,  $\gamma : w \rightarrow z$  for some  $v, w, u, z \in \mathbf{M}_S$ . We have a commutative diagram

$$\begin{array}{ccc}
F(F_S(v \square w)) & \xrightarrow{F(F_S(\beta \square \gamma))} & F(F_S(u \square z)) \\
\downarrow = & & \downarrow = \\
F(F_S(v) \otimes F_S(w)) & \xrightarrow{F(F_S(\beta) \otimes F_S(\gamma))} & F(F_S(u) \otimes F_S(z)) \\
\downarrow \tau_{F_S(v), F_S(w)} & & \downarrow \tau_{F_S(u), F_S(z)} \\
F(F_S(v)) \otimes F(F_S(w)) & \xrightarrow{F(F_S(\beta)) \otimes F(F_S(\gamma))} & F(F_S(u)) \otimes F(F_S(z)) \\
\downarrow \sigma_v \otimes \sigma_w & & \downarrow \sigma_u \otimes \sigma_z \\
F_f(v) \otimes F_f(w) & \xrightarrow{F_f(\beta) \otimes F_f(\gamma)} & F_f(u) \otimes F_f(z) \\
\downarrow = & & \downarrow = \\
F_f(v \square w) & \xrightarrow{F_f(\beta \square \gamma)} & F_f(u \square z)
\end{array}$$

Indeed, the first square from the top commutes by definition of  $F_S$ , the second square from the top is commutative according to the fact that  $\tau_{X,Y} : F(X \otimes Y) \rightarrow F(X) \otimes F(Y)$  is natural, the second square from the bottom is commutative, since instances of (\*) for  $\beta$  and  $\gamma$  are commutative and



finally the bottom square is commutative by definition of  $F_f$ . This proves that  $(*)$  is commutative for every morphism in  $\mathbf{Syn}_S$ .

Let  $\eta, \xi : v \rightarrow w$  be parallel morphisms in  $\mathbf{Syn}_S$ . Then commutativity of  $(*)$  for both  $\eta$  and  $\xi$  imply that

$$F(F_S(\eta)) = \sigma_w^{-1} \cdot F_f(\eta) \cdot \sigma_v, \quad F(F_S(\xi)) = \sigma_w^{-1} \cdot F_f(\xi) \cdot \sigma_v$$

If  $\mathcal{D}$  is a strict monoidal category, then  $F_f(v) = F_f(w)$  and

$$F_f(\eta) = 1_{F_f(v)} = 1_{F_f(w)} = F_f(\xi)$$

This last equality follows by decomposing each morphism in  $\mathbf{Syn}_S$  into the composition of arrows in  $\mathbf{A}_S$  and then by induction on complexity of an arrow in  $\mathbf{A}_S$ . Thus if  $\mathcal{D}$  is strict, we derive that  $F(F_S(\eta)) = F(F_S(\xi))$ . Therefore, in order to prove the theorem it suffices to construct a faithful monoidal functor  $F : \mathcal{C} \rightarrow \mathcal{D}$  into a strict monoidal category. For this consider the category  $\mathbf{End}(\mathcal{C}) = \mathbf{Fun}(\mathcal{C}, \mathcal{C})$  of endofunctors of  $\mathcal{C}$ . The functor (in infix notation)

$$\circ : \mathbf{End}(\mathcal{C}) \times \mathbf{End}(\mathcal{C}) \rightarrow \mathbf{End}(\mathcal{C})$$

that sends endofunctors  $F : \mathcal{C} \rightarrow \mathcal{C}$  and  $G : \mathcal{C} \rightarrow \mathcal{C}$  to their composition  $F \circ G$  makes  $\mathbf{End}(\mathcal{C})$  a strict monoidal category with  $1_{\mathcal{C}}$  serving as the unit. We define a functor  $\Phi : \mathcal{C} \rightarrow \mathbf{End}(\mathcal{C})$  by formula  $\Phi(X) = X \otimes (-)$  for object  $X$  in  $\mathcal{C}$  and  $\Phi(f) = f \otimes (-)$  for every morphism  $f$  in  $\mathcal{C}$ . Next we define  $\tau_{X,Y} : \Phi(X \otimes Y) \rightarrow \Phi(X) \circ \Phi(Y)$  for objects  $X, Y$  in  $\mathcal{C}$  by formula  $\tau_{X,Y} = \alpha_{X,Y,-}$ . Finally we define  $\phi : \Phi(I) \rightarrow 1_{\mathcal{C}}$  by formula  $\phi = l$ . A triple  $(\Phi, \tau, \phi)$  is a monoidal functor. Indeed, commutative diagrams asserting the fact that  $(\Phi, \tau, \phi)$  is monoidal are Mac Lane's pentagon, unit triangle and the first triangle in 2.2. The functor  $\Phi$  is faithful. Indeed, if we have  $\Phi(f) = \Phi(g)$  for some parallel morphisms  $f, g$  in  $\mathcal{C}$ , then this implies that  $f \otimes 1_I = g \otimes 1_I$  which implies that  $f = g$ .  $\square$

**Corollary 4.3.** *Let  $(\mathcal{C}, \otimes, I, \alpha, l, r)$  be a monoidal category. Then  $l_I = r_I$ .*

*Proof.* This follows from Theorem 4.2. We have  $l_I = F_{\emptyset}(l_e) = F_{\emptyset}(r_e) = r_I$ .  $\square$

As another consequence of Theorem 4.2 we obtain Mac Lane's strictness theorem.

**Theorem 4.4** (Mac Lane's strictness result). *Let  $\mathcal{C}$  be a monoidal category. Then there exists a strict monoidal category  $\mathbf{Strict}(\mathcal{C})$  and a strict monoidal functor  $F : \mathcal{C} \rightarrow \mathbf{Strict}(\mathcal{C})$  such that  $F$  is full, faithful and surjective on objects.*

*Proof.* According to [Mon19, Introduction] we may enlarge our base universe so that  $\mathcal{C}$  is a small category. Denote by  $I$  unit and by  $\otimes$  functor (as usual we use infix notation for it) in  $\mathcal{C}$ . Let  $S$  be a set of objects of  $\mathcal{C}$ . Let  $\mathbf{Syn}_S$  be the groupoid and  $F_S : \mathbf{Syn}_S \rightarrow \mathcal{C}$  be the functor defined above. Let  $X, Y$  be objects of  $\mathcal{C}$ . We write  $X \sim_{\mathbf{syn}} Y$  if and only if there exists an arrow  $\eta : v \rightarrow w$  in  $\mathbf{Syn}_S$  such that  $F_S(v) = X$  and  $F_S(w) = Y$ . Next suppose that  $f : X \rightarrow Y$  and  $g : Z \rightarrow T$  are morphisms of  $\mathcal{C}$ . Then we write  $f \equiv_{\mathbf{syn}} g$  if and only if there exist arrows  $\beta$  and  $\gamma$  in  $\mathbf{Syn}_S$  such that  $F_S(\beta) \cdot f = g \cdot F_S(\gamma)$ . Since  $\mathbf{Syn}_S$  is a groupoid, both  $\sim_{\mathbf{syn}}, \equiv_{\mathbf{syn}}$  are equivalence relations. Now the following assertions hold.

- (1) Suppose that  $f_1$  and  $f_2$  are morphisms in  $\mathcal{C}$  such that  $\text{cod}(f_1) \sim_{\mathbf{syn}} \text{dom}(f_2)$ . Then there exist  $g_1 \equiv_{\mathbf{syn}} f_1$  and  $g_2 \equiv_{\mathbf{syn}} f_2$  such that  $\text{cod}(g_1) = \text{dom}(g_2)$ .
- (2) If  $f_1 \equiv_{\mathbf{syn}} g_1$ ,  $f_2 \equiv_{\mathbf{syn}} g_2$  and compositions  $f_2 \cdot f_1$ ,  $g_2 \cdot g_1$  exist, then  $f_2 \cdot f_1 \equiv_{\mathbf{syn}} g_2 \cdot g_1$ .
- (3) If  $f$  is a morphism in  $\mathcal{C}$  and  $X \sim_{\mathbf{syn}} \text{dom}(f)$ ,  $Y \sim_{\mathbf{syn}} \text{cod}(f)$ , then there exists a unique morphism  $g : X \rightarrow Y$  in  $\mathcal{C}$  such that  $g \equiv_{\mathbf{syn}} f$ .
- (4) If  $X_1 \sim_{\mathbf{syn}} X_2$  and  $Y_1 \sim_{\mathbf{syn}} Y_2$ , then  $X_1 \otimes Y_1 \sim_{\mathbf{syn}} X_2 \otimes Y_2$ .
- (5) If  $f_1 \equiv_{\mathbf{syn}} g_1$  and  $f_2 \equiv_{\mathbf{syn}} g_2$ , then  $f_1 \otimes f_2 \equiv_{\mathbf{syn}} g_1 \otimes g_2$ .

Indeed, assertions (1), (4) and (5) hold since  $\mathbf{Syn}_S$  is a groupoid and assertions (2), (3) are consequences of Theorem 4.2. Let  $X$  be an object of  $\mathcal{C}$  and let  $f$  be a morphism in  $\mathcal{C}$ . Then we denote by  $[X]_{\sim_{\text{syn}}}$  and  $[f]_{\equiv_{\text{syn}}}$  their equivalence classes. Now we define a category  $\mathbf{Strict}(\mathcal{C})$ . Its objects are equivalence classes of  $\sim_{\text{syn}}$  and its morphisms are equivalence classes of  $\equiv_{\text{syn}}$ . Now if  $f : X \rightarrow Y$  is a morphism in  $\mathcal{C}$ , then  $[f]_{\equiv_{\text{syn}}}$  is a morphism in  $\mathbf{MorStrict}([X]_{\sim_{\text{syn}}}, [Y]_{\sim_{\text{syn}}})$ . Suppose that  $[f_1]_{\equiv_{\text{syn}}}$  and  $[f_2]_{\equiv_{\text{syn}}}$  are morphisms in  $\mathbf{Strict}(\mathcal{C})$  such that  $[\text{cod}(f_1)]_{\sim_{\text{syn}}} = [\text{dom}(f_2)]_{\sim_{\text{syn}}}$  i.e. they form a composable pair of morphisms in  $\mathbf{Strict}(\mathcal{C})$ . Then we define

$$[f_2]_{\equiv_{\text{syn}}} \cdot [f_1]_{\equiv_{\text{syn}}} = [g_2 \cdot g_1]_{\equiv_{\text{syn}}}$$

where  $g_1 \equiv_{\text{syn}} f_1$ ,  $g_2 \equiv_{\text{syn}} f_2$  and  $\text{cod}(g_1) = \text{dom}(g_2)$ . This is always possible by (1) and is well defined operation according to (2). Next the associativity of such defined composition follows from the associativity in  $\mathcal{C}$  and similarly for identities. Hence  $\mathbf{Strict}(\mathcal{C})$  is a category. We also have a functor  $F : \mathcal{C} \rightarrow \mathbf{Strict}(\mathcal{C})$  given by  $X \mapsto [X]_{\sim_{\text{syn}}}$  for every object  $X$  in  $\mathcal{C}$  and  $f \mapsto [f]_{\equiv_{\text{syn}}}$  for every morphism  $f$  in  $\mathcal{C}$ . According to (3) functor  $F$  is full and faithful. Clearly it is surjective on objects. Next according to (4) and (5) the following are well defined

$$[X]_{\sim_{\text{syn}}} \diamond [Y]_{\sim_{\text{syn}}} = [X \otimes Y]_{\sim_{\text{syn}}}, [f]_{\equiv_{\text{syn}}} \diamond [g]_{\equiv_{\text{syn}}} = [f \otimes g]_{\equiv_{\text{syn}}}$$

and these give rise to a functor  $\diamond : \mathbf{Strict}(\mathcal{C}) \times \mathbf{Strict}(\mathcal{C}) \rightarrow \mathbf{Strict}(\mathcal{C})$ . Moreover, by construction we derive that  $F \cdot F_S$  sends every arrow in  $\mathbf{Syn}_S$  to identity in  $\mathbf{Strict}(\mathcal{C})$ . This implies that

$$[\alpha_{X,Y,Z}]_{\equiv_{\text{syn}}} : [X]_{\sim_{\text{syn}}} \diamond ([Y]_{\sim_{\text{syn}}} \diamond [Z]_{\sim_{\text{syn}}}) \rightarrow ([X]_{\sim_{\text{syn}}} \diamond [Y]_{\sim_{\text{syn}}}) \diamond [Z]_{\sim_{\text{syn}}}$$

as well as

$$[l_X]_{\equiv_{\text{syn}}} : [I]_{\sim_{\text{syn}}} \diamond [X]_{\sim_{\text{syn}}} \rightarrow [X]_{\sim_{\text{syn}}}, [r_X]_{\equiv_{\text{syn}}} : [X]_{\sim_{\text{syn}}} \diamond [I]_{\sim_{\text{syn}}} \rightarrow [X]_{\sim_{\text{syn}}}$$

are identity morphisms in  $\mathbf{Strict}(\mathcal{C})$ . Hence  $\mathbf{Strict}(\mathcal{C})$  is a strict monoidal category with respect to  $\diamond$  and unit  $[I]_{\sim_{\text{syn}}}$ . By definition of  $F$  and  $\diamond : \mathbf{Strict}(\mathcal{C}) \times \mathbf{Strict}(\mathcal{C}) \rightarrow \mathbf{Strict}(\mathcal{C})$ , we derive that  $F$  is a strict monoidal functor.  $\square$

## 5. SYMMETRIC MONOIDAL CATEGORIES

**Definition 5.1.** Let  $\mathcal{C}$  be a monoidal category. We denote by  $(\otimes, \alpha, l, r, I)$  its monoidal structure. Let

$$\gamma_{X,Y} : X \otimes Y \rightarrow Y \otimes X$$

be an isomorphism defined and natural for every pair  $X, Y$  of objects in  $\mathcal{C}$ . Suppose that

$$\gamma_{Y,X} \cdot \gamma_{X,Y} = 1_{X \otimes Y}$$

for every pair  $X, Y$  of objects of  $\mathcal{C}$ . Assume also that the diagram

$$\begin{array}{ccc} X \otimes (Y \otimes Z) & \xrightarrow{\alpha_{X,Y,Z}} & (X \otimes Y) \otimes Z \\ \downarrow 1_X \otimes \gamma_{Y,Z} & & \downarrow \gamma_{X,Y} \otimes 1_Z \\ X \otimes (Z \otimes Y) & & (Y \otimes X) \otimes Z \\ \downarrow \alpha_{X,Y,Z} & & \downarrow \alpha_{Y,X,Z}^{-1} \\ (X \otimes Z) \otimes Y & \xrightarrow{\gamma_{X \otimes Z, Y}} & Y \otimes (X \otimes Z) \end{array}$$

is commutative for any objects  $X, Y, Z$  in  $\mathcal{C}$  and the triangle

$$\begin{array}{ccc}
 I \otimes X & \xrightarrow{\gamma_{I,X}} & X \otimes I \\
 \searrow l_X & & \swarrow r_X \\
 & X &
 \end{array}$$

is commutative for every object  $X$  in  $\mathcal{C}$ . Then we say that  $(\mathcal{C}, \otimes, \alpha, l, r, I, \gamma)$  is a *symmetric monoidal category* and  $\gamma$  is called a *symmetry*. Moreover, if  $\alpha, r, l, \gamma$  are identities, then  $(\mathcal{C}, \otimes, \alpha, l, r, I, \gamma)$  is called a *strict symmetric monoidal category*.

Let  $\mathcal{C}$  be a category. By abuse of language we say that  $\mathcal{C}$  is a symmetric monoidal category when we have a certain symmetric monoidal structure on  $\mathcal{C}$  in mind. Also when we deal with two or more symmetric monoidal categories we often use the same symbols to denote their symmetries. In each case it should be clear from the context how to distinguish monoidal structures that we consider.

**Definition 5.2.** Let  $\mathcal{C}$  and  $\mathcal{D}$  be symmetric monoidal categories. Suppose that  $(F, \tau, \phi)$  is a monoidal functor where  $F : \mathcal{C} \rightarrow \mathcal{D}$  is the underlying functor. Assume that the following square is commutative for every pair of objects  $X, Y$  in  $\mathcal{C}$ .

$$\begin{array}{ccc}
 F(X \otimes Y) & \xrightarrow{F(\gamma_{X,Y})} & F(Y \otimes X) \\
 \tau_{X,Y} \downarrow & & \downarrow \tau_{Y,X} \\
 F(X) \otimes F(Y) & \xrightarrow{\gamma_{F(X), F(Y)}} & F(Y) \otimes F(X)
 \end{array}$$

Then  $(F, \tau, \phi)$  is a *symmetric monoidal functor*.

If  $\mathcal{C}$  and  $\mathcal{D}$  are symmetric monoidal categories and  $(F, \tau, \phi)$  is a symmetric monoidal functor where  $F : \mathcal{C} \rightarrow \mathcal{D}$  is the underlying functor, then by the usual abuse of language we say that  $F : \mathcal{C} \rightarrow \mathcal{D}$  is a symmetric monoidal functor.

## 6. ALGEBRAIC STRUCTURES IN CATEGORIES OF PRESHEAVES

Notions like monoid, group, ring, actions of monoid etc. make sense in arbitrary category with finite products. The idea is that each of these algebraic structures can be described in terms of commutativity of certain sets of diagrams involving finite products. For reader's convenience and self-containment we discuss the case of a monoid in detail below. We indicate that our discussion can be effortlessly adapted to arbitrary finitary algebraic theory as defined in BOURCAUX.

**Remark 6.1.** Let  $\mathcal{C}$  be a category with finite products and  $(M, \mu, \eta)$  be a monoid in  $\mathcal{C}$ . Then actions of  $(M, \mu, \eta)$  and their morphisms constitute a category.

**Remark 6.2.** By imposing commutativity of certain diagrams we can similarly define modules over a ring in a category  $\mathcal{C}$  with finite products.

Let  $(M, \mu, \eta)$  be a monoid in a category  $\mathcal{C}$  with finite products. By the usual abuse of notation we often omit part of the data and say that  $M$  is a monoid in  $\mathcal{C}$ . Similar notational convention for groups, rings etc. in  $\mathcal{C}$ .

The category  $\widehat{\mathcal{C}}$  of presheaves on a locally small category  $\mathcal{C}$  is an example of a category with finite products by Corollary . However, for such categories the notion of a monoid can be rephrased differently. This is the content of the next result.

**Fact 6.3.** Let  $\mathcal{C}$  be a locally small category. Then there exists an isomorphism (identification) of categories

$$\mathbf{Mon}(\widehat{\mathcal{C}}) = \mathbf{Fun}(\mathcal{C}^{\text{op}}, \mathbf{Mon})$$

that sends each monoid  $(M, \mu, \eta)$  in  $\widehat{\mathcal{C}}$  to a contravariant functor given by formula

$$\mathcal{C} \ni X \mapsto (M(X), \mu_X, \eta_X) \in \mathbf{Mon}$$

*Proof.* Note that in order for triple  $(M, \mu, \eta)$  to be a monoid in  $\widehat{\mathcal{C}}$  certain diagrams (specified in the definition above) have to commute. This is equivalent with the fact that  $M$  is a presheaf,  $\mu, \eta$  are morphisms of presheaves and for every object  $X$  in  $\mathcal{C}$  the corresponding diagrams in  $\mathbf{Set}$  for  $(M(X), \mu_X, \eta_X)$  commutes. But these conditions are equivalent with the fact that

$$\mathcal{C} \ni X \mapsto (M(X), \mu_X, \eta_X) \in \mathbf{Mon}$$

defines a contravariant functor. Next if  $(M_1, \mu_1, \eta_1)$  and  $(M_2, \mu_2, \eta_2)$  are monoids in  $\widehat{\mathcal{C}}$  and  $f : M_1 \rightarrow M_2$  is a morphism of presheaves, then  $f$  is a morphism of monoids in  $\widehat{\mathcal{C}}$  if and only if for every object  $X$  of  $\mathcal{C}$  map  $f_X : M_1(X) \rightarrow M_2(X)$  is a morphism of monoids  $(M_1(X), \mu_{1X}, \eta_{1X})$  and  $(M_2(X), \mu_{2X}, \eta_{2X})$ .  $\square$

**Remark 6.4.** Actually the proof of Fact 6.3 works without any substantial modifications for any finitary algebraic theory and hence analogical identifications yields isomorphisms of categories

$$\mathcal{D}(\widehat{\mathcal{C}}) = \mathbf{Fun}(\mathcal{C}^{\mathrm{op}}, \mathcal{D})$$

for  $\mathcal{D} = \mathbf{Grp}, \mathbf{Ab}, \mathbf{Ring}, \mathbf{CRing}$ . By virtue of this identifications we interchangeably use terms: monoid (group, ring etc.) in  $\widehat{\mathcal{C}}$  and a presheaf of monoids (groups, rings etc.) on  $\mathcal{C}$ .

## 7. MONOIDS AND ACTIONS

**Definition 7.1.** Let  $\mathcal{C}$  be a monoidal category. A triple  $(M, \mu, \eta)$  consisting of an object  $M$  of  $\mathcal{C}$  and morphisms  $\mu : M \otimes M \rightarrow M, \eta : I \rightarrow M$  such that

$$\begin{array}{ccc} M \otimes M \otimes M & \xrightarrow{1_M \otimes \mu} & M \otimes M \\ \mu \otimes 1_M \downarrow & & \downarrow \mu \\ M \otimes M & \xrightarrow{\mu} & M \end{array} \quad \begin{array}{ccccc} M \otimes I & \xrightarrow{1_M \otimes \eta} & M \otimes M & \xleftarrow{\eta \otimes 1_M} & I \otimes M \\ & \searrow = & \downarrow \mu & \swarrow = & \\ & & M & & \end{array}$$

is called a *monoid in a monoidal category  $\mathcal{C}$* . A monoid object  $(M, \mu, \eta)$  in a symmetric monoidal category  $\mathcal{C}$  is a *commutative monoid in  $\mathcal{C}$*  if the triangle

$$\begin{array}{ccc} M \otimes M & \xrightarrow{s} & M \otimes M \\ & \searrow \mu & \swarrow \mu \\ & M & \end{array}$$

is commutative, where  $s : M \otimes M \rightarrow M \otimes M$  is the symmetry of  $\mathcal{C}$ .

**Definition 7.2.** Let  $\mathcal{C}$  be a monoidal category and let  $(M_1, \mu_1, \eta_1), (M_2, \mu_2, \eta_2)$  be monoids in  $\mathcal{C}$ . Then an arrow  $f : M_1 \rightarrow M_2$  in  $\mathcal{C}$  is a *morphism of monoids* if the following diagrams

$$\begin{array}{ccc} M_1 \otimes M_1 & \xrightarrow{f \otimes f} & M_2 \otimes M_2 \\ \mu_1 \downarrow & & \downarrow \mu_2 \\ M_1 & \xrightarrow{f} & M_2 \end{array} \quad \begin{array}{ccc} M_1 & \xrightarrow{f} & M_2 \\ \eta_1 \swarrow & & \searrow \eta_2 \\ & I & \end{array}$$

are commutative.

**Definition 7.3.** Let  $(M, \mu, \eta)$  be a monoid in a monoidal category  $\mathcal{C}$ . A (left) action of  $M$  on object  $X$  of  $\mathcal{C}$  consists of a morphism  $a : M \otimes X \rightarrow X$  that makes the following diagrams

$$\begin{array}{ccc} M \otimes M \otimes X & \xrightarrow{1_M \otimes a} & M \otimes X \\ \downarrow a & & \downarrow \mu \otimes 1_X \\ M \otimes X & \xrightarrow{a} & X \end{array} \quad \begin{array}{ccc} I \otimes X & \xrightarrow{\eta \otimes 1_X} & M \otimes X \\ & \searrow = & \downarrow a \\ & & X \end{array}$$

commutative.

**Definition 7.4.** Let  $(M, \mu, \eta)$  be a monoid in a monoidal category  $\mathcal{C}$ . Suppose that  $(X, a)$  and  $(Y, b)$  are object of  $\mathcal{C}$  equipped with actions of  $(M, \mu, \eta)$ . Then morphism  $f : X \rightarrow Y$  is a *morphism of actions* of  $(M, \mu, \eta)$  if the following diagram

$$\begin{array}{ccc} M \otimes X & \xrightarrow{1_M \otimes f} & M \otimes Y \\ \downarrow b & & \downarrow a \\ X & \xrightarrow{f} & Y \end{array}$$

is commutative.

## 8. COMONOIDS AND COACTIONS

**Definition 8.1.** Let  $\mathcal{C}$  be a monoidal category. A triple  $(C, \delta, \xi)$  consisting of an object  $C$  of  $\mathcal{C}$  and morphisms  $\delta : C \rightarrow C \otimes C$ ,  $\xi : C \rightarrow I$  such that

$$\begin{array}{ccc} C \otimes C \otimes C & \xleftarrow{1_C \otimes \delta} & C \otimes C \\ \uparrow \delta \otimes 1_C & & \uparrow \delta \\ C \otimes C & \xleftarrow{\delta} & C \end{array} \quad \begin{array}{ccc} C \otimes I & \xleftarrow{1_C \otimes \xi} & C \otimes C \\ & \searrow = & \uparrow \delta \\ & & C \end{array} \quad \begin{array}{ccc} C \otimes C & \xrightarrow{\xi \otimes 1_C} & I \otimes C \\ & \nwarrow = & \downarrow \delta \\ & & C \end{array}$$

is called a *comonoid* in a monoidal category  $\mathcal{C}$ . A comonoid object  $(C, \delta, \xi)$  in a symmetric monoidal category  $\mathcal{C}$  is a *cocommutative comonoid* in  $\mathcal{C}$  if the triangle

$$\begin{array}{ccc} C \otimes C & \xrightarrow{s} & C \otimes C \\ & \searrow \delta & \nearrow \delta \\ & C & \end{array}$$

is commutative, where  $s : C \otimes C \rightarrow C \otimes C$  is the symmetry of  $\mathcal{C}$ .

**Definition 8.2.** Let  $\mathcal{C}$  be a monoidal category and let  $(C_1, \delta_1, \xi_1)$ ,  $(C_2, \delta_2, \xi_2)$  be comonoids in  $\mathcal{C}$ . An arrow  $f : C_1 \rightarrow C_2$  in  $\mathcal{C}$  is a *morphism of comonoids* if the following diagrams

$$\begin{array}{ccc} C_1 \otimes C_1 & \xrightarrow{f \otimes f} & C_2 \otimes C_2 \\ \uparrow \delta_1 & & \uparrow \delta_2 \\ C_1 & \xrightarrow{f} & C_2 \end{array} \quad \begin{array}{ccc} C_1 & \xrightarrow{f} & C_2 \\ & \searrow \xi_1 & \nearrow \xi_2 \\ & I & \end{array}$$

are commutative.

**Definition 8.3.** Let  $(C, \delta, \xi)$  be a comonoid in a monoidal category  $\mathcal{C}$ . A (left) coaction of  $C$  on  $X$  in  $\mathcal{C}$  consists of a morphism  $c : X \rightarrow C \otimes X$  that makes the following diagrams

$$\begin{array}{ccc} C \otimes C \otimes X & \xleftarrow{1_C \otimes c} & C \otimes X \\ \uparrow c & & \uparrow \delta \otimes 1_X \\ C \otimes X & \xleftarrow{c} & X \end{array} \quad \begin{array}{ccc} I \otimes X & \xleftarrow{\xi \otimes 1_X} & C \otimes X \\ & \searrow = & \uparrow c \\ & & X \end{array}$$

commutative.

**Definition 8.4.** Let  $(C, \delta, \xi)$  be a comonoid in a monoidal category  $\mathcal{C}$ . Suppose that  $(X, c)$  and  $(Y, d)$  are object of  $\mathcal{C}$  equipped with coactions of  $(C, \delta, \xi)$ . Then morphism  $f : X \rightarrow Y$  is a *morphism of coactions* of  $(C, \delta, \xi)$  if the following diagram

$$\begin{array}{ccc} C \otimes X & \xrightarrow{1_C \otimes f} & C \otimes Y \\ \uparrow d & & \uparrow c \\ X & \xrightarrow{f} & Y \end{array}$$

is commutative.

## 9. BIALGEBRAS AND HOPF ALGEBRAS

**Definition 9.1.** Let  $\mathcal{C}$  be a symmetric monoidal category. Suppose that  $(B, \mu, \eta, \delta, \xi)$  is a quintuple consisting of an object  $B$  and morphisms of  $\mathcal{C}$  such that the following assertions hold.

- (1)  $(B, \mu, \eta)$  is a monoid in  $\mathcal{C}$ .
- (2)  $(B, \delta, \xi)$  is a comonoid in  $\mathcal{C}$ .
- (3) The following diagrams

$$\begin{array}{ccc} B \otimes B \otimes B \otimes B & \xrightarrow{1_B \otimes s \otimes 1_B} & B \otimes B \otimes B \otimes B \\ \uparrow \delta \otimes \delta & & \downarrow \mu \otimes \mu \\ B \otimes B & \xrightarrow{\mu} B \xrightarrow{\delta} B \otimes B & \end{array} \quad \begin{array}{ccc} I & \xrightarrow{\delta} & I \\ \eta \searrow & & \nearrow \xi \\ & B & \end{array}$$

$$\begin{array}{ccc} B \otimes B & \xrightarrow{\mu} & B \\ \downarrow \xi \otimes \xi & & \downarrow \xi \\ I \otimes I & \xrightarrow{\cong} & I \end{array} \quad \begin{array}{ccc} B & \xrightarrow{\delta} & B \otimes B \\ \uparrow \eta & & \uparrow \eta \otimes \eta \\ I & \xrightarrow{\cong} & I \otimes I \end{array}$$

are commutative, where  $s : B \otimes B \rightarrow B \otimes B$  is a symmetry.

Then we say that  $(B, \mu, \eta, \delta, \xi)$  is a *bialgebra* in a symmetric monoidal category  $\mathcal{C}$ .

**Definition 9.2.** Let  $\mathcal{C}$  be a symmetric monoidal category and let  $(B_1, \mu_1, \eta_1, \delta_1, \xi_1), (B_2, \mu_2, \eta_2, \delta_2, \xi_2)$  be bialgebras in  $\mathcal{C}$ . An arrow  $f : B_1 \rightarrow B_2$  in  $\mathcal{C}$  is a *morphism of bialgebras* if it is both a morphism of monoids and comonoids in  $\mathcal{C}$ .

## REFERENCES

- [BDR94] F. Borceux, B. Doran, and G.C. Rota. *Handbook of Categorical Algebra: Volume 2, Categories and Structures*. Cambridge Studies in Philosophy. Cambridge University Press, 1994.
- [Mac63] Saunders MacLane. Natural associativity and commutativity. *Rice Institute Pamphlet-Rice University Studies*, 49(4), 1963.
- [Mon19] Monygham. Categories of presheaves. *github repository: "Monygham/Pedo-mellon-a-minno"*, 2019.