

GEOMETRY OF k -FUNCTORS

1. INTRODUCTION

In this notes we present functorial approach to algebraic geometry. Our aim is to show that functorial and geometrical techniques are interrelated in a very efficient way.

2. k -FUNCTORS

Definition 2.1. The category $\mathbf{Fun}(\mathbf{Alg}_k, \mathbf{Set})$ of copresheaves on \mathbf{Alg}_k is called *the category of k -functors*.

If \mathfrak{X} and \mathfrak{Y} are k -functors, then we denote by $\mathrm{Mor}_k(\mathfrak{X}, \mathfrak{Y})$ the class of morphisms $\mathfrak{X} \rightarrow \mathfrak{Y}$ of k -functors.

Since the category of k -functors is a category of copresheaves, under assumptions specified in [Mon19a, section 5] for given k -functors $\mathfrak{X}, \mathfrak{Y}$ there exists an internal hom $\mathcal{M}\mathrm{or}_k(\mathfrak{X}, \mathfrak{Y})$. Let us discuss this important notion and also related ones. For details and proofs for general case we refer to [Mon19a, section 5].

Let \mathfrak{X} and \mathfrak{Y} be A -functors for some k -algebra A . Then we denote by $\mathrm{Mor}_A(\mathfrak{X}, \mathfrak{Y})$ the class of morphisms of A -functors $\mathfrak{X} \rightarrow \mathfrak{Y}$. For every A -algebra B and a morphism $\sigma : \mathfrak{X} \rightarrow \mathfrak{Y}$ of A -functors we denote by $\mathfrak{X}_B, \mathfrak{Y}_B, \sigma_B$ the restrictions $\mathfrak{X}|_{\mathbf{Alg}_B}, \mathfrak{Y}|_{\mathbf{Alg}_B}, \sigma|_{\mathbf{Alg}_B}$ of these entities to the category of B -algebras.

Fact 2.2. Let \mathfrak{X} and \mathfrak{Y} be k -functors. Assume that A is a k -algebra, B is an A -algebra, C is an B -algebra. Then the composition of maps of classes

$$\mathrm{Mor}_A(\mathfrak{X}_A, \mathfrak{Y}_A) \xrightarrow{\sigma \mapsto \sigma_B} \mathrm{Mor}_B(\mathfrak{X}_B, \mathfrak{Y}_B) \xrightarrow{\sigma \mapsto \sigma_C} \mathrm{Mor}_C(\mathfrak{X}_C, \mathfrak{Y}_C)$$

equals

$$\mathrm{Mor}_A(\mathfrak{X}_A, \mathfrak{Y}_A) \xrightarrow{\sigma \mapsto \sigma_C} \mathrm{Mor}_C(\mathfrak{X}_C, \mathfrak{Y}_C)$$

Proof. Left to the reader. □

Definition 2.3. Let \mathfrak{X} and \mathfrak{Y} be k -functors and suppose that for every k -algebra A the class $\mathrm{Mor}_A(\mathfrak{X}_A, \mathfrak{Y}_A)$ is a set. We define

$$\mathcal{M}\mathrm{or}_k(\mathfrak{X}, \mathfrak{Y})(A) = \mathrm{Mor}_A(\mathfrak{X}_A, \mathfrak{Y}_A)$$

for every k -algebra A . This is a k -functor, since for every k -algebra A and A -algebra B , we can compose a morphism $\sigma : \mathfrak{X}_A \rightarrow \mathfrak{Y}_A$ of k -functors with the forgetful functor $\mathbf{Alg}_B \rightarrow \mathbf{Alg}_A$ i.e. we have a map

$$\mathcal{M}\mathrm{or}_k(\mathfrak{X}, \mathfrak{Y})(A) \ni \sigma \mapsto \sigma_B \in \mathcal{M}\mathrm{or}_k(\mathfrak{X}, \mathfrak{Y})(B)$$

and these according to Fact 2.2 make $\mathcal{M}\mathrm{or}_k(\mathfrak{X}, \mathfrak{Y})$ a k -functor. The k -functor $\mathcal{M}\mathrm{or}_k(\mathfrak{X}, \mathfrak{Y})$ is called a *hom k -functor of \mathfrak{X} and \mathfrak{Y}* .

We define a k -functor $\mathbf{1}$ that assigns to every k -algebra a set with one element. For every k -algebra A the restriction $\mathbf{1}_A$ is a terminal object in the category of A -functors.

Fact 2.4. Let \mathfrak{X} be a k -functor. Suppose A is a k -algebra and $x \in \mathfrak{X}(A)$. Then x determines a morphism $\mathbf{1}_A \rightarrow \mathfrak{X}_A$ that for every A -algebra B with structural morphism $f : A \rightarrow B$ sends a unique element of $\mathbf{1}_A(B)$ to $\mathfrak{X}(f)(x) \in \mathfrak{X}_A(B)$. This gives rise to a bijection

$$\mathfrak{X}(A) \cong \text{Mor}_A(\mathbf{1}_A, \mathfrak{X}_A)$$

Proof. We left to the reader as an exercise. \square

Definition 2.5. Let \mathfrak{X} be a k -functor and A be a k -algebra. The set $\mathfrak{X}(A)$ is called *the set of A -points of \mathfrak{X}* .

Now let $\mathfrak{X}, \mathfrak{Y}$ be k -functors such that for every k -algebra A the class $\text{Mor}_A(\mathfrak{X}_A, \mathfrak{Y}_A)$ is a set. Suppose next that \mathfrak{U} is a k -functor and $\sigma : \mathfrak{U} \times \mathfrak{X} \rightarrow \mathfrak{Y}$ is a morphism of k -functors. Fix $x \in \mathfrak{U}(A)$. We denote by $i_x : \mathbf{1}_A \rightarrow \mathfrak{U}_A$ the morphism of A -functors corresponding to x by means of Fact 2.4. Since $\mathbf{1}_A$ is terminal A -functor, a morphism $\sigma_A \cdot (\mathbf{1}_{\mathfrak{X}_A} \times i_x)$ is isomorphic to a morphism $\tau_x : \mathfrak{X}_A \rightarrow \mathfrak{Y}_A$ of A -functors. Next $x \mapsto \tau_x$ gives rise to a morphism $\tau : \mathfrak{U} \rightarrow \text{Mor}_k(\mathfrak{X}, \mathfrak{Y})$ of k -functors and hence we have a map of classes

$$\text{Mor}_k(\mathfrak{U} \times \mathfrak{X}, \mathfrak{Y}) \ni \sigma \mapsto \tau \in \text{Mor}_k(\mathfrak{U}, \text{Mor}_k(\mathfrak{X}, \mathfrak{Y}))$$

Now we have the following result [Mon19a, Theorem 5.3].

Theorem 2.6. Let $\mathfrak{X}, \mathfrak{Y}$ be k -functors. Assume that for every k -algebra A the class $\text{Mor}_A(\mathfrak{X}_A, \mathfrak{Y}_A)$ is a set. Then the map

$$\text{Mor}_k(\mathfrak{U} \times \mathfrak{X}, \mathfrak{Y}) \rightarrow \text{Mor}_k(\mathfrak{U}, \text{Mor}_k(\mathfrak{X}, \mathfrak{Y}))$$

described above is a bijection natural in \mathfrak{U} .

3. ZARISKI LOCAL k -FUNCTORS AND ZARISKI SHEAVES

In this part we use notion of a Grothendieck topology on a category. For this notion we refer the reader to [Mon19b].

Definition 3.1. Let $\{f_i : X_i \rightarrow X\}_{i \in I}$ be a family of morphisms of k -schemes. We say that $\{f_i\}_{i \in I}$ is a *Zariski covering* of X if the following conditions are satisfied.

- (1) For every $i \in I$ morphism f_i is an open immersion of schemes.
- (2) Morphism $\coprod_{i \in I} X_i \rightarrow X$ induced by $\{f_i\}_{i \in I}$ is surjective.

The collection of all Zariski coverings on \mathbf{Sch}_k is a Grothendieck pretopology.

Definition 3.2. We call the Grothendieck topology generated by pretopology of Zariski coverings *the Zariski topology on \mathbf{Sch}_k* . A presheaf on \mathbf{Sch}_k that is a sheaf with respect to Zariski topology on \mathbf{Sch}_k is called a *Zariski sheaf*.

Let \mathfrak{X} be a presheaf on the category of k -schemes. Recall that by [Mon19b, Theorem 3.5] \mathfrak{X} is a Zariski sheaf if and only if for every k -scheme X and for every Zariski covering $\{f_i : X_i \rightarrow X\}$ of X the diagram

$$\mathfrak{X}(X) \xrightarrow{\langle \mathfrak{X}(f_i) \rangle_{i \in I}} \prod_{i \in I} \mathfrak{X}(X_i) \xrightleftharpoons[\langle \mathfrak{X}(f''_{ij}) \cdot pr_j \rangle_{(i,j)}]{\langle \mathfrak{X}(f'_{ij}) \cdot pr_i \rangle_{(i,j)}} \prod_{(i,j) \in I \times I} \mathfrak{X}(X_i \times_X X_j)$$

is a kernel of a pair of arrows, where for every $(i, j) \in I \times I$ morphisms f'_{ij} and f''_{ji} form a cartesian square

$$\begin{array}{ccc}
 X_i \times_X X_j & \xrightarrow{f''_{ij}} & X_j \\
 f'_{ij} \downarrow & & \downarrow f_j \\
 X_i & \xrightarrow{f_i} & X
 \end{array}$$

Now we repeat this definitions for k -algebras and k -functors.

Definition 3.3. Let $\{f_i : A \rightarrow A_i\}_{i \in I}$ be a family of morphisms of k -algebras. We say that $\{f_i\}_{i \in I}$ is a Zariski covering of A if the following conditions are satisfied.

- (1) For every $i \in I$ morphism $\text{Spec } f_i$ is an open immersion of schemes.
- (2) Morphism $\coprod_{i \in I} \text{Spec } A_i \rightarrow \text{Spec } A$ induced by $\{\text{Spec } f_i\}_{i \in I}$ is surjective.

The collection of all Zariski coverings on \mathbf{Alg}_k induces on its opposite category \mathbf{Aff}_k of affine k -schemes a Grothendieck pretopology.

Definition 3.4. We call the Grothendieck topology generated by pretopology of Zariski coverings the Zariski topology on \mathbf{Aff}_k . A k -functor that is a sheaf with respect to Zariski topology on \mathbf{Aff}_k is called a Zariski local k -functor.

Let \mathfrak{X} be a k -functor. Again by [Mon19b, Theorem 3.5] \mathfrak{X} is a Zariski local k -functor if and only if for every k -algebra A and for every Zariski covering $\{f_i : A \rightarrow A_i\}$ of A the diagram

$$\mathfrak{X}(A) \xrightarrow{\langle \mathfrak{X}(f_i) \rangle_{i \in I}} \prod_{i \in I} \mathfrak{X}(A_i) \xrightarrow[\langle \mathfrak{X}(f''_{ij}) \cdot pr_j \rangle_{(i,j)}]{\langle \mathfrak{X}(f'_{ij}) \cdot pr_i \rangle_{(i,j)}} \prod_{(i,j) \in I \times I} \mathfrak{X}(A_i \otimes_A A_j)$$

is a kernel of a pair of arrows, where for every $(i, j) \in I \times I$ morphisms f'_{ij} and f''_{ji} form a cocartesian square

$$\begin{array}{ccc}
 A & \xrightarrow{f_j} & A_j \\
 f_i \downarrow & & \downarrow f''_{ji} \\
 A_i & \xrightarrow{f'_{ij}} & A_i \otimes_A A_j
 \end{array}$$

Now we state the main result of this section.

Theorem 3.5. Let

$$\widehat{\mathbf{Sch}}_k \longrightarrow \text{the category of } k\text{-functors}$$

be the restriction of presheaves on \mathbf{Sch}_k to copresheaves on \mathbf{Alg}_k (k -functors) induced by the contravariant functor $\text{Spec} : \mathbf{Alg}_k \rightarrow \mathbf{Sch}_k$. Then it induces an equivalence of categories between Zariski sheaves on \mathbf{Sch}_k and Zariski local k -functors.

Proof. Note that \mathbf{Aff}_k with Zariski topology is a dense subsite ([Mon19b, definition 4.4]) of \mathbf{Sch}_k with Zariski topology. Hence the result is a special case of a more general theorem [Mon19b, Theorem 4.6]. \square

Proposition 3.6. Let $\sigma : \mathfrak{X} \rightarrow \mathfrak{Y}$ be a monomorphism of k -functors and \mathfrak{Y} be a Zariski local k -functor. Assume that for every k -algebra A and every morphism $\tau : \mathfrak{B}_{\text{Spec } A} \rightarrow \mathfrak{Y}$ of k -functors there exist a Zariski local k -functor \mathfrak{Z} that fits into a cartesian square

$$\begin{array}{ccc}
\mathfrak{Z} & \xrightarrow{\quad} & \mathfrak{X} \\
\downarrow & & \downarrow \sigma \\
\mathfrak{B}_{\text{Spec } A} & \xrightarrow{\tau} & \mathfrak{Y}
\end{array}$$

Then \mathfrak{X} is a Zariski local k -functor.

Proof. Suppose that A is a k -algebra and S is a covering sieve on A with respect to Zariski topology. Recall that by [Mon19b, page 2] we may consider S as a subcopresheaf of $\mathfrak{B}_{\text{Spec } A}$. Suppose that $\tau : \mathfrak{B}_{\text{Spec } A} \rightarrow \mathfrak{Y}$ and $m : S \rightarrow \mathfrak{X}$ are morphisms of k -functors such that $\sigma \cdot m$ is equal to the composition of $S \hookrightarrow \mathfrak{B}_{\text{Spec } A}$ with τ . Next there exists a Zariski local k -functor \mathfrak{Z} that fits into a cartesian square

$$\begin{array}{ccc}
\mathfrak{Z} & \xrightarrow{\tau'} & \mathfrak{X} \\
\downarrow \sigma' & & \downarrow \sigma \\
\mathfrak{B}_{\text{Spec } A} & \xrightarrow{\tau} & \mathfrak{Y}
\end{array}$$

of k -functors. By universal property of cartesian squares there exists a unique morphism $n : S \rightarrow \mathfrak{Z}$ of k -functors such that the diagram

$$\begin{array}{ccccc}
S & & \xrightarrow{m} & & \mathfrak{X} \\
& \searrow n & & \searrow \tau' & \\
& \mathfrak{Z} & \xrightarrow{\quad} & \mathfrak{X} & \\
& \downarrow \sigma' & & \downarrow \sigma & \\
& \mathfrak{B}_{\text{Spec } A} & \xrightarrow{\tau} & \mathfrak{Y} &
\end{array}$$

is commutative. Since \mathfrak{Z} is Zariski local, there exists a morphism $\rho : \mathfrak{B}_{\text{Spec } A} \rightarrow \mathfrak{Z}$ such that $\rho|_S = n$. Then $(\tau' \cdot \rho)|_S = \tau' \cdot n = m$ and hence matching family m admits an amalgamation. Since σ is a monomorphism, this suffices to prove that \mathfrak{X} is a Zariski local k -functor. \square

4. SCHEMES AND THEIR FUNCTORS OF POINTS

Let X be a k -scheme. We define a k -functor \mathfrak{P}_X by formula

$$\mathfrak{P}_X(A) = \text{Mor}_k(\text{Spec } A, X)$$

That is \mathfrak{P}_X is the restriction of the presheaf on \mathbf{Sch}_k represented by X to the category \mathbf{Alg}_k along the functor $\text{Spec} : \mathbf{Alg}_k \rightarrow \mathbf{Sch}_k$. Next if $f : X \rightarrow Y$ is a morphism of k -schemes, then \mathfrak{P}_f is the restriction of a morphism of presheaves on \mathbf{Sch}_k represented by f to the category of k -algebras along $\text{Spec} : \mathbf{Alg}_k \rightarrow \mathbf{Sch}_k$. Thus we have a functor

$$\mathbf{Sch}_k \xrightarrow{\mathfrak{P}} \text{the category of } k\text{-functors}$$

Fact 4.1. *Functor*

$$\mathbf{Sch}_k \xrightarrow{\mathfrak{P}} \text{the category of } k\text{-functors}$$

is full, faithful and its image consists of Zariski local k -functors. Moreover, \mathfrak{B} preserves limits.

Proof. Note that the presheaf h_X on \mathbf{Sch}_k represented by X is a Zariski sheaf. Indeed, this just rephrase standard fact that morphism of schemes can be glued in Zariski topology. Next according to Theorem 3.5 the functor $\text{Spec} : \mathbf{Alg}_k \rightarrow \mathbf{Sch}_k$ induces an equivalence between the category of Zariski sheaves and the category of local Zariski k -functors. Thus \mathfrak{P}_X is a local Zariski k -functor and \mathfrak{B} it is full and faithful. Note that Yoneda embedding $h : \mathbf{Sch}_k \rightarrow \widehat{\mathbf{Sch}}_k$ and the functor

$$\widehat{\mathbf{Sch}}_k \xrightarrow{\text{induced by Spec}} \text{the category of } k\text{-functors}$$

preserve limits. Thus their composition \mathfrak{B} also preserves limits. \square

Definition 4.2. Let X be a k -scheme. Then \mathfrak{P}_X is called *the k -functor of points of X* .

Finally note that for every k -algebra A we have an identification $\mathfrak{P}_{\text{Spec } A} = \text{Hom}_k(A, -)$ and this identification is natural with respect to A . In other words $\mathfrak{B} \cdot \text{Spec}$ is the (co)Yoneda embedding of \mathbf{Alg}_k into the category of k -functors.

Suppose now that A is a k -algebra and $\mathfrak{a} \subseteq A$ is an ideal. Then we define $V(\mathfrak{a}) = \text{Spec } A/\mathfrak{a}$ as a closed subscheme $\text{Spec } A$ induced by the quotient morphism $A \rightarrow A/\mathfrak{a}$. We define an open subscheme $D(\mathfrak{a}) = \text{Spec } A \setminus V(\mathfrak{a})$ of $\text{Spec } A$.

Definition 4.3. Let $\sigma : \mathfrak{X} \rightarrow \mathfrak{Y}$ be a morphism of k -functors. Assume that for every k -algebra A and every morphism $\tau : \mathfrak{B}_{\text{Spec } A} \rightarrow \mathfrak{Y}$ of k -functors there exist an ideal \mathfrak{a} in A and a morphism $\tau' : \mathfrak{B}_{D(\mathfrak{a})} \rightarrow \mathfrak{X}$ of k -functors such that the square

$$\begin{array}{ccc} \mathfrak{B}_{D(\mathfrak{a})} & \xrightarrow{\tau'} & \mathfrak{X} \\ \downarrow & & \downarrow \sigma \\ \mathfrak{B}_{\text{Spec } A} & \xrightarrow{\tau} & \mathfrak{Y} \end{array}$$

is cartesian. Then σ is an *open immersion of k -functors*.

Fact 4.4. The class of open immersions of k -functors is closed under base change and composition.

Proof. Left to the reader. \square

Definition 4.5. Let \mathfrak{X} be a k -functor and $\{\sigma_i : \mathfrak{X}_i \rightarrow \mathfrak{X}\}_{i \in I}$ be a family of open immersions. Then for every k -algebra A and $x \in \mathfrak{X}(A)$ we have a family of ideals $\{\mathfrak{a}_i\}_{i \in I}$ defined by cartesian squares

$$\begin{array}{ccc} \mathfrak{B}_{D(\mathfrak{a}_i)} & \xrightarrow{\tau'} & \mathfrak{X}_i \\ \downarrow & & \downarrow \sigma_i \\ \mathfrak{B}_{\text{Spec } A} & \xrightarrow{\tau} & \mathfrak{X} \end{array}$$

in which bottom vertical morphism $\tau : \mathfrak{B}_{\text{Spec } A} \rightarrow \mathfrak{X}$ corresponds to x . We say that $\{\sigma_i\}_{i \in I}$ is an *open cover of \mathfrak{X}* if for every k -scheme X and $x \in \mathfrak{X}(X)$ we have

$$\text{Spec } A = \bigcup_{i \in I} D(\mathfrak{a}_i)$$

or in other words $A = \sum_{i \in I} \mathfrak{a}_i$.

Theorem 4.6. *Let \mathfrak{X} be a k -functor. Then the following are equivalent.*

- (i) \mathfrak{X} is isomorphic with functor of points of some k -scheme.
- (ii) \mathfrak{X} is a Zariski local k -functor and there exists an open cover $\{\sigma_i : \mathfrak{B}_{X_i} \rightarrow \mathfrak{X}\}_{i \in I}$ such that $\{X_i\}_{i \in I}$ are k -schemes.
- (iii) \mathfrak{X} is a Zariski local k -functor and there exists an open cover $\{\sigma_i : \mathfrak{B}_{\text{Spec } A_i} \rightarrow \mathfrak{X}\}_{i \in I}$ such that $\{A_i\}_{i \in I}$ are k -algebras.

The proof depends on few lemmas.

Lemma 4.6.1. *Let $f : X \rightarrow Y$ be a morphism of k -schemes. Suppose that f is surjective morphism and an open immersion locally on X . Then \mathfrak{B}_f is a locally surjective morphism of Zariski local k -functors.*

Proof of the lemma. Let A be a k -algebra and $g : \text{Spec } A \rightarrow Y$ be an open immersion of schemes. Since f is surjective and an open immersion locally on X , there exist a Zariski cover $\{f_i : A \rightarrow A_i\}_{i \in I}$ and a family $\{g_i : \text{Spec } A_i \rightarrow X\}_{i \in I}$ of open immersions such that $f \cdot g_i = g \cdot \text{Spec } f_i$ for every $i \in I$. This implies that $\mathfrak{B}_f(g_i) = \mathfrak{B}_Y(f_i)(g)$ for every $i \in I$. Thus \mathfrak{B}_f is a locally surjective morphism of Zariski local k -functors. \square

The next lemma is a categorical formulation of the *recollement* technique [GD71, Chapitre 0, 4.1.7] and as such it is more convenient in our settings.

Lemma 4.6.2. *Let $X = \coprod_{i \in I} X_i, R = \coprod_{i,j \in I} R_{ij}$ be disjoint sums of k -schemes and let $p, q : R \rightarrow X$ be morphisms of k -schemes such that*

- (1) *For any $i, j \in I$ morphism $p|_{R_{ij}}$ induces an open immersion $R_{ij} \hookrightarrow X_i$ and morphism $q|_{R_{ij}}$ induces an open immersion $R_{ij} \hookrightarrow X_j$.*
- (2) *For every $i \in I$ morphisms $p|_{R_{ii}}$ and $q|_{R_{ii}}$ are equal and induce an isomorphisms $R_{ii} \rightarrow X_i$.*
- (3) *Triple (R, p, q) is an equivalence relation on X in the category of k -schemes.*

Then there exist a k -scheme Y and a morphism $f : X \rightarrow Y$ of k -schemes such that

$$\mathfrak{B}_R \xrightarrow[\mathfrak{B}_q]{\mathfrak{B}_p} \mathfrak{B}_X \xrightarrow{\mathfrak{B}_f} \mathfrak{B}_Y$$

is a cokernel of a pair $(\mathfrak{B}_p, \mathfrak{B}_q)$ in the category of Zariski local k -functors.

Proof of the lemma. We define first Y and f in the category of topological spaces and continuous maps. For this let Y be the quotient space of X by R and let f be the quotient topological map. Since topologically we have $f \cdot p = f \cdot q$, we deduce that $f_* p_* \mathcal{O}_R = f_* q_* \mathcal{O}_R$ and in addition we have a pair

$$f_* \mathcal{O}_X \xrightarrow[f_* q^\#]{f_* p^\#} f_* \cdot p_* \mathcal{O}_R = f_* \cdot q_* \mathcal{O}_R$$

of morphisms of sheaves of k -algebras on Y . We define a sheaf \mathcal{O}_Y of k -algebras to be a kernel of a pair $(f_* p^\#, f_* q^\#)$. This sheaf comes with the canonical morphism $f^\# : \mathcal{O}_Y \rightarrow f_* \mathcal{O}_X$ of sheaves of k -algebras. This makes Y into a ringed topological space and f into a morphism of ringed spaces such that

$$f \cdot p = f \cdot q$$

Now we show that Y is a k -scheme and f is a morphism of k -schemes. For every pair $i, j \in I$ we denote $p(R_{ij}) = X_{ij}$. This is an open subscheme of X_i and moreover, $X_{ii} = X_i$ for every $i \in I$. Next for every $i, j \in I$ we have a commutative triangle

$$\begin{array}{ccc} R_{ij} & \xrightarrow{s_{ij}} & R_{ji} \\ p_{ij} \searrow & & \swarrow q_{ij} \\ & X_{ij} & \end{array}$$

where s_{ij} is induced by symmetry on $X \times X$ and isomorphisms p_{ij} and q_{ij} are induced by p and q , respectively. In particular, $q_{ii} = p_{ii}$ for every $i \in I$. Since both p, q are open maps, we derive that f is an open map. Hence for every $i \in I$ a bijection $X_i \rightarrow f(X_i)$ induced by f is a topological isomorphism. Fix now $i \in I$ and denote $f(X_i)$ by Y_i . Note that this is an open subset of Y . Morphism $(f_* p^\#)_{Y_i} : \mathcal{O}_{X_i} \times \prod_{j \neq i} \mathcal{O}_{X_{ji}} \rightarrow \prod_{j \in I} \mathcal{O}_{R_{ij}} \times \prod_{j \neq i} \mathcal{O}_{R_{ji}}$ is given by

$$(a, (b_{ji})_{j \neq i}) \mapsto \left((p_{ij}^\#(a|_{X_{ij}}))_{j \in I}, (p_{ji}^\#(b_{ji}))_{j \neq i} \right)$$

and morphism $(f_* q^\#)_{Y_i} : \mathcal{O}_{X_i} \times \prod_{j \neq i} \mathcal{O}_{X_{ji}} \rightarrow \prod_{j \in I} \mathcal{O}_{R_{ij}} \times \prod_{j \neq i} \mathcal{O}_{R_{ji}}$ is given by

$$(a, (b_{ji})_{j \neq i}) \mapsto \left((q_{ij}^\#(a|_{X_{ij}}))_{j \in I}, (q_{ji}^\#(b_{ji}))_{j \neq i} \right)$$

Thus $(a, (b_{ji})_{j \neq i})$ lies in the kernel of $((f_* p^\#)_{Y_i}, (f_* q^\#)_{Y_i})$ if and only if

$$p_{ij}^\#(a|_{X_{ij}}) = q_{ji}^\#(b_{ji}), \quad q_{ij}^\#(a|_{X_{ij}}) = p_{ji}^\#(b_{ji})$$

for every $j \neq i$. This is equivalent with the system of equations

$$b_{ji} = \left((q_{ji}^\#)^{-1} \cdot p_{ij}^\# \right) (a|_{X_{ij}}), \quad b_{ji} = \left((p_{ji}^\#)^{-1} \cdot q_{ij}^\# \right) (a|_{X_{ij}})$$

for every $j \neq i$. Since

$$p_{ji}^\# \cdot (q_{ji}^\#)^{-1} = s_{ji}^\# = s_{ij}^\# = p_{ij}^\# \cdot (q_{ij}^\#)^{-1}$$

for every $i, j \in I$, we derive that the system of equations has a unique solution for every section of \mathcal{O}_{X_i} . This implies that $f_{Y_i}^\# : \mathcal{O}_{Y_i} \rightarrow \mathcal{O}_{f^{-1}(Y_i)}$ composed with the restriction $\mathcal{O}_{f^{-1}(Y_i)} \rightarrow \mathcal{O}_{X_i}$ is an isomorphism. Hence f induces an isomorphism of ringed spaces $X_i \cong Y_i$. Thus for every $i \in I$ ringed space Y_i is a k -scheme and f induces isomorphism of k -schemes $X_i \cong Y_i$. Note that

$$Y = f(X) = \bigcup_{i \in I} f(X_i) = \bigcup_{i \in I} Y_i$$

Therefore, Y is a k -scheme and $f : X \rightarrow Y$ is a morphism of k -schemes.

Now we verify that \mathfrak{B}_f is the quotient in the category of Zariski local k -functors. For this note that we proved above that f is open immersion of k -schemes locally on X and it is surjective. Thus by Lemma 4.6.1 we derive that \mathfrak{B}_f is a locally surjective morphism of Zariski local k -functors. Therefore ([Mon19b, Theorem 7.3]), it suffices to show that the square

$$\begin{array}{ccc} \mathfrak{B}_R & \xrightarrow{\mathfrak{B}_q} & \mathfrak{B}_X \\ \mathfrak{B}_p \downarrow & & \downarrow \mathfrak{B}_f \\ \mathfrak{B}_X & \xrightarrow{\mathfrak{B}_f} & \mathfrak{B}_Y \end{array}$$

is cartesian. Since \mathfrak{B} preserves limits (Fact 4.1), we derive that it suffices to check that

$$\begin{array}{ccc}
R & \xrightarrow{q} & X \\
p \downarrow & & \downarrow f \\
X & \xrightarrow{f} & Y
\end{array}$$

is cartesian square of k -schemes. This is clear since $R_{ij} = X_i \times_Y X_j$ for every $i, j \in I$ and hence

$$X \times_Y X = \left(\coprod_{i \in I} X_i \right) \times_Y \left(\coprod_{i \in I} X_i \right) = \coprod_{i, j \in I} (X_i \times_Y X_j) = \coprod_{i, j \in I} R_{ij} = R$$

Thus the result follows. \square

Proof of the theorem. If (i) holds, then we may assume that $\mathfrak{X} = \mathfrak{B}_Y$ for some k -scheme Y . Fact 4.1 states that \mathfrak{B}_Y is a Zariski local k -functor and clearly $1_{\mathfrak{B}_Y} : \mathfrak{B}_Y \rightarrow \mathfrak{B}_Y$ is an open cover. Thus (i) \Rightarrow (ii).

Every functor of points of a k -scheme admits open cover by functors of points of affine k -schemes. Indeed, it suffices to take open affine subschemes that cover given k -scheme and apply \mathfrak{B} . This implies that every open cover of a k -functor \mathfrak{X} by functors of points of k -schemes admits refinement by open cover of functors of points of affine k -schemes. Therefore, implication (ii) \Rightarrow (iii) holds.

Suppose that a k -functor \mathfrak{X} is Zariski local and $\{\sigma_i : \mathfrak{B}_{\text{Spec } A_i} \rightarrow \mathfrak{X}\}_{i \in I}$ is an open cover of \mathfrak{X} . Note that for every $i, j \in I$ there exist a k -scheme R_{ij} and open immersions $p_{ij} : R_{ij} \hookrightarrow \text{Spec } A_i$, $q_{ij} : R_{ij} \hookrightarrow \text{Spec } A_j$ such that the square

$$\begin{array}{ccc}
\mathfrak{B}_{R_{ij}} & \xrightarrow{\mathfrak{B}_{q_{ij}}} & \mathfrak{B}_{\text{Spec } A_j} \\
\mathfrak{B}_{p_{ij}} \downarrow & & \downarrow \sigma_j \\
\mathfrak{B}_{\text{Spec } A_i} & \xrightarrow{\sigma_i} & \mathfrak{X}
\end{array}$$

is cartesian. Consider k -scheme $X = \coprod_{i \in I} \text{Spec } A_i$ and morphism $\sigma : \mathfrak{B}_X \rightarrow \mathfrak{X}$ induced by $\{\sigma_i\}_{i \in I}$. Moreover, consider k -scheme $R = \coprod_{i, j \in I} R_{ij}$ and morphisms $p, q : R \rightarrow X$ induced by $\{p_{ij}\}_{i, j \in I}$ and $\{q_{ij}\}_{i, j \in I}$, respectively. Note that the square

$$\begin{array}{ccc}
\mathfrak{B}_R & \xrightarrow{\mathfrak{B}_q} & \mathfrak{B}_X \\
\mathfrak{B}_p \downarrow & & \downarrow \sigma \\
\mathfrak{B}_X & \xrightarrow{\sigma} & \mathfrak{X}
\end{array}$$

is cartesian and hence $(\mathfrak{B}_R, \mathfrak{B}_p, \mathfrak{B}_q)$ is an equivalence relation. By Lemma 4.6.2 there exist a k -scheme Y and a morphism $f : X \rightarrow Y$ such that

$$\mathfrak{B}_R \xrightarrow[\mathfrak{B}_q]{\mathfrak{B}_p} \mathfrak{B}_X \xrightarrow{\mathfrak{B}_f} \mathfrak{B}_Y$$

is a cokernel of $(\mathfrak{B}_p, \mathfrak{B}_q)$. Moreover, σ is locally surjective morphism of Zariski local k -functors and hence also

$$\mathfrak{B}_R \begin{array}{c} \xrightarrow{\mathfrak{B}_p} \\ \xrightarrow{\mathfrak{B}_q} \end{array} \mathfrak{B}_X \xrightarrow{\sigma} \mathfrak{X}$$

is a cokernel of $(\mathfrak{B}_p, \mathfrak{B}_q)$. Thus \mathfrak{B}_Y is isomorphic with \mathfrak{X} . This proves (iii) \Rightarrow (i). \square

5. REPRESENTABLE MORPHISMS OF k -FUNCTORS

Definition 5.1. Let $\sigma : \mathfrak{X} \rightarrow \mathfrak{Y}$ be a morphism of k -functors. Assume that for every k -algebra A and every morphism $\tau : \mathfrak{B}_{\text{Spec } A} \rightarrow \mathfrak{Y}$ of k -functors there exist a k -scheme X , a morphism $f : X \rightarrow \text{Spec } A$ and a morphism $\tau' : \mathfrak{B}_X \rightarrow \mathfrak{X}$ of k -functors such that the square

$$\begin{array}{ccc} \mathfrak{B}_X & \xrightarrow{\tau'} & \mathfrak{X} \\ \mathfrak{B}_f \downarrow & & \downarrow \sigma \\ \mathfrak{B}_{\text{Spec } A} & \xrightarrow{\tau} & \mathfrak{Y} \end{array}$$

is cartesian. Then σ is a *representable morphism of k -functors*.

Fact 5.2. The class of representable morphisms of k -functors is closed under base change and composition.

Proof. Left to the reader. \square

Proposition 5.3. Let $\sigma : \mathfrak{X} \rightarrow \mathfrak{Y}$ be a representable morphism of Zariski local k -functors. Fix a k -scheme Y and a morphism $\tau : \mathfrak{B}_Y \rightarrow \mathfrak{Y}$. Then there exist a k -scheme X , a morphism $f : X \rightarrow Y$ and a morphism $\tau' : \mathfrak{B}_X \rightarrow \mathfrak{X}$ such that the square

$$\begin{array}{ccc} \mathfrak{B}_X & \xrightarrow{\tau'} & \mathfrak{X} \\ \mathfrak{B}_f \downarrow & & \downarrow \sigma \\ \mathfrak{B}_Y & \xrightarrow{\tau} & \mathfrak{Y} \end{array}$$

is cartesian.

Proof. Let

$$\begin{array}{ccc} \mathfrak{Z} & \xrightarrow{\tau'} & \mathfrak{X} \\ \sigma' \downarrow & & \downarrow \sigma \\ \mathfrak{B}_Y & \xrightarrow{\tau} & \mathfrak{Y} \end{array}$$

be a cartesian square. According to [Mon19b, Theorem 2.12] k -functor \mathfrak{Z} is Zariski local. Suppose that $\{f_i : \text{Spec } A_i \rightarrow Y\}_{i \in I}$ is an open cover of Y . Then $\{\mathfrak{B}_{f_i} : \mathfrak{B}_{\text{Spec } A_i} \rightarrow \mathfrak{B}_Y\}_{i \in I}$ is an open cover of \mathfrak{B}_Y and hence its base change $\{\tau_i : \mathfrak{Z}_i \rightarrow \mathfrak{Z}\}_{i \in I}$ is an open cover of \mathfrak{Z} . Since σ is representable, we deduce that \mathfrak{Z}_i is a functor of points of some k -scheme for $i \in I$. Now by Theorem 4.6 we derive that there exists a k -scheme X such that \mathfrak{Z} is isomorphic with \mathfrak{B}_X . This proves the result. \square

Definition 5.4. Let $\sigma : \mathfrak{X} \rightarrow \mathfrak{Y}$ be a morphism of k -functors. Assume that for every k -algebra A and every morphism $\tau : \mathfrak{B}_{\text{Spec } A} \rightarrow \mathfrak{Y}$ of k -functors there exist an ideal \mathfrak{a} in A and morphism $\tau' : \mathfrak{B}_{V(\mathfrak{a})} \rightarrow \mathfrak{X}$ such that the square

$$\begin{array}{ccc}
\mathfrak{B}_{V(\mathfrak{a})} = \mathfrak{B}_{\mathrm{Spec} A/\mathfrak{a}} & \xrightarrow{\tau'} & \mathfrak{X} \\
\mathfrak{B}_{\mathrm{Spec} q} \downarrow & & \downarrow \sigma \\
\mathfrak{B}_{\mathrm{Spec} A} & \xrightarrow{\tau} & \mathfrak{Y}
\end{array}$$

is cartesian, where $q : A \rightarrow A/\mathfrak{a}$ is the quotient map. Then σ is a closed immersion of k -functors.

Fact 5.5. *The class of closed immersions of k -functors is closed under base change and composition.*

Proof. Left to the reader. □

Proposition 5.6. *Let $\sigma : \mathfrak{X} \rightarrow \mathfrak{Y}$ be a closed (open) immersion of k -functors. Fix a k -scheme Y and a morphism $\tau : \mathfrak{B}_Y \rightarrow \mathfrak{Y}$. Then there exist a k -scheme X , a closed (open) immersion $f : X \rightarrow Y$ of schemes and a morphism $\tau' : \mathfrak{B}_X \rightarrow \mathfrak{X}$ of k -functors such that the square*

$$\begin{array}{ccc}
\mathfrak{B}_X & \xrightarrow{\tau'} & \mathfrak{X} \\
\mathfrak{B}_f \downarrow & & \downarrow \sigma \\
\mathfrak{B}_Y & \xrightarrow{\tau} & \mathfrak{Y}
\end{array}$$

is cartesian.

Proof. According to Fact 5.5 (Fact 4.4) pullback $\mathfrak{X} \times_{\mathfrak{Y}} \mathfrak{B}_Y \rightarrow \mathfrak{B}_Y$ of σ along τ is a closed (open) immersion of k -functors. Since \mathfrak{B}_Y is a Zariski local k -functor by Fact 4.1 and closed (open) immersions are monomorphisms, we derive by Proposition 3.6 that a fiber-product $\mathfrak{X} \times_{\mathfrak{Y}} \mathfrak{B}_Y$ of σ and τ is a Zariski local k -functor. Since closed (open) immersions of k -functors are representable, we deduce by Proposition 5.3 that there exists a k -scheme X , a morphism $f : X \rightarrow Y$ of k -schemes and a morphism $\tau' : \mathfrak{B}_X \rightarrow \mathfrak{X}$ of k -functors such that the square

$$\begin{array}{ccc}
\mathfrak{X} \times_{\mathfrak{Y}} \mathfrak{B}_Y \cong \mathfrak{B}_X & \xrightarrow{\tau'} & \mathfrak{X} \\
\mathfrak{B}_f \downarrow & & \downarrow \sigma \\
\mathfrak{B}_Y & \xrightarrow{\tau} & \mathfrak{Y}
\end{array}$$

is cartesian and \mathfrak{B}_f is a closed (open) immersion of k -functors. Since the functor

$$\widehat{\mathrm{Sch}}_k \xrightarrow{\mathfrak{B}} \text{the category of } k\text{-functors}$$

preserves finite limits, it follows that for every open affine subset V of Y we have a cartesian square

$$\begin{array}{ccc}
\mathfrak{B}_{f^{-1}(V)} & \hookrightarrow & \mathfrak{B}_X \\
\mathfrak{B}_{f_V} \downarrow & & \downarrow \mathfrak{B}_f \\
\mathfrak{B}_V & \hookrightarrow & \mathfrak{B}_Y
\end{array}$$

where $f_V : f^{-1}(V) \rightarrow V$ is the restriction of f . Next as \mathfrak{B}_f is a closed (open) immersion and V is affine, we derive that f_V is a closed (open) immersion of schemes. Since this holds for every affine open subset V of Y , we deduce that f is a closed (open) immersion. \square

The next result is frequently used in the theory of *algebraic spaces*.

Proposition 5.7. *Let \mathfrak{Y} be a k -functor such that the diagonal $\mathfrak{Y} \rightarrow \mathfrak{Y} \times \mathfrak{Y}$ is representable. Then every morphism $\sigma : \mathfrak{X} \rightarrow \mathfrak{Y}$ of k -functors is representable.*

Proof. Fix a morphism of k -functors $\sigma : \mathfrak{X} \rightarrow \mathfrak{Y}$. Let Y be a k -scheme and let $\tau : \mathfrak{B}_Y \rightarrow \mathfrak{Y}$ be a morphism of k -functors. Consider the cartesian square

$$\begin{array}{ccc} \mathfrak{Z} & \xrightarrow{\tau'} & \mathfrak{X} \\ \sigma' \downarrow & & \downarrow \sigma \\ \mathfrak{B}_Y & \xrightarrow{\tau} & \mathfrak{Y} \end{array}$$

Then there exists a cartesian square

$$\begin{array}{ccc} \mathfrak{Z} & \xrightarrow{\quad} & \mathfrak{Y} \\ \downarrow & & \downarrow \text{diagonal} \\ \mathfrak{B}_Y \times \mathfrak{Y} & \xrightarrow{\tau \times \sigma} & \mathfrak{Y} \times \mathfrak{Y} \end{array}$$

Since the diagonal of \mathfrak{Y} is representable, we derive that \mathfrak{Z} is isomorphic with functor of points of some k -scheme. This finishes the proof. \square

6. TRANSPORTERS

Definition 6.1. Let X be a k -scheme. Suppose that there exists an open affine cover $X = \bigcup_{i \in I} X_i$ such that k -algebra $\Gamma(X_i, \mathcal{O}_{X_i})$ is free as a k -module. Then we say that X is a *locally free k -scheme*.

Next theorem is the main result of this section.

Theorem 6.2. *Let $j : \mathfrak{Y}' \rightarrow \mathfrak{Y}$ be a closed immersion of k -functors and X be a locally free k -scheme. Suppose that classes $\text{Mor}_A((\mathfrak{B}_X)_A, \mathfrak{Y}_A)$ are sets for every k -algebra A . Then classes $\text{Mor}_A((\mathfrak{B}_X)_A, \mathfrak{Y}'_A)$ are sets for every k -algebra A and the morphism*

$$\text{Mor}_k(1_{\mathfrak{B}_X}, j) : \text{Mor}_k(\mathfrak{B}_X, \mathfrak{Y}') \rightarrow \text{Mor}_k(\mathfrak{B}_X, \mathfrak{Y})$$

is a closed immersion of k -functors.

It is useful to isolate crucial steps in the argument. For this we proceed by proving some lemmas.

Lemma 6.2.1. *Suppose that A is a commutative ring. Let $j : \mathfrak{Y}' \rightarrow \mathfrak{Y}$ be a closed immersion of A -functors and X be an affine A -scheme such that $\Gamma(X, \mathcal{O}_X)$ is a free A -module. Assume that $\tau : \mathfrak{B}_X \rightarrow \mathfrak{Y}$ is a morphism of A -functors. Then there exists an ideal $\mathfrak{a} \subseteq A$ such that for every A -algebra B the restriction τ_B factors through j_B if and only if the structure morphism $f : A \rightarrow B$ of B satisfies $\mathfrak{a} \subseteq \ker(f)$.*

Proof of the lemma. Since j is a closed immersion of A -functors and X is affine k -scheme there exists an affine A -scheme X' , a closed immersion $j' : X' \rightarrow X$ of schemes and a cartesian square

$$\begin{array}{ccc}
\mathfrak{B}_{X'} & \longrightarrow & \mathfrak{Y}' \\
\mathfrak{B}_{j'} \downarrow & & \downarrow j \\
\mathfrak{B}_X & \xrightarrow{\tau} & \mathfrak{Y}
\end{array}$$

of A -functors. Next let B be an A -algebra with the structure morphism $f : A \rightarrow B$. Then τ_B factors through j_B if and only if the projection $\mathrm{Spec} B \times_{\mathrm{Spec} A} X \rightarrow X$ induced by f factors through X' . Let $A[X]$ be the A -algebra of global regular functions on X and let \mathfrak{J} be an ideal in $A[X]$ such that $A[X]/\mathfrak{J} = A[X']$ is the A -algebra of global regular functions of X' . With this notation we derive that the projection $\mathrm{Spec} B \times_{\mathrm{Spec} A} X \rightarrow X$ induced by f factors through X' if and only if the morphism $A[X] \rightarrow B \otimes_A A[X]$ induced by f sends every element of \mathfrak{J} to zero. Since $A[X]$ is a free A -module, we write $A[X] = A^{\oplus I}$ for some index set I . Then the morphism $A[X] \rightarrow B \otimes_A A[X]$ induced by f is just $f^{\oplus I} : A^{\oplus I} \rightarrow B^{\oplus I}$. We have $f^{\oplus I}(\mathfrak{J}) = 0$ if and only if $(pr_i^B \cdot f^{\oplus I})(\mathfrak{J}) = 0$ for every $i \in I$, where $pr_i^B : B^{\oplus I} \rightarrow B$ is the projection on i -th component. Pick $i \in I$ and consider the commutative diagram

$$\begin{array}{ccc}
A^{\oplus I} & \xrightarrow{f^{\oplus I}} & B^{\oplus I} \\
pr_i^A \downarrow & & \downarrow pr_i^B \\
A & \xrightarrow{f} & B
\end{array}$$

In the diagram pr_i^A is the projection on i -th component. Diagram implies that $(pr_i^B \cdot f^{\oplus I})(\mathfrak{J}) = 0$ for every $i \in I$ if and only if $(f \cdot pr_i^A)(\mathfrak{J}) = 0$ for every $i \in I$. This is equivalent with the condition that $f(a) = 0$ for ideal a in A generated by $\sum_{i \in I} pr_i^A(\mathfrak{J})$. Thus the lemma is proved. \square

Lemma 6.2.2. *Suppose that A is a commutative ring. Let $j : \mathfrak{Y}' \rightarrow \mathfrak{Y}$ be a closed immersion of A -functors and X be an A -scheme with open cover*

$$X = \bigcup_{i \in I} X_i$$

Assume that $\tau : \mathfrak{B}_X \rightarrow \mathfrak{Y}$ is a morphism of A -functors. Fix an A -algebra B . Then τ_B factors through j_B if and only if $(\tau|_{\mathfrak{B}_{X_i}})_B$ factors through j_B for every $i \in I$.

Proof of the lemma. If τ_B factors through j_B , then also $(\tau|_{\mathfrak{B}_{X_i}})_B$ factors through j_B for every $i \in I$. It suffices to prove the converse. So suppose that $(\tau|_{\mathfrak{B}_{X_i}})_B$ factors through j_B for every $i \in I$. Since j is a closed immersion of A -functors and X is an A -scheme, Proposition 5.6 implies that there exists a cartesian square

$$\begin{array}{ccc}
\mathfrak{B}_{X'} & \longrightarrow & \mathfrak{Y}' \\
\mathfrak{B}_{j'} \downarrow & & \downarrow j \\
\mathfrak{B}_X & \xrightarrow{\tau} & \mathfrak{Y}
\end{array}$$

where $j' : X' \rightarrow X$ is a closed immersion of A -schemes. For each $i \in I$ let $j'_i : j'^{-1}(X_i) \rightarrow X_i$ be the restriction of j' . We have the induced cartesian square

$$\begin{array}{ccc} \mathfrak{B}^{j'^{-1}}(X_i) & \longrightarrow & \mathfrak{Y}' \\ \mathfrak{B}_{j'_i} \downarrow & & \downarrow j \\ \mathfrak{B}_{X_i} & \xrightarrow{\tau_{\mathfrak{B}_{X_i}}} & \mathfrak{Y} \end{array}$$

Now $(\tau_{\mathfrak{B}_{X_i}})_B$ factors through j_B . This implies that $(\mathfrak{B}_{j'_i})_B$ admits a section for every $i \in I$. Then $(\mathfrak{B}_{j'_i})_B$ is an isomorphism for every $i \in I$. Thus $j'_i \times_{\text{Spec } A} 1_{\text{Spec } B}$ is an isomorphism for every $i \in I$ and hence $j' \times_{\text{Spec } A} 1_{\text{Spec } B}$ is an isomorphism of B -schemes. This means that τ_B factors through j_B . \square

Proof of the theorem. Let A be a k -algebra. The restriction functor $(-)|_{\mathbf{Alg}_A} = (-)_A$ preserves all closed immersions. Thus j_A is a closed immersion of A -functors and hence we derive that $j_A : \mathfrak{Y}'_A \rightarrow \mathfrak{Y}_A$ is a monomorphism of A -functors. Thus we have an injective map of classes

$$\text{Mor}_A(1_{(\mathfrak{B}_X)_A}, j_A) : \text{Mor}_A((\mathfrak{B}_X)_A, \mathfrak{Y}'_A) \hookrightarrow \text{Mor}_A((\mathfrak{B}_X)_A, \mathfrak{Y}_A)$$

Hence if $\text{Mor}_A((\mathfrak{B}_X)_A, \mathfrak{Y}_A)$ is a set, then $\text{Mor}_A((\mathfrak{B}_X)_A, \mathfrak{Y}'_A)$ is a set. All these facts imply that both internal homs

$$\text{Mor}_k(\mathfrak{B}_X, \mathfrak{Y}'), \text{Mor}_k(\mathfrak{B}_X, \mathfrak{Y})$$

exist and morphism $\text{Mor}_k(1_{\mathfrak{B}_X}, j)$ of k -functors is a monomorphism. Our task is to prove that it is a closed immersion. For this consider a k -algebra A and a morphism $\sigma : \mathfrak{B}_{\text{Spec } A} \rightarrow \text{Mor}_k(\mathfrak{B}_X, \mathfrak{Y})$ of k -functors that sends 1_A to some morphism $\tau : (\mathfrak{B}_X)_A \rightarrow \mathfrak{Y}_A$ of A -functors. Consider a cartesian square

$$\begin{array}{ccc} \mathfrak{U} & \longrightarrow & \text{Mor}_k(\mathfrak{B}_X, \mathfrak{Y}') \\ \downarrow & & \downarrow \text{Mor}_k(1_{\mathfrak{B}_X}, j) \\ \mathfrak{B}_{\text{Spec } A} & \xrightarrow{\sigma} & \text{Mor}_k(\mathfrak{B}_X, \mathfrak{Y}) \end{array}$$

Since $\text{Mor}_k(1_{\mathfrak{B}_X}, j)$ is a monomorphism, we may consider \mathfrak{U} as a k -subfunctor of $\mathfrak{B}_{\text{Spec } A}$. For every k -algebra B subset $\mathfrak{U}(B) \subseteq \text{Mor}_k(A, B) = \text{Mor}_k(\text{Spec } B, \text{Spec } A)$ consists of A -algebras B with structure morphisms $f : A \rightarrow B$ such that τ_B factors through $j_B : \mathfrak{Y}'_B \rightarrow \mathfrak{Y}_B$. Since X is a locally free k -scheme, we deduce that $(\mathfrak{B}_X)_A$ is a functor of points of a locally free A -scheme

$$\text{Spec } A \times_{\text{Spec } k} X$$

Pick an open affine cover $\bigcup_{i \in I} X_i$ of this A -scheme such that $\Gamma(X_i, \mathcal{O}_X)$ is a free A -module. Now Lemma 6.2.2 implies that τ_B factors through j_B if and only if $(\tau_{|X_i})_B$ factors through j_B for every $i \in I$. Next by Lemma 6.2.1 we deduce that $(\tau_{|X_i})_B$ factors through j_B for given $i \in I$ if and only if $f(\mathfrak{a}_i) = 0$ for some ideal $\mathfrak{a}_i \subseteq A$ independent of f . Thus \mathfrak{U} consists of all morphisms $f : A \rightarrow B$ of k -algebras such that $f(\mathfrak{a}) = 0$ where $\mathfrak{a} = \sum_{i \in I} \mathfrak{a}_i$. Therefore, $\mathfrak{U} \hookrightarrow \mathfrak{B}_{\text{Spec } A}$ is isomorphic with $\mathfrak{B}_{V(\mathfrak{a})} = \mathfrak{B}_{\text{Spec } A/\mathfrak{a}} \hookrightarrow \mathfrak{B}_{\text{Spec } A}$ induced by the quotient map $A \rightarrow A/\mathfrak{a}$ and hence $\text{Mor}_k(1_{\mathfrak{B}_X}, j)$ is a closed immersion of k -functors. \square

The Theorem 6.2 is a simple yet powerful result. Before giving any interesting applications we state its immediate consequence.

REFERENCES

- [GD71] Alexander Grothendieck and Jean Dieudonné. *Éléments de géométrie algébrique I*, volume 166 of *Grundlehren der Mathematischen Wissenschaften*. Springer-Verlag, 1971.
- [Mon19a] Monygham. Categories of presheaves. *github repository: "Monygham/Pedo-mellon-a-minno"*, 2019.
- [Mon19b] Monygham. Categories of sheaves. *github repository: "Monygham/Pedo-mellon-a-minno"*, 2019.