

BIAŁYNICKI-BIRULA FUNCTORS

1. INTRODUCTION

In this notes we study Białynicki-Birula functors. In the first section we prove some results concerning the forgetful functor $\mathbf{Rep}(\mathbf{M}) \rightarrow \mathbf{Rep}(\mathbf{G})$, where \mathbf{M} is an affine monoid k -scheme and \mathbf{G} is its group of units (we assume that \mathbf{G} is open and schematically dense in \mathbf{M}). These results will be used in the following sections.

We assume that k is a field. In these notes we use the following notational convention.

Remark 1.1. Since the Yoneda embedding $\mathbf{Sch}_k \hookrightarrow \widehat{\mathbf{Sch}_k}$ is full and faithful, we identify \mathbf{Sch}_k with the subcategory of $\widehat{\mathbf{Sch}_k}$ consisting of representable presheaves on \mathbf{Sch}_k . In particular, if X is a k -scheme, then we denote by the same symbol the presheaf representable by X .

2. TANNAKIAN FORMALISM FOR QUOTIENT STACKS

In this section we discuss an application of the main result of [Hall and Rydh, 2019]. For this we need to briefly discuss *algebraic stacks*, although for our purposes there is no need to use any technical details of this language. We refer the interested reader to the excellent exposition [Olsson, 2016] of this subject. We note the following facts.

- (1) An *algebraic stack* is a category fibered over \mathbf{Sch}_k satisfying certain extra conditions described in [Olsson, 2016, Definition 4.6.1] and [Olsson, 2016, Definition 8.1.4]. By [Olsson, 2016, Definition 8.2.1, Example 8.2.3] there are well defined notions of *locally noetherian*, *noetherian* and *excellent algebraic stacks*.
- (2) A *morphism of algebraic stacks* is a morphism of fibered categories over \mathbf{Sch}_k . If \mathcal{X} and \mathcal{Y} are algebraic stack, then we denote by $\mathrm{Mor}(\mathcal{X}, \mathcal{Y})$ the corresponding category of morphisms.
- (3) For every locally noetherian algebraic stack \mathcal{X} there exists an abelian monoidal category $\mathcal{Coh}(\mathcal{X})$ of coherent sheaves on \mathcal{X} ([Olsson, 2016, Definition 9.1.14]). If \mathcal{X} and \mathcal{Y} are locally noetherian algebraic stacks, then we denote by $\mathrm{Hom}_{r, \otimes, \cong}(\mathcal{Coh}(\mathcal{X}), \mathcal{Coh}(\mathcal{Y}))$ the category of right exact, monoidal functors $\mathcal{Coh}(\mathcal{X}) \rightarrow \mathcal{Coh}(\mathcal{Y})$ with natural isomorphism as morphisms.
- (4) If $f : \mathcal{X} \rightarrow \mathcal{Y}$ is a morphism of locally noetherian algebraic stacks, then f induces the functor $f^* : \mathcal{Coh}(\mathcal{Y}) \rightarrow \mathcal{Coh}(\mathcal{X})$ such that $f^* \in \mathrm{Hom}_{r, \otimes, \cong}(\mathcal{Coh}(\mathcal{X}), \mathcal{Coh}(\mathcal{Y}))$.
- (5) Let \mathbf{G} be a locally algebraic group k -scheme and let X be a k -scheme equipped with an action of \mathbf{G} . We consider \mathbf{Sch}_k as a Grothendieck site with respect to *fppf topology* ([Olsson, 2016, Example 2.1.14]). Next the quotient fibered category $[X/\mathbf{G}]$ ([Monygham, 2020b, Definition 9.5]) with respect to this topology is an algebraic stack by [Olsson, 2016, Example 8.1.12].
- (6) In (5) if k -scheme X is locally noetherian (noetherian, excellent), then $[X/\mathbf{G}]$ is a locally noetherian (noetherian, excellent) by [Olsson, 2016, Definition 8.2.1, Example 8.2.3] and [Olsson, 2016, Example 8.1.12].
- (7) In (5) if k -scheme X is locally noetherian, then there exists an equivalence of monoidal categories $\mathcal{Coh}([X/\mathbf{G}]) \cong \mathcal{Coh}_{\mathbf{G}}(X)$ ([Olsson, 2016, Exercise 9.H]). Moreover, this equivalence is functorial with respect to \mathbf{G} -equivariant morphism. That is if Y is another locally noetherian k -scheme with action of \mathbf{G} and $f : X \rightarrow Y$ is a \mathbf{G} -equivariant morphism, then f induces a morphism $[f/\mathbf{G}] : [X/\mathbf{G}] \rightarrow [Y/\mathbf{G}]$ by [Monygham, 2020b, Theorem 9.7] and the square

$$\begin{array}{ccc}
\mathfrak{Coh}([Y/\mathbf{G}]) & \xrightarrow{[f/\mathbf{G}]^*} & \mathfrak{Coh}([X/\mathbf{G}]) \\
\downarrow \cong & & \downarrow \cong \\
\mathfrak{Coh}_{\mathbf{G}}(Y) & \xrightarrow{f^*} & \mathfrak{Coh}_{\mathbf{G}}(X)
\end{array}$$

of categories and functors is commutative.

(8) If \mathbf{G} is smooth and affine over k , then $[X/\mathbf{G}]$ has *affine stabilizers*.

Remark 2.1. Let $\mathrm{Spec} k$ be a point equipped with the trivial action of a smooth and affine group \mathbf{G} . Then (7) together with [Monygham, 2020a, Example 4.7] imply that $\mathfrak{Coh}([\mathrm{Spec} k/\mathbf{G}])$ can be identified with the category $\mathbf{Repf}_{\mathbf{G}}$ of finite dimensional representations of \mathbf{G} .

Let us state the main result of [Hall and Rydh, 2019].

Theorem 2.2 ([Hall and Rydh, 2019, Theorem 1.1]). *Let \mathcal{X} be a noetherian algebraic stack with affine stabilizers. For every locally excellent algebraic stack \mathcal{T} the functor*

$$\mathrm{Mor}(\mathcal{X}, \mathcal{T}) \xrightarrow{f \mapsto f^*} \mathrm{Hom}_{r, \cong}(\mathfrak{Coh}(\mathcal{T}), \mathfrak{Coh}(\mathcal{X}))$$

is an equivalence of categories.

Keeping our previous remarks in mind we deduce the following result.

Corollary 2.3. *Let \mathbf{G} be an smooth affine group k -scheme and let X, Z be k -schemes equipped with an action of \mathbf{G} . Suppose that Z is noetherian and X is locally of finite type over k . Then*

$$\mathrm{Mor}([Z/\mathbf{G}], [X/\mathbf{G}]) \xrightarrow{f \mapsto f^*} \mathrm{Hom}_{r, \cong}(\mathfrak{Coh}([X/\mathbf{G}]), \mathfrak{Coh}([Z/\mathbf{G}]))$$

is an equivalence of categories.

Proof. Note that $[Z/\mathbf{G}]$ is a noetherian algebraic stack according to (5) and (6). It has affine stabilizers according to (8). Similarly by (5) $[X/\mathbf{G}]$ is an algebraic stack. Moreover, it is locally excellent according to the fact that X is locally excellent (it is locally of finite type over k and k is a field) and (6). Then by Theorem 2.2 we derive that the functor in the statement is an equivalence of categories. \square

Corollary 2.4. *Let \mathbf{G} be an smooth affine group k -scheme and let X, Z be k -schemes equipped with an action of \mathbf{G} . Suppose that Z is noetherian and X is locally of finite type over k . Then we have a bijection*

$$\{f : Z \rightarrow X \mid f \text{ is } \mathbf{G}\text{-equivariant}\} \xrightarrow{f \mapsto f^*} \{F \in \mathrm{Hom}_{r, \cong}(\mathfrak{Coh}_{\mathbf{G}}(X), \mathfrak{Coh}_{\mathbf{G}}(Z)) \mid F \cdot p_X^* = p_Z^*\}$$

where $p_X^* : \mathbf{Repf}(\mathbf{G}) \rightarrow \mathfrak{Coh}_{\mathbf{G}}(X)$ and $p_Z^* : \mathbf{Repf}(\mathbf{G}) \rightarrow \mathfrak{Coh}_{\mathbf{G}}(Z)$ are functors induced by \mathbf{G} -equivariant morphisms $p_X : X \rightarrow \mathrm{Spec} k$ and $p_Z : Z \rightarrow \mathrm{Spec} k$, respectively.

Proof. Since fppf topology is subcanonical, [Monygham, 2020b, Theorem 9.7] shows that there exists a bijection

$$\{f : Z \rightarrow X \mid f \text{ is } \mathbf{G}\text{-equivariant}\} \xrightarrow{f \mapsto [f/\mathbf{G}]} \{h : [Z/\mathbf{G}] \rightarrow [X/\mathbf{G}] \mid [p_X/\mathbf{G}] \cdot h = [p_Z/\mathbf{G}]\}$$

Corollary 2.3 implies that there exists a bijection

$$\{h : [Z/\mathbf{G}] \rightarrow [X/\mathbf{G}] \mid [p_X/\mathbf{G}] \cdot h = [p_Y/\mathbf{G}]\} \xrightarrow{h \mapsto h^*} \{F \in \text{Hom}_{r, \otimes, \cong}(\mathcal{Coh}([X/\mathbf{G}]), \mathcal{Coh}([Z/\mathbf{G}])) \mid F \cdot [p_X/\mathbf{G}]^* = [p_Z/\mathbf{G}]^*\}$$

Next (7) implies that there exists a bijection

$$\{F \in \text{Hom}_{r, \otimes, \cong}(\mathcal{Coh}([X/\mathbf{G}]), \mathcal{Coh}([Z/\mathbf{G}])) \mid F \cdot [p_X/\mathbf{G}]^* = [p_Z/\mathbf{G}]^*\} \longrightarrow \{F \in \text{Hom}_{r, \otimes, \cong}(\mathcal{Coh}_{\mathbf{G}}(X), \mathcal{Coh}_{\mathbf{G}}(Z)) \mid F \cdot p_X^* = p_Z^*\}$$

and for a \mathbf{G} -equivariant morphism $f : Z \rightarrow X$ the image of $[f/\mathbf{G}]^* : \mathcal{Coh}([X/\mathbf{G}]) \rightarrow \mathcal{Coh}([Z/\mathbf{G}])$ under this bijection is $f^* : \mathcal{Coh}_{\mathbf{G}}(X) \rightarrow \mathcal{Coh}_{\mathbf{G}}(Z)$. These imply that the map of classes

$$\{f : Z \rightarrow X \mid f \text{ is } \mathbf{G}\text{-equivariant}\} \xrightarrow{f \mapsto f^*} \{F \in \text{Hom}_{r, \otimes, \cong}(\mathcal{Coh}_{\mathbf{G}}(X), \mathcal{Coh}_{\mathbf{G}}(Z)) \mid F \cdot p_X^* = p_Z^*\}$$

is a bijection. □

Note that Corollary 2.4 relies on some assumptions regarding \mathbf{G} , X and Z . It is worth noting that Joachim Jelisiejew and the author were able to obtain a slightly more general (yet unpublished) result.

Theorem 2.5 ([Jelisiejew and Sienkiewicz, 2020, Theorem A.2]). *Let \mathbf{G} be an affine algebraic group over k . Let Z, X be k -schemes equipped with an action of \mathbf{G} and assume that X is quasi-compact and quasi-separated. Suppose that $F : \mathcal{Qcoh}_{\mathbf{G}}(X) \rightarrow \mathcal{Qcoh}_{\mathbf{G}}(Z)$ is a cocontinuous, monoidal functor such that $F \cdot p_X^* = p_Z^*$. Then there exists a unique \mathbf{G} -equivariant morphism $f : Z \rightarrow X$ such that $f^* = F$.*

3. RELATIONS BETWEEN REPRESENTATIONS OF A MONOID AND ITS GROUP OF UNITS

In this section we study the relation between the category $\mathbf{Rep}(\mathbf{M})$ of representations of an affine monoid k -scheme \mathbf{M} and the category $\mathbf{Rep}(\mathbf{G})$ of representations of its group of units \mathbf{G} . Let $i : k[\mathbf{M}] \rightarrow k[\mathbf{G}]$ be the morphism of k -bialgebras induced by $\mathbf{G} \hookrightarrow \mathbf{M}$. Let us first note the following elementary result.

Fact 3.1. *Assume that \mathbf{G} is open and schematically dense in \mathbf{M} . Then i is an injective morphism of k -algebras.*

Proof. This follows from [Görtz and Wedhorn, 2010, Proposition 9.19]. □

Fact 3.2. *The forgetful functor $\mathbf{Rep}(\mathbf{M}) \rightarrow \mathbf{Rep}(\mathbf{G})$ creates colimits and finite limits.*

Proof. This follows from [Monygham, 2020d, Theorem 14.3, Theorem 14.4] and the commutative triangle

$$\begin{array}{ccc} \mathbf{Rep}(\mathbf{M}) & \xrightarrow{\quad} & \mathbf{Rep}(\mathbf{G}) \\ & \searrow \quad \swarrow & \\ & \mathbf{Vect}_k & \end{array}$$

of functors. □

The theorem below characterizes representations of \mathbf{G} which are contained in the image of the forgetful functor $\mathbf{Rep}(\mathbf{M}) \rightarrow \mathbf{Rep}(\mathbf{G})$.

Theorem 3.3. *Assume that \mathbf{G} is open and schematically dense in \mathbf{M} . Let V be a \mathbf{G} -representation. Then the following are equivalent.*

- (i) V is in the image of the forgetful functor $\mathbf{Rep}(\mathbf{M}) \rightarrow \mathbf{Rep}(\mathbf{G})$.
- (ii) The coaction $d : V \rightarrow k[\mathbf{G}] \otimes_k V$ factors through $i \otimes_k 1_V : k[\mathbf{M}] \otimes_k V \hookrightarrow k[\mathbf{G}] \otimes_k V$.

Proof. In the proof we denote by $\Delta_{\mathbf{M}}$ and $\Delta_{\mathbf{G}}$ comultiplications of $k[\mathbf{M}]$ and $k[\mathbf{G}]$, respectively. We also denote by $\zeta_{\mathbf{M}}$ and $\zeta_{\mathbf{G}}$ counits of $k[\mathbf{M}]$ and $k[\mathbf{G}]$, respectively. According to Fact 3.1 i is an injective morphism of k -algebras.

Clearly (i) \Rightarrow (ii). We prove the converse. Suppose that (ii) holds. Let $c : V \rightarrow k[\mathbf{M}] \otimes_k V$ be a unique morphism such that $d = (i \otimes_k 1_V) \cdot c$. It suffices to prove that c is the coaction of the bialgebra $k[\mathbf{M}]$ on V . Observe that

$$\begin{aligned} (i \otimes_k i \otimes_k 1_V) \cdot (1_{k[\mathbf{M}]} \otimes_k c) \cdot c &= (i \otimes_k d) \cdot c = (1_{k[\mathbf{G}]} \otimes_k d) \cdot d = (\Delta_{\mathbf{G}} \otimes_k 1_V) \cdot d = \\ &= (\Delta_{\mathbf{G}} \otimes_k 1_V) \cdot ((i \otimes_k 1_V) \cdot c) = ((\Delta_{\mathbf{G}} \cdot i) \otimes_k 1_V) \cdot c = (i \otimes_k i \otimes_k 1_V) \cdot (\Delta_{\mathbf{M}} \otimes_k 1_V) \cdot c \end{aligned}$$

Since $i \otimes_k i \otimes_k 1_V$ is a monomorphism, we deduce that $(1_{k[\mathbf{M}]} \otimes_k c) \cdot c = (\Delta_{\mathbf{M}} \otimes_k 1_V) \cdot c$. Moreover, we have

$$(\zeta_{\mathbf{G}} \otimes_k 1_V) \cdot d = (\zeta_{\mathbf{G}} \otimes_k 1_V) \cdot ((i \otimes_k 1_V) \cdot c) = (\zeta_{\mathbf{M}} \otimes_k 1_V) \cdot c$$

and hence $(\zeta_{\mathbf{M}} \otimes_k 1_V) \cdot c$ is the canonical isomorphism $V \cong k \otimes_k V$. Thus c is the coaction of $k[\mathbf{M}]$ and $d = (i \otimes_k 1_V) \cdot c$. Therefore, V is in the image of $\mathbf{Rep}(\mathbf{M}) \rightarrow \mathbf{Rep}(\mathbf{G})$. \square

Theorem 3.4. *Assume that \mathbf{G} is open and schematically dense in \mathbf{M} . Then $\mathbf{Rep}(\mathbf{M})$ is a full subcategory of $\mathbf{Rep}(\mathbf{G})$ closed under subobjects and quotients.*

Proof. In the proof we denote by $\Delta_{\mathbf{M}}$ and $\Delta_{\mathbf{G}}$ comultiplications of $k[\mathbf{M}]$ and $k[\mathbf{G}]$, respectively. We also denote by $\zeta_{\mathbf{M}}$ and $\zeta_{\mathbf{G}}$ counits of $k[\mathbf{M}]$ and $k[\mathbf{G}]$, respectively. According to Fact 3.1 i is an injective morphism of k -algebras.

We first prove that $\mathbf{Rep}(\mathbf{M})$ is a full subcategory of $\mathbf{Rep}(\mathbf{G})$. For this consider \mathbf{M} -representations V, W and a their morphism $f : V \rightarrow W$ as \mathbf{G} -representations. Let c_V and c_W be coactions of $k[\mathbf{M}]$ on V and W , respectively. Our goal is to prove that f is a morphism of \mathbf{M} -representations. Consider the diagram

$$\begin{array}{ccc} k[\mathbf{G}] \otimes_k V & \xrightarrow{1_{k[\mathbf{G}]} \otimes_k f} & k[\mathbf{G}] \otimes_k W \\ \uparrow i \otimes_k 1_V & & \uparrow i \otimes_k 1_W \\ k[\mathbf{M}] \otimes_k V & \xrightarrow{\quad \quad \quad 1_{k[\mathbf{M}]} \otimes_k f \quad \quad \quad} & k[\mathbf{M}] \otimes_k W \\ \uparrow c_V & & \uparrow c_W \\ V & \xrightarrow{\quad \quad \quad f \quad \quad \quad} & W \end{array}$$

in which the outer square is commutative. Our goal is to prove that the bottom square is commutative. We have

$$(i \otimes_k 1_W) \cdot c_W \cdot f = (1_{k[\mathbf{G}]} \otimes_k f) \cdot (i \otimes_k 1_V) \cdot c_V = (i \otimes_k 1_W) \cdot (1_{k[\mathbf{M}]} \otimes_k f) \cdot c_V$$

Since $i \otimes_k 1_W$ is a monomorphism, we deduce that $c_W \cdot f = (1_{k[\mathbf{M}]} \otimes_k f) \cdot c_V$. Hence f is a morphism of \mathbf{M} -representations.

Next we prove that $\mathbf{Rep}(\mathbf{M})$ is a subcategory of $\mathbf{Rep}(\mathbf{G})$ that is closed under subquotients. Consider an \mathbf{M} -representation V and its quotient \mathbf{G} -representations $q : V \twoheadrightarrow W$. We show that W is a quotient \mathbf{M} -representation of V . Let c_V be the coaction of \mathbf{M} on V and let d_W be the coaction of \mathbf{G} on W . We have a commutative diagram

$$\begin{array}{ccc}
 k[\mathbf{G}] \otimes_k V & \xrightarrow{1_{k[\mathbf{G}]} \otimes_k q} & k[\mathbf{G}] \otimes_k W \\
 i \otimes_k 1_V \uparrow & & \uparrow d_W \\
 k[\mathbf{M}] \otimes_k V & & \\
 c_V \uparrow & & \\
 V & \xrightarrow{q} & W
 \end{array}$$

and hence $d_W(W) \subseteq k[\mathbf{M}] \otimes_k W$. Thus Theorem 3.3 implies that W is a representation of \mathbf{M} and q is a morphism of \mathbf{M} -representations. This shows that $\mathbf{Rep}(\mathbf{M})$ is a subcategory of $\mathbf{Rep}(\mathbf{G})$ closed under quotients. Next let $j : U \hookrightarrow V$ be a \mathbf{G} -subrepresentation of a \mathbf{M} -representation V . By what we proved above the cokernel $q : V \twoheadrightarrow W$ of j in $\mathbf{Rep}(\mathbf{G})$ is contained in $\mathbf{Rep}(\mathbf{M})$. Since both $\mathbf{Rep}(\mathbf{M})$ and $\mathbf{Rep}(\mathbf{G})$ are abelian and the forgetful functor $\mathbf{Rep}(\mathbf{M}) \rightarrow \mathbf{Rep}(\mathbf{G})$ is exact, we derive that the kernel of q in $\mathbf{Rep}(\mathbf{M})$ coincides with its kernel in $\mathbf{Rep}(\mathbf{G})$. Thus U is a \mathbf{M} -representation and $j : U \hookrightarrow V$ is a morphism of \mathbf{M} -representations. Hence $\mathbf{Rep}(\mathbf{M})$ is the category of $\mathbf{Rep}(\mathbf{G})$ closed under subobjects. \square

Theorem 3.5. *Assume that \mathbf{G} is open and schematically dense in \mathbf{M} . Let V be a \mathbf{G} -representation of \mathbf{G} . There exists an \mathbf{M} -representation W and a surjective morphism $q : V \twoheadrightarrow W$ of \mathbf{G} -representations such that for every \mathbf{M} -representation U and a morphism $f : V \rightarrow U$ of \mathbf{G} -representations there exists a unique morphism $\tilde{f} : W \rightarrow U$ of \mathbf{M} -representations making the triangle*

$$\begin{array}{ccc}
 V & \xrightarrow{f} & U \\
 q \downarrow & \nearrow \tilde{f} & \\
 W & &
 \end{array}$$

commutative.

Proof. Assume first that V is finite dimensional. Let \mathcal{K} be a set of \mathbf{G} -subrepresentations of V that consists of all $K \subseteq V$ such that V/K carries a structure of \mathbf{M} -representation. Clearly $\mathcal{K} = \emptyset$ because $\{0\} \in \mathcal{K}$. Since V is finite dimensional, there exists a finite subset $\{K_1, \dots, K_n\} \subseteq \mathcal{K}$ such that

$$\bigcap_{i=1}^n K_i = \bigcap_{K \in \mathcal{K}} K$$

Then a morphism

$$V / \left(\bigcap_{K \in \mathcal{K}} K \right) \ni v \mapsto (v \bmod K_i)_{1 \leq i \leq n} \in \bigoplus_{i=1}^n V/K_i$$

is a monomorphism and hence by Theorem 3.4 the quotient $W = V / (\bigcap_{K \in \mathcal{K}} K)$ is an \mathbf{M} -representation. Let $q : V \twoheadrightarrow W$ be the canonical epimorphism. Consider now a morphism $f : V \rightarrow U$ of \mathbf{G} -representations, where U is an \mathbf{M} -representation. Then $\text{im}(f)$ is a \mathbf{G} -subrepresentation of U and by Theorem 3.4 we derive that $\text{im}(f)$ is an \mathbf{M} -representation. This implies that $\ker(f)$ is in \mathcal{K} . Hence f factors through q . Thus there exists a unique morphism $\tilde{f} : W \rightarrow U$ of \mathbf{G} -representations such that $\tilde{f} \cdot q = f$. This completes the proof in case when V is finite dimensional.

Now consider the general V . Let \mathcal{F} be the set of all finite dimensional \mathbf{G} -representations of V . According to [Monygham, 2020d, Corollary 15.2] we deduce that $V = \operatorname{colim}_{F \in \mathcal{F}} F$. By the case considered above we deduce that for every F in \mathcal{F} there exists a universal morphism $q_F : F \rightarrow W_F$ of \mathbf{G} -representations into an \mathbf{M} -representation W_F . Note that if $F_1 \subseteq F_2$ are two elements of \mathcal{F} , then

$$\begin{array}{ccc} F_1 & \xrightarrow{q_{F_1}} & W_{F_1} \\ \downarrow & & \downarrow \\ F_2 & \xrightarrow{q_{F_2}} & W_{F_2} \end{array}$$

Thus $\{W_F\}_{F \in \mathcal{F}}$ together with morphisms $W_{F_1} \rightarrow W_{F_2}$ for $F_1 \subseteq F_2$ in \mathcal{F} form a diagram parametrized by the poset \mathcal{F} . The category $\mathbf{Rep}(\mathbf{M})$ has small colimits ([Monygham, 2020d, Corollary 14.5]) and we define $W = \operatorname{colim}_{F \in \mathcal{F}} W_F$. This is also a colimit of this diagram in the category $\mathbf{Rep}(\mathbf{G})$ by Fact 3.2. We also define $q = \operatorname{colim}_{F \in \mathcal{F}} q_F : V = \operatorname{colim}_{F \in \mathcal{F}} F \rightarrow W$. Since a colimit of a family of epimorphisms is an epimorphism, we derive that q is an epimorphism of \mathbf{G} -representations. Suppose now that $f : V \rightarrow U$ is a morphism of \mathbf{G} -representations and U is an \mathbf{M} -representation. Then $f|_F$ uniquely factors through q_F for every F in \mathcal{F} . Hence by universal property of colimits we derive that f factors through q in a unique way. This completes the proof. \square

4. BIALYNICKI-BIRULA FUNCTORS

In this section we fix an affine group k -scheme \mathbf{G} . Let \mathbf{M} be an affine monoid k -scheme with zero \mathbf{o} such that \mathbf{G} is its group of units.

Definition 4.1. Let X be a k -scheme equipped with an action of \mathbf{G} . For every k -scheme Y we define

$$\mathcal{D}_X(Y) = \{\gamma : \mathbf{M} \times_k Y \rightarrow X \mid \gamma \text{ is } \mathbf{G}\text{-equivariant}\}$$

This gives rise to a subfunctor \mathcal{D}_X of $\operatorname{Mor}_k(\mathbf{M} \times_k (-), X) : \mathbf{Sch}_k^{\operatorname{op}} \rightarrow \mathbf{Set}$. We call it *the Białynicki-Birula functor of X* .

Fact 4.2. Let X be a scheme equipped with an action of \mathbf{G} . Then \mathcal{D}_X is a Zariski sheaf.

Proof. This is a consequence of the fact that $\operatorname{Mor}_k(\mathbf{M} \times_k (-), X)$ is a Zariski sheaf and if we glue \mathbf{G} -equivariant morphisms, then the result is \mathbf{G} -equivariant. Indeed, this shows that \mathcal{D}_X is a Zariski subsheaf of $\operatorname{Mor}_k(\mathbf{M} \times_k (-), X)$. \square

Remark 4.3. Let X be a k -scheme equipped with an action of \mathbf{G} . Then there are canonical morphism of functors

$$\begin{array}{ccc} \mathcal{D}_X & \xrightarrow{i_X} & X \\ \downarrow s_X & & \downarrow r_X \\ & & X^{\mathbf{G}} \end{array}$$

which we define now. First let us explain that in the diagram X stands for the presheaf representable by the k -scheme X (Remark 1.1) and $X^{\mathbf{G}}$ denotes the functor of fixed points of X ([Monygham, 2020c, Definition 7.1]). Now fix k -scheme Y and $\gamma \in \mathcal{D}_X(Y)$, then we define

$$i_X(\gamma) = \gamma|_{\{e\} \times_k X} = \gamma \cdot \langle e, 1_X \rangle, \quad r_X(\gamma) = \gamma|_{\{\mathbf{o}\} \times_k X} = \gamma \cdot \langle \mathbf{o}, 1_X \rangle$$

where $e : \text{Spec } k \rightarrow \mathbf{M}$ is the unit of \mathbf{M} and $\mathbf{o} : \text{Spec } k \rightarrow \mathbf{M}$ is the zero. Next if $f : Y \rightarrow X$ is a morphism in $X^{\mathbf{G}}(Y)$, then we define

$$s_X(f) = f \cdot pr_Y$$

where $pr_Y : \mathbf{M} \times_k Y \rightarrow Y$ is the projection. Finally note that $r_X \cdot s_X = 1_{X^{\mathbf{G}}}$.

Remark 4.4. Let X be a k -scheme equipped with an action of \mathbf{G} . Then \mathbf{M} (actually the presheaf of monoids represented by \mathbf{M}) acts on \mathcal{D}_X . Indeed, fix k -scheme Y , $\gamma \in \mathcal{D}_X(Y)$ and $m : Y \rightarrow \mathbf{M}$. Then we define the product

$$m\gamma = \gamma \cdot \langle m, 1_Y \rangle$$

and this determines an action of \mathbf{M} on \mathcal{D}_X . Moreover, with respect to this action i_X is \mathbf{G} -equivariant and r_X, s_X are \mathbf{M} -equivariant ($X^{\mathbf{G}}$ is equipped with trivial action of \mathbf{M}).

Remark 4.5. Let X, Y be k -schemes equipped with actions of \mathbf{G} and let $f : X \rightarrow Y$ be a \mathbf{G} -equivariant morphism, then there exists a morphism of functors $\mathcal{D}_f : \mathcal{D}_X \rightarrow \mathcal{D}_Y$ given by

$$\mathcal{D}_f(\gamma) = f \cdot \gamma$$

for every element γ of the functor \mathcal{D}_X .

Let X be a k -scheme equipped with an action of \mathbf{G} . It is useful to discuss subfunctors of \mathcal{D}_X defined by closed \mathbf{G} -stable subschemes of X .

Theorem 4.6. Let X be a k -scheme equipped with an action of the group \mathbf{G} . Suppose that \mathbf{G} is open and schematically dense in \mathbf{M} . If $j : Z \hookrightarrow X$ is a closed \mathbf{G} -stable subscheme of X , then the square

$$\begin{array}{ccc} \mathcal{D}_Z & \xrightarrow{\mathcal{D}_j} & \mathcal{D}_X \\ i_Z \downarrow & & \downarrow i_X \\ Z & \xrightarrow{j} & X \end{array}$$

is cartesian in the category of presheaves on \mathbf{Sch}_k .

Proof. The fact that the square is commutative follows by examination of definitions in Remarks 4.3 and 4.5. Pick k -scheme Y , $f : Y \rightarrow Z$ and $\gamma \in \mathcal{D}_X(Y)$ such that $j \cdot f = i_X(\gamma)$. This is depicted in the diagram

$$\begin{array}{ccc} & \gamma & \\ & \downarrow i_X & \\ f \mapsto_j & j \cdot f = \gamma|_{\{e\} \times_k X} & \end{array}$$

Our goal is to show that there exists a unique \mathbf{G} -equivariant morphism $\eta : \mathbf{M} \times_k Y \rightarrow U$ such that $\mathcal{D}_j(\eta) = \gamma$ and $i_Z(\eta) = f$. This is depicted by the diagram

$$\begin{array}{ccc} \eta & \xrightarrow{\mathcal{D}_j} & \gamma = j \cdot \eta \\ r_U \downarrow & & \\ f = \eta|_{\{e\} \times_k X} & & \end{array}$$

It suffices to prove that γ factors through j . First note that the assumption $\gamma|_{\{e\} \times_k Y} = j \cdot f$ implies that

$$\gamma|_{\mathbf{G} \times_k Y} = j \cdot f \cdot pr_Y$$

where $pr_Y : \mathbf{G} \times_k Y \rightarrow Y$ is the projection. This implies that $\gamma|_{\mathbf{G} \times_k}$ factors through j . Consider scheme-theoretic preimage $\gamma^{-1}(Z)$. Then $\gamma^{-1}(Z)$ is a closed \mathbf{G} -stable (as an inverse image of a \mathbf{G} -stable closed subscheme under the \mathbf{G} -equivariant morphism) subscheme of $\mathbf{M} \times_k Y$, which contains $\mathbf{G} \times_k Y$. Since \mathbf{G} is open, schematically dense in \mathbf{M} and k is a field, we derive that $\mathbf{G} \times_k Y$ is open and schematically dense in $\mathbf{M} \times_k Y$. Thus $\gamma^{-1}(Z) = \mathbf{M} \times_k Y$ and hence γ factors through j . \square

In order to prove interesting result in the spirit of Theorem 4.6 which concerns open \mathbf{G} -stable subschemes, we need to assume that \mathbf{M} is a Kempf monoid.

Theorem 4.7. *Let X be a k -scheme equipped with an action of the group \mathbf{G} of units of a Kempf monoid \mathbf{M} . If $j : U \hookrightarrow X$ is an open \mathbf{G} -stable subscheme of X , then the square*

$$\begin{array}{ccc} \mathcal{D}_U & \xrightarrow{\mathcal{D}_j} & \mathcal{D}_X \\ r_U \downarrow & & \downarrow r_X \\ U^G & \xrightarrow{j^G} & X^G \end{array}$$

is cartesian in the category of presheaves on \mathbf{Sch}_k .

As we shall see this result follows from the following.

Lemma 4.7.1. *Let K be an algebraically closed field over k . Suppose that*

$$\mathbf{M}_K = \mathrm{Spec} K \times_k \mathbf{M}, \mathbf{G}_K = \mathrm{Spec} K \times_k \mathbf{G}$$

and let \mathbf{o}_K be the unique K -point of \mathbf{M}_K lying over \mathbf{o} . Let V be an open \mathbf{G}_K -stable subscheme of \mathbf{M}_K such that $\mathbf{o}_K \in V$. Then $V = \mathbf{M}_K$.

Proof of the lemma. Since \mathbf{M} is a Kempf monoid, there exists a closed embedding of monoids $v : \mathbb{A}_K^1 \hookrightarrow \mathbf{M}_K$ preserving zeros such that $v|_{\mathbb{G}_{m,K}} \subseteq \mathbf{G}_K$. Fix a point $p \in \mathbf{M}_K$ and let $u : \mathrm{Spec} k(p) \rightarrow \mathbf{M}_K$ be the associated morphism of K -schemes. Consider the composition

$$\mathbb{A}_{k(p)}^1 = \mathbb{A}_K^1 \times_K \operatorname{Spec} k(p) \xrightarrow{v \times_K u} \mathbf{M}_K \times_K \mathbf{M}_K \xrightarrow{\quad} \mathbf{M}_K$$

where the second morphism is the multiplication. Clearly h is $\mathbf{G}_{m,k(p)}$ -equivariant. Hence $h^{-1}(V)$ is an open $\mathbf{G}_{m,k(p)}$ -stable subscheme of $\mathbb{A}_{k(p)}^1$ containing zero of this monoid $k(p)$ -scheme (because $\mathbf{o}_K \in V$ by assumption). Since the only open $\mathbf{G}_{m,k(p)}$ -stable subscheme of $\mathbb{A}_{k(p)}^1$ containing zero is $\mathbb{A}_{k(p)}^1$, we derive that $h^{-1}(V) = \mathbb{A}_{k(p)}^1$. Thus $p \in V$. Since p is arbitrary point of \mathbf{M}_K , we derive that $V = \mathbf{M}_K$. \square

Proof of the theorem. The fact that the square is commutative follows by examination of definitions in Remarks 4.3 and 4.5. Pick k -scheme Y , $f \in U^G$ and $\gamma \in \mathcal{D}_X(Y)$ such that $j^G(f) = r_X(\gamma)$. This is depicted in the diagram

$$\begin{array}{ccc}
 & \gamma & \\
 & \downarrow r_X & \\
 f \xrightarrow{j^G} j \cdot f & = & \gamma|_{\{\mathbf{o}\} \times_k X}
 \end{array}$$

Our goal is to show that there exists a unique \mathbf{G} -equivariant morphism $\eta : \mathbf{M} \times_k Y \rightarrow U$ such that $\mathcal{D}_j(\eta) = \gamma$ and $r_U(\eta) = f$. This is depicted by the diagram

$$\begin{array}{ccc}
 \eta & \xrightarrow{\mathcal{D}_j} & \gamma = j \cdot \eta \\
 \downarrow r_U & & \\
 f = \eta|_{\{\mathbf{o}\} \times_k X} & &
 \end{array}$$

For this it suffices to prove that γ factors through j . Consider $W = \gamma^{-1}(U)$. Note that W is an open \mathbf{G} -stable (as an inverse image of a \mathbf{G} -stable open subscheme under the \mathbf{G} -equivariant morphism) subscheme of $\mathbf{M} \times_k Y$, which contains $\{\mathbf{o}\} \times_k Y$. Lemma 4.7.1 asserts that for every geometric point \bar{y} of Y we have $W_{\bar{y}} = \mathbf{M}_{k(\bar{y})}$, where $W_{\bar{y}}$ is the fiber over \bar{y} of the projection $\mathbf{M} \times_k Y \rightarrow Y$ restricted to W . Since W is open subscheme of $\mathbf{M} \times_k Y$, this implies that $W = \mathbf{M} \times_k Y$ and hence γ factors through j . \square

As we shall see below both Theorems are extremely useful properties of Białynicki-Birula functors. Now we introduce a formal version of this functor.

Definition 4.8. Let \mathbf{M} be an affine monoid k -scheme with zero \mathbf{o} and let \mathbf{G} be its group of units. For every $n \in \mathbb{N}$ let $\mathbf{M}_n \hookrightarrow \mathbf{M}$ be an n -th infinitesimal neighborhood of \mathbf{o} in \mathbf{M} . Let X be a k -scheme equipped with an action of \mathbf{G} . For every k -scheme Y we define

$$\widehat{\mathcal{D}}_X(Y) = \left\{ \{\gamma_n : \mathbf{M}_n \times_k Y \rightarrow X\}_{n \in \mathbb{N}} \mid \forall n \in \mathbb{N} \gamma_n \text{ is } \mathbf{G}\text{-equivariant and } \gamma_{n+1}|_{\mathbf{M}_n \times_k Y} = \gamma_n \right\}$$

This gives rise to a functor $\widehat{\mathcal{D}}_X$. We call it *the formal Białynicki-Birula functor of X* .

Remark 4.9. Let \mathbf{M} be an affine monoid k -scheme with zero \mathbf{o} and let \mathbf{G} be its group of units. Let X be a k -scheme equipped with an action of \mathbf{G} . Then there exists a canonical morphism of functors $\mathcal{D}_X \rightarrow \widehat{\mathcal{D}}_X$ given by

$$\gamma \mapsto \{\gamma|_{\mathbf{M}_n \times_k Y}\}_{n \in \mathbb{N}}$$

for every $\gamma \in \mathcal{D}_X(Y)$ and every k -scheme Y .

5. REPRESENTABILITY OF BIAŁYNICKI-BIRULA FUNCTOR FOR KEMPF MONOIDS

In this section we prove various results concerning representability of Białynicki-Birula functors.

Theorem 5.1. *Let \mathbf{M} be an affine monoid k -scheme with open and schematically dense group of units \mathbf{G} . Suppose that X is an affine k -scheme equipped with an action of \mathbf{G} . Then \mathcal{D}_X is representable and i_X is a closed immersion of k -schemes.*

Proof. Since X is an affine k -scheme, the action of \mathbf{G} on X corresponds to the coaction of $k[\mathbf{G}]$ by $c : \Gamma(X, \mathcal{O}_X) \rightarrow k[\mathbf{G}] \otimes_k \Gamma(X, \mathcal{O}_X)$. Note that c is a morphism of k -algebras. By Theorem 3.5 there exists a universal morphism $q : \Gamma(X, \mathcal{O}_X) \twoheadrightarrow W$ of \mathbf{G} -representations into a \mathbf{M} -representation. Let $I \subseteq \Gamma(X, \mathcal{O}_X)$ be the ideal generated by $\ker(q)$. Fix f in I . Then

$$f = \sum_{i=1}^n g_i \cdot f_i$$

where $g_i \in k[\mathbf{G}]$ and $f_i \in \ker(q)$ for $1 \leq i \leq n$. Then

$$c(f) = c\left(\sum_{i=1}^n g_i \cdot f_i\right) = \sum_{i=1}^n c(g_i) \cdot c(f_i) \subseteq \left(k[\mathbf{G}] \otimes_k \Gamma(X, \mathcal{O}_X)\right) \cdot \left(k[\mathbf{G}] \otimes_k \ker(q)\right) \subseteq k[\mathbf{G}] \otimes_k I$$

Thus $c(I) \subseteq k[\mathbf{G}] \otimes_k I$ and hence I is a \mathbf{G} -representation. Consider

$$X^+ = V(I) = \operatorname{Spec} \Gamma(X, \mathcal{O}_X)/I \hookrightarrow X$$

Since $\Gamma(X, \mathcal{O}_X)/I$ is the quotient \mathbf{G} -representation of W , we deduce by Theorem 3.5 that $\Gamma(X, \mathcal{O}_X)/I$ is a \mathbf{M} -representation. Hence X^+ is a k -scheme equipped with action of \mathbf{M} and $X^+ \hookrightarrow X$ is \mathbf{G} -equivariant. Suppose now that Y is an affine k -scheme. Then $\mathbf{M} \times_k Y$ is a \mathbf{M} -scheme with respect to the left-hand side action of \mathbf{M} and hence $\Gamma(\mathbf{M} \times_k Y, \mathcal{O}_{\mathbf{M} \times_k Y})$ is a \mathbf{M} -representation. Now Theorem 3.5 implies that if $\gamma : \mathbf{M} \times_k Y \rightarrow X$ is a \mathbf{G} -equivariant morphism, then a morphism $\gamma^\# : \Gamma(X, \mathcal{O}_X) \rightarrow \Gamma(\mathbf{M} \times_k Y, \mathcal{O}_{\mathbf{M} \times_k Y})$ of k -algebras and \mathbf{G} -representations factors through $q : \Gamma(X, \mathcal{O}_X) \twoheadrightarrow W$ and thus by construction of I we have

$$\begin{array}{ccccc} & & \gamma^\# & & \\ & \nearrow & & \searrow & \\ \Gamma(X, \mathcal{O}_X) & \twoheadrightarrow & \Gamma(X, \mathcal{O}_X)/I & \xrightarrow{f} & \Gamma(\mathbf{M} \times_k Y, \mathcal{O}_{\mathbf{M} \times_k Y}) \end{array}$$

for some morphism f of k -algebras and \mathbf{G} -representations. Since both $\Gamma(X, \mathcal{O}_X)/I$ and $\Gamma(\mathbf{M} \times_k Y, \mathcal{O}_{\mathbf{M} \times_k Y})$ are \mathbf{M} -representations and by Theorem 3.4 the subcategory $\mathbf{Rep}(\mathbf{M}) \subseteq \mathbf{Rep}(\mathbf{G})$ is full, we derive that f is a morphism of \mathbf{M} -representations. Thus f corresponds to a unique \mathbf{M} -equivariant morphism $\eta : \mathbf{M} \times_k Y \rightarrow X^+$ such that the diagram

$$\begin{array}{ccc} & & X^+ \\ & \nearrow \eta & \downarrow \\ \mathbf{M} \times_k Y & \xrightarrow{\gamma} & X \end{array}$$

is commutative. Now this result can be extended to an arbitrary k -scheme Y , since $\operatorname{Mor}_k(\mathbf{M} \times_k (-), X^+)$ is a Zariski sheaf and a morphism that is \mathbf{M} -equivariant locally on the domain is \mathbf{M} -equivariant. Thus for every h -scheme Y we have a bijection

$$\mathcal{D}_X(Y) \ni \gamma \mapsto \eta \in \{\mathbf{M}\text{-equivariant morphisms } \mathbf{M} \times_k Y \rightarrow X^+\}$$

Since we also have a bijection

$$\{\mathbf{M}\text{-equivariant morphisms } \mathbf{M} \times_k Y \rightarrow X^+\} \ni \eta \mapsto \eta \cdot \langle e, 1_{X^+} \rangle \in \operatorname{Mor}_k(Y, X^+)$$

and both this bijections are natural, we derive that \mathcal{D}_X is represented by X^+ and moreover, $i_X : \mathcal{D}_X \rightarrow X$ is a closed immersion $X^+ \hookrightarrow X$. \square

Corollary 5.2. *Let \mathbf{G} be a group k -scheme and \mathbf{M} be a Kempf monoid having \mathbf{G} as a group of units. Suppose that X is a k -scheme equipped with an action of \mathbf{G} such that there exists a family \mathcal{U} of open affine \mathbf{G} -stable open subschemes of X such that functors $\{U^\mathbf{G}\}_{U \in \mathcal{U}}$ form an open cover of $X^\mathbf{G}$. Then \mathcal{D}_X is representable.*

Proof. Note that \mathbf{G} is affine group k -scheme as a unit group of an affine monoid \mathbf{M} ([Monygham, 2020d, Proposition 12.4]). Moreover, \mathbf{M} is a Kempf monoid and hence \mathbf{G} is open and schematically dense in \mathbf{M} . By Theorem 5.1 each \mathcal{D}_U is representable by a k -scheme. On the other hand by Theorem 4.7 for each $U \in \mathcal{U}$ we have a cartesian square

$$\begin{array}{ccc} \mathcal{D}_U & \longrightarrow & \mathcal{D}_X \\ r_U \downarrow & & \downarrow r_X \\ U^{\mathbf{G}} & \longrightarrow & X^{\mathbf{G}} \end{array}$$

of functors. This implies that $\{\mathcal{D}_U \hookrightarrow \mathcal{D}_X\}_{U \in \mathcal{U}}$ is an open cover of \mathcal{D}_X as a pullback of an open cover $\{U^{\mathbf{G}} \hookrightarrow X^{\mathbf{G}}\}_{U \in \mathcal{U}}$. Hence Fact 4.2 and [Görtz and Wedhorn, 2010, Theorem 8.9] (or if you like [Monygham, 2019, Theorem 4.6]) imply that \mathcal{D}_X is representable. \square

Corollary 5.3. *Let \mathbf{G} be group k -scheme and \mathbf{M} be a Kempf monoid having \mathbf{G} as a group of units. Suppose that X is a locally linear \mathbf{G} -scheme. Then \mathcal{D}_X is representable.*

Proof. This is a consequence of Corollary 5.2. Indeed, X admits a cover \mathcal{U} by open \mathbf{G} -stable affine subschemes. Then $\{U^{\mathbf{G}}\}_{U \in \mathcal{U}}$ is an open cover of $X^{\mathbf{G}}$. \square

In order to prove more interesting result we need to recall some fact.

Remark 5.4. Let \mathbf{G} be a geometrically connected and locally algebraic group over k . It follows by [Monygham, 2020c, Theorem 7.2] that for every k -scheme X equipped with an action of \mathbf{G} there exists closed subscheme $X^{\mathbf{G}}$ of X representing the fixed point functor.

Theorem 5.5. *Let \mathbf{G} be a geometrically connected and locally algebraic group k -scheme and \mathbf{M} be a Kempf monoid having \mathbf{G} as a group of units. Suppose that X is a k -scheme equipped with action of \mathbf{G} . Then $\widehat{\mathcal{D}}_X$ is representable.*

REFERENCES

- [Görtz and Wedhorn, 2010] Görtz, U. and Wedhorn, T. (2010). *Algebraic Geometry: Part I: Schemes. With Examples and Exercises*. Advanced Lectures in Mathematics.
- [Hall and Rydh, 2019] Hall, J. and Rydh, D. (2019). Coherent tannaka duality and algebraicity of hom-stacks. *Algebra Number Theory*, 13(7):1633–1675.
- [Jelisiejew and Sienkiewicz, 2020] Jelisiejew, J. and Sienkiewicz, Ł. (2020). Białynicki-birula decomposition for reductive groups in positive characteristic. *arXiv*, pages arXiv–2006.
- [Monygham, 2019] Monygham (2019). Geometry of k -functors. *github repository: "Monygham/Pedo-mellon-a-minno"*.
- [Monygham, 2020a] Monygham (2020a). Algebraization of formal \mathbf{M} -schemes. *github repository: "Monygham/Pedo-mellon-a-minno"*.
- [Monygham, 2020b] Monygham (2020b). Fibered categories and equivariant objects. *github repository: "Monygham/Pedo-mellon-a-minno"*.
- [Monygham, 2020c] Monygham (2020c). Group schemes over field. *github repository: "Monygham/Pedo-mellon-a-minno"*.
- [Monygham, 2020d] Monygham (2020d). Monoid k -functors and their representations. *github repository: "Monygham/Pedo-mellon-a-minno"*.
- [Olsson, 2016] Olsson, M. (2016). *Algebraic Spaces and Stacks*. Colloquium Publications. American Mathematical Society.