

MAST20005/MAST90058: Week 9 Solutions

1. (a)

$$\mathbb{E}(\bar{X}_{..}) = \mathbb{E}\left(\frac{1}{I} \sum_{i=1}^I \bar{X}_{i.}\right) = \frac{1}{I} \sum_{i=1}^I \mu_i = \mu$$

(b)

$$\mathbb{E}(\bar{X}_{i.}^2) = \text{var}(\bar{X}_{i.}) + \mathbb{E}(\bar{X}_{i.})^2 = \frac{\sigma^2}{J} + \mu_i^2$$

(c)

$$\mathbb{E}(\bar{X}_{..}^2) = \text{var}(\bar{X}_{..}) + \mathbb{E}(\bar{X}_{..})^2 = \frac{\sigma^2}{IJ} + \mu^2$$

since we have

$$\begin{aligned} \text{var}(\bar{X}_{..}) &= \mathbb{E}((\bar{X}_{..} - \mu)^2) = \mathbb{E}\left(\left(I^{-1} \sum_{i=1}^I \bar{X}_{i.} - I^{-1} \sum_{i=1}^I \mu_i\right)^2\right) \\ &= \mathbb{E}\left[\left(I^{-1} \sum_{i=1}^I (\bar{X}_{i.} - \mu_i)\right)^2\right] = \frac{I\sigma^2}{I^2J} = \frac{\sigma^2}{IJ} \end{aligned}$$

(d)

$$\begin{aligned} \mathbb{E}(SS(T)) &= \mathbb{E}\left(\sum_{i=1}^I J(\bar{X}_{i.} - \bar{X}_{..})^2\right) = \mathbb{E}\left(\sum_{i=1}^I J\bar{X}_{i.}^2 + \sum_{i=1}^I J\bar{X}_{..}^2 - 2\sum_{i=1}^I J\bar{X}_{i.}\bar{X}_{..}\right) \\ &= \sum_{i=1}^I J\mathbb{E}(\bar{X}_{i.}^2) + IJ\mathbb{E}(\bar{X}_{..}^2) - 2IJ\mathbb{E}(\bar{X}_{..}) \\ &= \sum_{i=1}^I J\mathbb{E}(\bar{X}_{i.}^2) - IJ\mathbb{E}(\bar{X}_{..}^2) \\ &= \sum_{i=1}^I J\left(\frac{\sigma^2}{J} + \mu_i^2\right) - IJ\left(\frac{\sigma^2}{IJ} + \mu^2\right) \\ &= (I-1)\sigma^2 + J\sum_{i=1}^I (\mu_i^2 - \mu^2) \\ &= (I-1)\sigma^2 + J\sum_{i=1}^I (\mu_i - \mu)^2 \end{aligned}$$

as $\sum_{i=1}^I \mu_i \mu = I\mu^2$. Hence

$$\mathbb{E}(MS(T)) = \sigma^2 + \frac{J}{I-1} \sum_{i=1}^I (\mu_i - \mu)^2$$

as required. Note that we have used: $\sum_{i=1}^I (\mu_i - \mu)^2 = \sum_{i=1}^I (\mu_i^2 - 2\mu_i\mu + \mu^2) = \sum_{i=1}^I \mu_i^2 - 2I\mu^2 + I\mu^2 = \sum_{i=1}^I \mu_i^2 - I\mu^2 = \sum_{i=1}^I (\mu_i^2 - \mu^2)$.

(e) When H_0 is true, this is σ^2 , otherwise it is larger than σ^2 .

2.

$$F = \frac{MS(T)}{MS(E)} = \frac{2573.3}{1394.2} = 1.846$$

Under H_0 , $F \sim F_{4,15}$ so we reject H_0 if $F > 3.056$. Hence we cannot reject H_0 .

3. (a) This is the same as a test that the slope parameter is equal to zero (versus not zero). The test statistic for this is equal to $t = 0.75169/0.02846 = 26.41$. This is much larger than the critical value 2.66 (note: this is a two-sided test, so we need the 0.995 quantile of t_{60}). Thus, we reject the null hypothesis and conclude that there is strong evidence of an association between body and brain weights.
- (b) Under $H_0: \rho = 0$, we have the following

$$Z = \frac{\frac{1}{2} \ln \frac{1+R}{1-R}}{\sqrt{\frac{1}{n-3}}} \approx N(0, 1).$$

Therefore, we should reject H_0 if $|Z| > \Phi^{-1}(1 - \alpha/2)$.

- (c) From the R output we see that $r^2 = 0.9208$ (under **Multiple R-squared**), so we can calculate that $r = 0.9595$.
- (d) The observed value for the above test statistic is $z = 14.91 > 2.58$. Thus we reject the null hypothesis that there is no correlation between brain and body weight.
- (e) Both tests indicate strong evidence against the null hypothesis of no association between the body and brain weights of mammals.
4. (a) Let $A_0 = \{(\mu, \sigma^2) : -\infty < \mu < \infty, \sigma^2 = \sigma_0^2\}$ and $A_1 = \{(\mu, \sigma^2) : -\infty < \mu < \infty, \sigma^2 > 0\}$ be the set of parameter values consistent with H_0 and H_1 respectively. To maximise L under H_0 , we can show that we need $\hat{\mu} = \bar{x}$ and $\hat{\sigma}^2 = \sigma_0^2$, which gives:

$$L_0 = L(\hat{\mu}, \hat{\sigma}^2) = \left(\frac{1}{2\pi\sigma_0^2} \right)^{n/2} e^{-\frac{\sum_i (x_i - \bar{x})^2}{2\sigma_0^2}}.$$

To maximise L under H_1 , we can show that we need $\hat{\mu} = \bar{x}$ and $\hat{\sigma}^2 = n^{-1} \sum_i (x_i - \bar{x})^2$, which gives:

$$L_1 = L(\hat{\mu}, \hat{\sigma}^2) = \left[\frac{ne^{-1}}{2\pi \sum_i (x_i - \bar{x})^2} \right]^{n/2}.$$

- (b) The LRT statistic is,

$$\lambda = \frac{L_0}{L_1} = \left(\frac{w}{n} \right)^{n/2} e^{-w/2+n/2}$$

where $w = \sum_i (x_i - \bar{x})^2 / \sigma_0^2$. We reject H_0 if $\lambda \leq k$. Solving this inequality in w gives a solution of the form $w \leq c_1$ and $w \geq c_2$, where c_1 and c_2 are constants depending on k and n . These need to be selected so to achieve the desired significance level. (Note: since $w \sim \chi_{n-1}^2$, for convenience we typically set c_1 and c_2 as equal-tailed quantiles from this distribution, even if these do not exactly correspond to the c_1 and c_2 from the LRT.)