

ATLAS: machine-verified constructions of finite simple groups in Lean

Gerald Höhn

Department of Mathematics, Kansas State University
`gerald@monstrous-moonshine.de`

18 September 2026

Abstract

The ATLAS project constructs finite simple groups in Lean and proves their orders, simplicity, and structural properties. The source snapshot reported here covers eight family packages: prime cyclic groups, alternating groups, the classical types A, B, C, D , the split-octonion groups $G_2(q)$, and the small Ree groups ${}^2G_2(3^{2m+1})$ for $m \geq 1$. It also contains fifteen sporadic constructions, including all three Fischer simple groups. The models retain the codes, lattices, algebras, forms, and finite geometries from which the groups arise. The $G_2(q)$ model is the full automorphism group of the split octonions; the small Ree model is a seven-dimensional matrix group with a faithful doubly transitive action and an embedding into that octonion group. Comparison isomorphisms and equal-order nonisomorphism results relate the classical models. We describe the constructions, parameter ranges, dependencies, and recorded verification, and give per-group source-line dependency footprints. The project develops the construction side of the classification of finite simple groups, rather than its exhaustiveness proof, within a broader programme seeking geometric and vertex-algebraic explanations of sporadic symmetry.

Source and reproducibility. This article reports the source snapshot of 17 September 2026. The corresponding audited snapshot will be made public at <https://github.com/Moonshine-in-Kansas/atlas> after the final arXiv identifier has been assigned. The repository is not yet public at the time of this revision. Section 13 and Appendix A specify the source state, dependency pins, verification scope, and release procedure.

1 Introduction

The ATLAS project constructs finite simple groups in Lean [28], using Mathlib [24], and proves their orders, simplicity, and structural properties. Formalization results 3.1 and 3.2 summarize eight completed family packages and fifteen sporadic constructions in the reported snapshot. Their models retain the codes, lattices, forms, algebras, and geometries that produce the groups, together with actions and comparison maps for subsequent mathematics.

Classification has two complementary components: constructing the groups on the list and proving that every finite simple group occurs there. ATLAS develops the first component without assuming the exhaustiveness theorem of the classification of finite simple groups (CFSG). Its structural results and comparison maps are also intended as inputs to classification proofs. Section 2 explains the conditional recognition questions that connect the two components.

Three objectives guide the programme: geometric constructions of exceptional simple groups, formal proofs that preserve their mathematical structure, and an explanation of why

sporadic symmetries arise. The constructions combine established mathematics with the author’s source manuscripts; the contribution described here is their formal realization and the structural library it provides. Vertex operator algebras (VOAs) provide a principal direction for the third objective, illustrated by the public Monster work in Section 11. Geometric models also record more than a permutation action: the Fischer construction retains its algebra, rays, and incidence geometry, together with the arguments determining the group and its order.

The formalized constructions draw on the Hall–Janko manuscript [14] and the Conway–Parker algebra construction of the Fischer groups [13], alongside the classical literature cited below. The direct Monster-order work with Seysen [18] supplies a starting point for future formalization, not a completed entry in this release. Companion work on integral Albert algebras and the rank-26 lattice and Cayley-plane design develops the arithmetic and geometric direction [15, 17]. These source manuscripts and the present formalization have distinct publication and verification scopes.

The name recalls the *Atlas of Finite Groups* [5]. Our initial scope is constructions and structural theorems; complete character tables and maximal-subgroup lists are longer-term projects. Section 4 records the classical conventions and comparison results, and Sections 6–10 describe representative constructions. The author specifies the mathematics and directs AI-assisted implementation, as explained in Section 13. Appendix A identifies the source state and gives the complete table of proof-source footprints.

2 The mathematical content of an entry

For a family label X , let $\mathcal{A}_X$ be an explicitly specified parameter domain. An entry consists of a defined group $G_X(\lambda)$, for $\lambda \in \mathcal{A}_X$, and proofs of

$$|G_X(\lambda)| = N_X(\lambda), \quad G_X(\lambda) \text{ is simple}, \quad P_X(G_X(\lambda)), \quad (2.1)$$

together with finiteness and, for the nonabelian entries, explicit noncommutativity. Here P_X describes independently defined mathematical structure: an action, a code, a lattice, an algebra, or specified maps between such objects. Neither $\mathcal{A}_X$ nor P_X may incorporate the desired order or simplicity assertion as an unproved premise. For cyclic groups of prime order, commutativity is part of the intended conclusion.

Here *construction* refers to a specified mathematical model and proofs of its properties. The development uses classical mathematics in Lean, including classical choice; it need not provide an executable enumeration of every group element.

2.1 The chosen object and its full symmetry group

For a symmetry-group model, equality between a generated subgroup and the full automorphism group is a mathematical theorem. In a code or lattice construction, generation or a stabilizer argument establishes this equality before the order and simplicity assertions are attached to the full group.

Central quotients come with their maps and kernels. Thus Co_0 is distinguished from Co_1 , and a Hermitian isometry group from its scalar quotient. A different distinction is that between Fi_{24} and its derived subgroup Fi'_{24} . Abstract group isomorphisms, subgroup equalities, and equivariant identifications of actions are recorded with the corresponding data.

2.2 Conditional recognition and comparison

Conditional on CFSG, order determines a finite simple group up to isomorphism except for the pairs $\{\text{Alt}_8 \cong \text{PSL}_4(2), \text{PSL}_3(4)\}$ and $\{B_r(q), C_r(q)\}$ with $r \geq 3$ and q odd [20]. These pairs are distinguished by the number $k_2(G)$ of conjugacy classes of nonidentity involutions: the first has values 2 and 1, and the second r and $\lfloor r/2 \rfloor + 1$ [34, 4]. Consequently, assuming CFSG, the pair $(|G|, k_2(G))$ determines the isomorphism type of a finite simple group. The formal B/C counts and both nonisomorphism results are described in Section 4.4; the general recognition theorem and the $\text{Alt}_8/\text{PSL}_3(4)$ involution counts are proposed work.

For construction proofs, the relevant independence question concerns the hypotheses and dependency graph of each theorem. Background results and earlier constructions can be reused. A classification argument can also be non-circular when the property being deduced was not used in that argument, for example when it concerns smaller groups. The Monster order illustrates why precise attribution matters: Griess–Meierfrankenfeld–Segev, Corollary 3.7.3, uses substantial information about other sporadic groups [11], whereas Höhn–Seysen [18] gives a direct computational determination of the Monster and Baby Monster orders in the Leech-lattice setting. The former dependency is more specific than an invocation of the entire CFSG.

Recognition through an action or local geometry offers a natural interface: a classification argument produces the structure, and a comparison theorem identifies it with a constructed model. The actions, stabilizers, and maps retained here are intended to support such theorems.

3 Current constructions and audit scope

The catalogue records eight completed family packages and fifteen sporadic constructions [12]. In this article, *completed* means that the stated order, simplicity, and structural results have the recorded proofs and audits for the specified parameter domains; it does not mean that every property of the groups has been formalized. The verification scope is described in Section 13. Family labels count construction packages and can overlap up to isomorphism. Section 4 records the status of the comparison maps.

Formalization result 3.1 (Family packages). *The following family packages are completed. Here n is the degree of an alternating group or the matrix dimension of PSL_n ; r denotes Dynkin rank in the B, C, D families. For each finite field F below, put $q = |F|$.*

- (i) *For every prime p , the cyclic group C_p has order p and is simple and commutative.*
- (ii) *For every $n \geq 5$, the alternating group Alt_n is nonabelian simple and has order $n!/2$.*
- (iii) *Let F be any finite field, put $q = |F|$, and let $n \geq 2$. The central quotient $\text{PSL}_n(F) = \text{SL}_n(F)/Z(\text{SL}_n(F))$ has order*

$$|\text{PSL}_n(F)| = \frac{q^{n(n-1)/2} \prod_{i=2}^n (q^i - 1)}{\gcd(n, q - 1)}. \quad (3.1)$$

It is simple if and only if $(n, q) \notin \{(2, 2), (2, 3)\}$. For those admissible parameters it is also nonabelian.

- (iv) *For every finite field F and rank $r \geq 1$, both the projective quadratic model $B_r(F)$ and the symplectic quotient $C_r(F) = \text{PSp}_{2r}(F)$ have the order in (4.1). Each is simple exactly outside $(r, q) = (1, 2), (1, 3), (2, 2)$, and nonabelian at the simple parameters. The full nonsimple models at $(r, q) = (2, 2)$ and their derived subgroups are distinguished. The symplectic rank-zero model is trivial.*
- (v) *For every finite field F and $r \geq 4$, the projective split-orthogonal model $D_r(F)$ has the order in (4.2) with $\varepsilon = 1$ and is nonabelian simple in every characteristic, including*

$D_4(2)$.

- (vi) For every finite field F , with $q = |F|$, let $G_2(F) = \text{Aut}_F(\mathbb{O}_{\text{split}}(F))$ be the full group of multiplication-preserving linear automorphisms of the constructed split octonion algebra. Then

$$|G_2(F)| = q^6(q^6 - 1)(q^2 - 1). \quad (3.2)$$

It is simple if and only if $q > 2$, and noncommutativity is proved for every q . At $q = 2$, an explicitly defined normal subgroup has order 6048 and index two; its own simplicity and identification with $\text{PSU}_3(3)$ are not included in this result.

- (vii) Let F be a finite field of characteristic three with $|F| = q = 3^{2m+1}$, where $m \geq 1$. The group ${}^2G_2(F, m)$ generated by the Tits-twisted matrices in $\text{GL}_7(F)$ is nonabelian simple and has order

$$|{}^2G_2(F, m)| = q^3(q^3 + 1)(q - 1). \quad (3.3)$$

It acts faithfully and doubly transitively on a constructed set of $q^3 + 1$ points. The full point stabilizer is the soluble Borel subgroup of order $q^3(q - 1)$. Field transport and an injective homomorphism into the split-octonion group $G_2(F)$ are proved. The parameter $q = 3$ and a graph-automorphism fixed-point characterization are not part of this assertion.

The orthogonal packages retain intrinsic kernels, full centers, root generation, singular-line geometry, and field transport. Their domains include comparison models outside the primary label ranges in Table 2. For type A and G_2 , finite-field witnesses give existence at every admissible prime power; for the small Ree groups, a canonical field witness is supplied for every $m \geq 1$. The notation ${}^2G_2(F, m)$ follows the Lean model's explicit twist parameter; $|F| = 3^{2m+1}$ determines m uniquely.

Cyclic and alternating entries reuse Mathlib's principal order and simplicity results. The remaining packages combine general library theorems with the constructions described below. Mathieu parameters such as a dodecad or marked coordinates select objects in constructed finite structures; standard choices are supplied.

Formalization result 3.2 (Sporadic packages). *The development contains finite nonabelian simple groups with the constructions and orders in Table 1. These entries include the specified code, lattice, stabilizer, scalar-quotient, or algebraic interfaces.*

The auxiliary groups, notably Co_0 and the retained scalar covers, are not counted as additional simple groups. Nor are small isomorphisms between family members counted as new entries. The eleven sporadics outside that completed catalogue are $J_1, J_3, J_4, \text{He}, \text{Ru}, \text{O}'\text{N}, \text{HN}, \text{Ly}, \text{Th}, B$, and the Monster. The classical extension and the remaining uniform Lie-type programme are described next. The Tits group is a separate exceptional Lie-type entry, not an additional sporadic group.

4 Classical groups and comparison isomorphisms

4.1 Current classical development and notation

The untwisted families A, B, C, D have completed packages. The unitary 2A and minus-orthogonal 2D families and a common Chevalley framework are further construction tasks.

For rank r over a finite field F , the orthogonal constructions start from

$$Q_B(x, y, z) = \sum_{i=1}^r x_i y_i + z^2, \quad Q_D(x, y) = \sum_{i=1}^r x_i y_i.$$

Group	Order	Chosen construction
M_{11}	7 920	Point stabilizer in the dodecad model of M_{12}
M_{12}	95 040	Golay dodecad stabilizer
M_{22}	443 520	Pointwise stabilizer of two Golay coordinates
M_{23}	10 200 960	Stabilizer of one Golay coordinate
M_{24}	244 823 040	Full binary Golay code automorphism group
Co_1	4 157 776 806 543 360 000	Leech isometry group modulo $\{\pm I\}$
Co_2	42 305 421 312 000	Stabilizer of a Leech vector of squared norm 4
Co_3	495 766 656 000	Stabilizer of a Leech vector of squared norm 6
McL	898 128 000	Pointwise stabilizer of a Leech 2–2–3 triangle
HS	44 352 000	Pointwise stabilizer of a Leech 2–3–3 triangle
Suz	448 345 497 600	Full Eisenstein Hermitian group modulo six scalars
J_2	604 800	Full icosian Hermitian group modulo two signs
Fi'_{24}	1 255 205 709 190 661 721 292 800	Parity kernel in the constructed Fischer ray group
Fi_{23}	4 089 470 473 293 004 800	Centralizer of a marked involution modulo that involution
Fi_{22}	64 561 751 654 400	Common centralizer of two marked commuting involutions modulo their subgroup

Table 1: The fifteen completed sporadic constructions. A triangle label records half the three squared edge lengths. The groups are stabilizers of vectors or configurations, with the pointwise and setwise distinctions retained.

They construct the subgroups generated by Siegel transformations and divide by their scalar subgroups. In the completed parameter ranges, the elementary subgroups are identified in odd characteristic with the determinant/spinor kernels. In characteristic two, the B subgroup is the full quadratic isometry group and the split D subgroup is the intrinsic Dickson kernel. Their scalar subgroups are proved to be their full centers. The B geometric package has a faithful singular-line action in every positive rank, transitivity over finite fields, and primitivity from rank two; the split D package retains its faithful primitive singular-line action in rank at least four.

Throughout this section $q = p^f$ is a prime power, and r denotes *Dynkin rank*, not matrix dimension. We write Alt_m for an alternating group to distinguish it from the Lie-type symbol $A_r(q)$. The superscript in 2A_r denotes twisting; it is not a power of A_r . In twisted notation, q is the size of the fixed field: ${}^2A_r(q) = \text{PSU}_{r+1}(q)$ is constructed using a Hermitian space over $\mathbb{F}_{q^2}$. References writing ${}^2A_r(q^2)$ denote the same family with a different field convention. We use the fixed-field convention also for ${}^2D_r(q)$, ${}^2E_6(q)$, and ${}^3D_4(q)$.

Type	Classical model	Rank range	Exclusions for simplicity
$A_r(q)$	$\text{PSL}_{r+1}(q)$	$r \geq 1$	$(r, q) = (1, 2), (1, 3)$
$B_r(q)$	$\text{P}\Omega_{2r+1}(q)$	$r \geq 2$	$(r, q) = (2, 2)$
$C_r(q)$	$\text{PSp}_{2r}(q)$	$r \geq 3$	None in this rank range
$D_r(q)$	$\text{P}\Omega_{2r}^+(q)$	$r \geq 4$	None in this rank range
${}^2A_r(q)$	$\text{PSU}_{r+1}(q)$	$r \geq 2$	$(r, q) = (2, 2)$
${}^2D_r(q)$	$\text{P}\Omega_{2r}^-(q)$	$r \geq 4$	None in this rank range

Table 2: Classical families and proposed primary rank conventions, not the full implemented parameter domains. Types A, B, C, D have completed packages. The unitary and minus-orthogonal families remain to be constructed. The even-characteristic B/C comparison is proved for the retained models.

The symbols denote central quotients. For even q , type B_r is realized on a nonsingular

odd-dimensional quadratic space with one-dimensional polar radical; its symplectic comparison is proved. Exceptional small parameters require conventions for Ω as in Taylor [34, Chapters 8, 10–12].

The proposed primary ranges $r \geq 1, 2, 3, 4$ for A, B, C, D remove low-rank repetitions while keeping the comparison models in the library. In particular, B, C include all positive ranks, and D_2, D_3 are available. The odd-characteristic B_2 construction uses its symplectic C_2 image, and the geometric proof uses transport through a B_1 complement. Thus the maps $B_1 \cong A_1$ and $B_2 \cong C_2$ connect useful constructions. Even-characteristic B/C isomorphisms and isolated coincidences still give overlaps between primary labels.

For reference, the classical order formulae are

$$\begin{aligned} |A_r(q)| &= \frac{q^{r(r+1)/2}}{(r+1, q-1)} \prod_{i=2}^{r+1} (q^i - 1), \\ |{}^2A_r(q)| &= \frac{q^{r(r+1)/2}}{(r+1, q+1)} \prod_{i=2}^{r+1} (q^i - (-1)^i), \\ |B_r(q)| &= |C_r(q)| = \frac{q^{r^2}}{(2, q-1)} \prod_{i=1}^r (q^{2i} - 1), \end{aligned} \tag{4.1}$$

$$|D_r^\varepsilon(q)| = \frac{q^{r(r-1)}(q^r - \varepsilon)}{(4, q^r - \varepsilon)} \prod_{i=1}^{r-1} (q^{2i} - 1), \tag{4.2}$$

where $(a, b) = \gcd(a, b)$, $D_r^+ = D_r$, and $D_r^- = {}^2D_r$ [34, 37]. The A, B, C and split- D formulae are proved for the models in the ranges stated above; the unitary and minus- D formulae are targets for subsequent development.

4.2 Uniform coincidences and low ranks

Table 3 records the uniform identifications of central quotients, including auxiliary models below the primary rank ranges. Here 2A_1 abbreviates PSU_2 ; type A_1 has no nontrivial Dynkin-diagram twist.

The exterior-square representation produces the six-dimensional orthogonal model from a four-dimensional linear space. Related Klein-correspondence arguments give low-rank comparisons. In characteristic two, quotienting the polar radical of an odd-dimensional quadratic space gives a symplectic space. Over a finite field this has a converse lift [34, Chapters 11–12]. The completed untwisted comparisons implement these constructions.

4.3 Isolated coincidences across the classification list

Table 4 lists the isolated alternating and Lie-type coincidences, including simple derived groups at small parameters [5, 37, 38]. Its last column distinguishes proved maps from known isomorphisms awaiting formal comparison.

The binary orthogonal audit proves $B_2(2) \cong \text{Sym}_6$ and $B_2(2)' \cong \text{Alt}_6$ through `Atlas.Orthogonal.b2BinaryEquivSymmetric` and `Atlas.Orthogonal.b2BinaryDerivedEquivAlternating`. The full group has order 720 and its derived group order 360. The proved B_2/C_2 map identifies the full binary symplectic model, and its restriction to the derived subgroups is exported. The separate $\text{PSL}_2(9) \cong \text{Alt}_6$ comparison uses the projective line and the ten partitions of six letters into two triples. Composing these proved maps completes the order-360 row.

For the other small parameters, $G_2(2)'$ has index two in $G_2(2)$, and ${}^2G_2(3)'$ has index three in ${}^2G_2(3)$. The Tits group ${}^2F_4(2)'$ has index two in ${}^2F_4(2)$ but gives no further coincidence

Isomorphism	Parameters and formalization status
$A_1(q) \cong B_1(q) \cong C_1(q) \cong \text{PSU}_2(q)$	$B_1 \cong A_1$ and $C_1 \cong A_1$ proved over every field; the PSU_2 link remains planned. Simple for $q \geq 4$.
$B_2(q) \cong C_2(q)$	Proved over every finite field, including the nonsimple $q = 2$ case.
$D_2(q) \cong A_1(q) \times A_1(q)$	Proved over every field; the split rank-two model is not a simple entry.
$D_3(q) \cong A_3(q)$	Proved over every finite field for the split six-dimensional model.
${}^2D_2(q) \cong A_1(q^2)$	All q mathematically; four-dimensional minus-type comparison still planned.
${}^2D_3(q) \cong {}^2A_3(q)$	All q mathematically; six-dimensional minus-type comparison still planned.
$B_r(2^f) \cong C_r(2^f)$	Proved for $r \geq 1, f \geq 1$; primary-label overlaps persist for $r \geq 3$.

Table 3: Uniform identifications and their current proof status. For $D_2(2)$ we retain the central quotient of the two $\text{SL}_2(2)$ factors, not a possibly different derived-full-orthogonal convention. The table includes comparison models outside the primary rank ranges.

Order	Isomorphic simple groups	Status of comparison maps
60	$\text{Alt}_5 \cong A_1(4) \cong A_1(5)$	Both links with Alt_5 proved.
168	$A_1(7) \cong A_2(2)$	Proved through a common eight-point action.
360	$\text{Alt}_6 \cong A_1(9) \cong B_2(2)' \cong C_2(2)'$	All displayed links proved; the derived-group maps are explicitly exported.
504	$A_1(8) \cong {}^2G_2(3)'$	The $q = 3$ derived-subgroup package and this comparison remain to be formalized.
6 048	${}^2A_2(3) \cong G_2(2)'$	An index-two normal subgroup of $G_2(2)$ of order 6048 is constructed; its simplicity and the unitary comparison remain open in Lean.
20 160	$\text{Alt}_8 \cong A_3(2)$	Proved through the binary deleted permutation quadratic module.
25 920	${}^2A_3(2) \cong B_2(3) \cong C_2(3)$	$B_2 \cong C_2$ proved; the unitary link remains planned.

Table 4: Isolated cross-series coincidences and their formal status, up to the uniform identifications in Table 3. A prime denotes the derived subgroup. The status column records the available comparison maps.

in the table. The 26 sporadic groups supply no additional coincidences with each other or with the alternating and Lie-type simple groups [5, 37, 38]. Prime cyclic groups are separate; $\text{Alt}_3 \cong C_3$ lies outside the chosen alternating range. These tables do not classify nonsimple groups or central covers, and their exhaustiveness is not a Lean result here.

4.4 Equal orders and proved nonisomorphisms

The order collisions in Section 2 require *nonisomorphism* certificates, not additional comparison isomorphisms. Write $k_2(G)$ for the number of conjugacy classes of nonidentity involutions, not the number of involutions themselves. For every finite field of odd order and every positive rank, the formalized involution-class theorems give

$$k_2(B_r(q)) = r, \quad k_2(C_r(q)) = \lfloor r/2 \rfloor + 1. \quad (4.3)$$

Consequently, for $r \geq 3$ the constructed models satisfy

$$|B_r(q)| = |C_r(q)|, \quad B_r(q) \not\cong C_r(q).$$

The single exported theorem `Atlas.Comparisons.Classical.oddB_same_order_nonisomorphic_C` records both conclusions. Its assertion is the nonexistence of a `MulEquiv`, proved using the conjugacy-class invariance of k_2 .

Equal-order pair	k_2 values	Formal status
$B_r(q), C_r(q)$	$r, \lfloor r/2 \rfloor + 1$	Order equality and nonisomorphism proved for odd $q, r \geq 3$.
$\text{Alt}_8, \text{PSL}_3(4)$	2, 1	Nonisomorphism proved via Sylow 2-subgroup centers; these k_2 counts are not certified.

Table 5: The exceptional equal-order pairs. In the second row, the k_2 values are known mathematically, while the formal nonisomorphism uses a different invariant. Conditional on CFSG these two kinds exhaust the nonisomorphic simple-group order collisions [20, 34, 4].

The proved order-20 160 obstruction constructs the upper unitriangular subgroups in $\text{SL}_4(2)$ and $\text{SL}_3(4)$ and embeds them in the corresponding projective groups. Both images have order 64 and are Sylow 2-subgroups; symbolic commutation formulas show that their centers have orders 2 and 4, respectively. An isomorphism of the ambient groups would preserve this invariant. Transport through the independently proved $\text{PSL}_4(2) \cong \text{Alt}_8$ map gives the alternating-group obstruction. The optional involution-class counts remain separate: Alt_8 has the two even cycle types $2^2 1^4$ and 2^4 , whereas $\text{PSL}_3(4)$ has one involution class [34, Chapter 4]. Both nonisomorphism routes are unconditional, but neither a complete catalogue-wide collision theorem nor the conditional recognition theorem for $(|G|, k_2(G))$ is claimed as a Lean result.

4.5 Comparison maps and remaining targets

The comparisons marked as proved in Tables 3 and 4 are Lean `MulEquiv` objects: multiplication-preserving maps with proved inverses. The classical index exports `b1EquivA1`, `b2EquivC2`, `d2EquivProduct`, `d3EquivA3`, and `evenBEquivC` under `Atlas.Comparisons.Classical`. The symplectic rank-one map is `Atlas.Symplectic.rankOnePSL`. Field equivalences induce `bFieldEquiv` and `dFieldEquiv`.

The five isolated comparisons between the implemented families use natural actions or the deleted permutation quadratic module. Their order-60 argument also gives an elementary recognition theorem for nonabelian simple groups of that order. Unitary, twisted- D , and the

Representative modules (prefix <code>Mathlib.</code> omitted)	Mathematical input
<code>Data.Fintype.Card</code> <code>SetTheory.Cardinal.Finite</code> <code>GroupTheory.Index</code>	Finite types, cardinalities, subgroup indices, and counting identities.
<code>GroupTheory.QuotientGroup.Basic</code> <code>GroupTheory.Subgroup.Centralizer</code> <code>GroupTheory.Commutator.Basic</code>	Quotient groups, centralizers, commutators, and the maps used in central and derived-group constructions.
<code>GroupTheory.GroupAction.Primitive</code> <code>GroupTheory.GroupAction.MultipleTransitivity</code> <code>GroupTheory.GroupAction.Iwasawa</code>	Primitive and multiply transitive actions and the Iwasawa simplicity criterion.
<code>GroupTheory.SpecificGroups.Cyclic</code> <code>GroupTheory.SpecificGroups.Alternating</code> <code>GroupTheory.SpecificGroups.Alternating.Simple</code>	Existing cyclic and alternating group theory, including principal order and simplicity results.
<code>Data.ZMod.Basic</code> <code>FieldTheory.Finite.GaloisField</code>	Modular arithmetic and finite fields of prime-power order.
<code>LinearAlgebra.Matrix.GeneralLinearGroup.Card</code> <code>LinearAlgebra.Matrix.SpecialLinearGroup</code> <code>LinearAlgebra.Matrix.Transvection</code>	Matrix groups, general-linear-group cardinalities, and elementary matrices.
<code>LinearAlgebra.Matrix.ProjectiveSpecialLinearGroup</code> <code>LinearAlgebra.Projectivization.Action</code> <code>LinearAlgebra.Projectivization.PSL.PSL2</code>	Projective groups and actions, including the existing dimension-two simplicity theorem.
<code>LinearAlgebra.Basis.Basic</code> <code>LinearAlgebra.BilinearForm.DualLattice</code> <code>LinearAlgebra.QuadraticForm.Basic</code> <code>Algebra.Quaternion</code>	Bases, bilinear and quadratic forms, dual lattices, and quaternion arithmetic underlying the geometric constructions.

Table 6: Representative mathematical inputs from the pinned Mathlib dependency; they are not an exhaustive or minimal import list.

remaining exceptional links are further targets. Each comparison of actions or central covers needs its own maps, kernels, and equivariance statement; the group equivalences alone concern the simple quotients.

5 Lean, Mathlib, and the ATLAS library

Lean supplies dependent type theory, elaboration, and the proof kernel; Mathlib supplies mathematical definitions, theorems, and proof-producing automation. In particular, `Group`, `Nat.card`, and `IsSimpleGroup` are defined in `Mathlib.Algebra.Group.Defs`, `Mathlib.SetTheory.Cardinal.Finite`, and `Mathlib.GroupTheory.Subgroup.Simple`. Section 12 records their interpretation and the logical axioms used.

Table 6 lists representative mathematical inputs. Declaration dependencies identify the portion used by a particular proof; file imports make a broader interface available.

The cyclic entry uses `ZMod.card` and `isSimpleGroup_of_prime_card`; the alternating entry uses `nat_card_alternatingGroup` and `alternatingGroup.isSimpleGroup`. For PSL, the existing `rank_two_simple` theorem and `MulAction.IwasawaStructure.isSimpleGroup` support the general-rank arguments in Section 6. ATLAS develops the octonion multiplication and stabilizers, Golay–Leech structures, and Conway–Parker algebra on this foundation. Automation such as `ring` and `norm_num` produces proof terms checked by the kernel.

Shared mathematics can occur below public group interfaces. For example, McL and HS use `Atlas.Sporadic.Conway3.card`, while the Fischer entries share algebra and residue lemmas even when another Fischer group’s principal wrapper is absent from their dependencies. The source-line footprints in Appendix A count ATLAS declarations; their derivations also use Lean and Mathlib. The pinned environment is recorded there.

6 A uniform example: projective special linear groups

The projective linear family uses Mathlib’s matrix groups and central quotients, extending their theory by parameterized generation and simplicity arguments.

6.1 Order and the determinant quotient

The starting point is the standard count of ordered bases,

$$|\mathrm{GL}_n(q)| = \prod_{i=0}^{n-1} (q^n - q^i).$$

The scalar center of $\mathrm{GL}_n(q)$ has order $q - 1$. The center of $\mathrm{SL}_n(q)$ consists of the scalar matrices λI with $\lambda^n = 1$ and therefore has order $d = \gcd(n, q - 1)$, by cyclicity of $\mathbb{F}_q^\times$. These counts give (3.1). The development also proves a divisibility-safe identity

$$|\mathrm{PSL}_n(q)|(q - 1)d = \prod_{i=0}^{n-1} (q^n - q^i),$$

so that the natural-number division in the displayed order formula is known to be exact.

The determinant gives a stronger structural result. Over any field F and in positive dimension there is an isomorphism

$$\mathrm{PGL}_n(F)/\mathrm{PSL}_n(F) \cong F^\times / (F^\times)^n, \quad (6.1)$$

where $\mathrm{PSL}_n(F)$ on the left denotes its canonical image in $\mathrm{PGL}_n(F)$. Scaling a matrix changes its determinant by an n th power, so the map is well defined. If $\det(g) = u^n$, then $u^{-1}g$ has determinant one and represents the same projective transformation; this identifies the kernel. Surjectivity follows from that of the determinant. The formalized equivalence includes compatibility with matrix representatives. It is consequently usable independently of the finite order calculation.

6.2 Generation and simplicity

Elementary transvections are written $x_{ij}(a) = I + aE_{ij}$. The diagonal/transvection reduction available in Mathlib is completed by explicit two-coordinate identities, giving generation of $\mathrm{SL}_n(F)$ by transvections. For distinct i, j, k , the commutator identity

$$[x_{ik}(a), x_{kj}(b)] = x_{ij}(ab), \quad [g, h] = ghg^{-1}h^{-1}, \quad (6.2)$$

proves perfectness when $n \geq 3$. Generation and perfectness pass to the projective quotient.

For a line $L = Fv$, consider the subgroup of transformations

$$u_\varphi(x) = x + \varphi(x)v, \quad \varphi(v) = 0.$$

It is abelian, is normal in the stabilizer of L , and transforms covariantly under conjugation. Its projective images, as L varies, generate $\mathrm{PSL}_n(F)$. Together with the faithful primitive projective action, these facts fit the following criterion.

Proposition 6.1 (Iwasawa criterion). *Let a nontrivial perfect group G act faithfully and primitively on a set Ω with at least two points. Suppose that for some $\alpha \in \Omega$, the stabilizer G_α has an abelian normal subgroup A whose conjugates generate G . Then G is nonabelian simple.*

Proof. Let $1 \neq N \triangleleft G$. The N -orbits form a G -invariant partition. Faithfulness excludes the partition into singleton orbits for such N , and primitivity therefore makes N transitive. Hence $G = NG_\alpha$. In G/N , the image of every conjugate of A is consequently the image of A itself, because $A \triangleleft G_\alpha$. The conjugates generate G , so G/N is abelian. Perfectness implies $G/N = 1$, and $N = G$. A nontrivial perfect group cannot be abelian. $\square$

The existing formal criterion and projective-action infrastructure are reused. The added general-rank generation and perfectness arguments discharge its hypotheses. This proves simplicity for every field in dimension at least three. For finite fields in dimension two, the existing Mathlib theorem supplies simplicity when $q \geq 4$. The two exceptional cases are proved nonsimple using their projective actions and orders. The final family statement gives the exact exception set. Numerical examples follow by specialization.

7 The exceptional family $G_2(q)$ from split octonions

The first completed exceptional Lie-type family is constructed directly from the split octonion algebra. The formal model retains the algebra, its full linear automorphism group, and its singular-point geometry; see [37, Chapter 4] for the classical background. The uniform proof includes characteristics two and three.

7.1 The algebra and its full automorphism group

Write $\mathbb{O}_{\text{split}}(F) = \bigoplus_{i=1}^8 Fx_i$. The multiplication is the bilinear extension of the fixed integral split-octonion table used in the construction. In these coordinates, for $a = \sum a_i x_i$,

$$1 = x_4 + x_5, \quad \text{tr}(a) = a_4 + a_5, \quad N(a) = a_1 a_8 + a_2 a_7 + a_3 a_6 + a_4 a_5.$$

Conjugation is $\bar{a} = \text{tr}(a)1 - a$. The formal development proves both identity laws, bilinearity, the quadratic identity $a^2 - \text{tr}(a)a + N(a)1 = 0$, the conjugate-product identities, norm multiplicativity, and reversal of products under conjugation. These are polynomial identities over arbitrary commutative rings. The same algebra therefore applies in characteristic two.

The group $G_2(F)$ is the full subgroup of $\text{GL}(\mathbb{O}_{\text{split}}(F))$ preserving multiplication. It is represented by `Atlas.G2.Model`, using a generic construction of multiplication-preserving linear automorphisms that requires no associativity assumption. Preservation of the identity, trace, norm, and conjugation is proved from this definition. For example, surjectivity forces the identity to be fixed; for a nonscalar a , comparison of its quadratic identity before and after applying an automorphism forces the trace and norm coefficients to agree. Thus the invariants follow from preservation of multiplication. The natural eight-dimensional representation is faithful, and coefficientwise field isomorphisms transport the group and its action.

7.2 Explicit subgroups, full stabilizers, and order

Six root transformations are defined and verified directly from the multiplication table. Their additive laws and parameter injectivity are proved. Ordered products give a bijection $F^6 \rightarrow U$ with explicit multiplication and inverse formulas. The diagonal torus T is identified with $F^\times \times F^\times$, normalizes U , and intersects it trivially. Consequently,

$$|U| = q^6, \quad |T| = (q-1)^2, \quad |UT| = q^6(q-1)^2.$$

Together with explicit Weyl transformations, these maps are proved to generate the full octonion automorphism group.

The singular vectors are the nonzero a with $\text{tr}(a) = N(a) = 0$. Their number is $q^6 - 1$. Indeed, after eliminating one coordinate using the trace equation, the count reduces to pairs of three-vectors and a scalar satisfying $u \cdot v = t^2$: for each $u \neq 0$ and each t , the v -fibre has q^2 elements; for $u = 0$, one has $t = 0$ and q^3 choices of v . Including zero gives q^6 vectors. The projective singular-point set $\mathcal{P}$ therefore has size

$$|\mathcal{P}| = \frac{q^6 - 1}{q - 1}.$$

Coordinate reductions prove transitivity and determine the full stabilizers. The pointwise stabilizer of the opposite vectors x_1, x_8 is isomorphic to $\text{SL}_2(F)$; the stabilizer of x_1 has a parameter bijection with $F^\times \times \text{SL}_2(F)$. The full stabilizer P of the line Fx_1 is the explicit parabolic, with a bijection

$$F^\times \times F^5 \times \text{SL}_2(F) \longrightarrow P.$$

This bijection is a parameterization of the stabilizer. Since $|\text{SL}_2(F)| = q(q^2 - 1)$, orbit-stabilizer gives

$$|G_2(F)| = \frac{q^6 - 1}{q - 1} (q - 1) q^5 q(q^2 - 1),$$

which is (3.2). The order proof is independent of perfectness, simplicity, and the binary exception.

7.3 Singular-point geometry and simplicity

The action on $\mathcal{P}$ is faithful. Its point stabilizer has four orbits, with lengths

$$1, \quad q(q + 1), \quad q^3(q + 1), \quad q^5. \quad (7.1)$$

Coordinate reductions identify these strata as orbits. Their geometry also proves primitivity in every characteristic, including $q = 2$. The block-size argument leaves only the projective points of $W = \langle x_1, x_2, x_3 \rangle$ as a possible nontrivial proper block. But the linear span of products $W \cdot W$ is the distinguished line Fx_1 . Any automorphism stabilizing W must therefore fix that point, contradicting transitivity of a block stabilizer on a nontrivial block.

At Fx_1 , a pair of commuting root groups gives an abelian subgroup $R_0 \cong (F^2, +)$ of order q^2 , normal in the full point stabilizer. Transport defines such a subgroup at every singular point, with independence of the chosen transporter and covariance under $G_2(F)$. For $q > 2$, explicit root and Weyl relations show that their conjugates generate the full group. The argument chooses a field element distinct from 0 and 1 and applies also when $q = 3$. Torus commutators then give perfectness. The faithful primitive action and these local abelian groups satisfy Proposition 6.1, proving simplicity. Noncommutativity is established separately by two explicit root maps, even at $q = 2$. The positive simplicity proof does not use the numerical order of $G_2(F)$.

7.4 The binary exception and the family interface

For $F = \mathbb{F}_2$, let H be the normal closure of the long-root groups. The construction produces a second geometry: the two-dimensional singular subspaces on which all products vanish. There are exactly 63 such zero-product planes. Their completeness is checked by ordinary kernel reduction on 64×64 binary coordinate pairs; the group acting is still the full octonion automorphism group.

Two root involutions fix respectively 7 and 9 of these planes, so their permutation signs are $(-1)^{28} = 1$ and $(-1)^{27} = -1$. The resulting sign homomorphism is surjective and contains H in its kernel. Independently, primitivity of the singular-point action makes the nontrivial normal subgroup H transitive, and the explicit point-stabilizer generators give $[G_2(2) : H] \leq 2$. It follows that H is exactly the sign kernel. Using the already proved order,

$$|G_2(2)| = 12096, \quad [G_2(2) : H] = 2, \quad |H| = 6048.$$

Field transport gives the same conclusion for every field of cardinality two. This proves the negative case in the exact simplicity theorem. The present package does not prove $H = G_2(2)'$, simplicity of H , or its identification with $\text{PSU}_3(3)$; the conventional derived-group coincidence in Table 4 remains a separate comparison target.

The principal declarations in `Atlas.G2` are `card_Model`, `isSimple`, and `isSimple_iff`, with `exists_mul_ne_mul` for noncommutativity. The public `Atlas.typeG2_construction` package collects the results for the concrete model, and `Atlas.typeG2_prime_power` supplies finite fields for every prime power. The corresponding existence endpoints retain the concrete model as their witness. The evidence for order, positive simplicity, the binary exception, and the complete interface is separately scoped in Section 13.

8 Small Ree groups from twisted matrices

Let F be a finite field of characteristic three with $q = |F| = 3^{2m+1}$ and $m \geq 1$. The model `Atlas.ReeG2.ModelFm` is the subgroup of $\text{GL}_7(F)$ generated by three explicit root maps, a diagonal torus, and a Weyl involution. The coordinates follow Bäärnhielm's explicit matrix model [1, Section 3]; the geometric and simplicity arguments draw on Wilson's elementary constructions [36, 35]. Its order and simplicity are proved for this generated group, without an abstract Ree-group theorem or recognition by order.

8.1 Field twists and the root group

The twisting maps are the iterated Frobenius automorphisms

$$\theta(x) = x^{3^m}, \quad \sigma(x) = x^{3^{m+1}}.$$

The finite-field identity $x^{3^{2m+1}} = x$ gives $\theta\sigma = \sigma\theta = \text{id}$ and $\sigma^2(x) = x^3$. These identities, their compatibility with field equivalences, and the bound $q \geq 27$ are proved before the group arguments.

Write $S(a, b, c)$ for the product of the three root matrices in their specified order. Its entries recover a, b, c , and direct matrix multiplication gives

$$S(a, b, c)S(d, e, f) = S(a + d, b + e - a\theta(d)^3, c + f - db + a\theta(d)^3d - a^2\theta(d)^3).$$

Together with the inverse formula, this identifies the root subgroup U with the parameter set F^3 , so $|U| = q^3$. Projection to the first parameter has abelian image and abelian kernel, and hence U is soluble. The diagonal subgroup $H \cong F^\times$ normalizes U , with conjugation on parameters

$$(a, b, c) \mapsto \left(\frac{a\theta(\lambda)^3}{\lambda^2}, \frac{b\lambda}{\theta(\lambda)^3}, \frac{c}{\lambda} \right).$$

Unique factorization in $B = UH$, the equality $U \cap H = 1$, and the split projection $B \rightarrow H$ with kernel U prove that B is soluble and has order $q^3(q - 1)$.

8.2 The point geometry and the full stabilizer

The Ree–Tits point set consists of a point at infinity and the projective classes of the first rows of the matrices $S(a, b, c)$. Projective parameter injectivity gives $q^3 + 1$ points, and U acts regularly on the affine points. The matrix convention uses row vectors; inverse right multiplication supplies the corresponding left action.

The main geometric step is to prove preservation by the Weyl element and to identify the full stabilizer of infinity. Seven exterior-kernel relations and seven Tits-twisted bilinear equations characterize the point chart. Their invariance under negative coordinate reversal proves Weyl preservation. They also imply that the last affine coordinate vanishes only at the affine origin. The generated group is consequently doubly transitive.

Every generator preserves an explicit alternating cross product and the twisted bilinear relation. At infinity the relation recovers a distinguished two-dimensional subspace, and the cross product recovers the full flag. A matrix fixing infinity must therefore be triangular. Applying the opposite flag at the affine origin shows that the two-point stabilizer is diagonal; the invariant tensors force it to be exactly H . Regularity of U then gives the full one-point stabilizer B . This proves faithfulness as well as the order formula (3.3) by orbit–stabilizer. The stabilizer calculation completes the order proof.

8.3 Perfectness, simplicity, and comparison with octonions

For $q \geq 27$, torus commutators place all three root maps in the derived subgroup. Short characteristic-three matrix identities then place the Weyl involution and the minus-one torus element there. Weyl inversion supplies all torus squares. Since $q \equiv 3 \pmod{4}$, every nonzero scalar is a square or the negative of a square, so the whole torus lies in the derived subgroup. The group is therefore perfect.

Simplicity uses a separately proved general criterion: a nontrivial perfect group with a faithful quasiprimitive action and a soluble point stabilizer is simple. Indeed, any nontrivial normal subgroup N is transitive, so the stabilizer surjects onto G/N . This quotient is both soluble and perfect and is therefore trivial. The Ree action is primitive, and its full stabilizer is the soluble subgroup B . Explicit noncommuting root elements supply noncommutativity. This simplicity argument does not use the numerical group-order theorem.

Entrywise field equivalence induces a group isomorphism of the models. A separate comparison embeds the group into the split-octonion $G_2(F)$ of Section 7. In the fixed seven-coordinate basis, the scalar–vector identification with the eight-coordinate octonions is

$$(t, v) \longmapsto (v_0, v_1, v_2, t - v_3, t + v_3, -v_4, -v_5, -v_6).$$

In characteristic three the octonion product becomes

$$(s, v)(t, w) = (st - B(v, w), sw + tv + v \times w),$$

where B is the preserved symmetric form and $\times$ the preserved cross product. Extending inverse row multiplication while fixing the scalar coordinate thus gives an injective homomorphism into the full octonion automorphism group. This uses structural octonion results, not the order or simplicity of $G_2(F)$. No equality with the fixed points of an exceptional graph automorphism is asserted.

The principal declarations are `Atlas.ReeG2.order` and `Atlas.ReeG2.simple`. The public package `Atlas.typeReeG2_construction` and existence theorem `Atlas.exists_typeReeG2_parameter` use the same model. Specializations of the uniform order theorem at $q = 27, 243, 2187$ give, respectively,

$$10\,073\,444\,472, \quad 49\,825\,657\,439\,340\,552, \quad 239\,189\,910\,264\,352\,349\,332\,632.$$

These values specialize the uniform theorem to constructed finite fields. The public family excludes $q = 3$; the derived-group comparison with $\mathrm{PSL}_2(8)$ in Table 4 remains a separate task.

9 Sporadic groups from retained structures

The sporadic branch follows connected constructions. The code used to construct the Mathieu groups is retained in the Leech lattice, and the resulting lattice and its full symmetry group are retained in later stabilizer and scalar-centralizer constructions. Standard background for these relationships is provided by [6]; the formal models and recorded completion results are those of the ATLAS development [12].

9.1 Trio code, hexacode, and Golay code

The initial construction passes from a quadratic alphabet and its trio code to the additive Kleinian hexacode, and then to the binary Golay code by two explicit twists. The coordinate organization

$$3 \longrightarrow 3 \times 2 \longrightarrow (3 \times 2) \times 4$$

is retained along with markings, partitions, and recovery maps. The quadratic functions on the intermediate alphabets are part of the data. Any comparison with an $\mathbb{F}_4$ -linear hexacode is a separate mathematical statement.

The constructed binary code C is a doubly even self-dual $[24, 12, 8]$ code. Its weight-eight supports give the Witt design $S(5, 8, 24)$. The chosen model for M_{24} is the full coordinate automorphism group of this code, and equality with the automorphism group of its octad design is proved in the same ambient symmetric group. Sextet geometry supplies local structure: every tetrad has a unique sextet completion, there are 1771 sextets, and the full sextet stabilizer has order 138240. The subsequent global action arguments give order 244823040, five-transitivity, and simplicity.

The smaller Mathieu groups come from stabilizers. Fixing one or two coordinates gives M_{23} and M_{22} ; stabilizing a dodecad and then one of its points gives M_{12} and M_{11} . Their actions on complements or dodecads, the corresponding residual designs, and the embeddings relating these choices are part of the construction. In particular, a pointwise two-coordinate stabilizer is kept distinct from the larger setwise stabilizer.

9.2 The Leech lattice and Conway groups

The retained Golay code defines an integral lattice Λ . Its rank 24, evenness, positive definiteness, unimodularity, minimum squared norm 4, and generation by minimal vectors are proved from the construction. The group Co_0 is its full linear isometry group, and

$$\mathrm{Co}_1 = \mathrm{Co}_0 / \langle -I \rangle.$$

Finiteness and the relevant kernels are proved for these groups before the final simplicity arguments.

The shell counts for squared norms 4, 6, 8 are respectively

$$196560, \quad 16773120, \quad 398034000.$$

They arise from exhaustive coordinate descriptions of lattice vectors. The global order proof uses minimal vectors and orthogonal pairs. An explicitly constructed nonmonomial isometry,

together with the full monomial subgroup, generates Co_0 . The resulting order is

$$|\text{Co}_0| = 8\,315\,553\,613\,086\,720\,000.$$

The action on the 8292375 intrinsic crosses, and its primitivity, follow from the minimal-vector and orthogonal-pair analysis.

Vector stabilizers produce Co_2 and Co_3 . Pointwise triangle stabilizers produce McL and HS . These models retain their embeddings in the lattice isometry group. Additional geometry includes the 276-point action of Co_3 and the rank-three 100-point graph action of HS . The lattice representations and stabilizer descriptions allow these groups to be reused without reconstructing unrelated permutation models.

9.3 Eisenstein and icosian structures

The constructions of Suz and J_2 enrich the same Leech lattice with additional scalar structure. They illustrate why integral comparison maps matter: a rational isometry between two ambient spaces must be shown to map the lattices onto one another. The documented comparisons prove both integral containments.

For Suz , a constructed fixed-point-free isometry ρ satisfies

$$\rho^3 = I, \quad \rho^2 + \rho + I = 0.$$

Its full centralizer $C_E = C_{\text{Co}_0}(\rho)$ is identified with the full Hermitian isometry group of the corresponding Eisenstein lattice. Its center consists of six scalar isometries. The constructed simple quotient has order

$$|C_E|/6 = 2\,690\,072\,985\,600/6 = 448\,345\,497\,600.$$

The action on 232960 intrinsic frames and the thirteen suborbits of a frame stabilizer are proved from the lattice. The sign quotient $C_E/\{\pm I\}$ embeds into Co_1 and maps onto Suz with central kernel of order three. This keeps the embedded triple extension distinct from its simple quotient.

For J_2 , let $\mathcal{I}$ be the constructed icosian order over $\mathbb{Z}[(1 + \sqrt{5})/2]$, and let C_H be the full quaternion-linear Hermitian isometry group of the associated rank-three congruence lattice. The underlying integral lattice is identified with the retained Leech lattice. The 120 norm-one icosian units act by right scalars, and C_H is their full centralizer in Co_0 . The scalar action and the center of C_H are separate objects.

There are 37800 vectors of quaternionic Hermitian norm two, and each associated right quaternionic line contains exactly 120 of them. Thus there are 315 root lines. Each is orthogonal to ten others, and every orthogonal pair has a unique completing root line. This gives an intrinsic count of frames,

$$\frac{315 \cdot 10}{6} = 525.$$

The full coordinate-frame stabilizer has order 2304. Explicit quaternionic reflections generate the full group and prove frame transitivity, whence

$$|C_H| = 525 \cdot 2304 = 1209600, \quad |C_H/\{\pm I\}| = 604800. \quad (9.1)$$

The point stabilizer has suborbits of sizes 1, 10, 32, 32, 80, 160. Their proved geometry yields primitivity. Reflection relations yield perfectness, and the cyclic subgroup generated by a point reflection supplies the abelian normal subgroup in Proposition 6.1. The sign quotient is therefore nonabelian simple. This example exhibits the chain from arithmetic and local geometry to a full symmetry group, its order, and its simplicity.

10 The Fischer constructions

The development contains concrete models and proofs of order, simplicity, finiteness, and noncommutativity for Fi'_{24} , Fi_{23} , and Fi_{22} . Their construction packages also retain the algebra, reflecting rays, and residue geometry described below.

10.1 The formal models and their properties

The formal starting point is the explicit Conway–Parker coordinate algebra constructed from Golay and Parker data [13]. The algebra has dimension 783 over $E = \mathbb{Q}(\omega)$, where ω is a primitive cube root of unity; its coordinate index is the disjoint union of the 24 Golay coordinates and the 759 octads. A proved tensor identity supplies the reflecting automorphisms. Its normalized μ_3 -rays are three-element scalar orbits. The declaration `Atlas.Fischer.rootGeneratedRayGroup` defines the subgroup of permutations of the displayed reflecting rays generated by the constructed root involutions. Write F for this group. The simple model Fi'_{24} is the kernel of its parity homomorphism; the exported `derived` and `index` theorems identify that kernel with F' and prove $[F : F'] = 2$.

For marked commuting basic involutions t_i , the residue construction uses the centralizer of the marked set and divides by the subgroup it generates. In particular, the two fixed models are

$$G_{23} = C_F(t_i)/\langle t_i \rangle, \quad G_{22} = C_F(t_i, t_j)/\langle t_i, t_j \rangle \quad (i \neq j).$$

The chosen marked involutions are part of the construction. The quotient kernels have orders two and four, respectively. Both residue models are therefore specified group quotients.

For each Fischer entry, the public declarations `card`, `simple`, `finite`, and `exists_mul_ne_mul` establish the corresponding basic assertions. The expanded signatures contain no supplied order or simplicity hypothesis. The namespaces under `Atlas.Sporadic` are `Fischer24Prime`, `Fischer23`, and `Fischer22`. Construction and existence interfaces accompany these results.

The full ray group F has a primitive rank-three action with subdegrees 1, 31671, 275264. For its parity kernel Fi'_{24} , the proved action on the same 306936 rays is faithful and transitive; the recorded result does not include a rank-three assertion for this subgroup. The residue groups G_{23} and G_{22} have faithful primitive rank-three actions with point counts and subdegrees

	degree	subdegrees		
G_{23}	31671	1	3510	28160
G_{22}	3510	1	693	2816

The residue projections, their surjectivity, generation by distinguished involutions, and perfectness are also retained. These give structural interfaces for the residue groups.

The completion record also retains the nonsplit central double cover of G_{22} inside G_{23} and the perfect central triple cover of Fi'_{24} acting faithfully on the 783-dimensional E -algebra. The quotient's action is projective; no ordinary representation of the same degree is inferred for it. In the full semilinear group, odd symmetries invert the cubic scalars, so that scalar kernel is not central. These extension statements have their own statement and dependency certificates.

10.2 Ray counts and group orders

The finite-algebra construction produces three families of rays of sizes

$$24, \quad 759 \cdot 2^5, \quad \binom{24}{2} 2^{10},$$

whose sum is 306936. Frame geometry yields the largest group, and commuting residues yield the two smaller ones. The notation distinguishes the three-transposition group Fi_{24} from its derived subgroup Fi'_{24} , the sporadic simple group. A central triple cover, when retained at the algebra level, is a further distinct group. The factorizations of the orders proved for the formal models are

$$\begin{aligned} |\text{Fi}_{22}| &= 2^{17} 3^9 5^2 7 11 13, \\ |\text{Fi}_{23}| &= 2^{18} 3^{13} 5^2 7 11 13 17 23, \\ |\text{Fi}'_{24}| &= 2^{21} 3^{16} 5^2 7^3 11 13 17 23 29. \end{aligned}$$

The factor of two between $|\text{Fi}_{24}|$ and $|\text{Fi}'_{24}|$ is part of the derived-subgroup analysis.

11 The Monster and vertex operator algebras

Höhn–Seysen [18] determines the Monster and Baby Monster orders directly from computations with axes in the 196884-dimensional Griess algebra, using Seysen’s `mmgroup` implementation [31]. The Monster is generated by a subgroup of shape $2_+^{1+24}.\text{Co}_1$ and a triality element. Orbit and stabilizer calculations use the Golay code, Leech lattice, and Conway groups.

VOA arguments complement these calculations. The axes correspond, after normalization, to Ising vectors in the Moonshine module $V^\natural$. Orbifold and fusion arguments control involutions and their centralizers, while the framed VOA structure identifies $\text{Aut}(V^\natural)$ with the automorphisms of the Griess algebra preserving its invariant bilinear form. Combining the computed order with results of Carnahan and Borcherds, the paper proves that the constructed Monster is this full automorphism group and has exactly two conjugacy classes of involutions [18, Sections 2 and 5]. These are public mathematical results supporting future formalization; the Monster and Baby Monster are among the outstanding ATLAS entries.

This work continues the Moonshine programme of Borcherds [2] and Frenkel–Lepowsky–Meurman [7]. Borcherds’s proof of the Monstrous Moonshine conjectures [3] is a longer-term formalization goal, not a result of the present development. The Baby Monster vertex operator superalgebra provides a related source of sporadic symmetry [16]. Further examples include the 10-dimensional quadratic space over $\mathbb{F}_2$ recovered from the modules of $V_{\sqrt{2}E_8}^+$ in Shimakura’s treatment, with automorphism group $O_{10}^+(2)$ [32, Theorem 4.5], and Griess–Lam’s VOA existence proof of the Monster [10].

12 Formal statements and logical foundations

For a constructed carrier G , `Group G` supplies operations and proofs of the group laws. Subgroups and quotients obtain these from their ambient groups and proved normality. On finite types, `Nat.card G` is the number of elements; it is zero on infinite types. Finiteness is also exported as `Finite G`. `IsSimpleGroup G` asserts nontriviality and that every normal subgroup is trivial or the whole group. For the nonabelian entries, noncommutativity is proved separately.

For example, the fixed model `Atlas.Sporadic.Fischer22.Model` is the centralizer quotient G_{22} of Section 10. Its public statements are

```
finite :      Finite G22,
card :      Nat.card G22 = 64561751654400,
simple :      IsSimpleGroup G22,
exists_mul_ne_mul : ∃x, y ∈ G22, xy ≠ yx.
```

All four declarations lie in `Atlas.Sporadic.Fischer22`; the display suppresses printer-level universe and typeclass arguments.

The recorded dependency audits report only `propext` (propositional extensionality), `Classical.choice`, and `Quot.sound` (equality of related quotient representatives). They exclude `sorryAx`, project-specific mathematical axioms, and native-evaluator trust shortcuts. The assurance rests on Lean’s logical rules and a correct implementation of its kernel. The axiom report must be read together with the full theorem type and the definitions of its carrier, parameters, and instances: an axiom check alone does not establish that the formal statement expresses the intended mathematical claim [26].

13 AI-assisted development and verification

13.1 Mathematical specification and implementation

The author specifies the objects, theorems, proof architecture, and permitted dependencies. Codex, OpenAI’s coding agent, searches the pinned library and implements and repairs proofs against Lean’s elaborator and kernel. The author reports using ChatGPT, including GPT-6 Pro, for mathematical discussion and specifications, and Codex with GPT-6 Astra for Lean implementation. These names identify the reported development tools; they are not a controlled comparison of models or a measurement of autonomous performance.

The author directs and maintains ATLAS and is responsible for its mathematical claims and formal specifications. Mathematical sources and joint work are credited where used. Compiled statements and dependency reports support review of the results; reproducing the AI’s search is unnecessary for checking the resulting proof terms.

13.2 Recorded verification

Verification has three distinct aspects: the mathematical specification, the derivation of the formal statement, and the implementation of the checker. The recorded runs use the pinned Lean toolchain, incremental Lake builds, and inspection of compiled declarations. The recorded build of the affected root and audit targets reports 5 677 jobs. The full dependency audits traverse types, bodies, and inductive metadata and enforce the exclusions specified by each checkpoint.

The catalogue check validates 886 declaration references, including aliases and structural exports. The principal order-and-simplicity selection has 46 roots for 23 entries and a complete graph of 65 676 dependency nodes, with project source ranges recovered throughout. These are different scopes. Table 7 gives the corresponding source footprints; family, geometric, and comparison certificates record their own selections.

In particular, the G_2 and small Ree order audits exclude their perfectness and simplicity arguments. Their positive simplicity audits exclude the numerical group-order theorems. The Ree proofs also exclude untwisted G_2 results; its octonion embedding uses a separate structural scope. The source evidence records these boundaries.

The recorded builds retain cached dependency artifacts. A fresh source rebuild and an independent proof recheck remain outstanding for this snapshot. These are different checks: rebuilding tests the source and dependency setup, whereas a separate recheck reduces reliance on the original elaboration and compiled artifacts [26, 27]. Neither replaces review of the formal statements. Finite tables checked by ordinary kernel reduction are part of some proofs. The release’s `VERIFICATION.md` will collect the audit selections, commands, and limitations; any later checks will be dated and tied to the source snapshot they actually cover.

13.3 Public repository and reproducible release

The designated repository is <https://github.com/Moonshine-in-Kansas/atlas>. It will be made public only after the final arXiv identifier has been assigned. The planned tag `atlas-arxiv-v1` will identify the frozen source snapshot corresponding to the first arXiv version, together with its dependency pins, catalogue, verification records, Apache-2.0 license, and third-party notices. Continuing and unfinished development will remain separate from this published snapshot.

The release README and CITATION.cff will record the arXiv identifier, public source commit, and release tag. A version-specific archived software DOI will be added when the archive is published; no such identifier is assigned here. The source manifest will identify the relationship between the measured working tree in Appendix A and the public release. Changes to proof sources after that measurement require a new audit rather than an undocumented change to the snapshot. Publication metadata and any additional verification records can be added with their own dates.

14 Related formalization projects

The Odd Order project of Gonthier and collaborators formalized the Feit–Thompson theorem in Coq (now Rocq), using Mathematical Components [9]. Completed in 2012 and described in 2013, it proves that finite groups of odd order are soluble and supplies extensive reusable finite-group and representation theory. Mathlib supplies the cyclic, alternating, and rank-two PSL results used here, together with group-action and simplicity infrastructure [24, 25].

The Qiuzhen-CFSG project and FormaTheoria workflow report a Lean development whose public README lists the Odd Order and Bender–Suzuki theorems as completed [29]. Their reconstruction of interdependent theories from the literature complements the construction work here. This account follows the authors’ public description and does not report an independent rebuild of their project.

Two Utrecht theses from 2024 precede this work: Roxy van de Kuilen’s partial formalization of the first Janko group [22] and Erik van der Plas’s study of the Conway groups in Lean [30]. Their precise declaration interfaces merit comparison with the constructions presented here.

The public repository `KitaKen1/finite-simple-groups-lean` reports kernel-checked order and simplicity certificates for fourteen sporadic permutation groups [21]:

$$M_{11}, M_{12}, M_{22}, M_{23}, M_{24}, \text{HS}, J_2, J_1, \text{McL}, \text{Co}_3, \text{Suz}, \text{Co}_2, \text{Fi}_{22}, \text{He}.$$

Its README also lists nine individual Lie-type groups, including $G_2(4)$ and $\text{PSL}_3(4)$. External programs generate candidate certificates whose required properties are checked in Lean. This description follows the public repository consulted on 18 September 2026, without an independent audit of its proofs.

Parameterized families require constructions and proofs for all admissible ranks and fields. Root-group or geometric arguments provide one route; a uniformly verified algorithm could provide another. For individual groups, comparison isomorphisms could connect the permutation models with the code, lattice, and algebra models here, allowing their structural results to be combined.

15 Further development

The remaining eleven sporadic groups in Section 3 call for geometric constructions or verified computational arguments. The Hall–Janko, Fischer, Albert-algebra, and Cayley-plane source

manuscripts develop the geometric direction. For O’Nan’s group, Lee’s 2025 account records the absence at that time of a computer-free construction [23]; Jansen–Wilson give explicit modular constructions [19]. Formalizing such arguments requires proofs of the algorithms and checks of the finite data they use.

A common Lie-type framework should construct groups from root data, integral forms, and root operators, then establish generation, Bruhat decomposition, centers, orders, and simplicity. Steinberg [33] and Geck [8] provide sources. Beyond the completed families, the targets are the unitary and minus-orthogonal families, F_4, E_6, E_7, E_8 , and the graph–field twists 2E_6 and 3D_4 . The Suzuki and large Ree families

$${}^2B_2(2^{2e+1}), \quad {}^2F_4(2^{2e+1}) \quad (e \geq 1)$$

require exceptional characteristic-two endomorphisms. Their initial nonsimple parameters and the Tits group ${}^2F_4(2)'$ need separate statements [37]. Comparison with the existing G_2 and small Ree models, including the exceptional fixed-point description of the latter, is part of this programme.

For integration with classification arguments, priorities include centralizers of prime-order elements, Sylow and local subgroups, outer automorphisms, central extensions, and recognition geometries. The comparison maps of Section 4 allow these results to pass between the models encountered in different proofs.

Formalizing the Monster and Baby Monster results discussed in Section 11 requires both verified finite computations and infrastructure for graded spaces, vertex operations, lattice extensions, and the relevant subalgebras or commutants. These developments would connect the finite-group constructions to their vertex-algebraic sources.

A Source state and reproducibility

The measurements in this appendix refer to the working-tree source state recorded at 17 September 2026, 07:34:33 UTC, based on commit `ec06f2a7a6411d2a25fe1dddc5584e40f1a9da5` [12]. The accompanying manifest identifies the source files, their hashes and tracked status, the catalogue, and the dependency pins. It is part of the source identification: the base commit alone does not describe uncommitted material. The public release will preserve this binding and record its own commit and tag. The editorial revision of this article does not update these measurements.

A.1 Source inventory

The tracked physical inventory contains 2 295 Lean files under `Atlas/`, with 161 772 lines, and the root driver `Atlas.lean`, with 706 lines. The total is therefore **162 478 lines in 2 296 files**. This includes audit and helper modules, comments, and blank lines; it excludes untracked drafts, Mathlib, and verification-generated Lean files. The small Ree-specific modules, including finite-field support and the public package, account for 54 files and 3 470 lines, without shared generic infrastructure.

The physical inventory and file hashes are recorded in `SOURCE_SIZE.csv` and `SNAPSHOT.json` under `verification/status-audit/2026-09-17-complete-line-counts/`. These file sizes are distinct from the declaration-range footprints in Table 7, which count only source lines used by the selected principal results.

A.2 Build and audit records

The environment is pinned to Lean `leanprover/lean4:v4.34.0-rc2` and Mathlib revision `85e3a25e006c35636f0e53b0e9296caca2685bc0`. From the source root, with that toolchain installed, run

```
lake exe cache get
LEAN_NUM_THREADS=2 lake build
ruby docs/catalogue/manage.rb check
```

The default targets are `Atlas` and `Atlas.Audit`. The first command fetches compiled dependencies; these commands describe the cached-dependency build, not a fresh source rebuild of the entire dependency stack. Preserve `lean-toolchain`, `lakefile.toml`, and `lake-manifest.json`. The catalogue in `docs/catalogue/catalogue.json` maps claims to declarations and evidence. `verification/WORKFLOWS.md` and the companion `VERIFICATION.md` describe the full audits.

A.3 Per-group declaration footprints

Table 7 measures the source needed by the selected order and simplicity declarations for each completed entry. The global deduplicated ATLAS footprint is **88 856 lines**. This measures source dependencies, not proof-term size, model-token usage, or the amount of new mathematics.

Each column counts physical source lines in compiler-recorded declaration ranges reached through the selected theorem’s transitive type and proof dependencies, including interior comments and blank lines. Generated helpers use the recorded enclosing declaration’s range; a final position at column zero excludes that line. Ranges are merged by file, so column and row totals overlap. Lean core and external libraries are excluded.

For $\mathrm{PSL}_n(q)$, n is matrix dimension; r is rank for B, C, D . The type- A , symplectic, type- B , and G_2 simplicity roots are exact-range equivalences including their exceptional cases. The Ree root applies for $q = 3^{2m+1}$, $m \geq 1$. Isomorphic construction families have separate rows.

The measurement directory provides `CODE_COUNTS.csv`, `roots.json`, the extracted dependency graph, merged ranges, helper mappings, and source manifest. These reproduce the table arithmetically. The 46-root audit has 65 676 nodes, complete graph references, and resolved source ranges for every project dependency. Further root-level details are collected in `VERIFICATION.md`.

Group or family	Order footprint	Simplicity footprint	Union
C_p	5	12	12
Alt_n	4	4	7
$\text{PSL}_n(q)$	168	322	379
$B_r(q)$	4 121	7 898	8 009
$C_r(q)$	708	2 203	2 203
$D_r(q)$	4 290	2 466	5 408
$G_2(q)$	1 237	3 287	3 568
${}^2G_2(q)$	1 570	2 225	2 250
M_{11}	5 479	5 866	5 868
M_{12}	4 924	5 959	5 961
M_{22}	4 183	5 348	5 351
M_{23}	4 119	4 492	4 495
M_{24}	3 477	4 582	4 585
Co_1	10 826	12 334	12 346
Co_2	10 825	14 070	14 072
Co_3	12 137	16 423	16 423
McL	13 704	16 375	16 377
HS	13 447	14 771	14 776
Suz	19 727	21 320	21 374
J_2	6 989	7 954	8 184
Fi'_{24}	31 813	33 396	33 397
Fi_{23}	31 950	34 247	34 249
Fi_{22}	31 955	34 248	34 250
Global deduplicated union			88 856

Table 7: Compiler-derived ATLAS source-line dependency footprints for 46 principal order and simplicity declarations: eight families and fifteen sporadic constructions. Shared source lines are counted once within each union; different rows need not have disjoint dependencies.

References

- [1] H. B  arnhielm, Recognising the small Ree groups in their natural representations, *Journal of Algebra* **416** (2014), 139–166. doi:10.1016/j.jalgebra.2014.06.017; arXiv:1206.0411.
- [2] R. E. Borcherds, Vertex algebras, Kac–Moody algebras, and the Monster, *Proceedings of the National Academy of Sciences USA* **83** (1986), 3068–3071. doi:10.1073/pnas.83.10.3068.
- [3] R. E. Borcherds, Monstrous moonshine and monstrous Lie superalgebras, *Inventiones Mathematicae* **109** (1992), 405–444. doi:10.1007/BF01232032.
- [4] A. V. Borovik, *Orthogonal and symplectic black box groups, revisited*, 2001, Sections 2.5–2.6. arXiv:math/0110234.
- [5] J. H. Conway, R. T. Curtis, S. P. Norton, R. A. Parker, and R. A. Wilson, *Atlas of Finite Groups: Maximal Subgroups and Ordinary Characters for Simple Groups*, Clarendon Press, Oxford, 1985.
- [6] J. H. Conway and N. J. A. Sloane, *Sphere Packings, Lattices and Groups*, 3rd ed., Grundlehren der mathematischen Wissenschaften 290, Springer, New York, 1999. doi:10.1007/978-1-4757-6568-7.
- [7] I. Frenkel, J. Lepowsky, and A. Meurman, *Vertex Operator Algebras and the Monster*, Pure and Applied Mathematics 134, Academic Press, 1988.
- [8] M. Geck, *A Course on Lie algebras and Chevalley groups*, 2024; revised version 3, 2 October 2025. arXiv:2404.11472v3.
- [9] G. Gonthier et al., A machine-checked proof of the Odd Order Theorem, in *Interactive Theorem Proving*, LNCS 7998, Springer, 2013, pp. 163–179. doi:10.1007/978-3-642-39634-2_14. Source: <https://github.com/math-comp/odd-order>.
- [10] R. L. Griess, Jr. and C. H. Lam, A new existence proof of the Monster by VOA theory, *Michigan Mathematical Journal* **61** (2012), 555–573. doi:10.1307/mmj/1347040259; arXiv:1103.1414.
- [11] R. L. Griess, Jr., U. Meierfrankenfeld, and Y. Segev, A uniqueness proof for the Monster, *Annals of Mathematics* **130** (1989), 567–602.

- [12] G. Höhn, *ATLAS: constructive finite simple groups in Lean*, source snapshot and verification records, 17 September 2026. Planned public repository: <https://github.com/Moonshine-in-Kansas/atlas>. Release after assignment of this article’s final arXiv identifier; proposed tag `atlas-arxiv-v1`. Source identification and checking scope are given in Appendix A.
- [13] G. Höhn, *The Conway–Parker Algebra and the largest Fischer group*, manuscript in preparation, September 2026; mathematical source for the Fischer algebra formalization.
- [14] G. Höhn, *The Hall–Janko Simple Group and the Icosian Leech Lattice*, manuscript in preparation, September 2026.
- [15] G. Höhn, *Peirce–triviality gluing codes and a mass-assisted classification of definite integral Albert algebras*, manuscript in preparation, 2026.
- [16] G. Höhn, *Selbstduale Vertexoperatorsuperalgebren und das Babymonster*, Bonner Mathematische Schriften 286, 1996. arXiv:0706.0236.
- [17] G. Höhn, *The unique extremal threemodular lattice of rank 26, the generalized hexagon (2, 8), and the tight Cayley-plane 5-design*, manuscript in preparation, 2026.
- [18] G. Höhn and M. Seysen, *The Order of the Monster Finite Simple Group*, 2025. arXiv:2508.01037.
- [19] C. Jansen and R. A. Wilson, Two new constructions of the O’Nan group, *Journal of the London Mathematical Society* **56** (1997), 579–583. doi:10.1112/S0024610798005742.
- [20] W. Kimmerle, R. Lyons, R. Sandling, and D. N. Teague, Composition factors from the group ring and Artin’s theorem on orders of simple groups, *Proceedings of the London Mathematical Society* (3) **60** (1990), 89–122. doi:10.1112/plms/s3-60.1.89.
- [21] KitaKen1, *FiniteSimpleGroups: Lean/mathlib formalizations of finite simple groups*, software repository and README, consulted 18 September 2026. <https://github.com/KitaKen1/finite-simple-groups-lean>.
- [22] R. van de Kuilen, *The first Janko group J_1 : simplicity and formalization*, bachelor’s thesis, Utrecht University, 2024. <https://studenttheses.uu.nl/items/7f8263cb-453b-4e78-99e0-35135037c33e>.
- [23] M. Lee, Computation and the sporadic simple groups, in *Computational Group Theory*, Oberwolfach Report 27/2025, pp. 1411–1413. https://publications.mfo.de/bitstream/handle/mfo/4340/OWR_2025_27.pdf.
- [24] The mathlib Community, The Lean mathematical library, in *Proceedings of the 9th ACM SIGPLAN International Conference on Certified Programs and Proofs (CPP 2020)*, ACM, 2020, pp. 367–381. doi:10.1145/3372885.3373824; arXiv:1910.09336.
- [25] The mathlib Community, *Mathlib group-theory documentation*, including alternating groups and projective special linear groups. https://leanprover-community.github.io/mathlib4_docs/Mathlib/GroupTheory/SpecificGroups/Alternating.html.
- [26] The Lean development team, *Validating a Lean Proof*, Lean Language Reference, online documentation, consulted 18 September 2026. <https://lean-lang.org/doc/reference/latest/ValidatingProofs/>.
- [27] L. de Moura, *Postmortem for the Kernel Soundness Bug Hunt*, 24 August 2026, online article.
- [28] L. de Moura and S. Ullrich, The Lean 4 theorem prover and programming language, in *Automated Deduction—CADE 28*, Lecture Notes in Computer Science 12699, Springer, 2021, pp. 625–635. doi:10.1007/978-3-030-79876-5_37.
- [29] T. Nie, A. Zhang, Y. Tang, D. Testa, S.-T. Yau, P. Li, and Y. Zhou, *FormaTheoria: Constructing Large-Scale Lean Theories from Mathematical Literature—Toward the Formalization of the Classification of Finite Simple Groups*, 2026. arXiv:2608.10894. Source and project-status description: <https://github.com/Qiuzhen-CFSG/CFSG>, consulted 18 September 2026.
- [30] E. van der Plas, *Formalisation of the Finite Simple Conway Groups in Lean*, bachelor’s thesis, Utrecht University, 2024. <https://studenttheses.uu.nl/handle/20.500.12932/46756>.
- [31] M. Seysen, A fast implementation of the Monster group: The Monster has been tamed, *Journal of Computational Algebra* **9** (2024), Article 100012. doi:10.1016/j.jaca.2024.100012.
- [32] H. Shimakura, The automorphism group of the vertex operator algebra V_L^+ for an even lattice L without roots, *Journal of Algebra* **280** (2004), 29–57. doi:10.1016/j.jalgebra.2004.05.018; arXiv:math/0311141.
- [33] R. Steinberg, *Lectures on Chevalley Groups*, University Lecture Series 66, American Mathematical Society, 2016; original Yale lecture notes, 1967–1968.

- [34] D. E. Taylor, *The Geometry of the Classical Groups*, Sigma Series in Pure Mathematics 9, Heldermann Verlag, Berlin, 1992. Author's corrected text: <https://www.maths.usyd.edu.au/u/don/papers/gcg.pdf>.
- [35] R. A. Wilson, Another new approach to the small Ree groups, *Archiv der Mathematik* **94** (2010), 501–510. doi:10.1007/s00013-010-0130-4.
- [36] R. A. Wilson, A new construction of the Ree groups of type 2G_2 , *Proceedings of the Edinburgh Mathematical Society* **53** (2010), 531–542. doi:10.1017/S001309150800028X. Author's text: https://webpace.maths.qmul.ac.uk/r.a.wilson/pubs_files/ReeG2.pdf.
- [37] R. A. Wilson, *The Finite Simple Groups*, Graduate Texts in Mathematics 251, Springer, London, 2009. doi:10.1007/978-1-84800-988-2.
- [38] R. A. Wilson and collaborators, *ATLAS of Finite Group Representations*, online database; classical and exceptional small-group identifications consulted 15 September 2026. <https://brauer.maths.qmul.ac.uk/Atlas/>.