

Discrete Mathematics, 2016 Spring - HW 11

November 27, 2016

Instructor: Zsolt Pajor-Gyulai

Courant Institute of Mathematical Sciences, NYU

To get full credit in all of the problems, use rigorous justification and unless otherwise indicated, make sure that your solution reads as a perfect English sentence. You should only assume integers, operations and order relations as given. If you use a statement or a definition from the textbook, make sure to indicate it.

Section 37

- 8) Prove Proposition 37.4 in the textbook. Why is this proposition restricted to $n \geq 2$?
- 3) Solve the following equations for x in the \mathbb{Z}_n specified. Make sure to find all solutions.
 - (a) $342 \otimes x = 73$ in \mathbb{Z}_{1003} .
 - (b) $9 \otimes x = 4$ in \mathbb{Z}_{12} .
- 6) Prove that for all $a, b \in \mathbb{Z}_n$, $(a \ominus b) \oplus (b \ominus a) = 0$.
- 10) For ordinary integers, the following is true. If $ab = 0$, then $a = 0$ or $b = 0$. Note, however that in \mathbb{Z}_n ,

$$2 \otimes 5 = 0, \quad \text{but} \quad 2 \neq 0, 5 \neq 0.$$

For which values of $n \geq 2$ does the implication

$$a \otimes b = 0 \Leftrightarrow a = 0 \text{ or } b = 0$$

hold in \mathbb{Z}_n ?

Section 38

- 1) Solve $100x \equiv 74 \pmod{127}$. Make sure to find all solutions.
- 3) Solve the following system of congruences

$$3x \equiv 8 \pmod{10} \quad 2x + 4 \equiv 9 \pmod{11}$$

Section 39

- 10) Let a and b be integers. A *common multiple* of a and b is an integer n for which $a|n$ and $b|n$. We call an integer m the *least common multiple* of n provided (1) m is positive, (2) m is a common multiple of a and b , and (3) if n is any other positive common multiple of a and b , then $n \geq m$. For example, $\text{lcm}(24, 30) = 120$.

- (a) Develop a formula for the least common multiple of two positive integers in terms of their prime factorization.
- (b) Use your formula to show that if a and b are positive integers, then

$$ab = \gcd(a, b)\text{lcm}(a, b)$$

- 14) Let n be a positive integer and suppose we factor n into primes as follows:

$$n = p_1^{e_1} p_2^{e_2} \cdots p_t^{e_t},$$

where the p_j -s are distinct primes and the e_j -s are natural numbers.

- (a) Find a formula for the number of positive divisors of n .
 - (b) Recall that an integer n is called *perfect* if it equals the sum of all its divisors d with $1 \leq d < n$. For example 28 is perfect as $28 = 1 + 2 + 4 + 7 + 14$. Let a be a positive integer and prove that if $2^a - 1$ is prime, then $n = 2^{a-1}(2^a - 1)$ is perfect.
- 22) Prove that $\log_2 3$ is irrational.