

Discrete Mathematics, Section 002, Spring 2016

Lecture 3: Counterexamples, Boolean Algebra, Lists

Zsolt Pajor-Gyulai

zsolt@cims.nyu.edu

Courant Institute of Mathematical Sciences

September 14, 2016



Refresher question

Question

A logician just had a child. A friend of hers asks her if it's a boy or a girl. What is her answer?

Refresher question

Question

A logician just had a child. A friend of hers asks her if it's a boy or a girl. What is her answer?

Answer

Yes.

Outline

1 Counterexamples

2 Boolean Algebra

3 Lists

How do we show if a statement is false?

- Show that it implies a contradiction. \rightarrow Later

How do we show if a statement is false?

- Show that it implies a contradiction. \rightarrow Later
- Come up with a counterexample. \rightarrow Now.

How do we show if a statement is false?

- Show that it implies a contradiction. \rightarrow Later
- Come up with a counterexample. \rightarrow Now.

For example:

False claim

Let a and b be integers. If $a|b$ and $b|a$, then $a = b$.

How do we show if a statement is false?

- Show that it implies a contradiction. \rightarrow Later
- Come up with a counterexample. \rightarrow Now.

For example:

False claim

Let a and b be integers. If $a|b$ and $b|a$, then $a = b$.

If I can find just one pair a, b that are not equal but they divide each other, then I can refute the statement!

How do we show if a statement is false?

- Show that it implies a contradiction. \rightarrow Later
- Come up with a counterexample. \rightarrow Now.

For example:

False claim

Let a and b be integers. If $a|b$ and $b|a$, then $a = b$.

If I can find just one pair a, b that are not equal but they divide each other, then I can refute the statement!

$$a = 5, b = -5 \quad \rightarrow \quad 5 \cdot (-1) = -5, \quad (-5) \cdot (-1) = 5$$

This works!

How do we show if a statement is false?

Refuting a false if-then statement via a counterexample.

To disprove a statement of the form 'If A , then B ': Find an instance where A is true but B is false.

How do we show if a statement is false?

Refuting a false if-then statement via a counterexample.

To disprove a statement of the form 'If A , then B ': Find an instance where A is true but B is false.

- Easier than proving true statements in general.

How do we show if a statement is false?

Refuting a false if-then statement via a counterexample.

To disprove a statement of the form 'If A , then B ': Find an instance where A is true but B is false.

- Easier than proving true statements in general.
- Finding counterexamples can be tricky!

How do we show if a statement is false?

Refuting a false if-then statement via a counterexample.

To disprove a statement of the form 'If A , then B ': Find an instance where A is true but B is false.

- Easier than proving true statements in general.
- Finding counterexamples can be tricky!
- Try to create strange examples!

When you do not know whether a statement is true or false:

- Try proving it!

How do we show if a statement is false?

Refuting a false if-then statement via a counterexample.

To disprove a statement of the form 'If A , then B ': Find an instance where A is true but B is false.

- Easier than proving true statements in general.
- Finding counterexamples can be tricky!
- Try to create strange examples!

When you do not know whether a statement is true or false:

- Try proving it!
- When you get stuck, try to figure out what the problem is and whether it suggests that there should be a counterexample.

Example

False claim

Let a and b be integers. If $a|b$ and $b|a$, then $a = b$.

Proof attempt

Let a and b be integers with $a|b$ and $b|a$.

... Therefore $a = b$.



Example

False claim

Let a and b be integers. If $a|b$ and $b|a$, then $a = b$.

Proof attempt

Let a and b be integers with $a|b$ and $b|a$. Since $a|b$, there is an integer x such that $b = ax$. Since $b|a$, there is an integer y such that $a = by$ Therefore $a = b$. □

Example

False claim

Let a and b be integers. If $a|b$ and $b|a$, then $a = b$.

Proof attempt

Let a and b be integers with $a|b$ and $b|a$. Since $a|b$, there is an integer x such that $b = ax$. Since $b|a$, there is an integer y such that $a = by$ Therefore $a = b$. □

While trying to bridge the gap, it looks like we should show $x = y = 1$.

$$a = by \quad \rightarrow \quad a = (ax)y \quad \rightarrow \quad a = (xy)a$$

Example

False claim

Let a and b be integers. If $a|b$ and $b|a$, then $a = b$.

Proof attempt

Let a and b be integers with $a|b$ and $b|a$. Since $a|b$, there is an integer x such that $b = ax$. Since $b|a$, there is an integer y such that $a = by$ Therefore $a = b$. □

While trying to bridge the gap, it looks like we should show $x = y = 1$.

$$a = by \quad \rightarrow \quad a = (ax)y \quad \rightarrow \quad a = (xy)a$$

Only shows that $xy = 1$. \rightarrow $x = y = -1$ works too!

Example

False claim

Let a and b be integers. If $a|b$ and $b|a$, then $a = b$.

Proof attempt

Let a and b be integers with $a|b$ and $b|a$. Since $a|b$, there is an integer x such that $b = ax$. Since $b|a$, there is an integer y such that $a = by$ Therefore $a = b$. □

While trying to bridge the gap, it looks like we should show $x = y = 1$.

$$a = by \quad \rightarrow \quad a = (ax)y \quad \rightarrow \quad a = (xy)a$$

Only shows that $xy = 1$. \rightarrow $x = y = -1$ works too!

This would yield $a = -b$ and now it is easy to pick the counterexample.

Outline

1 Counterexamples

2 Boolean Algebra

3 Lists

Ordinary Algebra

Useful for reasoning about numbers:

$$x^2 - y^2 = (x - y)(x + y) \quad \text{Holds for any numbers } x \text{ and } y$$

Ordinary Algebra

Useful for reasoning about numbers:

$$x^2 - y^2 = (x - y)(x + y) \quad \text{Holds for any numbers } x \text{ and } y$$

- Letters stand for numbers.

Ordinary Algebra

Useful for reasoning about numbers:

$$x^2 - y^2 = (x - y)(x + y) \quad \text{Holds for any numbers } x \text{ and } y$$

- Letters stand for numbers.
- Operators are the usual ones: $+$, $-$, \cdot , $/$

Ordinary Algebra

Useful for reasoning about numbers:

$$x^2 - y^2 = (x - y)(x + y) \quad \text{Holds for any numbers } x \text{ and } y$$

- Letters stand for numbers.
- Operators are the usual ones: $+$, $-$, \cdot , $/$
- Value of an expression like $3x - 4$ depends on x .

Ordinary Algebra

Useful for reasoning about numbers:

$$x^2 - y^2 = (x - y)(x + y) \quad \text{Holds for any numbers } x \text{ and } y$$

- Letters stand for numbers.
- Operators are the usual ones: $+$, $-$, \cdot , $/$
- Value of an expression like $3x - 4$ depends on x .
- When e.g. $x = 1$, the value is $3(-1) - 4 = -1$.

Boolean Algebra

Useful dealing with statements.

- Letters stand for the values **True** or **False** . (Two values!)

Boolean Algebra

Useful dealing with statements.

- Letters stand for the values **True** or **False** . (Two values!)
- Operators: and (\wedge), or (\vee), not (\neg), etc.

Boolean Algebra

Useful dealing with statements.

- Letters stand for the values **True** or **False** . (Two values!)
- Operators: and (\wedge), or (\vee), not (\neg), etc.
- We can create and evaluate more complicated **Boolean expressions**:

$$\neg(x \wedge y) \quad (\neg x) \vee (\neg y)$$

Boolean Algebra

Useful dealing with statements.

- Letters stand for the values **True** or **False** . (Two values!)
- Operators: and (\wedge), or (\vee), not (\neg), etc.
- We can create and evaluate more complicated **Boolean expressions**:

$$\neg(x \wedge y) \quad (\neg x) \vee (\neg y)$$

- Values of expressions can be summarized in **Truth tables**.

Truth table of $x \wedge y$ (AND):

x	y	$x \wedge y$
True	True	True
True	False	False
False	True	False
False	False	False

Boolean Algebra

Useful dealing with statements.

- Letters stand for the values **True** or **False** . (Two values!)
- Operators: and (\wedge), or (\vee), not (\neg), etc.
- We can create and evaluate more complicated **Boolean expressions**:

$$\neg(x \wedge y) \quad (\neg x) \vee (\neg y)$$

- Values of expressions can be summarized in **Truth tables**.

Truth table of $x \vee y$ (OR):

x	y	$x \vee y$
True	True	True
True	False	True
False	True	True
False	False	False

Boolean Algebra

Useful dealing with statements.

- Letters stand for the values **True** or **False** . (Two values!)
- Operators: and (\wedge), or (\vee), not (\neg), etc.
- We can create and evaluate more complicated **Boolean expressions**:

$$\neg(x \wedge y) \quad (\neg x) \vee (\neg y)$$

- Values of expressions can be summarized in **Truth tables**.

Truth table of $\neg x$ (NOT):

x	$\neg x$
True	False
False	True

Identities

- In ordinary algebra, you cannot check

$$(x + y)^2 = x^2 + 2xy + y^2$$

for (infinitely many) every number x and y .

Identities

- In ordinary algebra, you cannot check

$$(x + y)^2 = x^2 + 2xy + y^2$$

for (infinitely many) every number x and y .

- However, look at

$$\neg(x \wedge y) \qquad (\neg x) \vee (\neg y)$$

x	y	$x \wedge y$	$\neg(x \wedge y)$	$\neg x$	$\neg y$	$(\neg x) \vee (\neg y)$
T	T	T	F	F	F	F
T	F	F	T	F	T	T
F	T	F	T	T	F	T
F	F	F	T	T	T	T

Identities

- In ordinary algebra, you cannot check

$$(x + y)^2 = x^2 + 2xy + y^2$$

for (infinitely many) every number x and y .

- However, look at

$$\neg(x \wedge y) \qquad (\neg x) \vee (\neg y)$$

x	y	$x \wedge y$	$\neg(x \wedge y)$	$\neg x$	$\neg y$	$(\neg x) \vee (\neg y)$
T	T	T	F	F	F	F
T	F	F	T	F	T	T
F	T	F	T	T	F	T
F	F	F	T	T	T	T

We say $\neg(x \wedge y)$ and $(\neg x) \vee (\neg y)$ are **logically equivalent**.

Truth table proof of logical equivalence

To show that two Boolean expressions are logically equivalent:
Construct a truth table showing the values of the two expressions for all possible values of the variables.

Truth table proof of logical equivalence

To show that two Boolean expressions are logically equivalent:
Construct a truth table showing the values of the two expressions for all possible values of the variables.

Basic algebraic properties of the operations \wedge , \vee , \neg :

- **(Commutative properties)**

$$x \wedge y = y \wedge x, \quad x \vee y = y \vee x.$$

- **(Associative properties)**

$$(x \wedge y) \wedge z = x \wedge (y \wedge z), \quad (x \vee y) \vee z = x \vee (y \vee z).$$

- **(Identity elements)**

$$x \wedge \text{True} = x, \quad x \vee \text{False} = x.$$

Truth table proof of logical equivalence

To show that two Boolean expressions are logically equivalent:
Construct a truth table showing the values of the two expressions for all possible values of the variables.

- **(Distributive properties)**

$$x \wedge (y \vee z) = (x \wedge y) \vee (x \wedge z), \quad x \vee (y \wedge z) = (x \vee y) \wedge (x \vee z).$$

- **(DeMorgan's Laws)**

$$\neg(x \wedge y) = (\neg x) \vee (\neg y), \quad \neg(x \vee y) = (\neg x) \wedge (\neg y).$$

- $\neg(\neg x) = x.$

- $x \wedge x = x$ and $x \vee x = x.$

- $x \wedge (\neg x) = \textit{False}$ and $x \vee (\neg x) = \textit{True}.$

Further logical operators

- The operations \wedge , \vee , \neg were created to replicate mathematicians use of the words *and*, *or*, *not*.

Further logical operators

- The operations \wedge , \vee , \neg were created to replicate mathematicians use of the words *and*, *or*, *not*.
- We also introduce *implies* (\rightarrow) and *if-and-only-if* (\leftrightarrow).

Further logical operators

- The operations \wedge , \vee , \neg were created to replicate mathematicians use of the words *and*, *or*, *not*.
- We also introduce *implies* (\rightarrow) and *if-and-only-if* (\leftrightarrow).

Models $A \Rightarrow B$, only false when A is true but B is not.

x	y	$x \rightarrow y$
True	True	True
True	False	False
False	True	True
False	False	True

Models $A \leftrightarrow B$, false when one is true but the other one is false.

x	y	$x \leftrightarrow y$
True	True	True
True	False	False
False	True	False
False	False	True

Outline

1 Counterexamples

2 Boolean Algebra

3 Lists

Collections

In this course we consider two types of collections:

- **Lists**: Ordered collections.

Collections

In this course we consider two types of collections:

- **Lists**: Ordered collections.
- **Sets**: Unordered collections.

Lists

A **list** is an ordered sequence of objects.

Lists

A **list** is an ordered sequence of objects.

For example:

$(1, 2, \mathbb{Z})$

Lists

A **list** is an ordered sequence of objects.

For example:

$$(1, 2, \mathbb{Z})$$

- The order in which elements appear matters.

$$(1, 2, 3) \quad \text{vs} \quad (3, 2, 1)$$

Lists

A **list** is an ordered sequence of objects.

For example:

$$(1, 2, \mathbb{Z})$$

- The order in which elements appear matters.

$$(1, 2, 3) \quad \text{vs} \quad (3, 2, 1)$$

- Elements in a list might be repeated.

$$(3, 3, 2)$$

Lists

A **list** is an ordered sequence of objects.

For example:

$$(1, 2, \mathbb{Z})$$

- The order in which elements appear matters.

$$(1, 2, 3) \quad \text{vs} \quad (3, 2, 1)$$

- Elements in a list might be repeated.

$$(3, 3, 2)$$

- **Length:** Number of elements in a list.

$$(1, 1, 2, 1) \text{ has length } 4.$$

Lists

A **list** is an ordered sequence of objects.

For example:

$$(1, 2, \mathbb{Z})$$

- The order in which elements appear matters.

$$(1, 2, 3) \quad \text{vs} \quad (3, 2, 1)$$

- Elements in a list might be repeated.

$$(3, 3, 2)$$


- **Length:** Number of elements in a list.

$$(1, 1, 2, 1) \text{ has length } 4.$$

- **Ordered pair:** A list of length two.


Example of lists:

- A point on the $x - y$ plane.



Example of lists:


- A point on the $x - y$ plane.



- A natural number, e.g. 172 is $\{1, 7, 2\}$.

Example of lists:


- A point on the $x - y$ plane.



- A natural number, e.g. 172 is $\{1, 7, 2\}$.
- An English word, e.g. $\{C, h, e, e, r, s\}$.

Example of lists:

- A point on the $x - y$ plane.



- A natural number, e.g. 172 is $\{1, 7, 2\}$.
- An English word, e.g. $\{C, h, e, e, r, s\}$.

Equality of lists

Two lists are **equal** provided they have the same length, and elements in the corresponding positions on the two lists are equal.

E.g. lists (a, b, c) and (x, y, z) are equal if

$$a = x, b = y, \text{ and } c = z.$$

Counting two element lists

Question

How many ordered pairs are there where the entries in the list may be any of the digits 1, 2, 3, and 4?

Counting two element lists

Question

How many ordered pairs are there where the entries in the list may be any of the digits 1, 2, 3, and 4?

Write out all possibilities:

(1, 1)	(1, 2)	(1, 3)	(1, 4)
(2, 1)	(2, 2)	(2, 3)	(2, 4)
(3, 1)	(3, 2)	(3, 3)	(3, 4)
(4, 1)	(4, 2)	(4, 3)	(4, 4)

Counting two element lists

Question

How many ordered pairs are there where the entries in the list may be any of the digits 1, 2, 3, and 4?

Write out all possibilities:

(1, 1)	(1, 2)	(1, 3)	(1, 4)
(2, 1)	(2, 2)	(2, 3)	(2, 4)
(3, 1)	(3, 2)	(3, 3)	(3, 4)
(4, 1)	(4, 2)	(4, 3)	(4, 4)

We wrote this in a manner ensuring we have neither repeated nor omitted any such lists.

Counting two element lists

Question

How many ordered pairs are there where the entries in the list may be any of the digits 1, 2, 3, and 4?

Write out all possibilities:

(1, 1)	(1, 2)	(1, 3)	(1, 4)
(2, 1)	(2, 2)	(2, 3)	(2, 4)
(3, 1)	(3, 2)	(3, 3)	(3, 4)
(4, 1)	(4, 2)	(4, 3)	(4, 4)

- First row contains all lists beginning with 1, second row contains the ones beginning with 2, etc.
- There are 4 such rows.
- Each row contains four lists.

Counting two element lists

Question

How many ordered pairs are there where the entries in the list may be any of the digits 1, 2, 3, and 4?

Write out all possibilities:

(1, 1)	(1, 2)	(1, 3)	(1, 4)
(2, 1)	(2, 2)	(2, 3)	(2, 4)
(3, 1)	(3, 2)	(3, 3)	(3, 4)
(4, 1)	(4, 2)	(4, 3)	(4, 4)

There are 16 such lists.

Counting two element lists

Question

How many ordered pairs are there where the entries in the list may be any of the digits $1, 2, 3, \dots, n$?

Counting two element lists

Question

How many ordered pairs are there where the entries in the list may be any of the digits $1, 2, 3, \dots, n$?

$$\begin{array}{cccc}
 (1, 1) & (1, 2) & \dots & (1, n) \\
 (2, 1) & (2, 2) & \dots & (2, n) \\
 \vdots & \vdots & \ddots & \vdots \\
 (n, 1) & (n, 2) & \dots & (n, n)
 \end{array}$$

Counting two element lists

Question

How many ordered pairs are there where the entries in the list may be any of the digits $1, 2, 3, \dots, n$?

$$\begin{array}{cccc} (1, 1) & (1, 2) & \dots & (1, n) \\ (2, 1) & (2, 2) & \dots & (2, n) \\ \vdots & \vdots & \ddots & \vdots \\ (n, 1) & (n, 2) & \dots & (n, n) \end{array}$$

There are n rows, each containing n elements.

Counting two element lists

Question

How many ordered pairs are there where the entries in the list may be any of the digits $1, 2, 3, \dots, n$?

$$\begin{array}{cccc}
 (1, 1) & (1, 2) & \dots & (1, n) \\
 (2, 1) & (2, 2) & \dots & (2, n) \\
 \vdots & \vdots & \ddots & \vdots \\
 (n, 1) & (n, 2) & \dots & (n, n)
 \end{array}$$

There are n rows, each containing n elements.

There are $n \cdot n = n^2$ such lists.

Counting two element lists

Question

How many ordered pairs are there for which there are n choices for the first element and m choices for the second one?

Counting two element lists

Question

How many ordered pairs are there for which there are n choices for the first element and m choices for the second one?

$$\begin{array}{cccc}
 (1, 1) & (1, 2) & \dots & (1, m) \\
 (2, 1) & (2, 2) & \dots & (2, m) \\
 \vdots & \vdots & \ddots & \vdots \\
 (n, 1) & (n, 2) & \dots & (n, m)
 \end{array}$$

Counting two element lists

Question

How many ordered pairs are there for which there are n choices for the first element and m choices for the second one?

$$\begin{array}{cccc}
 (1, 1) & (1, 2) & \dots & (1, m) \\
 (2, 1) & (2, 2) & \dots & (2, m) \\
 \vdots & \vdots & \ddots & \vdots \\
 (n, 1) & (n, 2) & \dots & (n, m)
 \end{array}$$

There are n rows, each containing m elements.

Counting two element lists

Question

How many ordered pairs are there for which there are n choices for the first element and m choices for the second one?

$$\begin{array}{cccc}
 (1, 1) & (1, 2) & \dots & (1, m) \\
 (2, 1) & (2, 2) & \dots & (2, m) \\
 \vdots & \vdots & \ddots & \vdots \\
 (n, 1) & (n, 2) & \dots & (n, m)
 \end{array}$$

There are n rows, each containing m elements.

There are $n \cdot m$ such lists.

Counting two element lists

Question

How many ordered pairs are there where the entries in the list may be any of the digits 1, 2, 3, 4, 5, but where the two numbers in the list are different?

Counting two element lists

Question

How many ordered pairs are there where the entries in the list may be any of the digits 1, 2, 3, 4, 5, but where the two numbers in the list are different?

—	(1, 2)	(1, 3)	(1, 4)	(1, 5)
(2, 1)	—	(2, 3)	(2, 4)	(2, 5)
(3, 1)	(3, 2)	—	(3, 4)	(3, 5)
(4, 1)	(4, 2)	(4, 3)	—	(4, 5)
(5, 1)	(5, 2)	(5, 3)	(5, 4)	—

Counting two element lists

Question

How many ordered pairs are there where the entries in the list may be any of the digits 1, 2, 3, 4, 5, but where the two numbers in the list are different?

—	(1, 2)	(1, 3)	(1, 4)	(1, 5)
(2, 1)	—	(2, 3)	(2, 4)	(2, 5)
(3, 1)	(3, 2)	—	(3, 4)	(3, 5)
(4, 1)	(4, 2)	(4, 3)	—	(4, 5)
(5, 1)	(5, 2)	(5, 3)	(5, 4)	—

- There are 5 rows, each containing 4 elements.

Counting two element lists

Question

How many ordered pairs are there where the entries in the list may be any of the digits 1, 2, 3, 4, 5, but where the two numbers in the list are different?

—	(1, 2)	(1, 3)	(1, 4)	(1, 5)
(2, 1)	—	(2, 3)	(2, 4)	(2, 5)
(3, 1)	(3, 2)	—	(3, 4)	(3, 5)
(4, 1)	(4, 2)	(4, 3)	—	(4, 5)
(5, 1)	(5, 2)	(5, 3)	(5, 4)	—

- There are 5 rows, each containing 4 elements.
- 5 choices for the first element and 4 for the second.

Counting two element lists

Question

How many ordered pairs are there where the entries in the list may be any of the digits 1, 2, 3, 4, 5, but where the two numbers in the list are different?

—	(1, 2)	(1, 3)	(1, 4)	(1, 5)
(2, 1)	—	(2, 3)	(2, 4)	(2, 5)
(3, 1)	(3, 2)	—	(3, 4)	(3, 5)
(4, 1)	(4, 2)	(4, 3)	—	(4, 5)
(5, 1)	(5, 2)	(5, 3)	(5, 4)	—

- There are 5 rows, each containing 4 elements.
- 5 choices for the first element and 4 for the second.
- The choices are not independent, but that doesn't matter.

Counting two element lists

Question

How many ordered pairs are there where the entries in the list may be any of the digits 1, 2, 3, 4, 5, but where the two numbers in the list are different?

—	(1, 2)	(1, 3)	(1, 4)	(1, 5)
(2, 1)	—	(2, 3)	(2, 4)	(2, 5)
(3, 1)	(3, 2)	—	(3, 4)	(3, 5)
(4, 1)	(4, 2)	(4, 3)	—	(4, 5)
(5, 1)	(5, 2)	(5, 3)	(5, 4)	—

- There are 5 rows, each containing 4 elements.
- 5 choices for the first element and 4 for the second.
- The choices are not independent, but that doesn't matter.

There are $5 \cdot 4 = 20$ such lists.

Theorem (Multiplication Principle)

Consider two-element lists for which there are n choices for the first element, and for each choice of the first element, there are m choices for the second element. Then the number of such lists is nm .

Theorem (Multiplication Principle)

Consider two-element lists for which there are n choices for the first element, and for each choice of the first element, there are m choices for the second element. Then the number of such lists is nm .

Proof.

Construct a chart of all possible lists. Each row of this chart contains all the two-element lists that begin with a particular element. Since there are n choices for the first element, there are n rows in the chart. Since, for each choice of the first element, there are m choices for the second element, we know that every row of the chart has m entries. Therefore the number of lists is

$$\underbrace{m + m + \cdots + m}_{n \text{ times}} = n \times m$$



Longer lists

Question

How many lists of length three are there where the entries in the list may be any of the digits 1, 2, 3?

Longer lists

Question

How many lists of length three are there where the entries in the list may be any of the digits 1, 2, 3?

(1, 1, 1)	(1, 1, 2)	(1, 1, 3)
(1, 2, 1)	(1, 2, 2)	(1, 2, 3)
(1, 3, 1)	(1, 3, 2)	(1, 3, 3)
(2, 1, 1)	(2, 1, 2)	(2, 1, 3)
(2, 2, 1)	(2, 2, 2)	(2, 2, 3)
(2, 3, 1)	(2, 3, 2)	(2, 3, 3)
(3, 1, 1)	(3, 1, 2)	(3, 1, 3)
(3, 2, 1)	(3, 2, 2)	(3, 2, 3)
(3, 3, 1)	(3, 3, 2)	(3, 3, 3)

Longer lists

Question

How many lists of length three are there where the entries in the list may be any of the digits 1, 2, 3?

(1, 1, 1)	(1, 1, 2)	(1, 1, 3)
(1, 2, 1)	(1, 2, 2)	(1, 2, 3)
⋮	⋮	⋮
(3, 2, 1)	(3, 2, 2)	(3, 2, 3)
(3, 3, 1)	(3, 3, 2)	(3, 3, 3)

Longer lists

Question

How many lists of length three are there where the entries in the list may be any of the digits 1, 2, 3?

(1, 1, 1)	(1, 1, 2)	(1, 1, 3)
(1, 2, 1)	(1, 2, 2)	(1, 2, 3)
⋮	⋮	⋮
(3, 2, 1)	(3, 2, 2)	(3, 2, 3)
(3, 3, 1)	(3, 3, 2)	(3, 3, 3)

- Every row of the chart corresponds to a fixed combination of the first two elements.
- Every row consists of 3 elements.
- How many rows are there in this chart?

Longer lists

Question

How many lists of length three are there where the entries in the list may be any of the digits 1, 2, 3?

(1, 1, 1)	(1, 1, 2)	(1, 1, 3)
(1, 2, 1)	(1, 2, 2)	(1, 2, 3)
⋮	⋮	⋮
(3, 2, 1)	(3, 2, 2)	(3, 2, 3)
(3, 3, 1)	(3, 3, 2)	(3, 3, 3)

- How many rows are there in this chart?

We have solved this problem! $\rightarrow 3 \times 3 = 9.$

Longer lists

Question

How many lists of length three are there where the entries in the list may be any of the digits 1, 2, 3?

(1, 1, 1)	(1, 1, 2)	(1, 1, 3)
(1, 2, 1)	(1, 2, 2)	(1, 2, 3)
⋮	⋮	⋮
(3, 2, 1)	(3, 2, 2)	(3, 2, 3)
(3, 3, 1)	(3, 3, 2)	(3, 3, 3)

- How many rows are there in this chart? $\rightarrow 3 \times 3 = 9$

There are $9 \times 3 = 3 \times 3 \times 3 = 27$ such lists.

Question

How many lists of length three are there where the entries in the list may be any of the digits 1, 2, 3, 4, 5 in which no number is repeated?

Question

How many lists of length three are there where the entries in the list may be any of the digits 1, 2, 3, 4, 5 in which no number is repeated?

We could write out the chart again but rather think:

- For the first element, we have 5 choices.

Question

How many lists of length three are there where the entries in the list may be any of the digits 1, 2, 3, 4, 5 in which no number is repeated?

We could write out the chart again but rather think:

- For the first element, we have 5 choices.
- For the second element, we have to choose from the 4 remaining.

Question

How many lists of length three are there where the entries in the list may be any of the digits 1, 2, 3, 4, 5 in which no number is repeated?

We could write out the chart again but rather think:

- For the first element, we have 5 choices.
- For the second element, we have to choose from the 4 remaining.
- For the third place, we have to choose from the 3 remaining.

Question

How many lists of length three are there where the entries in the list may be any of the digits 1, 2, 3, 4, 5 in which no number is repeated?

We could write out the chart again but rather think:

- For the first element, we have 5 choices.
- For the second element, we have to choose from the 4 remaining.
- For the third place, we have to choose from the 3 remaining.

There are $5 \times 4 \times 3 = 60$ such lists.

In general

How many lists are there of length k , in which each element of the list is selected from among n possibilities.

1 Repetitions allowed:

$$\underbrace{n \times n \times \cdots \times n}_{k \text{ times}} = n^k$$

2 Repetitions not allowed:

$$n \times (n - 1) \times (n - 2) \times \cdots \times n - (k - 1)$$

In general

How many lists are there of length k , in which each element of the list is selected from among n possibilities.

❶ Repetitions allowed:

$$\underbrace{n \times n \times \cdots \times n}_{k \text{ times}} = n^k$$

❷ Repetitions not allowed:

$$n \times (n - 1) \times (n - 2) \times \cdots \times n - (k - 1)$$

Is the second formula okay? What about $n = 2$, $k = 4$?

In general

How many lists are there of length k , in which each element of the list is selected from among n possibilities.

① Repetitions allowed:

$$\underbrace{n \times n \times \cdots \times n}_{k \text{ times}} = n^k$$

② Repetitions not allowed:

$$n \times (n-1) \times (n-2) \times \cdots \times n - (k-1)$$

when $k \leq n$, and 0 if $k > n$.

Is the second formula okay? What about $n = 2$, $k = 4$?

In general

Notation

The number

$$(n)_k = n(n-1)(n-2)\dots(n-k+1)$$

is called **falling factorial**.

In general

Notation

The number

$$(n)_k = n(n-1)(n-2)\dots(n-k+1)$$

is called **falling factorial**.

Theorem

The number of lists of length k whose elements are chosen from a pool of n possible elements is

$$= \begin{cases} n^k & \text{if repetitions are permitted} \\ (n)_k & \text{if repetitions are forbidden} \end{cases}$$