

Discrete Mathematics, Section 002, Spring 2016

Lecture 4: Factorials, Sets

Zsolt Pajor-Gyulai

`zsolt@cims.nyu.edu`

Courant Institute of Mathematical Sciences

September 19, 2016



Outline

- ## 2 Introduction to sets, Subsets

Factorials

A special case of what we did last time:

How many lists of length n can we make using n elements without repetition?

or alternatively

How many ways can we order n elements?

Factorials

A special case of what we did last time:

How many lists of length n can we make using n elements without repetition?

or alternatively

How many ways can we order n elements?

$$(n)_n = n(n-1)(n-2) \cdots 3 \cdot 2 \cdot 1$$

Factorials

A special case of what we did last time:

How many lists of length n can we make using n elements without repetition?

or alternatively

How many ways can we order n elements?

$$(n)_n = n(n-1)(n-2) \cdots 3 \cdot 2 \cdot 1$$

This occurs a lot, we call it **n -factorial** and denote it by $n!$.

Factorials

A special case of what we did last time:

How many lists of length n can we make using n elements without repetition?

or alternatively

How many ways can we order n elements?

$$(n)_n = n(n-1)(n-2) \cdots 3 \cdot 2 \cdot 1$$

This occurs a lot, we call it **n -factorial** and denote it by $n!$.

- Note that

$$n! = n \cdot (n-1)!$$

- Also

$$\begin{aligned}(n)_k &= n(n-1)\dots(n-k+1) = \\ &= \frac{n(n-1)\dots 2 \cdot 1}{(n-k)(n-k-1)\dots 2 \cdot 1} = \frac{n!}{(n-k)!}\end{aligned}$$

Factorials

Two special cases:

① $1! = 1$: Hardly suprising.

Factorials

Two special cases:

- 1 $1! = 1$: Hardly suprising.
- 2 $0! = 1$:

Factorials

Two special cases:

- ① $1! = 1$: Hardly suprising.
- ② $0! = 1$:
 - How many list of length zero can we form out of zero elements?

One, the empty list.

Factorials

Two special cases:

① $1! = 1$: Hardly suprising.

② $0! = 1$:

- How many list of length zero can we form out of zero elements?

One, the empty list.

- When $n = 1$, $n! = n(n-1)!$ becomes $1 = 1 \cdot 0!$

Factorials

Two special cases:

① $1! = 1$: Hardly suprising.

② $0! = 1$:

- How many list of length zero can we form out of zero elements?

One, the empty list.

- When $n = 1$, $n! = n(n - 1)!$ becomes $1 = 1 \cdot 0!$
- At the end of the day we make the definition like this, because it's convenient.

Product notation

Another way to write factorials:

$$n! = \prod_{k=1}^n k$$

- \prod stands for product.
- k is a dummy variable, a place holder that ranges from the lower value to the upper value.

Product notation

Another way to write factorials:

$$n! = \prod_{k=1}^n k$$

- \prod stands for product.
- k is a dummy variable, a place holder that ranges from the lower value to the upper value.

Another example,

$$\prod_{k=1}^5 (2k + 3) = (2 \cdot 1 + 3)(2 \cdot 2 + 3)(2 \cdot 3 + 3)(2 \cdot 4 + 3)(2 \cdot 5 + 3)$$

Product notation

Interpretation as a for loop to compute

$$\prod_{k=1}^n f_k$$

```
def evaluate():  
    prod = 1  
    for k in range(1,n+1):  
        prod *= f_k  
  
    return prod
```

Product notation

Further examples:

- Constant to the right of the product sign:

$$\prod_{k=1}^n 2 = 2^n$$

Product notation

Further examples:

- Constant to the right of the product sign:

$$\prod_{k=1}^n 2 = 2^n$$

- Empty product:

$$0! = \prod_{k=1}^0 k = 1$$

Outline

- 1 Factorials and products
- 2 Introduction to sets, Subsets

Last time we talked about lists.

What was a list?

A **list** is an ordered sequence of objects.

Last time we talked about lists.

What was a list?

A **list** is an ordered sequence of objects.

What is a set?

A **set** is a repetition-free, unordered collection of objects.

Last time we talked about lists.

What was a list?

A **list** is an ordered sequence of objects.

What is a set?

A **set** is a repetition-free, unordered collection of objects.

- An object is either a member of a set or not.

Last time we talked about lists.

What was a list?

A **list** is an ordered sequence of objects.

What is a set?

A **set** is a repetition-free, unordered collection of objects.

- An object is either a member of a set or not.
- There is no order to the members.

Last time we talked about lists.

What was a list?

A **list** is an ordered sequence of objects.

What is a set?

A **set** is a repetition-free, unordered collection of objects.

- An object is either a member of a set or not.
- There is no order to the members.
- Simplest way to specify: list elements.

$$\left\{2, 3, \frac{1}{2}\right\} \quad \left\{3, \frac{1}{2}, 2\right\} \quad \left\{2, 2, 3, \frac{1}{2}\right\}$$

Last time we talked about lists.

What was a list?

A **list** is an ordered sequence of objects.

What is a set?

A **set** is a repetition-free, unordered collection of objects.

- An object is either a member of a set or not.
- There is no order to the members.
- Simplest way to specify: list elements.

$$\left\{2, 3, \frac{1}{2}\right\} \quad \left\{3, \frac{1}{2}, 2\right\} \quad \left\{2, 2, 3, \frac{1}{2}\right\}$$

These are all the same set!

For example, sets of numbers:

- Naturals: \mathbb{N}

For example, sets of numbers:

- Naturals: \mathbb{N}
- Integers: \mathbb{Z}

For example, sets of numbers:

- Naturals: \mathbb{N}
- Integers: \mathbb{Z}
- Rationals: \mathbb{Q}

For example, sets of numbers:

- Naturals: \mathbb{N}
- Integers: \mathbb{Z}
- Rationals: \mathbb{Q}

Membership in a set

An object x that belongs to a set A is called an **element** of it.

Notation: $x \in A$

For example, $x \in \mathbb{Z}$ reads:

- "x is a member/element of \mathbb{Z} "

For example, sets of numbers:

- Naturals: \mathbb{N}
- Integers: \mathbb{Z}
- Rationals: \mathbb{Q}

Membership in a set

An object x that belongs to a set A is called an **element** of it.

Notation: $x \in A$

For example, $x \in \mathbb{Z}$ reads:

- "x is a member/element of \mathbb{Z} "
- "x is in \mathbb{Z} "

For example, sets of numbers:

- Naturals: \mathbb{N}
- Integers: \mathbb{Z}
- Rationals: \mathbb{Q}

Membership in a set

An object x that belongs to a set A is called an **element** of it.

Notation: $x \in A$

For example, $x \in \mathbb{Z}$ reads:

- "x is a member/element of \mathbb{Z} "
- "x is in \mathbb{Z} "
- "x is an integer"

Cardinality of a set

The **cardinality** of A , denoted by $|A|$, is the number of objects in the set.

Cardinality of a set

The **cardinality** of A , denoted by $|A|$, is the number of objects in the set.

For example,

$$\left| \left\{ 2, 3, \frac{1}{2} \right\} \right| = 3, \quad |\mathbb{Z}| = \infty$$

Cardinality of a set

The **cardinality** of A , denoted by $|A|$, is the number of objects in the set.

For example,

$$\left| \left\{ 2, 3, \frac{1}{2} \right\} \right| = 3, \quad |\mathbb{Z}| = \infty$$

A set A with $|A| \in \mathbb{N}$ is **finite**, otherwise it's **infinite**.

Empty set

The **empty set** \emptyset is a set with no elements.

Cardinality of a set

The **cardinality** of A , denoted by $|A|$, is the number of objects in the set.

For example,

$$\left| \left\{ 2, 3, \frac{1}{2} \right\} \right| = 3, \quad |\mathbb{Z}| = \infty$$

A set A with $|A| \in \mathbb{N}$ is **finite**, otherwise it's **infinite**.

Empty set

The **empty set** \emptyset is a set with no elements.

- The statement $x \in \emptyset$ is false.

Cardinality of a set

The **cardinality** of A , denoted by $|A|$, is the number of objects in the set.

For example,

$$\left| \left\{ 2, 3, \frac{1}{2} \right\} \right| = 3, \quad |\mathbb{Z}| = \infty$$

A set A with $|A| \in \mathbb{N}$ is **finite**, otherwise it's **infinite**.

Empty set

The **empty set** \emptyset is a set with no elements.

- The statement $x \in \emptyset$ is false.
- $|\emptyset| = 0$

Specifying sets

- List the elements between curly braces.

$\{3, 4, 9\}, \{table, chair, lamp\}$

Specifying sets

- List the elements between curly braces.

 $\{3, 4, 9\}, \{table, chair, lamp\}$

Clumsy for large sets, and impossible for infinite sets.

Specifying sets

- List the elements between curly braces.

$$\{3, 4, 9\}, \{table, chair, lamp\}$$

Clumsy for large sets, and impossible for infinite sets.

- Set-builder notation:

$$\{\text{dummy variable} : \text{conditions}\}$$

Specifying sets

- List the elements between curly braces.

$$\{3, 4, 9\}, \{table, chair, lamp\}$$

Clumsy for large sets, and impossible for infinite sets.

- Set-builder notation:

$$\{\text{dummy variable} : \text{conditions}\}$$

For example,

$$\mathbb{N} = \{x : x \in \mathbb{Z}, x \geq 0\}$$

This is a set of objects satisfying

- $x \in \mathbb{Z}$
- $x \geq 0$

Specifying sets

- Alternative set-builder notation:

$$\{\text{dummy variable} \in \text{set} : \text{conditions}\}$$

Specifying sets

- Alternative set-builder notation:

$$\{\text{dummy variable} \in \text{set} : \text{conditions}\}$$

For example,

$$\{x \in \mathbb{Z} : 2|x\}$$

This is the set of those integers that are divisible by 2.

Specifying sets

- Alternative set-builder notation:

$$\{\text{dummy variable} \in \text{set} : \text{conditions}\}$$

For example,

$$\{x \in \mathbb{Z} : 2|x\}$$

This is the set of those integers that are divisible by 2.

Now practice these on the worksheet!

Equality of sets

Definition

Two sets are equal if they have exactly the same elements.

Equality of sets

Definition

Two sets are equal if they have exactly the same elements.

To show $A = B$, we have the following template:

Let A and B be the sets.

- Suppose $x \in A \dots$ Therefore $x \in B$.
- Suppose $x \in B \dots$ Therefore $x \in A$.

Therefore $A = B$.

Proposition

The following two sets are equal:

$$E = \{x \in \mathbb{Z} : x \text{ is even}\},$$

$$F = \{x \in \mathbb{Z} : x = a + b \text{ where } a \text{ and } b \text{ are both odd}\}.$$

Proof.



Proposition

The following two sets are equal:

$$E = \{x \in \mathbb{Z} : x \text{ is even}\},$$

$$F = \{x \in \mathbb{Z} : x = a + b \text{ where } a \text{ and } b \text{ are both odd}\}.$$

Proof.

Let E and F as in the statement of the proposition. We seek to prove that $E = F$.

Suppose $x \in E$.

Therefore

$$x \in F.$$

Suppose $x \in F$.

Therefore

$$x \in E.$$



Proposition

The following two sets are equal:

$$E = \{x \in \mathbb{Z} : x \text{ is even}\},$$

$$F = \{x \in \mathbb{Z} : x = a + b \text{ where } a \text{ and } b \text{ are both odd}\}.$$

Proof.

Let E and F as in the statement of the proposition. We seek to prove that $E = F$.

Suppose $x \in E$. Therefore x is even, hence $2|x$ and so $x = 2y$ for some $y \in \mathbb{Z}$.

Therefore x is the sum of two odd numbers and so $x \in F$.

Suppose $x \in F$.

$x \in E$.

Therefore



Proposition

The following two sets are equal:

$$E = \{x \in \mathbb{Z} : x \text{ is even}\},$$

$$F = \{x \in \mathbb{Z} : x = a + b \text{ where } a \text{ and } b \text{ are both odd}\}.$$

Proof.

Let E and F as in the statement of the proposition. We seek to prove that $E = F$.

Suppose $x \in E$. Therefore x is even, hence $2|x$ and so $x = 2y$ for some $y \in \mathbb{Z}$. Note that $2y + 1$ and -1 are both odd and that $x = 2y = (2y + 1) + (-1)$. Therefore x is the sum of two odd numbers and so $x \in F$.

Suppose $x \in F$.

$x \in E$.

Therefore



Proposition

The following two sets are equal:

$$E = \{x \in \mathbb{Z} : x \text{ is even}\},$$

$$F = \{x \in \mathbb{Z} : x = a + b \text{ where } a \text{ and } b \text{ are both odd}\}.$$

Proof.

Let E and F as in the statement of the proposition. We seek to prove that $E = F$.

Suppose $x \in E$. Therefore x is even, hence $2|x$ and so $x = 2y$ for some $y \in \mathbb{Z}$. Note that $2y + 1$ and -1 are both odd and that $x = 2y = (2y + 1) + (-1)$. Therefore x is the sum of two odd numbers and so $x \in F$.

Suppose $x \in F$. **Therefore x is the sum of two odd integers. It was shown in Exercise 5.1 that x is then even.** Therefore $x \in E$. □

Subsets

Definition

Suppose A and B are sets. We say that A is a **subset** of B provided every element of A is also an element of B . We denote this by $A \subseteq B$.

Subsets

Definition

Suppose A and B are sets. We say that A is a **subset** of B provided every element of A is also an element of B . We denote this by $A \subseteq B$.

For example,

- $\{1, 2, 3\} \subseteq \{1, 2, 3, 4\}$

Subsets

Definition

Suppose A and B are sets. We say that A is a **subset** of B provided every element of A is also an element of B . We denote this by $A \subseteq B$.

For example,

- $\{1, 2, 3\} \subseteq \{1, 2, 3, 4\}$
- $A \subseteq A$ for every set A

Subsets

Definition

Suppose A and B are sets. We say that A is a **subset** of B provided every element of A is also an element of B . We denote this by $A \subseteq B$.

For example,

- $\{1, 2, 3\} \subseteq \{1, 2, 3, 4\}$
- $A \subseteq A$ for every set A
- $\emptyset \subseteq A$.
- If $B \subseteq A$ and $B \neq A, \emptyset$, we call it a **proper subset**.

Subsets

Definition

Suppose A and B are sets. We say that A is a **subset** of B provided every element of A is also an element of B . We denote this by $A \subseteq B$.

For example,

- $\{1, 2, 3\} \subseteq \{1, 2, 3, 4\}$
- $A \subseteq A$ for every set A
- $\emptyset \subseteq A$.
- If $B \subseteq A$ and $B \neq A, \emptyset$, we call it a **proper subset**.

To show that $A \subseteq B$:

Let $x \in A, \dots$ Therefore $x \in B$.

Subsets

Definition

A list of three integers (a, b, c) is called a **Pythagorean triple** provided $a^2 + b^2 = c^2$.

Subsets

Definition

A list of three integers (a, b, c) is called a **Pythagorean triple** provided $a^2 + b^2 = c^2$.

Proposition

Let P be the set of Pythagorean triples.

$$P = \{(a, b, c) : a, b, c \in \mathbb{Z} \text{ and } a^2 + b^2 = c^2\}$$

and

$$T = \{(p, q, r) : p = x^2 - y^2, q = 2xy, \\ \text{and } r = x^2 + y^2 \text{ where } x, y \in \mathbb{Z}\}$$

Then $T \subseteq P$.

$$T = \{(p, q, r) : p = x^2 - y^2, q = 2xy, \\ \text{and } r = x^2 + y^2 \text{ where } x, y \in \mathbb{Z}\}$$

For example $x = 3, y = 2,$

$$p = x^2 - y^2 = 9 - 4 = 5, \quad q = 2xy = 12, \quad r = x^2 + y^2 = 13$$

and $(p, q, r) \in P.$

$$T = \{(p, q, r) : p = x^2 - y^2, q = 2xy, \\ \text{and } r = x^2 + y^2 \text{ where } x, y \in \mathbb{Z}\}$$

Proof.

Let P and T as in the statement of the proposition.

Let $(p, q, r) \in T$.

Therefore $(p, q, r) \in P$.



$$T = \{(p, q, r) : p = x^2 - y^2, q = 2xy, \\ \text{and } r = x^2 + y^2 \text{ where } x, y \in \mathbb{Z}\}$$

Proof.

Let P and T as in the statement of the proposition.

Let $(p, q, r) \in T$. Therefore there are integers x and y such that $p = x^2 - y^2$, $q = 2xy$, and $r = x^2 + y^2$.

Therefore $(p, q, r) \in P$.



$$T = \{(p, q, r) : p = x^2 - y^2, q = 2xy, \\ \text{and } r = x^2 + y^2 \text{ where } x, y \in \mathbb{Z}\}$$

Proof.

Let P and T as in the statement of the proposition.

Let $(p, q, r) \in T$. Therefore there are integers x and y such that $p = x^2 - y^2$, $q = 2xy$, and $r = x^2 + y^2$. Note that p, q, r are integers because x and y are integers.

Therefore $(p, q, r) \in P$.



$$T = \{(p, q, r) : p = x^2 - y^2, q = 2xy, \\ \text{and } r = x^2 + y^2 \text{ where } x, y \in \mathbb{Z}\}$$

Proof.

Let P and T as in the statement of the proposition.

Let $(p, q, r) \in T$. Therefore there are integers x and y such that $p = x^2 - y^2$, $q = 2xy$, and $r = x^2 + y^2$. Note that p, q, r are integers because x and y are integers. We calculate

$$\begin{aligned} p^2 + q^2 &= (x^2 - y^2)^2 + (2xy)^2 = (x^4 - 2x^2y^2 + y^4) + 4x^2y^2 = \\ &= x^4 + 2x^2y^2 + y^4 = (x^2 + y^2)^2 = r^2 \end{aligned}$$

Therefore $(p, q, r) \in P$. □

Caveats

We have to be careful to make the distinction between

x vs. $\{x\}$

Caveats

We have to be careful to make the distinction between

x vs. $\{x\}$

and

\in vs. \subseteq

Caveats

We have to be careful to make the distinction between

x vs. $\{x\}$

and

\in vs. \subseteq

For example,

- $x \in \{x\}$
- But $x \subseteq \{x\}$ or $x = \{x\}$ are incorrect.

Question

How many subsets does a set have?

Question

How many subsets does a set have?

For example, what about $A = \{1, 2, 3\}$?

Number of elements	Subsets	Number
0	\emptyset	1
1	$\{1\}, \{2\}, \{3\}$	3
2	$\{1, 2\}, \{1, 3\}, \{2, 3\}$	3
3	$\{1, 2, 3\}$	1

Total: 8.

Question

How many subsets does a set have?

For example, what about $A = \{1, 2, 3\}$?

Number of elements	Subsets	Number
0	\emptyset	1
1	$\{1\}, \{2\}, \{3\}$	3
2	$\{1, 2\}, \{1, 3\}, \{2, 3\}$	3
3	$\{1, 2, 3\}$	1

Total: 8.

Alternatively

For every element, we have two choices independently of each other: include/not include.

Therefore $|A| = 2^3 = 8$.

Theorem

Let A be a finite set. The number of subsets of A is $2^{|A|}$.

Theorem

Let A be a finite set. The number of subsets of A is $2^{|A|}$.

Proof.

Let A be a finite set and let $n = |A|$. Let the n elements of A be named a_1, a_2, \dots, a_n . To each subset B of A , we can associate a list of length n ; each element of the list is one of the words "yes" or "no". The k th element of the list is "yes" precisely when $a_k \in B$. This establishes a one to one correspondence with yes/no lists of length n . The number of such lists is 2^n , so the number of subsets of A is 2^n where $n = |A|$. □

Theorem

Let A be a finite set. The number of subsets of A is $2^{|A|}$.

Proof.

Let A be a finite set and let $n = |A|$. Let the n elements of A be named a_1, a_2, \dots, a_n . To each subset B of A , we can associate a list of length n ; each element of the list is one of the words "yes" or "no". The k th element of the list is "yes" precisely when $a_k \in B$. This establishes a one to one correspondence with *yes/no* lists of length n . The number of such lists is 2^n , so the number of subsets of A is 2^n where $n = |A|$. □

This is a so-called **bijective** proof.

Power set

Definition

Let A be a set. The **power set** of A is the set of all subsets of A . We denote it by 2^A .

Power set

Definition

Let A be a set. The **power set** of A is the set of all subsets of A . We denote it by 2^A .

For example,

$$2^{\{1,2,3\}} = \{\emptyset, \{1\}, \{2\}, \{3\}, \{1,2\}, \{1,3\}, \{2,3\}, \{1,2,3\}\}$$

Power set

Definition

Let A be a set. The **power set** of A is the set of all subsets of A . We denote it by 2^A .

For example,

$$2^{\{1,2,3\}} = \{\emptyset, \{1\}, \{2\}, \{3\}, \{1,2\}, \{1,3\}, \{2,3\}, \{1,2,3\}\}$$

The notation is created so that

$$|2^A| = 2^{|A|}$$