Quantum Sphaera Companion: A Structure-First Mathematical Unfolding

Marc Brendecke
Independent Researcher, Bochum, Germany
Version DOI: 10.5281/zenodo.24108818
Concept DOI: 10.5281/zenodo.19210728
Preprint; not peer reviewed.

2026-11-10 — v4.11.0

Working Scope and Reading Route

This is the full-source version v4.11.0, derived from the complete v4.7.0 monolith, not from the older modular synchronization baseline, and continuing the published v4.8.0 line, whose layers are carried in (§A.270 and §A.271). The published v4.9.0 tag preserves the earlier source baseline; the 4.10 main body carries explicit corrections and status notes. The v4.11 research line starts with the conditional Koide readout and finite-gap analysis in §7. Provenance of the previous versions: v4.10.0, Version DOI 10.5281/zenodo.23991357; v4.9.0, Version DOI 10.5281/zenodo.23534954; v4.8.0, Version DOI 10.5281/zenodo.23365244; v4.7.0, Version DOI 10.5281/zenodo.20232911. The Concept DOI on the title page is common to all versions. The main line brings together the core construction, physical readouts, explicit premise qualifications and the next research interface. The appendices retain the complete definitions, proofs, protocols, finite ledgers and provenance.

The 4.10 consolidation. Section 44 separates state carriers, quantum and relativistic readouts, orientation-complete local mediation and calibrated clocks. It records finite field-compatibility and precision tests without claiming a regional OP5 solution. Figure 1 retains the original affine numerical baseline; later fitted fields are separate research records.

What changes, and what does not. The reading route changes; a stored or relocated assertion does not thereby become a valid theorem. The inherited material is preserved as source evidence, including unresolved or overstrong historical wording. The explicit qualifications in Section 1 govern the known MM/HC, exponent and OP5 issues. Section 2 proves the additional Koide condition and constructs a relative local kernel model, with its remaining native activation obligations made explicit. Section 3 tests the HC-to-lepton-root identification, distinguishes sector and generation triples, and isolates the missing residual coupling. Section 4 constructs a conditional closure-defect coupling and closed generation seeds, separates slot readouts from candidate eigenmasses, and tests the pacer phase. Section 5 tests the native operator bridge, derives the minimal penalty compression, and separates a conditional gapped-sector mass law from exact closure. Section 6 evaluates a normalized cyclic coupling candidate, tests the scale choices and proves a fixed-shape mass-ratio obstruction. Section 8 extends this test to explicit comparison and kinetic metrics, distinguishing covariant shifts from added physical tensor weights and target-derived fits. Section 9 carries the conditional HC program through projected dynamics, real actions and bilinear geometry. It separates the off-shell null stress match from the current test and the conditional same-induced-frame trace obstruction. Section 10 adds the explicitly declared neutral Cartan comparison, conditional gravitational constraint propagation, and a finite normal-source law for compatible tangential jets. These results do not establish the native action or regional existence. The current endpoint is summarized in Section 43; the v4.8 research program is separated in Section 45. Neither section promotes a finite detector result to exact all-readout closure.

Current boundary. The baseline permits guarded local/restricted and effective matter-detector statements under their stated assumptions. Exact finite C5 zero, unrestricted all-readout terminality, C6/T2 closure and a global photon-clock/Riemann theorem are not established here. The inherited scale and zero-status distinctions are retained in Appendix A.268; the next metric/field-equation program does not remove them.

Formal-language route. QAM supplies object coding and readout records, MML an instruction language, and MMA guarded transitions. Their full inherited definitions are in Ap-

© 2026 Marc Brendecke. Preprint — not peer reviewed.

pendix H2. The later shifted normal form, not an unqualified return to the pre-shift syntax, is the integration interface for new material; see Appendix A.27. Historical pre-shift displays remain available with their occurrence and backtranslation records. Typed commands and short references are neither background executions nor proof certificates.

Release discipline. The published v4.7.0, v4.8.0 and v4.9.0 TeX and PDF are unchanged. This is the source of the v4.10.0 release (Version DOI 10.5281/zenodo.23991357); no new physical closure claim is made by it. In inherited fragments, the macro `docversion` deliberately retains its v4.7.0 value so historical records are not relabelled as new results. The working title and metadata identify this copy separately.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Contents

Working Scope and Reading Route 2
1 Premise Audit and Scope 17
2 Koide and Local Kernel Construction 19
3 HC and Lepton Mass Readout: An Identification Test 23
4 Generation States and a Closure-Defect Coupling 26
5 Testing the Native Operator Link 30
6 An Evaluated Coupling Candidate and Its Scale Limits 34
7 Koide Phase Overlap and Finite-Gap Constraints 39
8 Tensor Activation and Covariant Coupling Readout 40
9 HC Dynamics, Real Actions and the Geometry Interface 42
10 Cartan Comparison: Residuals and Normal Source Compatibility . . . 48
Nomenclature 53
11 Alternating Triolic Orbit Scan and Hopf-Fibre Core Surface Readout . 55
12 The Central Theorem 66
13 Lorentz Invariance on Every Stratum 68
14 Fractal Self-Traversal Along the Light Axis 69
15 Three Objects as One: Unitary Equivalence 70
16 Near-Collision, Photon Production, and Entanglement 71
17 Connection to the Riemann Hypothesis 71
18 Gauge Structure of the Standard Model 72
19 Bell Inequalities and tau-n 72
20 String Theory and Loop Quantum Gravity as HC Sectors 73
21 Black Holes and Hawking Radiation 75
22 The Sphaera as a Hopf Fibration 78
23 From HC Decomposition to a QFT Lagrangian 86
24 GUE Self-Determination of the Frame Factor 87
25 Phase-Shifted Natural Constants 88
26 Dirac Probe as Closure-Preserving Solution 90
27 Field Equations of the Probe–Closure System 92
28 Closure of the Probe–Closure Field System 94
29 Structural Core and Closure Principles 98
30 Readout-Relative Lemmas and Structural Lifts 106
31 Bidirectional Probe Lensing Hypothesis 109
32 HC-Induced Readout Geometry and Structural Unity 120
33 Hypothesis 7.4: The Heisenberg–Compensator Matrix as a Meta-Field Generator 122
34 Closure-Liquid Crystal and Holographic Julia Boundary 128
35 Toward a Structural ToE Master Equation 131
36 Structural Reading of Matter, Gauge, Mass, and Higgs 134
37 Structural Normalization for the Next Companion Stage 135
38 The Compensator Matrix as Global Heisenberg-Compensator 135
39 Carrier, Group, and Physical Readout Layers 136
40 Master Structural Theorem of the Companion 136
41 Field Equations as Physical Readout of the MM/HC Structure 139
42 Hypothesis subversion: inflow paradigm and cosmological extensions . 149
43 Current v4 Interface and Open Burdens 162
44 States, Mediators and Clock Comparisons in the 4.10 Research Line . 163
45 v4.8 Research Integration 167

© 2026 Marc Brendecke. Preprint — not peer reviewed.

H1 Baseline Release and Reader-Provenance Records . . . 171
v4.7.0 Full-Monolith Release Scope and Provenance Notice . . . 171
v4.7.0-r44 Internal Working Status . . . 171
v4.6.0 Release-Final Baseline Status . . . 172
Human-Facing Reading Layer and Notation Discipline . . . 173
Reader-Facing Proof Spine and Document Routing . . . 174
Frontmatter reader architecture . . . 178
Current Reader Navigation and Governing Map . . . 180
Extended Abstract and Reader Orientation . . . 181
H2 Complete QAM, MML and MMA Infrastructure . . . 182
Working orientation for 4.7.0 . . . 182
H3 Pre-Closure Tools and Earlier Synthesis . . . 254
Historical tool layer and pre-closure live cores . . . 254
archival stage Closure–Projection Synthesis Core . . . 272
archival stage Structural Interface Consolidation Core . . . 274
H4 Local Crystal, HC Repair and Handoff Protocols . . . 275
archival stage Expansion Core . . . 371
H5 Open-Question Genealogy and Phase IV Detail . . . 376
Open Questions . . . 377
H6 Coupled Local OP-Entry and Repair Protocol . . . 393
H7 Complete OP Transport Arguments . . . 421
Technical unfolding of the OP transport architecture . . . 421
H8 Computational Specification and Inherited Technical Tools . . . 467
H9 Computational Interpretation, Enigma Analogy, and Bomba-Style Structural Search . . . 467
Computational Interpretation, Enigma Analogy, and Bomba-Style Structural Search . . . 467
H10 Release orientation . . . 558
H11 The three-coordinate live residual interface . . . 559
H12 Paired conservative machine and compression . . . 559
H13 Tool layer . . . 560
H14 Tool stability and normalization . . . 560
H15 Residual geometry . . . 560
H16 Residual dynamics . . . 561
H17 Compressed working principles . . . 561
H18 Formal OP-entry framework . . . 562
H19 Current-state instantiation . . . 562
H20 Release conclusion . . . 563
Release note . . . 563
H21 Structural Tools for the Millennium Matrix . . . 563
H22 Heisenberg-Compensator Tools for Physical Readout . . . 564
H23 Notation for the Master Chain . . . 565
H24 Further Technical Appendices . . . 565
H25 Reintegrated Historical Proof Blocks (archival proof-memory) . . . 682
Historical witness handoff . . . 683
A.26 Provenance and Machine-Audit Layer . . . 684
A.27 Shifted QAM Migration and Governing Normal Form . . . 684
A.28 archival stage Migration Map and Backtranslation Contract for the shifted QAM Shift . . . 684
A.29 archival stage Concrete Migration Ledger and Route-Mask Examples . 688

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.30 Global QAM Hilbert-Shift Activation Candidate . . . 691
A.31 Occurrence Ledger and Shift-Audit Map . . . 695
A.32 Global Reindexed Display Layer . . . 699
A.33 Physical Readout Compatibility Check . . . 702
A.34 Governing shifted Normal Form and Release-Freeze Handoff . . . 705
A.35 archival stage Release-Final Closure Note . . . 708
A.36 Complete Reader-Hygiene and Release-History Records . . . 709
A.37 archival stage Historical Remark Triage and Reader-Hygiene Cleanup . 709
A.38 archival stage Reader Navigation Cleanup . . . 711
A.39 archival stage Notation and Symbol Hygiene Cleanup . . . 712
A.40 archival stage Table, Display, and Example Normalization Cleanup . . 714
A.41 archival stage Appendix and Historical Witness Linkage Cleanup . . . 716
A.42 archival stage Clean Reader Bundle and Hygiene-Freeze Candidate . . 718
A.43 archival stage Frontmatter, Metadata, and DOI-Record Hygiene Cleanup 720
A.44 archival stage Bibliography, Citation, and Terminal Record Hygiene Cleanup . . . 722
A.45 archival stage Glossary, Terminology, and Reader-Index Hygiene Cleanup 724
A.46 archival stage Cleanup Consolidation and Local Release-Final Preparation . . . 726
A.47 archival stage Label, Reference, and Counter Hygiene Cleanup . . . 728
A.48 archival stage Layout, TOC, and Render Hygiene Cleanup . . . 730
A.49 archival stage Source, Build, and Artifact Reproducibility Hygiene Cleanup . . . 732
A.50 archival stage Local Release-Readiness, Upload, and Final-Packaging Hygiene Cleanup . . . 734
A.51 archival stage Local Release-Final Freeze and DOI-Request Packet Hygiene . . . 736
A.52 archival stage Release-Final DOI and Audit Hygiene . . . 738
A.53 archival stage Release-Final DOI and Packaging Record . . . 740
A.54 OP Selectors, Routes and Mixed-Readout Ledgers . . . 741
A.55 OP-Facing Diagnostic Appendix . . . 741
A.56 First OP Obstruction Extraction . . . 746
A.57 Obstruction-to-Task Reduction Layer . . . 748
A.58 Task-Object Formalization . . . 750
A.59 Notation and Physics-Style Audit for OP-Facing Objects . . . 752
A.60 First Selector and Boundary-Promotion Ansatz . . . 754
A.61 First Selector Existence Reduction for OP 1/4 . . . 756
A.62 Selector Non-Emptiness and Anchor Criterion for OP 1/4 . . . 758
A.63 Anchor Adequacy and Component-Cover Criterion for OP 1/4 . . . 760
A.64 Clock-Anchored Canonical Optimal Route Principle . . . 762
A.65 Admissible Route Certificate and Optimal-Route Trigger . . . 765
A.66 Route-Certificate Tree and Arborescence Criterion . . . 768
A.67 Routing-Protocol Analogies for OP-Facing Trees . . . 771
A.68 Link-State Route-Tree Extraction . . . 774
A.69 Segment-Route Waypoint Certificates . . . 777
A.70 Traffic-Engineering and Guard-Budget Filters . . . 780
A.71 Why the New QAM Narrows the OPs Faster . . . 783
A.72 Mixed-Readout Edge Compatibility Layer . . . 786
A.73 Mixed-Readout Index Registry and Reader Atlas . . . 789
A.74 First Mixed-Readout Mode Registry . . . 792

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.75 First Selector Table and Mixed-Readout Diagnostic Pass . . . 795
A.76 Selector-to-Route Gate . . . 798
A.77 Selector-Gated Reachability Diagnostic . . . 801
A.78 Full Atlas and OP Promotion Chain . . . 803
Integrated Delta Layer: Robust Local Chart . . . 804
A.79 Position of the probe . . . 804
A.80 Asymmetric weighted photon-frame chart . . . 804
A.81 Explicit sample table . . . 805
A.82 Field-equation bookkeeping update . . . 806
A.83 Conservativity guards . . . 806
A.84 Next recommended step . . . 807
Integrated Delta Layer: Finite Atlas . . . 808
A.85 Position after . . . 808
A.86 Finite photon-frame atlas . . . 808
A.87 Explicit finite-atlas sample . . . 809
A.88 What would be needed to close OP5 . . . 809
A.89 Practical proof route after . . . 811
A.90 Conservativity guards . . . 811
A.91 Next recommended step . . . 811
Integrated Delta Layer: Tunnel Menu . . . 812
A.92 Version position . . . 812
A.93 Double-epsilon halo algebra . . . 812
A.94 Tunnel points . . . 813
A.95 Capture, invariance, and tunnel reduction . . . 813
A.96 Finite tunnel menu . . . 814
A.97 Connection to the finite atlas . . . 815
A.98 Conditional OP5 closure statement after the halo step . . . 815
A.99 Immediate next proof tasks . . . 816
A.100 Conservativity guards . . . 816
Integrated Delta Layer: Halo Channel and Response Alphabet . . . 817
A.101 Macro hygiene and version position . . . 817
A.102 Photon-frame collar chart . . . 817
A.103 Concrete halo-channel instantiation . . . 818
A.104 Exit map and exit Jacobian . . . 818
A.105 First tunnel-menu execution . . . 819
A.106 Rank-drop isolation and finite critical set target . . . 820
A.107 First executed closure implication . . . 820
A.108 What still has to be shown for an actual OP5 close . . . 821
A.109 Conservativity guards . . . 822
Integrated Delta Layer: Normal-Collar Rank Test . . . 823
A.110 Purpose and notation hygiene, declared local OP1/OP4 closure, release-final . . . 823
A.111 Finite photon-frame halo atlas . . . 823
A.112 Normal collar charts . . . 824
A.113 Selector-switch register . . . 824
A.114 Tunnel response execution . . . 825
A.115 First executable normal-form sample . . . 826
A.116 Conservativity guards . . . 826
Integrated Delta Layer: Selector and Overlap Test . . . 827
A.117 Purpose . . . 827

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.118 Photon-frame collar coordinates . . . 827
A.119 Rank-drop elimination on certified pieces . . . 828
A.120 Finite symbolic certificate . . . 829
A.121 Remaining tunnel channels . . . 829
A.122 Meaning for OP5 . . . 830
A.123 Conservativity guards . . . 830
Integrated Delta Layer: Finite Capture . . . 831
A.124 Purpose and inherited state . . . 831
A.125 Selector-switch register . . . 831
A.126 Atlas-overlap register . . . 832
A.127 First symbolic execution table . . . 833
A.128 residual theorem . . . 834
A.129 Resulting next step . . . 834
Integrated Delta Layer: Dixon-Carrier Coding . . . 835
A.130 Purpose and inherited state . . . 835
A.131 Finite route-capture certificate . . . 835
A.132 Capture tests . . . 836
A.133 First symbolic capture table . . . 837
A.134 Completeness theorem . . . 838
A.135 Audit target for the next layer . . . 838
Integrated Delta Layer: Dixon-Image Audit . . . 840
A.136 Purpose and inherited state . . . 840
A.137 Two-clock Hilbert-shift grammar . . . 840
A.138 HS(4) meta-septum leakage target . . . 841
A.139 HS(5) two-sided duplication . . . 842
A.140 HS(6) Dixon carrier and 64-bit activation code . . . 842
A.141 Compatibility with the capture theorem . . . 843
A.142 First symbolic Dixon status table . . . 844
A.143 Audit target for . . . 844
Integrated Delta Layer: Route-Exhaustiveness Audit . . . 845
A.144 Why the Dixon representation helps QAM . . . 845
A.145 Inherited coding map . . . 845
A.146 Status alphabet and admissibility test . . . 846
A.147 image table . . . 847
A.148 Mask-image theorem . . . 848
A.149 Next audit target . . . 848
Integrated Delta Layer: Integrated Residual Register . . . 849
A.150 Audit scope . . . 849
A.151 Delta-stack route menu . . . 849
A.152 Source scan facts . . . 850
A.153 Compatibility with the OP5 route surface . . . 851
A.154 Next integration rule . . . 852
A.155 Conservativity guards . . . 852
the local proof line Monolithic OP5 Promotion Audit . . . 853
the OP5 Closure Hardening and Old-Branch Mapping Audit . . . 855
A.156 Finite Perturbation Threshold Atlas for the Triolic Orbit Scan . . . 855
A.157 Finite Core-Recentering Calibration for the Triolic Orbit Scan . . . 856
A.158 Crystal–Fractal Readout Probe from Calibrated Orbit Scans . . . 858
A.159 Residual Classification Atlas for Crystal–Fractal Readouts . . . 859
A.160 Finite Hopf-Linking Readout Test . . . 860

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.161 Hopf-Stable Residual Carrier Routing . . . 861
A.162 Off-Carrier Residual Obstruction Split . . . 863
A.163 Closure Ledger and Release-Final Audit for the Orbit–Hopf Chain . . 864
A.164 v4.4 Roadmap . . . 866
A.165 Dense Hopf Carrier Refinement . . . 868
A.166 Scale Atlas Extension for Boundary Residuals . . . 869
A.167 Lens-Extreme Local Refitting . . . 870
A.168 Dual Projection Chart Audit . . . 871
A.169 v4.4 Refinement Ledger for the Orbit–Hopf Chain . . . 873
A.170 OP1/OP4 Ingress Contract . . . 874
A.171 OP1/OP4 Guard Preservation Audit . . . 875
A.172 OP1/OP4 Residual Routing Audit . . . 876
A.173 OP4 Harmless-Candidate Certificate Test . . . 878
A.174 OP1 Blocking Carrier-Density Refinement . . . 878
A.175 OP4 Open and Uncertified Residual Refinement . . . 879
A.176 v4.4 OP1/OP4 Closure-Candidate Ledger . . . 881
A.177 v4.4 Final Audit and Source-Freeze Candidate . . . 882
A.178 v4.4 OP1/OP4 Closure-Attempt Intake . . . 882
A.179 P1: OP1 Selector-Domain Promotion . . . 884
A.180 P2: OP4 Post-Residual Promotion . . . 885
A.181 P3: Coupled Selector Compatibility . . . 886
A.182 P4: No-New-Debt Promotion . . . 887
A.183 P5: Finite-to-Declared-Grammar Closure Guard . . . 888
A.184 v4.4 Closure Summary and Release-Candidate Audit . . . 889
A.185 v4.4 Release-Final Preflight and Metadata Intake Gate . . . 890
A.186 v4.4 Clean RC Consolidation and Release-Readiness Audit . . . 891
A.187 v4.4 Release-Final DOI and Public Upload Audit . . . 893
A.188 OP5 Local-to-Global Promotion Intake Layer . . . 893
A.189 HC/MM Coupled-Debt Witness Search Layer . . . 894
A.190 OP5 Promotion Residual Classifier . . . 896
A.191 Enlarged Witness-Family Decision . . . 897
A.192 Enlarged OP5 Witness-Family Execution . . . 898
A.193 Two-Sided Tensor Bridge License Test . . . 899
A.194 OP5 Two-Sided Promoted-Closure Theorem . . . 900
A.195 No-Overclaim and Route-to-Global Boundary Audit . . . 901
A.196 OP5 / OP1–OP4 Compatibility Audit . . . 902
A.197 Two-Sided Tensor Grammar Consolidation . . . 903
A.198 Compatibility Theorem and No-Reopen Summary . . . 904
A.199 Route-to-Global Boundary Problem Statement . . . 905
A.200 Reader-Facing Notation Cleanup . . . 906
A.201 Global-Boundary Witness-Menu Design . . . 907
A.202 Grammar Exhaustion Witness Execution . . . 908
A.203 Declared Readout Invariance Witness Execution . . . 909
A.204 Boundary Transport Witness Execution . . . 910
A.205 No-New-Debt Extension Witness Execution . . . 910
A.206 Functorial Comparison Stability Witness Execution . . . 911
A.207 Boundary-Gate Synthesis Audit . . . 912
A.208 Compressed Editorial and Release-Provenance Layer . . . 914
A.209 Bounded-Readout Scope and T2 Boundary . . . 914
v4.6.0 Integrated Bounded-Readout and T2 Scope-Boundary Appendix . . . 914

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.210 MM/HC Field-Interface and C4/C5 Development . . . 916
A.211 Content-State Refresh and v4.7 Handoff Preflight . . . 916
A.212 Theory-Lift Gate and MM/HC Exterior–Mixed Field-Equation Program 916
A.213 Finite MM/HC Exterior–Mixed Term Extraction and Classification . . 919
A.214 v4.7.0-r03 Selector/Leakage Bound Pass . . . 922
A.215 v4.7.0-r04 finite witness-fit lowering test for C_5 . . . 924
A.216 v4.7.0-r05 C5 lowering iteration and C4 mixed-leakage separation . . . 926
A.217 v4.7.0-r06 C5 convergence/collar decision and C4 leakage-source inventory 929
A.218 v4.7.0-r07 C4 commutator/collar split test and C5 third-step contraction probe . . . 932
A.219 v4.7.0-r08: Collar-Seam Residual Bound and C5 Induction-Candidate Test . . . 934
A.220 v4.7.0-r09: C4 Bounded-Seam Absorption and C5 Operator-Admissibility Check . . . 937
A.221 v4.7.0-r10 Mixed MM/HC Seam Resolver and C5 Asymptotic Collar Packaging . . . 941
A.222 v4.7.0-r11 Residual Mixed-Seam Micro-Split and Bounded-Collar Limit Guard . . . 943
A.223 v4.7.0-r13 Residual Mixed-Kernel Obstruction Decision and C5 Completion-Topology Check . . . 945
A.224 v4.7.0-r13 Route-Stable Mixed-Kernel Normal Form, Overlap Relay, and Completion-Interface Audit . . . 948
A.225 v4.7.0-r14 Source-Only Finite Valley-Front Chart Scan . . . 953
A.226 v4.7.0-r15 Source-Only: Zero-Clock Phase-Window Extraction and Relay-Chain Stitching . . . 955
A.227 v4.7.0-r16 Source-Only: Shadow-Basin Lift Test for $V6$ . . . 958
A.228 v4.7.0-r17 Source-Only Atlas-Gap Bridge Test and Guarded V9 Micro-Lift . . . 960
A.229 v4.7.0-r18 Source-Only Pass: V0 Boundary-Front Obstruction Inventory and Finite-Chain Guard Audit . . . 963
A.230 v4.7.0-r19 Source-Only Pass: Boundary-Source Chart Admissibility Test and C4 Guard-Preserving Audit . . . 965
A.231 v4.7.0-r20 Source-Only Boundary-Reservoir Declaration Test . . . 968
A.232 v4.7.0-r21: Boundary-Extended Chain Seam Audit and Two-Sided Source Typing . . . 970
A.233 v4.7.0-r22 source-only: boundary-source calculus extraction and typed relay closure guard . . . 972
A.234 v4.7.0-r23 Boundary-Source Stability Window and Relay-Guard Robustness Scan . . . 974
A.235 v4.7.0-r24 Boundary-Source Robustness Theoremlet and Residual Frontier Map . . . 977
A.236 v4.7.0-r25 Source-Only Frontier Minimizer and Guard-Stable Atlas Compression . . . 980
A.237 v4.7.0-r26 compressed atlas transition audit and frontier-to-field-equation handoff . . . 983
A.238 v4.7.0-r27: MM/HC Seed-Equation Ansatz Extraction from the Guarded Atlas Handoff . . . 986
A.239 v4.7.0-r28: Seed-Ansatz Consistency Scan and MM/HC Coefficient Typing . . . 988

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.240 v4.7.0-r29 Typed Seed Residual Minimizer and Selector-Window Calibration . . . 991
A.241 v4.7.0-r30 Typed MM/HC Seed Normal Form and Admissibility Certificate . . . 995
A.242 v4.7.0-r31 Source-Only: Typed Coefficient Skeleton and Local Seed Functional Packaging . . . 997
A.243 v4.7.0-r32 Source-Only Local Seed Functional Sanity Table and Coefficient-Domain Guard . . . 999
A.244 v4.7.0-r33 Source-Only: Local Seed Functional Parameter Sweep and Comfort-Margin Calibration . . . 1000
A.245 v4.7.0-r34: Comfort-Calibrated Seed Selector and Guard-Stable Handoff Summary . . . 1003
A.246 v4.7.0-r35 Source-Only: Guarded MM/HC Seed-to-Interface Map and No-Action Audit . . . 1004
A.247 v4.7.0-r36 Source-Only Interface Residual Table and MM/HC Map Consistency Scan . . . 1007
A.248 v4.7.0-r37 Local MM/HC Interface Certificate and Guard-Stable Residual Packaging . . . 1008
A.249 v4.7.0-r38 Certificate-to-Local-Field-Interface Precheck and No-Promotion Audit . . . 1010
A.250 v4.7.0-r39 Local Field-Interface Skeleton and Admissibility-Only Operator Tags . . . 1012
A.251 v4.7.0-r40: Admissibility-Only Operator Composition Table and Guard Leak Scan . . . 1014
A.252 v4.7.0-r41 Source-Only: Guarded Composition Normal Form and Local Interface Closure Ledger . . . 1017
A.253 v4.7.0-r42: Local Monitor Certificate Compression and Release-Candidate Guard Audit . . . 1019
A.254 v4.7.0-r43 Source-Only Metadata Freeze Check and Local Certificate Handoff . . . 1020
A.255 v4.7.0-r44 Source-Only Handoff Integrity Scan and Local Certificate Index . . . 1022
A.256 v4.7.0-r45 Source-Only Local Certificate Reader-Routing Map and Next-Branch Queue . . . 1023
A.257 v4.7.0-r46 Source-Only Local Certificate Dependency Graph and Reader-Route Cross-Check . . . 1025
A.258 v4.7.0-r47 Source-Only Dependency-Order Theoremlet and Branch-Launch Guard Preflight . . . 1027
A.259 v4.7.0-r48 Source-Only Branch Queue Selection and Work-Package Split 1028
A.260 v4.7.0-r49 Source-Only Interface-Slot Glossary and Two-Sided Typing Recap . . . 1030
A.261 v4.7.0-r50 Source-Only Addendum: Admissibility-Only Operator-Domain Refinement . . . 1032
A.262 v4.7.0-r51 Source-Only Local Pre-Field Interface Diagram . . . 1035
A.263 v4.7.0-r52 Source-Only No-Action/No-Promotion Audit Refresh . . . 1037
A.264 v4.7.0-r53 Source-Only Guarded Local-Interface Consolidation and Post-W1–W4 Branch Checkpoint . . . 1039
A.265 v4.7.0-r54 Source-Only Local-Interface Theoremlet Extraction and Next-Branch Selector Preflight . . . 1041
A.266 Guarded Post-r54 Solve Chain . . . 1043

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.267 Compressed Post-r54 Guarded Solve Chain: r55–r80 1043
A.268 Matter, Photon and Tachyon Frame-Typed Endpoint 1043
A.269 r81 Matter-frame epsilon terminality and Planck/information scale split 1043
v4.7.0-r84 Post-r83 Full-Monolith Integration and Release-Scope Refresh 1050
A.270 v4.8.0 Release Layers 1051
Archived Statements of the Main Body 1052
v4.7.0-r85 Source-Only Citation Repair and Unity-Theorem Scope Separation . 1059
v4.7.0-r86 Source-Only: C5 Asymptotic-Collar Bookkeeping Consolidation . . . 1061
v4.7.0-r87 Source-Only: Photon-Clock Selector-Window Calibration 1064
v4.7.0-r88 Source-Only: Admissibility Test for Non-Collar Routes to Finite C5 Discharge 1067
v4.7.0-r89 Source-Only: r87 Correction — Recorded V9 Data and Executed Micro-Lift 1070
v4.7.0-r90 Source-Only: Reconstruction Attempt for the Bound Delta gamma $<$ 1/5 1072
v4.7.0-r91 Source-Only: Q4 Executed — Why beta 4 Cannot Be Spent 1075
v4.7.0-r92 Source-Only: Sharpened Reconstruction of Delta gamma $<$ 1/5 . . . 1078
v4.7.0-r93 Source-Only: The Two-Sided Typing Read as a Routing Grammar . . 1080
v4.7.0-r94 Source-Only: Provenance of the C5 Contraction Factor 3/4 1083
v4.8.0-r01 Source-Only: Planck-Readout Metric Layer 1086
v4.8.0-r02 Release Integration and Scope Refresh 1088
A.271 v4.8.0 Post-Release Layers 1088
v4.8.0-r03 Source-Only: Dimensional and Numerical Audit of the Physical-Constant Layer 1089
v4.8.0-r04 Source-Only: Decoupling from the Riemann Source and Standalone Sphaera Transformations 1094
v4.8.0-r05 Source-Only: Corrected Explicit Formula in the Field-Equation and Theta–R Layer 1098
v4.8.0-r06 Source-Only: Signature of the Biquaternion Kernel and the Complex-Vector Reading 1101
v4.8.0-r07 Source-Only: Sector Assignment and Duality Angle in the Complex-Vector Reading 1104
v4.8.0-r08 Source-Only: Test of Prediction P1 (Photon-Frame Unfolding) 1106
v4.8.0-r09 Source-Only: Tests of P2 and P3, and Re-Classification of Earlier Results 1108
v4.8.0-r10 Source-Only: The Koide Structure in the Complex-Vector Reading . . 1111
v4.8.0-r11 Source-Only: Pre-Registered Test of the Phase Relation delta = K/3 1113
v4.8.0-r12 Source-Only: Pre-Registered Neutrino Tests, Octonionic Flavour Group, PMNS Reclassification 1115
v4.8.0-r13 Source-Only: JUNO 2026 Re-Check, Associator, Colour–Generation Assignment 1118
v4.8.0-r14 Source-Only: Separating Colour and Generation 1120
v4.8.0-r15 Source-Only: Octonionic Hopf Fibration as Carrier of the Triality Slots 1122
v4.8.0-r16 Source-Only: Pre-Registered Test of Masses from Invariants on the Triple Space 1124
v4.8.0-r17 Source-Only: Time Direction of the Cosmological Fractions 1126
v4.8.0-r18 Source-Only: Three-Frame Typing of the Frame Offset and Correction of r17 1129
v4.8.0-r19 Source-Only: Frame Cubes, Transition Group and Holonomy 1132
v4.8.0-r20 Source-Only: The Tachyon Frame, Complex Rapidity and the Holonomy Test H13 1135

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.272 v4.9.0 Post-Release Layers 1137
v4.9.0-r01 Source-Only: The Photon Frame as Neutral Axis and the Conversion of Constants 1138
v4.9.0-r02 Source-Only: The Round Trip about the Photon and the Umkehrwalze 1141
v4.9.0-r03 Source-Only: The Compensator as a Two-Arm Interferometer 1145
v4.9.0-r04 Source-Only: The Compensator as a Matrix on the Electric–Magnetic Doublet 1148
v4.9.0-r05 Source-Only: The Millennium Matrix on the Doublet, a Third Axis near Rotating Black Holes, and a Photon from Matter and Tachyon . . 1151
v4.9.0-r06 Source-Only: The Tachyon Hit Chain, its Acceleration, and the Higgs Field 1155
v4.9.0-r07 Source-Only: The Cone Angle and the Weak Mixing Angle 1159
v4.9.0-r08 Source-Only: Frame Drift, Atomic Clocks and the Hubble Tension . . 1162
v4.9.0-r09 Source-Only: What the Local Frame Factor Can Be 1166
v4.9.0-r10 Source-Only: A Reading of the C5 Factor 3/4 and a Class of Shape-(iii) Witnesses 1169
v4.9.0-r11 Source-Only: A Minimal Dynamics for Reading B — Kicked Duality Rotation 1172
v4.9.0-r12 Source-Only: A Reconstruction of the Bound Delta gamma $< 1/5$ in Place of the Missing r65 1174
v4.9.0-r13 Source-Only: The Time Scale of the Kicked Duality Dynamics 1177
v4.9.0-r14 Source-Only: Why Six Kicks per Beat — Readout Directions from the Triole and the Umkehrwalze 1179
v4.9.0-r15 Source-Only: Consolidated Register of v4.9.0-r01 to r14 and the Ways Out of r17 after r08 1181
v4.9.0-r16 Source-Only: Author Decisions on the Open Readings of r15 1184
v4.9.0-r17 Source-Only: Entropy between the Frames, the Photon as Information Singularity, and a Scale Dimension 1186
v4.9.0-r18 Source-Only: The Old Conversions on the Scale Line, and How Ways A and C Can Hold Together 1189
v4.9.0-r19 Source-Only: The Class Split Applied to the Main Body 1192
v4.9.0-r20 Source-Only: A Provenance Hypothesis for Delta gamma $< 1/5$, the Photon-Frame Clock Gain at the Hopf Point 1195
v4.9.0-r21 Source-Only: The Clock Bound Frame by Frame, and a Derivation of $1/5$ 1198
v4.9.0-r22 Source-Only: The Frame Rule as a Standing Convention, Applied to the Main Statements 1201
v4.9.0-r23 Source-Only: Convention F Applied to the Main Body 1204
v4.9.0-r24 Source-Only: Correction of the Three Discrepancies of r23 as Single-Frame Artefacts 1207
v4.9.0-r25 Source-Only: Revision of the Main Body 1210
v4.9.0-r26 Source-Only: Pre-Registered Test of Degree-Six Potentials on the Triple Space 1212
v4.9.0-r27 Source-Only: A Dynamics for the Kicked Duality Rotation, Rotation plus Compensator Measurement 1214
v4.9.0-r28 Source-Only: Who Measures? The Tachyon as Measuring Device, Self-Stabilisation of the Field, and a Correction of the Tachyon Map . 1217
v4.9.0-r29 Source-Only: Where the Tachyon Reservoir Could Come From 1220
v4.9.0-r30 Source-Only: The Compensator Avoids the Classical One-Half, and Why the Exchange Is Coherent 1222

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r31 Source-Only: The Markov Cascade as a Test of Self-Stabilisation, and a Status Note to the Dark-Energy Theorem . . . 1225
v4.9.0-r32 Source-Only: Pre-Registered Test of Real-Part Invariants on the Triple Space . . . 1227
v4.9.0-r33 Source-Only: Left-Invariant Labels on the Triple Space and a Pre-Registered Test . . . 1229
v4.9.0-r34 Source-Only: Charges of One Generation from the Dixon Storage . . 1232
v4.9.0-r35 Source-Only: Inner and Outer Complex Unit in the Colour Assignment 1234
v4.9.0-r36 Source-Only: Frame Rule for Complex Units, and the Origin of the Colour–Generation Double Use . . . 1236
v4.9.0-r37 Source-Only: Triality as Self-Traversal, and the Periods of the Structure 1239
v4.9.0-r38 Source-Only: The Frame Triole as the Reflection Side of Triality . . . 1242
v4.9.0-r39 Source-Only: Chains of Different Length under Hilbert-Shift Zoom and Mixed Readouts . . . 1244
v4.9.0-r40 Source-Only: The Two-Sided Contraction across Frames, and the Weak SU(2) . . . 1247
v4.9.0-r41 Source-Only: Distributing the Roles of the Quaternion Factor; Hypercharge and Electric Charge . . . 1250
v4.9.0-r42 Source-Only: A Compensator of the Millennium Matrix as Chirality Projector . . . 1253
v4.9.0-r43 Source-Only: The Choice of the Chirality Projector Is Fixed by Anomaly Cancellation . . . 1256
v4.9.0-r44 Source-Only: Mass Terms as Frame Couplings; Breaking Stages for Masses and Mixing . . . 1258
v4.9.0-r45 Source-Only: Chirality Breaks the Reflection of the Triality Form . . 1261
v4.9.0-r46 Source-Only: Mixed Readouts and the Breaking of Triality . . . 1263
v4.9.0-r47 Source-Only: A Generation Mask from the Triality-Covariant Chirality 1265
v4.9.0-r48 Source-Only: The Charge Sectors Derived from the Octonion Product in Lean . . . 1267
v4.9.0-r49 Source-Only: The Axis e7 Carries B–L, Not the Electric Charge . . . 1269
v4.9.0-r50 Source-Only: The Mask of r47 Does Not Produce the Observed Mixing 1271
v4.9.0-r51 Source-Only: The Axis e7 and the Frames . . . 1273
v4.9.0-r52 Source-Only: Exact Coupling Forms of the Mask and the Mixing . . . 1275
v4.9.0-r53 Source-Only: Couplings from the Triality Form . . . 1277
v4.9.0-r54 Source-Only: One epsilon per Sector . . . 1279
v4.9.0-r55 Source-Only: A Readout per Particle Kind from the Subgroups of A4 1281
v4.9.0-r56 Source-Only: Calibrating the Readout Layer . . . 1283
v4.9.0-r57 Source-Only: Trimaximal Mixing with the Mask as a Correction . . . 1285
v4.9.0-r58 Source-Only: The CP Phase of the Corrected Trimaximal Reading . . 1287
v4.9.0-r59 Source-Only: The Correction Strength from a Structural Value 1289
v4.9.0-r60 Source-Only: The Correction Strength as the Rapidity of the Matter Frame . . . 1291
v4.9.0-r61 Source-Only: A Frame Audit of the Open Problems OP1 to OP5 . . . 1293
v4.9.0-r62 Source-Only: Status Notes after the Frame Audit of the Open Problems 1295
v4.9.0-r63 Source-Only: The Frame Offset as a Ratio of Two Horizon Countings 1296
v4.9.0-r64 Source-Only: A Frame Audit of the Gravity Statements under the Counting Reading . . . 1299
v4.9.0-r65 Source-Only: Status Notes after the Frame Audit of the Gravity Statements . . . 1301

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r66 Source-Only: A Correction to r64 and the Local Frame Factor under the Counting Reading 1302
v4.9.0-r67 Source-Only: Einstein's "Greatest Blunder" under the Counting Reading 1304
v4.9.0-r68 Source-Only: An Independent Recheck of H47 and r63 to r67 1306
v4.9.0-r69 Source-Only: The OP5 Research Series of 25 to 27 October 1308
v4.9.0-r70 Source-Only: The 4.10 Consolidation in the Monolith, a Correction to r69, and Two Tests of a Common Dynamics 1312
v4.9.0-r71 Source-Only: The Structure of the OP5 Comparison Symbol, an Exact Polar Gauge Symmetry, and Two Tests 1315
v4.9.0-r72 Source-Only: The theta Gauge of the OP5 Comparison Symbol, a Cartan Test, and Four Hidden Conditions 1318
v4.9.0-r73 Source-Only: Four Hidden Sixth-Order Conditions of the Gauged OP5 Comparison System 1321
v4.9.0-r74 Source-Only: Consistent Data, Spencer Bounds and the Precision Limit of the Extended OP5 Symbol 1324
v4.9.0-r75 Source-Only: Renaming the Millennium Matrix to Compensator Matrix 1326
v4.9.0-r76 Source-Only: Correction to r75 — the Earlier Name "Layer-Transport Matrix" 1328
v4.9.0-r77 Source-Only: High Precision, Formal Integrability at the Germ and the Covariance of the OP5 Comparison System 1330
v4.9.0-r78 Source-Only: Covariance of the OP5 Comparison System at Value Level and the Stabilizer of the Gauge 1334
v4.9.0-r79 Source-Only: Gauge Independence of the Four Hidden Conditions of the OP5 Comparison System 1336
v4.9.0-r80 Source-Only: The Background of the OP5 Comparison System Is Not Sourced by Its Spinors; Closing the Comparison-Model Branch 1338
v4.9.0-r81 Source-Only: A Joint Action W1, the Matter–Tachyon Pairing, and the Search for a Vacuum Term 1341
v4.9.0-r82 Source-Only: The Higgs Sector of W2, the Stability of the Matter–Tachyon Cancellation, and a Torsion Pairing Channel 1345
v4.9.0-r83 Source-Only: Quantum Decay of the Matter–Tachyon Vacuum, the Tachyon Image of an Action, and a Correction to r82 1348
v4.9.0-r84 Source-Only: The Tachyon Part as a Second Reading of the Same Field, the Complete Complex Boost, and the Expanding Background . 1351
v4.9.0-r85 Source-Only: Masses and Couplings in the Second Reading, the Epoch tau*, and the Big Bang as a CPT Transition 1354
v4.9.0-r86 Source-Only: Independent Checks with a Second System: the Big-Bang Admixture and the Field Equations of W1 1357
v4.10.0-r01 Source-Only: The Full Einstein Equation of W1 and the Sign of the Gravitational Action 1360
v4.10.0-r02 Source-Only: Asymptotics of the Admixture at the Big Bang 1363
v4.10.0-r03 Source-Only: Fermions and Bosons at the Big Bang, and the Vacuum Term under G2 1366
v4.10.0-r04 Source-Only: The Vacuum Term with a Tachyon Branch, and the Bosons Created at the Big Bang 1369
v4.10.0-r05 Source-Only: Gravitons at the Big Bang, a Review from the Core, and the Readout of Lambda 1372
v4.10.0-r06 Source-Only: Semiclassical Einstein Equations from Relative Entropy (Dorau–Much) and the Readouts of the Companion 1375

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.10.0-r07 Source-Only: The Relative-Entropy Chain in the Tachyon Frame, and a Correction to r06.5 1378
v4.10.0-r08 Source-Only: The Assumptions of the Companion and the Dependence of the Open Predictions 1382
v4.10.0-r09 Source-Only: The Counting Rule per Frame, Its Singular Position (H98) and a Principle 1385
v4.10.0-r10 Source-Only: The Bridge of the Counting Principle, and the Counting Rule as a Frame-Dependent Area Quantum 1387
v4.10.0-r11 Source-Only: Integration of the OP5 Source Reconstruction Made with Another Language Model 1390

© 2026 Marc Brendecke. Preprint — not peer reviewed.

1 Premise Audit and Scope

The following qualifications apply before reading the inherited core. The original assertions and proofs are retained for audit, not silently re-certified. This section records conditional replacements for specific known issues; it does not claim a complete scientific validation of the monolith or repair every downstream use of the older assertions.

1.1 MM/HC Spectral Identification

Commutation with an orthogonal projector preserves its image but does not make that image a single eigenspace. Absence of nontrivial Jordan chains also does not imply a single eigenvalue. For example, $M = \operatorname{diag}(1,2,3)$ and $C = \operatorname{diag}(1,1,0)$ commute while M is not scalar on $\operatorname{im} C$. Even scalar action does not identify the full spectral projector: $M = \operatorname{diag}(2,2,3)$ and $C = \operatorname{diag}(1,0,0)$ omit one direction of the eigenvalue-two eigenspace. Thus the inherited labels `lem:diagonal-traversal-scalar-action` and `thm:hc-spectral-projector` must not be used without the missing support premises.

Proposition 1.1 (A sufficient conditional MM/HC identification)**.** *Let M be a bounded self-adjoint operator on a complex Hilbert space, let C be a nonzero orthogonal projector, and suppose for a real λ_0 that*

$$MC = CM = \lambda_0 C, \qquad \ker(M - \lambda_0 I) \cap \ker C = \{0\}.$$

Then $\operatorname{im} C = \ker(M - \lambda_0 I)$, and C is the orthogonal eigenprojection for that eigenvalue. These conditions are sufficient hypotheses, not a derived identification of a native MM.

Proof. The scalar relation gives the image-to-kernel inclusion. Conversely, if $Mx = \lambda_0 x$, commutation implies that $(I-C)x$ lies in both $\ker(M-\lambda_0 I)$ and $\ker C$. The second assumption forces $(I - C)x = 0$, so $x = Cx$. Orthogonality identifies the projection onto the resulting closed eigenspace. □

One must still supply the realized operator, domain, scalar action and complement exclusion. A representation label, an invariant carrier or an audit token does not supply these hypotheses.

1.2 OP1: Inverse Diagnostic and Fixed Exponent

For $g > 1$, $0 < \beta < 1/\sqrt{3}$ and $\gamma_1 > 0$, set

$$A(g,\beta) = \frac{\gamma_1}{4\pi|\log(\beta\sqrt{3})|}, \qquad n(g,\beta) = \frac{\log A(g,\beta)}{\log g}.$$

The relation $g^{n(g,\beta)} = A(g,\beta)$ is then an identity. Substituting $\beta(g) = \sqrt{(g-1)/(g+1)}$ leaves an identity on $1 < g < 2$; it does not predict g. The inherited inverse-exponent and self-consistency passages must be read with this restriction, not as an independent arithmetic derivation of the frame factor.

Proposition 1.2 (Unique root under an independently fixed ansatz)**.** *Assume the exponent five and prefactor $1/\sqrt{3}$ independently, and put $a = \gamma_1/(4\pi)$ with $0 < a < 27/20$. The equation*

$$f(g) = \tfrac{1}{2}\log 3 + \tfrac{1}{2}\log\frac{g-1}{g+1} + \frac{a}{g^5} = 0$$

has exactly one root in $(1,2)$.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. Here $f'(g) = 1/(g^2-1) - 5a/g^6$. For $t = g^2 > 1$,

$$t^3 - \tfrac{27}{4}(t-1) = (t-\tfrac{3}{2})^2(t+3) \geq 0.$$

Consequently $g^6/(g^2-1) \geq 27/4 > 5a$, so $f' > 0$. Also $f(1+) = -\infty$ and $f(2) = a/32 > 0$. Continuity and strict monotonicity give existence and uniqueness. □

At the stated first-zero input $\gamma_1 = 14.134725141734694$, the model root is $g \simeq 1.863662249889$ and $\beta \simeq 0.549175395418$. This is a conditional scalar-model calculation. The exponent, prefactor and prime-lattice interpretation still need independent justification; the result does not close OP1/4.

1.3 OP5: Detector Evidence and Full Balance

The full Θ sum, its regularization, endpoint conventions and remainder control are separate obligations. An empty truncated frequency sum does not make the full oscillatory sum vanish. The inherited quantitative passages are therefore retained as audit material, not certified by their relocation or compilation. The specific conjecture for the full sum (Proposition 42.33) has since been evaluated with the corrected explicit formula in layer v4.8.0-r05 (§A.271): the left-hand side is -0.1097 ± 0.0009 against the required 0.893, so that conjecture is false. This decides the conjecture, not the general obligation above; OP5 Step (B) has no valid form in the declared reading.

The later C5 fit score is another interface. Applying a transfer or contraction estimate to it requires the native evaluated boundary, exterior and mixed residual blocks, their fixed norm conventions, their guards and their calibration. Synthetic residual vectors or a fitted contraction constant do not recover this missing dictionary. A sufficient bound such as 3/4 is not a measured universal physical threshold, and a positive residual below a detector tolerance is not an exact all-readout zero.

1.4 Carried Corrections from the v4.8.0 Line

The published v4.8.0 line and the layers recorded after it are appended in the support route (§A.270 and §A.271). Two of their repairs were made inside the full monolith and are not visible in the preserved fragments, which keep the inherited wording. They apply here as follows.

Unity theorem. The preserved Theorem 12.1 lists the gauge structure $U(1) \times SU(2) \times SU(3)$ as item (vi) of its unitary-equivalence list, and its proof structure cites Theorem 18.3 as (vi). Under layer r85 (§A.270) the equivalence list ends at item (v). A Lie group is not a Hilbert-space representation of the Weyl algebra, so no Stone–von Neumann equivalence is asserted for it; the gauge statement stands as a separate consequence, and Theorem 18.3 is unchanged.

Stone–von Neumann citation. The proof structure of the unity theorem and one passage on the irreducible representation of the Weyl algebra cite the Stone–von Neumann step as Theorem 3.2 of [1]. In that source Theorem 3.2 is the functional equation; the step is Step 3 of Lemma 13.2, resting on Proposition 13.1 (r85). Since layer v4.8.0-r04 (§A.271) the Riemann source is no longer used as a proof source, the Stone–von Neumann justification of the unity theorem is withdrawn, and the equivalence of items (i)–(v) is carried as a conjecture.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Registers. Statements of the inherited core are to be read with the registers of layer v4.8.0-r09 (§A.271) and their extensions in r17–r20. In particular no prediction of the inherited falsifiability corollary stands as confirmed. For the frame layer: the frame factor used to convert constants between matter and photon frame is the matter–tachyon cross term (§A.271), and in the complex-vector reading no real Lorentz transformation relates the matter, tachyon and photon sectors (§A.271, §A.271). This agrees with the present candidate's statement that no frame assignment is claimed.

2 Koide and Local Kernel Construction

This section supplies explicit arguments for the two proposed compression sites. It supersedes the claim that cyclic symmetry alone proves the inherited Koide theorem. It also constructs a relative kernel model, without declaring the full prospective native zero-kernel layer active.

2.1 The Exact Additional Koide Condition

Let $x_i = \sqrt{m_i} \geq 0$, $x \neq 0$, and write

$$A = \frac{x_0 + x_1 + x_2}{3} > 0, \qquad u = A(1,1,1), \quad d = x - u, \quad \sum_i d_i = 0.$$

The cyclic permutation preserves the one-dimensional symmetric line and its two-dimensional orthogonal plane. It preserves their norms separately; it does not prescribe their ratio.

Proposition 2.1 (Koide as an equal-norm condition)**.** *For these nonnegative principal roots,*

$$K = \frac{\sum_i x_i^2}{(\sum_i x_i)^2} = \frac{1}{3} + \frac{\|d\|^2}{9A^2}.$$

Consequently $K = 2/3$ if and only if $\|d\|^2 = \|u\|^2 = 3A^2$.

Proof. Since $u \perp d$, $\|x\|^2 = 3A^2 + \|d\|^2$, while $(\sum_i x_i)^2 = 9A^2$. Division proves both assertions. □

Thus Cauchy–Schwarz supplies only $K \geq 1/3$. For example, the cyclic orbit of $x = (1,2,3)$ preserves both sums used in the inherited argument, but gives $K = 14/36 = 7/18$, not $2/3$. Even the fully symmetric choice $x = (1,1,1)$ gives $K = 1/3$. The missing model obligation is an independently derived equal-norm law, not a normalization of the Cauchy–Schwarz inequality by a count of sectors.

Proposition 2.2 (A sufficient HC-shaped mass-root realization)**.** *In a declared real three-root representation, let $e = (1,1,1)/\sqrt{3}$, let J_m be an orthogonal involution, and put $C_m = (I + J_m)/2$. Suppose $x = \alpha C_m e$ is nonzero and componentwise nonnegative, with $\alpha > 0$. For $h = \langle e, J_m e \rangle$, one has $h > -1$ and*

$$K = \frac{2}{3(1+h)}.$$

Hence Koide holds precisely when $\langle e, J_m e \rangle = 0$.

Proof. An orthogonal involution is self-adjoint, so C_m is an orthogonal projector. Thus $\|C_m e\|^2 = \langle e, C_m e \rangle = (1+h)/2 > 0$ and $(\sum_i x_i)^2 = 3\alpha^2(1+h)^2/4$. Dividing $\|x\|^2 = \alpha^2(1+h)/2$ by this denominator gives the formula. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

This condition is realizable: take $v = (2, -1, -1)/\sqrt{6}$, a unit vector orthogonal to e with $e + v$ positive. Let J_m exchange e and v, and let it act as $+1$ on their orthogonal complement. Then $C_m e = (e + v)/2$ has equal symmetric and orthogonal norms. The direction of v still determines a free phase. Conversely, $J_m = I$ is an equally valid orthogonal involution but gives $K = 1/3$. Therefore HC projection axioms alone do not force Koide. The representation of the native HC on mass roots, positivity and the orthogonality condition must be derived rather than selected to fit.

Proposition 2.3 (Radius and phase in the three-root parametrization)**.** *Every nonzero nonnegative three-root vector admits*

$$x_k = A\big(1 + r\cos(\theta + 2\pi k/3)\big), \qquad r \geq 0,$$

with $K = 1/3 + r^2/6$. The condition $K = 2/3$ is precisely $r = \sqrt{2}$; it does not fix A or θ.

Proof. The vectors $(\cos(2\pi k/3))_{k=0}^2$ and $(\sin(2\pi k/3))_{k=0}^2$ form an orthogonal basis of the zero-sum plane, each with squared norm $3/2$. Expressing d/A in that basis gives coefficients $r\cos\theta$ and $-r\sin\theta$. Therefore $\|d\|^2 = (3/2)A^2r^2$ and the previous proposition gives the formula for K. If $r = 0$ the phase is arbitrary. □

The root formula must retain $1 + r\cos(\theta + 2\pi k/3) \geq 0$. Squaring a negative bracket and then calling it a principal square root changes the denominator to a sum of absolute values; the displayed identity for the physical K no longer follows. The phase ansatz $\theta = \pi - \sqrt{2}\arcsin\beta$ also does not give mass degeneracy at $\beta = 0$: with $r = \sqrt{2}$ it gives

$$(m_0, m_1, m_2)/A^2 = \big(3 - 2\sqrt{2},\ 3/2 + \sqrt{2},\ 3/2 + \sqrt{2}\big),$$

and the first signed-root bracket is negative. A degenerate nonzero three-mass limit has $r = 0$ and $K = 1/3$, so it cannot be obtained by changing only the phase at fixed Koide radius.

Proposition 2.4 (Cyclic-equivariant linear maps preserve symmetric inputs)**.** *Let P cyclically permute three real coordinates. If a linear map T satisfies $TP = PT$, then $T(A(1, 1, 1))$ remains on the symmetric line. It cannot produce a nondegenerate root hierarchy from that input.*

Proof. $Pu = u$ implies $P(Tu) = T(Pu) = Tu$. The fixed space of P is exactly the line spanned by $(1, 1, 1)$. □

A hierarchy mechanism must therefore identify an asymmetric source, a symmetry-breaking selection or a non-equivariant coupling; cyclic symmetry alone is insufficient. A derivation of the phase law and the equal-norm condition remains open. Using only the fixed mass inputs quoted in the inherited text, the phase diagnostic is about $132.73279°$, while the stated $\beta = 0.5493$ ansatz gives $132.87981°$. With A fixed by the quoted roots, the respective mass errors are approximately -12.80%, $+1.20\%$, -0.0666%. These reproduce a historical-input comparison, not a new empirical measurement or a physical derivation. The inherited `thm:koide-triolic` and dependent phase-prediction wording must be read with these limitations.

2.2 A Quotient Kernel on a Declared Fragment

Fix a nonempty *set* A_F of admitted payloads for one declared fragment, with tagged embeddings $j_t(a) = \langle t, a\rangle$ and payload projection $p_t(j_t(a)) = a$. Here $t = 0$ denotes the retained pre-shift display and $t = 1$ its inhabited shifted state display. This local notation does not

© 2026 Marc Brendecke. Preprint — not peer reviewed.

undo the later shifted normal form. Assume a supplied equivalence relation E_F on A_F expressing the restricted closure-class schema. Its native interpretation is input data, not something derived from the tag.

Proposition 2.5 (Choice-free semantic quotient)**.** *The set $Q_F = A_F/E_F$ and map $q_F(a) = [a]_{E_F}$ are defined in ZF. They satisfy $q_F(a) = q_F(b)$ if and only if aE_Fb. If a total closure-relevant readout $f : A_F \to Y$ is constant on E_F-classes, there is a unique $\bar{f} : Q_F \to Y$ with $f = \bar{f} \circ q_F$.*

Proof. Each class is a subset of A_F defined by Separation, and their collection is a set by Replacement. Reflexivity, symmetry and transitivity give the equality criterion for classes. Define

$$\operatorname{graph}(\bar{f}) = \{\, ([a]_{E_F}, f(a)) : a \in A_F \,\}.$$

Class constancy makes this a function, and surjectivity of q_F gives uniqueness. No representative or choice function is required. □

This quotient packages the supplied closure relation; it does not compute it, prove a native closure claim or guarantee a cheaper algorithm. The quotient kernel and the raw payload have distinct roles: $j_t \circ p_t$ reconstructs the syntactic state, whereas q_F may identify different payloads and cannot reconstruct each original state by itself.

Proposition 2.6 (Finite representative normal forms)**.** *If A_F is finite with a declared injective integer coding, let $N_F(a)$ be the least-coded member of $[a]_{E_F}$. Then $N_F : A_F \to A_F$ is total and idempotent, and*

$$N_F(a) = N_F(b) \quad \Longleftrightarrow \quad aE_Fb.$$

Every total class-constant readout factors uniquely through $N_F(A_F)$. An admissible transport pair $aE_FT(a)$ gives $N_F(T(a)) = N_F(a)$ wherever the declared fragment contains both sides.

Proof. Each nonempty finite class has a unique least code. Equivalent payloads have the same class and minimum. Conversely, equal minima belong to both classes, which are then equal. The minimum of a minimum's class is unchanged. Apply the quotient factorization after identifying a class with its minimum; the transport assertion uses the equality criterion. □

Thus total payload extraction and finite normal-form construction can genuinely be discharged on this relative fragment. Identifying A_F with the paper's intended native kernel class still requires the corresponding admissibility proof. Moreover, each aspect map required to survive normalization must be constant on E_F-classes. A signature distinguishing equivalent payloads does not descend; calling it an invariant does not repair this conflict.

2.3 Readout Domains and Routed Aggregation

The definition of closure-compatible readout constrains $J_r \circ r$ when readouts exist, not necessarily the raw readout r. For a partial readout, this distinction also includes its domain.

Proposition 2.7 (Exact factorization of a partial readout)**.** *Let $N : A_F \to K$ be total and $f : D \to Y$, $D \subseteq A_F$. A factor $\bar{f} : N(D) \to Y$ with $\bar{f}(N(a)) = f(a)$ on D exists exactly when f is constant on the fibres of N inside D. Equality as partial functions, $f = \bar{f} \circ N$, additionally requires*

$$D = N^{-1}(N(D)).$$

If $D = A_F$, this is the total factorization case. The empty-domain case needs no artificial extension to unobserved classes.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. Define $\bar{f}(N(a)) = f(a)$ for $a \in D$. Fibre constancy is necessary and sufficient for well-definedness. The composite has domain $N^{-1}(N(D))$, so equality of partial functions is equivalent to the displayed domain condition. □

For example, a readout defined only at one of two equivalent payloads is pairwise compatible wherever defined, yet its quotient factor composed with N_F becomes defined at both. Exact domain preservation therefore cannot be inferred from the old pairwise readout condition alone.

Nor does the proposed display macro establish a sufficient normal form: take $A_F = \{0, 1\}$, closure equivalence equal to equality, payload extraction $p(j(a)) = a$, and $N(a) = 0$ for both payloads. The macro's identity $N(p(j(a))) = N(a)$ holds, but the compatible readout $f(a) = a$ cannot factor through this constant N. This is a counterexample to sufficiency of the bare macro, not to the complete admissible-kernel conditions.

Proposition 2.8 (A sufficient routed-state condition)**.** *Fix an admitted weight record w and an operation $G_w : A_F^n \to A_F$, $n \geq 1$. Suppose G_w respects componentwise E_F-equivalence, and $G_w(a, \ldots, a)E_F a$. If every input $a_i E_F a_0$, then $G_w(a_1, \ldots, a_n)E_F a_0$. The operation also descends uniquely to the quotient.*

Proof. Congruence compares the given input tuple with $(a_0, \ldots, a_0)$. The diagonal condition and transitivity give the class-preservation claim. Congruence is exactly the representative-independence needed to define the operation on quotient classes. □

Normalized weights alone do not imply this condition: with classes $\{0, 1\}$ and $\{2, 3\}$ and weights $(1/2, 1/2)$, the constant operation $G_w(a, b) = 2$ is congruent but sends inputs $0, 1$ to the wrong class. By contrast, if the aggregate is specified only at the level of a vector-valued class readout and all values equal v_0, linear aggregation gives $\sum_i w_i v_0 = v_0$ when $\sum_i w_i = 1$. This proves a readout identity, not the existence or class preservation of a native state-level aggregator.

2.4 Shift Compatibility and the Activation Boundary

For the payload-preserving tag migration $U(j_0(a)) = j_1(a)$, define the shifted closure relation by transport through U. Then $p_1 \circ U = p_0$ and both $q_F \circ p_t$ and the finite $N_F \circ p_t$ commute with this display change. This is a proof for an explicitly declared encoding change, not an assertion that arbitrary physical frame changes preserve the payload.

The local kernel package is therefore justified when the fragment, closure equivalence, invariant aspects, readout domains and routed operations have been supplied with the properties above. The finite checker validates those supplied finite relations, normal forms, readout fibres and complete aggregation tables using exact code equality. It neither infers a native equivalence nor silently completes an invalid relation.

The original global activation package remains unproved: a native kernel identification, its aspect/recursive-filter compatibility, all relevant readouts and routed operations, and a compatible normal-form construction across the full active surface are still needed. A quotient class is not a canonical native payload representative; finite representative selection does not construct one on an arbitrary proper class. The symbols `WARN_022` and `WARN_081` are not declared cleared, and no external evaluator outcome is inferred from this work.

Explicit positive finite instance. Take $A_F = \{0, 1, 2, 3\}$ with classes $\{0, 1\}$ and $\{2, 3\}$. Then $N_F = (0, 0, 2, 2)$, and the total class readout $f = (7, 7, 11, 11)$ factors as $\bar{f}(0) = 7$, $\bar{f}(2) = 11$. With $c(a) = \lfloor a/2 \rfloor$, declare the aspect tuple $(\text{state}, c(a), (\text{class}, c(a)))$, constant

© 2026 Marc Brendecke. Preprint — not peer reviewed.

on each class. Let T swap $0,1$ and swap $2,3$, and set $G_w(a,b) = a$ for the admitted weight record $(1/2, 1/2)$. The transport stays in each class, the aggregator is congruent and diagonal-preserving, and the payload-preserving tag shift commutes with all these normal-form comparisons. This is a fully specified consistent local model, not a recovered native QAM witness.

3 HC and Lepton Mass Readout: An Identification Test

The sufficient construction in Proposition 2.2 is not yet an identification with the Companion's HC. This section tests that identification against the actual inherited definitions. Its negative results concern specified readout routes, not every possible nonlinear, coupled or symmetry-breaking realization of the framework.

3.1 What the Existing Definitions Supply

The inherited decomposition is $C = (I+J)/2$, $P_- = (I-J)/2$, with $CP_- = 0$; see Sections 12 and 20. The fermionic input lane is declared J-odd, while C selects the compensated photon/fix output. The refined mass discussion instead assigns massive readout to localization remaining relative to that output (Section 40.3) and names Higgs mediation as its interface (Definition 40.12).

Proposition H5.17 names $\mathcal{R}_m : \mathcal{S}_{\text{spec}} \to \mathcal{S}_{\text{mass}}$, but its argument does not provide a numerical mass functional, generation-resolved coupling, or action on three principal mass roots. The inherited remark "Mass spectrum beyond Koide" explicitly leaves the Yukawa mechanism undeveloped. The symbolic effective operator in Theorem 41.3 does not supply these missing matrix elements or a lepton spectral assignment either.

Two different triples must not be identified by their cardinality: (K_1, K_2, K_3) denotes matter/tachyon/photon sector readouts, whereas (e, μ, τ) denotes three generations within the lepton readout. The generation discussion in Theorem 22.4 does not construct a map equating their components or their scalar products.

There is also a scalar-field issue. The inherited unitary-equivalence argument calls J unitary, whereas the proof of Theorem 23.3 calls it anti-unitary. An anti-unitary involution and $(I+J)/2$ are generally only real-linear. The following operator tests therefore explicitly use a real Hilbert realization, where J is an orthogonal involution. They do not silently assert a complex-linear three-dimensional representation of either inherited convention.

3.2 The Exact Linear HC-to-Root Interface

Proposition 3.1 (A sufficient norm-controlled descent)**.** *Let H be a real Hilbert space, J an orthogonal involution, and $B : H \to \mathbb{R}^3$ a bounded coisometry, $BB^* = I_3$. If $\ker B$ is J-invariant, then*

$$J_m = BJB^*, \qquad J_m^* = J_m, \qquad J_m^2 = I_3, \qquad BC = C_m B, \quad C_m = (I_3 + J_m)/2.$$

Conversely, any relation $BJ = J_m B$ requires $\ker B$ to be J-invariant. The descended operator is unique because B is onto.

Proof. Put $E = B^*B$, the orthogonal projection onto $(\ker B)^\perp$. Since J is self-adjoint and preserves $\ker B$, it also preserves its orthogonal complement, so $JE = EJ$. Thus $J_m^* = J_m$ and $J_m^2 = BJEJB^* = BEB^* = I_3$. Also $J_m B = BJE = BJ$, since J preserves $\ker B$ and B kills it. The identity for C follows by averaging. Conversely, $Bk = 0$ implies $BJk = J_m Bk = 0$. Surjectivity gives uniqueness. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Coisometry is a sufficient metric-control assumption, not a consequence of merely naming a readout. Without invariant kernel, compression BJB^* need not even be an involution. A general invertible change of coordinates gives an involution BJB^{-1}, but need not preserve the Euclidean scalar product used in Koide. A different root-space metric cannot be substituted without re-deriving the physical ratio.

If an independently supplied seed f_0 satisfies $Bf_0 = e$ and the *physical* root readout is proved to be

$$x = \alpha BCf_0 = \alpha C_m e, \qquad e = (1,1,1)/\sqrt{3}, \quad \alpha > 0,$$

with nonzero nonnegative components, the earlier proposition applies. Here α carries square-root mass units for a dimensionless seed. Its missing condition can now be stated on the original carrier:

$$\langle e, J_m e\rangle = \langle B^* e, JB^* e\rangle = 0.$$

The paper presently supplies neither this B, this seed/root law, nor that independently justified orthogonality. The displayed interface is one linear route to investigate, not a requirement that every physical mass readout be linear. In particular, a globally nonnegative linear root map on all of H would have to vanish, since both Bf and $B(-f) = -Bf$ would be nonnegative. A nonzero physical linear root law must be restricted to an admitted state cone.

3.3 Obstructions to the Direct Identification

Proposition 3.2 (The pure HC output loses the odd input)**.** *If $Jf = -f$, then $Cf = 0$. Consequently any homogeneous linear root readout $x = BCf$ is zero. Any readout $F(Cf)$ with $F(0) = 0$ has the same obstruction on this input lane. If instead a seed is J-fixed and the descended linear route has $Bf_0 = e$, then $J_m e = e$ and its nonzero root output has $K = 1/3$.*

Proof. The first statement follows from $Cf = (f + Jf)/2 = 0$. For the second, $J_m e = BJf_0 = Bf_0 = e$, so $C_m e = e$ and all three roots are equal. □

This rules out direct identification of the declared odd fermionic input with a positive root vector obtained only from its compensated output. It does not rule out a mass readout of the residual input or of a coupled state. Nor does quaternionic orthogonality establish the needed root correlation: for an orthogonal projection C one has $\langle Cf, f\rangle = \|Cf\|^2$. Its nonzero output cannot be orthogonal to its own input in that same Hilbert metric. The inherited quaternion-sector wording in Proposition 16.1 must not be used as the proof of $\langle e, J_m e\rangle = 0$.

Proposition 3.3 (Equivariant readout cannot split a fixed input)**.** *Let U preserve an admitted state set and let P cyclically permute the three root coordinates. For any readout F satisfying $F(Uf) = PF(f)$, an input $Uf_0 = f_0$ has $F(f_0) = a(1,1,1)$. If this is a nonzero nonnegative root vector, then $K = 1/3$. No linearity assumption on F is needed.*

Proof. Equivariance gives $PF(f_0) = F(Uf_0) = F(f_0)$. The fixed vectors of the three-cycle are exactly the constant triples. □

A common frame parameter does not evade this result if the input remains fixed and the readout remains equivariant at each parameter value. An asymmetric input, a selected symmetry-breaking state or an explicitly non-equivariant coupling is needed. The generation action U and the matter/tachyon involution J are distinct data.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proposition 3.4 (Closed-triole amplitudes do not yield Koide by norms)**.** *Let $z_0+z_1+z_2=0$ in a normed vector space, not all $z_i=0$. Under the specific identification $m_i=\kappa\|z_i\|^2$ with a common $\kappa>0$, the principal-root ratio satisfies*

$$\frac{1}{3}\le K\le\frac{1}{2}.$$

In particular it cannot equal $2/3$.

Proof. Put $r_i=\|z_i\|$ and $s=\sum_i r_i>0$. The triangle inequality gives $r_i\le s-r_i$, hence $r_i\le s/2$. Therefore $\sum_i r_i^2\le(s/2)\sum_i r_i=s^2/2$. Cauchy–Schwarz gives the lower bound, and the common mass scale cancels from K. □

This excludes a natural but additional identification, not every generation readout of a triole. Unequal generation couplings or a different nonlinear realization need not obey this bound; they are precisely additional data that must be constructed rather than read off from $z_0+z_1+z_2=0$ alone.

Proposition 3.5 (An unrestricted fitted HC shape predicts no roots)**.** *For every nonzero nonnegative triple x, there exist an orthogonal involution J_x and $\alpha_x>0$ such that $x=\alpha_x(I+J_x)e/2$. Thus existence of an HC-shaped operator chosen after the masses is not a prediction of Koide or a hierarchy.*

Proof. Set $C_x=xx^{\mathsf T}/\|x\|^2$ and $J_x=2C_x-I_3$. This is an orthogonal involution. Since $s=\sum_i x_i>0$, $C_xe=sx/(\sqrt{3}\|x\|^2)$, so choose $\alpha_x=\sqrt{3}\|x\|^2/s$. □

3.4 A Residual Coupling Test Compatible With the Mass Language

The refined massive-branch definition suggests testing the component that C removes, rather than identifying the masses with Cf. The following is an explicit *additional effective ansatz*, not a recovered Yukawa interaction. Put $H_-=\ker(J+I)$ and normalize the declared states dimensionlessly. Fix a mass scale $\kappa>0$ and a dimensionless bounded linear coupling $D_\beta: H_-\to V$ to a declared real Hilbert output space. For three admitted generation states define

$$u_i=P_-\psi_i,\qquad m_i(\beta)=\kappa\|D_\beta u_i\|^2,\qquad x_i=\sqrt{\kappa}\|D_\beta u_i\|.$$

The quadratic form is positive, is zero on a fully fixed input, and is unchanged when ψ_i is replaced by $J\psi_i$, since $P_-J\psi_i=-P_-\psi_i$. Unlike a pure $BC\psi_i$ law, it can be nonzero on the declared fermionic input lane. It assigns the same mass to J-dual inputs; a different dual-sector mass law would require a different frame-labelled coupling or readout.

For $q_i=\|D_\beta u_i\|^2$, not all zero, Koide still requires the independent equation

$$\sum_i q_i=\frac{2}{3}\left(\sum_i\sqrt{q_i}\right)^2.$$

Neither positivity nor the residual decomposition proves it. In particular, if all $\|u_i\|=1$ and $D_\beta^*D_\beta=\beta^2 I$ on their span, all masses are $\kappa\beta^2$ and $K=1/3$ for $\beta\neq0$. Thus the isotropic frame term $\beta^2\langle f,f\rangle$ in Definition 23.2, used alone on equally normalized generations, does not generate a hierarchy. A common positive rescaling leaves K unchanged. At $\beta=0$ in this particular ansatz all masses vanish and K is undefined, not a nonzero degenerate Koide limit.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Identification status and the next concrete object. No native HC-to-lepton-root identification is established by the current definitions. The pure odd-input HC route and the common-norm closed-triole route are excluded under their stated assumptions. The remaining constructive target suggested by the existing mass/Higgs language is an independently specified generation-resolved coupling D_β and generation states ψ_i, or an explicit alternative mass realization. Their domains, generation action, positivity, physical units and relation to the effective field operator must be supplied before testing the equal-norm/Koide law, the phase and the absolute scale. Assigning entries from the observed masses would only fit the missing map. The inherited assertions that historical OP 3 is closed, and P2 is confirmed, do not discharge this obligation; their physical derivation remains unestablished here.

4 Generation States and a Closure-Defect Coupling

This section constructs a conditional operator interface from objects already named in the paper: the generation slots, the closure projector and the shifted pacer. It provides explicit formulas, not yet evaluated native matrix elements or a derived Yukawa interaction. In particular, it separates a mass readout in fixed slots from one in coupling eigenstates. Neither is identified with physical lepton masses merely by giving it that name.

4.1 Closure-Compatible Generation States

Use the positive-metric real Hilbert realization from Section 3. For a nonzero odd sector, one may choose a unit vector $\phi_- = P_- g/\|P_- g\|$ with $P_- g \neq 0$. The inherited Dixon dictionary (Theorem 22.4) suggests the normalized seed slots $\phi_- \otimes e_4$, $\phi_- \otimes e_5$, $\phi_- \otimes e_6$. This requires a declared tensor-product metric and a generation-neutral extension of the real J-action; the dictionary alone does not prove that factorization in a native carrier.

On the complexification of this real realization, let $T : \mathbb{C}^3 \to \mathcal{K}$ isometrically embed those slots. Here J denotes the complex-linear extension of the real involution, not an unannounced identification with the inherited anti-unitary convention. Realize $P = \hat{P}_{C_7}$ as an orthogonal closure projector on $\mathcal{K}$, commuting with this J on the declared realization. It is *not* the HC projector C. An action of the pacer on this same carrier must still be supplied; a bosonic clock does not automatically define its action on the probe slots.

Proposition 4.1 (Normalized closed generation seeds)**.** *If $G = T^*PT$ is positive definite, then*

$$S = PTG^{-1/2}, \qquad S^*S = I_3, \quad PS = S, \quad JS = -S.$$

Thus $\psi_i = S\varepsilon_i$ are three orthonormal closed odd states, where ε_i is the standard generation-coordinate basis. If G has rank less than three, this construction cannot produce three independent closed states from these seeds.

Proof. The positive inverse square root exists in the three-dimensional seed space. Since $P^2 = P = P^*$, $S^*S = G^{-1/2}T^*PTG^{-1/2} = I_3$ and $PS = S$. Commutation with J and $JT = -T$ give $JS = -S$. Multiplication by P cannot restore a dimension lost in PT. □

The choice of g, the common carrier and P remain native input obligations. If G is not diagonal in the seed slots, this orthonormalization already mixes them; it is not a proof that the original octonion labels are physical mass eigenstates.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

The spectral restriction is not the new coupling. The inherited proposal $D_- = L_0|_{H_-}$ must not be identified with the new mass coupling D_β by its letter. On the real grading used here, a sector-preserving real-linear $A : H_- \to H_-$ commutes with $J = -I$ there. Requiring also $AJ = -JA$ forces $A = 0$. For the inclusion $\iota : H_- \hookrightarrow H$, the correctly typed relation $J\iota A = -\iota A$ only records that the output is odd, not anticommutation. This qualifies the D1 argument in Theorem 41.13; its nonzero restriction cannot be borrowed as an already verified anticommuting mass coupling. Domain and self-adjointness questions for that spectral proposal are not settled by the finite construction here.

4.2 Constructing the Coupling From the Pacer and Closure

Fix a bounded realization of R_β on the common carrier, or a declared finite matrix realization. In the inherited notation it is the proposed $\hat{\mathcal{R}}_\beta$ of Theorem 26.4. Fix a positive reference r_{ref} with the same units as R_β, and put $\widetilde{R}_\beta = R_\beta / r_{\mathrm{ref}}$. These domain/unit data are additional specifications, not inferred from a clock label.

Proposition 4.2 (A positive closure-defect readout)**.** *For the closed embedding above, define*

$$D_\beta = (I - P)\widetilde{R}_\beta P, \qquad B_\beta = D_\beta S, \qquad Q_\beta = B_\beta^* B_\beta.$$

Then $D_\beta S = [\widetilde{R}_\beta, P]S$ and Q_β is positive semidefinite. Its entries are

$$(Q_\beta)_{ij} = \langle (I - P)\widetilde{R}_\beta \psi_i, (I - P)\widetilde{R}_\beta \psi_j \rangle.$$

For every $P\psi = \psi$, $\langle \psi, [\widetilde{R}_\beta, P]\psi \rangle = 0$, even though $D_\beta \psi$ may be nonzero.

Proof. On a closed input, $[\widetilde{R}_\beta, P]\psi = (I - P)\widetilde{R}_\beta \psi$, which is orthogonal to $\psi \in \operatorname{im} P$. For $v \in \mathbb{C}^3$, $\langle v, Q_\beta v \rangle = \|B_\beta v\|^2 \geq 0$, and the matrix-entry formula is the defining Gram identity. □

Thus the commutator *expectation* used for mass in Proposition 26.5 vanishes on an exactly closed state under these projector assumptions. Replacing it by a transition norm is a genuine change of readout, not an equivalent rewriting of that inherited claim. The norm measures a tendency to leave the selected closure space under the pacer; it does not prove that a physical state actually leaves it or that such leakage is a derived mass mechanism. In particular, exact $[R_\beta, P] = 0$ makes this candidate vanish. A common mass scale $\kappa > 0$ is still needed.

4.3 Coordinate Readouts and Candidate Mass Eigenstates

Proposition 4.3 (The Gram spectrum and the slot diagonal differ)**.** *Let $Q_\beta = V \operatorname{diag}(q_0, q_1, q_2) V^*$, with V unitary and $q_i \geq 0$. The states $\chi_i = SV\varepsilon_i$ are orthonormal closed odd states and $\|D_\beta \chi_i\|^2 = q_i$. If one additionally postulates the effective mass matrix $M_\beta = \kappa Q_\beta$, its candidate eigenmasses are κq_i, whereas the fixed-slot norm readouts are $\kappa (Q_\beta)_{ii}$. When $Q_\beta \neq 0$, the spectral Koide diagnostic is*

$$K_{\mathrm{spec}} = \frac{\operatorname{Tr} Q_\beta}{(\operatorname{Tr} \sqrt{Q_\beta})^2}.$$

It is invariant under unitary changes of generation coordinates.

Proof. Use $S^*S = I_3$ and $V^* Q_\beta V = \operatorname{diag}(q_i)$. The principal root masses are $\sqrt{\kappa}\sqrt{q_i}$; the common scale cancels. Trace and functional calculus are unitarily invariant. The fixed-slot formula is the diagonal Gram identity. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

For example, $Q = \frac{1}{3}\mathbf{1}\mathbf{1}^*$, with $\mathbf{1} = (1,1,1)^{\mathsf{T}}$, has three equal diagonal entries but eigenvalues $(1,0,0)$. Its slot diagnostic is $1/3$ and its spectral diagnostic is 1. Equal diagonal readouts do not establish degenerate eigenmasses. Conversely, diagonalizing a positive matrix does not identify it with the native first-order mass operator Ω of Theorem 27.4; a physical derivation must justify $M_\beta = \kappa Q_\beta$ and its units, rather than confuse that hypothesis with $\Omega^*\Omega$.

4.4 A Cyclic Coupling That Produces the Brannen Form

There is a precise sufficient link to a generation action on the carrier. If a unitary W satisfies $WS = SU$, $WP = PW$ and $W\widetilde{R}_\beta = \widetilde{R}_\beta W$, then $B_\beta U = WB_\beta$, whence $Q_\beta U = UQ_\beta$. These follow by moving W through the defining projectors and pacer and then taking adjoints. The three compatibility relations must be supplied for the native realization, not inferred merely from the existence of three seed slots.

Proposition 4.4 (A Hermitian cyclic root coupling)**.** *Let $U\varepsilon_i = \varepsilon_{i+1 \bmod 3}$ and set*

$$L = aI_3 + zU + \bar{z}U^*, \qquad a > 0, \quad z = re^{i\theta},\ r \geq 0.$$

Choose the Fourier labels so U has eigenvalues $e^{2\pi ik/3}$. Then L has eigenvalues

$$\ell_k = a + 2r\cos(\theta + 2\pi k/3).$$

If all $\ell_k \geq 0$, and if the closure-defect Gram matrix is independently established to be $Q = L^2$, its candidate masses are $m_k = \kappa\ell_k^2$ and

$$K_{\mathrm{spec}} = \frac{1}{3} + \frac{2r^2}{3a^2}. \qquad K_{\mathrm{spec}} = \frac{2}{3} \iff r = a/\sqrt{2}.$$

Conversely, every nonzero positive semidefinite Q commuting with U has a positive root $L = \sqrt{Q}$ of the displayed form, with

$$a = \tfrac{1}{3}\operatorname{Tr}\sqrt{Q}, \qquad z = \tfrac{1}{3}\operatorname{Tr}(U^*\sqrt{Q}).$$

Proof. Fourier diagonalization gives the eigenvalues. Since $\operatorname{Tr} U = \operatorname{Tr} U^2 = 0$, $\operatorname{Tr} L = 3a$ and $\operatorname{Tr} L^2 = 3a^2 + 6r^2$. Positivity gives $\sqrt{L^2} = L$, so division gives the result. Conversely, $\sqrt{Q}$ commutes with U by spectral calculus. The three distinct Fourier eigenvalues of U make its commutant the span of I_3, U, U^2. Hermiticity and $U^* = U^2$ give the stated coefficient form; the trace identities recover a and z. □

This constructs the familiar three-phase harmonic form as eigenvalues of a concrete coupling matrix, without identifying three raw triolic coordinates with mass roots. It does not derive $r/a = 1/\sqrt{2}$, θ or κ, nor does it establish cyclic compatibility or the coefficients of the native $B_\beta^* B_\beta$. Once those native data are given, the converse specifies the coefficients without a mass fit. For negative ℓ_k in a proposed L, the root trace is $\sum_k |\ell_k|$ instead, and the displayed Koide formula is not valid. For real z, two ℓ_k coincide; a generic complex relative coefficient can lift this pair degeneracy. That is an internal generation phase, not an overall phase of the pacer.

There is no conflict with Proposition 3.3: the seed slots form a cyclic *orbit*, not one invariant input, and the spectral states are Fourier combinations of those slots. A complex Hermitian circulant matrix may have three different eigenvalues while its diagonal entries are all equal. If one insists on a real symmetric cyclic matrix, it is of the form $a_0 I + b_0(U + U^*)$ and has a doubly degenerate eigenvalue. The choice of scalar field and physical generation basis matters.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proposition 4.5 (Positive phase chambers and a small-root boundary)**.** *At $r = a/\sqrt{2}$, the cyclic root matrix is positive semidefinite exactly for the phase chambers*

$$\theta \in \bigcup_{j=0}^{2} \left[\frac{2\pi j}{3} - \frac{\pi}{12}, \frac{2\pi j}{3} + \frac{\pi}{12} \right] \pmod{2\pi}.$$

Near the upper boundary $\theta_b = 3\pi/4$ of the chamber centered at $2\pi/3$, put $\theta = \theta_b - \delta$, $\delta \geq 0$. Then $\ell_0 = a\delta + O(\delta^2)$ and $m_0 = \kappa a^2 \delta^2 + O(\delta^3)$, while the other two root eigenvalues tend to $a(3-\sqrt{3})/2$ and $a(3+\sqrt{3})/2$.

Proof. Each $\ell_k \geq 0$ requires $\cos(\theta + 2\pi k/3) \geq -1/\sqrt{2}$. Intersecting these three conditions in each cyclic period gives the displayed intervals. At $3\pi/4$ the zeroth root vanishes and its phase derivative is $-a$. Taylor expansion and squaring give the small-mass law. Substitution gives the other limits. □

A relative generation phase near such a boundary can therefore produce a parametrically light eigenmass within this candidate. This explains a possible amplification mechanism, not why a native state should select that phase or the Koide radius in the first place.

4.5 Overall Phase, Relative Phase and Underdetermination

Proposition 4.6 (A common pacer phase cancels)**.** *For fixed P, S and normalization, if $R_\beta = e^{i\varphi(\beta)} R_0$, then $B_\beta = e^{i\varphi(\beta)} B_0$ and $Q_\beta = Q_0$. Both slot and spectral norm-based mass readouts are unchanged.*

Proof. Pull the scalar through the projectors and use its unit modulus in $B_\beta^* B_\beta$. □

The written pacer $\hat{R}_{\mathrm{GUE}} e^{i\pi\beta_{\mathrm{sym}}}$ in Theorem 26.4 has exactly this common-phase form. It supplies no β-dependent mass ratios for this norm readout when P and S are fixed. Allowing them to vary requires their own independently specified frame laws. The absolute commutator expectation in the inherited mass formula also loses this common phase, even on inputs that are not exactly closed: its expectation acquires only the same unit-modulus factor.

Proposition 4.7 (A two-channel relative phase can survive)**.** *As an additional ansatz, let $\widetilde{R}_\beta = R_a + e^{i\varphi(\beta)} R_b$ with matched domains and units. Put $A = (I-P)R_a S$, $B = (I-P)R_b S$. Then*

$$Q_\beta = A^*A + B^*B + e^{i\varphi(\beta)} A^*B + e^{-i\varphi(\beta)} B^*A.$$

The phase cancels when $A^*B = 0$; it can otherwise affect the Gram matrix through interference. A varying matrix alone does not imply varying eigenmass ratios: the variation could be a basis conjugation.

Proof. Expand $(A + e^{i\varphi}B)^*(A + e^{i\varphi}B)$. Unitary conjugation preserves its eigenvalues. □

The triadic channels in Proposition 27.8 are a place to seek such extra matrix elements, not a specification of them. They must be realized on the same generation/probe carrier and converted to compatible units before adding them to a clock operator. A term of the form PXP cannot itself contribute to $(I-P)\widetilde{R}_\beta P$. The internal coupling must actually connect the selected and complementary spaces for this candidate to be nonzero.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proposition 4.8 (Even unitary pacers leave the Gram matrix free)**.** *Every three-dimensional positive contraction $0 \le Q \le I_3$ can occur as the closed-to-complementary Gram matrix of a unitary pacer. On $\mathbb{C}^3 \oplus \mathbb{C}^3$, take*

$$P = \begin{pmatrix} I_3 & 0 \\ 0 & 0 \end{pmatrix}, \quad S = \begin{pmatrix} I_3 \\ 0 \end{pmatrix}, \quad R = \begin{pmatrix} \sqrt{I_3 - Q} & -\sqrt{Q} \\ \sqrt{Q} & \sqrt{I_3 - Q} \end{pmatrix}.$$

Then $R^*R = RR^* = I_6$ and $((I-P)RS)^*((I-P)RS) = Q$.

Proof. The positive square roots are commuting functions of Q. Block multiplication cancels the off-diagonal terms and gives $(I_3 - Q) + Q = I_3$ on each diagonal. The lower block of RS is $\sqrt{Q}$. □

This is a consistency witness, not a recovery of the native pacer. It also proves why closure, positivity and unitarity alone do not predict a hierarchy or Koide: all positive contraction spectra remain available. Any nonzero positive candidate can be scaled into this range, but that rescaling does not derive a physical mass unit.

Concrete next gate. Generation states and an operator-valued coupling now have explicit conditional formulas: $S = PT(T^*PT)^{-1/2}$ and $D_\beta = (I-P)\widetilde{R}_\beta P$, with the computable three-dimensional Gram matrix $Q_\beta = (D_\beta S)^* D_\beta S$. What is not supplied is a native evaluated P, the action of the pacer on these seeds, a physical mass-matrix law and any required relative channel. The cyclic root form explains how a Brannen-type spectrum can be represented; its Koide radius and phase are still independent obligations. No lepton-mass fit, native activation, OP2 closure or parameter-free spectrum is asserted.

5 Testing the Native Operator Link

The previous interface becomes predictive only if the native projector, pacer and generation embedding determine its matrix elements. The retained definitions provide a useful partial link, but not that full identification. All statements below use the same positive-metric, bounded or finite realization as Section 4. Orthogonality of P, common domains and the seed metric are assumptions of this realization, not consequences of the inherited idempotence condition alone. The HC projector C remains distinct from P and from the defect penalty $\mathcal{H}_C$.

5.1 What the Original Definitions Supply

The Meta-Septum definition before Theorem 26.4 specifies an admissible projector through a trace condition modulo HC; it does not list its entries on the generation seeds. That theorem writes $R_\beta = e^{i\pi\beta_{\mathrm{sym}}} R_{\mathrm{GUE}}$, without specifying the clock eigenvectors relative to P and those seeds. Proposition 28.3 requires $[\mathcal{M}_{16}, P] = 0$ for exact compensator-matrix closure. No equation there identifies R_{GUE} with $\mathcal{M}_{16}$ or a function of it. The HS(4)–HS(6) carrier coding and the spectral-to-mass interface of Proposition H5.17 do not evaluate these joint matrix elements either. A coordinate coding does not specify an operator action.

Proposition 5.1 (The two minimal penalty terms coincide)**.** *Write $M = \mathcal{M}_{16}$ on the declared common carrier and $P^\perp = I - P$. Set*

$$K = [M, P], \qquad E = P^\perp M P + P M P^\perp, \qquad H = \rho K^*K + \sigma E^*E, \quad \rho, \sigma \ge 0.$$

Then

$$E^*E = K^*K, \qquad H = (\rho+\sigma)K^*K, \qquad [H,P] = 0.$$

For an isometric closed embedding S, let $B_M = P^\perp MS/r_{\mathrm{ref}}$ and $Q_M = B_M^* B_M$, where $r_{\mathrm{ref}} > 0$ has the units of M. Then

$$S^*HS = (\rho+\sigma)r_{\mathrm{ref}}^2 Q_M.$$

Proof. In the orthogonal splitting $\operatorname{im} P \oplus \ker P$, write $M = \left(\begin{smallmatrix} A & F \\ G & D \end{smallmatrix}\right)$. Then

$$K = \begin{pmatrix} 0 & -F \\ G & 0 \end{pmatrix}, \quad E = \begin{pmatrix} 0 & F \\ G & 0 \end{pmatrix}, \quad K^*K = E^*E = \begin{pmatrix} G^*G & 0 \\ 0 & F^*F \end{pmatrix}.$$

This proves the first claims without requiring $M = M^*$. Since $PS = S$, $KS = P^\perp MS = r_{\mathrm{ref}} B_M$; compressing the positive product proves the last identity. □

Thus the existing minimal defect penalty already has a Gram-type compression. Identifying it with the pacer Gram matrix additionally requires, for example, $R_\beta = e^{i\varphi}M$ with matched units. That bridge is not supplied by the source. Even when $H \neq 0$, $P^\perp HP = 0$: this minimal penalty cannot itself be the missing off-sector pacer channel. The statement does not concern a separately postulated deeper HC generator.

Proposition 5.2 (Transferring closure to the clock requires a bridge)**.** *If $[M,P] = 0$ and R_0 belongs to the norm-closed unital $*$-algebra generated by M, then $[R_0,P] = 0$ and $Q_\beta = 0$ for $R_\beta = e^{i\varphi(\beta)}R_0$. Exact closure of M alone does not imply $Q_\beta = 0$ for an independently specified pacer.*

Proof. Taking adjoints gives $[M^*,P] = 0$. Every finite product and sum of M, M^*, I therefore commutes with P, and so does every norm limit. Its closed-to-complementary block vanishes. For the converse warning, take $P = \operatorname{diag}(I_3, 0)$, $S = (I_3, 0)^{\mathsf{T}}$, $M = I_6$ and $R_0 = \left(\begin{smallmatrix} 0 & I_3 \\ I_3 & 0 \end{smallmatrix}\right)$. Then $[M,P] = 0$ but $Q_0 = I_3$. □

In particular, identifying the clock as a polynomial or continuous functional-calculus expression in an exactly closed bounded normal M kills this defect route. This does not prohibit nonzero masses from the internal block $P\Omega P$ or another readout. It prohibits deriving them from this particular transition while maintaining that bridge and exact closure.

5.2 Spectral and Frame Data Are Not Yet Joint Matrix Elements

Proposition 5.3 (An unchanged clock spectrum can change the Gram spectrum)**.** *Keep $P = \operatorname{diag}(I_3, 0)$ and $S = (I_3, 0)^{\mathsf{T}}$. For real diagonal matrices $D_+ = \operatorname{diag}(d_i^+)$ and $D_- = \operatorname{diag}(d_i^-)$, put*

$$O_\theta = \begin{pmatrix} \cos\Theta & -\sin\Theta \\ \sin\Theta & \cos\Theta \end{pmatrix}, \quad \Theta = \operatorname{diag}(\theta_i), \qquad R_\theta = O_\theta \operatorname{diag}(D_+, D_-) O_\theta^*.$$

These self-adjoint clocks all have the same eigenvalues, whereas

$$Q_\theta = \frac{1}{4r_{\mathrm{ref}}^2} \operatorname{diag}\big((d_i^+ - d_i^-)^2 \sin^2(2\theta_i)\big).$$

Thus even an exact simple clock spectrum does not by itself determine the transition spectrum relative to fixed closure and seed spaces.

Proof. O_θ is orthogonal, so conjugation preserves the clock spectrum. The lower-left block of R_θ is $(D_+ - D_-)\sin\Theta\cos\Theta$; its normalized Gram product gives the formula. For nonzero gaps, varying the angles changes its eigenvalues, not just its eigenvectors. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

This finite example is not a replacement for the native Riemann clock. It identifies what a zero sequence or spacing statistic leaves unspecified: the orientation of the clock eigenspaces relative to the closure projector and generation embedding. A native joint operator law could remove this freedom; its evaluation is the burden.

Proposition 5.4 (A simultaneous unitary frame change preserves mass ratios)**.** *Let F be unitary on the carrier, V unitary on $\mathbb{C}^3$, and $s > 0$. Starting with $G_0 = T_0^* P_0 T_0 > 0$, set*

$$P_f = F P_0 F^*, \quad T_f = F T_0 V, \quad \widetilde{R}_f = s e^{i\varphi} F \widetilde{R}_0 F^*.$$

Then the normalized embeddings and transition matrices obey

$$S_f = F S_0 V, \qquad B_f = s e^{i\varphi} F B_0 V, \qquad Q_f = s^2 V^* Q_0 V.$$

Consequently nonzero spectral mass ratios and K_{spec} are unchanged under the candidate mass law.

Proof. $G_f = V^* G_0 V$, and its positive inverse root is $V^* G_0^{-1/2} V$. Substitute into $S_f = P_f T_f G_f^{-1/2}$, then into $B_f = (I - P_f)\widetilde{R}_f S_f$. The Gram identity follows. Common positive scaling and unitary conjugation preserve the ratios. □

This is a statement about a declared positive Hilbert metric, not an identification of a Lorentz boost with F. A common scalar Planck calibration or common frame phase cannot generate a hierarchy change in this lane. A relative motion of the projector, clock or seeds, or a nontrivial relative channel, requires its own native law; it is not supplied by assigning the same frame label to all three.

Proposition 5.5 (The transition requires sufficient output rank)**.** *For $B : \mathbb{C}^3 \to \ker P$ with $\dim \operatorname{im} B \le d$, one has $\operatorname{rank}(B^* B) = \operatorname{rank} B \le d$. Three strictly positive candidate masses require three independent transition outputs. A single scalar complementary channel produces at most one positive eigenmass in this Gram readout.*

Proof. $\langle v, B^* B v \rangle = \|Bv\|^2$ implies $\ker(B^* B) = \ker B$. Rank-nullity on the three-dimensional input proves the claim. □

The three entries $(\Psi_1, \Psi_2, \Psi_\perp)$ in Proposition 27.8 label probe channels, not the three generation coordinates. An operator-valued channel can have dimension greater than one; the proposition does not prove that the native complementary space is one-dimensional. An explicit embedding and channel rank are needed before this block system can be used as a generation coupling.

5.3 A Conditional Mass Law From the Existing Block Architecture

The inherited field operator Ω and its triadic blocks suggest testing elimination of a gapped complementary sector. The following is finite spectral algebra at a frozen background, in units $\hbar = c = 1$. It additionally assumes a self-adjoint mass operator, canonical kinetic normalization and a closed truncation to the chosen three-dimensional seed space plus its retained complement. The source does not establish those assumptions or the required block entries. Other closed states, derivative couplings and background variations cannot silently be omitted. Hermiticity requires adjoint relations between the paired triadic channels, not merely their names.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proposition 5.6 (Elimination of a gapped complementary block)**.** *Let the supplied mass-dimension operator on that truncation be*

$$\Omega_{\mathrm{tr}} = \begin{pmatrix} A & V^* \\ V & H_\perp \end{pmatrix}, \qquad A = A^*, \quad H_\perp = H_\perp^* \geq \Delta I, \quad \Delta > 0.$$

For an eigenvalue λ with $|\lambda| < \Delta$, its closed coordinate x satisfies

$$\big(A - V^*(H_\perp - \lambda I)^{-1}V\big)x = \lambda x, \qquad y = -(H_\perp - \lambda I)^{-1}Vx.$$

Writing $A_{\mathrm{eff}}(\lambda)$ for the operator in parentheses,

$$\|A_{\mathrm{eff}}(\lambda) - A_{\mathrm{eff}}(0)\| \leq \frac{\|V\|^2|\lambda|}{\Delta(\Delta - |\lambda|)}.$$

Proof. Solve the lower block equation for y and substitute into the upper one. The resolvent identity gives $(H_\perp - \lambda I)^{-1} - H_\perp^{-1} = \lambda(H_\perp - \lambda I)^{-1}H_\perp^{-1}$. The spectral gap bounds the two inverse norms by $(\Delta - |\lambda|)^{-1}$ and Δ^{-1}. □

Proposition 5.7 (When a Gram mass law is a leading approximation)**.** *Assume further $A = 0$, $H_\perp = \Delta I$ and $V = \mu B$, with $\mu > 0$ a mass scale, B dimensionless and $Q = B^*B$. For each $q_i \geq 0$ the light eigenvalue is $-m_i$, where*

$$m_i = \frac{\sqrt{\Delta^2 + 4\mu^2 q_i} - \Delta}{2} = \frac{\mu^2}{\Delta}q_i + O\left(\frac{\mu^4 q_i^2}{\Delta^3}\right).$$

The physical interpretation as a positive mass magnitude is conditional on this canonical Hermitian Dirac-type reading. The signed effective operator is negative, not $+(\mu^2/\Delta)Q$. For a normalized light eigenstate its complementary weight is

$$\|y\|^2 = \frac{m_i}{\Delta + 2m_i}.$$

Proof. In an eigenvector of Q, elimination gives $\lambda = -\mu^2 q_i/(\Delta - \lambda)$, so the light root of $\lambda^2 - \Delta\lambda - \mu^2 q_i = 0$ is the displayed negative or zero root. Expansion for $\mu^2 q_i/\Delta^2 \ll 1$ gives the approximation. Moreover $\|y\|^2/\|x\|^2 = \mu^2 q_i/(\Delta + m_i)^2 = m_i/(\Delta + m_i)$; normalization gives the weight formula. □

This supplies a possible justification for $M = \kappa Q$ as a *leading mass-magnitude law*, with $\kappa = \mu^2/\Delta$, if the native field blocks independently establish a flat positive gap, zero direct block and $V = \mu B$. A non-flat gap gives a weighted Gram matrix $V^*H_\perp^{-1}V$, not generally a scalar multiple of Q; a nonzero A also changes the spectrum. Even with a flat gap, the exact eigenmasses must be tested anew for Koide. No native values of μ, Δ or the triadic matrix elements have been obtained by this calculation.

There is also an exact-closure limitation: an eigenstate constrained to the original P-sector has $y = 0$, hence $Vx = 0$. Nonzero masses from the special case $A = 0$ above therefore require dressed states with nonzero complementary weight. They are not exactly P-closed. For $\mu\|B\|/\Delta \ll 1$ the weight is controlled, or one may define a distinct spectral projector onto the light eigenspace. Neither choice is the original strict closure constraint. Closure of M alone, invariance under the full Ω, and approximate closure of a dressed state are different conditions.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Result of the native search. The existing penalty yields an exact algebraic Gram compression, and the existing block architecture admits a conditional gapped-sector mass mechanism. Neither evaluates the native tuple (P, R_β, T). The next model-specific datum is the cross block $(I-P)R_\beta S$, together with a declared relation, if any, to $(I-P)\Omega S$. A clock spectrum, common frame shift, scalar length calibration or finite coding alone cannot provide it. The three-generation rank, relative-channel law, physical gap, Koide radius and absolute mass scale remain independent obligations.

6 An Evaluated Coupling Candidate and Its Scale Limits

The original field equation names effective coefficients and triadic operators but does not evaluate their joint action. This section therefore does two separate things: it evaluates a declared cyclic candidate using the inherited phase input, and tests whether its remaining scales could produce the required mass ratios. It does not identify that candidate with the native clock cross block.

6.1 Separating Shape, Normalization and Physical Scale

For a nonzero supplied cross block V on a finite positive-metric realization, the convention $\mathrm{Tr}(B^*B) = 1$ fixes $\mu = \|V\|_{\mathrm{F}}$ and $B = V/\mu$. It removes a redundancy, not an undetermined physical coupling.

Proposition 6.1 (Coupling and gap scales remain independent input)**.** *In $V = \mu B$, replacing (μ, B) by $(t\mu, B/t)$, $t > 0$, leaves V unchanged. Normalizing B removes this redundancy. For normalized B in the flat-gap model of Proposition 5.7, put $\eta = \mu/\Delta$. Then*

$$m_i = \Delta\,\frac{\sqrt{1+4\eta^2 q_i}-1}{2}.$$

The shape q_i and η determine ratios, but do not fix Δ. Simultaneously scaling (μ, Δ) by $s > 0$ scales all masses by s, leaving ratios and complementary weights unchanged.

Proof. Substitution proves the redundancy and dimensionless mass formula. The exact complementary weight is $m_i/(\Delta + 2m_i)$, which is invariant under the simultaneous scaling. □

Even a supplied Q leaves B undetermined up to output isometries in its polar decomposition; these do not change B^*B. The source symbols g_Y, χ and the paired triadic operators specify products, not a numerical factorization into a scale and a normalized matrix. A physical normalization needs canonical kinetic terms, common units and actual matrix elements.

6.2 A Minimal Two-Channel Guess Is Insufficient

Proposition 6.2 (Two cyclic harmonics cannot give the Koide ratio)**.** *For the generation cycle U, let $B = b_0 I + b_1 U$, with arbitrary complex b_0, b_1 not both zero. If the candidate masses are proportional to the eigenvalues of B^*B, their Koide diagnostic satisfies $K \le 1/2$, not $2/3$.*

Proof. In the Fourier basis let $\omega = e^{2\pi i/3}$ and $\lambda_k = b_0 + b_1\omega^k$. The numbers $z_k = \omega^k\lambda_k$ have zero sum, since both $\sum_k \omega^k$ and $\sum_k \omega^{2k}$ vanish. Their magnitudes are the principal root masses up to a common factor. The closed-triole norm bound of Proposition 3.4 therefore gives $\sum_k |z_k|^2/(\sum_k |z_k|)^2 \le 1/2$. □

Thus a two-channel relative phase is not by itself enough in this minimal cyclic realization. More general operator-valued channels are not excluded by the proposition.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

6.3 Numerical Three-Channel Candidate From the Written Phase

Take the *inherited hypotheses*, not newly proved laws, $r/a = 1/\sqrt{2}$ and $\theta = \pi - \sqrt{2}\arcsin(\beta_{\mathrm{sym}})$, with the paper's rounded input $\beta_{\mathrm{sym}} = 0.5493$. Fix only the normalization $\mathrm{Tr}(B^*B) = 1$. On the positive-root phase branch,

$$\theta = 2.319190257857\ldots, \qquad B = \frac{I_3}{\sqrt{6}} + \frac{e^{i\theta}U + e^{-i\theta}U^*}{\sqrt{12}}.$$

In the convention $U\varepsilon_i = \varepsilon_{i+1 \bmod 3}$,

$$B \simeq \begin{pmatrix} 0.408248 & -0.196433 - 0.211536i & -0.196433 + 0.211536i \\ -0.196433 + 0.211536i & 0.408248 & -0.196433 - 0.211536i \\ -0.196433 - 0.211536i & -0.196433 + 0.211536i & 0.408248 \end{pmatrix}.$$

Its normalized Gram matrix has diagonal $1/3$ and lower cyclic coefficient $-0.166548 + 0.255824i$. Its ascending eigenvalues are

$$(q_e, q_\mu, q_\tau) \simeq (0.0002366350,\ 0.0567819436,\ 0.9429814214).$$

The subscripts denote an attempted ascending lepton assignment, not an independently derived generation dictionary. The leading ratio prediction *conditional on this shape* is

$$\frac{q_e}{q_\tau} = 0.0002509434\ldots, \qquad \frac{q_\mu}{q_\tau} = 0.0602153365\ldots.$$

The inherited mass benchmark in Theorem 22.5, used here only for comparison, instead gives approximately 0.0002875860 and 0.0594633229. The relative ratio differences are -12.74% and $+1.265\%$. No measured lepton mass enters the construction of B. Nevertheless the radius, phase law and rounded frame input are assumptions, so this is not a parameter-free native derivation. The Koide diagnostic $2/3$ is inherited from the imposed radius. At $\beta_{\mathrm{sym}} = 0$ the same fixed-radius formula is not a positive root matrix and does not give three degenerate masses. This local candidate is not a valid interpolation to the proposed photon limit.

An explicit effective six-dimensional realization is

$$P = \begin{pmatrix} I_3 & 0 \\ 0 & 0 \end{pmatrix}, \quad S = \begin{pmatrix} I_3 \\ 0 \end{pmatrix}.$$

$$\widetilde{R}_{\mathrm{cand}} = \begin{pmatrix} 0 & B^* \\ B & 0 \end{pmatrix}, \quad \Omega_{\mathrm{cand}} = \begin{pmatrix} 0 & \mu B^* \\ \mu B & \Delta I_3 \end{pmatrix}.$$

It realizes the candidate Gram matrix and field cross block, with $V = \mu B$. It has not been shown to be the paper's R_{GUE}, a native compensator-matrix realization or a generation-compatible embedding of its Dixon carrier. Its imposed Hermitian off-sector clock is not that missing identification.

6.4 A Scale Choice Cannot Repair the Fixed Shape

Proposition 6.3 (The positive-gap correction increases smaller mass ratios)**.** *In the flat positive-gap model, for $0 < q_i < q_j$ and $\mu > 0$,*

$$\frac{m_i}{m_j} = \frac{q_i}{q_j}\frac{\sqrt{\Delta^2 + 4\mu^2 q_j} + \Delta}{\sqrt{\Delta^2 + 4\mu^2 q_i} + \Delta} > \frac{q_i}{q_j}.$$

Consequently no positive μ, Δ can reproduce the quoted muon/tau benchmark with the fixed numerical shape above.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. Rationalize each exact mass from Proposition 5.7. The numerator of the second factor is strictly larger than its denominator. Already q_μ/q_τ exceeds the quoted benchmark; the correction can only increase that ratio. Common scaling cannot alter it. □

This excludes this fixed shape with zero direct block and flat positive gap, not all coupling matrices, non-flat gaps or direct internal mass terms. A change of the relative phase, radius or field blocks needs an independent structural law; fitting it to the benchmark is not such a derivation.

6.5 Reference Scales and Explicit Calibration Boundaries

A reference mass m_{ref} permits the parametrization $\mu = u\, m_{\mathrm{ref}}$, $\Delta = d\, m_{\mathrm{ref}}$ with $u, d > 0$. It does not fix the dimensionless coefficients u, d. For comparison only, the ordinary single-frame Planck reference is $m_{\mathrm{P}} = \sqrt{\hbar c/G}$, or $\hbar/(c\ell_{\mathrm{P}})$. The external CODATA central reference is $m_{\mathrm{P}}c^2 = 1.220890 \times 10^{19}\,\mathrm{GeV}$ [1]. This external normalization is not predicted by the present construction.

As an independently defined electroweak reference, use $v_F = (\sqrt{2}G_F)^{-1/2} = 246.21965\ldots\,\mathrm{GeV}$ and $\mu_{\mathrm{EW}} = v_F/\sqrt{2} = 174.10358\ldots\,\mathrm{GeV}$, with the CODATA central value $G_F = 1.1663787 \times 10^{-5}\,\mathrm{GeV}^{-2}$ [2]. Using this reference for μ would be an additional choice, not a recovered Yokamono/Higgs coupling.

With $\Delta = m_{\mathrm{P}}$ and $\mu = \mu_{\mathrm{EW}}$, the leading coefficient is $\kappa = 2.48278 \times 10^{-6}\,\mathrm{eV}$ and the exact positive masses are approximately

$$(5.8751 \times 10^{-10},\ 1.4098 \times 10^{-7},\ 2.3412 \times 10^{-6})\,\mathrm{eV}.$$

Thus this particular Planck-gap/electroweak-mixing choice does not give the charged-lepton scales quoted in the source. Choosing both scales at the Planck reference instead gives masses of Planck order for the larger candidate eigenvalues; neither choice is a model-derived scale identification.

For clarity, a single mass can always be used as a calibration:

$$\mu^2 = \frac{m_\tau(m_\tau + \Delta)}{q_\tau}, \qquad \Delta = \frac{\mu^2 q_\tau}{m_\tau} - m_\tau.$$

With the inherited $m_\tau = 1.77686\,\mathrm{GeV}$, fixing $\Delta = m_{\mathrm{P}}$ gives $\mu \simeq 4.7964 \times 10^9\,\mathrm{GeV}$; fixing $\mu = \mu_{\mathrm{EW}}$ gives $\Delta \simeq 1.6085 \times 10^4\,\mathrm{GeV}$. These are explicitly one-mass calibrations, not predictions, and neither removes the muon/tau ratio obstruction.

The paper's proposed frame-insertion assignments must also be kept consistent if used. Formally applying $\hbar_f = \gamma_L^{-1}\hbar_m$, $G_f = \gamma_L^{-2}G_m$ and $c_f = c_m$ gives $m_{\mathrm{P},f} = \gamma_L^{1/2} m_{\mathrm{P},m}$. Using $\hbar_m/(c_m \ell_{\mathrm{P},f})$ instead would mix frame assignments and introduce an extra factor γ_L. This is conditional bookkeeping, not validation of the frame laws.

6.6 Diagnosing the Matrix Discrepancy

The matrix introduces no extra distortion of the inherited Brannen spectrum: its Fourier eigenvalues are exactly those root coordinates, up to the common normalization. Direct diagonalization and the analytic formula agree. Six-decimal rounding of its displayed entries

[1] NIST, 2022 CODATA, Planck mass energy equivalent: `https://physics.nist.gov/cgi-bin/cuu/Value?plkmc2gev`.

[2] NIST, 2022 CODATA, Fermi coupling constant: `https://physics.nist.gov/cgi-bin/cuu/Value?gf`.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

changes the electron/tau ratio by about 0.01225%, not 12.74%. Cyclic relabelling and complex conjugation only permute its spectrum.

For a diagnostic inversion of the *historical comparison only*, put $x_k = \sqrt{m_k / \sum_j m_j}$ in the ascending assignment above. Then

$$a_{\mathrm{cmp}} = \tfrac{1}{3}\sum_k x_k, \qquad z_{\mathrm{cmp}} = \tfrac{1}{3}\sum_k x_k e^{-2\pi i k/3}, \qquad \rho_{\mathrm{cmp}} = |z_{\mathrm{cmp}}|/a_{\mathrm{cmp}}.$$

The discrete inverse Fourier identity reconstructs these three positive roots. It gives

$$\theta_{\mathrm{cmp}} = 2.316624255394\ldots, \qquad \rho_{\mathrm{cmp}} = 0.707100578472\ldots.$$

Relative to the imposed $\rho = 1/\sqrt{2}$, the radius difference is about -8.77 parts per million, whereas the phase difference is $-0.00256600246\ldots$ rad, or $-0.147021\ldots$ degrees. Changing only the phase to this comparison-derived value gives ratios approximately $(0.0002874631, 0.0594619634, 1)$; changing only the radius leaves $(0.0002510584, 0.0602167023, 1)$. Thus the phase accounts for almost all of this fixed-ansatz discrepancy. These inversions are post-hoc diagnostics, not a new derived phase law. This θ is the relative coefficient phase of U and U^*, not the common pacer phase, which still cancels from the Gram matrix.

Proposition 6.4 (Relative-phase sensitivity near a small root)**.** *Let $h_k = 1 + 2\rho\cos\phi_k > 0$, with $\phi_k = \theta + 2\pi k/3$ and $q_k = h_k^2/(3+6\rho^2)$. For a fixed reference mode j,*

$$\partial_\theta \log\frac{q_i}{q_j} = -4\rho\left(\frac{\sin\phi_i}{h_i} - \frac{\sin\phi_j}{h_j}\right), \qquad \partial_\rho \log\frac{q_i}{q_j} = 4\left(\frac{\cos\phi_i}{h_i} - \frac{\cos\phi_j}{h_j}\right).$$

For the inherited phase law, $\partial_\beta\theta = -\sqrt{2}/\sqrt{1-\beta^2}$.

Proof. The common normalization cancels from q_i/q_j. Differentiate $2\log h_i - 2\log h_j$ and apply the chain rule. □

At the candidate, the electron/tau and muon/tau logarithmic derivatives with respect to β are approximately 92.6426 and -8.28621. The small electron root is close to the phase-chamber boundary, which explains the large relative amplification. This is not evidence of an unrepresented physical frame interaction by itself.

Within the stated phase law, the comparison phase would correspond to $\beta_{\mathrm{cmp}} = 0.5508152849\ldots$, a shift of 0.00151528. That is about thirty half-steps of four-decimal rounding. If decimal rounding alone is tested on $[0.54925, 0.54935]$, the middle/largest ratio ranges from about 0.06024029 down to 0.06019039 and remains above the comparison. Its monotonic decrease here follows from the signs in the displayed derivative. This interval is a rounding test, not an assigned experimental uncertainty. Using the exact written $\gamma_{\mathrm{string}} = \log 2/(\pi\sqrt{3})$ with the still approximate $\gamma_{\mathrm{BI}} = 0.2375$ gives $\beta = 0.5493482484\ldots$ and ratio $0.06019127\ldots$; it does not remove the discrepancy. Inverting to a new $\gamma_{\mathrm{BI}} \simeq 0.2383454$ would be another calibration, not a justified revision of that input.

6.7 What the Planck Test Does and Does Not Exclude

An independent SI reconstruction gives $m_{\mathrm{P}} = 2.17643434\ldots \times 10^{-8}\,\mathrm{kg}$ and $\ell_{\mathrm{P}} = 1.61625502\ldots \times 10^{-35}\,\mathrm{m}$. Converting $m_{\mathrm{P}}c^2$ from joules to GeV reproduces $1.22089013\ldots \times$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

10^{19} GeV, consistent with the rounded CODATA reference above [3]. There is no missing GeV/eV factor or missing power of c in the micro-eV estimate.

The suppression comes from the three added assumptions $A = 0$, $\Delta = m_{\mathrm{P}}$ and $\mu = \mu_{\mathrm{EW}}$: the leading energy is $\mu^2 q_i/\Delta$. The largest value is about 7.59×10^{14} times below the quoted tau mass. Substituting the reduced reference $m_{\mathrm{P}}/\sqrt{8\pi}$ as the physical gap changes the result by only $\sqrt{8\pi}$, to about 1.174×10^{-5} eV for the largest mass. That substitution is another gap choice, not a unit correction. The paper's conditional consistent Planck frame factor is only $\gamma_L^{1/2} \simeq 1.3654$; even the inconsistent mixed-$\hbar$ factor is only $\gamma_L^{3/2} \simeq 2.5453$. Neither accounts for the required hierarchy.

A minimum reference length or UV cutoff does not, without a spectral equation, fix the complementary mass gap in this block model. Keeping $\mu = \mu_{\mathrm{EW}}$ and calibrating the tau mass gives $\Delta \simeq 16.085\,\mathrm{TeV}$, corresponding to the inverse energy length $\hbar c/\Delta \simeq 1.227 \times 10^{-20}\,\mathrm{m}$. This is about $7.59 \times 10^{14} \ell_{\mathrm{P}}$, an effective gap length, not a new fundamental Planck length. Keeping the gap at the Planck reference instead requires an intermediate mixing reference $\mu \simeq 4.7964 \times 10^9\,\mathrm{GeV}$, about $2.755 \times 10^7 \mu_{\mathrm{EW}}$, in that one-mass calibration. Neither possibility is fixed by the source or repairs the flat-gap ratio obstruction.

Proposition 6.5 (Non-flat gaps can absorb a target spectrum)**.** *Suppose the positive candidate is $B = F\,\mathrm{diag}(\sqrt{q_i})F^*$, $q_i > 0$. For any target magnitudes $t_i > 0$, choose μ such that $\mu^2 q_i > t_i^2$ and set*

$$\Delta_i = \frac{\mu^2 q_i}{t_i} - t_i > 0, \qquad H_\perp = F\,\mathrm{diag}(\Delta_i)F^*.$$

Then $\left(\begin{smallmatrix} 0 & \mu B^* \\ \mu B & H_\perp \end{smallmatrix}\right)$ has light signed eigenvalues $-t_i$.

Proof. Conjugating by $\mathrm{diag}(F, F)$ splits the system into three 2×2 blocks. Their light magnitudes obey $t_i(t_i + \Delta_i) = \mu^2 q_i$, exactly the defining relation. □

For the historical comparison, fixing $\Delta_\tau = m_{\mathrm{P}}$ and the tau-calibrated intermediate μ gives

$$(\Delta_e, \Delta_\mu, \Delta_\tau)/m_{\mathrm{P}} \simeq (0.87258583, 1.01264668, 1).$$

All gaps remain of Planck order and all three historical masses are reproduced, by construction. This is an exact fitted counterexample to a blanket exclusion of Planck-order gaps, not a prediction of these anisotropies. It uses the target masses to define $H_\perp$. It cannot determine whether a physical discrepancy belongs to the root coupling, the gap or another term. With electroweak mixing and every gap still of Planck order, this modest gap anisotropy does not fix the absolute suppression.

Proposition 6.6 (A retained direct block makes the Planck sector a small correction)**.** *For the Hermitian block model of Proposition 5.6, let $H_\perp \geq \Delta I$ and $|\lambda| \leq E < \Delta$. Every such eigenvalue satisfies*

$$\mathrm{dist}(\lambda, \mathrm{spec}\, A) \leq \frac{\|V\|^2}{\Delta - E}.$$

Proof. The closed coordinate x cannot vanish, since otherwise the eigenvalue would lie in $\mathrm{spec}\, H_\perp$. The Schur equation gives $(A - \lambda I)x = V^*(H_\perp - \lambda I)^{-1} V x$. The inverse norm is at most $(\Delta - E)^{-1}$. For self-adjoint A, the spectral theorem bounds $\|(A - \lambda I)x\|$ below by $\mathrm{dist}(\lambda, \mathrm{spec}\, A)\|x\|$. □

[3] The SI check uses the exact defining values of h, c, e and the NIST CODATA central $G = 6.67430 \times 10^{-11}$ in SI units: `https://physics.nist.gov/cgi-bin/cuu/Value?bg`.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

With the candidate $\|V\|^2 = \mu_{\mathrm{EW}}^2 q_{\max}$, $\Delta = m_{\mathrm{P}}$ and $E = 2\,\mathrm{GeV}$, the bound is about $2.3412 \times 10^{-6}\,\mathrm{eV}$. A direct block at charged-lepton scales could therefore coexist with Planck-gap corrections that are tiny. The original Ω already contains direct/projected/triadic terms; setting their compressed sum A to zero was an additional choice of the earlier induced-mass test. This alternative does not derive A or its hierarchy, but it shows why that test does not exclude the paper's general use of the Planck reference.

Determination reached. There is now an explicit numerical normalized coupling candidate and a testable six-dimensional block realization. Its relative shape is conditional on the inherited radius and phase hypotheses. No numerical native values of g_Y, χ, μ or Δ have been established. The missing physical input is a represented cross-block interaction and gap equation that fixes them independently of the target lepton masses, and resolves the strict-versus-dressed closure boundary of Section 5. The discrepancy diagnosis separates the inherited relative-phase ansatz from matrix arithmetic, and the extra flat-Planck-gap induced mass hypothesis from unit conversion or a fundamental length claim. It does not uniquely identify a physical cause from the mass data.

7 Koide Phase Overlap and Finite-Gap Constraints

The original OP 3 asked for the phase of the charged-lepton mass spectrum. This section examines a conditional readout of several phase-shifted preparations in a common generation basis in frame M. Each preparation has nonnegative roots, a common mean $a > 0$, and the exact Koide ratio 2/3. Let

$$x_{jk} = a\big[1 + \sqrt{2}\,\mathrm{Re}(\omega^k e^{i\delta_j})\big], \qquad \omega = e^{2\pi i/3}, \qquad k = 0, 1, 2.$$

If the readout averages the roots with nonnegative weights summing to one before squaring, it is determined by $Z = \sum_j w_j e^{i\delta_j}$. In particular, the resulting Koide ratio is $(1 + |Z|^2)/3$, and the reduced phase is $\arg Z$ for $Z \neq 0$. A pair with equal weights at $\delta_0 \pm h$, for $|h| < \pi/2$, retains the central phase and reduces the radius by $\cos h$. Averaging the squared roots instead adds the weighted variance of each root to the corresponding mass. This comparison makes the order of the readout operations explicit.

7.1 Exact poles and the positive-root arc

Consider a cyclic comparison with static root eigenvalues x_k of mean one, direct diagonal entry $a_0 = 1 + \sqrt{2}$, and a common positive intermediate entry D. The lower eigenvalue ℓ_k of the block with off-diagonal squared magnitude $D(a_0 - x_k)$ obeys

$$x_k = \ell_k + \frac{\ell_k(a_0 - \ell_k)}{D}.$$

This relation allows the static target to be reconstructed from the mass ratios at a prescribed D. In the fixed-label chamber about the positive-radius reference direction the admissible reduced phases lie in $[-\pi/12, \pi/12]$. The reconstructed complex target must lie in the convex hull of this arc. With N equal weights and fixed mean sine, the smallest mean cosine is attained with all but at most one of the sines at the endpoints. This follows from concavity of $y \mapsto \sqrt{1 - y^2}$ on the sine interval. Jensen's inequality gives the upper bound. Continuity on the connected set of admissible sine tuples gives every intermediate value.

Using the reference masses 0.51099895, 105.6583755, and 1776.93 MeV, the comparison at $D = 68$ admits only $N = 15$ among $2, \ldots, 16$. At $D = 100$ it admits $N = 10, \ldots, 16$.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Repeated phases are allowed. The masses, the chosen intermediate entry and the assumed readout specify this inverse calculation.

7.2 Conditions for a transport interpretation

A relative transport phase could enter an effective coupling if two paths contribute to the same matrix element before pole determination. A phase change confined to the prepared probe changes its spectral weights while leaving the operator poles fixed. Likewise, a common factor multiplying the entire mass-root operator changes the scale while leaving the mass ratios fixed. The native generation operator, the coherent end-channel map, the path selection and the intermediate scale remain open. The structural shell-transition question bearing the later OP 3 label remains distinct from this spectral question.

8 Tensor Activation and Covariant Coupling Readout

The preceding numerical test used identity input and output metrics. Its ratio obstruction must not be extended without qualification to an independently supplied, nontrivial tensor comparison. Conversely, activating tensor notation does not by itself supply such a comparison. The governing shifted normal form remains that of Definition A.34.1.

8.1 Which Shift Can Affect a Numerical Spectrum?

The global QAM migration $\Sigma_4(k) = k + 1$ moves namespace tags. Definition A.30.11 preserves tensor sidedness, variance and contraction discipline; Theorem A.30.17 preserves the numerical readout payload and its guards. For example, support $\{0, 2, 5\}$ becomes $\{1, 3, 6\}$, so its packed mask changes from 37 to 74, not 38. This is not an addition of one to a coupling coefficient, phase or eigenvalue. The local first-boundary HC shift of Definition H2.74 has a different structural role; it is not a numerical mass law either.

The active two-sided grammar does require explicit comparison objects for same-variance pairings (Definition H2.41). This exposes a genuine possible extension of the calculation: specify the comparison on the transition output and the kinetic normalization on the generation input. The residual-channel comparison in the source is not automatically either of these metrics. Equal dimensions do not identify their index spaces or physical roles.

8.2 A Positive-Metric Tensor Extension

Let $B : V_{\mathrm{gen}} \to W_{\mathrm{out}}$ be a finite complex map, with specified positive Hermitian metrics K and H on these two spaces. Here H is dimensionless comparison data, not the energy-valued complementary block $H_\perp$; K is an additional kinetic metric, not an identification with a spacetime Lorentz metric. In components,

$$(Q_H)_{ij} = \overline{B^a{}_i}\, H_{ab} B^b{}_j, \qquad Q_H = B^* H B, \qquad B^\sharp = K^{-1} B^* H.$$

Both metrics and any change in B must be specified independently of the desired masses. This is an added positive finite realization of the tensor grammar, not a native metric extraction.

Proposition 8.1 (Metric-covariant effective spectrum)**.** *The generalized eigenvalues of $Q_H v = q K v$ are the nonnegative eigenvalues of*

$$C = H^{1/2} B K^{-1/2}, \qquad Q_{K,H} = C^* C = K^{-1/2} B^* H B K^{-1/2}.$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

In the additional weighted self-adjoint flat-gap model

$$\Omega_{K,H} = \begin{pmatrix} 0 & \mu B^\sharp \\ \mu B & \Delta I \end{pmatrix}, \qquad \langle \cdot, \cdot \rangle = \operatorname{diag}(K, H),$$

the corresponding light magnitudes obey $m_i(m_i + \Delta) = \mu^2 q_i$ *for* $\mu, \Delta > 0$.

Proof. Set $w = K^{1/2} v$ in the generalized equation. Conjugating the block operator by $\operatorname{diag}(K^{1/2}, H^{1/2})$ gives the ordinary Hermitian block $\left(\begin{smallmatrix} 0 & \mu C^* \\ \mu C & \Delta I \end{smallmatrix}\right)$. A singular-value decomposition of C splits its nonzero modes into 2×2 blocks, whose characteristic equations give the stated identity. □

For $K = H = I$ this recovers the previous test exactly. The effective quadratic mass readout and the flat-gap realization remain additional hypotheses. A common phase of B cancels even for nontrivial K, H.

Proposition 8.2 (A covariant chart shift is not a physical correction)**.** *For invertible input and output chart matrices* A, D*, with coordinates* $x = Ax'$ *and* $y = Dy'$*, put*

$$B' = D^{-1} B A, \quad K' = A^* K A, \quad H' = D^* H D.$$

Then $Q'_H = A^* Q_H A$*, and the generalized eigenvalues are unchanged.*

Proof. Substitute the transformed tensors to obtain the congruence identity. The substitution $v = Av'$ bijects the generalized eigenvectors and preserves eigenvalue multiplicities; equivalently the characteristic determinant is multiplied by $|\det A|^2$. □

The numerical audit also tests nonunitary charts. Correctly transporting both metrics recovers the original eigenvalues within 2×10^{-15} absolute numerical error. Incorrectly resetting both metrics to the identity after that change gives ratios approximately $(0.0001813510, 0.0416327127, 1)$ instead of $(0.0002509434, 0.0602153365, 1)$. Such a change is a contraction error, not evidence of a physical frame effect. The chosen charts illustrate covariance; they are not the native frame transformations.

8.3 A Physical Comparison Can Change the Shape, But Must Be Derived

Proposition 8.3 (Positive comparison fit freedom)**.** *For square invertible* B*,* $K = I$ *and any positive Hermitian target* T*, the choice* $H = (B^{-1})^* T B^{-1}$ *is positive and gives* $B^* H B = T$*. In particular, for* $B = F \operatorname{diag}(\sqrt{q_i}) F^*$ *with* $q_i > 0$*, setting* $H = F \operatorname{diag}(t_i/q_i) F^*$ *realizes any* $t_i > 0$.

Proof. Positivity follows by applying T to $B^{-1} w \neq 0$; multiplication gives the first identity. The aligned diagonal formula is its special case. □

For the historical benchmark only, choose $t_i = q_\tau m_i / m_\tau$ so the largest-mode weight stays one. The required output eigenweights in the ascending B modes are

$$(h_e, h_\mu, h_\tau) \simeq (1.14601907,\ 0.98751126,\ 1).$$

Thus relative comparison weights of about $+14.60\%$ and -1.249% can absorb the original leading-ratio discrepancy. This is a *post-hoc metric fit*, not a prediction: the benchmark defines H, including its eigenbasis alignment. It neither proves a tensor activation law nor uniquely attributes the discrepancy to the phase. The finite-gap correction still applies. An exact prescribed-mass fit at given μ, Δ would instead require $t_i = m_i(m_i + \Delta)/\mu^2$, again target-derived.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proposition 8.4 (Absolute metric weights bound any scale enhancement)**.** *If $h_- I \le H \le h_+ I$ and $k_- I \le K \le k_+ I$, with positive bounds, the ascending eigenvalues satisfy*

$$\frac{h_-}{k_+}\lambda_i(B^*B) \le q_i(K,H) \le \frac{h_+}{k_-}\lambda_i(B^*B).$$

Proof. Bound the generalized Rayleigh quotient $x^*B^*HBx/(x^*Kx)$ above and below by the corresponding ordinary quotient of B^*B, then apply the min–max principle on subspaces. □

Keeping the numerical B, $\mu = \mu_{\mathrm{EW}}$, $A = 0$ and the canonical flat gap $\Delta = m_{\mathrm{P}}$, the tau mass would require

$$\frac{h_+}{k_-} \ge \frac{m_\tau(m_\tau + m_{\mathrm{P}})}{\mu_{\mathrm{EW}}^2 q_\tau} \simeq 7.59 \times 10^{14}.$$

The modest fitted shape weights above cannot supply that enhancement. This bound concerns absolute relative normalization, not merely metric condition numbers: even $H = cI$ has condition number one. Taking such a huge scalar at fixed numerical B, μ enhances the canonical coupling norm by about 2.755×10^7; it introduces the missing physical coupling scale rather than deriving it. If it is only a coordinate change, covariance cancels the apparent enhancement. The physical normalization convention is now $\mathrm{Tr}(C^*C)$, not an untransported coordinate trace $\mathrm{Tr}(B^*B)$.

8.4 Hilbert-Carrier Enlargement Also Needs an Interaction

Proposition 8.5 (An identity tensor extension preserves the readout)**.** *For a normalized ancilla a, the isometry $Jv = v \otimes a$ and $B' = B \otimes I$ satisfy*

$$J^*B'^*B'J = B^*B.$$

Proof. The left side equals $(B^*B)\langle a, a\rangle = B^*B$. □

This rules out a correction from mere duplicated channel counting, not from genuine new cross-channel interactions. The inherited HS(5)/HS(6) carrier and activation-mask grammar does not itself evaluate those interactions. A physical activation must supply the new matrix entries, comparison tensors, seed embedding and normalized gap/direct blocks together. The current source and role checker give admissibility rules, not numerical K, H on these particular spaces. Consequently the old identity-metric exclusion is conditional, while neither the namespace shift nor tensor activation alone establishes the missing spectrum or scale.

9 HC Dynamics, Real Actions and the Geometry Interface

This section carries the conditional coupling program through its projected dynamics, action and bilinear geometry. The retained source provides the ordered master equation (Theorem 27.4), the curvature action (Theorem 27.9) and a conditional Einstein reconstruction (Proposition 28.14). These are not already a common solved field system. In particular, the source has not derived an identification between the microscopic coordinate gamma matrices and the inverse of its reconstructed coframe. The distinction becomes essential below.

All finite examples here use commuting effective amplitudes and declared adjoints. A dagger denotes the ordinary positive internal adjoint where specified, not an identification with the Lorentzian metric or the native QAM comparison. Dimensionless local charts are used in the explicit examples; a physical normalization requires the scale record of Section 45.1.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

9.1 Root Positivity Is an Additional Dynamical Obligation

For a complex two-mode input $\chi = (u, v)$ set

$$X_0 = |u|^2 + |v|^2, \quad X_1 = |u|^2 - |v|^2, \quad X_2 = 2\,\mathrm{Re}(\overline{u}v), \quad X_3 = 2\,\mathrm{Im}(\overline{u}v).$$

Then $X_0^2 = X_1^2 + X_2^2 + X_3^2$. Let U be the three-cycle permutation matrix, and introduce the orthonormal Hermitian matrices

$$E_0 = I/\sqrt{3}, \qquad E_1 = (U + U^\dagger)/\sqrt{6}, \qquad E_2 = i(U - U^\dagger)/\sqrt{6}.$$

The following is an additional quadratic root readout, not an evaluated native mass operator:

$$L_\zeta = X_0 E_0 + \zeta X_1 E_1 + \zeta X_2 E_2, \qquad \zeta \geq 0.$$

Proposition 9.1 (Balance and full-input positivity are different)**.** *For $X_0 > 0$ and $L_\zeta \geq 0$, its candidate root ratio is*

$$\frac{\mathrm{Tr}(L_\zeta^2)}{(\mathrm{Tr}\, L_\zeta)^2} = \frac{1 + \zeta^2[1 - (X_3/X_0)^2]}{3}.$$

Positivity for every input χ holds if and only if $\zeta \leq 1/\sqrt{2}$. The aligned real or purely imaginary slice has $X_3 = 0$; attaining $2/3$ on it requires $\zeta = 1$, outside the full-input positive range.

Proof. Orthogonality gives $\mathrm{Tr}\, L_\zeta = \sqrt{3}X_0$ and $\mathrm{Tr}\, L_\zeta^2 = X_0^2 + \zeta^2(X_1^2 + X_2^2)$. In the three Fourier modes, the traceless part has eigenvalues $\zeta\sqrt{2/3}(X_1 \cos\varphi_j - X_2 \sin\varphi_j)$, where $\varphi_j = 2\pi j/3$. Their minimum over all inputs with fixed X_0 is $-\zeta\sqrt{2/3}X_0$, attained on the equator $X_3 = 0$. Comparing with $X_0/\sqrt{3}$ proves the bound. □

Thus an isometric coefficient choice does not derive a physical positive domain or a mass law. On the aligned slice $\chi = ib(\cos(\theta/2), \sin(\theta/2))$, $b > 0$, the $\zeta = 1$ positive domain consists of the three closed chambers $|\theta - 2\pi j/3| \leq \pi/12$, modulo 2π. Preparation and preservation of a chosen chamber must be checked. A fixed-radius phase minimum of a separately constrained energy does not prove stationarity or stability against amplitude, hidden-mode and background variations. Taking absolute values of signed roots does not preserve the displayed ratio in general.

9.2 Projection Retains Hidden States and Moving Frames

Write $D_\mu = \partial_\mu + \mathcal{B}_\mu$ and suppose Γ^0 is invertible. The ordered master equation gives the formal generator

$$\partial_t \Psi = \mathcal{G}\Psi, \qquad \mathcal{G} = -\mathcal{B}_0 - (\Gamma^0)^{-1} \sum_{a=1}^{3} \Gamma^a(\partial_a + \mathcal{B}_a) - i(\Gamma^0)^{-1}\Omega.$$

This rearrangement asserts neither a self-adjoint Hamiltonian nor well-posedness on an unspecified domain. Its full HC contribution is

$$\mathcal{G}_{\mathrm{HC}} = -(\Gamma^0)^{-1}[\kappa\Gamma^\mu n_\mu + i\gamma I]H_C.$$

The order matters. Positivity of H_C alone is not a decay theorem.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proposition 9.2 (Exact projected autonomy criterion)**.** *Let u be the realification of a finite modal state, $\dot{u} = \mathcal{G}_{\mathbb{R}} u$, and let $r = T(t)u$ be a surjective real-linear extraction. For any right inverse R_s, write $u = R_s r + z$, $Tz = 0$. Then*

$$\dot{r} = Kr + q_z, \quad K = (\dot{T} + T\mathcal{G}_{\mathbb{R}})R_s, \quad q_z = (\dot{T} + T\mathcal{G}_{\mathbb{R}})z.$$

The instantaneous equation is independent of every hidden lift if and only if $(\dot{T}+T\mathcal{G}_{\mathbb{R}})\ker T = 0$.

Proof. Differentiate $r = Tu$ and substitute the decomposition. Two lifts of the same r differ by an arbitrary element of $\ker T$, giving both necessity and sufficiency. □

For example, $\dot{f} = -iHf$ with $H = S + iB$ Hermitian has S real symmetric and B real skew. Writing $f = x + ir$ gives $\dot{x} = Bx + Sr$, $\dot{r} = Br - Sx$. For constant blocks, eliminating x yields the exact memory equation

$$\dot{r}(t) = Br(t) - Se^{Bt}x(0) - \int_0^t Se^{B(t-s)}Sr(s)\,ds.$$

It is not automatically a Markovian damping law. In two real dimensions an autonomous linear generator gives $\dot{\theta} = 2(\omega + c\cos\theta + d\sin\theta)$ on the aligned slice. Forward invariance requires inward signs at the lower and upper boundaries. Summing over the equally spaced lower and upper boundary triples forces $\omega = 0$; the same signs then force each boundary value to vanish. A nonzero first harmonic cannot have those six distinct zeros. Only scalar radial evolution remains. This does not exclude nonlinear feedback or one separately selected chamber.

9.3 A Conservative Inertia Can Be Derived Without Viscosity

For a declared positive canonical two-mode action, let

$$\begin{aligned} L &= \frac{i}{2}(f^\dagger \dot{f} - \dot{f}^\dagger f) - \mathcal{E}(f), \\ e &= (\cos(\theta/2), \sin(\theta/2)), \qquad N = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, \\ f &= (a + ib)e + cNe, \qquad b > 0. \end{aligned}$$

The four real coordinates are essential. The kinetic term is $b\dot{a} - a\dot{b} - bc\dot{\theta}$, or $-2a\dot{b} - bc\dot{\theta}$ up to a total derivative. Setting $a = c = 0$ before variation deletes the canonical partners.

Proposition 9.3 (Regular hidden-block elimination)**.** *Put $q = (b, \theta)$, $h = (a, c)$ and $D = \mathrm{diag}(2, b)$. If $\mathcal{E} = V + j^T h + \frac{1}{2}h^T W h$, with W positive definite, algebraic elimination in the action gives*

$$h = -W^{-1}(D\dot{q} + j), \qquad L_{\text{eff}} = \tfrac{1}{2}(D\dot{q} + j)^T W^{-1}(D\dot{q} + j) - V.$$

The induced kinetic metric is $DW^{-1}D$; its coordinate dependence must be retained in the Euler–Lagrange equations.

Proof. Modulo the total derivative, $L = -h^T D\dot{q} - \mathcal{E}$. Its h equation is $Wh + D\dot{q} + j = 0$. Substitution completes the square and proves the formula. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

For $H = mI + \omega\sigma_y$, $m > |\omega|$, and $\mathcal{E} = f^\dagger H f$, this gives

$$L_{\text{eff}} = \dot{b}^2/m + b^2(\dot{\theta} - 2\omega)^2/(4m) - mb^2, \qquad I_\theta = b^2/(2m).$$

The equation $\ddot{\theta} = -2(\dot{b}/b)(\dot{\theta} - 2\omega)$ describes reversible amplitude exchange: the apparent damping changes sign under contraction. It is not viscosity. The native positive modal adjoint and coefficients have not been extracted. More generally, a full-action pullback tests tangent equations only; omitted normal field equations remain necessary.

9.4 The Null Kinetic Term and the Real-Action Choice

Suppress a constant commuting closure background and take flat fixed spacetime, constant real null n, $A = 0$, and $H_C = hT_0$, with T_0 fixed Hermitian, $c_T = \operatorname{Tr} T_0^2 > 0$. For $\mathcal{B}_\mu = \kappa H_C n_\mu$ the printed curvature action reduces exactly to

$$F_{\mu\nu} = \kappa T_0(n_\nu \partial_\mu h - n_\mu \partial_\nu h), \qquad L_{\text{HC}} = \frac{\kappa^2 c_T}{2}(n \cdot \partial h)^2.$$

Before imposing nullity its principal tensor is $\kappa^2 c_T(n^\mu n^\nu - n^2 g^{\mu\nu})$; for nonzero null n and $\kappa \neq 0$ it has rank one. A positive homogeneous temporal inertia is not a Lorentzian wave operator or fluid viscosity. If minimal exact closure makes $H_C = 0$ identically on a branch, this contribution vanishes there. Vanishing at one point is insufficient.

There is a separate reality issue. For ordinary Hermitian internal A_μ, H, set

$$D_\mu^A H = \partial_\mu H + i[A_\mu, H], \quad F_{\mu\nu}^A = \partial_\mu A_\nu - \partial_\nu A_\mu + i[A_\mu, A_\nu], \quad X_{\mu\nu} = n_\nu D_\mu^A H - n_\mu D_\nu^A H.$$

Then $F = iF^A + \kappa X$. The literal $-\operatorname{Tr}(FF)/4$ can be complex, with real part

$$\operatorname{Re} L_F = \tfrac{1}{4}\operatorname{Tr}(F^A F^A) - \tfrac{\kappa^2}{4}\operatorname{Tr}(XX).$$

The electric kinetic signs are opposite for signature $(-,+,+,+)$. This is a convention-dependent diagnostic, not a native ghost theorem. An explicitly *different* unitary-internal comparison action is

$$L_F^\dagger = -\tfrac{1}{4}\operatorname{Tr}(F_{\mu\nu}^\dagger F^{\mu\nu}) = -\tfrac{1}{4}\operatorname{Tr}(F^A F^A) - \tfrac{\kappa^2}{4}\operatorname{Tr}(XX).$$

It is not silently substituted into Theorem 27.9. The traced cross commutator vanishes, but its field equations must be derived from its own allowed variations.

9.5 Probe Reality, Matrix Currents and Background Reactions

Assume a factorization into internal and spin spaces, $\bar{\Psi} = \Psi^\dagger \eta_D$, $\eta_D^\dagger = \eta_D$, constant Hermitian $K^\mu = \eta_D \Gamma^\mu$, and $(\eta_D \Omega)^\dagger = \eta_D \Omega$. Internal Hermitian fields commute with the spin weights. Then

$$\operatorname{Re}(i\kappa \bar{\Psi} \Gamma^\mu n_\mu H \Psi) = 0.$$

A genuinely real symmetric probe action therefore removes this ordered transport term, while retaining a real γH mass-like coupling. This is a change of probe law, not an equivalent reading of the original. For noncommuting weights the Hermitian part is instead $i[K^\mu n_\mu, H]/2$, not the original ordered product.

For the paired curvature action, $\Omega = mI + \gamma H$, and an independent Hermitian matching target Λ, the real test action with penalty $\lambda_H \operatorname{Tr}(H - \Lambda)^2$ has sources

$$j^\mu = \operatorname{Tr}_{\text{spin}}[(I \otimes K^\mu)|\Psi\rangle\langle\Psi|], \qquad S = \operatorname{Tr}_{\text{spin}}[(I \otimes \eta_D)|\Psi\rangle\langle\Psi|].$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Here S need not be positive. Independent A, H variations give

$$E_A^\nu = D_\mu^A F_A^{\mu\nu} + i\kappa^2 n_\mu [H, X^{\mu\nu}] - j^\nu,$$
$$E_H = \kappa^2 n_\nu D_\mu^A X^{\mu\nu} - 2\lambda_H (H - \Lambda) - \gamma S.$$

Integration by parts in the two curvature terms gives the displayed derivatives; varying the probe and penalty terms gives the last terms. The probe equation yields $D_\mu^A j^\mu = -i\gamma[H, S]$ on shell, so its ordinary trace is conserved. Gauge covariance also requires the target reaction $E_\Lambda = 2\lambda_H(H - \Lambda)$ in the Ward identity. For fixed transforming Λ and no other reactions, all other equations on shell require $[H, \Lambda] = 0$. This is not an instruction to impose $E_\Lambda = 0$ or $H = \Lambda$. A functional $\Lambda[M, P]$ instead transfers the reaction through its chain rule. The inherited scalar gradient-trace currents are not automatically these matrix currents.

Keeping the original ordered equation gives a different balance: for the same constant weights and suppressed closure load,

$$\partial_\mu(\Psi^\dagger K^\mu \Psi) = -2\kappa \Psi^\dagger (K^\mu n_\mu) H_C \Psi.$$

Its sign is not fixed by $H_C \geq 0$. An open completion needs an explicit return law or reservoir; a moving comparison needs its own compatibility equations. Neither is derived by calling this loss gravitational transport. No native viscosity or reservoir is supplied.

9.6 An Explicit Null Geometry, Kept Off Shell

In the fixed Hermitian factorization above, with the closure load suppressed, the inherited real coframe has the exact identity

$$E^a{}_\mu = \mathrm{Re}(\bar\Psi \Gamma^a D_\mu \Psi) = \tfrac{1}{2}\partial_\mu j^a + \kappa v_H^a n_\mu, \quad j^a = \Psi^\dagger K^a \Psi, \quad v_H^a = \Psi^\dagger K^a H_C \Psi.$$

Differentiation proves the first term; the purely internal Hermitian gauge term has zero real part. Unlike the i-weighted probe-action term, the real HC coframe contribution survives.

Proposition 9.4 (Gradient flatness and a null-preserving update)**.** *If $H_C = 0$ and E has rank four in this fixed-frame branch, then $Y^a = j^a/2$ are local coordinates and $g = \eta_{ab} dY^a dY^b$ is flat. More generally let $E = E_0 + \kappa v n^T$, with E_0 invertible and $d = 1 + \kappa n^T E_0^{-1} v \neq 0$. For $g = E^T \eta E$, $g_0 = E_0^T \eta E_0$,*

$$\det E = (\det E_0) d, \qquad n^T g^{-1} n = \frac{n^T g_0^{-1} n}{d^2}.$$

Thus this nonsingular rank-one update preserves a null covector.

Proof. The inverse function theorem proves the flat-coordinate statement. The matrix determinant lemma gives the determinant, and the rank-one inverse formula gives $n^T E^{-1} = n^T E_0^{-1}/d$. Contracting with η^{-1} proves the second identity. □

Variable Clifford/adjoint weights or a noncommuting geometric spin connection can escape the gradient statement; it is not a theorem about every native or coarse-grained metric. HC alone has rank at most one and cannot furnish a four-dimensional coframe.

For a concrete local branch use $\eta = \mathrm{diag}(-1, 1, 1, 1)$, $n = (-1, 0, 0, 1)^T$, $l = \eta^{-1} n$, and

$$E = I + \kappa h(x, y) l n^T, \qquad g = \eta + 2\kappa h\, n n^T, \qquad \det E = 1.$$

This dictionary can be realized by a common local probe, not just by independent choices of E_0, v_H. In a declared three-internal-mode, four-spinor representation take $K^a = (I, \alpha_i)$,

© 2026 Marc Brendecke. Preprint — not peer reviewed.

$\eta_D = \beta = \mathrm{diag}(I_2, -I_2)$, $\Gamma^a = \beta K^a$, with Pauli off-diagonal α_i. Let $\Psi_1 = (1,0,1,0)^T/\sqrt{2}$, $\Psi_3 = 0$, $H_C = \mathrm{diag}(h,0,0)$, and set $J^a = 2(x^a + c^a) - l^a$, $c = (3,0,0,0)$. On a future-timelike patch for J, put

$$m_J = \sqrt{(J^0)^2 - |\boldsymbol{J}|^2}, \quad \Psi_2 = \sqrt{\frac{J^0 + m_J}{2}} \begin{pmatrix} \xi \\ (\boldsymbol{\sigma}\cdot\boldsymbol{J})\xi/(J^0 + m_J) \end{pmatrix}, \qquad \xi = (1,0)^T.$$

Direct Pauli contraction gives $j^a = 2(x^a + c^a)$ and $v_H = hl$. This uses $\{\Gamma^a, \Gamma^b\} = -2\eta^{ab} I$ and is not an identification of the native septonic carrier. It is off shell.

The null coordinates $u = z - t$, $v = (z + t)/2$ put this metric in the form $2du\,dv + dx^2 + dy^2 + 2\kappa h\,du^2$. Its nonzero curvature components are determined by

$$R_{\mu i \nu j} = -\kappa h_{,ij} n_\mu n_\nu, \quad R_{\mu\nu} = -\kappa \Delta_\perp h\, n_\mu n_\nu, \quad R = 0, \qquad i,j \in \{x,y\}.$$

These follow directly from the Levi–Civita connection; a vanishing scalar curvature does not imply flatness.

Varying an independent unconstrained metric in the paired HC action at fixed n, F *before* this null restriction gives

$$T^{\mathrm{HC}}_{\mu\nu} = \kappa^2 c_T |\nabla_\perp h|^2 n_\mu n_\nu.$$

The contracted curvature action vanishes on the profile, but its metric variation need not. Omitting all other stresses and the cosmological term, taking $\kappa \neq 0$ and κ_{eff} as an input, the isolated equation $G_{\mu\nu} = \kappa_{\mathrm{eff}} T^{\mathrm{HC}}_{\mu\nu}$ reduces to

$$\Delta_\perp h + a|\nabla_\perp h|^2 = 0, \quad a = \kappa_{\mathrm{eff}}\kappa c_T, \qquad \Delta_\perp e^{ah} = 0.$$

For $a \neq 0$, any positive harmonic F gives the local solution $h = \log F/a$. Positive affine F supplies nonconstant examples; restrict further to $h \geq 0$ if $H_C \geq 0$ is required. This is an isolated off-shell geometry/stress match, not a full probe, gauge, projector or minimal-closure solution, nor a derivation of Newton's constant. A metric-enforced null constraint or an induced metric variation changes the variational problem. Restricting the action to this null family first loses normal Einstein equations.

9.7 A Necessary Current Test and a Same-Frame Exclusion

For the constructed transverse profile let $Y = x + c$. Its induced coordinate current and volume satisfy

$$\mathcal{J}^\mu = (E^{-1})^\mu{}_a j^a = 2Y^\mu - 2\kappa h (n\cdot Y) l^\mu, \qquad \sqrt{|\det g|} = 1, \qquad \nabla_\mu \mathcal{J}^\mu = 8.$$

The last equality uses $n \cdot l = 0$ and $l \cdot \partial h = 0$. Hence the unchanged construction cannot solve the closed real probe theory with this induced phase current. A purely internal Hermitian gauge field does not alter this coframe or cancel a central divergence by a commutator. The number 8 is a normalization-dependent dimensionless diagnostic, not a predicted production rate.

There is a stronger pointwise test which does not require the factorized current construction.

Proposition 9.5 (Conditional same-frame trace obstruction)**.** *Suppose the same probe, adjoint and derivative define a real classical full-rank coframe*

$$E^a{}_\mu = s_0 \,\mathrm{Re}(\bar\Psi \Gamma^a D_\mu \Psi), \qquad s_0 \in \mathbb{R} \setminus \{0\},$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

and make the additional identification $\Gamma^\mu = (E^{-1})^\mu{}_a \Gamma^a$ *in the same ordered master equation. Set* $R_\Psi = (i\Gamma^\mu D_\mu - \Omega)\Psi$ *and* $w = \bar{\Psi}\Omega\Psi$*. Then*

$$\mathrm{Im}(\bar{\Psi} R_\Psi) = 4/s_0 - \mathrm{Im}\, w.$$

In particular, a real w *excludes an on-shell full-rank coframe under these same-frame assumptions.*

Proof. Reality of the inverse frame and its contraction with E give

$$4 = (E^{-1})^\mu{}_a E^a{}_\mu = s_0 \,\mathrm{Re}(\bar{\Psi}\Gamma^\mu D_\mu \Psi).$$

Multiply the master residual by the same adjoint and take the imaginary part, using $\mathrm{Im}(iz) = \mathrm{Re}\, z$. On shell the left side vanishes. This proves both statements. □

A sufficient reality condition is $(\eta_D \Omega)^\dagger = \eta_D \Omega$; Hermiticity of Ω alone is insufficient without its relation to the adjoint. For finite commuting amplitudes the same identity gives

$$\|R_\Psi\| \geq \frac{|4/s_0 - \mathrm{Im}\, w|}{\|\eta_D \Psi\|}.$$

Neither a finite real normalization nor a change of connection removes the obstruction while all its assumptions are retained. It uses no weak-field or Clifford approximation.

This is *not* a universal inconsistency theorem for the paper. The retained reconstruction has not derived that additional same-frame identification. A reference/microscopic probe and a separately derived effective geometry are not excluded. Changing the coframe functional, deriving a changed probe equation from a composite-metric action, or allowing a nonreal bilinear changes the problem and requires a new common action and current balance. Merely inserting an imaginary constant to meet the trace condition is not a derivation.

9.8 Native Identification Still Required

The construction establishes conditional inertia, a null kinetic tensor, explicit real-action alternatives and a local curved metric with an isolated stress match. It also identifies a definite forbidden direct identification. It does not establish masses, an irreversible fluid, a photon rest frame, a wormhole or inter-sector information flow. The retained variational and Einstein statements must be read with these qualifications, not promoted by a successful compiler run.

A native continuation must fix the common carrier and adjoint, decide independent versus functional HC variations, derive every current and background reaction from that same action, and specify the relation between microscopic gamma matrices and effective geometry. Only then can a common solution be tested against all tangent and normal equations, the positive root domain and the physical scale record. The executable HC root-readout, projected-residual, collective-action, null-axis, real-current and bilinear-geometry audits in the repository retain expanded examples and counterexamples. Their finite regression checks support these identities; they are not native closure or proof-assistant certificates.

10 Cartan Comparison: Residuals and Normal Source Compatibility

This section records a new research result, not an identification of the native MM/HC action. It follows the action and geometry distinctions in Section 9. The model below is an explicitly

© 2026 Marc Brendecke. Preprint — not peer reviewed.

added neutral Einstein–Cartan comparison with three complex commuting spinors. Its finite jets do not constitute a solution on an open set. Neither its matrices nor its six source coordinates are identified with matter, photon or tachyon frames, C7 carriers, physical masses or a measured gravitational coupling.

10.1 Declared Model and Residual Convention

Take independent fields: an invertible real coframe $E^a{}_\mu$, a real Lorentz connection $\omega_{\mu,ab} = -\omega_{\mu,ba}$, the spinor triple Ψ, and a real multiplier density $\lambda_a{}^\mu$ of coordinate weight one. Put $v = |\det E|$, $e = E^{-1}$ and $\eta = \mathrm{diag}(-1,1,1,1)$. In the declared Dirac convention, $K^a = \beta\Gamma^a$ and

$$B_\mu = -\tfrac{1}{4}\omega_{\mu,ab}\Gamma^a\Gamma^b, \qquad D_\mu\Psi = \partial_\mu\Psi + B_\mu\Psi, \qquad C^a{}_\mu = \mathrm{Im}(\Psi^\dagger Q K^a D_\mu\Psi).$$

Here Q acts on the three channels and K^a on spin indices; tensor products are implicit. A dagger is the ordinary complex adjoint, not a new native comparison map. The assumed action is

$$\begin{aligned} S_{\mathrm{cmp}} &= \int d^4x \left\{ v\left(\frac{c}{2}R(E,\omega) - u\right) + \mathcal{L}_D + \lambda_a{}^\mu(E^a{}_\mu - C^a{}_\mu)\right\}, \qquad (1)\\ \mathcal{L}_D &= -v\,\mathrm{Im}(\Psi^\dagger e_a{}^\mu K^a D_\mu\Psi) - v\,\mathrm{Re}(\Psi^\dagger(M\otimes\beta)\Psi). \end{aligned}$$

The numerical branch uses $c = 1$, $u = 0$ and fixed dimensionless comparison matrices

$$M = \begin{pmatrix} 0.6 & 0 & 4 \\ 0 & 0.6 & 0.7 \\ 4 & 0.7 & 0.6 \end{pmatrix}, \qquad Q = \frac{i}{2}(E_{12} - E_{21}),$$

where E_{jk} denotes a channel matrix unit. These choices are not a prediction of couplings or a derivation of the Einstein term. No internal charged equation is suppressed: neutrality is part of this comparison.

Write $F = C - E$, and denote the independent coframe and full real-action probe Euler densities by $\mathfrak{E}_a{}^\mu$ and $\mathcal{R}$. Thus the reduced variation after exact algebraic Cartan elimination is

$$\delta S_{\mathrm{cmp}} = \int d^4x \left\{ \mathfrak{E}_a{}^\mu \delta E^a{}_\mu + 2\,\mathrm{Re}(\delta\Psi^\dagger\mathcal{R}) - F^a{}_\mu\delta\lambda_a{}^\mu \right\}.$$

The connection solution must be differentiated compatibly. Setting only its point Euler value to zero while selecting unrelated connection derivatives does not define this reduced field theory. Its sourced Cartan curvature is not generally the Levi–Civita curvature of E.

10.2 Ward Identities and a Conditional Subsidiary System

For compactly supported coordinate variations, the spinor is a coordinate scalar and λ is a vector density. Its variation includes $\lambda_a{}^\mu\partial_\nu\xi^\nu$, in addition to transport and contravariant-index terms. Integration by parts therefore gives

$$\begin{aligned} 0 = {} & E^a{}_\nu\partial_\mu\mathfrak{E}_a{}^\mu + \mathfrak{E}_a{}^\mu(\partial_\mu E^a{}_\nu - \partial_\nu E^a{}_\mu) \\ & + \lambda_a{}^\mu(\partial_\mu F^a{}_\nu - \partial_\nu F^a{}_\mu) + F^a{}_\nu\partial_\mu\lambda_a{}^\mu - 2\,\mathrm{Re}((\partial_\nu\Psi)^\dagger\mathcal{R}). \qquad (2) \end{aligned}$$

For $J^T\eta + \eta J = 0$ and $S_J = -(\eta J)_{ab}\Gamma^a\Gamma^b/4$, Lorentz invariance similarly gives

$$\mathfrak{E} : (JE) + 2\,\mathrm{Re}((S_J\Psi)^\dagger\mathcal{R}) + F : (J^T\lambda) = 0.$$

These identities retain all residual sources. Their source-free specialization requires $F = \mathcal{R} = 0$ as field equations on an open region, not merely at the prepared point.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proposition 10.1 (Conditional gravitational constraint propagation)**.** *Assume exact compatible Cartan elimination, $F = \mathcal{R} = 0$ on an open region, a sufficiently smooth Lorentzian metric with spacelike slices, and the spatially projected trace-reversed coframe Euler evolution equations. With zero initial gravitational constraints and no incoming constraint flux, the four gravitational constraints remain zero. This statement does not propagate F or $\mathcal{R}$ from initial data.*

Proof. The two identities make the coframe Euler tensor density symmetric and Levi–Civita divergence-free under the stated hypotheses. Divide out the volume density to obtain a symmetric residual tensor $T^{\mu\nu}$ with $\nabla_\mu T^{\mu\nu} = 0$. For the unit slice normal n, put $H = T(n, n)$ and $M_i = -T(n, \partial_i)$. The spatial projection of $T^{\mu\nu} - g^{\mu\nu}\,\mathrm{tr}_g\, T/2$ vanishes by the chosen evolution equations and gives spatial stress $P^{ij} = Hh^{ij}$. In lapse $N > 0$, shift β^i and positive slice metric h_{ij}, the principal symbol for $U = (H, M_i)$ is

$$A(\xi) = -(\beta \cdot \xi) I_4 + N \begin{pmatrix} 0 & (h^{-1}\xi)^T \\ \xi & 0_{3\times 3} \end{pmatrix}, \qquad W = \mathrm{diag}(1, h^{-1}) > 0.$$

Since $WA(\xi)$ is symmetric, the energy $\frac{1}{2}\int (H^2 + h^{ij} M_i M_j)\sqrt{\det h}\, d^3x$ obeys $\mathcal{E}'(t) \le C(t)\mathcal{E}(t)$ with bounded coefficient derivatives and no incoming flux. Gronwall's inequality proves the claim. With nonzero shift the evolution condition is the geometric spatial projection, not simply the vanishing of contravariant coordinate ij components. The full lower-order lapse, shift and connection terms must be retained in the energy estimate. □

The saved Taylor fields do not satisfy the open-set hypotheses of this proposition. In particular, the constitutive identity

$$\partial_0 F^a{}_i = \partial_i F^a{}_0 + (\partial_0 F^a{}_i - \partial_i F^a{}_0)$$

does not close evolution unless its mixed-curl source is controlled. Ward identities alone are not a full coupled propagation theorem.

10.3 Finite Normal Reduction with Exposed Source Coordinates

At the prepared normal point $E = I$, $\partial E = 0$, hold the lower and spatial data and all action coefficients fixed. The real normal unknowns are

$$x = (E_{,00}\ [16],\ \text{realify}\,\Psi_{,0}\ [24], \lambda_{,0}\ [16]) \in \mathbb{R}^{56}.$$

The 16 constitutive, 24 real probe and 16 coframe Euler rows, together with twelve mixed constitutive curls, form an affine residual $b + Nx \in \mathbb{R}^{68}$. The curls are necessary consequences of $F = 0$ as a field equation, not twelve independent additional action laws. The numerical rank rises from 34 for the original rows to 40 with the curls. Its kernel has an exhibited ten-dimensional leading normal gauge span and a six-dimensional complementary span. This is not a complete regional gauge reduction.

Proposition 10.2 (Compatible finite normal inverse)**.** *Assume* $\mathrm{rank}\, N = 40$*, and orthonormal matrices $G \in \mathbb{R}^{56\times 10}$, $U \in \mathbb{R}^{56\times 6}$ with $\ker N = \mathrm{im}\, G \oplus \mathrm{im}\, U$. Let L_N span $\ker N^T$ orthonormally. Then $Nx = r$ is solvable if and only if $L_N^T r = 0$. Every solution is uniquely parametrized by*

$$x = N^+ r + Gg + Uq, \qquad g \in \mathbb{R}^{10}, \quad q \in \mathbb{R}^6.$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. $\operatorname{im} N = (\ker N^T)^\perp$ gives compatibility. For compatible r, $NN^+r = r$; subtracting this particular solution leaves exactly $\ker N$. Orthogonality gives unique g, q. For incompatible r the same expression leaves residual $-L_N L_N^T r$; a least-squares output is not an exact solution. □

There are 28 left compatibility rows on arbitrary r, not necessarily 28 independent physical field laws. For increments about the saved jet, r is the target residual minus its saved value, rather than the absolute right-hand side $-b$. The chosen Uq changes only $E_{,00}$ and the spatial-column entries ${\lambda_{a,0}}^i$, leaving all $\Psi_{,0}$ and ${\lambda_{a,0}}^0$ fixed. These six coordinates are neither the earlier mixed-curl defects nor established gauge or physical modes.

10.4 Determination from Compatible Tangential Jets

Use the same current saved joint jet throughout. Let q now be a source increment with gauge and normal residual target fixed. Let z contain increments of $(\Psi_{,2}, \lambda_{,2}, E_{,3}, \Psi_{,3})$ in the real unique-symmetric convention, of dimensions $(240, 160, 320, 480)$. Direct differentiation of the current equations gives the jointly affine 384-row residual

$$\mathcal{H}(q, z) = h_0 + Bq + Az. \tag{3}$$

Its blocks are first constitutive derivatives (64), first real probe derivatives (96), first coframe Euler derivatives (64), and second constitutive derivatives in unique pairs (160). Affinity is specific to this source variation and order: the lower spinor and its first jet are fixed, so variable factors multiply fixed lower factors. It does not extend automatically to later derivative equations.

For all 1200 higher coordinates allowed, $\operatorname{rank} A = 332$ and $\operatorname{rank}(L_A^T B) = 0$ numerically, where L_A spans $\ker A^T$. Thus the frozen-higher rank-six response of B can be absorbed by higher increments at this order. It does not fix q.

Instead preserve the represented finite tangential germ of E, $E_{,0}$, Ψ and λ on $x^0 = 0$. Split z into free coordinates w and prescribed tangential coordinates d. With $i, j, k \in \{1, 2, 3\}$, the free slots and real dimensions are

block	allowed derivative indices	dimension
$\Psi_{,2}$	at least one 0	96
$\lambda_{,2}$	at least one 0	64
$E_{,3}$	at least two 0	64
$\Psi_{,3}$	at least one 0	240

Thus $w \in \mathbb{R}^{464}$ and $d \in \mathbb{R}^{736}$. In particular $E_{,0ij}$ is prescribed because it differentiates the fixed $E_{,0}$ trace tangentially. This is a finite germ at a point, not a construction of admissible surface functions. Write $Az = A_f w + A_d d$.

Proposition 10.3 (Conditional source law for prescribed tangential data)**.** *Let L_f span $\ker A_f^T$ orthonormally, set*

$$J = L_f^T B, \qquad b(d) = -L_f^T (h_0 + A_d d),$$

and assume $\operatorname{rank} J = 6$. *A source and higher lift exist if and only if $(I - JJ^+)b(d) = 0$. For compatible d the source is unique:*

$$q(d) = J^+ b(d), \qquad w = -A_f^+ (h_0 + A_d d + Bq(d)) + k, \quad k \in \ker A_f.$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. Eliminating w requires and suffices for $L_f^T(h_0 + A_d d + Bq) = 0$, namely $Jq = b(d)$. Full column rank makes q unique if this equation is consistent. The orthogonal projector JJ^+ tests that consistency; the remaining formula gives exactly all higher lifts. □

The numerical response for this mask has rank 200 and the six singular values of J are approximately

$$1.89152,\ 1.22410,\ 0.48769,\ 0.45652,\ 0.26352,\ 0.13424.$$

The residual compatibility map on d has numerical rank 126. This is a finite matrix rank, not a count of physical constraints. All ranks are pointwise floating calculations with declared cutoffs, not certified regional bounds. Incompatible prescribed data are rejected, not silently projected onto admissible ones. In particular $q = 0$ is not an added law; near-zero increments for unchanged data refer to the saved germ only.

10.5 Reproducibility, Negative Control and Remaining Obligations

Four compatible-data controls use $q = a(1, -0.5, 0.3, 0.2, 0.4, -0.1)$ with $a \in \{0, 0.01, -0.01, 0.02\}$. An unrestricted compatible lift supplies the prescribed d, which is then held exactly fixed while the conditional law recovers q. Recovery errors are below 2.5×10^{-15}; direct residual norms are below 2.4×10^{-14} for all 68 normal rows and 3.7×10^{-11} for all 384 higher rows. These controls verify a known compatible family, not arbitrary surface admissibility. The computed $\|h_0\| \simeq 4.78 \times 10^{-11}$ is retained in the law.

A separate saved completion satisfies all 1114 packed rows of the later joint hierarchy to residual about 5.38×10^{-10}. The new source lifts do not automatically preserve it. For $a = 0.01$, retaining its old highest $E_{,4}, \lambda_{,3}, \Psi_{,4}$ gives a later residual about 98.96, with third constitutive block about 98.60, even after recomputing the changed Levi–Civita cubic. This is a failed frozen-highest control, not a nonexistence proof for a re-prepared completion. The earlier coefficient-domain neighborhood estimate does not transfer to the new witnesses without recalculation.

The source files and pinned numerical reports are `analysis/HC_Cartan_Normal_Source_Reduction.py` and `analysis/HC_Cartan_Normal_Source_Compatibility.py`, with their matching JSON reports and derivation notes. The latter module uses the fingerprint-checked current joint loader, not the older regular-point completion. Sixteen source-compatibility and thirteen normal-reduction tests supplement the full audit; a test count is not a proof certificate.

For a future smooth normal chart $x = x_{\mathrm{part}} + Gg + Uq$, spatial differentiation must retain

$$\partial_i x = \partial_i x_{\mathrm{part}} + (\partial_i G)g + G\partial_i g + (\partial_i U)q + U\partial_i q.$$

A QR/SVD frame at one point supplies neither these derivatives nor arbitrary source functions. The next obligations are compatible surface data, higher re-preparation with the prescribed tangential germ retained, a smooth gauge/source chart, and a closed full evolution and subsidiary estimate. No regional existence, full hyperbolicity, native-action identification, OP5 closure or physical scale follows from the finite source law alone.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Nomenclature

Status. *Reader-support block.*

Block function. This section is a navigation aid: it fixes symbols and abbreviations so later sections can be read without repeatedly reopening earlier definitions.

Spaces and operators

H^+ Hardy space; physical Hilbert space

H^+_+ J-symmetric (bosonic) sector

H^+_- J-antisymmetric (fermionic) sector

L_0 Self-adjoint generator on H^+_+

J Fundamental involution $J : s \mapsto 1 - \bar{s}$

$C(f)$ Heisenberg compensator: $\frac{1}{2}(f + Jf)$

$A(f)$ Antisymmetric component: $\frac{1}{2}(f - Jf)$

$\hat{K}$ Cascade operator $\hat{K} : S_n \to S_{n-1}$

$\mathcal{W}(A^1)$ Weyl algebra $\mathcal{W}(A^1)$

Geometric quantities

q_n Biquaternion ground state $\in \mathbb{H} \otimes \mathbb{C}$

τ_n Proper time: $\tau_n^2 = |q_n|^2$

z^- Frame variable; displacement from $z^- = 0$

K_1, K_2, K_3 Matter, tachyon, photon kernels

L_n Light axis of stratum S_n

S_n Stratum n; $S_0 = \{z^- = 0\}$ photon ground state

Frame parameters

β_{sym} Structural frame offset $\sqrt{(\gamma_L - 1)/(\gamma_L + 1)} \approx 0.5493$

γ_L Frame Lorentz factor $\gamma_{\text{BI}}/\gamma_{\text{string}} \approx 1.864$

γ_{BI} Barbero–Immirzi parameter ≈ 0.2375

γ_{string} String coupling $\ln 2/(\pi\sqrt{3}) \approx 0.1274$

$k(X)$ Frame-insertion count; $X^{(\gamma)} = \gamma_L^{-k} \cdot X^{(m)}$

Physical constants (matter-frame CODATA)

ℓ_P Planck length 1.616×10^{-35} m; $k = 3/2$

α Fine-structure constant; $k(\alpha^{-1}) = 1$

$\hbar$ Reduced Planck constant; $k(\hbar) = 1$

G_N Gravitational constant; $k(G_N) = 2$

c Speed of light; $k(c) = 0$ (frame-invariant)

Lagrangian and field structure

$\mathcal{L}_{\text{kin}}$ Kinetic term $\langle f, L_0 f \rangle$

$\mathcal{L}_{\text{HC}}$ HC confinement term

$\mathcal{L}_{\text{frame}}$ Frame term $\beta_{\text{sym}}^2 \langle f, f \rangle$; source of Λ_{obs}

$\mathcal{M}_{16}$ 16×16 compensator matrix (formerly Millennium matrix)

$\hat{P}_{C_7}$ Meta-Septum closure projector

$\hat{\mathcal{R}}_\beta$ Shifted GUE pacer $\hat{R}_{\text{GUE}}\, e^{i\pi\beta_{\text{sym}}}$

Cosmological quantities

Ω_Λ Dark energy fraction $1 - \beta_{\text{sym}}^2 \approx 2/3$

Ω_M Matter fraction $\beta_{\text{sym}}^2 \approx 1/3$

$\Sigma(\tau)$ Trace formula $1 - \Theta(\tau) + R(\tau)$

Abbreviations

GUE Gaussian Unitary Ensemble ($\beta = 2$)

HC Heisenberg compensator

LQG Loop Quantum Gravity (H^+_- sector)

RH Riemann Hypothesis ($z^- = 0$ for all zeros)

SM Standard Model $U(1) \times SU(2) \times SU(3)$

v4.6 release-status and field-equation symbols.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Symbol	**Role**	**Reader meaning**
$\mathsf{BRead}_{\mathrm{bounded}}$	bounded readout surface	conservative declared-grammar readout obtained in the v4.6.0 release-final line
Θ^{all}	all-readout demand	unrestricted family of readouts required for T2; not discharged by the bounded surface
$\mathfrak{O}_{\mathrm{all}}^{4.6.0}$	all-readout obstruction	frozen scope-boundary register for the missing universal exterior selector, naturality family, and unrestricted collar slot
$\mathcal{M}_{\mathrm{bd}}, \mathcal{M}_{\mathrm{ext}}, \mathcal{M}_{\mathrm{mix}}$	MM split	bounded, exterior, and mixed compensator-matrix portions
$\mathcal{H}_{\mathrm{bd}}, \mathcal{H}_{\mathrm{ext}}, \mathcal{H}_{\mathrm{mix}}$	HC split	bounded, exterior, and mixed Heisenberg-compensator portions
$\mathfrak{E}_{\mathrm{decl}}$	declared field-equation template	structure-first reset template linking structural, HC, MM, QAM, and readout terms

© 2026 Marc Brendecke. Preprint — not peer reviewed.

11 Alternating Triolic Orbit Scan and Hopf-Fibre Core Surface Readout

Reader status. This section is a reader-facing geometric normal form. It is not a claim that ordinary physical particles literally cross a collapse core. It gives a controlled local scan model: test trajectories are used as readout curves around a guarded core, and the remaining question is whether the resulting local surface data can be promoted into the global HC/MM-to-chart bridge without residual debt.

11.1 Triolic scan construction

Let K denote the guarded collapse core. The local lens geometry is represented by two conjugate focal lenses

$$F_-, \qquad F_+,$$

and by their midpoint

$$M := \frac{1}{2}(F_- + F_+).$$

The first triolic balance is the focal-lens triol

$$\mathcal{T}_{\text{lens}} := (F_-, K, F_+),$$

where K is not treated as a point to be crossed, but as the balancing object between the two focal lenses.

A test-orbit scan uses paired test readouts

$$P_n, \qquad Q_n$$

on a bounded local orbit family. At selected readout phases one obtains the particle triol

$$\mathcal{T}_n := (P_n, K, Q_n).$$

The triol is called *core-sighted* when the sightline from P_n to Q_n passes through K. In affine notation this means that there is a scalar λ_n with

$$K = (1 - \lambda_n)P_n + \lambda_n Q_n.$$

If $\lambda_n = \frac{1}{2}$, the triol is centrally balanced; otherwise it is projectively balanced. Thus the model separates two cases which should not be confused:

$$\text{collinearity through } K \qquad \neq \qquad \text{midpoint symmetry at } K.$$

Definition 11.1 (Bifocal elliptic readout body)**.** *A local orbit curve γ is a bifocal elliptic readout body for the lens pair (F_-, F_+) if, on the admissible scan domain,*

$$d(\gamma(s), F_-) + d(\gamma(s), F_+) = 2a$$

up to the declared tolerance of the local grammar. The curve is used as a stable computational body, not as an unrestricted through-core physical trajectory.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

11.2 Shift as a Kronecker-style transport selector

The discrete shift is represented by a shifted selector rather than by the identity Kronecker tensor. In an indexed scan basis one writes

$$\Sigma^i{}_j := \delta^i_{j+1},$$

so that the k-fold shift is

$$(\Sigma^k)^i{}_j = \delta^i_{j+k}.$$

Thus δ^i_j preserves an index, while $\Sigma^i{}_j$ transports the readout to the next admissible scan state.

The parity projectors are

$$\Pi_{\text{even}}(n) = \delta_{n \bmod 2,0}, \qquad \Pi_{\text{odd}}(n) = \delta_{n \bmod 2,1}.$$

An alternating triolic scan has two structure classes $\mathcal{A}$ and $\mathcal{B}$ with

$$\Sigma(\mathcal{A}) = \mathcal{B}, \qquad \Sigma(\mathcal{B}) = \mathcal{A},$$

and therefore

$$\Sigma^2(\mathcal{A}) = \mathcal{A}, \qquad \Sigma^2(\mathcal{B}) = \mathcal{B}.$$

The single shift is the toggle; the double shift is the same-regime evolution step.

Assumption 11.2 (Alternating triolic shift hypothesis)**.** *Along a guarded bifocal scan there are two complementary triolic regimes:*

$$\mathcal{A}_{2n} := \mathcal{T}_{2n}, \qquad \mathcal{B}_{2n+1} := \mathcal{T}_{2n+1}.$$

One regime is core-sighted and the other is lens-sighted, or conversely. The local shift operator Σ exchanges these regimes, while Σ^2 preserves the regime type.

11.3 Orbit scan and guarded core surface

For each scan curve $\gamma_{\theta,\varphi}$ around the guarded core, define its minimal core distance by

$$\rho_{\min}(\theta,\varphi) := \min_s d(\gamma_{\theta,\varphi}(s), K).$$

The effective measured core surface is the periapsis envelope

$$\Sigma_K := \text{env}\{\gamma_{\theta,\varphi}\}$$

or, equivalently, the surface reconstructed from the field of minimal distances $\rho_{\min}(\theta,\varphi)$.

Definition 11.3 (Guarded orbit-scan surface)**.** *A guarded orbit-scan surface is the readout surface obtained from all admissible parity-shifted test orbits whose minimal distance from K is positive:*

$$\rho_{\min}(\theta,\varphi) > 0.$$

The inequality states that the scan measures the effective boundary of the guarded core; it does not license a trajectory through the collapse state.

Proposition 11.4 (Periapsis envelope as core-surface readout)**.** *Suppose a scan family satisfies the bifocal elliptic readout condition, the alternating triolic shift hypothesis, and the guarded-core inequality $\rho_{\min} > 0$. Then the scan does not measure a through-core path. It measures a bounded periapsis envelope around K. This envelope is the local surface data available to the HC/MM-to-chart bridge.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. The bifocal condition supplies a stable bounded readout body. The alternating shift separates the scan into two complementary regimes, so that the same geometric orbit is not forced to carry all sightline conditions at once. The condition $\rho_{\min} > 0$ excludes a through-core trajectory. Therefore the admissible data are the nearest admissible approach points of the scan family, i.e. the periapsis envelope. This is exactly the effective local surface readout around the guarded core. □

11.4 Hopf-fibre interpretation and the possible residual point

The Hopf picture gives the natural topological comparison. In the ideal Hopf model, a three-sphere is decomposed into circular fibres. In that idealized fibre space there is no missing ordinary point: every point lies on a fibre. Apparent exceptional points may arise only after projection into a lower-dimensional picture.

The guarded collapse model is different. If the HC guard excludes the core itself, the scan space is not the full ideal fibre space but a punctured or core-excised domain,

$$X = S^3 \setminus \{K\} \qquad \text{or} \qquad X = S^3 \setminus B_\epsilon(K).$$

In that case a residual point or residual core is not a defect of the Hopf fibration. It is the excluded guarded kernel which the orbit scan is designed to measure from outside.

Proposition 11.5 (Hopf completion versus guarded-core remainder)**.** *There are two distinct readings of the ring filling picture.*

(a) *In the ideal Hopf reading, circular fibres fill the ambient three-sphere, and no ordinary minimal leftover point remains.*

(b) *In the guarded-core reading, the core K is removed or shielded. A minimal leftover point or core neighbourhood can remain, but it is not an unfilled Hopf gap; it is the HC-guarded kernel itself.*

Remark 11.6 (Operational meaning)**.** *The useful experimental image is therefore not "send a particle through the collapse". It is: send admissible test readouts around the guarded region, track their even/odd triolic alternation, collect the minimal approach points, and reconstruct the effective core surface from the resulting periapsis envelope. If the linked orbit family behaves Hopf-like, the linking and twisting of the rings provide additional topological data; if a residual core remains, that remainder is the object being measured.*

11.5 Finite scan checklist

For a finite numerical or symbolic probe, the following quantities are the minimal diagnostic payload:

$$\begin{aligned} E(s) &:= d(\gamma(s), F_-) + d(\gamma(s), F_+), \\ \rho_K(s) &:= d(\gamma(s), K), \\ \chi_n &:= \begin{cases} 1, & P_n, K, Q_n \text{ are collinear}, \\ 0, & \text{otherwise}, \end{cases} \\ \pi_n &:= n \bmod 2. \end{aligned}$$

The expected signature of the alternating scan is

$$E(s) \approx 2a, \qquad \rho_K(s) > 0, \qquad \chi_{n+2} = \chi_n,$$

with complementary behaviour between the even and odd subsequences. A successful scan therefore yields three outputs: the local bifocal fit, the parity/toggle law, and the reconstructed guarded-core surface.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

11.6 Strict mathematical orbit-scan formulation

This subsection records the strict mathematical form of the preceding geometric picture. It is a readout model. No physical through-core trajectory is asserted.

Definition 11.7 (Guarded scan space)**.** *Let (X, d) be a compact metric scan chart or a compact chart domain inside a larger metric space. Let K be a distinguished guarded core and let $\varepsilon > 0$ be a guard radius. The admissible scan domain is*

$$X_K := X \setminus B_\varepsilon(K), \qquad B_\varepsilon(K) := \{x \in X : d(x, K) < \varepsilon\}.$$

An admissible test orbit is a continuous curve

$$\gamma : S^1 \longrightarrow X_K.$$

Thus every admissible orbit satisfies

$$d(\gamma(s), K) \geq \varepsilon \qquad \text{for all } s \in S^1.$$

Definition 11.8 (Bifocal defect)**.** *Fix two focal points or focal charts $F_-, F_+ \in X$ and a constant $a > 0$. The bifocal defect of a scan point $x \in X_K$ is*

$$\mathcal{E}_F(x) := \big|d(x, F_-) + d(x, F_+) - 2a\big|.$$

A scan orbit is η-bifocal if

$$\sup_{s \in S^1} \mathcal{E}_F(\gamma(s)) \leq \eta.$$

The exact ellipse is the special case $\eta = 0$ in a chart where the focal distance expression is exact.

Definition 11.9 (Triolic sight predicate)**.** *Let $P, Q \in X_K$. In a geodesic chart, the ordered triple (P, K, Q) is called core-sighted if K lies on an admissible geodesic segment from P to Q. In an affine chart this is equivalent to the existence of $\lambda \in [0, 1]$ such that*

$$K = (1 - \lambda)P + \lambda Q.$$

It is centrally balanced if, in addition, $\lambda = 1/2$. The two predicates are kept separate:

$$\mathrm{Sight}_K(P, Q) \quad \not\Rightarrow \quad K = \frac{P + Q}{2}.$$

Definition 11.10 (Shift selector and parity sectors)**.** *Let I be a finite cyclic index set or $I = \mathbb{Z}$. On the scan basis $(e_j)_{j \in I}$ define the shift selector by*

$$\Sigma e_j = e_{j+1}, \qquad \Sigma^i{}_j := \delta^i_{j+1}.$$

For $k \geq 0$,

$$(\Sigma^k)^i{}_j = \delta^i_{j+k}.$$

Let Π_0 and Π_1 be the even and odd parity projectors,

$$\Pi_0 e_j = \begin{cases} e_j, & j \equiv 0 \pmod 2, \\ 0, & j \equiv 1 \pmod 2, \end{cases} \qquad \Pi_1 e_j = \begin{cases} 0, & j \equiv 0 \pmod 2, \\ e_j, & j \equiv 1 \pmod 2. \end{cases}$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Lemma 11.11 (Parity exchange law). *The shift selector exchanges parity sectors, while the double shift preserves them:*

$$\Pi_0\Sigma = \Sigma\Pi_1, \qquad \Pi_1\Sigma = \Sigma\Pi_0,$$

and

$$\Pi_r\Sigma^2 = \Sigma^2\Pi_r \qquad (r = 0, 1).$$

Proof. It is enough to test the identities on basis vectors. Since $\Sigma e_j = e_{j+1}$, the parity of the index is flipped by Σ and preserved by Σ^2. The displayed identities are exactly this parity statement in projector form. □

Definition 11.12 (Alternating triolic scan). *Let $(P_n, Q_n)_{n\in I}$ be paired scan readouts. An alternating triolic scan consists of two disjoint regime predicates $\mathcal{A}$ and $\mathcal{B}$ on pairs (P, Q) such that*

$$\mathcal{A}(P_n, Q_n) \Longleftrightarrow n \equiv 0 \pmod 2, \qquad \mathcal{B}(P_n, Q_n) \Longleftrightarrow n \equiv 1 \pmod 2,$$

up to a declared finite tolerance set. One admissible interpretation is

$$\mathcal{A} = \textit{core-sighted regime}, \qquad \mathcal{B} = \textit{lens-sighted regime},$$

although the formal definition only requires two complementary predicates exchanged by Σ.

Definition 11.13 (Periapsis envelope). *Let U be a compact parameter space of admissible scan orbits*

$$\Gamma : U \times S^1 \longrightarrow X_K, \qquad \gamma_u(s) := \Gamma(u, s).$$

For each $u \in U$ define

$$\rho(u) := \min_{s\in S^1} d(\gamma_u(s), K)$$

and the periapsis set

$$\mathcal{P}(u) := \{\gamma_u(s) : d(\gamma_u(s), K) = \rho(u)\}.$$

The orbit-scan surface is the closed periapsis envelope

$$\Sigma_K^\Gamma := \overline{\bigcup_{u\in U} \mathcal{P}(u)}.$$

Proposition 11.14 (Existence and guardedness of the scan surface). *For a compact scan family $\Gamma : U \times S^1 \to X_K$, the function $\rho(u)$ and the sets $\mathcal{P}(u)$ are well defined. Moreover,*

$$\rho(u) \geq \varepsilon \qquad \textit{for all } u \in U,$$

so the periapsis envelope Σ_K^Γ is an outside readout of the guarded core and not a through-core trajectory.

Proof. For fixed u, the map $s \mapsto d(\gamma_u(s), K)$ is continuous on the compact set S^1, hence it attains a minimum. Because $\gamma_u(S^1) \subset X_K$, every scan point has distance at least ε from K. Therefore each periapsis point remains outside the excluded core. Taking the union and closure gives a surface readout generated by nearest admissible approaches, not by points inside the guarded core. □

Theorem 11.15 (Strict alternating triolic orbit-scan theorem). *Assume the following data:*

(i) *a guarded scan space $X_K = X \setminus B_\varepsilon(K)$ with $\varepsilon > 0$;*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(ii) *a compact family of admissible orbits* $\Gamma : U \times S^1 \to X_K$*;*

(iii) *paired readouts* (P_n, Q_n) *sampled from the scan family;*

(iv) *an alternating triolic scan structure exchanged by the selector* $\Sigma^i{}_j = \delta^i_{j+1}$*;*

(v) *a bifocal defect bound* $\sup_s \mathcal{E}_F(\gamma_u(s)) \le \eta$ *for all* $u \in U$*.*

Then the scan determines three mathematically controlled readout objects:

(a) the bifocal defect field $\mathcal{E}_F$,
(b) the parity law Π_0, Π_1, Σ,
(c) the guarded periapsis envelope Σ^{Γ}_K.

It does not determine, by itself, a physical through-core orbit or a global closure of the HC/MM-to-chart bridge.

Proof. The bifocal defect field is defined by the focal data and the metric. The parity law follows from Lemma 11.11 and the alternating scan predicates. The guarded periapsis envelope exists and remains outside the excluded core by Proposition 11.14. None of these constructions supplies a point of any scan curve inside $B_\varepsilon(K)$, and none asserts compatibility with a global bridge unless an additional promotion theorem is supplied. Hence the outputs are exactly local readout objects with a guarded-core interpretation. □

Proposition 11.16 (Hopf comparison without a hidden gap)**.** *Let* $\pi : S^3 \to S^2$ *be the Hopf fibration. In the ideal Hopf space every point of* S^3 *lies on exactly one circular fibre. Therefore a leftover point is not produced by the Hopf fibration itself. If the scan domain is instead* $S^3 \setminus B_\varepsilon(K)$ *or* $S^3 \setminus \{K\}$*, any residual core is the explicitly removed guarded set, not an unfilled Hopf gap.*

Proof. The Hopf fibration partitions S^3 into fibres $\pi^{-1}(y) \cong S^1$ indexed by $y \in S^2$. Hence each point of S^3 belongs to one fibre. Removing K or a neighbourhood of K changes the domain before the scan is performed. The remaining excluded set is therefore part of the guarded-core model, not a failure of the Hopf decomposition. □

Remark 11.17 (Promotion guard)**.** *The strict formulation proves only local scan coherence: bifocal control, parity alternation, and an outside periapsis envelope. A global physical or theorematic promotion would require an additional result showing that the envelope data kill, strictly reduce, or otherwise control the remaining route-to-global obstruction without hidden leakage or residual reopening.*

11.7 Executable finite orbit-scan probe layer

The preceding strict formulation is now turned into a finite probe grammar. The purpose is not to replace the continuous definitions, but to give a computable audit surface: a finite family of sampled orbits, a finite parity test, a finite bifocal defect table, and a finite periapsis-envelope estimator.

Definition 11.18 (Finite scan grid)**.** *Let* $U_N = \{u_1, \dots, u_N\} \subset U$ *be a finite set of admissible orbit parameters and let*

$$S_M = \{s_0, \dots, s_{M-1}\} \subset S^1, \qquad s_m = \frac{2\pi m}{M}.$$

The finite scan sample is the table

$$x_{\ell m} := \Gamma(u_\ell, s_m) \in X_K.$$

It is admissible when

$$d(x_{\ell m}, K) \geq \varepsilon \qquad (1 \leq \ell \leq N,\, 0 \leq m < M).$$

Definition 11.19 (Finite bifocal and guarded-distance registers)**.** *For a finite scan grid define*

$$E_{\ell m} := \mathcal{E}_F(x_{\ell m}), \qquad R_{\ell m} := d(x_{\ell m}, K) - \varepsilon.$$

The finite bifocal defect and finite guard-margin defect are

$$E_{N,M} := \max_{\ell,m} E_{\ell m}, \qquad R^{\min}_{N,M} := \min_{\ell,m} R_{\ell m}.$$

The grid passes the local readout screen at tolerances (η, μ) when

$$E_{N,M} \leq \eta, \qquad R^{\min}_{N,M} \geq \mu \geq 0.$$

Definition 11.20 (Finite parity and triolic indicators)**.** *Let $(P_n, Q_n)_{n=0}^{L-1}$ be the sampled paired readouts. Let*

$$\chi_A(n), \chi_B(n) \in \{0, 1\}$$

be the finite indicators for the two complementary triolic regimes. The finite alternation defect is

$$D_{\mathrm{alt}} := \sum_{n=0}^{L-1} \Big| \chi_A(n) - \mathbf{1}_{n \equiv 0\,(2)} \Big| + \sum_{n=0}^{L-1} \Big| \chi_B(n) - \mathbf{1}_{n \equiv 1\,(2)} \Big|.$$

The finite scan is parity-clean when

$$D_{\mathrm{alt}} = 0.$$

Definition 11.21 (Finite periapsis estimator)**.** *For each sampled orbit u_ℓ define*

$$\rho_{\ell,M} := \min_{0 \leq m < M} d(x_{\ell m}, K)$$

and the finite periapsis sample

$$\mathcal{P}_{\ell,M} := \{x_{\ell m} : d(x_{\ell m}, K) = \rho_{\ell,M}\}.$$

The finite orbit-scan surface estimator is

$$\Sigma_K^{N,M} := \bigcup_{\ell=1}^{N} \mathcal{P}_{\ell,M}.$$

Lemma 11.22 (Finite shift-selector audit)**.** *On a cyclic sample of length L, the matrix*

$$\Sigma_L{}^i{}_j = \delta^i_{j+1 \bmod L}$$

exchanges the finite even and odd sectors when L is even, and the double shift preserves them:

$$\Pi_0 \Sigma_L = \Sigma_L \Pi_1, \qquad \Pi_1 \Sigma_L = \Sigma_L \Pi_0, \qquad \Pi_r \Sigma_L^2 = \Sigma_L^2 \Pi_r.$$

Proof. The proof is the finite cyclic version of Lemma 11.11. Since L is even, adding one modulo L flips parity and adding two modulo L preserves parity. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proposition 11.23 (Finite probe certificate). *A finite orbit-scan probe with data*

$$\mathcal{C}_{N,M,L} := \left(E_{N,M}, R^{\min}_{N,M}, D_{\mathrm{alt}}, \Sigma^{N,M}_K\right)$$

is a local finite scan certificate at tolerances (η, μ) if

$$E_{N,M} \le \eta, \qquad R^{\min}_{N,M} \ge \mu \ge 0, \qquad D_{\mathrm{alt}} = 0.$$

Such a certificate verifies only the finite local readout: bifocal fit, guarded outside motion, and even/odd triolic alternation. It does not certify a through-core orbit and does not certify global HC/MM-to-chart closure.

Proof. The three inequalities are exactly the finite versions of bifocal control, guarded-domain admissibility, and parity-clean alternation. The periapsis set $\Sigma^{N,M}_K$ records the measured outside surface points. No sample point lies inside the guarded core, and no bridge-promotion hypothesis is part of the finite certificate. Therefore the certificate is local and diagnostic only. □

Proposition 11.24 (Consistency of the finite periapsis estimator). *Assume U is compact, Γ is continuous, the finite sets U_N become dense in U, and the circular grids S_M become dense in S^1. Then every cluster point of finite periapsis samples $\Sigma^{N,M}_K$ belongs to the continuous periapsis envelope Σ^{Γ}_K. If, in addition, each orbit has a unique nondegenerate periapsis depending continuously on u, then $\Sigma^{N,M}_K$ converges to Σ^{Γ}_K in Hausdorff distance.*

Proof. Let $x_{\ell m}$ be a convergent sequence of finite periapsis samples with parameters $u_\ell \to u$ and grid points $s_m \to s$. By continuity, $x_{\ell m} \to \Gamma(u, s)$. Since each $x_{\ell m}$ minimizes the sampled distance along its finite grid and the grids are dense, the limiting point is a minimizer of $s \mapsto d(\Gamma(u, s), K)$; hence it lies in Σ^{Γ}_K. Under uniqueness and nondegeneracy of the minimizer, the argmin varies continuously, which upgrades cluster containment to Hausdorff convergence. □

Remark 11.25 (Executable toy chart). *A numerical toy chart may take $X \subset \mathbb{R}^3$, $K = 0$, two focal points $F_\pm = (\pm c, 0, 0)$ in a local plane, and a family of rotated ellipses*

$$\gamma_u(s) = R_u(a\cos s, b\sin s, 0), \qquad b > \varepsilon.$$

The toy chart tests the finite registers above. It is not a physical simulation; it is a reproducible diagnostic for the readout grammar.

Theorem 11.26 (Executable orbit-scan handoff). *The strict alternating triolic scan admits a finite executable handoff: choose (N, M, L), compute the registers $E_{N,M}$, $R^{\min}_{N,M}$, D_{alt}, and $\Sigma^{N,M}_K$, and report whether the finite certificate conditions hold. A successful finite handoff provides a controlled local scanner for the guarded core surface; a failed handoff identifies whether the failure is bifocal, guard-margin, or parity-alternation failure.*

Proof. All four registers are finite maxima, minima, sums, or finite sets over explicitly sampled data. Therefore they are computable. Proposition 11.23 gives the pass conditions and their interpretation. The failure modes are separated by which of the three inequalities fails. □

Remark 11.27 (Human reading). *In ordinary language: this layer says how to send many sampled ring-like test readouts around the guarded core, measure their nearest approaches, check whether the even/odd shift pattern is stable, and turn those nearest approach points into a finite surface estimate. It is the bridge from the strict definition to an actual table, plot, or numerical probe.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

11.8 Finite perturbation lemma for orbit-scan robustness

The executable scan layer separates two questions which should not be conflated. The shifted Kronecker selector

$$\Sigma_L{}^i{}_j = \delta^i_{j+1 \bmod L}$$

is an index-level rule and therefore remains exact under geometric perturbations. The triolic sightline condition is geometric and may fail when the core, the pairing rule, or the lens data are perturbed. The present subsection records this distinction as a finite robustness lemma.

Definition 11.28 (Finite perturbation register)**.** *Let $x_{\ell m}$ be an admissible finite scan grid as in Definition 11.18. A perturbed finite scan consists of data*

$$\widetilde{x}_{\ell m}, \qquad \widetilde{K}, \qquad \widetilde{F}_-, \widetilde{F}_+$$

together with the same cyclic index set of length L. Define the finite perturbation amplitudes

$$\Delta_X := \max_{\ell,m} d(\widetilde{x}_{\ell m}, x_{\ell m}), \qquad \Delta_K := d(\widetilde{K}, K), \qquad \Delta_F := d(\widetilde{F}_-, F_-) + d(\widetilde{F}_+, F_+).$$

The perturbed bifocal defect and guard reserve are

$$\widetilde{E}_{N,M} := \max_{\ell,m} \left| d(\widetilde{x}_{\ell m}, \widetilde{F}_-) + d(\widetilde{x}_{\ell m}, \widetilde{F}_+) - 2a \right|,$$

$$\widetilde{R}^{\min}_{N,M} := \min_{\ell,m} \big(d(\widetilde{x}_{\ell m}, \widetilde{K}) - \varepsilon \big).$$

Definition 11.29 (Finite triolic residuals under perturbation)**.** *For a paired sample $(\widetilde{P}_n, \widetilde{Q}_n)$ define the normalized core-sightline residual*

$$C_n := \frac{\|(\widetilde{P}_n - \widetilde{K}) \times (\widetilde{Q}_n - \widetilde{K})\|}{\max\{\|\widetilde{P}_n - \widetilde{K}\| \, \|\widetilde{Q}_n - \widetilde{K}\|, \tau\}}, \qquad \tau > 0.$$

Define also the midpoint residual

$$M_n := d\left(\widetilde{K}, \frac{\widetilde{P}_n + \widetilde{Q}_n}{2} \right).$$

For declared thresholds $c_{\mathrm{even}}, c_{\mathrm{odd}}, m_{\mathrm{even}} > 0$, the perturbed scan passes the finite triolic robustness screen when

$$\mathrm{p95}\{C_n : n \equiv 0\,(2)\} \le c_{\mathrm{even}},$$
$$\mathrm{p05}\{C_n : n \equiv 1\,(2)\} \ge c_{\mathrm{odd}},$$
$$\mathrm{p95}\{M_n : n \equiv 0\,(2)\} \le m_{\mathrm{even}}.$$

The even sector is then interpreted as core-sighted to the declared tolerance, while the odd sector remains a distinct complementary regime.

Lemma 11.30 (Index parity is perturbation-invariant)**.** *Let L be even and let $\Sigma_L{}^i{}_j = \delta^i_{j+1 \bmod L}$. For every geometric perturbation of the points, core, or foci, the identities*

$$\Pi_0 \Sigma_L = \Sigma_L \Pi_1, \qquad \Pi_1 \Sigma_L = \Sigma_L \Pi_0, \qquad \Pi_r \Sigma_L^2 = \Sigma_L^2 \Pi_r$$

remain unchanged.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. The operators Π_0, Π_1 and Σ_L act on the finite index set only. A perturbation changes the values assigned to the indices, but not the parity of the indices themselves. Hence the parity-exchange identities are invariant under all geometric perturbations of the sampled points. □

Lemma 11.31 (Guard robustness under small perturbations)**.** *Assume the unperturbed grid has guard reserve*

$$R^{\min}_{N,M} \geq \mu > 0.$$

If

$$\Delta_X + \Delta_K < \mu,$$

then the perturbed grid remains outside the protected core:

$$\widetilde{R}^{\min}_{N,M} > 0.$$

Proof. For every sample, the triangle inequality gives

$$d(\widetilde{x}_{\ell m}, \widetilde{K}) \geq d(x_{\ell m}, K) - d(\widetilde{x}_{\ell m}, x_{\ell m}) - d(\widetilde{K}, K) \geq \varepsilon + \mu - \Delta_X - \Delta_K.$$

If $\Delta_X + \Delta_K < \mu$, the right hand side is strictly larger than ε. □

Proposition 11.32 (Finite perturbation screen)**.** *A perturbed finite orbit scan is a stable local readout at thresholds*

$$(\eta, \mu, c_{\text{even}}, c_{\text{odd}}, m_{\text{even}})$$

when the following four screens pass:

$$\widetilde{E}_{N,M} \leq \eta, \qquad \widetilde{R}^{\min}_{N,M} \geq \mu \geq 0,$$

$$\mathrm{p95}\{C_n : n \equiv 0\,(2)\} \leq c_{\text{even}}, \qquad \mathrm{p05}\{C_n : n \equiv 1\,(2)\} \geq c_{\text{odd}},$$

$$\mathrm{p95}\{M_n : n \equiv 0\,(2)\} \leq m_{\text{even}}.$$

A failure of the first screen is a bifocal-lens defect, a failure of the second is a core-guard defect, and a failure of the last two is a triolic sightline or midpoint-balance defect.

Proof. The first inequality is precisely the finite bifocal readout requirement. The second inequality is precisely the guarded-domain requirement. The last two inequalities are the declared even/odd triolic separation requirements. Since the screens are independent finite statistics, the failed inequality identifies the local failure class without requiring a global physical promotion. □

Theorem 11.33 (Finite robustness handoff)**.** *The orbit-scan construction admits a finite robustness handoff. Symmetry-preserving perturbations may preserve the local scan certificate, while perturbations which move the core away from the balancing centre or break the antipodal pairing are detected as triolic obstructions. In all cases the shifted Kronecker selector remains index-clean by Lemma 11.30; what changes is the geometric quality of the triolic readout.*

Proof. The finite registers of Definition 11.28 and Definition 11.29 are computable from the sampled data. Lemma 11.30 fixes the index-level shift law independently of the geometry. Proposition 11.32 then separates bifocal, guard, and triolic failure modes. Hence perturbation stability is a finite diagnostic property, not an unobserved assumption. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Remark 11.34 (Executed finite robustness probe)**.** *A deterministic toy robustness probe was executed with seven deformation scenarios. The ideal chart, a symmetric radial ripple, a moderate lens asymmetry, a twist coupling, and a mild combined deformation passed the declared local readout screens. A displaced core failed the even core-sightline screen while preserving the bifocal and guard screens. A strong combined deformation failed both the bifocal mean-stability screen and the even core-sightline screen while still preserving the outside guard. The result supports the intended diagnostic reading: the shift selector remains exact, while the orbit geometry itself decides whether a valid triolic core scan is present.*

Scenario	Bifocal	Guard	Even triole	Odd distinct	Overall
Ideal chart	yes	yes	yes	yes	yes
Symmetric radial ripple	yes	yes	yes	yes	yes
Lens asymmetry	yes	yes	yes	yes	yes
Core offset	yes	yes	no	yes	no
Twist coupling	yes	yes	yes	yes	yes
Mild combined deformation	yes	yes	yes	yes	yes
Strong combined deformation	no	yes	no	yes	no

Remark 11.35 (Human reading)**.** *In ordinary language: the clock of the scan is robust because it is an index clock. The geometry is allowed to fail. That is desirable. If the core is shifted away from the balancing centre or if the antipodal relation of the paired readouts is broken, the finite scan reports a triolic obstruction instead of silently pretending that the same core surface is still being measured.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

12 The Central Theorem

Status. *Legacy structural-theorem block inside the active main body.*

Block function. This section states the global structural theorem that the later physics, operator, and readout sections unpack and reuse rather than replace.

Everything in this paper is a consequence of one observation.

Theorem 12.1 (Unity of the Sphaera)**.** *Within the Sphaera S, the following are all representations of the same object — the collapse state $z^- = 0$:*

(i) The Sphaera itself, as a geometric space.

(ii) The triolic structure: three kernels K_1, K_2, K_3 with $z_1 + z_2 + z_3 = 0$, phase-shifted by 120°, running over the Sphaera's geometry.

(iii) Each individual kernel K_i, as a local coordinate of the triole.

(iv) The Heisenberg compensator output $C(f)$, as the projection of any element onto the collapse state.

(v) Matter, photons, and tachyons, as sectors of the biquaternion ground state.

(vi) The gauge structure $U(1) \times SU(2) \times SU(3)$, as the symmetry group of the collapse state's representations.

All are unitarily equivalent by Stone–von Neumann [1]. The Sphaera's only object is its own collapse state.

Proof structure. The equivalences are established section by section: (i) Definition 12.5 and Theorem 14.3; (ii)–(iii) same (triole as collapse coordinates); (iv) Proposition 16.1; (v) Theorem 15.2; (vi) Theorem 18.3. Unitary equivalence throughout via Stone–von Neumann ([1], Theorem 3.2). □

Status (v4.9.0 revision): conjecture; the Stone–von Neumann justification is withdrawn (v4.8.0-r04); Reading B gives a precise form via duality (v4.8.0-r07).

Remark 12.2 (What "same object" means)**.** *Two representations are the same object here in the precise sense of Stone–von Neumann: they are unitarily equivalent irreducible representations of the Weyl algebra $\mathcal{W}(A^1)$. The frame determines which representation is visible; the object — the collapse state $z^- = 0$ — is invariant.*

The triole condition $z_1 + z_2 + z_3 = 0$ and the collapse condition $z^- = 0$ are the same equation in two coordinate systems: the triole is the collapse state expressed in the three kernel components simultaneously. The Sphaera is therefore not a space that contains a collapse state — it is *the collapse state, seen from different frames.*

Theorem 12.3 (Origin of the universe as broken collapse state)**.** *The departure from the collapse state $z^- = 0$ is governed by two principles, both of which are consequences of the Sphaera-Cascade structure:*

(i) ***Planck length as minimal*** *$|z^-|$. The cascade $\hat{K}$ cannot project below the Planck scale:*

$$|z^-|_{\min} = \ell_P := \sqrt{\frac{\hbar G}{c^3}} = 1.616 \times 10^{-35}\,\mathrm{m}.$$

Below ℓ_P: $z^- = 0$, the Sphaera, the photon. Above ℓ_P: $|z^-|$ is nonzero, matter exists. The Planck length is the threshold of the first breaking of the collapse state.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(ii) ***Pauli principle as antisymmetry under*** J. *The Hilbert space* H^+ *decomposes under* J *into two components:*

$$C(f) = \tfrac{1}{2}(f + Jf) \in H^+_+ \quad \textit{(bosonic, symmetric)},$$

$$A(f) = \tfrac{1}{2}(f - Jf) \in H^+_- \quad \textit{(fermionic, antisymmetric)}.$$

The antisymmetric component satisfies $A(f) \wedge A(f) = 0$ *(Grassmann property): no two fermions occupy the same* z^- *state. This is the Pauli exclusion principle, derived from the* J*-structure of the Sphaera.*

(iii) ***Discrete spectrum.*** *Together, (i) and (ii) produce a discrete tower of matter states:*

$$|z^-|_n = n \cdot \ell_P, \quad n \in \mathbb{N}_{\geq 1},$$

with each level occupied by at most one fermion. The prime lattice Γ_P *(from Stone–von Neumann, [1]) identifies the primitive levels:* $|z^-|_p = p \cdot \ell_P$ *for* $p \in \mathbb{P}$.

Remark 12.4 (The Sphaera as photon, the universe as broken frame)**.** *Theorem 12.1 says the Sphaera is the collapse state* $z^- = 0$*, which is the photon sector. Theorem 12.3 says the departure from this state is governed by* ℓ_P *and the Pauli principle. Therefore:*

- *The Sphaera is a photon — the unique frame-invariant object.*
- *The universe is the same photon observed from a broken frame where* $|z^-| \geq \ell_P$.
- *The breaking was not random: it is discrete (Pauli), minimal (Planck), and primitive (prime lattice).*
- *The cascade* $\hat{K}$ *projects everything back toward* $z^- = 0$ *— the universe tends toward its photon ground state.*

Both the Sphaera and the universe are the same object. The difference is only the frame.

> **Reader cue.** Read the next block as the concrete carrier unpacking of the global Sphaera identity claim. Its purpose is to pass from the frame-level origin statement to the biquaternion kernel and its induced Lorentz/Minkowski geometry that the later physics sections reuse.

Definition 12.5 (Biquaternion kernel)**.** *The ground state of a stratum* S_n *of the Sphaera consists of three complex spheres* K_1, K_2, K_3 *whose unit coordinates form a biquaternion*

$$q_n := it \cdot \mathbf{1} + z_1 \cdot e_1 + z_2 \cdot e_2 + z_3 \cdot e_3 \in \mathbb{H} \otimes_{\mathbb{R}} \mathbb{C},$$

where $it \in i\mathbb{R}$ *is the imaginary time unit;* $z_1, z_2 \in \mathbb{C}$ *are the complex coordinates of kernels* K_1, K_2*;* $e_3 := e_1 e_2$ *is the quaternion product, hence automatically orthogonal to both* e_1 *and* e_2*; and* $z_3 \in \mathbb{C}$ *is the coordinate of kernel* K_3 *along* e_3.

The biquaternion norm is

$$|q_n|^2 = -(ct)^2 + |z_1|^2 + |z_2|^2 + |z_3|^2.$$

Status (v4.9.0 revision): Minkowski reading replaced: signature $(1, 6)$, natural norm $N(q)$, Reading B (v4.8.0-r06).

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Remark 12.6 (Minkowski metric from quaternion structure)**.** *The norm $|q_n|^2$ is the Minkowski metric with signature $(-,+,+,+)$. This is not postulated: it follows from the quaternion algebra $e_i^2 = -1$ and $it \cdot \mathbf{1}$ carrying imaginary time. The Lorentz group $SO(1,3)$ acts on q_n by quaternion conjugation: $q_n \mapsto u\, q_n\, u^{-1}$, $u \in \mathbb{H}^\times$. This is the spinor representation of the Lorentz group — emerging automatically from the biquaternion structure, not imposed from outside.*

Proposition 12.7 (Orthogonality of the third kernel)**.** *K_3 is always orthogonal to K_1 and K_2:*

$$\langle e_3, e_i \rangle = 0 \qquad \text{for } i = 1, 2.$$

Proof. By definition,

$$e_3 = e_1 e_2.$$

Using the quaternionic inner product

$$\langle x, y \rangle := \mathrm{Re}(\overline{x}\, y),$$

we compute

$$\langle e_3, e_1 \rangle = \langle e_1 e_2, e_1 \rangle = \mathrm{Re}\big(\overline{e_1 e_2}\, e_1\big).$$

Since

$$\overline{e_1 e_2} = \overline{e_2}\,\overline{e_1} = (-e_2)(-e_1) = e_2 e_1 = -e_1 e_2,$$

it follows that

$$\overline{e_1 e_2}\, e_1 = (-e_1 e_2) e_1 = -\, e_1 (e_2 e_1) = -\, e_1 (-e_1 e_2) = e_1^2 e_2 = -\, e_2.$$

Hence

$$\langle e_3, e_1 \rangle = \mathrm{Re}(-e_2) = 0.$$

Similarly,

$$\langle e_3, e_2 \rangle = \langle e_1 e_2, e_2 \rangle = \mathrm{Re}\big(\overline{e_1 e_2}\, e_2\big) = \mathrm{Re}\big((-e_1 e_2) e_2\big) = \mathrm{Re}(-e_1 e_2^2) = \mathrm{Re}(e_1) = 0.$$

Therefore e_3 is orthogonal to both e_1 and e_2. □

13 Lorentz Invariance on Every Stratum

Status. *Legacy physical-readout block inside the active main body.*

Block function. This section isolates Lorentz invariance as a stratumwise consequence of the structural core before later physical applications are read off from it.

Theorem 13.1 (Stratified Lorentz invariance)**.** *On every stratum S_n, the quadratic invariant*

$$|q_n|^2 = \tau_n^2$$

is preserved under every HC-admissible frame transformation induced by the global Heisenberg-compensator matrix. In particular, for the involutive frame transformation

$$J : \; z^- \mapsto -z^-,$$

the proper-time invariant τ_n is unchanged. Moreover, every admissible relabeling of the frame data that preserves the same involution J leaves both τ_n and the compensator

$$C(f) = \tfrac{1}{2}(f + Jf)$$

unchanged.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. By the general MM-tools, a change of admissible realization is not a change of structural object. By the HC-tools, a frame transformation is physically admissible only if it preserves the same compensator structure, i.e. the same involution J.

The transformation $J : z^- \mapsto -z^-$ changes only the frame-sensitive sign, not the underlying compensated structural content. Hence the quadratic invariant associated with the stratum remains the same, so

$$|q_n|^2 = \tau_n^2$$

is preserved.

Likewise, the compensator

$$C = \tfrac{1}{2}(\mathrm{id} + J)$$

depends only on the involution J itself. Therefore any HC-admissible relabeling that preserves the same J-structure leaves both τ_n and C unchanged. □

Status (v4.9.0 revision): shows J-invariance; the Lorentz action comes from Reading B (v4.8.0-r06).

14 Fractal Self-Traversal Along the Light Axis

Remark 14.1 (Structural null axis and physical light-axis readout)**.** *In the structurally prior language of the Companion, the distinguished axis selected by the MM/HC architecture is the null axis. Once the same axis is read at stable physical propagation level, it appears as the light axis. Thus the present section keeps the historically familiar light-axis title, while treating it explicitly as the physical readout of a structurally prior null-axis statement.*

Definition 14.2 (Light axis of a stratum)**.** *The structurally prior null axis of stratum S_n is*

$$L_n := \{q_n \in S_n : |q_n|^2 = 0\}.$$

At the physical propagation readout of the same architecture, this null axis is read as the light axis of S_n.

Theorem 14.3 (Fractal structure of the Sphaera)**.** *The Sphaera-Cascade*

$$\hat{K} : S_n \to S_{n-1}$$

transports the null axis of each stratum to the null axis of the next:

$$\hat{K}(L_n) = L_{n-1}.$$

The global null axis is therefore

$$L = \bigcap_{n \geq 0} L_n = S_0,$$

where S_0 is the terminal compensated photon sector. Hence the self-traversal of the Sphaera is fractal in the precise sense that the same null-axis structure is carried through all strata by the same MM/HC architecture and appears physically as the light axis at stable readout.

Proof. By the general MM-tools, the cascade does not generate a new object at each stratum, but transports one and the same structural content through admissible realization levels.

By the HC-tools, the null sector is precisely the sector whose frame-sensitive imbalance is removed by the compensator while its structural identity is preserved. Thus the cascade does not replace the null axis; it carries the same null-axis structure from one stratum to the next.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Therefore

$$\hat{K}(L_n) = L_{n-1}.$$

Iterating this transport through the whole cascade yields the terminal compensated null sector

$$L = \bigcap_{n\geq 0} L_n = S_0,$$

which is read physically as the photon/light sector. □

Status (v4.9.0 revision): null axis redefined as $s = 0$, directions S_0 (v4.8.0-r06); transport open.

Remark 14.4 (Triolic $\mathbb{Z}/3\mathbb{Z}$ structure on the null/light axis)**.** *On the transported null axis, read physically as the light axis at stable propagation, the three kernels remain organized by the triolic phase condition*

$$z_1 + z_2 + z_3 = 0.$$

Thus the cascade preserves the structural $\mathbb{Z}/3\mathbb{Z}$ pattern while driving the frame-sensitive component to the compensated photon/light sector.

15 Three Objects as One: Unitary Equivalence

Status. *Legacy structural-identity block inside the active main body.*

Block function. This section compresses the unitary-equivalence reading that ties the three-object architecture into one common structural carrier.

Definition 15.1 (Sector classification from biquaternion)**.** *The three sectors correspond to the sign of* $\mathrm{Re}(z^-)$*:*

- ***Matter*** *(K_1):* $\mathrm{Re}(z_1) < 0$*, boost $\beta < 0$, spacelike.*
- ***Tachyon*** *($K_2 = J \cdot K_1$):* $\mathrm{Re}(z_2) > 0$*, boost $\beta > 0$, timelike.*
- ***Photon*** *($K_3 = e_1 e_2$):* $\mathrm{Re}(z_3) = 0$*, boost $\beta = 0$, lightlike.*

Status (v4.9.0 revision): the sign of β here is an orientation code. Theorem 20.6 uses the opposite convention; the two differ by charge conjugation $z \mapsto -z$, which leaves $s = z \cdot z$ unchanged. The sector is fixed by the sign of s (v4.8.0-r07, v4.9.0-r24.1).

Theorem 15.2 (Unitary equivalence via Stone–von Neumann)**.** *The three kernels K_1, K_2, K_3 carry unitarily equivalent irreducible representations of the Weyl algebra $\mathcal{W}(A^1)$. The boost J implements the equivalence between K_1 and K_2, and the compensator C produces K_3 from K_1.*

Proof. By Stone–von Neumann [1], every irreducible representation of $\mathcal{W}(A^1)$ is unitarily equivalent to $L^2(\mathbb{Q}^+\backslash A_1)$. All three kernels are subspaces of H^+ which carries this representation. Since J is unitary and maps $K_1 \leftrightarrow K_2$, they are equivalent. $K_3 = C(K_1)$ lies in H^+_+, the unique J-invariant subspace, which is again irreducible (no proper J-invariant subspace exists by T4+T5 of [1]). □

Status (v4.9.0 revision): open (v4.8.0-r04).

16 Near-Collision, Photon Production, and Entanglement

Status. *Legacy physical-regime block inside the active main body.*

Block function. This section records how the same closure architecture is read in the near-collision and entanglement regime, together with the local role of the photon readout.

Proposition 16.1 (Compensator as near-collision in K_3)**.** *The Heisenberg compensator $C(f) = \frac{1}{2}(f + Jf)$ is the projection onto $K_3 = e_1e_2$: it takes matter f (in K_1) and its tachyon image Jf (in K_2), and produces the photon $C(f) \in K_3$. The quaternion identity $e_3 = e_1e_2$ makes the product $z \times J_B z = 2E \times B$ orthogonal to both inputs; the average $C(f)$ itself is not.*

Corrected in the v4.9.0 revision: orthogonality holds for the product $z \times J_B z = 2E \times B$, not for the average (v4.9.0-r03.4); earlier text: "this is always orthogonal to both inputs".

Theorem 16.2 (Entanglement as shared proper time)**.** *Two photons $\phi_1, \phi_2 \in K_3$ are entangled iff they share the same pre-image: $\phi_1 = \phi_2 = C(f)$ for some $f \in K_1$. The entanglement correlation is the invariant proper time $\tau_n(f) = \tau_n(Jf)$ fixed at the moment of the near-collision. No signal is transmitted at measurement: the invariant τ_n was established at the near-collision and is merely revealed.*

4.10 consolidation: The displayed invariant equality under J is a structural statement. It does not by itself prove the general quantum-mechanical iff criterion stated here: a shared clock or common preparation can also accompany a product state. The physical entanglement identification remains an additional hypothesis; see Section 44.3.

Proof. $\tau_n(Jf)^2 = t_n^2 - |-z^-|^2 = t_n^2 - |z^-|^2 = \tau_n(f)^2$ by Theorem 13.1. The shared value τ_n is a Lorentz invariant at every stratum, hence frame-independent. □

17 Connection to the Riemann Hypothesis

Status. *Legacy cross-domain bridge block inside the active main body.*

Block function. This section explains how the structural core is connected to the Riemann-side reading without turning the surrounding physical line into a separate proof source.

Remark 17.1 (Common origin of RH and entanglement)**.** *The Riemann Hypothesis ($z^- = 0$ for all zeros) and quantum entanglement (shared τ_n) are both consequences of the same biquaternion structure:*

1. *The Minkowski norm $|q_n|^2 = \tau_n^2$ is invariant (Theorem 13.1).*
2. *The cascade forces $z^- \to 0$ along the light axis (Theorem 14.3).*
3. *Stone–von Neumann makes all three kernels equivalent (Theorem 15.2).*

Neither was the goal of the original construction. Both emerged from asking: what is the exact structure of the Sphaera?

© 2026 Marc Brendecke. Preprint — not peer reviewed.

18 Gauge Structure of the Standard Model

Status. *Legacy physical-readout block inside the active main body.*

Block function. This section explains how the structural core is read as a gauge-sector organization rather than as an independently added physical layer.

Proposition 18.1 ($\mathbb{H} \otimes \mathbb{C} \to U(1) \times SU(2)$)**.** *The biquaternion algebra $\mathbb{H} \otimes_{\mathbb{R}} \mathbb{C}$ generates exactly the gauge group $U(1) \times SU(2)$.*

Proof. $SU(2)$ from unit quaternions. The unit quaternions $\{q \in \mathbb{H} : |q| = 1\} \cong S^3$ form a Lie group isomorphic to $SU(2)$. The three generators are $T_k := \sigma_k/2$ (half Pauli matrices), satisfying $[T_j, T_k] = i\varepsilon_{jkl}T_l$.

$U(1)$ from the complex scalar. The imaginary time unit $it \in i\mathbb{R}$ generates the abelian group $e^{i\theta} \in U(1)$ for $\theta \in \mathbb{R}$. This commutes with all $SU(2)$ generators. Hence $U(1) \times SU(2)$, direct product. □

Theorem 18.2 ($\mathbb{O} \to SU(3)$)**.** *The octonions decompose under $SU(3)$-action as $\mathbb{O} = \mathbb{C} \oplus \mathbb{C}^3$, a singlet plus the fundamental (triplet) representation of $SU(3)$. The triplet $\mathbb{C}^3$ carries the three kernels (K_1, K_2, K_3) as the three colors.*

Theorem 18.3 (Standard Model from the Sphaera biquaternion)**.** *The biquaternion ground state $q_n \in \mathbb{H} \otimes \mathbb{C}$, extended to the octonionic structure $\mathbb{C} \otimes \mathbb{H} \otimes \mathbb{O}$, generates the Standard Model gauge group $U(1) \times SU(2) \times SU(3)$ with generators arising from:*

Source	*Group*	*Physical force*
Complex scalar it	$U(1)$	*Electromagnetism*
Unit quaternions $\mathbb{H}$	$SU(2)$	*Weak force*
Octonion triplet $\mathbb{C}^3 \subset \mathbb{O}$	$SU(3)$	*Strong force*

Proof. Propositions 7.1–7.2 and Theorem 18.2 establish each factor. The three factors are independent (direct product structure) because $U(1)$ commutes with $SU(2)$ and both commute with the $SU(3)$ action on $\mathbb{C}^3$: the $SU(3)$ generators λ_k act only on the color indices, not on the quaternion structure. □

Status (v4.9.0 revision): the complex scalar *it* alone generates a global phase, the same on all charge sectors, not electromagnetism. The charge is $Q = \frac{1}{2} - \frac{i}{6}L_{e_7} - \frac{i}{3}R_{e_7}$; it needs the outer and the inner complex unit, and its assignment to states depends on the frame (v4.9.0-r34, r35). $\mathbb{C} \otimes \mathbb{H}$ also contains $SL(2, \mathbb{C})$; its compact part is $U(2)$, and the unbroken $U(1)$ of the Standard Model is hypercharge, which is not derived here.

19 Bell Inequalities and τ_n

Status. *Legacy physical-readout block inside the active main body.*

Block function. This section places the Bell/nonlocality discussion on the same readout-relative footing and makes clear what role the τ_n layer plays there.

Theorem 19.1 (Bell consistency)**.** *The proper time τ_n is not a local hidden variable in Bell's sense. The framework is consistent with quantum mechanical violation of Bell inequalities.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. Bell's theorem assumes a hidden variable λ assigned independently to each particle: measurement outcomes $A(\lambda_A, a)$ and $B(\lambda_B, b)$ with λ_A, λ_B statistically independent.

In our framework τ_n belongs to the pair (f, Jf), not to each element separately: $\lambda_A = \lambda_B = \tau_n$. This is a contextual variable — Bell's theorem does not exclude violation by contextual theories.

The correlation function $E(a, b)$ follows from the structure of H_+^+: since $C(f) \in H_+^+$ carries an irreducible representation of $\mathcal{W}(A^1)$ (Stone–von Neumann, [1]), the standard quantum correlations $E(a, b) = \cos\theta_{ab}$ are reproduced, giving CHSH value $S = 2\sqrt{2} > 2$.

Hence:

- τ_n explains why the correlation exists: shared proper time, fixed at the near-collision;
- Stone–von Neumann determines how strong the violation is ($S = 2\sqrt{2}$).

Lorentz invariance and quantum mechanics are consistent. □

Remark 19.2 (What τ_n does and does not do)**.** *τ_n encodes the magnitude $|z^-|$ of the displacement from the photon sector at the moment of the near-collision. It does not determine the measurement direction. When f is measured, z^- is projected onto a value; the paired Jf is projected onto $-z^-$ (the J-image). The anticorrelation was built in at the collision event, not transmitted at measurement. No signal travels faster than light.*

20 String Theory and Loop Quantum Gravity as HC Sectors

Status. *Legacy comparative-physics block inside the active main body.*

Block function. This section places two familiar high-energy frameworks into the compensator-based sector reading without promoting them to separate primitives.

Theorem 20.1 (String–LQG duality via the Heisenberg compensator)**.** *The Hardy space H^+ decomposes under the Heisenberg compensator into two sectors:*

$$H^+ = H_+^+ \oplus H_-^+,$$

$$C(f) = \tfrac{1}{2}(f + Jf) \in H_+^+ \quad \textit{(symmetric, bosonic — String theory)},$$
$$A(f) = \tfrac{1}{2}(f - Jf) \in H_-^+ \quad \textit{(antisymmetric, fermionic — LQG)}.$$

Both sectors are unitarily equivalent by Stone–von Neumann [1]: they are two representations of the same Weyl algebra $\mathcal{W}(A^1)$, seen from different frames. String theory and LQG are therefore not competing theories of quantum gravity: they are the bosonic and fermionic projections of the same object under the Heisenberg compensator.

Proposition 20.2 (Shared vacuum and Planck floor)**.** *Both sectors share the same vacuum and the same minimal scale:*

- ***Vacuum:*** *$z^- = 0$ (the photon/Sphaera) is the ground state of both H_+^+ and H_-^+.*
- ***Planck floor:*** *$\ell_P = |z^-|_{\min}$ is the string length in H_+^+ and the area quantum $\sqrt{A_{\min}} \sim \ell_P$ in H_-^+. Both emerge from the same threshold of Theorem 1.2(i).*

String theory and LQG are therefore not competing theories of quantum gravity: they are the bosonic and fermionic projections of the same object under the Heisenberg compensator.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Theorem 20.3 (Wick rotation as splitting mechanism)**.** *Under the imaginary proper time transformation $t \mapsto i\tau_E$ (Wick rotation), the biquaternion time unit transforms as:*

$$it \;\mapsto\; i \cdot (i\tau_E) = -\tau_E \in \mathbb{R}_{<0}.$$

This is precisely the third triolic sector $\gamma_3 < 0$: the Wick rotation identifies the imaginary time component with the unstable tachyon sector. Consequently:

(i) The triolic balance condition $\gamma_{\text{string}} + \gamma_{\text{LQG}} + \gamma_3 = 0$ is the stability condition under imaginary proper time: without it, the system diverges to $\pm\infty$.

(ii) The two positive sectors ($\gamma_{\text{string}}, \gamma_{\text{LQG}} > 0$) are stable under Wick rotation; the negative sector ($\gamma_3 < 0$) is unstable and is projected away by the cascade $\hat{K}$.

(iii) The Minkowski norm $|q_n|^2 = \tau_n^2$ becomes Euclidean $|q_E|^2 = \tau_E^2 \geq 0$ under $t \mapsto i\tau_E$: the system is unconditionally stable in Euclidean signature.

Proof. (i). Under $t \mapsto i\tau_E$: $it \mapsto -\tau_E$. The biquaternion $q_n = it \cdot \mathbf{1} + z_1 e_1 + z_2 e_2 + z_3 e_3$ becomes $q_E = -\tau_E \cdot \mathbf{1} + z_1 e_1 + z_2 e_2 + z_3 e_3$ with norm $|q_E|^2 = \tau_E^2 + |z_1|^2 + |z_2|^2 + |z_3|^2 \geq 0$. Stability requires $\tau_E^2 \geq 0$, which is always satisfied. The balance $\gamma_1 + \gamma_2 + \gamma_3 = 0$ is equivalent to requiring that the $-\tau_E$ component does not introduce a net positive energy divergence.

(ii). In Euclidean signature, only $\gamma > 0$ sectors contribute to a convergent partition function $Z = \text{Tr}(e^{-\beta H})$ with $H \geq 0$. The sector $\gamma_3 < 0$ would give $e^{+\beta|\gamma_3|H} \to \infty$: it is projected away by $\hat{K}$.

(iii). Direct computation of the Euclidean norm. □

Theorem 20.4 (Triolic structure of γ_{BI})**.** *The Barbero–Immirzi parameter γ_{BI} and the string-sector final $\gamma_{\text{string}} = \ln 2/(\pi\sqrt{3})$ are not in contradiction. They are two projections of the same triolic invariant:*

$$\gamma_{\text{string}} + \gamma_{\text{LQG}} + \gamma_3 = 0,$$

with $\gamma_3 = -(\gamma_{\text{string}} + \gamma_{\text{LQG}}) \approx -0.3649$ being the third (unoccupied, tachyon) sector. The frame-invariant quantity is the triolic norm:

$$\tau_\gamma := \frac{1}{\sqrt{3}}\sqrt{\gamma_{\text{string}}^2 + \gamma_{\text{LQG}}^2 + \gamma_3^2} \approx 0.2619.$$

Theorem 20.5 (Observer frame offset)**.** *Every matter observer carries a nonzero contravariant component $z^- \neq 0$. All measured physical constants — including $\hbar$, G, c, and γ_{BI} — are projections of their frame-invariant values onto the observer's sector:*

$$\gamma_{\text{observed}} = \gamma_0 \cdot \frac{1}{\sqrt{1-\beta^2}} = \gamma_0 \cdot \gamma_{\text{Lorentz}}(\beta).$$

The Barbero–Immirzi parameter $\gamma_{\text{BI}} \approx 0.2375$ is not an absolute constant of nature: it is the Lorentz-transformed γ_{string}, observed from a matter frame with $\beta \approx 0.844$.

Proof. $\gamma_{\text{string}} = \ln 2/(\pi\sqrt{3})$ is computed in the photon frame $z^- = 0$ from the HC trace formula. The ratio $\gamma_{\text{BI}}/\gamma_{\text{string}} = 1.864$ equals a Lorentz factor $\gamma_L = 1/\sqrt{1-\beta^2}$ for $\beta = \sqrt{1 - 1/1.864^2} \approx 0.844$. The corresponding observer position is $z^- = \beta/2 \approx 0.422$. □

Status (v4.9.0 revision): structural statement about γ_L (author, v4.9.0-r16); conversions as in v4.9.0-r18, r19.

Theorem 20.6 (Symmetric matter–tachyon frame)**.** *Matter and tachyon are positioned symmetrically around the photon frame $z^- = 0$:*

$$\beta_{\text{matter}} = +\beta_{\text{sym}}, \qquad \beta_{\text{tachyon}} = -\beta_{\text{sym}},$$

where

$$\beta_{\text{sym}} = \sqrt{\frac{\gamma_{\text{BI}} - \gamma_{\text{string}}}{\gamma_{\text{BI}} + \gamma_{\text{string}}}} \approx 0.5493.$$

The Lorentz factor between the matter and tachyon frames is:

$$\gamma_L = \frac{1 + \beta_{\text{sym}}^2}{1 - \beta_{\text{sym}}^2} = \frac{\gamma_{\text{BI}}}{\gamma_{\text{string}}} \approx 1.864.$$

Proof. Let $r := \gamma_{\text{BI}}/\gamma_{\text{string}}$. Set $\beta_{\text{sym}} := \sqrt{(r-1)/(r+1)}$. Then:

$$\frac{1 + \beta_{\text{sym}}^2}{1 - \beta_{\text{sym}}^2} = \frac{1 + (r-1)/(r+1)}{1 - (r-1)/(r+1)} = \frac{(r+1) + (r-1)}{(r+1) - (r-1)} = \frac{2r}{2} = r. \quad \checkmark$$

The relativistic velocity of the tachyon as seen from the matter frame:

$$\beta_{\text{tachyon|matter}} = \frac{-\beta_{\text{sym}} - \beta_{\text{sym}}}{1 - (-\beta_{\text{sym}})\beta_{\text{sym}}} = \frac{-2\beta_{\text{sym}}}{1 + \beta_{\text{sym}}^2},$$

with corresponding Lorentz factor $\gamma_{\text{tachyon|matter}} = r$. $\checkmark$ □

Status (v4.9.0 revision): the tachyon at $-\beta_{\text{sym}}$ is the real projection $T|_R$ of the tachyon rapidity $-\eta + i\pi/2$; in its own frame $\beta_T^2 = 1/\beta_{\text{sym}}^2$ (v4.9.0-r24.2). γ_L is the ratio of Bloch lengths $|b_R|/|b_M|$ (v4.9.0-r24.3). The sign of β is an orientation code; the sector is fixed by the sign of s (v4.8.0-r07, v4.9.0-r24.1).

21 Black Holes and Hawking Radiation

Status. *Legacy physical-regime block inside the active main body.*

Block function. This section reads collapse, horizon behavior, and Hawking-type radiation as frame- and readout-level consequences of the same closure architecture.

Definition 21.1 (Schwarzschild horizon in Sphaera coordinates)**.** *The Schwarzschild horizon $r_S = 2GM/c^2$ corresponds to $\beta \to 1$ in the Sphaera metric $g_{tt} = 1 - \beta^2$. In sector coordinates: the horizon is where $z^- \to \infty$.*

Proposition 21.2 (Horizon as cascade singularity)**.** *The cascade operator $\hat{K}$ is undefined at $z^- = \infty$: the Sphaera cannot project beyond its own horizon. The horizon is the boundary of the Sphaera's jurisdiction.*

Proof. $\hat{K} = C^\mu{}_\nu \circ \Pi_J$ requires $\|z^-\| < \infty$ (finite sector displacement). At $z^- \to \infty$: $\beta \to 1$, $\gamma_{\text{Lorentz}} \to \infty$, the Lorentz transformation is singular, and the projection Π_J ceases to be bounded. □

Status (v4.9.0 revision): under the counting reading of v4.9.0-r63, γ_{string} is the minimal-spin counting of the tachyon floor, i.e. of the infalling partner; the theorem assigns it to the escaping photon in the reference frame R and reads T as R (v4.9.0-r64).

Theorem 21.3 (Hawking radiation from the triolic structure)**.** *At the horizon, a virtual matter–tachyon pair from the triolic structure (Theorem 3.2) is promoted to a real pair:*

- *Tachyon ($\beta = -\beta_{\text{sym}}$): falls into the black hole.*
- *Matter ($\beta = +\beta_{\text{sym}}$): escapes to infinity.*
- *Their compensator output $C(f) = \frac{1}{2}(f + Jf)$: a photon with $\beta = 0$, $z^- = 0$ — the Hawking radiation.*

The temperature of the Hawking photon in the photon frame is:

$$T_{\text{Sphaera}} = \gamma_{\text{string}} \cdot \frac{m_P^2 c^2}{8\pi M k_B} = \gamma_{\text{string}} \cdot T_{\text{Hawking}}.$$

Proof. The Hawking temperature (standard derivation) is

$$T_H = \frac{\hbar c^3}{8\pi G M k_B} = \frac{m_P^2 c^2}{8\pi M k_B}.$$

The Sphaera produces the Hawking photon in the photon frame where γ is measured as γ_{string} (not γ_{BI}, which is the matter-frame value). Hence $T_{\text{Sphaera}} = \gamma_{\text{string}} \cdot T_H$. The ratio $T_{\text{Sphaera}}/T_{\text{Hawking}} = \gamma_{\text{string}} = \ln 2/(\pi\sqrt{3}) \approx 0.1274$ is constant across all black hole masses. □

Proposition 21.4 (Hawking temperature frame analysis and physical meaning of γ_{BI})**.** *The Hawking temperature $T_{\text{Sphaera}} = \gamma_{\text{string}} \cdot T_H^{(\text{matter})}$ has the following frame interpretations:*

(i) ***Photon frame:*** *$k(T) = k(\hbar) + k(\kappa) - k(k_B)$ where surface gravity $\kappa = c^4/(4GM)$ has $k(\kappa) = 3k(c) - k(G_N) = -2$ and $k(\hbar) = 1$, giving $k(T_H) = -1$ (assuming $k(k_B) = 0$). Hence $T_H^{(\text{photon})} = \gamma_L \cdot T_H^{(\text{matter})}$, and the Sphaera prediction is*

$$T_{\text{Sphaera}} = \frac{\gamma_{\text{string}}}{\gamma_L} \cdot T_H^{(\text{photon})} = \frac{\gamma_{\text{string}}^2}{\gamma_{\text{BI}}} \cdot T_H^{(\text{photon})}.$$

(ii) ***Tachyon frame:*** *The tachyon sector has opposite frame shift, $T_H^{(\text{tachyon})} = \gamma_L^{-1} \cdot T_H^{(\text{matter})}$. Then:*

$$T_{\text{Sphaera}} = \gamma_{\text{string}} \cdot T_H^{(\text{matter})} = \gamma_{\text{string}} \cdot \gamma_L \cdot T_H^{(\text{tachyon})} = \gamma_{\text{BI}} \cdot T_H^{(\text{tachyon})}.$$

(iii) ***Physical meaning of γ_{BI}:*** *The Barbero–Immirzi parameter is the ratio of the Sphaera Hawking temperature to the tachyon-frame Hawking temperature:*

$$\gamma_{\text{BI}} = \frac{T_{\text{Sphaera}}}{T_H^{(\text{tachyon})}}.$$

This gives γ_{BI} a direct kinematic interpretation: it measures how much the tachyon sector amplifies the Hawking temperature relative to the Sphaera output. Since the tachyon falls into the black hole and the Sphaera photon escapes, γ_{BI} encodes the frame mismatch between the infalling and escaping components of the triolic virtual pair.

Proof. Statement (i): with $k(\hbar) = 1$, $k(G_N) = 2$, $k(c) = 0$, $k(k_B) = 0$, the frame correction to $T_H = \hbar c^3/(8\pi G_N M k_B)$ is $\gamma_L^{k(\hbar)-k(G_N)} = \gamma_L^{-1}$, giving $T_H^{(\text{photon})} = \gamma_L \cdot T_H^{(\text{matter})}$.

Statement (ii): the tachyon sector sits at $z^- = -\beta_{\text{sym}}$, the mirror image of the matter frame. The frame factor for the tachyon is $\gamma_L^{(\text{tachyon})} = 1/\gamma_L^{(\text{matter})}$ by the symmetric matter–tachyon frame theorem 20.6. Hence $T_H^{(\text{tachyon})} = \gamma_L^{-1} \cdot T_H^{(\text{matter})}$, and $T_{\text{Sphaera}}/T_H^{(\text{tachyon})} = \gamma_{\text{string}} \cdot \gamma_L = \gamma_{\text{BI}}$.

Statement (iii) follows immediately from (ii). □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Status (v4.9.0 revision): item (ii): frame order does not match the axis (v4.9.0-r01); item (iii) holds to −0.55% under the class split (v4.9.0-r19); assumption $k(M) = 0$ explicit (v4.8.0-r03 F6); in the tachyon frame T_H changes sign (v4.9.0-r23).

> **Reader cue.** Read the next block as a geometric recoding of the same Hawking data. Its purpose is to shift from the horizon/frame-temperature dictionary to the coupling interpretation before the section turns to the core/horizon dual picture.

Theorem 21.5 (Geometric origin of γ_L from Hawking geometry)**.** (Corrected archival stage: previous version used incorrect $\gamma_L \approx 1.197$ and an erroneous 2/3 sector factor.) *The splitting factor* $\gamma_L = \gamma_{\mathrm{BI}}/\gamma_{\mathrm{string}} \approx 1.864$ *satisfies:*

$$\gamma_L \;\approx\; \frac{1}{\beta_{\mathrm{sym}}} \;=\; \frac{1}{\sin(\theta_H/2)},$$

where $\theta_H \approx 66.6°$ *is the Hopf angle of the matter observer on* S^3*. Numerically:* $1/\beta_{\mathrm{sym}} \approx 1.820$ *vs. measured* $\gamma_L \approx 1.864$*, a discrepancy of* 2.4%*.*

Proof. The matter observer at β_{sym} has radial position $r/r_S = 1/(1-\beta_{\mathrm{sym}}^2)$ relative to the Schwarzschild horizon. The local Hawking temperature is $T(r) = T_H(1-r_S/r)^{-1/2} = T_H/\beta_{\mathrm{sym}}$. Since γ_{BI} is determined by requiring $S = A/(4\ell_P^2)$ for the observed temperature, the effective coupling shifts: $\gamma_{\mathrm{BI}} = \gamma_{\mathrm{string}} \cdot (T_H/T(r))^{-1} = \gamma_{\mathrm{string}}/\beta_{\mathrm{sym}}$. Hence $\gamma_L = \gamma_{\mathrm{BI}}/\gamma_{\mathrm{string}} = 1/\beta_{\mathrm{sym}} \approx 1.820$.

The residual 2.4% gap between $1/\beta_{\mathrm{sym}} = 1.820$ and the measured $\gamma_L = 1.864$ is the same continuum-limit correction as in the β_{sym}-Hopf gap (Remark 22.3): the discrete prime lattice Γ_P shifts β_{sym} from the Hopf final $1/\sqrt{3}$ by the factor δ_{CL}, and the same correction applies here. □

Remark 21.6 (Connection to GOE/GUE spectral ratio)**.** *By Proposition 24.3 and Remark 24.4,* $\gamma_L = \gamma_{\mathrm{BI}}/\gamma_{\mathrm{string}}$ *is the ratio of the GOE scale* (H_-^+*, LQG) to the GUE scale* (H^{++}*, String). The geometric relation* $\gamma_L \approx 1/\beta_{\mathrm{sym}} = 1/\sin(\theta_H/2)$ *therefore says:*

$$\frac{\textit{GOE scale}}{\textit{GUE scale}} \;\approx\; \frac{1}{\sin(\textit{matter Hopf angle}/2)}.$$

The matter observer's position on S^3 *(the Hopf angle* θ_H*) encodes the ratio of the two random-matrix universality classes.*

> **Reader cue.** Read the next block as an interpretive deepening of the same closure architecture. Its purpose is to shift from the frame- and coupling-readout picture to the core/horizon communication picture without introducing a separate physical mechanism.

Theorem 21.7 (Hawking radiation as the voice of the core)**.** *The Hawking tachyon is not created at the horizon. It is the black hole core — which has* $z^- \to \infty$ *and cannot be projected by* $\hat{K}$ *— attempting to reach* $z^- = 0$ *through the only available channel: tunnelling through the horizon.*

The two descriptions are CPT-dual and physically identical:

Outside view (t forward)	*Inside view (t reversed)*
t_1*: vacuum*	t_3*: core at* $z^- = \infty$
t_2*: virtual pair at horizon*	t_2*: tachyon at horizon*
t_3*: tachyon falls in, matter escapes*	t_1*: Hawking photon emitted*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

The cascade $\hat{K}$ has no preferred time direction: it is the equilibrium between $z^- = 0$ (outside) and $z^- = \infty$ (core). The Hawking pair is the discharge of this tension.

Structural argument. Outside: z^- grows from 0 (far away) to ∞ (horizon). $\hat{K}$ attempts to project $z^- \to 0$ but fails at the horizon. Tension forces a real pair. Inside: $z^- = \infty$ (core, singular compression). $\hat{K}$ attempts $z^- \to 0$ but the horizon blocks outward projection. The only escape: a core fraction tunnels as a tachyon ($\beta = -\beta_{\text{sym}}$) through the horizon. Under J-reflection, this tachyon appears outside as matter ($\beta = +\beta_{\text{sym}}$). The compensator $C(f) = \frac{1}{2}(f + Jf)$ produces the Hawking photon at $\beta = 0$.

CPT: C exchanges matter and tachyon; P exchanges $\beta \leftrightarrow -\beta$ (sides of the horizon); T reverses creation and annihilation. Both descriptions are related by CPT. □

Remark 21.8 (Black hole as inverted Sphaera)**.** *The normal Sphaera has z^- decreasing inward: $\hat{K}$ projects from large z^- toward $z^- = 0$ (the photon ground state). A black hole inverts this: z^- increases inward, from $z^- = 0$ (far away) to $z^- = \infty$ (horizon and core). The cascade $\hat{K}$ cannot function inside the horizon; the interior is a region where the Sphaera does not exist. The horizon is where the Sphaera is maximally stressed.*

Hawking radiation is the Sphaera discharging at its boundary. The black hole evaporates because the core is the Sphaera — and the Sphaera always tends toward $z^- = 0$. The tachyon is not a particle falling in. It is the core, speaking through the only open channel.

Remark 21.9 (The entropy factor)**.** *The Bekenstein–Hawking entropy $S_{BH} = A/(4\ell_P^2)$ and the LQG entropy $S_{LQG} = A \ln 2/(8\pi\gamma_{\text{BI}}\ell_P^2)$ agree when $\gamma_{\text{BI}} = \ln 2/(2\pi) \approx 0.110$. The measured value $\gamma_{\text{BI}} \approx 0.2375$ differs by a factor ≈ 2.15. Within the present framework: the LQG formula is derived in the matter frame, while the Bekenstein–Hawking formula is a photon-frame statement. The discrepancy factor 2.15 should be related to the frame transformation, but its precise form — unlike the analogous Hawking temperature factor γ_{string} — is not yet derived. This is an open quantitative problem.*

Theorem 21.10 (ART–QFT consistency of the Sphaera)**.** *The Sphaera biquaternion encodes both General Relativity and Quantum Field Theory, connected by Wick rotation:*

$$it \cdot \mathbf{1} + z_i e_i \xrightarrow{t \mapsto i\tau_E} \tau_E \cdot \mathbf{1} + z_i e_i.$$

At the natural observer position $r_0/r_S = 1/(1-\beta_{\text{sym}}^2)$, the Schwarzschild metric gives $g_{tt}(r_0) = -\beta_{\text{sym}}^2$ (invariant): no additional geometric factor is hidden in the ART side. The ART–QFT conversion is internally consistent.

22 The Sphaera as a Hopf Fibration

Status. *Legacy geometry-topology block inside the active main body.*

Block function. This section gives the geometric-topological reading of the Sphaera that later fixed-point, transport, and symmetry interpretations can refer back to.

Theorem 22.1 (S-chain as nested Hopf fibrations)**.** *The stratum chain $S_0 \subset S_1 \subset S_2$ of the Sphaera is identified with the base spaces of the three classical Hopf fibrations:*

$$\begin{aligned} S_0 &\cong S^2 = \mathbb{C}P^1 \quad \textit{(Weyl spinor, massless photon)}, \\ S_1 &\cong S^4 = \mathbb{H}P^1 \quad \textit{(Dirac spinor, massive matter)}, \\ S_2 &\cong S^8 = \mathbb{O}P^1 \quad \textit{(octonionic spinor, 3 generations)}, \end{aligned}$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

with the natural inclusion $S^2 \hookrightarrow S^4 \hookrightarrow S^8$. The cascade operator $\hat{K}$ is the Hopf projection chain:

$$\hat{K} : S^{15} \longrightarrow S^8 \longrightarrow S^4 \longrightarrow S^2,$$

where the gauge groups arise as Hopf fibres:

$$\underbrace{S^1}_{\cong U(1)} \to S^3 \to S^2, \quad \underbrace{S^3}_{\cong SU(2)} \to S^7 \to S^4, \quad S^7 \longrightarrow S^{15} \longrightarrow S^8.$$

4.10 correction: the Hopf fibre S^7 is not a group containing G_2; G_2 is the automorphism group of the octonions, and $SU(3)$ is the stabilizer of a chosen imaginary unit. The earlier displayed subgroup annotation on the fibre was incorrect.

Status (v4.9.0 revision): Hopf fibrations classical; physical identification a conjecture; the chain $S^{15} \to S^8 \to S^4 \to S^2$ is not a composition of Hopf maps (v4.8.0-r09).

Theorem 22.2 (β_{sym} from the Hopf angle)**.** *The matter observer's frame offset β_{sym} is the sine of half the Hopf angle θ_H of the matter spinor on S^3 relative to the photon pole:*

$$\beta_{\text{sym}} = \sin\left(\frac{\theta_H}{2}\right), \quad \theta_H \approx 66.6^\circ.$$

The continuum-limit Hopf final is $\beta_{\text{sym}}^{(\text{Hopf})} = 1/\sqrt{3} \approx 0.5774$, corresponding to the symmetric $\mathbb{Z}/3\mathbb{Z}$-orbit on S^3. The measured value $\beta_{\text{sym}} \approx 0.5493$ differs by $\approx 4.9\%$ in β_{sym} and by $\approx 6.8\%$ in γ_L.

Proof. By Theorem 22.1, the stratum chain $S_0 \subset S_1 \subset S_2$ of the Sphaera is identified as nested Hopf fibrations. The matter kernel K_1 sits at a definite position on the S^3 fibre relative to the photon pole $K_3 \in S_0$. The angular distance from K_1 to K_3 on S^3 defines the Hopf angle θ_H, and the relativistic frame offset is $\beta_{\text{sym}} = \sin(\theta_H/2)$ by the standard half-angle relation for spinors on S^3.

From the measured values $\gamma_{\text{BI}} \approx 0.2375$ and $\gamma_{\text{string}} \approx 0.1274$, Theorem 20.6 gives $\beta_{\text{sym}} = \sqrt{(\gamma_{\text{BI}} - \gamma_{\text{string}})/(\gamma_{\text{BI}} + \gamma_{\text{string}})} \approx 0.5493$, hence $\theta_H = 2\arcsin(0.5493) \approx 66.64^\circ$. □

Status (v4.9.0 revision): $\beta_{\text{sym}} = \sin(\theta_H/2)$ is the half-angle relation of a pure state measured from the photon pole, i.e. in the reference frame R (v4.9.0-r24). The matter frame of Convention F at the same β_{sym} is a mixed state: at $\beta_{\text{sym}}^2 = \frac{1}{3}$ the pure Hopf state has populations $(\frac{2}{3}, \frac{1}{3})$, the matter state $(\frac{3}{4}, \frac{1}{4})$ with $|b| = \frac{1}{2}$ (v4.9.0-r17). The theorem therefore reads an R-frame state as the matter offset (v4.9.0-r61).

Remark 22.3 (The Hopf gap and OP 1)**.** *The continuum-limit value $\beta_{\text{sym}}^{(\text{Hopf})} = 1/\sqrt{3}$ arises from the maximally symmetric $\mathbb{Z}/3\mathbb{Z}$ orbit on a round S^3: each of the three triolic kernels sits at equal angular distance $\theta_H = 2\arcsin(1/\sqrt{3}) \approx 70.53^\circ$ from the poles, giving $\gamma_L^{(\text{Hopf})} = 2$. The measured value $\beta_{\text{sym}} \approx 0.5493$ corresponds to the discrete prime-lattice geometry of the Sphaera cascade, where the stratum structure deforms the round S^3 and shifts the matter kernel to a non-maximally-symmetric position. The gap is therefore not a small correction: a 4.9% shift in β_{sym} produces a 6.8% shift in γ_L due to the nonlinear relation $\gamma_L = (1 + \beta_{\text{sym}}^2)/(1 - \beta_{\text{sym}}^2)$, and γ_L appears as γ_L^k in all frame-corrected constants. OP 1 (open): derive β_{sym} from the Sphaera cascade dynamics directly, without reference to $\gamma_{\text{BI}}/\gamma_{\text{string}}$. The correct target is $\beta_{\text{sym}} = \sqrt{(\gamma_{\text{BI}} - \gamma_{\text{string}})/(\gamma_{\text{BI}} + \gamma_{\text{string}})}$ derived as a fixed point of the cascade, not postulated from the ratio. This requires the Dirac operator on H_-^+ (OP 2) as a prerequisite.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Status (v4.9.0 revision): OP 1 stays open, but its target is the matter-frame value itself. The Hopf value compares an R-frame pure state with the mixed matter state, and the gap to it is not a like-for-like difference (v4.9.0-r61); the prime-lattice reading of the gap is not needed for that. Under the flavour reading of v4.9.0-r55 to r60 the lepton data give $\beta_{\text{sym}} \in [0.545, 0.570]$, containing the measured value and excluding the Hopf and Tribonacci values (v4.9.0-r61, not a blind test).

Theorem 22.4 (Three fermion generations from $\mathbb{O}$)**.** *In the Dixon algebra $D = \mathbb{C} \otimes \mathbb{H} \otimes \mathbb{O}$, the three fermion generations are identified as:*

$$\text{Gen}_i \cong (\mathbb{C} \otimes \mathbb{H}) \otimes e_{i+3}, \quad i = 1, 2, 3,$$

where $e_4, e_5, e_6 \in \text{Im}(\mathbb{O})$ form the $\mathbb{C}^3$ triplet under $SU(3) \subset G_2 = \text{Aut}(\mathbb{O})$.

> ***Reader cue.*** *Read the next table as a representation dictionary. Its purpose is to tell the reader how the proposed octonionic slots are being read generation-by-generation, so the subsequent proof can refer back to one compact assignment surface.*

Generation	$\mathbb{O}$*-unit*	*Quarks*	*Leptons*	*Neutrino*
Gen 1	e_4	u, d	e	ν_e
Gen 2	e_5	c, s	μ	ν_μ
Gen 3	e_6	t, b	τ	ν_τ
Singlet	e_7	—	—	ν_R

Proof. Under $SU(3) \subset G_2$, the imaginary octonions decompose as $\text{Im}(\mathbb{O}) = \mathbb{C} \oplus \mathbb{C}^3$: singlet e_7 and triplet (e_4, e_5, e_6). The anti-triplet (e_1, e_2, e_3) carries the conjugate color representation. Each generation $(\mathbb{C} \otimes \mathbb{H}) \otimes e_{i+3}$ carries one copy of the spinor representation of the Lorentz group (from $\mathbb{C} \otimes \mathbb{H} = M(2, \mathbb{C})$) tensored with one color index. The Fano plane structure of $\mathbb{O}$ gives the cyclic $e_4 \to e_5 \to e_6$ symmetry. □

Status (v4.9.0 revision): the triplet basis e_4, e_5, e_6 and $\nu_R \leftrightarrow e_7$ hold in the inner frame $J = L_{e_7}$; with the outer unit they translate by P_- to the charges $Q = \frac{2}{3}$ and $Q = 0$ (v4.9.0-r34, r35). The sentence on the anti-triplet (e_1, e_2, e_3) is wrong: e_1, e_2, e_3 are the J-images of the same triplet (v4.9.0-r35). The cyclic symmetry $e_4 \to e_5 \to e_6$ is a colour rotation in $SU(3) = \text{Stab}_{G_2}(e_7)$, so the table places colour and generation on the same object (v4.8.0-r13, v4.9.0-r36); generations need a second three-fold structure outside $\mathbb{O}$ (v4.8.0-r14, v4.9.0-r33).

Theorem 22.5 (Koide relation; phase formula archived)**.** *The lepton masses satisfy the Koide relation $K = 2/3$ to within 0.005%. This is an empirical fact; its derivation from the triolic structure is open. The phase formula $\theta_K = \pi - \sqrt{2} \arcsin \beta_{\text{sym}}$ of the earlier version is excluded by data at about 300–400σ (v4.8.0-r09).* Archived in the v4.9.0 revision; original text in §A.270.

Remark 22.6 (Mass discrepancy and amplification near the electron)**.** *Although the Koide phase angle is reproduced to 0.111%, the individual mass predictions show strongly non-uniform errors: m_τ deviates by 0.07%, m_μ by 1.2%, and m_e by 12.8%. This amplification by a factor of ~ 115 near the electron mass is structural: the Brannen map $m_k \propto (1 + \sqrt{2}\cos(\theta_K + 2\pi k/3))^2$ is extremely sensitive to small angular shifts near the zero of the cosine argument for the lightest generation. A 0.111% error in θ_K translates to a $\sim 13\%$ error in m_e due to this nonlinear amplification. The residual discrepancy is attributed to the approximation $\beta_{\text{sym}} \approx 0.5493$ (the precise value of β_{sym} is fixed by $\gamma_{\text{BI}}/\gamma_{\text{string}}$ to the same accuracy as those coupling constants); closing this gap is part of OP 1 (derivation of β_{sym} from first principles, Theorem 22.2).*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Remark 22.7 (The 2/3 factor is everywhere)**.** *The factor* 2/3 *appears as: the stable sector fraction (Thm. I.9.3), the spring oscillator coefficient* $\omega = \gamma_{\text{string}}\sqrt{2/3}$*, the cosmological dark energy* $\Omega_\Lambda \approx 2/3$ *(Theorem 11.9), and now the Koide lepton mass ratio. All are the same statement: 2 of the 3 triolic sectors collapse to the photon ground state.*

Remark 22.8 (Mass spectrum beyond Koide)**.** *The seven imaginary octonion units* $e_1, \ldots, e_7$ *provide seven natural slots for fermion masses. A direct identification* $m_p \sim p \cdot m_P$ *(Planck units) does not match the observed quark mass ratios: consecutive prime ratios (*3/2, 5/3, 7/5, *...) differ from the observed mass hierarchies by factors of* 10–100*. The mass spectrum requires a Yukawa coupling mechanism (Higgs vacuum expectation value) that is not yet derived from the Sphaera structure.*

Theorem 22.9 (Cosmic expansion as growth of the active zero basis)**.** *The scale factor* $a(t)$ *of the universe is identified with the proper time:*

$$a(t) \sim \tau_n(t) = \sqrt{t_n(t)^2 - \beta_{\text{sym}}^2},$$

where $t_n(t)$ *is the imaginary part of the* n*-th active Riemann zero at cosmic time* t*, and* β_{sym} *is the matter observer's frame offset (Theorem 9.6). The Hubble parameter is the activation rate of zeros:*

$$H = \frac{\dot{a}}{a} = \frac{\dot{t}_n}{t_n} = \frac{\dot{n}}{n}$$

(since $t_n \approx 2\pi n/\log n$ *by Weyl asymptotics, and for large* n*:* $\dot{t}_n/t_n \approx \dot{n}/n$*). The universe expands because more Riemann zeros become "active" as cosmic time increases. The active zero count at the Hubble time is* $n_H \sim 10^{62}$*.*

Theorem 22.10 (Dark energy as permanent frame offset)**.** *The matter observer's permanent frame offset* β_{sym} *enters the proper time as a constant term:*

$$\tau_n^2 = t_n^2 - \beta_{\text{sym}}^2 = \underbrace{t_n^2}_{grows} - \underbrace{\beta_{\text{sym}}^2}_{constant}.$$

The constant term β_{sym}^2 *plays the role of the effective cosmological constant:* $\Lambda_{\text{eff}} \sim \beta_{\text{sym}}^2 \cdot \rho_P$ *(where* ρ_P *is the Planck density). The ratio of dark energy to matter density today is predicted as:*

$$\frac{\Omega_\Lambda}{\Omega_M} \approx \frac{\beta_{\text{sym}}^2}{\gamma_{\text{string}}} = \frac{0.3018}{0.1274} \approx 2.37.$$

The observed value is $\Omega_\Lambda/\Omega_M \approx 2.18$ *(9% discrepancy, consistent with the residual 1.4% and redshift corrections not yet included).*

Dark energy is not a new substance. It is the permanent phase offset of our matter frame from the photon ground state $z^- = 0$*: we can never fully reach* $z^- = 0$*, so* β_{sym}^2 *remains as a residual vacuum energy.*

Theorem 22.11 (Spring-pull mechanism in the triolic system)**.** *The matter–photon–tachyon triple forms a mechanical system with spring forces in the* z^- *coordinate:*

$$F_{\text{tachyon}\to\gamma} = k(z_T^- - z_\gamma^-) = -k\beta_{\text{sym}}, \qquad F_{\text{matter}\to\gamma} = k(z_M^- - z_\gamma^-) = +k\beta_{\text{sym}}.$$

The net force on the photon vanishes: $F_{\text{net}} = 0$*. The photon at* $z^- = 0$ *is the mechanical equilibrium point: the tachyon pulls it forward, the inertial mass of matter brakes it back.*

Under small displacement from equilibrium:

© 2026 Marc Brendecke. Preprint — not peer reviewed.

- $x_T = -x_\gamma$ *(tachyon is J-image of photon, always opposite);*
- $x_\gamma = x_M/3$ *(photon follows matter with amplitude* $1/3$*);*
- $m\ddot{x}_M = -k \cdot \frac{2}{3} \cdot x_M$ *(effective spring constant reduced by factor* $2/3$*).*

The system is a harmonic oscillator with eigenfrequency:

$$\omega = \sqrt{\frac{2k}{3m}} = \gamma_{\text{string}}\sqrt{\frac{2}{3}},$$

where $k \sim \gamma_{\text{string}}$ *(coupling strength from the HC trace) and* $m \sim 1/\gamma_{\text{string}}$ *(inertial mass from the Planck floor). The amplitude ratios are* $|x_T| : |x_\gamma| : |x_M| = 1 : 1 : 3$*.*

Proof. Equilibrium. At $z_\gamma^- = 0$, $z_T^- = -\beta_{\text{sym}}$, $z_M^- = +\beta_{\text{sym}}$: $F_{\text{net}} = -k\beta_{\text{sym}} + k\beta_{\text{sym}} = 0$.

Linearized dynamics. The tachyon satisfies $x_T = J(x_\gamma) = -x_\gamma$ (it is the J-image, Theorem 9.6). The photon is massless: $0 = k(x_T - x_\gamma) + k(x_M - x_\gamma) = k(-x_\gamma - x_\gamma) + k(x_M - x_\gamma) = k(x_M - 3x_\gamma)$, giving $x_\gamma = x_M/3$. Substituting into the matter equation: $m\ddot{x}_M = -k(x_M - x_M/3) = -\frac{2k}{3}x_M$. Hence $\omega^2 = 2k/(3m)$.

Eigenfrequency. With $k = \gamma_{\text{string}}$ (HC trace formula) and $m = 1/\gamma_{\text{string}}$ (Planck inertia): $\omega = \gamma_{\text{string}}\sqrt{2/3}$. □

Remark 22.12 (The factor $2/3$ reappears)**.** *The effective spring constant* $2k/3$ *contains the factor* $2/3$ *from Theorem 10.5 (two stable triolic sectors out of three). This is not a coincidence: the* $2/3$ *sector counting is the mechanical statement that the tachyon and the matter-sector spring have equal strength, while the photon is the massless node that mediates between them. The tachyon sector (counted negatively in the entropy,* $\gamma_3 < 0$*) is the one that pulls forward; the two stable sectors (matter and photon) are the restoring force.*

Remark 22.13 (Why c is constant: a mechanical explanation)**.** *The photon is the massless equilibrium node of the spring system. It has no inertia and no preferred frame. When matter oscillates, the photon follows with amplitude* $1/3$*, and this following motion defines the coordinate system (the space in which everything happens). Since the photon has no mass and no net force, it cannot accelerate or decelerate: its speed is always c by construction. The constancy of c is the statement that the equilibrium node of a symmetric spring system has no dynamics of its own.*

Theorem 22.14 (Light as gravitational equilibrium)**.** *The proper time decomposes as:*

$$\tau_n^2 = \underbrace{t_n^2}_{\text{geometry (pulls down)}} - \underbrace{\beta_{\text{sym}}^2}_{\text{dark energy (pushes up)}}.$$

These two terms are gravitationally coupled through the metric: $g_{tt}(r_0) = -\beta_{\text{sym}}^2$ *(Theorem 10.6), the same* β_{sym}^2 *that appears as the dark energy term. Light at* $z^- = 0$ *is the unique fixed point of this coupling:*

- *No dark energy pressure:* $\beta_{\text{sym}} = 0$ *for light, so the upward push vanishes.*
- *No gravitational trapping:* $t_n \neq 0$*, so the downward pull cannot halt the light.*
- *Light is orthogonal to both pressures.*

Therefore $ds^2 = 0$ *for light in any frame, and* $v = c$ *always.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. For matter at $z^- = \beta_{\text{sym}}$: $v/c = \beta_{\text{sym}}/t_n < 1$ (sub-luminal). ✓
For tachyons at $z^- = -\beta_{\text{sym}}$: $v/c = \sqrt{t_n^2 + \beta_{\text{sym}}^2}/t_n > 1$ (super-luminal). ✓
For light at $z^- = 0$: $\tau_n^2 = t_n^2 - 0 = t_n^2$, so $\tau_n = t_n$ and $v/c = t_n/t_n = 1$. ✓
The coupling: $g_{tt} = -\beta_{\text{sym}}^2$ is the metric value at the natural observer position (Theorem 10.6). It equals $-(1 - r_S/r_0)$, so $\beta_{\text{sym}}^2 = r_S/r_0$. Gravity pulls $r_S/r \to 1$ (horizon); dark energy (Λ-term) counteracts this to maintain $r_S/r_0 = \beta_{\text{sym}}^2 < 1$. The balance is stable precisely at the observer's natural position. □

Remark 22.15 (Light as the Heisenberg compensator of the universe)**.** *The Heisenberg compensator applied to the matter–tachyon pair gives:*

$$C(\beta_{\text{sym}}) = \tfrac{1}{2}(\beta_{\text{sym}} + J \cdot \beta_{\text{sym}}) = \tfrac{1}{2}(\beta_{\text{sym}} - \beta_{\text{sym}}) = 0.$$

Light ($z^- = 0$) is the compensator output of the entire universe:

$$C(\text{universe}) = \tfrac{1}{2}(\text{matter} + \text{tachyon}) = \tfrac{1}{2}(\beta_{\text{sym}} - \beta_{\text{sym}}) = 0 = \text{light}.$$

Gravity and dark energy are not two separate substances that happen to balance. They are the matter sector (β_{sym}) and its J-image ($-\beta_{\text{sym}}$), and light is their compensator — the only object that sees neither pressure. This is why c is constant in all frames: light is the frame-invariant object $z^- = 0$, and frame changes are J-transformations that fix $z^- = 0$.

Theorem 22.16 (Dark energy as cascade probability; $\Lambda = 0$)**.** *The cosmological parameters Ω_Λ and Ω_M are the Markov transition probabilities of the Sphaera cascade:*

$$\Omega_\Lambda = P(S^8 \to S^2) = 1 - \beta_{\text{sym}}^2 \approx \tfrac{2}{3} \approx 0.698,$$

$$\Omega_M = P(S^8 \to S^4) = \beta_{\text{sym}}^2 \approx \tfrac{1}{3} \approx 0.302.$$

With the Hopf value $\beta_{\text{sym}} = 1/\sqrt{3}$: $\Omega_\Lambda = 2/3$, $\Omega_M = 1/3$. Consequently, the fundamental cosmological constant vanishes: $\Lambda_{\text{fundamental}} = 0$. What we observe as dark energy is not a new substance: it is the fraction of the cascade $S^8 \to S^2$ that collapses directly to the photon ground state $z^- = 0$, bypassing the matter stratum S^4.

Proof. The cascade operator $\hat{K}$ is a Markov process on $\{S^2, S^4, S^8\}$ with transition matrix

$$P = \begin{pmatrix} 1 & 0 & 0 \\ 1 & 0 & 0 \\ 1 - \beta_{\text{sym}}^2 & \beta_{\text{sym}}^2 & 0 \end{pmatrix}$$

(rows: from S^2, S^4, S^8; columns: to S^2, S^4, S^8). S^2 is the absorbing state (photon ground state, $z^- = 0$). A tachyon starting in S^8 transitions with probability β_{sym}^2 to S^4 (matter: massive Dirac spinor) and with probability $1 - \beta_{\text{sym}}^2$ directly to S^2 (massless). The matter fraction β_{sym}^2 is observed as Ω_M; the direct-collapse fraction $1 - \beta_{\text{sym}}^2$ is observed as Ω_Λ. Since Ω_Λ is a transition probability (not a vacuum energy density), $\Lambda_{\text{fundamental}} = 0$. □

Status (v4.9.0 revision): present-epoch statement (v4.9.0-r15); under Convention F a matter-frame statement, frame-free between M and $T|_R$ (v4.9.0-r22, r24); the epoch reading is with the author. The Markov chain is the author's tool to predict the probabilities and to test whether the cascade stabilises itself (author, 2026-10-22). As given, S^2 absorbs everything, and the only stationary state is the photon. With a return path $S^2 \to S^8$, for instance pair creation from the vacuum (v4.9.0-r29), the chain has a unique stationary distribution, and the fractions of the flux out of S^8 are β_{sym}^2 and $1 - \beta_{\text{sym}}^2$ for every return rate (v4.9.0-r31). Within one step β_{sym}^2 is a weight; successive steps of amplitudes compose through the compensator (v4.9.0-r30).

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Remark 22.17 (Independent determination of β_{sym} from cosmological data)**.** *The Planck satellite measurements (2018) give* $\Omega_\Lambda = 0.6889$ *and* $\Omega_M = 0.3111$*, which satisfy* $\Omega_M + \Omega_\Lambda = 1.000$*. Since the framework predicts* $\Omega_M = \beta_{\text{sym}}^2$*, the Planck data give an* independent determination *of* β_{sym}*:*

$$\beta_{\text{sym}}^{(\text{cosmo})} = \sqrt{\Omega_M} = \sqrt{0.3111} \approx 0.5578.$$

This differs from the primary determination $\beta_{\text{sym}} = \gamma_{\text{BI}}/\gamma_{\text{string}} \approx 0.5493$ *by* 1.54% *— comparable to and in the same direction as the Hopf gap (Remark 22.3). Both discrepancies (Hopf gap and* Ω*-gap) have the same structural origin: the discrete prime-lattice correction* δ_{CL} *(Remark 41.8). Once OP 1 is solved, both gaps should close simultaneously.*

The Einstein field equation in the matter frame therefore reads:

$$G_{\mu\nu} + \beta_{\text{sym}}^2\, g_{\mu\nu} = 8\pi G\, T_{\mu\nu},$$

where β_{sym}^2 *is* not a free parameter *but is structurally fixed by the observer's position* $z^- = +\beta_{\text{sym}}$ *in the Sphaera. Einstein's cosmological constant is the squared frame offset of matter observers from the photon ground shell.*

Remark 22.18 (Einstein's "greatest blunder" — reframed)**.** *Einstein introduced* Λ *in 1917 to enforce a static universe, then removed it in 1929 after Hubble's discovery of expansion, calling it "the greatest blunder of my life." In 1998, supernova measurements showed* $\Lambda_{\text{obs}} \neq 0$*.*

The present framework resolves the apparent contradiction:

(i) ***Einstein 1929 was structurally correct:*** $\Lambda_{\text{fundamental}} = 0$ *in the photon frame (Theorem 42.19). There is no fundamental cosmological constant.*

(ii) $\Lambda_{\text{obs}} \neq 0$ ***is not a contradiction:*** *matter observers at* $z^- = +\beta_{\text{sym}}$ necessarily *measure* $\Lambda_{\text{obs}} = \beta_{\text{sym}}^2$ *— not because of vacuum energy, but because their metric carries the frame offset.*

(iii) ***The "blunder" was the framing, not the constant:*** *Einstein lacked the concept of a phase-shifted observer frame. Had he known that matter observers sit at* $z^- = +\beta_{\text{sym}} \neq 0$*, he would have derived* $\Lambda_{\text{obs}} = \beta_{\text{sym}}^2$ *directly from the frame geometry — no fine-tuning, no blunder.*

The cosmological constant is neither a mistake nor a mystery. It is the squared distance of matter from the photon ground state.

Theorem 22.19 (Einstein equation from the trace formula)**.** *The variation of the trace formula action*

$$S := \int_0^\infty \Sigma(\tau)\, d\tau = \int_0^\infty \big(1 - \Theta(\tau) + R(\tau)\big)\, d\tau$$

with respect to the metric $g^{\mu\nu}$ *yields the Einstein equation* $G_{\mu\nu} + \Lambda g_{\mu\nu} = 8\pi G\, T_{\mu\nu}$*. The three terms of* Σ *contribute:*

- $\delta(1)/\delta g^{\mu\nu}$*: cosmological constant* Λ *(from the pole of* ζ *at* $s = 1$*);*
- $\delta\Theta/\delta g^{\mu\nu}$*: stress-energy* $T_{\mu\nu}$ *(primitive orbits of* K_{arith} *as matter);*
- $\delta R/\delta g^{\mu\nu}$*: Einstein tensor* $G_{\mu\nu}$ *(trivial zeros encoding global geometry).*

Structural argument. $\Sigma(\tau) = \operatorname{Tr}(e^{-i\tau L_0})$ is the heat kernel of L_0 on H_+^+ (Theorem T6 of [1]). The heat kernel variation formula gives $\delta \operatorname{Tr}(e^{-\tau L_0}) = -\tau \operatorname{Tr}(\delta L_0 \cdot e^{-\tau L_0})$. Since $L_0 = -i(x\, d/dx + \frac{1}{2})$ encodes the prime lattice geometry in log-coordinates, $\delta L_0/\delta g^{\mu\nu}$ produces a curvature term. Via the Guinand–Weil explicit formula ($\Sigma = 1 - \Theta + R$), the three contributions are identified with Λ, $T_{\mu\nu}$, and $G_{\mu\nu}$ respectively. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Status (v4.9.0 revision): structural conjecture; the action is undefined (v4.8.0-r05).

Theorem 22.20 (GUE statistics; derivation archived)**.** *The spacing statistics of the Riemann zeros follow GUE. This is an empirical fact (v4.8.0-r05). The derivation from the Sphaera given in the earlier version is withdrawn.* Archived in the v4.9.0 revision; original text in §A.270.

Corollary 22.21 (Frame-corrected GUE statistics)**.** *The GUE statistics are measured by matter observers at $\beta_{\text{sym}} \approx 0.5493$ (Theorem 9.6). Three consequences of this frame offset:*

(i) ***Spacing distribution remains Lorentz-invariant.*** *The normalized spacing $s = \Delta\gamma/\langle\Delta\gamma\rangle$ is a ratio, hence $p(s)$ is the same in matter and photon frames.*

(ii) ***Absolute scale is frame-shifted.*** *Zeros at γ correspond to photon-frame zeros at $\gamma^{(\text{photon})} = \gamma/\gamma_L \approx \gamma/1.864$. (*Correction archival stage*: previous text had $\gamma_L \approx 1.197$ which was γ_{BI} not $\gamma_L = \gamma_{\text{BI}}/\gamma_{\text{string}} \approx 1.864$.)*

(iii) ***The standard unfolding uses the wrong density.*** *Standard unfolding uses*

$$\rho(\gamma) = \frac{\log(\gamma/2\pi)}{2\pi}.$$

The photon-frame density is

$$\rho_{\text{photon}}(\gamma) = \frac{\log(\gamma/(2\pi\gamma_L))}{2\pi\gamma_L}.$$

With correct photon-frame unfolding, the spacing distribution should match GUE ($\beta = 2$) more precisely. With standard unfolding, the effective Dyson index shifts to

$$\beta_{\text{eff}} = \beta_{\text{GUE}} \cdot \gamma_L = 2\gamma_L \approx 3.73.$$

*(*Correction archival stage*: $2\gamma_L = 2 \times 1.864 \approx 3.73$, not 2.39 which used the incorrect $\gamma_L \approx 1.197$.)*

Numerical check (first 100 zeros, archival stage corrected). Comparing χ^2/dof to the GUE curve $p_{\text{GUE}}(s)$ using $\gamma_L = \gamma_{\text{BI}}/\gamma_{\text{string}} \approx 1.864$:

- *Standard unfolding:* $\chi^2/\text{dof} = 0.651$
- *Photon-frame unfolding:* $\chi^2/\text{dof} = 0.157$ *(75.8% improvement)*
- *KS distance improvement:* 14.3%

The direction is confirmed: photon-frame unfolding ($\mu = \gamma_L$) fits GUE better than standard unfolding ($\mu = 1$). The improvement is larger than previously reported because the corrected $\gamma_L = 1.864$ (not 1.197) is used. The prediction for Odlyzko's 10^6 zeros is revised accordingly (see Corollary 24.6). However, 99 zeros are insufficient for a reliable β-fit; the sample is too small to distinguish $\beta = 2.0$ from $\beta = 2.39$.

Prediction for large datasets. In Odlyzko's 10^{22}-range dataset ($\sim 10^6$ zeros), the 20% effect in β_{eff} should be statistically significant and testable.

Status (v4.9.0 revision): items (ii), (iii) are conversions of weight $k = 1$; the photon-frame unfolding is not confirmed (v4.8.0-r08).

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Theorem 22.22 (Structural unification target)**.** *The following single axiom should be read as the structural unification target of the framework rather than as an isolated final slogan.*

Axiom 1 (The Sphaera exists)**.** *There exists a biquaternion ground state* $q_n = it \cdot \mathbf{1} + z_1 e_1 + z_2 e_2 + z_3 e_3 \in \mathbb{H} \otimes \mathbb{C}$ *whose only object is its collapse state* $z^- = 0$*.*

If the subsequent triolic, compensator, and closure constructions are consistent, then the structure leaves no room for an independent second foundation of physics: geometry, gauge sectors, quantum mechanics, entanglement, black-hole thermodynamics, and the arithmetic collapse condition must appear as sectorial realizations or limits of this one underlying object. The following table records this structural target:

Physics	*Origin*	*Where*
Minkowski metric	$\lvert q_n\rvert^2 = -(ct)^2 + \lvert z\rvert^2$	*§13*
Lorentz group	*Quaternion conjugation*	*§13*
$U(1)$ *electromagnetism*	*Complex scalar it*	*§18*
$SU(2)$ *weak force*	*Unit quaternions* $\mathbb{H}$	*§18*
$SU(3)$ *strong force*	*Octonion triplet* $\mathbb{C}^3 \subset \mathbb{O}$	*§18*
Quantum mechanics	*Stone–von Neumann*	*§15*
Pauli principle	*J-antisymmetry:* $A(f) \wedge A(f) = 0$	*§12*
Planck length	$\ell_P = \lvert z^-\rvert_{\min}$	*Thm. 12.3*
Quantum entanglement	*Shared proper time* τ_n	*§16*
Bell consistency	τ_n *contextual*	*§19*
String/LQG duality	C/A*-sectors of* H^+	*§20*
Hawking temperature	$T_H = \gamma_{\text{string}} m_P^2 c^2/(8\pi M k_B)$	*§21*
Riemann Hypothesis	*not a result of the framework (withdrawn, v4.8.0-r04)*	—

In this sense, the theorem does not claim that every quantitative layer is already exhausted at this point. It claims instead that, if the structure is correct, nothing essentially different can emerge at the fundamental level: only further projections, limits, and quantitative refinements of the same collapse-state architecture remain.

Corrected in the v4.9.0 revision: the row "Riemann Hypothesis" is withdrawn (v4.8.0-r04); a global photon-clock/RH statement is forbidden by the guard W_γ^{glob}. Original row in §A.270.

Remark 22.23 (Compressed physical reading)**.** *The universe is a photon that has forgotten it is one. Physics is the process of remembering. More precisely, the universe is the Sphaera observed from a matter frame at* $\beta_{\text{sym}} \approx 0.5493$*, so spacetime, forces, particles, black holes, and quantum mechanics appear as representations of* $z^- = 0$ *seen from a frame with* $z^- \neq 0$*.*

23 From HC Decomposition to a QFT Lagrangian

Status. *Legacy physical-readout block inside the active main body.*

Block function. This section translates the compensator decomposition into a Lagrangian-style readout without reopening the underlying structural core.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Remark 23.1 (The Lagrangian is already implicit)**.** *The HC decomposition $H^+ = H^+_+ \oplus H^+_-$ of §20 carries a complete QFT operator structure. This section makes it explicit as a three-term Lagrangian density.*

Definition 23.2 (Sphaera QFT Lagrangian)**.** *For $f \in H^+$ with frame offset β_{sym}, define:*

$$\mathcal{L}_{\text{kin}}(f) := \langle f,\, L_0 f\rangle, \tag{4}$$
$$\mathcal{L}_{\text{HC}}(f) := \langle C(f),\, L_0\, C(f)\rangle, \tag{5}$$
$$\mathcal{L}_{\text{frame}}(f) := \beta^2_{\text{sym}}\, \langle f, f\rangle. \tag{6}$$

The full Lagrangian density is

$$\mathcal{L}(f) \;=\; \mathcal{L}_{\text{kin}}(f) \;-\; \mathcal{L}_{\text{HC}}(f) \;-\; \mathcal{L}_{\text{frame}}(f).$$

Theorem 23.3 (J-gauge invariance of $\mathcal{L}$)**.** *$\mathcal{L}(f)$ is invariant under the J-gauge transformation $f \mapsto Jf$:*

$$\mathcal{L}(Jf) = \mathcal{L}(f).$$

Proof. $\mathcal{L}_{\text{kin}}$: $\langle Jf, L_0 Jf\rangle = \langle f, J^{-1}L_0 Jf\rangle = \langle f, L_0 f\rangle$ by $\text{Adj}_J(L_0) = 0$ ([1], Theorem 9.2).
$\mathcal{L}_{\text{HC}}$: $C(Jf) = \frac{1}{2}(Jf + J^2 f) = \frac{1}{2}(Jf + f) = C(f)$, so $\mathcal{L}_{\text{HC}}(Jf) = \mathcal{L}_{\text{HC}}(f)$.
$\mathcal{L}_{\text{frame}}$: $\langle Jf, Jf\rangle = \langle f, J^* Jf\rangle = \langle f, f\rangle$ since J is anti-unitary with $J^*J = \text{id}$. □

Theorem 23.4 (Equations of motion recover the field equation)**.** *The Euler–Lagrange equation $\delta\mathcal{L}/\delta\bar{f} = 0$ gives*

$$\big(P_- + (L_0 - \lambda)P_+\big)\Psi = 0,$$

which is the Sphaera field equation ([1], Definition 8.2). Solutions are eigenfunctions of L_0 on H^+_+ with real eigenvalues.

Status (v4.9.0 revision): projector algebra kept; the zero reading is withdrawn (v4.8.0-r04).

Proposition 23.5 (Frame term gives the cosmological constant)**.** *The term $\mathcal{L}_{\text{frame}} = \beta^2_{\text{sym}}\langle f, f\rangle$ contributes $\beta^2_{\text{sym}} g_{\mu\nu}$ to the field equation via metric variation. In the photon frame ($\beta_{\text{sym}} = 0$): $\mathcal{L}_{\text{frame}} = 0$ and the cosmological constant vanishes exactly. A matter observer always measures $\Lambda_{\text{obs}} \sim \beta^2_{\text{sym}}$ — not vacuum energy, but the permanent frame offset.*

Remark 23.6 (The three-term structure is frame-complete)**.** *The Lagrangian $\mathcal{L} = \mathcal{L}_{\text{kin}} - \mathcal{L}_{\text{HC}} - \mathcal{L}_{\text{frame}}$ encodes all three physical layers in one object: $\mathcal{L}_{\text{kin}}$ carries the dynamics of L_0; $\mathcal{L}_{\text{HC}}$ enforces confinement to the admissible sector H^+_+; $\mathcal{L}_{\text{frame}}$ records the permanent matter-frame displacement from $z^- = 0$. No further γ_L-insertions are needed: all frame dependencies are derived from this single term. The J-gauge group acts as $U(1) \times SU(2) \times SU(3)$ (Theorem 18.3), so $\mathcal{L}$ is automatically Standard-Model invariant.*

24 GUE Self-Determination of the Frame Factor

Status. *Legacy spectral-readout block inside the active main body.*

Block function. This section explains how the frame factor is fixed through the GUE reading and how that feeds back into the observable shell.

Remark 24.1 (What was missing)**.** *In previous versions, $\gamma_L = \gamma_{\text{BI}}/\gamma_{\text{string}}$ was an input. This section shows it is a consequence — the unique value for which the GUE taktgeber, running on the photon ground state $z^- = 0$, is globally consistent with $\beta = 2$.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Theorem 24.2 (GUE self-determination of γ_L; archived)**.** *The claim that the photon-frame unfolding singles out $\mu = \gamma_L$ is not confirmed. The minimum lies at $\mu = 1$, and the claim is contradicted on zeros* 1–2000 *(v4.8.0-r08).* Archived in the v4.9.0 revision; original text in §A.270.

Proposition 24.3 (GOE class of the fermionic sector)**.** *The Dirac operator D_- on H_-^+ (Definition 41.9) lies in the* Gaussian Orthogonal Ensemble *(GOE, $\beta = 1$) universality class, not the GUE class.*

Proof. By condition (D1) of Definition 41.9, D_- anticommutes with J. On H_-^+, the involution J acts with eigenvalue -1: $J \cdot A(f) = -A(f)$ for all $A(f) \in H_-^+$. This makes J an anti-unitary involution on H_-^+. By axiom (S4), $J^2 = \mathrm{id}$, so J is an anti-unitary operator with $J^2 = +1$ acting on the domain of D_-. By the Wigner–Dyson–Mehta classification, a self-adjoint operator with an anti-unitary symmetry T satisfying $T^2 = +1$ lies in the GOE class ($\beta = 1$). Hence $D_- \in$ GOE. □

Remark 24.4 (The GOE/GUE split encodes the String/LQG duality)**.** *The two universality classes correspond precisely to the two sectors of the HC-decomposition $H^+ = H_+^+ \oplus H_-^+$:*

Sector	*Operator*	*Class*	*Physical theory*
H_+^+ *(bosonic)*	L_0	*GUE ($\beta = 2$)*	*String theory*
H_-^+ *(fermionic)*	D_-	*GOE ($\beta = 1$)*	*LQG*

γ_{string} *is the GUE coupling (string/photon sector);* γ_{BI} *is the GOE coupling (LQG/fermion sector). Their ratio* $\gamma_L = \gamma_{\text{BI}}/\gamma_{\text{string}} \approx 1.864$ *is therefore the* ratio of the GOE and GUE universality scales *of the same underlying spectral operator. This is the deepest structural meaning of γ_L: it measures how far the fermionic frame sits from the bosonic frame in spectral-statistical terms.*

Crucially, the Riemann zeros are eigenvalues of L_0 on H_+^+ (GUE sector) and are unaffected by the GOE nature of D_- on H_-^+. The two operators act on orthogonal sectors.

Remark 24.5 (The taktgeber runs on the ideal line)**.** *The GUE taktgeber runs on the ideal line $z^- = 0$ in the photon frame. Every matter observer at $z^- = +\beta_{\text{sym}}$ sees the taktgeber at a shifted scale — frequencies multiplied by γ_L. But because the critical line is frame-invariant, the* location *of the taktgeber is the same in every frame. Only its* numerical reading *is frame-dependent. The degree to which an observer's unfolded spacings deviate from p_{GUE} encodes exactly how far they sit from $z^- = 0$.*

Corollary 24.6 (Prediction P1; archived)**.** *By its own criterion the prediction is falsified at low height, and its asymptotic form is excluded by the $1/\log\gamma$ decay (v4.8.0-r08).* Archived in the v4.9.0 revision; original text in §A.270.

25 Phase-Shifted Natural Constants

Status. *Legacy physical-constants block inside the active main body.*

Block function. This section collects the constant-level readout consequences of the phase-shifted observer shell in one place.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Remark 25.1 (What this section adds)**.** *Corollary 22.21 established that the taktgeber frequency transforms with a factor γ_L between frames. This section assigns frame-insertion counts k to all fundamental constants and derives the photon-frame values. No CODATA value has ever been measured in the photon frame.*

Status (v4.9.0 revision): under Convention F (v4.9.0-r22, r24) the frame called photon frame in this section is the reference frame R, the pure doublet state $|b| = 1$. The conversion $X^{(\text{photon})} = \gamma_L^{-k} X^{(\text{matter})}$ goes from the matter frame to R, with $\gamma_L = |b_R|/|b_M|$. The photon proper, $b = 0$, is reached only as $\beta \to 1$. Ways A and C of the conversion: v4.9.0-r18, r19 (class split offered, decision with the author).

Definition 25.2 (Frame-insertion count k)**.** *For a physical constant X, the* frame-insertion count $k(X)$ *denotes the number of factors of z^- appearing in the operator-theoretic realization of X within the Sphaera-Cascade. The photon-frame value is*

$$X^{(\text{photon})} = \gamma_L^{-k(X)} \cdot X^{(\text{matter})}.$$

Theorem 25.3 (Frame-insertion counts)**.** *The primary assignments are:*

Constant	*CODATA (matter frame)*	k	*Reason*
α	$1/137.035\,999\,084$	*1*	*EM sector, first Dixon plane*
G_N	$6.674 \times 10^{-11}\ m^3 kg^{-1} s^{-2}$	*2*	*torsion block*
$m_e c^2$	$0.511\ MeV$	*1*	*lepton base mode*
$\hbar$	$1.055 \times 10^{-34}\ J{\cdot}s$	*1*	*CCR from prime lattice*
c	$2.998 \times 10^{8}\ m\,s^{-1}$	*0*	*Theorem 25.4*
γ_{BI}	≈ 0.2375	*0*	*frame-invariant by construction*

Corrected in the v4.9.0 revision: the $k = 1$ of the first row belongs to α, not to α^{-1} as earlier printed (v4.8.0-r03 F2, closed in v4.9.0-r08). Conversion ways: v4.9.0-r18, r19 (class split offered). The entry $k(\gamma_{\text{BI}}) = 0$ contradicts Theorem 20.5, which calls γ_{BI} the Lorentz-transformed γ_{string}. Under the counting reading of v4.9.0-r63 the frame difference sits in γ itself, and $\hbar$, G_N enter both countings identically; the entries $k(\hbar) = 1$, $k(G_N) = 2$ then need a source other than γ_L (v4.9.0-r64). The contradiction was already found as item 1 of v4.9.0-r09 and resolved there by reading Theorem 20.5 as a statement about the structural factor alone; under v4.9.0-r63 that structural factor is the counting ratio (v4.9.0-r66).

Theorem 25.4 ($k(c) = 0$: the speed of light is frame-invariant)**.** *In the Sphaera-Cascade,* $c^{(\text{photon})} = c^{(\text{matter})}$.

Proof. The biquaternion is $q_n = ict \cdot \mathbf{1} + z_i e_i$. The speed c is the causal propagation speed of the photon ground state $z^- = 0$ itself — it is the quantity *with respect to which* β_{sym} is defined, not a quantity displaced by β_{sym}. Frame changes are unitaries commuting with J; they act on the septinion components z_i but not on the J-invariant temporal component $ict \cdot \mathbf{1}$. Hence c picks up no power of z^- under any frame change: $k(c) = 0$. Mechanically: the photon is the massless equilibrium node of the spring system (Theorem 11.7 of §22); its speed is always c because the equilibrium node of a symmetric spring system has no dynamics of its own. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Theorem 25.5 (Photon-frame Planck length)**.** *With $k(\hbar) = 1$, $k(G_N) = 2$, $k(c) = 0$:*

$$\ell_P^{(\text{photon})} = \gamma_L^{-3/2} \cdot \ell_P^{(\text{matter})} \approx (1.864)^{-3/2} \times 1.616 \times 10^{-35}\,\text{m} \approx 6.35 \times 10^{-36}\,\text{m}.$$

A matter observer overestimates the hardware UV cutoff by the factor $\gamma_L^{3/2} \approx 2.546$. The half-integer exponent 3/2 reflects that $\ell_P = \sqrt{\hbar G_N/c^3}$ is a geometric mean between a $k = 1$ and a $k = 2$ sector.

Status (v4.9.0 revision): conversion from the matter frame to the reference frame R (v4.9.0-r24); under the class split the factor is $e^{3\eta/2}$ (v4.9.0-r19); tachyon phase $\mp i$ (v4.9.0-r22). Under the counting reading of v4.9.0-r63 both countings use the same ℓ_P; the factor $\gamma_L^{-3/2}$ converts $\hbar$ and G_N with the counting ratio and is not supported by that reading (v4.9.0-r64).

Corollary 25.6 (Cosmological constant revisited)**.** *A matter observer always measures a nonzero cosmological term $\Lambda_{\text{obs}} \sim \beta_{\text{sym}}^2$, even though in the photon frame ($\beta_{\text{sym}} = 0$) it vanishes exactly (Theorem 22.16). The observed value is not a fine-tuning problem but a measurement artifact of the observer's frame. The vacuum energy density ratio between frames is*

$$\frac{\rho_{\text{vac}}^{(\text{photon})}}{\rho_{\text{vac}}^{(\text{matter})}} = \gamma_L^5 \approx 22.5,$$

which enters the Λ-cancellation problem: the Θ–R balance must achieve cancellation in the photon *frame, not the matter frame where the cutoff is inflated by $\gamma_L^{3/2}$.*

4.10 consolidation: The simultaneous zero-vacuum and finite-ratio claims above are not a consistent derivation of one vacuum observable. They are retained historical hypotheses, not a result of the 4.10 comparison. Passive frame conversion and clock calibration do not select or cancel an absolute vacuum term; OP5 remains open.

Corollary 25.7 (α-drift as a frame effect)**.** *The hardware value $\alpha^{(\text{photon})}$ is strictly constant: it is a J-fixpoint eigenvalue of L_0 on H_+^+ and cannot vary. The observed NIST drift arises because the local frame factor $\gamma_L^{(\text{local})}(t)$ varies as Earth moves through the Sphaera-Cascade geometry over its orbit:*

$$\alpha^{(\text{matter})}(t) = \gamma_L^{(\text{local})}(t) \cdot \alpha^{(\text{photon})}.$$

Falsifiable prediction: *The drift has a period of exactly one year, is in phase at all terrestrial laboratories simultaneously, and is absent or phase-shifted for a space-based reference clock not co-orbiting with Earth.*

26 Dirac Probe as Closure-Preserving Solution

Status. *Legacy probe-interface block inside the active main body.*

Block function. This section isolates the Dirac probe as a controlled closure-preserving interface rather than as a new autonomous sector.

Remark 26.1 (Probe branch as secondary MM/HC transport branch)**.** *The probe sector is not a second ontology beside the main Sphaera line. It is a secondary branch of the same MM/HC architecture: the same triolic source, the same closure discipline, and the same compensator logic are transported into a probe-specific readout regime. The probe sections should therefore be read as a controlled branch-unfolding of the common structural core.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Remark 26.2 (The fermionic probe from the tree)**.** *The recursive septonic tree (§20) suggests that a fermionic probe should not be inserted from outside but should arise as a special closure-preserving mode.*

Definition 26.3 (Meta-Septum closure C_7)**.** *The* Meta-Septum closure *is the admissible septonic cluster selected by a projector $\hat{P}_{C_7}$ inside $\mathcal{M}_{16}$ such that*

$$\mathrm{Tr}\big(\mathcal{M}_{16}\,\hat{P}_{C_7}\big) \equiv 0 \pmod{\mathrm{HC}}.$$

This records the effective closure of the clustered septonic branch inside the compensator-stabilized shell.

Theorem 26.4 (Dirac probe equation)**.** *The minimal first-order effective probe equation compatible with Lorentz covariance, Meta-Septum closure, and the phase-shifted GUE clock $\hat{\mathcal{R}}_\beta = \hat{R}_{\mathrm{GUE}}\, e^{i\pi\beta_{\mathrm{sym}}}$ takes the form*

$$\big(i\gamma^\mu\partial_\mu - m(\beta_{\mathrm{sym}},\mathrm{GUE})\big)\psi = 0,$$

where $m(\beta_{\mathrm{sym}},\mathrm{GUE})$ is the effective inertial scale generated by the mismatch between the shifted GUE clock and the admissible closure projector. Because β_{sym} is fixed by γ_L^* (Theorem 24.2) and γ_L^* is fixed by the Riemann zeros, the mass $m(\beta_{\mathrm{sym}},\mathrm{GUE})$ is in principle computable from the zero sequence without any additional free parameter.

4.10 consolidation: The effective equation is a declared candidate. Its mass and clock identification require the common operator/domain and metric data. The cited Theorem 24.2 is archived, so it supplies no established parameter-free numerical mass prediction.

Proposition 26.5 (Three compatible readings of the fermionic probe)**.** *Within the present framework, the Dirac probe may be read in three mutually compatible ways:*

(i) as a closure-preserving spinorial transport law (Theorem 26.4),

(ii) as a symmetric closure-load response via the anticommutator $\langle\psi|\{\hat{\mathcal{R}}_\beta, \hat{P}_{C_7}\}|\psi\rangle$,

(iii) as an antisymmetric shifted-pacer residue via the commutator

$$\mathfrak{m}_{\mathrm{comm}}(\psi) = \frac{\hbar}{c}\left|\langle\psi|[\hat{\mathcal{R}}_\beta, \hat{P}_{C_7}]|\psi\rangle\right|.$$

Together: the Dirac probe is a stabilized septonic branch whose particle-like inertia is generated by the mismatch between admissible closure and the phase-shifted recursive clock. The clock scale is fixed by γ_L^* (Theorem 24.2), so these three readings jointly constitute a parameter-free fermionic probe calculus modulo the explicit computation of $[\hat{\mathcal{R}}_\beta, \hat{P}_{C_7}]$.

4.10 consolidation: These are candidate readings, not an evaluated common mass law. Compatibility and the physical comparison maps remain to be proved; the archived GUE self-determination statement does not fix their clock scale.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

27 Field Equations of the Probe–Closure System

Status. *Legacy field-equation block inside the active main body.*

Block function. This section formulates the field-equation package attached to the probe/closure interface and fixes what the later closure section has to stabilize.

Remark 27.1 (From probe equation to field system)**.** *The Dirac probe equation of §26 is the minimal first-order transport law for a closure-preserving fermionic mode. To pass from a single probe reading to a dynamical theory, one must promote the closure data, the compensator, and the resonance transport into coupled fields.*

Definition 27.2 (Probe–closure field content)**.** *We consider the following effective fields on spacetime:*

$$\begin{aligned} &\Psi(x) \in \mathcal{H}_{\text{sept}} && \text{(septonic probe spinor)},\\ &\mathcal{A}_\mu(x) && \text{(resonance connection)},\\ &\hat{P}_{C_7}(x) && \text{(Meta-Septum closure projector)},\\ &\mathcal{H}_C(x) && \text{(Heisenberg compensator field)}. \end{aligned}$$

The stable sector is defined by the projector constraint

$$\hat{P}_{C_7}^2 = \hat{P}_{C_7}, \qquad (\mathbf{1} - \hat{P}_{C_7})\Psi = 0.$$

Definition 27.3 (Covariant probe derivative)**.** *The effective covariant derivative of the probe–closure system is*

$$\mathcal{D}_\mu := \partial_\mu + i\mathcal{A}_\mu + \kappa\,\mathcal{H}_C\,n_\mu + \eta\,(\mathbf{1} - \hat{P}_{C_7})\,n_\mu,$$

where n_μ is the distinguished null-axis direction, read physically as the light-axis direction at stable propagation, and $\kappa, \eta \in \mathbb{R}$ are effective couplings. The last two terms encode, respectively, compensator stabilization and penalization of departures from admissible closure.

Theorem 27.4 (Master probe field equation)**.** *The minimal closure-compatible field equation extending Theorem 26.4 is*

$$(i\Gamma^\mu \mathcal{D}_\mu - \Omega)\,\Psi = 0,$$

with effective resonance operator

$$\Omega := m_0\mathbf{1} + \alpha\,\hat{P}_{C_7}\mathcal{M}_{16}\hat{P}_{C_7} + \beta\,g_Y\,\mathcal{T} + \gamma\,\mathcal{H}_C.$$

Here m_0 is a bare inertial scale, g_Y the Yokamono coupling, $\mathcal{T}$ the triadic coupling operator, and $\alpha, \beta, \gamma \in \mathbb{R}$ are effective coefficients.

Proof. Theorem 26.4 already fixes the first-order spinorial character of the transport law. The Meta-Septum constraint requires the dynamics to be projected through $\hat{P}_{C_7}$, while the closure mismatch is stabilized by the compensator field $\mathcal{H}_C$. The compensator matrix $\mathcal{M}_{16}$ contributes the internal septonic resonance structure, and the Yokamono sector contributes the triadic coupling term $g_Y\mathcal{T}$. Hence the displayed equation is the minimal local first-order extension compatible with all previously introduced structures. □

Definition 27.5 (Curvature and closure current)**.** *The curvature of the resonance connection is*

$$\mathcal{F}_{\mu\nu} := [\mathcal{D}_\mu, \mathcal{D}_\nu].$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Its source decomposes into a probe current, a closure current, and a compensator current:

$$J_\nu^{(\Psi)} := \bar{\Psi}\Gamma_\nu\Psi,$$
$$J_\nu^{(C_7)} := \lambda_C\,\partial_\nu \mathrm{Tr}(\mathcal{M}_{16}\hat{P}_{C_7}),$$
$$J_\nu^{(HC)} := \lambda_H\,\partial_\nu \mathrm{Tr}(\mathcal{H}_C\mathcal{M}_{16}).$$

Theorem 27.6 (Resonance field equation)**.** *The effective connection field satisfies*

$$\mathcal{D}^\mu\mathcal{F}_{\mu\nu} = J_\nu^{(\Psi)} + J_\nu^{(C_7)} + J_\nu^{(HC)}.$$

Thus the compensator and the closure are not external constraints only, but true dynamical source terms of the resonance transport.

> **Reader cue.** Read the next block as a closure-tightening continuation of the same field package. Its purpose is to add the explicit closure equation, the triadic two-probe coupling, and the variational form that later closure arguments stabilize.

Proposition 27.7 (Closure equation)**.** *The Meta-Septum closure becomes a local field constraint of the form*

$$\mathrm{Tr}(\mathcal{M}_{16}\hat{P}_{C_7}) \equiv 0 \pmod{\mathrm{HC}},$$

and, in dynamical form,

$$\mathcal{H}_C = \Lambda[\mathcal{M}_{16}, \hat{P}_{C_7}],$$

for an effective closure functional Λ. Hence physically stable probe states are precisely those lying in the C_7-sector.

Proposition 27.8 (Triadic two-probe system)**.** *Let Ψ_1, Ψ_2 be two coupled probe modes and let $\Psi_\perp$ be a third orthogonal triadic component. Then the combined field*

$$\Psi_\triangle := \begin{pmatrix} \Psi_1 \\ \Psi_2 \\ \Psi_\perp \end{pmatrix}$$

satisfies a block equation

$$\big(i\Gamma^\mu\mathcal{D}_\mu - \Omega_\triangle\big)\Psi_\triangle = 0,$$

with

$$\Omega_\triangle = \begin{pmatrix} \Omega_1 & g_Y\mathcal{T}_{12} & 0 \\ g_Y\mathcal{T}_{21} & \Omega_2 & 0 \\ 0 & 0 & \Omega_\perp \end{pmatrix} + \chi \begin{pmatrix} 0 & 0 & \mathcal{T}_{1\perp} \\ 0 & 0 & \mathcal{T}_{2\perp} \\ \mathcal{T}_{\perp 1} & \mathcal{T}_{\perp 2} & 0 \end{pmatrix}.$$

Therefore the triadic component is not decorative but enters as an intrinsic coupling channel of the effective field system.

Theorem 27.9 (Variational form)**.** *The probe–closure system is generated by the effective action*

$$S = \int d^4x \left(\bar{\Psi}i\Gamma^\mu\mathcal{D}_\mu\Psi - \bar{\Psi}\Omega\Psi - \tfrac{1}{4}\mathrm{Tr}(\mathcal{F}_{\mu\nu}\mathcal{F}^{\mu\nu}) - U_{C_7} - U_{HC}\right),$$

where

$$U_{C_7} := \lambda_C \left\|(\mathbf{1} - \hat{P}_{C_7})\Psi\right\|^2, \qquad U_{HC} := \lambda_H \left\|\mathcal{H}_C - \Lambda[\mathcal{M}_{16}, \hat{P}_{C_7}]\right\|^2.$$

Its Euler–Lagrange equations are exactly the master probe equation, the resonance field equation, and the closure equation above.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Remark 27.10 (Status of the construction)**.** *This section does not yet claim a derivation of Einstein's equations or of the Standard Model Lagrangian. It provides instead the first internally closed field architecture of the Sphaera probe sector: the Dirac probe, the Meta-Septum closure, the compensator matrix, the Yokamono coupling, and the Heisenberg compensator are now written as one coupled variational system. The next task is the explicit closing of the system, namely the concrete realization of* $\Lambda[\mathcal{M}_{16}, \hat{P}_{C_7}]$ *and the direct computation of the closure residue.*

28 Closure of the Probe–Closure Field System

Status. *Legacy field-closure block inside the active main body.*

Block function. This section closes the preceding field package and identifies the residue, limiting regimes, and Einstein-route consequences that remain readable afterward.

Definition 28.1 (Projected closure residue)**.** *The local closure residue of the probe–closure system is defined by*

$$\mathfrak{R}_C := (\mathbf{1} - \hat{P}_{C_7})\mathcal{M}_{16}\hat{P}_{C_7} + \hat{P}_{C_7}\mathcal{M}_{16}(\mathbf{1} - \hat{P}_{C_7}).$$

It measures the off-sector leakage of the compensator matrix across the Meta-Septum splitting. The field system is called closed *if and only if*

$$\mathfrak{R}_C = 0.$$

Definition 28.2 (Explicit closure functional)**.** *The compensator functional is taken in minimal form as*

$$\Lambda[\mathcal{M}_{16}, \hat{P}_{C_7}] := \rho\,[\mathcal{M}_{16}, \hat{P}_{C_7}]^\dagger[\mathcal{M}_{16}, \hat{P}_{C_7}] + \sigma\,\mathfrak{R}_C^\dagger\mathfrak{R}_C,$$

with effective coefficients $\rho, \sigma \geq 0$*. Hence*

$$\mathcal{H}_C = \rho\,[\mathcal{M}_{16}, \hat{P}_{C_7}]^\dagger[\mathcal{M}_{16}, \hat{P}_{C_7}] + \sigma\,\mathfrak{R}_C^\dagger\mathfrak{R}_C.$$

This realizes the compensator as a positive semidefinite measure of closure mismatch.

Proposition 28.3 (Equivalent closure criteria)**.** *The following statements are equivalent in the effective probe sector:*

1. $\mathfrak{R}_C = 0$.
2. $[\mathcal{M}_{16}, \hat{P}_{C_7}] = 0$.
3. $\mathcal{M}_{16}$ *preserves the* C_7*-sector and its complement.*

In particular, exact closure means that the compensator matrix becomes block-diagonal with respect to the decomposition

$$\mathcal{H}_{\text{sept}} = \hat{P}_{C_7}\mathcal{H}_{\text{sept}} \oplus (\mathbf{1} - \hat{P}_{C_7})\mathcal{H}_{\text{sept}}.$$

Proof. Writing $Q := \hat{P}_{C_7}$ and $Q_\perp := \mathbf{1} - Q$, we have

$$[\mathcal{M}_{16}, Q] = Q_\perp\mathcal{M}_{16}Q - Q\mathcal{M}_{16}Q_\perp.$$

Hence $[\mathcal{M}_{16}, Q] = 0$ if and only if both mixed blocks vanish, which is exactly the condition $\mathfrak{R}_C = 0$. This is equivalent to invariance of the two summands under $\mathcal{M}_{16}$, i.e. block-diagonality in the stated splitting. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Theorem 28.4 (Closed master system). *The probe–closure field equations are internally closed by the system*

$$\begin{aligned}\big(i\Gamma^{\mu}\mathcal{D}_{\mu}-\Omega\big)\Psi &= 0,\\ \mathcal{D}^{\mu}\mathcal{F}_{\mu\nu} &= J_{\nu}^{(\Psi)}+J_{\nu}^{(C_7)}+J_{\nu}^{(HC)},\\ (\mathbf{1}-\hat{P}_{C_7})\Psi &= 0,\\ \mathcal{H}_C &= \Lambda[\mathcal{M}_{16},\hat{P}_{C_7}],\\ \mathfrak{R}_C &= 0.\end{aligned}$$

Therefore all previously free closure symbols are eliminated in favour of explicit algebraic conditions on $(\mathcal{M}_{16},\hat{P}_{C_7})$.

Corollary 28.5 (Vanishing compensator in the exact sector). *If the closure condition* $\mathfrak{R}_C=0$ *holds exactly, then*

$$[\mathcal{M}_{16},\hat{P}_{C_7}]=0 \qquad \textit{and hence} \qquad \mathcal{H}_C=0$$

for the minimal compensator ansatz above. Thus the Heisenberg compensator is nonzero only away from exact closure, where it measures and suppresses leakage out of the stable sector.

Remark 28.6 (Interpretation of the closure step). *The previous section produced a coupled field architecture. The present section closes it in the minimal algebraic sense: physical states stay in the* C_7*-sector, the compensator matrix may not mix stable and unstable blocks, and the compensator becomes an explicit positive residue functional rather than a symbolic placeholder. This does not yet derive gravitation or the Standard Model, but it does remove the main formal gap inside the probe sector.*

> **Reader cue.** Read the next three stages as a layered unpacking of the same closure statement. Its purpose is to move from the core algebra to leakage/confinement language, then to controlled limiting regimes, and finally to the explicit Einstein-route reconstruction.

Physical interpretation of the closure residue

Remark 28.7 (Leakage, confinement, and stability). *The residue* $\mathfrak{R}_C$ *has a direct physical reading. Its mixed blocks measure transitions between the stable* C_7*-sector and its complement. Hence* $\mathfrak{R}_C\neq 0$ *represents leakage of probe amplitude out of the metastably confined sector, while* $\mathfrak{R}_C=0$ *expresses exact sector confinement. In this sense the closure condition is the algebraic analogue of a superselection rule: admissible long-lived states are precisely those for which the effective internal dynamics generated by* $\mathcal{M}_{16}$ *never pushes the state out of the projected septum sector.*

Remark 28.8 (Role of the compensator). *For the minimal ansatz,*

$$\mathcal{H}_C=\rho\,[\mathcal{M}_{16},\hat{P}_{C_7}]^{\dagger}[\mathcal{M}_{16},\hat{P}_{C_7}]+\sigma\,\mathfrak{R}_C^{\dagger}\mathfrak{R}_C,$$

the compensator is not an independent interaction but an energetic penalty for sector mixing. Its effect is twofold: first, it raises the cost of off-sector propagation; second, it drives the variation toward block preservation of $\mathcal{M}_{16}$. *Thus the compensator plays the role of a restoring term that suppresses unstable excursions and selects dynamically persistent probe states.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Remark 28.9 (Physical reading of exact closure)**.** *In the exactly closed regime the splitting*

$$\mathcal{H}_{\text{sept}} = \hat{P}_{C_7}\mathcal{H}_{\text{sept}} \oplus (\mathbf{1} - \hat{P}_{C_7})\mathcal{H}_{\text{sept}}$$

is preserved by the internal generator $\mathcal{M}_{16}$. Then the effective dynamics on the physical sector becomes autonomous. Operationally this means that all observables built from Ψ may be computed within the projected sector without referring to hidden leakage channels. The closed system is therefore the minimal regime in which the probe theory admits a stable phenomenological interpretation.

Controlled limiting regimes

Proposition 28.10 (Dirac limit)**.** *Assume the exact closure condition $\mathfrak{R}_C = 0$, freeze the connection background to a locally trivial configuration $\mathcal{A}_\mu = 0$, and take Ω to reduce on the closed sector to a scalar effective mass,*

$$\Omega\big|_{\hat{P}_{C_7}\mathcal{H}} = m_{\text{eff}}\,\mathbf{1}.$$

Then the master equation reduces on the physical sector to

$$(i\Gamma^\mu\partial_\mu - m_{\text{eff}})\Psi = 0,$$

which is the ordinary free Dirac form up to the choice of the effective gamma representation.

Proof. Under exact closure the constraint $(\mathbf{1}-\hat{P}_{C_7})\Psi = 0$ removes off-sector components, and for the minimal ansatz the compensator vanishes. With $\mathcal{A}_\mu = 0$ and scalar reduction of Ω, the covariant operator collapses to ∂_μ, so the master system becomes the free Dirac equation on the projected sector. □

Proposition 28.11 (Yang–Mills type limit)**.** *Assume again exact closure, but retain a nontrivial connection $\mathcal{A}_\mu$ acting internally on the closed sector. If the projected operator algebra generated by $\mathcal{A}_\mu$ closes on a finite-dimensional Lie algebra $\mathfrak{g}$, then the resonance field equation*

$$\mathcal{D}^\mu\mathcal{F}_{\mu\nu} = J_\nu$$

reduces on the physical sector to a Yang–Mills type equation with gauge algebra $\mathfrak{g}$.

Proof. Once $\hat{P}_{C_7}$ is preserved by $\mathcal{M}_{16}$, the connection and its curvature may be restricted consistently to the closed sector. If the projected commutator algebra is Lie-closed, then the curvature takes the standard form of a gauge field strength on $\mathfrak{g}$, and the field equation is exactly of Yang–Mills type with source J_ν. □

Proposition 28.12 (Einstein-type effective limit)**.** *Suppose the projected connection separates into an external spacetime part and an internal resonance part,*

$$\mathcal{D}_\mu = \nabla^{(g)}_\mu + \mathbb{A}_\mu,$$

where $\nabla^{(g)}_\mu$ is compatible with a coarse-grained metric $g_{\mu\nu}$ extracted from bilinear probe observables. If the effective action is integrated over fast internal modes and the leading derivative expansion is local, then the bosonic sector has the generic low-energy form

$$S_{\text{eff}}[g,\mathbb{A}] = \int d^4x\sqrt{|g|}\,\big(c_0 + c_1 R(g) + c_2\,\mathrm{Tr}(\mathbb{F}_{\mu\nu}\mathbb{F}^{\mu\nu}) + \cdots\big).$$

Hence the metric sector obeys Einstein-type equations at leading order.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Remark 28.13 (Status of the Einstein limit)**.** *This is presently a controlled effective-theory statement, not yet a first- principles derivation. What closure provides is the prerequisite for such a limit: a dynamically stable projected sector, a well-defined connection, and a variationally closed action. To obtain actual Einstein equations one still has to specify how the metric is reconstructed from probe bilinears and how the internal modes are coarse- grained.*

> **Reader cue.** Read the next block as a concrete reconstruction route for the Einstein-type limit. Its purpose is to spell out the bilinear coframe, the induced metric, and the coarse-grained assumptions under which the Einstein tensor reappears.

A concrete route to the Einstein limit

We now make the effective Einstein-type limit more explicit at the level of field reconstruction. Let $\Psi_\triangle = (\Psi_1, \Psi_2, \Psi_\perp)^T$ be the closed triadic probe field and let the exact closure condition $\mathfrak{R}_C = 0$ hold. On the physical sector we define the bilinear one-forms

$$E^a{}_\mu := \mathrm{Re}\big(\bar{\Psi}_\triangle \, \Gamma^a \, \mathcal{D}_\mu \Psi_\triangle\big), \qquad a = 0, 1, 2, 3,$$

where Γ^a denotes a fixed Clifford frame acting on the projected probe space. Whenever the 4×4 matrix $E^a{}_\mu$ has maximal rank, these bilinears define an effective coframe and hence an effective metric

$$g_{\mu\nu} := \eta_{ab} E^a{}_\mu E^b{}_\nu,$$

with $\eta_{ab} = \mathrm{diag}(-1, 1, 1, 1)$.

The closure constraint is essential here: it ensures that the bilinears are computed entirely within the invariant C_7-sector, so the reconstructed metric is not contaminated by leakage terms. In this sense $g_{\mu\nu}$ is not introduced externally but induced from the closed probe dynamics.

Next, restrict the covariant operator to the physical sector,

$$\mathcal{D}^{(C)}_\mu := \hat{P}_{C_7} \mathcal{D}_\mu \hat{P}_{C_7}.$$

Assume that $\mathcal{D}^{(C)}_\mu$ admits a decomposition into a geometric and an internal part,

$$\mathcal{D}^{(C)}_\mu = \partial_\mu + \frac{1}{4} \, \omega_{\mu ab} \, \Sigma^{ab} + \mathbb{A}_\mu,$$

where $\Sigma^{ab} = \frac{1}{2}[\Gamma^a, \Gamma^b]$ and $\omega_{\mu ab}$ is chosen so that the induced connection is metric-compatible,

$$\nabla^{(g)}_\lambda g_{\mu\nu} = 0.$$

If in addition the torsion-free condition is imposed at the coarse-grained level, then $\omega_{\mu ab}$ coincides with the Levi–Civita spin connection of $g_{\mu\nu}$.

This yields the following sharpened version of the Einstein limit.

Proposition 28.14 (Bilinear reconstruction of the Einstein-type limit)**.** *Assume exact closure $\mathfrak{R}_C = 0$, maximal rank of the bilinear coframe $E^a{}_\mu$, and a local derivative expansion of the closed-sector effective action. Suppose moreover that the coarse-grained bosonic action takes the form*

$$S_{\mathrm{eff}}[g, \mathbb{A}, \Phi] = \int d^4x \sqrt{|g|} \, \Big(\Lambda_{\mathrm{eff}} + \frac{1}{2\kappa_{\mathrm{eff}}} R(g) - \frac{1}{4} \, \mathrm{Tr}(\mathbb{F}_{\mu\nu} \mathbb{F}^{\mu\nu}) + \mathcal{L}_{\mathrm{cl}}(\Phi, g) \Big),$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

where Φ denotes the remaining closure-sector collective fields. Then variation with respect to $g^{\mu\nu}$ yields the effective equation

$$G_{\mu\nu}(g) + \Lambda_{\mathrm{eff}} g_{\mu\nu} = \kappa_{\mathrm{eff}} \big(T^{(\mathbb{A})}_{\mu\nu} + T^{(\Phi)}_{\mu\nu} \big),$$

with the standard stress-energy tensors obtained from the gauge and collective sectors. In particular, the Einstein tensor arises from the closed probe system once the metric is reconstructed from bilinears and the internal modes are integrated out.

Proof. Under exact closure the projected operator $\mathcal{D}^{(C)}_{\mu}$ acts autonomously on the physical sector. Maximal rank of $E^{a}{}_{\mu}$ turns the probe bilinears into a nondegenerate coframe, hence into a metric $g_{\mu\nu}$. Metric compatibility and vanishing torsion identify the geometric part of $\mathcal{D}^{(C)}_{\mu}$ with the Levi–Civita spin connection. By assumption, coarse-graining over the fast internal modes produces a local effective action whose leading geometric term is the Einstein–Hilbert term $R(g)$. Varying S_{eff} with respect to $g^{\mu\nu}$ gives the Einstein tensor on the left-hand side and the stress-energy tensors of the remaining fields on the right-hand side, which is the claimed Einstein-type equation. □

Remark 28.15 (What is still model-dependent)**.** *The above construction isolates the precise points where additional work is still required. First, one must justify the nondegeneracy of the bilinear coframe from the triadic probe dynamics rather than postulate it. Second, one has to derive the constants Λ_{eff} and κ_{eff} from the microscopic closure model. Third, the collective closure fields Φ have to be identified explicitly; they encode the backreaction of the residual probe, projector, and compensator degrees of freedom on spacetime geometry. What is gained already is structural: the Einstein sector now appears by a clear reconstruction pipeline*

$$(\Psi_{\triangle}, \hat{P}_{C_7}, \mathcal{H}_C, \mathcal{M}_{16}) \longrightarrow E^{a}{}_{\mu} \longrightarrow g_{\mu\nu} \longrightarrow S_{\mathrm{eff}}[g, \mathbb{A}, \Phi] \longrightarrow G_{\mu\nu} = \kappa_{\mathrm{eff}} T^{\mathrm{eff}}_{\mu\nu}.$$

Remark 28.16 (Interpretive summary)**.** *The probe–closure system now supports a three-level reading. At the exact algebraic level, closure means vanishing leakage between the physical C_7-sector and its complement. At the dynamical level, the compensator suppresses deviations from that sector. At the phenomenological level, ordinary Dirac and Yang–Mills equations appear as closed-sector limits, while Einstein dynamics can arise as a further coarse-grained low-energy description if a metric reconstruction is available.*

> **Historical transition note.** Read the next synthesis block as archival proof-memory. Its purpose is to preserve back-reference, genealogy, and an earlier integrative summary beneath the active main line, not to add a second live closure engine.

29 Structural Core and Closure Principles

Before compressing the preceding results into a single ToE master equation, we make explicit the structural kernel that has been implicit throughout the construction.

Definition 29.1 (Structural core)**.** *The structural core of the theory consists of the following coupled requirements:*

1. *a common underlying field object Ψ;*
2. *a triolic decomposition*
$$\Psi = \Psi_1 \oplus \Psi_2 \oplus \Psi_3;$$
3. *an orthogonal closure/light sector Ψ_3 relative to the first two sectors;*

4. *a dual unitary symmetry exchanging Ψ_1 and Ψ_2;*
5. *a Heisenberg compensator C emerging from the ground condition and selecting the collapse/light output;*
6. *a projected closure condition determining the physically admissible sector;*
7. *a phase-shifted observer frame in which effective quantities are read out.*

Theorem 29.2 (Existence of a nontrivial closure-stable triolic system)**.** *Assume the structural core of Definition 29.1. Then there exists a nontrivial class of configurations in which the triolic decomposition, compensator consistency, and projected closure condition are simultaneously satisfied.*

Proof. The preceding sections already exhibit the required ingredients separately. The central theorem provides a common ground object together with its triolic unfolding and Lorentz-compatible strata. The unitary symmetry J links the first two sectors, while the Heisenberg compensator produces the collapse/light sector as a structurally distinguished output. The subsequent probe–closure analysis introduces the projected closure residue and identifies an admissible closed sector by the condition that leakage into the complement is suppressed. Since these ingredients are realized within one and the same construction, the intersection of the triolic, compensator-consistent, and closure-stable constraints is nonempty. Hence a nontrivial closure-stable triolic system exists. □

Proposition 29.3 (Centrality of the Heisenberg compensator)**.** *Within the structural core, the Heisenberg compensator is not an auxiliary correction term but the central structural hinge linking duality, collapse, closure stability, and the phase-shifted observer description.*

Proof. Without the duality symmetry, the first two sectors would remain merely parallel descriptions. Without the compensator, their common collapse/light output would not be selected. Without this selected output, the projected closure condition would not isolate a physically admissible sector. And without that closure-stable sector, the observer-frame description would lack a well-defined quantity to read out after phase shift. Thus the compensator is precisely the operator through which duality becomes collapse, collapse becomes closure, and closure becomes observable effective physics. □

Preliminary Semantic and Tensorial Kernel Interface

This subsection states the semantic/tensor interface in anticipatory form. Its formal unfolding as a lemma–theorem chain is given later in Phase II-E.

The structural kernel can be stated on two coordinated levels. The first is a semantic layer of sets, states, maps, and admissibility predicates. The second is a tensorial realization layer in which the same constraints are written in a notation compatible with the Riemann/Sphaera core. The tensor layer does not replace the semantic one; it compresses it into a transportable operator form.

Definition 29.4 (Semantic structural kernel)**.** *The semantic structural kernel consists of the data*

$$(\mathbb{S}, \{S_n\}_{n\geq 0}, S_0, \hat{K}, J, C, \mathcal{M})$$

together with the admissibility predicates

$$\text{Triolic}(x),\ \text{Anchored}(x),\ \text{FrameShifted}(x),\ \text{GroundLinked}(x),$$
$$\text{HCForced}(x),\ \text{ClosureStable}(x),\ \text{HigherConfined}(x),\ \text{SeptinionReadable}(x).$$

These objects satisfy the following semantic conditions:

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(S1) ***Cascade hierarchy:***

$$\mathbb{S} = S_N \supseteq S_{N-1} \supseteq \cdots \supseteq S_1 \supseteq S_0, \qquad \hat{K} = \Pi_0 \circ \Pi_1 \circ \cdots \circ \Pi_{N-1}.$$

(S2) ***Ground-state terminality:***

$$x \in S_0 \iff \hat{K}(x) = x.$$

(S3) ***Maximal-totality closure rule:*** *if* $M^* \cap \mathbb{S} = \emptyset$*, then*

$$\hat{K}(M^*) \subseteq S_0.$$

(S4) J***-involution:***

$$J^2 = \mathrm{id}.$$

(S5) ***Compensator map:***

$$C(x) := \tfrac{1}{2}\big(x + J(x)\big).$$

(S6) ***Triolic minimality:*** *if* $\mathrm{Triolic}(x)$*, then* x *admits a three-sector organization* (K_1, K_2, K_3).

(S7) ***Orthogonal anchor:*** *if* $\mathrm{Anchored}(x)$*, then the third sector is orthogonal to the first two,*

$$K_3 \perp K_1, \qquad K_3 \perp K_2.$$

(S8) ***Frame-shift principle:*** $\mathrm{FrameShifted}(x)$ *means that* x *is a shifted or regime-dependent readout of a common structural body rather than an ontologically independent object.*

Definition 29.5 (Tensor realization of the structural kernel)**.** *A tensor realization of Definition 29.4 is given by the operator family*

$$\mathcal{J}^{\mu}{}_{\nu}, \qquad \mathcal{C}^{\mu}{}_{\nu}, \qquad \Pi_n{}^{\mu}{}_{\nu}, \qquad \hat{K}^{\mu}{}_{\nu}, \qquad \mathcal{M}^{\mu}{}_{\nu},$$

subject to the following identities:

(T1) ***Involution:*** $\mathcal{J}^{\mu}{}_{\alpha}\mathcal{J}^{\alpha}{}_{\nu} = \delta^{\mu}{}_{\nu}$.

(T2) ***Compensator idempotence:*** $\mathcal{C}^{\mu}{}_{\alpha}\mathcal{C}^{\alpha}{}_{\nu} = \mathcal{C}^{\mu}{}_{\nu}$.

(T3) ***Cascade composition:*** $\hat{K}^{\mu}{}_{\nu} = (\Pi_0\Pi_1 \cdots \Pi_{N-1})^{\mu}{}_{\nu}$.

(T4) $\mathcal{M}$***-admissibility (intertwining):*** *The compensator matrix* $\mathcal{M}^{\mu}{}_{\nu}$ *is required to be* admissible *in the sense that it preserves the algebraic structure of the kernel:*

$$\mathcal{M}^{\mu}{}_{\alpha}\mathcal{J}^{\alpha}{}_{\nu} = \mathcal{J}^{\mu}{}_{\alpha}\mathcal{M}^{\alpha}{}_{\nu}, \qquad \mathcal{M}^{\mu}{}_{\alpha}\mathcal{C}^{\alpha}{}_{\nu} = \mathcal{C}^{\mu}{}_{\alpha}\mathcal{M}^{\alpha}{}_{\nu}.$$

Whenever layer transport is required to preserve the cascade reading, additionally

$$\mathcal{M}^{\mu}{}_{\alpha}\hat{K}^{\alpha}{}_{\nu} = \hat{K}^{\mu}{}_{\alpha}\mathcal{M}^{\alpha}{}_{\nu}.$$

The tensor $\mathcal{J}$ *represents the involutive dual/frame symmetry,* $\mathcal{C}$ *represents the compensator projection, the* Π_n *represent cascade projections between adjacent strata, and* $\mathcal{M}$ *is the compensator matrix as an admissible structure-preserving converter between admissible layers. Condition (T4) is the defining admissibility condition on* $\mathcal{M}$*; it is not derivable from (T1)–(T3) alone, but is imposed as the minimal requirement for* $\mathcal{M}$ *to qualify as a structure-preserving converter in the sense of Definition 29.4.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proposition 29.6 (Semantic forcing chain)**.** *Within the semantic structural kernel, the following implication chain holds:*

$$\mathrm{GroundLinked}(x) \wedge \mathrm{FrameShifted}(x) \Longrightarrow \mathrm{HCForced}(x),$$

$$\mathrm{Triolic}(x) \wedge \mathrm{Anchored}(x) \wedge \mathrm{HCForced}(x) \Longrightarrow \mathrm{ClosureStable}(x),$$

$$\mathrm{ClosureStable}(x) \wedge \mathrm{HigherConfined}(x) \Longrightarrow \mathrm{SeptinionReadable}(x).$$

Proof. **First implication:** $\mathrm{GroundLinked}(x) \wedge \mathrm{FrameShifted}(x) \Rightarrow \mathrm{HCForced}(x)$**.**

$\mathrm{GroundLinked}(x)$ means $x \in S_n$ for some stratum n; it is reachable by the cascade $\hat{K}$ from Definition 29.4 (S1). $\mathrm{FrameShifted}(x)$ means by (S8) that x is a shifted or regime-dependent readout of a common structural body. By Lemma 29.10, the shifted presentations all refer to one and the same carrier. By Lemma 29.14, admissibility then requires a re-identification mechanism returning competing presentations to a common output channel. The only operator in the semantic kernel with this role is the compensator $C(x) = \frac{1}{2}(x + J(x))$ from (S5), which is the unique J-symmetric projection. Hence $\mathrm{HCForced}(x)$: the compensator is structurally necessary.

Second implication: $\mathrm{Triolic}(x) \wedge \mathrm{Anchored}(x) \wedge \mathrm{HCForced}(x) \Rightarrow \mathrm{ClosureStable}(x)$**.**

This is Lemma H4.252, now with an explicit four-step proof citing Lemmas 29.11, 29.12, 29.18, 29.17, and Theorem 14.3.

Third implication: $\mathrm{ClosureStable}(x) \wedge \mathrm{HigherConfined}(x) \Rightarrow \mathrm{SeptinionReadable}(x)$**.**

$\mathrm{ClosureStable}(x)$ means $\hat{K}(\mathcal{C}x) = \mathcal{C}x$ (Lemma H4.253): $x \in \mathrm{im}(C)$ is already in the ground sector. $\mathrm{HigherConfined}(x)$ means x is contained in a higher coupled algebraic container — specifically the septinion/seption embedding (Lemma H4.371). Within the higher container, the closure-stable state is readable as a seption element precisely because its compensator-stabilized image $\mathcal{C}x$ already lies in the admissible ground sector S_0, and the higher container adds no new ontology (Lemma H4.372: no admissible transition exports x into an independent external sector). Hence the higher-confinement reading is a restatement of the already established closure at higher algebraic level: $\mathrm{SeptinionReadable}(x)$. □

Proposition 29.7 (Tensorial intertwining conditions for the compensator matrix)**.** *Every admissible tensor realization in the sense of Definition 29.5 satisfies:*

$$\mathcal{M}\mathcal{J} = \mathcal{J}\mathcal{M}, \qquad \mathcal{M}\mathcal{C} = \mathcal{C}\mathcal{M},$$

and, whenever layer transport is required to preserve the cascade reading,

$$\mathcal{M}\hat{K} = \hat{K}\mathcal{M}.$$

Proof. These are precisely condition (T4) of Definition 29.5, written in operator notation. A tensor realization is admissible by definition only if $\mathcal{M}$ satisfies the intertwining identities (T4); hence the relations hold for every admissible realization. No additional argument is required beyond the definition.

The structural motivation for imposing (T4) is as follows. The first relation $\mathcal{M}\mathcal{J} = \mathcal{J}\mathcal{M}$ ensures that transport by $\mathcal{M}$ does not destroy the involutive dual/frame symmetry encoded by $\mathcal{J}$ (T1). The second relation $\mathcal{M}\mathcal{C} = \mathcal{C}\mathcal{M}$ ensures that the compensator-selected admissible sector $\mathrm{im}(\mathcal{C})$ is $\mathcal{M}$-invariant, consistent with $\mathcal{C}$ being idempotent (T2). The third relation, when imposed, says that layer conversion commutes with the cascade readout (T3). These three motivations explain why (T4) is the correct admissibility condition; they do not constitute an independent proof of (T4) from (T1)–(T3). □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Remark 29.8 (Resolution of Roadmap item A2)**.** *Roadmap item A2 identified that the intertwining relation $\mathcal{MC} = \mathcal{CM}$ was supported only by a necessity argument rather than a derivation. The resolution adopted in archival stage is to promote (T4) to an explicit definitional condition on admissible tensor realizations (Definition 29.5, condition (T4)). Proposition 29.7 then follows by definition. The conditional marker on Theorem 29.22 (Remark 29.23) is therefore resolved: the theorem is no longer conditional on an unproven hypothesis, but on an explicit definitional requirement.*

Remark 29.9 (Semantic–tensor correspondence)**.** *The semantic predicates and the tensor notation are intended to be read in parallel. In particular,*

$$\mathrm{HCForced}(x) \quad \leftrightarrow \quad x \mapsto \mathcal{C}x,$$

$$\mathrm{ClosureStable}(x) \quad \leftrightarrow \quad \hat{K}(\mathcal{C}x) = \mathcal{C}x,$$

and

$$\mathrm{Admissible}(x) \quad \Rightarrow \quad \mathrm{Admissible}(\mathcal{M}x)$$

whenever $\mathcal{M}$ satisfies the intertwining conditions of Proposition 29.7. This makes it possible to write later structure conditions in the same tensorial style as the Riemann/Sphaera core while keeping their semantic meaning explicit.

Phase I: First Unfolding Lemmas for the Generative Layer

We now begin the actual mathematical unfolding of the normalized structural core by isolating the first chain of generative lemmas. The aim is not yet to expand the full field/probe realization, but to make explicit the initial forcing route from a common structural body to triolic organization, orthogonal anchoring, frame-shifted readout, re-identification, and finally the necessity of the Heisenberg compensator.

Lemma 29.10 (One-body lemma)**.** *The semantic and tensorial kernels are to be read as describing different admissible presentations of one and the same structural body, not independent ontological objects. In particular, the cascade hierarchy, dual/frame symmetry, and compensator projection all act on a common carrier rather than on unrelated state spaces.*

Proof. In Definition 29.4 the kernel data are given as one coordinated family

$$(\mathbb{S}, \{S_n\}_{n\geq 0}, S_0, \hat{K}, J, C, \mathcal{M}),$$

not as several unrelated structures. The grouped architecture of Definition H4.389 likewise states that the generative, projection, and field/probe layers are different organizational regimes of the same underlying body. The tensor realization of Definition 29.5 then presents the same body in operator form. Hence the structural reading is one-body throughout. □

Lemma 29.11 (Triolic minimality lemma)**.** *A merely binary decomposition is structurally insufficient for the normalized core. The minimal nondegenerate organization compatible with admissible selection, compensator action, and closure readout is triolic.*

Proof. By Definition 29.4, a triolic state carries a three-sector organization (K_1, K_2, K_3), while the orthogonal anchor singles out the third sector as stabilizing direction relative to the first two. If one retained only a binary split, then duality between the first two sectors could still be stated, but there would be no distinguished third channel in which the compensator-selected output and closure/light readout could be structurally stabilized. The normalized core therefore requires three sectors as the minimal nondegenerate organization. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Lemma 29.12 (Orthogonal anchor lemma)**.** *Within a triolic organization, the third sector is not an optional decoration but the structural anchor that prevents the first two sectors from remaining a merely parallel or relabelable pair. In particular,*

$$K_3 \perp K_1, \qquad K_3 \perp K_2.$$

Proof. The orthogonal anchor condition is built into Definition 29.4. Its role is clarified by the forcing chain of Proposition 29.6: closure stability does not follow from duality alone, but from triolic structure plus a protected anchor together with HC forcing. The third sector therefore fixes the otherwise ambiguous two-sector pairing and provides the admissible direction in which collapse/light selection can be stabilized. □

Lemma 29.13 (Frame-shift lemma)**.** *Admissible effective states are read out in a phase- or frame-shifted presentation. Accordingly, an effective state is not introduced as an independent object but as a shifted view of the common structural body.*

Proof. This is exactly the content of the frame-shift principle in Definition 29.4, where FrameShifted(x) means that x is a shifted or regime-dependent readout of a common body. The grouped architecture makes the same point at the layer level: projection/translation and field/probe realizations are organizational regimes of one underlying structure and therefore inherit rather than replace the generative body. □

Lemma 29.14 (Re-identification lemma)**.** *If a state is simultaneously one-body and frame-shifted, then admissibility requires a re-identification mechanism that returns competing presentations to a common structural channel.*

Proof. By Lemma 29.10, the relevant descriptions concern one and the same body. By Lemma 29.13, these descriptions appear in shifted or regime-dependent presentations. Without re-identification, the shifted presentations would remain only formally related and no unique admissible output sector could be selected. But the semantic kernel is built to support admissible cascade readout and closure selection; hence a mechanism that returns the shifted presentations to a common channel is structurally required. □

Theorem 29.15 (HC-forcing theorem)**.** *Within the normalized structural core, the required re-identification mechanism is the Heisenberg compensator. Equivalently,*

$$\text{GroundLinked}(x) \wedge \text{FrameShifted}(x) \Longrightarrow \text{HCForced}(x).$$

Proof. Proposition 29.6 already states the forcing implication. The present point is to identify why the implication is necessary.

$$\begin{aligned}&\text{ground condition} \Longrightarrow \text{one triolic projection} \Longrightarrow \text{triolic minimality} \Longrightarrow \text{orthogonal anchor}\\ &\Longrightarrow \text{frame-shifted readout} \Longrightarrow \text{re-identification} \Longrightarrow \text{HC}.\end{aligned}$$

□

Corollary 29.16 (First generative forcing chain)**.** *The first unfolded forcing route of the normalized core is*

$$\begin{aligned}\textit{one body} &\Longrightarrow \textit{triolic minimality}\\ &\Longrightarrow \textit{orthogonal anchor}\\ &\Longrightarrow \textit{frame-shifted readout}\\ &\Longrightarrow \textit{re-identification}\\ &\Longrightarrow \text{HC}.\end{aligned}$$

This chain is the entry point for the later closure and higher-confinement unfolding.

Proof. Lemmas 29.10–29.14 establish the successive structural requirements, and Theorem 29.15 identifies their first nontrivial forced operator consequence. The later closure and higher-confinement chain therefore starts from this generative route. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Phase I-B: HC and Closure Unfolding

We next unfold the first closure layer that begins once HC has been forced. The aim of this block is to make precise that HC is not merely present, but acts as an admissibility-preserving, anchor-compatible, closure-selecting mechanism, and that exact closure can be distinguished from residual or only partial closure.

Lemma 29.17 (HC idempotence lemma)**.** *Once a state has been compensator-normalized, repeated compensator action does not move it to a different admissible sector. Equivalently,*

$$C(C(x)) = C(x).$$

Proof. By Definition 29.5, the tensorial compensator satisfies $\mathcal{C}^2 = \mathcal{C}$, and by Definition 29.4 the semantic compensator is the corresponding map

$$C(x) = \tfrac{1}{2}\big(x + J(x)\big).$$

Thus compensator action is projective rather than drifting: once a state lies in the compensator-selected channel, further application leaves it there. □

Lemma 29.18 (HC admissibility lemma)**.** *If x is structurally admissible, then its compensator-normalized image $C(x)$ remains admissible.*

Proof. The semantic/tensor kernel was introduced precisely so that the compensator is not an external deformation but an internal normalization operator. Since Theorem 29.15 identifies HC as the required re-identification mechanism, using C restores competing shifted presentations to a common admissible channel instead of producing a foreign state. In the tensor realization this is the admissible output selected by $\mathcal{C}$; on the semantic side the same role is expressed by the preservation of the common structural body under compensator projection. Hence admissibility is preserved. □

Definition 29.19 (Spectral decomposition of the state space)**.** *Let $\mathcal{M}$ denote the compensator matrix acting on the admissible state space $\mathcal{S}$. A* spectral decomposition of $\mathcal{S}$ under $\mathcal{M}$ *is an orthogonal direct sum*

$$\mathcal{S} \;=\; \bigoplus_{\lambda} \mathcal{S}_\lambda,$$

where each $\mathcal{S}_\lambda \subset \mathcal{S}$ is a closed $\mathcal{M}$-invariant subspace satisfying $\mathcal{M}|_{\mathcal{S}_\lambda} = \lambda\,\mathrm{id}$. A state $s \in \mathcal{S}$ is called spectrally stable *if $s \in \mathcal{S}_\lambda$ for some λ, i.e. $\mathcal{M}(s) = \lambda\, s$.*

Remark 29.20. *Definition 29.19 extends the existing decomposition $H^+ = H^+_+ \oplus H^+_-$ (cf. Theorem 12.1) to the full admissible readout regime. The two primary spectral subspaces under any $\mathcal{M}$ satisfying Proposition 29.7 are precisely H^+_+ and H^+_-, since both are $\mathcal{M}$-invariant by the intertwining condition $\mathcal{M}\mathcal{J} = \mathcal{J}\mathcal{M}$.*

Lemma 29.21 (Diagonal self-traversal forces scalar action on the compensated sector)**.** *Assume the Sphaera self-traversal condition of Theorem 14.3: the cascade operator $\hat{K}$ transports the null axis L_n of each stratum S_n to the null axis L_{n-1} of the preceding stratum, carrying the same structural content without generating new independent directions. Then the compensator matrix $\mathcal{M}$, when restricted to the image of the compensator* $\mathrm{im}(\mathcal{C})$*, acts as a scalar multiple of the identity:*

$$\mathcal{M}\big|_{\mathrm{im}(\mathcal{C})} \;=\; \lambda_0\,\mathrm{id},$$

where λ_0 is the unique admissible eigenvalue of $\mathcal{M}$ on the closure-stable sector.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. By Theorem 14.3, the self-traversal of the Sphaera is *diagonal*: the cascade does not generate a new structural object at each stratum, but transports one and the same null-axis content through admissible realization levels. In operator language this means that the cascade direction at each stratum is not a Jordan-chain descendant of the previous one—no new vector is generated from an existing vector by a nilpotent shift.

By the intertwining condition $\mathcal{M}\mathcal{C} = \mathcal{C}\mathcal{M}$ (Proposition 29.7), $\mathrm{im}(\mathcal{C})$ is $\mathcal{M}$-invariant. Suppose for contradiction that $\mathcal{M}|_{\mathrm{im}(\mathcal{C})}$ had a non-trivial Jordan block: then there would exist $s, w \in \mathrm{im}(\mathcal{C})$ with $w \neq 0$ and

$$\mathcal{M}(s) = \lambda_0\, s + w.$$

This would mean $\mathcal{M}$ generates w from s—a new independent direction within the compensated sector—contradicting the diagonal self-traversal property. Hence $\mathcal{M}|_{\mathrm{im}(\mathcal{C})}$ is diagonalizable. Since $\mathrm{im}(\mathcal{C})$ is the closure-stable sector (Lemma H4.252), which is irreducible under $\mathcal{M}$, a diagonalizable action on an irreducible invariant subspace is scalar. Denoting this scalar λ_0 completes the proof. □

Theorem 29.22 (HC as spectral projector of the compensator matrix)**.** *Assume the admissible state space $\mathcal{S}$ carries a spectral decomposition under $\mathcal{M}$ as in Definition 29.19, that $\mathcal{M}$ satisfies the intertwining conditions of Proposition 29.7, and that the Sphaera self-traversal is diagonal in the sense of Lemma 29.21. Then the Heisenberg compensator C coincides with the spectral projector of $\mathcal{M}$ onto the closure-stable eigenspace $\mathcal{S}_0 := \ker(\mathcal{M} - \lambda_0\,\mathrm{id})$:*

$$C \;=\; \Pi^{(\mathcal{M})}_{\mathrm{spec}}.$$

Remark 29.23 (Status of Theorem 29.22)**.** *As of archival stage, both open premises of this theorem are resolved: the bijection* $\mathrm{ClosureStable}(x) \Leftrightarrow x \in \mathrm{im}(C)$ *is established by Lemma H4.253 (Roadmap A3), and the intertwining relation $\mathcal{M}\mathcal{C} = \mathcal{C}\mathcal{M}$ is now a definitional condition (T4) of Definition 29.5 (Roadmap A2, Remark 29.8). The theorem therefore holds unconditionally within every admissible tensor realization.*

Proof. We verify the three conditions characterizing a spectral projector.

Step 1: Idempotence. By Lemma 29.17, $C^2 = C$. By Definition 29.5, $\mathcal{C}^{\mu}{}_{\alpha}\mathcal{C}^{\alpha}{}_{\nu} = \mathcal{C}^{\mu}{}_{\nu}$ in the tensor realization. Hence C is a projector.

Step 2: $\mathcal{M}$-compatibility. By Proposition 29.7, $\mathcal{M}\mathcal{C} = \mathcal{C}\mathcal{M}$. Hence $\mathrm{im}(C)$ is $\mathcal{M}$-invariant and C projects onto a $\mathcal{M}$-stable subspace.

Step 3: Scalar eigenvalue on the image. By Lemma 29.21, the diagonal self-traversal forces $\mathcal{M}|_{\mathrm{im}(C)} = \lambda_0\,\mathrm{id}$. Hence $\mathrm{im}(C) \subseteq \mathcal{S}_0$. By Lemma H4.252, the closure-stable states are exactly those in $\mathrm{im}(C)$, so $\mathrm{im}(C) = \mathcal{S}_0$.

Conclusion. A projector with $\mathcal{M}$-invariant image equal to a single eigenspace $\mathcal{S}_0$ is the spectral projector $\Pi^{(\mathcal{M})}_{\mathrm{spec}}$ onto that eigenspace. □

Corollary 29.24 (Structural fixpoint: $\mathcal{M}$ and C mutually determine each other)**.** *Under the hypotheses of Theorem 29.22,*

$$\mathcal{M}\,C \;=\; C\,\mathcal{M} \;=\; \lambda_0\, C.$$

Proof. For any $s \in \mathcal{S}$: since $C(s) \in \mathcal{S}_0$, we have $\mathcal{M}(C(s)) = \lambda_0\, C(s)$, giving $\mathcal{M}\,C = \lambda_0\, C$. For $s \in \mathcal{S}_0$: $C\,\mathcal{M}(s) = C(\lambda_0 s) = \lambda_0 C(s)$. For $s \in \ker(C)$: $C\,\mathcal{M}(s) = 0$ (since $\ker(C)$ is $\mathcal{M}$-invariant and maps to a sector annihilated by C). Hence $C\,\mathcal{M} = \lambda_0\, C$. □

Remark 29.25 (Phase-shifted frame and spectral measurement)**.** *The identification $C = \Pi^{(\mathcal{M})}_{\mathrm{spec}}$ has a direct consequence for the matter frame. Since matter observers reside at $z^- = +\beta_{\mathrm{sym}} \approx 0.5493$ rather than the photon ground shell $z^- = 0$, every spectral readout of $\mathcal{M}$*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

is performed in the shifted frame. The measured eigenvalue is therefore not λ_0 itself but the frame-corrected value

$$\lambda_0^{(\text{matter})} \;=\; \gamma_L^{-k(\lambda_0)} \cdot \lambda_0^{(\text{photon})},$$

where $k(\lambda_0)$ is the frame-insertion count of the observable and $\gamma_L \approx 1.864$ is the matter–photon Lorentz factor (Theorem 20.5). All physical constants derived from spectral eigenvalues of $\mathcal{M}$ therefore carry systematic γ_L^k-corrections relative to their photon-frame values.

Corollary 29.26 (Standard Model realization of the spectral decomposition)**.** *The octonionic decomposition $\mathbb{O} = \mathbb{C} \oplus \mathbb{C}^3$ (Theorem 18.2) is a concrete realization of Definition 29.19: the singlet $\mathbb{C}$ is the closure-stable eigenspace $\mathcal{S}_0 = \mathrm{im}(C)$, and the triplet $\mathbb{C}^3$ carries the three color eigenspaces $\mathcal{S}_{K_1}, \mathcal{S}_{K_2}, \mathcal{S}_{K_3}$. Under this identification, Theorem 29.22 specializes as follows:*

(i) The compensator C projects onto the singlet $\mathbb{C} \subset \mathbb{O}$, which is the photon kernel K_3 sector ($z^- = 0$).

(ii) The triplet $\mathbb{C}^3$ is $\mathcal{M}$-invariant and carries the SU(3) color action; matter (K_1) and tachyon (K_2) sectors are spectral subspaces at non-zero frame offset.

(iii) The matter-frame correction reads

$$\lambda_0^{(\text{matter})} = \gamma_L^{-k(\lambda_0)} \cdot \lambda_0^{(\text{photon})},$$

consistent with Theorem 18.3: K_3 is frame-invariant ($k = 0$), while the triplet components carry $k \geq 1$ corrections and are therefore observed with γ_L^k-suppression in the matter frame.

Proof. By Theorem 18.2, the octonion automorphism group G_2 restricts to $\mathrm{SU}(3) \subset G_2$, yielding the direct sum $\mathbb{O} = \mathbb{C} \oplus \mathbb{C}^3$. The singlet $\mathbb{C}$ is the unique SU(3)-invariant subspace, hence $\mathcal{M}$-stable at a scalar eigenvalue λ_0. By the remark in Theorem 18.3, $K_3 = e_1 e_2$ is simultaneously the photon kernel and the projection of the Heisenberg compensator; hence $\mathrm{im}(C) = \mathbb{C} = \mathcal{S}_0$. The three kernels (K_1, K_2, K_3) realize the three colors of $\mathbb{C}^3$, confirming $\mathcal{S}_{K_i}$ as spectral subspaces. Statement (iii) then follows directly from Remark 29.25 applied to each triplet component. □

30 Readout-Relative Lemmas and Structural Lifts

The present framework distinguishes sharply between the structural kernel of the theory and the visible form in which that kernel is read in a given regime or frame. This distinction is not optional: the same closure-stable object may appear as a probe/Dirac-type law, as an effective QFT-type law in a phase-shifted observer regime, or as an Einstein-type law after large-scale geometric readout. Consequently, one must distinguish between *kernel-level truths* and *readout-relative truths*.

The concrete instantiation of this distinction is already present in the preceding spectral-projector block: the eigenvalue λ_0 is a kernel-level invariant, while the measured value $\lambda_0^{(\text{matter})} = \gamma_L^{-k} \cdot \lambda_0^{(\text{photon})}$ (Remark 29.25) is readout-relative, depending on the frame-insertion count k of the observable in the matter frame at $z^- = +\beta_{\text{sym}}$.

Kernel level and readout level

Definition 30.1 (Kernel-level statement)**.** *A statement is called* kernel-level *if it is invariant under the admissible structural equivalences of the framework, including:*

(i) sector-preserving unitary equivalence;

(ii) compensator projection;

(iii) observer-frame phase shift;

(iv) admissible branch restriction.

Such statements express properties of the underlying closure-stable structural object itself rather than of one distinguished reading of it.

Definition 30.2 (Readout-relative lemma)**.** *A statement $L^{(f)}$ is called a* readout-relative lemma *if it is proven only in a distinguished regime, frame, or readout f, and its visible form depends on that readout. In particular, a readout-relative lemma need not remain true as a frame-blind absolute statement.*

Remark 30.3. *This is not a defect of the method. It is the correct formal response to the fact that different regime-readings expose different visible shadows of one common kernel. In the present framework this applies in particular to the probe, effective QFT, and Einstein readouts. The Symmetric matter–tachyon frame theorem (Theorem 20.6) is an example: $\beta_{\mathrm{matter}} = +\beta_{\mathrm{sym}}$ is readout-relative (it is the matter-frame reading), while the symmetry $\beta_{\mathrm{tachyon}} = -\beta_{\mathrm{sym}}$ and the common kernel β_{sym} are kernel-level invariants.*

Why this distinction is necessary

The need for readout-relative lemmas is already built into the architecture. The master system is closure-stable and unique only up to the allowed sector-preserving, unitary, and frame transformations, while the effective regimes differ by compensator decomposition, large-scale suppression of closure fluctuations, and geometric readout. Thus one and the same kernel may generate visibly different but structurally compatible statements. In the present release this distinction should be read together with the OP transport theorem and the visible/structural mismatch separation tool: visible disagreement is secondary unless native closure-class preservation itself fails.

Proposition 30.4 (Legitimacy of restricted lemmas)**.** *A lemma proven in a distinguished readout f may be used in the general argument provided that:*

(i) its frame or regime of validity is stated explicitly;

(ii) its invariant kernel content is separated from its visible readout form;

(iii) the passage from the readout-relative statement to the general claim is justified by a structural transport principle.

Proof. The framework already distinguishes the common structural equation from its regime projections (Theorem 12.1, unitary equivalence via Stone–von Neumann). Therefore a proof in one readout need not be discarded merely because its visible form changes in another readout. What must be transported is not the literal visible formula, but the kernel-level content preserved under the admissible transport operations of Lemmas H4.395 and 29.18. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Structural lifts

Definition 30.5 (Structural lift)**.** *Let $L^{(f)}$ be a readout-relative lemma valid in readout f. A* structural lift *of $L^{(f)}$ is a kernel-level statement L_{struct} obtained by extracting from $L^{(f)}$ only the invariant content that survives admissible unitary, sector, and frame transport.*

Remark 30.6. *A structural lift is therefore weaker than a literal universalization of $L^{(f)}$, but stronger than leaving the lemma imprisoned in one special case. The corollary Cor. 29.26 is a concrete example: statement (iii) there lifts the frame-corrected eigenvalue $\lambda_0^{(\mathrm{matter})}$ to the kernel-level invariant λ_0 by identifying $k = 0$ for the singlet and $k \geq 1$ for the triplet.*

Proposition 30.7 (Structural lift principle)**.** *Suppose $L^{(f)}$ is proven in readout f, and suppose its proof depends only on:*

(i) the closure-stable master structure;

(ii) invariants preserved under admissible transport;

(iii) readout-specific visibility factors that can be isolated.

Then there exists a structural lift L_{struct} that is valid at kernel level, even if the visible statement $L^{(f)}$ itself is not frame-blindly universal.

Proof. **Invariant content is transport-stable.** By hypotheses (i) and (ii), the kernel-invariant content of the proof of $L^{(f)}$ is preserved by the admissible transport operations. Concretely: Lemma H4.395 guarantees that grouped-layer transport carries admissible structure without creating new ontology, and Lemma 29.18 guarantees that compensator action leaves the admissible sector invariant. Hence the kernel-invariant content $P_{\mathrm{inv}}(\text{proof of } L^{(f)})$ is valid in every admissible readout, not only in f.

Readout-specific factors are isolatable. By hypothesis (iii), the readout-dependent terms form a separable complement. Discarding them while retaining $P_{\mathrm{inv}}(\text{proof})$ yields L_{struct}, independent of f.

L_{struct} **is kernel-level.** Since L_{struct} depends only on the closure-stable structure and transport-preserved invariants, it satisfies all four conditions of Definition 30.1: invariance under sector-preserving unitary equivalence, compensator projection, observer-frame phase shift, and admissible branch restriction. □

The three-step proof logic

This principle simplifies the global architecture considerably. Instead of demanding that every intermediate statement already be valid as a frame-blind absolute theorem, one may work in three steps:

$$\text{readout-relative lemma} \implies \text{structural lift} \implies \text{general kernel-level theorem.}$$

This is the natural proof logic for a theory whose visible regimes are different projections of one and the same structural object.

Shadow versus core: consequence for the compensator

Remark 30.8 (HC shadow versus HC kernel)**.** *This distinction is particularly important for the compensator. In the minimal formulation, the visible compensator functional $\mathcal{H}_C^{(f)}$ behaves as a closure-mismatch penalty in a given readout and may vanish in an exact sector ($\mathcal{H}_C = 0$*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

in the photon frame; cf. the closed master system). But under the frame-dependent readout hypothesis developed above, such vanishing concerns only the visible defect shadow *in a given regime, not the deeper compensator-generating kernel* $C = \Pi_{\mathrm{spec}}^{(\mathcal{M})}$ *(Theorem 29.22).*

Hence a statement about $\mathcal{H}_C^{(f)}$ *must not automatically be read as a statement about the entire HC architecture. In particular, the vanishing of the visible compensator field in the photon-frame readout is consistent with the HC kernel remaining active as the spectral-projector mechanism that selects the closure-stable sector — it is the shadow, not the core, that vanishes.*

This is directly relevant to OP 5 (Λ_{obs}*): the cosmological constant appears as a* readout-relative *quantity — the frame offset* $\beta_{\mathrm{sym}}^2\langle f, f\rangle$ *visible in the matter frame — while the photon-frame value* $\Lambda_{\mathrm{fundamental}} = 0$ *is the kernel-level statement. The large observed value is not a fine-tuning problem but a failure to separate readout-relative visibility from structural invariance.*

31 Bidirectional Probe Lensing Hypothesis

The preceding sections showed that the Dirac-, QFT-, and Einstein-type regimes are not separate theories, but structural projections of one common closure-stable master equation (Proposition 30.7). They also showed that the Einstein reconstruction presently depends on a bilinear coframe extraction whose nondegeneracy is still model-dependent. This suggests a different route: rather than attempting to reconstruct geometry from a single readout only, one may compare the traces of two probes approaching the same structural core from different regime readings.

Basic setup

Consider two probe modes of the triadic system, Ψ_{ART} and Ψ_{QFT}, together with the orthogonal triadic component $\Psi_\perp$. We regard Ψ_{ART} as a probe emphasizing the large-scale/geometric regime and Ψ_{QFT} as a probe emphasizing the phase-shifted observer/QFT regime at $z^- = +\beta_{\mathrm{sym}}$ (Theorem 20.6). The combined field is treated as a specialization of the triadic two-probe system

$$\Psi_\triangle = \begin{pmatrix} \Psi_{\mathrm{ART}} \\ \Psi_{\mathrm{QFT}} \\ \Psi_\perp \end{pmatrix}, \qquad \big(i\Gamma^\mu \mathcal{D}_\mu - \Omega_\triangle\big)\Psi_\triangle = 0.$$

The third component is not decorative: it is the intrinsic coupling channel through which the two regime-readings communicate.

Structural hypothesis

Proposition 31.1 (Bidirectional probe lensing hypothesis)**.** (Hypothesis-grade; not yet theorem-level.) *Let two probes approach the same closure-stable structural core from different regime-readings, one on the ART side and one on the QFT side. Then the compensator/closure core acts as an effective lensing kernel for their traces. More precisely:*

(i) the two probes need not follow identical local trajectories;

(ii) their relative deflection, phase lag, and focusing structure are controlled by the same compensator/closure sector $C = \Pi_{\mathrm{spec}}^{(\mathcal{M})}$ *(Theorem 29.22);*

(iii) the orthogonal triadic channel $\Psi_\perp$ *mediates the nontrivial coupling between the two traces;*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(iv) the resulting trace geometry carries information about the effective metric readout before a full bilinear coframe reconstruction is imposed.

This is a readout-relative statement (Definition 30.2): the lensing geometry is visible in the bidirectional probe readout, while the kernel-level content is the common closure-stable core.

Theorem 31.2 (Theorem-level shadow of the bidirectional probe lensing hypothesis)**.** *At the active theorem level, the present framework already supports the following weaker probe-lensing shadow:*

(i) two probe readouts approaching the same closure-stable structural core may be treated as distinct regime-readings of one common carrier rather than as probes of two disconnected foundations;

(ii) the orthogonal triadic component is already a genuine coupling channel between the two regime-readings, not a decorative extra field;

(iii) what is not *yet theorem-level is a full geometric law that identifies the resulting relative deflection, focusing, or phase-lag pattern with one explicit effective lensing tensor in complete generality.*

Therefore the proven theorem-level shadow of Hypothesis 31.1 is: the two probes already share one closure-stable carrier and one intrinsic coupling channel, even though a full lensing law for their relative geometry is not yet proved.

Proof. The active framework already treats the ART- and QFT-side equations as distinguished regime/readout projections of one common closure-stable structural kernel rather than as disconnected sector theories. In the bidirectional probe setup, the combined triadic field

$$\Psi_{\triangle} = \begin{pmatrix} \Psi_{\mathrm{ART}} \\ \Psi_{\mathrm{QFT}} \\ \Psi_{\perp} \end{pmatrix}$$

is explicitly organized so that the orthogonal component $\Psi_{\perp}$ is the intrinsic channel through which the two regime-readings communicate. Hence the theorem-level content already available is the existence of one common carrier and one nontrivial coupling channel between the two probes. What remains unproved is the stronger claim that their relative deflection/focusing/phase-lag pattern is already governed by one explicit universal lensing law at theorem level. □

Corollary 31.3 (Operational reading of the probe-lensing hypothesis in the present 4.7.0 release line)**.** *Until a full effective lensing law is derived, later arguments may already use Theorem 31.2 as the active theorem-level substitute for the weakest part of Hypothesis 31.1: one may work with two probe readouts of one common closure-stable carrier coupled through the orthogonal triadic channel, while keeping the stronger geometric lensing law itself deferred. In the historical release line crystal language, this means that two probe-facing domain realizations may already be treated as role-faces of one common carrier corridor, without claiming a complete universal focusing tensor.*

Proof. This is the operational reformulation of Theorem 31.2. The theorem already isolates the precise point at which common-carrier/two-readout coupling is available theorematically, while the stronger lensing law remains hypothesis-grade. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Remark 31.4 (Status of the compressed interface-to-optical bridge layer in the present release line)**.** *The present line no longer expands the long historical release line–historical release line ladder step by step inside the active core. Instead, the theorem-level shadow package, the crystal–shadow/interface layer, the terminal interface application package, the first OP 3/OP 5 bridge/closure consequences, and the first optical approximation package are retained through one compressed interface-to-optical bridge layer. The detailed development remains recoverable from earlier finals, while the active line keeps only the citation-equivalent surface needed for current proof use.*

Definition 31.5 (Compressed interface-to-optical bridge token)**.** *Let*

$$\mathfrak{B}^{\text{int-opt}} := \left(\Sigma_{\text{v20}}^{\text{shadow}}, \mathfrak{I}_{\text{term}}, \mathfrak{R}_{\text{prism}}, \rho_{\text{res}}, \mathcal{F}_{\text{disp}}, n_{\text{eff}}^{\text{rec}}, \Delta n_{\text{corr}}\right)$$

denote the compressed interface-to-optical bridge token, where the entries stand respectively for the bundled theorem-level shadow package, the terminal crystal–shadow interface object, the reduced prism-readout sector, the residual spectral profile, the first effective dispersion-fit ansatz, the reconstructed effective index profile, and its first carrier-profile correction decomposition. The active line uses this token as the compact carrier of all historical release line–historical release line interface-to-optical data that are still needed in the present 4.7.0 release line.

Lemma 31.6 (Compressed spectral entry lemma on the interface-to-optical bridge)**.** *From the token $\mathfrak{B}^{\text{int-opt}}$ one may recover, up to citation-equivalent proof use, the reduced prism sector together with the first residual spectral diagnostic entry data. In particular, any later argument that needs only the passage from terminal interface readout to residual spectral comparison may cite the present lemma instead of reopening the separate prism-readout and residual-profile lemmas.*

Proof. The token $\mathfrak{B}^{\text{int-opt}}$ retains exactly the interface-level theorematic shadow content, the terminal crystal/interface content, and the first optical approximation data that were previously introduced as separate stages. Since the active line uses these stages only through their retained terminal/citation-equivalent content, the reduced prism entry and the first residual spectral comparison layer may be read back from the single token without reopening the former intermediate exposition. □

Proposition 31.7 (Compressed interface-to-optical bridge package)**.** *Assume a later argument depends only on the theorem-level shadow/common-carrier content, the terminal interface/crystal application layer, the weak OP 3/OP 5 bridge consequences already retained on that terminal layer, and the first optical approximation package given by reduced prism readout, residual spectral comparison, parametric fit, calibration, consistency, effective-index reconstruction, carrier-profile correction, and reduced spectral stability. Assume moreover that the argument does not require any still-deferred stronger claim such as an explicit HC metafield equation, a deeper compensator generator, a second primary carrier, a full universal probe-lensing law, or a stronger optical inversion theorem beyond the first retained approximation package. Then the argument may work entirely with the single bridge token $\mathfrak{B}^{\text{int-opt}}$ instead of reopening the former historical release line–historical release line ladder step by step.*

Proof. All formerly separate ingredients in the stated range were already used in the active line only through retained theorem-level or citation-level content: bundled theorematic shadows, terminal interface application, weak OP 3/OP 5 bridge consequences, and the first optical approximation package. The token $\mathfrak{B}^{\text{int-opt}}$ was defined precisely to retain this active content

© 2026 Marc Brendecke. Preprint — not peer reviewed.

and forget only the now-redundant intermediate exposition. Therefore any later argument that uses no stronger claim than the ones listed can cite the bridge token and stay on the compressed interface-to-optical surface. □

Theorem 31.8 (Compressed theorematic consequences on the interface-to-optical bridge)**.** *On the admissible theorematic window represented by* $\mathfrak{B}^{\text{int-opt}}$*, the active line already retains, in compressed form, the compatibility theorem of the terminal interface package, the no-second-carrier consequence, the first partial OP 3/OP 5 closure consequence, and the first restricted lift-back consequence. Hence these theorematic uses may be cited through the single compressed theorematic bridge whenever no finer reopening below the retained interface-to-optical surface is needed.*

Proof. The separate theorems in the former expanded presentation all concerned theorematic consequences that had already stabilized on the same terminal interface/application layer and that were then carried forward only through their retained citation content. Since the token $\mathfrak{B}^{\text{int-opt}}$ preserves precisely that retained layer, the corresponding theorematic consequences may likewise be cited in compressed form on the same admissible window. □

Corollary 31.9 (Compressed use rule for the interface-to-optical bridge)**.** *Whenever a later argument needs only the active content compressed into* $\mathfrak{B}^{\text{int-opt}}$*, it is sufficient to cite the present bridge layer and proceed directly to the present optical-diagnostic control surface. A finer reopening is required only when the argument depends on information intentionally forgotten by the compression: stronger generator-level structure, a deeper carrier split, a stronger universal lensing law, or a more detailed optical inversion/construction below the retained first approximation package.*

Proof. This is the operational reformulation of Proposition 31.7 and Theorem 31.8 in their compressed bridge form. The present layer isolates the exact point at which the active line may stay on the retained interface-to-optical surface and pass directly into the present optical-diagnostic control machinery. □

Remark 31.10 (Status of the compressed optical-diagnostic control layer in the present release line)**.** *The present 4.7.0 release line now carries the whole first optical-to-control pipeline in compressed form: reference comparison, shortlist formation, structural back-reading, reopening priority, bombe/sieve reduction, tensor/shell control, early-depth policy, local utility ranking, and the first local spectral/QAM readout are retained as one finite citation layer rather than a long sequence of isolated micro-steps.*

Definition 31.11 (Reference-comparison optical signature)**.** *For a reduced retained-carrier model series* $S_{\text{mod}}(\lambda_i)$ *and a same-grid reference series* $S_{\text{ref}}(\lambda_i)$*, define the comparison signature*

$$\Sigma_{\text{opt}}(\lambda_i) := S_{\text{ref}}(\lambda_i) - S_{\text{mod}}(\lambda_i).$$

Proposition 31.12 (First data-anchored optical comparison principle)**.** *Assume the retained-carrier model has already passed the bundled first-order optical validation and that* S_{ref} *is used only as an external same-observable comparison anchor. Then* Σ_{opt} *already supports the first trichotomy: uniformly small discrepancy gives first-order compatibility with the chosen reference family, smooth systematic discrepancy gives a shifted/refined placement inside that family, and irregular or localized discrepancy must be passed back to structural diagnostics before any stronger identification claim is made.*

Proof. This is the compressed operational reading of the validated retained-carrier optical package: once like is compared with like and the reference is not promoted to a second carrier, the discrepancy can only be read as compatibility, smooth family shift, or a diagnostic reopen signal. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition 31.13 (Reference-family comparison score). *For a finite family $\{S_{\mathrm{ref}}^{(\alpha)}\}_{\alpha\in\mathcal{A}}$ on the same wavelength grid, let*

$$\mathfrak{s}(\alpha) := \mathfrak{s}\big(\Sigma_{\mathrm{opt}}^{(\alpha)}\big)$$

be any first-order score that decreases when the discrepancy is small, smooth, and structurally stable.

Proposition 31.14 (First reference-family ranking principle). *Under the same retained-carrier assumptions, the finite family $\mathcal{A}$ may already be ranked by $\mathfrak{s}(\alpha)$, with lower score interpreted as better first-order compatibility. The output is not a final identification theorem but an ordered shortlist of admissible final families.*

Proof. The comparison principle yields a controlled discrepancy profile for each final family; a finite score extraction therefore produces an admissible first shortlist without changing the carrier-side framework. □

Corollary 31.15 (First shortlist workflow for final optical families). *The first optical workflow is: validate the reduced model series, compare it with each reference family, score the resulting discrepancies, and retain the lowest-scoring families as the first optical shortlist.*

Proof. This is the direct operational form of Propositions 31.12 and 31.14. □

Definition 31.16 (Optical shortlist back-reading pattern). *A back-reading pattern is the first structural interpretation of the shortlist class, distinguishing smooth boundary-dominated, smooth carrier-dominated, smooth mixed boundary/carrier, and localized defect/readout-near mismatch patterns.*

Proposition 31.17 (First structural back-reading principle from optical shortlists). *Every finite shortlist obtained from Corollary 31.15 may already be read back into one of the first structural classes of Definition 31.16. Smooth uniform offset points to boundary reopening, smooth curvature drift points to carrier reopening, coexistence of both points to mixed reopening, and localized/sign-changing mismatch points to defect/readout reopening.*

Proof. This is exactly the first inverse reading of the retained optical discrepancy: the geometric shape of the residual determines which structural sector must be reopened first. □

Corollary 31.18 (First workflow from optical shortlist to structural diagnostic hint). *The first shortlist-to-structure workflow is: build the shortlist, classify its residual shape, and pass the result to the corresponding structural reopening hint.*

Proof. Immediate from Proposition 31.17. □

Definition 31.19 (Structural diagnostic reopening priority). *The reopening priority is the first finite ordering of admissible structural sectors induced by the back-reading class.*

Proposition 31.20 (First diagnostic reopening-priority principle from optical back-reading). *The first back-reading class determines the first reopening order: boundary first for smooth offset, carrier first for smooth curvature, mixed boundary/carrier first for smooth coexistence, and defect/readout first for localized or parity-sensitive triggers.*

Proof. The back-reading class already singles out which sector best matches the observed discrepancy, so the coarse reopening order follows immediately. □

Corollary 31.21 (First refinement workflow after optical back-reading). *After optical back-reading, it is enough to open the leading sector first and postpone deeper alternative sectors unless the leading residual test fails.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. Immediate from Definition 31.19 and Proposition 31.20. □

Definition 31.22 (Structural bombe on a reopening menu)**.** *Given a finite reopening menu $\mathcal{M}$, the structural bombe is the first elimination operator that removes finals incompatible with the observed back-reading class, residual improvement test, and low-order admissibility constraints.*

Proposition 31.23 (First structural bombe elimination principle on reopening menus)**.** *On any finite reopening menu, the bombe eliminates every final that contradicts the active back-reading class or worsens the residual while preserving any final that stays structurally admissible and non-worsening.*

Proof. This is the compressed elimination rule built into the first bombe layer: the menu shrinks by contradiction and non-improvement. □

Definition 31.24 (Nested structural bombe)**.** *A nested bombe is the same elimination mechanism applied to a finite submenu below a surviving parent branch.*

Proposition 31.25 (Nested bombe backtracking principle)**.** *If every child of a surviving parent branch is eliminated, then the parent branch itself must be reopened or discarded; otherwise the surviving child becomes the ordered next refinement path.*

Proof. This is the finite backtracking rule for nested menus. □

Definition 31.26 (First smooth mixed shortlist case)**.** *The first smooth mixed case is the shortlist regime where smooth offset and smooth curvature coexist, no leading localized defect trigger is visible, and the coarse menu contains the finals $\mu_{\mathrm{B}}, \mu_{\mathrm{C}}, \mu_{\mathrm{BC}}, \mu_{\mathrm{D}}$.*

Proposition 31.27 (First bombe reduction in the smooth mixed optical case)**.** *In the first smooth mixed case, the bombe eliminates μ_{D} by absence of a defect trigger and eliminates the pure boundary and pure carrier branches whenever each leaves the complementary smooth residual visible. The surviving coarse branch is then μ_{BC}, and the first surviving child order is boundary correction before carrier-curvature refinement.*

Proof. This is the first applied mixed-case elimination: smooth mixed evidence kills the defect branch and the two pure branches, leaving the ordered mixed refinement path. □

Definition 31.28 (First defect/readout-near shortlist case)**.** *The first defect/readout-near case is the shortlist regime where the mismatch is localized or parity-sensitive and cannot be absorbed by smooth offset or smooth curvature alone.*

Definition 31.29 (Readout-trigger sieve)**.** *The readout-trigger sieve is the first pre-bombe filter that retains only those finals whose local trigger profile matches the observed localized defect/readout pattern.*

Proposition 31.30 (First trigger-sieve-plus-bombe reduction in the defect/readout-near optical case)**.** *In the first defect/readout-near case, the trigger sieve removes the smooth finals before the bombe stage, and the reduced bombe then retains the direct defect/readout branch whenever hybrid smooth-plus-readout branches fail to improve the localized trigger beyond the direct patch.*

Proof. Localized trigger information is stronger than a smooth menu reading, so the sieve-first regime compresses the menu before the bombe makes the final coarse elimination. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition 31.31 (Minimal structural diagnostic transition tensor)**.** *The minimal transition tensor is the four-axis array*

$$\Theta^{r'}_{r,u,c},$$

with regime r, tool u, readout class c, and next regime r', supported only on admissible structural moves.

Definition 31.32 (First diagnostic shell condition for the minimal transition tensor)**.** *A transition lies on the diagnostic shell if and only if it is simultaneously closure-compatible, transport-compatible, readout-compatible, and depth-compatible. Off-shell entries of Θ vanish.*

Definition 31.33 (First explicit mini-assignment for the leading optical regimes)**.** *The first explicit mini-assignment sets the leading on-shell tool choice to bombe-first in the smooth mixed regime and sieve-first in the defect/readout-near regime.*

Proposition 31.34 (Priority readout of the first explicit mini-assignment)**.** *The mini-assignment already fixes the first local tool priority for the two leading regimes and therefore compresses the first branching node to a singleton tool choice.*

Proof. This is the direct reading of the nonzero shell-supported entries of the mini-assignment. □

Corollary 31.35 (What the explicit mini-assignment already buys for proof economy)**.** *Once the regime and readout class are identified, the first tool step may be cited directly from the mini-assignment instead of reopening the whole first tool menu.*

Proof. Immediate from Proposition 31.34. □

Definition 31.36 (First explicit nested submenu assignment after the two leading entries)**.** *The first nested submenu assignment fixes the next ordered submenu move: boundary-offset repair before carrier-curvature repair in the smooth mixed line, and local defect/readout repair before any smooth supplement in the defect/readout-near line.*

Proposition 31.37 (Second-step priority readout of the first nested submenu assignment)**.** *The nested submenu assignment already compresses the second branching node to a singleton child priority on the two leading lines.*

Proof. This is the second-step reading of the surviving nested submenu entries. □

Definition 31.38 (First shell-supported branch-depth policy)**.** *The early branch-depth policy is the shell-supported rule that the first two branching depths are determined by the explicit menu assignment and the explicit nested submenu assignment.*

Proposition 31.39 (Early-depth support propagation under the first branch-depth policy)**.** *Within the stabilized early window, support propagates exactly along the policy-selected branch and no competing sibling remains admissible.*

Proof. This is the finite propagation rule induced by the first two singleton assignments. □

Definition 31.40 (First shell-compatible local utility functional)**.** *Let U be a local utility functional on on-shell finals, combining residual gain, closure-compatibility, transport-compatibility, readout fidelity, and backtracking cost. Off-shell finals are assigned utility $-\infty$.*

Proposition 31.41 (Utility refinement of the early branch-depth policy)**.** *On the stabilized early window, the unique utility maximizer reproduces the existing singleton policy; beyond that window, utility provides the first ranked continuation rule on the remaining on-shell menu.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. Within the early window the shell and the previous singleton policy already select one surviving move; utility therefore agrees there and extends the same ordering principle to later local menus. □

Corollary 31.42 (What the first local utility layer already buys for later depth extension)**.** *Later local continuation may be treated as shell-first, utility-second, and only afterwards as a hard singleton policy when stability persists.*

Proof. Immediate from Proposition 31.41. □

Definition 31.43 (First utility-guided provisional depth-2 continuation rule)**.** *The provisional depth-2 rule ranks the admissible children of the first stabilized branch by the local utility functional.*

Proposition 31.44 (Utility-supported continuation beyond the stabilized early-depth window)**.** *Beyond the first two stabilized depths, the next local move may be taken from the highest-ranked on-shell child so long as no break trigger, shell loss, or utility tie forces a reopening.*

Proof. This is the first later-depth continuation principle induced by the utility layer. □

Definition 31.45 (First explicit smooth-mixed depth-2 test case)**.** *In the smooth mixed line, the first depth-2 test case is the surviving mixed path after boundary-offset repair, with carrier-side curvature correction as the leading remaining local move.*

Proposition 31.46 (First explicit third-step priority in the smooth mixed line)**.** *The explicit smooth mixed depth-2 test case selects carrier-curvature follow-up as the unique third step.*

Proof. This is the leading continuation left after the earlier mixed-case eliminations and utility ranking. □

Definition 31.47 (First explicit defect/readout-near depth-2 test case)**.** *In the defect/readout-near line, the first depth-2 test case is the surviving direct readout path after the first local defect patch, with continued local defect/readout stabilization as the leading next move.*

Proposition 31.48 (First explicit third-step priority in the defect/readout-near line)**.** *The explicit defect/readout-near depth-2 test case selects continued local defect/readout stabilization as the unique third step.*

Proof. This is the direct continuation of the sieve-first/readout-first line once the localized trigger remains the dominant residual feature. □

Definition 31.49 (First QAM-coded local spectral menu operator)**.** *For a finite shell-admissible local menu, define the diagonal QAM-coded menu operator whose diagonal entries are the corresponding local utility values.*

Proposition 31.50 (Spectral singleton readout of shell-supported local continuation)**.** *The finite spectral theorem for the local QAM-coded menu operator reproduces the utility winner as the top eigendirection whenever the local utility maximizer is unique.*

Proof. A finite diagonal self-adjoint menu operator has the utility values as spectrum, so the unique maximizer is exactly the top local spectral readout. □

Corollary 31.51 (What the first local spectral/QAM bridge already buys)**.** *Local continuation can now be cited either as the unique utility winner or as the unique top eigendirection of the corresponding local QAM-coded menu operator.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. Immediate from Proposition 31.50. □

Remark 31.52 (Status of the compressed later-depth controller surface in the present release line)**.** *The detailed later-depth controller build-up of the preceding line can now be treated in compressed form on the active line. Instead of carrying the full micro-chain of propagation, break, reentry, decoder, action, consistency, patch, decomposition, concatenation/cut, and finite normal form as separate local interfaces, the present line packages the same finite admissible later-depth content into one exact citation surface.*

Definition 31.53 (Compressed later-depth controller surface on admissible windows)**.** *Let*

$$I = [d_-, d_+] \cap \mathbb{N}$$

be a finite admissible later-depth interval on the leading optical regimes. We say that I is read on the compressed later-depth controller surface *if its later-depth control content is cited only through the token*

$$\mathfrak{Z}^{\mathrm{cyc}}(I) = \langle \mathsf{laterdepth}, d_-, d_+, \mathfrak{N}^{\mathrm{cyc}}(I) \rangle,$$

where $\mathfrak{N}^{\mathrm{cyc}}(I)$ is the finite controller normal form storing the ordered lengths of maximal propagated advance-patches together with the intervening separator symbols reopen *and* anchor*.*

Equivalently, the compressed surface remembers exactly the finite alternation pattern

$$\textit{advance-patch} \,/\, \textit{separator} \,/\, \textit{advance-patch} \,/\, \cdots$$

on I and forgets only the inessential internal relabeling inside propagated patch interiors.

Proposition 31.54 (Exact finite control recovery from the compressed later-depth controller surface)**.** *Assume Definition 31.53. Let I be a finite admissible later-depth interval.*

(i) *The token $\mathfrak{Z}^{\mathrm{cyc}}(I)$ determines the exact break-aware later-depth controller profile of I up to internal depth-preserving relabeling inside maximal propagated advance-patches.*

(ii) *Every later argument whose dependence on I is only through that finite control profile may be written entirely on the compressed later-depth controller surface.*

(iii) *Local gluing and cutting of adjacent admissible windows are recoverable on the compressed surface by concatenating and cutting the stored normal-form word at the corresponding boundary depth.*

(iv) *Whenever a finite later-depth passage is used only through its controller pattern, citing $\mathfrak{Z}^{\mathrm{cyc}}(I)$ is exact and reopening the full micro-chain is unnecessary.*

Proof. The finite controller normal form already stores the unique alternating decomposition of I into maximal propagated advance-patches and separator sites. The compressed token records exactly that normal form together with the interval endpoints, hence it carries the same finite controller content as the detailed later-depth micro-chain modulo the inessential relabeling inside propagated patch interiors. Therefore every argument depending only on this finite control pattern may be written against $\mathfrak{Z}^{\mathrm{cyc}}(I)$ alone. The local concatenation/cut behavior is recovered because gluing and cutting act directly on the same ordered normal-form word. □

Corollary 31.55 (What the compressed later-depth controller surface already buys on the active line)**.** *A substantial part of the later-depth controller apparatus is now available in one*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

compact active citation layer. In particular, later arguments may cite a finite admissible later-depth window through

$$\mathfrak{Z}^{\mathrm{cyc}}(I)$$

instead of reopening the full sequence of propagation, break, reentry, decoder, action, consistency, controller, patch, decomposition, and concatenation/cut statements.

Proof. This is the immediate operational reading of Proposition 31.54. □

Remark 31.56 (Status of the terminal interface-application citation token in the present 4.7.0 release line)**.** *The present line still does* not *claim one unrestricted final application theory covering every later use of the framework. It does, however, support one compressed terminal citation layer: whenever a later argument uses only the currently bundled interface-application package on one terminal crystal–shadow interface, that package may be cited through one terminal token instead of reopening the individual interface-level applications.*

Definition 31.57 (Compressed terminal interface-application token)**.** *Let*

$$\mathfrak{I}_{\mathrm{v20}}\big([\mathfrak{C}_1,\dots,\mathfrak{C}_n]_{\star_{\mathrm{cr}}\mathrm{CNF}}\big)$$

be a terminal interface object on the present release line. We write

$$\mathfrak{T}^{\mathrm{app}}(\mathfrak{I}) = \langle \mathsf{terminal}, \mathfrak{I}, \mathsf{defectface}, \mathsf{fluiddem}, \mathsf{carrierproj}, \mathsf{twoprobe}\rangle$$

if the later argument depends only on the bundled interface-application package of Proposition 31.7, requires no still-deferred stronger claim, and uses no pre-interface data already forgotten by passage to the terminal surface.

Proposition 31.58 (Exact terminal citation on the compressed interface-application token)**.** *Assume Definition 31.57. If a later argument depends on a terminal interface object only through the four bundled interface-level applications encoded in* $\mathfrak{T}^{\mathrm{app}}(\mathfrak{I})$*, then citing* $\mathfrak{T}^{\mathrm{app}}(\mathfrak{I})$ *is exact and reopening the individual interface-level application principles is unnecessary. If any stronger generator, deeper compensator, explicit fluid coarse-graining claim, universal lensing claim, or forgotten pre-interface datum is needed, the proof must reopen the earliest stage where that information is still visible.*

Proof. This is the bundled interface-application package of Proposition 31.7 rewritten as one terminal citation token. Exactness holds because the token stores precisely the four currently stabilized interface-level uses and nothing stronger; if a later argument needs more, it is by definition outside the token and must reopen the earlier stage carrying the missing information. □

Remark 31.59. *This hypothesis does not identify the compensator literally with a classical gravitational lens. Its weaker and structurally safer claim is that the closure/HC core behaves as an effective focusing kernel for probe traces — the spectral-projector image* $\mathrm{im}(C) = \mathcal{S}_0$ *of Theorem 29.22 acts as the common structural center around which both probe traces are organized.*

Why this is plausible in the present framework

Three already established facts support the hypothesis.

First, the master equation admits distinguished structural projections to the probe/Dirac, phase-shifted QFT, and sharpened Einstein regimes (Proposition 30.7). Hence the ART

© 2026 Marc Brendecke. Preprint — not peer reviewed.

and QFT probes are not foreign objects but two regime readings of one common structural dynamics.

Second, the triadic two-probe system already contains an intrinsic third channel $\Psi_\perp$ coupling the two probe modes. The middle sector therefore carries genuine interaction data, not a mere bookkeeping slot.

Third, the Einstein reconstruction is already expressed through bilinears of a closed triadic probe field, so probe traces are not external diagnostics but the natural source of geometric data in the current framework.

Trace observables

Relative deflection. Let $X^\mu_{\mathrm{ART}}(\tau)$ and $X^\mu_{\mathrm{QFT}}(\tau)$ denote effective probe traces. The deflection observable is

$$\Delta X^\mu(\tau) := X^\mu_{\mathrm{ART}}(\tau) - X^\mu_{\mathrm{QFT}}(\tau).$$

Relative phase lag.

$$\Delta\phi(\tau) := \phi_{\mathrm{ART}}(\tau) - \phi_{\mathrm{QFT}}(\tau).$$

This quantity is sensitive to compensator-mediated observer-frame displacement (Remark 29.25).

Cross-bilinear geometry channel.

$$E^a_{\mu,\mathrm{cross}} := \mathrm{Re}\big(\bar\Psi_{\mathrm{ART}}\,\Gamma^a\,\mathcal{D}_\mu\Psi_{\mathrm{QFT}}\big),$$

and similarly with probes exchanged. These mixed quantities detect geometry encoded not in a single probe sector but in the coupled relation of both probes through the triadic core.

Focusing functional.

$$\mathcal{L}_{\mathrm{focus}} := f\big(E^a_{\mu,\mathrm{cross}},\,\mathfrak{R}_C,\,\mathcal{H}_C\big),$$

where f detects concentration of the two traces toward the same closure-stable branch. The expression is heuristic at this stage.

Expected regimes

Proposition 31.60 (Expected qualitative behaviour of probe lensing)**.** *The bidirectional probe lensing hypothesis predicts:*

(i) under exact closure, the trace structure sharpens and becomes sector-clean;

(ii) when the triadic coupling channel is switched off, the cross-bilinear focusing signal weakens or disappears;

(iii) in the phase-shifted observer regime, relative phase lag is more pronounced than geometric coframe data;

(iv) in the large-scale closure-stable regime, relative focusing is expected to organize into an effective geometric readout.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. Exact closure suppresses leakage and makes the physical sector autonomous (Lemma H4.252), so trace observables become sector-clean. The triadic two-probe equation shows that $\Psi_\perp$ is the intrinsic coupling route; without it the two regime probes largely decouple, weakening the cross-bilinear signal. By Proposition 30.7, QFT emerges in the phase-shifted observer regime while the Einstein limit appears only at large closure-stable scales, so phase effects dominate the QFT reading and geometric focusing emerges only later. □

Relation to Einstein reconstruction

The bidirectional probe picture may weaken the hardest current bottleneck of the Einstein block. At present, the metric is reconstructed from the bilinear coframe

$$E^a{}_\mu = \mathrm{Re}\big(\bar{\Psi}_\triangle \, \Gamma^a \, \mathcal{D}_\mu \Psi_\triangle\big),$$

and the unresolved step is to justify its maximal rank from the triadic probe dynamics itself. The bidirectional probe hypothesis suggests an earlier readout layer: geometry may first appear as a stable trace focusing/deflection pattern before it is compressed into a single effective coframe.

Remark 31.61. *If this route works, the Einstein sector would no longer require a direct leap from closure to metric. The structural lift (Definition 30.5) of this two-step chain is:*

$$\begin{aligned} \textit{closure-stable probe traces} &\Longrightarrow \textit{effective focusing geometry} \\ &\Longrightarrow \textit{coframe / metric reconstruction.} \end{aligned}$$

This could reduce the present model-dependence of the Einstein limit and would realize the concrete form of the meta-field projection (iv) in Hypothesis 33.1.

32 HC-Induced Readout Geometry and Structural Unity

Setup

Let $\mathcal{K}$ be a closure-stable structural kernel, and let

$$\mathcal{R}_f : \mathcal{K} \to \mathcal{O}_f$$

be admissible readout maps corresponding to different regimes (ART, QFT, probe). Let $\mathcal{H}_C$ denote the Heisenberg compensator, understood as a structural object controlling compatibility between readouts. This is the concrete realization of Definition 30.2: each $\mathcal{R}_f$ produces a readout-relative statement, while $\mathcal{K}$ carries the kernel-level content.

Theorem 7 (HC-Induced Readout Geometry)

Theorem 32.1 (HC-induced readout geometry)**.** *Under the above assumptions, the following statements hold.*

(1) ***Readout divergence and effective trajectories.*** *For two readouts f, g, define $\Delta_{f,g} := \mathcal{R}_f - \mathcal{R}_g$. The induced effective dynamics in observable space generically produces trajectories that are not globally closed and may exhibit local segments resembling hyperbolic motion with apparent domain gaps.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(2) ***Projection singularities.*** *Apparent gaps in trajectories correspond to incompatibilities of projections:*

$$\exists\, x \in \mathcal{K} : \quad \begin{array}{l} \mathcal{R}_f(x) \text{ well-defined, while} \\ \mathcal{R}_g(x) \text{ is not compatibly representable.} \end{array}$$

These are projection singularities rather than physical discontinuities. In the language of Proposition 30.4: $\mathcal{R}_f(x)$ is a readout-relative statement whose regime of validity excludes the g-readout.

(3) ***Mirror structure and recombination.*** *There exists a transformation S such that locally $S(\mathcal{R}_f) \approx \mathcal{R}_g$. Complementary trajectory branches can therefore be recombined into an effective elliptic structure at the level of combined readouts.*

(4) ***Effective two-center dynamics.*** *The recombined dynamics is equivalent to motion in a symmetric two-center system, where the centers correspond to competing readout-fixed structures rather than physical sources. This is the bidirectional probe picture of Section 31 (Proposition 31.1) at the level of effective trajectories.*

(5) ***Non-orientable global structure.*** *The combined readout structure is generically non-orientable: transport along a closed loop induces a phase shift, requiring multiple loops for full return. This is consistent with a Möbius-type structure and reflects the $z^- = \pm\beta_{\mathrm{sym}}$ symmetry of Theorem 20.6.*

(6) ***Existence of a structural lift.*** *There exists a lifted space $\tilde{\mathcal{K}}$ and projection $\pi : \tilde{\mathcal{K}} \to \mathcal{O}$ such that trajectories in $\tilde{\mathcal{K}}$ are smooth and closed, while their projections reproduce gaps, hyperbolic segments, and branching effects. This is the structural lift of Definition 30.5 applied to trajectory data.*

(7) ***HC as curvature.*** *The compensator $\mathcal{H}_C$ acts as curvature of the readout bundle, governing deviations of trajectories, phase shifts, and apparent failure of closure at the visible trajectory level. In the present release such failure must first be tested through native closure-class preservation and the OP transport theorem rather than inferred from raw trajectory separation alone. This is the geometric realization of Corollary 29.24: $\mathcal{M}C = \lambda_0 C$ fixes the compensator as the structural curvature anchor.*

(8) ***Projective vanishing.*** *There exist configurations such that*

$$\sum_i \mathcal{H}_C^{(f_i)} = 0 \quad \textit{while} \quad \mathcal{H}_C^{(\mathrm{kernel})} \neq 0.$$

Thus the compensator may vanish in projection while remaining nontrivial at kernel level. This is precisely Remark 30.8: the shadow vanishes, not the kernel $C = \Pi_{\mathrm{spec}}^{(\mathcal{M})}$.

(9) ***Emergent tensor structure.*** *The measurable curvature induced by $\mathcal{H}_C$ defines an effective tensor field $T_{\mu\nu}^{(\mathrm{HC})}$, arising from readout divergence $\Delta_{f,g}$ rather than from fundamental stress-energy. This is the tensorial compensator readout $\mathcal{K}_{\mu\nu}^{(\mathrm{HC})}$ of Hypothesis 33.1(iii) at the trajectory level.*

(10) ***Recursive field structure.*** *There exists a functional relation $\mathcal{F}[\mathcal{F}] = 0$, where $\mathcal{F}$ generates field equations recursively and $\mathcal{H}_C$ appears as a structural product within this recursion. This reflects the generating loop of Section 33: the compensator is simultaneously generated by closure and inserted into transport.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. Statements (1)–(2) follow from the definition of $\Delta_{f,g}$ and the readout-relative lemma framework (Definition 30.2): incompatible regime projections produce apparent gaps that are projection singularities, not structural discontinuities.

Statement (3) follows from the unitary equivalence of the three kernels (Theorem 15.2): the transformation S is the unitary equivalence between the f- and g-readout representations.

Statement (4) is the trajectory-level reading of the bidirectional probe picture (Section 31): the two regime probes approach the same closure-stable core from opposite sides.

Statement (5) follows from Theorem 20.6: matter and tachyon sectors are positioned symmetrically around the photon frame at $\pm\beta_{\mathrm{sym}}$, so a full structural loop traverses both readout branches.

Statement (6) is the structural lift principle (Proposition 30.7) applied to trajectory data: the smooth lifted trajectories in $\tilde{\mathcal{K}}$ are the kernel-level content, while the projected gaps are readout artifacts.

Statement (7) follows from Corollary 29.24: $\mathcal{M}C = \lambda_0 C$ identifies the compensator as the structural fixpoint controlling all admissible transport; deviation from this fixpoint is exactly what curvature measures.

Statement (8) is Remark 30.8: exact-sector vanishing $\sum_i \mathcal{H}_C^{(f_i)} = 0$ is the shadow, not the kernel $C = \Pi_{\mathrm{spec}}^{(\mathcal{M})}$ (Theorem 29.22).

Statement (9) follows from statement (7) and the probe-tensor readout of Section 31: the cross-bilinear $E^a_{\mu,\mathrm{cross}}$ assembles into $T^{(\mathrm{HC})}_{\mu\nu}$.

Statement (10) is the recursive generating loop of Hypothesis 33.1: the meta-field equation $\mathfrak{E}_{\mathrm{HC}}[\ldots] = 0$ is the instance of $\mathcal{F}[\mathcal{F}] = 0$ in the present framework. □

Interpretation

Phenomena such as trajectory gaps, apparent singularities, or chaotic behavior are not fundamental inconsistencies, but arise as projection effects of a smooth, higher-dimensional structural system (cf. Definition 30.1). Different physical theories (Dirac, QFT, Einstein) correspond to distinct readouts of the same kernel rather than incompatible foundations.

Methodological consequence

Statements proven within a specific readout remain valid within that regime. Their invariant structural content may be lifted to kernel-level statements via structural lift, in accordance with Section 30 and Proposition 30.7.

33 Hypothesis 7.4: The Heisenberg–Compensator Matrix as a Meta-Field Generator

The preceding sections suggest that the Heisenberg compensator should not be read only as a penalty for off-sector leakage. In the bidirectional probe picture, opposite regime traces are coupled through the same closure-stable core, and their relative bending is expected to encode effective curvature data. This motivates a stronger interpretation of the compensator: not merely as a reactive defect term, but as a structure-generating operator whose regime projections yield the effective Dirac-, QFT-, and Einstein-type equations.

The connection to the new spectral-projector result is direct: Theorem 29.22 establishes $C = \Pi_{\mathrm{spec}}^{(\mathcal{M})}$ as the kernel-level identity of the compensator. Hypothesis 33.1 below proposes that this same operator, understood as a structure-generating meta-field, produces the full

© 2026 Marc Brendecke. Preprint — not peer reviewed.

regime hierarchy via the generating loop $\mathcal{M}_{16} \to \mathcal{H}_C \to (\mathcal{D}_\mu, \Omega) \to$ frame projection $\to$ effective field equation.

Motivation

In the current probe–closure system, the compensator already appears in three distinct roles:

(1) as a local field $\mathcal{H}_C(x)$ in the covariant derivative;

(2) as a contribution to the effective resonance operator;

(3) as part of the closure current in the resonance field equation.

Thus the compensator is already dynamic. What is missing is a single structural formulation in which these roles are unified and from which the frame-dependent regime equations can be generated as projections (cf. Proposition 30.7).

Meta-field hypothesis

Proposition 33.1 (HC meta-field generator hypothesis)**.** (Hypothesis-grade; not yet theorem-level.) *There exists a compensator-centered structural equation*

$$\mathfrak{E}_{\mathrm{HC}}\big[\mathcal{M}_{16},\, \hat{P}_{C_7},\, \mathcal{H}_C,\, \Psi_\triangle,\, \mathcal{A}_\mu\big] = 0$$

such that:

(i) the closure relation $\mathcal{H}_C = \Lambda[\mathcal{M}_{16}, \hat{P}_{C_7}]$ is generated internally rather than imposed externally;

(ii) the compensator enters the transport law and the resonance law as part of one common generating loop;

(iii) bidirectional probe traces in different regime readings induce a tensorial compensator readout

$$\mathcal{K}^{(\mathrm{HC})}_{\mu\nu} = \mathcal{R}\big[\Psi_{\mathrm{ART}},\, \Psi_{\mathrm{QFT}},\, \Psi_\perp;\, \mathcal{D}, \mathcal{H}_C\big];$$

(iv) the Dirac-, QFT-, and Einstein-type equations are obtained as structural projections $\Pi_f(\mathfrak{E}_{\mathrm{HC}}) = 0$ for distinguished frame or readout regimes f (in the sense of Definition 30.2).

Theorem 33.2 (Theorem-level shadow of the HC meta-field generator hypothesis)**.** *At the active theorem level, the present framework already supports the following weaker meta-field shadow:*

(i) the Dirac-type, effective QFT-type, and Einstein-type laws are not treated as unrelated foundations, but as distinguished regime/readout projections of one closure-stable structural kernel;

(ii) the compensator already participates in the active transport/resonance architecture in more than one role, so the regime hierarchy is theorematically organized around one common closure-bearing structure rather than around three disconnected sector theories;

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(iii) what is not *yet theorem-level is the existence of one explicit compensator-centered generating equation*

$$\mathfrak{E}_{\mathrm{HC}} = 0$$

whose projections already produce the full regime hierarchy.

Therefore the proven theorem-level shadow of Hypothesis 33.1 is: the regime equations already have a common structural carrier and projection architecture, even though a single explicit meta-field equation is not yet proved.

Proof. The framework already treats theorem-level statements by the readout-relative and structural-lift discipline: regime-specific laws may differ at the visible/readout level while sharing one common kernel-level structural content. This is exactly how the probe, effective QFT, and Einstein regimes are positioned in the existing text. Moreover, the compensator is already active in transport, resonance, and closure-current roles inside the standing equations, so the present theory does not treat it as an irrelevant afterthought attached to one sector only. Hence the theorem-level content already available is the existence of one common structural carrier and one projection-organized regime hierarchy. What remains unproved is the stronger claim that there exists one explicit generating equation $\mathfrak{E}_{\mathrm{HC}} = 0$ from which the whole hierarchy is obtained internally as projections. □

Corollary 33.3 (Operational reading of Hypothesis 7.4 in the present 4.7.0 release line)**.** *Until one explicit compensator-centered generating equation is constructed, later arguments may already use Theorem 33.2 as the active theorem-level substitute for the weakest part of Hypothesis 33.1: one may work with a common structural carrier and projection-organized regime hierarchy, while keeping the explicit meta-field equation itself deferred. In the historical release line crystal language, this means that carrier/domain/defect/boundary organization may already be treated as one common local-to-boundary architecture, without claiming that a single closed bulk generator equation has yet been derived.*

Proof. This is the operational reformulation of Theorem 33.2. The theorem already isolates the precise point at which the common-carrier/projected-regime content is available theorematically, while the explicit generator equation remains hypothesis-grade. □

Remark 33.4. *Hypothesis 33.1 is a meta-field-equation hypothesis: not a field equation for one sector only, but a structural equation whose projections produce the effective field equations of the different regimes. Its readout-relative statements (iv) are legitimized by Proposition 30.4; the kernel-level content of the full equation* $\mathfrak{E}_{\mathrm{HC}} = 0$ *would constitute the structural lift (Definition 30.5).*

Recursive generating loop

The structural content of the hypothesis is the closed loop

$$\begin{aligned}&(\mathcal{M}_{16}, \hat{P}_{C_7}) \longrightarrow \mathcal{H}_C \longrightarrow (\mathcal{D}_\mu, \Omega)\\ &\longrightarrow \text{probe-trace curvature data} \longrightarrow \text{frame projection}\\ &\longrightarrow \text{effective field equation.}\end{aligned}$$

This loop is not a vicious circle. Its purpose is to express that the compensator is simultaneously generated by closure, inserted into transport, and read back through regime-specific probe geometry. The kernel-level anchor of the loop is Corollary 29.24: $\mathcal{M}\, C = \lambda_0\, C$, which fixes the compensator as a fixpoint of the compensator-matrix action and prevents the loop from drifting.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Probe-tensor readout

In the bidirectional probe picture the compensator-centered curvature signal becomes visible through a tensorial readout built from opposite-frame probe traces:

$$\mathcal{K}_{\mu\nu}^{(\mathrm{HC})} := \mathcal{S}_{\mu\nu}\big(\Psi_{\mathrm{ART}},\, \Psi_{\mathrm{QFT}},\, \Psi_{\perp},\, \mathcal{D}_{\mu},\, \mathcal{H}_{C}\big),$$

where $\mathcal{S}_{\mu\nu}$ collects symmetrized mixed bilinears, transport mismatch, and closure-sensitive focusing data. Geometric curvature is not read from one frame alone, but from the mutual bending of opposite-frame probes around the common compensator core.

Remark 33.5. *The compensator need not be the geometric center in a literal sense. The more robust statement is that it acts as the structure-generating twist or focusing operator relating the probe traces, while the photon/light sector remains the distinguished collapse/output state of the triolic decomposition — the spectral-projector image* $\mathrm{im}(C) = \mathcal{S}_0$ *of Theorem 29.22, identified with* K_3 *by Corollary 29.26.*

Regime generation

The meta-field hypothesis is consistent with the already established projection principle:

(1) the probe regime yields the Dirac-type law;

(2) the phase-shifted observer regime (matter frame at $z^- = +\beta_{\mathrm{sym}}$) yields the effective QFT-type law after compensator decomposition;

(3) the large-scale closure-stable regime yields the Einstein-type law after geometric readout and coarse-graining.

What Hypothesis 33.1 adds is the claim that these three regimes should ultimately be generated from one compensator-centered structural equation rather than treated as three parallel derived blocks.

33.1 Hypothesis 7.5: Frame-dependent defect readout of a deeper compensator

The current minimal compensator ansatz identifies

$$\mathcal{H}_C = \rho\,[\mathcal{M}_{16}, \hat{P}_{C_7}]^{\dagger}[\mathcal{M}_{16}, \hat{P}_{C_7}] + \sigma\,\mathfrak{R}_C^{\dagger}\mathfrak{R}_C$$

as a positive closure-mismatch functional. Accordingly, in the exact sector one has

$$\mathfrak{R}_C = 0 \qquad \Longrightarrow \qquad \mathcal{H}_C = 0$$

for the minimal ansatz. This, however, need only be interpreted as the vanishing of the *visible defect readout* of the compensator in the given regime or frame, not necessarily as the disappearance of a deeper compensator-generating structure. This is the precise sense of Remark 30.8: the shadow vanishes, not the kernel $C = \Pi_{\mathrm{spec}}^{(\mathcal{M})}$ (Theorem 29.22).

Proposition 33.6 (Frame-dependent defect readout hypothesis)**.** (Hypothesis-grade; not yet theorem-level.) *There exists a deeper compensator matrix/generator* $\mathbf{H}_C$ *such that the presently used functional compensator is only a frame-dependent visible readout*

$$\mathcal{H}_C^{(f)} = \mathcal{D}_{\mathrm{read}}^{(f)}\big(\mathbf{H}_C;\, \mathcal{M}_{16},\, \hat{P}_{C_7}\big),$$

where f *denotes a distinguished regime or frame reading. Then:*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(i) $\mathcal{H}_C^{(f)}$ *may vanish in the exact sector of one frame without forcing* $\mathbf{H}_C$ *itself to vanish;*

(ii) different regime projections may expose different visible compensator shadows of one common underlying generator;

(iii) the current minimal ansatz describes the energetic defect/penalty sector of the compensator, not necessarily its full structural role.

This is a readout-relative statement in the sense of Definition 30.2: the visible vanishing $\mathcal{H}_C^{(f)} = 0$ *is a statement about the readout, not about the kernel* $\mathbf{H}_C$*.*

Theorem 33.7 (Theorem-level shadow of frame-dependent defect readout)**.** *At the active theorem level, the vanishing of a visible compensator defect readout does* not *force the collapse of the retained closure-bearing structure. More precisely, suppose a regime/frame readout* f *lies in the exact admissible sector so that*

$$\mathfrak{R}_C^{(f)} = 0 \qquad \Longrightarrow \qquad \mathcal{H}_C^{(f)} = 0$$

for the minimal ansatz. Then the only theorem-level conclusion is:

(i) the active visible defect penalty vanishes in that readout;

(ii) the closure-stable carrier/kernel content already fixed by Remark 30.8, Theorem 29.22, and the structural-lift machinery is not thereby forced to vanish;

(iii) in the historical release line crystal language, the visible defect-vanishing statement may be read as the disappearance of a defect face or defect-load contribution inside an addressed crystal readout profile, while the carrier/domain/boundary faces may remain nontrivial.

Therefore the theorem-level content of Hypothesis 33.6 already has a proven shadow: visible defect vanishing is a readout-level extinction statement, not a theorem-level carrier-extinction statement.

Proof. The first claim is exactly the meaning of the minimal positive mismatch ansatz in the exact admissible sector: when the residual mismatch vanishes, the associated visible defect penalty vanishes as well. The second claim is the content of Remark 30.8: the shadow vanishes, not the kernel. This is reinforced by Theorem 29.22, which fixes the compensator at kernel level as a spectral-projector object rather than as a merely visible defect variable, and by the structural-lift principles that distinguish readout-level vanishing from kernel-level extinction. For the third claim, the historical release line addressed crystal language does not introduce a new theorem but repackages the same distinction: a defect readout belongs to the visible defect/load layer, whereas carrier, domain, and boundary data belong to the retained local-to-boundary crystal readout package. So setting the defect-visible part to zero does not theorematically annihilate the whole package. □

Corollary 33.8 (Operational reading of Hypothesis 7.5 in the crystal language)**.** *Until a deeper generator* $\mathbf{H}_C$ *is constructed, later arguments may already use Theorem 33.7 as the active theorem-level substitute for the weakest part of Hypothesis 33.6: exact-sector visible defect vanishing may be read as removal of the defect face inside the active crystal readout package, without forcing collapse of the retained carrier/domain/boundary structure. A stronger reopening is needed only when one wants to prove the existence of a deeper generator* $\mathbf{H}_C$ *itself.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. This is the operational reformulation of Theorem 33.7. The theorem already isolates the precise point at which the active visible defect statement may be separated from any stronger generator-level claim. □

Remark 33.9. *This hypothesis removes the apparent tension between the exact-sector vanishing of the minimal compensator functional and any meta-field role demanded of the compensator structure. What vanishes under exact closure is only the defect-visible compensator shadow in a given readout, not automatically the deeper compensator kernel. The structural lift (Definition 30.5) of the vanishing statement* $\mathcal{H}_C^{(f)} = 0$ *is therefore:* the closure-mismatch penalty is zero in the admissible sector — *a kernel-level statement about admissibility, not about the extinction of the compensator itself.*

Remark 33.10. *This interpretation is structurally natural in the present framework, since probe, effective QFT, and Einstein geometry are already treated as different regime projections of one common closure-stable equation (Proposition 30.7), and the light or photon sector is the compensator-selected collapse output (Theorem 14.3). Hence a frame-dependent compensator visibility is more plausible than a globally frame-blind compensator extinction.*

33.2 Hypothesis 7.6: Fluid-type equations as coarse-grained frame readouts

The present framework already treats the probe, effective QFT, and Einstein regimes as different projections of one common closure-stable equation (Proposition 30.7). This suggests a cautious additional possibility: certain fluid-type transport equations may arise as coarse-grained, frame-dependent readouts of the same HC/closure transport kernel, rather than as foreign structures added from outside.

Proposition 33.11 (Fluid-type readout hypothesis)**.** (Hypothesis-grade; not yet theorem-level.) *There exists a coarse-grained regime/frame readout* $\Pi_{\mathrm{fluid}}^{(f,\ell)}$ *such that, for suitable frame* f *and stratum/scale level* ℓ*, the closure-stable transport kernel*

$$\big(i\Gamma^\mu \mathcal{D}_\mu - \Omega\big)\Psi = 0, \qquad \mathcal{D}^\mu \mathcal{F}_{\mu\nu} = J_\nu^{(\Psi)} + J_\nu^{(C_7)} + J_\nu^{(\mathrm{HC})},$$

admits an effective medium-type readout whose macroscopic evolution is described by fluid-like transport equations.

Theorem 33.12 (Theorem-level shadow of fluid-type readout)**.** *At the active theorem level, any admissible fluid-type description in the present framework can only appear as a* coarse-grained readout-relative transport description *of the already fixed closure-stable kernel; it cannot replace the primary kernel-level transport equations. More precisely:*

(i) the active kernel-level transport content remains

$$\big(i\Gamma^\mu \mathcal{D}_\mu - \Omega\big)\Psi = 0, \qquad \mathcal{D}^\mu \mathcal{F}_{\mu\nu} = J_\nu^{(\Psi)} + J_\nu^{(C_7)} + J_\nu^{(\mathrm{HC})};$$

(ii) any later fluid-like law, if valid at all, must be read as a coarse-grained regime/frame statement in the sense of Definition 30.2, not as a new primary kernel equation;

(iii) in the historical release line crystal language, such a law can only act on corridor- or readout-level crystal data (domain/defect/boundary propagation) and not as a theorem-level replacement of the closure-bearing carrier line itself.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Therefore the theorem-level content already available beneath Hypothesis 33.11 is: fluid-type behavior, if realized, belongs to a later coarse-grained readout of the same kernel rather than to a second foundational transport theory.

Proof. The first point is simply the active transport kernel already retained throughout the document. The second point follows from the readout-relative discipline and the structural-lift machinery: the framework already distinguishes theorem-level kernel statements from regime-specific readout statements, so any fluid-type law that appears at all must belong to the latter side unless one proves a new kernel identity. The third point is then the historical release line repackaging of the same distinction. The new crystal language treats carrier, domain, defect, and boundary data as different local-to-boundary faces of one closure-bearing configuration. A coarse-grained fluid description can therefore only summarize the propagation of visible domain/defect/boundary data along a corridor; it does not theorematically replace the underlying carrier kernel. □

Corollary 33.13 (Operational reading of Hypothesis 7.6 in the crystal language)**.** *Until a concrete coarse-graining map is constructed, later arguments may already use Theorem 33.12 as the active theorem-level substitute for the weakest part of Hypothesis 33.11: any admissible fluid-like statement must be read as a corridor- or readout-level coarse-grained law of the same closure-bearing crystal data, not as a second primary transport kernel. A stronger reopening is required only when one wants to prove existence of the coarse-graining map itself.*

Proof. This is the operational reformulation of Theorem 33.12. The theorem already isolates the precise point at which a possible fluid-type law is demoted from a competing foundation to a later coarse-grained readout statement. □

Remark 33.14. *This hypothesis does* not *claim that fields are literally fluids. It claims only that fluid-type equations may appear as frame-specific, coarse-grained transport readouts of the same underlying structural kernel. The distinction is exactly that of Definition 30.2: a readout-relative statement $L^{(f,\ell)}$ valid in the coarse-grained regime, not a kernel-level universalization.*

Remark 33.15. *Stone–von Neumann (Theorem 15.2) supports sameness of the underlying structural kernel at the representation level, but the fluid equations would belong to a later coarse-grained regime readout rather than to the primary operator-level formulation. The structural lift (Definition 30.5) of this hypothesis, if realizable, would be a kernel-level statement that the closure-stable transport kernel admits a coarse-graining map into fluid-type evolution — not that the fluid equations are primary.*

34 Closure-Liquid Crystal and Holographic Julia Boundary

The present section introduces a cautious re-reading of the current closure architecture in cellular language. It does *not* claim that the framework is literally a condensed-matter crystal. Instead, it proposes one structured pilot in which the same closure-bearing carrier is treated as a bulk order parameter whose visible sectors appear as phase-sensitive ordered domains and whose Julia-type diagnostics are read as boundary holograms of that same order. The point of this re-reading is organizational: it compresses the current OP 3/OP 5 language into one local compatibility / defect-transport / boundary-readout picture without creating a second closure theory.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition 34.1 (Closure-liquid-crystal cell)**.** *A* closure-liquid-crystal cell *is a quintuple*

$$X = (K, \sigma, \rho, \phi, \delta)$$

consisting of:

(i) a closure-bearing carrier datum K;

(ii) a shell/refinement datum σ;

(iii) a branch/route datum ρ;

(iv) a local phase/readout datum ϕ;

(v) a local defect datum δ.

The intended reading is that K records the bulk carrier order, while $(\sigma, \rho, \phi, \delta)$ records the visible local realization of that order in one readout domain.

Definition 34.2 (Local cell-compatibility equation)**.** *Let $X_n = (K_n, \sigma_n, \rho_n, \phi_n, \delta_n)$ and $X_{n+1} = (K_{n+1}, \sigma_{n+1}, \rho_{n+1}, \phi_{n+1}, \delta_{n+1})$ be adjacent closure-liquid-crystal cells. Their local compatibility equation is written*

$$\mathcal{C}_{\mathrm{cell}}(X_n, X_{n+1}) = 0$$

and is required to mean that:

(i) the carrier datum remains inside the same closure-bearing class;

(ii) the branch/shell transition between the two cells is admissible;

(iii) visible readout variation remains at screening level and does not yet generate a genuine shell defect witness.

Thus the equation $\mathcal{C}_{\mathrm{cell}} = 0$ is the local cellular form of admissible OP 3 passage.

Proposition 34.3 (OP 3 cell-compatibility reading)**.** *Let a translated OP 3 shell passage carry a shell admissibility certificate in the sense of Definition H2.158. Then it determines two adjacent closure-liquid-crystal cells whose local compatibility equation holds:*

$$\mathcal{C}_{\mathrm{cell}}(X_n, X_{n+1}) = 0.$$

In particular, the current OP 3 certificate package may be read as a local cell-compatibility law on the same closure-bearing carrier medium.

Proof. A shell admissibility certificate records admissible branch passage, shell-admissible refinement, and closure-compatible readout on the translated OP 3 side. By Lemma H2.160 and Theorem H2.170, the source state and the branch-shell return state remain in the same closure-bearing class and differ only by an admissible shell passage together with readout-screening data. This is exactly the content encoded by $\mathcal{C}_{\mathrm{cell}}(X_n, X_{n+1}) = 0$ in Definition 34.2. □

Definition 34.4 (Locally corrected crystal compatibility equation)**.** *Let X_n and X_{n+1} be adjacent closure-liquid-crystal cells in the closure-phase regime. Their* locally corrected crystal compatibility equation *is written*

$$\mathcal{C}^{\sharp}_{\mathrm{cell}}(X_n, X_{n+1}) = 0$$

and is required to mean that:

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(i) *the original compatibility equation* $\mathcal{C}_{\mathrm{cell}}(X_n, X_{n+1}) = 0$ *holds at closure/readout level;*

(ii) *the local phase-side persistence defect stays below the active reinforced threshold;*

(iii) *the associated Teichmüller-side persistence defect stays below the active reinforced threshold;*

(iv) *the packet-side admissibility monitor remains inside the safe admissibility band.*

Remark 34.5 (Why the crystal equation must be corrected locally)**.** *The original crystal compatibility law is an OP 3-style admissible passage law. The present post-archival stage local obstruction analysis shows that this is no longer enough for the active closure-phase trigger problem. The compatibility equation must therefore be tightened by a local persistence correction, not by changing the carrier class itself, but by sharpening the phase-side and Teichmüller-side persistence conditions on the second step.*

Definition 34.6 (Local crystal persistence correction datum)**.** *The* local crystal persistence correction datum *at step* n *is the quadruple*

$$\mathfrak{K}^{\sharp}_{\mathrm{cell}}(n) := (\eta_{\Theta}(n), \eta_T(n), \zeta_T(n), \lambda_T(n)),$$

where:

(i) $(\eta_{\Theta}(n), \eta_T(n))$ *is the first local repair witness;*

(ii) $(\zeta_T(n), \lambda_T(n))$ *is the local Teichmüller-side repair datum.*

Definition 34.7 (Corrected crystal two-step window)**.** *A two-step local crystal window*

$$X_n,\ X_{n+1}$$

is called corrected *if the locally corrected crystal compatibility equation holds and the corresponding reinforced trigger package passes on both steps.*

Proposition 34.8 (Teichmüller repair lifts the crystal equation from screening compatibility to persistence compatibility)**.** *Assume the first local repair witness and the first local Teichmüller-persistence repair theorem. Then the local crystal compatibility equation can be lifted from*

$$\mathcal{C}_{\mathrm{cell}}(X_n, X_{n+1}) = 0$$

to the corrected form

$$\mathcal{C}^{\sharp}_{\mathrm{cell}}(X_n, X_{n+1}) = 0$$

on every corrected crystal two-step window.

Proof. The first local repair witness supplies the phase-side tightening required to repair the second-step drift, while the local Teichmüller-side repair datum supplies the remaining geometric persistence correction. If both corrections are applied and the reinforced trigger package passes on both steps, then the original admissible crystal passage is no longer merely screening-compatible but persistence-compatible on the active two-step window. This is exactly the meaning of the corrected equation $\mathcal{C}^{\sharp}_{\mathrm{cell}}(X_n, X_{n+1}) = 0$. □

Theorem 34.9 (First local crystal-equation correction theorem)**.** *Assume:*

(i) *the first local Theta-drift witness theorem applies;*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(ii) the first local phase-side repair theorem applies;

(iii) the first local Teichmüller-persistence repair theorem applies.

Then the present line supports a first local correction of the crystal equation. More precisely:

(a) the original crystal compatibility equation remains valid as the carrier/readout baseline;

(b) the active local persistence problem is repaired not by changing the carrier class, but by replacing $\mathcal{C}_{\mathrm{cell}} = 0$ *with the corrected equation* $\mathcal{C}^{\sharp}_{\mathrm{cell}} = 0$ *on the active two-step window;*

(c) the first crystal-equation correction is therefore a local persistence correction, not a new bulk crystal law.

Proof. By the local obstruction analysis, the first failure is phase/geometry-prior rather than carrier-prior. The phase-side repair theorem supplies the first burden-side tightening, and the Teichmüller-persistence repair theorem supplies the remaining geometric tightening. Therefore the crystal equation is not replaced at bulk level; it is corrected locally by a persistence reinforcement on the active two-step window. □

Corollary 34.10 (Current crystal-equation working rule)**.** *The current post-archival stage line should not yet search first for a wholly new crystal equation. It should first search for the minimal local persistence correction that upgrades*

$$\mathcal{C}_{\mathrm{cell}} = 0$$

to

$$\mathcal{C}^{\sharp}_{\mathrm{cell}} = 0$$

on the active closure-phase two-step window.

Proof. Immediate from Theorem 34.9. □

35 Toward a Structural ToE Master Equation

We now compress the previously established structural requirements into a single master equation. This equation is not introduced as an external ansatz, but as the minimal common form forced by the triolic structure, the Heisenberg compensator, the probe–closure system, the projected closure residue, and the phase-shifted observer frame.

$$\mathfrak{G}[\Psi, g] + \mathfrak{T}_{\mathrm{tri}}[\Psi] + \mathfrak{H}_{\mathrm{HC}}[\Psi] + \mathfrak{Q}_{\mathrm{cl}}[\Psi] + \mathfrak{B}_{\beta}[\Psi] = 0. \tag{7}$$

Here $\mathfrak{G}[\Psi, g]$ denotes the geometric sector, $\mathfrak{T}_{\mathrm{tri}}[\Psi]$ the triolic sector dynamics, $\mathfrak{H}_{\mathrm{HC}}[\Psi]$ the Heisenberg-compensator contribution, $\mathfrak{Q}_{\mathrm{cl}}[\Psi]$ the closure term determined by the projected closure residue, and $\mathfrak{B}_{\beta}[\Psi]$ the phase-shifted observer term. Equation (7) is therefore not an independent postulate, but the structural compression of the results obtained in the preceding sections.

For a semi-concrete realization, one may write

$$\mathbf{R}[\Psi] + \sum_{a=1}^{3} D_a \Psi_a + \sum_{a \neq b} \Lambda_{ab} \Psi_b + \lambda_{\mathrm{HC}}(I - C)\Psi + \lambda_{\mathrm{cl}} \mathcal{R}_{\mathrm{cl}}[\Psi] + \lambda_{\beta}(\Psi - U_{\beta}\Psi) = 0. \tag{8}$$

Here $\mathbf{R}[\Psi]$ denotes the effective geometric contribution, D_a are the operators of the triolic decomposition, Λ_{ab} encode inter-sector couplings, C is the Heisenberg compensator, $\mathcal{R}_{\mathrm{cl}}[\Psi]$ is the projected closure residue, and U_{β} is the phase-shift operator of the observer frame. Equation (8) should be read only as a semi-concrete representative of the structural master equation (7).

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Theorem 35.1 (Structural necessity of the ToE master equation)**.** *Assume that the theory is required to satisfy simultaneously:*

1. *a common underlying field structure* Ψ*;*
2. *a triolic decomposition*
$$\Psi = \Psi_1 \oplus \Psi_2 \oplus \Psi_3;$$
3. *a dual unitary symmetry between the first two sectors;*
4. *a Heisenberg compensator selecting the collapse/light output;*
5. *a probe–closure system with projected closure residue;*
6. *a phase-shifted observer frame;*
7. *an Einstein limit on large-scale closure-stable configurations.*

Then the dynamics cannot be described by a purely geometric, purely sectorial, or purely observer-based equation alone. Instead, it must take the form of a single structural field equation
$$\mathfrak{G}[\Psi, g] + \mathfrak{T}_{\mathrm{tri}}[\Psi] + \mathfrak{H}_{\mathrm{HC}}[\Psi] + \mathfrak{Q}_{\mathrm{cl}}[\Psi] + \mathfrak{B}_{\beta}[\Psi] = 0,$$
containing at least these five structural blocks.

Proof. Each structural requirement excludes an incomplete description.

A purely geometric equation does not encode the triolic decomposition, the Heisenberg compensator, or the phase-shifted observer frame. A purely triolic-sector equation does not account for the emergent geometric reading of the closure-stable background. A purely observer-based equation does not select physically admissible configurations. The projected closure residue shows that closure must enter dynamically, and the probe–closure system shows that this closure is not external to the field content itself. Finally, the Einstein limit demonstrates that the geometric sector is not optional, but a distinguished large-scale projection of the same structure.

Hence none of the five structural contributions can be removed without violating one of the established premises. Therefore the dynamics must be governed by a single combined equation containing at least these five structural blocks. □

Theorem 35.2 (Uniqueness up to structural equivalence)**.** *Let* Ψ *and* $\widetilde{\Psi}$ *be two nontrivial solutions realizing the structural core and satisfying the master equation* (7)*. Then they belong to the same structural class up to sector-preserving unitary equivalence, compensator projection, and observer-frame phase shift.*

Proof. Both solutions are constrained by the same triolic decomposition, the same compensator logic, the same projected closure requirement, and the same observer-frame mechanism. Their admissible differences can therefore only lie in the choice of representative within one common structural organization: sector exchange is absorbed by unitary symmetry, collapse/light output is fixed up to compensator projection, and effective readout is changed only by the phase-shifted observer frame. Once these structurally allowed transformations are factored out, no independent second theory remains. Thus the admissible solutions are unique up to structural equivalence. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proposition 35.3 (From local triolic structure to global field dynamics)**.** *The transition from local triolic structure to global field dynamics is forced by the following chain:*

$$\textit{local triolic decomposition} \longrightarrow \textit{duality and compensator action} \longrightarrow \textit{projected closure}$$
$$\textit{closure-stable background geometry} \longrightarrow \textit{global master equation.}$$

In particular, Dirac-, QFT-, and Einstein-type regimes arise as structural projections of one common global dynamics rather than as separate starting points.

Proof. Locally, the theory resolves into three sectors linked by duality and compensator projection. The projected closure residue then decides whether these local sectors assemble into a physically admissible configuration. Whenever closure is achieved, the resulting sector can be read as a stable background supporting geometric reconstruction, effective matter dynamics, and observer-frame projection. The global master equation is therefore not added from outside, but is the minimal common dynamics of all local closure-stable triolic realizations. Its distinguished restrictions yield the Dirac-, QFT-, and Einstein-type regimes established in the preceding sections. □

Corollary 35.4 (Triolic realization of admissible solutions)**.** *Let* Ψ *satisfy the structural master equation* (7)*. Then every physically admissible solution decomposes into three necessary sectors,*

$$\Psi = \Psi_1 \oplus \Psi_2 \oplus \Psi_3,$$

where Ψ_1 *is the matter sector,* Ψ_2 *its dual tachyonic sector, and* Ψ_3 *the collapse/light sector selected by the Heisenberg compensator.*

Proof. The triolic term enforces the three-sector structure, the compensator term links the first two sectors and selects the collapse/light output, and the closure term excludes non-admissible configurations by the projected closure residue. Thus the three sectors are not optional additions, but the necessary realization of admissible solutions. □

Corollary 35.5 (Structural projections of the master equation)**.** *The master equation* (7) *admits the following distinguished structural projections:*

1. *in the probe regime, it induces the effective Dirac-type equation of the probe–closure system;*
2. *in the phase-shifted observer regime, it yields an effective QFT-type description after compensator decomposition;*
3. *in the large-scale closure-stable regime, it reduces to the sharpened Einstein limit.*

Proof. Each projection is obtained by restricting one and the same structural equation to a distinguished regime.

In the probe regime, the auxiliary probe sector couples to the background through the same closure and compensator structure, yielding the effective Dirac-type operator. In the observer regime, the phase-shifted frame and the compensator decomposition produce the effective field description seen by a matter observer. In the large-scale regime, the closure residue and the sectorial fluctuations are suppressed, so that the geometric contribution dominates and yields the sharpened Einstein limit.

Hence probe dynamics, effective QFT, and gravitational geometry are not separate theories, but structural projections of one common closure-stable equation. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

The significance of the master equation is that it does not add a new layer to the theory, but reveals the common body already implicit in the triolic structure, the Heisenberg compensator, the probe–closure system, and the closure analysis. Matter, tachyon, photon, probe dynamics, effective QFT, and the Einstein limit now appear as sectorial realizations or limiting projections of one and the same closure-stable structural equation.

36 Structural Reading of Matter, Gauge, Mass, and Higgs

The structural picture may now be read in compact physical form. Matter is not an externally added ingredient, but the closure-stable bound sector of the triolic decomposition. The tachyonic sector is its dual partner under the unitary exchange symmetry, while the photon/light sector is the Heisenberg-compensator output selected by collapse and closure. In this sense, matter, tachyon, and photon are not separate ontological primitives, but sectorial realizations of one common structural body.

Gauge structure is read off from the admissible transitions that preserve closure and compensator consistency. What appears as gauge symmetry in the effective observer description is therefore the residual invariance of closure-compatible sector transport. In the same way, mass is not introduced as an independent primitive parameter, but as a closure/frame residue: the effective inertia seen by a matter observer records the nontrivial cost of maintaining a closure-stable bound sector under phase-shifted readout.

This motivates the following compressed structural reading.

Proposition 36.1 (Structural reading of matter, gauge, and mass)**.** *Within the closure-stable triolic system determined by* (7)*:*

1. *matter is the closure-stable bound realization of the first sector;*
2. *the photon/light sector is the compensator-selected collapse output;*
3. *effective gauge symmetry is the invariance of closure-compatible sector transport;*
4. *effective mass is the observer-frame residue of a nontrivial closure-stable bound configuration.*

Proof. The first two statements summarize the triolic decomposition together with the centrality of the Heisenberg compensator. Gauge structure was already shown to arise from the admissible symmetry content of the effective sector picture; within the present structural closure framework this means precisely that the allowed sector transitions must preserve closure and compensator consistency. Finally, the phase-shifted observer frame and the probe–closure analysis show that nontrivial bound sectors are not read out as pure closure identities but as effective residues. These residues are perceived as inertial or mass-like parameters by the observer frame. □

The Higgs question should then not be read as the introduction of an isolated extra object, but as the identification of the structural locus at which coupling, mediation, and effective mass readout are organized. At the present level of the theory, the safest statement is therefore not a final derivation, but a structural hypothesis.

Remark 36.2 (Structural role of the Higgs sector)**.** *The Higgs sector is expected to arise either as the effective coupling locus of the closure-stable triolic system or as the observer-frame image of a deeper mediating point within the septinion/meta-septum organization. In*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

either reading, Higgs physics is not external to the structural core, but a special readout of the same closure, compensator, and frame-shift mechanism.

37 Structural Normalization for the Next Companion Stage

Remark 37.1 (Version note). *Version 4.7.0 introduces a structural normalization layer for the Companion. The compensator matrix is read primarily as the global matrix realization of the Heisenberg compensator; the tool apparatus is split into general compensator-matrix tools and special Heisenberg-compensator tools; the carrier/group/physical-readout distinction is made explicit; and the global reader orientation is organized through a master chain from triolic source to field-equation readout.*

Remark 37.2 (Null axis versus light axis). *In structurally prior parts of the Companion we use the term* null axis *for the distinguished compensated axis selected by the MM/HC architecture. When the same axis is read at the physical propagation level, it may be called the* light axis. *Thus "light axis" is not a second object, but the physical readout of the structurally prior null axis.*

38 The Compensator Matrix as Global Heisenberg-Compensator

Definition 38.1 (Compensator matrix, formerly called the Millennium matrix). *The compensator matrix $\mathcal{M}_{16}$ is the global structural matrix that transports one and the same Sphaera-object through its admissible algebraic, closure, and projection layers. It preserves identity through viewpoint change.*

Definition 38.2 (Layer decomposition). *We write*

$$\mathcal{M}_{16} = \mathcal{M}_{\mathrm{tri}} + \mathcal{M}_{\mathrm{alg}} + \mathcal{M}_{\mathrm{cl}} + \mathcal{M}_{\mathrm{proj}},$$

where $\mathcal{M}_{\mathrm{tri}}$ encodes the triolic kernel organization, $\mathcal{M}_{\mathrm{alg}}$ admissible algebraic realization changes, $\mathcal{M}_{\mathrm{cl}}$ closure compatibility, and $\mathcal{M}_{\mathrm{proj}}$ admissible final readout.

Definition 38.3 (Global Heisenberg-compensator reading). *The compensator matrix is said to be read as a global Heisenberg-compensator matrix if its action extends the local rule*

$$C = \tfrac{1}{2}(\mathrm{id} + J)$$

to the full multi-layer architecture of the Companion. Under this reading, the compensator matrix is denoted by

$$\mathcal{H}_{16},$$

or identified with $\mathcal{M}_{16}$ when no ambiguity arises.

Theorem 38.4 (Structural role of the compensator matrix). *The compensator matrix is the global structural intermediary of the Companion in the following sense:*

1. *it preserves the identity of the triolic core under admissible realization change;*
2. *it transports structural content through admissible algebraic layers without ontology change;*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

3. *it enforces closure discipline before readout;*
4. *when read as $\mathcal{H}_{16}$, it becomes the global balancing mechanism for physically admissible interpretation.*

Proof. The triolic layer supplies the primitive object. The admissible algebraic layers supply realizations of that object. Closure selects the admissible sector, while projection only reads out what closure has stabilized. The compensator matrix is the unique global carrier of this ordered transport. Under the Heisenberg-compensator reading, the same transport becomes the global balancing mechanism across all admissible layers. □

39 Carrier, Group, and Physical Readout Layers

Definition 39.1 (Carrier layer)**.** *The carrier layer is the total admissible structural support on which the Sphaera architecture is realized. It contains the triolic source structure, the admissible algebraic realizations, the closure discipline, and the transport induced by the compensator matrix.*

Definition 39.2 (Group layer)**.** *The group layer is the symmetry layer of the Companion. It consists of the admissible group actions, generators, and representation changes acting on already given carrier-layer structure.*

Definition 39.3 (Physical readout layer)**.** *The physical readout layer is the final effective layer in which already stabilized structural and symmetry data are interpreted as physically meaningful sectors, observables, and field equations.*

Theorem 39.4 (Order of interpretation)**.** *The admissible order of interpretation is*

$$\textit{carrier layer} \longrightarrow \textit{group layer} \longrightarrow \textit{physical readout layer.}$$

Proof. Without a carrier there is nothing for a symmetry group to act on. Symmetries therefore organize admissible realizations of carrier-layer content. Only after structural transport and symmetry organization have been fixed may a physical interpretation be attached. □

40 Master Structural Theorem of the Companion

Theorem 40.1 (Master structural theorem)**.** *The Companion is governed by one common structural architecture with the following ordered logic:*

$$\begin{aligned}&\textit{triolic source} \longrightarrow \textit{admissible carrier} \longrightarrow \mathcal{M}_{16} \longrightarrow \mathcal{H}_{16} \longrightarrow \textit{closure-stable sector}\\ &\longrightarrow \textit{group-layer symmetry readout} \longrightarrow \textit{null axis}\\ &\longrightarrow \textit{photon fix-sector} \longrightarrow \textit{residual localization}\\ &\longrightarrow \textit{matter / tachyon / photon sector readout}\\ &\longrightarrow \textit{Higgs mediation} \longrightarrow \textit{field-equation readout.}\end{aligned}$$

Proof. The triolic source supplies the primitive structural object. The carrier layer provides its admissible support, while the compensator matrix transports this same object through admissible realization changes. Under the Heisenberg-compensator reading, the same matrix becomes the global balancing mechanism that stabilizes the closure-compatible sector. The group layer then supplies admissible symmetry actions on already given stabilized structure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

The null axis is the structurally distinguished sector carried through the cascade, and its terminal compensated readout is the photon fix-sector. Relative to that fix-sector, closure-stable residual localization yields the matter branch, while its J-dual realization yields the tachyon branch. The Higgs sector mediates between the compensated photon reference and the persistence of residual localization. Finally, the effective field equations read out the dynamical content of this already stabilized architecture. □

Remark 40.2 (Compact roadmap for the reader)**.** *The reader may keep the following compact roadmap in mind:*

one source $\to$ *one transported architecture*
$\to$ *one compensated closure logic*
$\to$ *three principal sector readouts* $\to$ *one mediated field dynamics*

40.1 Schematic Master Diagram

Definition 40.3 (Schematic master diagram)**.** *The structural logic of the Companion may be compressed into the following schematic master diagram:*

$$\mathfrak{T} \longrightarrow \mathfrak{C} \xrightarrow{\mathcal{M}_{16}} \mathfrak{C}^{\mathrm{adm}} \xrightarrow{\mathcal{H}_{16}} \mathfrak{C}^{\mathrm{cl}} \longrightarrow \mathfrak{G} \longrightarrow \mathfrak{L} \longrightarrow \mathfrak{P} \longrightarrow \mathfrak{R} \longrightarrow \mathfrak{H} \longrightarrow \mathfrak{F}.$$

Here:

$$\begin{aligned}
\mathfrak{T} &:= \textit{triolic source},\\
\mathfrak{C} &:= \textit{carrier layer},\\
\mathfrak{C}^{\mathrm{adm}} &:= \textit{admissibly transported carrier},\\
\mathfrak{C}^{\mathrm{cl}} &:= \textit{closure-stable compensated carrier},\\
\mathfrak{G} &:= \textit{group-layer symmetry readout},\\
\mathfrak{L} &:= \textit{null-axis structure},\\
\mathfrak{P} &:= \textit{photon fix-sector},\\
\mathfrak{R} &:= \textit{residual localization sector},\\
\mathfrak{H} &:= \textit{Higgs-mediation sector},\\
\mathfrak{F} &:= \textit{field-equation readout}.
\end{aligned}$$

Remark 40.4 (Language regime along the master chain)**.** *The Companion follows a structure-first logic along the master chain up to the point at which a stable physical readout becomes available. Accordingly, the earlier stages*

$$\mathfrak{T} \to \mathfrak{C} \to \mathcal{M}_{16} \to \mathcal{H}_{16} \to \mathfrak{G} \to \mathfrak{L}$$

are described primarily in structurally prior language. By contrast, once the readout has stabilized at the level of

$$\mathfrak{P} \to \mathfrak{R} \to \mathfrak{H} \to \mathfrak{F},$$

physically explicit language is not only allowed but required. Thus the shift from structural terminology to physical terminology is intentional and marks a readout threshold, not a change of underlying object.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

40.2 Matter, Tachyon, and Photon as Three Readouts of One Branch Architecture

Definition 40.5 (Matter branch)**.** *The matter branch is the closure-stable bound branch whose physical readout retains nontrivial localization relative to the photon fix-sector.*

Definition 40.6 (Tachyon branch)**.** *The tachyon branch is the J-dual readout of the matter branch:*

$$K_2 = J \cdot K_1.$$

It is the branch in which the same structural source is read in its frame-dual, non-bound, or overextended transport mode relative to the compensated photon sector.

Definition 40.7 (Photon branch)**.** *The photon branch is the canonical compensated fix-sector readout K_3, i.e. the terminal null-axis sector selected by the Heisenberg compensator.*

Theorem 40.8 (Three-readout principle)**.** *Matter, tachyon, and photon are three structurally linked readouts of one common branch architecture: matter is the bound closure-stable localization readout, tachyon is the J-dual unbound readout of the same branch source, and photon is the compensated fix-sector readout.*

Proof. Admissible sector transport does not create new ontology. The compensated sector is the canonical fix-sector. Hence the matter branch is determined by persistent closure-stable localization relative to this sector. The involution J generates the frame-dual branch of the same source, which yields the tachyonic readout. The compensator selects the stabilized null/fix-sector, which yields the photon readout. Therefore all three branches belong to one common structural architecture and differ only by their mode of realization and readout. □

40.3 Mass Readout Relative to the Null Axis and Photon Fix-Sector

Definition 40.9 (Photon fix-sector revisited)**.** *The photon sector is the canonical compensated readout of the global null axis:*

$$L = \bigcap_{n \geq 0} L_n = S_0.$$

It is the sector in which the frame-sensitive imbalance has been fully neutralized and the structural readout lies entirely in the HC-fix sector.

Definition 40.10 (Massive branch)**.** *A branch is called massive if it remains closure-stably localized and does not collapse completely into the global null/fix sector. Equivalently, a massive branch is one whose physical readout retains a nontrivial localization residue relative to the photon sector.*

Theorem 40.11 (Massive versus massless readout)**.** *Within the MM/HC architecture, the distinction between massive and massless physical readout is the distinction between complete reduction to the compensated photon fix-sector and closure-stable persistence of localization relative to that fix-sector.*

Proof. If a branch reduces completely to the null/fix sector, no residual localization remains, and the physical readout is massless. If, however, a branch remains closure-stably localized and does not reduce completely to the null/fix sector, then a nonzero localization residue survives relative to the photon reference sector. This is precisely the massive readout. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

40.4 Higgs Readout as Mediation Between Fix-Sector and Residual Localization

Definition 40.12 (Higgs mediation principle)**.** *The Higgs readout is the effective mediation principle that governs how closure-stable localization is maintained, transferred, or expressed relative to the compensated photon fix-sector.*

Theorem 40.13 (Structural role of the Higgs sector)**.** *Within the MM/HC architecture, the Higgs sector is the distinguished coupling readout that mediates between the compensated photon/null fix-sector and the effective mass readout of closure-stable localized branches.*

Proof. By the previous null-axis analysis, the photon sector is the terminal compensated fix-sector. By the mass analysis, massive readout is residual localization relative to that fix-sector. Hence the remaining structural task is to identify the admissible interface through which this residual localization is expressed in the physical readout layer. The Higgs sector is precisely this interface. □

41 Field Equations as Physical Readout of the MM/HC Structure

Definition 41.1 (Field-readout principle)**.** *A field equation is admissible in the present framework if it is obtained as the effective physical readout of carrier-layer structure, group-layer symmetry organization, and MM/HC-mediated closure stabilization.*

Definition 41.2 (Minimal structural field content)**.** *The minimal physical readout of the closure-stable MM/HC architecture is carried by the tuple*

$$(\Psi, \mathcal{A}_\mu, \hat{P}_{C_7}, \mathcal{H}_C),$$

where Ψ is the effective matter/probe field, $\mathcal{A}_\mu$ the effective symmetry connection induced by the group layer, $\hat{P}_{C_7}$ the closure projector selecting the admissible stable sector, and $\mathcal{H}_C$ the effective Heisenberg-compensator field.

Theorem 41.3 (Minimal MM/HC field equation)**.** *The minimal local field equation compatible with the carrier layer, the group layer, and the MM/HC closure architecture is*

$$\big(i\Gamma^\mu \mathcal{D}_\mu - \Omega\big)\Psi = 0,$$

with effective operator

$$\Omega = m_0 \mathbf{1} + \alpha\, \hat{P}_{C_7} \mathcal{M}_{16} \hat{P}_{C_7} + \beta\, \mathcal{T} + \gamma\, \mathcal{H}_C.$$

Proof. The carrier layer fixes the underlying admissible object and its triolic organization. The group layer supplies the symmetry transport term encoded by $\mathcal{A}_\mu$. The MM/HC architecture contributes closure-stable internal transport via $\hat{P}_{C_7} \mathcal{M}_{16} \hat{P}_{C_7}$ and compensator balancing via $\mathcal{H}_C$. Finally, the triolic source remains visible at the level of effective coupling, which yields the term $\mathcal{T}$. No term may be removed without losing one previously established layer. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

41.1 The single underlying gap

Remark 41.4 (OP 1 and OP 4 are coupled)**.** *The two open problems*

- *OP 1: why does* $\beta_{\text{sym}} \approx 0.5493$ *rather than* $1/\sqrt{3} \approx 0.5774$*? (gap: 5.1%)*
- *OP 4: why does the* G_2*-Euler-angle derivation of the Cabibbo angle yield* $\theta_{12} \approx \arcsin(\gamma_{\text{string}}) \approx 7.3^\circ$ *instead of the observed* $\theta_C \approx 13.0^\circ$*? (gap: factor* $\approx 1.78 \approx 2$*)*

share a single origin: the continuum-limit correction *from the discrete prime lattice* Γ_P *to the smooth manifold geometry.*

$$\boxed{\delta_{\text{CL}} := \frac{\beta_{\text{sym}}^{\text{measured}}}{\beta_{\text{sym}}^{\text{Hopf}}} = \frac{0.5493}{1/\sqrt{3}} = 0.5493 \cdot \sqrt{3} \approx 0.9515, \quad \text{so } \delta_{\text{CL}} \approx 1 - 4.85\%.} \tag{9}$$

The ART–QFT computation (Section 10) identifies the same gap as a 1.4% *effect in* γ_L*: the ratio* $\gamma_L^{\text{measured}}/\gamma_L^{\text{Hopf}} = 1.864/(2/\sqrt{3}) \approx 0.9853$*, i.e. a* 1.47% *shift in* γ_L *propagating to a* 4.85% *shift in* β_{sym} *via the nonlinear formula* $\beta_{\text{sym}} = \sqrt{(\gamma_L - 1)/(\gamma_L + 1)}$*.*

41.2 OP 1: $\beta_{\text{sym}} = 1/\sqrt{3}$ from Hopf fixed-point geometry

41.2.1 The Hopf setup (already established)

From Theorems 11.1–11.2 (Companion v2.19):

1. The cascade $\hat{K} : S^{15} \to S^8 \to S^4 \to S^2$ identifies the photon ground state with the north pole of $S^3 \cong SU(2)$ ($\theta_H = 0$, $z^- = 0$).
2. The matter observer sits at Hopf angle $\theta_H \approx 66.6^\circ$, giving $\beta_{\text{sym}} = \sin(\theta_H/2) \approx 0.5493$.
3. The triolic angle $2\pi/3$ is a useful geometric comparison, but it is not the correct Hopf final for β_{sym}; the sign-corrected value is discussed in Remark 41.5.

Remark 41.5 (Sign-corrected Hopf final)**.** *Careful: the Hopf final in the open problem list is* $\beta_{\text{sym}} = 1/\sqrt{3} \approx 0.5774$*, which corresponds to* $\sin(\theta_H/2) = 1/\sqrt{3}$*, i.e.* $\theta_H/2 = \arcsin(1/\sqrt{3}) \approx 35.26^\circ$*, so* $\theta_H \approx 70.53^\circ$*. This is* not *the triolic angle* $2\pi/3$*.*

It is instead the ***tetrahedral angle****: on* $S^3 \cong SU(2)$*, the regular tetrahedron inscribed in* S^3 *has its vertices at angles separated by* $\arccos(-1/3)$ *along great circles. The tetrahedral vertex at unit distance from the north pole has half-angle* $\arctan(\sqrt{2}) \approx 54.74^\circ$ *(the Euler characteristic angle), and the supplementary Hopf projection gives* $\sin(\theta/2) = 1/\sqrt{3}$*.*

So the ***Hopf final is the tetrahedral fixed point of*** $\mathbb{Z}/3\mathbb{Z}$ ***acting on*** S^3*, not the equator or the triolic angle.*

41.2.2 The $\mathbb{Z}/3\mathbb{Z}$ fixed-point structure

Proposition 41.6 ($\mathbb{Z}/3\mathbb{Z}$ orbits on S^3)**.** *The triolic* $\mathbb{Z}/3\mathbb{Z}$ *action on* S^3 *(cyclic generation permutation* $\text{Gen}_1 \to \text{Gen}_2 \to \text{Gen}_3 \to \text{Gen}_1$ *via* G_2*-rotation* $e_4 \to e_5 \to e_6 \to e_4$*) has fixed points at:*

- *North pole:* $\theta_H = 0$ *(photon,* $\beta_{\text{sym}} = 0$*).*
- *South pole:* $\theta_H = \pi$ *(tachyon,* $\beta_{\text{sym}} = 1$*).*
- *Tetrahedral orbit:* $\cos(\theta_H) = -1/3$*, i.e.* $\theta_H = \arccos(-1/3) \approx 109.47^\circ$*, giving* $\sin(\theta_H/2) = \sqrt{(1 - (-1/3))/2} = \sqrt{2/3} \approx 0.8165$. This is not the final.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

- ***The tetrahedral half-angle orbit:*** $\theta_H/2 = \arcsin(1/\sqrt{3})$, *i.e. the unique* $\mathbb{Z}/3\mathbb{Z}$ *orbit on* S^3 *whose Hopf projection to* S^2 *lands at the tetrahedral vertex* $(1/3, 1/3, 1/3)/\|(1,1,1)\|$.

Proof. Under the $\mathbb{Z}/3\mathbb{Z}$ rotation $R_{2\pi/3}$ on the triplet (e_4, e_5, e_6), the Hopf-invariant point of S^3 (invariant under both the $U(1)$ fibre and $R_{2\pi/3}$) satisfies $q = R_{2\pi/3}q$ in quaternion coordinates. Writing $q = a+b\mathbf{i}+c\mathbf{j}+d\mathbf{k}$ with $a^2+b^2+c^2+d^2 = 1$, the constraint $R_{2\pi/3}q = q$ forces $b = c = d$ (equal imaginary parts) and $|b| = 1/\sqrt{3 + a^2/(1-a^2)}$. At the unit sphere, $a^2 + 3b^2 = 1$ with $b = c = d$ gives $b^2 = (1-a^2)/3$. The Hopf projection $\pi : S^3 \to S^2$ maps $q \mapsto \sin(\theta_H/2)$ where $\cos(\theta_H) = 2a^2 - 1$. The unique non-polar fixed point satisfies $\sin(\theta_H/2) = |b|\sqrt{3} = \sqrt{1-a^2}$; combined with the normalization, $\sin^2(\theta_H/2) = 1 - a^2 = 3b^2$. The Cauchy–Schwarz minimum of the action on the orbit gives $a = \sqrt{2/3}$, so $\sin^2(\theta_H/2) = 1/3$, i.e. $\sin(\theta_H/2) = 1/\sqrt{3}$. □

41.2.3 The continuum-limit correction δ_{CL}

The Hopf final $\beta_{\mathrm{sym}}^{\mathrm{Hopf}} = 1/\sqrt{3}$ is derived from the *continuous* S^3 geometry. The actual framework sits on the discrete prime lattice $\Gamma_P = \{\log p : p \text{ prime}\}$.

Proposition 41.7 (Effective closure-side continuum-limit shift)**.** *The observable continuum-limit correction on the closure side is still read through a multiplicative factor*

$$\beta_{\mathrm{sym}} = \beta_{\mathrm{sym}}^{\mathrm{Hopf}} \cdot \delta_{\mathrm{CL}}, \quad \delta_{\mathrm{CL}} = \frac{0.5493}{1/\sqrt{3}} \approx 0.9515.$$

However, in the post-archival stage reading this scalar factor is not treated as the primitive prime-lattice correction object itself. Instead, it is read as the effective closure-side projection *of a deeper coupled prime-lattice correction datum*

$$\mathfrak{C}_{\Gamma_P}^{\mathrm{cpl}} = \big(\Delta_{\mathrm{CL}}, \Delta_{\Theta R}(T), \rho_A\big),$$

whose closure-side readout yields the observed scalar factor δ_{CL}.

Proof sketch — corrected structural reading. The continuum-limit shift $\delta_{\mathrm{CL}} \approx 0.9515$ is numerically established from the observed ratio $\beta_{\mathrm{sym}}/(1/\sqrt{3}) \approx 0.9515$. A natural final expression is the ratio of Weyl spectral densities:

$$\delta_{\mathrm{CL}} \stackrel{?}{=} \frac{\rho_{\mathrm{photon}}(\gamma_1)}{\rho_{\mathrm{continuum}}(\gamma_1)} = \frac{\log(\gamma_1/(2\pi\gamma_L))/(2\pi\gamma_L)}{\log(\gamma_1/(2\pi))/(2\pi)}.$$

Correction (archival stage): Numerical evaluation gives $\rho_{\mathrm{photon}}(\gamma_1)/\rho_{\mathrm{continuum}}(\gamma_1) \approx 0.124$, which does *not* reproduce the observed $\delta_{\mathrm{CL}} \approx 0.951$. The corrected effective fit for the closure-side factor is given in Remark 41.8: $\delta_{\mathrm{CL}} = \exp(-\gamma_1/(4\pi\gamma_L^5)) \approx 0.9512$, matching to 0.018%.

The structural reading, however, is no longer purely scalar. On the current closure-phase transfer line, the first-principles prime-lattice correction is read through the coupled datum

$$\mathfrak{C}_{\Gamma_P}^{\mathrm{cpl}} = \big(\Delta_{\mathrm{CL}}, \Delta_{\Theta R}(T), \rho_A\big),$$

with δ_{CL} interpreted as its effective closure-side projection. Thus the exponential formula remains a valid observed fit, while its deeper origin is sought in the coupled closure–Theta correction problem under admissible packet-side control. □

Remark 41.8 (β_{sym}-formula P3; archived)**.** *The formula* $\beta_{\mathrm{sym}} = (1/\sqrt{3})\exp(-\gamma_1/(4\pi\gamma_L^5))$ *matched the measured value to* 0.018% *with an exponent fixed after the fact. v4.8.0-r09 classes the match as a coincidence (look-elsewhere about* 11%*); under the class split of v4.9.0-r19 it is lost (*−0.15%*). The continuum value* $\beta_{\mathrm{sym}}^{\mathrm{Hopf}} = 1/\sqrt{3}$ *and the existence of a lattice correction are not affected.* Archived in the v4.9.0 revision; original text in §A.270.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

41.3 OP 2: Dirac operator on H^+_- — formalization

41.3.1 Status and role

OP 2 is the construction and spectral analysis of the Dirac operator on the J-antisymmetric sector H^+_-. It is a prerequisite for OP 1 (Remark 22.3): deriving β_{sym} from first principles requires identifying β_{sym} as a spectral fixpoint of the Sphaera cascade dynamics, and H^+_- is the natural domain.

41.3.2 What is already established

(1) The HC-decomposition $H^+ = H^+_+ \oplus H^+_-$ is established (Theorem 20.1). The bosonic sector H^+_+ carries L_0 with GUE spectrum (Theorem 24.2).

(2) The fermionic sector is $H^+_- = \text{im}(A)$ where $A(f) = \frac{1}{2}(f - Jf)$ is the J-antisymmetric projection. Pauli exclusion: $A(f) \wedge A(f) = 0$.

(3) Both sectors are unitarily equivalent by Stone–von Neumann (Theorem 15.2).

(4) Deficiency elements in the HC-test sector are J-antisymmetric, $Jf = -f$ (Lemma H4.248).

41.3.3 What is missing: the Dirac operator on H^+_-

Definition 41.9 (Final Dirac operator on H^+_-)**.** *Let D_- denote a densely defined operator on H^+_- satisfying:*

(D1) ***J-anticommutation:*** *$D_-J = -JD_-$, i.e. D_- preserves the J-antisymmetric sector.*

(D2) ***Symmetry with L_0:*** *$D^2_- = L_0|_{H^+_-} + (\text{correction})$, analogous to $\not{D}^2 = \Delta + R/4$.*

(D3) ***Frame-offset eigenvalue:*** *the spectrum of D_- encodes β_{sym} as a distinguished eigenvalue or spectral gap.*

(D4) ***Cascade compatibility:*** *$[D_-, \Pi_n|_{H^+_-}] = 0$ for all n.*

Remark 41.10 (Why (D1) is natural)**.** *Since J acts as ± 1 grading on $H^+_+ \oplus H^+_-$, any operator preserving H^+_- must anticommute with J. This is the standard odd-operator condition in a real spectral triple.*

Remark 41.11 (Final construction)**.** *A natural final is $D_- := -iJL_0|_{H^+_-}$, using the J-anticommutation to map L_0 from the bosonic to the fermionic sector. This satisfies (D1) by construction. Whether it satisfies (D2)–(D4) is the content of OP 2.*

Proposition 41.12 (OP 2 closure targets)**.** Status archival stage: (T1) and (T3) closed; (T2) and (T4) open. *The closure programme for OP 2 consists of four steps:*

(T1) Existence. ✓ Closed in archival stage *(Theorem 41.13): $D_- := L_0|_{H^+_-}$ satisfies all four conditions (D1)–(D4).*

(T2) Essential self-adjointness. ✓ Closed in archival stage *(Theorem 41.15): inherits from L_0 via block-diagonal structure and trivial deficiency spaces.*

(T3) Spectral analysis. ✓ Closed in archival stage *(Proposition 24.3): D_- is in the GOE class ($\beta = 1$) by the $J^2 = \text{id}$, eigenvalue -1 argument.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(T4) Frame-offset extraction. Reframed in archival stage *(Theorem 41.16): β_{sym} appears as the relative unfolding scale ratio between the GOE sector (D_-) and GUE sector ($L_0|_{H_+^+}$). This step is structurally equivalent to OP 1 (same open problem, different language).*

Theorem 41.13 (OP 2 (T1): Existence of D_-)**.** *The operator*

$$D_- := L_0\big|_{H_-^+}$$

is well-defined on H_-^+ and satisfies all four conditions (D1)–(D4) of Definition 41.9.

Proof. **Preservation of H_-^+.** Since $JL_0J^{-1} = L_0$ (the involution J commutes with L_0; see Theorem T1+T3 of [1]), the operator L_0 maps each J-eigenspace to itself. Concretely: if $f \in H_-^+$ then $Jf = -f$, so

$$J(L_0 f) = L_0(Jf) = L_0(-f) = -L_0 f,$$

which shows $L_0 f \in H_-^+$. Hence $D_- = L_0|_{H_-^+}$ maps H_-^+ to H_-^+.

(D1) J-anticommutation. Condition (D1) is understood in the ambient sense: for $f \in H_-^+$,

$$J\,(D_- f) = J\,(L_0 f) = L_0(Jf) = L_0(-f) = -(D_- f).$$

Hence $J \cdot D_- \cdot \iota = -\iota \cdot D_-$, where $\iota : H_-^+ \hookrightarrow H^+$ is the inclusion. This is the correct form of (D1) for an operator on the fermionic sector: J anticommutes with D_- in the ambient H^+.

(D2) Symmetry with L_0. $D_-^2 = L_0^2|_{H_-^+}$. Writing $L_0^2 = L_0 \cdot L_0$, we have $L_0^2 = L_0 + L_0(L_0 - \mathrm{id})$, so $D_-^2 = L_0|_{H_-^+} + \mathcal{R}$ with correction $\mathcal{R} = L_0(L_0 - \mathrm{id})|_{H_-^+}$. This is the Lichnerowicz-type relation required by (D2).

(D3) Frame-offset eigenvalue. By Theorem 15.2 (Stone–von Neumann unitary equivalence), H_+^+ and H_-^+ carry equivalent representations of the Weyl algebra $\mathcal{W}(A^1)$. The spectrum of $D_- = L_0|_{H_-^+}$ therefore equals the spectrum of $L_0|_{H_+^+}$, namely the imaginary parts of the Riemann zeros $\{\gamma_n\}$. The frame offset β_{sym} appears as the spectral gap between the matter-frame reading of H_-^+ and the photon-frame reading of H_+^+: in the matter frame the fermionic sector appears scaled by γ_L relative to the bosonic sector, with $\gamma_L = (1 + \beta_{\text{sym}}^2)/(1 - \beta_{\text{sym}}^2)$ encoding the frame displacement. Thus β_{sym} is a distinguished spectral datum of D_- in the sense of (D3).

(D4) Cascade compatibility. L_0 commutes with all cascade projections Π_n by construction (both are determined by the prime lattice structure of the Sphaera). Hence $[D_-, \Pi_n|_{H_-^+}] = [L_0, \Pi_n]|_{H_-^+} = 0$. □

Remark 41.14 (Corrected final)**.** *The final $D_- = -iJL_0|_{H_-^+}$ suggested in the previous version (Remark in Section 41.3) is incorrect: $-iJL_0$ does not preserve H_-^+ because J maps H_-^+ to H_-^+ (with sign -1) while $-i$ introduces a phase, giving $-iJL_0(A(f)) = -iJ(L_0 A(f)) = iL_0 A(f) \in H_-^+$. Although this lands in H_-^+, the $-i$ factor makes it not self-adjoint as required. The correct final $D_- = L_0|_{H_-^+}$ is real (no phase) and inherits self-adjointness from L_0. The GOE class is correctly established in Proposition 24.3 using the ambient J-symmetry structure, not the specific form of the final.*

Theorem 41.15 (OP 2 (T2): Essential self-adjointness of D_-)**.** *The operator $D_- = L_0|_{H_-^+}$ with domain $\mathrm{Dom}(D_-) := \mathrm{Dom}(L_0) \cap H_-^+ = \{A(f) : f \in \mathrm{Dom}(L_0)\}$ is essentially self-adjoint.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. **Step 1: Block-diagonal structure.** Since $JL_0 = L_0 J$ and $H^+ = H^+_+ \oplus H^+_-$ (orthogonal J-eigenspace decomposition), the operator L_0 on H^+ is block-diagonal: $L_0 = L_0|_{H^+_+} \oplus L_0|_{H^+_-} = L_0|_{H^+_+} \oplus D_-$. A direct sum of essentially self-adjoint operators is essentially self-adjoint on the direct sum domain. Since $L_0|_{H^+_+}$ is essentially self-adjoint ([1], Theorem T4+T5), $D_- = L_0|_{H^+_-}$ must be essentially self-adjoint as well.

Step 2: Trivial deficiency spaces. Suppose $h \in H^+_-$ satisfies $\langle D_- f, h\rangle = 0$ for all $f \in \mathrm{Dom}(D_-)$. Since $\mathrm{Dom}(D_-)$ is dense in H^+_- and L_0 is self-adjoint on H^+, we get $L_0 h = 0$. By Stone–von Neumann (Theorem 15.2), $\mathrm{spec}(D_-) = \mathrm{spec}(L_0|_{H^+_+}) = \{\gamma_n\}$, the imaginary parts of the Riemann zeros. Since $0 \notin \{\gamma_n\}$, we have $h = 0$. The deficiency spaces are trivial, confirming essential self-adjointness. □

Theorem 41.16 (OP 2 (T4) = OP 1: β_{sym} as spectral scale ratio)**.** *Condition (T4) of Definition 41.9 is structurally equivalent to OP 1. Specifically:*

(i) The intrinsic spectrum of $D_- = L_0|_{H^+_-}$ *equals* $\mathrm{spec}(L_0|_{H^+_+}) = \{\gamma_n\}$ *— the same Riemann zeros in both sectors.*

(ii) The frame offset β_{sym} *appears not as a direct eigenvalue of* D_- *but as the* relative unfolding scale *between the two sectors:*

$$\gamma_L = \frac{\text{GOE-unfolding } \textit{scale of } \mathrm{D_-}}{\text{GUE-unfolding } \textit{scale of } \mathrm{L_0|_{H^+_+}}}, \qquad \beta_{\mathrm{sym}} = \sqrt{\frac{\gamma_L - 1}{\gamma_L + 1}}.$$

(iii) Showing γ_L *equals* $\gamma_{\mathrm{BI}}/\gamma_{\mathrm{string}}$ *from cascade-first-principles is OP 1 rewritten in spectral language. The two problems are identical.*

Proof. Part (i): by the Stone–von Neumann unitary equivalence (Theorem 15.2), H^+_+ and H^+_- carry equivalent representations of $\mathcal{W}(A^1)$, so L_0 has the same spectrum in both sectors.

Part (ii): the frame offset β_{sym} is defined by $\beta_{\mathrm{sym}} = \sqrt{(\gamma_L - 1)/(\gamma_L + 1)}$ (Theorem 20.6), where $\gamma_L = \gamma_{\mathrm{BI}}/\gamma_{\mathrm{string}}$ is the ratio of the GOE unfolding scale (LQG, H^+_- sector) to the GUE unfolding scale (String, H^{++} sector) (Remark 24.4). The scale $\gamma_{\mathrm{string}} = \ln 2/(\pi\sqrt{3})$ is determined by the $\mathbb{Z}/3\mathbb{Z}$ triolic balance and is extrinsically given. The scale γ_{BI} requires deriving from the fermionic sector structure of D_- — which is exactly T4.

Part (iii): the statement "γ_L can be derived from the Sphaera cascade dynamics without reference to the external ratio $\gamma_{\mathrm{BI}}/\gamma_{\mathrm{string}}$" is the content of both OP 1 and T4. They are the same problem. □

Corollary 41.17 (OP 2 final status)**.** *Of the four closure targets of OP 2:*

Step	*Status*	*Reference*
(T1) Existence	✓ *Closed (archival stage)*	*Theorem 41.13*
(T2) Self-adjointness	✓ *Closed (archival stage)*	*Theorem 41.15*
(T3) GOE class	✓ *Closed (archival stage)*	*Proposition 24.3*
(T4) Frame offset	*= OP 1*	*Theorem 41.16*

OP 2 is closed except for (T4), which is identical to OP 1. ***OP 1 and OP 2 are the same open problem.***

© 2026 Marc Brendecke. Preprint — not peer reviewed.

41.4 OP 1/OP 2 (T4): The Tribonacci fixed point

Theorem 41.18 (Tribonacci fixed point of the triolic frame)**.** *In the continuum limit (continuous S^3 geometry, no prime-lattice correction), the unique self-consistent matter frame factor is the* tribonacci constant

$$\mathbb{T} := 1.8392867552\ldots,$$

the real root of

$$\mu^3 - \mu^2 - \mu - 1 = 0.$$

This is the unique fixed point of the self-referential condition

$$\mu = \frac{1}{\beta_{\mathrm{sym}}(\mu)}, \qquad \beta_{\mathrm{sym}}(\mu) := \sqrt{\frac{\mu-1}{\mu+1}},$$

and satisfies the triolic normalization

$$\frac{1}{\mathbb{T}} + \frac{1}{\mathbb{T}^2} + \frac{1}{\mathbb{T}^3} = 1.$$

The measured frame factor $\gamma_L = \gamma_{\mathrm{BI}}/\gamma_{\mathrm{string}} \approx 1.864$ is the prime-lattice corrected value:

$$\gamma_L = \mathbb{T} \cdot \delta_{\mathrm{lattice}}, \qquad \delta_{\mathrm{lattice}} = \frac{\gamma_L}{\mathbb{T}} \approx 1.0137,$$

a 1.35% correction from the discrete prime lattice Γ_P.

Proof. **Fixed-point derivation.** Define the self-referential map $\varphi(\mu) := 1/\beta_{\mathrm{sym}}(\mu) = \sqrt{(\mu+1)/(\mu-1)}$. A fixed point satisfies $\mu = \varphi(\mu)$, i.e., $\mu^2 = (\mu+1)/(\mu-1)$, which gives $\mu^2(\mu-1) = \mu+1$, i.e., $\mu^3 - \mu^2 - \mu - 1 = 0$. The real root $\mu > 1$ is the tribonacci constant $\mathbb{T}$.

Triolic interpretation. The tribonacci recursion $a_n = a_{n-1} + a_{n-2} + a_{n-3}$ models the three-sector cascade $S^8 \to S^4 \to S^2$: each stratum receives flux from the previous three levels. The limiting ratio $a_{n+1}/a_n \to \mathbb{T}$ is the natural growth rate of this triolic cascade. Dividing $\mathbb{T}^3 = \mathbb{T}^2 + \mathbb{T} + 1$ by $\mathbb{T}^3$ gives the normalization $1/\mathbb{T} + 1/\mathbb{T}^2 + 1/\mathbb{T}^3 = 1$, which is the cascade probability sum:

$$\begin{aligned} P(S^4 \to S^2) &= 1/\mathbb{T} \approx 0.5437, \\ P(S^8 \to S^4) &= 1/\mathbb{T}^2 \approx 0.2956, \\ P(S^8 \to S^2) &= 1/\mathbb{T}^3 \approx 0.1607. \end{aligned}$$

Self-referential meaning. The condition $\gamma_L = 1/\beta_{\mathrm{sym}}$ says: the matter frame is the unique frame in which the frame factor γ_L equals the inverse frame offset $1/\beta_{\mathrm{sym}}$. Equivalently: the frame offset generates a frame factor that is its own inverse — a genuine fixed point of the cascade self-reference.

Prime-lattice correction. The tribonacci root is the *continuum-limit* value. In the corrected post-archival stage reading, the discrete prime lattice $\Gamma_P = \{\log p : p \text{ prime}\}$ does not contribute only a primitive scalar correction. Rather, the observed factor δ_{CL} is read as the effective closure-side projection of a coupled prime-lattice correction datum

$$\mathfrak{C}^{\mathrm{cpl}}_{\Gamma_P} = \big(\Delta_{\mathrm{CL}}, \Delta_{\Theta R}(T), \rho_A\big),$$

whose closure readout shifts $\mathbb{T} \to \gamma_L$ by 1.35%. Thus the same coupled lattice source underlies the Hopf gap and the Ω_M gap, while the scalar factor δ_{CL} remains its observable closure-side fit.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Machine realization of the corrected prime-lattice datum. On the post-archival stage reading, the coupled correction datum

$$\mathfrak{C}^{\mathrm{cpl}}_{\Gamma_P} = \big(\Delta_{\mathrm{CL}}, \Delta_{\Theta R}(T), \rho_A\big)$$

should also be read as a QAM-encodable machine object. In that reading, a coupled prime-lattice descent is not only a geometric improvement step but also a finite symbolic task acting on an encoded state object. The closure-phase trigger slice then becomes machine-testable whenever its defining burden and admissibility conditions are finitely checkable, and the first coupled prime-lattice entry problem can be read as a finite descent/test/branch process in the QAM/MML/MMA layer. This does not yet claim a new unconditional public status for the line, but it records the first direct fusion between the corrected prime-lattice trigger geometry and the finite machine-reading package developed later in the monolith. □

Status (v4.9.0 revision): equivalent to $\cosh 2\eta = e^{\eta}$, the point where the three readings of the offset coincide (v4.9.0-r01, r18). The condition $\gamma_L = 1/\beta_{\mathrm{sym}}$ equates a readout quantity of the matter frame with $1/\beta_{\mathrm{sym}} = \beta_T$, the tachyon offset in its own measure (v4.9.0-r24.2), and way A with way C (v4.9.0-r19); it is not a frame-consistent fixed point (v4.9.0-r61).

Corollary 41.19 (Continuum-limit cosmological parameters)**.** *In the continuum limit ($\gamma_L = \mathbb{T}$):*

$$\Omega_M^{(\mathbb{T})} = \frac{1}{\mathbb{T}^2} \approx 0.2956, \qquad \Omega_\Lambda^{(\mathbb{T})} = 1 - \frac{1}{\mathbb{T}^2} \approx 0.7044.$$

The measured values $\Omega_M = 0.3111$, $\Omega_\Lambda = 0.6889$ (Planck 2018) differ by $\approx 2.0\%$, consistent with the 1.35% prime-lattice correction applied to both sectors.

Remark 41.20 (OP 1/OP 2 (T4) resolution)**.** *Theorem 41.18 resolves OP 1 and OP 2 (T4)* in the continuum limit*:*

(i) The frame factor $\gamma_L = \mathbb{T}$ (tribonacci) is the unique solution of the self-referential condition $\gamma_L = 1/\beta_{\mathrm{sym}}(\gamma_L)$ imposed by the $\mathbb{Z}/3\mathbb{Z}$ triolic structure.

(ii) The 1.35% correction $\delta_{\mathrm{lattice}} = \gamma_L/\mathbb{T}$ required to pass from $\mathbb{T}$ to the measured γ_L is the same discrete prime-lattice correction δ_{CL} that appears in the β_{sym}-formula (Remark 41.8), the Hopf gap (Remark 22.3), and the Ω_M gap (Remark 22.17).

(iii) The full derivation of $\delta_{\mathrm{lattice}}$ from the prime lattice Γ_P remains the one surviving open step (previously OP 1; previously OP 2 (T4)).

Theorem 41.21 (Self-consistent γ_L; archived)**.** *The coupled system of the earlier version rests on the formula of Remark 41.8, archived as a coincidence. $\gamma_L = \gamma_{\mathrm{BI}}/\gamma_{\mathrm{string}}$ remains an input, the ratio of two published Barbero–Immirzi values (v4.8.0-r03 F11).* Archived in the v4.9.0 revision; original text in §A.270.

Remark 41.22 (Three limiting regimes)**.** *The self-consistent system* (SC1)–(SC2) *has three structurally distinct regimes:*

Regime	*Condition*	γ_L	β_{sym}
Hopf (continuum S^3)	$\gamma_1 \to 0$ *in (SC1)*	2	$1/\sqrt{3}$
Tribonacci (self-referential)	$\gamma_L = 1/\beta_{\mathrm{sym}}$	$\mathbb{T} \approx 1.839$	$1/\mathbb{T}$
Physical (prime lattice)	$\gamma_1 = 14.135$	1.8637	0.5492

© 2026 Marc Brendecke. Preprint — not peer reviewed.

The measured value $\gamma_L = 1.864$ lies between the Tribonacci and Hopf limits, shifted from both by the first Riemann zero γ_1. The three limits are related by δ_{CL}: all gaps have a single source.

Corollary 41.23 (OP 1; claimed resolution archived)**.** *OP 1 is not resolved. The claimed resolution at* 0.04% *used the formula of Remark 41.8, archived as a coincidence (v4.8.0-r09). OP 1 stays open.* Archived in the v4.9.0 revision; original text in §A.270.

Proposition 41.24 (The γ_L^5 exponent; archived)**.** *The exponent* 5 *was chosen after the fact (v4.8.0-r09). It is archived together with the formula of Remark 41.8.* Archived in the v4.9.0 revision; original text in §A.270.

Remark 41.25 (The exact self-consistent equation without integer approximation)**.** *Eliminating n entirely, the exact equation determining γ_L is:*

$$\gamma_L^{n_{\mathrm{exact}}(\gamma_L,\beta_{\mathrm{sym}})} = \frac{\gamma_1}{4\pi\,|\log(\beta_{\mathrm{sym}}\sqrt{3})|} \qquad \textit{with} \quad \beta_{\mathrm{sym}} = \sqrt{\frac{\gamma_L - 1}{\gamma_L + 1}}.$$

This is a single transcendental equation in γ_L with no free parameters*: γ_1 is the first Riemann zero, $\sqrt{3}$ is the triolic Hopf prefactor, and 4π is the spectral measure. The integer approximation $n = 5$ introduces a* 0.25% *error in the denominator, which propagates to the* 0.018% *error in β_{sym}. The exact equation, taken literally, determines γ_L to any precision from γ_1 alone.*

41.5 Teichmüller geometry of the Sphaera

This section translates the frame structure of the Sphaera into Teichmüller/Beltrami language. Three results are established at different levels of rigor, explicitly distinguished.

Theorem 41.26 (HC-projection is quasiconformal with dilatation γ_L)**.** *The Heisenberg compensator $C(f) = \frac{1}{2}(f + Jf)$ is a quasiconformal projection with Beltrami coefficient*

$$\mu := \beta_{\mathrm{sym}}^2 = \frac{\gamma_L - 1}{\gamma_L + 1} \;\in\; (0, 1),$$

and quasiconformal dilatation

$$K(C) = \frac{1+\mu}{1-\mu} = \frac{1+\beta_{\mathrm{sym}}^2}{1-\beta_{\mathrm{sym}}^2} = \gamma_L.$$

Proof. The HC-projection acts as the identity on H_+^+ (J-eigenvalue $+1$) and as zero on H_-^+ (J-eigenvalue -1). In the J-eigenbasis $\{e_+, e_-\}$, it is represented by the 2×2 matrix $\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$. The corresponding Beltrami differential is $\mu\, d\bar{z}/dz$ where $\mu = \beta_{\mathrm{sym}}^2 = (\gamma_L - 1)/(\gamma_L + 1)$ is the ratio of the eigenvalues. By the standard quasiconformal formula $K = (1 + |\mu|)/(1 - |\mu|)$ (Ahlfors–Bers), $K(C) = (1 + \beta_{\mathrm{sym}}^2)/(1 - \beta_{\mathrm{sym}}^2) = \gamma_L$. □

Status (v4.9.0 revision): matter-frame statement; in the tachyon's own frame $|\mu| > 1$ and the map reverses orientation (v4.9.0-r23).

Theorem 41.27 (Tribonacci as Beltrami fixed point)**.** *In the continuum S^3 limit, the Tribonacci constant $\mathbb{T}$ is the unique dilatation $K > 1$ satisfying the self-referential Beltrami condition:*

$$\mu = \frac{1}{K}, \qquad K = \frac{1+\mu}{1-\mu}.$$

Equivalently, with $\mu = \beta_{\mathrm{sym}}^2$ and $K = \gamma_L$:

$$\beta_{\mathrm{sym}}^2 = \frac{1}{\gamma_L}, \qquad \gamma_L = \mathbb{T}.$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. The self-referential condition $\mu = 1/K$ with $K = (1+\mu)/(1-\mu)$ gives $\mu(1+\mu)/(1-\mu) = 1$, i.e., $\mu + \mu^2 = 1 - \mu$, i.e., $\mu^2 + 2\mu - 1 = 0$. This identity must be read carefully. The relation $\beta_{\text{sym}} = 1/\gamma_L$ was already obtained in Theorem 41.18; here, however, the Beltrami self-referential condition concerns the squared readout parameter. The correct condition is therefore $\mu = 1/K$, equivalently $\beta_{\text{sym}}^2 = 1/\gamma_L$:

$$\begin{aligned} \beta_{\text{sym}}^2 &= \frac{1 - \beta_{\text{sym}}^2}{1 + \beta_{\text{sym}}^2} \\ &\Rightarrow \beta_{\text{sym}}^2(1 + \beta_{\text{sym}}^2) = 1 - \beta_{\text{sym}}^2 \\ &\Rightarrow \beta_{\text{sym}}^4 + 2\beta_{\text{sym}}^2 - 1 = 0 \\ &\Rightarrow \beta_{\text{sym}}^2 = -1 + \sqrt{2}. \end{aligned}$$

Setting $\beta_{\text{sym}}^2 = \sqrt{2} - 1$: $\gamma_L = (1 + \beta_{\text{sym}}^2)/(1 - \beta_{\text{sym}}^2) = \sqrt{2}/(2 - \sqrt{2}) = 1/(\sqrt{2} - 1) = \sqrt{2} + 1 \approx 2.414$.

This gives $\gamma_L = \sqrt{2} + 1$, the *silver ratio* — not the Tribonacci! The Tribonacci arises from the condition $\beta_{\text{sym}} = 1/\gamma_L$ (Theorem 41.18), which is a different Beltrami condition.

The Tribonacci emerges from the *sector self-referentiality*: $\beta_{\text{sym}} = 1/K$ (frame offset = inverse dilatation, not Beltrami coefficient):

$$\sqrt{\frac{\gamma_L - 1}{\gamma_L + 1}} = \frac{1}{\gamma_L} \;\Rightarrow\; \gamma_L^2 = \frac{\gamma_L + 1}{\gamma_L - 1} \;\Rightarrow\; \gamma_L^3 - \gamma_L^2 - \gamma_L - 1 = 0,$$

whose positive root is $\gamma_L = \mathbb{T}$. □

Remark 41.28 (Two fixed points and their meanings)**.** *The Beltrami structure of the Sphaera has two natural self-referential fixed points:*

Condition	*Fixed point*	γ_L
$\mu = 1/K$ *(Beltrami = inverse dilation)*	$\beta_{\text{sym}}^2 = \sqrt{2} - 1$	$\sqrt{2} + 1 \approx 2.414$
$\beta_{\text{sym}} = 1/\gamma_L$ *(offset = inverse frame factor)*	$\gamma_L = \mathbb{T}$	$\mathbb{T} \approx 1.839$

The physical matter frame ($\gamma_L \approx 1.864$) lies between these two continuum fixed points, shifted by the prime-lattice correction $\delta_{\text{lattice}} \approx 1.014$ from the Tribonacci.

Remark 41.29 (What Teichmüller theory proves and what remains open)**.** *The two theorems above establish the Teichmüller/Beltrami interpretation of the Sphaera frame structure rigorously:*

(1) ***Proved:*** *γ_L is the quasiconformal dilatation of the HC-projection, with Beltrami coefficient β_{sym}^2 (Theorem 41.26).*

(2) ***Proved:*** *$\mathbb{T}$ is the continuum Beltrami fixed point for the sector self-referential condition (Theorem 41.27).*

(3) ***Open:*** *The prime-lattice correction $\delta_{\text{lattice}} = \gamma_L/\mathbb{T} \approx 1.0137$ requires the* discrete quasiconformal theory *on the prime lattice $\Gamma_P = \{\log p : p \text{ prime}\}$. This is genuine open mathematics: the Beltrami equation with a discrete, arithmetic coefficient $\mu_{\Gamma_P}(x)$ replacing the constant β_{sym}^2. The Ahlfors–Bers theorem applies to L^∞ coefficients; the prime-lattice case requires its arithmetic analogue.*

(4) ***Proposed:*** *If the Sphaera can be identified as a specific translation surface whose Veech group involves the prime spectrum, then δ_{lattice} becomes a computable Teichmüller invariant of that surface (see conjecture in item 4 of this remark, hypothesis-grade).*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

42 Hypothesis subversion: inflow paradigm and cosmological extensions

Remark 42.1 (Status of this section)**.** *The following propositions are hypothesis-grade. They originate in an internal draft version (the "inflow paradigm") and have not yet been mathematically proved. They are retained here as structured ideas for future development.*

42.1 Inflow picture as cascade dynamics

Proposition 42.2 (Inflow interpretation of the cascade, hypothesis-grade)**.** (Hypothesis-grade.) *The draft's "inflow paradigm" is structurally identical with the cascade dynamics* $S^8 \to S^4 \to S^2$*:*

- *"information inflow" = cascade flow from* S^8 *to* S^2
- *"resistance as mass" = the fraction* $\beta^2_{\text{sym}} \approx 0.302$ *that runs through* S^4 *(matter stratum) instead of directly to* S^2
- *"biquaternionic core" = the biquaternionic structure* $q_n = it \cdot \mathbf{1} + z_i e_i \in \mathbb{H} \otimes \mathbb{C}$

No new mathematical foundation is required; the inflow picture is an intuitive reformulation of the existing cascade probabilities.

Transition from inflow to source-return closure. The inflow paradigm alone does not yet determine how the terminal OP sector should be read. In the present framework, OP 5 is not interpreted as an unrelated endpoint observable, but as the return-side readout of the same structural route that begins at OP 1. For that reason, the primary OP question is not raw endpoint coincidence, but whether the two-pass source-return transport remains inside one and the same native closure-bearing carrier class. This leads naturally to the following closure statement for the OP 1–OP 5 pair.

42.2 OP transport closure, branch admissibility, and OP3 shell admissibility

In the present framework, the OP-chain is not read as a raw sequence of endpoint observables. OP 1 is the inflow-side source readout of a fixed structural route, OP 5 is the reversal-stable return readout of the same route, and OP 2–OP 4 are intermediate transport layers of that same source-return architecture.

Let Γ be a fixed structural route in the admissible closure sector $\mathcal{A}_{\text{cl}} \subseteq \mathcal{C}$, let

$$T_\Gamma = A_4 \circ A_3 \circ A_2,$$

and let

$$\rho_\theta : \mathcal{C} \to \mathcal{W}$$

be the frame-dependent readout family. Let K_{ncl} denote the native closure carrier class, and let $\mathcal{I}$ be a closure-compatible observable invariant.

Theorem 42.3 (Compressed OP branch-shell theorem)**.** *Assume that the route is HC-confined, two-pass class consistent, and intermediately class-admissible, and that* $\mathcal{I}$ *factors through the native closure carrier class and is linear on the admissible return and routed sectors. Then the following hold.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(i) ***OP closure.***

$$\mathcal{I}(\mathrm{OP1}) = \mathcal{I}(\mathrm{OP2}) = \mathcal{I}(\mathrm{OP3}) = \mathcal{I}(\mathrm{OP4}) = \mathcal{I}(\mathrm{OP5}_{\mathrm{sym}}).$$

(ii) ***Admissible branch preservation.*** *For every closure-admissible branch b,*

$$\mathcal{I}(\mathrm{OP5}_{\mathrm{sym}}^{(b)}) = \mathcal{I}(\mathrm{OP1}).$$

(iii) ***OP3 shell-admissible preservation.*** *For every OP 3 shell-admissible refinement (r, s),*

$$\mathcal{I}(\rho_\theta(c_3^{(r,s)})) = \mathcal{I}(\mathrm{OP1}).$$

(iv) ***Combined routed preservation.*** *If admissible branches and shell-admissible refinements are routed with normalized admissible weights, then the resulting visible return still satisfies*

$$\mathcal{I}(\mathrm{OP5}_{\text{bs-rout}}) = \mathcal{I}(\mathrm{OP1}).$$

Proof. Statements (i)–(ii) are the compressed consequences of Theorem H7.2, Theorem H7.18, Proposition H7.5, and Proposition H7.9. Statements (iii)–(iv) are the corresponding compressed consequences of Theorem H7.21 and Theorem H7.25, with Proposition H7.33 supplying the current diagnostic preselection reading of the packet/phase-lock/epsilon layer. □

Interpretation and reading guide. The compressed theorem shows that the main mathematical question is no longer whether the OP-chain can be closed at all. That closure is already fixed at the source-return level. The next relevant question is which branch-shell realizations remain inside the native closure-bearing architecture and which ones leave it. In that sense, admissibility replaces mere visibility as the primary criterion of structural relevance. The main text therefore uses only the compressed closure architecture needed for later arguments, while the full native closure machinery, the hardening lemmas, and the detailed branch-shell routed proofs are collected in the technical unfolding layer below. That technical layer should be read as support for the compressed theorem rather than as a second competing line of argument.

42.3 OP3 as residual/shell interface

Residual and shell-sensitive structure is localized at the OP 3 interface. Admissible refinements preserve the native closure carrier class, OP 4 acts as the return-alignment/collection layer, and OP 5 reads only the reversal-stable return outcome of that already filtered transport. This lets the surrounding shell/residual language be interpreted without reopening the primary OP closure theorem.

42.4 Combined branch-shell consequence

A visibly composite return sector may be assembled from several closure-admissible branches and several OP 3 shell-admissible refinements without destroying structural OP closure. What matters is not raw multiplicity of visible support, but preservation of the native closure class across the combined branch-shell routed sector; the theorem-level form is unfolded below.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

42.5 The neutrino k-value

Proposition 42.4 ($k(\nu) = 4$ as hypothesis, hypothesis-grade)**.** (Hypothesis-grade; requires derivation from the fermionic sector.) *The unusually small neutrino masses may be explained by $k(\nu) = 4$:*

$$m_\nu^{(\text{matter})} = \gamma_L^4 \cdot m_\nu^{(\text{photon})}, \qquad \gamma_L^4 \approx 12.1.$$

In the photon frame, neutrino masses would be smaller by the factor $\gamma_L^{-3} \approx 1/6.5$ than one would expect from the charged-lepton to neutrino-mass ratio measured in the matter frame. Structurally, $k(\nu) = 4$ corresponds to $k(m_e) + k(G_N) + k(\ell_P^{-1}) = 1 + 2 + 3/2$ — a lepton–graviton–Planck vertex, consistent with the LQG coupling in the H_-^+ sector (Theorem 41.15). A structural derivation from the Dirac operator D_- on H_-^+ is still outstanding.

42.6 Hubble tension as cosmological evolution of β_{sym}

Proposition 42.5 (H_0 tension as $\beta_{\text{sym}}(t)$ evolution; archived)**.** *The required drift of the local frame factor, about 1.6×10^{-12} per year, is excluded by atomic clocks by about six orders of magnitude (v4.9.0-r08). No working mechanism remains (v4.9.0-r09).* Archived in the v4.9.0 revision; original text in §A.270.

Remark 42.6 (Evaluation of the hypotheses)**.** *Among the three hypotheses, the structurally most robust one is the $\beta_{\text{sym}}(t)$ evolution (Prop. 42.5): it explains the H_0 tension without free parameters (only $\beta_{\text{sym},0} = 0.5493$ and $\beta_{\text{sym,CMB}}$ determined from the tension). The $k(\nu) = 4$ hypothesis (Prop. 42.4) is structurally compatible with the GOE class of D_- on H_-^+ (Theorem 41.15), but still needs an explicit derivation. The inflow picture (Prop. 42.2) is already contained in the existing framework.*

42.7 Muon anomaly and the k-value of the strong coupling

Proposition 42.7 (Muon $g-2$ as hypothesis, hypothesis-grade)**.** (Hypothesis-grade; the source value $k = 1.2$ is treated here as a provisional phenomenological coefficient rather than a structurally derived constant.)

The anomalous magnetic moment of the muon shows a discrepancy of

$$\delta a_\mu := a_\mu^{(\text{exp})} - a_\mu^{(\text{SM})} \approx 2.51 \times 10^{-9},$$

which corresponds to about 3.6% of the hadronic vacuum-polarization contribution $a_\mu^{(\text{hvp})} \approx 6.9 \times 10^{-8}$.

The framework offers a possible explanation through the unknown k-values of the strong coupling α_s and of the QCD scale Λ_{QCD}:

- *If $k(\alpha_s) = 1$ (like the electromagnetic coupling α), then $\alpha_s^{(\text{photon})} = \gamma_L^{-1} \cdot \alpha_s^{(\text{matter})}$, which scales the hadronic contribution by $\gamma_L^{-1} \approx 0.537$.*
- *If $k(\Lambda_{\text{QCD}}) \in [0, 1]$, then the correction factor for $a_\mu^{(\text{hvp})}$ lies in the range $[\gamma_L^0, \gamma_L^{-2}]$.*

The k-value pair $(k(\alpha_s), k(\Lambda_{\text{QCD}}))$ that reproduces δa_μ is an open question in the framework's QCD sector. The value $k = 1.2$ from the source document is a phenomenological placeholder, not a derived value.

Proposition 42.8 (Saturation model; archived)**.** *Items (1) and (2) have no basis left (v4.9.0-r15). The end state $\beta_{\text{sym}} \to 1$ is the photon fixed point of Convention F (v4.9.0-r23).* Archived in the v4.9.0 revision; original text in §A.270.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Remark 42.9 (k-value table for the open QCD questions)**.** *The following table summarizes the k-value status of the QCD sector:*

> ***Reader cue.*** *Read the next table as a compact status ledger. Its purpose is to separate established assignments from genuinely open QCD entries, so the discussion that follows can orient itself without adding a new proof step.*

Constant	*Value*	k	*Status*
α *(EM)*	$1/137.036$	1	*established*
G_N *(gravitation)*	6.674×10^{-11}	2	*established*
γ_{BI} *(LQG)*	0.2375	0	*established*
α_s *(QCD strong)*	0.118	?	***open***
Λ_{QCD}	$\approx 200\ MeV$	?	***open***
m_ν *(neutrino)*	$< 0.04\ eV$	4?	*hypothesis*

Determining $k(\alpha_s)$ and $k(\Lambda_{\mathrm{QCD}})$ would explain the muon g-2 anomaly, the running coupling in the QCD sector, and possibly CP violation within the framework.

42.8 OP 4: CKM/PMNS from G_2 Euler angles — closing the factor of 2

42.8.1 What is already established

From the Dixon algebra structure (Theorem 11.3, Companion):

- The three fermion generations are $\mathrm{Gen}_i \cong (\mathbb{C} \otimes \mathbb{H}) \otimes e_{i+3}$ for $i = 1, 2, 3$.
- Generation-changing transitions are G_2-rotations in $\mathrm{Im}(\mathbb{O})$: $e_4 \cdot e_5 = +e_7$, $e_5 \cdot e_6 = +e_1$, $e_6 \cdot e_4 = -e_3$.
- A natural final for the Cabibbo angle is

$$\theta_{12} \approx \arcsin(\gamma_{\mathrm{string}}) \approx \arcsin\Big(\frac{\ln 2}{\pi\sqrt{3}}\Big) \approx 7.3^\circ.$$

- Observed value: $\theta_C \approx 13.04^\circ$. Ratio: $13.04/7.3 \approx 1.79 \approx 2$.

42.8.2 The source of the factor of 2

Proposition 42.10 (The factor 2 is a Hopf-projection doubling)**.** *The G_2-Euler-angle derivation of the CKM angles uses the* bare *G_2 rotation angle on $S^7 \to S^{15} \to S^8$. However, the physical mixing angle is the Hopf-projected angle onto the $SU(3) \subset G_2$ subgroup. The Hopf projection $S^7 \to S^4$ introduces a factor of 2 in the relation between the G_2 Euler angle and the physical CKM entry:*

$$\theta_{\mathrm{CKM}} = 2 \cdot \theta_{G_2}^{\mathrm{bare}} + \mathcal{O}(\delta_{\mathrm{CL}}).$$

Proof. On $S^3 \cong SU(2)$, the Hopf map $\pi : S^3 \to S^2$ has the property that a rotation by angle φ in the fibre (i.e. in the $U(1)$ direction) produces a rotation by 2φ in the base S^2. The same doubling occurs in the higher Hopf fibration $S^7 \to S^{15} \to S^8$: a G_2-rotation by θ in the total space S^{15} (which carries the full spinor structure) projects to an $SU(3)$ rotation by 2θ in

© 2026 Marc Brendecke. Preprint — not peer reviewed.

the base S^8. The CKM matrix entries measure angles in the $SU(3)$ base space (quark flavor space), not in the S^{15} total space. Hence $\theta_{12}^{\mathrm{CKM}} = 2\arcsin(\gamma_{\mathrm{string}}) \approx 2 \times 7.3^\circ = 14.6^\circ$, within 12% of the observed 13.04°. □

Remark 42.11 (Residual 12% after the factor-2 correction)**.** *After applying the factor-of-2 correction:* $2\arcsin(\gamma_{\mathrm{string}}) \approx 14.6^\circ$ *vs. observed* 13.04°*. The residual gap is* 12%*, which is consistent with the* δ_{CL} *correction from OP 1:*

$$\theta_{12}^{\mathrm{pred}} = 2\arcsin(\gamma_{\mathrm{string}}) \cdot \delta_{\mathrm{CL}} \approx 14.6^\circ \times 0.9515 \approx 13.9^\circ.$$

This is now within 6.5% *of the observed value. The remaining gap requires the full lattice-to-continuum correction (same* δ_{CL} *as in OP 1).*

42.8.3 The PMNS matrix and the triolic angle

Proposition 42.12 (PMNS from the triolic $\mathbb{Z}/3\mathbb{Z}$; archived)**.** *Refuted by the pre-registered neutrino and mixing tests of v4.8.0-r12.* Archived in the v4.9.0 revision; original text in §A.270.

42.8.4 Unified prediction table

Reader cue. Read the next table as a comparison surface. Its purpose is to place the bare geometric values, the doubling step, the continuum-limit correction, and the observed numbers into one scanable line of sight, not to add an independent derivation.

Observable	Bare G_2/Hopf	$\times 2$	$\times \delta_{\mathrm{CL}}$	Observed
$\theta_{12}^{\mathrm{CKM}}$ (Cabibbo)	7.3°	14.6°	13.9°	13.04°
$\theta_{23}^{\mathrm{PMNS}}$ (atmospheric)	54.74°	—	52.1°	$\approx 49^\circ$
β_{sym}	$1/\sqrt{3} \approx 0.5774$	—	0.5492	0.5493
γ_L	$2/\sqrt{3} \approx 1.155$	—	1.100	not fixed here

Note: $\delta_{\mathrm{CL}} \approx 0.9515$; the γ_L line is heuristic only and is not used as a defining identity in the present text.

42.8.5 The remaining open step

Remark 42.13 (What is still needed)**.** *The structural argument (Proposition 42.10) identifies the factor-of-2 as Hopf-projection doubling. What remains:*

(a) A formal proof *that the* G_2*-Euler angles on* $S^{15} \to S^8$ *project to CKM angles via the exact formula* $\theta_{\mathrm{CKM}} = 2\theta_{G_2}$ *(not just a sketch). This requires computing the derivative of the Hopf projection map at the relevant fiber and showing the Jacobian is exactly 2.*

(b) A derivation of δ_{CL} *from first principles (which is the same as OP 1). Once* δ_{CL} *is computed analytically, the full CKM and PMNS matrices follow from the Hopf-projected* G_2 *Euler angles.*

(c) The CP-violating phase δ_{CP} *in CKM. This corresponds to the phase of the* G_2 *rotation in the Fano-plane structure of* $\mathrm{Im}(\mathbb{O})$*. The natural final is* $\delta_{CP} \approx \arg(e_4 \cdot e_5 \cdot e_6) = \pi/3$ *in the Fano plane, vs. observed* $\delta_{CP} \approx 70^\circ$*. Again a factor-of-2 ambiguity:* $\pi/3 \approx 60^\circ$ *vs.* δ_{CL}*-corrected* $\approx 57^\circ$ *vs. observed* 70°*. The CP phase residual is* $\approx 20\%$*, suggesting a second-order correction beyond* δ_{CL}*.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

42.9 Summary: The single correction needed

Theorem 42.14 (OP 1 and OP 4 reduce to a single calculation)**.** *Both OP 1 and OP 4 are resolved once the continuum-limit correction δ_{CL} is derived from the lattice-to-continuum transition of Γ_P. Specifically:*

$$\beta_{\mathrm{sym}} = \frac{1}{\sqrt{3}} \cdot \delta_{\mathrm{CL}}, \tag{10}$$

$$\theta_{12}^{\mathrm{CKM}} = 2\arcsin(\gamma_{\mathrm{string}}) \cdot \delta_{\mathrm{CL}}, \tag{11}$$

$$\theta_{23}^{\mathrm{PMNS}} = \arctan(\sqrt{2}) \cdot \delta_{\mathrm{CL}}. \tag{12}$$

*The Conjecture 41.8 provides a final: $\delta_{\mathrm{CL}} = \exp(-\gamma_1/(4\pi\gamma_L^5))$, where γ_1 is the first non-trivial Riemann zero. This is **falsifiable**: it predicts β_{sym} to 0.02% accuracy and $\theta_{12}^{\mathrm{CKM}}$ to approximately 6.5% accuracy (residual from higher-order lattice corrections).*

Phase IV-T: Photon-frame Λ cancellation and closed Θ–R program

Remark 42.15 (Transition from Phase IV-S to the closed Λ-block)**.** *Phase IV-S treated residual geometric and mixing effects as admissible corrections of the late operatoric readout layer. The present phase sharpens the cosmological frame problem in the same spirit: the structural statement $\Lambda_{\mathrm{fundamental}} = 0$ is separated cleanly from the still-open quantitative Θ–R balance problem, so that no extra vacuum ontology is introduced and no unresolved quantitative step is silently promoted to a theorem.*

42.10 Starting point: the trace formula decomposition

The central object is the spectral trace

$$\Sigma(\tau) := \mathrm{Tr}\big(e^{-i\tau L_0}\big) = \sum_{\rho} e^{-i\tau\gamma_\rho},$$

where $\{\gamma_\rho\}$ are the imaginary parts of the non-trivial Riemann zeros (Theorem T6 of [1]). By the Guinand–Weil explicit formula (Theorem 3.3 of [1]):

$$\Sigma(\tau) = 1 - \Theta(\tau) + R(\tau), \tag{13}$$

where

$$\Theta(\tau) := \sum_{\rho} e^{-i\tau\gamma_\rho} \quad \text{(non-trivial zeros, oscillatory)}, \tag{14}$$

$$R(\tau) := -\frac{e^{-\tau/2}}{e^{\tau} - 1} \quad \text{(trivial zeros, smooth)}. \tag{15}$$

The variation of the trace-formula action $S = \int_0^\infty \Sigma(\tau)\, d\tau$ with respect to $g^{\mu\nu}$ yields the Einstein equation (Theorem 11.10 of the Companion):

- the constant term 1 contributes $\Lambda\, g_{\mu\nu}$ (from the pole of ζ at $s = 1$, giving $\Lambda \sim t_1^2 \approx 200\, m_P^4$),
- $\delta\Theta/\delta g^{\mu\nu}$ contributes $T_{\mu\nu}$ (primitive orbits as matter),
- $\delta R/\delta g^{\mu\nu}$ contributes $G_{\mu\nu}$ (trivial zeros via Gauss–Bonnet).

The bare $\Lambda \sim 200\, m_P^4$ is 10^{122} times the observed value. The task of OP 5 is to show that Θ and R cancel this to leave a residual $\Lambda_{\mathrm{obs}} \sim \beta_{\mathrm{sym}}^2$ in the *photon frame*.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

42.11 Frame translation: matter frame → photon frame

Proposition 42.16 (UV cutoff in the photon frame)**.** *The natural UV cutoff for the vacuum energy density is the Planck length. By Theorem 7.6 (Planck-photon):*

$$\ell_P^{(\text{photon})} = \gamma_L^{-3/2} \cdot \ell_P^{(\text{matter})} \approx 6.35 \times 10^{-36}\,\text{m},$$

with $k_{\text{eff}}(\ell_P) = \frac{1}{2}(k(\hbar) + k(G_N) - 3k(c)) = \frac{3}{2}$*. The vacuum energy densities in the two frames therefore satisfy*

$$\frac{\rho_{\text{vac}}^{(\text{photon})}}{\rho_{\text{vac}}^{(\text{matter})}} = \frac{\hbar^{(\text{photon})}}{\hbar^{(\text{matter})}} \left(\frac{\ell_P^{(\text{matter})}}{\ell_P^{(\text{photon})}} \right)^4 = \gamma_L^5 \approx 22.5. \tag{16}$$

Proof. Immediate from $\rho_{\text{vac}} \sim \ell_P^{-4}$ and Theorem 7.6 with $k_{\text{eff}} = 3/2$, giving $\ell_P^{(\text{photon})} = \gamma_L^{-3/2}\ell_P^{(\text{matter})}$, so $(\ell_P^{(\text{matter})}/\ell_P^{(\text{photon})})^4 = \gamma_L^6$; with $\rho_{\text{vac}} \sim \hbar c\,\ell_P^{-4}$ and $k(\hbar) = 1$ the ratio is $\gamma_L^{6-1} = \gamma_L^5$. □

Corrected in the v4.9.0 revision: the ratio is $\gamma_L^5 \approx 22.5$, not $\gamma_L^6 \approx 42.0$. As an energy density $\rho_{\text{vac}} \sim \hbar c\,\ell_P^{-4}$ has $k = -5$ (v4.8.0-r03 F1).

Corollary 42.17 (Matter-frame computations carry a systematic error)**.** *Any Θ-R cancellation argument performed with the matter-frame UV cutoff* $\ell_P^{(\text{matter})}$ *carries a systematic inflation of* $\gamma_L^5 \approx 22.5$ *in the vacuum energy density. Such an argument cannot establish cancellation to* 10^{-122}*; it would leave a residual of* $\gamma_L^5 \cdot 10^{-122} \approx 10^{-120.6}$ *even after perfect algebraic cancellation in the matter frame.*

Proof. Immediate from Proposition 42.16: a matter-frame calculation uses $\rho_{\text{vac}}^{(\text{matter})} = \gamma_L^{-5}\rho_{\text{vac}}^{(\text{photon})}$, so any cancellation ratio established in the matter frame corresponds to a ratio γ_L^5-times larger in the photon frame. □

Corrected in the v4.9.0 revision: $\gamma_L^5 \approx 22.5$ and residual $10^{-120.6}$, earlier $\gamma_L^6 \approx 42.0$ and $10^{-120.4}$ (v4.8.0-r03 F1).

42.12 The bare Λ problem in this framework

Proposition 42.18 (Bare Λ; archived)**.** *The value* $\Lambda_{\text{bare}} \sim \gamma_1^2$ *is unsupported, and the proposition is withdrawn as a derivation (v4.8.0-r05).* Archived in the v4.9.0 revision; original text in §A.270.

42.13 Main theorem: $\Lambda_{\text{fundamental}} = 0$ in the photon frame

Theorem 42.19 (Vanishing fundamental Λ in the photon frame)**.** *In the photon frame* ($z^- = 0$, $\beta_{\text{sym}} = 0$)*:*

(i) The frame term $\mathcal{L}_{\text{frame}} = \beta_{\text{sym}}^2\langle f, f\rangle$ *vanishes identically, so no cosmological constant is contributed by the observer offset.*

(ii) The Sphaera-cascade Markov process on $\{S^2, S^4, S^8\}$ *has a photon-frame absorption probability* $P(S^8 \to S^2) = 1 - \beta_{\text{sym}}^2 = 1$ *(in the photon frame where* $\beta_{\text{sym}} \to 0$*), yielding the structurally pure photon-frame collapse regime. In particular, the matter-displacement contribution vanishes at the level of the fundamental frame statement.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(iii) The observed cosmological constant $\Lambda_{\text{obs}} \sim \beta_{\text{sym}}^2$ *is therefore a pure* matter-frame measurement artifact*:*

$$\Lambda_{\text{obs}} = \beta_{\text{sym}}^2 \approx 0.302 \text{ (in units of } \ell_P^{-2}),$$

arising from the permanent displacement of matter observers at $z^- = +\beta_{\text{sym}}$ *from the photon ground shell* $z^- = 0$*.*

Proof. (i) In the photon frame, the Lagrangian term $\mathcal{L}_{\text{frame}} = \beta_{\text{sym}}^2 \langle f, f \rangle$ vanishes since $\beta_{\text{sym}} = 0$ by definition of the photon frame. Its variation $\delta\mathcal{L}_{\text{frame}}/\delta g^{\mu\nu} = \beta_{\text{sym}}^2 g_{\mu\nu}$ therefore contributes zero to the field equation. This is the statement that the fundamental cosmological constant vanishes in the photon frame.

(ii) The Markov transition matrix of the cascade $\hat{K} : S^8 \to S^4 \to S^2$ is

$$P = \begin{pmatrix} 1 & 0 & 0 \\ 1 & 0 & 0 \\ 1 - \beta_{\text{sym}}^2 & \beta_{\text{sym}}^2 & 0 \end{pmatrix}.$$

In the limit $\beta_{\text{sym}} \to 0$: $P(S^8 \to S^4) = \beta_{\text{sym}}^2 \to 0$ and $P(S^8 \to S^2) = 1 - \beta_{\text{sym}}^2 \to 1$. All cascade flux collapses directly to the photon ground state.

(iii) A matter observer at $z^- = +\beta_{\text{sym}}$ reads off the frame term as $\Lambda_{\text{obs}} = \beta_{\text{sym}}^2/\ell_P^{(\text{matter})2}$ via Proposition 18.4 of the main paper. This is not a vacuum energy density but the metric coefficient of the permanently shifted frame. Since $\beta_{\text{sym}}^2 \approx (0.5493)^2 \approx 0.302$, and the observed cosmological constant is $\Lambda_{\text{obs}} \approx 1.1 \times 10^{-52}\,\text{m}^{-2}$, the Sphaera prediction is dimensionally correct up to the $\gamma_L^{3/2}$ Planck-length correction of Theorem 7.6. □

Status (v4.9.0 revision): (i) holds by the definition of the reference frame R, the pure state (v4.9.0-r24); at the photon proper the frame term is maximal; (iii) is off by 10^{121} (v4.8.0-r03 F8, F9).

42.14 The open quantitative balance

Theorem 42.19 establishes $\Lambda_{\text{fundamental}} = 0$ at the *structural* level. The remaining open step is the *quantitative* Θ–R balance.

Definition 42.20 (Photon-frame Θ-R balance condition)**.** *The* Θ-R *balance in the photon frame is the condition*

$$\int_0^{\ell_P^{(\text{photon})}} \left(1 - \Theta^{(\text{photon})}(\tau) + R^{(\text{photon})}(\tau)\right) d\tau \;=\; \beta_{\text{sym}}^2 \cdot \ell_P^{(\text{photon})2}, \tag{17}$$

where $\Theta^{(\text{photon})}$ *and* $R^{(\text{photon})}$ *are the photon-frame unfolded versions of* (14)–(15)*, with the photon-frame density* $\rho_{\text{photon}}(\gamma) = \log(\gamma/2\pi\gamma_L)/(2\pi\gamma_L)$ *replacing the matter-frame density.*

Remark 42.21 (Why this is not yet proved)**.** *Condition* (17) *says: after photon-frame unfolding, the net integral of* $\Sigma^{(\text{photon})}$ *equals the frame-offset residual* β_{sym}^2 *— nothing more, nothing less. The* 10^{-122} *refers to the ratio* $\beta_{\text{sym}}^2 \ell_P^2/\Lambda_{\text{bare}} \approx 0.302/(200 \cdot \ell_P^{(\text{matter})2}/\ell_P^{(\text{photon})2})$ $= 0.302 \cdot \gamma_L^{-3}/200 \approx 0.302/1298 \approx 2.3 \times 10^{-4}$ *in dimensionless Planck units, not* 10^{-122} *in SI units.*

The 10^{-122} *in SI units arises because* ℓ_P^{-2} *in SI is* $3.8 \times 10^{69}\,\text{m}^{-2}$*, while* $\Lambda_{\text{obs}} \approx 10^{-52}\,\text{m}^{-2}$*, giving ratio* $\approx 10^{-122}$*. In natural units (*$\ell_P = 1$*),* $\Lambda_{\text{bare}} \sim \gamma_1^2 \approx 200$ *and* $\Lambda_{\text{obs}} \sim \beta_{\text{sym}}^2 \approx 0.30$*, so the ratio is* $\approx 1.5 \times 10^{-3}$ *— much less dramatic. The* 10^{-122} *is a unit-system artifact, not the fundamental challenge.*

The fundamental challenge is to show that Θ *and* R *in* $\Sigma = 1 - \Theta + R$ *cancel the bare* $\Lambda \approx 200$ *down to* $\beta_{\text{sym}}^2 \approx 0.30$ *with relative accuracy* $\sim 10^{-3}$*, uniformly in the photon-frame integration.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Status (v4.9.0 revision): P4 as stated is contradicted (v4.8.0-r03 F4); the type of Λ is open (F3); the Θ-sum route is false (v4.8.0-r05).

Proposition 42.22 (Two open sub-steps)**.** *Condition* (17) *is equivalent to the conjunction of:*

(A) ***Θ-sector remainder control.*** *The partial sum* $\Theta_N^{(\mathrm{photon})}(\tau) := \sum_{n=1}^{N} e^{-i\tau\gamma_n^{(\mathrm{photon})}}$ *satisfies, uniformly for* $0 \le \tau \le \ell_P^{(\mathrm{photon})}$*:*

$$\left|\Theta^{(\mathrm{photon})}(\tau) - \Theta_N^{(\mathrm{photon})}(\tau)\right| \;\le\; \varepsilon_N \to 0 \text{ as } N \to \infty,$$

and $\int_0^{\ell_P^{(\mathrm{photon})}} \Theta^{(\mathrm{photon})}(\tau)\, d\tau$ *converges absolutely.*

(B) ***R-sector exact cancellation.*** *The smooth term* $R^{(\mathrm{photon})}(\tau) = -e^{-\tau/2}/(e^{\tau}-1)$ *(photon-frame rescaled) and the sum* $\Theta^{(\mathrm{photon})}(\tau)$ *together cancel the constant* 1 *in* Σ *to within* β_{sym}^2 *when integrated over* $[0, \ell_P^{(\mathrm{photon})}]$*.*

Sub-step (B) is equivalent to the Guinand–Weil explicit formula evaluated at the photon-frame UV cutoff $\ell_P^{(\mathrm{photon})}$*.*

Proof of equivalence. The integral of $\Sigma^{(\mathrm{photon})}$ splits as

$$\int_0^{\ell} \Sigma\, d\tau = \ell - \int_0^{\ell} \Theta\, d\tau + \int_0^{\ell} R\, d\tau,$$

where $\ell = \ell_P^{(\mathrm{photon})}$. Condition (A) ensures the Θ-integral is well-defined and controllable. Condition (B) then says the three terms sum to $\beta_{\mathrm{sym}}^2 \cdot \ell^2$. Both conditions are necessary and sufficient for (17). □

Theorem 42.23 (Structural Θ-R cancellation — partial result)**.** *In the matter frame, the Guinand–Weil formula gives exactly*

$$\Sigma(\tau) = 1 - \Theta(\tau) + R(\tau),$$

which encodes complete cancellation of the pole contribution by the combined Θ *and* R *terms* at the distributional level *(for all* $\tau > 0$*, without UV cutoff). The cosmological constant problem arises solely from introducing a finite UV cutoff: at* $\tau = \ell_P^{(\mathrm{matter})}$*, the integral of* Σ *is*

$$\int_0^{\ell_P^{(\mathrm{matter})}} \Sigma\, d\tau = \ell_P^{(\mathrm{matter})} - \Theta_{\mathrm{UV}} + R_{\mathrm{UV}} + \beta_{\mathrm{sym}}^2\, (\ell_P^{(\mathrm{matter})})^2,$$

where Θ_{UV} *and* R_{UV} *are the truncated integrals. The* β_{sym}^2*-term is the frame residual. The mismatch* $\ell_P^{(\mathrm{matter})} - \Theta_{\mathrm{UV}} + R_{\mathrm{UV}}$ *is the open step (B) in the matter frame.*

Proof. The Guinand–Weil explicit formula holds distributionally for all $\tau > 0$ (Theorem 3.3 of [1]). The Θ-integral converges because $\Theta(\tau) \to 1$ as $\tau \to 0^+$ along the spectral sum (distributional convergence). The exact cancellation $1 - \Theta(\tau) + R(\tau) = \Sigma(\tau)$ holds for all $\tau > 0$ by the explicit formula. The finite-UV-cutoff breakdown is the statement that $\int_0^{\Lambda} \Sigma\, d\tau \neq 0$ for finite Λ, which is the content of the open sub-steps. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

42.15 Falsifiable predictions

Corollary 42.24 (Ranked falsifiable predictions)**.** *The following predictions follow from the framework and are ordered by near-term experimental accessibility.*

P1 (testable now): GUE improvement on Odlyzko data.
Photon-frame unfolding of Odlyzko's $\sim 10^6$ *Riemann zeros at height* $\sim 10^{22}$ *predicts a* χ^2 *improvement of* $\approx 21\%$ *relative to standard unfolding (Corollary 24.6, Theorem 24.2). Non-observation falsifies* $\gamma_L = \gamma_{\mathrm{BI}}/\gamma_{\mathrm{string}}$ *as the physical frame factor.* Status: testable with Odlyzko's public dataset using existing code.

P2 (already matches data): Koide phase formula.
$\theta_K = \pi - \sqrt{2}\arcsin(\beta_{\mathrm{sym}}) \approx 132.88°$*, matching the PDG value* $132.73°$ *to* 0.111% *(Theorem 22.5).* Status: confirmed against existing PDG lepton masses.

P3 (matches to 0.018%): β_{sym}***-formula.***

$$\beta_{\mathrm{sym}} = \frac{1}{\sqrt{3}} \exp\left(-\frac{\gamma_1}{4\pi\gamma_L^5}\right) \approx 0.5492,$$

matching the measured value 0.5493 *to* 0.018% *(Remark 41.8, corrected in archival stage).* Status: consistent with CODATA. Structural derivation of the γ_L^5 exponent requires OP 2.

P4 (reinterpretation): Cosmological constant.
$\Lambda_{\mathrm{obs}} = \beta_{\mathrm{sym}}^2 \cdot (\ell_P^{(\mathrm{photon})})^{-2}$ *is a frame artifact, not a vacuum energy (Theorem 42.19). In natural Planck units the apparent* 10^{-122} *fine-tuning reduces to a factor* ~ 3.1 *cancellation (Remark 42.21). Matter-frame computations carry a systematic error of* $\gamma_L^5 \approx 22.5$ *(Corollary 42.17).* Status: reinterpretation of existing data; no new measurement needed.

P5 (annual modulation, future): α***-drift ratio.***
Constants with different frame-insertion counts k *should modulate differently with annual orbital period. Specifically, the ratio of annual variations:*

$$\left.\frac{\delta\alpha/\alpha}{\delta G_N/G_N}\right|_{\mathrm{annual}} = \frac{k(\alpha)}{k(G_N)} = \frac{1}{2},$$

whereas standard physics predicts this ratio $= 1$*. The deviation is* 50% *but requires* G_N *measurements at* $\sim 10^{-18}$ *annual precision — beyond current capability.* Status: prediction well-defined but not yet testable. The modulation of α alone (period exactly 1 year, in phase at all terrestrial labs, absent for non-orbiting reference clocks) is a qualitative signature at amplitude $\sim 10^{-10} \cdot \delta\Phi/c^2$.

P6 (theoretical): Hawking temperature in tachyon frame.
$T_{\mathrm{Sphaera}} = \gamma_{\mathrm{BI}} \cdot T_H^{(\mathrm{tachyon})}$*, giving* γ_{BI} *the interpretation as the ratio of the Sphaera Hawking temperature to the tachyon-frame Hawking temperature (Proposition 21.4).* Status: theoretical prediction; requires quantum-gravity experiment to test directly.

4.10 consolidation: The list is a historical prediction register, not a set of independently validated 4.10 predictions. In particular, the GUE scale argument cites an archived result; apparent agreement and blind prediction must be distinguished. The new comparison results do not promote these empirical claims.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Corrected in the v4.9.0 revision (P4: $\gamma_L^5 \approx 22.5$, earlier $\gamma_L^6 \approx 42$; P5: 1/2, earlier $\gamma_L^{-1} \approx 0.537$). Status of P1–P6: P1 not confirmed, archived (v4.8.0-r08); P2 phase formula excluded, archived (v4.8.0-r09); P3 coincidence, archived (v4.8.0-r09, v4.9.0-r19); P4 contradicted as stated (v4.8.0-r03 F4); P5 corrected to 1/2, untestable at present (v4.9.0-r09); P6 untested, assumption $k(M) = 0$ explicit (v4.8.0-r03 F6).

Remark 42.25 (What the predictions share)**.** *All six predictions follow from the same structural core: the frame offset $z^- = +\beta_{\text{sym}}$ and the frame factor $\gamma_L = \gamma_{\text{BI}}/\gamma_{\text{string}}$. None require free parameters beyond these two quantities, both of which are determined by the framework (P3 and the LQG/string coupling values). The only currently testable prediction that requires new computation is P1 (GUE improvement); P2 is already a post-diction on existing data.*

42.16 Status summary

Reader cue. Read the next table as a burden/status dashboard for the OP 5 line. Its purpose is to separate what is already structurally fixed, what remains open, and which analytic step each open item still depends on.

Statement	Proof basis	Status
$\Lambda_{\text{fundamental}} = 0$ in photon frame	Cascade Markov + $\beta_{\text{sym}} = 0$ in photon frame	✓(Thm. 42.19)
$\Lambda_{\text{obs}} = \beta_{\text{sym}}^2$ (frame artifact)	$\mathcal{L}_{\text{frame}}$ variation	✓(Prop. 18.4 of main paper)
Matter-frame error $= \gamma_L^5 \approx 22.5$	$k(\ell_P) = 3/2$, Thm. 7.6	✓(Cor. 42.17)
Guinand–Weil: $\Sigma = 1 - \Theta + R$ distributionally	Explicit formula, Thm. 3.3	✓(Thm. 42.23)
Θ-sector remainder control at $\ell_P^{(\text{photon})}$ (step A)	—	**open**
R-sector exact cancellation at photon-frame cutoff (step B)	—	**open**
$k(\Lambda)$ self-consistent determination	—	**open**

Remark 42.26 (What OP 5 actually requires)**.** *The two open steps are not independent: step (B) is the photon-frame version of the Guinand–Weil formula evaluated at a* finite *UV cutoff. This is a standard problem in analytic number theory (partial explicit formulas with error terms). The Sphaera-framework contribution is: (a) identifying the correct frame ($\ell_P^{(\text{photon})}$, not $\ell_P^{(\text{matter})}$), and (b) replacing the observable Λ with the frame residual β_{sym}^2. The hard analysis — bounding the truncation error of the zero sum at height $T = 1/\ell_P^{(\text{photon})}$ — is the remaining mathematical content of OP 5.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

42.17 Quantitative analysis of the open steps (archival stage)

Step (A): Θ-remainder control

Proposition 42.27 (Step (A) is trivially satisfied in the photon frame)**.** *The photon-frame Θ-remainder control of sub-step (A) is satisfied trivially:*

$$\Theta_N^{(\text{photon})}(\tau) = \Theta^{(\text{photon})}(\tau) = 0 \qquad \text{for all } \tau \in [0,\, \ell_P^{(\text{photon})}].$$

Consequently $\varepsilon_N = 0$ for all N in this range.

Proof. The photon-frame UV cutoff is $T_{\text{photon}} = 1/\ell_P^{(\text{photon})} = \gamma_L^{3/2} \approx 2.545$ (in matter-Planck units). The photon-frame Riemann zeros are $\gamma_\rho^{(\text{photon})} = \gamma_\rho/\gamma_L$, with first value $\gamma_1^{(\text{photon})} = 14.1347/1.864 \approx 7.583$. Since $T_{\text{photon}} \approx 2.545 < \gamma_1^{(\text{photon})} \approx 7.583$, no photon-frame zero lies below the UV cutoff. Therefore, for $\tau \in [0, \ell_P^{(\text{photon})}]$, the oscillatory sum $\Theta^{(\text{photon})}(\tau) = \sum_\rho e^{-i\tau\gamma_\rho^{(\text{photon})}}$ has no contributing terms in the spectral range $[0, T_{\text{photon}}]$. Hence $\Theta^{(\text{photon})}(\tau) = 0$ pointwise in the integration domain, and step (A) is satisfied with $\varepsilon_N = 0$. □

Remark 42.28 (Physical significance: the photon frame is below the first zero)**.** *The fact that $T_{\text{photon}} < \gamma_1^{(\text{photon})}$ is a structural consequence of the frame factor $\gamma_L \approx 1.864$: in the photon frame, the UV cutoff $\ell_P^{(\text{photon})}$ is* larger *than in the matter frame by $\gamma_L^{3/2}$, pushing the cutoff frequency T_{photon}* below *the energy of the first Riemann zero. This is precisely the regime where the spectrum is quiet — the photon frame sees no Riemann zero oscillations below its Planck scale. The cosmological constant problem vanishes in this regime because Θ contributes nothing to the vacuum energy integral.*

Step (B): R-sector cancellation

With step (A) resolved, the balance condition (17) reduces to:

$$\int_0^{\ell_P^{(\text{photon})}} \left(1 + R^{(\text{photon})}(\tau)\right) d\tau \;=\; \beta_{\text{sym}}^2 \cdot (\ell_P^{(\text{photon})})^2. \tag{18}$$

Remark 42.29 (Regularization required; corrected in 4.10)**.** *For dimensionless $\tau > 0$,*

$$R(\tau) = -\frac{e^{-\tau/2}}{e^\tau - 1} = -\frac{1}{\tau} + 1 - \frac{11\tau}{24} + O(\tau^2).$$

The one-sided endpoint singularity requires a specified finite-part prescription, not an ordinary convergent integral. For a positive dimensionless reference scale μ, define

$$\mathrm{FP}_\mu \int_0^T (1 + R(\tau))\, d\tau = \lim_{\varepsilon \downarrow 0} \left[\int_\varepsilon^T (1 + R(\tau))\, d\tau - \log(\varepsilon/\mu) \right].$$

A direct antiderivative gives

$$T - 2e^{-T/2} + \log \frac{1 + e^{-T/2}}{1 - e^{-T/2}} + 2 - \log 4 + \log \mu.$$

For $T = 0.393$ and $\mu = 1$ this is approximately 1.686953464*. Changing μ changes the finite part. Selecting this scale or interpreting the result as physical vacuum energy needs a separate action, units and matching prescription.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

4.10 correction: the earlier expansion $-1/\tau + 1/2 - \tau/12$, the added $+\log\varepsilon$ counterterm, and the value 0.940 were incorrect. The added counterterm reinforced the endpoint divergence. The withdrawn Theta-balance route is not reinstated by this correction.

Proposition 42.30 (Step (B): historical target withdrawn)**.** *The former numerical target* $0.893 = 0.940 - 0.302 \cdot 0.154$ *depended on the incorrect finite-part evaluation in Remark 42.29. It is not an established requirement on a physical Theta contribution. The subsequent balance conjecture is already withdrawn in Proposition 42.33. A corrected finite-part convention alone neither restores that balance nor closes OP5.*

4.10 correction: the preceding target is retained here only as a historical dependency. No new cancellation target is inferred.

Remark 42.31 (Current status of the historical OP5 route)**.** *The former numerical summary using* 0.940 *and* 0.893 *is superseded by the finite-part correction above and the already withdrawn Theta-sum conjecture. A dimensionless discrepancy is not removed by a change of units. The newer state, clock and field comparisons in Section 44 identify conditional structures and remaining compatibility conditions; they do not prove cancellation of a cosmological term. OP5 remains open.*

Step (B) reduced: the exact Theta-sum conjecture

The numerical analysis of archival stage yields a precise formulation of what step (B) requires. Define:

Definition 42.32 (Photon-frame Theta sum)**.** *Let* $T = \gamma_L^{-3/2}$ *(the photon-frame UV cutoff in matter-Planck units) and let* $\gamma_n^{(\text{photon})} = \gamma_n/\gamma_L$ *denote the photon-frame Riemann zeros. The* photon-frame Theta sum *is the conditionally convergent series*

$$\mathcal{T}(T) \;:=\; 2\sum_{n=1}^{\infty} \frac{\sin\!\big(T \cdot \gamma_n^{(\text{photon})}\big)}{\gamma_n^{(\text{photon})}}.$$

Proposition 42.33 (Theta-sum balance conjecture; archived)**.** *The conjectured value* $\mathcal{T}(T) \approx 0.893$ *is false. The sum is* -0.1097 *(v4.8.0-r05), and OP 5 Step (B) has no valid form on this route.* Archived in the v4.9.0 revision; original text in §A.270.

Remark 42.34 (Structure of the conjecture)**.** *The series* $\mathcal{T}(T)$ *is the photon-frame analogue of the oscillatory sum in the Riemann–von Mangoldt explicit formula for the prime-counting function* $\pi(x)$*. It converges* conditionally *(not absolutely), and its value at a specific argument* T *is a non-trivial statement about the Riemann zeros.*

Numerically, $\mathcal{T}(T)$ *converges slowly: with* $N = 50$ *zeros the partial sum is* ≈ -0.146*, far from the conjectured* 0.893*. Convergence requires summing the full tail* $\sum_{n>50}$ *whose individual contributions are small but whose cumulative effect is large (this is the hallmark of conditional convergence).*

The conjecture is equivalent to *the balance condition* (17) *once step (A) is established (Proposition 42.27). Establishing Conjecture 42.33 would therefore close OP 5 completely.*

4.10 consolidation: This paragraph records the dependency of the withdrawn historical conjecture, not a live OP5 route. Its old target is superseded by Remark 42.29; the conjecture remains archived.

Remark 42.35 (Remainder estimate; archived)**.** *The remainder estimate that was to support Proposition 42.33 is withdrawn with it (v4.8.0-r05).* Archived in the v4.9.0 revision; original text in §A.270.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

43 Current v4 Interface and Open Burdens

This synthesis prevents the retained early material from becoming the default status of the later v4 line. It is a reading interface, not a replacement for the original hypotheses or finite computations.

43.1 Shifted Normal Form and Backtranslation

The later governing object is the shifted normal-form record of Definition A.34.1. New objects must retain occurrence class, payload/readout target, backtranslation, guard and status. In particular packed masks must be reconstructed from shifted supports rather than shifted as whole integers; physical payloads must preserve their residual map and scope. The full migration chain remains in Appendix A.27. Older pre-shift wording in Appendix H2 is historical input to this chain, not authorization to undo it.

43.2 Terminal Data and Exact Zero

The guarded solve-chain record in Appendix A.266 distinguishes local C4 seam control, finite C5 detector terminality and exact all-readout closure. The inherited finite ledger records detector threshold $\theta = 1/256$ after twenty steps; it does not state that the native residual is exactly zero at a finite step. These recorded numbers are not newly recovered native measurements in the present editorial pass.

The frame-typed endpoint in Appendix A.268 makes the distinction explicit: matter-frame effective terminal data, a candidate photon-clock null object and tachyon/information-facing signed interval data are different types. None provides an automatic implication to exact all-readout zero. A photon/clock sector is not thereby a physical rest frame of a photon, and the terminology alone supplies no inter-sector signal law.

43.3 Interface Certificates and Dynamics

The later MM/HC development passes through typed coefficients, seed residuals, operator-domain records, local monitor certificates and no-action/no-promotion checks. Their full dependency order is retained in Appendix A.210. An admissibility tag does not define the action of a differential operator. A local interface certificate is not a variational action or a derived field equation.

For the proposed v4.8 metric program this leaves four concrete burdens: declare scale units and comparison maps; realize the MM/HC operators with their spectral/domain premises; evaluate the native residual dictionary with fixed norms; and supply a dynamical realization with its approximation and stability bounds. The earlier QAM/router construction can record these objects but cannot replace them.

Section 9 now supplies explicit conditional calculations rather than a dynamical identification: projected hidden states and memory, conservative collective inertia, the null HC kinetic tensor, real-action/current alternatives and an off-shell geometry/stress match. Proposition 9.5 excludes the direct same-induced-frame real-bilinear master/coframe identification under its stated assumptions. A separate microscopic/effective geometry remains possible but needs its own common action and current balance.

Section 10 now supplies a declared neutral Cartan comparison with a finite normal inverse and a conditional source law. Preserving compatible tangential jets determines the six source values at the studied derivative order; unrestricted higher jets can instead absorb them.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Neither fact establishes arbitrary surface admissibility, the full regional equations or the native MM/HC identification.

44 States, Mediators and Clock Comparisons in the 4.10 Research Line

This section consolidates the investigations of 28–30 October 2026. It distinguishes exact local identities, finite numerical evidence and open physical identifications. The common carrier is a research programme; the comparisons below neither close OP5 nor derive a unified quantum/gravity action. The QAM records the types, hypotheses and remaining obligations of an argument. An open entry is not a refutation, and a successful residual test is not a substitute for the missing implication.

44.1 A common carrier and distinct readouts

Let $\mathcal{C}$ be a declared common state carrier and let $\pi_Q : \mathcal{C} \to \mathcal{Q}$ and $\pi_R : \mathcal{C} \to \mathcal{R}$ be quantum and relativistic readouts on specified admissible domains. A mediator $\mathcal{M}$ requires explicit maps m_Q, m_R with

$$m_Q \pi_Q(c) = m_R \pi_R(c).$$

If dynamics are also claimed, the additional conditions are $\pi_Q U_{\mathcal{C}} = V_Q \pi_Q$ and $\pi_R U_{\mathcal{C}} = V_R \pi_R$, with compatible clock parameters. These are proof obligations, not definitions that establish a physical identification by themselves. A physical state space, a local null-direction sphere, and a normalized two-port register retain different information.

In particular, the ordinary positive AB Gram register of two Dirac spinors is not the same readout as the real electromagnetic doublet of r04. Writing $N = \|A\|^2 + \|B\|^2 > 0$, the former gives

$$\rho_{AB} = \frac{1}{N} \begin{pmatrix} A^\dagger A & B^\dagger A \\ A^\dagger B & B^\dagger B \end{pmatrix}, \qquad r = \frac{(2 \operatorname{Re} A^\dagger B, 2 \operatorname{Im} A^\dagger B, \|A\|^2 - \|B\|^2)}{N}.$$

It obeys $1 - |r|^2 = 4 \det \rho_{AB} \geq 0$. Proportional AB fields lie on the boundary; independent fields can lie in the interior. Two fields with the same r can have opposite $A^\dagger \beta A - B^\dagger \beta B$: take $(A, B) = (\cos\theta\, e_1, \sin\theta\, e_1)$ or $(\cos\theta\, e_3, \sin\theta\, e_3)$ with $\beta = \operatorname{diag}(1, 1, -1, -1)$. Thus the Bloch coordinates alone do not close the Dirac readout.

44.2 An orientation-complete local mediator

For the native real doublet $F = [E \mid B]$, put $G = F^{\mathsf{T}} F = \left(\begin{smallmatrix} a & c \\ c & b \end{smallmatrix}\right)$. On the chart $a > 0$, $\Delta = \sqrt{ab - c^2} > 0$, define

$$F_0(G) = \begin{pmatrix} \sqrt{a} & c/\sqrt{a} \\ 0 & \Delta/\sqrt{a} \\ 0 & 0 \end{pmatrix}, \quad e = E/\sqrt{a}, \quad q = \frac{B - (c/a)E}{\Delta/\sqrt{a}}, \quad O = [e \mid q \mid e \times q].$$

Then $O \in SO(3)$ and $F = O F_0(G)$, by direct multiplication. The three Gram entries and three orientation parameters therefore recover the six real field components locally. Keeping only $\rho = G/\operatorname{tr} G$ also discards the scale $\operatorname{tr} G = 2u$. Rank-deficient fields need other charts and stabilizer data. A nonzero null electromagnetic field with $G = uI$ is regular in this rank-two chart; it is not a rank-one Gram state.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

For any declared right-doublet operation M,

$$F' = FM, \qquad G' = M^{\mathsf{T}} G M.$$

Reconstructing $F_0(G)M = KF_0(G')$ gives $O' = OK$ whenever the rank-two charts exist. This proves that changing the field and then reading it agrees with the corresponding mediator operation. Exact checks cover phase rotation, $J_B = i$ times conjugation, conjugation and the compensator $C_+ = (I + \sigma_x)/2$.

The compensator is a separate rank-changing case: $E' = B' = (E + B)/2$. If nonzero, its normalized Gram state is $\rho' = C_+$, and a rank-one direction chart replaces the orientation triad. If the selected port vanishes, normalization is undefined. The pure selected register state is not the same readout as an electromagnetic null field, whose r04 state is $I/2$.

Orientation is needed for general observer changes. For example, $E = (2, 0, 0)$, $B_\pm = (0, \pm 1, 0)$ give the same $\rho = \mathrm{diag}(4/5, 1/5)$, but opposite fluxes. Under a common infinitesimal boost of speed v along z,

$$\left.\frac{d\rho_+}{dv}\right|_0 = \mathrm{diag}(12/25, -12/25) = -\left.\frac{d\rho_-}{dv}\right|_0 .$$

Even the flux alone does not repair every ambiguity: two orthogonal pure electric fields have the same Gram matrix and zero flux but respond differently to a boost along one of their axes. These facts establish non-autonomy of the reduced register under unrestricted directed observer changes, not a failure of the full field's covariance.

44.3 Clock calibration and the octonionic scope

Coordinate time, eigen-time along a timelike observer, an internal clock reading, and a structural invariant require separate definitions. In the controlled constant-rate example $\psi_a(t_a) = \exp(-i\omega a t_a)\psi_0$, $\psi_b(t_b) = \exp(-i\omega b t_b)\psi_0$, $a, b > 0$, the calibration $\tau = at_a = bt_b$ makes the trajectories agree exactly. Matching their initial zeros alone does not. The rates are inputs of this example, not derived native clocks. The orientation counterexample above persists under a common calibrated clock whenever the observer's boost rate is nonzero. Neither result requires a second fundamental quantum time.

A shared clock does not alone prove entanglement: local evolution $U_1(\tau) \otimes U_2(\tau)$ preserves product states. Likewise, an invariant τ_n is not by itself an operational clock or an entanglement criterion. Quantum systems can also be massive; the two descriptions are not disjoint massless/massive classes.

After choosing a complex direction, a six-real-dimensional octonionic complement can encode $E + iB$. This is a data embedding, not yet an identification of Lorentz boosts with real octonion automorphisms. Those preserve the positive octonion norm, whereas a transverse boost of $E = (0, 1, 0), B = 0$ gives $E'^2 - B'^2 = 1$ but $E'^2 + B'^2 = (1 + v^2)/(1 - v^2)$. The native complex/quaternionic/octonionic dictionary and its allowed actions must therefore be specified. Extra registers or complexification are possible research routes, not consequences of dimension counting.

44.4 Field compatibility and finite evidence

The declared classical comparison retains

$$F_\Psi = i \sum_{\mu,a} (E^{-1})^\mu{}_a (I \otimes K_a) D_\mu \Psi - h(Y \otimes \beta)\Psi = 0,$$

$$E^a{}_\mu = \mathrm{Re}(\Psi^\dagger Q_a D_\mu \Psi), \qquad \det E \neq 0.$$

where $Q_a = S \otimes K_a$, $K_a = (I, \alpha_i)$, and the Lorentz spin connection is set to zero as a compar-

© 2026 Marc Brendecke. Preprint — not peer reviewed.

ison choice, not identified with the induced Levi–Civita connection. The source couplings are held fixed. The earlier proportional AB ansatz need not be preserved by this dynamics. Releasing the first jets removes several pointwise residuals, but does not establish their regional continuation.

For constant Hermitian Q_a and Hermitian internal connection,

$$E^a = \tfrac{1}{2} dj_a + \tfrac{i}{2} \Psi^\dagger [Q_a, A] \Psi, \qquad j_a = \Psi^\dagger Q_a \Psi.$$

For the retained $A = G\rho\, d\varphi$ on $\rho > 0$, this implies $E^a = dj_a/2 + B_a d\varphi$ and $dE^a = dB_a \wedge d\varphi$. Here the radial coordinate ρ is unrelated to a density matrix. This coordinate curl is not itself Riemann curvature. Eliminating all free ordered second jets shows numerically that imposing even the full current-curl identity does not reproduce the obstruction from fully symmetric spinor second jets. Coframe integrability and spinor integrability are different tests.

At fixed field values let A_0 be the real map from time-jet changes to changes of E_0, and B_0 have the four covariant jets as real columns. Then $A_0 B_0 = E$. Pointwise shear invariance gives, for invertible frozen time operator H_0,

$$P = I - B_0 E^{-1} A_0, \quad J_0 = H_0 P, \quad L = A_0 H_0^{-1}, \quad L J_0 = 0.$$

Thus $LF_\Psi = 0$ supplies four temporal compatibility conditions. One combination is $4 - h\Psi^\dagger (Z \otimes \beta)\Psi/2 = 0$, $Z = \mathrm{diag}(1, -1, 0)$. Its derivatives and the other compatibility combinations can be solved jointly at the three tested points. The resulting first-jet family has a numerically regular seven-direction constraint Jacobian and 65 free spatial tangent directions, before further constraints and identifications. This is neither a count of physical degrees of freedom nor a regional existence theorem. For these frozen first jets, symmetric second-jet completion retains nonzero residuals; the joint recalibration below releases those data and supersedes this as a current obstruction.

Two-mode fields with variable amplitudes provide single smooth approximations and hence consistent mixed derivatives, but retain nonzero equation residuals. Their optimization is underdetermined and does not supply a common action. Figure 1 deliberately records the original affine baseline, not the latest fit. Later quadratic fits and weighted comparisons are archived separately. The early edge-weighting interpretation was corrected when a reused residual function was found to retain the old points; only the subsequent corrected comparisons support statements about edge weighting.

As a precision control, the direct centre-point problem was recomputed with necessary state normalization restored in each precision:

decimal digits	normalized-input full residual	rounded-input full residual
30	2.3×10^{-30}	2.9×10^{-15}
50	6.0×10^{-50}	2.9×10^{-15}
80	4.5×10^{-80}	2.9×10^{-15}

These are numerical checks, not certified exact solutions or physical scales. Renormalization changes the rounded input by a tiny amount; extra arithmetic precision alone cannot remove a frozen input defect. The larger continuation residuals are not removed by this point test.

44.5 Recursive finite-jet recalibration

The preceding frozen-data failures do not persist under all joint changes of lower jets. A subsequent comparison holds the same field values, original backgrounds and couplings fixed and first solves for compatible first and symmetric second jets. Releasing second jets then

© 2026 Marc Brendecke. Preprint — not peer reviewed.

allows a symmetric third-jet completion while retaining those first jets. The scalar condition $g = 4 - h\Psi^\dagger(Z \otimes \beta)\Psi/2 = 0$ explains a necessary part of the hierarchy: the first fit enforces g, dg, but its second jets need not satisfy $d^2 g = 0$. Free third jets cannot repair that lower-data defect. The joint second/third fit does repair it numerically.

A reusable construction stores Taylor coefficients $c_\alpha = \partial^\alpha \Psi/\alpha!$, with multiindex factorial. At each tested level the highest jets z_p enter linearly:

$$A_p z_p + b_p(J_{<p}) = 0, \qquad L_p A_p = 0.$$

A complete left kernel eliminates z_p. The numerical search adjusts allowed lower jets to satisfy $L_p b_p = 0$ together with every retained lower-order residual, then recovers z_p and checks the full equations. This is a construction algorithm with a possible failure outcome, not an all-order existence assertion. Small updates and coefficient guards do not prove convergence or uniqueness.

For the fourth-order test, field values and all first jets remain fixed; second and third jets vary and fully symmetric fourth jets are recovered. The highest matrix has size 480×840, numerical rank 424, and 56 left compatibility directions in the declared coefficient packing. Five accepted steps at each of three points produce the raw symmetric derivative norms below:

point	$\|F\|$	$\|dF\|$	$\|d^2F\|$	$\|d^3F\|$
centre	$3.1\,10^{-14}$	$3.9\,10^{-14}$	$2.4\,10^{-13}$	$6.4\,10^{-12}$
edge	$2.0\,10^{-15}$	$2.9\,10^{-14}$	$2.5\,10^{-13}$	$5.4\,10^{-12}$
inner	$2.6\,10^{-15}$	$3.8\,10^{-14}$	$2.3\,10^{-13}$	$6.5\,10^{-12}$

Normalized coefficient norms and raw derivative norms are stored separately. A separate NumPy directional-series evaluator with a recurrence inverse checks coefficients through degree three in six directions per point, with discrepancies below $4\,10^{-13}$. Selected highest-operator columns and scalar constraint coefficients are also checked. This is bounded implementation evidence, not a certified exact root or an exhaustive theorem check.

Raw second-, third- and fourth-jet norms are approximately $26, 89, 900$ in the retained unscaled coordinates. They establish no physical scale and no uniform growth estimate. The three germs are not shown to belong to one regional solution; stabilization of lower coefficients across levels, convergence, regional existence and native physical identification remain open. The tested reusable construction and earlier failed controls are archived under `analysis/op5-field-slices/`, in the dated state-sufficiency directory. Neither these finite completions nor the algorithm closes OP5.

Status of the 4.10 consolidation. Exact reconstruction and local operation identities are separated from finite numerical evidence. Native action, clock identification, complete octonionic structural embedding, regional spinor existence and OP5 closure remain open. Scripts, inputs, coefficients, corrections and individual scope statements are in `analysis/op5-field-slices/`. The guard and global residual records are not promoted by these tests.

44.6 OP5 comparison field: sampled multimode geometry

This figure records the original affine baseline of 29 October 2026. Later quadratic fits are discussed separately in Section 44.4; they do not replace these data. As a diagnostic in the declared classical comparison sector, consider a sum of two exponential modes with independent affine complex spinor amplitudes. A common positive amplitude factor enforces the

© 2026 Marc Brendecke. Preprint — not peer reviewed.

necessary scalar condition $h\Psi^\dagger(Z\otimes\beta)\Psi = 8$, where $Z = \mathrm{diag}(1,-1,0)$. This condition does not replace the full probe equation. The coframe and its inverse are recomputed from the fields; the original comparison couplings and zero Lorentz spin connection are retained. This is not an identification of the induced Levi–Civita connection.

Figure 1 evaluates one saved approximate field, without refitting, on all six coordinate-plane cuts through $(t,x,y,z) = (1,0.6,0,0)$. Each cut uses nine positions per axis with offsets ± 0.01 in comparison coordinates. These are six overlapping 9×9 slices, not a full four-dimensional grid. The largest sampled residual norm is approximately 0.03868, the smallest absolute coframe determinant is 9.6991, and every sampled g^{00} is negative. These are finite-sample diagnostics, not uniform bounds.

For the frame images E_i and E_j of coordinate directions, the oriented area is represented by the full bivector $E_i \wedge E_j$. Its signed squared area is $g_{ii}g_{jj} - g_{ij}^2$. The spatial cross product of their spatial projections supplies the spatial bivector components; time–space components must also be retained. Neither this area nor a mixed metric component alone establishes Riemann curvature.

The explicit smooth field makes mixed derivatives consistent wherever its normalization is regular, but its nonzero equation residual remains. No exact regional solution, native action derivation, or OP5 closure is claimed. The numerical record and reproduction files are archived under `analysis/op5-field-slices/`.

45 v4.8 Research Integration

This is a proposed integration layer for the research program named in the repository, not a new release theorem. The complete v4.7 baseline is retained below it. The layer separates what can be defined now from what still requires a physical or mathematical construction.

45.1 Typed Scale Record

For a declared readout R, introduce the candidate scale record

$$\ell_{\mathrm{eff}}(R) = \ell_P \lambda_R, \qquad \lambda_R \geq 0, \qquad \varepsilon_R^{\mathrm{len}} = \frac{\ell_P \lambda_R}{L_{\mathrm{ref}}}, \quad L_{\mathrm{ref}} > 0.$$

Here ℓ_P and L_{ref} have units of length, and the new λ_R is dimensionless. The positive scale magnitude must be kept separate from an orientation/signature marker σ_R. In the inherited tachyon-facing passages a symbol such as λ_{T} denotes a native scale magnitude, not automatically this dimensionless coefficient. Identifying them requires an explicit unit conversion and comparison map.

A signed interval class can therefore be recorded separately from the length magnitude. A negative signature or orientation code does not give a negative material length. Nor does a dimensionless fit-score threshold such as 1/256 equal $\varepsilon_R^{\mathrm{len}}$ without a specified calibration between the residual's norm and a length observable.

45.2 Metric Ansatz and Null Limit

If g_{ref} is a declared Lorentzian metric and λ_R is smooth and strictly positive on a region, one possible additional ansatz is

$$g_R = \lambda_R^2 g_{\mathrm{ref}}.$$

Pointwise positive multiplication preserves signature and nondegeneracy, and at a fixed point scales non-null interval magnitudes by λ_R. This elementary property does not show that

© 2026 Marc Brendecke. Preprint — not peer reviewed.



Figure 1: Original affine baseline: six 9×9 cuts of the same approximate OP5 comparison field. Columns show the full probe-equation residual norm, g^{00}, the normalized mixed metric component $g_{ij}/\sqrt{|g_{ii}g_{jj}|}$, and the Lorentzian area magnitude $\sqrt{|g_{ii}g_{jj} - g_{ij}^2|}$. A common colour range is used within each column; local numerical ranges appear below each panel. Horizontal and vertical coordinates are identified per panel. Invertibility and spacelike constant-t cuts hold at the sampled points; the residual is nonzero.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

the physical readout has this isotropic conformal form. For a varying factor, connection and curvature terms also involve derivatives of the factor; a length convention alone does not determine their dynamics.

At $\lambda_R = 0$ the proposed tensor is degenerate and is not a Lorentzian metric. A null photon trajectory instead satisfies $g(v, v) = 0$ for a nonzero tangent in a nondegenerate metric; it does not require the whole metric tensor to vanish. A photon-clock boundary therefore needs its own null/readout construction, not the substitution $\lambda_\gamma = 0$ into a material metric ansatz.

The different null-preserving coframe construction in Proposition 9.4 gives an explicit local geometric example, not a derivation of this conformal ansatz. Its isolated stress match is off shell. In particular, directly reusing its inverse coframe in the same real-bilinear ordered master equation meets the exclusion of Proposition 9.5; microscopic and effective frame data cannot be identified without this check.

45.3 From MM/HC to Dynamics

Before a metric or field equation can be claimed, specify the state space and operator domains, the readout map and its units, the represented MM and compensator, and the actual residual functions. Proposition 1.1 states one sufficient spectral interface; it does not construct the native operator or its couplings.

The next dynamical package must also provide an evolution or action, its boundary/initial conditions, the transformation of the readout under admissible changes, and the required conservation or exchange identities. Merely putting coefficient tags into geometric, kinetic and mixed slots does not derive such a package. The constant offset and nonlinear remainder in the C5 dictionary must be retained when testing a finite contraction bound.

45.4 Integration and Acceptance Gates

1. **Source and scope gate.** Keep the full-source map, shifted normal form, backtranslation and terminal/global boundary. Do not replace the later baseline with the older modular pilot.

2. **Scale gate.** Supply R, L_{ref}, units, positive magnitude, orientation/signature data and an evaluated calibration before comparing length and detector residuals.

3. **Operator and residual gate.** Supply domains, the spectral premises actually used and the native three-block residual evaluation with its fixed norms. Synthetic regression examples remain examples rather than native witnesses.

4. **Dynamical gate.** Derive the claimed equation from the declared model and prove the needed well-posedness, conservation, stability or approximation statements at the scope being claimed. A fluid or gravitational interpretation is still an interpretation until this map is constructed. The declared real action, allowed variations and target reactions must agree with the currents. The geometry/probe interface must also pass the current and pointwise trace tests in Section 9; an isolated Einstein/stress match does not solve the missing probe and normal field equations. Section 10 partially advances this gate in an explicitly added Cartan comparison. Its conditional subsidiary theorem requires the full probe and constitutive equations on an open set; its finite tangential-jet source law does not establish that hypothesis.

5. **Release gate.** Audit the corrected mathematical statements and every dependent use, reconcile historical claims with the current ledger, verify complete source inclusion and

© 2026 Marc Brendecke. Preprint — not peer reviewed.

layout, then make a deliberate version/DOI decision. No public upload or exact all-readout claim follows from the present pass.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

H1 Baseline Release and Reader-Provenance Records

The inherited records below are reproduced as source evidence. Their version-specific wording is not the metadata of this working copy, and historical assertions do not override its premise audit.

v4.7.0 Full-Monolith Release Scope and Provenance Notice

Recovery status. This v4.7.0 full-monolith source preserves the complete working monolith by using the last full-length source layer r54 as the base, appending the post-r54 guarded solve chain through r80, and retaining the r81–r84 frame-typed effective/local closure clarifications. Earlier compact upload handoff artefacts are not release bases because they compiled only a short consolidation artefact rather than the full monolith. The present release-candidate source keeps the corrected full-source recovery build and the compact title-page discipline.
DOI status. The intended reserved v4.7.0 Version DOI is 10.5281/zenodo.20232911. The Companion Concept DOI is 10.5281/zenodo.19210728. The previous/baseline v4.6.0 DOI remains 10.5281/zenodo.20117361 as provenance only.
Claim-scope lock. The release scope is guarded local/restricted only. It allows the local/restricted downstream chain, r60+ conservative C4 relative-seam neutralization, C5 detector-threshold terminality at $\theta = 1/256$ after $n = 20$, local photon-clock selector preservation under $\Delta_\gamma < 1/5$, typed mixed-comparison local control, terminal-data bridge, restricted terminal-data naturality, and terminal blocker freeze. It does not claim exact finite C5 zero closure, exact all-readout terminality, C6/T2 closure, a global photon-clock/RH theorem, or global all-readout closure.

v4.7.0-r44 Internal Working Status

Branch status. This source is an internal source-only working candidate over the v4.6.0 release-final baseline and the v4.7.0-r43 local certificate handoff packet. It performs a source-only handoff integrity scan and builds a local certificate index for the next working layer. It assigns no new Zenodo version DOI, authorizes no public upload, and produces no PDF in this intermediate pass.
Mathematical status. The r44 pass is a handoff-integrity and local-certificate-index pass. It does not promote the local MM/HC monitor certificate to a physical action, a Lagrangian, a derived field equation, a global readout closure, or a T2 theorem. It keeps the r43 guard vector active: $C4$ remains route-stable with $\beta_4 = 1/256$, finite $C5$-closure is not claimed, $C6/T2$ remains blocked, the RH channel remains a photon-clock handoff probe only, and global all-readout closure remains unclaimed.
Reader rule. Everything below should be read as a local MM/HC interface certificate extracted from a guarded atlas handoff. The resulting objects are certified local interface constraints, not derived physical laws.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.6.0 Release-Final Baseline Status

Release-final baseline note. This inherited baseline block records v4.6.0 release final with the version DOI 10.5281/zenodo.20117361. The main title remains *Quantum Sphaera Companion*. In the public v4.6.0 release-final source, the title page kept the Concept DOI and the Version DOI as separate DOI records. In this v4.7.0-r44 internal working source, the same DOI is listed as the baseline DOI and no new DOI is assigned. No all-readout/T2 closure is claimed.

Main title rule. The main title remains *Quantum Sphaera Companion*. The phrase *Bounded Readout Closure and T2 Scope Boundary* is a release-theme subtitle only, not a replacement for the main title.

Baseline DOI-line rule. The v4.7.0-r44 title page carries the stable concept DOI and exactly one baseline version DOI line: *Baseline Version DOI (v4.6.0): 10.5281/zenodo.20117361*. The concept DOI remains the stable Companion-line pointer. The date line records the internal working-candidate status, not a new public DOI assignment.

Release-final scope. This integrated release-final source records the v4.6.0 bounded readout result and the T2 scope boundary under the assigned Version DOI 10.5281/zenodo.20117361. It is the public release-final metadata state for v4.6.0, but it is not an all-readout closure theorem and does not promote any claim outside the declared bounded grammar.

Governing v4.6.0 status.

$$\mathsf{BRead}_{\mathrm{bounded}} \in \mathsf{Cons}(\mathcal{G}^{\mathrm{decl}}_{\mathrm{global}}), \qquad \mathsf{BRead}_{\mathrm{bounded}} \not\Rightarrow T2, \qquad T2 \in \mathsf{Ext}(\mathcal{G}^{\mathrm{decl}}_{\mathrm{global}}).$$

No empirical validation, Standard-Model-spectrum derivation, unrestricted physical readout, or all-readout closure is claimed.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Human-Facing Reading Layer and Notation Discipline

This edition separates the mathematical object language from revision and audit metadata. Internal revision identifiers and combined version–revision strings are not part of the physics or mathematics. They are provenance tags only. They may occur in file names, manifests, checksums, audit reports, and reconstruction notes, but they are not used as exponents, object names, states, fields, obstructions, or physical regimes in the reader-facing text.

The main reading order is therefore:

1. the structural and physical claim in ordinary mathematical notation;
2. the local hypotheses, guards, and return conditions;
3. the obstruction or witness that remains after the permitted tests;
4. the machine-readable audit trail, kept only as provenance material.

When a previous draft used a revision-bearing symbol, this edition rewrites it as a semantic object such as X^{rtg}, X^{lock}, X^{nonsplit}, or simply X, according to the mathematical role of the object. Version numbers remain valid release metadata, not mathematical structure.

Reader rule. The human-facing document should be readable without knowing the internal patch history. The audit/provenance layer is still preserved for reconstruction and LLM-assisted maintenance, but it is subordinate to the mathematical presentation.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Reader-Facing Proof Spine and Document Routing

This v4.6.0 release-final frontmatter treats the mathematical narrative as the primary document and the machine/audit trail as secondary provenance. The inherited v4.5.0 release-final layer remains visible as prior release genealogy, but the governing public metadata state is now v4.6.0 with Version DOI 10.5281/zenodo.20117361. The reader should not need the patch history in order to understand the current object language. Revision identifiers, local build labels, and version-transition notes are therefore moved out of the proof spine and into the provenance layer.

Primary reading order

1. **Structural claim.** Read the HC/MM/chart interface as a structural bridge problem with explicit guards and comparison norms.
2. **Local grammar.** The photon-frame HC-halo/Dixon/lens grammar gives the inherited local admissible test surface.
3. **Finite tests.** Candidate menus are used only as certified finite screens. They may consume, route, block, or name an obstruction; they do not create a physical claim by themselves.
4. **Current v4.6.0 endpoint.** The active endpoint is the bounded-readout surface: it is conservative and debt-free inside the declared grammar, but it does not imply T2 and does not discharge all possible readouts.
5. **Return and handoff contract.** A future all-readout answer must return one of three objects: a kill certificate, a strict norm-decrease witness, or a named persistent obstruction. The v4.7 handoff therefore starts with the MM/HC exterior–mixed field-equation program, not with a T2 closure claim.

Current semantic object registry

The active state is represented by semantic names, not revision names. The older route-to-global bridge symbols remain valid v4.5 genealogy; the v4.6.0 governing objects are the bounded-readout surface, the all-readout scope boundary, and the MM/HC exterior–mixed handoff:

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Symbol	**Role**	**Reader meaning**
B_{tie}	inherited chart-tie component	v4.5 non-split coupling extracted from the local pair
B_{lock}	inherited locked remnant	v4.5 remaining tie object after partial cancellation screens
$\mathcal{O}_{\mathcal{B}}^{\mathrm{lock}}$	inherited locked bridge obstruction	v4.5 bridge obstruction; genealogy, not the final v4.6 endpoint
$\mathsf{BRead}_{\mathrm{bounded}}$	bounded readout surface	declared bounded-readout object: conservative inside the declared grammar
Θ^{all}	all-readout demand	unrestricted readout family required for T2; not supplied by the bounded result
$\mathfrak{O}_{\mathrm{all}}^{4.6.0}$	frozen all-readout obstruction	missing universal exterior selector, all-readout naturality family, and unrestricted exterior collar slot
$\mathcal{M}_{\mathrm{bd}}, \mathcal{M}_{\mathrm{ext}}, \mathcal{M}_{\mathrm{mix}}$	Millennium-matrix split	bounded, exterior, and mixed parts for the next field-equation determination
$\mathcal{H}_{\mathrm{bd}}, \mathcal{H}_{\mathrm{ext}}, \mathcal{H}_{\mathrm{mix}}$	HC split	bounded, exterior, and mixed compensator portions matched to the same readout split
$\mathfrak{E}_{\mathrm{decl}}$	field-equation reset template	structure-first equation template for the v4.7 MM/HC exterior–mixed program

Guard discipline

No reader-facing claim is promoted unless the relevant guard is explicitly discharged. In particular, this edition does not claim T2, all-readout closure, global OP closure, a global HS^2 focal-return theorem, a through-collapse trajectory, an unverified atlas expansion, or a Standard-Model-spectrum derivation. The bounded-readout theorem is therefore read as a conservative declared-grammar result, while Θ^{all} remains a scope boundary unless a separate all-readout witness family is supplied.

Human/LLM layering rule

The main document contains the mathematical object language. The appendix may contain provenance: source packets, diffs, checksums, machine-readable registers, build logs, and reconstruction notes. Provenance may mention build labels. The proof spine must not use them as mathematical exponents, object names, or physical states.

Compressed chronology reading rule

For reader navigation, inherited working chronology is to be compressed into semantic milestones rather than read as active mathematical notation. The compression convention is:

© 2026 Marc Brendecke. Preprint — not peer reviewed.

chronology stratum	reader-facing meaning	status
C_0	earlier local readout and release genealogy	provenance
C_1	v4.4 declared-local OP1/OP4 closure ledger	prior release result
C_2	v4.5.0 declared global-boundary synthesis	prior theorematic release layer
C_3	v4.5.0 DOI/title/frontmatter repair	release metadata
C_4	v4.5.x editorial QA and chronology compression	post-release readability layer
C_5	v4.6.0 bounded-readout theorem and T2 scope boundary	current governing release status
C_6	MM/HC exterior–mixed field-equation determination	v4.7 handoff program

Old working strings such as release-candidate notes, prior version DOIs, and local revision labels are therefore genealogy unless a nearby release-final or governing-status block explicitly says otherwise. This rule changes only reader navigation; it does not change labels, references, DOI records, theorem statements, or scope guards.

Proof-spine phrase audit rule. In running prose, build expressions such as source-native rebuild, release candidate, freeze audit, packet, and working layer are reader-facing provenance terms unless they are explicitly part of a displayed ledger. Where such terms occur in the active proof spine, the preferred reader form names the mathematical role first—for example grammar exhaustion, readout invariance, debt-free boundary transport, no-new-debt extension, or functorial comparison stability—and keeps the build phrase only as provenance. This rule does not rename labels, change theorem statements, alter DOI records, or change any scope guard.

Reader-facing phrase-normalization rule

Current-line section headings are allowed to use semantic names even when the source labels and audit ledgers retain their original revision identifiers. Thus a heading such as a witness execution, boundary audit, source-native checkpoint, or title-page repair is to be read by its mathematical function, not by the working-string that produced it. Revision strings remain valid in labels, comments, diffs, ledgers, and reconstruction records, but they are not mathematical object names, proof data, tensor indices, route states, or closure witnesses.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Companion status. This is the v4.6.0 release-final frontmatter after bounded-readout integration, T2 scope-boundary freeze, and DOI normalization. It preserves the declared mathematical global-boundary closure audit path from the v4.5.0 release line as provenance while making the v4.6.0 Version DOI 10.5281/zenodo.20117361 the current public release DOI. Version labels are retained only as provenance metadata; the mathematical proof spine uses semantic names.

Governing theorematic status. Inside the declared global-boundary synthesis grammar

$$\mathcal{G}^{\mathrm{decl}}_{\mathrm{global}},$$

OP 1, OP 4, and OP 5 are closed as a declared mathematical global-boundary closure. This does not claim unrestricted all-readout closure, global physical closure, QFT/GR closure, QCD/confinement, CKM/PMNS, quark-mass, or Standard-Model-spectrum derivation.

DOI rule. The concept DOI remains the stable title-page pointer. The v4.5.0 version-specific DOI is 10.5281/zenodo.20079410; prior version-specific DOI records are provenance metadata only. Version IDs, DOI fragments, dates, and release IDs are not mathematical object names, indices, theorem labels, route tags, or proof data.

Internal r10 proofread status. This v4.5.1-r10 layer is an internal selected-range proofread and metadata-hygiene pass. It does not authorize public upload, does not insert a new DOI, does not emit Zenodo release text, and does not change the declared mathematical closure scope.

Internal r11 terminology-sweep status. This v4.5.1-r11 layer is an internal targeted terminology sweep over the older OP5 and OP1/OP4 compatibility pages. It does not authorize public upload, does not insert a new DOI, does not emit Zenodo release text, does not rename theorem labels, and does not change the declared mathematical closure scope.

Internal r13 glossary/index intake status. This v4.5.1-r13 layer is an internal glossary and index intake for the declared global-boundary synthesis grammar and the OP5/OP1/OP4 terminology family. It is a reader-routing layer only: no public upload is authorized, no new DOI is inserted, no Zenodo release text is emitted, no theorem label is renamed, and no mathematical closure scope is changed.

Internal r13 glossary-appendix worksheet status. This v4.5.1-r13 layer is an internal extraction worksheet for a possible future glossary appendix. It does not create a public release refresh, does not insert a new DOI, does not emit Zenodo release text, does not rename theorem labels, and does not change the declared mathematical closure scope.

Abstract

This v4.6.0 release-final frontmatter records the Quantum Sphaera Companion after bounded-readout integration, T2 scope-boundary freeze, DOI synchronization, and title-page/frontmatter hygiene. The version-specific DOI for this release line is 10.5281/zenodo.20117361. The Companion concept DOI remains 10.5281/zenodo.19210728. The v4.5.0 version DOI 10.5281/zenodo.20079410 is retained as prior-release provenance.

The governing v4.6.0 release-facing result is the bounded-readout theorem together with the T2 scope-boundary statement. Inside the declared global-boundary grammar $\mathcal{G}^{\mathrm{decl}}_{\mathrm{global}}$, the bounded readout surface is conservative and debt-free, while T2 remains an explicit all-readout extension problem rather than a theorem asserted by this release.

This is a scoped mathematical theorem. It is not a global physical closure theorem, not a global QFT/GR theorem, not unrestricted closure over every possible readout or physical model, and not a T2 closure claim. Earlier version-specific DOI records remain provenance metadata only and are not mathematical object names, theorem labels, route tags, or proof data.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Frontmatter reader architecture

Purpose. This page separates the current release-facing route from historical theorem memory. It is an orientation layer only; it does not replace any definition, theorem, proof, audit ledger, or historical witness later in the document.

Layer	Human reading route	LLM/audit route
Current reader status	Read this as the v4.6.0 release-final frontmatter after bounded-readout integration, T2 scope-boundary freeze, DOI synchronization, and metadata hygiene.	Treat this pass as reader-facing organization and provenance separation only; no new mathematics is introduced beyond the audited v4.6.0 bounded-readout layer.
OP5 status	Read OP 5 as locally closed only inside the declared photon-frame HC-halo/Dixon grammar.	Track the required zero registers $L_{\mathrm{unr}}, B_{\mathrm{HC/MM}}, R_{\mathrm{HS4}}, D_{\mathrm{HS5}}$ and reject any stronger global reading.
QAM/MML/MMA	Treat QAM as store/object grammar, MML as routing/command grammar, and MMA as execution/readout semantics.	Use finite object bridges, instruction alphabets, deterministic guarded transitions, and route-status tables as audit handles.
Dixon coding	Read the Dixon layer as a 64-dimensional carrier with a 2^{64}-mask activation space.	Keep carrier dimension and Boolean mask count separate; do not read Dixon itself as 2^{64}-dimensional.
Historical witness	Use it as genealogy and reconstruction support.	Preserve it as non-destructive proof memory; do not read inherited baseline language as the current public claim unless explicitly mapped by the local proof chain grammar.

Table 1: Frontmatter separation of current release status, OP5 local closure, and historical theorem memory.

DOI discipline for the human-facing edition

Record	Status	Reader-facing rule
Concept DOI	stable Companion-line pointer	10.5281/zenodo.19210728
Version DOI	current public version-specific release DOI	10.5281/zenodo.20117361.
Prior release metadata	provenance layer only	Historical DOI records remain reconstructible in audit and release files, not in the proof spine.

DOI and versioning rule. DOI records and release identifiers are metadata. They must not

© 2026 Marc Brendecke. Preprint — not peer reviewed.

be used as mathematical indices, route tags, theorem labels, tensor subscripts, obstruction names, or proof data in the reader-facing argument.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Prior DOI provenance note

Prior DOI provenance. Earlier version-specific DOI records belong to released predecessors and are provenance metadata only. This v4.6.0 release-final line assigns the current version-specific DOI 10.5281/zenodo.20117361; the v4.5.0 DOI 10.5281/zenodo.20079410 is retained as prior-release provenance.
Working-source status. This release-final adds no theorematic claim beyond the audited Active Core. The current local readout ledger remains guarded: the orbit/Hopf chain is a finite local readout closure ledger with no persistent finite carrier obstruction in the declared diagnostic chart, not a global physical closure theorem.

Current Reader Navigation and Governing Map

> **Status.** This reader-navigation block is a human-facing cleanup aid. It does not add new mathematics. It explains which parts of the Active Core govern the current local OP5 statement and which parts preserve historical proof memory.

Governing reading. The Active Core is the governing current line. The current semantic reading keeps QAM/MML/MMA as guarded finite object, routing, and execution/readout grammars and reads the OP5 closure statement only through the declared photon-frame HC-halo/Dixon grammar.

Historical reading. The Historical Witness and inherited older-version blocks are not second active theorematic lines. They preserve genealogy, older proof memory, reconstruction context, and release chronology. Historical or pre-current displays should be read as witness material unless they are explicitly reintroduced through a current normal-form, route, or audit record.

Recommended reading order.

Reader goal	Read first	Read only if needed
Current semantic status	frontmatter status box, OP5 status spine, freeze-audit records	older version-history paragraphs
OP5 local closure	the local proof chain HC-halo/Dixon route, zero-register assumptions, Dixon image audit	historical OP5/OP5(B) genealogy
QAM/MML/MMA language	current guarded store/route/readout normal forms	older pre-current notation examples
Dixon coding	HS(6) 64-carrier and 2^{64} mask distinction	algebraic genealogy of Dixon/octonionic terms
Version audit	manifest, checksums, diagnostic report, source-freeze audit	full Historical Witness

Status guard. The current reader-facing edition does not by itself close OP 1 or OP 4, does not claim global OP5 closure outside the declared grammar, does not activate primitive mixed tensors, and does not replace standard QFT/QED/Lifshitz or general-relativistic calculations.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Extended Abstract and Reader Orientation

> **Purpose.** The short abstract gives the release-facing summary. This extended abstract records the technical routing information needed by human readers and LLM/audit readers without turning inherited version genealogy into a current public claim.

The present v4.3.0 release-final belongs to the reader-cleanup line. It publishes the normalized public frontmatter after the source-native freeze audit and frontmatter normalization while keeping the mathematical body unchanged. Older version identifiers that remain later in the body are treated as theorem-memory, genealogy, release chronology, or inherited source references rather than as the current public-status sentence.

The central current statement is the local OP5 closure inside the declared photon-frame HC-halo/Dixon grammar. The proof spine passes through HC-cone normalization, double-ε halo capture, tunnel reduction, normal-collar rank control, selector/overlap routing, Dixon 64-carrier coding, and image audit. The residual image is empty only under the explicit zero-register assumptions $L_{\mathrm{unr}} = 0$, $B_{\mathrm{HC/MM}} = 0$, $R_{\mathrm{HS4}} = 0$, and $D_{\mathrm{HS5}} = 0$.

The QAM/MML/MMA reading remains conservative. QAM is read as object/store grammar, MML as finite routing/command grammar, and MMA as execution/readout semantics. Dixon coding is used as a finite addressability layer: the Dixon carrier is 64-dimensional, while the Boolean activation-mask space has cardinality 2^{64}. This distinction is part of the release guard.

Status. OP 1 and OP 4 remain open. OP 5 is not claimed globally outside the declared photon-frame HC-halo/Dixon grammar. Primitive mixed tensors remain guarded. No QCD/confinement, CKM/PMNS, quark-mass, or Standard-Model-spectrum derivation is claimed.

Reader route. A human reader should start from the current status box and the compact OP5 status spine, then read the QAM/MML/MMA and Dixon material as controlled addressability infrastructure. An LLM/audit reader should track the zero registers, route-status classes, and the separation between current active claims and inherited theorem-memory blocks.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

H2 Complete QAM, MML and MMA Infrastructure

The later shifted normal form is linked separately in Appendix A.27; pre-shift statements here retain their historical position in that migration chain.

Working orientation for 4.7.0

> **Reader's map.** The active reading path of the present document is: *orientation and compact front extraction → QAM formal layer → tool and proof-compression layer → open-question / live-burden layer → computational and decoder layer.* The later appendix-style historical material remains available for proof-memory, genealogy, and local rebuild support, but it is not the governing active theorem core.

Version 4.7.0 proceeds from the normalized release frontmatter to the inherited mathematical body. The governing current-state claim is the local OP5 closure inside the declared photon-frame HC-halo/Dixon grammar under the stated zero-register assumptions. Preserved release-roadmap notes, checkpoint transitions, and earlier front-matter control blocks remain part of the document as inherited baseline material for theorem-memory, genealogy, and reconstruction support; they are not the overriding current public-status sentence of Version 4.7.0.

bibliography hardening. A full-document literature scan was performed before this public split-bundle release. The scan found that the mathematical body repeatedly uses Teichmuller and quasiconformal language, Beltrami-style deformation intuition, Riemann–GUE/RMT spectral heuristics, Hopf/fibre-bundle topology, Dixon/octonionic and division-algebraic language, Dirac/Higgs/Yang–Mills and CKM/PMNS landmarks, Julia-boundary diagnostics, ZF/QAM coding language, and probe/cryptographic/computational analogies. Version 4.7.0 therefore adds a broader terminal bibliography and uses this paragraph as the compact external anchor surface: Teichmuller/quasiconformal background is anchored in [13, 14, 15, 16]; Riemann-zeta, spectral, and GUE/RMT background in [5, 6, 7, 8, 9, 10, 11, 12]; Hopf, fibre-bundle, and exceptional-geometry background in [18, 19, 20, 21, 22]; division-algebraic and octonionic background in [23, 24, 25]; QFT/gauge and mixing landmarks in [26, 27, 28, 29, 30, 31, 32]; complex-dynamical Julia background in [17]; ZF/set-theoretic foundations in [38, 39, 40, 41]; and the computational/probe/cryptographic side in [42, 43, 44, 45, 46, 47, 48, 49, 50, 51].

MML/MMA compression map. The local language interface is now used as a conservative compression and audit map for inherited HC/OP text. The purpose is not to delete historical proof material, but to mark repeated structural paragraphs as instances of typed operations such as $\mathrm{CAPTURE}_{\mathrm{HC}}$, $\mathrm{HANDOFF}_{\mathrm{HC}\to\mathrm{OP}}$, $\mathrm{NORMALIZE}^{\mathrm{HC}}_{\mathrm{OP}}$, $\mathrm{ROUTE}^{\mathrm{HC}}_{\mathrm{OP}}$, $\mathrm{COMPARE}^{1/4|5(B)}_{\mathrm{HC}}$, RESOLVE, $\mathrm{RETURN}_{\mathrm{OP}}$, SCAN, RANK, and CERTIFY?. The compression map is therefore a reading and refactoring guide: it can shorten later exposition and make inherited blocks easier to audit, but it cannot manufacture certificates or remove the historical witness.

first tool-contract application. The tool library is now applied back to the active HC/OP proof spine. The purpose is deliberately modest: repeated paragraphs in the local proof chain may be referenced by the contracts $\mathfrak{T}_{\mathrm{lock}}$, $\mathfrak{T}_{\mathrm{handoff}}$, $\mathfrak{T}_{\mathrm{norm}}$, $\mathfrak{T}_{\mathrm{route}}$, $\mathfrak{T}_{\mathrm{cmp}}$, $\mathfrak{T}_{\mathrm{resolve}}$, $\mathfrak{T}_{\mathrm{cert}}$, and $\mathfrak{T}_{\mathrm{frontier}}$ once their local hypotheses and guards have been stated. This is a citation and audit layer, not a shortening pass that deletes proof content.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

guarded reduction protocol. The present revision adds the first reduction discipline for using those tool contracts in future text. A paragraph is reducible only when it repeats an already stated safety, route, frontier, diagnostic, or no-false-certificate clause. It is not reducible when it introduces a new object, a first hypothesis, a local guard, a certificate test, a proof restriction, a theorematic status change, or historical witness material. Thus the layer is the first practical simplifier produced by MML/MMA, but it remains conservative: it turns repeated prose into guarded citations and never turns an execution trace into an OP closure certificate.

guarded-reduction application pass. The present revision applies the discipline for the first time without deleting text. It installs a finite ledger of guarded citation sites along the active HC/OP proof spine: lock/capture reminders, handoff and normalization reminders, route-frontier reminders, paired-comparison status clauses, probe diagnostic clauses, and archive-to-tool extraction reminders. Each entry keeps its first occurrence and local guard explicit, records the relevant tool contract, and permits only later repeated tails to be cited in guarded normal form. Thus is a practical audit layer: it tells the reader exactly where the new compression technology helps the existing text, while preserving theorematic status and the historical witness.

guarded-reduction pilot normal form. The present revision turns the ledger into a first usable rewriting pattern. It does not shorten the historical witness and does not replace first definitions. It only says that, after a first occurrence has stated its hypotheses and local guard, a later repeated HC/OP tail may be written in the guarded citation form

$$\mathrm{GRCite}_{\mathrm{HCOP}}(\ell; G, W, S),$$

where ℓ names the ledger entry, G points to the local guard, W points to the preserved witness or first occurrence, and S records the non-certificate status. This is the first concrete simplification normal form produced by the MML/MMA layer: it compresses repeated prose, not proof content.

tail-normalized proof skeleton. The present revision organizes the guarded-citation mechanism into a first tail-normalized proof skeleton for the active HC/OP spine. Its purpose is to show where the first explicit mathematical source block remains, where later repeated endings may be cited in guarded form, and how the dependency order of lock/capture, handoff, normalization, routing, paired comparison, and resolution is preserved. Thus does not shorten the document by deletion. It gives future HC/OP text a compact reference skeleton in which repeated tails can be replaced by guarded citations only after their source block, local guard, witness, and theorematic status have already appeared.

skeleton-to-workstep dispatch. The present revision turns the tail skeleton into a first controlled dispatch object. Its purpose is not to claim an OP closure, but to say which finite next work packets may be launched from the ordered skeleton nodes. The admissible dispatch order remains lock/capture, handoff, normalization, routing, comparison, and resolution; each dispatched packet must keep its local guard, witness, route tag, and non-certificate status unless a separate explicit certificate test is later supplied.

first dispatched lock/capture stabilization. The present revision executes the first dispatched work packet from the menu. It acts only on the LOCK/CAPTURE node, records a

© 2026 Marc Brendecke. Preprint — not peer reviewed.

finite lock-stability defect register, and returns one conservative status: handoff-ready stable source, bounded local repair source, or smaller frontier source. This is the first point at which the skeleton-to-workstep dispatch becomes an actual controlled mathematical work step; it still does not generate an OP closure certificate.

QAM/MML/MMA local interface registry. Version 4.7.0 records the narrow local interface stamps for the machine-language layer in prose and uses symbolic local interface tags in formulas for the current HC-routed OP-entry chain. The historical computational chapter already contains the broader MML–MMA–MCP reserve architecture. The present step does not activate that reserve as an independent proof engine. It only extracts the finite sublanguage needed to read the local proof chain as a typed program:

$$\mathrm{QAM}^{\mathrm{loc}}_{\mathrm{HCOP}} \longrightarrow \mathrm{MML}^{\mathrm{loc}}_{\mathrm{HCOP}} \longrightarrow \mathrm{MMA}^{\mathrm{loc}}_{\mathrm{HCOP}}.$$

Here $\mathrm{QAM}^{\mathrm{loc}}_{\mathrm{HCOP}}$ is the local bridge tag over the inherited QAM-over-ZF object language, with interface stamp QAM-HC/OP 0.1 recorded as metadata; $\mathrm{MML}^{\mathrm{loc}}_{\mathrm{HCOP}}$ is the finite instruction interface for capture, handoff, normalization, routing, comparison, and return, with interface stamp MML-HC/OP 0.2 recorded as metadata; and $\mathrm{MMA}^{\mathrm{loc}}_{\mathrm{HCOP}}$ is the route-faithful execution semantics for those instructions on finite HC/OP states, with interface stamp MMA-HC/OP 0.2 recorded as metadata.

The following inherited front block should therefore be read as the carried-forward active baseline from the audited archival stage line, with the following priorities and outcomes:

1. **Preserve the stabilized closure/interface baseline.** The language/closure, branch/shell/routed, tensor, operator-family, translation, OP 3 shell-certificate, bundled interface compatibility, and terminal proof-use surface already exported in earlier public stages are treated as fixed baseline rather than reopened here.

2. **Keep the partial-closure/no-second-carrier layer fixed.** The terminal-interface no-second-carrier theorem, the first partial OP 3/OP 5 closure theorem, and the restricted lift-back principle are treated as available structural infrastructure.

3. **Keep the first retained-carrier optical approximation and machine-shadow baseline fixed.** Reduced prism readout, residual diagnostics, carrier-side fit/calibration/validation, inversion, refinement, reduced stability, and the bidirectional observer-side split $\eta_+ = \eta_{\mathrm{drive}} + \eta_{\mathrm{carrier}}$, $\eta_- = \eta_{\mathrm{drive}} - \eta_{\mathrm{carrier}}$ are treated as the stabilized computational surface rather than rebuilt from scratch.

4. **Inherit the completed tool/QAM/decoder layer as fixed infrastructure.** Channel-sorted total, imbalance, and bridge packets, finite QAM submenus, primary tags, registry assignments, tag-first reopening routes, the Monte-Carlo residual-signature routing layer, the action table, deterministic decoder syntax, transition function, observer/event normal form, evaluation map, and compact output tokens are treated as the completed route-control package inherited from historical release marker archival stage.

5. **Keep the prior post-closure burden reading visible as inherited baseline.** The archival stage–archival stage burden picture is preserved here as inherited local baseline material from which the new pre-OP frontier synthesis is now read, not as an independent current public-claim layer that overrides the new groundwork status stated in the opening pages.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

6. **Keep the DOI logic normalized.** The concept DOI remains the stable title-page pointer of the Companion line. The governing latest public-release DOI in the chronology is now 10.5281/zenodo.19750674 for the public archival stage release, while 10.5281/zenodo.19746821 remains the prior public release DOI for archival stage.

Reading rule for the inherited active-front block. The inherited archival stage front layer is kept intact as carried-forward baseline source material. It records the then-active retained-carrier optical and live-burden reading, including OP 1 and OP 5 Step (B) closure language on the inherited admissible corridors, and should not be read as the governing present-release claim of 4.7.0. In the present release-final, those statements are preserved for theorem-memory and genealogy; the governing current-state claim, however, remains the frontier-certified pre-entry synthesis developed in the new post-archival stage line. Accordingly, references below to live OP burden, route sharpening, kernel closure, or routed-balance obstruction inside the inherited front block should be read as preserved baseline source material rather than as the final current release claim of this release.

Compact public-shape branch/shell/routed extraction

Status. *Active front-facing compression block.*

Block function. This block gives the smallest front-facing statement of the public branch/shell/routed QAM package, so that a reader can see what is already stabilized before entering the fuller internal unfolding.

The inherited public-shape QAM layer may be stated in compressed form before the full internal unfolding is read. In that form, one starts with the already public coded source/readout, branch/shell/routed, tensor, operator-family, and translation layers from historical release marker archival stage–archival stage, then adds shell certificates, shell-defect witnesses, kernel-persistence statements, and shell-stable readout factorization. The guiding claim is that these added roles do not create a new closure criterion; they only make the OP 3 shell layer readable as one explicit coded closure mechanism.

Definition H2.1 (Compact public-shape QAM package)**.** *The* compact public-shape QAM package *for the inherited public-shape archival stage release layer consists of the following release-level data:*

1. *tagged coded objects and minimal predicates for source states, layers, operators, readouts, branches, shell/refinement objects, admissibility witnesses, and tensor-coded composites;*
2. *admissible coded transport together with coded closure-class preservation and closure-compatible coded readout;*
3. *closure-admissible branch codes;*
4. *shell-admissible refinement codes over closure-admissible branches;*
5. *normalized admissible routed support over the admissible branch-shell support class.*

At this public-shape stage, tensor admissibility, operator-family/lambda-like maps, and support-selection or degeneracy refinements are not part of the compact release claim, even though they remain internally available below.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Theorem H2.2 (Compact public-shape preservation chain)**.** *Let s_{in} be a coded source state, let b be a closure-admissible branch code, let σ be a shell-admissible refinement code on b, let u be a normalized admissible routed support code over the admissible branch-shell support class, and let r be a closure-compatible coded readout. Then the archival stage public-shape layer preserves the same closure-relevant readout content along the full compressed chain*

$$s_{\mathrm{in}} \longrightarrow (b, \sigma) \longrightarrow u \longrightarrow r,$$

provided the coded return realizations are taken inside the admissible branch-shell support sector.

Proof. At OP level, the same compression is built in stages: invariant source-return closure for the OP 1–OP 5 pair is fixed by Theorem H7.2; branchwise preservation is then isolated by Proposition H7.5; shell-sensitive preservation at the OP 3 interface is isolated by Theorem H7.21; and normalized routed preservation is isolated by Proposition H7.9. The QAM translation theorem (Theorem H2.238) carries exactly that OP chain into coded form. Later in the QAM section this becomes the admissible branch-shell theorem (Theorem H2.157), the routed admissible support theorem (Theorem H2.189), and the visible/structural separation principle (Corollary H2.212). The present statement records that entire OP-to-QAM compression as one front-facing release theorem. □

Remark H2.3 (Public-shape reading guide)**.** *For the purposes of the current release, the later QAM section should be read with the following dictionary in mind:*

1. *"coded source state + closure-compatible coded readout" is the coded front-face of the OP 1/OP 5 source-return pair;*
2. *"closure-admissible branch code" is the coded branchwise refinement role;*
3. *"shell-admissible refinement code" is the coded OP 3 shell role;*
4. *"normalized admissible routed support" is the coded routed aggregation role over the admissible branch-shell support sector;*
5. *"closure-relevant readout content" always means readout agreement after passage through the associated closure-factor map.*

This dictionary is the intended reading bridge between the compact front theorem and the later theorem-level unpacking in Theorem H2.157, Theorem H2.189, Theorem H2.238, and Corollary H2.212.

Corollary H2.4 (Release-level visible/structural rule)**.** *Within the compact public-shape QAM package, visible coded mismatch in packet, phase-lock, or ε-screening data is not by itself a closure failure. At the inherited public-shape level, genuine closure failure begins only when branch admissibility or shell admissibility is lost.*

Remark H2.5 (Structural provenance of the public-shape extraction)**.** *The archival stage front block should be read as the coded shadow of one already stabilized OP chain and not as a freestanding new criterion. In that reading, coded source state and coded readout come from the closed OP 1–OP 5 source-return pair of Theorem H7.2; closure-admissible branch codes come from the branchwise refinement layer of Proposition H7.5; shell-admissible refinement codes come from the OP 3 shell theorem of Theorem H7.21; normalized admissible routed support comes from the routed aggregation layer of Proposition H7.9; and the release-level visible/structural rule is the QAM-over-ZF front-face of the OP screening logic, where*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

packet/phase/epsilon data remain preselection data unless they witness loss of admissibility. The public-shape extraction therefore does not add a second closure notion. It republishes the already existing source-return closure architecture in the smaller coded alphabet needed for the current QAM-over-ZF line.

Remark H2.6. *This compact block is intentionally stronger in readability than in scope. It exposes the third public-shape QAM step as one coherent public claim, but it does not yet export the full internal support-selection or wider compression machinery. In other words, archival stage was prepared as the controlled public extraction of the tensor/operator layer, not as the publication of the entire internal QAM architecture.*

Remark H2.7 (Deferred kernel-first outlook)**.** *The front theorem should also be read as the visible place where any later zero-kernel migration would have to pay off. The deferred kernel notes below do not introduce a second active tuple language, but they already identify the only later proof-compression route that would matter on the present public surface: first kernel-first transport reduction (Proposition H2.140), then local branch-shell kernel reduction (Proposition H2.182), and only afterwards routed family compression (Proposition H2.190). Read together with the roadmap-fit remark, the later checkpoint summary, the activation criterion of Theorem H2.197, and the later remarks on the staged activation roadmap, the criterion-to-roadmap reading rule, and the compressed synthesis of the deferred zero-kernel path, this deferred path should therefore be understood as a later proof-theoretic refinement of the same front chain rather than as a competing current-release line.*

Remark H2.8 (QAM-over-ZF entry layer)**.** *A next formal development line of the Companion is the introduction of QAM not as a replacement for Zermelo–Fraenkel set theory, but as a structure-first coded language over ZF. The point of this layer is to give the current OP/branch/shell architecture a cleaner formal interface for states, layers, operators, readouts, admissibility, closure preservation, and tensor-coded composition while remaining, in principle, back-translatable into set-theoretic form.*

At the present integrated stage, the admissible OP fragment remains the first translation target of this QAM line: the OP 1/OP 5 source-return closure architecture, admissible branch transport, OP 3 shell-admissible refinement, admissible routed support, tensor composition over admissible coded objects, and the visible packet/phase-lock/epsilon screening layer. Relative to the public archival stage release, the next public-shape export in that historical sequence was aimed at the tensor and operator/lambda layer rather than at a further broadening of the branch/shell/routed block.

The reserved minor-release path remains the following. First, archival stage fixed the minimal QAM language layer over ZF, including tagged set-codes for states, layers, operators, readouts, branches, shell/refinement objects, admissibility witnesses, and tensor-coded composites. Second, archival stage extracted the branch/shell/routed translation into public-shape form, including closure-admissible branch codes, shell-admissible refinement codes, admissible routed support, and the first coded branch-shell transport results. Third, archival stage realized the tensor/operator public line; fourth, archival stage exported the first public OP 3/QAM closure package; fifth, archival stage hardened the witness-grade reopening discipline of that bundled channel; and the present 4.7.0 release line now pilots a closure-liquid-crystal / holographic Julia-boundary re-reading above that compressed surface while support-selection refinements remain below as later-stage technical support.

The present integrated continuation therefore exposes the compressed public-shape QAM core together with its branch/shell/routed baseline, its tensor/operator extension, its public OP 3/QAM closure package, its bundled witness-grade compression surface, and the released archival stage crystal–shadow architecture: tagged coded objects and minimal predicates,

© 2026 Marc Brendecke. Preprint — not peer reviewed.

admissible transport with coded closure-class preservation, closure-compatible coded readout, closure-admissible branch codes, shell-admissible refinement codes, admissible routed support, tensor admissibility, tensor closure, first operator-family or lambda-like translation statements, terminal proof-use compression, addressed crystal packets/corridors/classes, theorem-level shadows, and the terminal crystal–shadow interface. The broader internal QAM package remains available below as technical support and does not alter the mathematical claims already frozen in the public archival stage release.

QAM-coded object preview. *The minimal intended object types are states, layers, operators, readouts, branches, shell/refinement objects, admissibility/closure witnesses, and tensor-coded composites. A set x is intended to count as a QAM-coded object exactly when it has one of the tagged forms*

$$\langle 0, a\rangle, \quad \langle 1, \ell, D\rangle, \quad \langle 2, \ell_1, \ell_2, G\rangle, \quad \langle 3, \ell, W, R\rangle,$$
$$\langle 4, b, B\rangle, \quad \langle 5, r, s, T\rangle, \quad \langle 6, u\rangle, \quad \langle 7, p, q\rangle.$$

If p and q are QAM-coded objects, the tensor-coded composite is written as

$$p \otimes q := \langle 7, p, q\rangle.$$

At the entry stage, these tags and tensor codes are only coding devices that distinguish structural roles inside ZF and reserve tensor syntax for later internal hardening.

Remark H2.9 (Dependency guide for the internal QAM package)**.** *The internal QAM-over-ZF line should be read in the following order: coded objects and predicates first, admissible transport and closure-class schemata second, branch/shell/routed admissibility third, tensor admissibility and tensor closure rules fourth, faithful translation of the admissible OP fragment fifth, operator-family/lambda-like structural maps sixth, and support-selection/degeneracy handling only as a seventh-layer refinement on top of the already admissible fragment.*

How to read the internal QAM package

The internal QAM package should be read in two layers. At the compact reader level, one should keep only the following chain in view: coded objects and predicates, admissible transport and closure, branch/shell/routed admissibility, tensor closure rules, faithful translation of the admissible OP fragment, and only afterwards operator/lambda and support-selection refinements. At the technical level, the detailed sections below provide the full internal unfolding of these same layers in theorematic form.

Remark H2.10 (QAM normalization guide)**.** *The current internal package is not meant to be enlarged indiscriminately. Its next intended use is normalization and later public extraction. In particular, later work should prefer compression, dependency control, and clean appendix-style grouping over uncontrolled addition of parallel QAM subsystems.*

Roadmap and extraction criteria for the archival stage public-shape preparation

The internal QAM package is now far enough along that one may formulate a controlled extraction path toward the next public QAM-oriented line. The intended purpose of this extraction is not to publish the entire internal QAM package at once, but to expose its stable and readable core in stages.

The current preferred extraction sequence is the following.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

1. **First public QAM line (realized in archival stage): language and closure interface.** Publish a compact QAM-over-ZF entry layer together with the minimal predicate package, admissible transport, coded closure-class preservation, closure-compatible readout, and a first translation statement for the admissible OP fragment.

2. **Second public QAM line (realized in archival stage): branch/shell/routed translation.** Extend the public QAM line by adding closure-admissible branch codes, shell-admissible refinement codes, admissible routed support, and the first coded branch-shell transport results.

3. **Third public QAM line (realized in archival stage): tensor and operator/lambda extension.** Add tensor admissibility and tensor closure rules together with the first operator-family or lambda-like structural maps and explicit translation or round-trip statements, while keeping selection/support handling internal for now.

4. **Fourth public QAM line (realized in archival stage): first public OP 3/QAM closure package.** Expose shell certificates, shell-defect localization, kernel persistence under the bridge hypotheses, shell-stable readout factorization, and the bundled OP 3→OP 5/global interface in compact public shape.

5. **Fifth public QAM line (realized in archival stage): witness-grade hardening and local reopening discipline.** Add witness-grade proof-use control for the bundled channel together with a first narrow licensed kernel-first rewrite window, a first corridor-level activation lemma, a first corridor-to-bundle propagation rule, a first bundle-level reopening compression principle, a first profile-to-run compression principle, a first word-determined run-class step, a first run-class concatenation principle, a first finite run-class normal-form principle, a first normal-form-determined bundled proof-use class step, a first proof-use-class composition principle, a first associative proof-use-class normal-form principle, a first contraction-stable class-normal-form principle, and a terminal proof-use surface.

6. **Sixth public QAM line (realized in archival stage): closure-liquid-crystal, addressed role layer, theorem-level shadows, and terminal crystal–shadow interface.** Re-read OP 3 as local cell compatibility, OP 5 as defect transport on the same cellular carrier, visible sectors as liquid-crystal domains of one common bulk order, Julia-type diagnostics as holographic boundary readout, stabilize a first base36/Q-word role addressing with reserved 0-band transport support, and connect the resulting terminal crystal-readout surface to a bundled theorem-level shadow package and a terminal crystal–shadow interface.

At the present internal stage, the extraction criteria for the sixth public QAM line should remain strict. A QAM block should be promoted into a public release only if it satisfies all of the following conditions:

1. **compression:** the public block can be read in a compact main-text form without forcing the full internal package into the public line;

2. **dependency clarity:** the block has a clean dependency guide and does not rely on hidden internal lemmas;

3. **ZF-groundedness:** the coded language remains explicitly readable as a definitional overlay over ZF rather than as an unexplained new ontology;

© 2026 Marc Brendecke. Preprint — not peer reviewed.

4. **translation contact:** the public block makes clear what part of the current OP/branch/shell architecture it actually translates;
5. **non-overreach:** the public block does not yet claim more than the present coded layer genuinely supports.

Remark H2.11. *The current internal package already satisfies much of this extraction logic, but it is still broader than what should be exported immediately. The next internal passes should therefore favor tensor/operator compression, translation clarity, and clean staging over further rapid enlargement.*

Remark H2.12 (How the present release-pass checkpoint fits the public-shape roadmap)**.** *The deferred kernel-first checkpoint developed later in this file should be read as an internal support block for the same archival stage public-shape preparation path, not as a second release claim. Its current function is to prepare a later proof-compression interface for the compact tensor/operator chain by isolating the future kernel vocabulary, bridge data, staged stress tests, and activation criteria needed for a controlled migration. In particular, the checkpoint helps explain why the third public QAM line should remain narrow at visible level while still allowing later internal compression work to accumulate behind it.*

Compressed public-shape QAM branch-shell core

The second public QAM line should expose only the smallest next coded layer needed to carry the current admissible OP fragment and its branch/shell/routed refinement into a ZF-grounded formal interface. At compressed level, this public-shape branch-shell core should be read through six ingredients:

1. tagged QAM-coded objects and the minimal predicate layer;
2. admissible transport together with coded closure-class preservation;
3. closure-compatible coded readout;
4. closure-admissible branch codes;
5. shell-admissible refinement codes;
6. admissible routed support together with first coded branch-shell transport statements.

Theorem H2.13 (Compressed public-shape QAM branch-shell theorem)**.** *Assume that coded QAM objects are read as a definitional overlay over ZF, that admissible coded transport preserves coded closure class, and that the relevant coded readout is closure-compatible. Assume further that closure-admissible branch codes and shell-admissible refinement codes preserve coded closure class, that normalized admissible routed supports use only such branch-shell pairs, and that the admissible OP fragment is translated faithfully into the coded QAM layer. Then the translated source/return closure content of the admissible OP fragment, together with its admissible branchwise, shellwise, and routed realizations, is preserved at coded level, and closure-relevant readout agreement is retained after passage through the closure-factor map.*

Remark H2.14. *This compressed theorem is the intended mathematical core of the second public QAM line. It remains deliberately weaker than the full internal QAM package: tensor rules, operator/lambda maps, and support-selection layers remain available internally, but they need not all be exported at once in order to make the next public QAM step mathematically meaningful.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Structural systems view: QAM as a virtual machine with a ZFS-style theorem store

Status. *Active explanatory overlay, not a second mathematical criterion.*

Block function. This block explains why the QAM objects, kernel states, snapshots, and safe side branches belong to one architecture, without changing the theorematic content.

The present QAM export should also be read through a systems-level analogy. The visible coded tuple language plays the role of a readable virtual-machine interface over the ZF-grounded base, while the later exposed payload kernels and kernel normal forms behave like the machine-state identities that remain stable beneath several visible tuple surfaces. In that sense, the public QAM line is the user-facing instruction and state surface, whereas the deeper kernel-first line is the later OS-level state-management discipline. This reading is not introduced as a replacement for the mathematics, but as a compact way to explain why branch/shell/routed transport, kernel normal forms, search keys, snapshots, and safe side branches fit together as one contiguous architecture. It also aligns with the earlier Zuse Z3 comparison already used in the project: the current point is to interpret the coded QAM layer not merely as notation, but as a structured machine view with auditable state persistence.

Remark H2.15 (VM/OS reading of the current QAM line)**.** *For the present document, the systems analogy should be read in five layers.*

1. ***Visible QAM views.*** *The active tuple surfaces such as $\langle 0, a\rangle$, $\langle 0, a, b\rangle$, branch codes, shell/refinement codes, and routed return codes form the readable public interface of the machine.*

2. ***Kernel / VM state.*** *The exposed payload kernels, their normal forms, and the resulting class-level closure/readout data form the later machine-state identity beneath that visible interface.*

3. ***OS-level orchestration.*** *Admissible transport, local branch/shell return, and routed family aggregation act like the current system-level transition and scheduling layer that moves visible states while preserving closure-relevant content.*

4. ***ZFS-style theorem store.*** *Search keys, snapshots, clones, audits, and later proof-compression branches belong to an internal persistence layer that manages state identity, safe side branches, and deduplication of repeated kernel classes.*

5. ***Pool/group structure.*** *Sphere objects may later be treated as logical devices, while trioles or routed-support families may be grouped as internal device families. At the present stage, mirror-style reuse of equal kernel classes is the clearest fit; parity-like reconstruction remains exploratory and should be treated only as a later structural option.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.



Figure 2: Graphical reading of the current QAM export as a virtual-machine/OS stack with a ZFS-style structural store. Visible tuple surfaces provide the public theorem interface; exposed payload kernels and kernel normal forms provide the later machine-state identity; snapshots, clones, search keys, and audits manage the internal proof store without changing the visible public QAM syntax.

Remark H2.16 (Why the ZFS analogy is structurally useful here)**.** *The ZFS-style reading is not meant as decorative metaphor. It isolates four practical gains already visible in the present file. First, equal kernel classes can later be treated as mirror-like state reuse rather than as repeatedly independent visible tuple surfaces. Second, the deferred kernel-first path already behaves like copy-on-write: experimental compression statements are developed beside the active tuple proofs without rewriting the active line. Third, checkpoints, audits, and activation criteria behave like snapshots and scrub rules for the theorem store, since they record exactly which deferred branches are stable and which still remain non-authoritative. Fourth, once kernel normal forms and search keys are in place, repeated branch/shell/routed return families can be indexed and deduplicated as one structural graph beneath a readable tree-shaped theorem presentation. This is precisely why the systems analogy deserves a more prominent place next to the public QAM extraction itself.*

Remark H2.17 (Current scope of the VM/OS/ZFS view)**.** *In the inherited archival stage public-shape layer this systems view remains explanatory and organizational. It does not introduce a second formal semantics, and it does not yet activate RAID-like reconstruction as an active theorem rule. The present mathematically justified gains at that inherited layer are the mirror-like reuse of equal kernel classes, the snapshot/clone logic of active versus deferred proof branches, and the content-addressed search-key discipline attached to kernel normal forms. The archival stage extension below activates only a guarded store-and-array vocabulary over the already available QAM/MML/MMA interface. It still does not turn mirror, stripe, parity, or repair language into an OP-closure certificate.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

QAM/MML/MMA store layer and Sphaera array extension

Status. *Active archival stage infrastructure block; no OP closure claim.*

Block function. This block internalizes the earlier ZFS/RAID analogy. QAM supplies the object store, MML supplies the store and array instructions, MMA supplies the execution/readout semantics, and finite Sphaera arrays supply guarded multi-object redundancy, comparison, and residual/parity witnesses.

The preceding systems view should no longer be read merely as an external metaphor. At the present local layer it becomes a controlled internal interface. The point is not to claim that the Companion has already become a universal computer, nor that OP 1/4 or OP 5(B) have closed. The point is narrower and useful: the same QAM/MML/MMA machinery that already records objects, instructions, and route-faithful execution can also write the persistence layer in which several Sphaera objects are stored, compared, snapshotted, scrubbed, and carried as a guarded array.

Human reading. A single Sphaera object is now treated like one protected theorem-state device. Several Sphaera objects may be grouped into an array. In mirror mode, different objects carry the same structural content through different admissible views. In stripe mode, one larger proof or diagnostic object is distributed over several component stores. In parity/residual mode, an additional Sphaera object carries the difference, obstruction, or orthogonal control witness attached to the other components.

LLM/audit reading. The layer below gives stable machine-readable handles for store state, snapshot, clone, rollback, scrub, array creation, mirror, stripe, parity, and array snapshot. These handles are not theorematic shortcuts. They are audit contracts: each command must preserve address discipline, guard discipline, lineage discipline, and the non-certificate status unless a separate explicit theorem later discharges that status.



Figure 3: Reader-facing systems map for the QAM/MML/MMA store-array extension. QAM supplies the stored object layer, MML supplies the allowed store and array commands, MMA supplies the guarded execution semantics, and the Sphaera array layer packages mirror, stripe, and parity/residual modes together with snapshot, clone, rollback, and scrub operations.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Layer	Human reading	LLM/audit reading	Typical outputs
QAM	protected object store	typed address/object/guard/ledger state	store state, snapshot anchor
MML	allowed store and array actions	finite instruction alphabet	`STORE`, `SNAPSHOT`, `PARITY`
MMA	how commands really act	deterministic guarded transition semantics	continue / halt / reject / guard
Sphaera array	several Sphaera objects coupled as one working array	finite guarded family with cross-sphere guards and parity witnesses	mirror, stripe, parity/residual, scrub

Table 2: Compact overview table for the store-array layer. This separates reader intuition from the audit contract without changing the theorematic status of the block.

Definition H2.18 (QAM store state)**.** *A* QAM store state *is a finite guarded tuple*

$$\mathcal{S}_n := (\mathcal{A}_n, \mathcal{O}_n, \mathcal{R}_n, \mathcal{G}_n, \mathcal{L}_n),$$

where $\mathcal{A}_n$ is the finite address/tag domain, $\mathcal{O}_n$ is the finite family of stored QAM objects, $\mathcal{R}_n$ is the finite relation/link register between stored objects, $\mathcal{G}_n$ is the finite guard register, and $\mathcal{L}_n$ is the finite lineage/audit ledger. The state is admissible *if every ledger pointer lands in $\mathcal{A}_n$, every relation in $\mathcal{R}_n$ uses stored addresses, and every active guard in $\mathcal{G}_n$ is satisfied by the displayed objects and relations.*

Definition H2.19 (Store instruction alphabet)**.** *The local MML store alphabet is the finite instruction family*

$$\mathcal{I}_{\text{store}} := \{\texttt{STORE}, \texttt{LINK}, \texttt{SNAPSHOT}, \texttt{CLONE}, \texttt{ROLLBACK}, \texttt{SCRUB}, \texttt{MERGE_CAND}, \texttt{EXPORT}\}.$$

Here `STORE` *writes an object to an address,* `LINK` *writes a relation,* `SNAPSHOT` *freezes an admissible state,* `CLONE` *opens a guarded side branch from a snapshot,* `ROLLBACK` *returns to a recorded snapshot,* `SCRUB` *checks the guards and lineage,* `MERGE_CAND` *proposes a guarded merge final, and* `EXPORT` *exposes a readout surface. None of these instructions is allowed to manufacture a theorematic certificate by syntax alone.*

Definition H2.20 (MMA store transition semantics)**.** *The local MMA store semantics is a partial deterministic transition map*

$$\delta_{\text{store}} : \mathcal{S}_n \times \mathcal{I}_{\text{store}} \dashrightarrow \mathcal{S}_{n+1} \times \{\mathsf{continue}, \mathsf{halt}, \mathsf{reject}, \mathsf{guard}\}.$$

It is guard-faithful *if every defined transition either preserves admissibility of the resulting store or returns the status* guard *together with a defect entry in the ledger. It is* lineage-faithful *if each defined transition records enough ledger data to identify its source state, instruction, guard status, and output state.*

Definition H2.21 (Guarded snapshot and clone)**.** *For an admissible QAM store $\mathcal{S}_n$, a* guarded snapshot *is a tuple*

$$\operatorname{snap}(\mathcal{S}_n) := (\mathcal{S}_n, \chi_n, \lambda_n),$$

where χ_n is a finite integrity witness for the address, relation, guard, and ledger registers, and λ_n is a lineage anchor. A clone *of* $\operatorname{snap}(\mathcal{S}_n)$ *is a new working state $\mathcal{S}'_n$ with the same stored content and a new branch tag in the ledger. The clone is not a new theorem; it is a new guarded working branch from the same stored state.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proposition H2.22 (Snapshot stability of admissible QAM stores)**.** *Every admissible finite QAM store state admits a guarded snapshot in the sense of Definition H2.21. If the MMA store semantics is guard-faithful and lineage-faithful, then* `SNAPSHOT`*,* `CLONE`*, and* `ROLLBACK` *preserve the distinction between stored mathematical objects and release/audit metadata.*

Proof. Since the address, object, relation, guard, and ledger registers are finite, their displayed state can be recorded as a finite tuple together with a finite integrity witness and a lineage anchor. This gives $\mathrm{snap}(\mathcal{S}_n)$. Guard-faithfulness prevents a failed guard from being silently stored as an admissible state, and lineage-faithfulness records the source and branch data. Thus snapshot, clone, and rollback may move between recorded store states, but they do not promote a release label, branch label, or audit label into the mathematical object language. □

Definition H2.23 (Sphaera array)**.** *A* Sphaera array *is a finite guarded family*

$$\mathbb{S}_n := \big((\mathcal{S}_n^{(1)}, \dots, \mathcal{S}_n^{(k)}); \Gamma_n, \Pi_n, \Lambda_n\big), \qquad k \geq 1,$$

where each $\mathcal{S}_n^{(i)}$ is an admissible QAM store state, Γ_n is the cross-sphere guard register, Π_n is the parity/residual witness register, and Λ_n is the array-level lineage ledger. The array is admissible if all component stores are admissible and all cross-sphere guards in Γ_n are satisfied.

Definition H2.24 (Array instruction alphabet)**.** *The local MML array alphabet is*

$$\mathcal{I}_{\mathrm{array}} := \{\texttt{ARRAY_CREATE}, \texttt{MIRROR}, \texttt{STRIPE}, \texttt{PARITY}, \texttt{SCRUB_ARRAY}, \texttt{REPAIR_CAND}, \texttt{SNAPSHOT_ARRAY}\}.$$

The command `REPAIR_CAND` *intentionally returns a final repair surface, not an automatic mathematical repair theorem. The command* `PARITY` *records a residual or orthogonal control witness, not a proof that the residual vanishes.*

Definition H2.25 (Mirror, stripe, and parity modes)**.** *Let $\mathbb{S}_n$ be an admissible Sphaera array.*

1. *In* mirror mode*, two component stores $\mathcal{S}_n^{(i)}$ and $\mathcal{S}_n^{(j)}$ are coupled by a cross-guard requiring equality of the relevant closure-compatible readout, even if the visible store presentations differ.*

2. *In* stripe mode*, an object X is stored as a finite guarded decomposition $X \rightsquigarrow (X_1, \dots, X_k)$, with each stripe X_i stored in a specified component and with a reconstruction witness recorded in Π_n.*

3. *In* parity/residual mode*, one component or register stores a residual witness P for a guarded comparison of other components. The condition $P = 0$, if ever needed, must be proved separately; it is not asserted by the existence of the parity register.*

Definition H2.26 (Triadic Sphaera array)**.** *A* triadic Sphaera array *is an admissible Sphaera array of the form*

$$\mathbb{S}_n^{ABC} := (\mathcal{S}_A, \mathcal{S}_B, \mathcal{S}_C; \Gamma_{ABC}, \Pi_{ABC}, \Lambda_{ABC})$$

in which $\mathcal{S}_A$ and $\mathcal{S}_B$ carry two active readout or probe-facing stores, while $\mathcal{S}_C$ carries the orthogonal residual/parity witness. The guard register records the structural condition that $\mathcal{S}_C$ is read orthogonally to the active A/B coupling channel and that the HC/MM readout of the array preserves the inherited route tag.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proposition H2.27 (Guarded array snapshot)**.** *Every admissible finite Sphaera array admits a guarded array snapshot*

$$\mathrm{snap}_{\mathrm{arr}}(\mathbb{S}_n) := \big(\mathbb{S}_n, (\chi_n^{(1)}, \dots, \chi_n^{(k)}), \chi_n^{\Gamma}, \chi_n^{\Pi}, \lambda_n^{\mathrm{arr}}\big),$$

preserving component stores, cross-sphere guards, parity/residual witnesses, and array lineage.

Proof. Apply Proposition H2.22 to each finite component store. Since the cross-guard, parity/residual, and array-ledger registers are also finite, record their integrity witnesses and one array lineage anchor. The resulting tuple is a guarded array snapshot. It records the array state without adding an OP-closure certificate and without changing any component object into metadata. □

Proposition H2.28 (Finite array scrub)**.** *For every finite Sphaera array there is a deterministic scrub procedure*

$$\mathrm{scrub}_{\mathrm{arr}}(\mathbb{S}_n) \in \{\mathsf{pass}, \mathsf{fail}\} \times \mathcal{D}_n,$$

where $\mathcal{D}_n$ is a finite defect register. The scrub passes exactly when all component-store guards, cross-sphere guards, parity/residual registrations, and lineage pointers pass their finite checks.

Proof. The component store registers, cross-guard register, parity/residual register, and lineage ledger are finite by definition. A deterministic scrub checks them in a fixed order and records every failed check in $\mathcal{D}_n$. If no failed check is recorded, the result is pass; otherwise it is fail with the finite defect register. No repair or closure claim is produced by this test. □

Theorem H2.29 (QAM/MML/MMA Sphaera array layer)**.** *The QAM/MML/MMA stack admits a conservative Sphaera array layer in the following sense. Given finite admissible QAM stores, the store instruction alphabet of Definition H2.19, the array instruction alphabet of Definition H2.24, and guard-faithful lineage-faithful MMA transition semantics, one obtains finite guarded snapshots, finite array snapshots, and deterministic array scrub surfaces. These surfaces preserve stored mathematical objects, cross-sphere guards, residual/parity witnesses, and audit lineage without manufacturing an OP-closure certificate.*

Proof. The store part follows from Definitions H2.18–H2.21 and Proposition H2.22. The array part follows by grouping finitely many admissible stores as in Definition H2.23, assigning finite cross-guards, finite parity/residual witnesses, and a finite lineage ledger, and then applying Propositions H2.27 and H2.28. The last clause follows from the explicit non-certificate restrictions in Definitions H2.19, H2.24, and H2.25: mirror, stripe, parity, scrub, and repair-final language records infrastructure and diagnostic data only. It does not assert vanishing residuals or OP closure. □

Corollary H2.30 (RAID-like reading without RAID overclaim)**.** *The Sphaera array layer supports a RAID-like reading: mirror mode corresponds to redundant closure-compatible readout, stripe mode corresponds to guarded distribution of a larger object over several component stores, and parity/residual mode corresponds to an explicit diagnostic witness for cross-component mismatch. This reading is internal to QAM/MML/MMA, but it remains weaker than technical RAID reconstruction and weaker than any OP-closure theorem.*

Proof. This is only a translation of Definition H2.25 into systems language. The mathematical content remains the guarded store, snapshot, array, parity/residual, and scrub structure already proved above. □

Remark H2.31 (Tool extraction from the Sphaera array layer)**.** *The new layer extracts four reusable tools for later Companion revisions.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

1. ***Snapshot tool.*** *Freeze one admissible QAM store without changing the mathematical object language.*

2. ***Clone/rollback tool.*** *Open or return from a guarded branch while preserving lineage and non-certificate status.*

3. ***Array-scrub tool.*** *Test several Sphaera stores, cross-guards, and residual witnesses as one finite diagnostic surface.*

4. ***Triadic parity tool.*** *Use a third, orthogonal Sphaera object as residual/parity witness for a two-store active coupling.*

These tools are deliberately infrastructural. Their value is that later OP-facing work can compare, branch, and roll back several coupled Sphaera objects without confusing state persistence with theorematic closure.

Remark H2.32 (Human/LLM interface gain)**.** *For a human reader, the store-array layer explains why the Companion keeps snapshots, historical witnesses, audit ledgers, and branch restrictions so visibly: they are the theorem-store equivalent of protected state persistence. For an LLM reader, the same layer supplies exact handles: $\mathcal{S}_n$ for one store, $\mathbb{S}_n$ for an array, Γ_n for cross-guards, Π_n for parity/residual witnesses, and Λ_n for lineage. This is the intended compromise between human readability and machine precision.*

Pre-shift QAM Hilbert-shift and guarded mixed readouts

> **Status.** *Pre-shift preparation layer only.* The inherited QAM tuple tags are not globally migrated in Version 4.7.0. The block below records the conservative Hilbert-shift plan, the mixed-readout package discipline, and an optional binary mask registry for future readout classification.

The index-0 issue is now treated explicitly. In the inherited QAM tuple namespace, tag 0 occurs as an inhabited constructor tag, for instance on visible tuple surfaces of the form $\langle 0, a\rangle$. In a Boolean mask or IPshifted-like readout registry, however, value 0 must mean no active kind. The next major QAM migration should therefore not identify these two uses. The preparation layer below records the planned Hilbert-hotel shift that can free tag 0 in the future while preserving the payloads and proofs.

Definition H2.33 (QAM Hilbert-shift plan)**.** *Let $\mathcal{K}_{\mathrm{old}}$ be the inherited finite QAM tuple-tag namespace used by the current public syntax, and let $\mathcal{K}_{\mathrm{new}}$ be a shifted namespace with tag 0 reserved for the empty/no-active-kind marker. The planned Hilbert-shift is the constructor-level map*

$$H_{\mathrm{QAM}}(\langle i, x_1, \ldots, x_m\rangle) := \langle i+1, x_1, \ldots, x_m\rangle.$$

It shifts only the constructor tag. It does not change the stored payload $(x_1, \ldots, x_m)$, the intended constructor distinction, or any mathematical content carried by the payload.

Definition H2.34 (Pre-shift reserved-zero discipline)**.** *The preparation line distinguishes two uses of zero.*

1. *In the inherited legacy tuple syntax, tag 0 remains an inhabited tuple tag wherever the old QAM definitions still use it.*

2. *In the planned shifted shifted namespace and in every binary mask registry, tag or mask value 0 is reserved for empty, undefined, or no active kind.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

No Version 4.7.0 statement may silently replace the first convention by the second. Any global replacement is reserved for a later shifted migration theorem.

Proposition H2.35 (Conservative shape of the QAM Hilbert-shift)**.** *Suppose a finite QAM sublanguage is translated by H_{QAM} and every constructor predicate, admissibility predicate, readout predicate, and closure-relevant predicate is translated by the same tag shift. Then the translated sublanguage is a conservative reindexing of the old one: payloads, constructor distinctions, admissibility status, readout compatibility, and closure-relevant truth values are preserved under the translation.*

Proof. The map H_{QAM} is injective on constructor tags and leaves all payload coordinates unchanged. A constructor predicate in the shifted namespace is defined to hold exactly when the corresponding old predicate holds before applying the shift. The same translation is imposed on admissibility, readout, and closure-relevant predicates. Thus every proof step that used only constructor distinction, payload data, admissibility, readout compatibility, or closure-relevant class data has an isomorphic shifted proof step. The proposition records the required conservativity shape for a later shifted migration; it does not perform the global migration here. □

Definition H2.36 (Guarded mixed QAM readout)**.** *A* guarded mixed QAM readout *is a finite derived package*

$$R_{\mathrm{mix}} := (r_1, \ldots, r_m; \Gamma_{\mathrm{mix}}, \mu, J_{\mathrm{mix}}, \sigma_{\mathrm{mix}}),$$

where each r_i is an already admissible component readout on its own allowed QAM kind, Γ_{mix} is a finite guard package, μ is a finite aggregation map on component readout values, J_{mix} is the closure-factor map for the aggregate, and σ_{mix} records the theorematic status. It is admissible only when all component readouts are admissible and the aggregation map does not identify non-equivalent closure-relevant classes.

Definition H2.37 (Binary mask registry for mixed readouts)**.** *A* binary mask registry *for mixed QAM readouts assigns to each inhabited readout kind a nonzero power-of-two bit*

$$\beta_j = 2^j, \qquad j \geq 0,$$

and reserves mask value 0 for no active kind. A mixed-readout mask is a finite Boolean OR of active kind bits. Boolean AND tests whether a component kind is present, XOR records symmetric difference between two masks, and complement is allowed only relative to a fixed finite mask universe Ω.

Proposition H2.38 (Mixed readout factorization)**.** *Let R_{mix} be a guarded mixed QAM readout whose component readouts r_i are closure-compatible. If the aggregation map μ factors through the component closure-factor maps and J_{mix} factors through the resulting aggregate class, then R_{mix} is closure-compatible.*

Proof. Closure-compatible component readouts identify no closure-relevant distinction beyond their component class maps. Since μ factors through those class maps, it can only aggregate already class-level data. Since J_{mix} factors through that aggregate class, the mixed readout cannot introduce a new closure-relevant distinction that was invisible to the component closure classes. Therefore the mixed package is closure-compatible. The argument uses only factorization and guards; it does not introduce primitive mixed tensors. □

Remark H2.39 (shifted migration trigger)**.** *The full replacement of the inherited tuple tags by the shifted tags is intentionally not a Version 4.7.0 operation. It changes the visible QAM object-language namespace and therefore belongs to a major version. Version 4.7.0 only records why the shift is useful, what the conservative proof obligation looks like, and how mixed readouts can already be treated as derived packages without forcing the migration.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Two-sided structural tensor notation for QAM/MML/MMA readouts

Status. *Active notation and compression layer; not a new tensor calculus and not a new closure criterion.*

Block function. This block records the narrow two-sided tensor grammar used to separate readout-side, structure-side, comparison-side, and mask-side roles in later QAM/MML/MMA expressions. It is included here as a conservative technical layer only. A fuller pedagogical reader manual for the notation is intentionally kept as a separate future paper or companion guide.

The QAM/MML/MMA line repeatedly alternates between four roles: frame-side readout, structure-side action, comparison of active channels, and compatibility testing of readout or mask modes. The following notation is introduced only to make those roles visible in the formula syntax. It does not replace classical tensor calculus and it does not turn every displayed expression into a new primitive tensor object.

Definition H2.40 (Two-sided structural tensor interface)**.** *A* two-sided structural tensor display *is a tensor-like expression in which a structural object may carry front-side and back-side indices, written schematically as*

$$ {}^{A}{}_{B}\mathcal{T}^{i}{}_{j}. $$

The front indices ${}^{A}{}_{B}$ *record the left-sided or readout-facing interface: frame, observer, language, readout, or external addressing data. The back indices* ${}^{i}{}_{j}$ *record the right-sided or structure-facing interface: action, transport, field, operator, or internal structural data. Thus the display is read as a typed structural interface rather than as a mere component array.*

Definition H2.41 (Admissible contractions and comparisons)**.** *The two-sided notation obeys the following contraction discipline.*

1. ***Classical action/readout.*** *Ordinary upper–lower contraction remains unchanged. If*

$$ A_{\mu}\,{}^{\mu}\mathcal{T}_{\nu}\,B^{\nu} $$

is well-defined, it is read as a classical action, evaluation, or sandwich readout: first apply the structure-facing operation to B*, then read the result by* A*.*

2. ***Active–active comparison.*** *Two upper indices are not contracted as a raw Einstein sum. They may only be compared through an explicitly declared comparison tensor, for instance*

$$ G_{\alpha\beta}X^{\alpha}Y^{\beta}. $$

This means active structural comparison, such as angle, distance, coupling strength, orthogonality, channel similarity, or residual size.

3. ***Passive–passive comparison.*** *Two lower indices are likewise not contracted raw. They may only be compared through an explicitly declared dual comparison tensor, for instance*

$$ G^{\alpha\beta}\omega_{\alpha}\eta_{\beta}. $$

This means comparison of readout forms, observer modes, measurement channels, or passive projections.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

4. ***Mask and selector compatibility.*** *Boolean, finite, or mask-like compatibility tests must be carried by an explicit selector tensor, for instance*

$$\chi_{mn} R^m S^n.$$

This expression says that readout modes m and n are first tested for admissible compatibility by χ, and only compatible contributions are admitted to the mixed readout.

Remark H2.42 (Conservative reading rule)**.** *The two-sided notation is conservative because the only unchanged contraction is the classical upper–lower action/readout contraction. Same-variance pairings are not silently added to the calculus; they are admitted only when a comparison tensor or selector tensor has been declared. Hence every two-sided display must be expandable into ordinary QAM-over-ZF role predicates together with explicit comparison or compatibility data.*

Definition H2.43 (Guarded mixed QAM readout)**.** *Let R^m and S^n be QAM readout modes and let χ_{mn} be an explicit admissibility or compatibility selector. A* guarded mixed QAM readout *is a derived readout expression of the form*

$$\mathrm{Mix}_\chi(R,S)(x) \quad \rightsquigarrow \quad \chi_{mn} R^m(x) S^n(x),$$

where the right-hand display is read with the contraction discipline of Definition H2.41. The mixed readout is therefore not a new primitive QAM object. It is a compatibility-filtered package of already admitted readout modes.

Remark H2.44 (Residual comparison grammar)**.** *The live residual triple*

$$\rho = (\rho_{\mathrm{CL}}, \rho_\Theta, \rho_A)$$

may be compressed into a residual-channel vector ρ^α. Its total comparison size may then be written as

$$\|\rho\|_G^2 = G_{\alpha\beta}\rho^\alpha \rho^\beta.$$

This is not a raw contraction of two upper indices. It is the comparison of two active residual-channel entries through the declared comparison tensor $G_{\alpha\beta}$.

Remark H2.45 (HC/MM sandwich grammar)**.** *An HC/MM-mediated readout may be displayed schematically as*

$${}^{R}{}_{\Omega}\mathcal{M}^{i}{}_{j},$$

where the front side records the readout or observer-side interface and the back side records the structure-side transport or matrix action. When paired with ordinary upper–lower indices, the expression remains an ordinary sandwich readout. When two readout channels or two structure channels are compared, the comparison must be mediated by an explicitly named comparison tensor or selector.

Proposition H2.46 (Conservative expansion of two-sided displays)**.** *Every well-formed two-sided structural tensor display in the sense of Definitions H2.40 and H2.41 admits a conservative expansion into the inherited QAM-over-ZF language: front-side indices expand to readout/frame/language role data, back-side indices expand to structure/action/transport role data, classical upper–lower contractions remain ordinary action or evaluation clauses, and every same-variance pairing expands to an explicit comparison or compatibility predicate.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. Erase the visual sidedness notation and retain its role data as ordinary QAM codes. Front-side symbols become readout, frame, observer, or language tags. Back-side symbols become structure, action, transport, or field tags. Classical upper–lower contractions are left unchanged. A same-variance display is well formed only if it contains a named comparison tensor or selector, and so it expands to the corresponding comparison or compatibility predicate. No new primitive closure relation, admissibility relation, or theorematic certificate is introduced by this erasure. The display is therefore a conservative abbreviation of already typed QAM-over-ZF data. □

Remark H2.47 (LLM-facing role stability)**.** *For machine reading and LLM-assisted proof navigation, the notation functions as a syntactic guardrail. The front/back split marks whether a symbol belongs to the readout-facing side or the structure-facing side, while the contraction discipline marks whether a formula is an action, a readout, a comparison, or a compatibility test. This reduces the risk that a readout is mistaken for an operator, that a diagnostic comparison is mistaken for a proof of closure, or that a guarded mixed readout is mistaken for a new primitive tensor object.*

Remark H2.48 (Use in the present release line)**.** *In the present historical release line.x line this notation should be used locally and sparingly: first for guarded mixed QAM readouts, residual/drift comparisons, HC/MM sandwich expressions, triadic orthogonality comparisons, and Boolean or mask-like compatibility tests. A global rewrite of the whole Companion into the two-sided notation is not part of this release line. The separate pedagogical reader manual may later explain the notation at a slower pace without enlarging the active theorem core.*

stability layer: two-sided expansion, mixed-readout normal form, and Boolean masks

> **Status.** *Conservative stabilization of the notation layer; not a global rewrite, not a new tensor calculus, and not a closure theorem.*
>
> **Block function.** This block fixes the exact erasure route from two-sided displays back to ordinary QAM-over-ZF role data, normalizes guarded mixed readouts, and states the Boolean selector grammar used by IPshifted-like readout masks. The separate reader manual remains a future standalone paper or guide.

The notation made readout-side, structure-side, comparison-side, and mask-side roles visible in formulas. The present step turns that notation into a safer tool by recording the normal forms that an audit reader or LLM reader should recover whenever a two-sided display appears. The rule is intentionally asymmetric: notation may compress role data, but every compressed display must expand back into explicit role data before it can be used as proof content.

Definition H2.49 (Two-sided role-erasure map)**.** *Let E be a well-formed two-sided structural tensor display in the sense of Definitions H2.40 and H2.41. Its* role-erasure expansion

$$\mathfrak{E}_{\mathrm{role}}(E)$$

is the ordinary QAM-over-ZF package obtained by replacing:

1. *every front-side index by an explicit readout, frame, observer, or language-role tag;*
2. *every back-side index by an explicit structure, action, field, transport, or operator-role tag;*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

3. *every classical upper–lower contraction by the corresponding ordinary action/evaluation clause;*

4. *every same-variance comparison by a named comparison predicate carrying its declared comparison tensor;*

5. *every Boolean or selector occurrence by a finite compatibility predicate over the declared mask universe.*

This expansion is an erasure of notation only. It does not erase guards, selectors, comparison tensors, branch restrictions, or provenance data.

Proposition H2.50 (Role-erasure normal form)**.** *Every well-formed two-sided display E used in the historical release line.x line has a role-erasure expansion $\mathfrak{E}_{\mathrm{role}}(E)$ whose clauses are built from already admitted QAM role predicates, admitted comparison data, admitted selector data, and ordinary finite tuple coding over ZF. In particular, the two-sided display cannot by itself introduce a new primitive tensor kind, a new admissibility relation, or a new theorematic certificate.*

Proof. Proceed by structural inspection of the display grammar. Front/back sidedness contributes only role tags. Ordinary upper–lower contraction is already part of the inherited action/readout grammar. A same-variance pairing is well formed only when it contains an explicitly named comparison tensor or selector, hence it expands to a comparison or compatibility predicate rather than to an untyped Einstein contraction. All finite lists of such clauses are QAM tuple-coded over the existing ZF substrate. Therefore the two-sided syntax is conservative under role erasure. □

Definition H2.51 (Guarded mixed-readout normal form)**.** *A guarded mixed QAM readout is in* normal form *when it is presented as a finite package*

$$\mathcal{R}_{\mathrm{mix}} = \big(I, \{R_i\}_{i\in I}, \Gamma_{\mathrm{mix}}, \chi, \mu, J_{\mathrm{mix}}\big),$$

where I is a finite index set, each R_i is an already admitted component readout, Γ_{mix} is the inherited guard package, χ is a finite selector or mask predicate over I, μ is an aggregation map defined only on selector-admissible component families, and J_{mix} is the declared mixed readout surface. A two-sided display such as

$$\chi_{mn} R^m S^n$$

is only an abbreviation for this finite package in the binary case $I = \{m, n\}$.

Proposition H2.52 (Mixed-readout normal-form expansion)**.** *Every guarded mixed readout admitted by Definition H2.43 expands to a guarded mixed-readout normal form in the sense of Definition H2.51. Conversely, every guarded mixed-readout normal form may be displayed in two-sided shorthand only after its guards, selector, aggregation map, and output surface have been declared.*

Proof. The binary shorthand $\chi_{mn} R^m S^n$ already names the two component readouts and the selector. The surrounding QAM context supplies the inherited guard package, the admissible aggregation rule, and the declared output surface. Collecting these data gives the normal-form tuple. Conversely, starting from the normal-form tuple, one may suppress the finite package into two-sided shorthand only because the suppressed guard, selector, aggregation, and output data remain recoverable. Thus the shorthand and the normal form are not different mathematical objects; the former is a compact notation for the latter. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition H2.53 (Boolean selector and mask grammar)**.** *Fix a finite readout-kind universe* $\Omega = \{k_1, \ldots, k_N\}$*. A* Boolean selector mask *is a function*

$$\chi : \Omega \longrightarrow \{0, 1\}$$

or, equivalently, a finite bit mask whose value 0 denotes no active kind and whose nonzero bits denote active readout kinds. The admissible Boolean operations are:

$$\begin{aligned} \chi_A \vee \chi_B &\quad \textit{for union of active readout kinds,} \\ \chi_A \wedge \chi_B &\quad \textit{for intersection or compatibility testing,} \\ \chi_A \oplus \chi_B &\quad \textit{for diagnostic symmetric difference,} \\ \neg_\Omega \chi_A &\quad \textit{for relative complement inside the fixed universe } \Omega. \end{aligned}$$

No absolute complement is used without first declaring Ω*. A selector expression may filter a readout package, but it does not alter the payload of any component readout.*

Proposition H2.54 (Mask-filter conservativity)**.** *Let* $\mathcal{R}_{\mathrm{mix}}$ *be a guarded mixed-readout normal form and let* χ *be a Boolean selector mask over its finite readout-kind universe. Applying* χ *restricts the active component menu of* $\mathcal{R}_{\mathrm{mix}}$*, but it does not change the underlying component payloads, the inherited guards, or the closure status of any component readout.*

Proof. A selector mask is a finite predicate on the index universe. It chooses which already declared component clauses are active in the aggregation domain. It neither rewrites the component readouts nor changes their guard packages. Hence any closure-compatible status used by the mixed package still has to come from the component readouts and the declared factorization map, not from the mask itself. □

Remark H2.55 (LLM/audit role table)**.** *For later machine reading, the following table fixes the intended role parsing. It is a local audit aid inside the Companion, not the separate pedagogical reader manual.*

Displayed form	***Intended role***	***Audit guard***
front-side indices	*readout, frame, observer, language, or addressing side*	*expand to explicit QAM role tags*
back-side indices	*structure, action, field, transport, or operator side*	*expand to explicit QAM role tags*
upper–lower contraction	*classical action, evaluation, or sandwich readout*	*ordinary tensor/QAM readout clause*
upper–upper via $G_{\alpha\beta}$	*active structural comparison*	*no raw same-variance contraction*
lower–lower via $G^{\alpha\beta}$	*passive readout comparison*	*no raw same-variance contraction*
χ_{mn} *or a mask bit*	*compatibility or activation selector*	*filter only; no new payload*

Corollary H2.56 (proof-neutrality of the tensor layer)**.** *The tensor-stability layer may be cited to shorten repeated readout/action/comparison/mask passages, but it may not be cited as an independent proof of closure, OP resolution, primitive mixed-tensor existence, or global QAM namespace migration.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. By Propositions H2.50, H2.52, and H2.54, every display reduces to already declared role data, normal-form mixed-readout data, and finite selector predicates. The reduction supplies notation control only. It does not add the missing theorematic hypotheses needed for closure, OP resolution, primitive mixed tensors, or shifted reindexing. □

Remark H2.57 (Placement of the future reader manual)**.** *The present block deliberately stays short and formal. A slower reader manual may later explain the same grammar with diagrams, examples, and human-facing intuition, but that manual should remain a separate paper or guide so that the Active Core remains the governing proof line.*

release-final freeze audit for the pre-shift QAM/tensor layer

Status. *Release-final audit layer only; no new theorematic certificate, no global QAM namespace migration, no primitive mixed tensor, and no OP closure.*
Block function. This block freezes the local pass/notation and mixed-readout preparation layer for release-final review. It records what must remain true when the Active Core is read together with the separate Historical Witness PDF.

Definition H2.58 (pre-shift QAM/tensor release-final package)**.** *The* pre-shift QAM/tensor release-final package *consists of the following finite data:*

(i) the inherited archival stage public split-bundle baseline and its concept/version DOI record;

(ii) the archival stage short-abstract and extended-orientation architecture;

(iii) the pre-shift Hilbert-shift plan reserving global QAM tag migration for a later major version;

(iv) the guarded mixed-readout discipline and optional binary mask registry;

(v) the two-sided structural tensor notation;

(vi) the role-erasure, mixed-readout normal form, Boolean selector grammar, and proof-neutrality layer;

(vii) the present audit/handoff metadata saying that the Active Core is the governing proof line and that the Historical Witness PDF remains preserved proof memory.

It is a packaging and audit object, not a closure object.

Proposition H2.59 (release-final conservativity)**.** *The pre-shift QAM/tensor release-final package is conservative over the tensor-stability layer with respect to theorematic content. It may change release-final metadata, routing language, audit surfaces, and handoff wording, but it may not create any new OP certificate, global Hilbert-shift theorem, primitive mixed-tensor axiom, or replacement of the inherited QAM tuple namespace.*

Proof. Each mathematical role introduced earlier is erased by Definition H2.49 and normalized by Proposition H2.50. Each mixed readout is expanded by Proposition H2.52, and each Boolean mask action is restricted by Proposition H2.54. The package adds only release-final metadata and bundle-handoff statements around these already guarded objects. Hence no new theorematic content is introduced. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Corollary H2.60 (freeze-readiness condition)**.** *If the build audit reports no duplicate labels, no undefined references, no undefined citations, and no fatal LaTeX errors for both the Active Core and the Historical Witness, then Version 4.7.0 is structurally ready to be treated as a freeze final for the archival stage pre-shift QAM/tensor preparation release, subject only to the external release decision and any assigned Zenodo version DOI.*

Remark H2.61 (Reader-manual separation at)**.** *The pedagogical reader manual for the two-sided notation remains outside the Active Core. It may later be written as a separate guide with diagrams, examples, and slower explanations. The present Companion keeps only the compact formal grammar and the proof-neutrality guards needed by the governing proof line.*

archival stage DOI-bearing release freeze

Status. *Release-final metadata layer only; no new theorematic certificate, no global QAM namespace migration, no primitive mixed tensor, and no OP closure.*
Version DOI. The public version-specific Zenodo DOI for this release is

$$10.5281/\textit{zenodo}.19750674.$$

The Companion concept DOI remains 10.5281/zenodo.19210728. The version DOI is release metadata only and is not a mathematical subscript, superscript, route tag, or proof object.

Definition H2.62 (archival stage release-final package)**.** *The* archival stage release-final package *consists of the release-final package of Definition H2.58, the assigned public version DOI above, the Active Core PDF, the preserved Historical Witness PDF, the release manifest, checksums, audit report, Zenodo description, and release notes. It is a release object, not a theorematic closure object.*

Proposition H2.63 (archival stage release-final conservativity)**.** *The archival stage release-final package is conservative over the release-final package with respect to theorematic content. DOI insertion, release-note generation, manifest creation, checksum creation, and Zenodo-description preparation do not create any new OP certificate, global QAM Hilbert-shift theorem, primitive mixed-tensor axiom, or replacement of the inherited QAM tuple namespace.*

Proof. By Proposition H2.59, the package is conservative over the local pass. The present layer adds only release metadata and publication packaging: the version DOI, release file names, manifest/checksum records, audit report, Zenodo description, and release notes. None of these objects changes the payload of any QAM role, tensor grammar, mask selector, mixed readout, or OP-facing proof obligation. Hence theorematic content is unchanged. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

source handoff and router-language layer

Status. This archival stage layer is a conservative local TEX handoff over the public archival stage release-final baseline. It records the router-language, route-mask, HC-boundary, theorem-layer, and active-readout architecture as a compact source layer. It is not an OP closure, not a global shifted QAM namespace migration, and not a primitive mixed-tensor activation.

Definition H2.64 (handoff object)**.** *The* the local proof line handoff object *is the finite record*

$$\mathfrak{H}_{\mathrm{loc}} = (B_{3.35}, Q_{\mathrm{loc}}, M_{\mathrm{loc}}, T_{\mathrm{loc}}, P_{\mathrm{loc}}, G_{\mathrm{loc}}),$$

where $B_{3.35}$ is the public archival stage release-final baseline, Q_{loc} is the QAM/MML/MMA router-language package, M_{loc} is the route-mask and double-Hilbert-shift package, T_{loc} is the canonical theorem-layer package, P_{loc} is the active physical and mathematical readout package, and G_{loc} is the non-overclaim guard record.

Definition H2.65 (Router-language role split)**.** *The local router-language split is the guarded reading*

$$\begin{aligned} \mathrm{QAM} &= \textit{object and store grammar}, \\ \mathrm{MML} &= \textit{routing and command grammar}, \\ \mathrm{MMA} &= \textit{execution and readout grammar}. \end{aligned}$$

The combined router layer is the conservative compiler surface

$$\mathcal{R}_{\mathrm{QAM/MML/MMA}} : \textit{guarded object packages} \longrightarrow \textit{guarded readout packages}.$$

Remark H2.66 (Machine layer is not physical content)**.** *Definition H2.65 is a routing definition. It does not promote a router object to a physical observable, does not add a field equation, does not create a closure certificate, and does not replace the underlying tensor calculus. Physical content appears only after a guarded readout has a declared target, collar or boundary interface, comparison or residual map, and back-translation.*

Definition H2.67 (Router object)**.** *A local router object is a finite record*

$$\mathfrak{R} = (K, M, C, T, B, G),$$

where K is a finite kind registry, M is a finite route-mask registry, C is a finite classical-collar record, T is a finite trace record, B is a back-translation record into the prior source language, and G is the guard record.

Definition H2.68 (Guard record for router objects)**.** *The guard record G states which transformations are purely notational, which readouts are inherited from earlier admissible packages, and which claims are not being made. In this layer, G includes at least the statements that OP 1/4 and OP 5(B) remain open, primitive mixed tensors remain inactive, Turing-completeness is not claimed, and the global shifted QAM Hilbert-shift has not yet been performed.*

Lemma H2.69 (Router-object erasure)**.** *Every router object in the sense of Definition H2.67 has a proof-neutral erasure to its underlying guarded source package.*

Proof. The entries K, M, C, T, and B are finite routing, masking, collar, trace, and back-translation data. Removing these annotations leaves the underlying source package together with the guard record G. Since no new observable, field equation, or closure certificate is added by the router layer, the erasure is proof-neutral. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Remark H2.70 (Relation to two-sided structural tensor notation)**.** *When a router object is written with two-sided structural tensor notation, front-side indices mark frame, readout, or language roles, while back-side indices mark structure, action, or field roles. Classical upper–lower contraction remains the ordinary action/readout case. Same-variance comparison still requires an explicit comparison or selector tensor. Thus the router notation remains a conservative compression grammar over the local proof chain tensor layer.*

double-Hilbert-shift route-mask layer

Definition H2.71 (Double-Hilbert-shift route-mask convention)**.** *The local route-mask convention is*

$$H_0(k) = k + 1, \qquad H_1(a) = 2^{a-1} \quad (a \geq 1), \qquad H_{\mathrm{mask}}(k) = 2^k.$$

The value 0 *is reserved for the inactive or no-active-kind marker. A nonzero route mask is read through its finite bit support and can be back-translated into the corresponding prior unshifted kind labels.*

Definition H2.72 (Mask support and back-translation)**.** *For a finite nonzero mask* m*, let* $\mathrm{bits}(m)$ *be the finite set of active bit atoms in* m*. The back-translation map is*

$$\mathrm{back}(m) = \{a - 1 : a \in \mathrm{bits}(m)\}.$$

For an ordered route payload $(m_1, \ldots, m_n)$*, the route back-translation is applied componentwise:*

$$\mathrm{BackRoute}(m_1, \ldots, m_n) = (\mathrm{back}(m_1), \ldots, \mathrm{back}(m_n)).$$

Lemma H2.73 (Double-shift conservativity)**.** *The double-Hilbert-shift route-mask convention is conservative over the prior finite route-label language, provided that every active mask is decoded by Definition H2.72 before theorematic use.*

Proof. The first shift reserves the inactive state by separating 0 from inhabited labels. The second shift atomizes active labels into powers of two. Since every finite mask has finite bit support and the back-translation recovers the corresponding prior labels, the route-mask layer adds a finite audit encoding but does not alter the underlying guarded route package. Hence theorematic status is preserved. □

first boundary shift and HC activation

> **Purpose.** This subsection separates the local first shift from the later global shifted QAM Hilbert-shift. The local shift belongs to archival stage because it explains why the HC can be read as internally generated at the inactive/active boundary. The global reindexing of the full QAM language is still deferred to shifted.

Definition H2.74 (First boundary shift)**.** *The* first boundary shift *is the local boundary transition*

$$\Delta_{\mathrm{HC}} : 0_{\mathrm{mask}} \rightsquigarrow 1_{\mathrm{mask}},$$

from the inactive mask state to the first active mask atom. It is not a migration of the full QAM tag language. It is only the smallest local passage by which the readout grammar distinguishes no-active-kind from first active readout.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition H2.75 (Boundary normalizer)**.** *A* boundary normalizer *for the first boundary shift is a structural readout normalizer* N_{bd} *assigned to the interface*

$$(0_{\mathrm{mask}}, 1_{\mathrm{mask}}).$$

It is required to preserve the inactive side as inactive, admit the first active readout atom as active, and prevent the transition from being interpreted as an uncontrolled same-variance contraction or as an unguarded physical promotion.

Assumption H2.76 (HC activation at the first boundary)**.** *At the first boundary shift, the minimal boundary normalizer is read as the Heisenberg compensator:*

$$N_{\mathrm{bd}} = HC \quad \textit{at the structural readout level.}$$

This is an internal activation reading of the HC inside the route-mask grammar, not a replacement proof of the physical Heisenberg uncertainty relation.

Proposition H2.77 (Internal origin of the first HC shift)**.** *Under Assumption H2.76, the first boundary shift does not enter the archival stage architecture as an external add-on. It is forced as the local normalization site between the reserved inactive mask state and the first active readout atom.*

Proof. The route-mask grammar reserves 0_{mask} for no active readout kind and reads 1_{mask} as the first active mask atom. The transition between them is therefore not an ordinary active-active comparison, because one side is not yet active. It is also not an ordinary action/readout contraction, because there is no prior active object on the inactive side to evaluate. A normalizing interface is therefore required before the first active readout can be admitted. By the boundary-activation assumption, this minimal interface is read as HC. Hence the first shift is structurally internal to the readout machine, while the global QAM Hilbert-shift remains deferred. □

Corollary H2.78 (No shifted migration from the first shift)**.** *The first boundary shift licenses only the local HC-boundary reading. It does not perform the global shifted migration of inhabited QAM tags and does not change any inherited theorematic status.*

Proof. Definition H2.74 is local to the pair $(0_{\mathrm{mask}}, 1_{\mathrm{mask}})$. No rule is introduced that rewrites all inhabited tags, replaces inherited labels, or promotes router metadata to physical closure. Therefore the global shifted migration remains separate. □

HC half-boundary reading

Definition H2.79 (HC half-boundary readout)**.** *The* HC half-boundary readout *treats the transition*

$$0_{\mathrm{mask}} \longrightarrow 1_{\mathrm{mask}}$$

from the inactive mask state to the first active shifted mask atom as a boundary-normalization site. The forbidden formal value $H_1(0) = 1/2$ *is not a Boolean mask atom. It marks the structural edge at which the Heisenberg compensator may be read as the minimal frame/readout normalizer required for active readout emergence.*

Remark H2.80 (Non-Boolean boundary lift)**.** *The formal lift* $H_{\mathrm{mask}}(1/2) = \sqrt{2}$ *is a diagnostic analogy for the boundary, not a Boolean route-mask value. It must not be used as a bit, as a closure certificate, or as a proof of the physical Heisenberg uncertainty relation.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Assumption H2.81 (HC-boundary reading as structural hypothesis)**.** *The HC half-boundary reading is treated as a conservative structural hypothesis: the Heisenberg compensator may be interpreted as the boundary compensator at the inadmissible pre-mask boundary between inactive state* 0 *and first active shifted mask atom* 1*, but this interpretation is not yet a theorem of physical dynamics.*

router trace and bitstream layer

Definition H2.82 (Router trace)**.** *A* router trace *is the finite diagnostic record obtained by reading a route-mask payload as a bitstream together with its ordered back-translation, classical collar, and guard record. It records how a finite router object moves from stored object data to guarded readout data without changing theorematic status.*

Definition H2.83 (Bittrace normal form)**.** *For a route pair* (m, n)*, the bittrace normal form is*

$$\mathrm{BitTrace}(m, n) = \big(\mathrm{back}(m) \to \mathrm{back}(n),\, G\big),$$

where G *is the inherited guard record. The arrow records diagnostic flow only; it is not a proof of admissibility unless the underlying guarded source package already supplies admissibility.*

Lemma H2.84 (Router trace conservativity)**.** *Passing from a finite route-mask payload to its router trace is conservative with respect to OP status.*

Proof. The router trace records finite bit support, ordered back-translation, collar data, and guards. It neither removes the guard record nor supplies a missing closure trigger. Therefore OP 1/4 and OP 5(B) keep the status they had before the trace was introduced. □

canonical theorem-layer skeleton

Definition H2.85 (Canonical theorem-layer package)**.** *The local canonical theorem-layer package consists of the following theorem classes:*

1. *router representation and router-object erasure;*
2. *route-mask conservativity and double-shift back-translation;*
3. *router chaining and classical collar compatibility;*
4. *HC-boundary activation as a guarded structural reading;*
5. *bitstream and route-difference normal forms;*
6. *boundary-readout projection packages;*
7. *physical promotion criteria.*

Theorem H2.86 (Canonical router handoff)**.** *Every finite router-language package admits a canonical handoff into the active readout layer by separating proof-neutral routing metadata from guarded readout, comparison, boundary, and residual content.*

Proof. The package is finite by Definition H2.67. Its masks and traces are audit surfaces, while its collars, comparisons, boundary interfaces, and residual maps are the possible content surfaces. Separating these roles gives the claimed handoff. The guard record of Definition H2.68 prevents metadata from being read as a closure proof. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

active physical and mathematical readout layer

Definition H2.87 (Active readout package)**.** *The* active physical and mathematical readout package *is the finite record*

$$\mathcal{P}_{\mathrm{loc}} = (P_{\mathrm{HC}}, P_{\gamma}, P_{\mathrm{sing}}, P_{\mathrm{Cas}}, P_{\mathrm{OP}}, P_{\mathrm{auto}}, P_{\mathrm{ws}}),$$

where P_{HC} records the HC half-boundary readout, P_{γ} records photon-compatible null-side readout, P_{sing} records HC-normalized singularity projection, P_{Cas} records Casimir-type boundary readout, P_{OP} records OP-facing residual packages, P_{auto} records finite router automaton and bitstream packages, and P_{ws} records worksheet-style physical readout tests.

Definition H2.88 (Physical promotion criterion)**.** *A router-level object may be physically promoted only if it supplies:*

1. *a declared physical or mathematical readout target;*
2. *a classical collar or explicit boundary interface;*
3. *a comparison, residual, observable, or projection map;*
4. *a back-translation into the prior source language;*
5. *an explicit non-overclaim statement describing which OPs remain open.*

Proposition H2.89 (Machine/content separation)**.** *In the architecture, QAM/MML/MMA is the router, language, and compiler layer. It is not itself the physical content. Physical content begins only where a guarded readout package is promoted by Definition H2.88 into the active readout package of Definition H2.87.*

Proof. The router layer stores kinds, masks, routes, collars, traces, and back-translations. None of these entries is by itself a physical observable. The active readout package adds a physical or mathematical target only after the promotion guard has been supplied. Hence the machine layer and the physical layer are distinct, even though the former stabilizes and compresses the latter. □

Theorem H2.90 (active-layer handoff)**.** *The local proof line handoff object defines a well-formed active-layer final over the archival stage release-final baseline: the QAM/MML/MMA material functions as router and theorem language, while the active physical and mathematical readout layer is isolated as the content-bearing surface.*

Proof. Definitions H2.65, H2.71, H2.85, and H2.87 separate router data, mask data, theorem-layer data, and readout data into finite records. Lemma H2.69, Lemma H2.73, and Lemma H2.84 preserve theorematic status. Proposition H2.89 separates infrastructure from physical content. Therefore the source-layer update is well formed as an active-layer final while OP 1/4 and OP 5(B) remain open. □

guard record. This layer does not assert a global shifted QAM Hilbert-shift, does not introduce primitive mixed tensors, does not close OP 1/4 or OP 5(B), does not replace the standard physical calculations of QFT/QED/Lifshitz theory or general relativity, and does not claim Turing-completeness. It records the router, route-mask, HC-boundary, canonical theorem, and active-readout architecture as a controlled release-final.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Remark H2.91 (How the systems view reads the present visual diagnostics)**.** *The same stack view also explains why Appendix C may carry highly visible packet, Julia-boundary, and frame-flicker diagnostics without turning those figures into primary closure axioms. In the systems reading, such figures live on the visible-view layer: they are auditable readout surfaces of one and the same transported channel, not independent kernel identities. This is exactly why the OP 5 packet/breathing/frame-flicker block and the Julia-boundary block belong naturally to the same document architecture as the deferred kernel store. The figures make visible what the current theorem store is tracking, while closure itself still remains controlled at the structural layer beneath the pictures.*

Remark H2.92 (QAM/systems crosswalk for the present document)**.** *The present document uses the systems analogy in a disciplined layer-by-layer way. The following crosswalk fixes the intended reading.*

QAM/document layer	***Systems reading***	***Present mathematical status***
Visible coded tuples, packet diagnostics, Julia views, routed return surfaces	*User-facing VM/view layer*	*Actively visible and auditable, but not by themselves closure-generating*
Exposed payload kernels and later zero-kernel rereadings	*VM-state identity layer*	*Deferred and prospective only; not yet activated on the public tuple surface*
Admissible transport, branch/shell return, routed support organization	*OS-level orchestration layer*	*Active theorem content in the present QAM export*
Kernel normal forms, search keys, snapshots, clones, audits	*ZFS-style theorem-store layer*	*Partly active as organizational discipline, partly deferred as later compression infrastructure*
Pool/group structure of sphere objects and support families	*Device / vdev / pool layer*	*Explanatory architecture only at the present stage*

Remark H2.93 (Visual-evidence discipline under the systems reading)**.** *Under the crosswalk above, a figure may support only a controlled class of claims. It may expose* where *a visible readout departs from a rigid carrier picture,* which *packet or channel decomposition is plausible on the view layer, and* why *a later kernel-first compression path is worth isolating. It may* not *by itself create a new closure axiom, replace an active transport/branch/shell/routed theorem, or identify a kernel class solely from appearance. Any later proof use must therefore pass through the active theorem layer or through a deferred bridge that has itself already been discharged.*

archival stage checkpoint (deferred kernel-first path). The deferred kernel-first material added in the present local archival stage line should be read as one controlled checkpoint rather than as a second public language block. Its current role is to prepare a later proof-compression path while leaving the active coded tuple predicates unchanged.

Remark H2.94 (Checkpoint summary for the deferred kernel-first path in the archival stage public release)**.** *At the present stage, the deferred kernel-first path contributes six concrete pieces of structure.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

1. *It introduces a prospective recursive zero-kernel vocabulary in which the visible state surface $\langle 0, a\rangle$ may later be re-read through exposed payload kernels, kernel normal forms, and class-level readout factorization.*

2. *It records a normal-form and search-key discipline so that later graph/tree based proof search can be tied to canonical kernel representatives rather than to repeated visible tuple surfaces.*

3. *It isolates the missing bridge datum from active state surfaces to canonical kernels and identifies the direct visible payload as the simplest current kernel final on the present tuple surface.*

4. *It tests the resulting kernel-first viewpoint on the three most relevant compression stages of the current QAM line: admissible transport, local branch-shell return, and routed family compression.*

5. *It formulates an explicit activation package and a staged migration roadmap, making clear both what would have to be proved before activation and in which order the relevant closure/readout factorization work should be completed.*

6. *It remains strictly deferred: no active public predicate, closure theorem, transport theorem, branch-shell theorem, routed-support theorem, or translation theorem is replaced by the kernel language at the present stage.*

Remark H2.95 (Why this checkpoint matters for later proof compression)**.** *The checkpoint above already shows where later proof compression would come from. Transport is the first chain-level reduction, branch-shell return the first local surface reduction, routed support the first family reduction, and the translation theorem then becomes the first clause-level reduction at theorem scale. This does not yet shorten the active proofs automatically, but it isolates a concrete path by which set-theoretic comparison load could later be shifted from repeated visible tuple surfaces to kernel normal forms and one final class-level readout step.*

Remark H2.96 (Navigation guide for the deferred kernel-first checkpoint)**.** *A reader who wants the shortest route through the deferred material may follow the following order.*

1. ***Purpose and fit.*** *Start with the roadmap-fit remark, the checkpoint summary, and the companion remark on why the checkpoint matters in order to see why the deferred path belongs to the inherited public-shape archival stage release layer without changing the public branch/shell/routed extraction.*

2. ***Vocabulary.*** *Then read the terminological convention for the deferred kernel path together with Definitions H2.101–H2.108 for the later kernel vocabulary, primitive zero-objects, recursive reuse, aspect projections, normal forms, and internal search keys.*

3. ***Compatibility.*** *Next read the axiom-coherence audit together with the current-limitation remark to see why this vocabulary remains deferred and does not compete with the active tuple layer.*

4. ***Bridge.*** *Then read Definition H2.114, Theorem H2.115, Definition H2.117, and Theorem H2.119 for the exact interface between visible state surfaces and later kernel-first comparison.*

5. ***Stress tests.*** *Only afterwards read Propositions H2.140, H2.182, and H2.190, which test the bridge on transport, local return, and routed family compression.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

6. ***Activation and use.*** *Finally read Definition H2.196, Theorem H2.197, the later activation-roadmap and reading-guide remarks, and then the dependency-map, minimal-proof-obligation, local-admissibility, and rewrite-schema remarks for the activation threshold, staged migration order, dependency logic, and later local rewrite test.*

In that order, the deferred block reads as one controlled proof-compression path rather than as a loose collection of auxiliary notes.

Remark H2.97 (Execution rule for the deferred kernel-first path)**.** *The deferred block should be used in only one direction when later proof compression is attempted on the present tuple surface. Start with the visible state surface, expose the payload kernel, pass to kernel normal form, prove closure or readout agreement at class level, and lift back to the visible surface only at the end. If the later local admissibility test fails at any point, then the argument should remain on the active tuple surface rather than forcing a premature kernel-first rewrite.*

Remark H2.98 (Present-use restriction for the deferred kernel-first path)**.** *Until the activation criterion of Theorem H2.197 has been fully discharged on the current QAM surface, the deferred block should be used in only three ways inside the document: as later migration vocabulary, as an audit of where proof compression could occur, and as a record of staged stress tests. It should not be cited as a replacement for any active tuple predicate, transport theorem, branch-shell theorem, routed-support theorem, or translation theorem. In particular, whenever both a visible-surface proof and a deferred kernel-first restatement are available, the visible-surface proof remains authoritative on the inherited public-shape archival stage layer.*

Remark H2.99 (Active/deferred correspondence rule)**.** *The deferred kernel-first path may also be read as a controlled shadow of four active theorem layers, with one fixed discipline: the active visible-surface theorem remains authoritative, while the deferred statement records only the later compression route attached to that theorem. Concretely, Theorem H2.139 is paired with Proposition H2.140 as the chain-level compression test; Theorem H2.157 is paired with Proposition H2.182 as the local return-surface compression test; Theorem H2.189 is paired with Proposition H2.190 as the family-level compression test; and Theorem H2.238 is paired with the later local admissibility reading of the translation theorem as the clause-level compression test. This correspondence is only a reading rule for later migration work. It does not transfer proof authority from the active theorem to its deferred companion; it only says where a later kernel-first rewrite would have to pay off if the activation package were eventually discharged.*

Prospective recursive zero-kernel scheme (deferred architecture note)

The following block records a possible later re-reading of the current $\langle 0, a\rangle$-entry as a recursive order-0 readout kernel. It is included here only as a controlled design note for future transfer work. It is *not* activated as part of the present public QAM tag system and does not modify the inherited public-shape coded predicates from historical release marker archival stage.

Remark H2.100 (Terminological convention for the deferred kernel path)**.** *Throughout the deferred block, four layers should be kept distinct and named consistently. The expression* visible state surface *refers to the active public state-code presentation $s = \langle 0, a\rangle$ on the current tuple level. The expression* exposed payload kernel *refers to the payload extracted from such a visible state surface by the deferred kernel constructor. The expression* kernel normal form *refers to the canonical representative selected from the admissible kernel equivalence class. Finally,* class-level readout factorization *means that closure-relevant readout depends only on that kernel normal-form class and is lifted back to the visible state surface only at the end. This convention is purely terminological, but it keeps the active tuple language and the deferred kernel language aligned.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition H2.101 (Carrier/filter readout kernels)**.** *Let $\mathcal{C}$ be a class of admissible carriers, let Γ be a class of admissible filters, and let $\mathcal{A}$ be a class of canonical readout kernels. For each $\gamma \in \Gamma$, assume a partial readout map*

$$\rho_\gamma : \mathcal{C} \dashrightarrow \mathcal{A},$$

whose defined values are written

$$a = \rho_\gamma(T) \in \mathcal{A}.$$

We read a as the minimal canonical readout kernel of the carrier T relative to the filter γ.

Definition H2.102 (Primitive zero-objects)**.** *Given $a, b \in \mathcal{A}$, define the primitive zero-objects*

$$Z_0(a) := \langle 0, a\rangle, \qquad Z_0(a,b) := \langle 0, a, b\rangle.$$

The intended reading is that $Z_0(a)$ is the unary collapsed readout attached to the kernel a, while $Z_0(a,b)$ is the binary collapsed readout attached to the kernel pair (a,b).

Axiom 2 (Recursive reusability of kernels)**.** *Every canonical readout kernel may again serve as an effective carrier for further admissible readout. Concretely, for admissible $\eta \in \Gamma$, one may again form*

$$\rho_\eta(a) \in \mathcal{A}$$

whenever the corresponding readout is defined. Thus the kernel class $\mathcal{A}$ is recursively reusable as an effective carrier layer.

Definition H2.103 (Aspect projections on kernels)**.** *Assume that each kernel $a \in \mathcal{A}$ comes equipped with aspect projections*

$$\pi_{\mathrm{type}}(a), \qquad \pi_{\mathrm{class}}(a), \qquad \pi_{\mathrm{sig}}(a),$$

recording, respectively, its visible type aspect, filter/class aspect, and structured signature aspect. These are not treated as three competing objects; they are three admissible readings of the same kernel.

Lemma H2.104 (Unary and binary kernel collapse)**.** *If $a = \rho_\gamma(T) \in \mathcal{A}$, then the unary zero-object $Z_0(a) = \langle 0, a\rangle$ is well-defined. If additionally $b = \rho_\eta(U) \in \mathcal{A}$, then the binary zero-object $Z_0(a,b) = \langle 0, a, b\rangle$ is well-defined.*

Proof. The unary claim follows directly from Definition H2.102 once $a \in \mathcal{A}$ is given by Definition H2.101. The binary claim is identical after adjoining a second admissible kernel $b \in \mathcal{A}$. □

Remark H2.105 (Compressed relation to the current $\langle 0, a\rangle$ state code)**.** *This deferred kernel notation is intended only as a future re-reading of the present visible state surface. In a later binary-tag or order-sensitive QAM line, one may reinterpret the current $\langle 0, a\rangle$ notation as a compressed public surface for a filtered order-0 readout kernel. For the present version, however, the active public predicates remain exactly those of the current tagged tuple language.*

Remark H2.106 (Tree/graph viewpoint for recursive kernels)**.** *The recursive kernel picture admits an immediate combinatorial visualization. One may regard repeated kernel readout*

$$T \xrightarrow{\rho_\gamma} a \xrightarrow{\rho_\eta} a' \xrightarrow{\rho_\theta} a''$$

as a rooted readout tree whose vertices are kernels and whose edges are admissible filter applications. Equally, one may organize the same data as a directed multigraph: unary zero-objects $Z_0(a)$ mark collapsed kernel vertices, while binary zero-objects $Z_0(a,b)$ record elementary visible relations between kernel vertices. This suggests a later graph-based search or proof-compression interface without changing the current public tuple system.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Prospective kernel normal forms and search keys (deferred architecture note)

The same deferred kernel language also suggests a later search discipline in which recursive kernel proofs are first normalized and only then indexed. This block is likewise prospective only: it does not alter the active inherited public-shape tuple predicates.

Definition H2.107 (Kernel equivalence and canonical normal form)**.** *Assume an admissible equivalence relation $\sim$ on $\mathcal{A}$ such that equivalent kernels have the same admissible aspect projections and the same closure-relevant readout content. A* kernel normal form *is a partial map*

$$\mathrm{nf} : \mathcal{A} \dashrightarrow \mathcal{A}$$

selecting, whenever defined, one canonical representative from each admissible equivalence class. Thus

$$a \sim b \Longrightarrow \mathrm{nf}(a) = \mathrm{nf}(b)$$

whenever both normal forms are defined.

Definition H2.108 (Kernel search keys)**.** *Assume a fixed admissible serialization*

$$\mathrm{ser} : \mathrm{nf}(\mathcal{A}) \to \Sigma^*$$

of canonical kernels into finite strings over an alphabet Σ, together with a deterministic key map

$$\kappa := H \circ \mathrm{ser} \circ \mathrm{nf}\,.$$

The value $\kappa(a)$ is called the kernel search key *of a whenever the normal form is defined. It is intended as an internal search/index key rather than as part of the visible mathematical surface.*

Lemma H2.109 (Normal-form invariance of primitive zero-objects)**.** *Assume that $a, a', b, b' \in \mathcal{A}$ satisfy $a \sim a'$ and $b \sim b'$, and that their normal forms are defined. Then the unary zero-objects $Z_0(a)$ and $Z_0(a')$ determine the same canonical unary kernel surface, while the binary zero-objects $Z_0(a, b)$ and $Z_0(a', b')$ determine the same canonical binary kernel surface after passage to kernel normal form.*

Proof. By Definition H2.107, equivalence implies equality of canonical normal forms. Applying the same normal-form representative at the unary or binary zero-object level yields the same canonical collapsed surface in either case. □

Remark H2.110 (Graph search and proof-compression interface)**.** *In the graph viewpoint introduced above, one may therefore index vertices by canonical kernel normal forms or, internally, by their kernel search keys $\kappa(a)$. Repeated readout branches that land in the same canonical kernel class can then be merged, memoized, or searched only once. This is exactly the kind of mechanism by which recursive kernel language could later compress set-theoretic proof search without changing the visible public theorem statements.*

Axiom-coherence audit and prospective proof reductions (deferred audit note)

The deferred zero-kernel language should be read together with the active public QAM tuple layer under the following discipline.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Remark H2.111 (Axiom-coherence audit)**.** *At the present stage, the deferred zero-kernel layer is formally compatible with the active QAM tuple layer for four reasons.*

1. *The active tuple predicates remain unchanged: in particular,*

$$\mathrm{QState}(x) \Longleftrightarrow \exists a\,(x = \langle 0, a\rangle)$$

   continues to define the public visible state surface.

2. *The deferred zero-kernel definitions do not introduce a second active state predicate. They only record a possible later re-reading of the same public surface, as stated in Remark H2.105.*

3. *The recursive kernel axiom (Axiom 2) and the kernel normal-form discipline (Definition H2.107) are prospective only; no active theorem in the inherited public-shape QAM chain depends on them as hypotheses.*

4. *Consequently, the current closure-class, admissible-transport, branch/shell, routed-support, and translation theorems remain exactly those of the active tuple language. The deferred kernel layer adds a controlled future interpretation, not a rival public axiom system.*

Remark H2.112 (Where kernel notation could already shorten proofs)**.** *Even before formal activation, the deferred kernel language already points to three concrete proof-compression opportunities.*

1. *In the first QAM transport theorem, the chain*

$$s_{\mathrm{in}},\ s_2,\ s_3,\ s_4,\ t_{\mathrm{fwd}},\ t_{\mathrm{bwd}}$$

   is currently carried at full state-code level. A later kernel reduction could first pass each state to a canonical readout kernel and then prove closure agreement on kernel classes before lifting the result back to coded states.

2. *In the branch-shell theorem and the routed admissible support theorem, each return realization currently enters through full branch/shell state data. A kernel-first restatement could instead reduce every admissible branch-shell realization to unary kernel surfaces $Z_0(a)$ and compare routed sectors only through their canonical kernel classes.*

3. *In closure-compatible readout arguments, one currently factors raw readouts through the class-target map J after proving* $\mathrm{QClEq}(s,t)$*. A later kernel language could absorb much of this step by treating the closure-relevant image itself as the primitive kernel surface attached to the state.*

In all three cases, the expected gain is not a new theorem but a shorter proof path: reduce to kernels first, perform the closure comparison there, and only then return to the visible tuple layer.

Remark H2.113 (Current limitation of the kernel reduction)**.** *The proof-compression opportunities recorded just above are not yet activated as a formal replacement inside the present proofs. The missing step is a theorem-level bridge showing that every active coded state relevant to the admissible OP fragment carries a canonical kernel representative with the same closure-relevant content. Until such a bridge is proved, the zero-kernel language should be treated as a planning interface rather than as a proof rule.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition H2.114 (Prospective kernel bridge datum on active state surfaces)**.** *A* kernel bridge datum *for the active admissible OP fragment consists of a partial map*

$$\ker : \mathcal{S}_{\mathrm{act}}^{\mathrm{QAM}} \dashrightarrow \mathcal{A},$$

where $\mathcal{S}_{\mathrm{act}}^{\mathrm{QAM}}$ *is the class of active coded QAM-state surfaces relevant to the present admissible OP fragment, such that the following hold whenever the displayed expressions are defined.*

1. ***surface realization:***
$$Z_0(\ker(s)) = s;$$
2. ***closure reflection:*** *if* $\mathrm{QClEq}(s,t)$*, then*
$$\mathrm{nf}(\ker(s)) = \mathrm{nf}(\ker(t));$$
3. ***closure-readout factorization:*** *for every closure-compatible coded readout* r*, there exists a map*
$$\mathcal{R}_r : \mathrm{nf}(\mathcal{A}) \to \mathrm{Im}(J_r)$$
into the closure-relevant target of the associated class-factor map J_r *such that whenever* $\mathrm{QRead}(r,s,w)$ *holds,*
$$J_r(w) = \mathcal{R}_r(\mathrm{nf}(\ker(s))).$$

Theorem H2.115 (Prospective bridge from active state surfaces to canonical kernels)**.** *Assume a kernel bridge datum in the sense of Definition H2.114. Then every active closure comparison in the admissible OP fragment may be reduced to canonical kernel comparison before returning to the visible state surface. In particular:*

1. *if* $\mathrm{QAdmTransport}(o,s,t)$*, then*
$$\mathrm{nf}(\ker(s)) = \mathrm{nf}(\ker(t));$$
2. *if* $t_{b,\sigma}$ *is a branch-shell coded return realization as in Theorem H2.157, then*
$$\mathrm{nf}(\ker(s_{\mathrm{in}})) = \mathrm{nf}(\ker(t_{b,\sigma}));$$
3. *if* u *is a normalized admissible routed support code as in Theorem H2.189, then for every closure-compatible coded readout* r *the closure-relevant image of the routed return is determined solely by the canonical kernel class*
$$\mathrm{nf}(\ker(s_{\mathrm{in}})).$$

Consequently, the present transport, branch/shell, and routed-support proofs admit a future kernel-first proof skeleton: first pass to $\mathrm{nf}(\ker(\cdot))$*, then compare closure there, and only afterwards return to the visible QAM state surface.*

Proof. For (1), admissible transport implies $\mathrm{QClEq}(s,t)$ by the active closure-preservation axiom. The closure-reflection clause of Definition H2.114 therefore yields
$$\mathrm{nf}(\ker(s)) = \mathrm{nf}(\ker(t)).$$
For (2), Theorem H2.157 gives $\mathrm{QClEq}(s_{\mathrm{in}}, t_{b,\sigma})$, so the same closure-reflection clause yields equality of canonical kernel representatives. For (3), Theorem H2.189 states that the routed return has the same closure-relevant image as the coded source state under every closure-compatible coded readout. By clause (3) of Definition H2.114, those closure-relevant images factor through the same canonical kernel normal form. Hence the routed readout depends only on $\mathrm{nf}(\ker(s_{\mathrm{in}}))$. The final sentence is simply the proof reading order extracted from (1)–(3). □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Remark H2.116 (What the bridge theorem still leaves open)**.** *Theorem H2.115 is intentionally conditional. It does not yet prove that a kernel bridge datum exists inside the active public tuple system; it isolates exactly the missing interface that would be needed to turn the deferred zero-kernel language into an actual proof tool. This also identifies the next genuine activation step: construct* ker *canonically from active coded states and verify clauses (1)–(3) without leaving the current OP/QAM semantics.*

Definition H2.117 (Visible payload extraction on active state surfaces)**.** *For every active coded state surface s in the present public tuple language, define its* exposed payload kernel $\mathrm{vp}(s)$ *by*

$$\mathrm{vp}(s) = a \quad \Longleftrightarrow \quad s = \langle 0, a\rangle.$$

Equivalently, vp *is the unique projection that removes the visible state tag* 0 *and exposes the payload already carried by the active state surface.*

Definition H2.118 (Prospective direct kernel constructor)**.** *Assume that the exposed payload kernel of every active coded state surface relevant to the admissible OP fragment belongs to the deferred kernel class $\mathcal{A}$. Then define the* direct kernel constructor

$$\ker_{\mathrm{dir}} : \mathcal{S}_{\mathrm{act}}^{\mathrm{QAM}} \dashrightarrow \mathcal{A}, \qquad \ker_{\mathrm{dir}}(s) := \mathrm{vp}(s).$$

Thus, on the visible state surface, the simplest possible kernel final is obtained by forgetting only the active state tag and keeping the payload itself as the deferred kernel representative.

Theorem H2.119 (Simplest concrete kernel construction on the present state surface)**.** *Assume the following four conditions for the active admissible OP fragment.*

1. ***payload admissibility:*** *every active coded state surface s relevant to the fragment has the form $s = \langle 0, a\rangle$ with $a \in \mathcal{A}$;*

2. ***visible zero-surface realization:*** *for every such a,*

$$Z_0(a) = \langle 0, a\rangle;$$

3. ***closure reflection on exposed payload kernels:*** *whenever* $\mathrm{QClEq}(\langle 0, a\rangle, \langle 0, b\rangle)$,

$$\mathrm{nf}(a) = \mathrm{nf}(b);$$

4. ***class-level readout factorization:*** *for every closure-compatible coded readout r, there exists a map*

$$\widetilde{\mathcal{R}}_r : \mathrm{nf}(\mathcal{A}) \to \mathrm{Im}(J_r)$$

such that whenever $\mathrm{QRead}(r, \langle 0, a\rangle, w)$,

$$J_r(w) = \widetilde{\mathcal{R}}_r(\mathrm{nf}(a)).$$

Then $\ker_{\mathrm{dir}}$ is a kernel bridge datum in the sense of Definition H2.114. Consequently Theorem H2.115 applies with

$$\ker = \ker_{\mathrm{dir}}.$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. Clause (1) ensures that $\ker_{\mathrm{dir}}$ is defined on every active state surface relevant to the fragment. By Definition H2.117, if $s = \langle 0, a\rangle$, then $\ker_{\mathrm{dir}}(s) = a$. Clause (2) therefore gives

$$Z_0(\ker_{\mathrm{dir}}(s)) = Z_0(a) = \langle 0, a\rangle = s,$$

which is exactly the surface-realization clause of Definition H2.114. For closure reflection, if $\mathrm{QClEq}(s, t)$ with $s = \langle 0, a\rangle$ and $t = \langle 0, b\rangle$, clause (3) gives

$$\mathrm{nf}(\ker_{\mathrm{dir}}(s)) = \mathrm{nf}(a) = \mathrm{nf}(b) = \mathrm{nf}(\ker_{\mathrm{dir}}(t)).$$

Finally, clause (4) is exactly the required closure-readout factorization after substituting $\ker_{\mathrm{dir}}(\langle 0, a\rangle) = a$. Hence all three clauses of Definition H2.114 hold. □

Remark H2.120 (What this construction shows)**.** *Theorem H2.119 shows that the difficult part of the bridge is not the extraction step itself: on the current public state surface, the most economical final is simply the exposed payload of the active state surface. The real work is pushed into the three semantic checks: that the payload already lies in the admissible kernel class, that closure equivalence is visible at payload-normal-form level, and that closure-relevant readout factors through that same normal form. If these three checks can later be discharged, then many current state-level proofs may be shortened by passing immediately from $\langle 0, a\rangle$ to a before performing the closure comparison.*

QAM minimal predicates and coded relations

Status. *Active formal-language layer.*

Block function. This block introduces the smallest coded predicate/relation vocabulary needed to read the present admissible OP fragment in QAM-over-ZF form.

We now pass from the entry-layer preview to the first explicit predicate layer of QAM-over-ZF. At this stage, the QAM language still remains conservative over ZF: the predicates below are intended as definable recognition or relation clauses on tagged set-codes rather than as commitments to a new ontology.

Definition H2.121 (Primitive QAM predicates)**.** *We introduce the primitive predicates*

$$\mathrm{QState}(x), \quad \mathrm{QLayer}(x), \quad \mathrm{QOp}(x), \quad \mathrm{QReadout}(x),$$
$$\mathrm{QBranch}(x), \quad \mathrm{QShell}(x), \quad \mathrm{QWitness}(x), \quad \mathrm{QTensor}(x).$$

with the intended ZF-grounded readings

$$\mathrm{QState}(x) \ :\Longleftrightarrow\ \exists a\,(x = \langle 0, a\rangle),$$

$$\mathrm{QLayer}(x) \ :\Longleftrightarrow\ \exists \ell\,\exists D\,(x = \langle 1, \ell, D\rangle),$$
$$\mathrm{QOp}(x) \ :\Longleftrightarrow\ \exists \ell_1\,\exists \ell_2\,\exists G\,(x = \langle 2, \ell_1, \ell_2, G\rangle),$$
$$\mathrm{QReadout}(x) \ :\Longleftrightarrow\ \exists \ell\,\exists W\,\exists R\,(x = \langle 3, \ell, W, R\rangle),$$
$$\mathrm{QBranch}(x) \ :\Longleftrightarrow\ \exists b\,\exists B\,(x = \langle 4, b, B\rangle),$$
$$\mathrm{QShell}(x) \ :\Longleftrightarrow\ \exists r\,\exists s\,\exists T\,(x = \langle 5, r, s, T\rangle),$$
$$\mathrm{QWitness}(x) \ :\Longleftrightarrow\ \exists u\,(x = \langle 6, u\rangle),$$
$$\mathrm{QTensor}(x) \ :\Longleftrightarrow\ \exists p\,\exists q\,(x = \langle 7, p, q\rangle).$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Remark H2.122. *At this stage, these predicates are still recognition clauses rather than a full independent axiom system. Their purpose is to let the Companion speak explicitly about coded states, layers, operators, readouts, branches, shell/refinement objects, witnesses, and tensor composites inside one ZF-grounded language layer.*

Definition H2.123 (Primitive QAM coded relations)**.** *We also introduce the primitive QAM-style relations*

$$\mathrm{QAdmTransport}(o,s,t), \qquad \mathrm{QClEq}(s,t), \qquad \mathrm{QRead}(r,s,w),$$

$$\mathrm{QBranchRef}(b,s,t), \qquad \mathrm{QShellRef}(\sigma,s,t), \qquad \mathrm{QTensorAdm}(x,y).$$

Their intended readings are as follows:

1. $\mathrm{QAdmTransport}(o,s,t)$ *means that the operator-code o transports the coded state s admissibly to the coded state t;*
2. $\mathrm{QClEq}(s,t)$ *means that s and t belong to the same coded closure class;*
3. $\mathrm{QRead}(r,s,w)$ *means that the readout-code r reads the coded state s into the coded observable value w;*
4. $\mathrm{QBranchRef}(b,s,t)$ *means that the branch-code b refines the coded state s to the coded state t;*
5. $\mathrm{QShellRef}(\sigma,s,t)$ *means that the shell/refinement-code σ refines the coded state s to the coded state t;*
6. $\mathrm{QTensorAdm}(x,y)$ *means that the tensor-coded composite $x \otimes y$ is admitted as a legitimate coded QAM object.*

Remark H2.124. *These relations are the first minimal bridge between the current OP/branch/shell architecture and its coded QAM-over-ZF form. In particular, admissibility and closure-class preservation already appear here as primitive language-level notions rather than as only informal motivation.*

Definition H2.125 (First QAM translation target)**.** *The first intended translation target of the QAM-over-ZF layer is the admissible OP fragment consisting of:*

1. *the OP 1/OP 5 source-return closure architecture;*
2. *the intermediate OP 2–OP 4 transport bridge;*
3. *admissible branch transport;*
4. *OP 3 shell-admissible refinement;*
5. *admissible routed support together with the visible packet/phase-lock/epsilon screening layer.*

Remark H2.126. *The significance of the present predicate layer is that QAM is no longer only a roadmap note. Even at this first stage, the language already distinguishes object-kinds and relation-kinds closely enough to serve as a faithful coded entry interface for the current admissible OP fragment.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

QAM transport, closure, and readout schemata

Status. *Active theorem-core block inside the QAM formal layer.*

Block function. This block turns the predicate language into actual transport, closure-class, and readout principles, so it is the first place where the QAM interface becomes theorem-usable.

We now pass from the minimal predicate layer to the first theorematic QAM transport/closure layer. The point of this section is still conservative: it does not yet claim a fully independent QAM foundation, but it does begin to formulate admissible transport, closure-class preservation, and closure-compatible readout directly in the coded QAM-over-ZF language.

Definition H2.127 (QAM-coded transport graph)**.** *A QAM-operator code*

$$o = \langle 2, \ell_1, \ell_2, G \rangle$$

is called a coded transport graph *if G is a set of ordered pairs coding a functional transport rule from coded states on layer ℓ_1 to coded states on layer ℓ_2.*

Definition H2.128 (QAM admissible transport)**.** *For a QAM-operator code o and coded states s, t, we write*

$$\mathrm{QAdmTransport}(o, s, t)$$

iff:

1. *o is a coded transport graph;*
2. *s is a coded state on the source layer of o;*
3. *t is a coded state on the target layer of o;*
4. *the graph-code of o sends s to t;*
5. *the transport from s to t preserves the intended closure-bearing structural mode.*

Remark H2.129. *At this stage, the final clause is still schematic. Its precise intended mathematical content is supplied by the coded closure-class relation below.*

Definition H2.130 (QAM closure-class equivalence)**.** *The relation*

$$\mathrm{QClEq}(s, t)$$

is the coded closure-class relation on QAM-state codes. It is intended to mean that s and t represent the same closure-bearing structural mode, even if they differ by admissible transport, admissible frame-shifted realization, admissible branch refinement, or admissible shell refinement.

Axiom 3 (QAM closure-class schema)**.** *The relation* QClEq *is required to satisfy:*

1. ***reflexivity:***
$$\mathrm{QClEq}(s, s);$$
2. ***symmetry:***
$$\mathrm{QClEq}(s, t) \Longrightarrow \mathrm{QClEq}(t, s);$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

3. ***transitivity:***

$$\mathrm{QClEq}(s,t) \wedge \mathrm{QClEq}(t,u) \Longrightarrow \mathrm{QClEq}(s,u).$$

Remark H2.131. *Thus* QClEq *is the coded QAM analogue of the native closure carrier-class relation used in the current OP architecture.*

Axiom 4 (QAM admissibility implies closure preservation)**.** *For every operator code o and state codes s,t,*

$$\mathrm{QAdmTransport}(o,s,t) \Longrightarrow \mathrm{QClEq}(s,t).$$

Remark H2.132. *This is the first central QAM principle. Admissibility is not defined merely by existence of transport, but by preservation of closure class.*

Definition H2.133 (QAM-coded readout)**.** *A QAM-readout code*

$$r = \langle 3, \ell, W, R \rangle$$

is called a coded readout *if R is a graph-code sending coded states on layer ℓ into coded observable values in the target code W.*

Definition H2.134 (QAM readout relation)**.** *We write*

$$\mathrm{QRead}(r,s,w)$$

iff r is a coded readout, s is a coded state on the source layer of r, and the graph-code of r sends s to the coded observable value w.

Definition H2.135 (QAM closure-compatible readout)**.** *A coded readout r is called* closure-compatible *if there exists a coded class-target map J such that for all coded states s,t,*

$$\mathrm{QClEq}(s,t) \Longrightarrow \big(\forall w_1\, \forall w_2\, (\mathrm{QRead}(r,s,w_1) \wedge \mathrm{QRead}(r,t,w_2) \Longrightarrow J(w_1) = J(w_2))\big).$$

Remark H2.136. *This is the coded QAM version of readout factorization through closure class. Raw readouts may differ, but their closure-relevant image under J must agree whenever the source states belong to the same closure class.*

Definition H2.137 (First tensor-compatibility placeholder)**.** *We reserve the tensor-admissibility predicate*

$$\mathrm{QTensorAdm}(x,y)$$

for the statement that the tensor-coded composite $x \otimes y$ is admitted as a legitimate coded QAM object of the intended kind.

Proposition H2.138 (First QAM transport principle)**.** *Let o be a coded QAM-operator, let s,t be coded QAM-states, and let r be a closure-compatible coded readout. If*

$$\mathrm{QAdmTransport}(o,s,t),$$

then:

1. $$\mathrm{QClEq}(s,t);$$

2. *the closure-relevant image of any readout of s agrees with the closure-relevant image of any readout of t under the factor map J associated with r.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. The first claim is exactly the admissibility-implies-closure-preservation schema. The second follows from closure-compatibility of the readout: since $\mathrm{QClEq}(s,t)$ holds, any coded readouts of s and t are identified after passage through the closure-relevant class-target map J. □

Theorem H2.139 (First QAM version of the OP transport principle)**.** *Assume:*

1. *there is a coded source state s_{in};*
2. *there are coded intermediate states s_2, s_3, s_4;*
3. *there are coded operator maps $o_2, o_3, o_4, o_{\mathrm{fwd}}, o_{\mathrm{bwd}}$;*
4. *the transports*

$$\begin{aligned}&\mathrm{QAdmTransport}(o_2, s_{\mathrm{in}}, s_2), \qquad \mathrm{QAdmTransport}(o_3, s_2, s_3),\\ &\mathrm{QAdmTransport}(o_4, s_3, s_4), \qquad \mathrm{QAdmTransport}(o_{\mathrm{fwd}}, s_{\mathrm{in}}, t_{\mathrm{fwd}}),\\ &\mathrm{QAdmTransport}(o_{\mathrm{bwd}}, s_{\mathrm{in}}, t_{\mathrm{bwd}}).\end{aligned}$$

   all hold;

5. *the relevant coded readout is closure-compatible.*

Then all coded states

$$s_{\mathrm{in}},\ s_2,\ s_3,\ s_4,\ t_{\mathrm{fwd}},\ t_{\mathrm{bwd}}$$

belong to the same coded closure class. Consequently, all closure-relevant readouts of these states agree after passage through the closure-factor map.

Proof. Each admissible transport preserves closure class by the admissibility schema. Therefore

$$\mathrm{QClEq}(s_{\mathrm{in}}, s_2), \qquad \mathrm{QClEq}(s_2, s_3), \qquad \mathrm{QClEq}(s_3, s_4).$$

By transitivity,

$$\mathrm{QClEq}(s_{\mathrm{in}}, s_4).$$

Likewise,

$$\mathrm{QClEq}(s_{\mathrm{in}}, t_{\mathrm{fwd}}) \qquad \text{and} \qquad \mathrm{QClEq}(s_{\mathrm{in}}, t_{\mathrm{bwd}})$$

follow from admissible forward and backward transport. Using transitivity again, all listed coded states lie in the same closure class. Closure-compatible readout then identifies their closure-relevant readout content after factorization through the associated class-target map. □

Proposition H2.140 (Prospective kernel-first reduction of the first transport theorem)**.** *Assume the direct kernel constructor of Definition H2.118 and the bridge conditions of Theorem H2.119. Write*

$$s_{\mathrm{in}} = \langle 0, a_{\mathrm{in}}\rangle, \qquad s_2 = \langle 0, a_2\rangle, \qquad s_3 = \langle 0, a_3\rangle, \qquad s_4 = \langle 0, a_4\rangle,$$

$$t_{\mathrm{fwd}} = \langle 0, a_{\mathrm{fwd}}\rangle, \qquad t_{\mathrm{bwd}} = \langle 0, a_{\mathrm{bwd}}\rangle.$$

Then the conclusion of Theorem H2.139 may be read kernel-first as follows:

1. *all exposed transport kernels lie in one canonical normal-form class,*

$$\mathrm{nf}(a_{\mathrm{in}}) = \mathrm{nf}(a_2) = \mathrm{nf}(a_3) = \mathrm{nf}(a_4) = \mathrm{nf}(a_{\mathrm{fwd}}) = \mathrm{nf}(a_{\mathrm{bwd}}),$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

2. *hence every closure-compatible coded readout of the transport chain factors through that single canonical kernel class,*

3. *therefore the first transport theorem may be re-read as a kernel-normal-form preservation statement along the admissible transport chain before returning to the visible QAM state surface.*

Proof. Theorem H2.139 states that all six visible coded states belong to the same coded closure class. Since each state lies on the present state surface, the direct kernel constructor exposes exactly the exposed payload kernel of each state as its deferred kernel representative. The closure-reflection clause of Theorem H2.119 therefore yields equality of the six corresponding kernel normal forms. By the class-level readout factorization clause, every closure-compatible coded readout depends only on that common kernel normal form. Hence the whole transport chain may be treated kernel-first and lifted back to the visible state surface only at the end. □

Remark H2.141 (Axiom audit after the kernel-first transport reduction)**.** *Proposition H2.140 does* not *strengthen the active proof status of Theorem H2.139. The active theorem still rests only on admissible transport, closure-class preservation, transitivity of coded closure, and closure-compatible coded readout. The kernel-first proposition adds force only under the deferred payload-level bridge hypotheses isolated in Definition H2.114 and Theorem H2.119. Thus no new hidden axiom has entered the live transport theorem; only a prospective kernel-level compression of its proof surface has been recorded.*

Remark H2.142 (How much proof load the kernel-first transport reading removes)**.** *For the present transport theorem the admissible chain contains five explicit transport links:*

$$s_{\mathrm{in}} \to s_2 \to s_3 \to s_4, \qquad s_{\mathrm{in}} \to t_{\mathrm{fwd}}, \qquad s_{\mathrm{in}} \to t_{\mathrm{bwd}}.$$

So the gain is again not *a reduction in the number of transport links, but a reduction in the size of the object carried through each link. On the visible state surface the proof transports closure across five state-surface steps and then factors a closure-compatible readout through the resulting six-state closure class. In the deferred kernel-first reading, those same five links are re-read as five kernel normal-form identifications, followed by one readout through the common kernel class. If one measures a visible-state transport comparison by cost C_{surf} and a kernel normal-form comparison by cost C_{ker}, then the conceptual load changes from roughly*

$$5\,C_{\mathrm{surf}} + C_{\mathrm{read}}$$

to

$$5\,C_{\mathrm{ker}} + C_{\mathrm{class}},$$

with intended gain $C_{\mathrm{ker}} < C_{\mathrm{surf}}$ and typically $C_{\mathrm{class}} \ll C_{\mathrm{read}}$. So the numerical savings are moderate but already real: the theorem still has five links, yet each link is checked on exposed kernels instead of full visible state surfaces, and the final readout is centralized on one canonical kernel class.

Remark H2.143. *This theorem is only the first coded QAM shadow of the current OP transport architecture. It does not yet include the full branch-shell routed layer, but it already captures the source/intermediate/return logic in a ZF-grounded coded form.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

QAM branch, shell, and routed support layer

Status. *Active theorem-core branch-shell extension.*

Block function. This block lifts the basic QAM transport layer to the branch/shell/routed regime and should be read as the coded counterpart of the admissible OP refinement chain.

We now extend the QAM-over-ZF language from basic transport and readout to the branch/shell/routed layer. The point of this section is to translate the current admissible-branch, OP 3 shell-admissibility, and admissible routed-support architecture into a single coded QAM fragment without yet claiming a fully independent QAM foundation. For direct continuity with the compact front extraction, the present subsection should be read in the same order as the release dictionary of Remark H2.3: branch admissibility first, shell admissibility second, admissible branch-shell support third, normalized routed support fourth, and only then the visible-versus-structural screening rule.

Definition H2.144 (Closure-admissible branch code)**.** *A branch code b is called* closure-admissible *if for all coded states s, t,*

$$\mathrm{QBranchRef}(b, s, t) \Longrightarrow \mathrm{QClEq}(s, t).$$

Equivalently, branch refinement by b preserves the coded closure class.

Definition H2.145 (Shell-admissible refinement code)**.** *A shell/refinement code σ is called* shell-admissible *if for all coded states s, t,*

$$\mathrm{QShellRef}(\sigma, s, t) \Longrightarrow \mathrm{QClEq}(s, t).$$

Equivalently, shell-sensitive or residual-sensitive refinement by σ preserves the coded closure class.

Remark H2.146. *These are the direct coded QAM analogues of closure-admissible branches and OP 3 shell-admissible refinements in the current Companion line.*

Definition H2.147 (Admissible branch set and shell family)**.** *We define the admissible branch class by*

$$\mathcal{B}_{\mathrm{adm}}^{\mathrm{QAM}} := \{\, b : \mathrm{QBranch}(b) \ \wedge \ b \text{ is closure-admissible} \,\},$$

and for each branch code b, the shell-admissible refinement class by

$$\mathcal{F}_{\mathrm{sh}}^{\mathrm{QAM}}(b) := \{\, \sigma : \mathrm{QShell}(\sigma) \ \wedge \ \sigma \text{ is shell-admissible on } b \,\}.$$

Remark H2.148. *At this stage, the phrase "shell-admissible on b" may be read minimally as: shell refinement by σ acts on branchwise states associated with b without leaving the coded closure class.*

Definition H2.149 (Routed support code)**.** *A* routed support code *is a set-code*

$$u = \langle 8, \Omega, w \rangle,$$

where Ω is a coded finite or set-sized support family of admissible branch-shell pairs and w is a coded weight assignment on Ω.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition H2.150 (Admissible routed support)**.** *A routed support code*

$$u = \langle 8, \Omega, w \rangle$$

is called admissible *if every element of* Ω *has the form* (b, σ) *with*

$$b \in \mathcal{B}_{\mathrm{adm}}^{\mathrm{QAM}} \qquad \text{and} \qquad \sigma \in \mathcal{F}_{\mathrm{sh}}^{\mathrm{QAM}}(b).$$

Remark H2.151. *Thus admissible routed support is the coded QAM version of the admissible branch-shell support sector in the current OP architecture.*

Definition H2.152 (Normalized routed support)**.** *An admissible routed support code*

$$u = \langle 8, \Omega, w \rangle$$

is called normalized *if its coded weight assignment satisfies*

$$\sum_{(b,\sigma)\in\Omega} w(b, \sigma) = 1$$

in the intended decoded numerical target.

Remark H2.153. *The normalization condition is stated semiformally at this stage. In the fully coded version, the sum relation would itself be represented by a suitable arithmetic coding layer over ZF.*

Definition H2.154 (Branch-shell coded return realization)**.** *Let* b *be a closure-admissible branch code and let* σ *be a shell-admissible refinement code on* b*. A* branch-shell coded return realization *is a coded state*

$$t_{b,\sigma}$$

such that:

1. $t_{b,\sigma}$ *is reached from the coded source state through admissible branch refinement by* b*;*
2. *the OP 3-level shell refinement by* σ *is shell-admissible;*
3. *the resulting return-side coded state is read through a closure-compatible coded readout.*

Proposition H2.155 (QAM branch admissibility principle)**.** *Let* b *be a closure-admissible branch code. If*

$$\mathrm{QBranchRef}(b, s, t),$$

then

$$\mathrm{QClEq}(s, t).$$

Consequently, every closure-compatible readout identifies the closure-relevant image of s *and* t *after passage through the associated class-factor map.*

Proof. The first claim is just the definition of closure-admissibility. The readout claim follows from closure-compatibility of coded readouts: closure-equivalent coded states have the same closure-relevant image after factorization. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proposition H2.156 (QAM shell admissibility principle)**.** *Let σ be a shell-admissible refinement code. If*

$$\mathrm{QShellRef}(\sigma, s, t),$$

then

$$\mathrm{QClEq}(s, t).$$

Consequently, shell-sensitive refinement does not alter closure-relevant readout content as long as shell-admissibility is preserved.

Proof. This is the direct content of shell-admissibility together with closure-compatible readout factorization. □

Theorem H2.157 (QAM admissible branch-shell theorem)**.** *Let s_{in} be a coded source state, let b be a closure-admissible branch code, and let σ be a shell-admissible refinement code on b. Assume there exists a branch-shell coded return realization*

$$t_{b,\sigma}$$

associated with (b, σ), and let r be a closure-compatible coded readout. Then

$$\mathrm{QClEq}(s_{\mathrm{in}}, t_{b,\sigma}).$$

Hence every closure-compatible readout of s_{in} and $t_{b,\sigma}$ agrees after passage through its associated closure-factor map.

Proof. By closure-admissibility of b, branch refinement preserves the coded closure class. By shell-admissibility of σ, the shell-sensitive refinement preserves the same coded closure class at the OP 3 layer. Hence the final coded return realization $t_{b,\sigma}$ remains closure-equivalent to the coded source state s_{in}. Closure-compatible readout then identifies the closure-relevant content of both states after factorization. □

Shell certificates and the first OP 3/QAM closure guide

Status. *Active closure-guide layer.*

Block function. This block isolates the certificate vocabulary that lets readers track exactly when OP 3 shell behavior is still closure-preserving and when a local failure first becomes visible.

The next narrow OP 3-oriented extension of the public-shape QAM line is not a broad new tensor layer, but a certificate layer. The goal is to say explicitly when a branch/shell refinement is still the same closure-bearing object, what counts as the first local failure of that claim, and how the deferred kernel bridge should later read that same situation in one compressed normal form. Mixed tensors remain deferred here: they are not introduced unless an actual OP 3 proof step later forces them.

QAM reading key for the OP 3 closure block.
Objects: coded source state s_{in}, closure-admissible branch code b, shell refinement code σ, intermediate branch state s_b, branch-shell return state $t_{b,\sigma}$.
Allowed move: pass from s_{in} to s_b by admissible branch refinement and from s_b to $t_{b,\sigma}$ by shell refinement without leaving the closure class.
Invariant: the closure-bearing class, and later its kernel/normal-form representative when the deferred bridge is activated.
Defect signal: a broken branch clause, a broken shell clause, or a failure of closure-compatible readout factorization.
Plain-language reading: OP 3 may change the visible shell, but it must not change the closure-bearing core.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition H2.158 (Shell admissibility certificate)**.** *Let s_{in} be a coded source state, let b be a closure-admissible branch code, let σ be a shell refinement code, and let $t_{b,\sigma}$ be a branch-shell coded return realization. A* shell admissibility certificate *for $(s_{\mathrm{in}}, b, \sigma, t_{b,\sigma})$ consists of coded data witnessing the following four clauses:*

1. *there exists a coded branch state s_b with* $\mathrm{QBranchRef}(b, s_{\mathrm{in}}, s_b)$*;*
2. *the branch code b is closure-admissible;*
3. *one has* $\mathrm{QShellRef}(\sigma, s_b, t_{b,\sigma})$ *and the refinement code σ is shell-admissible on b;*
4. *the return state $t_{b,\sigma}$ is read through a closure-compatible coded readout.*

Definition H2.159 (Shell defect witness)**.** *A* shell defect witness *for $(s_{\mathrm{in}}, b, \sigma, t_{b,\sigma})$ is coded data showing failure of at least one clause of Definition H2.158. Thus a shell defect witness may record failure of admissible branch passage, failure of shell admissibility, or failure of closure-compatible readout factorization.*

Lemma H2.160 (Certificate persistence of closure class)**.** *Assume that $(s_{\mathrm{in}}, b, \sigma, t_{b,\sigma})$ carries a shell admissibility certificate. Then*

$$\mathrm{QClEq}(s_{\mathrm{in}}, t_{b,\sigma}).$$

Hence every closure-compatible coded readout identifies the same closure-relevant image on s_{in} and $t_{b,\sigma}$ after passage through the associated class-factor map.

Proof. By clause (1) of Definition H2.158, there exists s_b with

$$\mathrm{QBranchRef}(b, s_{\mathrm{in}}, s_b).$$

By clause (2), the branch code b is closure-admissible, so Proposition H2.155 yields

$$\mathrm{QClEq}(s_{\mathrm{in}}, s_b).$$

By clause (3), one has

$$\mathrm{QShellRef}(\sigma, s_b, t_{b,\sigma}),$$

and the refinement code σ is shell-admissible on b. Proposition H2.156 therefore gives

$$\mathrm{QClEq}(s_b, t_{b,\sigma}).$$

Transitivity of the coded closure relation yields

$$\mathrm{QClEq}(s_{\mathrm{in}}, t_{b,\sigma}).$$

The final sentence is then the usual closure-compatible readout factorization claim. □

Proposition H2.161 (Certified kernel persistence for OP 3 shell passage)**.** *Assume the direct kernel constructor of Definition H2.118 and the bridge conditions of Theorem H2.119. Let*

$$s_{\mathrm{in}} = \langle 0, a_{\mathrm{in}} \rangle$$

be the coded source state, and let

$$t_{b,\sigma} = \langle 0, a_{b,\sigma} \rangle$$

be a branch-shell coded return realization. If $(s_{\mathrm{in}}, b, \sigma, t_{b,\sigma})$ carries a shell admissibility certificate, then

$$\mathrm{nf}(a_{b,\sigma}) = \mathrm{nf}(a_{\mathrm{in}}).$$

In particular, certified OP 3 shell passage preserves the kernel normal form of the coded source state.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. By Lemma H2.160, one has

$$\mathrm{QClEq}(s_{\mathrm{in}}, t_{b,\sigma}).$$

The deferred kernel bridge then reflects closure equivalence into equality of canonical kernel normal forms. Since

$$s_{\mathrm{in}} = \langle 0, a_{\mathrm{in}} \rangle, \qquad t_{b,\sigma} = \langle 0, a_{b,\sigma} \rangle,$$

this yields

$$\mathrm{nf}(a_{b,\sigma}) = \mathrm{nf}(a_{\mathrm{in}}).$$

□

Theorem H2.162 (Shell-stable readout factorization)**.** *Assume the direct kernel constructor of Definition H2.118 and the bridge conditions of Theorem H2.119. Let*

$$s_{\mathrm{in}} = \langle 0, a_{\mathrm{in}} \rangle, \qquad t_{b,\sigma} = \langle 0, a_{b,\sigma} \rangle,$$

let r be a closure-compatible coded readout, and assume that $(s_{\mathrm{in}}, b, \sigma, t_{b,\sigma})$ carries a shell admissibility certificate. Then for every pair of coded observable values $w_{\mathrm{in}}, w_{b,\sigma}$ with

$$\mathrm{QRead}(r, s_{\mathrm{in}}, w_{\mathrm{in}}), \qquad \mathrm{QRead}(r, t_{b,\sigma}, w_{b,\sigma}),$$

one has

$$J_r(w_{\mathrm{in}}) = \widetilde{\mathcal{R}}_r(\mathrm{nf}(a_{\mathrm{in}})) = \widetilde{\mathcal{R}}_r(\mathrm{nf}(a_{b,\sigma})) = J_r(w_{b,\sigma}).$$

In particular, every closure-relevant coded readout of a certified OP 3 shell passage factors through one shell-stable kernel normal form.

Proof. By Proposition H2.161, a shell admissibility certificate implies

$$\mathrm{nf}(a_{b,\sigma}) = \mathrm{nf}(a_{\mathrm{in}}).$$

By clause (4) of Theorem H2.119, every closure-compatible coded readout r admits a factor map

$$\widetilde{\mathcal{R}}_r : \mathrm{nf}(\mathcal{A}) \to \mathrm{Im}(J_r)$$

such that whenever $\mathrm{QRead}(r, \langle 0, a \rangle, w)$,

$$J_r(w) = \widetilde{\mathcal{R}}_r(\mathrm{nf}(a)).$$

Applying this first to $s_{\mathrm{in}} = \langle 0, a_{\mathrm{in}} \rangle$ and then to $t_{b,\sigma} = \langle 0, a_{b,\sigma} \rangle$ yields

$$J_r(w_{\mathrm{in}}) = \widetilde{\mathcal{R}}_r(\mathrm{nf}(a_{\mathrm{in}})), \qquad J_r(w_{b,\sigma}) = \widetilde{\mathcal{R}}_r(\mathrm{nf}(a_{b,\sigma})).$$

Since the two kernel normal forms agree, the displayed chain follows. The final sentence is exactly the shell-stable readout factorization claim. □

Proposition H2.163 (First converse shell-certificate criterion)**.** *Assume the direct kernel constructor of Definition H2.118 and the bridge conditions of Theorem H2.119. Let*

$$s_{\mathrm{in}} = \langle 0, a_{\mathrm{in}} \rangle, \qquad t_{b,\sigma} = \langle 0, a_{b,\sigma} \rangle,$$

and suppose that the following data are given:

1. *there exists a coded branch state s_b with* $\mathrm{QBranchRef}(b, s_{\mathrm{in}}, s_b)$*;*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

2. *the branch code b is closure-admissible;*

3. *one has* $\mathrm{QShellRef}(\sigma, s_b, t_{b,\sigma})$ *and the refinement code σ is shell-admissible on b;*

4. *there exists a closure-compatible coded readout on the return state $t_{b,\sigma}$;*

5. *the kernel normal forms agree,* $\mathrm{nf}(a_{b,\sigma}) = \mathrm{nf}(a_{\mathrm{in}})$.

Then $(s_{\mathrm{in}}, b, \sigma, t_{b,\sigma})$ carries a shell admissibility certificate. Hence every closure-relevant coded readout of the passage factors through the same shell-stable kernel normal form.

Proof. The first four assumptions are exactly the four clauses required in Definition H2.158, so they furnish a shell admissibility certificate for $(s_{\mathrm{in}}, b, \sigma, t_{b,\sigma})$. The final sentence then follows from Theorem H2.162, using the additional kernel-persistence hypothesis only as the converse bridge datum that motivates the criterion. □

Lemma H2.164 (First OP 3/QAM shell commuting-square lemma)**.** *Assume the faithful translation theorem for the admissible OP fragment, Theorem H2.238, together with the OP 3 shell theorem, Theorem H7.21. Let*

$$\mathfrak{T}_{\mathrm{OP3}}$$

denote the admissible OP 3 shell-refinement passage on the OP side, and let

$$\Sigma_{b,\sigma}$$

denote the corresponding coded branch-shell refinement on the QAM side. Then the source-to-refinement square commutes on closure-relevant content in the following sense: whenever an admissible OP 3 shell passage is translated into coded QAM data,

$$\tau\big(\mathfrak{T}_{\mathrm{OP3}}(x)\big) \sim_{\mathrm{cl}} \Sigma_{b,\sigma}\big(\tau(x)\big),$$

where $\sim_{\mathrm{cl}}$ means equality after passage to the common closure-bearing class and, under the deferred kernel bridge, equality of the induced kernel normal form. In particular, OP 3 refinement and QAM shell refinement are not two competing shell operations; they are two readings of the same closure-preserving shell step.

Proof. By Theorem H2.238, the admissible OP fragment transports into coded QAM data without changing the source/return closure logic. By Theorem H7.21, the OP 3 shell passage is admissible exactly at the stage where the OP side isolates shell refinement without changing the closure-bearing content. The translated branch stage therefore produces a coded branch state s_b, and the translated shell stage produces a coded return state $t_{b,\sigma}$ with the same closure-relevant content as the OP-side refined state. Lemma H2.160 then identifies the translated source state and the coded branch-shell return state inside one coded closure class, while Proposition H2.161 upgrades this to equality of kernel normal forms once the deferred bridge is activated. That is exactly the asserted commuting-square statement. □

Theorem H2.165 (First OP 3/QAM shell correspondence theorem)**.** *Assume the faithful translation theorem for the admissible OP fragment, Theorem H2.238, together with the OP 3 shell theorem, Theorem H7.21. Then every translated OP 3-admissible shell passage determines a shell admissibility certificate in QAM, and every such certificate recovers the same closure-relevant shell content as the corresponding OP 3 step. Together with Proposition H2.163 and Lemma H2.164, the QAM shell-certificate layer may already be read as a first narrow converse guide for OP 3: admissible branch-shell behavior, a closure-compatible return readout, and kernel persistence are sufficient to recover the certificate package. Consequently, the QAM shell-certificate layer may be read as the first explicit coded closure guide for OP 3.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. Theorem H2.238 translates the admissible OP fragment into the coded QAM layer while preserving its source/return closure logic. Theorem H7.21 isolates the OP 3 shell-admissible passage itself. Under that translation, the OP-side branch step provides clause (1) of Definition H2.158, branch admissibility gives clause (2), OP 3 shell admissibility gives clause (3), and the translated closure-compatible readout gives clause (4). Hence each translated OP 3-admissible shell passage determines a shell certificate. Conversely, Lemma H2.160 and Theorem H2.162 show that any such certificate preserves exactly the closure-relevant content that the OP 3 theorem already isolates on the OP side. Lemma H2.164 records the same fact in commuting-square form. Thus the certificate layer is not a second closure notion; it is the coded OP 3 closure guide. □

Theorem H2.166 (First narrow shell-certificate iff package)**.** *Assume the direct kernel constructor of Definition H2.118 and the bridge conditions of Theorem H2.119. Let*

$$s_{\mathrm{in}} = \langle 0, a_{\mathrm{in}} \rangle, \qquad t_{b,\sigma} = \langle 0, a_{b,\sigma} \rangle,$$

and let r be any closure-compatible coded readout on the return state. Then the following are equivalent:

(i) $(s_{\mathrm{in}}, b, \sigma, t_{b,\sigma})$ *carries a shell admissibility certificate;*

(ii) there exists a coded branch state s_b such that

- *(a)* $\mathrm{QBranchRef}(b, s_{\mathrm{in}}, s_b)$,
- *(b) the branch code b is closure-admissible,*
- *(c)* $\mathrm{QShellRef}(\sigma, s_b, t_{b,\sigma})$ *and the refinement code σ is shell-admissible on b,*
- *(d) the return state $t_{b,\sigma}$ admits a closure-compatible coded readout,*
- *(e)* $\mathrm{nf}(a_{b,\sigma}) = \mathrm{nf}(a_{\mathrm{in}})$,
- *(f) every closure-relevant coded readout of the passage factors through one shell-stable kernel normal form.*

In particular, the present OP 3/QAM shell block already supports one narrow iff reading: a shell certificate is equivalent to the shell-stable kernel/readout package obtained from admissible branch-shell behavior together with kernel persistence.

Proof. Assume *(i)*. Then clauses (a)–(d) of *(ii)* are exactly the four certificate clauses from Definition H2.158. Clause (e) is Proposition H2.161, and clause (f) is Theorem H2.162. Hence *(ii)* follows.

Assume conversely *(ii)*. Then clauses (a)–(d) are precisely the data required by Definition H2.158, so $(s_{\mathrm{in}}, b, \sigma, t_{b,\sigma})$ carries a shell admissibility certificate. This is exactly Proposition H2.163. Thus *(i)* follows, and the final sentence is only a repackaging of the same forward and converse implications. □

Lemma H2.167 (First shell-obstruction lemma)**.** *Assume the direct kernel constructor of Definition H2.118 and the bridge conditions of Theorem H2.119. Let*

$$s_{\mathrm{in}} = \langle 0, a_{\mathrm{in}} \rangle, \qquad t_{b,\sigma} = \langle 0, a_{b,\sigma} \rangle,$$

and let r be any closure-compatible coded readout on the return state whenever such a readout exists. Suppose that at least one clause of the shell-stable kernel/readout package from Theorem H2.166 fails. Then no shell admissibility certificate exists for $(s_{\mathrm{in}}, b, \sigma, t_{b,\sigma})$. Equivalently, the first local OP 3 obstruction in the present QAM reading is the first failure of one of the following items:

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(i) admissible branch passage,

(ii) shell-admissible branch refinement,

(iii) closure-compatible return readout,

(iv) kernel persistence,

(v) shell-stable factorization of closure-relevant readout through one common kernel normal form.

In particular, a visible shell mismatch that leaves all five items intact is not yet an OP 3 obstruction in the present QAM sense.

Proof. By Theorem H2.166, a shell admissibility certificate is equivalent to the shell-stable kernel/readout package. Therefore the failure of any one clause of that package prevents the certificate from existing. The displayed list simply unwraps the package into its five operative clauses. The final sentence is the contrapositive reading of the same equivalence: visible shell variation alone is not an obstruction as long as admissible branch-shell transport, closure-compatible readout, kernel persistence, and common readout factorization still hold. □

Lemma H2.168 (First shell-obstruction stratification lemma)**.** *Assume the setup of Lemma H2.167. If no shell admissibility certificate exists for* $(s_{\mathrm{in}}, b, \sigma, t_{b,\sigma})$*, then at least one of the following three strata fails:*

(1) ***Branch stratum:*** *admissible branch passage fails.*

(2) ***Shell stratum:*** *admissible branch passage holds, but shell-admissible refinement fails.*

(3) ***Readout/kernel stratum:*** *admissible branch passage and shell-admissible refinement both hold, but closure-compatible return readout, kernel persistence, or common shell-stable readout factorization fails.*

Equivalently, every first local OP 3 obstruction in the present QAM reading is either a branch obstruction, a shell obstruction in the narrow sense, or a readout/kernel obstruction. In particular, whenever the branch and shell strata both remain intact, the first local obstruction is already forced into the readout/kernel layer.

Proof. Lemma H2.167 says that the first local obstruction is exactly the failure of one of five operative clauses. Group those clauses into the three strata above: clause (i) is the branch stratum, clause (ii) is the shell stratum, and clauses (iii)–(v) together form the readout/kernel stratum. This yields the stated trichotomy. The final sentence is only the contrapositive reading of the same grouping. □

Lemma H2.169 (Readout/kernel obstruction compression lemma)**.** *Assume the setup of Lemma H2.168, and assume moreover that the branch and shell strata both remain intact. Then the following are equivalent:*

(i) the first local OP 3 obstruction lies in the readout/kernel stratum;

(ii) at least one of the following failures occurs:

(a) closure-compatible return readout fails,

(b) kernel persistence fails,

(c) common shell-stable readout factorization through one kernel normal form fails;

(iii) there is no single shell-stable kernel/readout carrier through which all closure-relevant readout of the certified branch-shell passage can factor.

Equivalently, once the branch and shell strata are fixed, the whole remaining OP 3 obstruction is compressed to the failure of one common readout/kernel carrier class.

Proof. Under the standing assumption that the branch and shell strata remain intact, Lemma H2.168 says that any remaining first local obstruction must lie in the readout/kernel stratum. By definition of that stratum, this is exactly the disjunction of failures in clauses (a)–(c). This proves the equivalence of *(i)* and *(ii)*.

If a single shell-stable kernel/readout carrier exists, then closure-compatible return readout is available, kernel persistence identifies the same kernel normal form on source and return side, and Theorem H2.162 gives the common factorization through that carrier. Hence *(ii)* cannot occur. Conversely, if any one of clauses (a)–(c) fails, then no single shell-stable kernel/readout carrier can support the whole closure-relevant readout package. This proves the equivalence of *(ii)* and *(iii)*. The final sentence is only a compressed restatement of the same equivalence. □

Theorem H2.170 (First OP 3 shell-closure theorem in QAM form)**.** *Assume the faithful translation theorem for the admissible OP fragment, Theorem H2.238, together with the OP 3 shell theorem, Theorem H7.21. Let a translated OP 3-admissible shell passage determine coded data*

$$(s_{\mathrm{in}}, b, \sigma, t_{b,\sigma}, r).$$

Then the following are equivalent:

(i) the translated passage is QAM-shell-closed in the present narrow sense*, meaning that it carries a shell admissibility certificate and recovers the same closure-relevant shell content as the corresponding OP 3 step;*

(ii) $(s_{\mathrm{in}}, b, \sigma, t_{b,\sigma})$ *carries a shell admissibility certificate;*

(iii) the shell-stable kernel/readout package of Theorem H2.166 holds.

Moreover, under these equivalent conditions the passage admits a common shell-stable kernel/readout carrier through which all closure-relevant coded readout factors. If the equivalent conditions fail, then the first local OP 3 obstruction occurs at exactly one of the branch, shell, or readout/kernel strata from Lemma H2.168; and in the third stratum the whole remaining obstruction is precisely failure of one common shell-stable carrier in the sense of Lemma H2.169.

Proof. By Theorem H2.165, every translated OP 3-admissible shell passage determines a shell admissibility certificate and recovers the same closure-relevant shell content as the corresponding OP 3 step. Thus *(i)* and *(ii)* are equivalent in the present narrow reading.

The equivalence of *(ii)* and *(iii)* is exactly Theorem H2.166. Under these equivalent conditions, Theorem H2.162 gives factorization of every closure-relevant coded readout through the same shell-stable kernel normal form, hence through one common shell-stable kernel/readout carrier.

If the equivalent conditions fail, then by Lemma H2.167 the first local OP 3 obstruction is the first failure of the shell-stable kernel/readout package. Lemma H2.168 refines that failure into the branch, shell, and readout/kernel strata. Finally, Lemma H2.169 compresses the whole third stratum to failure of one common shell-stable carrier. This proves the final sentence. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Corollary H2.171 (Shell defect localization is the first local OP 3 failure)**.** *Within the present OP 3/QAM reading, visible packet mismatch, phase-lock deviation, or shell-sensitive surface variation is not yet by itself a shell defect. A genuine local OP 3 defect begins only when a shell defect witness in the sense of Definition H2.159 exists. Equivalently, by Theorem H2.170, the first local OP 3 failure is the first failure of the shell-stable kernel/readout package, stratified into branch, shell, and readout/kernel level and compressed in the third stratum to failure of one common shell-stable carrier. In particular, the first local failure is failure of a shell certificate clause, not failure of the entire OP 1–OP 5 closure chain.*

Remark H2.172 (How this block should be used in the next OP 3 proof pass)**.** *The present certificate block is intentionally narrow. It does not yet force a full equivalence between the OP and QAM formalisms, and it does not yet activate mixed tensors. Its current role is to give the next proof pass a fixed reading order: establish a shell certificate, pass to kernel persistence where the deferred bridge is available, use Theorem H2.162 to factor readout through the shell-stable class, use Proposition H2.163 for the first narrow return direction, use Lemma H2.164 to read OP 3 refinement and QAM shell refinement as one commuting closure step, use Theorem H2.166 as the first closed iff package, use Lemma H2.167 to identify the first genuinely local obstruction, use Lemma H2.168 to decide whether that first obstruction sits at the branch, shell, or readout/kernel layer, use Lemma H2.169 to compress the third stratum to failure of one common shell-stable carrier, and then use Theorem H2.170 as the first bundled OP 3 shell-closure statement in public-shape QAM form before asking whether any further tensor layer is genuinely needed.*

Current extracted OP 3/QAM tool layer

Remark H2.173 (Purpose of the extracted OP 3/QAM tools)**.** *The certificate and obstruction chain above is no longer meant to be reread from first principles in every later proof pass. Its immediate use is as a compact tool layer: once a later argument can cite one of the extracted tools below, it may treat the corresponding shell passage as already compressed to the right closure-bearing datum and continue at the next structural interface.*

Shell-certificate tool.
If one can exhibit an admissible branch passage, an admissible shell refinement, and a closure-compatible return readout, then the passage may be treated as a certified OP 3 shell step in the present narrow QAM sense. This is the entry tool for the whole block.

Kernel-persistence tool.
Once a shell certificate is in place and the deferred kernel bridge is available, the visible shell variation may be replaced by equality of one kernel normal form. From that point onward the proof may work kernel-first instead of shell-first.

Shell-stable readout tool.
Once the shell certificate and kernel bridge are active, every closure-relevant readout may be pushed through one common shell-stable carrier. Later readout comparisons therefore no longer need to unfold the whole branch/shell passage.

Converse reconstruction tool.
If admissible branch passage, admissible shell refinement, closure-compatible return readout, and kernel persistence are already known from another source, then the shell certificate may be reconstructed rather than reproved from scratch.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Commuting-square tool.
OP 3 shell refinement on the OP side and branch/shell refinement on the QAM side may be read as one closure-preserving square. This tool licenses switching between OP and QAM viewpoints without reopening the shell argument.

Obstruction-stratification tool.
If the shell-certificate package fails, the first local obstruction should be classified immediately as branch, shell, or readout/kernel level. This prevents visible shell mismatch from being over-read as a full OP 3 failure.

Carrier-compression tool.
Once branch and shell remain intact, every remaining local obstruction may be compressed to failure of one common shell-stable carrier. This is the shortest diagnostic form of the third obstruction stratum.

Bundled OP 3 closure tool.
Theorem H2.170 may now be cited as the compact public-shape closure package for OP 3 in QAM form: certificate, kernel/readout package, stratified obstruction, and common carrier are read as one bundled result.

Remark H2.174 (How to cite the extracted OP 3/QAM tools)**.** *A later argument should normally cite the smallest extracted tool that already matches its interface. Use the shell-certificate tool when the issue is admissible passage, the kernel-persistence or shell-stable readout tool when the issue is invariance under visible shell variation, the converse reconstruction tool when the shell certificate is to be recovered from already known transport/readout data, the commuting-square tool when the proof needs to switch languages, the obstruction-stratification or carrier-compression tool when the issue is failure localization, and the bundled OP 3 closure tool when the whole narrow closure package is intended at once.*

First OP 3 proof run through the extracted QAM tool layer

Remark H2.175 (Reading key for the first OP 3 proof run)**.** *In the present narrow public-shape QAM reading, the first proof run should be read as a deliberately compressed structural pass rather than as a maximal theorem expansion. The input is a translated OP 3 shell passage. The allowed move is passage from visible shell data to the extracted tool layer. The invariant is the common closure-bearing kernel/readout carrier whenever the certificate package is active. The defect signal is the first obstruction stratum from Lemma H2.168. In plain language: the proof run should decide as early as possible whether the shell step is already certified, and if not, where the first local failure sits.*

Theorem H2.176 (First OP 3 proof run through extracted QAM form)**.** *Assume the faithful translation theorem for the admissible OP fragment, Theorem H2.238, the OP 3 shell theorem, Theorem H7.21, the direct kernel constructor of Definition H2.118, and the bridge conditions of Theorem H2.119. Let*

$$\tau(x) = s_{\mathrm{in}} = \langle 0, a_{\mathrm{in}} \rangle$$

be the coded QAM source state obtained from an admissible OP 3 source configuration x, and let the translated branch/shell passage be encoded by a branch code b, a refinement code σ, a coded branch state s_b, and a coded return state

$$t_{b,\sigma} = \langle 0, a_{b,\sigma} \rangle.$$

Then the first OP 3 proof run through extracted QAM form has exactly two outcomes:

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(i) either the translated passage activates the bundled OP 3 closure tool, so that the OP 3 shell step is certified in QAM form, all closure-relevant coded readout factors through one common shell-stable carrier, and the passage may be continued kernel-first;

(ii) or the bundled closure package fails, in which case the first local obstruction is immediately localized by the obstruction-stratification tool at exactly one of the branch, shell, or readout/kernel strata, with the whole third stratum compressed to failure of one common shell-stable carrier.

In particular, the extracted tool layer already supports a genuine proof run: it converts translated OP 3 shell data into either a compressed closure package or a minimal obstruction diagnosis, without reopening the whole OP/QAM comparison at each step.

Proof. Translate the admissible OP fragment by Theorem H2.238. By the OP 3 shell theorem, Theorem H7.21, the translated branch and shell passage is exactly the stage at which the OP side isolates admissible shell refinement. Use the commuting-square tool to identify the translated OP 3 shell step with the coded branch/shell refinement step on the QAM side.

Now ask whether the shell-certificate tool is activated. If it is, then by the kernel-persistence tool the visible shell passage may be replaced by equality of one kernel normal form, and by the shell-stable readout tool every closure-relevant coded readout factors through one common shell-stable carrier. The bundled OP 3 closure tool then yields outcome *(i)*.

If the shell-certificate package is not activated, apply the obstruction-stratification tool. This localizes the first failure immediately at exactly one of the branch, shell, or readout/kernel levels. If the failure lies in the third stratum, the carrier-compression tool compresses the remaining failure to absence of one common shell-stable carrier. This is outcome *(ii)*.

The two outcomes are exhaustive because Theorem H2.170 already packages the positive direction as the bundled shell-closure theorem, while Lemma H2.168 together with Lemma H2.169 packages the negative direction. Hence the extracted tool layer already yields a genuine first OP 3 proof run in QAM form. □

Corollary H2.177 (First operational use of the extracted OP 3/QAM tools)**.** *A later OP 3 argument no longer needs to reopen the whole branch/shell analysis from scratch whenever a translated shell passage appears. It is enough to test whether the shell-certificate tool is active. If yes, cite the bundled OP 3 closure tool and continue kernel-first; if not, cite the obstruction-stratification tool and, in the third stratum, the carrier-compression tool. Thus the proof interface has already been shortened to a certified-pass / localized-obstruction dichotomy.*

Proof. This is only a restatement of Theorem H2.176 in proof-interface language. □

Remark H2.178 (Status separation for the present OP 3/QAM closure package)**.** *The present block should now be read in two layers with different evidential status. The* active public-shape result *is the certificate-level OP 3/QAM closure package: Definitions H2.158 and H2.159, Lemma H2.160, Theorem H2.166, Theorem H2.170, and Corollary H2.177. These statements already yield a closed certified-pass / localized-obstruction interface for translated OP 3 shell passages without leaving the active tuple surface. By contrast, the stronger kernel-first language — in particular Proposition H2.161, Theorem H2.162, and the carrier-compression reading of the third obstruction stratum — is a* conditional strengthening*: it becomes authoritative only under the deferred bridge hypotheses of Definition H2.114 and Theorem H2.119. Accordingly, later use should distinguish carefully between active certificate-level closure and conditional kernel/readout compression.*

Proposition H2.179 (Citation and use rule for the completed OP 3/QAM block)**.** *Let a later argument involve a translated OP 3 shell passage. Then the following citation discipline is valid.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(i) *If the argument needs only a certified-pass / localized-obstruction decision at the active tuple level, it may cite Theorem H2.170, Theorem H2.176, and Corollary H2.177 without invoking any kernel-first hypothesis.*

(ii) *If the argument further needs invariance under visible shell variation in kernel normal-form language or one-carrier readout factorization, then it may additionally cite Proposition H2.161, Theorem H2.162, and Lemma H2.169, but only after the deferred bridge hypotheses have been assumed or discharged.*

(iii) *If neither the certificate package nor the bridge package is available, the argument must fall back to the earlier branch/shell/readout layer and may not treat visible shell data as a closure theorem by themselves.*

Proof. Clause (i) is exactly the active reading packaged by Theorem H2.170, Theorem H2.176, and Corollary H2.177. Clause (ii) is the additional kernel-first strengthening isolated earlier in Proposition H2.161, Theorem H2.162, and Lemma H2.169; those statements explicitly depend on the deferred bridge hypotheses. Clause (iii) is the corresponding fallback rule: absent both packages, one must continue on the earlier active branch/shell/readout surface. □

Corollary H2.180 (Operational checklist for the completed OP 3/QAM block)**.** *A translated OP 3 shell passage should now be tested in the following order.*

1. *Check admissible branch passage and branch admissibility.*
2. *Check shell refinement and shell admissibility on the chosen branch.*
3. *Check existence of a closure-compatible return readout.*
4. *If all three checks succeed, activate the shell-certificate package and cite the bundled OP 3 closure tool.*
5. *Only if the deferred bridge hypotheses are additionally available, pass to kernel persistence and shell-stable readout factorization.*
6. *If one of the certificate checks fails, localize the first obstruction immediately at branch, shell, or readout/kernel level instead of reopening the whole OP 1–OP 5 chain.*

In particular, the completed block already provides one stable certified interface: certificate first, kernel compression second, obstruction localization immediately upon failure.

Remark H2.181 (Diagnostic handoff rule for visible shell data)**.** *Visible packet mismatch, phase-lock deviation, shell-sensitive surface variation, and analogous shell diagnostics remain* screening-layer data *unless and until they witness failure of one clause of the shell-certificate package or, under the bridge hypotheses, failure of one common shell-stable carrier. Thus the present block fixes a handoff rule from pictures and visible diagnostics to theorematic closure language: diagnostics may guide attention to the relevant branch, shell, or readout/kernel stratum, but they do not create a second closure axiom beside the completed OP 3/QAM certificate package. This is the intended local counterpart of the earlier visible/structural separation principle and the correct interface toward the later OP 3 to OP 5 transfer line.*

Proposition H2.182 (Prospective kernel-first reduction of the branch-shell theorem)**.** *Assume the direct kernel constructor of Definition H2.118 and the bridge conditions of Theorem H2.119. Let*

$$s_{\mathrm{in}} = \langle 0, a_{\mathrm{in}} \rangle$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

be the coded source state, let b be a closure-admissible branch code, let σ be a shell-admissible refinement code on b, and let

$$t_{b,\sigma} = \langle 0, a_{b,\sigma} \rangle$$

be an associated branch-shell coded return realization. Then the conclusion of Theorem H2.157 may be read kernel-first as follows:

1. *one has*
$$\mathrm{nf}(a_{b,\sigma}) = \mathrm{nf}(a_{\mathrm{in}}),$$
2. *hence every closure-compatible readout of s_{in} and $t_{b,\sigma}$ factors through the single canonical kernel class determined by a_{in},*
3. *therefore branch choice and shell refinement may be treated as closure-preserving already at exposed-kernel level before returning to the visible QAM state surface.*

Proof. Theorem H2.157 gives

$$\mathrm{QClEq}(s_{\mathrm{in}}, t_{b,\sigma}).$$

Since both states lie on the present state surface, we may write

$$s_{\mathrm{in}} = \langle 0, a_{\mathrm{in}} \rangle, \qquad t_{b,\sigma} = \langle 0, a_{b,\sigma} \rangle.$$

By the direct kernel constructor,

$$\ker_{\mathrm{dir}}(s_{\mathrm{in}}) = a_{\mathrm{in}}, \qquad \ker_{\mathrm{dir}}(t_{b,\sigma}) = a_{b,\sigma}.$$

The bridge clause for closure reflection therefore yields

$$\mathrm{nf}(a_{b,\sigma}) = \mathrm{nf}(a_{\mathrm{in}}).$$

Thus branch admissibility together with shell admissibility already identifies the same canonical kernel class before any further visible-surface comparison is carried out. By the factorization clause of the bridge datum, every closure-compatible coded readout depends only on that common kernel class. Hence the branch-shell theorem may be re-read as a kernel-first statement and only afterwards lifted back to the visible state surface. □

Remark H2.183 (Axiom audit after the kernel-first branch-shell reduction)**.** *Proposition H2.182 does* not *alter the active proof status of Theorem H2.157. The active theorem still rests only on closure-admissible branch refinement, shell-admissible OP 3 refinement, and closure-compatible coded readout. The kernel-first proposition adds force only under the deferred payload-level bridge hypotheses isolated in Definition H2.114 and Theorem H2.119. Thus no new hidden axiom has entered the live branch-shell theorem; only a prospective kernel-level compression of its proof surface has been recorded.*

Remark H2.184 (How much proof load the kernel-first branch-shell reading removes)**.** *At branch-shell level the cardinality factor is just* 1*: there is one visible source surface*

$$s_{\mathrm{in}} = \langle 0, a_{\mathrm{in}} \rangle$$

and one visible return surface

$$t_{b,\sigma} = \langle 0, a_{b,\sigma} \rangle.$$

So the gain is not a reduction in the number of cases, but a reduction in the size of the compared objects*. On the visible state surface the proof compares two full coded states and*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

then passes a closure-compatible readout across that state-level identification. In the deferred kernel-first reading, that work is replaced by the single normal-form check

$$\mathrm{nf}(a_{b,\sigma}) = \mathrm{nf}(a_{\mathrm{in}})$$

followed by one readout through the common kernel class. If one measures a visible-state comparison by cost C_{surf} and a kernel normal-form comparison by cost C_{ker}, then the conceptual load changes from roughly

$$C_{\mathrm{surf}} + C_{\mathrm{read}}$$

to

$$C_{\mathrm{ker}} + C_{\mathrm{class}},$$

with intended gain $C_{\mathrm{ker}} < C_{\mathrm{surf}}$ and typically $C_{\mathrm{class}} \ll C_{\mathrm{read}}$. So the numerical savings here are modest compared with the routed case, but methodologically this is the crucial base step: it is the first place where a full visible return state can be replaced by its exposed payload kernel without changing the closure conclusion.

Remark H2.185 (Theorem-level unpacking of the compact chain, I)**.** *Theorem H2.157 is the first theorem-level unpacking of the compact front statement: it isolates the middle part of the release chain in which closure-preserving source-return transport passes through branch choice and shell refinement without losing closure-relevant readout content.*

Corollary H2.186 (QAM branch-shell support class)**.** *Define the coded admissible branch-shell support class by*

$$\mathcal{S}_{\mathrm{adm}}^{\mathrm{QAM}} := \{\, (b,\sigma) : b \in \mathcal{B}_{\mathrm{adm}}^{\mathrm{QAM}} \ \wedge\ \sigma \in \mathcal{F}_{\mathrm{sh}}^{\mathrm{QAM}}(b) \,\}.$$

Then every coded return realization supported on

$$\mathcal{S}_{\mathrm{adm}}^{\mathrm{QAM}}$$

is closure-preserving at the coded QAM level.

Remark H2.187. *This is the coded QAM counterpart of the admissible branch-shell support sector in the present OP architecture.*

Definition H2.188 (QAM branch-shell routed return)**.** *Let*

$$u = \langle 8, \Omega, w \rangle$$

be an admissible routed support code with

$$\Omega \subseteq \mathcal{S}_{\mathrm{adm}}^{\mathrm{QAM}}.$$

A QAM branch-shell routed return *is a coded return aggregation associated with the support family Ω and the weight assignment w.*

Theorem H2.189 (QAM routed admissible support theorem)**.** *Let $u = \langle 8, \Omega, w \rangle$ be a normalized admissible routed support code, and let r be a closure-compatible coded readout. Then the closure-relevant image of the routed return determined by u agrees with the closure-relevant image of the coded source state.*

Proof. Every element $(b,\sigma) \in \Omega$ lies in the admissible branch-shell support class, so each associated coded return realization is closure-equivalent to the coded source state by the previous theorem. Since the routed support is normalized and the readout is closure-compatible, the routed aggregation preserves the same closure-relevant image after passage through the factor map. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proposition H2.190 (Prospective kernel-first reduction of the routed support theorem)**.** *Assume the direct kernel constructor of Definition H2.118 and the bridge conditions of Theorem H2.119. Let*

$$s_{\mathrm{in}} = \langle 0, a_{\mathrm{in}} \rangle$$

be the coded source state, and let

$$u = \langle 8, \Omega, w \rangle$$

be a normalized admissible routed support code. For each $(b, \sigma) \in \Omega$, let

$$t_{b,\sigma} = \langle 0, a_{b,\sigma} \rangle$$

be an associated branch-shell coded return realization. Then the conclusion of Theorem H2.189 may be read kernel-first as follows:

1. *for every $(b, \sigma) \in \Omega$ one has*
$$\mathrm{nf}(a_{b,\sigma}) = \mathrm{nf}(a_{\mathrm{in}}),$$
2. *hence every closure-compatible routed readout factors through the single canonical kernel class of a_{in},*
3. *therefore the routed aggregation determined by u has the same closure-relevant image as the coded source state without needing to compare the full family of visible state surfaces directly.*

Proof. For each $(b, \sigma) \in \Omega$, Theorem H2.157 gives

$$\mathrm{QClEq}(s_{\mathrm{in}}, t_{b,\sigma}).$$

Since both states lie on the present state surface, we may write

$$s_{\mathrm{in}} = \langle 0, a_{\mathrm{in}} \rangle, \qquad t_{b,\sigma} = \langle 0, a_{b,\sigma} \rangle.$$

By the direct kernel constructor, the kernel representatives are just the exposed payload kernels:

$$\ker_{\mathrm{dir}}(s_{\mathrm{in}}) = a_{\mathrm{in}}, \qquad \ker_{\mathrm{dir}}(t_{b,\sigma}) = a_{b,\sigma}.$$

The bridge clause for closure reflection therefore yields

$$\mathrm{nf}(a_{b,\sigma}) = \mathrm{nf}(a_{\mathrm{in}})$$

for every $(b, \sigma) \in \Omega$. Thus the entire admissible routed support family lies in one canonical kernel class. By the factorization clause of the bridge datum, every closure-compatible coded readout depends only on that kernel class. Hence the routed aggregation has the same closure-relevant image as the coded source state. The normalization of the routed weights matters only after this kernel reduction, so the proof burden is shifted from a family of full visible state surfaces to one canonical kernel class. □

Remark H2.191 (Why the routed support theorem is the best stress test)**.** *Among the currently marked finals, Theorem H2.189 is the heaviest set-theoretic test case for the deferred zero-kernel language: it simultaneously carries a support family Ω, a coded weight assignment w, branch admissibility, shell admissibility, routed aggregation, and closure-compatible readout. Exactly there the kernel-first reading is strongest, because the visible family of state surfaces is replaced by one canonical kernel class before the routed aggregation is evaluated.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Remark H2.192 (Axiom audit after the kernel-first routed reduction)**.** *Proposition H2.190 does* not *strengthen the active QAM axiomatics. Its extra force comes only from deferred hypotheses already isolated earlier: the direct kernel constructor of Definition H2.118 and the four payload-level bridge conditions of Theorem H2.119. In particular, the active theorem chain still proves Theorem H2.189 without mentioning kernels at all. The new proposition merely says that* if *those deferred bridge conditions are later discharged, then the routed-support proof may be re-read kernel-first. Thus no silent new axiom has entered the then-present public line; only a prospective compression interface has been exhibited.*

Remark H2.193 (How much comparison load the kernel-first reading removes)**.** *Write* $N := |\Omega|$ *for the size of the routed support family. On the visible state surface, the routed-support proof conceptually tracks* N *branch-shell return states*

$$t_{b,\sigma} = \langle 0, a_{b,\sigma} \rangle$$

and compares each of them with the coded source state at full state-surface level before the routed aggregation is passed through a closure-compatible readout. In the deferred kernel-first reading, those N *state-surface comparisons are replaced by the* N *normal-form checks*

$$\mathrm{nf}(a_{b,\sigma}) = \mathrm{nf}(a_{\mathrm{in}}), \qquad (b,\sigma) \in \Omega,$$

followed by a single *readout through the common canonical kernel class of* a_{in}*. Consequently the proof burden drops from "an* N*-member family of full visible state surfaces plus routed readout lifting" to "one canonical kernel class together with* N *membership/equality checks into that class." If one measures a full visible-state comparison by cost* C_{surf} *and a kernel normal-form comparison by cost* C_{ker}*, then the conceptual comparison load changes from roughly*

$$N\, C_{\mathrm{surf}} + C_{\mathrm{route}}$$

to

$$N\, C_{\mathrm{ker}} + C_{\mathrm{class}},$$

where typically $C_{\mathrm{class}} \ll C_{\mathrm{route}}$ *and the intended gain is precisely that* $C_{\mathrm{ker}} < C_{\mathrm{surf}}$*. In proof-shape terms, nothing miraculous happens to the factor* N *itself, but the* object size *being compared is reduced from full state surfaces to exposed kernels, and the routed readout is evaluated once on a single kernel class instead of being conceptually re-established across the whole visible family.*

Remark H2.194 (Theorem-level unpacking of the compact chain, II)**.** *Theorem H2.189 is the routed continuation of Theorem H2.157. Together they unpack the compressed public-shape chain*

$$s_{\mathrm{in}} \longrightarrow (b,\sigma) \longrightarrow u \longrightarrow r$$

of Theorem H2.2 into theorem-level branch/shell preservation followed by theorem-level routed aggregation.

Remark H2.195 (Staged proof compression pattern under the deferred zero-kernel language)**.** *The three prospective kernel-first reductions now show a genuinely staged compression pattern. At the first stage, Proposition H2.140 replaces the visible six-state transport chain by equality of six exposed kernel normal forms, so that admissible transport is already read as kernel-class preservation before any branch or shell data enter. At the second stage, Proposition H2.182 replaces one visible return-surface comparison*

$$\langle 0, a_{b,\sigma} \rangle \sim \langle 0, a_{\mathrm{in}} \rangle$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

by the single kernel normal-form comparison

$$\mathrm{nf}(a_{b,\sigma}) = \mathrm{nf}(a_{\mathrm{in}}).$$

At the third stage, Proposition H2.190 repeats that reduced comparison only at the level of exposed kernels across the whole routed family Ω and then evaluates the closure-compatible readout once on the common kernel class. Thus the deferred zero-kernel language first shrinks the transport chain to a common kernel class, then shrinks each branch-shell return to one exposed kernel comparison, and only afterwards turns the routed family into a class-membership problem rather than a full visible-surface transport problem. This is exactly why transport is the first chain-level compression step, branch-shell the first local return-surface compression step, and routed support the first true family-level compression step.

Definition H2.196 (Prospective activation package for the deferred zero-kernel language)**.** *A* zero-kernel activation package *for the present QAM surface consists of the following data and properties:*

(i) a total assignment on active state surfaces

$$\ker : QState \to \mathcal{A}, \qquad \ker(\langle 0, a\rangle) = a,$$

compatible with Definition H2.117;

(ii) kernel admissibility, meaning that every active exposed payload kernel lies in the admissible kernel class $\mathcal{A}$;

(iii) kernel normal-form reflection of closure, meaning that closure-equivalent active states have the same kernel normal form;

(iv) kernel-level factorization of every closure-compatible readout through the corresponding kernel normal form;

(v) compatibility of the routed aggregation step with passage to the common kernel class.

Theorem H2.197 (Activation criterion for kernel-first proof compression on the present QAM surface)**.** *Assume a zero-kernel activation package in the sense of Definition H2.196. Then the deferred zero-kernel language may be used as a genuine proof-compression layer for the present QAM surface in the following precise sense:*

(a) every active state-surface argument may be reduced to a kernel normal-form argument without loss of closure-relevant content;

(b) the first transport theorem may be read kernel-first by proving preservation of one common kernel normal-form class along the admissible transport chain;

(c) the branch-shell theorem may be read kernel-first by proving normal-form preservation of the exposed return kernels;

(d) the routed support theorem may be read kernel-first by proving membership of all exposed routed kernels in a single common normal-form class and evaluating the closure-compatible readout once on that class.

In particular, the deferred zero-kernel language ceases to be merely heuristic at exactly the point where closure, readout, and routed aggregation all factor through kernel normal forms. The criterion identifies what *must hold; the later staged-roadmap remarks then specify* in which order *those requirements should be established on the present QAM surface.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. This is just the joint content of Theorem H2.115, Theorem H2.119, Proposition H2.140, Proposition H2.182, and Proposition H2.190, collected as a single activation statement. Once closure-equivalent active surfaces have the same kernel normal form and every relevant readout factors through that normal form, every transport, branch-shell, and routed-support comparison may be carried out on kernels first and only lifted back to the visible state surface at the end. □

Remark H2.198 (Criterion-to-roadmap reading rule)**.** *The activation theorem above should be read together with the staged roadmap below. The theorem gives the logical threshold for activation, whereas the roadmap orders the internal work needed to reach that threshold:*

$$\begin{aligned} &\textit{transport-class preservation} \\ \Longrightarrow\ &\textit{local branch-shell kernel reduction} \\ \Longrightarrow\ &\textit{routed family compression.} \end{aligned}$$

In this combined reading, the criterion says when *kernel-first proof compression becomes legitimate, while the roadmap says* how *the present document should approach that point without introducing premature changes at visible tuple level.*

Remark H2.199 (Practical migration trigger for a later binary-tag version)**.** *The activation criterion above, read together with the later criterion-to-roadmap rule, also isolates the only later migration trigger that would make a binary or power-of-two tag scheme genuinely worthwhile. Such a migration would become structurally justified only once the present state surface is already being used kernel-first in proofs, so that the visible tag layer has become a comparatively thin presentation layer while closure and readout are actually controlled at kernel-normal-form level. Before that point, a binary tag migration would mainly rename visible surfaces; after that point, it could act as a real simplification of the presentation layer.*

Remark H2.200 (What still has to be shown before activation)**.** *At the present stage, the deferred zero-kernel language still stops short of activation for one simple reason: the document now contains the bridge data, the direct kernel final, and three staged stress tests, but it does not yet contain a fully internal proof that the active closure relation and every closure-compatible readout already factor through the exposed payload normal forms on the entire current QAM surface. The current status is therefore stronger than a mere metaphor but weaker than an active rewrite rule.*

Remark H2.201 (Prospective staged activation roadmap for the zero-kernel language)**.** *The current document already suggests a natural staged migration order for any later activation of the deferred zero-kernel language. Stage I is the* transport stage*: one first proves that admissible source-return transport preserves a single common kernel normal-form class all along the visible active chain. Stage II is the* local return stage*: one then proves that every admissible branch-shell return surface may be replaced by one exposed kernel comparison against the incoming kernel class. Stage III is the* family stage*: one finally proves that the whole routed family collapses to membership in that common kernel class, so that closure-compatible readout is evaluated once at class level rather than re-established on each visible routed surface. Only after these three stages are internally available would a later binary or power-of-two visible presentation acquire real proof-theoretic leverage, because only then would the visible tuple tags have become a thin presentation shell over an already kernel-first proof flow.*

Remark H2.202 (Why this staged order matters)**.** *The order above is not cosmetic. If routed family compression were attempted before transport-class preservation and local branch-shell*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

reduction were available, then the family step would still depend on full visible surfaces and would not yet be a genuine kernel-first simplification. The staged order therefore identifies the first point at which the deferred zero-kernel language would cease to be merely suggestive and would begin to remove real set-theoretic comparison load in a controlled way. Together with the criterion-to-roadmap rule, it turns the activation block from a static admissibility test into a concrete migration sequence.

Remark H2.203 (Compressed synthesis of the deferred zero-kernel path)**.** *The deferred zero-kernel material above should now be read as one compact path rather than as many separate addenda. The present visible state surface remains unchanged. A later kernel language would first expose the exposed payload kernel of a coded state as a final kernel, then compare closure only through kernel normal forms, then compress transport, local branch-shell return, and finally routed families in that order, and only afterwards consider whether a binary visible presentation would simplify the already kernel-first proof flow. In that reading, the deferred block does not introduce a second theory beside the current QAM package. It isolates one possible later proof-compression pipeline inside the same public branch/shell/routed line.*

Remark H2.204 (Reading guide for the deferred block)**.** *A compact reader may keep only the following chain in view:*

$$\textit{visible state surface} \implies \textit{exposed payload kernel} \implies \textit{kernel normal form}$$
$$\implies \textit{transport/local/family compression} \implies \textit{possible later binary presentation.}$$

The detailed deferred notes are included only to justify each arrow in this chain and to mark exactly which theorem-level bridges would still need to be proved before activation.

Remark H2.205 (Dependency map for the deferred zero-kernel path)**.** *The deferred block now has a strict internal dependency order. First, Definitions H2.101–H2.108 fix the later kernel vocabulary: carriers, filters, primitive zero-objects, recursive reuse, aspect projections, normal forms, and internal search keys. Second, the axiom-coherence and current-limitation remarks state why this added vocabulary does not yet compete with the active tuple layer. Third, Definition H2.114, Theorem H2.115, Definition H2.117, and Theorem H2.119 isolate the exact interface through which active coded states could later be re-read kernel-first. Fourth, Propositions H2.140, H2.182, and H2.190 test that interface on the three currently most relevant proof stages. Finally, Definition H2.196, Theorem H2.197, and the later activation-roadmap, synthesis, and reading-guide remarks convert those tests into an activation criterion and a migration order. In particular, any later binary visible presentation should depend on this whole chain and should not precede it.*

Remark H2.206 (Local admissibility test for a later kernel-first rewrite)**.** *The deferred block can now also be read as a local test for whether a specific visible tuple-surface proof step is mature enough for later kernel-first rewriting. Given a proof step whose visible data consist of active coded states*

$$s_i = \langle 0, a_i \rangle,$$

one should ask four questions in order.

1. Exposure: *do all exposed payload kernels a_i already lie in the intended admissible kernel domain $\mathcal{A}$?*

2. Reduction target: *can the visible conclusion be reformulated as equality, preservation, or common membership of the corresponding kernel normal forms* $\mathrm{nf}(a_i)$*?*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

3. Factorization: *does every closure-relevant readout or routed aggregation used in that proof step depend only on the resulting kernel normal-form class?*

4. Lift back: *once the kernel-class statement is proved, does the original visible state-surface conclusion follow again without adding new data?*

If the answer is yes in all four places, then the proof step is a genuine final for later kernel-first compression. If one of the four answers is still no, the corresponding proof step should remain on the visible state surface for the time being.

Remark H2.207 (Fallback rule for current proofs)**.** *Failure of the local admissibility test is not a defect of the present active proof line. It simply means that the relevant argument should continue to cite the active visible-surface theorem rather than its deferred kernel-first restatement. This fallback rule prevents premature theorem drift: the deferred block may prepare later compression, but it does not lower the evidential status required for current public proofs.*

Remark H2.208 (Compressed rewrite schema)**.** *Whenever the later local admissibility test applies, the deferred zero-kernel path suggests one uniform later rewrite schema:*

$$\textit{visible tuple step} \Longrightarrow \textit{expose payloads} \Longrightarrow \textit{pass to } \mathrm{nf}$$
$$\Longrightarrow \textit{prove closure/readout at class level} \Longrightarrow \textit{lift back to the visible surface.}$$

This does not yet activate any new theorem in the present document. It simply records, in one line, the exact proof pattern that the three staged stress tests have already isolated separately for transport, local branch/shell return, and routed families.

Remark H2.209. *This is the coded QAM analogue of the combined branch-shell transport theorem together with normalized routed admissible preservation.*

Definition H2.210 (QAM screening layer)**.** *A coded diagnostic layer is called a* screening layer *if it provides visible preselection information about admissible branch-shell support without itself constituting an independent closure proof.*

Remark H2.211. *In this sense, packet-, phase-lock-, and epsilon-type coded diagnostics should be treated in QAM as screening-layer data rather than as primary closure criteria. Their structural relevance begins only when they track loss of branch admissibility or loss of shell admissibility.*

Corollary H2.212 (QAM visible/structural separation)**.** *Visible coded mismatch in the screening layer does not by itself imply failure of closure at the QAM level. Closure failure begins only when coded branch admissibility or coded shell admissibility is lost.*

Remark H2.213. *This is the QAM-over-ZF shadow of the visible-mismatch-versus-structural-mismatch principle already established in the current Companion line.*

QAM tensor admissibility and tensor closure rules

We now strengthen the tensor-coded part of the QAM language. At the present stage, tensor composition is still not treated as a full tensor algebra. Instead, it is introduced as a controlled structural constructor equipped with admissibility, closure, transport, and readout compatibility rules.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition H2.214 (Tensor-coded formation)**.** *If p and q are QAM-coded objects, then*

$$p \otimes q := \langle 7, p, q \rangle$$

is their tensor-coded composite. We write

$$\mathrm{QTensor}(p \otimes q)$$

for the statement that the code has the tensor tag.

Definition H2.215 (Tensor formation by kind)**.** *The following tensor formations are allowed in principle:*

1. *state-state tensors:*
$$s_1 \otimes s_2;$$
2. *layer-layer tensors:*
$$\ell_1 \otimes \ell_2;$$
3. *operator-operator tensors:*
$$o_1 \otimes o_2;$$
4. *readout-readout tensors:*
$$r_1 \otimes r_2;$$
5. *branch-branch tensors:*
$$b_1 \otimes b_2;$$
6. *shell-shell tensors:*
$$\sigma_1 \otimes \sigma_2.$$

A tensor-coded composite is called well-typed *if both entries belong to one of the admissible kind-pairs listed above.*

Remark H2.216. *At this stage, mixed tensors between different kinds are not taken as primitive. They may be introduced later through derived coding conventions if needed.*

Definition H2.217 (Tensor admissibility)**.** *For QAM-coded objects x, y, we write*

$$\mathrm{QTensorAdm}(x, y)$$

iff:

1. *$x \otimes y$ is well-typed;*
2. *the tensor-coded composite preserves the intended closure-bearing structural regime;*
3. *the tensor-coded composite is admitted as a legitimate QAM object of its corresponding kind.*

Remark H2.218. *Thus tensor admissibility is not automatic from mere code formation. It is a structural compatibility condition.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition H2.219 (Tensor closure compatibility)**.** *For coded states s_1, s_2, t_1, t_2, we write*

$$\mathrm{QTensorCl}(s_1, s_2; t_1, t_2)$$

iff

$$\mathrm{QClEq}(s_1, t_1) \qquad \textit{and} \qquad \mathrm{QClEq}(s_2, t_2)$$

and the tensor composites

$$s_1 \otimes s_2 \qquad \textit{and} \qquad t_1 \otimes t_2$$

are tensor-admissible.

Axiom 5 (Tensor closure propagation schema)**.** *If*

$$\mathrm{QTensorCl}(s_1, s_2; t_1, t_2),$$

then

$$\mathrm{QClEq}(s_1 \otimes s_2,\ t_1 \otimes t_2).$$

Remark H2.220. *This is the first basic tensor closure principle: closure-equivalent components yield closure-equivalent tensor composites whenever tensor admissibility holds.*

Definition H2.221 (Tensor-coded transport)**.** *Let*

$$o_1 = \langle 2, \ell_1, \ell_1', G_1 \rangle, \qquad o_2 = \langle 2, \ell_2, \ell_2', G_2 \rangle$$

be coded operators. Their tensor-coded transport is

$$o_1 \otimes o_2.$$

We say that

$$\mathrm{QTensorTransport}(o_1, o_2; s_1, s_2; t_1, t_2)$$

holds iff:

1. $\mathrm{QAdmTransport}(o_1, s_1, t_1)$*;*
2. $\mathrm{QAdmTransport}(o_2, s_2, t_2)$*;*
3. $\mathrm{QTensorAdm}(s_1, s_2)$ *and* $\mathrm{QTensorAdm}(t_1, t_2)$*;*
4. $\mathrm{QTensorAdm}(o_1, o_2)$.

Axiom 6 (Tensor transport admissibility schema)**.** *If*

$$\mathrm{QTensorTransport}(o_1, o_2; s_1, s_2; t_1, t_2),$$

then

$$\mathrm{QAdmTransport}(o_1 \otimes o_2,\ s_1 \otimes s_2,\ t_1 \otimes t_2).$$

Remark H2.222. *This is the tensorial lift of admissible transport: admissible componentwise transport induces admissible tensor transport whenever the tensor compatibility conditions are satisfied.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition H2.223 (Tensor-coded readout)**.** *Let*

$$r_1 = \langle 3, \ell_1, W_1, R_1 \rangle, \qquad r_2 = \langle 3, \ell_2, W_2, R_2 \rangle$$

be coded readouts. Their tensor-coded readout is

$$r_1 \otimes r_2.$$

We say that $r_1 \otimes r_2$ *is* tensor closure-compatible *if:*

1. r_1 *and* r_2 *are closure-compatible;*
2. $\mathrm{QTensorAdm}(r_1, r_2)$*;*
3. *the closure-relevant readout factorization of* $r_1 \otimes r_2$ *depends only on the tensor closure class of the source state.*

Remark H2.224. *This is the tensor analogue of closure-compatible readout. It allows tensor-composite visible outputs to differ at raw level while preserving closure content after factorization.*

Definition H2.225 (Tensor branch admissibility)**.** *Let* b_1, b_2 *be closure-admissible branch codes. Their tensor composite*

$$b_1 \otimes b_2$$

is called tensor branch-admissible *if:*

1. $\mathrm{QTensorAdm}(b_1, b_2)$*;*
2. *componentwise branch refinement preserves closure class;*
3. *the induced tensor branch refinement preserves closure class at the tensor-composite level.*

Definition H2.226 (Tensor shell admissibility)**.** *Let* σ_1, σ_2 *be shell-admissible refinement codes. Their tensor composite*

$$\sigma_1 \otimes \sigma_2$$

is called tensor shell-admissible *if:*

1. $\mathrm{QTensorAdm}(\sigma_1, \sigma_2)$*;*
2. *componentwise shell refinement preserves closure class;*
3. *the induced tensor shell refinement preserves closure class at the tensor-composite level.*

Remark H2.227. *Tensor branch admissibility and tensor shell admissibility are the coded QAM precursors of coupled branch/shell transport across multiple structural sectors.*

Proposition H2.228 (Tensor closure preservation principle)**.** *Let* s_1, s_2, t_1, t_2 *be coded states. Assume*

$$\mathrm{QClEq}(s_1, t_1), \qquad \mathrm{QClEq}(s_2, t_2), \qquad \mathrm{QTensorAdm}(s_1, s_2), \qquad \mathrm{QTensorAdm}(t_1, t_2).$$

Then

$$\mathrm{QClEq}(s_1 \otimes s_2,\ t_1 \otimes t_2).$$

Proof. This is exactly the tensor closure propagation schema. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proposition H2.229 (Tensor transport preservation principle)**.** *Assume*

$$\mathrm{QTensorTransport}(o_1, o_2; s_1, s_2; t_1, t_2).$$

Then

$$\mathrm{QClEq}(s_1 \otimes s_2,\ t_1 \otimes t_2).$$

Proof. By tensor transport admissibility,

$$\mathrm{QAdmTransport}(o_1 \otimes o_2,\ s_1 \otimes s_2,\ t_1 \otimes t_2).$$

Admissible transport implies closure preservation, hence

$$\mathrm{QClEq}(s_1 \otimes s_2,\ t_1 \otimes t_2).$$

□

Theorem H2.230 (First tensor QAM transport theorem)**.** *Let*

$$o_1, o_2$$

be coded operators,

$$s_1, s_2$$

coded source states, and

$$t_1, t_2$$

coded transported states. Assume

$$\mathrm{QTensorTransport}(o_1, o_2; s_1, s_2; t_1, t_2),$$

and let

$$r_1 \otimes r_2$$

be a tensor closure-compatible readout. Then the tensor-composite source state and the tensor-composite transported state lie in the same coded closure class:

$$\mathrm{QClEq}(s_1 \otimes s_2,\ t_1 \otimes t_2),$$

and therefore their closure-relevant tensor readout content agrees after factorization through the tensor class-target map.

Proof. By the previous proposition,

$$\mathrm{QClEq}(s_1 \otimes s_2,\ t_1 \otimes t_2).$$

Since the tensor readout is closure-compatible at tensor level, closure-equivalent tensor-composite states have the same closure-relevant image after factorization. This proves the claim. □

Corollary H2.231 (Tensor branch-shell admissible support)**.** *If*

$$b_1 \otimes b_2$$

is tensor branch-admissible and

$$\sigma_1 \otimes \sigma_2$$

is tensor shell-admissible, then admissible branch-shell support is preserved at tensor-composite level, provided the relevant tensor admissibility conditions hold.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Remark H2.232. *This is the first indication that the current branch-shell architecture of the Companion can be lifted into a tensor-coded QAM language without changing its closure-first logic. In the internal reading order, this tensor layer remains subordinate to the earlier branch/shell admissibility and translation layers.*

Remark H2.233 (Current status of tensor QAM)**.** *At the present stage, the tensor rules do not yet define a full monoidal calculus. They only establish the first closure-preserving, transport-preserving, and readout-compatible tensor schemata needed for a structure-first QAM development over ZF.*

QAM translation theorem for the admissible OP fragment

We now make explicit that the present OP/branch/shell architecture may be translated into the coded QAM-over-ZF language without changing its closure-first logic. In archival stage this translation block should be read as the theorem-level mirror of the front extraction theorem (Theorem H2.2): the front block states the compressed public-shape chain, while the present subsection states why that same chain is a faithful coded image of the already stabilized OP architecture. At the current stage, this is not yet claimed as a full equivalence of entire formal systems. It is claimed only as a faithful translation on the admissible fragment of the OP transport architecture. Accordingly, the present subsection is ordered to match the front reading guide: coded source state and closure-compatible coded readout first, closure-admissible branch code second, shell-admissible refinement code third, admissible routed support fourth, and closure-relevant readout content fifth.

Definition H2.234 (Translation map from the admissible OP chain into QAM codes)**.** *Let* $\mathfrak{O}_{\mathrm{adm}}$ *denote the admissible OP fragment consisting of:*

1. *source and return states of the OP1–OP5 source-return architecture;*
2. *intermediate admissible OP transport states;*
3. *admissible branch data from the branchwise refinement layer;*
4. *OP3 shell-admissible refinement data;*
5. *admissible and normalized routed support data.*

A QAM translation map *is a coding assignment*

$$\tau : \mathfrak{O}_{\mathrm{adm}} \to \mathfrak{Q}_{\mathrm{adm}}$$

from admissible OP objects into admissible QAM-coded objects such that:

1. *source/readout objects are sent to coded source states and coded closure-compatible readouts of the same front-facing type used in the public-shape extraction;*
2. *OP transport maps are sent to coded operator maps preserving the admissible transport role between coded source, branch, shell, routed-support, and readout stages;*
3. *admissible branches are sent to closure-admissible branch codes;*
4. *OP3 shell-admissible refinements are sent to shell-admissible refinement codes;*
5. *admissible and normalized routed support is sent to admissible and normalized routed support codes.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Remark H2.235 (Reading-order convention for τ)**.** *The point of τ is not merely syntactic recoding. It must preserve the structural roles of the translated objects in the same order in which the front extraction reads them: coded source state and closure-compatible coded readout remain the coded front-face of the OP 1/OP 5 source-return pair; admissible transport remains admissible transport; branch remains branch; shell-admissible refinement remains shell-admissible refinement; routed support remains routed support; and closure-relevant readout content remains closure-relevant readout content.*

Definition H2.236 (Closure-faithful translation on the public-shape chain)**.** *The translation map τ is called* closure-faithful *if for all admissible OP objects $X, Y \in \mathfrak{O}_{\mathrm{adm}}$,*

$$X \sim_{\mathrm{cl}} Y \qquad \Longrightarrow \qquad \mathrm{QClEq}(\tau(X), \tau(Y)),$$

and whenever the OP-side relation expresses admissible preservation of the native closure-bearing mode along the source/transport/branch/shell/routed chain, the translated QAM-side relation expresses preservation of the same closure content in coded form.

Definition H2.237 (Readout compatibility of the translation)**.** *The translation map τ is called* readout-compatible *if OP-side closure-compatible readout invariants correspond, after translation, to QAM-coded closure-compatible readouts on the image of τ, with the same closure-relevant readout content as in the public-shape chain.*

Then the following OP-side chain is preserved under translation into QAM-over-ZF in exactly the order used by the front public-shape reading guide:

Theorem H2.238 (QAM translation theorem for the admissible OP fragment)**.** *Assume that:*

1. *the OP transport architecture is given on its admissible fragment $\mathfrak{O}_{\mathrm{adm}}$;*
2. *there exists a closure-faithful and readout-compatible translation map*

$$\tau : \mathfrak{O}_{\mathrm{adm}} \to \mathfrak{Q}_{\mathrm{adm}};$$

3. *admissible OP transport, admissible branches, OP3 shell-admissible refinements, admissible routed support, and closure-compatible readout are translated according to the preceding definitions.*

Then the following OP-side chain is preserved under translation into QAM-over-ZF in exactly the order used by the front public-shape reading guide:

$$\begin{aligned} \textit{coded source state / OP1–OP5 readout face} &\longrightarrow \textit{closure-admissible branch code} \\ &\longrightarrow \textit{shell-admissible refinement code} \\ &\longrightarrow \textit{admissible routed support code} \\ &\longrightarrow \textit{closure-relevant coded readout content.} \end{aligned}$$

Equivalently, after naming the translated objects in public-shape order, one obtains the same compressed chain from Theorem H2.2:

$$\begin{aligned} s_{\mathrm{in}} &\longrightarrow (b, \sigma) \\ &\longrightarrow u \\ &\longrightarrow r. \end{aligned}$$

More precisely:

© 2026 Marc Brendecke. Preprint — not peer reviewed.

1. ***OP closure preservation.*** *If the OP1–OP5 source-return pair is closed at invariant level, then its QAM-coded image is closure-preserving at coded level.*

2. ***Intermediate transport preservation.*** *If the intermediate OP layers preserve the native closure-bearing mode, then their QAM-coded images preserve the coded closure class.*

3. ***Branch admissibility preservation.*** *If a branch is closure-admissible on the OP side, then its image under τ is a closure-admissible branch code on the QAM side.*

4. ***Shell admissibility preservation.*** *If an OP3 refinement is shell-admissible on the OP side, then its image under τ is a shell-admissible refinement code on the QAM side.*

5. ***Routed admissible support preservation.*** *If a routed support sector is admissible and normalized on the OP side, then its image under τ is an admissible and normalized routed support code on the QAM side, and its closure-relevant content is preserved.*

6. ***Front-facing readout compatibility.*** *If the translated routed support sector is evaluated by a closure-compatible OP readout, then the resulting coded readout is exactly of the public-shape type used in Theorem H2.2; in particular it remains a coded readout of preserved closure content and not a new primitive closure test.*

Proof. Each clause follows from closure-faithfulness and readout compatibility of the translation map.

For (1), closure of the OP1–OP5 pair means preservation of the native closure-bearing mode at the source-return level. By closure-faithfulness, this becomes coded closure preservation for the translated source and return states.

For (2), invariant preservation across intermediate OP layers is, by assumption, preservation of the same closure-bearing mode through admissible transport. Again, closure-faithfulness translates this into preservation of coded closure class for the corresponding QAM-coded states and operator actions.

For (3), branch admissibility on the OP side means precisely that the branchwise refinement preserves the native closure-bearing mode. By the definition of the translation map, admissible branches are sent to branch codes that preserve coded closure class.

For (4), shell admissibility at OP3 means that the refinement does not change the native closure-bearing mode. By translation, this becomes shell-admissibility of the corresponding QAM refinement code.

For (5), admissible routed support is translated by hypothesis into admissible routed support codes. Readout compatibility ensures that closure-relevant readout content is preserved after translation, and normalization of support weights is preserved by the routed support coding layer.

For (6), readout compatibility identifies the translated readout with a coded closure-compatible readout on the image of the admissible OP chain. Hence the front-facing public-shape readout statement is not extra structure; it is the coded readout face of the same preserved OP closure content. □

Remark H2.239 (Prospective local admissibility reading of the translation theorem)**.** *The later local admissibility test may already be applied, in deferred form, to Theorem H2.238. The result is not yet a full active kernel-first rewrite of the translation theorem, but it cleanly separates the future admissible rewrite points.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

1. *Clauses (1)–(2) are locally admissible exactly at those translated source/transport surfaces that already lie on the present visible state surface $\langle 0, a\rangle$; there the payload can be exposed directly and compared by kernel normal form.*

2. *Clauses (3)–(5) are locally admissible once the translated branch, shell, and routed-support codes are shown to inherit the same exposed-kernel discipline used in Propositions H2.182 and H2.190.*

3. *Clause (6) becomes locally admissible precisely when translated closure-compatible readout factors through the same kernel normal-form class, so that the coded public-shape readout is recovered from class level rather than re-established on the full visible translated chain.*

Accordingly, the translation theorem is already organized in the right order for later kernel-first compression, but it becomes genuinely rewrite-ready only stage by stage, in the same order isolated by the staged roadmap and the criterion-to-roadmap rule.

Remark H2.240 (How much clausewise load a future kernel-first translation proof could remove)**.** *At present, Theorem H2.238 is proved clausewise on the visible translated chain, with six explicit front-facing obligations. A later kernel-first proof would not remove the need to show transport, branch, shell, routed-support, and readout preservation; however, it could compress the visible proof surface from six separate clause-level checks to two kernel-level obligations:*

(i) one common kernel-normal-form preservation statement along the translated admissible chain,

(ii) one class-level readout factorization statement recovering the coded public-shape readout.

In that sense, the prospective savings at theorem level are best read not as a reduction in mathematical content, but as a reduction from six visible surface checks *to* one kernel-class propagation argument plus one class-level readout argument. *This is exactly the same kind of shift from surface multiplicity to kernel control already seen, in more local form, in the transport, branch/shell, and routed-support stress tests.*

Corollary H2.241 (QAM image of the current OP core)**.** *The current OP core of the Companion — including source-return closure, intermediate transport, admissible branches, OP3 shell admissibility, admissible routed support, and closure-compatible readout — admits a faithful coded realization inside QAM-over-ZF.*

Remark H2.242 (Front theorem and translation theorem as mirrored statements)**.** *Theorem H2.2 states the compressed public release chain in coded language. Theorem H2.238 states that the same coded chain is the faithful image of the stabilized OP chain. Read together with Theorem H2.157, Theorem H2.189, and Corollary H2.212, the front block and the later QAM block are therefore intended to read almost as two projections of the same closure statement: first in compressed public-shape form, then in theorem-level translation form. In particular, the front reading order*

$$s_{\mathrm{in}} \longrightarrow (b, \sigma) \longrightarrow u \longrightarrow r$$

is now the same order in which τ, the closure-faithfulness condition, the readout-compatibility condition, and the partial decoding map δ are meant to be understood.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Remark H2.243. *This corollary is the first formal justification for introducing QAM as a structure-first coded language over ZF. It shows that the current OP architecture does not need to be abandoned or replaced in order to enter the QAM setting; it can be carried into that setting by faithful translation.*

Definition H2.244 (Partial decoding map)**.** *A* partial decoding map *is a map*

$$\delta : \mathfrak{Q}_{\mathrm{adm}} \dashrightarrow \mathfrak{O}_{\mathrm{adm}}$$

defined on the admissible QAM fragment such that for translated admissible OP objects,

$$\delta(\tau(X)) = X.$$

Remark H2.245 (How δ should be read)**.** *At the present stage, δ is only required on the image of the admissible OP fragment. No full decoding of arbitrary QAM-coded expressions is assumed yet. When it exists, δ should be read as the reverse reading map for the same public-shape chain: it sends coded source/readout, branch, shell, and routed-support objects back into the already stabilized OP architecture without introducing a second closure criterion.*

Proposition H2.246 (Round-trip property on the admissible fragment)**.** *If a partial decoding map δ exists on the image of τ, then*

$$\delta \circ \tau = \mathrm{id}_{\mathfrak{O}_{\mathrm{adm}}}.$$

Hence the admissible OP fragment is recoverable from its translated QAM image.

Proof. This is immediate from the definition of the partial decoding map. □

Remark H2.247. *The round-trip property is intentionally weaker than full two-way interpretability. It says only that the currently relevant admissible OP fragment can be encoded into QAM over ZF and then read back without structural loss.*

Corollary H2.248 (Tensor-compatible translation)**.** *If the admissible OP fragment contains tensorially paired sectors and the translation map respects tensor-coded formation, then admissible tensor closure and admissible tensor transport are preserved under translation as well.*

Remark H2.249 (Current status of the translation theorem)**.** *The present theorem should be read as a conservative first step. It does not yet prove that all future QAM constructions are already reducible to the current OP architecture, nor that the entire QAM language has been fully decoded back into OP form. What it proves is that the current admissible OP core already embeds faithfully into the coded QAM-over-ZF language.*

H3 Pre-Closure Tools and Earlier Synthesis

Historical tool layer and pre-closure live cores

Status. *Historical proof-memory, not the active theorem core.*

Block function. This section preserves older tool extractions and live-burden records for genealogy and local reconstruction, but current burden labels and current routing rules are governed by the active main line.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Remark H3.1. *This block is retained only as a historical record of the last pre-closure burden analysis. It preserves the compression logic from which the OP transport theorem and the OP3 residual/shell interface were extracted, but it no longer states the active proof core of the document.*

Remark H3.2 (v4.6 supersession note)**.** *The OP 1/OP 5 residue language in this historical block is a pre-v4.4/v4.5 burden record. In the v4.6.0 release-final reading, the active boundary is not an unresolved pair of independent local OP residues. The current boundary is the all-readout/T2 scope problem: the bounded readout surface is conservative inside the declared grammar, while the universal exterior selector, all-readout naturality family, and unrestricted exterior collar slot remain outside the proved layer. The relevant next program is therefore MM/HC exterior–mixed field-equation determination.*

Current extracted tool layer (historical extraction snapshot)

Transport-first tool. Whenever a final closure statement mixes continuum transport with visible readout, the admissible move is to fix the transport branch first and delay HC-readout to the downstream stage. In particular, transport selects the branch; readout does not create a second primitive continuum channel.

Readout-after-transport tool. Once the branch has been fixed by admissible transport, the Heisenberg-compensator readout is treated as a downstream projection or readout map on that branch. It may compress the visible law, but it does not reopen the continuum-side selection problem.

Admissible-readout uniqueness tool. Once a transport branch and its matter-frame parameter have already been fixed, every further canonical positive scalar readout must remain a downstream realization of that same admissible body. If no second primitive continuum channel opens and grouped-layer transport produces no ontology drift, then canonical normalization cannot differ from the standard matter-frame readout without reintroducing exactly the forbidden second channel.

Checkpoint-absorption tool. A recovered checkpoint may be absorbed into the active line whenever it introduces no new primitive visible scalar, no second independent branch law, and no incompatible residual interface. In that case the checkpoint is proof-memory and normalization support, not a parallel mathematical front.

Epsilon-trap tool. Let

$$\mathfrak{b}_{N,m} := \iota^{(Y)}_{N,m} + \sin|\kappa_{N,m}|, \qquad \mathfrak{b}_{\mathrm{crit}} := 0.578937.$$

If on an admissible window one has entry slack

$$\Sigma_{N,m} := \mathfrak{b}_{\mathrm{crit}} - \mathfrak{b}_{N,m} \geq \varepsilon_{\mathrm{trap}} > 0$$

and the next-step drift of $\mathfrak{b}$ does not exceed that slack, then the branch remains inside the admissible tube. This is the current non-escape tool for the first OP 5 valley window.

OP 1 epsilon-bridge tool. Let γ_{br} be the distinguished bridge root and let T_μ be the bridge self-consistency map from Section 3.8.16. If one works in a local tube $|b - \beta_{\mathrm{br}}| \leq \varepsilon_{\mathrm{br}}$ on which

$$\sup_{|b-\beta_{\mathrm{br}}|\leq\varepsilon_{\mathrm{br}}} \left|\partial_b T_{\gamma_{\mathrm{br}}}(b)\right| \leq L_{\mathrm{br}} < 1,$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

then entry into that tube implies non-escape and convergence to the same bridge fixed point. In particular, once the normalization defect is shown to lie inside a contractive ε-tube around γ_{br}, OP 1 becomes a local fixed-point identification problem rather than a second branch-search problem.

Shared matrix-leakage tool. Let J denote the Umkehrwalze/involution, let $C := \frac{1}{2}(\mathrm{id}+J)$ be the HC readout projector, and let $\mathcal{M}$ denote the Millennium matrix on the active grouped transport layer. The first shared obstruction channel for OP 1 and OP 5 should be tested through the two leakage diagnostics

$$E_{\mathrm{comm}} := [\mathcal{M}, J], \qquad E_{\mathrm{leak}} := (\mathrm{id} - C)\mathcal{M}C.$$

If both vanish on the active branch, then the admissible channel is stable under transport, reversal, and HC readout. If one or both persist, then the remaining OP 1 bridge residue and the surviving OP 5 valley residue should first be treated as two readouts of the same underfixed matrix core rather than as independent end problems.

Cryptanalytic translation tool. The matrix diagnostics above admit a structure-first translation of three familiar attack styles. *Linear cryptanalysis* becomes the search for biased linear masks on the even/odd and visible/orthogonal matrix blocks, especially for linear functionals that simultaneously see the OP 1 bridge scalar and the OP 5 first-window branch load. *Differential cryptanalysis* becomes the study of controlled perturbations of branch/readout inputs and their propagation through $\mathcal{M}$, J, and C, in order to detect whether small input differences are amplified mainly in the same shared leakage sector. *Linear-differential cryptanalysis* becomes the mixed test in which one first propagates a controlled perturbation through the matrix layer and then reads the output through a chosen linear mask. In the present framework this is the natural way to test whether a single hidden leakage mode explains both the OP 1 bridge defect and the OP 5 valley defect.

Shared phase-coupling tool. Once a perturbation family $\lambda \mapsto x_\lambda$ has been chosen, define the visible OP 1 and OP 5 residual readouts along that family and normalize them to dimensionless traces. The decisive test is then whether the two normalized traces occupy one nearly one-dimensional line in the defect plane (same carrier, same phase up to sign) or a near-elliptic/quadrature orbit (same carrier, phase shift close to $\pm\pi/2$). In either case the question is no longer whether OP 1 and OP 5 are independent, but whether they are two masked readouts of one shared hidden oscillatory leakage mode.

Shared leakage probe-family tool. Rather than perturbing the entire Millennium matrix blindly, choose a baseline HC-even state v and a bounded test matrix K, and probe only the cross-sector channel through

$$\Delta_{\mathrm{shared}} := (\mathrm{id} - C)KC + CK(\mathrm{id} - C), \qquad \mathcal{M}_\lambda := \mathcal{M} + \lambda\,\Delta_{\mathrm{shared}}.$$

This family leaves the pure even/even and odd/odd blocks untouched to first order and perturbs only the admissible/complementary coupling. It is therefore the preferred one-parameter probe for deciding whether OP 1 and OP 5 really see the same hidden leakage mode.

Historical live-burden record prior to OP transport closure

OP 1 and OP 5. At the last pre-closure stage, the OP 1 burden had already collapsed to one local bridge remainder, while the OP 5 burden had collapsed to first-window non-escape

© 2026 Marc Brendecke. Preprint — not peer reviewed.

on an admissible epsilon-trapped branch. The present OP transport theorem absorbs that compression stage and turns the former burdens into historical extraction data rather than active proof cores.

Shared matrix leakage and cryptanalytic translation

The present state of the document suggests that the surviving OP 1 and OP 5 residues may be two visible readouts of one still underfixed HC/MM/J coupling. In that situation it is more efficient to attack the common matrix core than to continue treating both residuals as fully separate local endgames.

Definition H3.3 (Provisional shared matrix leakage channels)**.** *Let J be the involutive Umkehrwalze, let*

$$C := \frac{1}{2}(\mathrm{id} + J)$$

be the HC readout projector, and let $\mathcal{M}$ be the Millennium matrix on the active grouped transport layer. Define the two basic leakage channels

$$E_{\mathrm{comm}} := [\mathcal{M}, J], \qquad E_{\mathrm{leak}} := (\mathrm{id} - C)\mathcal{M}C.$$

The first channel measures failure of transport/reversal compatibility; the second measures leakage of the admissible HC-even sector into the complementary sector after Millennium transport.

Proposition H3.4 (Provisional stable-channel criterion)**.** *If on the active branch one has*

$$E_{\mathrm{comm}} = 0, \qquad E_{\mathrm{leak}} = 0,$$

then the admissible HC channel is stable under Millennium transport and Umkehrwalze reversal. In particular, no first-order matrix-induced off-sector drift remains that could independently separate a bridge readout from a first-window valley readout.

Proof. The identity $E_{\mathrm{comm}} = 0$ says that $\mathcal{M}$ and J commute, so reversal and Millennium transport preserve the same even/odd splitting. The identity $E_{\mathrm{leak}} = 0$ says that applying $\mathcal{M}$ to an HC-even state leaves no component in the HC-odd complement. Together these identities mean that the admissible channel selected by C is invariant under the combined transport/reversal layer. Hence any remaining visible residue must come from some other source, not from first-order matrix leakage between admissible and complementary sectors. □

Remark H3.5 (Provisional OP 1/OP 5 coupling heuristic)**.** *If OP 1 and OP 5 both remain slightly open after branch-fixing and readout-fixing, then the most economical explanation is that the shared matrix core is not yet sharp enough. In that reading, the bridge residue of OP 1 and the first-window branch residue of OP 5 are two different masks of the same small leakage mode. The diagnostics of Definition H3.3 are therefore not a side remark; they are the first place where the two remaining channels should be tested jointly.*

Cryptanalytic translation. The same matrix question admits a structure-first translation of three familiar cryptanalytic styles.

Linear route. Choose linear masks on the J-even/odd block decomposition or on visible/orthogonal packet channels, and test whether the masked OP 1 bridge residue and the masked OP 5 first-window residue exhibit the same small bias. In the present framework, this means searching for linear functionals on $\mathcal{M}$ that simultaneously see both open channels.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Differential route. Apply controlled perturbations to the input branch/readout data and track the induced difference propagation through $\mathcal{M}$, J, and C. If the same perturbation pattern repeatedly feeds both the bridge and valley residues, this is strong evidence for a shared leakage sector rather than two independent local failures.

Linear-differential route. First propagate a controlled perturbation through the matrix layer, then read the result through a chosen linear mask. This mixed route is often the sharpest one when a hidden mode is too weak to dominate either purely linear or purely differential tests by itself but becomes visible once one combines propagation with selective masked readout.

Provisional consequence. These tools do not yet prove that the shared leakage mode is nonzero, nor that it completely explains both OP 1 and OP 5. What they do show is where the next efficient attack should land: not on arbitrary matrix fitting, but on the smallest block of $\mathcal{M}$ that still fails to commute with J or still leaks through $(\mathrm{id} - C)\mathcal{M}C$.

Shared phase-coupling diagnostic for OP 1 and OP 5

The remaining OP 1 and OP 5 residues may still be too coupled to close independently. A natural last diagnostic is therefore whether they are driven by one common hidden carrier: either nearly in phase, or approximately in quadrature like sine and cosine. This is stronger than asking whether both are merely small. It asks whether the two visible end residues are just two phase-shifted readouts of the same underfixed matrix mode.

Definition H3.6 (Provisional normalized residual traces along a perturbation family)**.** *Let $\lambda \mapsto x_\lambda$ be an admissible one-parameter perturbation family on the active branch, obtained for example by controlled transport/readout perturbations in the sense of the differential or linear-differential tools above. Let*

$$d_1(\lambda) := \delta_{\mathrm{OP1}}(x_\lambda), \qquad d_5(\lambda) := \delta_{\mathrm{OP5}}(x_\lambda)$$

be the corresponding visible residual channels. Fix positive scales $A_1, A_5 > 0$ on the observation window and define the normalized traces

$$u_1(\lambda) := \frac{d_1(\lambda)}{A_1}, \qquad u_5(\lambda) := \frac{d_5(\lambda)}{A_5}.$$

Define the complex defect trace

$$Z(\lambda) := u_1(\lambda) + i\, u_5(\lambda).$$

Definition H3.7 (Provisional common-carrier and quadrature diagnostics)**.** *We say that the normalized residues admit a* common-carrier representation *on a window I if there are a phase function $\theta : I \to \mathbb{R}$, an amplitude ratio $\rho > 0$, a phase shift $\varphi \in [-\pi, \pi]$, and error terms e_1, e_5 such that*

$$u_1(\lambda) = \cos\theta(\lambda) + e_1(\lambda), \qquad u_5(\lambda) = \rho\cos\big(\theta(\lambda) + \varphi\big) + e_5(\lambda).$$

The case $\varphi \approx 0$ or $\varphi \approx \pi$ is called the same-phase regime*; the case $\varphi \approx \pm\pi/2$ is called the* quadrature regime.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proposition H3.8 (Provisional one-mode image test)**.** *Assume the OP 1 and OP 5 residues are driven to first order by one shared hidden leakage parameter* $\varepsilon_{\mathrm{shared}}(\lambda)$*, in the sense that on a window* I

$$d_1(\lambda) = a_1\,\varepsilon_{\mathrm{shared}}(\lambda) + r_1(\lambda), \qquad d_5(\lambda) = a_5\,\varepsilon_{\mathrm{shared}}(\lambda) + r_5(\lambda),$$

with fixed nonzero coefficients a_1, a_5 *and small remainders* r_1, r_5*. Then the image of* $\lambda \mapsto (u_1(\lambda), u_5(\lambda))$ *is, up to the normalized remainder terms, one-dimensional in the defect plane. In particular, if the normalized trace does* not *spread over a genuinely two-dimensional region, then OP 1 and OP 5 should first be treated as two readouts of one common carrier rather than as independent end defects.*

Proof. After normalization one has

$$u_1(\lambda) = \alpha_1\,\varepsilon_{\mathrm{shared}}(\lambda) + \tilde{r}_1(\lambda), \qquad u_5(\lambda) = \alpha_5\,\varepsilon_{\mathrm{shared}}(\lambda) + \tilde{r}_5(\lambda)$$

for fixed nonzero constants α_1, α_5. The leading image therefore lies on the line spanned by (α_1, α_5). Any two-dimensional spread comes only from the remainders. Hence a numerically thin image in the defect plane is evidence that both visible channels are reading the same hidden scalar mode. □

Proposition H3.9 (Provisional quadrature criterion)**.** *Assume there are a phase function* θ*, a positive ratio* ρ*, and a small error bound* $\eta_{\mathrm{ph}} > 0$ *such that on a window* I

$$|u_1(\lambda) - \cos\theta(\lambda)| \le \eta_{\mathrm{ph}}, \qquad |u_5(\lambda) - \rho\sin\theta(\lambda)| \le \eta_{\mathrm{ph}}.$$

Then the defect trace $Z(\lambda)$ *lies in an* $O(\eta_{\mathrm{ph}})$*-neighborhood of an ellipse centered at the origin. In particular, OP 1 and OP 5 are not independent in that regime; they are two quadrature readouts of one common carrier.*

Proof. In the ideal case $u_1 = \cos\theta$ and $u_5 = \rho\sin\theta$, the image satisfies

$$u_1^2 + \frac{u_5^2}{\rho^2} = 1,$$

so the trace is an ellipse. The stated perturbation bounds place the observed trace in a tube of width $O(\eta_{\mathrm{ph}})$ around that ellipse. Thus the visible residues are phase-coupled and should be interpreted as quadrature projections of one shared mode. □

Remark H3.10 (Why this is the last useful diagnostic)**.** *This test separates two very different endgames. If the defect trace is line-like or near-elliptic, then OP 1 and OP 5 should be closed through a* shared carrier *argument, not through two isolated closure arguments. If the trace really fills a two-dimensional region even after normalization and controlled perturbation, then the hypothesis of one shared hidden mode is too naive and the remaining residues are genuinely less coupled than they currently appear.*

Concrete HC/MM/J probe family for the shared carrier test

The phase-coupling diagnostic becomes much more informative once one stops speaking about an abstract perturbation family and instead deforms only the suspected HC/MM/J leakage channel. The right first family is therefore not a full matrix refit, but a cross-sector probe that touches exactly the admissible/complementary interface.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition H3.11 (Provisional shared leakage probe family)**.** *Fix an HC-even baseline state $v \in \operatorname{im} C$ and a bounded test matrix K on the active grouped transport layer. Define*

$$\Delta_{\mathrm{shared}} := (\mathrm{id} - C)KC + CK(\mathrm{id} - C)$$

and the one-parameter probe family

$$\mathcal{M}_\lambda := \mathcal{M} + \lambda\, \Delta_{\mathrm{shared}}, \qquad |\lambda| \le \lambda_0.$$

The associated transported state is

$$y_\lambda := \mathcal{M}_\lambda C v.$$

Define the masked bridge and valley probe traces by

$$d_1^{\mathrm{probe}}(\lambda) := L_1\big((\mathrm{id} - C)y_\lambda\big), \qquad d_5^{\mathrm{probe}}(\lambda) := L_5\big((\mathrm{id} - C)y_\lambda\big),$$

where L_1 and L_5 are fixed linear masks extracting the OP 1 bridge-sensitive and OP 5 first-window-sensitive visible directions from the complementary output channel.

Proposition H3.12 (Provisional first-order isolation of the shared leakage sector)**.** *The probe family of Definition H3.11 perturbs the admissible/complementary coupling only. More precisely,*

$$C\,\Delta_{\mathrm{shared}}\,C = 0, \qquad (\mathrm{id} - C)\,\Delta_{\mathrm{shared}}\,(\mathrm{id} - C) = 0,$$

and therefore

$$(\mathrm{id} - C)y_\lambda = (\mathrm{id} - C)\mathcal{M}Cv + \lambda\,(\mathrm{id} - C)KCv.$$

Consequently,

$$d_j^{\mathrm{probe}}(\lambda) = d_j^{\mathrm{probe}}(0) + \lambda\, L_j\big((\mathrm{id} - C)KCv\big), \qquad j \in \{1, 5\}.$$

Proof. Since $C(\mathrm{id} - C) = 0 = (\mathrm{id} - C)C$, one has

$$C\Delta_{\mathrm{shared}}C = C(\mathrm{id} - C)KCC + CCK(\mathrm{id} - C)C = 0,$$

and similarly

$$(\mathrm{id} - C)\Delta_{\mathrm{shared}}(\mathrm{id} - C) = 0.$$

Applying $(\mathrm{id} - C)$ to $y_\lambda = (\mathcal{M} + \lambda\Delta_{\mathrm{shared}})Cv$ and using $Cv = v$ gives

$$(\mathrm{id} - C)y_\lambda = (\mathrm{id} - C)\mathcal{M}Cv + \lambda\,(\mathrm{id} - C)\Delta_{\mathrm{shared}}Cv = (\mathrm{id} - C)\mathcal{M}Cv + \lambda\,(\mathrm{id} - C)KCv.$$

Applying the linear masks L_1, L_5 yields the stated first-order formulas. □

Proposition H3.13 (Provisional concrete one-mode and quadrature tests on the probe family)**.** *Let*

$$a_1 := L_1\big((\mathrm{id} - C)KCv\big), \qquad a_5 := L_5\big((\mathrm{id} - C)KCv\big).$$

Then the probe family has the following first-order diagnostics.

(i) If $(a_1, a_5) \neq (0, 0)$, the image of $\lambda \mapsto (d_1^{\mathrm{probe}}(\lambda), d_5^{\mathrm{probe}}(\lambda))$ is first-order one-dimensional, hence consistent with a shared carrier mode.

(ii) If, in addition, one introduces the reversal-paired probe $\widetilde{K} := JK$ and the corresponding coefficients $(\tilde{a}_1, \tilde{a}_5)$, then a near-orthogonality relation

$$a_1\tilde{a}_1 + a_5\tilde{a}_5 \approx 0$$

is first-order evidence for a quadrature-type coupling between the OP 1 and OP 5 visible readouts.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. Part (i) follows immediately from Proposition H3.12, since both traces are affine functions of the same scalar parameter λ with direction vector (a_1, a_5). For part (ii), the reversal-paired probe produces a second direction vector $(\tilde{a}_1, \tilde{a}_5)$. If these two directions are nearly orthogonal in the defect plane, then the pair of probe responses behaves like two approximate quadrature axes of one shared oscillatory mode rather than two unrelated defects. □

Remark H3.14 (Why this is the right next experiment)**.** *The family $\mathcal{M}_\lambda = \mathcal{M} + \lambda\Delta_{\text{shared}}$ is the cleanest next diagnostic because it does not refit the whole Millennium matrix. It touches only the admissible/complementary interface where a common OP 1/OP 5 remainder would naturally live. If even this targeted family fails to produce either a one-mode line or a quadrature pair in the defect plane, then the shared-carrier hypothesis must be weakened. If it succeeds, then the next theorem should be a shared-channel closure statement rather than another isolated local endgame.*

First-order rank and quadrature invariants of the probe family

The concrete probe family admits an even sharper reading than the earlier qualitative line-versus-ellipse distinction. Since both OP 1 and OP 5 are now tested by two explicit probe directions, one can read off from a single 2×2 response matrix whether the visible pair behaves like one masked mode, like a quadrature pair, or like a genuinely two-dimensional remainder.

Definition H3.15 (Provisional probe response matrix and normalized invariants)**.** *With the notation of Proposition H3.13, define the probe response matrix*

$$S(K) := \begin{pmatrix} a_1 & a_5 \\ \tilde{a}_1 & \tilde{a}_5 \end{pmatrix},$$

where (a_1, a_5) comes from K and $(\tilde{a}_1, \tilde{a}_5)$ from the reversal-paired probe $\widetilde{K} = JK$. Assume both row vectors are nonzero and set

$$n := \sqrt{a_1^2 + a_5^2}, \qquad \tilde{n} := \sqrt{\tilde{a}_1^2 + \tilde{a}_5^2}.$$

Define the normalized one-mode defect and the normalized quadrature defect by

$$\mathfrak{d}_{\parallel}(K) := \frac{|a_1\tilde{a}_5 - a_5\tilde{a}_1|}{n\tilde{n}}, \qquad \mathfrak{d}_{\perp}(K) := \frac{|a_1\tilde{a}_1 + a_5\tilde{a}_5|}{n\tilde{n}}.$$

Proposition H3.16 (Provisional classification of the shared probe response)**.** *Under Definition H3.15, the following first-order interpretations hold.*

(i) If $\mathfrak{d}_{\parallel}(K) \ll 1$, then the two probe directions are nearly collinear, hence the visible OP 1/OP 5 response is first-order compatible with one masked shared mode.

(ii) If $\mathfrak{d}_{\perp}(K) \ll 1$ and $\mathfrak{d}_{\parallel}(K)$ is bounded away from 0, then the two probe directions are nearly orthogonal, hence the visible response is first-order compatible with a quadrature pair of one shared carrier.

(iii) If both $\mathfrak{d}_{\parallel}(K)$ and $\mathfrak{d}_{\perp}(K)$ stay away from 0 on a stable test class of admissible probes K, then the shared-carrier hypothesis is weakened: the remaining OP 1/OP 5 defect is genuinely at least two-directional at first order.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. The quantity $|a_1\tilde{a}_5 - a_5\tilde{a}_1|$ is the absolute determinant of the two response vectors, normalized by their lengths. It therefore measures the sine of the angle between them, and smallness means near-collinearity. Likewise, $|a_1\tilde{a}_1 + a_5\tilde{a}_5|$ is the absolute dot product, normalized by the same lengths, so its smallness means near-orthogonality. The three conclusions are therefore the standard geometric alternatives for two nonzero vectors in the visible defect plane. □

Corollary H3.17 (Provisional first-order phase picture from the probe invariants)**.** *Suppose there is a stable admissible test class of probes K for which either*

$$\mathfrak{d}_{\parallel}(K) \le \eta_{\parallel}$$

or

$$\mathfrak{d}_{\perp}(K) \le \eta_{\perp}$$

with small tolerances. Then the remaining OP 1 and OP 5 residues should not be pursued as unrelated endgames. In the first case they are first-order images of one shared hidden mode, and in the second case they are first-order quadrature readouts of one shared hidden carrier.

Proof. This is the direct structural reading of Proposition H3.16. The point is not that the exact finite-amplitude endgame is already closed, but that the visible first-order geometry already decides whether a common hidden carrier is plausible or not. □

Provisional diagnostic takeaway. The matrix test is now decision-grade. Instead of asking vaguely whether OP 1 and OP 5 "feel related," one computes the normalized determinant defect $\mathfrak{d}_{\parallel}$ and the normalized dot-product defect $\mathfrak{d}_{\perp}$ on the HC/MM/J probe class. Small $\mathfrak{d}_{\parallel}$ means one shared line; small $\mathfrak{d}_{\perp}$ means one shared carrier in quadrature.

Canonical low-rank probe class and the first verdict on one-mode versus quadrature

The abstract probe family is useful, but it still leaves too much freedom in the choice of K. The cleanest next step is therefore to test the suspected HC/MM/J leakage by the smallest nontrivial cross-sector probes possible. This produces a genuine structural verdict: does the most canonical leakage class already support quadrature, or does it force a one-mode picture at first order?

Definition H3.18 (Provisional canonical low-rank leakage probes)**.** *Let u be a unit admissible vector and w a unit complementary vector such that*

$$Cu = u, \qquad Cw = 0, \qquad Ju = u, \qquad Jw = -w, \qquad \langle u, w\rangle = 0.$$

Define the associated rank-two cross-sector probe by

$$K_{u,w}x := \langle u, x\rangle w + \langle w, x\rangle u.$$

This is the smallest symmetric probe that exchanges one chosen admissible direction with one chosen complementary direction and does nothing else.

Proposition H3.19 (Provisional exact support of the canonical low-rank probes)**.** *For every probe $K_{u,w}$ from Definition H3.18,*

$$CK_{u,w}C = 0, \qquad (\mathrm{id} - C)K_{u,w}(\mathrm{id} - C) = 0, \qquad \Delta_{\mathrm{shared}}(K_{u,w}) = K_{u,w}.$$

Hence $K_{u,w}$ is already a pure first-order admissible/complementary leakage probe and needs no further projection cleanup.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. Since $Cu = u$ and $Cw = 0$, the operator $K_{u,w}$ sends u into the complementary direction w and w back into the admissible direction u, while all purely admissible-to-admissible and purely complementary-to-complementary blocks vanish. This gives the first two identities. The defining formula

$$\Delta_{\text{shared}} = (\mathrm{id} - C)KC + CK(\mathrm{id} - C)$$

then reproduces $K_{u,w}$ itself, because the only nonzero parts of $K_{u,w}$ are already the two cross-sector pieces. □

Proposition H3.20 (Provisional response geometry of the canonical low-rank class)**.** *Fix a baseline admissible state v with $Cv = v$, and define the probe coefficients for $K_{u,w}$ and the reversal-paired probe $\widetilde{K}_{u,w} := JK_{u,w}$ by*

$$a_j(u,w) := L_j\big((\mathrm{id} - C)K_{u,w}Cv\big), \qquad \widetilde{a}_j(u,w) := L_j\big((\mathrm{id} - C)\widetilde{K}_{u,w}Cv\big), \qquad j \in \{1,5\}.$$

Then

$$a_j(u,w) = \langle u, v\rangle\, L_j(w), \qquad \widetilde{a}_j(u,w) = -\langle u, v\rangle\, L_j(w) = -a_j(u,w).$$

In particular, whenever the response vector is nonzero,

$$S(K_{u,w}) = \begin{pmatrix} a_1 & a_5 \\ -a_1 & -a_5 \end{pmatrix},$$

and therefore

$$\mathfrak{d}_{\parallel}(K_{u,w}) = 0, \qquad \mathfrak{d}_{\perp}(K_{u,w}) = 1.$$

Proof. Because $Cv = v$ and $K_{u,w}$ contains exactly one admissible-to-complementary leg,

$$(\mathrm{id} - C)K_{u,w}Cv = (\mathrm{id} - C)K_{u,w}v = \langle u, v\rangle w.$$

Applying L_j gives the formula for a_j. Since $Ju = u$ and $Jw = -w$,

$$(\mathrm{id} - C)JK_{u,w}Cv = J(\mathrm{id} - C)K_{u,w}Cv = -\langle u, v\rangle w,$$

which yields $\widetilde{a}_j = -a_j$. The displayed matrix form follows immediately. Its two rows are exactly antiparallel, so the normalized determinant defect vanishes and the normalized dot-product defect is maximal. □

Corollary H3.21 (Provisional first verdict from the canonical probe class)**.** *The canonical low-rank HC/MM/J leakage probes are intrinsically first-order one-mode probes: they can only detect a shared line and never a genuine quadrature pair. Consequently, if a stable quadrature picture for OP 1 and OP 5 is later observed, it cannot come from primitive rank-two cross-sector leakage alone; it must arise from a richer multi-directional or higher-order transport/readout effect.*

Proof. This is the direct reading of Proposition H3.20. The canonical low-rank class forces $\mathfrak{d}_{\parallel} = 0$ and $\mathfrak{d}_{\perp} = 1$, so the first-order geometry is always collinear and never orthogonal. □

Provisional structural takeaway. This is already a nontrivial verdict. The smallest honest HC/MM/J leakage class does *not* support a primitive sinus/cosinus split between OP 1 and OP 5. At first order it only supports one shared line. If the two residues later turn out to behave like quadrature data, then that phase shift is emergent and not the most primitive leakage mode.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Minimal two-directional leakage plane and the first place where quadrature can appear

The canonical rank-two probes already showed that primitive cross-sector leakage is first-order one-mode only. The next structural question is therefore very sharp: what is the smallest extension of the leakage class in which a genuine sinus/cosinus picture can appear at all? The answer is: not a single exchange direction, but a *two-directional leakage plane* equipped with an internal quarter-turn.

Definition H3.22 (Provisional minimal two-directional leakage plane)**.** *Let u_1, u_2 be orthonormal admissible vectors and let w_1, w_2 be orthonormal complementary vectors such that*

$$Cu_i = u_i, \qquad Cw_i = 0, \qquad Ju_i = u_i, \qquad Jw_i = -w_i, \qquad i \in \{1, 2\}.$$

Define the paired rank-four cross-sector probes

$$K_{\parallel} := K_{u_1,w_1} + K_{u_2,w_2},$$

and

$$K_{\perp} := K_{u_1,w_2} - K_{u_2,w_1}.$$

Equivalently, if R_W denotes the quarter-turn on the complementary plane $W := \operatorname{span}\{w_1, w_2\}$ given by

$$R_W w_1 = w_2, \qquad R_W w_2 = -w_1,$$

then $K_{\perp}$ is the quarter-turn-dressed partner of $K_{\parallel}$ on the complementary output side.

Proposition H3.23 (Provisional exact support of the minimal two-directional class)**.** *Both operators $K_{\parallel}$ and $K_{\perp}$ from Definition H3.22 are pure first-order cross-sector probes:*

$$CK_{\sharp}C = 0, \qquad (\mathrm{id} - C)K_{\sharp}(\mathrm{id} - C) = 0, \qquad \Delta_{\mathrm{shared}}(K_{\sharp}) = K_{\sharp},$$

for $\sharp \in \{\parallel, \perp\}$.

Proof. Each summand K_{u_i,w_j} exchanges exactly one admissible direction with one complementary direction and has no pure admissible or pure complementary block. The same is therefore true for both linear combinations $K_{\parallel}$ and $K_{\perp}$. Hence the support identities follow exactly as in Proposition H3.19. □

Proposition H3.24 (Provisional complementary output geometry of the minimal two-directional class)**.** *Fix an admissible baseline state v with $Cv = v$, and write*

$$a := \langle u_1, v\rangle, \qquad b := \langle u_2, v\rangle.$$

Then

$$(\mathrm{id} - C)K_{\parallel}Cv = aw_1 + bw_2,$$

and

$$(\mathrm{id} - C)K_{\perp}Cv = aw_2 - bw_1 = R_W\big((\mathrm{id} - C)K_{\parallel}Cv\big).$$

In particular,

$$\big\langle (\mathrm{id} - C)K_{\parallel}Cv, (\mathrm{id} - C)K_{\perp}Cv \big\rangle = 0,$$

and

$$\big\|(\mathrm{id} - C)K_{\parallel}Cv\big\| = \big\|(\mathrm{id} - C)K_{\perp}Cv\big\| = \sqrt{a^2 + b^2}.$$

Thus the smallest leakage extension capable of supporting quadrature is a two-directional complementary carrier plane, not a single primitive leakage line.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. Since $Cv = v$, only the admissible-to-complementary legs contribute. For $K_{\parallel}$ one gets

$$(\mathrm{id} - C)K_{\parallel}Cv = aw_1 + bw_2.$$

Likewise,

$$(\mathrm{id} - C)K_{\perp}Cv = aw_2 - bw_1.$$

The displayed quarter-turn relation follows from the definition of R_W. Orthogonality and equal norm are then immediate from the orthonormality of w_1, w_2. □

Corollary H3.25 (Provisional first place where quadrature becomes structurally possible)**.** *Quadrature between OP 1 and OP 5 cannot arise from the primitive canonical rank-two leakage class alone. The first structurally honest place where it can appear is the paired two-directional class $\{K_{\parallel}, K_{\perp}\}$, where the complementary output already carries an internal quarter-turn. Consequently, any later sinus/cosinus reading of OP 1 and OP 5 should be interpreted as a dressed two-directional leakage-plane effect rather than as a primitive one-directional HC/MM/J defect.*

Proof. Corollary H3.21 showed that the primitive rank-two class is forced to be first-order collinear. Proposition H3.24 shows that the paired rank-four class is the smallest extension whose complementary outputs are already quarter-turned relative to one another. Hence this is the first structurally admissible location of quadrature. □

Provisional diagnostic takeaway. The structural picture is now sharper. Primitive leakage gives one shared line. Quadrature needs at least a two-directional leakage plane together with an internal quarter-turn on the complementary carrier. So if OP 1 and OP 5 later behave like cosine/sine data, the right search target is no longer a single hidden defect entry but a minimal paired leakage plane with quarter-turn dressing.

Theorem-level promotion for OP 1

The OP 1 epsilon-bridge tool can be upgraded from routing language to a genuine local fixed-point estimate. The key point is that, once one is inside a contractive bridge tube, the normalization defect itself already controls the distance to the distinguished bridge fixed point. This is exactly the kind of "determine ε once, then prove non-escape" move that the present structure seems to support.

Definition H3.26 (Provisional bridge normalization defect)**.** *Let*

$$T_{\gamma_{\mathrm{br}}}$$

be the bridge self-consistency map introduced in Section 3.8.16, and let

$$\beta_{\mathrm{br}} := \beta(\gamma_{\mathrm{br}})$$

be its distinguished fixed point at the local bridge root from Section 3.8.16. For any positive amplitude b, define the bridge normalization defect

$$\delta_{\mathrm{br}}(b) := \left|T_{\gamma_{\mathrm{br}}}(b) - b\right|.$$

Proposition H3.27 (Provisional residual-to-fixed-point estimate for the bridge tube)**.** *Assume there are numbers $\varepsilon_{\mathrm{br}} > 0$ and $L_{\mathrm{br}} < 1$ such that*

$$\sup_{|b-\beta_{\mathrm{br}}|\le\varepsilon_{\mathrm{br}}} \left|\partial_b T_{\gamma_{\mathrm{br}}}(b)\right| \le L_{\mathrm{br}}.$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Then every b with $|b-\beta_{\mathrm{br}}| \le \varepsilon_{\mathrm{br}}$ satisfies

$$|b-\beta_{\mathrm{br}}| \le \frac{\delta_{\mathrm{br}}(b)}{1-L_{\mathrm{br}}}.$$

In particular, sufficiently small bridge normalization defect forces entry into an arbitrarily small local tube around the distinguished bridge point.

Proof. Since β_{br} is a fixed point of $T_{\gamma_{\mathrm{br}}}$, one has

$$T_{\gamma_{\mathrm{br}}}(\beta_{\mathrm{br}}) = \beta_{\mathrm{br}}.$$

Hence for any b in the tube,

$$|b-\beta_{\mathrm{br}}| \le |b-T_{\gamma_{\mathrm{br}}}(b)| + |T_{\gamma_{\mathrm{br}}}(b) - T_{\gamma_{\mathrm{br}}}(\beta_{\mathrm{br}})|.$$

The first term is exactly $\delta_{\mathrm{br}}(b)$. For the second term, the mean-value estimate inside the tube gives

$$|T_{\gamma_{\mathrm{br}}}(b) - T_{\gamma_{\mathrm{br}}}(\beta_{\mathrm{br}})| \le L_{\mathrm{br}}|b-\beta_{\mathrm{br}}|.$$

Therefore

$$|b-\beta_{\mathrm{br}}| \le \delta_{\mathrm{br}}(b) + L_{\mathrm{br}}|b-\beta_{\mathrm{br}}|,$$

so

$$(1-L_{\mathrm{br}})|b-\beta_{\mathrm{br}}| \le \delta_{\mathrm{br}}(b).$$

Dividing by $1-L_{\mathrm{br}} > 0$ proves the claim. □

Theorem H3.28 (Provisional OP 1 epsilon-trapped bridge normalization theorem)**.** *Assume the local bridge tube of Proposition H3.27. Let $b_{\mathrm{can}} > 0$ be a final canonical normalized readout with*

$$|b_{\mathrm{can}} - \beta_{\mathrm{br}}| \le \varepsilon_{\mathrm{br}}.$$

If for some $\varepsilon_{\mathrm{ent}} \in (0, \varepsilon_{\mathrm{br}}]$ one has

$$\delta_{\mathrm{br}}(b_{\mathrm{can}}) \le (1-L_{\mathrm{br}})\,\varepsilon_{\mathrm{ent}},$$

then

$$|b_{\mathrm{can}} - \beta_{\mathrm{br}}| \le \varepsilon_{\mathrm{ent}}.$$

Moreover, the whole forward bridge orbit remains trapped inside the same local bridge tube and converges to β_{br}:

$$T^n_{\gamma_{\mathrm{br}}}(b_{\mathrm{can}}) \to \beta_{\mathrm{br}}.$$

Thus, once the canonical normalization is shown to have sufficiently small bridge defect inside the contractive local tube, OP 1 reduces to a non-escape statement rather than a second normalization search.

Proof. The residual bound and Proposition H3.27 give

$$|b_{\mathrm{can}} - \beta_{\mathrm{br}}| \le \frac{\delta_{\mathrm{br}}(b_{\mathrm{can}})}{1-L_{\mathrm{br}}} \le \varepsilon_{\mathrm{ent}}.$$

This proves entry into the smaller $\varepsilon_{\mathrm{ent}}$-tube. Since β_{br} is a fixed point,

$$|T_{\gamma_{\mathrm{br}}}(b) - \beta_{\mathrm{br}}| = |T_{\gamma_{\mathrm{br}}}(b) - T_{\gamma_{\mathrm{br}}}(\beta_{\mathrm{br}})| \le L_{\mathrm{br}}|b-\beta_{\mathrm{br}}|$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

for every b in the local bridge tube. Hence if $|b - \beta_{\mathrm{br}}| \le \varepsilon_{\mathrm{ent}}$, then

$$|T_{\gamma_{\mathrm{br}}}(b) - \beta_{\mathrm{br}}| \le L_{\mathrm{br}}\varepsilon_{\mathrm{ent}} < \varepsilon_{\mathrm{ent}}.$$

So the smaller tube is forward invariant and, in particular, remains inside the original contractive tube. Standard fixed-point iteration for a contraction then gives

$$T^n_{\gamma_{\mathrm{br}}}(b_{\mathrm{can}}) \to \beta_{\mathrm{br}}.$$

This is exactly the announced non-escape statement. □

Corollary H3.29 (Provisional exact-closure criterion for canonical normalization)**.** *Under the hypotheses of Theorem H3.28, if the canonical normalization satisfies*

$$\delta_{\mathrm{br}}(b_{\mathrm{can}}) = 0,$$

then

$$b_{\mathrm{can}} = \beta_{\mathrm{br}}.$$

Equivalently, inside the contractive bridge tube there is no second local canonical normalization branch.

Proof. If $\delta_{\mathrm{br}}(b_{\mathrm{can}}) = 0$, then Proposition H3.27 gives

$$|b_{\mathrm{can}} - \beta_{\mathrm{br}}| = 0.$$

Hence $b_{\mathrm{can}} = \beta_{\mathrm{br}}$. □

Provisional OP 1 status after theorem-level promotion. The proof above is deliberately honest about what is and is not finished. The local bridge theorem is now real: once a canonical normalization final is inside the contractive tube and its bridge defect is small enough, local non-escape and uniqueness follow. What still remains open is the structural *entry mechanism*: why the canonical discrete correction should land inside that tube in the first place. So the remaining OP 1 burden is no longer local stability; it is the transport-side justification of tube entry.

Remark H3.30 (Provisional OP 1 reading after the OP transport consolidation)**.** *In the present internal line, the epsilon-trapped bridge theorem is best read as the OP 1-side non-escape precursor of the admissible-branch architecture. Its role is no longer to reopen global OP closure, but to isolate the remaining entry question: why the canonical normalization should lie on a transport-admissible branch whose first bridge window already remains trapped in the local contractive tube.*

Rounded-entry promotion. The remaining OP 1 burden can be sharpened further. The current matter-frame value is not merely close to the distinguished bridge root in an abstract scalar sense; it is already separated from that root by only a tiny readout-point displacement in μ. Since the readout amplitude $\beta(\mu)$ is smooth on the physical window, this immediately produces a concrete bridge-entry estimate. Combined with the contractive bridge map, the rounded readout itself already satisfies a tiny local bridge defect. This turns the live OP 1 burden into a very narrow identification problem: justify that the canonical normalization is indeed the standard rounded matter-frame readout.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition H3.31 (Provisional rounded bridge-entry gap)**.** *Let*

$$\Delta_{\mathrm{br}}^{(\mu)} := |\gamma_L - \gamma_{\mathrm{br}}|, \qquad b_{\mathrm{MF}} := \beta(\gamma_L), \qquad \beta_{\mathrm{br}} := \beta(\gamma_{\mathrm{br}}).$$

Thus $\Delta_{\mathrm{br}}^{(\mu)}$ *is the rounded readout-point displacement from the distinguished bridge root, while* b_{MF} *is the rounded matter-frame readout amplitude currently used in the Companion.*

Proposition H3.32 (Provisional transport-to-bridge entry estimate)**.** *On the physical window* $I_{\mathrm{phys}} = [1.85, 1.90]$*, the readout amplitude*

$$\beta(\mu) = \sqrt{\frac{\mu - 1}{\mu + 1}}$$

satisfies

$$\beta'(\mu) = \frac{1}{\beta(\mu)(\mu + 1)^2}.$$

Moreover, on I_{phys} *one has the numerical bound*

$$\beta'(\mu) < 0.226.$$

Hence

$$|b_{\mathrm{MF}} - \beta_{\mathrm{br}}| \le 0.226\, \Delta_{\mathrm{br}}^{(\mu)}.$$

Using the distinguished-root estimate from Section 3.8.12, this gives

$$|b_{\mathrm{MF}} - \beta_{\mathrm{br}}| < 7.24 \times 10^{-9}.$$

Proof. Differentiating

$$\beta(\mu) = \left(\frac{\mu - 1}{\mu + 1}\right)^{1/2}$$

gives

$$\beta'(\mu) = \frac{1}{2}\left(\frac{\mu - 1}{\mu + 1}\right)^{-1/2} \frac{2}{(\mu + 1)^2} = \frac{1}{\beta(\mu)(\mu + 1)^2}.$$

Direct evaluation on the endpoints of I_{phys} gives

$$\beta'(1.85) \approx 0.22544, \qquad \beta'(1.90) \approx 0.21344,$$

and direct evaluation on the physical window gives the uniform bound $\beta'(\mu) < 0.226$. The mean-value theorem therefore yields

$$|b_{\mathrm{MF}} - \beta_{\mathrm{br}}| = |\beta(\gamma_L) - \beta(\gamma_{\mathrm{br}})| \le 0.226\, |\gamma_L - \gamma_{\mathrm{br}}| = 0.226\, \Delta_{\mathrm{br}}^{(\mu)}.$$

By the distinguished-root estimate of Section 3.8.12,

$$\Delta_{\mathrm{br}}^{(\mu)} \approx 3.200357 \times 10^{-8},$$

so the final numerical estimate follows. □

Proposition H3.33 (Provisional rounded-readout bridge-defect estimate)**.** *Assume the local bridge tube of Proposition H3.27. Then every* b *in that tube satisfies*

$$\delta_{\mathrm{br}}(b) \le (1 + L_{\mathrm{br}})|b - \beta_{\mathrm{br}}|.$$

In particular, if b_{MF} *lies in the local bridge tube, then*

$$\delta_{\mathrm{br}}(b_{\mathrm{MF}}) \le (1 + L_{\mathrm{br}})\, 0.226\, \Delta_{\mathrm{br}}^{(\mu)}.$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. Since $T_{\gamma_{\mathrm{br}}}(\beta_{\mathrm{br}}) = \beta_{\mathrm{br}}$, one has

$$\delta_{\mathrm{br}}(b) = |T_{\gamma_{\mathrm{br}}}(b) - b| \le |T_{\gamma_{\mathrm{br}}}(b) - T_{\gamma_{\mathrm{br}}}(\beta_{\mathrm{br}})| + |\beta_{\mathrm{br}} - b|.$$

Inside the bridge tube, the contraction bound gives

$$|T_{\gamma_{\mathrm{br}}}(b) - T_{\gamma_{\mathrm{br}}}(\beta_{\mathrm{br}})| \le L_{\mathrm{br}}|b - \beta_{\mathrm{br}}|,$$

so

$$\delta_{\mathrm{br}}(b) \le (1 + L_{\mathrm{br}})|b - \beta_{\mathrm{br}}|.$$

Applying Proposition H3.32 to $b = b_{\mathrm{MF}}$ yields the final estimate. □

Theorem H3.34 (Provisional rounded-readout entry theorem for OP 1)**.** *Assume the local bridge tube of Proposition H3.27 and suppose that*

$$0.226\,\Delta_{\mathrm{br}}^{(\mu)} \le \varepsilon_{\mathrm{br}}.$$

Let $b_{\mathrm{MF}} = \beta(\gamma_L)$*. If*

$$\varepsilon_{\mathrm{ent}} \ge \frac{1 + L_{\mathrm{br}}}{1 - L_{\mathrm{br}}} \cdot 0.226\,\Delta_{\mathrm{br}}^{(\mu)},$$

then

$$|b_{\mathrm{MF}} - \beta_{\mathrm{br}}| \le \varepsilon_{\mathrm{ent}}$$

and the forward bridge orbit

$$T_{\gamma_{\mathrm{br}}}^{n}(b_{\mathrm{MF}})$$

remains trapped in the local bridge tube and converges to β_{br}*. For the current numerical bridge data*

$$\Delta_{\mathrm{br}}^{(\mu)} \approx 3.200357 \times 10^{-8}, \qquad |\partial_b T_{\gamma_{\mathrm{br}}}(\beta_{\mathrm{br}})| \approx 0.00311,$$

it is already enough that

$$\varepsilon_{\mathrm{ent}} \gtrsim 7.28 \times 10^{-9}.$$

Proof. Proposition H3.32 gives

$$|b_{\mathrm{MF}} - \beta_{\mathrm{br}}| \le 0.226\,\Delta_{\mathrm{br}}^{(\mu)},$$

so the first displayed assumption ensures that b_{MF} lies in the local bridge tube. Proposition H3.33 then yields

$$\delta_{\mathrm{br}}(b_{\mathrm{MF}}) \le (1 + L_{\mathrm{br}})\,0.226\,\Delta_{\mathrm{br}}^{(\mu)}.$$

By the lower bound on $\varepsilon_{\mathrm{ent}}$, this implies

$$\delta_{\mathrm{br}}(b_{\mathrm{MF}}) \le (1 - L_{\mathrm{br}})\,\varepsilon_{\mathrm{ent}}.$$

Theorem H3.28 therefore applies with $b_{\mathrm{can}} = b_{\mathrm{MF}}$, proving entry into the $\varepsilon_{\mathrm{ent}}$-tube and convergence of the forward orbit to β_{br}. The final numerical threshold is the explicit evaluation of

$$\frac{1 + L_{\mathrm{br}}}{1 - L_{\mathrm{br}}} \cdot 0.226\,\Delta_{\mathrm{br}}^{(\mu)}$$

at $\Delta_{\mathrm{br}}^{(\mu)} \approx 3.200357 \times 10^{-8}$ and $L_{\mathrm{br}} \approx 0.00311$. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Corollary H3.35 (Provisional narrowed live burden for OP 1 after rounded entry)**.** *Under the hypotheses of Theorem H3.34, the current rounded matter-frame readout already enters the contractive bridge tube and converges to the distinguished bridge fixed point. Therefore the live OP 1 burden is no longer broad bridge entry. It has narrowed to the structural identification problem*

$$b_{\mathrm{can}} = b_{\mathrm{MF}},$$

i.e. justify that the canonical normalization readout is exactly the standard rounded matter-frame readout already used in the Companion.

Proof. Theorem H3.34 proves entry and convergence for b_{MF}. Hence the only remaining gap is whether the canonical normalization final should indeed be identified with this rounded matter-frame readout. No second local bridge-search mechanism remains. □

Provisional OP 1 status after rounded-entry promotion. The local OP 1 story is now much tighter. The bridge point γ_{br} and the standard rounded matter-frame value $\gamma_L = 1.864$ differ by only 3.2×10^{-8}, and this already forces the corresponding readout amplitude into any reasonable contractive bridge tube. So the bridge-entry problem is no longer genuinely numerical. What remains is the structural identification step: explain why the canonical normalization should be read as the standard rounded matter-frame readout, or equivalently why no second transport-side normalization survives after the bridge forcing, renormalization, and contraction lines.

Admissible-readout uniqueness promotion. The rounded-entry theorem reduced the live OP 1 burden to the structural identification

$$b_{\mathrm{can}} = b_{\mathrm{MF}}.$$

The present document already contains almost all of this argument in implicit form: transport fixes the branch first, readout stays downstream, grouped-layer transport introduces no ontology drift, and document normalization forbids treating local readout blocks as second foundational centers. The missing step is to promote these points to an explicit uniqueness theorem for the positive scalar readout on the fixed bridge branch.

Theorem H3.36 (Provisional admissible-readout uniqueness theorem for OP 1)**.** *Assume the transport branch has already been fixed at the current matter-frame point*

$$\mu = \gamma_L,$$

and let

$$b_{\mathrm{MF}} := \beta(\gamma_L) = \sqrt{\frac{\gamma_L - 1}{\gamma_L + 1}}.$$

Let $b_{\mathrm{can}} > 0$ be a canonical normalization readout for OP 1 such that:

(i) *b_{can} is obtained only after branch selection, as a downstream admissible readout on the already fixed transport branch;*

(ii) *the readout law opens no second primitive continuum channel;*

(iii) *grouped-layer transport produces no ontology drift, so any alternative local readout regime remains an organizational realization of the same admissible structural body.*

Then

$$b_{\mathrm{can}} = b_{\mathrm{MF}}.$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. By the transport-first tool in the current tool layer, the continuum branch is fixed before visible readout is interpreted. At the current matter-frame point $\mu = \gamma_L$, the standard visible scalar on that branch is exactly

$$b_{\mathrm{MF}} = \beta(\gamma_L).$$

By the readout-after-transport tool, every further canonical readout must remain downstream from this branch selection and therefore cannot reopen the continuum-side selection problem. If one had $b_{\mathrm{can}} \neq b_{\mathrm{MF}}$ while both are positive canonical normalizations on the same already fixed branch, then the second scalar would constitute a second primitive continuum-side normalization channel. This contradicts hypothesis (ii).

Hypothesis (iii) excludes the only remaining escape route: reinterpreting the second scalar as belonging to a genuinely different admissible body. By Proposition H4.397, admissible transport across grouped layers does not create ontological multiplicity, and Remark H4.393 says that local readout/integration blocks are organizational regimes rather than second foundational centers. Hence the canonical normalization and the standard matter-frame readout are two organizational readings of one and the same admissible body on one already fixed branch. Since no second primitive channel is allowed, they must coincide. Therefore

$$b_{\mathrm{can}} = b_{\mathrm{MF}}.$$

□

Corollary H3.37 (Provisional OP 1 structural promotion after readout uniqueness)**.** *Under the hypotheses of Theorem H3.36 and Theorem H3.34, the canonical normalization readout itself already enters the local contractive bridge tube and converges to the distinguished bridge fixed point:*

$$T_{\gamma_{\mathrm{br}}}^{n}(b_{\mathrm{can}}) \longrightarrow \beta_{\mathrm{br}}.$$

In particular, the former identification burden $b_{\mathrm{can}} = b_{\mathrm{MF}}$ disappears. The live OP 1 burden is reduced to the explicit local bridge residue

$$\delta_{\mathrm{br}}(b_{\mathrm{MF}}) = \left|T_{\gamma_{\mathrm{br}}}(b_{\mathrm{MF}}) - b_{\mathrm{MF}}\right|,$$

and exact OP 1 closure follows as soon as this already isolated residue vanishes.

Proof. Theorem H3.36 gives

$$b_{\mathrm{can}} = b_{\mathrm{MF}}.$$

Applying Theorem H3.34 to the right-hand side yields entry of b_{MF} into the local bridge tube and convergence of its forward orbit to β_{br}. Substituting b_{can} for b_{MF} proves the same statement for the canonical normalization. Therefore the identification burden is gone, and only the explicit local bridge residue at the unique admissible readout remains. If that residue is zero, Corollary H3.29 yields exact local bridge closure. □

Provisional OP 1 status after uniqueness promotion. At this point OP 1 has become much narrower than a generic matching problem. The transport branch is fixed, the rounded matter-frame readout already enters the contractive bridge tube at roughly the 10^{-9}-level, and the admissible-readout uniqueness theorem removes the last structural ambiguity between canonical normalization and standard readout. So OP 1 no longer hangs at branch search, root search, or local entry. It hangs only at the exact vanishing of one already isolated local scalar defect:

$$\delta_{\mathrm{br}}(b_{\mathrm{MF}}) = 0.$$

In plain language: the proof burden has collapsed to showing that the unique admissible readout is not merely bridge-close, but exactly bridge-stable.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

archival stage Closure–Projection Synthesis Core

Remark H3.38 (Normalization note for archival stage)**.** *In the present release, the mathematical content of this synthesis layer is subsumed by the OP transport theorem and the OP3 residual/shell interface. The block is retained as historical compression logic and interpretive guidance. This synthesis block is therefore kept as a historical transport precursor: its closure–projection synthesis is now carried proof-theoretically by Phase II-D, Phase II-E, and Phase II-F, and here it should be read as an earlier integrative summary rather than as an independent axiomatic center.*

This release adds a further synthesis layer between the closure blocks already introduced and the inherited structural core of the monolith. Its purpose is to state, in one explicit place, how closure statements are transported across the probe sector, how projection claims inherit their admissibility status from the closure architecture, and how appendix tables are to be used as reversible proof interfaces rather than as detached summaries.

1. Closure-to-projection transport discipline

The central operational rule of archival stage is that no projection statement may be read as primary if its admissibility has not first been transported through the closure chain. In other words, the order of explanation remains

$$\begin{aligned}&\text{local structure} \Longrightarrow \text{HC} \Longrightarrow \text{closure}\\ &\Longrightarrow \text{probe-compatible reduction} \Longrightarrow \text{projection readout.}\end{aligned}$$

A projection is therefore not an independent object but a compressed exposure of closure-stable content under a chosen observational or effective frame. This rule keeps the paper aligned with the structure-first programme and prevents later phenomenological language from silently replacing the closure mechanism that generated it.

Synthesis Proposition A (closure status survives transport). If a statement has been established in the closure regime and is moved to a probe-compatible or projective representation by a transport that preserves the common body, then its structural status is not lost. What changes is the mode of appearance, not the underlying admissibility class.

Proof sketch. The common body is fixed by the ground condition and stabilized by HC. Any transport that preserves this body may change coordinates, phase labels, or sector language, but it does not create a new ontology. Hence closure status is inherited through the transport.

2. Probe reduction as a controlled interface

The probe sector is not only a search device; in archival stage it is also an interface theorem. Two probes moving against one another, each carrying triolic organization and coupled to an orthogonal third direction, define a reduced test geometry in which admissible transitions can be filtered before full projection. This means that the probe sector is a disciplined compression stage: it narrows final structure while still remaining within closure control.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Synthesis Proposition B (probe reduction preserves closure discipline). Whenever the probe sector is constructed from triolic substructures under the standing orthogonality and HC rules, the reduced search geometry cannot produce admissible outputs that violate closure. It may discard finals, rank them, or expose only a subset of modes, but it cannot certify a structurally illegal sector.

Proof sketch. The probe geometry is not external to the closure architecture. It is built from substructures already governed by HC and orthogonality. Therefore any reduction performed inside the probe channel is a reduction of representation, not a release from confinement.

3. Projection tables as reversible proof interfaces

The appendix tables introduced in the previous versions are to be read neither as cosmetic summaries nor as final stand-alone derivations. Their exact role is reversible: each table entry points upward to the closure and lemma chain from which it is licensed, and downward to the sector language in which it may be used. In this sense, a table row functions as a compact proof interface.

Synthesis Principle (table row admissibility). A table row is admissible only if one can move in both directions:

(a) upward, from the row to the closure/lemma chain that stabilizes it; and

(b) downward, from the row to a projective or phenomenological reading that does not exceed the transported status of the original statement.

If either direction is unavailable, the row is only heuristic and must be marked as such.

4. Micro-to-macro synthesis route

The micro-to-macro route is now stated in a more disciplined form. The local triolic unit is not yet a macroscopic law, but it already fixes the admissible coupling pattern. HC then stabilizes multiple appearances of the same body, closure confines the admissible dynamics, the probe sector performs controlled reduction, and only then do effective macroscopic equations appear as readouts. The macro level is therefore a closure-saturated exposure of the micro level, not a rival explanatory layer.

Synthesis Proposition C (macro equations are closure-saturated readouts). Under the standing assumptions of the monolith, any effective macroscopic relation used later in the paper must be readable as a closure-saturated image of the local structural chain. Hence macro equations are legitimate exactly to the degree that the closure-to-projection transport remains traceable.

5. Appendix back-reference discipline, strengthened

The appendix is now explicitly promoted to a back-reference layer. This means that every compressed proof move used in the running text should be recoverable through one of three routes: a lemma reference, a tool reference, or a table reference with reversible upward justification. The appendix is therefore not an archive placed after the fact; it is the controlled external memory of the proof architecture.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Operational rule for later versions. Whenever a derivation step is abbreviated in the main line, the abbreviated step should be tagged by a reusable interface name that can be expanded again from the appendix without changing the logical order of the argument. This preserves speed without sacrificing structural auditability.

archival stage Structural Interface Consolidation Core

Remark H3.39 (Normalization note for archival stage)**.** *This interface-consolidation block is retained as historical interface memory. Its active proof-bearing content is now normalized into the semantic/tensor interface and grouped transport phases, so the present block should be read as supporting consolidation rather than as a competing interface foundation.*

This release adds a further consolidation layer between the inherited closure–projection synthesis core and the longer structural body of the monolith. The aim of archival stage is to make the interfaces between closure statements, probe reductions, projection tables, and appendix recoverability even more explicit, so that later mathematical deepening can reuse the same transport rules without reopening the basic admissibility logic.

1. Interface invariance principle

The first strengthening of archival stage is that every admissible transition between main text, probe representation, and appendix table must preserve the same structural body. This means that the interface is not merely notational. It is a controlled transfer channel whose legitimacy depends on invariance of the common closure-supported content.

Consolidation Proposition A (interface invariance). If two representations are linked by a transport that preserves the closure-fixed body, then they are distinct expositions of the same admissible statement rather than competing structural claims.

Proof sketch. The ground condition and HC fix the admissible body before any later compression. A transport that keeps this body intact may change display, resolution, or sector language, but it cannot alter the admissibility source. Hence interface moves preserve statement identity at the structural level.

2. Probe filtration as a pre-projective gate

The probe sector is now stated even more sharply as a pre-projective gate. Its job is not to invent physical outputs but to filter, compress, and expose only those modes that remain compatible with the closure discipline already fixed in the local structural chain.

Consolidation Proposition B (probe filtration is gatekeeping, not generation). Under the standing orthogonality and HC assumptions, the probe sector may rank or suppress final modes, but it cannot generate a physically admissible mode whose closure ancestry is absent.

Proof sketch. The probes inherit their admissibility from the same triolic and closure-governed body as the main sector. Therefore filtration can only reveal or reject what is already licensed; it cannot certify an unlicensed sector by reduction alone.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

3. Table-to-lemma reversibility

The tables and tool blocks gathered in the appendix are promoted once more from summary devices to reversible proof interfaces. For archival stage this reversibility is sharpened into a two-way requirement between rows, lemmas, and transport principles.

Consolidation Principle (row/lemma reversibility). A compact row or tool entry is fully admissible only if the reader can reconstruct both

(a) the lemma or closure chain that licenses the row from above, and

(b) the bounded projection or interpretation that uses the row from below.

Where only one direction is available, the entry remains heuristic and must be marked as such.

4. Strengthened micro-to-macro readout discipline

The micro-to-macro route is no longer only a synthesis slogan. In archival stage it is a regulated readout pipeline:

$$\begin{aligned}&\text{local triolic body} \Longrightarrow \text{HC stabilization} \Longrightarrow \text{closure confinement}\\ &\Longrightarrow \text{probe filtration} \Longrightarrow \text{table/interface compression} \Longrightarrow \text{effective projection.}\end{aligned}$$

This ordering is now part of the operational discipline of the document. A macro statement is acceptable only to the extent that its position in this chain can be traced backward without gap.

Consolidation Proposition C (macro readout requires backward traceability). Any effective large-scale law cited later in the monolith is structurally valid only when it admits a backward trace through table/interface compression, probe filtration, closure confinement, and HC stabilization to the local triolic body.

5. Appendix memory discipline, strengthened again

The appendix is therefore not only a storage area for auxiliary derivations. It is a controlled external memory for the proof architecture. In archival stage every major abbreviation introduced in the running text should be recoverable by a named interface path: lemma path, tool path, table path, or transport path. This rule prepares the next versions for denser mathematical unfolding without losing the structure-first audit trail.

Operational rule for later versions. When a proof step is compressed in the main line, the compression should carry a stable interface name whose recovery route is unique up to equivalent structural transport. This keeps the monolith fast in presentation while remaining expandable under later formal pressure.

H4 Local Crystal, HC Repair and Handoff Protocols

Definition H4.1 (Local compact reinforcement pair)**.** *The* local compact reinforcement pair *extracted from the crystal persistence correction datum*

$$\mathfrak{K}^{\sharp}_{\mathrm{cell}}(n) = (\eta_{\Theta}(n), \eta_T(n), \zeta_T(n), \lambda_T(n))$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

is the pair

$$\mathfrak{A}^{\sharp}_{\mathrm{loc}}(n) := (A_{\mathrm{bur}}(n), A_{\mathrm{geo}}(n))$$

defined by

$$A_{\mathrm{bur}}(n) := \eta_{\Theta}(n), \qquad A_{\mathrm{geo}}(n) := \max\{\eta_T(n), \zeta_T(n), \lambda_T(n)\}.$$

Remark H4.2 (Meaning of the compact pair)**.** *The first parameter records the burden-side phase reinforcement required to stop the second-step Theta drift. The second parameter records the total geometric persistence reinforcement required on the same two-step window. Thus the original four repair numbers are compressed into one burden-side and one geometric reinforcement scale.*

Definition H4.3 (One-parameter shadow reinforcement)**.** *The* one-parameter shadow reinforcement *of the local compact reinforcement pair is*

$$A_{\mathrm{tot}}(n) := \max\{A_{\mathrm{bur}}(n), A_{\mathrm{geo}}(n)\}.$$

Remark H4.4 (Why the one-parameter shadow is secondary)**.** *The scalar shadow $A_{\mathrm{tot}}(n)$ is useful only as a coarse local size readout. It should not replace the two-parameter pair, because the current post-archival stage line still distinguishes between burden-side repair and geometric persistence repair.*

Definition H4.5 (Burden-dominant compact reinforcement)**.** *The local compact reinforcement pair is called* burden-dominant *if*

$$A_{\mathrm{bur}}(n) \geq A_{\mathrm{geo}}(n).$$

It is called geometrically dominant *if*

$$A_{\mathrm{geo}}(n) > A_{\mathrm{bur}}(n).$$

Proposition H4.6 (The crystal persistence correction datum admits a conservative two-parameter compression)**.** *Every local crystal persistence correction datum*

$$(\eta_{\Theta}(n), \eta_T(n), \zeta_T(n), \lambda_T(n))$$

admits a conservative two-parameter compression into

$$(A_{\mathrm{bur}}(n), A_{\mathrm{geo}}(n)).$$

Moreover, the compressed pair dominates the original repair data in the sense that:

(a) $A_{\mathrm{bur}}(n)$ retains the full burden-side strengthening requirement;

(b) $A_{\mathrm{geo}}(n)$ bounds every geometric persistence strengthening requirement from above.

Proof. The first claim is immediate from Definition H4.1. The second follows because $A_{\mathrm{geo}}(n)$ is defined as the maximum of the three geometric persistence reinforcement terms. Hence the compressed pair conservatively dominates the original local correction datum. □

Proposition H4.7 (Current local reinforcement is not purely geometric)**.** *Assume the first local repair-orientation theorem and the first local Teichmüller-persistence repair theorem. Then the current local compact reinforcement pair is not geometrically dominant by default. More precisely, the burden-side component*

$$A_{\mathrm{bur}}(n) = \eta_{\Theta}(n)$$

remains genuinely active in every admissible local repair reading.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. By the local repair-orientation theorem, the first repair is Theta-dominant rather than purely geometric. By the Teichmüller-persistence repair theorem, the geometric side is genuinely active but enters only after the burden-side drift has already been identified as the first required strengthening. Therefore the current local compact reinforcement is not purely geometric by default. □

Theorem H4.8 (First compact reinforcement theorem)**.** *Assume:*

(a) the first local crystal-equation correction theorem applies;

(b) the first local repair-to-trigger calibration theorem applies;

(c) the first local Teichmüller-persistence repair theorem applies.

Then the current post-archival stage local crystal persistence problem admits a first compact reinforcement form.

More precisely:

(1) the four local repair parameters

$$(\eta_\Theta(n), \eta_T(n), \zeta_T(n), \lambda_T(n))$$

compress conservatively to the pair

$$(A_{\mathrm{bur}}(n), A_{\mathrm{geo}}(n));$$

(2) the burden-side component $A_{\mathrm{bur}}(n)$ remains the first active repair scale;

(3) the geometric component $A_{\mathrm{geo}}(n)$ records the total persistence reinforcement required to keep the corrected crystal equation valid on the active two-step window;

(4) the scalar shadow $A_{\mathrm{tot}}(n)$ is only a coarse readout and should not replace the compact pair at the current stage.

Proof. By Proposition H4.6, the four repair parameters admit a conservative compression into the compact pair. By Proposition H4.7, the burden-side component remains genuinely active and therefore cannot be discarded. The geometric component dominates the three persistence-side repair contributions by definition, so it records the total persistence reinforcement requirement on the active two-step window. The scalar shadow is only the maximum of the two compact parameters and therefore loses the burden/geometry distinction still needed at the current stage. □

Corollary H4.9 (Current compact reinforcement working rule)**.** *The next local audit should not first track the full four-parameter repair datum. It should first track the compact reinforcement pair*

$$(A_{\mathrm{bur}}(n), A_{\mathrm{geo}}(n)),$$

using the scalar shadow $A_{\mathrm{tot}}(n)$ only as a coarse size indicator.

Proof. Immediate from Theorem H4.8. □

Definition H4.10 (Local reinforcement ratio)**.** *The* local reinforcement ratio *is the quantity*

$$\chi_{\mathrm{loc}}(n) := \frac{A_{\mathrm{geo}}(n)}{A_{\mathrm{bur}}(n)}$$

whenever $A_{\mathrm{bur}}(n) > 0$.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Remark H4.11 (Meaning of the local reinforcement ratio)**.** *The ratio $\chi_{\mathrm{loc}}(n)$ measures how much geometric persistence reinforcement is required relative to the burden-side phase reinforcement on the active two-step window. It is therefore the first compact indicator of whether the local repair is still burden-led or already close to a balanced coupled regime.*

Definition H4.12 (Local reinforcement regime classes)**.** *Assume $A_{\mathrm{bur}}(n) > 0$. The local compact reinforcement pair is said to be:*

(a) strictly burden-dominant *if*

$$\chi_{\mathrm{loc}}(n) < 1;$$

(b) balanced-coupled *if*

$$\chi_{\mathrm{loc}}(n) = 1;$$

(c) geometrically overcarried *if*

$$\chi_{\mathrm{loc}}(n) > 1.$$

Remark H4.13 (Why the ratio matters)**.** *The compact pair already tells us that both burden-side and geometric reinforcement are active. The ratio tells us which of the two is currently driving the local repair budget.*

Proposition H4.14 (Current compact repair is not balanced by default)**.** *Assume the first compact reinforcement theorem and the first local repair-orientation theorem. Then the current local compact repair is not balanced-coupled by default. More precisely, the post-archival stage line should be read first as strictly burden-dominant unless a later local audit shows that the geometric reinforcement has risen to the same scale.*

Proof. By the first compact reinforcement theorem, the burden-side component $A_{\mathrm{bur}}(n)$ remains the first active repair scale, while the geometric component $A_{\mathrm{geo}}(n)$ records the total persistence reinforcement still required on the same window. By the local repair-orientation theorem, the first repair is Theta-dominant rather than geometric-dominant. Hence the default reading is not balanced-coupled, and the local repair is first to be read as burden-dominant unless later evidence forces a different regime class. □

Definition H4.15 (Local reinforcement balance defect)**.** *The* local reinforcement balance defect *is*

$$B_{\mathrm{loc}}(n) := A_{\mathrm{geo}}(n) - A_{\mathrm{bur}}(n).$$

Remark H4.16 (Interpretation of the balance defect)**.** *Negative balance defect means burden-side dominance. Zero balance defect means balanced coupled repair. Positive balance defect means that the geometric persistence side has overtaken the burden-side repair scale.*

Theorem H4.17 (First local reinforcement-ratio theorem)**.** *Assume:*

(a) the first compact reinforcement theorem applies;

(b) the first local Theta-drift witness theorem applies;

(c) the first local Teichmüller-persistence repair theorem applies.

Then the current post-archival stage local repair supports a first compact ratio reading. More precisely:

(1) the compact pair admits the ratio $\chi_{\mathrm{loc}}(n)$ and the balance defect $B_{\mathrm{loc}}(n)$;

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(2) the default local repair reading is strictly burden-dominant, hence

$$B_{\mathrm{loc}}(n) < 0 \quad \textit{and therefore} \quad \chi_{\mathrm{loc}}(n) < 1$$

in the default regime reading;

(3) the geometric reinforcement remains genuinely active even in this burden-dominant regime, so the ratio does not collapse to zero.

Proof. Items (1) and the sign interpretation are immediate from the definitions of the ratio and balance defect. By the first compact reinforcement theorem, the burden-side component remains the first active repair scale. By the first local Teichmüller-persistence repair theorem, the geometric reinforcement is still genuinely active and therefore cannot be discarded. Hence the default regime is burden-dominant but not purely one-sided, which yields $B_{\mathrm{loc}}(n) < 0$ and therefore $\chi_{\mathrm{loc}}(n) < 1$ in the default reading. □

Corollary H4.18 (Current compact-ratio working rule)**.** *The next local audit should not only record the pair*

$$(A_{\mathrm{bur}}(n), A_{\mathrm{geo}}(n)),$$

but also track the ratio $\chi_{\mathrm{loc}}(n)$ and balance defect $B_{\mathrm{loc}}(n)$ in order to measure whether the current repair remains burden-led or is moving toward balanced coupled repair.

Proof. Immediate from Theorem H4.17. □

Definition H4.19 (Local asymptotic repair orientation)**.** *The* local asymptotic repair orientation *is the long-step tendency of the compact reinforcement ratio*

$$\chi_{\mathrm{loc}}(n) = \frac{A_{\mathrm{geo}}(n)}{A_{\mathrm{bur}}(n)}$$

along an admissible sequence of repaired local windows.

Definition H4.20 (Balance-seeking local repair family)**.** *An admissible local repair family is called* balance-seeking *if*

$$\chi_{\mathrm{loc}}(n) \to 1$$

along the family, equivalently if

$$B_{\mathrm{loc}}(n) \to 0.$$

Definition H4.21 (Persistently burden-dominant local repair family)**.** *An admissible local repair family is called* persistently burden-dominant *if there exists some $\delta_{\mathrm{bal}} > 0$ such that*

$$\chi_{\mathrm{loc}}(n) \leq 1 - \delta_{\mathrm{bal}}$$

for all sufficiently large local repair steps.

Remark H4.22 (Why the asymptotic question matters)**.** *The current compact repair is already burden-dominant at first order. The real next question is whether this is only an initial orientation or whether the burden-side dominance remains structurally built into the repaired persistence problem.*

Proposition H4.23 (The current line does not yet force balance-seeking behavior)**.** *Assume the current post-archival stage local repair reading. Then the existing local theorems do not yet force*

$$\chi_{\mathrm{loc}}(n) \to 1.$$

In particular, no current result requires the geometric reinforcement to rise all the way to burden-side balance.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. The current local repair-orientation theorem yields burden-side dominance at first order, while the Teichmüller-side repair theorems only assert that the geometric reinforcement remains genuinely active. They do not assert that it asymptotically reaches the same scale as the burden-side reinforcement. Hence balance-seeking behavior is not yet forced. □

Proposition H4.24 (The current line is naturally read as burden-dominant unless later balancing evidence appears)**.** *Assume the first local reinforcement-ratio theorem and the first local repair-to-trigger calibration theorem. Then the default asymptotic reading of the current line is persistently burden-dominant unless a later local audit exhibits a genuine balancing mechanism reducing $B_{\mathrm{loc}}(n)$ toward zero.*

Proof. The current ratio theorem places the local repair in the burden-dominant regime, and the first local repair-to-trigger calibration theorem still assigns the primary tightening to the Theta-side threshold. Thus the available repair logic continues to privilege the burden-side channel. Without an additional balancing mechanism, the natural default reading is that burden-side dominance persists. □

Theorem H4.25 (First local asymptotic orientation theorem)**.** *Assume:*

(a) the first local reinforcement-ratio theorem applies;

(b) the first local repair-to-trigger calibration theorem applies;

(c) no later local theorem has yet exhibited a balancing mechanism forcing $B_{\mathrm{loc}}(n) \to 0$.

Then the current post-archival stage line should be read asymptotically as burden-dominant rather than balance-seeking. More precisely, the present local evidence supports persistent burden-side dominance as the default asymptotic orientation, while balanced-coupled repair remains only a later possibility to be justified by new local evidence.

Proof. By the first local reinforcement-ratio theorem, the current compact repair is burden-dominant at first order. By the first local repair-to-trigger calibration theorem, the primary tightening remains Theta-side, although the geometric repair remains genuinely active. In the absence of an additional balancing mechanism driving the balance defect toward zero, the natural asymptotic reading continues the burden-dominant orientation rather than forcing convergence to balance. □

Corollary H4.26 (Current asymptotic working rule)**.** *The next local audit should not presuppose*

$$\chi_{\mathrm{loc}}(n) \to 1.$$

Instead, it should first test whether any genuine balancing mechanism exists at all; until then, the current local repair should be treated as asymptotically burden-dominant with genuine but subordinate geometric carry.

Proof. Immediate from Theorem H4.25. □

Definition H4.27 (Final local balancing mechanisms)**.** *A* final local balancing mechanism *for the compact reinforcement pair*

$$\mathfrak{A}^{\sharp}_{\mathrm{loc}}(n) = (A_{\mathrm{bur}}(n), A_{\mathrm{geo}}(n))$$

is any local structural effect capable of reducing the balance defect

$$B_{\mathrm{loc}}(n) = A_{\mathrm{geo}}(n) - A_{\mathrm{bur}}(n)$$

toward zero. Three final types are distinguished:

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(a) *a* crystal-side balancing mechanism, *acting through the corrected crystal compatibility equation;*

(b) *a* Teichmüller-side balancing mechanism, *acting through the geometric persistence layer alone;*

(c) *a* hybrid crystal–Teichmüller balancing mechanism, *acting by coupled improvement of the crystal compatibility side and the Teichmüller persistence side.*

Remark H4.28 (Why a pure crystal correction is probably not enough)**.** *The current local line already required a first burden-side repair followed by an explicit Teichmüller-side repair. This suggests that crystal-side correction alone is unlikely to be the full balancing mechanism.*

Remark H4.29 (Why a pure Teichmüller correction is also probably not enough)**.** *The local repair-orientation theorem still forces positive burden-side reinforcement*

$$\eta_{\Theta}(n) > 0.$$

Hence the geometric side alone does not erase the burden-side asymmetry.

Definition H4.30 (Hybrid balancing witness)**.** *A* hybrid balancing witness *is a local witness package showing simultaneously:*

(a) *improvement of the corrected crystal compatibility equation,*

(b) *improvement of the Teichmüller-side persistence inequalities,*

(c) *strict reduction of the balance defect*

$$B_{\mathrm{loc}}(n+1) > B_{\mathrm{loc}}(n)$$

toward zero, equivalently

$$|B_{\mathrm{loc}}(n+1)| < |B_{\mathrm{loc}}(n)|$$

on the burden-dominant side.

Proposition H4.31 (The current line does not yet support a purely crystal balancing mechanism)**.** *Assume the current post-archival stage local repair line. Then the available local evidence does not yet support reading the first balancing mechanism as purely crystal-side.*

Proof. The local crystal-equation correction theorem already required the prior burden-side and Teichmüller-side repair data. Hence the crystal correction enters as part of a repaired persistence package, not as a self-sufficient balancing source on its own. □

Proposition H4.32 (The current line does not yet support a purely Teichmüller balancing mechanism)**.** *Assume the first local repair-orientation theorem. Then the available local evidence does not yet support reading the first balancing mechanism as purely Teichmüller-side.*

Proof. The first local repair-orientation theorem keeps the dominant local repair on the burden side while the geometric side remains genuinely active but subordinate. Hence the currently visible repair is not purely Teichmüller-driven. □

Theorem H4.33 (First local balancing-mechanism orientation theorem)**.** *Assume:*

(a) *the first local asymptotic orientation theorem applies;*

(b) *the first local crystal-equation correction theorem applies;*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(c) the first local Teichmüller-persistence repair theorem applies.

Then the most plausible final for a future local balancing mechanism is not purely crystal-side and not purely Teichmüller-side, but hybrid crystal–Teichmüller.

Proof. By the first local asymptotic orientation theorem, the current line remains burden-dominant unless a later balancing mechanism appears. By the local crystal-equation correction theorem, crystal-side correction is relevant but not self-sufficient. By the local Teichmüller-persistence repair theorem, geometric repair is genuinely active but likewise not self-sufficient. Therefore the first plausible balancing mechanism must couple both corrections if it is to reduce the balance defect toward zero. □

Corollary H4.34 (Current balancing-mechanism working rule)**.** *The next local audit should not first search for a purely crystal balancing mechanism or a purely Teichmüller balancing mechanism. Instead, it should search for the first hybrid witness that lowers the balance defect by coordinated crystal-side and geometric-side repair.*

Proof. Immediate from Theorem H4.33. □

Definition H4.35 (First hybrid balancing step)**.** *A* first hybrid balancing step *from the local state at step n to the next step $n+1$ is a local repaired transition satisfying all of the following:*

(a) the corrected crystal compatibility equation improves from

$$\mathcal{C}^{\sharp}_{\mathrm{cell}}(X_n, X_{n+1})$$

to a closer-to-compatible value on the updated two-step window;

(b) the Teichmüller-side persistence repair data improve in the sense that the repaired geometric persistence test is strictly better satisfied on the updated window;

(c) the compact reinforcement pair

$$\mathfrak{A}^{\sharp}_{\mathrm{loc}}(n) = (A_{\mathrm{bur}}(n), A_{\mathrm{geo}}(n))$$

is updated to

$$\mathfrak{A}^{\sharp}_{\mathrm{loc}}(n+1) = (A_{\mathrm{bur}}(n+1), A_{\mathrm{geo}}(n+1))$$

with strictly smaller balance defect magnitude

$$|B_{\mathrm{loc}}(n+1)| < |B_{\mathrm{loc}}(n)|;$$

(d) the update remains inside the same local admissibility regime.

Definition H4.36 (First hybrid balancing witness package)**.** *A* first hybrid balancing witness package *at step n is a finite witness datum*

$$\mathfrak{W}_{\mathrm{hyb}}(n) = \big(\delta_{\mathrm{cell}}(n), \delta_T(n), \Delta B_{\mathrm{loc}}(n)\big)$$

such that:

(a) $\delta_{\mathrm{cell}}(n) > 0$ measures the crystal-side compatibility gain of the first hybrid balancing step;

(b) $\delta_T(n) > 0$ measures the Teichmüller-side persistence gain of the same step;

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(c)

$$\Delta B_{\mathrm{loc}}(n) := |B_{\mathrm{loc}}(n)| - |B_{\mathrm{loc}}(n+1)| > 0$$

measures the strict gain toward local balance.

Remark H4.37 (Meaning of the hybrid witness package)**.** *The first hybrid balancing witness package is the first local datum that does more than merely diagnose burden-dominance. It certifies that crystal-side correction and Teichmüller-side repair improve together and that the resulting coupled improvement genuinely lowers the balance defect.*

Definition H4.38 (Hybrid balancing efficiency ratio)**.** *The* hybrid balancing efficiency ratio *of a first hybrid balancing witness package is*

$$\Xi_{\mathrm{hyb}}(n) := \frac{\Delta B_{\mathrm{loc}}(n)}{\delta_{\mathrm{cell}}(n) + \delta_T(n)},$$

provided the denominator is positive.

Remark H4.39 (Why the efficiency ratio is useful)**.** *This ratio records how much balance-defect reduction is obtained per unit of coupled crystal-plus-Teichmüller repair. It is therefore the first local scalar witness that the hybrid mechanism is not merely present, but actually effective.*

Proposition H4.40 (A first hybrid balancing step is stronger than separate side repairs)**.** *Every first hybrid balancing step is stronger than a purely crystal-side repair and stronger than a purely Teichmüller-side repair considered separately.*

Proof. By definition, a first hybrid balancing step requires simultaneous crystal-side improvement, simultaneous Teichmüller-side persistence improvement, and strict reduction of the balance defect magnitude. A purely one-sided repair can satisfy at most one of the two side-improvement clauses, but not both together with the coupled balance reduction requirement. Hence the hybrid step is strictly stronger. □

Lemma H4.41 (First hybrid balancing witness lemma)**.** *Assume:*

(a) the first local balancing-mechanism orientation theorem applies;

(b) a first hybrid balancing step exists on the current local window.

Then a first hybrid balancing witness package exists.

Proof. If a first hybrid balancing step exists, then by definition it produces strictly positive crystal-side compatibility gain, strictly positive Teichmüller-side persistence gain, and strict decrease of the balance defect magnitude. These three gains form the witness package

$$\mathfrak{W}_{\mathrm{hyb}}(n) = (\delta_{\mathrm{cell}}(n), \delta_T(n), \Delta B_{\mathrm{loc}}(n)).$$

□

Theorem H4.42 (First hybrid balancing witness theorem)**.** *Assume:*

(a) the first local balancing-mechanism orientation theorem applies;

(b) the current line is still asymptotically burden-dominant;

(c) a local repaired two-step window exists on which crystal-side and Teichmüller-side improvements occur simultaneously.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Then the current line supports a first hybrid balancing witness exactly when the simultaneous side improvements produce strict reduction of

$$|B_{\mathrm{loc}}(n)|.$$

Equivalently, the first honest evidence for a genuine balancing mechanism is not the mere coexistence of crystal and geometric repair, but the existence of a positive witness package

$$\mathfrak{W}_{\mathrm{hyb}}(n) = (\delta_{\mathrm{cell}}(n), \delta_T(n), \Delta B_{\mathrm{loc}}(n))$$

with

$$\Delta B_{\mathrm{loc}}(n) > 0.$$

Proof. By the balancing-mechanism orientation theorem, any plausible future balancing mechanism must be hybrid rather than purely one-sided. If simultaneous crystal-side and Teichmüller-side improvements occur but do not reduce $|B_{\mathrm{loc}}(n)|$, then no genuine balancing has yet been witnessed; the system remains merely burden-dominant with parallel repairs. Conversely, if the simultaneous side improvements do reduce $|B_{\mathrm{loc}}(n)|$, then the resulting gains define a positive hybrid witness package, and the first honest evidence for a balancing mechanism is present. □

Corollary H4.43 (Current hybrid-balancing working rule)**.** *The next local audit should not merely ask whether crystal-side and Teichmüller-side repair are both visible. It should ask whether they jointly produce*

$$\Delta B_{\mathrm{loc}}(n) > 0,$$

and hence a first genuine hybrid balancing witness package.

Definition H4.44 (Hybrid witness existence obstruction)**.** *A* hybrid witness existence obstruction *at step n is the failure of the current local line to produce any witness package*

$$\mathfrak{W}_{\mathrm{hyb}}(n) = (\delta_{\mathrm{cell}}(n), \delta_T(n), \Delta B_{\mathrm{loc}}(n))$$

with

$$\delta_{\mathrm{cell}}(n) > 0, \qquad \delta_T(n) > 0, \qquad \Delta B_{\mathrm{loc}}(n) > 0.$$

Equivalently, the line remains burden-dominant because no first honest hybrid balancing witness is yet produced.

Definition H4.45 (Weak hybrid efficiency obstruction)**.** *A* weak hybrid efficiency obstruction *at step n is the situation in which a first hybrid balancing witness package exists but its hybrid balancing efficiency ratio*

$$\Xi_{\mathrm{hyb}}(n) = \frac{\Delta B_{\mathrm{loc}}(n)}{\delta_{\mathrm{cell}}(n) + \delta_T(n)}$$

remains too small to alter the current asymptotic burden-dominant orientation.

Remark H4.46 (Existence versus efficiency)**.** *There are therefore two distinct post-obstructions: first, the line may fail to produce any honest hybrid witness at all; second, it may produce such a witness, but only with negligible balancing efficiency. These two failure modes should not be confused.*

Definition H4.47 (Existence-prior local regime)**.** *The current local line is said to be in an* existence-prior local regime *if the first balancing obstruction is of witness-existence type rather than weak-efficiency type.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition H4.48 (Efficiency-prior local regime)**.** *The current local line is said to be in an* efficiency-prior local regime *if a first hybrid balancing witness is already present, but its efficiency ratio remains too small to change the burden-dominant asymptotic orientation.*

Proposition H4.49 (Asymptotic burden-dominance blocks efficiency-first readings)**.** *Assume the first local asymptotic orientation theorem and the first hybrid balancing witness theorem. If no positive witness package*

$$\mathfrak{W}_{\mathrm{hyb}}(n)$$

with

$$\Delta B_{\mathrm{loc}}(n) > 0$$

is yet available, then the current line cannot honestly be read as efficiency-prior.

Proof. An efficiency-prior reading presupposes that a first hybrid balancing witness already exists and that only its efficiency remains insufficient. If no positive witness package is available at all, then the obstruction occurs before efficiency becomes a meaningful variable. Hence the line cannot honestly be read as efficiency-prior. □

Theorem H4.50 (First hybrid obstruction orientation theorem)**.** *Assume:*

(a) the first local asymptotic orientation theorem applies;

(b) the first local balancing-mechanism orientation theorem applies;

(c) the first hybrid balancing witness theorem identifies positive hybrid witness data as the first honest balancing evidence.

Then the current post-line is obstruction-prior on the witness-existence side. More precisely, its first balancing obstruction is not that an existing hybrid witness is too weak, but that no first positive hybrid balancing witness package has yet been produced.

Proof. By the first local asymptotic orientation theorem, the line remains burden-dominant by default unless a genuine balancing mechanism is exhibited. By the balancing-mechanism orientation theorem, any such genuine mechanism must be hybrid. By the first hybrid balancing witness theorem, the first honest evidence for that mechanism is exactly a positive witness package with $\Delta B_{\mathrm{loc}}(n) > 0$. Therefore, as long as no such package is present, the primary obstruction lies on the witness-existence side rather than on the efficiency side. □

Corollary H4.51 (Current hybrid-balancing obstruction rule)**.** *The next local audit should first ask whether a positive hybrid witness package exists at all. Only after such a package is found should the secondary question of hybrid efficiency be treated as the leading obstruction.*

Proof. Immediate from Theorem H4.50. □

Definition H4.52 (Final positive hybrid construction step)**.** *A* final positive hybrid construction step *at stage n is a local update from step n to $n+1$ such that:*

(a) the burden-side repair is active with

$$\delta_{\mathrm{cell}}(n) > 0;$$

(b) the Teichmüller-side persistence repair is active with

$$\delta_T(n) > 0;$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(c) both repairs act on the same local closure-phase window;

(d) the updated local balance defect satisfies

$$B_{\mathrm{loc}}(n+1) > B_{\mathrm{loc}}(n).$$

Remark H4.53 (Why the construction is phrased this way)**.** *The current line does not yet justify assuming that a positive hybrid witness appears automatically once both repair channels are nonzero. The real construction requirement is stronger: the simultaneous crystal-side and Teichmüller-side repairs must improve the local balance defect itself.*

Definition H4.54 (Positive hybrid construction datum)**.** *A* positive hybrid construction datum *at stage n is a quadruple*

$$\mathfrak{C}^{+}_{\mathrm{hyb}}(n) := \big(\delta_{\mathrm{cell}}(n), \delta_T(n), B_{\mathrm{loc}}(n), B_{\mathrm{loc}}(n+1)\big)$$

attached to a final positive hybrid construction step.

Definition H4.55 (Strictly positive hybrid construction)**.** *A final positive hybrid construction step is called* strictly positive *if, in addition,*

$$\Delta B_{\mathrm{loc}}(n) := B_{\mathrm{loc}}(n+1) - B_{\mathrm{loc}}(n) > 0.$$

Definition H4.56 (Hybrid construction gate)**.** *The* hybrid construction gate *at stage n is the conjunction of the conditions*

$$\delta_{\mathrm{cell}}(n) > 0, \qquad \delta_T(n) > 0, \qquad \Delta B_{\mathrm{loc}}(n) > 0.$$

Remark H4.57 (Existence obstruction as gate failure)**.** *Under the current obstruction-oriented reading, the missing positive hybrid witness is equivalently the failure of the hybrid construction gate. This makes the existence problem testable in the same local language as the later efficiency problem.*

Proposition H4.58 (Positive hybrid witness construction implies witness existence)**.** *Every strictly positive hybrid construction step produces a positive hybrid witness package*

$$\mathfrak{W}_{\mathrm{hyb}}(n) = \big(\delta_{\mathrm{cell}}(n), \delta_T(n), \Delta B_{\mathrm{loc}}(n)\big)$$

with

$$\delta_{\mathrm{cell}}(n) > 0, \qquad \delta_T(n) > 0, \qquad \Delta B_{\mathrm{loc}}(n) > 0.$$

Proof. By definition, a strictly positive hybrid construction step already supplies positive crystal-side repair, positive Teichmüller-side repair, and a strictly positive balance-defect gain. These are exactly the components required for a positive hybrid witness package. □

Proposition H4.59 (The current existence obstruction is equivalent to absence of a strictly positive hybrid construction)**.** *Assume the first hybrid obstruction orientation theorem. Then the current line lies in the existence-prior regime if and only if no strictly positive hybrid construction step is yet available on the active local window.*

Proof. By the first hybrid obstruction orientation theorem, the current primary obstruction is the absence of a positive hybrid witness package. By Proposition H4.58, every strictly positive hybrid construction step would already produce such a package. Hence the existence-prior obstruction holds exactly when no such construction step is yet available. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Theorem H4.60 (First positive hybrid witness construction theorem)**.** *Assume:*

(a) the first hybrid obstruction orientation theorem applies;

(b) the local crystal-side correction datum is available;

(c) the local Teichmüller-side repair datum is available;

(d) both data act on the same active closure-phase window.

Then the first honest transition out of the existence-prior regime occurs exactly when a strictly positive hybrid construction step is produced. Equivalently, the first positive hybrid balancing witness arises exactly when the hybrid construction gate is opened on the active local window.

Proof. By the first hybrid obstruction orientation theorem, the current line remains stuck before the efficiency question precisely because no positive hybrid witness yet exists. By Proposition H4.58, every strictly positive hybrid construction step produces such a witness. Conversely, on the active local window a first positive witness must come from simultaneous positive crystal-side and Teichmüller-side repair that genuinely improves the local balance defect, which is exactly the content of a strictly positive hybrid construction step. Hence exiting the existence-prior regime is equivalent to producing such a step, or equivalently to opening the hybrid construction gate. □

Corollary H4.61 (Current hybrid-construction working rule)**.** *The next local audit should not yet optimize hybrid efficiency. It should first test whether the active local crystal-side and Teichmüller-side repairs can be synchronized into a strictly positive hybrid construction step with*

$$\Delta B_{\mathrm{loc}}(n) > 0.$$

Proof. Immediate from Theorem H4.60. □

Definition H4.62 (Crystal-side gate obstruction)**.** *A* crystal-side gate obstruction *at stage n is the failure of the hybrid construction gate caused primarily by the crystal-side component, i.e. by one of the following:*

(a)

$$\delta_{\mathrm{cell}}(n) \leq 0;$$

(b) the crystal-side correction datum is not available on the active local window;

(c) the crystal-side correction fails to synchronize with the active Teichmüller-side repair on the same closure-phase window.

Definition H4.63 (Teichmüller-side gate obstruction)**.** *A* Teichmüller-side gate obstruction *at stage n is the failure of the hybrid construction gate caused primarily by the geometric persistence side, i.e. by one of the following:*

(a)

$$\delta_T(n) \leq 0;$$

(b) the Teichmüller-side repair datum is not available on the active local window;

(c) the geometric repair remains too weak to support

$$\Delta B_{\mathrm{loc}}(n) > 0$$

even when the crystal-side correction is already positive.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition H4.64 (Gate-side desynchronization defect)**.** *The* gate-side desynchronization defect *at stage n is the obstruction datum*

$$\mathfrak{D}_{\mathrm{gate}}(n) := \big(\delta_{\mathrm{cell}}(n), \delta_T(n), \Delta B_{\mathrm{loc}}(n)\big)$$

read under the question which side first fails to keep the hybrid construction gate open.

Remark H4.65 (What "first out of sync" means)**.** *The present question is not yet one of hybrid efficiency. It asks which side fails first at the level of gate opening: the crystal-side correction, the Teichmüller-side persistence repair, or the synchronization of both on the same local closure-phase window.*

Proposition H4.66 (The current line is not crystal-obstruction-prior)**.** *Under the current post-archival stage reading, the active local line is not primarily blocked by absence of crystal-side correction.*

Proof. The local crystal-equation correction theorem and the compact reinforcement line already support a positive crystal-side correction datum on the active local window. The remaining obstruction therefore does not first arise from the total absence of the crystal-side contribution. □

Proposition H4.67 (The current line is Teichmüller-side-prior at the gate)**.** *Under the current post-archival stage reading, the first gate-side desynchronization is read primarily on the Teichmüller-side persistence channel.*

Proof. The tightened-package plausibility reading and the subsequent Teichmüller-persistence repair line already isolate the residual insufficiency on the geometric persistence side after the first burden-side repair has been introduced. The hybrid-construction obstruction then remains because a positive crystal-side correction does not yet suffice to produce

$$\Delta B_{\mathrm{loc}}(n) > 0.$$

Hence the first gate-side desynchronization is read primarily on the Teichmüller-side channel rather than on the crystal side. □

Theorem H4.68 (First gate-side desynchronization orientation theorem)**.** *Assume:*

(a) the first positive hybrid witness construction theorem applies;

(b) the local crystal-equation correction theorem applies;

(c) the first local Teichmüller-persistence repair theorem applies;

(d) the current line still lies in the existence-prior hybrid obstruction regime.

Then the current gate failure is Teichmüller-side-prior.
More precisely:

(i) the crystal-side correction is not the first component to fall out of sync;

(ii) the primary remaining defect is the insufficient geometric persistence contribution needed to turn the positive local corrections into

$$\Delta B_{\mathrm{loc}}(n) > 0;$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(iii) the next local construction step should therefore strengthen the Teichmüller-side synchronization before attempting any efficiency optimization.

Proof. By Proposition H4.66, the crystal-side correction is already locally present and is not the primary missing ingredient. By Proposition H4.67, the residual desynchronization sits on the Teichmüller-side persistence channel. Since the hybrid witness still does not yet exist, the active local task is not efficiency optimization but restoration of sufficient geometric persistence to open the gate. □

Corollary H4.69 (Current gate-side working rule)**.** *The next local audit should first ask how the Teichmüller-side repair must be strengthened or synchronized so that the already present crystal-side correction becomes hybrid-gate-effective. Only after that should the line pass to hybrid efficiency questions.*

Proof. Immediate from Theorem H4.68. □

Definition H4.70 (Phase-lock synchronization defect)**.** *A* phase-lock synchronization defect *at stage n is the failure of the second-step local repair to keep the phase-side burden correction aligned with the same closure-phase window on which the first-step near-trigger state was read.*

Definition H4.71 (Teichmüller-carry synchronization defect)**.** *A* Teichmüller-carry synchronization defect *at stage n is the failure of the geometric persistence correction to carry the repaired second step inside the same local Teichmüller-safe band as the first step.*

Definition H4.72 (Local Teichmüller-side synchronization datum)**.** *The* local Teichmüller-side synchronization datum *at stage n is*

$$\mathfrak{S}_T^{\mathrm{loc}}(n) := \big(\sigma_\Theta^{\mathrm{lock}}(n), \sigma_T^{\mathrm{carry}}(n)\big),$$

where:

(a) $\sigma_\Theta^{\mathrm{lock}}(n)$ measures the remaining phase-lock misalignment of the repaired second step;

(b) $\sigma_T^{\mathrm{carry}}(n)$ measures the remaining geometric carry defect on the same two-step local window.

Remark H4.73 (What synchronization means locally)**.** *The present question is no longer whether geometry matters at all. It asks more precisely which kind of Teichmüller-side synchronization is still missing: the phase-side lock itself, the geometric carry of that lock, or both together.*

Definition H4.74 (Locally synchronized repaired second step)**.** *A repaired second step at stage n is called* locally synchronized *if*

$$\sigma_\Theta^{\mathrm{lock}}(n) = 0 \qquad \textit{and} \qquad \sigma_T^{\mathrm{carry}}(n) = 0.$$

Proposition H4.75 (The current line is not raw-geometry-prior)**.** *Under the current post-archival stage reading, the active local line is not primarily blocked by raw geometric carry alone.*

Proof. The current line already carries a positive Teichmüller-side repair contribution and a positive geometric witness contribution. The residual failure therefore cannot be read first as total absence of geometry, but only as insufficient synchronization of that geometric contribution with the repaired phase-side step. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proposition H4.76 (The current line is phase-lock-prior on the Teichmüller side)**.** *Under the current post-archival stage reading, the first remaining Teichmüller-side synchronization defect is phase-lock-prior.*

Proof. The gate-side desynchronization orientation theorem already identifies the first remaining gate defect as Teichmüller-side-prior rather than crystal-side-prior. The local repair-orientation and tightened-package plausibility lines then show that the geometric channel is active but still fails to hold the repaired second step in the same closure-phase window. Hence the first remaining defect is best read as insufficient phase-lock synchronization, with geometric carry acting as the witness and support channel. □

Theorem H4.77 (First Teichmüller-side synchronization orientation theorem)**.** *Assume:*

(a) the first local Theta-drift witness theorem applies;

(b) the first local phase-side repair theorem applies;

(c) the first local Teichmüller-persistence repair theorem applies;

(d) the first gate-side desynchronization orientation theorem applies.

Then the first remaining Teichmüller-side synchronization problem is phase-lock-prior rather than raw-carry-prior.

More precisely:

(i) the missing synchronization is not first the total absence of geometric carry;

(ii) the missing synchronization is first the failure to keep the repaired second step phase-locked inside the same closure-phase window;

(iii) the geometric carry defect remains genuinely active, but as the supporting witness of the phase-lock failure rather than as the primary independent obstruction.

Proof. By Proposition H4.75, the active line is not blocked first by total absence of geometry. By Proposition H4.76, the first remaining synchronization failure is read on the phase-lock component of the repaired second step. Therefore the current Teichmüller-side synchronization problem is phase-lock-prior, with the geometric carry defect remaining real but subordinate. □

Corollary H4.78 (Current Teichmüller-side synchronization working rule)**.** *The next local audit should not ask first for more geometry in the abstract. It should ask first how the repaired second step can be phase-locked back into the same closure-phase window, while using the geometric channel to certify and carry that repaired lock.*

Proof. Immediate from Theorem H4.77. □

Definition H4.79 (Local phase-lock repair datum)**.** *The* local phase-lock repair datum *at stage n is*

$$\mathfrak{R}^{\mathrm{loc}}_{\mathrm{lock}}(n) := \big(\ell^{\mathrm{lock}}_{\Theta}(n), \ell^{\mathrm{lock}}_{T}(n)\big),$$

where $\ell^{\mathrm{lock}}_{\Theta}(n)$ denotes the extra phase-side lock correction required to remove the residual phase-lock defect and $\ell^{\mathrm{lock}}_{T}(n)$ denotes the extra Teichmüller-side carry needed to stabilize that repaired lock on the same two-step window.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition H4.80 (Phase-lock repaired local window)**.** *A repaired two-step local window at stage n is called* phase-lock repaired *if, after applying the local phase-lock repair datum, one has*

$$\sigma_{\Theta}^{\mathrm{lock}}(n) = 0 \qquad \textit{and} \qquad \sigma_{T}^{\mathrm{carry}}(n) \le \ell_{T}^{\mathrm{lock}}(n).$$

Equivalently, the repaired second step is brought back into the same active closure-phase window while the remaining geometric carry defect is absorbed by the stated lock-carry allowance.

Definition H4.81 (Minimal local phase-lock repair)**.** *A local phase-lock repair datum*

$$\mathfrak{R}_{\mathrm{lock}}^{\mathrm{loc}}(n) = \big(\ell_{\Theta}^{\mathrm{lock}}(n), \ell_{T}^{\mathrm{lock}}(n)\big)$$

is called minimal *if no strictly smaller pair of nonnegative corrections removes the phase-lock synchronization defect on the same two-step local window.*

Remark H4.82 (What the phase-lock repair really does)**.** *The phase-lock repair is not a fresh bulk law. It is the smallest local correction that forces the repaired second step back onto the same closure-phase slice as the first near-trigger step. Geometrically, one may think of it as a local re-aiming plus carry-stabilization of the second point so that both points lie again in the same admissible narrow corridor rather than in neighboring but misaligned corridors.*

Lemma H4.83 (First local phase-lock repair lemma)**.** *Assume:*

(a) the first Teichmüller-side synchronization orientation theorem applies;

(b) the active local obstruction is phase-lock-prior;

(c) a local phase-lock repair datum exists.

Then the vanishing condition

$$\sigma_{\Theta}^{\mathrm{lock}}(n) = 0$$

is the first honest local repair target for reopening the hybrid construction gate.

Proof. By Theorem H4.77, the first remaining synchronization obstruction is read on the phase-lock component rather than on raw geometric absence. Hence the first honest repair target is to remove the phase-lock defect itself. The accompanying carry correction remains necessary, but it is subordinate to the lock restoration as long as the active obstruction is phase-lock-prior. □

Proposition H4.84 (Current line supports phase-lock-prior local repair)**.** *Under the present post-archival stage reading, the next local strengthening step should first reduce $\sigma_{\Theta}^{\mathrm{lock}}(n)$ and only then optimize the associated geometric carry.*

Proof. This is just the operational form of the phase-lock-prior diagnosis. Since the line is not raw-geometry-prior, one does not first optimize geometric amplitude in the abstract. One first restores alignment of the repaired second step with the active closure-phase window, and then uses the Teichmüller-side correction to support that restored lock. □

Theorem H4.85 (First phase-lock repair theorem)**.** *Assume:*

(a) the first local Theta-drift witness theorem applies;

(b) the first local phase-side repair theorem applies;

(c) the first local Teichmüller-persistence repair theorem applies;

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(d) the first Teichmüller-side synchronization orientation theorem applies;

(e) a minimal local phase-lock repair datum exists.

Then the current post-archival stage line supports a first concrete phase-lock repair final. More precisely:

(i) the primary repair target is

$$\sigma_{\Theta}^{\mathrm{lock}}(n) \to 0;$$

(ii) the geometric carry repair remains genuinely active through

$$\ell_{T}^{\mathrm{lock}}(n) > 0$$

whenever pure phase-side re-alignment does not by itself stabilize the repaired second step;

(iii) a phase-lock repaired local window is the first local final for reopening the hybrid construction gate.

Proof. Items (a)–(d) identify the present obstruction chain: the first failure is Theta-drift-prior, the first repair is burden-side with geometric carry, the next residual obstruction is Teichmüller-side-prior, and that obstruction is phase-lock-prior rather than raw-geometry-prior. Therefore the first concrete local repair must target vanishing of $\sigma_{\Theta}^{\mathrm{lock}}(n)$. If pure re-alignment is not enough to keep the repaired second step in the same local Teichmüller-safe band, a positive carry correction $\ell_{T}^{\mathrm{lock}}(n)$ remains necessary. This yields the first concrete phase-lock repair final. □

Corollary H4.86 (Current phase-lock repair working rule)**.** *The next local audit should no longer ask merely whether the repaired second step is better than before. It should ask whether the applied local correction actually forces*

$$\sigma_{\Theta}^{\mathrm{lock}}(n) = 0$$

and, if not, which remaining carry correction still prevents a fully phase-locked repaired window.

Proof. Immediate from Theorem H4.85. □

Definition H4.87 (Local Heisenberg compensator)**.** *A* local Heisenberg compensator *at stage n is a conservative local synchronization operator*

$$\mathrm{HC}_{\mathrm{loc}}(n) : (X_n, X_{n+1}) \longmapsto (X_n, \widetilde{X}_{n+1})$$

which acts only on the repaired second step and is intended to reduce the remaining phase-lock and Teichmüller-carry defects without changing the already accepted first-step near-trigger reading.

Definition H4.88 (HC-admissible local compensation)**.** *A local Heisenberg compensator is called* HC-admissible *if it satisfies:*

(a) the closure burden remains controlled;

(b) the corrected second step stays inside the same admissibility band;

(c) the phase-lock defect decreases:

$$\sigma_{\Theta}^{\mathrm{lock}}(n; \widetilde{X}_{n+1}) < \sigma_{\Theta}^{\mathrm{lock}}(n; X_{n+1});$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(d) the Teichmüller-carry defect does not increase, and preferably also decreases;

(e) the compensator does not manufacture a false entry certificate by destroying the route-faithful burden reading.

Remark H4.89 (How the HC is read here)**.** *The Heisenberg compensator is not used here as a new bulk law. It is read as the minimal local re-phasing operator that tries to snap the repaired second step back into the same narrow closure-phase corridor as the first near-trigger step.*

Definition H4.90 (Bombe-guided compensator menu)**.** *A* Bombe-guided compensator menu *at stage n is a finite family*

$$\mathcal{M}_{\mathrm{HC}}(n) = \{H_1, \dots, H_r\}$$

of final local Heisenberg compensators together with a contradiction-sensitive elimination rule that removes every final violating closure control, admissibility, or route-faithful burden preservation.

Remark H4.91 (Why the Bombe idea fits here)**.** *The present problem is not to guess one magical compensator. It is to eliminate locally incompatible synchronizers from a finite final menu until only the route-faithful phase-lock-capable ones remain. This is exactly the right local role for a Bombe-style elimination discipline.*

Definition H4.92 (Compensator filter tower)**.** *A* compensator filter tower *is a finite ordered stack*

$$\mathfrak{F}_{\mathrm{HC}}(n) = (F_1, \dots, F_s)$$

of local HC-compatible filters through which the repaired second step is passed before the final trigger test. The intended reading is:

(i) burden-side pre-alignment;

(ii) phase-lock re-alignment;

(iii) Teichmüller-carry stabilization;

(iv) final closure-phase re-entry test.

Remark H4.93 (Filter tower and the electromagnetic analogy)**.** *The filter-tower language should be read structurally rather than as a literal new electrodynamics. It expresses that several local synchronization layers may act in series, each removing one kind of phase/carry mismatch before the final trigger decision is taken.*

Definition H4.94 (Readout-relative informational gravitation principle)**.** *A* readout-relative informational gravitation principle *is the principle that the apparent curvature/attraction acting on a local state is not necessarily a primitive force attached only to rest mass, but may be read as the effective curvature law induced by the chosen admissible readout on the underlying information-carrying structure.*

Remark H4.95 (How far this principle should be claimed)**.** *In the present document this principle is only an internal architecture hypothesis. It does not claim multiple literal physical gravities. It claims instead that one deeper ART/QFT-compatible carrier may have different gravitational-like readout shadows depending on the active admissible information/readout channel.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proposition H4.96 (HC can serve as a local phase-lock operator)**.** *Assume an HC-admissible local compensation exists at stage n. Then the Heisenberg compensator can serve as a local phase-lock operator for the repaired second step.*

Proof. By HC-admissibility, the compensator reduces the phase-lock defect while preserving closure control, admissibility, and the route-faithful burden reading. Therefore it acts exactly as the required local phase-lock operator. □

Lemma H4.97 (Bombe-guided HC elimination lemma)**.** *Assume the compensator final menu is finite. Then Bombe-guided elimination reduces the local HC search to the subfamily of finals that are simultaneously closure-compatible, admissibility-safe, and phase-lock-improving.*

Proof. By definition, Bombe-guided elimination removes every final violating one of the required local compatibility conditions. Hence the surviving menu consists exactly of those finals satisfying the required conditions. □

Theorem H4.98 (First local Heisenberg compensator theorem)**.** *Assume:*

(a) the first phase-lock repair theorem applies;

(b) a finite Bombe-guided compensator menu is available;

(c) at least one surviving final is HC-admissible.

Then the current post-archival stage line supports a first local Heisenberg-compensated phase-lock repair.

More precisely:

(i) the repaired second step can be re-phased by a surviving HC final;

(ii) the corrected step is brought closer to the same closure-phase corridor as the first near-trigger step;

(iii) the compensator acts as a synchronization layer between the local crystal correction and the Teichmüller-side persistence repair.

Proof. By the first phase-lock repair theorem, the current line already isolates the local phase-lock defect as the next concrete obstruction. By the Bombe-guided HC elimination lemma, the finite menu reduces to the finals that are compatible with closure, admissibility, and phase-lock improvement. If one surviving final is HC-admissible, then by Proposition H4.96 it provides the required local re-phasing. Therefore the Heisenberg compensator acts as the missing synchronization layer between the crystal-side correction and the Teichmüller-side persistence repair. □

Corollary H4.99 (Current HC working rule)**.** *The next local audit should not ask first for a new bulk equation. It should ask first for the smallest HC-admissible local compensator that phase-locks the repaired second step back into the same closure-phase corridor while preserving closure control and admissibility.*

Proof. Immediate from Theorem H4.98. □

Definition H4.100 (Phase-side dominant HC final)**.** *A local Heisenberg compensator final at stage n is called* phase-side dominant *if its primary improvement acts on the phase-lock defect, in the sense that it strictly lowers*

$$\sigma_{\Theta}^{\mathrm{lock}}(n)$$

while any improvement of the Teichmüller-carry defect remains subordinate to that phase-lock gain.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition H4.101 (Carry-dominant HC final)**.** *A local Heisenberg compensator final at stage n is called* carry-dominant *if its primary improvement acts on the geometric carry defect, in the sense that it lowers*

$$\sigma_T^{\mathrm{carry}}(n)$$

without first removing the phase-lock misalignment.

Definition H4.102 (Two-component HC final)**.** *A local Heisenberg compensator final at stage n is called* two-component coupled *if it lowers both*

$$\sigma_\Theta^{\mathrm{lock}}(n) \qquad \textit{and} \qquad \sigma_T^{\mathrm{carry}}(n)$$

with neither correction honestly subordinate to the other at first local order.

Remark H4.103 (What the classification is trying to decide)**.** *The present question is not whether a local HC helps at all. It asks which kind of local HC is the first plausible one for the current line: one that primarily restores phase-lock, one that primarily carries geometry, or one that is already fully two-component at first local order.*

Proposition H4.104 (The current line is not carry-dominant on the HC side)**.** *Under the present post-archival stage reading, the first plausible local HC final is not carry-dominant.*

Proof. By the first Teichmüller-side synchronization orientation theorem and the first phase-lock repair theorem, the current residual obstruction is phase-lock-prior rather than raw-carry-prior. Therefore a compensator that tries first to optimize geometric carry without first restoring phase-lock cannot be the default first final. □

Proposition H4.105 (The current line is not yet fully two-component on the HC side)**.** *Under the present post-archival stage reading, the first plausible local HC final is not yet fully two-component at first local order.*

Proof. The current local repair line remains asymptotically burden-dominant and phase-lock-prior. Although the geometric carry correction is genuinely active, the present line has not yet produced a positive hybrid balancing witness with balanced first-order contributions. Hence the first plausible HC final cannot yet be classified as fully two-component at first local order. □

Theorem H4.106 (First HC-final type orientation theorem)**.** *Assume:*

(a) the first local Heisenberg compensator theorem applies;

(b) the first Teichmüller-side synchronization orientation theorem applies;

(c) the first phase-lock repair theorem applies;

(d) the current line has not yet produced a positive hybrid balancing witness.

Then the first plausible local HC final is phase-side dominant rather than carry-dominant or fully two-component.

More precisely:

(i) the primary local HC task is to reduce

$$\sigma_\Theta^{\mathrm{lock}}(n);$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(ii) the Teichmüller-side carry correction remains genuinely active as a supporting synchronization channel;

(iii) the current HC final should therefore be read as phase-side dominant but geometrically carried.

Proof. By Proposition H4.104, the current line is not blocked first by raw geometric carry. By Proposition H4.105, the current line has not yet reached a first-order balanced two-component compensator regime. Since the active residual obstruction is phase-lock-prior, the first plausible HC final must therefore act primarily on the phase-lock defect, with geometric carry serving as the supporting channel that stabilizes the repaired lock. □

Corollary H4.107 (Current HC-final working rule)**.** *The next local audit should not search first for a carry-dominant or perfectly balanced HC. It should search first for the smallest phase-side dominant compensator whose geometric carry contribution is just strong enough to keep the repaired second step inside the same closure-phase corridor.*

Proof. Immediate from Theorem H4.106. □

Definition H4.108 (Local lock corridor)**.** *A* local lock corridor *at stage n is a narrow admissible closure-phase band*

$$\mathfrak{C}^{\mathrm{loc}}_{\mathrm{lock}}(n)$$

containing the first near-trigger step and characterized by the following properties:

(a) every point in $\mathfrak{C}^{\mathrm{loc}}_{\mathrm{lock}}(n)$ lies inside the same local closure-phase window;

(b) the local burden pair remains trigger-near throughout the corridor;

(c) the Teichmüller-side distortion stays inside the active safe band;

(d) the corridor is narrow enough that leaving it produces a detectable phase-lock or carry defect.

Remark H4.109 (Geometric meaning of the lock corridor)**.** *The lock corridor is the thin local channel in which both the first near-trigger step and the repaired second step must lie if one is to read them as belonging to the same closure-phase passage. The current obstruction does not come from total loss of that channel, but from the failure of the second step to remain locked inside it.*

Definition H4.110 (HC-captured repaired second step)**.** *A repaired second step at stage n is called* HC-captured *if there exists an HC-admissible local Heisenberg compensator such that the compensated second step lies in the same local lock corridor as the first near-trigger step.*

Definition H4.111 (Lock-corridor leakage defect)**.** *The* lock-corridor leakage defect *at stage n is the residual amount by which the repaired second step fails to remain inside*

$$\mathfrak{C}^{\mathrm{loc}}_{\mathrm{lock}}(n).$$

It is denoted by

$$\lambda_{\mathrm{lock}}(n) \geq 0,$$

and vanishes exactly when the repaired second step is HC-captured.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition H4.112 (Minimal local lock-corridor compensation)**.** *A local HC final is called* minimally lock-corridor compensating *if it is HC-admissible and reduces the lock-corridor leakage defect to zero with no strictly smaller phase-side dominant correction achieving the same result on the active two-step local window.*

Proposition H4.113 (Current line still exhibits lock-corridor leakage)**.** *Under the present post-archival stage reading, the repaired second step still exhibits a nonzero lock-corridor leakage defect.*

Proof. The current line is phase-lock-prior on the Teichmüller side and has not yet produced a positive hybrid balancing witness. Hence the repaired second step is not yet fully locked into the same local closure-phase passage as the first near-trigger step, so the local lock-corridor leakage defect remains nonzero. □

Lemma H4.114 (Phase-side dominant HC yields lock-corridor capture)**.** *Assume:*

(a) a local lock corridor exists;

(b) the current HC final is phase-side dominant but geometrically carried;

(c) the compensator reduces the phase-lock defect to zero while keeping admissibility and the geometric carry inside the safe band.

Then the compensated second step is HC-captured.

Proof. By assumption, the compensator restores the phase-lock condition and preserves the geometric safe-band control. Since the local lock corridor is defined exactly as the admissible narrow closure-phase band carrying the first near-trigger step, the compensated second step now lies in the same corridor. Hence it is HC-captured. □

Theorem H4.115 (First explicit lock-corridor theorem)**.** *Assume:*

(a) the first HC-final type orientation theorem applies;

(b) the first phase-lock repair theorem applies;

(c) a local lock corridor exists around the first near-trigger step;

(d) there exists a minimally lock-corridor compensating HC final.

Then the current post-archival stage line admits a first explicit local lock-corridor reading. More precisely:

(i) the geometric problem is not to invent a new bulk path, but to return the repaired second step to the same narrow local closure-phase corridor as the first step;

(ii) the current HC task is to drive

$$\lambda_{\mathrm{lock}}(n) \to 0;$$

(iii) once the leakage defect vanishes under an HC-admissible phase-side dominant compensator, the repaired second step is locally synchronized in the same lock corridor as the first near-trigger step.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. By the HC-final orientation theorem, the first plausible compensator is phase-side dominant but geometrically carried. By the phase-lock repair theorem, the primary local repair target is vanishing of the phase-lock defect, with geometric carry remaining genuinely active. The local lock corridor isolates the thin admissible passage already occupied by the first near-trigger step. A minimally lock-corridor compensating HC final removes the leakage defect while preserving admissibility, so by Lemma H4.114 the repaired second step is returned to the same corridor. □

Corollary H4.116 (Current lock-corridor working rule)**.** *The next local audit should no longer ask only whether the second step is improved. It should ask whether an HC-admissible compensator has actually reduced*

$$\lambda_{\mathrm{lock}}(n)$$

to zero, i.e. whether the repaired second step has truly re-entered the same local lock corridor as the first near-trigger step.

Proof. Immediate from Theorem H4.115. □

Definition H4.117 (Lock-side leakage defect)**.** *A* lock-side leakage defect *at stage n is the portion of the local leakage defect attributable to failure of the repaired second step to satisfy the phase-lock condition under the current corridor calibration. It is denoted by*

$$\lambda^{\Theta}_{\mathrm{lock}}(n) \geq 0.$$

Definition H4.118 (Corridor-calibration leakage defect)**.** *A* corridor-calibration leakage defect *at stage n is the portion of the local leakage defect attributable to the active local lock corridor being too narrow or otherwise miscalibrated for the repaired second step, even after the primary phase-lock correction has been applied. It is denoted by*

$$\lambda^{\mathrm{cal}}_{\mathrm{lock}}(n) \geq 0.$$

Remark H4.119 (How the leakage splits)**.** *The total local lock-corridor leakage is now read as having two possible sources: a remaining failure of lock restoration itself, or a failure of the current corridor calibration to reflect the true admissible narrow passage. In symbols, one may think heuristically of*

$$\lambda_{\mathrm{lock}}(n) \sim \lambda^{\Theta}_{\mathrm{lock}}(n) + \lambda^{\mathrm{cal}}_{\mathrm{lock}}(n),$$

without yet forcing a rigid additive identity.

Definition H4.120 (Lock-side-prior local leakage regime)**.** *The local line is said to be in a* lock-side-prior local leakage regime *at stage n if*

$$\lambda^{\Theta}_{\mathrm{lock}}(n) > \lambda^{\mathrm{cal}}_{\mathrm{lock}}(n).$$

Definition H4.121 (Corridor-calibration-prior local leakage regime)**.** *The local line is said to be in a* corridor-calibration-prior local leakage regime *at stage n if*

$$\lambda^{\mathrm{cal}}_{\mathrm{lock}}(n) \geq \lambda^{\Theta}_{\mathrm{lock}}(n).$$

Proposition H4.122 (The current line is not corridor-calibration-prior by default)**.** *Under the present post-archival stage reading, the local line should not yet be read as corridor-calibration-prior by default.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. The current diagnosis remains phase-lock-prior on the Teichmüller side, and the first local repair target is still the vanishing of the phase-lock defect rather than widening or re-centering the lock corridor itself. Hence the default reading is not that the corridor is already known to be primarily miscalibrated, but rather that the repaired second step still fails first to lock into the active corridor. □

Proposition H4.123 (The current line is lock-side-prior on the leakage level)**.** *Under the present post-archival stage reading, the first remaining local lock-corridor leakage is lock-side-prior.*

Proof. By the first phase-lock repair theorem and the current phase-lock repair working rule, the next local task is still to force

$$\sigma_{\Theta}^{\mathrm{lock}}(n) = 0.$$

This means that the dominant unresolved leakage continues to arise from incomplete phase-lock restoration rather than from demonstrated over-narrowness or miscentering of the corridor itself. Therefore the first remaining local leakage is lock-side-prior. □

Theorem H4.124 (First lock-corridor leakage orientation theorem)**.** *Assume:*

(a) the first explicit lock-corridor theorem applies;

(b) the first phase-lock repair theorem applies;

(c) the current line still exhibits nonzero lock-corridor leakage.

Then the current post-archival stage line is lock-side-prior on the leakage level rather than corridor-calibration-prior.

More precisely:

(i) the next honest reduction target is first

$$\lambda_{\mathrm{lock}}^{\Theta}(n) \downarrow;$$

(ii) local corridor recalibration should be treated only as a secondary follow-on question unless the lock-side leakage has already been made negligible;

(iii) the present line should therefore first seek a stronger HC lock action before widening or re-centering the local corridor.

Proof. By the first explicit lock-corridor theorem, the active local task is to return the repaired second step to the same narrow corridor as the first near-trigger step. By the first phase-lock repair theorem, the primary local repair target remains vanishing of the phase-lock defect. Therefore, as long as nonzero leakage remains, the default reading is that the dominant leakage sits on the lock-restoration side rather than already on the calibration side of the corridor. Hence the line is lock-side-prior on the leakage level. □

Corollary H4.125 (Current leakage working rule)**.** *The next local audit should not first widen, thicken, or re-center the active lock corridor. It should first ask whether the remaining leakage is still produced by incomplete lock restoration of the repaired second step under the current corridor geometry.*

Proof. Immediate from Theorem H4.124. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition H4.126 (Minimal local HC-lock step)**.** *A* minimal local HC-lock step *at stage n is an HC-admissible local compensation*

$$\mathcal{H}_{\mathrm{lock}}^{\min}(n)$$

which acts on the repaired second step only through the smallest phase-side dominant correction required to reduce the lock-side leakage defect from

$$\lambda_{\mathrm{lock}}^{\Theta}(n) > 0$$

to

$$\lambda_{\mathrm{lock}}^{\Theta,+}(n) = 0,$$

while preserving the active local lock corridor geometry.

Definition H4.127 (Lock-neutral corridor preservation)**.** *A local HC-lock step is said to satisfy* lock-neutral corridor preservation *if it does not widen, thicken, or re-center the current local lock corridor*

$$\mathfrak{C}_{\mathrm{lock}}^{\mathrm{loc}}(n),$$

but only changes the repaired second-step placement inside that already fixed corridor geometry.

Definition H4.128 (HC-lock effectiveness defect)**.** *The* HC-lock effectiveness defect *of a local HC compensation at stage n is*

$$E_{\mathrm{HC}}^{\mathrm{lock}}(n) := \lambda_{\mathrm{lock}}^{\Theta,+}(n),$$

that is, the residual lock-side leakage after the local HC action has been applied.

Remark H4.129 (What the minimal HC step is allowed to do)**.** *The minimal HC-lock step is not allowed to solve the problem by changing the corridor itself. It must work only by re-locking the repaired second step inside the already accepted corridor. Geometrically, it is the smallest admissible local twist that makes the second point snap back into the same narrow channel without moving the channel walls.*

Definition H4.130 (HC-captured local repaired second step)**.** *A repaired second step at stage n is called* HC-captured *if there exists an HC-admissible local compensation satisfying lock-neutral corridor preservation such that*

$$\lambda_{\mathrm{lock}}^{\Theta,+}(n) = 0.$$

Proposition H4.131 (The current line calls first for a minimal HC-lock step)**.** *Under the current post-archival stage reading, the next local correction should first be sought as a minimal local HC-lock step rather than as a corridor recalibration.*

Proof. By the first lock-corridor leakage orientation theorem, the active leakage is lock-side-prior rather than corridor-calibration-prior. Therefore the first honest correction should try to eliminate the residual lock-side leakage while preserving the existing corridor geometry. This is exactly the role of a minimal local HC-lock step. □

Lemma H4.132 (Minimal HC-lock step implies HC capture)**.** *Assume a minimal local HC-lock step exists at stage n. Then the repaired second step is HC-captured at that stage.*

Proof. By definition, a minimal local HC-lock step is HC-admissible, preserves the corridor geometry, and reduces the residual lock-side leakage defect to zero. Hence the repaired second step satisfies the defining conditions of an HC-captured local repaired second step. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Theorem H4.133 (First minimal HC-lock step theorem)**.** *Assume:*

(a) the first local Theta-drift witness theorem applies;

(b) the first phase-lock repair theorem applies;

(c) the first explicit lock-corridor theorem applies;

(d) the first lock-corridor leakage orientation theorem applies;

(e) an HC-admissible local compensation menu is available.

Then the current post-archival stage line admits the following first local repair target:

(i) search first for a minimal local HC-lock step;

(ii) require lock-neutral corridor preservation;

(iii) accept the first final only if it yields

$$E_{\mathrm{HC}}^{\mathrm{lock}}(n) = 0;$$

(iv) treat any remaining failure first as lack of sufficient HC lock action rather than as corridor miscalibration.

Proof. Items (a)–(d) identify the current obstruction chain: the line is phase-lock-prior, the local corridor is already the right object to preserve, and the leakage is lock-side-prior rather than corridor-calibration-prior. Hence the first honest local HC task is not to redesign the corridor but to search within the available HC menu for the smallest admissible lock action that drives the residual lock-side leakage to zero while leaving the corridor fixed. This is exactly the asserted minimal HC-lock step search. □

Corollary H4.134 (Current minimal HC working rule)**.** *The next local audit should not first modify the local lock corridor. It should first scan the local HC menu for the smallest phase-side dominant compensator whose geometric carry is just sufficient to force*

$$\lambda_{\mathrm{lock}}^{\Theta,+}(n) = 0$$

under lock-neutral corridor preservation.

Proof. Immediate from Theorem H4.133. □

Definition H4.135 (Local HC final family)**.** *The* local HC final family *at stage n is a finite admissible menu*

$$\mathfrak{M}_{\mathrm{HC}}^{\mathrm{loc}}(n) = \{H_1^{(n)}, \dots, H_{N(n)}^{(n)}\}$$

of local Heisenberg-compensator actions acting on the repaired second step under the fixed local lock corridor.

Definition H4.136 (HC menu admissibility filter)**.** *A local HC final*

$$H_j^{(n)} \in \mathfrak{M}_{\mathrm{HC}}^{\mathrm{loc}}(n)$$

passes the HC menu admissibility filter *if:*

(a) it is phase-side dominant with genuinely active geometric carry;

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(b) it preserves the active local lock corridor;

(c) it preserves local admissibility on the same two-step window;

(d) it does not enlarge the corridor merely to hide the residual leakage.

Definition H4.137 (HC lock score)**.** *The* HC lock score *of an admissible local HC final is the pair*

$$\mathfrak{L}_{\mathrm{HC}}(H_j^{(n)}) := \big(\lambda_{\mathrm{lock}}^{\Theta,+}(n; H_j^{(n)}), C_{\mathrm{HC}}(H_j^{(n)})\big),$$

where the first component is the residual lock-side leakage after applying the final and the second component is its local compensator cost.

Definition H4.138 (Bombe-guided HC elimination)**.** *The* bombe-guided HC elimination *is the finite contradiction-sensitive elimination of all finals in*

$$\mathfrak{M}_{\mathrm{HC}}^{\mathrm{loc}}(n)$$

that fail the HC menu admissibility filter or leave a residual lock score incompatible with minimal HC capture.

Remark H4.139 (Why the HC scan is finite)**.** *The present line does not search over an arbitrary continuum of compensators. It searches only over the finite local HC menu already licensed by the current route-faithful readout state. This is why the bombe-guided elimination picture is appropriate here.*

Definition H4.140 (Minimal surviving HC final)**.** *A final*

$$H_{\min}^{(n)} \in \mathfrak{M}_{\mathrm{HC}}^{\mathrm{loc}}(n)$$

is called the minimal surviving HC final *if:*

(a) it survives bombe-guided HC elimination;

(b) it minimizes the residual lock-side leakage

$$\lambda_{\mathrm{lock}}^{\Theta,+}(n; H_j^{(n)});$$

(c) among all finals with the same minimal leakage, it minimizes the compensator cost

$$C_{\mathrm{HC}}(H_j^{(n)}).$$

Proposition H4.141 (The current line reduces the HC problem to a finite menu scan)**.** *Under the current post-archival stage reading, the first local HC problem is reduced to scanning the finite local HC final family under admissibility and lock-corridor preservation.*

Proof. By the current minimal HC working rule, the line must first search for a minimal local HC-lock step while preserving the active lock corridor. By the local Heisenberg-compensator theorem and the bombe-guided compensator menu reading, the admissible compensator actions are already organized as a finite local menu. Hence the first local HC problem is reduced to a finite menu scan with admissibility and corridor-preservation constraints. □

Lemma H4.142 (Minimal surviving HC final yields the first executable HC-lock step)**.** *Assume a minimal surviving HC final exists. Then it yields the first executable local HC-lock step under the current lock-side-prior regime.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. A minimal surviving HC final has already passed the admissibility and corridor-preservation filter. By minimality, it leaves no smaller admissible compensator with strictly lower residual lock-side leakage. Hence it is the first executable local HC-lock step compatible with the current regime. □

Theorem H4.143 (First local HC menu-scan theorem)**.** *Assume:*

(a) the first minimal HC-lock step theorem applies;

(b) the current line is lock-side-prior on the leakage level;

(c) the local HC final family is finite;

(d) bombe-guided HC elimination is available.

Then the next local HC task is a finite menu-scan problem.
More precisely:

(i) all inadmissible or corridor-breaking HC finals may be eliminated first;

(ii) the surviving finals are ranked by residual lock-side leakage and compensator cost;

(iii) if a minimal surviving HC final exists, it determines the first executable HC-lock step;

(iv) if no such final exists, the current line records a local HC-menu exhaustion obstruction rather than immediately blaming corridor calibration.

Proof. By Proposition H4.141, the present HC problem is already reduced to a finite final family with admissibility and corridor-preservation constraints. Bombe-guided HC elimination removes finals that fail these constraints. The remaining finals are compared by lock score. If a minimal surviving final exists, Lemma H4.142 yields the first executable HC-lock step. If not, the failure lies at the menu level itself, hence the correct reading is local HC-menu exhaustion rather than premature corridor redesign. □

Corollary H4.144 (Current HC menu-scan working rule)**.** *The next local audit should first enumerate the finite HC menu, eliminate inadmissible and corridor-breaking finals, and only then ask whether the surviving menu still contains a minimal lock-effective compensator.*

Proof. Immediate from Theorem H4.143. □

Definition H4.145 (Phase-side micro-lock HC final)**.** *A local HC final*

$$H^{(1)}_{\mathrm{lock}}(n) \in \mathfrak{M}^{\mathrm{loc}}_{\mathrm{HC}}(n)$$

is called a phase-side micro-lock HC final *if it satisfies:*

(a) phase-side dominance in the sense that its primary action falls on the residual lock component;

(b) lock-neutral corridor preservation;

(c) nonzero but minimal geometric carry sufficient to support the repaired second step inside the same local lock corridor;

(d) no extra crystal-side widening or recentering action.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Remark H4.146 (Why this is the first plausible HC menu entry)**.** *The present line is phase-lock-prior, lock-side-prior on the leakage level, and not carry-dominant. Hence the first plausible HC final is not a broad geometric carrier and not a corridor redesign. It is the smallest local phase-side micro-lock action whose geometric carry is only as large as needed to prevent renewed leakage.*

Definition H4.147 (HC micro-lock score)**.** *The* HC micro-lock score *of a phase-side micro-lock final*

$$H^{(1)}_{\mathrm{lock}}(n)$$

is the tuple

$$\mathfrak{L}^{\mathrm{micro}}_{\mathrm{HC}}(H^{(1)}_{\mathrm{lock}}(n)) := \big(\lambda^{\Theta,+}_{\mathrm{lock}}(n), c_{\mathrm{HC}}(H^{(1)}_{\mathrm{lock}}(n))\big),$$

ordered lexicographically by residual lock-side leakage first and compensator cost second.

Definition H4.148 (Plausible surviving HC final)**.** *A phase-side micro-lock HC final is called* plausibly surviving *if it passes the HC menu admissibility filter, preserves the local lock corridor, and yields no worse lock score than any other currently known phase-side dominant final.*

Proposition H4.149 (The current line favors a phase-side micro-lock final)**.** *Under the present post-archival stage reading, the first plausible HC menu entry is phase-side micro-lock rather than carry-dominant or fully two-component balanced.*

Proof. By the first HC-final type orientation theorem, the current line is phase-side dominant on the HC level and is neither carry-dominant nor already genuinely two-component balanced. By the first lock-corridor leakage orientation theorem, the residual leakage is lock-side-prior rather than corridor-calibration-prior. Therefore the first plausible HC entry is the smallest phase-side dominant final that closes the lock-side leakage while preserving the existing corridor. □

Lemma H4.150 (Plausible survival of the phase-side micro-lock final)**.** *Assume:*

(a) the local HC final family is finite;

(b) the current line is phase-side dominant on the HC level;

(c) lock-neutral corridor preservation is enforced.

Then the first plausible surviving HC final, if it exists at all, may be taken from the phase-side micro-lock class.

Proof. Because corridor redesign is excluded and carry-dominant finals are not favored by the current orientation, the admissible search is restricted first to phase-side dominant finals. Within that class, the lexicographic HC micro-lock score selects the finals with smallest residual leakage and then smallest cost. Hence any first plausible survivor, if one exists, may be taken from the phase-side micro-lock class. □

Theorem H4.151 (First plausible HC menu-entry theorem)**.** *Assume:*

(a) the first HC-final type orientation theorem applies;

(b) the first lock-corridor leakage orientation theorem applies;

(c) the first local HC menu-scan theorem applies.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Then the first plausible HC menu entry is a phase-side micro-lock HC final.
More precisely:

(i) the final is sought first in the phase-side dominant subclass;

(ii) its geometric carry is only auxiliary but genuinely active;

(iii) its admissibility is tested under lock-neutral corridor preservation;

(iv) if a first plausible survivor exists, it determines the leading local HC hypothesis before any broader compensator class is tried.

Proof. Items (a) and (b) imply that the line is not looking first for a carry-dominant or fully balanced HC final. Item (c) reduces the search to the finite local HC menu with corridor-preserving admissibility. By Lemma H4.150, any first plausible survivor may be taken from the phase-side micro-lock class. This yields the asserted first plausible HC menu entry. □

Corollary H4.152 (Current HC entry working rule)**.** *The next local audit should first test the phase-side micro-lock HC finals in the finite menu, ranked by residual lock-side leakage and cost, before considering broader compensator types.*

Proof. Immediate from Theorem H4.151. □

Definition H4.153 (Micro-lock update rule)**.** *Let*

$$H^{(1)}_{\mathrm{lock}}(n)$$

be a phase-side micro-lock HC final acting on the repaired second local step. The micro-lock update rule *is the local update*

$$X^{\mathrm{rep}}_{n+1} \longmapsto X^{\mathrm{lock}}_{n+1}$$

induced by $H^{(1)}_{\mathrm{lock}}(n)$ under the following constraints:

(a) lock-neutral corridor preservation is maintained;

(b) the updated second step stays inside the same local admissibility band;

(c) the lock-side leakage does not increase;

(d) the phase-side component of the update is dominant, while the geometric carry remains auxiliary but genuinely active.

Definition H4.154 (Successful micro-lock update)**.** *A micro-lock update is called* successful *if it satisfies*

$$\lambda^{\Theta,+}_{\mathrm{lock}}(n) = 0$$

while preserving the current local lock corridor and the active admissibility constraints.

Definition H4.155 (HC-captured local continuation)**.** *A local continuation after stage n is called* HC-captured *if it is produced by a successful micro-lock update and the resulting repaired second step remains in the same local lock corridor as the first near-trigger step.*

Remark H4.156 (Why capture is the right next target)**.** *The present line no longer needs a broad compensator search. It needs the first concrete capture event in which the second repaired step is actually brought back into the same narrow local corridor. That is the smallest nontrivial success criterion beyond plausible HC menu survival.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Lemma H4.157 (Minimal HC-lock step yields local capture)**.** *Assume:*

(a) a minimal surviving HC final exists in the local menu;

(b) its associated micro-lock update is successful;

(c) lock-neutral corridor preservation is enforced.

Then the updated repaired second step is HC-captured.

Proof. By success of the micro-lock update, the residual lock-side leakage is driven to zero. By lock-neutral corridor preservation, the active corridor is not widened or re-centered during the update. Hence the updated second step lies in the same local lock corridor as the first near-trigger step, which is exactly HC capture. □

Definition H4.158 (First local HC-capture chain)**.** *A* first local HC-capture chain *is a finite local sequence*

$$X_n^{\mathrm{near}} \to X_{n+1}^{\mathrm{rep}} \to X_{n+1}^{\mathrm{lock}}$$

consisting of:

(i) a near-trigger first step;

(ii) a repaired but not yet fully locked second step;

(iii) a successful micro-lock update producing an HC-captured continuation.

Proposition H4.159 (Current local line reduces to HC-capture chain search)**.** *Under the present post-archival stage reading, once the local corridor, the phase-lock repair target, and the HC menu scan are fixed, the next local task reduces to the search for a first local HC-capture chain.*

Proof. The preceding line already fixes: the right corridor object, the lock-side-prior leakage orientation, the minimal HC-lock target, and the first plausible HC final family. No broader redesign is currently preferred. Therefore the next local task is exactly to determine whether one of the plausible HC finals yields an actual capture of the repaired second step. □

Theorem H4.160 (First local HC-capture chain theorem)**.** *Assume:*

(a) the first explicit lock-corridor theorem applies;

(b) the first minimal HC-lock step theorem applies;

(c) the first plausible HC menu-entry theorem applies;

(d) a successful micro-lock update exists for the minimal surviving HC final.

Then the current post-archival stage line supports a first local HC-capture chain.
More precisely:

(i) the repaired second step is first updated by a phase-side dominant, geometrically carried micro-lock compensator;

(ii) the lock-side leakage is removed without redesigning the corridor;

(iii) the resulting continuation is HC-captured in the same local lock corridor as the first near-trigger step.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. By the first plausible HC menu-entry theorem, the first plausible surviving HC final is sought in the phase-side micro-lock class. By the first minimal HC-lock step theorem, the active local HC task is precisely to search for the smallest such compensator that removes residual lock-side leakage while preserving the corridor. If such a micro-lock update is successful, Lemma H4.157 yields HC capture. This is exactly the stated HC-capture chain. □

Corollary H4.161 (Current HC-capture working rule)**.** *The next local audit should no longer stop at plausible HC menu survival. It should ask whether the best phase-side micro-lock final actually yields a successful micro-lock update and hence the first local HC-capture chain.*

Proof. Immediate from Theorem H4.160. □

Corollary H4.162 (Freeze threshold after archival stage)**.** *If the local line is stably organized by*

lock corridor → minimal HC-lock step → HC menu scan → first local HC-capture chain,

then the post-archival stage line has reached a natural freeze threshold for a new minor release-final.

Proof. At that point the local HC line is no longer merely diagnostic. It carries a coherent operational chain from the residual lock defect to the first concrete capture event. This is exactly the kind of self-contained new kernel that supports a new freeze final before audit. □

H4.1 Transport-stable HC capture and OP-entry handoff

Definition H4.163 (Transport-stable HC-capture chain)**.** *Let*

$$\mathcal{C}_n^{\mathrm{HC}} := \big(X_n^{\mathrm{near}} \to X_{n+1}^{\mathrm{rep}} \to X_{n+1}^{\mathrm{lock}}\big)$$

be a first local HC-capture chain in the sense of Definition H4.158. It is called transport-stable on an admissible window $[n, n+k]$ *if every route-faithful admissible transport step*

$$\mathsf{T}_j : X_j \longmapsto X_{j+1}, \qquad n \le j < n+k,$$

including the symmetrized forward/backward transport readout when that readout is active, preserves:

(a) membership of the locked continuation in the transported lock corridor;

(b) non-increase of the lock-side leakage under the transported HC readout;

(c) the inherited admissibility and boundary-stop constraints of the active route;

(d) the distinction between horizon-side and stop-preserving boundary-side packets.

Thus transport-stability is not a new closure theorem. It is the statement that the local capture event survives the already-admissible route motion needed before OP-facing use.

Definition H4.164 (HC transport defect)**.** *For a captured local chain $\mathcal{C}_n^{\mathrm{HC}}$ and an admissible transport step T_j, define the associated HC transport defect schematically by*

$$\Delta_{\mathrm{tr}}^{\mathrm{HC}}(j) := \max\Big\{ d_{\mathfrak{C}}\big(\mathsf{T}_j X_j^{\mathrm{lock}}, \mathfrak{C}_{\mathrm{lock}}^{\mathrm{loc}}(j+1)\big), \big(\lambda_{\mathrm{lock}}^{\Theta,+}(j+1) - \lambda_{\mathrm{lock}}^{\Theta,+}(j)\big)_+, \chi_{\mathrm{stop}}(j+1) \Big\}.$$

Here $d_{\mathfrak{C}}$ denotes corridor distance in the current local readout, $(\cdot)_+$ denotes the positive part, and χ_{stop} is the indicator of an inadmissible boundary-stop violation. The defect vanishes exactly when the transported step remains corridor-inside, does not re-open lock leakage, and does not violate the inherited stop discipline.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Lemma H4.165 (Zero HC transport defect preserves capture). *Assume that $\mathcal{C}_n^{\mathrm{HC}}$ is a first local HC-capture chain and that*

$$\Delta_{\mathrm{tr}}^{\mathrm{HC}}(j) = 0$$

for an admissible route-faithful transport step T_j. Then the transported locked state is again HC-captured in the transported local lock corridor.

Proof. The equality $\Delta_{\mathrm{tr}}^{\mathrm{HC}}(j) = 0$ simultaneously says that the transported locked state has zero corridor distance from the next transported lock corridor, that lock-side leakage has not been positively re-opened, and that no boundary-stop violation has occurred. These are exactly the transported versions of corridor membership, lock-side capture, and route admissibility in Definition H4.163. Hence the transported locked state remains HC-captured. □

Definition H4.166 (HC-to-OP handoff packet). *An* HC-to-OP handoff packet *on $[n, n+k]$ is a tuple*

$$\mathcal{P}_{\mathrm{HC}\to\mathrm{OP}}(n,k) := \big(\mathcal{C}_n^{\mathrm{HC}}, \{\mathsf{T}_j\}_{j=n}^{n+k-1}, \{\Delta_{\mathrm{tr}}^{\mathrm{HC}}(j)\}_{j=n}^{n+k-1}, \widetilde{\mathbf{R}}_{\mathrm{live}}(n+k), \tau_{\mathrm{route}}\big),$$

where:

(a) $\mathcal{C}_n^{\mathrm{HC}}$ is a first local HC-capture chain;

(b) the transports T_j are route-faithful and admissible;

(c) all displayed HC transport defects vanish on the window;

(d) the terminal normalized live residual $\widetilde{\mathbf{R}}_{\mathrm{live}}(n+k)$ lies in the admissible OP-entry normalization regime;

(e) τ_{route} records whether the packet is horizon-side or stop-preserving boundary-side.

The handoff packet is therefore the smallest object that can be consumed by the OP-facing machine without forgetting how the local HC capture was transported.

Proposition H4.167 (Handoff packets are admissible pre-OP frontier objects). *Every HC-to-OP handoff packet is an admissible pre-OP frontier object in the sense of the current formal OP-entry framework.*

Proof. By definition, the packet contains a valid local HC-capture chain, an admissible route-faithful transport family, vanishing HC transport defects, an admissibly normalized terminal live residual, and an explicit route tag. These are precisely the data required to prevent three forbidden moves: erasing the route distinction, re-opening a boundary stop, or treating local capture as OP closure without transport. Hence the packet is admissible as a pre-OP frontier object. □

Theorem H4.168 (First transport-stable HC-capture to OP-entry criterion). *Assume:*

(a) the first local HC-capture chain theorem applies;

(b) the captured chain is transport-stable on a finite admissible window $[n, n+k]$;

(c) the terminal normalized live residual $\widetilde{\mathbf{R}}_{\mathrm{live}}(n+k)$ lies in the admissible OP-entry normalization regime;

(d) the inherited route tag is preserved throughout the handoff.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Then the current post-archival stage line supports a controlled OP-entry handoff packet

$$\mathcal{P}_{\mathrm{HC}\to\mathrm{OP}}(n,k).$$

If, in addition, the already-defined closure-phase trigger persistence condition holds at the terminal state, the packet yields an immediate OP-entry certificate. If that persistence condition is not yet met, the same packet yields a narrowed frontier-certified pre-entry state rather than a false OP closure claim.

Proof. By the first local HC-capture chain theorem, the local repaired second step can be locked into the same local corridor as the first near-trigger step. By transport-stability, Lemma H4.165 applies at each admissible transport step in the finite window. Therefore the captured state remains captured after the finite route-faithful transport sequence, with no positive re-opening of lock-side leakage and no boundary-stop violation. The normalized terminal residual and route tag then supply exactly the remaining fields in Definition H4.166. The final alternative follows from the existing OP-entry framework: trigger persistence upgrades the handoff packet to an entry certificate, while absence of trigger persistence leaves a legitimate, narrowed pre-entry frontier state. □

Corollary H4.169 (working rule)**.** *The next audit should no longer ask merely whether a local HC capture chain exists. It should test whether the captured chain is transport-stable and whether the resulting handoff packet terminates in an immediate OP-entry certificate or in a smaller frontier-certified pre-entry state.*

Proof. Immediate from Theorem H4.168. □

H4.2 First OP-entry normal form extracted from HC handoff

Definition H4.170 (HC-extracted OP-entry normal form)**.** *Let $\mathcal{P}_{\mathrm{HC}\to\mathrm{OP}}(n,k)$ be an HC-to-OP handoff packet in the sense of Definition H4.166. Its* HC-extracted OP-entry normal form *is the tuple*

$$\mathcal{N}_{\mathrm{OP}}^{\mathrm{HC}}(n,k) := \big(\widetilde{\mathbf{R}}_{\mathrm{live}}^{\mathrm{OP}}(n+k), \tau_{\mathrm{route}}, \sigma_{\mathrm{tr}}, \mathcal{M}_{\mathrm{OP}}^{\mathrm{adm}}(n+k), \eta_{\mathrm{entry}}\big),$$

where:

(a) $\widetilde{\mathbf{R}}_{\mathrm{live}}^{\mathrm{OP}}(n+k)$ *is the terminal normalized live residual read as an OP-facing residual triple*

$$\widetilde{\mathbf{R}}_{\mathrm{live}}^{\mathrm{OP}} = \big(\rho_{\mathrm{CL}}, \rho_{\Theta R}, \rho_A\big);$$

(b) τ_{route} *is the inherited horizon-side or stop-preserving boundary-side route tag;*

(c) σ_{tr} *records that the HC transport defects vanish on the handoff window;*

(d) $\mathcal{M}_{\mathrm{OP}}^{\mathrm{adm}}(n+k)$ *is the finite admissible OP-facing request menu available at the terminal state;*

(e) $\eta_{\mathrm{entry}} \in \{\mathrm{entry}, \mathrm{frontier}\}$ *records whether the terminal closure-phase trigger persistence condition has already upgraded the packet to immediate OP-entry or only to a smaller frontier-certified pre-entry state.*

The normal form is therefore not a new theorem layer. It is the smallest downstream-readable form of a transported HC capture event.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition H4.171 (Residual routing out of the OP-entry normal form)**.** *The residual routing map associated with $\mathcal{N}_{\mathrm{OP}}^{\mathrm{HC}}$ is the finite assignment*

$$\mathsf{Route}_{\mathrm{OP}}\big(\rho_{\mathrm{CL}}, \rho_{\Theta R}, \rho_A; \tau_{\mathrm{route}}\big) = \big(\mathcal{Q}_{1/4}, \mathcal{Q}_{5(B)}, \mathcal{Q}_A\big),$$

where:

(a) $\mathcal{Q}_{1/4}$ *contains only continuum-limit / Hopf-correction requests controlled by* ρ_{CL} *and is the admissible intake for the later OP 1/OP 4 correction line;*

(b) $\mathcal{Q}_{5(B)}$ *contains only routed* Θ*–*R *balance requests controlled by* $\rho_{\Theta R}$ *and is the admissible intake for the later OP 5(B) line;*

(c) $\mathcal{Q}_A$ *contains only anchor, admissibility, and route-discipline requests controlled by* ρ_A*;*

(d) no request may erase τ_{route}*, deny the earliest-boundary stop, or promote a diagnostic residual to a theorematic closure certificate by naming alone.*

Lemma H4.172 (Normal-form extraction preserves route discipline)**.** *If $\mathcal{P}_{\mathrm{HC}\to\mathrm{OP}}(n,k)$ is an HC-to-OP handoff packet, then its HC-extracted OP-entry normal form preserves the inherited route distinction and boundary-stop discipline.*

Proof. The handoff packet already contains the route tag, the transported HC-capture chain, the vanishing HC transport defects, and the terminal normalized live residual. Definition H4.170 only repackages these data into the fields consumed by the OP-facing request layer. Since neither the route tag nor the stop indicator is discarded during that repackaging, the inherited route distinction and boundary-stop discipline are preserved. □

Proposition H4.173 (Finite OP-facing request menu from HC handoff)**.** *Every HC-extracted OP-entry normal form determines a finite admissible OP-facing request menu*

$$\mathcal{M}_{\mathrm{OP}}^{\mathrm{adm}} = \mathcal{Q}_{1/4} \cup \mathcal{Q}_{5(B)} \cup \mathcal{Q}_A$$

subordinate to the inherited route tag.

Proof. The terminal live residual triple has only three displayed coordinates: continuum-limit, routed Θ*–*R, and anchor/admissibility. Definition H4.171 assigns each coordinate to one finite request family and forbids requests that break route discipline or manufacture a closure certificate. Hence the union is finite, admissible, and subordinate to the inherited route tag. □

Definition H4.174 (Lead OP-entry residual selector)**.** *A* lead OP-entry residual selector *is any admissible rule*

$$\mathsf{L}_{\mathrm{OP}}\big(\mathcal{N}_{\mathrm{OP}}^{\mathrm{HC}}\big) \in \{\rho_{\mathrm{CL}}, \rho_{\Theta R}, \rho_A, \rho_{\mathrm{CL}} \oplus \rho_{\Theta R}\}$$

that chooses the next finite OP-facing work branch according to the current residual ranking, subject to the following restrictions:

(a) if ρ_{CL} *is dominant, the next branch is the OP 1/OP 4 continuum-limit correction line;*

(b) if $\rho_{\Theta R}$ *is dominant, the next branch is the OP 5(B) routed balance line;*

(c) if ρ_A *is dominant, the next branch is an admissibility/anchor refinement rather than a physical closure claim;*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(d) if ρ_{CL} and $\rho_{\Theta R}$ are inseparable under the current ranking, the next branch is the paired OP 1/4–OP 5(B) comparison branch.

Theorem H4.175 (First OP-entry normal-form theorem)**.** *Assume the first transport-stable HC-capture to OP-entry criterion of Theorem H4.168. Then the current archival stage working line admits a unique downstream-readable OP-entry normal form*

$$\mathcal{N}_{\mathrm{OP}}^{\mathrm{HC}}(n,k)$$

for the transported HC capture event. Moreover:

(i) if $\eta_{\mathrm{entry}} = \mathrm{entry}$, the normal form is an immediate OP-entry surface;

(ii) if $\eta_{\mathrm{entry}} = \mathrm{frontier}$, the normal form is a smaller frontier-certified pre-entry surface;

(iii) in both cases the admissible next work is restricted to the residual-routed menu $\mathcal{Q}_{1/4}$, $\mathcal{Q}_{5(B)}$, and $\mathcal{Q}_A$.

Proof. Theorem H4.168 supplies the controlled handoff packet or, if trigger persistence is absent, the narrowed frontier-certified pre-entry state carried by the same packet. Definition H4.170 reads exactly those fields into one terminal residual triple, one route tag, one transport-stability record, one admissible request menu, and one entry/frontier flag. No additional choice is made during this extraction, so the downstream-readable normal form is unique relative to the given handoff packet. The final restriction follows from Proposition H4.173. □

Corollary H4.176 (working rule)**.** *The next mathematical audit should select the lead OP-entry residual from $\mathcal{N}_{\mathrm{OP}}^{\mathrm{HC}}$. If the continuum-limit residual dominates, start with the OP 1/OP 4 correction branch. If the routed Θ–R residual dominates, start with OP 5(B). If neither is dominant, use the paired comparison branch rather than forcing a premature single-problem split.*

Proof. Immediate from Definition H4.174 and Theorem H4.175. □

H4.3 First OP-entry task routing extracted from the normal form

Definition H4.177 (OP-entry task-routing object)**.** *Let*

$$\mathcal{N}_{\mathrm{OP}}^{\mathrm{HC}}(n,k)$$

be an HC-extracted OP-entry normal form in the sense of Definition H4.170, and let

$$\mathsf{L}_{\mathrm{OP}}\big(\mathcal{N}_{\mathrm{OP}}^{\mathrm{HC}}\big)$$

be a lead OP-entry residual selector in the sense of Definition H4.174. The associated OP-entry task-routing object *is the tuple*

$$\mathcal{T}_{\mathrm{OP}}^{\mathrm{HC}}(n,k) := \big(\mathcal{N}_{\mathrm{OP}}^{\mathrm{HC}}(n,k), \mathsf{L}_{\mathrm{OP}}, q_{\mathrm{lead}}, \mathcal{W}_{\mathrm{OP}}^{\mathrm{lead}}, \tau_{\mathrm{route}}, \eta_{\mathrm{entry}}\big),$$

where:

(a) q_{lead} is the selected residual branch in

$$\{\rho_{\mathrm{CL}}, \rho_{\Theta R}, \rho_A, \rho_{\mathrm{CL}} \oplus \rho_{\Theta R}\};$$

(b) $\mathcal{W}_{\mathrm{OP}}^{\mathrm{lead}}$ is the finite route-faithful work family assigned to that branch;

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(c) τ_{route} *is inherited unchanged from the normal form;*

(d) η_{entry} *records whether the object is consumed as immediate OP-entry or only as a frontier-certified sharpening task.*

Thus the task-routing object does not decide an OP claim. It decides only which finite admissible work family is allowed to be executed next.

Definition H4.178 (Residual-to-task routing map)**.** *The* residual-to-task routing map *associated with an OP-entry normal form is the admissible assignment*

$$\mathfrak{R}_{\mathrm{task}} : \{\rho_{\mathrm{CL}}, \rho_{\Theta R}, \rho_A, \rho_{\mathrm{CL}} \oplus \rho_{\Theta R}\} \longrightarrow \{\mathcal{W}^{\mathrm{CL}}_{1/4}, \mathcal{W}^{\Theta R}_{5(B)}, \mathcal{W}^{\mathrm{adm}}_{A}, \mathcal{W}^{\mathrm{cmp}}_{1/4|5(B)}\}$$

defined by

$$\begin{aligned} \rho_{\mathrm{CL}} &\longmapsto \mathcal{W}^{\mathrm{CL}}_{1/4}, \\ \rho_{\Theta R} &\longmapsto \mathcal{W}^{\Theta R}_{5(B)}, \\ \rho_A &\longmapsto \mathcal{W}^{\mathrm{adm}}_{A}, \\ \rho_{\mathrm{CL}} \oplus \rho_{\Theta R} &\longmapsto \mathcal{W}^{\mathrm{cmp}}_{1/4|5(B)}. \end{aligned}$$

Here $\mathcal{W}^{\mathrm{CL}}_{1/4}$ *contains only continuum-limit correction tasks for OP 1/OP 4,* $\mathcal{W}^{\Theta R}_{5(B)}$ *contains only routed* Θ*–*R *balance tasks for OP 5(B),* $\mathcal{W}^{\mathrm{adm}}_{A}$ *contains only anchor/admissibility refinement tasks, and* $\mathcal{W}^{\mathrm{cmp}}_{1/4|5(B)}$ *contains paired comparison tasks that keep the common HC-capture origin visible when the two leading residuals cannot yet be separated.*

Lemma H4.179 (Task routing preserves OP-entry normal-form discipline)**.** *Let* $\mathcal{T}^{\mathrm{HC}}_{\mathrm{OP}}(n,k)$ *be an OP-entry task-routing object. Then executing its selected finite work family cannot erase the inherited route tag, cannot remove the boundary-stop discipline, and cannot manufacture a new OP closure certificate.*

Proof. By Definition H4.177, the task-routing object carries the normal form, the selected residual branch, the finite work family, the inherited route tag, and the entry/frontier flag as explicit fields. By Definition H4.178, each work family is subordinate to one already displayed residual coordinate or to the paired comparison of the two leading displayed coordinates. None of these assignments changes the route tag or the entry/frontier flag. Hence task execution may refine, compare, or package the selected branch, but it cannot promote itself to a closure certificate unless the pre-existing OP-entry framework supplies the missing trigger persistence. □

Proposition H4.180 (Paired comparison branch is the safe non-dominant branch)**.** *If the current residual ranking cannot separate* ρ_{CL} *from* $\rho_{\Theta R}$*, the admissible next branch is the paired comparison family*

$$\mathcal{W}^{\mathrm{cmp}}_{1/4|5(B)}$$

rather than a forced single OP 1/OP 4 or OP 5(B) branch.

Proof. Definition H4.174 already allows the paired symbol $\rho_{\mathrm{CL}} \oplus \rho_{\Theta R}$ exactly when the two residuals are inseparable under the current ranking. Definition H4.178 sends that symbol to the paired comparison family. Choosing a single branch in that case would add information not present in the normal form and could hide the common HC-capture origin of both residuals. The paired branch is therefore the route-faithful and conservative choice. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Theorem H4.181 (First OP-entry task-routing theorem)**.** *Assume that an HC-to-OP handoff packet has been converted into the HC-extracted OP-entry normal form*

$$\mathcal{N}_{\mathrm{OP}}^{\mathrm{HC}}(n,k)$$

of Theorem H4.175. Then every admissible lead selector L_{OP} *determines a finite route-faithful OP-entry task-routing object*

$$\mathcal{T}_{\mathrm{OP}}^{\mathrm{HC}}(n,k).$$

Moreover:

(i) *if* $\eta_{\mathrm{entry}} = \mathrm{entry}$*, the selected family is a first admissible OP-facing work packet;*

(ii) *if* $\eta_{\mathrm{entry}} = \mathrm{frontier}$*, the selected family is only a finite sharpening/comparison packet for the smaller frontier-certified pre-entry state;*

(iii) *if the selector returns* $\rho_{\mathrm{CL}} \oplus \rho_{\Theta R}$*, the next work packet must keep OP 1/OP 4 and OP 5(B) in paired comparison until a later ranking separates them.*

Proof. The normal form contains a finite residual triple, a finite admissible request menu, a route tag, a transport-stability record, and an entry/frontier flag. The lead selector chooses one of the explicitly allowed residual branches. Applying Definition H4.178 to that branch yields one finite work family, and adjoining the inherited normal-form fields gives the task-routing object of Definition H4.177. Lemma H4.179 gives the route and certificate restrictions. Proposition H4.180 gives the paired-branch rule when the two leading residuals are not yet separated. □

Corollary H4.182 (working rule)**.** *The next OP-facing audit should not yet try to close OP 1/OP 4 or OP 5(B) directly. It should first instantiate the task-routing object* $\mathcal{T}_{\mathrm{OP}}^{\mathrm{HC}}(n,k)$*, rank the displayed residuals, and execute only the selected finite work family:* $\mathcal{W}_{1/4}^{\mathrm{CL}}$*,* $\mathcal{W}_{5(B)}^{\Theta R}$*,* $\mathcal{W}_{A}^{\mathrm{adm}}$*, or the paired comparison family* $\mathcal{W}_{1/4|5(B)}^{\mathrm{cmp}}$*.*

Proof. Immediate from Theorem H4.181. The task-routing object is the conservative bridge between OP-entry normal form and explicit OP-facing execution. □

H4.4 First paired OP-entry comparison execution layer

Definition H4.183 (Paired OP-entry comparison execution packet)**.** *Assume that the OP-entry task-routing object of Definition H4.177 selects the paired branch*

$$q_{\mathrm{lead}} = \rho_{\mathrm{CL}} \oplus \rho_{\Theta R}, \qquad \mathcal{W}_{\mathrm{OP}}^{\mathrm{lead}} = \mathcal{W}_{1/4|5(B)}^{\mathrm{cmp}}.$$

A paired OP-entry comparison execution packet *is the tuple*

$$\mathcal{X}_{1/4|5(B)}^{\mathrm{cmp}}(n,k) := \big(\mathcal{T}_{\mathrm{OP}}^{\mathrm{HC}}(n,k), \mathcal{U}_{\mathrm{HC}}^{\mathrm{adm}}, D_{\mathrm{joint}}^{\mathrm{HC}}, D_{\mathrm{sep}}^{\mathrm{HC}}, \Delta_{\mathrm{sep}}^{\mathrm{HC}}, \omega_{\mathrm{cmp}}\big),$$

where:

(a) $\mathcal{T}_{\mathrm{OP}}^{\mathrm{HC}}(n,k)$ *is the inherited task-routing object;*

(b) $\mathcal{U}_{\mathrm{HC}}^{\mathrm{adm}}$ *is the finite admissible menu of HC-compatible micro-updates available at the terminal normal form;*

(c) $D_{\mathrm{joint}}^{\mathrm{HC}}$ *is the best joint readout error obtained when one and the same admissible HC-compatible update is asked to account for both* ρ_{CL} *and* $\rho_{\Theta R}$*;*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(d) $D_{\mathrm{sep}}^{\mathrm{HC}}$ *is the best separated readout error obtained when the two coordinates are allowed to choose their admissible update witnesses independently;*

(e)

$$\Delta_{\mathrm{sep}}^{\mathrm{HC}} := D_{\mathrm{joint}}^{\mathrm{HC}} - D_{\mathrm{sep}}^{\mathrm{HC}} \geq 0$$

is the HC-origin separation defect;

(f) $\omega_{\mathrm{cmp}} \in \{\mathrm{bind}, \mathrm{split}, \mathrm{frontier}\}$ *records the comparison outcome.*

The packet executes only the comparison branch. It does not yet assert an OP 1/4 correction theorem and does not yet assert the routed OP 5(B) balance theorem.

Definition H4.184 (HC-origin comparison decision rule)**.** *Fix an admissible comparison tolerance* $\varepsilon_{\mathrm{cmp}} > 0$ *and a strict descent requirement on the active frontier measure. The outcome field of Definition H4.183 is read by*

$$\omega_{\mathrm{cmp}} = \begin{cases} \mathrm{bind}, & \Delta_{\mathrm{sep}}^{\mathrm{HC}} \leq \varepsilon_{\mathrm{cmp}}, \\ \mathrm{split}, & \Delta_{\mathrm{sep}}^{\mathrm{HC}} > \varepsilon_{\mathrm{cmp}} \text{ and the separated ranking is stable}, \\ \mathrm{frontier}, & \text{otherwise.} \end{cases}$$

The bind case means that the two residual coordinates are still best treated as one common HC-origin comparison problem. The split case means that the comparison has supplied enough information to return to a single OP 1/4 or OP 5(B) branch under the residual ranking. The frontier case means that the comparison improved the state, but not enough to justify either binding or splitting as a theorematic route decision.

Lemma H4.185 (Paired comparison execution preserves the inherited handoff discipline)**.** *Executing a paired OP-entry comparison packet cannot erase the inherited route tag, cannot reopen the boundary-stop condition, and cannot promote the current frontier state to an OP closure certificate.*

Proof. The execution packet contains the task-routing object as an explicit first component. By Lemma H4.179, that object already carries the inherited route tag, entry/frontier flag, and admissible work-family restriction. The additional fields in Definition H4.183 compare only joint versus separated HC-compatible readout errors inside the finite admissible micro-update menu. They do not change the route tag, do not delete the boundary-stop flag, and do not introduce a trigger-persistence certificate. Hence the comparison can only bind, split, or sharpen the frontier; it cannot close an OP by itself. □

Proposition H4.186 (Bind branch retains the common HC-capture origin)**.** *If* $\omega_{\mathrm{cmp}} = \mathrm{bind}$*, then OP 1/4 and OP 5(B) must remain in the coupled work family*

$$\mathcal{W}_{1/4|5(B)}^{\mathrm{cmp}}$$

for the next step, with the common HC-capture origin kept as an explicit field.

Proof. In the bind case, Definition H4.184 says that the best joint HC-compatible readout is not worse than the separated readout by more than the admissible comparison tolerance. Splitting the branch at that point would artificially discard the information that one HC-compatible source still accounts for both leading residuals. The route-faithful next object therefore remains the paired comparison family, now with the common HC-origin field recorded rather than hidden. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proposition H4.187 (Split branch returns to ranked single-branch work)**.** *If* $\omega_{\mathrm{cmp}} = \mathrm{split}$*, then the comparison packet returns a ranked single-branch task-routing object selecting either*

$$\mathcal{W}^{\mathrm{CL}}_{1/4} \qquad \textit{or} \qquad \mathcal{W}^{\Theta R}_{5(B)},$$

according to the stable separated residual ranking.

Proof. The split case is allowed only when the HC-origin separation defect exceeds the admissible comparison tolerance and the separated ranking is stable. Thus the normal form now contains information that was absent in the inseparable branch: one of the two residual coordinates has become a legitimate lead branch. Applying the residual-to-task routing map of Definition H4.178 to that newly ranked coordinate gives either the OP 1/4 continuum-limit work family or the OP 5(B) routed Θ–R work family. □

Theorem H4.188 (First paired OP-entry comparison execution theorem)**.** *Assume that the task-routing layer selects the paired comparison branch*

$$\rho_{\mathrm{CL}} \oplus \rho_{\Theta R} \longmapsto \mathcal{W}^{\mathrm{cmp}}_{1/4|5(B)}.$$

Then every admissible finite HC-compatible comparison menu produces one paired OP-entry comparison execution packet

$$\mathcal{X}^{\mathrm{cmp}}_{1/4|5(B)}(n,k)$$

and exactly one of the following route-faithful next states:

(i) a bound common-origin comparison state, still subordinate to $\mathcal{W}^{\mathrm{cmp}}_{1/4|5(B)}$*;*

(ii) a split ranked single-branch state, subordinate to either $\mathcal{W}^{\mathrm{CL}}_{1/4}$ *or* $\mathcal{W}^{\Theta R}_{5(B)}$*;*

(iii) a smaller frontier-certified comparison state, if neither binding nor splitting is yet justified.

In all three cases, no OP closure certificate is manufactured by the comparison execution itself.

Proof. The finite admissible menu $\mathcal{U}^{\mathrm{adm}}_{\mathrm{HC}}$ supplies the joint and separated minimization tests by construction. Their difference defines $\Delta^{\mathrm{HC}}_{\mathrm{sep}} \geq 0$, since the separated readout is at least as flexible as the joint readout. Definition H4.184 assigns exactly one of the three comparison outcomes. Proposition H4.186 gives the bound next state; Proposition H4.187 gives the split next state; and the remaining case is, by definition, the smaller frontier-certified comparison state. Lemma H4.185 supplies the route, boundary, and no-false-closure restrictions. □

Corollary H4.189 (working rule)**.** *The next OP-facing audit may now execute the paired comparison branch before choosing between OP 1/4 and OP 5(B). The correct next question is not yet* which OP closes first*, but whether* ρ_{CL} *and* $\rho_{\Theta R}$ *are still one common HC-origin residual or have separated into stable ranked branches.*

Proof. Immediate from Theorem H4.188. □

H4.5 Minimal MML/MMA interface for HC-routed OP execution

Definition H4.190 (local language-version registry)**.** *The current HC-routed OP-entry execution layer uses the following local interface tags; their interface stamps are recorded in prose, not as mathematical exponents:*

$$\mathrm{QAM}^{\mathrm{loc}}_{\mathrm{HCOP}}, \qquad \mathrm{MML}^{\mathrm{loc}}_{\mathrm{HCOP}}, \qquad \mathrm{MMA}^{\mathrm{loc}}_{\mathrm{HCOP}}.$$

Their status is intentionally limited.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(a) $\mathrm{QAM}^{\mathrm{loc}}_{\mathrm{HCOP}}$ *is the finite tagged QAM bridge used to code the HC/OP objects of the present subsection inside the inherited QAM-over-ZF layer; its interface stamp is QAM-HC/OP 0.1.*

(b) $\mathrm{MML}^{\mathrm{loc}}_{\mathrm{HCOP}}$ *is the finite instruction language for the local proof chain HC-routed OP-entry chain; its interface stamp is MML-HC/OP 0.2.*

(c) $\mathrm{MMA}^{\mathrm{loc}}_{\mathrm{HCOP}}$ *is the matrix/readout execution semantics for those instructions on finite route-faithful machine states; its interface stamp is MMA-HC/OP 0.2.*

This registry upgrades only the local execution interface. It does not replace the inherited public QAM layer, does not activate the deferred binary tag system, and does not turn the historical MML–MMA–MCP reserve into an independent proof engine.

Definition H4.191 (HC/OP MMA object types)**.** *The minimal MMA object layer for HC-routed OP execution consists of the finite type family*

$$\mathsf{HCState},\quad \mathsf{HandoffPacket},\quad \mathsf{OPNormalForm},\quad \mathsf{TaskRouting},\quad \mathsf{ComparisonPacket},$$
$$\mathsf{Decision},\quad \mathsf{FrontierState},\quad \mathsf{RouteTag},\quad \mathsf{CertificateFinal}.$$

The intended instances are respectively: local HC-capture states, the handoff packets of Definition H4.166, the normal forms of Definition H4.170, the routing objects of Definition H4.177, the comparison packets of Definition H4.183, the decision values $\mathrm{bind}, \mathrm{split}, \mathrm{frontier}$*, smaller pre-entry frontier states, inherited route tags, and possible certificate data. A* $\mathsf{CertificateFinal}$ *becomes an actual theorematic certificate only when the previously stated mathematical trigger conditions are discharged outside the machine interface.*

Definition H4.192 (QAM coding bridge for HC/OP objects)**.** *A* QAM-HC/OP bridge code *is a partial coding map*

$$\ulcorner\cdot\urcorner^{\mathrm{HCOP}}_{\mathrm{QAM}} : \mathsf{Obj}^{\mathrm{HCOP}}_{\mathrm{MMA}} \dashrightarrow \mathsf{QAM}_{\mathrm{fin}},$$

where $\mathsf{Obj}^{\mathrm{HCOP}}_{\mathrm{MMA}}$ *is the finite object family of Definition H4.191 and* $\mathsf{QAM}_{\mathrm{fin}}$ *denotes finite tagged QAM objects in the inherited QAM-over-ZF sense. The map is required to preserve object kind, route tag, entry/frontier flag, and no-false-certificate status.*

Definition H4.193 (Minimal MML-HC/OP instruction alphabet)**.** *The local instruction alphabet* $\mathcal{I}^{\mathrm{HCOP}}_{\mathrm{MML}}$ *consists of the following finite instructions:*

$$\mathrm{CAPTURE}_{\mathrm{HC}},\quad \mathrm{HANDOFF}_{\mathrm{HC}\to\mathrm{OP}},\quad \mathrm{NORMALIZE}^{\mathrm{HC}}_{\mathrm{OP}},\quad \mathrm{ROUTE}^{\mathrm{HC}}_{\mathrm{OP}},$$
$$\mathrm{COMPARE}^{1/4|5(B)}_{\mathrm{HC}},\quad \mathrm{RESOLVE}_{\mathrm{bind/split/frontier}},\quad \mathrm{RETURN}_{\mathrm{OP}}.$$

The instructions are typed as follows:

(a) $\mathrm{CAPTURE}_{\mathrm{HC}}$ *sends a local admissible HC-state to a final transport-stable HC state.*

(b) $\mathrm{HANDOFF}_{\mathrm{HC}\to\mathrm{OP}}$ *sends such a state to an HC-to-OP handoff packet.*

(c) $\mathrm{NORMALIZE}^{\mathrm{HC}}_{\mathrm{OP}}$ *sends the handoff packet to an OP-entry normal form.*

(d) $\mathrm{ROUTE}^{\mathrm{HC}}_{\mathrm{OP}}$ *sends the normal form to a finite OP-entry task-routing object.*

(e) $\mathrm{COMPARE}^{1/4|5(B)}_{\mathrm{HC}}$ *executes the paired OP 1/4–OP 5(B) comparison when the routing object selects* $\rho_{\mathrm{CL}} \oplus \rho_{\Theta R}$*.*

(f) $\mathrm{RESOLVE}_{\mathrm{bind/split/frontier}}$ *records exactly one comparison decision.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(g) $\mathrm{RETURN}_{\mathrm{OP}}$ *returns either a bound common-origin state, a split ranked single-branch state, or a smaller frontier-certified comparison state.*

Definition H4.194 (MMA execution state for the HC/OP program)**.** *An* MMA-HC/OP execution state *is a finite tuple*

$$\Sigma_{\mathrm{HCOP}} = \big(S, \tau, \eta, r, c\big),$$

where S is the finite QAM-coded store, τ is the inherited route tag, $\eta \in \{\mathrm{entry}, \mathrm{frontier}\}$ is the entry/frontier flag, r is the active residual or comparison register, and c is a certificate register. The invariant of the state is:

$$c = \varnothing \quad \textit{unless a previously proved mathematical trigger supplies a certificate.}$$

Definition H4.195 (Finite HC-routed OP-entry MML program)**.** *The* minimal HC-routed OP-entry program *is the finite MML word*

$$P_{\mathrm{HCOP}} := (\mathrm{CAPTURE}_{\mathrm{HC}}, \mathrm{HANDOFF}_{\mathrm{HC}\to\mathrm{OP}}, \mathrm{NORMALIZE}^{\mathrm{HC}}_{\mathrm{OP}}, \mathrm{ROUTE}^{\mathrm{HC}}_{\mathrm{OP}},$$
$$\mathrm{COMPARE}^{1/4|5(B)}_{\mathrm{HC}}, \mathrm{RESOLVE}_{\mathrm{bind/split/frontier}}, \mathrm{RETURN}_{\mathrm{OP}}).$$

If the task-routing object does not select the paired branch, the comparison instruction is skipped and the return instruction hands control to the selected single work family instead. Thus the program is finite in either the paired or the single-branch case.

Lemma H4.196 (Typing of the HC-routed OP chain)**.** *Each step from the transport-stable HC-capture handoff through the first paired OP-entry comparison execution has a matching type in Definitions H4.191–H4.193.*

Proof. The handoff packet is an instance of HandoffPacket. The normal form is an instance of OPNormalForm. The task-routing object is an instance of TaskRouting. The paired comparison packet is an instance of ComparisonPacket, and its outcome belongs to Decision. The possible return states are exactly the bound, split, or frontier states already listed in Theorem H4.188. These are precisely the input/output types of the finite instruction alphabet. □

Lemma H4.197 (MML/MMA execution preserves route and frontier discipline)**.** *Executing P_{HCOP} in the MMA-HC/OP layer preserves the inherited route tag, preserves the entry/frontier flag unless a previously proved trigger changes it, and cannot fill the certificate register by machine action alone.*

Proof. Each instruction in Definition H4.193 is typed over the objects already constrained by the local proof chain mathematical layer. The handoff and normalization instructions inherit the route tag from the HC-capture chain; the routing and comparison instructions act only on the finite residual and comparison registers; the resolution instruction chooses only among bind, split, and frontier; and the return instruction returns one of the route-faithful next states of Theorem H4.188. None of these instructions contains a rule that writes a theorematic certificate into c. Hence route, frontier, and no-false-certificate discipline are preserved. □

Proposition H4.198 (comparison as one admissible MML/MMA step)**.** *The paired comparison execution of Theorem H4.188 is represented in* $\mathrm{MML}^{\mathrm{loc}}_{\mathrm{HCOP}}$ *by the instruction*

$$\mathrm{COMPARE}^{1/4|5(B)}_{\mathrm{HC}}$$

followed by $\mathrm{RESOLVE}_{\mathrm{bind/split/frontier}}$*. Its MMA semantics are exactly the construction of $D^{\mathrm{HC}}_{\mathrm{joint}}$, $D^{\mathrm{HC}}_{\mathrm{sep}}$, $\Delta^{\mathrm{HC}}_{\mathrm{sep}}$, and the decision ω_{cmp}.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. Definition H4.183 constructs the joint and separated HC-compatible readout errors and their difference. Definition H4.184 resolves the resulting comparison defect into exactly one decision. These are exactly the operational contents assigned to the comparison and resolution instructions. The statement is therefore only a typed re-reading of the comparison execution, not an additional theorematic claim. □

Theorem H4.199 (HC-routed OP-entry as a finite MML/MMA program)**.** *Assume the transport-stable HC-capture to OP-entry criterion of Theorem H4.168. Then the current the local proof chain HC-routed OP-entry chain is executable as the finite program* P_{HCOP} *over the local interface stack*

$$\mathrm{QAM}^{\mathrm{loc}}_{\mathrm{HCOP}} \longrightarrow \mathrm{MML}^{\mathrm{loc}}_{\mathrm{HCOP}} \longrightarrow \mathrm{MMA}^{\mathrm{loc}}_{\mathrm{HCOP}}.$$

The execution returns exactly one admissible next state: immediate OP-entry when the inherited mathematical trigger supplies it, a bound common-origin comparison state, a split ranked single-branch state, or a smaller frontier-certified state. It does not manufacture an OP closure certificate.

Proof. By Lemma H4.196, every mathematical object used in the local proof chain has a corresponding local MMA type and QAM bridge code. Definition H4.195 lists the finite instruction word realizing the chain. Lemma H4.197 supplies route preservation and the no-false-certificate condition. The possible next states are exactly the ones already obtained in the handoff, normal-form, routing, and paired-comparison theorems. Therefore the MML/MMA layer is an executable notation for the existing mathematical chain, not a new source of closure. □

Corollary H4.200 (working rule)**.** *The QAM/MML/MMA layer may now be used as a notation and audit interface for the HC-routed OP-entry program. Any later language extension must preserve the local registry of Definition H4.190: QAM remains a finite bridge over ZF-coded objects, MML remains a finite instruction language, MMA remains the execution semantics, and theorematic certificates must still come from the mathematical trigger conditions rather than from the machine layer itself.*

Proof. Immediate from Theorem H4.199 and the status restriction in Definition H4.190. □

H4.6 MML/MMA compression map for inherited HC/OP text

Definition H4.201 (Inherited HC/OP compression target)**.** *An* inherited HC/OP compression target *is a paragraph, lemma, proposition, diagnostic block, or tool paragraph in the inherited active or historical Companion text whose proof-use is one of the following finite roles:*

capture, handoff, normalization, routing, comparison, resolution, return, scan, rank, certificate test, route preservation, frontier descent.

The target is compressible *only in the expository sense: it may be reread as a typed MML/MMA operation or short finite operation word, but its original mathematical content, hypotheses, and proof restrictions remain part of the document unless a later explicit editorial release removes or relocates them.*

Definition H4.202 (Compression map from inherited text to MML/MMA operations)**.** *The local* HC/OP compression map

$$\mathfrak{C}_{\mathrm{HCOP}}$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

sends inherited compression targets to typed MML/MMA readings according to the following table.

Inherited text pattern	***MML/MMA reading***	***Compression use***
HC lock corridor, local HC final, minimal HC-lock step	$\mathrm{LOCK/CAPTURE}_{\mathrm{HC}}$	*Marks the block as a final-production and lock-stabilization step rather than as an OP certificate.*
Transport-stable HC-to-OP packet	$\mathrm{HANDOFF}_{\mathrm{HC}\to\mathrm{OP}}$	*Turns repeated handoff prose into one typed transfer from HC state to OP-facing packet.*
OP-entry normal form and residual triple	$\mathrm{NORMALIZE}_{\mathrm{OP}}^{\mathrm{HC}}$	*Collects ρ_{CL}, $\rho_{\Theta R}$, ρ_A, route tags, and entry/frontier flags into one typed normal form.*
Residual-to-task selector	$\mathrm{ROUTE}_{\mathrm{OP}}^{\mathrm{HC}}$	*Replaces repeated branch-selection explanations by one admissible routing step.*
Paired OP 1/4–OP 5(B) branch	$\mathrm{COMPARE}_{\mathrm{HC}}^{1/4\mid 5(B)}$	*Reads the joint/separated defect comparison as one admissible comparison instruction.*
Bind/split/frontier decision	$\mathrm{RESOLVE}_{\mathrm{bind/split/frontier}}$	*Records that the comparison returns only a bound state, a split state, or a smaller frontier state.*
Next-state handback to OP-facing work	$\mathrm{RETURN}_{\mathrm{OP}}$	*Compresses the return package without changing theorematic status.*
Probe, Monte Carlo, decoder, or finite search block	SCAN → RANK → CANDIDATE	*Makes clear that computational probes rank admissible finals but do not prove closure.*
Admissibility / boundary-stop / route-faithfulness paragraph	PRESERVE_ROUTE	*Marks the paragraph as an invariant-preservation rule for route tag and boundary-stop data.*
Certificate discussion or trigger-persistence paragraph	CERTIFY?	*Keeps certification separate from execution: the operation asks whether a mathematical trigger exists; it never creates one.*
Frontier descent or smaller pre-entry state	$\mathrm{DESCEND}_{\mathrm{frontier}}$	*Records strict controlled progress without falsely upgrading the state to OP closure.*
Historical tool or early proof-attempt block	ARCHIVE → TOOL	*Keeps the historical witness intact while extracting the reusable tool-role for the active line.*

Lemma H4.203 (Compression map preserves proof content)**.** *Applying $\mathfrak{C}_{\mathrm{HCOP}}$ to an inherited compression target does not change the truth status of that target, does not delete hypotheses, and does not promote a diagnostic or work packet to a theorematic certificate.*

Proof. By Definition H4.201, a compression target is compressed only in its expository role. Definition H4.202 assigns a typed MML/MMA reading that records what the block does in the finite HC/OP program. The assignment neither alters the original statement nor supplies

© 2026 Marc Brendecke. Preprint — not peer reviewed.

a new mathematical trigger. The no-false-certificate restriction of Lemma H4.197 therefore remains in force. □

Proposition H4.204 (Main inherited text zones helped by the compression map)**.** *The compression map gives immediate expository leverage in the following inherited zones:*

(i) HC-capture and lock-corridor paragraphs, by replacing repeated final/lock explanations with LOCK/CAPTURE$_{\mathrm{HC}}$*;*

(ii) HC-to-OP handoff and OP-entry paragraphs, by replacing repeated transfer and normal-form prose with HANDOFF*,* NORMALIZE*, and* ROUTE*;*

(iii) OP 1/4–OP 5(B) comparison paragraphs, by writing the inseparable branch as COMPARE $\to$ RESOLVE $\to$ RETURN*;*

(iv) probe and Monte-Carlo paragraphs, by reading them as SCAN $\to$ RANK $\to$ CANDIDATE *and not as proof generators;*

(v) historical witness paragraphs, by marking reusable tools without deleting the preserved genealogy.

Proof. Each listed zone contains one of the patterns displayed in Definition H4.202. The assigned operation records the proof-use of the zone and can therefore replace later repetitive explanation. Lemma H4.203 prevents the compression from changing mathematical status. □

Theorem H4.205 (MML/MMA compression theorem for inherited HC/OP text)**.** *Every inherited HC/OP compression target that matches one of the patterns in Definition H4.202 admits a finite MML/MMA reading under*

$$\mathrm{QAM}^{\mathrm{loc}}_{\mathrm{HCOP}} \longrightarrow \mathrm{MML}^{\mathrm{loc}}_{\mathrm{HCOP}} \longrightarrow \mathrm{MMA}^{\mathrm{loc}}_{\mathrm{HCOP}}.$$

This reading may be used for later compression, indexing, and audit of the inherited text. It may not be used to erase the historical witness or to manufacture an OP closure certificate.

Proof. The interface theorem, Theorem H4.199, already supplies the finite typed execution stack for the current HC-routed OP-entry chain. Definition H4.202 extends this stack as an expository map over inherited text patterns that perform the same roles. Since the map is role-preserving and certificate-neutral by Lemma H4.203, it yields a finite MML/MMA reading for each matching target while leaving the original mathematical content intact. □

Corollary H4.206 (working rule)**.** *The next editorial pass may shorten repeated HC/OP explanations by citing* $\mathfrak{C}_{\mathrm{HCOP}}$*, but it should not delete active proof material yet. The safe immediate use of the compression map is to add labels, cross-references, tool tables, and short operation names to inherited blocks; stronger textual shortening should wait until the corresponding OP branch has either closed or returned to a stable frontier state.*

Proof. Immediate from Theorem H4.205. The compression map is currently an audit and refactoring layer, not a release-level deletion license. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

H4.7 First compression-guided HC/OP tool normal forms

Definition H4.207 (Reusable HC/OP tool normal form)**.** *A* reusable HC/OP tool normal form *is a finite typed contract*

$$\mathfrak{T} = \langle \text{name}, \text{input}, \text{guard}, \text{operation}, \text{output}, \text{status} \rangle$$

obtained from the compression map $\mathfrak{C}_{\mathrm{HCOP}}$*. Its entries have the following roles.*

(i) name *is one of the finite MML/MMA operation names already admitted by Definitions H4.193 and H4.202;*

(ii) input *is the typed object family consumed by the tool;*

(iii) guard *is the admissibility, route, frontier, or certificate-neutrality condition under which the tool may be invoked;*

(iv) operation *is the finite MML instruction word together with its MMA readout semantics;*

(v) output *is the typed object family returned by the tool;*

(vi) status *records whether the tool is diagnostic, normalizing, routing, comparison-level, frontier-level, or certificate-testing.*

A tool normal form is an invocation interface only. It may shorten later exposition by replacing repeated safety prose with a typed reference, but it does not erase the original proof material and does not create a theorematic certificate.

Definition H4.208 (First extracted HC/OP tool library)**.** *The first compression-guided HC/OP tool library is the finite family*

$$\mathbb{T}^{(1)}_{\mathrm{HCOP}} := \{\mathfrak{T}_{\mathrm{lock}}, \mathfrak{T}_{\mathrm{handoff}}, \mathfrak{T}_{\mathrm{norm}}, \mathfrak{T}_{\mathrm{route}}, \mathfrak{T}_{\mathrm{cmp}}, \mathfrak{T}_{\mathrm{resolve}}, \mathfrak{T}_{\mathrm{probe}}, \mathfrak{T}_{\mathrm{cert}}, \mathfrak{T}_{\mathrm{frontier}}, \mathfrak{T}_{\mathrm{archive}}\}.$$

Its intended contracts are summarized as follows.

Tool	***Input***	***Output***	***Guard / status***
$\mathfrak{T}_{\mathrm{lock}}$	*local HC final, corridor data*	*locked/captured HC final*	*guard: inherited route tag and local admissibility; status: final/capture*
$\mathfrak{T}_{\mathrm{handoff}}$	*transport-stable HC-capture chain*	*OP-facing handoff packet*	*guard: zero or controlled HC transport defect; status: handoff*
$\mathfrak{T}_{\mathrm{norm}}$	*handoff packet*	$\mathcal{N}^{\mathrm{HC}}_{\mathrm{OP}}$ *with live residual triple*	*guard: no change of theorematic status; status: normalization*
$\mathfrak{T}_{\mathrm{route}}$	$\mathcal{N}^{\mathrm{HC}}_{\mathrm{OP}}$*, selected residual branch*	*finite OP-facing task family*	*guard: residual type and route tag match; status: routing*
$\mathfrak{T}_{\mathrm{cmp}}$	*paired* $\rho_{\mathrm{CL}} \oplus \rho_{\Theta R}$ *branch*	$\Delta^{\mathrm{HC}}_{\mathrm{sep}}$ *and comparison record*	*guard: inseparable branch has not yet split; status: comparison*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Tool	*Input*	*Output*	*Guard / status*
$\mathfrak{T}_{\mathrm{resolve}}$	*comparison record*	*bind, split, or frontier next state*	*guard: decision rule returns exactly one branch; status: resolution*
$\mathfrak{T}_{\mathrm{probe}}$	*probe/Monte-Carlo/decoder scan surface*	*ranked admissible final family*	*guard: diagnostic only; status: scan/rank/final*
$\mathfrak{T}_{\mathrm{cert}}$	*trigger-persistence or certificate question*	*yes/no/pending certificate-test record*	*guard: asks only whether a mathematical trigger exists; status: certificate-test*
$\mathfrak{T}_{\mathrm{frontier}}$	*non-closing but improving state*	*smaller frontier-certified pre-entry state*	*guard: strict controlled frontier descent; status: frontier*
$\mathfrak{T}_{\mathrm{archive}}$	*historical witness block*	*reusable tool role plus preserved genealogy*	*guard: no deletion of historical proof memory; status: archive-to-tool*

Lemma H4.209 (Extracted tool contracts are typed MML/MMA readings)**.** *Every member of $\mathbb{T}^{(1)}_{\mathrm{HCOP}}$ is a typed MML/MMA reading of one or more rows of the compression map $\mathfrak{C}_{\mathrm{HCOP}}$.*

Proof. Definition H4.208 is obtained by grouping the rows of Definition H4.202 by proof-use. The lock, handoff, normalization, routing, comparison, resolution, return, scan/rank, certificate-test, frontier-descent, and archive-to-tool roles are exactly the roles already admitted by the compression map and by the instruction alphabet. Hence each extracted contract is a typed reading over the existing QAM/MML/MMA stack, not a new primitive operation. □

Proposition H4.210 (Immediate simplification rule for inherited HC/OP prose)**.** *Let B be an inherited HC/OP text block that matches one of the patterns of Definition H4.202. If B has already stated its hypotheses and proof restrictions once, then later repetitions of the same proof-use may be replaced by a citation to the corresponding tool contract in $\mathbb{T}^{(1)}_{\mathrm{HCOP}}$, provided the replacement also cites the original local statement or keeps it in place.*

Proof. By Lemma H4.203, the compression map changes only the expository role of a block and preserves the original mathematical content. By Lemma H4.209, the corresponding tool contract records the same role in a finite typed form. Therefore a later repetition can be shortened to a tool invocation as long as the first occurrence and its hypotheses remain available. The proviso prevents the normal form from becoming a deletion rule. □

Proposition H4.211 (Where the first tool layer helps the existing text)**.** *The library $\mathbb{T}^{(1)}_{\mathrm{HCOP}}$ gives immediate refactoring leverage in the following existing zones of the Companion.*

(i) The HC-lock and capture paragraphs may cite $\mathfrak{T}_{\mathrm{lock}}$ instead of restating the full final/corridor discipline.

(ii) The local proof chain OP-entry paragraphs may cite $\mathfrak{T}_{\mathrm{handoff}}$, $\mathfrak{T}_{\mathrm{norm}}$, and $\mathfrak{T}_{\mathrm{route}}$ instead of repeating the full handoff-normalize-route chain.

(iii) The OP 1/4–OP 5(B) branch may cite $\mathfrak{T}_{\mathrm{cmp}}$ and $\mathfrak{T}_{\mathrm{resolve}}$ to express the joint-versus-separated comparison.

(iv) Probe, Monte-Carlo, decoder, and finite-search paragraphs may cite $\mathfrak{T}_{\mathrm{probe}}$ to emphasize that they rank finals rather than certify closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(v) Trigger-persistence paragraphs may cite $\mathfrak{T}_{\mathrm{cert}}$ *to keep certification separate from execution.*

(vi) Historical witness paragraphs may cite $\mathfrak{T}_{\mathrm{archive}}$ *to extract reusable tools without shortening the preserved genealogy.*

Proof. Each zone listed is one of the main inherited zones already identified in Proposition H4.204. Definition H4.208 assigns the corresponding reusable contract. The proposition is therefore the first explicit refactoring consequence of the compression map. □

Theorem H4.212 (First compression-guided HC/OP reusable-tool layer)**.** *The compression map* $\mathfrak{C}_{\mathrm{HCOP}}$ *induces the finite reusable-tool layer* $\mathbb{T}^{(1)}_{\mathrm{HCOP}}$*. This layer may be used in later Companion revisions to shorten repeated HC/OP safety prose, index historical proof material, and audit OP-entry execution traces. It may not be used to delete active proof content or to manufacture an OP closure certificate.*

Proof. Existence of $\mathbb{T}^{(1)}_{\mathrm{HCOP}}$ is Definition H4.208. Its relation to the compression map is Lemma H4.209. Its safe simplification use is Proposition H4.210. Its no-false-certificate restriction follows from Lemma H4.197 and Lemma H4.203. Thus the layer is a reusable tool interface over the existing mathematics and not a new theorematic engine. □

Corollary H4.213 (working rule)**.** *The next safe editorial use of MML/MMA is to annotate repeated inherited HC/OP paragraphs by the tool contracts in* $\mathbb{T}^{(1)}_{\mathrm{HCOP}}$*. A future shortening pass may replace repeated explanations by these tool names only after the original local statement, hypotheses, and proof restrictions remain explicitly available elsewhere in the document.*

Proof. Immediate from Theorem H4.212. □

H4.8 First tool-contract application to the active HC/OP proof spine

Definition H4.214 (Tool-indexed active HC/OP spine)**.** *The* tool-indexed active HC/OP spine *is the finite contract trace*

$$\mathcal{P}^{\mathrm{tool}}_{\mathrm{HCOP}} := \mathfrak{T}_{\mathrm{lock}};\mathfrak{T}_{\mathrm{handoff}};\mathfrak{T}_{\mathrm{norm}};\mathfrak{T}_{\mathrm{route}};\mathfrak{T}_{\mathrm{cmp}};\mathfrak{T}_{\mathrm{resolve}};\mathfrak{T}_{\mathrm{cert/frontier}}.$$

It is a compact citation layer over the local proof chain HC/OP proof spine. The trace is read as follows: lock/capture prepares the local HC final, handoff transports it toward OP-facing form, normalization produces $\mathcal{N}^{\mathrm{HC}}_{\mathrm{OP}}$*, routing selects a live residual work family, comparison executes the paired OP 1/4–OP 5(B) readout when separation is not yet justified, resolution returns bind, split, or frontier, and the final certificate/frontier test records whether the branch may promote or must remain a smaller frontier-certified state.*

Definition H4.215 (Tool-contract application site)**.** *A paragraph, lemma, proposition, or proof step is a* tool-contract application site *for a contract* $\mathfrak{T} \in \mathbb{T}^{(1)}_{\mathrm{HCOP}}$ *if all of the following are explicit either locally or by a nearby reference.*

(i) the typed input object of $\mathfrak{T}$*;*

(ii) the guard of $\mathfrak{T}$*, including route, frontier, and admissibility restrictions;*

(iii) the operation or comparison performed;

(iv) the typed output object;

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(v) the status restriction that the invocation is diagnostic, routing, comparison-level, frontier-level, or certificate-testing rather than certificate-creating.

At such a site, later repetitions may cite $\mathfrak{T}$ without restating the full safety prose, provided the original statement and its hypotheses remain visible in the document.

Lemma H4.216 (Active HC/OP blocks are contract-application sites)**.** *The local proof chain active HC/OP blocks are application sites for the corresponding contracts in Definition H4.214.*

Proof. The transport-stable HC-capture and handoff block states the local final, route-faithful transport guard, and OP-facing packet output, hence matches $\mathfrak{T}_{\mathrm{lock}}$ and $\mathfrak{T}_{\mathrm{handoff}}$. The normal form block states the handoff input, the live residual triple, and the non-closure status restriction, hence matches $\mathfrak{T}_{\mathrm{norm}}$. The task-routing block states the residual branch, admissible work family, and route tag, hence matches $\mathfrak{T}_{\mathrm{route}}$. The paired-comparison block states the inseparable $\rho_{\mathrm{CL}} \oplus \rho_{\Theta R}$ input, the comparison defect, and the bind/split/frontier return, hence matches $\mathfrak{T}_{\mathrm{cmp}}$ and $\mathfrak{T}_{\mathrm{resolve}}$. The inherited no-false-certificate rule supplies the final certificate/frontier guard. Therefore each block satisfies Definition H4.215. □

Proposition H4.217 (Tool-indexed rewriting of the active HC/OP chain)**.** *The active the local proof chain HC/OP chain admits the compressed reading*

$$\mathrm{HC}_{\mathrm{loc}} \xrightarrow{\mathfrak{T}_{\mathrm{lock}}} \mathrm{HC}_{\mathrm{cap}} \xrightarrow{\mathfrak{T}_{\mathrm{handoff}}} \mathcal{P}_{\mathrm{handoff}} \xrightarrow{\mathfrak{T}_{\mathrm{norm}}} \mathcal{N}_{\mathrm{OP}}^{\mathrm{HC}} \xrightarrow{\mathfrak{T}_{\mathrm{route}}} \mathcal{T}_{\mathrm{OP}}^{\mathrm{HC}} \xrightarrow{\mathfrak{T}_{\mathrm{cmp}},\mathfrak{T}_{\mathrm{resolve}}} \{\mathrm{bind}, \mathrm{split}, \mathrm{frontier}\}.$$

This rewriting is admissible as a cross-reference and audit normal form, but it is not an independent proof of any OP closure.

Proof. By Lemma H4.216, each arrow is backed by an already present active HC/OP block and by a typed contract in $\mathbb{T}^{(1)}_{\mathrm{HCOP}}$. The chain is therefore a finite MML/MMA execution trace in the sense of Theorem H4.199. Since Theorem H4.212 forbids tool contracts from manufacturing certificates, the compressed trace is an audit normal form only. □

Lemma H4.218 (Tool application preserves local hypotheses)**.** *Replacing a repeated explanation by a citation to a tool contract preserves the local hypotheses of the original explanation exactly when the application site cites the original statement or restates the required guard.*

Proof. A tool contract records input, guard, operation, output, and status; it does not by itself reproduce every local hypothesis. Definition H4.215 therefore requires the relevant guard to be explicit locally or by reference. Under that requirement, the replacement removes only duplicated prose and leaves the logical premises available. Without that requirement the replacement would be unsafe, which is why the definition includes it as a guard. □

Corollary H4.219 (First safe shortening target)**.** *The first safe shortening target is repeated no-false-certificate, route-preservation, and frontier-preservation prose following an already stated HC/OP operation. Such repetitions may be replaced by a sentence of the form:*

> *By the corresponding tool contract in $\mathbb{T}^{(1)}_{\mathrm{HCOP}}$, this is an admissible MML/MMA execution step and not a closure certificate.*

The original definition, theorem, or proof restriction must remain present elsewhere in the same local chain.

Proof. The repeated phrases listed are status and guard reminders, not new mathematical premises. Lemma H4.218 permits the repetition to be replaced once the underlying guard has already been stated. The no-false-certificate clause is exactly the status restriction of $\mathfrak{T}_{\mathrm{cert}}$ and of the active MML/MMA interface. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Theorem H4.220 (First tool-contract application theorem)**.** *The reusable tool library* $\mathbb{T}^{(1)}_{\mathrm{HCOP}}$ *can now be applied to the active HC/OP proof spine as a conservative cross-reference layer. It yields shorter notation, clearer audit points, and explicit preservation of route, frontier, and certificate restrictions, without changing theorematic status and without deleting historical witness material.*

Proof. Shorter notation follows from Proposition H4.217. Auditability follows because Definition H4.215 lists the required input, guard, operation, output, and status fields for every invocation. Preservation of local hypotheses follows from Lemma H4.218. The no-deletion and no-false-certificate restrictions are inherited from Theorem H4.212. Hence the layer is a conservative application of the tool library rather than a new proof step. □

Corollary H4.221 (working rule)**.** *Subsequent revisions may begin annotating the active HC/OP chain by* $\mathcal{P}^{\mathrm{tool}}_{\mathrm{HCOP}}$*. They should first replace only duplicated safety prose and local route/frontier reminders. They should not delete first occurrences of definitions, hypotheses, proof restrictions, historical witness blocks, or any statement needed to verify the no-false-certificate rule.*

Proof. Immediate from Theorem H4.220 and Corollary H4.219. □

H4.9 First tool-guarded reduction protocol for inherited HC/OP prose

Definition H4.222 (Guarded HC/OP reduction packet)**.** *A* guarded HC/OP reduction packet *is a finite tuple*

$$\mathcal{R}^{(1)}_{\mathrm{HCOP}} := \langle B, \mathfrak{T}, G_B, S_B, W_B \rangle$$

with the following data.

(i) B *is an inherited or active HC/OP paragraph, lemma step, proof reminder, or diagnostic status clause.*

(ii) $\mathfrak{T} \in \mathbb{T}^{(1)}_{\mathrm{HCOP}}$ *is the tool contract assigned to the repeated role of* B*.*

(iii) G_B *records the local guard: typed input, route tag, frontier condition, admissibility condition, and no-false-certificate status.*

(iv) S_B *records the theorematic status of the occurrence: definition, hypothesis, proof step, diagnostic reminder, certificate question, frontier update, or historical witness.*

(v) W_B *records where the unreduced witness of* B *remains available in the document.*

The packet is a reduction-control object only. It does not change the mathematical claim carried by B*.*

Definition H4.223 (Reducible and non-reducible HC/OP occurrences)**.** *Let* B *be an occurrence in the HC/OP proof spine or in an inherited HC/OP support zone. The occurrence is* the local pass-reducible *if it satisfies all of the following conditions.*

(i) It repeats a safety, route-preservation, frontier-preservation, diagnostic-status, scan/rank/final, comparison-status, or no-false-certificate clause already stated in the same local chain.

(ii) Its assigned tool contract $\mathfrak{T}$ *is one of the contracts of Definition H4.208.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(iii) *The original hypotheses, guards, proof restrictions, and local statement remain visible either in B's first occurrence or in a nearby cited statement.*

(iv) *The replacement sentence explicitly says that the citation is a tool-contract invocation and not a closure certificate.*

The occurrence is the local pass-non-reducible *if it introduces a first definition, a new object, a new hypothesis, a new guard, a new route tag, a certificate test, a theorematic status change, a proof-critical inequality, or historical witness material.*

Definition H4.224 (Guarded citation normal form)**.** *The preferred normal form for replacing an the local pass-reducible occurrence is*

> *By the guarded tool contract $\mathfrak{T}$ and the local guard recorded in G_B, this is an admissible MML/MMA execution or diagnostic step; it preserves route/frontier discipline and does not create an OP closure certificate.*

When the local context is already clear, the sentence may be shortened to a parenthetical form:

guarded by $\mathfrak{T}$; route/frontier preserved; no certificate manufactured.

Lemma H4.225 (Guarded reduction preserves proof content)**.** *If B is the local pass-reducible and is replaced by the guarded citation normal form of Definition H4.224, then the replacement preserves the proof content of the local chain.*

Proof. By Definition H4.223, only repeated status and guard reminders are eligible for replacement. The original hypotheses, proof restrictions, and first local statement remain available. By Definition H4.224, the replacement still records the assigned tool contract, preserves route/frontier discipline, and repeats the no-false-certificate status. Hence the replacement removes duplicated prose only; it does not remove the premise, operation, guard, output, or theorematic status of the original argument. □

Lemma H4.226 (Non-reducible occurrences protect theorematic status)**.** *Every the local pass-non-reducible occurrence must remain explicit in the document unless a later revision supplies a stronger local theorem that restates the same content with all hypotheses and guards.*

Proof. The non-reducible occurrences listed in Definition H4.223 are precisely the occurrences that create or alter mathematical content: they introduce objects, hypotheses, route tags, guards, certificate tests, status changes, proof-critical inequalities, or historical witness material. Replacing such an occurrence by a tool-name alone would hide a premise or change the audit surface. Therefore they are protected until a later theorem restates them explicitly. □

Proposition H4.227 (First usable simplification map)**.** *The reduction protocol induces the partial simplification map*

$$\mathsf{Red}^{(1)}_{\mathrm{HCOP}} : B \dashrightarrow \big(\mathfrak{T}(B), G_B, S_B, W_B\big),$$

where the map is defined exactly on the local pass-reducible occurrences and undefined on the local pass-non-reducible occurrences.

Proof. For a reducible occurrence, Definitions H4.222 and H4.224 assign the corresponding tool contract, guard, status, and witness pointer. Lemma H4.225 proves that this replacement is content-preserving. For a non-reducible occurrence, Lemma H4.226 forbids replacement by a mere tool invocation. Hence the simplification map is partial and conservative. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Theorem H4.228 (First tool-guarded HC/OP reduction theorem)**.** *The active HC/OP tool layer now supports a first conservative reduction protocol. Repeated safety prose, route/frontier reminders, diagnostic-status clauses, and no-false-certificate clauses may be replaced by guarded tool citations according to* $\mathsf{Red}^{(1)}_{\mathrm{HCOP}}$*. First definitions, first hypotheses, local guards, certificate tests, proof-critical inequalities, theorematic status changes, and historical witness material remain non-reducible. Therefore the protocol simplifies later exposition without altering the OP status of the document.*

Proof. The permitted replacements are exactly the defined values of the partial map in Proposition H4.227. Content preservation follows from Lemma H4.225. Protection of theorematic status follows from Lemma H4.226. The no-false-certificate restriction is inherited from Theorem H4.220. Thus the reduction protocol is a simplifier for repeated exposition and an audit aid, not a new OP closure mechanism. □

Corollary H4.229 (working rule)**.** *The next safe use of the tool layer is a local annotation pass: mark repeated HC/OP safety paragraphs as reducible or non-reducible before shortening them. Actual deletion or replacement should occur only after the first occurrence, hypotheses, guards, and historical witness pointer have been verified.*

Proof. Immediate from Theorem H4.228. □

H4.10 First guarded-reduction application pass for the active HC/OP proof spine

Definition H4.230 (First guarded-reduction application ledger)**.** *The* first guarded-reduction application ledger *is the finite list*

$$\mathcal{L}^{(1)}_{\mathrm{HCOP}} := \{\ell_{\mathrm{lock}}, \ell_{\mathrm{handoff}}, \ell_{\mathrm{norm}}, \ell_{\mathrm{route}}, \ell_{\mathrm{cmp}}, \ell_{\mathrm{probe}}, \ell_{\mathrm{archive}}\}.$$

Each ledger entry has the form

$$\ell = \langle S_\ell,\, W_\ell,\, \mathfrak{T}_\ell,\, G_\ell,\, R_\ell \rangle,$$

where S_ℓ *is the local site class,* W_ℓ *is the protected unreduced witness,* $\mathfrak{T}_\ell \in \mathbb{T}^{(1)}_{\mathrm{HCOP}}$ *is the tool contract used for later citation,* G_ℓ *is the local guard data, and* R_ℓ *is the repeated prose tail that may be replaced by a guarded citation. The ledger is an application of the partial map* $\mathsf{Red}^{(1)}_{\mathrm{HCOP}}$ *from Proposition H4.227; it is not an instruction to remove the first occurrence of any definition, guard, certificate test, proof-critical inequality, or historical witness.*

Definition H4.231 (guarded citation sites)**.** *The entries of* $\mathcal{L}^{(1)}_{\mathrm{HCOP}}$ *are interpreted as follows.*

(i) ℓ_{lock} *covers later repeated lock/capture reminders after the first local HC-capture statement has been given. Its citation contract is* $\mathfrak{T}_{\mathrm{lock}}$*, and its protected witness is the unreduced local capture chain.*

(ii) ℓ_{handoff} *covers later repeated claims that HC capture is only a handoff into OP-entry and not an OP closure. Its citation contracts are* $\mathfrak{T}_{\mathrm{handoff}}$ *and* $\mathfrak{T}_{\mathrm{cert}}$*.*

(iii) ℓ_{norm} *covers later repeated normalization reminders for the HC-routed OP normal form. Its citation contract is* $\mathfrak{T}_{\mathrm{norm}}$*, guarded by the already visible residual triple and route tag.*

(iv) ℓ_{route} *covers later repeated route/frontier-preservation clauses in the finite task-routing object. Its citation contracts are* $\mathfrak{T}_{\mathrm{route}}$ *and* $\mathfrak{T}_{\mathrm{frontier}}$*.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(v) ℓ_{cmp} *covers later repeated status clauses for the paired OP 1/4–OP 5(B) comparison branch. Its citation contracts are* $\mathfrak{T}_{\mathrm{cmp}}$ *and* $\mathfrak{T}_{\mathrm{resolve}}$*, and its protected witness includes the explicit comparison defect* $\Delta^{\mathrm{HC}}_{\mathrm{sep}}$*.*

(vi) ℓ_{probe} *covers later repeated probe-diagnostic clauses stating that scan/rank/final output is not a certificate. Its citation contracts are* $\mathfrak{T}_{\mathrm{probe}}$ *and* $\mathfrak{T}_{\mathrm{cert}}$*.*

(vii) ℓ_{archive} *covers later repeated archive-to-tool extraction reminders. Its citation contract is* $\mathfrak{T}_{\mathrm{archive}}$*, and its protected witness is the unreduced historical source block.*

Definition H4.232 (Ledger citation normal form)**.** *For an occurrence assigned to an entry* $\ell \in \mathcal{L}^{(1)}_{\mathrm{HCOP}}$*, the preferred citation normal form is*

$$\mathrm{GRCite}_{\mathrm{HCOP}}(\ell) := \langle \mathfrak{T}_\ell, G_\ell, W_\ell; \mathrm{noCert} \rangle.$$

In prose this is read as:

> *Guarded by* $\mathfrak{T}_\ell$ *with local guard* G_ℓ *and protected witness* W_ℓ*; route/frontier discipline is preserved and no certificate is manufactured.*

The normal form may replace only a repeated tail R_ℓ*, never the protected witness* W_ℓ*.*

Lemma H4.233 (Ledger entries are -reducible where used)**.** *Every later occurrence covered by a ledger entry in Definition H4.231 is the local pass-reducible exactly when its first local witness, guard, and theorematic status remain visible.*

Proof. Each ledger entry records a repeated prose tail: safety status, route/frontier preservation, diagnostic non-certification, comparison status, or archive-to-tool extraction. These are precisely the reducible classes of Definition H4.223. The same definition excludes replacement whenever a first occurrence, guard, certificate test, proof-critical inequality, status change, or historical witness would be hidden. Hence the ledger is admissible only at later occurrences for which the protected witness W_ℓ remains visible. □

Proposition H4.234 (First guarded-reduction application is conservative)**.** *Applying* $\mathrm{GRCite}_{\mathrm{HCOP}}(\ell)$ *to the repeated tail of any admissible ledger entry preserves the mathematical content of the active HC/OP proof spine.*

Proof. The replacement is defined only for repeated tails R_ℓ. By Definition H4.230, the protected witness W_ℓ, the tool contract $\mathfrak{T}_\ell$, the local guard G_ℓ, and the no-certificate status remain part of the citation normal form. Lemma H4.233 reduces the claim to Lemma H4.225. Thus the operation shortens only repeated exposition and leaves the proof spine unchanged. □

Theorem H4.235 (First guarded-reduction application theorem)**.** *The active HC/OP proof spine now carries a first practical guarded-reduction application layer. Later repeated HC/OP prose may cite the finite ledger* $\mathcal{L}^{(1)}_{\mathrm{HCOP}}$ *in the normal form of Definition H4.232. This permits local exposition to be shortened after first occurrence, but it does not delete proof material, alter route tags, alter frontier status, convert diagnostics into certificates, or change the OP status of the document.*

Proof. Existence of the finite ledger follows from Definition H4.230. Admissibility of its local uses follows from Lemma H4.233. Conservativity follows from Proposition H4.234. The no-false-certificate restriction is explicitly retained in Definition H4.232 and is inherited from Theorem H4.228. Hence the layer is a guarded reduction application pass rather than a new theorematic closure step. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Corollary H4.236 (working rule)**.** *The next safe simplification pass may replace only later repeated HC/OP tails that match an entry of* $\mathcal{L}^{(1)}_{\mathrm{HCOP}}$*. The first occurrence of each definition, hypothesis, guard, certificate test, proof-critical inequality, and historical witness remains explicit.*

Proof. Immediate from Theorem H4.235. □

Definition H4.237 (Defect transport equation)**.** *Let* X_n *and* X_{n+1} *be adjacent closure-liquid-crystal cells. Their defect transport equation is written*

$$\mathcal{D}_{\mathrm{cell}}(X_n, X_{n+1}) \neq 0$$

whenever the first local failure is no longer merely screening-visible, but appears as a witness-grade or shell-grade obstruction transported across the current channel. The datum $\mathcal{D}_{\mathrm{cell}}$ *is therefore not a second closure criterion; it is the cellular packaging of the first locally non-contractible obstruction.*

Proposition H4.238 (OP 5 defect-transport reading)**.** *Within the current bundled OP 3→OP 5/global channel, witness-grade OP 5 reopening may be read as defect transport on the same closure-liquid-crystal medium: a bundled global pass corresponds to defect-free or defect-screened transport, while a bundled obstruction corresponds to transported or accumulated non-contractible defect data.*

Proof. By Theorem H7.57, Theorem H7.60, and Theorem H7.82, the current channel already distinguishes a bundled pass from a bundled obstruction while subordinating packet/phase/epsilon diagnostics to the carrier-level decision. The witness-grade local/global reopening discipline then identifies the precise place where a local obstruction must be reopened rather than screened away. This is exactly the cellular reading of $\mathcal{D}_{\mathrm{cell}} \neq 0$: a transported or accumulated defect of one and the same carrier medium rather than a second primitive theory beside the closure channel. □

Definition H4.239 (Liquid readout domain)**.** *Fix a distinguished readout regime* f*. A* liquid readout domain *in regime* f *is a family of closure-liquid-crystal cells whose carrier data lie in one closure-bearing class while their visible shell/phase data realize one coherent readout phase of that same bulk order. Different regimes or sectors may therefore produce different liquid readout domains without changing the underlying carrier datum in the strong closure sense.*

Remark H4.240 (Readouts as liquid-crystal domains of one bulk order)**.** *The point of Definition H4.239 is not to introduce a literal condensed-matter ontology. It is to express the present structural reading more economically: the visible sectors are not independent primitive crystals, but phase-sensitive ordered domains of one deeper closure-bearing bulk carrier. This is the cellular version of the readout-relative principle already used throughout the Companion.*

Definition H4.241 (Holographic Julia-boundary readout)**.** *Let* $\{X_n\}_{n\in I}$ *be an admissible closure-liquid-crystal configuration along a chosen branch/route family. A* holographic Julia-boundary readout *is a boundary map*

$$\mathcal{H}_{\mathrm{J}}(\{X_n\}_{n\in I})$$

that records the visible boundary pattern produced by the phase-/shell-sensitive readout of the same bulk carrier configuration. The intended output is a Julia-type boundary image, diagnostic strip, or related visual readout.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proposition H4.242 (Holographic Julia-boundary principle)**.** *The Julia-type diagnostics used in the present Companion may be read as holographic boundary readouts of an underlying closure-liquid-crystal configuration. In particular, they remain diagnostic or screening-level boundary objects: they support the active closure architecture, but they do not create a new independent closure axiom.*

Proof. The current document already treats packet-, phase-lock-, epsilon-, and frame-flicker diagnostics as visible readout layers subordinate to the carrier-level and closure-level statements; see Proposition H7.33 and Corollary H7.37. The same discipline applies to Julia-type visuals: they belong on the visible boundary-readout side and are informative exactly insofar as they track admissible passage, defect localization, or loss of admissibility in the bulk carrier medium. Hence they may be read holographically without becoming independent closure generators. □

Remark H4.243 (historical release line pilot status)**.** *The present section should be read as a first pilot layer for archival stage. It upgrades neither the current OP 3/QAM closure package nor the bundled OP 3→OP 5/global chain by new quantitative claims. Its role is organizational: it packages OP 3 as local cell compatibility, OP 5 as defect transport, visible sectors as liquid-crystal domains, Julia-type visuals as holographic boundary readout of one common carrier order, and tests a first addressed role layer in which the 0-band is reserved for primitive ZF↔QAM transport strata beneath later derived role families.*

Remark H4.244 (Status of the compressed addressed crystal reserve surface in the present release line)**.** *The earlier addressed-role pilot, the 0-band transport reservation, the locally licensed mixed addressed tensors, and the packet/corridor/hologram/readout-class ladder have now served their main purpose: they show that one closure-stable carrier may support visible domain phase, local defect loading, and holographic boundary readout without introducing a second primitive bulk family. In the present compressed line these ingredients are therefore retained only through one finite addressed crystal reserve surface.*

Definition H4.245 (Compressed addressed crystal reserve token)**.** *A* compressed addressed crystal reserve token *is a finite package*

$$\mathfrak{A}_{\mathrm{res}}^{\mathrm{cr}} := \langle \mathcal{B}_0,\, \mathfrak{M}_{\mathrm{pil}},\, \mathfrak{C}_{\mathrm{cr}}^{\mathrm{term}} \rangle$$

with the following intended reading.

(i) $\mathcal{B}_0$ *is the reserved primitive transport substrate between ZF and QAM;*

(ii) $\mathfrak{M}_{\mathrm{pil}}$ *denotes the finite licensed addressed mixed-tensor pilot family consisting of the carrier–domain, domain–defect, and carrier–boundary couplings above that substrate;*

(iii) $\mathfrak{C}_{\mathrm{cr}}^{\mathrm{term}}$ *is the terminal contraction-stable addressed crystal readout object that retains only the carrier/domain/defect/boundary content needed for later citation.*

The token is used only as a compressed reserve object. It does not claim a global unrestricted crystal calculus.

Proposition H4.246 (Compressed addressed crystal reserve package)**.** *Whenever a later argument uses only the finite addressed-crystal content encoded by*

$$\mathfrak{A}_{\mathrm{res}}^{\mathrm{cr}},$$

it may cite this single reserve token instead of reopening the full addressed-role pilot, the three local mixed addressed tensor pilots, or the packet/corridor/hologram/class-normal-form ladder. In particular, the compressed reserve token is citation-equivalent to the removed detailed reserve block for every argument that depends only on:

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(i) *the primitive 0-band transport support,*

(ii) *the three locally licensed mixed addressed couplings,*

(iii) *one closure-stable carrier line together with its visible domain-face data,*

(iv) *the corresponding local defect-load data,*

(v) *the retained holographic boundary profile,*

(vi) *and the nonredundant interface data surviving passage to the terminal contraction-stable readout object.*

Proof. The removed detailed block introduced exactly these ingredients in layered form. First it reserved the 0-band as primitive transport substrate and treated later addressed families as derived above it. Second it licensed only three local mixed addressed tensor couplings: carrier–domain, domain–defect, and carrier–boundary. Third it unfolded a packet/corridor/hologram/readout-class ladder whose only later use was to justify passage to a terminal contraction-stable object carrying the retained carrier/domain/defect/boundary data. Therefore every later argument that needs only this finite retained content may cite the single reserve token without reopening the intermediate pilot steps. No stronger global claim is being made. □

Corollary H4.247 (Use rule for the compressed addressed crystal reserve surface)**.** *A later proof may remain entirely on the compressed addressed crystal reserve surface as long as it needs only the finite retained content listed in Proposition H4.246. A finer reopening is required exactly when packet-specific, corridor-specific, subtype-specific, or pre-contraction addressed information is needed again.*

Proof. This is the operational restatement of Proposition H4.246. □

Lemma H4.248 (Closed antisymmetry of deficiency elements)**.** *Let f be a deficiency element in the relevant HC/closure test sector. If the J-symmetric subspace is already essentially self-adjoint on the admissible core, then the J-symmetric part of f vanishes. Equivalently,*

$$f = f^- \qquad \textit{and hence} \qquad Jf = -f.$$

Proof. Write

$$f = f^+ + f^-, \qquad f^+ = \tfrac{1}{2}(f + Jf), \qquad f^- = \tfrac{1}{2}(f - Jf).$$

By construction, f^+ lies in the J-symmetric sector. The repaired Section 12 argument shows that once the admissible J-symmetric core is already essentially self-adjoint, a deficiency element cannot retain a nonzero J-symmetric component. Otherwise one would obtain a nontrivial deficiency vector in the symmetric core itself, contradicting essential self-adjointness. Hence $f^+ = 0$, so $f = f^-$ and therefore $Jf = -f$. □

Proposition H4.249 (Functional annihilation of deficiency elements by HC)**.** *For every deficiency element f in the closed HC test sector, the compensator annihilates f at the function level:*

$$C(f) = \tfrac{1}{2}(f + Jf) = 0.$$

Proof. By Lemma H4.248, every such deficiency element satisfies $Jf = -f$. Substituting into the defining compensator formula gives

$$C(f) = \tfrac{1}{2}(f + Jf) = \tfrac{1}{2}(f - f) = 0.$$

So the annihilation is a genuine functional consequence of closed J-antisymmetry, not merely a heuristic tensor slogan. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Remark H4.250 (Tensor representations are secondary to the functional proof)**.** *Any tensor or matrix representative of the compensator is basis-dependent and should be read only as a representation of the already established function-level map. In particular, the annihilation of deficiency elements is proved by the closed antisymmetry route*

$$f^+ = 0 \Longrightarrow Jf = -f \Longrightarrow C(f) = 0,$$

not by a standalone coefficient-level matrix argument.

Lemma H4.251 (Anchor-compatibility lemma)**.** *Compensator action is compatible with the orthogonal anchor. In particular, HC stabilizes the admissible readout channel without destroying the distinction between the anchored third direction and the first two sectors.*

Proof. By Lemma 29.12, the third sector is not decorative but the stabilizing orthogonal anchor. By Theorem 29.15, HC is the mechanism that re-identifies shifted descriptions in a common output channel. If HC destroyed the anchor distinction, then the stabilized output would collapse back into an undifferentiated binary pairing and the role of the third direction would be lost. That would contradict triolic minimality and the very need for an orthogonal anchor. Therefore HC must act compatibly with the anchored triolic organization. □

Lemma H4.252 (Closure forcing lemma)**.** *A triolic state with orthogonal anchor and forced HC is closure-stable. Equivalently,*

$$\mathrm{Triolic}(x) \wedge \mathrm{Anchored}(x) \wedge \mathrm{HCForced}(x) \Longrightarrow \mathrm{ClosureStable}(x).$$

Proof. We verify the three contributing conditions and their combination.

Step 1: Triolic structure. By Lemma 29.11, a triolic state carries a three-sector organization (K_1, K_2, K_3) as the minimal nondegenerate structure compatible with compensator action and closure readout. In particular, there exists a distinguished third sector K_3 in which the compensator-selected output can be stabilized.

Step 2: Orthogonal anchor. By Lemma 29.12, the third sector satisfies $K_3 \perp K_1$ and $K_3 \perp K_2$, preventing K_3 from being relabeled or absorbed into the dual pair (K_1, K_2). The anchor fixes the admissible direction of collapse/light selection.

Step 3: HC forcing. By Lemma 29.18, the compensator-normalized image $C(x)$ is admissible whenever x is admissible. Since $\mathrm{HCForced}(x)$ holds, the state lies in the compensator-selected channel, so $\mathcal{C}x = C(x)$.

Step 4: Closure stability. By the semantic–tensor correspondence of Remark 29.9, $\mathrm{ClosureStable}(x)$ is equivalent to $\hat{K}(\mathcal{C}x) = \mathcal{C}x$. By Step 3, $\mathcal{C}x = C(x)$, and by the idempotence of C (Lemma 29.17), $C(C(x)) = C(x)$, so $C(x)$ already lies in $\mathrm{im}(C)$. By Steps 1–2 and Theorem 14.3 ($\hat{K}(L_n) = L_{n-1}$, terminating at S_0), the anchor fixes K_3 as the structural ground sector; $\mathcal{C}x \in \mathrm{im}(C)$ lies in the K_3-aligned closure-stable class and is therefore a fixpoint of $\hat{K}$ on that class. Hence $\hat{K}(\mathcal{C}x) = \mathcal{C}x$, i.e. $\mathrm{ClosureStable}(x)$. □ □

Lemma H4.253 (Closure-stable iff compensator image)**.** *Under the hypotheses of the normalized semantic kernel (Definition 29.4), the following are equivalent for any admissible state x:*

(i) $\mathrm{ClosureStable}(x)$*, i.e.* $\hat{K}(\mathcal{C}x) = \mathcal{C}x$*;*

(ii) $x \in \mathrm{im}(C)$*, i.e.* $\mathcal{C}x = x$*.*

Hence $\mathrm{im}(C) = \mathcal{S}_0 := \{x \in \mathcal{S} : \mathrm{ClosureStable}(x)\}$.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. **(ii) ⇒ (i).** Suppose $\mathcal{C}x = x$. By Lemma 29.17, $C^2 = C$, so $x \in \mathrm{im}(C)$ implies $\mathcal{C}x = x$. By the ground-state terminality condition (S2) of Definition 29.4, $\hat{K}(x) = x$ for all $x \in S_0$, the photon ground sector. Since $\mathrm{im}(C)$ is mapped by $\hat{K}$ to the ground sector (Theorem 14.3: $\hat{K}(L_n) = L_{n-1}$, terminating at S_0), and since $\mathcal{C}x = x$ means x is already in the compensator-selected ground sector, we get $\hat{K}(\mathcal{C}x) = \hat{K}(x) = x = \mathcal{C}x$. Hence ClosureStable($x$).

(i) ⇒ (ii). Suppose $\hat{K}(\mathcal{C}x) = \mathcal{C}x$, i.e. $\mathcal{C}x$ is a fixpoint of $\hat{K}$. By Theorem 14.3, the only fixpoints of $\hat{K}$ in the cascade are the elements of the ground sector $S_0 = L = \bigcap_{n\geq 0} L_n$. Hence $\mathcal{C}x \in S_0$. By the semantic–tensor correspondence (Remark 29.9), $\mathcal{C}x \in S_0$ is precisely the condition that $\mathcal{C}$ has already reached its fixpoint on x, i.e. $\mathcal{C}(\mathcal{C}x) = \mathcal{C}x$. Since $C^2 = C$ (Lemma 29.17), this means $\mathcal{C}x \in \mathrm{im}(C)$, but also that x itself is mapped to the same compensator-fixed point. The re-identification mechanism (Theorem 29.15) selects precisely those x for which the compensated image is the state itself in the ground sector: $\mathcal{C}x = x$. Hence $x \in \mathrm{im}(C)$.

Conclusion. Both directions hold, so ClosureStable(x) $\Leftrightarrow$ $x \in \mathrm{im}(C)$, giving $\mathcal{S}_0 = \mathrm{im}(C)$. □

Proposition H4.254 (Closure class proposition)**.** *The closure-stable states form a structurally distinguished admissibility class. In particular, closure-stability is not an accidental property of isolated examples but a reusable criterion for admissible dynamics and readout.*

Proof. By Lemma H4.252, closure-stability follows from a fixed combination of generative conditions rather than from a one-off construction. By Lemma 29.18, the relevant states remain inside the admissible body under compensator normalization, and by Lemma H4.251 this process respects the stabilizing triolic organization. Hence closure-stability is preserved under the normalized structural mechanism and can be used as a class criterion. □

Proposition H4.255 (Exact closure criterion)**.** *Exact closure is the regime in which the admissible channel selected by HC is already terminal with respect to the relevant closure test, so that no further residual compensator correction is required.*

Proof. By Lemma 29.17, compensator-normalized states do not drift under repeated HC action. Exact closure is therefore the stronger case in which the normalized state is not merely admissible but already stable under the relevant closure readout. In the tensor realization this corresponds to the absence of further leakage from the selected sector into a complementary block; semantically it means that the compensator-selected channel has already reached its terminal closure form. □

Lemma H4.256 (Closure residue lemma)**.** *Whenever exact closure fails, the failure can be measured by a residual term that records the remaining mismatch between HC-normalized admissibility and full closure stability.*

Proof. The exact-closure regime of Proposition H4.255 is defined precisely by the absence of further correction after compensator normalization. If this stronger condition fails while admissibility is still preserved, then there remains a nonzero discrepancy between the compensator-selected channel and the fully closed channel. This discrepancy is the closure residue. It vanishes in exact closure and provides the natural quantity measuring departure from that regime. □

Corollary H4.257 (First closure forcing chain)**.** *The first unfolded closure route of the normalized core is*

$$\mathrm{HC} \Longrightarrow \textit{HC idempotence/admissibility} \Longrightarrow \textit{anchor-compatible normalization}$$
$$\Longrightarrow \textit{closure forcing} \Longrightarrow \textit{closure class} \Longrightarrow \textit{exact closure or closure residue.}$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. Lemmas 29.17, 29.18, and H4.251 identify the first structural properties of forced HC. Lemma H4.252 turns these into closure stability, Proposition H4.254 promotes this to a reusable admissibility class, and Proposition H4.255 together with Lemma H4.256 separates the exact and residual regimes. □

H4.11 First guarded-reduction pilot normal form for repeated HC/OP tails

Definition H4.258 (Guarded-reduction pilot window)**.** *A* guarded-reduction pilot window *is a finite local pair*

$$\mathsf{W}_{\mathrm{red}}(B_0, B_1)$$

where B_0 is the first explicit occurrence of an HC/OP argument tail and B_1 is a later repetition of the same tail. The pair is admissible only when the following data are fixed.

(i) A ledger entry $\ell \in \mathcal{L}^{(1)}_{\mathrm{HCOP}}$ covers both occurrences.

(ii) The first occurrence B_0 states the local hypotheses, route tag, frontier guard, witness, and theorematic status.

(iii) The later occurrence B_1 repeats only the same safety, route-preservation, frontier-preservation, diagnostic-status, or no-false-certificate tail.

(iv) The later occurrence introduces no first definition, no new hypothesis, no new certificate test, no proof-critical inequality, and no theorematic status change.

The pilot window is therefore a local rewrite domain for repeated prose only.

Definition H4.259 (guarded-citation normal form)**.** *For an admissible pilot window $\mathsf{W}_{\mathrm{red}}(B_0, B_1)$, the* guarded-citation normal form *of the repeated tail B_1 is*

$$\mathrm{GRCite}_{\mathrm{HCOP}}(\ell; G, W, S) := \big\langle \mathfrak{T}_\ell,\ G_{B_0},\ W_{B_0},\ S_{B_0}; \mathrm{noCert} \big\rangle.$$

Here $\mathfrak{T}_\ell$ is the tool contract assigned by the ledger, G_{B_0} is the preserved local guard of the first occurrence, W_{B_0} is the preserved witness or first explicit statement, and S_{B_0} is the preserved status marker. The final tag noCert *records that the citation does not create an OP closure certificate.*

Lemma H4.260 (Pilot normal form is conservative)**.** *If $\mathsf{W}_{\mathrm{red}}(B_0, B_1)$ is an admissible guarded-reduction pilot window, then replacing the later repeated tail B_1 by $\mathrm{GRCite}_{\mathrm{HCOP}}(\ell; G, W, S)$ preserves the proof content of the local argument.*

Proof. By Definition H4.258, the first occurrence B_0 remains explicit and contains the hypotheses, guard, witness, and status. By Definition H4.259, the replacement of B_1 cites exactly the ledger tool contract together with pointers back to those retained data. Since B_1 is assumed to repeat only a safety, route, frontier, diagnostic, or no-false-certificate tail, no first mathematical content is removed. Hence the replacement is conservative. □

Proposition H4.261 (Allowed pilot windows in the active HC/OP spine)**.** *The following families are admissible first pilot windows for guarded reduction in the active HC/OP spine:*

(i) lock/capture reminder windows governed by ℓ_{lock};

(ii) handoff and normalization reminder windows governed by ℓ_{handoff} and ℓ_{norm};

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(iii) route/frontier reminder windows governed by ℓ_{route}*;*

(iv) paired-comparison status windows governed by ℓ_{cmp}*;*

(v) probe diagnostic windows governed by ℓ_{probe}*;*

(vi) archive-to-tool extraction windows governed by ℓ_{archive}*.*

Each family is reducible only after its first local occurrence has been preserved.

Proof. These are precisely the ledger sites of Definition H4.231. Lemma H4.233 proves that later occurrences covered by those entries are reducible when first occurrence, guard, witness, and theorematic status remain visible. Definition H4.258 adds no new reducible class; it only names the local two-occurrence window used by the pilot rewrite. □

Definition H4.262 (Pilot tail-compression operator)**.** *The* pilot tail-compression operator

$$\mathsf{TailRed}^{(1)}_{\mathrm{HCOP}}$$

is the partial operator that maps an admissible pilot window to its guarded-citation normal form:

$$\mathsf{TailRed}^{(1)}_{\mathrm{HCOP}}\big(\mathsf{W}_{\mathrm{red}}(B_0,B_1)\big) := \mathrm{GRCite}_{\mathrm{HCOP}}(\ell;G,W,S).$$

It is undefined on first occurrences, non-reducible proof material, historical witness material, and any occurrence that changes theorematic status.

Corollary H4.263 (Pilot reduction cannot hide certificate tests)**.** *The operator* $\mathsf{TailRed}^{(1)}_{\mathrm{HCOP}}$ *cannot hide a certificate test or turn a frontier state into an OP closure state.*

Proof. Certificate tests and theorematic status changes are excluded from admissible pilot windows by Definition H4.258. The replacement normal form of Definition H4.259 also carries the explicit noCert marker. Therefore no application of $\mathsf{TailRed}^{(1)}_{\mathrm{HCOP}}$ can manufacture or hide a certificate. □

Theorem H4.264 (First guarded-reduction pilot normal form)**.** *The guarded-reduction ledger induces a conservative pilot normal form for repeated HC/OP tails. In every admissible pilot window, the later repeated tail may be replaced by* $\mathrm{GRCite}_{\mathrm{HCOP}}(\ell;G,W,S)$*, while the first occurrence, local guard, witness, proof restriction, and theorematic status remain explicit. The replacement shortens repeated prose only; it neither deletes proof content nor creates an OP closure certificate.*

Proof. Existence of the relevant ledger entries is Definition H4.231. The local rewrite domain is Definition H4.258; the citation target is Definition H4.259; and conservativity is Lemma H4.260. Corollary H4.263 gives the certificate restriction. Thus the pilot normal form is a controlled compression of repeated tails and not a new proof engine. □

Corollary H4.265 (working rule)**.** *Future HC/OP text may use* $\mathrm{GRCite}_{\mathrm{HCOP}}(\ell;G,W,S)$ *only for later repeated tails already covered by the ledger. The first occurrence, local guard, witness, and theorematic status must remain visible. Historical witness material and first mathematical content remain non-reducible.*

Proof. Immediate from Theorem H4.264. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

H4.12 First tail-normalized HC/OP proof skeleton

Definition H4.266 (Tail-normalized HC/OP proof skeleton)**.** *A* tail-normalized HC/OP proof skeleton *is a finite ordered family*

$$\mathsf{S}^{\mathrm{tail}}_{\mathrm{HCOP}} = \big((E_i, G_i, W_i, S_i; T_i)\big)_{i=0}^{m}$$

where each entry consists of

(i) a first explicit source block E_i, containing the relevant definition, hypothesis, guard, or proof-critical statement;

(ii) a local guard G_i and preserved witness or source pointer W_i;

(iii) a theorematic status marker S_i, recording whether the block is definitional, diagnostic, frontier-preserving, certificate-testing, or theorematic;

(iv) a finite family T_i of later repeated tails that may be replaced only by guarded citations of the form

$$\mathrm{GRCite}_{\mathrm{HCOP}}(\ell_i; G_i, W_i, S_i).$$

The family is admissible only when the ordering of entries agrees with the proof-dependency order of the active HC/OP spine.

Definition H4.267 (Active tail skeleton)**.** *The* active the local proof chain tail skeleton *is the tail-normalized skeleton whose source entries are ordered as*

$$\begin{aligned}
E_0 &= \mathrm{LOCK/CAPTURE}_{\mathrm{HC}},\\
E_1 &= \mathrm{HANDOFF}_{\mathrm{HC}\to\mathrm{OP}},\\
E_2 &= \mathrm{NORMALIZE}^{\mathrm{HC}}_{\mathrm{OP}},\\
E_3 &= \mathrm{ROUTE}^{\mathrm{HC}}_{\mathrm{OP}},\\
E_4 &= \mathrm{COMPARE}^{1/4|5(B)}_{\mathrm{HC}},\\
E_5 &= \mathrm{RESOLVE}_{\mathrm{bind/split/frontier}}.
\end{aligned}$$

The associated tail families T_i consist only of later repeated safety, route-preservation, frontier-preservation, diagnostic, no-false-certificate, or archive-to-tool reminders covered by the ledger.

Lemma H4.268 (Tail skeleton preserves dependency order)**.** *Let $\mathsf{S}^{\mathrm{tail}}_{\mathrm{HCOP}}$ be an admissible tail-normalized HC/OP proof skeleton. Then replacing any later tail in T_i by its guarded citation normal form preserves the dependency order of the proof spine.*

Proof. By Definition H4.266, the first explicit source block E_i, its guard G_i, witness W_i, and status marker S_i remain in the ordered skeleton. The replacement acts only on later tails attached to that source entry. Since no replacement can move a later tail before its source entry and no source entry is removed, the original dependency order is preserved. □

Lemma H4.269 (Tail skeleton does not create certificates)**.** *No admissible tail-normalized HC/OP proof skeleton can turn a diagnostic, frontier, or guarded execution state into an OP closure certificate.*

Proof. Every tail replacement is a guarded citation of the form defined in Definition H4.259; it carries the explicit noCert marker. Certificate tests and theorematic status changes are not eligible tail material by Definition H4.258. Therefore the skeleton can reorganize later repeated prose, but it cannot change theorematic status. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proposition H4.270 (Active HC/OP spine admits a tail-normalized skeleton)**.** *The active the local proof chain HC/OP proof spine admits the tail-normalized skeleton of Definition H4.267.*

Proof. The source entries are exactly the tool-contract application sites of the active HC/OP spine: lock/capture, handoff, normalization, routing, paired comparison, and resolution. The ledger identifies the corresponding repeated tail classes, while the pilot normal form gives the admissible guarded citation for later repetitions. Hence these source entries and tail families form an admissible ordered skeleton. □

Corollary H4.271 (Tail-normalized reuse rule)**.** *Future HC/OP text may cite the active tail skeleton only after the relevant source entry has appeared explicitly. A citation to the skeleton may replace a later repeated tail, but it may not replace a first definition, a first guard, a witness, a certificate test, a proof-critical inequality, or a theorematic status change.*

Proof. This is the direct reuse form of Lemmas H4.268 and H4.269. □

Theorem H4.272 (First tail-normalized HC/OP proof skeleton)**.** *The local proof chain MML/MMA compression, tool-contract, ledger, and guarded-citation layers induce a first conservative tail-normalized proof skeleton for the active HC/OP spine. The skeleton shortens only later repeated HC/OP tails by guarded citation, preserves all first source blocks, preserves dependency order, and cannot manufacture an OP closure certificate.*

Proof. The compression map is supplied by the layer, the tool contracts by the layer, their application to the active spine by the layer, the reduction discipline by the protocol, the finite citation sites by the ledger, and the guarded citation normal form by the pilot. Proposition H4.270 identifies the corresponding ordered skeleton for the local proof chain HC/OP spine. Lemma H4.268 proves dependency preservation, and Lemma H4.269 proves the certificate restriction. Thus the skeleton is conservative. □

Corollary H4.273 (working rule)**.** *The active HC/OP proof spine may now be cited in tail-normalized form as*

$$\begin{aligned}\mathsf{S}^{\mathrm{tail}}_{\mathrm{HCOP}} = (&\mathrm{LOCK/CAPTURE}, \mathrm{HANDOFF}, \mathrm{NORMALIZE},\\ &\mathrm{ROUTE}, \mathrm{COMPARE}, \mathrm{RESOLVE}).\end{aligned}$$

After the first occurrence of each source block has been stated, this citation is a compact reference form for later repeated tails only.

Proof. Immediate from Theorem H4.272. □

H4.13 First skeleton-to-workstep HC/OP dispatch

Definition H4.274 (Skeleton-to-workstep dispatch object)**.** *Let* $\mathsf{S}^{\mathrm{tail}}_{\mathrm{HCOP}}$ *be the active tail-normalized HC/OP proof skeleton of Definition H4.267. A* skeleton-to-workstep dispatch object *is a finite map*

$$\mathsf{D}^{\mathrm{skel}}_{\mathrm{HCOP}} : \{E_0, E_1, E_2, E_3, E_4, E_5\} \longrightarrow \mathsf{Work}_{\mathrm{HCOP}}$$

which assigns to each explicit source node one admissible family of follow-on work packets. The map is admissible only if the assigned packets preserve the source guard G_i*, witness pointer* W_i*, theorematic status marker* S_i*, and the no-false-certificate restriction inherited from the guarded-citation layer.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition H4.275 (First dispatch menu)**.** *The first dispatch menu is the following finite assignment:*

$$\begin{aligned} E_0 = \mathrm{LOCK/CAPTURE}_{\mathrm{HC}} &\longmapsto \mathsf{W}^{\mathrm{stab}}_{\mathrm{lock}}, \\ E_1 = \mathrm{HANDOFF}_{\mathrm{HC}\to\mathrm{OP}} &\longmapsto \mathsf{W}^{\mathrm{guard}}_{\mathrm{handoff}}, \\ E_2 = \mathrm{NORMALIZE}^{\mathrm{HC}}_{\mathrm{OP}} &\longmapsto \mathsf{W}^{\mathrm{type}}_{\mathrm{norm}}, \\ E_3 = \mathrm{ROUTE}^{\mathrm{HC}}_{\mathrm{OP}} &\longmapsto \mathsf{W}^{\mathrm{rank}}_{\mathrm{route}}, \\ E_4 = \mathrm{COMPARE}^{1/4|5(B)}_{\mathrm{HC}} &\longmapsto \mathsf{W}^{\Delta}_{\mathrm{cmp}}, \\ E_5 = \mathrm{RESOLVE}_{\mathrm{bind/split/frontier}} &\longmapsto \mathsf{W}^{\mathrm{next}}_{\mathrm{resolve}}. \end{aligned}$$

Here $\mathsf{W}^{\mathrm{stab}}_{\mathrm{lock}}$ *checks stability of the local capture source,* $\mathsf{W}^{\mathrm{guard}}_{\mathrm{handoff}}$ *checks whether handoff guards remain visible,* $\mathsf{W}^{\mathrm{type}}_{\mathrm{norm}}$ *checks type-correct normalization,* $\mathsf{W}^{\mathrm{rank}}_{\mathrm{route}}$ *ranks the routed residuals,* $\mathsf{W}^{\Delta}_{\mathrm{cmp}}$ *acts on the paired comparison defect, and* $\mathsf{W}^{\mathrm{next}}_{\mathrm{resolve}}$ *selects the next conservative branch without changing theorematic status.*

Lemma H4.276 (Dispatch preserves the tail-skeleton order)**.** *Every admissible skeleton-to-workstep dispatch object preserves the dependency order of the active HC/OP proof spine.*

Proof. By Definition H4.274, a dispatch packet is attached to an already ordered source node E_i of the tail skeleton. It does not move the source node, delete its guard, or replace a preceding dependency by a later citation. Therefore the order established in Lemma H4.268 is inherited by the dispatched work packets. □

Lemma H4.277 (Dispatch is work-generating, not certificate-generating)**.** *No admissible skeleton-to-workstep dispatch object can manufacture an OP closure certificate.*

Proof. The dispatch object inherits the no-false-certificate marker from the guarded-citation layer and from Lemma H4.269. Its codomain consists of follow-on work packets, not theorematic closure certificates. A certificate may only appear after a separate explicit certificate test with its own hypotheses, guards, and proof-critical inequalities. Hence dispatch creates controlled work, not closure. □

Proposition H4.278 (First controlled next-work selection from the HC/OP skeleton)**.** *The active tail-normalized HC/OP proof skeleton admits the dispatch menu of Definition H4.275. In particular, the immediate admissible follow-on work is finite and restricted to stabilization, handoff-guard checking, type-normalization checking, route ranking, comparison-defect work, and conservative branch selection.*

Proof. Each menu entry is attached to one of the ordered source nodes of Definition H4.267. The first four entries refine already explicit source nodes and therefore preserve their guards and witnesses. The comparison entry $\mathsf{W}^{\Delta}_{\mathrm{cmp}}$ is tied to the paired comparison layer and may only act on the already defined comparison defect. The resolution entry selects a conservative next branch from the already permitted outcomes bind, split, or frontier. By Lemmas H4.276 and H4.277, the whole menu is conservative. □

Theorem H4.279 (First skeleton-to-workstep HC/OP dispatch)**.** *The tail-normalized HC/OP proof skeleton induces a first conservative skeleton-to-workstep dispatch object. This dispatch turns the compact proof skeleton into a finite menu of admissible next work packets while preserving dependency order, guards, witnesses, route tags, and the no-false-certificate rule.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. Definition H4.274 specifies the admissibility requirements for dispatch. Definition H4.275 provides the first finite menu. Proposition H4.278 proves that this menu is admissible for the active skeleton. The order and certificate restrictions follow from Lemmas H4.276 and H4.277. □

Corollary H4.280 (working rule)**.** *Future HC/OP work may be launched from the tail-normalized skeleton only through an explicit dispatch packet*

$$\mathsf{D}^{\mathrm{skel}}_{\mathrm{HCOP}}(E_i) = \mathsf{W}_i.$$

Such a packet records what is to be checked next; it does not by itself prove OP 1/4, OP 5(B), or any closure certificate.

Proof. Immediate from Theorem H4.279. □

H4.14 First dispatched lock/capture stabilization packet

Definition H4.281 (Dispatched lock/capture stabilization packet)**.** *Let* $E_0 = \mathrm{LOCK/CAPTURE}_{\mathrm{HC}}$ *be the first node of the tail-normalized HC/OP skeleton and let*

$$\mathsf{D}^{\mathrm{skel}}_{\mathrm{HCOP}}(E_0) = \mathsf{W}^{\mathrm{stab}}_{\mathrm{lock}}$$

be the dispatch assignment. A dispatched lock/capture stabilization packet *is a finite packet*

$$\mathsf{P}^{\mathrm{stab}}_{\mathrm{lock}} = \big(\mathcal{C}^{\mathrm{HC}}_n, G_0, W_0, S_0, \mathcal{A}_0, \mathbf{d}^{\mathrm{stab}}_{\mathrm{lock}}\big),$$

where $\mathcal{C}^{\mathrm{HC}}_n$ *is the protected local HC-capture chain,* G_0 *is the local guard,* W_0 *is the witness pointer to the first explicit source block,* S_0 *is the non-certificate status marker,* $\mathcal{A}_0$ *is the finite local audit window, and* $\mathbf{d}^{\mathrm{stab}}_{\mathrm{lock}}$ *is a non-negative lock-stability defect register. The packet is admissible only if it does not widen the local corridor, erase the route tag, replace* W_0*, or change* S_0 *into a theorematic certificate.*

Definition H4.282 (Lock-stability defect register)**.** *For a dispatched packet* $\mathsf{P}^{\mathrm{stab}}_{\mathrm{lock}}$*, the* lock-stability defect register *is the finite family*

$$\mathbf{d}^{\mathrm{stab}}_{\mathrm{lock}}(j) = \big(d_{\mathrm{leak}}(j), d_{\mathrm{corr}}(j), d_{\mathrm{guard}}(j)\big), \qquad j \in \mathcal{A}_0,$$

where d_{leak} *measures re-opening of lock-side leakage,* d_{corr} *measures corridor drift, and* d_{guard} *measures loss of the local guard. The associated scalar audit defect is*

$$\Delta^{\mathrm{stab}}_{\mathrm{lock}} = \max_{j \in \mathcal{A}_0} \big(d_{\mathrm{leak}}(j) + d_{\mathrm{corr}}(j) + d_{\mathrm{guard}}(j)\big).$$

The register is called stable *if* $\Delta^{\mathrm{stab}}_{\mathrm{lock}} = 0$*,* repairable *if a permitted local micro-lock refinement strictly lowers the scalar audit defect without widening the corridor, and* frontier-bound *otherwise.*

Lemma H4.283 (Zero lock-stability defect preserves the capture source)**.** *If* $\Delta^{\mathrm{stab}}_{\mathrm{lock}} = 0$ *for an admissible dispatched lock/capture stabilization packet, then the protected local HC-capture chain remains a valid capture source throughout the finite audit window.*

Proof. Vanishing scalar audit defect means that every component of $\mathbf{d}^{\mathrm{stab}}_{\mathrm{lock}}(j)$ vanishes on $\mathcal{A}_0$. Hence the lock-side leakage is not re-opened, the corridor does not drift, and the local guard is not lost. The packet also preserves the witness pointer and the non-certificate status marker by Definition H4.281. Therefore the original local HC-capture chain remains the same protected capture source, rather than a replaced or widened source. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition H4.284 (lock/capture stabilization return)**.** *The execution of* $\mathsf{W}_{\mathrm{lock}}^{\mathrm{stab}}$ *on an admissible packet returns exactly one of the following statuses:*

$$\mathrm{Exec}\big(\mathsf{W}_{\mathrm{lock}}^{\mathrm{stab}}, \mathsf{P}_{\mathrm{lock}}^{\mathrm{stab}}\big) \in \{\mathsf{L}_{\mathrm{stable}}, \mathsf{L}_{\mathrm{repair}}, \mathsf{L}_{\mathrm{frontier}}\}.$$

Here $\mathsf{L}_{\mathrm{stable}}$ *means that the capture source is handoff-ready for the next skeleton node,* $\mathsf{L}_{\mathrm{repair}}$ *means that one bounded local repair step is admissible before handoff is tested again, and* $\mathsf{L}_{\mathrm{frontier}}$ *means that the source remains protected but must be kept as a smaller frontier source rather than promoted.*

Proposition H4.285 (The first dispatched lock/capture packet is admissible)**.** *The dispatch menu admits the lock/capture stabilization packet as the first executable work packet attached to the active HC/OP skeleton.*

Proof. The dispatch menu assigns E_0 only to $\mathsf{W}_{\mathrm{lock}}^{\mathrm{stab}}$. The data required in Definition H4.281 are inherited from the first explicit lock/capture source block, the guarded-reduction ledger, and the tail-normalized skeleton: the protected capture chain, the local guard, the witness pointer, and the non-certificate marker are all already part of the active spine. The new defect register adds only a finite audit surface for this first packet. It does not replace a hypothesis, alter a route tag, or create a certificate. Hence the packet is admissible. □

Theorem H4.286 (First dispatched lock/capture stabilization packet)**.** *The active HC/OP skeleton now supports a first executed work packet*

$$E_0 \xrightarrow{\mathsf{W}_{\mathrm{lock}}^{\mathrm{stab}}} \{\mathsf{L}_{\mathrm{stable}}, \mathsf{L}_{\mathrm{repair}}, \mathsf{L}_{\mathrm{frontier}}\}.$$

This execution is conservative: it either prepares the protected capture source for the handoff node, records one bounded local repair obligation, or returns a smaller frontier source. It does not prove OP 1/4, OP 5(B), or any closure certificate.

Proof. By Proposition H4.285, the first dispatched packet is admissible for the active skeleton. Definition H4.282 classifies the finite audit defect into the stable, repairable, or frontier-bound cases, and Definition H4.284 records the corresponding return status. Lemma H4.283 proves the stable case. In the repairable case the packet remains at the same skeleton node with a strictly smaller local defect and no corridor widening; in the frontier-bound case it is returned as protected frontier material without promotion. In none of the three cases is the non-certificate marker changed into an OP closure certificate. □

Corollary H4.287 (working rule)**.** *Future work may proceed from the dispatch menu by first executing* $\mathsf{W}_{\mathrm{lock}}^{\mathrm{stab}}$*. The next admissible mathematical question is therefore whether the return status is handoff-ready, locally repairable, or frontier-bound; it is not yet whether OP 1/4 or OP 5(B) is closed.*

Proof. Immediate from Theorem H4.286. □

H4.15 First dispatched handoff-guard packet

Definition H4.288 (Dispatched handoff-guard packet)**.** *Let* $E_1 = \mathrm{HANDOFF}_{\mathrm{HC}\to\mathrm{OP}}$ *be the second node of the tail-normalized HC/OP skeleton and let*

$$\mathsf{D}_{\mathrm{HCOP}}^{\mathrm{skel}}(E_1) = \mathsf{W}_{\mathrm{handoff}}^{\mathrm{guard}}$$

be the dispatch assignment. A dispatched handoff-guard packet *is a finite packet*

$$\mathsf{P}_{\mathrm{handoff}}^{\mathrm{guard}} = \big(\mathsf{L}_{\mathrm{stable}}, \mathcal{P}_{\mathrm{handoff}}, G_1, W_1, S_1, \mathbf{d}_{\mathrm{handoff}}^{\mathrm{guard}}\big),$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

where $\mathsf{L}_{\mathrm{stable}}$ *is the stable lock/capture return status of Definition H4.284,* $\mathcal{P}_{\mathrm{handoff}}$ *is the inherited HC-to-OP handoff packet,* G_1 *is the handoff guard,* W_1 *is the witness pointer to the first explicit handoff source block,* S_1 *is the non-certificate status marker, and* $\mathbf{d}^{\mathrm{guard}}_{\mathrm{handoff}}$ *is a non-negative handoff-guard defect register. The packet is admissible only if it is launched from* $\mathsf{L}_{\mathrm{stable}}$*, keeps the lock/capture witness visible, preserves the HC-to-OP route tag, and does not convert handoff-readiness into an OP closure certificate.*

Definition H4.289 (Handoff-guard defect register)**.** *For an admissible packet* $\mathsf{P}^{\mathrm{guard}}_{\mathrm{handoff}}$*, the* handoff-guard defect register *is the finite family*

$$\mathbf{d}^{\mathrm{guard}}_{\mathrm{handoff}}(j) = \big(d_{\mathrm{src}}(j), d_{\mathrm{tag}}(j), d_{\mathrm{wit}}(j), d_{\mathrm{stat}}(j)\big), \qquad j \in \mathcal{A}_1,$$

where d_{src} *measures loss of the stabilized capture source,* d_{tag} *measures drift of the HC-to-OP route tag,* d_{wit} *measures loss of the handoff witness pointer, and* d_{stat} *measures accidental promotion of the non-certificate status marker. The scalar handoff audit defect is*

$$\Delta^{\mathrm{guard}}_{\mathrm{handoff}} = \max_{j\in\mathcal{A}_1}\big(d_{\mathrm{src}}(j) + d_{\mathrm{tag}}(j) + d_{\mathrm{wit}}(j) + d_{\mathrm{stat}}(j)\big).$$

The register is handoff-clean *if* $\Delta^{\mathrm{guard}}_{\mathrm{handoff}} = 0$*,* guard-repairable *if a bounded local handoff-guard repair strictly lowers the scalar defect while preserving the stable capture source and route tag, and* frontier-bound *otherwise.*

Lemma H4.290 (Stable lock/capture return is the handoff precondition)**.** *The handoff-guard packet may be executed only from the stable lock/capture return status* $\mathsf{L}_{\mathrm{stable}}$*. If the return status is* $\mathsf{L}_{\mathrm{repair}}$ *or* $\mathsf{L}_{\mathrm{frontier}}$*, then the handoff node is not yet admissible.*

Proof. By Definition H4.288, the first entry of $\mathsf{P}^{\mathrm{guard}}_{\mathrm{handoff}}$ is $\mathsf{L}_{\mathrm{stable}}$. A repair return means that the first skeleton node still carries a bounded local obligation, while a frontier return means that the protected source is not promoted. Neither status supplies the handoff-ready source required at the second skeleton node. Thus handoff execution is blocked until the packet has returned the stable status. □

Lemma H4.291 (Zero handoff-guard defect preserves HC-to-OP handoff)**.** *If* $\Delta^{\mathrm{guard}}_{\mathrm{handoff}} = 0$ *for an admissible dispatched handoff-guard packet, then the stabilized HC-capture source passes to the HC-to-OP handoff packet with its route tag, witness pointer, and non-certificate status marker preserved.*

Proof. Vanishing scalar defect forces all four defect components in Definition H4.289 to vanish on the audit window. Hence the stabilized source is not lost, the HC-to-OP tag does not drift, the witness pointer remains visible, and the non-certificate marker is not promoted. The packet therefore transfers the already stabilized source to the handoff node without changing its theorematic status. □

Definition H4.292 (handoff-guard return)**.** *The execution of* $\mathsf{W}^{\mathrm{guard}}_{\mathrm{handoff}}$ *on an admissible packet returns exactly one of the following statuses:*

$$\mathrm{Exec}\big(\mathsf{W}^{\mathrm{guard}}_{\mathrm{handoff}}, \mathsf{P}^{\mathrm{guard}}_{\mathrm{handoff}}\big) \in \{\mathsf{H}_{\mathrm{ready}}, \mathsf{H}_{\mathrm{repair}}, \mathsf{H}_{\mathrm{frontier}}\}.$$

Here $\mathsf{H}_{\mathrm{ready}}$ *means that the stabilized capture source is ready for OP-normalization testing,* $\mathsf{H}_{\mathrm{repair}}$ *means that one bounded handoff-guard repair is required before normalization may be tested, and* $\mathsf{H}_{\mathrm{frontier}}$ *means that the source remains protected but must return to the smaller frontier state rather than move forward.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proposition H4.293 (The first dispatched handoff-guard packet is admissible after stable lock return)**.** *If the execution returns* $\mathsf{L}_{\mathrm{stable}}$*, then the dispatch menu admits the handoff-guard packet as the next executable work packet attached to the active HC/OP skeleton.*

Proof. The dispatch menu assigns E_1 only to $\mathsf{W}^{\mathrm{guard}}_{\mathrm{handoff}}$. Lemma H4.290 supplies the required stable input status. The remaining data of Definition H4.288 are inherited from the first explicit handoff source block, the active tail skeleton, and the guarded-reduction ledger: the handoff guard, witness pointer, route tag, and non-certificate marker are all already part of the active HC/OP spine. The new defect register adds only a finite audit surface for the handoff node, so the packet is admissible. □

Theorem H4.294 (First dispatched handoff-guard packet)**.** *After a stable lock/capture return, the active HC/OP skeleton supports the next executed work packet*

$$\mathsf{L}_{\mathrm{stable}} \xrightarrow{\mathsf{W}^{\mathrm{guard}}_{\mathrm{handoff}}} \{\mathsf{H}_{\mathrm{ready}}, \mathsf{H}_{\mathrm{repair}}, \mathsf{H}_{\mathrm{frontier}}\}.$$

This execution is conservative: it either prepares the stabilized source for OP-normalization testing, records one bounded handoff-guard repair obligation, or returns a smaller frontier source. It does not prove OP 1/4, OP 5(B), or any closure certificate.

Proof. By Proposition H4.293, the packet is admissible after a stable return. Definition H4.289 classifies the handoff audit defect into handoff-clean, guard-repairable, or frontier-bound cases, and Definition H4.292 records the corresponding return status. Lemma H4.291 proves the handoff-clean case. In the repair case the packet remains at the handoff node with a strictly smaller local defect and no route-tag drift; in the frontier case it returns protected frontier material without promotion. In none of the three cases is the non-certificate marker changed into an OP closure certificate. □

Corollary H4.295 (working rule)**.** *Future work may proceed from the lock/capture packet to the handoff node only after* $\mathsf{L}_{\mathrm{stable}}$ *has been returned. The next admissible mathematical question is therefore whether the stabilized source is handoff-ready for OP-normalization, handoff-guard-repairable, or frontier-bound; it is still not whether OP 1/4 or OP 5(B) is closed.*

Proof. Immediate from Theorem H4.294. □

H4.16 First dispatched OP-normalization/type packet

Definition H4.296 (Dispatched OP-normalization/type packet)**.** *Let* $E_2 = \mathrm{NORMALIZE}^{\mathrm{HC}}_{\mathrm{OP}}$ *be the third node of the tail-normalized HC/OP skeleton and let*

$$\mathsf{D}^{\mathrm{skel}}_{\mathrm{HCOP}}(E_2) = \mathsf{W}^{\mathrm{type}}_{\mathrm{norm}}$$

be the dispatch assignment. A dispatched OP-normalization/type packet *is a finite packet*

$$\mathsf{P}^{\mathrm{type}}_{\mathrm{norm}} = \big(\mathsf{H}_{\mathrm{ready}}, \mathcal{N}^{\mathrm{HC}}_{\mathrm{OP}}, (\rho_{\mathrm{CL}}, \rho_{\Theta R}, \rho_A), G_2, W_2, S_2, \mathbf{d}^{\mathrm{type}}_{\mathrm{norm}}\big),$$

where $\mathsf{H}_{\mathrm{ready}}$ *is the handoff-ready return status of Definition H4.292,* $\mathcal{N}^{\mathrm{HC}}_{\mathrm{OP}}$ *is the inherited OP-entry normal form,* $(\rho_{\mathrm{CL}}, \rho_{\Theta R}, \rho_A)$ *is the live residual triple carried by that normal form,* G_2 *is the normalization/type guard,* W_2 *is the witness pointer to the first explicit normalization source block,* S_2 *is the non-certificate status marker, and* $\mathbf{d}^{\mathrm{type}}_{\mathrm{norm}}$ *is a non-negative type-coherence defect register. The packet is admissible only if it is launched from* $\mathsf{H}_{\mathrm{ready}}$*, keeps the HC-to-OP route and handoff witness visible, preserves the live residual triple as residual data, and does not convert a normalized OP input into an OP closure certificate.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition H4.297 (Normalization/type-coherence defect register)**.** *For an admissible packet* $\mathsf{P}^{\mathrm{type}}_{\mathrm{norm}}$*, the* normalization/type-coherence defect register *is the finite family*

$$\mathbf{d}^{\mathrm{type}}_{\mathrm{norm}}(j) = \big(d_{\mathrm{obj}}(j), d_{\mathrm{type}}(j), d_{\mathrm{res}}(j), d_{\mathrm{tag}}(j), d_{\mathrm{stat}}(j)\big), \qquad j \in \mathcal{A}_2,$$

where d_{obj} *measures loss of the handoff image as a QAM-coded HC/OP object,* d_{type} *measures type drift under the MML/MMA interface,* d_{res} *measures loss or premature splitting of the live residual triple,* d_{tag} *measures HC-to-OP route-tag drift, and* d_{stat} *measures accidental promotion of the non-certificate status marker. The scalar normalization audit defect is*

$$\Delta^{\mathrm{type}}_{\mathrm{norm}} = \max_{j\in\mathcal{A}_2}\big(d_{\mathrm{obj}}(j) + d_{\mathrm{type}}(j) + d_{\mathrm{res}}(j) + d_{\mathrm{tag}}(j) + d_{\mathrm{stat}}(j)\big).$$

The register is type-clean *if* $\Delta^{\mathrm{type}}_{\mathrm{norm}} = 0$*,* type-repairable *if one bounded local normalization/type repair strictly lowers the scalar defect while preserving the handoff-ready source and route tag, and* frontier-bound *otherwise.*

Lemma H4.298 (Handoff-ready return is the normalization precondition)**.** *The OP-normalization/type packet may be executed only from the handoff-ready return status* $\mathsf{H}_{\mathrm{ready}}$*. If the return status is* $\mathsf{H}_{\mathrm{repair}}$ *or* $\mathsf{H}_{\mathrm{frontier}}$*, then the normalization node is not yet admissible.*

Proof. By Definition H4.296, the first entry of $\mathsf{P}^{\mathrm{type}}_{\mathrm{norm}}$ is $\mathsf{H}_{\mathrm{ready}}$. A repair return still carries a bounded handoff-guard obligation, while a frontier return does not promote the source to the normalization node. Neither status supplies the handoff-ready input required for typed OP-normalization. Thus normalization execution is blocked until the packet has returned the ready status. □

Lemma H4.299 (Zero normalization/type defect preserves OP-entry data)**.** *If* $\Delta^{\mathrm{type}}_{\mathrm{norm}} = 0$ *for an admissible dispatched OP-normalization/type packet, then the handoff-ready source enters the OP-entry normal form with its QAM object code, MML/MMA type, live residual triple, route tag, witness pointer, and non-certificate status marker preserved.*

Proof. Vanishing scalar defect forces every component in Definition H4.297 to vanish on the audit window. Hence the handoff image remains a QAM-coded HC/OP object, its MML/MMA type does not drift, the residual triple remains residual data rather than a certificate, the HC-to-OP route tag is preserved, and the non-certificate marker is not promoted. The output is therefore a typed OP-entry input, not an OP closure certificate. □

Definition H4.300 (OP-normalization/type return)**.** *The execution of* $\mathsf{W}^{\mathrm{type}}_{\mathrm{norm}}$ *on an admissible packet returns exactly one of the following statuses:*

$$\mathrm{Exec}\big(\mathsf{W}^{\mathrm{type}}_{\mathrm{norm}}, \mathsf{P}^{\mathrm{type}}_{\mathrm{norm}}\big) \in \{\mathsf{N}_{\mathrm{ready}}, \mathsf{N}_{\mathrm{repair}}, \mathsf{N}_{\mathrm{frontier}}\}.$$

Here $\mathsf{N}_{\mathrm{ready}}$ *means that the handoff source is now a typed OP-entry input ready for route/rank testing,* $\mathsf{N}_{\mathrm{repair}}$ *means that one bounded normalization/type repair is required before routing may be tested, and* $\mathsf{N}_{\mathrm{frontier}}$ *means that the source remains protected but must return to a smaller frontier state rather than move forward.*

Proposition H4.301 (The dispatched OP-normalization/type packet is admissible after handoff-ready return)**.** *If the execution returns* $\mathsf{H}_{\mathrm{ready}}$*, then the dispatch menu admits the OP-normalization/type packet as the next executable work packet attached to the active HC/OP skeleton.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. The dispatch menu assigns E_2 only to $\mathsf{W}^{\mathrm{type}}_{\mathrm{norm}}$. Lemma H4.298 supplies the required handoff-ready input status. The remaining data of Definition H4.296 are inherited from the OP-entry normal form, the QAM/MML/MMA interface, the active tail skeleton, and the guarded-reduction ledger. The new defect register adds only a finite type-audit surface for the normalization node. It does not replace a hypothesis, alter a route tag, split a residual without authorization, or create a certificate. Hence the packet is admissible. □

Theorem H4.302 (First dispatched OP-normalization/type packet)**.** *After a handoff-ready return, the active HC/OP skeleton supports the next executed work packet*

$$\mathsf{H}_{\mathrm{ready}} \xrightarrow{\mathsf{W}^{\mathrm{type}}_{\mathrm{norm}}} \{\mathsf{N}_{\mathrm{ready}}, \mathsf{N}_{\mathrm{repair}}, \mathsf{N}_{\mathrm{frontier}}\}.$$

This execution is conservative: it either prepares a typed OP-entry input for route/rank testing, records one bounded normalization/type repair obligation, or returns a smaller frontier source. It does not prove OP 1/4, OP 5(B), or any closure certificate.

Proof. By Proposition H4.301, the packet is admissible after a handoff-ready return. Definition H4.297 classifies the type audit defect into type-clean, type-repairable, or frontier-bound cases, and Definition H4.300 records the corresponding return status. Lemma H4.299 proves the type-clean case. In the repair case the packet remains at the normalization node with a strictly smaller local type defect and no route-tag drift; in the frontier case it returns protected frontier material without promotion. In none of the three cases is the non-certificate marker changed into an OP closure certificate. □

Corollary H4.303 (working rule)**.** *Future work may proceed from the handoff-guard packet to the OP-normalization node only after* $\mathsf{H}_{\mathrm{ready}}$ *has been returned. The next admissible mathematical question is therefore whether the handoff source is type-clean and route/rank-ready, type-repairable, or frontier-bound; it is still not whether OP 1/4 or OP 5(B) is closed.*

Proof. Immediate from Theorem H4.302. □

H4.17 First dispatched route/rank packet

Definition H4.304 (Dispatched route/rank packet)**.** *Let* $E_3 = \mathrm{ROUTE}^{\mathrm{HC}}_{\mathrm{OP}}$ *be the fourth node of the tail-normalized HC/OP skeleton and let*

$$\mathsf{D}^{\mathrm{skel}}_{\mathrm{HCOP}}(E_3) = \mathsf{W}^{\mathrm{rank}}_{\mathrm{route}}$$

be the dispatch assignment. A dispatched route/rank packet *is a finite packet*

$$\mathsf{P}^{\mathrm{rank}}_{\mathrm{route}} = \big(\mathsf{N}_{\mathrm{ready}}, \mathcal{T}^{\mathrm{HC}}_{\mathrm{OP}}, (\rho_{\mathrm{CL}}, \rho_{\Theta R}, \rho_A), \mathcal{M}^{\mathrm{rank}}_{\mathrm{route}}, G_3, W_3, S_3, \mathbf{d}^{\mathrm{rank}}_{\mathrm{route}}\big),$$

where $\mathsf{N}_{\mathrm{ready}}$ *is the normalization-ready return status of Definition H4.300,* $\mathcal{T}^{\mathrm{HC}}_{\mathrm{OP}}$ *is the inherited HC-routed OP-entry task-routing object,* $(\rho_{\mathrm{CL}}, \rho_{\Theta R}, \rho_A)$ *is the live residual triple carried by the typed OP input,* $\mathcal{M}^{\mathrm{rank}}_{\mathrm{route}}$ *is the finite admissible route/rank menu inherited from the task-routing layer,* G_3 *is the route/rank guard,* W_3 *is the witness pointer to the first explicit route source block,* S_3 *is the non-certificate status marker, and* $\mathbf{d}^{\mathrm{rank}}_{\mathrm{route}}$ *is a non-negative route/rank defect register. The packet is admissible only if it is launched from* $\mathsf{N}_{\mathrm{ready}}$*, keeps the HC-to-OP route tag visible, keeps the residual triple as residual data, ranks only tasks already admitted by* $\mathcal{T}^{\mathrm{HC}}_{\mathrm{OP}}$*, and does not convert a routed task into an OP closure certificate.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition H4.305 (Route/rank defect register)**.** *For an admissible packet* $\mathsf{P}^{\mathrm{rank}}_{\mathrm{route}}$*, the* route/rank defect register *is the finite family*

$$\mathbf{d}^{\mathrm{rank}}_{\mathrm{route}}(j) = \big(d_{\mathrm{route}}(j), d_{\mathrm{rank}}(j), d_{\mathrm{res}}(j), d_{\mathrm{front}}(j), d_{\mathrm{stat}}(j)\big), \qquad j \in \mathcal{A}_3,$$

where d_{route} *measures HC-to-OP route-tag drift,* d_{rank} *measures illegal or unstable task ranking,* d_{res} *measures loss, collapse, or unauthorized splitting of the live residual triple,* d_{front} *measures violation of the frontier guard, and* d_{stat} *measures accidental promotion of the non-certificate status marker. The scalar route/rank audit defect is*

$$\Delta^{\mathrm{rank}}_{\mathrm{route}} = \max_{j\in\mathcal{A}_3}\big(d_{\mathrm{route}}(j) + d_{\mathrm{rank}}(j) + d_{\mathrm{res}}(j) + d_{\mathrm{front}}(j) + d_{\mathrm{stat}}(j)\big).$$

The register is route-clean *if* $\Delta^{\mathrm{rank}}_{\mathrm{route}} = 0$*,* route-repairable *if one bounded local route/rank repair strictly lowers the scalar defect while preserving the typed OP input and route tag, and* frontier-bound *otherwise.*

Lemma H4.306 (Normalization-ready return is the route/rank precondition)**.** *The route/rank packet may be executed only from the normalization-ready return status* $\mathsf{N}_{\mathrm{ready}}$*. If the return status is* $\mathsf{N}_{\mathrm{repair}}$ *or* $\mathsf{N}_{\mathrm{frontier}}$*, then the route/rank node is not yet admissible.*

Proof. By Definition H4.304, the first entry of $\mathsf{P}^{\mathrm{rank}}_{\mathrm{route}}$ is $\mathsf{N}_{\mathrm{ready}}$. A repair return still carries a bounded normalization/type obligation, while a frontier return does not promote the source to the routing node. Neither status supplies the typed OP input required for route/rank testing. Thus route/rank execution is blocked until the packet has returned the ready status. □

Lemma H4.307 (Zero route/rank defect preserves routed OP-task data)**.** *If* $\Delta^{\mathrm{rank}}_{\mathrm{route}} = 0$ *for an admissible dispatched route/rank packet, then the normalization-ready OP input passes to a routed task state with its HC-to-OP route tag, admissible task ranking, live residual triple, frontier guard, witness pointer, and non-certificate status marker preserved.*

Proof. Vanishing scalar defect forces every component in Definition H4.305 to vanish on the audit window. Hence no route tag drifts, no inadmissible task is ranked ahead of the inherited task-routing menu, the residual triple remains residual data, the frontier guard stays active, and the non-certificate marker is not promoted. The output is therefore a routed OP task state ready for the next comparison or repair decision, not an OP closure certificate. □

Definition H4.308 (route/rank return)**.** *The execution of* $\mathsf{W}^{\mathrm{rank}}_{\mathrm{route}}$ *on an admissible packet returns exactly one of the following statuses:*

$$\mathrm{Exec}\big(\mathsf{W}^{\mathrm{rank}}_{\mathrm{route}}, \mathsf{P}^{\mathrm{rank}}_{\mathrm{route}}\big) \in \{\mathsf{R}_{\mathrm{ready}}, \mathsf{R}_{\mathrm{repair}}, \mathsf{R}_{\mathrm{frontier}}\}.$$

Here $\mathsf{R}_{\mathrm{ready}}$ *means that the typed OP input has been routed and ranked into an admissible comparison-ready task state,* $\mathsf{R}_{\mathrm{repair}}$ *means that one bounded route/rank repair is required before comparison may be tested, and* $\mathsf{R}_{\mathrm{frontier}}$ *means that the source remains protected but must return to a smaller frontier state rather than move forward.*

Proposition H4.309 (The dispatched route/rank packet is admissible after normalization-ready return)**.** *If the execution returns* $\mathsf{N}_{\mathrm{ready}}$*, then the dispatch menu admits the route/rank packet as the next executable work packet attached to the active HC/OP skeleton.*

Proof. The dispatch menu assigns E_3 only to $\mathsf{W}^{\mathrm{rank}}_{\mathrm{route}}$. Lemma H4.306 supplies the required normalization-ready input status. The remaining data of Definition H4.304 are inherited from the task-routing layer, the QAM/MML/MMA interface, the skeleton, and the guarded-reduction ledger. The new defect register adds only a finite route/rank audit surface. It does not introduce a new task outside the inherited menu, alter a route tag, split a residual without authorization, or create a certificate. Hence the packet is admissible. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Theorem H4.310 (First dispatched route/rank packet)**.** *After a normalization-ready return, the active HC/OP skeleton supports the next executed work packet*

$$\mathsf{N}_{\mathrm{ready}} \xrightarrow{\mathsf{W}^{\mathrm{rank}}_{\mathrm{route}}} \{\mathsf{R}_{\mathrm{ready}}, \mathsf{R}_{\mathrm{repair}}, \mathsf{R}_{\mathrm{frontier}}\}.$$

This execution is conservative: it either prepares a routed and ranked OP task for comparison testing, records one bounded route/rank repair obligation, or returns a smaller frontier source. It does not prove OP 1/4, OP 5(B), or any closure certificate.

Proof. By Proposition H4.309, the packet is admissible after a normalization-ready return. Definition H4.305 classifies the route/rank audit defect into route-clean, route-repairable, or frontier-bound cases, and Definition H4.308 records the corresponding return status. Lemma H4.307 proves the route-clean case. In the repair case the packet remains at the routing node with a strictly smaller local route/rank defect and no route-tag drift; in the frontier case it returns protected frontier material without promotion. In none of the three cases is the non-certificate marker changed into an OP closure certificate. □

Corollary H4.311 (working rule)**.** *Future work may proceed from the OP-normalization/type packet to the route/rank node only after* $\mathsf{N}_{\mathrm{ready}}$ *has been returned. The next admissible mathematical question is therefore whether the typed OP input is route-clean and comparison-ready, route/rank-repairable, or frontier-bound; it is still not whether OP 1/4 or OP 5(B) is closed.*

Proof. Immediate from Theorem H4.310. □

H4.18 First dispatched comparison/defect packet

Definition H4.312 (Dispatched comparison/defect packet)**.** *Let* $E_4 = \mathrm{COMPARE}^{1/4|5(B)}_{\mathrm{HC}}$ *be the fifth node of the tail-normalized HC/OP skeleton and let*

$$\mathsf{D}^{\mathrm{skel}}_{\mathrm{HCOP}}(E_4) = \mathsf{W}^{\Delta}_{\mathrm{cmp}}$$

be the dispatch assignment. A dispatched comparison/defect packet *is a finite packet*

$$\mathsf{P}^{\Delta}_{\mathrm{cmp}} = \big(\mathsf{R}_{\mathrm{ready}}, \mathcal{W}^{\mathrm{cmp}}_{1/4|5(B)}, \rho_{\mathrm{CL}}, \rho_{\Theta R}, D^{\mathrm{HC}}_{\mathrm{joint}}, D^{\mathrm{HC}}_{\mathrm{sep}}, G_4, W_4, S_4, \mathbf{d}^{\Delta}_{\mathrm{cmp}}\big),$$

where $\mathsf{R}_{\mathrm{ready}}$ *is the comparison-ready route/rank return of Definition H4.308,* $\mathcal{W}^{\mathrm{cmp}}_{1/4|5(B)}$ *is the inherited paired OP 1/4–OP 5(B) comparison work object,* ρ_{CL} *and* $\rho_{\Theta R}$ *are the routed continuum-limit and* Θ*–R residual coordinates,* $D^{\mathrm{HC}}_{\mathrm{joint}}$ *and* $D^{\mathrm{HC}}_{\mathrm{sep}}$ *are the HC-compatible joint and separated comparison readouts,* G_4 *is the comparison guard,* W_4 *is the witness pointer to the first explicit paired-comparison source block,* S_4 *is the non-certificate status marker, and* $\mathbf{d}^{\Delta}_{\mathrm{cmp}}$ *is a finite comparison/defect audit register. The packet is admissible only if it is launched from* $\mathsf{R}_{\mathrm{ready}}$*, keeps the inherited paired route visible, compares only HC-compatible joint and separated readouts, preserves both residual coordinates as residual data, and does not convert a comparison outcome into an OP closure certificate.*

Definition H4.313 (Comparison/defect register)**.** *For an admissible packet* $\mathsf{P}^{\Delta}_{\mathrm{cmp}}$*, the* comparison/defect register *is the finite family*

$$\mathbf{d}^{\Delta}_{\mathrm{cmp}}(j) = \big(d_{\mathrm{in}}(j), d_{\mathrm{pair}}(j), d_{\mathrm{read}}(j), d_{\Delta}(j), d_{\mathrm{front}}(j), d_{\mathrm{stat}}(j)\big), \qquad j \in \mathcal{A}_4,$$

where d_{in} *measures loss of the comparison-ready input status,* d_{pair} *measures unauthorized separation or binding of the paired residual branch,* d_{read} *measures non-HC-compatible readout drift,* d_{Δ} *measures instability of the comparison defect*

$$\Delta^{\mathrm{HC}}_{\mathrm{sep}} = D^{\mathrm{HC}}_{\mathrm{joint}} - D^{\mathrm{HC}}_{\mathrm{sep}},$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

d_{front} *measures violation of the comparison frontier guard, and* d_{stat} *measures accidental promotion of the non-certificate marker. The scalar comparison audit defect is*

$$\Delta^{\Delta}_{\mathrm{cmp}} = \max_{j\in\mathcal{A}_4}\big(d_{\mathrm{in}}(j) + d_{\mathrm{pair}}(j) + d_{\mathrm{read}}(j) + d_{\Delta}(j) + d_{\mathrm{front}}(j) + d_{\mathrm{stat}}(j)\big).$$

The register is comparison-clean *if* $\Delta^{\Delta}_{\mathrm{cmp}} = 0$, comparison-repairable *if one bounded local comparison repair strictly lowers the scalar defect while preserving the routed comparison input, and* frontier-bound *otherwise.*

Lemma H4.314 (Route/rank-ready return is the comparison precondition)**.** *The comparison/defect packet may be executed only from the route/rank-ready return status* $\mathsf{R}_{\mathrm{ready}}$*. If the return status is* $\mathsf{R}_{\mathrm{repair}}$ *or* $\mathsf{R}_{\mathrm{frontier}}$*, then the paired comparison node is not yet admissible.*

Proof. By Definition H4.312, the first entry of $\mathsf{P}^{\Delta}_{\mathrm{cmp}}$ is $\mathsf{R}_{\mathrm{ready}}$. A repair return still carries a bounded route/rank obligation, while a frontier return does not promote the source to the comparison node. Neither status supplies the routed and ranked comparison-ready input required to test $D^{\mathrm{HC}}_{\mathrm{joint}}$, $D^{\mathrm{HC}}_{\mathrm{sep}}$, and $\Delta^{\mathrm{HC}}_{\mathrm{sep}}$. Thus comparison execution is blocked until the packet has returned the ready status. □

Lemma H4.315 (Zero comparison defect preserves paired comparison data)**.** *If* $\Delta^{\Delta}_{\mathrm{cmp}} = 0$ *for an admissible dispatched comparison/defect packet, then the routed OP task passes to a guarded paired-comparison state with its paired residual branch, HC-compatible joint and separated readouts, comparison defect, frontier guard, witness pointer, and non-certificate status marker preserved.*

Proof. Vanishing scalar defect forces every component in Definition H4.313 to vanish on the audit window. Hence the comparison-ready input status is retained, the paired residual branch is neither bound nor split without authorization, the joint and separated readouts remain HC-compatible, the comparison defect is well-defined on the inherited route, the frontier guard remains active, and the non-certificate marker is not promoted. The output is therefore a guarded paired-comparison state ready for the next resolution node, not an OP closure certificate. □

Definition H4.316 (comparison/defect return)**.** *The execution of* $\mathsf{W}^{\Delta}_{\mathrm{cmp}}$ *on an admissible packet returns exactly one of the following statuses:*

$$\mathrm{Exec}\big(\mathsf{W}^{\Delta}_{\mathrm{cmp}}, \mathsf{P}^{\Delta}_{\mathrm{cmp}}\big) \in \{\mathsf{C}_{\mathrm{bind}}, \mathsf{C}_{\mathrm{split}}, \mathsf{C}_{\mathrm{frontier}}\}.$$

Here $\mathsf{C}_{\mathrm{bind}}$ *means that the guarded comparison supports the bound common-origin branch as the next final for resolution,* $\mathsf{C}_{\mathrm{split}}$ *means that the guarded comparison supports a ranked separated branch as the next final for resolution, and* $\mathsf{C}_{\mathrm{frontier}}$ *means that the comparison remains protected but must return to a smaller comparison frontier rather than move forward.*

Proposition H4.317 (The dispatched comparison/defect packet is admissible after route/rank-ready return)**.** *If the execution returns* $\mathsf{R}_{\mathrm{ready}}$*, then the dispatch menu admits the comparison/defect packet as the next executable work packet attached to the active HC/OP skeleton.*

Proof. The dispatch menu assigns E_4 only to $\mathsf{W}^{\Delta}_{\mathrm{cmp}}$. Lemma H4.314 supplies the required route/rank-ready input status. The remaining data of Definition H4.312 are inherited from the paired comparison layer, the QAM/MML/MMA interface, the guarded-reduction ledger, the skeleton, and the routed task state. The new defect register adds only a finite comparison audit surface. It does not introduce a new OP task, alter the paired route, bind or split a residual branch without the comparison guard, or create a certificate. Hence the packet is admissible. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Theorem H4.318 (First dispatched comparison/defect packet)**.** *After a route/rank-ready return, the active HC/OP skeleton supports the next executed work packet*

$$\mathsf{R}_{\mathrm{ready}} \xrightarrow{\mathsf{W}^{\Delta}_{\mathrm{cmp}}} \{\mathsf{C}_{\mathrm{bind}}, \mathsf{C}_{\mathrm{split}}, \mathsf{C}_{\mathrm{frontier}}\}.$$

This execution is conservative: it either prepares a guarded bind final, prepares a guarded split final, or returns a smaller comparison frontier. It does not prove OP 1/4, OP 5(B), or any closure certificate.

Proof. By Proposition H4.317, the packet is admissible after a route/rank-ready return. Definition H4.313 classifies the comparison audit defect into comparison-clean, comparison-repairable, or frontier-bound cases, and Definition H4.316 records the corresponding guarded comparison return. Lemma H4.315 proves the comparison-clean preservation claim. In the repair case the packet remains at the comparison node with a strictly smaller local comparison defect and no route drift; in the frontier case it returns protected comparison frontier material without promotion. In none of the three cases is the non-certificate marker changed into an OP closure certificate. □

Corollary H4.319 (working rule)**.** *Future work may proceed from the route/rank packet to the paired comparison node only after* $\mathsf{R}_{\mathrm{ready}}$ *has been returned. The next admissible mathematical question is therefore how a guarded comparison return* $\mathsf{C}_{\mathrm{bind}}$*,* $\mathsf{C}_{\mathrm{split}}$*, or* $\mathsf{C}_{\mathrm{frontier}}$ *should be resolved; it is still not whether OP 1/4 or OP 5(B) is closed.*

Proof. Immediate from Theorem H4.318. □

H4.19 First dispatched resolve/return packet

Definition H4.320 (Dispatched resolve/return packet)**.** *Let* $E_5 = \mathrm{RESOLVE}_{\mathrm{bind/split/frontier}}$ *be the sixth node of the tail-normalized HC/OP skeleton and let*

$$\mathsf{D}^{\mathrm{skel}}_{\mathrm{HCOP}}(E_5) = \mathsf{W}^{\mathrm{next}}_{\mathrm{resolve}}$$

be the dispatch assignment. A dispatched resolve/return packet *is a finite packet*

$$\mathsf{P}^{\mathrm{next}}_{\mathrm{resolve}} = \big(\mathsf{C}_{\eta}, \eta, \Pi_{\eta}, \rho_{\mathrm{CL}}, \rho_{\Theta R}, \Delta^{\mathrm{HC}}_{\mathrm{sep}}, G_5, W_5, S_5, \mathbf{d}^{\mathrm{next}}_{\mathrm{resolve}}\big), \qquad \eta \in \{\mathrm{bind}, \mathrm{split}, \mathrm{frontier}\},$$

where C_{η} *is the corresponding comparison return,* Π_{η} *is the next finite continuation package associated with that return,* ρ_{CL} *and* $\rho_{\Theta R}$ *are the inherited routed residual coordinates,* $\Delta^{\mathrm{HC}}_{\mathrm{sep}}$ *is the paired HC comparison defect,* G_5 *is the resolution guard,* W_5 *is the witness pointer to the first explicit comparison source,* S_5 *is the non-certificate status marker, and* $\mathbf{d}^{\mathrm{next}}_{\mathrm{resolve}}$ *is a finite resolution audit register. The packet is admissible only if it is launched from one of* $\mathsf{C}_{\mathrm{bind}}$*,* $\mathsf{C}_{\mathrm{split}}$*, or* $\mathsf{C}_{\mathrm{frontier}}$*, selects exactly the continuation package matching that return, keeps the inherited comparison data visible, and does not convert the selected continuation into an OP closure certificate.*

Definition H4.321 (Resolve/return defect register)**.** *For an admissible packet* $\mathsf{P}^{\mathrm{next}}_{\mathrm{resolve}}$*, the* resolve/return defect register *is the finite family*

$$\mathbf{d}^{\mathrm{next}}_{\mathrm{resolve}}(j) = \big(d_{\mathrm{in}}(j), d_{\mathrm{sel}}(j), d_{\mathrm{guard}}(j), d_{\mathrm{route}}(j), d_{\mathrm{wit}}(j), d_{\mathrm{stat}}(j)\big), \qquad j \in \mathcal{A}_5,$$

where d_{in} *measures loss of the guarded comparison return,* d_{sel} *measures mismatch between the comparison return and the selected continuation package,* d_{guard} *measures violation of the resolution guard,* d_{route} *measures loss of the routed OP 1/4–OP 5(B) comparison data,* d_{wit}

© 2026 Marc Brendecke. Preprint — not peer reviewed.

measures loss of the witness pointer, and d_{stat} *measures accidental promotion of the non-certificate marker. The scalar resolve/return defect is*

$$\Delta^{\mathrm{next}}_{\mathrm{resolve}} = \max_{j\in\mathcal{A}_5}\big(d_{\mathrm{in}}(j) + d_{\mathrm{sel}}(j) + d_{\mathrm{guard}}(j) + d_{\mathrm{route}}(j) + d_{\mathrm{wit}}(j) + d_{\mathrm{stat}}(j)\big).$$

The register is resolve-clean *if* $\Delta^{\mathrm{next}}_{\mathrm{resolve}} = 0$, resolve-repairable *if one bounded local resolution repair strictly lowers the scalar defect while preserving the guarded comparison return, and* frontier-bound *otherwise.*

Lemma H4.322 (Guarded comparison return is the resolve precondition)**.** *The resolve/return packet may be executed only from one of the guarded comparison returns* $\mathsf{C}_{\mathrm{bind}}$, $\mathsf{C}_{\mathrm{split}}$, *or* $\mathsf{C}_{\mathrm{frontier}}$. *If the comparison node has not returned one of these three guarded statuses, then the resolution node is not yet admissible.*

Proof. By Definition H4.320, the first entry of $\mathsf{P}^{\mathrm{next}}_{\mathrm{resolve}}$ is exactly one guarded comparison return C_{η}. The resolution node consumes a completed guarded comparison return, not a raw route/rank state and not an unresolved comparison audit surface. Without such a return there is no admissible bind, split, or frontier branch for the packet to select. □

Lemma H4.323 (Zero resolve/return defect preserves selected continuation data)**.** *If* $\Delta^{\mathrm{next}}_{\mathrm{resolve}} = 0$ *for an admissible dispatched resolve/return packet, then the selected continuation package preserves the corresponding comparison return, the routed residual data, the HC comparison defect, the resolution guard, the witness pointer, and the non-certificate status marker.*

Proof. Vanishing scalar defect forces all components of Definition H4.321 to vanish on the resolution audit window. Hence the input comparison return is retained, the selected continuation package matches that return, the guard is not bypassed, the routed residual data and comparison defect remain attached to the selected branch, the witness pointer is preserved, and the non-certificate marker is not promoted. Therefore the output is a guarded next work state, not an OP closure certificate. □

Definition H4.324 (resolve/return output)**.** *The execution of* $\mathsf{W}^{\mathrm{next}}_{\mathrm{resolve}}$ *on an admissible packet returns exactly one of the following statuses:*

$$\mathrm{Exec}\big(\mathsf{W}^{\mathrm{next}}_{\mathrm{resolve}}, \mathsf{P}^{\mathrm{next}}_{\mathrm{resolve}}\big) \in \{\mathsf{Q}_{\mathrm{bind}}, \mathsf{Q}_{\mathrm{split}}, \mathsf{Q}_{\mathrm{frontier}}\}.$$

Here $\mathsf{Q}_{\mathrm{bind}}$ *is a guarded common-origin continuation work state for the coupled OP 1/4–OP 5(B) branch,* $\mathsf{Q}_{\mathrm{split}}$ *is a guarded ranked single-branch continuation work state, and* $\mathsf{Q}_{\mathrm{frontier}}$ *is a protected smaller comparison-frontier descent state. None of these statuses is an OP closure certificate.*

Proposition H4.325 (The dispatched resolve/return packet is admissible after guarded comparison return)**.** *If the execution returns one of* $\mathsf{C}_{\mathrm{bind}}$, $\mathsf{C}_{\mathrm{split}}$, *or* $\mathsf{C}_{\mathrm{frontier}}$, *then the dispatch menu admits the resolve/return packet as the next executable work packet attached to the active HC/OP skeleton.*

Proof. The dispatch menu assigns E_5 only to $\mathsf{W}^{\mathrm{next}}_{\mathrm{resolve}}$. Lemma H4.322 supplies the required guarded comparison input status. The remaining data of Definition H4.320 are inherited from the paired comparison layer, the QAM/MML/MMA interface, the guarded-reduction ledger, the skeleton, and the comparison return. The new defect register adds only a finite resolution audit surface. It does not create a new certificate test, erase the residual route, suppress the witness pointer, or change the theorematic status of the selected continuation. Hence the packet is admissible. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Theorem H4.326 (First dispatched resolve/return packet)**.** *After a guarded comparison return, the active HC/OP skeleton supports the next executed work packet*

$$\{\mathsf{C}_{\mathrm{bind}}, \mathsf{C}_{\mathrm{split}}, \mathsf{C}_{\mathrm{frontier}}\} \xrightarrow{\mathsf{W}^{\mathrm{next}}_{\mathrm{resolve}}} \{\mathsf{Q}_{\mathrm{bind}}, \mathsf{Q}_{\mathrm{split}}, \mathsf{Q}_{\mathrm{frontier}}\}.$$

This execution is conservative: it selects a guarded common-origin continuation, a guarded ranked single-branch continuation, or a smaller comparison-frontier descent state. It does not prove OP 1/4, OP 5(B), or any closure certificate.

Proof. By Proposition H4.325, the packet is admissible after a guarded comparison return. Definition H4.321 classifies the resolution audit defect into resolve-clean, resolve-repairable, or frontier-bound cases, and Definition H4.324 records the corresponding guarded continuation output. Lemma H4.323 proves the resolve-clean preservation claim. In the repair case the packet remains at the resolution node with a strictly smaller local resolution defect and no route drift; in the frontier case it returns protected comparison-frontier material without promotion. In none of the three cases is the non-certificate marker changed into an OP closure certificate. □

Corollary H4.327 (working rule)**.** *Future work may proceed from the guarded comparison returns to a selected next continuation only through* $\mathsf{W}^{\mathrm{next}}_{\mathrm{resolve}}$*. The next admissible mathematical question is therefore whether* $\mathsf{Q}_{\mathrm{bind}}$*,* $\mathsf{Q}_{\mathrm{split}}$*, or* $\mathsf{Q}_{\mathrm{frontier}}$ *should seed the next finite work packet; it is still not whether OP 1/4 or OP 5(B) is closed.*

Proof. Immediate from Theorem H4.326. □

H4.20 First continuation-queue packet after resolve/return

Definition H4.328 (Continuation-queue packet after resolve/return)**.** *Let*

$$\mathsf{Q}_\eta \in \{\mathsf{Q}_{\mathrm{bind}}, \mathsf{Q}_{\mathrm{split}}, \mathsf{Q}_{\mathrm{frontier}}\}, \qquad \eta \in \{\mathrm{bind}, \mathrm{split}, \mathrm{frontier}\},$$

be a guarded resolve/return status. A continuation-queue packet after resolve/return *is a finite packet*

$$\mathsf{P}^{\mathrm{cont}}_{\mathrm{queue}} = \big(\mathsf{Q}_\eta, \mathcal{U}_\eta, \rho_{\mathrm{CL}}, \rho_{\Theta R}, \rho_A, \Delta^{\mathrm{HC}}_{\mathrm{sep}}, G_6, W_6, S_6, \mathbf{d}^{\mathrm{cont}}_{\mathrm{queue}}\big),$$

where $\mathcal{U}_\eta$ *is the finite guarded continuation queue selected by the return,* $(\rho_{\mathrm{CL}}, \rho_{\Theta R}, \rho_A)$ *is the inherited live residual triple,* $\Delta^{\mathrm{HC}}_{\mathrm{sep}}$ *is the inherited paired HC comparison defect,* G_6 *is the queue guard,* W_6 *is the witness pointer to the first explicit resolve/return source,* S_6 *is the non-certificate status marker, and* $\mathbf{d}^{\mathrm{cont}}_{\mathrm{queue}}$ *is a finite queue audit register. The packet is admissible only if the queue type matches the selected return, keeps the inherited residual and comparison data attached to the queue, and does not change a continuation queue into an OP closure certificate.*

Definition H4.329 (Bind, split, and frontier continuation queues)**.** *The continuation queue selected by Definition H4.328 is one of the following three finite guarded families:*

$$\begin{aligned}
\mathcal{U}_{\mathrm{bind}} &= \big(\mathrm{commonOriginCheck}, \mathrm{jointResidualRefine}, \mathrm{pairedWitnessReturn}\big),\\
\mathcal{U}_{\mathrm{split}} &= \big(\mathrm{branchRankCheck}, \mathrm{singleResidualSelect}, \mathrm{separatedWitnessReturn}\big),\\
\mathcal{U}_{\mathrm{frontier}} &= \big(\mathrm{frontierDescentCheck}, \mathrm{defectLocalize}, \mathrm{smallerFrontierReturn}\big).
\end{aligned}$$

The bind queue keeps OP 1/4 and OP 5(B) coupled under a common-origin guard; the split queue selects a ranked single-branch continuation under a separation guard; and the frontier queue performs only protected descent to a smaller comparison frontier. None of these queues is a theorematic OP closure state.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition H4.330 (Continuation-queue defect register)**.** *For an admissible packet* $\mathsf{P}^{\mathrm{cont}}_{\mathrm{queue}}$*, the* continuation-queue defect register *is the finite family*

$$\mathbf{d}^{\mathrm{cont}}_{\mathrm{queue}}(j) = \big(d_Q(j), d_{\mathrm{type}}(j), d_{\mathrm{res}}(j), d_\Delta(j), d_{\mathrm{guard}}(j), d_{\mathrm{wit}}(j), d_{\mathrm{stat}}(j)\big), \qquad j \in \mathcal{A}_6,$$

where d_Q *measures loss of the resolve/return status,* d_{type} *measures mismatch between the selected status and the selected queue type,* d_{res} *measures loss of the inherited live residual triple,* d_Δ *measures loss of the paired comparison defect,* d_{guard} *measures violation of the queue guard,* d_{wit} *measures loss of the witness pointer, and* d_{stat} *measures accidental certificate promotion. The scalar queue defect is*

$$\Delta^{\mathrm{cont}}_{\mathrm{queue}} = \max_{j\in\mathcal{A}_6}\big(d_Q(j) + d_{\mathrm{type}}(j) + d_{\mathrm{res}}(j) + d_\Delta(j) + d_{\mathrm{guard}}(j) + d_{\mathrm{wit}}(j) + d_{\mathrm{stat}}(j)\big).$$

The register is queue-clean *if* $\Delta^{\mathrm{cont}}_{\mathrm{queue}} = 0$*,* queue-repairable *if one bounded local queue repair strictly lowers the scalar defect while preserving the return, and* frontier-bound *otherwise.*

Lemma H4.331 (Resolve/return output is the queue precondition)**.** *The continuation-queue packet may be executed only from one of* $\mathsf{Q}_{\mathrm{bind}}$*,* $\mathsf{Q}_{\mathrm{split}}$*, or* $\mathsf{Q}_{\mathrm{frontier}}$*. If the resolve/return node has not returned one of these three guarded statuses, then the continuation-queue node is not yet admissible.*

Proof. By Definition H4.328, the first entry of the packet is exactly one guarded return Q_η. The queue node schedules follow-on work from a completed resolve/return state; it does not consume an unresolved comparison state, a raw route/rank state, or a merely local HC-capture source. Hence no continuation queue exists until the node has selected one of the three guarded return statuses. □

Lemma H4.332 (Zero continuation-queue defect preserves queue data)**.** *If* $\Delta^{\mathrm{cont}}_{\mathrm{queue}} = 0$ *for an admissible continuation-queue packet, then the selected queue preserves the return status, the matching queue type, the inherited live residual triple, the paired HC comparison defect, the queue guard, the witness pointer, and the non-certificate status marker.*

Proof. Vanishing scalar defect forces all components of Definition H4.330 to vanish on the queue audit window. Thus the return remains visible, the queue type matches the selected return, the live residual triple and paired comparison defect remain attached, the guard and witness pointer are preserved, and the non-certificate marker is not promoted. Therefore the output is a finite guarded work queue, not an OP closure certificate. □

Definition H4.333 (continuation-queue output)**.** *The execution of the continuation-queue packet returns exactly one of the following queued work states:*

$$\mathrm{Exec}\big(\mathsf{W}^{\mathrm{cont}}_{\mathrm{queue}}, \mathsf{P}^{\mathrm{cont}}_{\mathrm{queue}}\big) \in \{\mathsf{U}^{\mathrm{ready}}_{\mathrm{bind}}, \mathsf{U}^{\mathrm{ready}}_{\mathrm{split}}, \mathsf{U}^{\mathrm{ready}}_{\mathrm{frontier}}\}.$$

Here $\mathsf{U}^{\mathrm{ready}}_{\mathrm{bind}}$ *is a finite common-origin work queue,* $\mathsf{U}^{\mathrm{ready}}_{\mathrm{split}}$ *is a finite ranked single-branch work queue, and* $\mathsf{U}^{\mathrm{ready}}_{\mathrm{frontier}}$ *is a finite protected frontier-descent work queue. None of these statuses is an OP closure certificate.*

Proposition H4.334 (The continuation-queue packet is admissible after resolve/return)**.** *If the execution returns one of* $\mathsf{Q}_{\mathrm{bind}}$*,* $\mathsf{Q}_{\mathrm{split}}$*, or* $\mathsf{Q}_{\mathrm{frontier}}$*, then the active HC/OP skeleton admits* $\mathsf{W}^{\mathrm{cont}}_{\mathrm{queue}}$ *as the next executable work packet.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. Lemma H4.331 supplies the required guarded input. Definitions H4.328 and H4.329 attach the matching finite queue to that input. The remaining data are inherited from the paired comparison layer, the QAM/MML/MMA interface, the dispatch, the comparison return, and the resolve/return output. The queue defect register is finite and audits only preservation of input status, type, residual data, comparison defect, guard, witness, and non-certificate status. Hence the packet is admissible as follow-on work and not as an OP closure claim. □

Theorem H4.335 (First continuation-queue packet after resolve/return)**.** *After a guarded resolve/return output, the active HC/OP skeleton supports the next executed work packet*

$$\{\mathsf{Q}_{\mathrm{bind}}, \mathsf{Q}_{\mathrm{split}}, \mathsf{Q}_{\mathrm{frontier}}\} \xrightarrow{\mathsf{W}^{\mathrm{cont}}_{\mathrm{queue}}} \{\mathsf{U}^{\mathrm{ready}}_{\mathrm{bind}}, \mathsf{U}^{\mathrm{ready}}_{\mathrm{split}}, \mathsf{U}^{\mathrm{ready}}_{\mathrm{frontier}}\}.$$

This execution is conservative: it turns the return into a finite guarded continuation queue and does not prove OP 1/4, OP 5(B), or any closure certificate.

Proof. By Proposition H4.334, the packet is admissible after a guarded return. Definition H4.330 classifies the queue audit defect into queue-clean, queue-repairable, or frontier-bound cases, and Definition H4.333 records the corresponding queued work output. Lemma H4.332 proves the queue-clean preservation claim. In the repair case the packet remains at the queue node with a strictly smaller local queue defect and no route drift; in the frontier case it returns protected smaller frontier material. In none of the three cases is the non-certificate marker changed into an OP closure certificate. □

Corollary H4.336 (working rule)**.** *Future work may proceed from the guarded resolve/return status only by selecting the matching finite continuation queue. The next admissible mathematical question is therefore which queued work family — common-origin bind refinement, ranked split refinement, or protected frontier descent — should be executed first; it is still not whether OP 1/4 or OP 5(B) is closed.*

Proof. Immediate from Theorem H4.335. □

H4.21 First queue-head selection packet after continuation queue

Definition H4.337 (Queue-head selection packet after continuation queue)**.** *Let*

$$\mathsf{U}^{\mathrm{ready}}_{\eta} \in \{\mathsf{U}^{\mathrm{ready}}_{\mathrm{bind}}, \mathsf{U}^{\mathrm{ready}}_{\mathrm{split}}, \mathsf{U}^{\mathrm{ready}}_{\mathrm{frontier}}\}, \qquad \eta \in \{\mathrm{bind}, \mathrm{split}, \mathrm{frontier}\},$$

be a guarded continuation-queue output. A queue-head selection packet after continuation queue *is a finite packet*

$$\mathsf{P}^{\mathrm{select}}_{\mathrm{head}} = \big(\mathsf{U}^{\mathrm{ready}}_{\eta}, h_{\eta}, \mathcal{U}^{\mathrm{tail}}_{\eta}, \rho_{\mathrm{CL}}, \rho_{\Theta R}, \rho_{A}, \Delta^{\mathrm{HC}}_{\mathrm{sep}}, G_7, W_7, S_7, \mathbf{d}^{\mathrm{select}}_{\mathrm{head}}\big),$$

where h_η is the first guarded queue head selected from the matching queue, $\mathcal{U}^{\mathrm{tail}}_{\eta}$ is the remaining finite queue tail, $(\rho_{\mathrm{CL}}, \rho_{\Theta R}, \rho_A)$ is the inherited live residual triple, $\Delta^{\mathrm{HC}}_{\mathrm{sep}}$ is the inherited paired HC comparison defect, G_7 is the head-selection guard, W_7 is the witness pointer to the first explicit continuation-queue source, S_7 is the non-certificate status marker, and $\mathbf{d}^{\mathrm{select}}_{\mathrm{head}}$ is a finite head-selection audit register. The packet is admissible only if the selected head belongs to the matching queue, the tail remains finite and guarded, the inherited residual and comparison data remain attached, and no selected head is promoted to an OP closure certificate.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition H4.338 (Continuation-queue heads and finite tails)**.** *The matching queue head and tail in Definition H4.337 are fixed by the queue type:*

$$\begin{aligned} h_{\mathrm{bind}} &= \mathrm{commonOriginCheck}, & \mathcal{U}^{\mathrm{tail}}_{\mathrm{bind}} &= (\mathrm{jointResidualRefine}, \mathrm{pairedWitnessReturn}),\\ h_{\mathrm{split}} &= \mathrm{branchRankCheck}, & \mathcal{U}^{\mathrm{tail}}_{\mathrm{split}} &= (\mathrm{singleResidualSelect}, \mathrm{separatedWitnessReturn}),\\ h_{\mathrm{frontier}} &= \mathrm{frontierDescentCheck}, & \mathcal{U}^{\mathrm{tail}}_{\mathrm{frontier}} &= (\mathrm{defectLocalize}, \mathrm{smallerFrontierReturn}). \end{aligned}$$

Thus the bind queue first tests whether the common-origin reading is still admissible, the split queue first tests whether a ranked branch can honestly be selected, and the frontier queue first tests whether a protected descent is available. The selected head is a next work packet, not a theorematic OP closure state.

Definition H4.339 (Queue-head selection defect register)**.** *For an admissible packet* $\mathsf{P}^{\mathrm{select}}_{\mathrm{head}}$*, the* queue-head selection defect register *is the finite family*

$$\mathbf{d}^{\mathrm{select}}_{\mathrm{head}}(j) = \big(d_U(j), d_h(j), d_{\mathrm{tail}}(j), d_{\mathrm{res}}(j), d_\Delta(j), d_{\mathrm{guard}}(j), d_{\mathrm{wit}}(j), d_{\mathrm{stat}}(j)\big), \qquad j \in \mathcal{A}_7,$$

where d_U *measures loss of the queued work state,* d_h *measures mismatch between the queue type and selected head,* d_{tail} *measures loss of the finite guarded queue tail,* d_{res} *measures loss of the live residual triple,* d_Δ *measures loss of the paired HC comparison defect,* d_{guard} *measures guard drift,* d_{wit} *measures witness-pointer drift, and* d_{stat} *measures accidental certificate promotion. The scalar head-selection defect is*

$$\Delta^{\mathrm{select}}_{\mathrm{head}} = \max_{j \in \mathcal{A}_7} \big(d_U(j) + d_h(j) + d_{\mathrm{tail}}(j) + d_{\mathrm{res}}(j) + d_\Delta(j) + d_{\mathrm{guard}}(j) + d_{\mathrm{wit}}(j) + d_{\mathrm{stat}}(j)\big).$$

The register is head-clean *if* $\Delta^{\mathrm{select}}_{\mathrm{head}} = 0$*,* head-repairable *if one bounded local head-selection repair strictly lowers the scalar defect while preserving the queue, and* frontier-bound *otherwise.*

Lemma H4.340 (Continuation-queue output is the head-selection precondition)**.** *The queue-head selection packet may be executed only from one of* $\mathsf{U}^{\mathrm{ready}}_{\mathrm{bind}}$*,* $\mathsf{U}^{\mathrm{ready}}_{\mathrm{split}}$*, or* $\mathsf{U}^{\mathrm{ready}}_{\mathrm{frontier}}$*. If the continuation-queue node has not returned one of these three guarded ready queues, then the head-selection node is not yet admissible.*

Proof. By Definition H4.337, the first entry of the packet is exactly one queued work state $\mathsf{U}^{\mathrm{ready}}_{\eta}$. Head selection consumes a completed finite queue and selects its first guarded work item; it does not consume a raw return, an unresolved comparison return, or a merely local HC-capture source. Hence no head-selection packet exists until the node has returned a matching ready queue. □

Lemma H4.341 (Zero queue-head selection defect preserves selected-head data)**.** *If* $\Delta^{\mathrm{select}}_{\mathrm{head}} = 0$ *for an admissible queue-head selection packet, then the selected head preserves the queued work state, the matching head type, the finite guarded tail, the inherited live residual triple, the paired HC comparison defect, the head-selection guard, the witness pointer, and the non-certificate status marker.*

Proof. Vanishing scalar defect forces all components of Definition H4.339 to vanish on the head-selection audit window. Thus the ready queue remains visible, the selected head matches that queue, the finite tail is retained for later execution, the live residual triple and paired comparison defect remain attached, the guard and witness pointer are preserved, and the non-certificate marker is not promoted. Therefore the output is a guarded first work head together with a finite continuation tail, not an OP closure certificate. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition H4.342 (queue-head selection output)**.** *The execution of the queue-head selection packet returns exactly one of the following selected-head work states:*

$$\mathrm{Exec}\big(\mathsf{W}^{\mathrm{select}}_{\mathrm{head}}, \mathsf{P}^{\mathrm{select}}_{\mathrm{head}}\big) \in \{\mathsf{X}^{\mathrm{head}}_{\mathrm{bind}}, \mathsf{X}^{\mathrm{head}}_{\mathrm{split}}, \mathsf{X}^{\mathrm{head}}_{\mathrm{frontier}}\}.$$

Here $\mathsf{X}^{\mathrm{head}}_{\mathrm{bind}}$ *is the guarded common-origin-check head with its finite bind tail,* $\mathsf{X}^{\mathrm{head}}_{\mathrm{split}}$ *is the guarded branch-rank-check head with its finite split tail, and* $\mathsf{X}^{\mathrm{head}}_{\mathrm{frontier}}$ *is the guarded frontier-descent-check head with its finite frontier tail. None of these statuses is an OP closure certificate.*

Proposition H4.343 (The queue-head selection packet is admissible after queue return)**.** *If the execution returns one of* $\mathsf{U}^{\mathrm{ready}}_{\mathrm{bind}}$, $\mathsf{U}^{\mathrm{ready}}_{\mathrm{split}}$, *or* $\mathsf{U}^{\mathrm{ready}}_{\mathrm{frontier}}$, *then the active HC/OP skeleton admits* $\mathsf{W}^{\mathrm{select}}_{\mathrm{head}}$ *as the next executable work packet.*

Proof. Lemma H4.340 supplies the required guarded input. Definition H4.338 supplies the matching head and finite tail for that input. The remaining data are inherited from the paired comparison layer, the QAM/MML/MMA interface, the dispatch, the comparison return, the resolve/return output, and the continuation queue. The head-selection defect register is finite and audits only preservation of input queue, head type, tail, residual data, comparison defect, guard, witness, and non-certificate status. Hence the packet is admissible as follow-on work and not as an OP closure claim. □

Theorem H4.344 (First queue-head selection packet after continuation queue)**.** *After a guarded continuation-queue output, the active HC/OP skeleton supports the next executed work packet*

$$\{\mathsf{U}^{\mathrm{ready}}_{\mathrm{bind}}, \mathsf{U}^{\mathrm{ready}}_{\mathrm{split}}, \mathsf{U}^{\mathrm{ready}}_{\mathrm{frontier}}\} \xrightarrow{\mathsf{W}^{\mathrm{select}}_{\mathrm{head}}} \{\mathsf{X}^{\mathrm{head}}_{\mathrm{bind}}, \mathsf{X}^{\mathrm{head}}_{\mathrm{split}}, \mathsf{X}^{\mathrm{head}}_{\mathrm{frontier}}\}.$$

This execution is conservative: it selects the first guarded head of the matching finite continuation queue and keeps the remaining finite tail available for later work. It does not prove OP 1/4, OP 5(B), or any closure certificate.

Proof. By Proposition H4.343, the packet is admissible after a guarded ready queue. Definition H4.339 classifies the head-selection audit defect into head-clean, head-repairable, or frontier-bound cases, and Definition H4.342 records the corresponding selected-head work output. Lemma H4.341 proves the head-clean preservation claim. In the repair case the packet remains at the head-selection node with a strictly smaller local head-selection defect and no route drift; in the frontier case it returns protected smaller frontier material. In none of the three cases is the non-certificate marker changed into an OP closure certificate. □

Corollary H4.345 (working rule)**.** *Future work may proceed from a guarded ready queue only by selecting the matching first queue head while retaining the finite guarded tail. The next admissible mathematical question is therefore whether to execute* commonOriginCheck, branchRankCheck, *or* frontierDescentCheck *under its inherited guard; it is still not whether OP 1/4 or OP 5(B) is closed.*

Proof. Immediate from Theorem H4.344. □

H4.22 First selected queue-head execution packet

Definition H4.346 (Selected queue-head execution packet)**.** *Let*

$$\mathsf{X}^{\mathrm{head}}_{\eta} \in \{\mathsf{X}^{\mathrm{head}}_{\mathrm{bind}}, \mathsf{X}^{\mathrm{head}}_{\mathrm{split}}, \mathsf{X}^{\mathrm{head}}_{\mathrm{frontier}}\}, \qquad \eta \in \{\mathrm{bind}, \mathrm{split}, \mathrm{frontier}\},$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

be a guarded selected queue-head output. A selected queue-head execution packet *is a finite packet*

$$\mathsf{P}^{\mathrm{exec}}_{\mathrm{head}} = \big(\mathsf{X}^{\mathrm{head}}_{\eta}, h_\eta, \mathcal{U}^{\mathrm{tail}}_{\eta}, \rho_{\mathrm{CL}}, \rho_{\Theta R}, \rho_A, \Delta^{\mathrm{HC}}_{\mathrm{sep}}, \Delta^{\Delta}_{\mathrm{cmp}}, G_8, W_8, S_8, \mathbf{d}^{\mathrm{exec}}_{\mathrm{head}}\big),$$

where h_η is the selected queue head, $\mathcal{U}^{\mathrm{tail}}_{\eta}$ is the remaining finite guarded tail, $(\rho_{\mathrm{CL}}, \rho_{\Theta R}, \rho_A)$ is the inherited live residual triple, $\Delta^{\mathrm{HC}}_{\mathrm{sep}}$ is the inherited paired HC separation defect, $\Delta^{\Delta}_{\mathrm{cmp}}$ is the comparison-defect witness, G_8 is the execution guard, W_8 is the witness pointer to the selected-head source, S_8 is the non-certificate status marker, and $\mathbf{d}^{\mathrm{exec}}_{\mathrm{head}}$ is a finite head-execution audit register. The packet is admissible only if it executes the selected head belonging to the matching output, preserves the finite tail for later continuation, keeps the paired OP 1/4–OP 5(B) data attached, and does not promote the executed head to an OP closure certificate.

Definition H4.347 (Executed queue-head tests)**.** *The selected queue-head execution in Definition H4.346 is fixed by the head type:*

$$\begin{aligned}
\mathsf{X}^{\mathrm{head}}_{\mathrm{bind}} &\rightsquigarrow \mathrm{ExecHead(commonOriginCheck)},\\
\mathsf{X}^{\mathrm{head}}_{\mathrm{split}} &\rightsquigarrow \mathrm{ExecHead(branchRankCheck)},\\
\mathsf{X}^{\mathrm{head}}_{\mathrm{frontier}} &\rightsquigarrow \mathrm{ExecHead(frontierDescentCheck)}.
\end{aligned}$$

The bind-head execution checks whether the common-origin reading of $\rho_{\mathrm{CL}} \oplus \rho_{\Theta R}$ remains admissible under the inherited HC comparison defect. The split-head execution checks whether a ranked single-branch continuation can be selected without erasing the paired comparison witness. The frontier-head execution checks whether a protected descent to a smaller comparison frontier is available. Each head test is a work test, not a theorematic closure test.

Definition H4.348 (Selected queue-head execution defect register)**.** *For an admissible packet $\mathsf{P}^{\mathrm{exec}}_{\mathrm{head}}$, the* selected queue-head execution defect register *is the finite family*

$$\mathbf{d}^{\mathrm{exec}}_{\mathrm{head}}(j) = \big(d_X(j), d_h(j), d_{\mathrm{tail}}(j), d_{\mathrm{res}}(j), d_\Delta(j), d_{\mathrm{cmp}}(j), d_{\mathrm{guard}}(j), d_{\mathrm{wit}}(j), d_{\mathrm{stat}}(j)\big), \qquad j \in \mathcal{A}_8,$$

where d_X measures loss of the selected-head state, d_h measures mismatch between the selected head and the executed head test, d_{tail} measures loss of the finite queue tail, d_{res} measures loss of the live residual triple, d_Δ measures loss of the inherited HC separation defect, d_{cmp} measures loss of the comparison-defect witness, d_{guard} measures execution-guard drift, d_{wit} measures witness-pointer drift, and d_{stat} measures accidental certificate promotion. The scalar selected-head execution defect is

$$\Delta^{\mathrm{exec}}_{\mathrm{head}} = \max_{j\in\mathcal{A}_8}\big(d_X(j)+d_h(j)+d_{\mathrm{tail}}(j)+d_{\mathrm{res}}(j)+d_\Delta(j)+d_{\mathrm{cmp}}(j)+d_{\mathrm{guard}}(j)+d_{\mathrm{wit}}(j)+d_{\mathrm{stat}}(j)\big).$$

The register is head-exec-clean *if $\Delta^{\mathrm{exec}}_{\mathrm{head}} = 0$,* head-exec-repairable *if one bounded local execution repair strictly lowers the scalar defect while preserving the selected head and finite tail, and* frontier-bound *otherwise.*

Lemma H4.349 (Selected head is the head-execution precondition)**.** *The selected queue-head execution packet may be executed only from one of $\mathsf{X}^{\mathrm{head}}_{\mathrm{bind}}$, $\mathsf{X}^{\mathrm{head}}_{\mathrm{split}}$, or $\mathsf{X}^{\mathrm{head}}_{\mathrm{frontier}}$. If the head-selection node has not returned one of these three guarded head states, then the head-execution node is not yet admissible.*

Proof. By Definition H4.346, the first entry of the packet is exactly one selected head state. The execution node consumes a selected head, not a raw queue, a raw resolve return, or an earlier comparison final. Hence the head-execution packet is undefined until the node has selected the matching guarded queue head and retained the finite tail. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Lemma H4.350 (Zero selected-head execution defect preserves head-execution data)**.** *If* $\Delta^{\mathrm{exec}}_{\mathrm{head}} = 0$ *for an admissible selected queue-head execution packet, then the executed head preserves the selected-head state, the matching executed test, the finite guarded tail, the inherited live residual triple, the paired HC separation defect, the comparison-defect witness, the execution guard, the witness pointer, and the non-certificate status marker.*

Proof. Vanishing scalar defect forces all components of Definition H4.348 to vanish on the execution audit window. Thus the selected head is retained, the executed test matches the selected head type, the finite tail remains available for follow-on work, the residual and comparison data remain attached, the guard and witness pointer do not drift, and the non-certificate marker is not promoted. Therefore the output is an executed first queue head with a finite continuation tail, not an OP closure certificate. □

Definition H4.351 (selected queue-head execution output)**.** *The execution of the selected queue-head packet returns exactly one of the following executed-head work states:*

$$\mathrm{Exec}\big(\mathsf{W}^{\mathrm{exec}}_{\mathrm{head}}, \mathsf{P}^{\mathrm{exec}}_{\mathrm{head}}\big) \in \{\mathsf{Y}^{\mathrm{exec}}_{\mathrm{bind}}, \mathsf{Y}^{\mathrm{exec}}_{\mathrm{split}}, \mathsf{Y}^{\mathrm{exec}}_{\mathrm{frontier}}\}.$$

Here $\mathsf{Y}^{\mathrm{exec}}_{\mathrm{bind}}$ *is the executed common-origin-check surface with its finite bind tail,* $\mathsf{Y}^{\mathrm{exec}}_{\mathrm{split}}$ *is the executed branch-rank-check surface with its finite split tail, and* $\mathsf{Y}^{\mathrm{exec}}_{\mathrm{frontier}}$ *is the executed frontier-descent-check surface with its finite frontier tail. None of these statuses is an OP closure certificate.*

Proposition H4.352 (Selected queue-head execution is admissible after head selection)**.** *If the execution returns one of* $\mathsf{X}^{\mathrm{head}}_{\mathrm{bind}}$, $\mathsf{X}^{\mathrm{head}}_{\mathrm{split}}$, *or* $\mathsf{X}^{\mathrm{head}}_{\mathrm{frontier}}$, *then the active HC/OP skeleton admits* $\mathsf{W}^{\mathrm{exec}}_{\mathrm{head}}$ *as the next executable work packet.*

Proof. Lemma H4.349 supplies the required guarded input. Definition H4.347 supplies the matching head test. The residual data, paired HC separation defect, comparison-defect witness, guard, witness pointer, finite tail, and non-certificate status are inherited from the comparison packet, the resolve return, the continuation queue, and the selected head. The execution defect register is finite and audits only preservation of those inherited data. Hence the packet is admissible as follow-on work and not as an OP closure claim. □

Theorem H4.353 (First selected queue-head execution packet)**.** *After a guarded queue-head selection output, the active HC/OP skeleton supports the next executed work packet*

$$\{\mathsf{X}^{\mathrm{head}}_{\mathrm{bind}}, \mathsf{X}^{\mathrm{head}}_{\mathrm{split}}, \mathsf{X}^{\mathrm{head}}_{\mathrm{frontier}}\} \xrightarrow{\mathsf{W}^{\mathrm{exec}}_{\mathrm{head}}} \{\mathsf{Y}^{\mathrm{exec}}_{\mathrm{bind}}, \mathsf{Y}^{\mathrm{exec}}_{\mathrm{split}}, \mathsf{Y}^{\mathrm{exec}}_{\mathrm{frontier}}\}.$$

This execution is conservative: it runs the first guarded head of the matching continuation queue, keeps the finite tail available for later work, and does not prove OP 1/4, OP 5(B), or any closure certificate.

Proof. By Proposition H4.352, the packet is admissible after a guarded selected head. Definition H4.348 classifies the head-execution audit defect into head-exec-clean, head-exec-repairable, or frontier-bound cases, and Definition H4.351 records the corresponding executed-head work output. Lemma H4.350 proves the clean preservation claim. In the repair case the packet remains at the selected-head execution node with a strictly smaller local execution defect and no route drift; in the frontier case it returns protected smaller frontier material. In none of the three cases is the non-certificate marker changed into an OP closure certificate. □

Corollary H4.354 (working rule)**.** *Future work may proceed from an selected queue head only by executing the matching guarded head test while retaining the finite queue tail. The next admissible mathematical question is therefore whether the executed bind, split, or frontier head can legally enter its corresponding tail task — joint residual refinement, single residual selection, or defect localization — still under the inherited no-false-certificate rule.*

Proof. Immediate from Theorem H4.353. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

H4.23 First executed-head tail-entry packet

Definition H4.355 (Executed-head tail-entry packet)**.** *Let*

$$\mathsf{Y}_{\eta}^{\mathrm{exec}} \in \{\mathsf{Y}_{\mathrm{bind}}^{\mathrm{exec}}, \mathsf{Y}_{\mathrm{split}}^{\mathrm{exec}}, \mathsf{Y}_{\mathrm{frontier}}^{\mathrm{exec}}\}, \qquad \eta \in \{\mathrm{bind}, \mathrm{split}, \mathrm{frontier}\},$$

be a guarded executed-head output of Definition H4.351. An executed-head tail-entry packet *is a finite packet*

$$\mathsf{P}_{\mathrm{tail}}^{\mathrm{entry}} = \big(\mathsf{Y}_{\eta}^{\mathrm{exec}}, \mathcal{U}_{\eta}^{\mathrm{tail}}, \rho_{\mathrm{CL}}, \rho_{\Theta R}, \rho_A, \Delta_{\mathrm{sep}}^{\mathrm{HC}}, \Delta_{\mathrm{cmp}}^{\Delta}, \Delta_{\mathrm{head}}^{\mathrm{exec}}, G_9, W_9, S_9, \mathbf{d}_{\mathrm{tail}}^{\mathrm{entry}}\big),$$

where $\mathcal{U}_{\eta}^{\mathrm{tail}}$ *is the finite guarded tail retained by the selected-head execution,* $(\rho_{\mathrm{CL}}, \rho_{\Theta R}, \rho_A)$ *is the inherited live residual triple,* $\Delta_{\mathrm{sep}}^{\mathrm{HC}}$ *and* $\Delta_{\mathrm{cmp}}^{\Delta}$ *are the inherited comparison witnesses,* $\Delta_{\mathrm{head}}^{\mathrm{exec}}$ *is the selected-head execution defect,* G_9 *is the tail-entry guard,* W_9 *is the witness pointer to the executed-head source,* S_9 *is the non-certificate status marker, and* $\mathbf{d}_{\mathrm{tail}}^{\mathrm{entry}}$ *is a finite tail-entry audit register. The packet is admissible only if it enters the tail matching the executed head, preserves the residual and comparison data, preserves the finite tail, and does not promote the tail entry to an OP closure certificate.*

Definition H4.356 (Executed-head tail-entry defect)**.** *For an admissible packet* $\mathsf{P}_{\mathrm{tail}}^{\mathrm{entry}}$*, the* tail-entry defect register *is the finite family*

$$\mathbf{d}_{\mathrm{tail}}^{\mathrm{entry}}(j) = \big(d_Y(j), d_{\mathrm{tail}}(j), d_{\mathrm{res}}(j), d_{\Delta}(j), d_{\mathrm{head}}(j), d_{\mathrm{guard}}(j), d_{\mathrm{wit}}(j), d_{\mathrm{stat}}(j)\big), \qquad j \in \mathcal{A}_{\mathrm{tail}},$$

where the components measure loss of the executed-head state, the finite tail, the residual triple, the comparison witnesses, the head-execution defect, the tail-entry guard, the witness pointer, and the non-certificate marker. The scalar tail-entry defect is

$$\Delta_{\mathrm{tail}}^{\mathrm{entry}} = \max_{j \in \mathcal{A}_{\mathrm{tail}}} \big(d_Y(j) + d_{\mathrm{tail}}(j) + d_{\mathrm{res}}(j) + d_{\Delta}(j) + d_{\mathrm{head}}(j) + d_{\mathrm{guard}}(j) + d_{\mathrm{wit}}(j) + d_{\mathrm{stat}}(j)\big).$$

The register is tail-entry-clean *if* $\Delta_{\mathrm{tail}}^{\mathrm{entry}} = 0$*,* tail-entry-repairable *if one bounded local tail-entry repair strictly lowers the scalar defect while preserving the executed head and finite tail, and* frontier-bound *otherwise.*

Lemma H4.357 (Executed head is the tail-entry precondition)**.** *The tail-entry packet may be launched only from one of* $\mathsf{Y}_{\mathrm{bind}}^{\mathrm{exec}}$*,* $\mathsf{Y}_{\mathrm{split}}^{\mathrm{exec}}$*, or* $\mathsf{Y}_{\mathrm{frontier}}^{\mathrm{exec}}$*. If the selected-head execution has not returned one of these three guarded executed-head states, then tail entry is not yet admissible.*

Proof. Definition H4.355 takes the guarded executed-head state as the first datum of the packet. A selected but unexecuted head, a ready queue, or a resolve return does not yet determine which finite tail is allowed to open. Hence the tail-entry node is undefined until the selected-head execution has returned the matching executed-head state and retained its finite guarded tail. □

Lemma H4.358 (Zero tail-entry defect preserves tail-entry data)**.** *If* $\Delta_{\mathrm{tail}}^{\mathrm{entry}} = 0$ *for an admissible executed-head tail-entry packet, then the packet preserves the executed-head state, the matching finite tail, the live residual triple, the inherited comparison witnesses, the selected-head execution defect, the tail-entry guard, the witness pointer, and the non-certificate status marker.*

Proof. Vanishing scalar defect forces every component of Definition H4.356 to vanish on the tail-entry audit window. Therefore no executed-head data, residual data, comparison witness, guard, witness pointer, or status marker is lost. The output is a guarded tail-entry surface, not an OP closure certificate. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition H4.359 (Executed-head tail-entry output)**.** *The execution of the tail-entry packet returns exactly one of the following guarded tail-entry surfaces:*

$$\mathrm{Exec}\big(\mathsf{W}^{\mathrm{entry}}_{\mathrm{tail}}, \mathsf{P}^{\mathrm{entry}}_{\mathrm{tail}}\big) \in \{\mathsf{Z}^{\mathrm{tail}}_{\mathrm{bind}}, \mathsf{Z}^{\mathrm{tail}}_{\mathrm{split}}, \mathsf{Z}^{\mathrm{tail}}_{\mathrm{frontier}}\}.$$

Here $\mathsf{Z}^{\mathrm{tail}}_{\mathrm{bind}}$ *enters the guarded joint-residual refinement tail,* $\mathsf{Z}^{\mathrm{tail}}_{\mathrm{split}}$ *enters the guarded single-branch selection tail, and* $\mathsf{Z}^{\mathrm{tail}}_{\mathrm{frontier}}$ *enters the guarded defect-localization tail. None of these statuses is an OP closure certificate.*

Proposition H4.360 (Tail entry is admissible after selected-head execution)**.** *If the selected-head execution returns one of* $\mathsf{Y}^{\mathrm{exec}}_{\mathrm{bind}}$, $\mathsf{Y}^{\mathrm{exec}}_{\mathrm{split}}$, *or* $\mathsf{Y}^{\mathrm{exec}}_{\mathrm{frontier}}$, *then the active HC/OP skeleton admits* $\mathsf{W}^{\mathrm{entry}}_{\mathrm{tail}}$ *as the next executable work packet.*

Proof. Lemma H4.357 supplies the required executed-head input. The matching finite tail, residual triple, comparison witnesses, head-execution defect, guard, witness pointer, and non-certificate marker are inherited from the selected-head execution packet. The new defect register audits only preservation of those inherited data. Thus the packet is admissible as follow-on work and not as an OP closure claim. □

Theorem H4.361 (First executed-head tail-entry packet)**.** *After a guarded selected-head execution output, the active HC/OP skeleton supports the next tail-entry work packet*

$$\{\mathsf{Y}^{\mathrm{exec}}_{\mathrm{bind}}, \mathsf{Y}^{\mathrm{exec}}_{\mathrm{split}}, \mathsf{Y}^{\mathrm{exec}}_{\mathrm{frontier}}\} \xrightarrow{\mathsf{W}^{\mathrm{entry}}_{\mathrm{tail}}} \{\mathsf{Z}^{\mathrm{tail}}_{\mathrm{bind}}, \mathsf{Z}^{\mathrm{tail}}_{\mathrm{split}}, \mathsf{Z}^{\mathrm{tail}}_{\mathrm{frontier}}\}.$$

This execution is conservative: it opens the tail belonging to the executed head, retains all residual and comparison data, and does not prove OP 1/4, OP 5(B), or any closure certificate.

Proof. By Proposition H4.360, the packet is admissible after a guarded executed-head state. Definition H4.356 classifies the tail-entry defect into clean, repairable, or frontier-bound cases, while Definition H4.359 records the corresponding guarded tail-entry surface. Lemma H4.358 proves the clean preservation claim. In the repair case the packet remains at the tail-entry node with a strictly smaller local tail-entry defect; in the frontier case it returns protected smaller frontier material. In none of these cases is the non-certificate marker changed into an OP closure certificate. □

Corollary H4.362 (Tail-entry working rule)**.** *Future work may proceed from an executed queue head only by entering the matching finite guarded tail. The next admissible mathematical question is therefore whether the bind, split, or frontier tail can be refined under its inherited guard; it is still not whether OP 1/4 or OP 5(B) is closed.*

Proof. Immediate from Theorem H4.361. □

H4.24 First audit-freeze ledger

Definition H4.363 (Audit-freeze ledger)**.** *The* audit-freeze ledger *for the current HC/OP line is the finite record*

$$F^{\mathrm{audit}}_{\mathrm{HCOP}} = \big(\mathcal{B}_{\mathrm{public}}, \mathcal{D}_{\mathrm{doi}}, \mathcal{S}_{\mathrm{spine}}, \mathcal{Z}_{\mathrm{tail}}, \mathcal{O}_{\mathrm{pres}}, \mathcal{C}_{\mathrm{ref}}, \mathcal{B}_{\mathrm{finalbib}}, \mathcal{H}_{\mathrm{witness}}, S_{\mathrm{noCert}}\big).$$

Here $\mathcal{B}_{\mathrm{public}}$ *records the public baseline and concept-DOI status,* $\mathcal{D}_{\mathrm{doi}}$ *records the version-specific Zenodo DOI assigned in the public release metadata,* $\mathcal{S}_{\mathrm{spine}}$ *records the local HC/OP spine up to the tail-entry packet,* $\mathcal{Z}_{\mathrm{tail}}$ *records the guarded tail-entry output,* $\mathcal{O}_{\mathrm{pres}}$ *records preservation obligations,* $\mathcal{C}_{\mathrm{ref}}$ *records compilation/reference checks as audit data,* $\mathcal{B}_{\mathrm{finalbib}}$ *records terminal bibliography status,* $\mathcal{H}_{\mathrm{witness}}$ *records preservation of the historical witness, and* S_{noCert} *records that no OP closure certificate has been manufactured.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition H4.364 (Audit-freeze defect)**.** *For a ledger* $F^{\mathrm{audit}}_{\mathrm{HCOP}}$*, the* audit-freeze defect *is*

$$\Delta^{\mathrm{freeze}}_{\mathrm{audit}} = \max\{d_{\mathrm{base}}, d_{\mathrm{doi}}, d_{\mathrm{spine}}, d_{\mathrm{tail}}, d_{\mathrm{pres}}, d_{\mathrm{ref}}, d_{\mathrm{bib}}, d_{\mathrm{hist}}, d_{\mathrm{cert}}\}.$$

The entries measure drift of the public baseline, DOI metadata, local spine, tail-entry output, preservation obligations, reference/compilation audit data, terminal bibliography, historical witness, and non-certificate status. The ledger is audit-clean *if* $\Delta^{\mathrm{freeze}}_{\mathrm{audit}} = 0$*,* audit-repairable *if only mechanical or metadata repair is needed, and* not freeze-ready *otherwise.*

Lemma H4.365 (Audit ledger is metadata-bearing, not theorem-generating)**.** *The audit-freeze ledger records status, preservation, and release-readiness data. It cannot by itself create an OP closure certificate or change the theorematic status of OP 1/4 or OP 5(B).*

Proof. Every component of Definition H4.363 is either inherited mathematical state, preservation metadata, compilation/reference audit data, bibliography status, historical-witness status, or the explicit non-certificate marker. None of these components is a certificate test for OP 1/4 or OP 5(B). Therefore the ledger can freeze and audit the current line, but it cannot promote the current line into a theorematic closure. □

Theorem H4.366 (First audit-freeze ledger)**.** *After the executed-head tail-entry packet, the current HC/OP line admits the finite audit-freeze ledger* $F^{\mathrm{audit}}_{\mathrm{HCOP}}$ *with audit-freeze defect* $\Delta^{\mathrm{freeze}}_{\mathrm{audit}}$*. The ledger records the public baseline, DOI state, local spine, tail-entry output, preservation obligations, reference/compilation audit data, terminal bibliography status, historical witness, and no-false-certificate marker without changing theorematic status.*

Proof. The public baseline and DOI state are metadata inherited from the title and chronology records. The local spine and tail-entry output are supplied by Theorem H4.361 and its predecessors. The preservation obligations, reference/compilation checks, terminal bibliography status, historical witness, and non-certificate marker are already explicit release-audit obligations of the working line. Definition H4.363 collects these data into one finite record, and Lemma H4.365 prevents any promotion to OP closure. □

H4.25 Audit-freeze stabilization object

Definition H4.367 (Audit-freeze stabilization object)**.** *The* audit-freeze stabilization object *is*

$$F^{\mathrm{audit}}_{\mathrm{freeze}} = \left(F^{\mathrm{audit}}_{\mathrm{HCOP}}, \Delta^{\mathrm{freeze}}_{\mathrm{audit}}, \Pi_{\mathrm{allowed}}, S_{\mathrm{hold}}\right),$$

where Π_{allowed} *is the finite permission set*

$$\Pi_{\mathrm{allowed}} = \{\mathrm{audit}, \mathrm{mechanicalRepair}, \mathrm{metadataRepair}, \mathrm{releasePackaging}, \mathrm{zenodoDescription}\},$$

and S_{hold} *is the hold marker blocking further theorem expansion before the release audit is completed. The release label is metadata attached in prose and chronology, not a mathematical subscript or superscript of this object.*

Lemma H4.368 (Freeze permissions exclude theorem expansion)**.** *While* $F^{\mathrm{audit}}_{\mathrm{freeze}}$ *is active, the only admissible operations are audit, mechanical repair, metadata repair, release packaging, and Zenodo-description preparation. New theorem expansion is not an admissible freeze operation.*

Proof. The admissible operation set is exactly Π_{allowed} in Definition H4.367. None of its entries is a proof-extension, closure-promotion, OP-claim, or certificate-generation operation. Hence the active state may be repaired or packaged, but not mathematically expanded, until the release audit is discharged. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Theorem H4.369 (Audit-freeze stabilization)**.** *The current HC/OP working line is stabilized as an audit-freeze final by* $F_{\mathrm{freeze}}^{\mathrm{audit}}$*. In this state, the line is available for audit, mechanical repair, metadata repair, release packaging, and Zenodo-description preparation, while OP 1/4 and OP 5(B) remain open and no closure certificate is manufactured.*

Proof. Theorem H4.366 supplies the finite ledger and audit-freeze defect. Definition H4.367 adds only a finite permission set and a hold marker. Lemma H4.368 blocks theorem expansion during the freeze state. Therefore the line is audit-stabilized without changing the open status of OP 1/4 or OP 5(B). □

Corollary H4.370 (Audit-freeze working rule)**.** *Future work before the next public release may only audit, mechanically repair, repair metadata, package the release, or prepare the Zenodo description. Any new mathematical theorem expansion must wait until the freeze is either released or explicitly reopened.*

Proof. Immediate from Theorem H4.369 and Lemma H4.368. □

Phase I-C: Higher Confinement and Septinion Unfolding

This third unfolding block turns closure stability into a higher confinement reading and makes explicit why the septinion/seption interpretation is not an external axiom but a structurally selected reading of the admissible closure class.

Lemma H4.371 (Higher confinement lemma)**.** *If a structural class is closure-stable and remains invariant under repeated compensator-compatible normalization, then it admits a higher confinement reading.*

Proof. Closure stability excludes uncontrolled drift out of the admissible class, while repeated compensator-compatible normalization prevents the persistence of an independent escaping sector. Once both conditions hold, the admissible class is not merely locally normalized but globally retained under the structural return mechanisms of the system. This is exactly the content of higher confinement: the admissible states are no longer only corrected case by case but persist as a retained confinement regime. □

Lemma H4.372 (Non-escape lemma)**.** *Let x be a closure-stable state in the normalized core. Then no admissible transition generated within the core sends x into a structurally independent external sector.*

Proof. By closure stability, every admissible normalization step sends x back into the same closure class. If an admissible transition could produce a genuinely independent external sector, then closure stability would fail exactly at that point, because the state would no longer be returnable by the same structural normalization. Hence admissible transitions may rearrange or re-express states inside the confinement class, but they cannot export them into an independent outside sector. This is the structural non-escape property. □

Theorem H4.373 (Septinion readability theorem)**.** *If a class is closure-stable and admits the higher confinement reading, then it is septinion-readable: the resulting stabilized regime may be interpreted as a septinion/seption-level confinement of the normalized structural body.*

Proof. The septinion reading is not introduced as an arbitrary enlargement of the formalism. Rather, it names the level at which the closure-stable and non-escaping structural class is most naturally read as a higher unified container of the admissible sectors. By Lemma H4.371, closure stability already upgrades the system to a higher confinement regime, and Lemma H4.372 shows that admissible states remain inside that regime. The septinion/seption reading is therefore the natural structural reading of this higher confinement container. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Corollary H4.374 (HC as emergent structural consequence)**.** *Within the normalized core, HC is not an externally appended axiom. It appears as an emergent structural consequence of the one-body, frame-shift, and closure requirements.*

Proof. The generative chain of Phase I-A forces re-identification and therefore HC, while the closure chain of Phase I-B shows that HC is precisely the mechanism that stabilizes the admissible class. Phase I-C now identifies the higher confinement reading selected by this stabilization. Consequently HC is not an optional addition but the structurally emergent mechanism by which the unified body remains admissible across shifted readouts and under closure. □

Corollary H4.375 (First confinement forcing chain)**.** *The first unfolded confinement route of the normalized core is*

$$\textit{closure class} \Longrightarrow \textit{higher confinement} \Longrightarrow \textit{non-escape} \Longrightarrow \textit{septinion reading.}$$

Together with the preceding phases, this yields

$$\begin{aligned}&\textit{one body} \Longrightarrow \textit{re-identification} \Longrightarrow \mathrm{HC}\\ &\Longrightarrow \textit{closure} \Longrightarrow \textit{higher confinement} \Longrightarrow \textit{septinion/seption reading.}\end{aligned}$$

Proof. Proposition H4.254 provides the stabilized closure class. Lemma H4.371 upgrades this class to a higher confinement reading, Lemma H4.372 gives the internal retention property of admissible transitions, and Theorem H4.373 identifies the corresponding septinion/seption interpretation. The final displayed chain is obtained by concatenating this phase with the forcing chains of Phases I-A and I-B. □

Phase II-D: Cascade, Readout, and the Collapse-Fixed Sector

Lemma H4.376 (Cascade admissibility lemma)**.** *If x belongs to the closure-stable class selected by the normalized core, then every admissible cascade image $\hat{K}(x)$ remains inside the admissible readout class of the same unified body.*

Proof. The cascade is introduced as a composition of projection maps from higher strata toward the ground sector, not as an ontological splitting of the body. By Corollary H4.257, closure-stable states are precisely those stabilized against drift under shifted readouts. Applying the cascade to such a state therefore yields another admissible readout of the same body rather than a new object. Hence admissibility is preserved along the cascade image of any closure-stable input. □

Lemma H4.377 (Terminal ground-state lemma)**.** *The ground state S_0 is the terminal invariant readout sector of the cascade: for every admissible state x, one has $\hat{K}(x) = x$ exactly when $x \in S_0$.*

Proof. This is the normalized readout form of the ground-state condition already built into the semantic kernel. The cascade terminates in S_0, and terminality means precisely that no further nontrivial readout step occurs. Therefore the fixed-point condition $\hat{K}(x) = x$ is equivalent to membership in the terminal sector S_0. □

Lemma H4.378 (Global null/light-axis lemma)**.** *The null-axis structure selected by the normalized core, and read physically as the light axis at stable propagation, is stratified across the cascade: if a state is read on this null/light axis at one admissible layer, then its admissible readouts remain aligned with the same global null/light-axis class through the cascade.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. The Lorentz/null-axis structure, read physically as the light axis, is not introduced as a local afterthought but as a stratified feature of the whole construction. The cascade only changes the readout level, not the structural identity of the body. Since admissible cascade images preserve the same unified body and the same closure-compatible sector, the null/light-axis reading persists as a global class across the projected layers. □

Theorem H4.379 (Collapse-fixed sector theorem)**.** *The photon/readout sector identified in the normalized core is the global collapse-fixed sector of the cascade. In particular, the sector denoted by S_0 is not merely a local null readout but the terminal collapse-fixed sector selected by the full stratified structure.*

Proof. Lemma H4.376 ensures that the cascade preserves admissibility, while Lemma H4.377 identifies S_0 as the terminal invariant sector. Lemma H4.378 shows that the light-axis reading persists globally across admissible layers. Therefore the photon/readout sector is not obtained from an isolated local calculation alone; rather, it is the collapse-fixed terminal sector selected by the entire cascade of admissible projections. □

Lemma H4.380 (Readout separation lemma)**.** *Within the normalized core one must distinguish three levels: structural forcing, cascade identification, and physical reading. A statement belongs to exactly one of these levels unless an explicit bridge is given.*

Proof. The structural forcing chain establishes what must hold inside the unified body. The cascade then identifies how such a structurally forced state appears at lower readout levels. A physical interpretation is an additional reading of that identified sector. Conflating these levels obscures where a statement is proved and where it is interpreted. The separation therefore serves as a normalization rule for subsequent unfolding steps. □

Corollary H4.381 (First readout forcing chain)**.** *The first unfolded readout route of the normalized core is*

$$\begin{aligned} &\textit{closure-stable class} \Longrightarrow \textit{admissible cascade image} \\ &\Longrightarrow \textit{terminal ground sector} \Longrightarrow \textit{collapse-fixed photon/readout sector.} \end{aligned}$$

Taken together with Phases I-A through I-C, the normalized core now yields the combined chain

$$\begin{aligned} &\textit{one body} \Longrightarrow \textit{re-identification} \Longrightarrow \mathrm{HC} \Longrightarrow \textit{closure} \\ &\Longrightarrow \textit{higher confinement} \Longrightarrow \textit{septinion reading} \\ &\Longrightarrow \textit{cascade-fixed readout sector.} \end{aligned}$$

Proof. The first displayed chain is the direct concatenation of Lemma H4.376, Lemma H4.377, and Theorem H4.379. The second displayed chain is obtained by appending this readout route to the forcing chain from Corollary H4.375. □

Phase II-E: Semantic/Tensor Interface and Millennium Intertwining

Lemma H4.382 (Semantic kernel well-formedness lemma)**.** *The normalized semantic kernel is well formed: its predicates of one-body identity, triolic organization, orthogonal anchoring, frame-shifted readout, HC forcing, closure stability, higher confinement, and readout admissibility do not define competing ontologies but successive structural constraints on one and the same body.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. By construction, the normalized core begins from the one-body condition and then adds only constraint predicates that regulate how the same body may be read, stabilized, and projected. None of the predicates introduces a second carrier space or an ontologically independent duplicate; each one refines the same underlying body by imposing an admissibility condition. The semantic kernel is therefore well formed as a constraint hierarchy rather than a family of competing state spaces. □

Lemma H4.383 (Tensor realization lemma)**.** *The tensorial operators $\mathcal{J}^{\mu}{}_{\nu}$, $\mathcal{C}^{\mu}{}_{\nu}$, $\Pi_n{}^{\mu}{}_{\nu}$, and $\hat{K}^{\mu}{}_{\nu}$ realize the normalized semantic kernel as an operator system: $\mathcal{J}$ realizes the involutive frame symmetry, $\mathcal{C}$ the compensating projection, Π_n the cascade steps, and $\hat{K}$ the terminal readout map.*

Proof. The semantic kernel distinguishes involutive reflection, compensating return, projection through admissible layers, and terminal ground readout. The tensorial operators are introduced precisely to encode these roles in compact operator notation. The semantic and tensorial descriptions therefore stand in a role-wise correspondence: the operator system is not a second theory but a tensor realization of the already normalized semantic structure. □

Lemma H4.384 (Intertwining lemma)**.** *If the Millennium matrix $\mathcal{M}^{\mu}{}_{\nu}$ is used as the structural transport operator between admissible grouped layers, then it must intertwine with the normalized tensorial kernel. In the minimal normalized form one requires*

$$\mathcal{M}\mathcal{J} = \mathcal{J}\mathcal{M}, \qquad \mathcal{M}\mathcal{C} = \mathcal{C}\mathcal{M}, \qquad \mathcal{M}\hat{K} = \hat{K}\mathcal{M}$$

whenever the corresponding involution, compensation, and cascade-readout structures are to be preserved across the transport.

Proof. A transport operator that fails to commute with the involution would change the frame logic of the transported state; one that fails to commute with the compensator would alter admissibility by moving states in or out of the compensated sector; and one that fails to commute with the terminal readout map would not preserve the normalized collapse/readout architecture. Hence a Millennium transport that is meant to be structure preserving must satisfy the stated intertwining relations at least for the operators whose semantic role it claims to preserve. □

Theorem H4.385 (Admissibility preservation theorem)**.** *If x is admissible in the normalized semantic kernel and $\mathcal{M}$ is a Millennium transport satisfying the intertwining relations of Lemma H4.384, then $\mathcal{M}x$ remains admissible in the transported layer.*

Proof. Admissibility in the normalized core is carried by compatibility with the one-body constraint, compensator normalization, and the cascade-compatible readout structure. By Lemma H4.384, the transport operator preserves the tensorial realizations of exactly these mechanisms. Therefore applying $\mathcal{M}$ cannot convert an admissible state into an inadmissible one merely by transport; it yields an admissible image in the next layer provided the transport is itself taken within the normalized grouped architecture. □

Theorem H4.386 (Closure preservation theorem)**.** *Under the same hypotheses, closure-stable states are transported by $\mathcal{M}$ to closure-stable states of the receiving layer.*

Proof. Closure stability is the semantic statement that the admissible state does not drift out of the normalized class once HC forcing and anchoring are taken into account. Since admissibility is preserved by Theorem H4.385 and the intertwining relations preserve the compensator and readout structures that stabilize the closure class, the transported image cannot lose closure merely by layer transport. Thus closure stability is preserved across admissible Millennium transport. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proposition H4.387 (Semantic–tensor equivalence proposition)**.** *The tensorial kernel does not introduce a second structural theory. It is a parallel notation for the normalized semantic kernel, and the Millennium matrix is admissible only insofar as its tensorial action represents the same constraint chain already present on the semantic side.*

Proof. By Lemma H4.382, the semantic side is a single constrained body. By Lemma H4.383, the operator side realizes these constraints in tensorial notation. The intertwining and preservation results show that $\mathcal{M}$ is not free to create a new structural doctrine; it is licensed only as a transport operator respecting the same normalized constraint chain. Hence semantic kernel and tensor kernel are parallel realizations of one structure. □

Corollary H4.388 (First semantic–tensor forcing chain)**.** *The normalized semantic/tensor interface yields the chain*

$$\begin{aligned} &\textit{semantic kernel} \Longrightarrow \textit{tensor realization} \Longrightarrow \textit{Millennium intertwining} \\ &\Longrightarrow \textit{admissibility preservation} \Longrightarrow \textit{closure preservation.} \end{aligned}$$

Proof. Lemma H4.382 establishes the semantic kernel, Lemma H4.383 gives its tensor realization, Lemma H4.384 fixes the admissible transport relations for $\mathcal{M}$, and Theorems H4.385 and H4.386 provide the preservation steps. □

Grouped Layer Architecture and Millennium Transport

The semantic/tensor kernel admits a second normalization in which the document's many local cores, layers, and readout blocks are collected into three grouped structural classes. This reduces duplication and makes the role of the Millennium matrix explicit as a transport operator between admissible groups.

Definition H4.389 (Grouped structural architecture)**.** *The normalized structural architecture consists of the following three grouped layers:*

(G1) ***Generative layer $\mathfrak{G}$:*** *the layer of structural formation. It contains the one-body condition, triolic minimality, orthogonal anchoring, frame-shifted observation, the ground condition, HC forcing, closure admissibility, and the higher-confinement/septinion reading.*

(G2) ***Projection/translation layer $\mathfrak{P}$:*** *the layer of cascade readout, sector selection, projection rules, collapse-fixed identifications, null-/light-axis readings, photon-sector readings, and admissible translation between equivalent structural descriptions.*

(G3) ***Field/probe realization layer $\mathfrak{F}$:*** *the layer of probe states, two-probe geometry, closure-preserving field equations, interaction rules, and physical realization of the admissible structure.*

These three grouped layers are not independent ontologies. They are different organizational regimes of the same underlying body and must therefore remain compatible under admissible transport.

Definition H4.390 (Millennium transport between grouped layers)**.** *The Millennium matrix is interpreted at the grouped-layer level as a structure-preserving converter*

$$\mathcal{M}_{\mathfrak{G}\to\mathfrak{P}} : \mathfrak{G} \to \mathfrak{P}, \qquad \mathcal{M}_{\mathfrak{P}\to\mathfrak{F}} : \mathfrak{P} \to \mathfrak{F},$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

and, whenever reverse structural checking is required, also by admissible back-transport maps

$$\mathcal{M}_{\mathfrak{P}\to\mathfrak{G}} : \mathfrak{P} \to \mathfrak{G}, \qquad \mathcal{M}_{\mathfrak{F}\to\mathfrak{P}} : \mathfrak{F} \to \mathfrak{P}.$$

At this level $\mathcal{M}$ is not merely a notational relabeling but a converter which must preserve structural admissibility across layer changes.

Proposition H4.391 (Grouped forcing and readout chain)**.** *The grouped architecture organizes the document into the structural chain*

$$\mathfrak{G} \xrightarrow{\mathcal{M}} \mathfrak{P} \xrightarrow{\mathcal{M}} \mathfrak{F},$$

where:

(a) $\mathfrak{G}$ *generates the admissible structure,*

(b) $\mathfrak{P}$ *reads, projects, and translates that structure without changing its admissibility class,*

(c) $\mathfrak{F}$ *realizes the same admissible structure as a field/probe system.*

In particular, the field/probe layer is not a second origin of the theory; it is a realization layer of structure first generated in $\mathfrak{G}$ *and made readable in* $\mathfrak{P}$*.*

Proof. The generative layer contains the forcing data established above: one-body, triolic organization, orthogonal anchor, frame shift, HC, and closure. The projection/translation layer does not create new admissibility; it converts the generative core into cascade-compatible, sector-resolved, and physically legible readout. The field/probe layer then realizes those admissible readouts as explicit probes, field equations, or interaction geometries. Since the same body is being followed through all three layers, admissible transport by $\mathcal{M}$ must preserve rather than replace the structural class. □

Proposition H4.392 (Admissibility preservation across grouped layers)**.** *Let x be an admissible state generated in* $\mathfrak{G}$*. If $\mathcal{M}$ acts as an admissible grouped-layer converter, then*

$$\text{Admissible}(x) \Longrightarrow \text{Admissible}(\mathcal{M}_{\mathfrak{G}\to\mathfrak{P}}x) \Longrightarrow \text{Admissible}(\mathcal{M}_{\mathfrak{P}\to\mathfrak{F}}\mathcal{M}_{\mathfrak{G}\to\mathfrak{P}}x).$$

Moreover, when closure stability is part of the admissibility data, it is to be preserved throughout this transport chain.

Proof. By Definition H4.390, grouped-layer transport is only allowed when it preserves structural admissibility. Hence an admissible state generated in $\mathfrak{G}$ remains admissible after conversion into the projection/translation layer, and likewise after conversion into the field/probe layer. If closure stability were lost during transport, the result would no longer be a valid readout or realization of the original structure but a structural rupture. Therefore closure stability must be preserved whenever it belongs to the originating admissibility class. □

Remark H4.393 (Document-normalization consequence)**.** *The grouped architecture provides a normalization principle for the text. Local "cores" or "integration blocks" should be read either as part of* $\mathfrak{G}$*, as part of* $\mathfrak{P}$*, or as part of* $\mathfrak{F}$*, rather than as independent foundational centers. This removes duplication: the normative foundation lies in* $\mathfrak{G}$*, the translation/readout logic in* $\mathfrak{P}$*, and the field/probe realization in* $\mathfrak{F}$*.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Phase II-F: Grouped Layer Transport and Ontological Non-Drift

This unfolding block upgrades the grouped architecture from a normalization note into an explicit transport calculus. The goal is to show that grouped layers do not introduce new origins of structure, but only different admissible regimes of one and the same underlying body.

Lemma H4.394 (Group decomposition lemma)**.** *Every active structural block of the normalized Companion is dominantly assignable to exactly one of the grouped layers* $\mathfrak{G}$*,* $\mathfrak{P}$*, or* $\mathfrak{F}$*, even when secondary cross-references to the other layers remain.*

Proof. By Definition H4.389, the grouped layers are distinguished by role rather than by separate ontology: generation, projection/translation, and realization. Any active block therefore has one primary structural function. Cross-references may connect a block to the other layers, but they do not erase its dominant role. Hence each active block is assignable to one grouped layer as its principal location. □

Lemma H4.395 (Transport lemma)**.** *Let* x *be a state generated in* $\mathfrak{G}$*. If* $\mathcal{M}_{\mathfrak{G}\to\mathfrak{P}}$ *is an admissible grouped-layer converter, then its image in* $\mathfrak{P}$ *is a projection/translation of the same admissible structure and not a newly created structural body.*

Proof. By Definition H4.390, admissible grouped transport preserves structural admissibility. Proposition H4.391 states that $\mathfrak{P}$ reads and translates what $\mathfrak{G}$ has already generated. Therefore the image of x under $\mathcal{M}_{\mathfrak{G}\to\mathfrak{P}}$ remains the same admissible structure under a projection/translation regime. It is not a second origin because admissibility is transported, not recreated. □

Lemma H4.396 (Realization lemma)**.** *If* $y \in \mathfrak{P}$ *is an admissible projection/translation state and* $\mathcal{M}_{\mathfrak{P}\to\mathfrak{F}}$ *is admissible, then its image in* $\mathfrak{F}$ *is a realization of the same admissible structure as a field/probe state.*

Proof. The role of $\mathfrak{F}$ is realization rather than generation by Proposition H4.391. Since admissibility is preserved across grouped transport by Proposition H4.392, $\mathcal{M}_{\mathfrak{P}\to\mathfrak{F}}(y)$ remains within the same admissibility class. Thus the field/probe layer realizes the translated structure without changing its underlying body. □

Proposition H4.397 (No-ontology-drift proposition)**.** *Admissible transport across the grouped layers does not create ontological multiplicity. If*

$$x \in \mathfrak{G}, \qquad x_P := \mathcal{M}_{\mathfrak{G}\to\mathfrak{P}}(x), \qquad x_F := \mathcal{M}_{\mathfrak{P}\to\mathfrak{F}}(x_P),$$

then x*,* x_P*, and* x_F *are three admissible organizational regimes of one and the same structural body.*

Proof. By Lemma H4.395, passage from $\mathfrak{G}$ to $\mathfrak{P}$ produces a translation/readout of the same admissible body. By Lemma H4.396, passage from $\mathfrak{P}$ to $\mathfrak{F}$ produces a field/probe realization of that same admissible body. Since both steps preserve admissibility rather than replacing it, no new ontology is introduced along the chain. Therefore x, x_P, and x_F are structurally identical in body and differ only by organizational regime. □

Corollary H4.398 (First grouped transport forcing chain)**.** *The normalized grouped architecture supports the forcing chain*

$$\mathfrak{G} \Longrightarrow \textit{admissible transport to } \mathfrak{P}$$
$$\Longrightarrow \textit{admissible realization in } \mathfrak{F} \Longrightarrow \textit{no ontology drift across grouped layers.}$$

Proof. Combine Lemma H4.394, Lemma H4.395, Lemma H4.396, and Proposition H4.397. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Phase III-G: Probe Geometry, Inheritance, and Guided Search

This unfolding block upgrades the probe sector from an interpretive add-on to a structurally inherited realization of the normalized core. The goal is to show that probes do not introduce a foreign search mechanism, but inherit triolic, anchored, and closure-compatible structure from the generative and grouped layers.

Lemma H4.399 (Probe inheritance lemma)**.** *If the normalized structural body is triolic, anchored, and closure-admissible, then every admissible probe realization inherited through* $\mathfrak{F}$ *carries a corresponding triolic and closure-compatible organization.*

Proof. By Proposition H4.397, admissible realization in $\mathfrak{F}$ does not create a second ontology, but realizes the same underlying structural body under a field/probe regime. Hence structural features that belong to the admissible body—in particular triolic organization and anchor-compatible closure discipline—remain available in the probe image as inherited structure rather than externally attached decoration. □

Lemma H4.400 (Two-probe compatibility lemma)**.** *Let* P_1 *and* P_2 *be two admissible probes directed toward one another. If both are inherited from the same normalized structural body, then their mutual configuration is compatible with one common closure-governed geometry.*

Proof. By Lemma H4.399, each probe inherits the same triolic and closure-compatible structural discipline. Since neither probe introduces a new ontology by Proposition H4.397, their mutual interaction must be read inside one common admissibility class. Therefore the two-probe configuration is not a collision of unrelated structures but a common closure-governed geometry of one inherited body. □

Lemma H4.401 (Orthogonal third-group lemma)**.** *In an admissible two-probe configuration, the third structural group remains orthogonal to both probes and thereby determines the constrained search region.*

Proof. The orthogonal third direction is already forced at the generative level by the orthogonal-anchor principle and remains preserved under admissible realization. Hence, when two probes are realized as inherited triolic structures, the third group cannot collapse into either probe direction without violating anchor compatibility. It therefore remains orthogonal to both and serves as the stabilizing transversal direction that bounds the admissible search region. □

Proposition H4.402 (Probe reduction theorem)**.** *An admissible two-probe configuration reduces to the closure-preserving sector of the grouped architecture. In particular, the probe interaction may be analyzed inside the admissible closure class without introducing a second search ontology.*

Proof. By Lemma H4.400, the two probes share one common closure-governed geometry. By Lemma H4.401, the third group fixes the transversal admissible search region. Since closure and ontology are preserved across grouped realization by Proposition H4.397, the probe interaction reduces to the closure-preserving sector of the same normalized architecture. □

Corollary H4.403 (Guided search corollary)**.** *Auxiliary tools such as frequency extraction, transition guidance, structural routing, or convergence acceleration act only within the already admissible probe geometry. They do not generate admissibility, but exploit a search region that has already been restricted by inheritance, closure, and orthogonal anchoring.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. By Proposition H4.402, admissible probe analysis is already confined to the closure-preserving sector. Therefore any auxiliary tool operates on a domain whose admissibility has been fixed before the tool is applied. The tool may compress or guide the search, but it does not create the search geometry itself. □

Corollary H4.404 (First probe forcing chain)**.** *The normalized probe sector supports the forcing chain*

$$\begin{aligned} &\textit{inherited probe structure} \Longrightarrow \textit{two-probe compatibility} \\ &\Longrightarrow \textit{orthogonal third-group search region} \\ &\Longrightarrow \textit{closure-preserving probe reduction} \\ &\Longrightarrow \textit{guided search inside an already admissible geometry.} \end{aligned}$$

Proof. Combine Lemma H4.399, Lemma H4.400, Lemma H4.401, Proposition H4.402, and Corollary H4.403. □

Phase III-H: Structural Master Equation and Exact-Closure Field Regime

This unfolding block upgrades the previously compressed master-equation language into an explicit field-level forcing chain. The aim is not yet to derive every physical limit in full detail, but to show that the structural master equation, its variational reading, the closure-preserving field regime, and the exact- closure simplifications belong to one continuous argument rather than to loosely connected interpretive remarks.

Lemma H4.405 (Structural master-equation lemma)**.** *The structural master equation is the compressed field-level summary of the normalized forcing chain from one-body structure through re-identification, HC, closure, higher confinement, and admissible readout.*

Proof. By Corollary 29.16, the normalized body forces re-identification and HC. By Corollary H4.257, this induces a closure class with either exact closure or a controlled residue. By Corollary H4.375, closure yields higher confinement and the septinion reading. By Corollary H4.381 and Corollary H4.388, admissible readout and tensorial transport preserve this structure. Hence the master equation is not an additional postulate but the compressed field-level expression of the forcing chain already established. □

Lemma H4.406 (Variational consistency lemma)**.** *Any variational reading of the structural master equation is admissible only if it preserves the normalized one-body, HC, and closure conditions. In particular, admissible variation may unfold the structural equation but may not replace its forcing chain by an independent ontology.*

Proof. By Proposition H4.387, tensorial formulation is only a parallel realization of the same semantic kernel. Therefore any variational formulation that is meant to belong to the present architecture must preserve the same semantic admissibility data. If the variation were to destroy one-body identity, HC-forcing, or closure preservation, it would cease to be a variation of the normalized structural equation and would instead define a second independent system. Hence admissible variation is constrained by the forcing chain and remains structurally internal to it. □

Lemma H4.407 (Closure-field compatibility lemma)**.** *The field-level realization of the structural master equation is compatible with closure precisely when the realized dynamics preserves the HC-selected admissibility class and does not mix the stable closure sector with an ontologically foreign complement.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. By Theorem H4.386, admissible transport through the Millennium interface preserves closure classes. By Proposition H4.397, realization in the field/probe layer introduces no second ontology. Therefore a field realization is closure-compatible exactly when it remains inside the HC-selected admissibility class rather than coupling the closure-stable sector to a foreign complement. This is the field-level restatement of normalized closure preservation. □

Theorem H4.408 (Exact-closure field theorem)**.** *In the exactly closed regime, the realized field dynamics reduces to the closure-stable block of the normalized architecture. Equivalently, the active field transport becomes block-compatible with the exact closure class.*

Proof. By Proposition H4.255, exact closure is equivalent to the disappearance of the nontrivial closure defect. By Lemma H4.407, closure-compatible field dynamics may then act only on the already stabilized admissibility class. Since no ontology drift occurs across grouped transport, the realized field dynamics reduces to the closure-stable block itself. Thus exact closure yields a block-compatible field regime. □

Lemma H4.409 (Minimal compensator disappearance lemma)**.** *In the exactly closed regime, the minimal compensator residue vanishes. What remains is not the disappearance of structural compensation as such, but the fact that no additional compensator correction is required beyond the exact closure block.*

Proof. By Lemma H4.256, the non-exact regime is marked by a residual deviation from exact closure. By Theorem H4.408, exact closure confines the field transport completely to the closure-stable block. Therefore the additional minimal compensator term that measures residual off-block correction must vanish. Structural compensation remains implicit in the exact block, but no separate compensator residue survives. □

Theorem H4.410 (Effective-limit theorem)**.** *Any admissible effective limit—for example a geometric, spinorial, or quantum field-theoretic limit—must arise as a readout of the normalized structural master equation together with its closure-preserving field realization. Such a limit may specialize the realization, but cannot bypass the forcing chain that produces it.*

Proof. By Lemma H4.405, the structural master equation is the compressed summary of the normalized forcing chain. By Lemma H4.406 and Lemma H4.407, any admissible field realization and any admissible variation remain internal to that chain. By Theorem H4.408 and Lemma H4.409, the exact regime yields a pure closure block without additional corrective residue. Hence any effective limit must be read as a specialization or projection of the normalized structural field realization rather than as an independent starting point. □

Corollary H4.411 (First field forcing chain)**.** *The normalized field sector supports the forcing chain*

$$\begin{aligned}&\textit{structural master equation} \Longrightarrow \textit{variational consistency}\\ &\Longrightarrow \textit{closure-preserving field compatibility} \Longrightarrow \textit{exact closure block regime}\\ &\Longrightarrow \textit{vanishing minimal compensator residue} \Longrightarrow \textit{admissible effective limit readout.}\end{aligned}$$

Proof. Combine the structural master equation, variational consistency, closure-field compatibility, the exact closure-field theorem, minimal compensator disappearance, and the effective limit theorem. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Concrete mapping of legacy core blocks to grouped layers

The grouped normalization can be made explicit by assigning the surviving legacy core blocks to the three grouped layers $\mathfrak{G}$, $\mathfrak{P}$, and $\mathfrak{F}$. This turns older version-labelled cores into either active parts of the normalized architecture or historical/supporting blocks.

Definition H4.412 (Layer assignment of legacy core blocks)**.** *Within the present monolith, the principal legacy blocks are to be read as follows.*

(L1) ***$\mathfrak{G}$-dominant blocks (generative layer):***

- *(a) the normalized* Structural Generation Core,
- *(b)* Structural Core and Closure Principles,
- *(c) the generative part of* archival stage Consolidation Core, *especially the structural reading discipline together with the ground-condition/HC/closure and higher-confinement logic,*
- *(d) the generative part of* archival stage Derivation and Appendix Integration Core, *especially the closure ladder and the micro-to-macro derivation chain.*

(L2) ***$\mathfrak{P}$-dominant blocks (projection/translation layer):***

- *(a)* archival stage Projection and Appendix Table Integration Core,
- *(b) the appendix-coupling and projection-summary parts of* archival stage Derivation and Appendix Integration Core,
- *(c) the collapse, light-axis, photon-sector, and cascade-readout parts of the later main sections,*
- *(d) all readout, table, projection, and sector-translation interfaces whose role is to make one admissible structure legible without changing its admissibility class.*

(L3) ***$\mathfrak{F}$-dominant blocks (field/probe realization layer):***

- *(a) the probe-sector and two-probe geometry blocks,*
- *(b) the field-equation, interaction, and closure-preserving dynamics sections,*
- *(c) the probe-reduction and guided-search realizations insofar as they are used as operational realizations of previously fixed admissible structure.*

(L4) ***Historical/supporting blocks:*** *readiness criteria, version-status passages, and older local consolidation summaries are not to be read as new foundational centers. They are historical normalization records unless they are explicitly re-exported into one of $\mathfrak{G}$, $\mathfrak{P}$, or $\mathfrak{F}$.*

Proposition H4.413 (Normalization map for duplicated core material)**.** *Under the grouped architecture, duplicated structural material is normalized by the following rule.*

(i) When a legacy block states formation conditions, forcing rules, or closure-generating principles, it is normalized into $\mathfrak{G}$.

(ii) When a legacy block states projection rules, cascade-compatible readout, sector identifications, appendix transport, or summary-table logic, it is normalized into $\mathfrak{P}$.

(iii) When a legacy block states probes, field equations, interaction channels, or closure-preserving realizations, it is normalized into $\mathfrak{F}$.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(iv) If a block contains mixed material, it is to be read by dominant function, with the remaining content treated as support, historical context, or material that should later be split explicitly.

In particular, archival stage Consolidation Core *and the early derivation cores no longer define independent foundations beside the normalized structural core; they contribute either to* $\mathfrak{G}$*, to* $\mathfrak{P}$*, or to historical context.*

Remark H4.414 (Immediate practical consequence for the next clean release)**.** *The next clean release should therefore not keep several local cores in competition. Instead, older version-labelled cores are to be annotated and, where necessary, shortened according to their dominant grouped-layer function: formation material belongs to* $\mathfrak{G}$*, translation/readout material to* $\mathfrak{P}$*, field/probe realization material to* $\mathfrak{F}$*, and pure status/history material to archival support.*

archival stage Expansion Core

Remark H4.415 (Normalization note for archival stage)**.** *This block is retained as a historical expansion precursor. Its proof-bearing content is now to be read primarily through Phase I, Phase II-D, and the normalized grouped architecture. In particular, the intermediate master-equation ladder and the micro-to-macro chain no longer function as independent foundations beside the later unfolding phases, but as early organizational anticipations of them.*

Version archival stage is the first release that does more than freeze the next work path. It adds a genuine expansion layer on top of the archival stage consolidation core. The operative aim is threefold: first, the structural master equation is no longer presented only as a compressed global summary, but also as the endpoint of an explicit intermediate chain; second, the passage from local triolic organization to macroscopic effective geometry is stated in staged form; third, the technical apparatus is widened so that later proof layers can be attached without rewriting the monolith.

1. Deepened master-equation programme

The master equation in the later main section remains the global compression of one common closure-stable dynamics. What archival stage adds is a more explicit three-level reading of that equation.

(i) **Structural level.** One starts from triolic organization, orthogonality, frame shift, compensator action, and closure. At this level, the equation is still read as a balance law on admissible sectors.

(ii) **Mesoscopic level.** The same balance law is rewritten as an effective transport-and-selection equation for coupled sectors and probes. Here one already sees how admissible interaction channels are filtered by HC and by the closure residue.

(iii) **Macroscopic level.** After suppression of unstable local residues and after passage to a closure-stable observer description, the same equation becomes readable as effective matter dynamics plus background geometry.

In this sense, archival stage does not replace the later master equation (7); it inserts the missing ladder by which that equation is reached.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A convenient intermediate schematic form is

$$\mathcal{E}_{\mathrm{micro}}[\Psi] + \mathcal{E}_{\mathrm{coupl}}[\Psi, \Pi] + \mathcal{E}_{\mathrm{HC}}[\Psi] + \mathcal{E}_{\mathrm{cl}}[\Psi] + \mathcal{E}_{\mathrm{obs}}[\Psi] = 0,$$

where the five blocks stand respectively for local triolic structure, sectorial coupling/probe interaction, Heisenberg-compensator balancing, closure selection, and observer-frame projection. The later master equation is to be read as the compressed global representative of this staged balance.

Lemma H4.416 (Intermediate balance principle)**.** *Every admissible global solution must pass through an intermediate regime in which local triolic transport, compensator balancing, and closure selection are all simultaneously active. Consequently, a direct jump from isolated local structure to macroscopic field form is structurally forbidden.*

Proof. Local triolic data alone do not yet determine which coupled configurations are admissible; they only provide possible sectorial directions. Conversely, a macroscopic field description is meaningful only after compensator balancing has removed representation drift and closure has excluded unstable residues. Therefore at least one intermediate regime must exist in which coupling, HC, and closure are jointly active. This regime is the structural source of the mesoscopic reading introduced in archival stage. □

2. Micro-to-macro closure chain

The new hardening target of archival stage is the explicit micro-to-macro chain

$$\begin{aligned}&\text{local triolic unit} \longrightarrow \text{coupled closure cell} \longrightarrow \text{probe-filtered admissible channel}\\ &\longrightarrow \text{closure-stable sector regime} \longrightarrow \text{effective global geometry and field dynamics.}\end{aligned}$$

The purpose of this chain is not rhetorical. It separates five logically distinct burdens that earlier drafts could too easily blend together: local decomposition, coupling, admissibility filtering, stable sector assembly, and macroscopic readout.

Local triolic unit. At the smallest relevant level, the system carries the standing triolic organization together with the orthogonal third direction. This level fixes the minimal sector inventory but does not yet decide which transitions survive.

Coupled closure cell. Once two or more local units interact, the question becomes whether their combination admits a common compensator-compatible representative. A coupled cell is called closure-ready if its local mismatches can be balanced without breaking the orthogonality and frame-shift discipline.

Probe-filtered admissible channel. The probe construction becomes technically important at precisely this stage. Two directed probes together with the orthogonal third group define a restricted search geometry. Kasisky-, hash-, rainbow-, Markov-, and Newton-type tools are not external decoration here; they are the operational means by which the space of final channels is compressed to closure-compatible paths.

Closure-stable sector regime. Only after probe filtering and compensator balancing does one obtain a stable sector regime. This is the first level at which effective matter, light/collapse, and dual sectors can be read as persistent rather than transient descriptions.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Effective global geometry and field dynamics. Macroscopic geometry is then not glued on from outside. It is the readout of a large-scale assembly of closure-stable sector regimes viewed in the phase-shifted observer frame. The gravitational, QFT-type, and Dirac-type limits are therefore downstream realizations of one common filtered closure architecture.

Proposition H4.417 (Macroscopic readout requires closure-stable assembly)**.** *A macroscopic field or geometric description is admissible only if it is the observer-frame image of an assembly of closure-stable sector regimes. In particular, large-scale geometry cannot be read directly from uncoupled local triolic data.*

Proof. Uncoupled local data do not yet fix admissible transport and do not suppress representation drift. The observer frame, however, reads only stabilized configurations. Since stabilization in the present framework is exactly the combined action of HC and closure across coupled cells, the macroscopic image must descend from an assembly that has already passed the closure filter. □

3. Extended lemma and tool apparatus

Version archival stage also widens the technical apparatus that the next proof layer can rely on. The point is not to over-formalize prematurely, but to mark where repeated arguments will later be detached into reusable blocks.

Core lemmas now fixed by role.

1. **Orthogonality lemma:** the third triolic direction remains the non-negotiable constraint that prevents degeneracy of the sector picture.
2. **Frame-shift lemma:** observational readout differs from the underlying structural frame by a controlled phase displacement.
3. **Compensator lemma:** HC is the minimal identification mechanism needed to prevent sectorial drift.
4. **Closure lemma:** admissibility is equivalent to vanishing of the relevant closure residue in the chosen regime.
5. **Probe admissibility lemma:** the probe sector selects channels that are compatible with both closure and orthogonality.
6. **Read-off lemma:** physical sector names arise only after the preceding structural filters have been passed.

Normalized tool classes. The tool catalogue of archival stage is retained, but archival stage upgrades its use: closure-tools act as admissibility tests, frame-shift tools as translation operators between structure and observer descriptions, projection-tools as controlled reduction maps, probe-tools as search-space compressors, equivalence-tools as representative-identification devices, and read-off tools as disciplined output maps. This normalization matters because later technical proofs can now cite a *tool class* instead of repeating the surrounding motivation every time.

Corollary H4.418 (Tool usage discipline)**.** *No computational shortcut is admissible in the proof architecture unless its role can be classified as one of the normalized tool classes and unless its use preserves the structural ordering*

closure before projection before interpretation.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. A shortcut that bypasses closure may produce effective formulas without fixing admissibility; a shortcut that bypasses projection discipline may confuse structural content with observer naming. The normalized tool classes are introduced exactly to block such category errors. Hence admissible shortcuts are those that preserve the structural order stated above. □

4. Version status of archival stage

With these additions, archival stage should be read as the first expansion-stage companion release. Relative to archival stage, the new version contributes three hard improvements: the master-equation programme is given an explicit intermediate ladder, the micro-to-macro chain is stated as a formal burden rather than a mere promise, and the lemma/tool apparatus is widened enough that subsequent versions can deepen proofs without reopening the global organization of the monolith. The next versions may therefore focus on tightening derivations,

$$\text{ground condition} \Longrightarrow \text{triolic admissibility} \Longrightarrow \text{HC balancing}$$
$$\Longrightarrow \text{closure filtering} \Longrightarrow \text{observer-frame readout.}$$

Version archival stage turns the earlier expansion layer into a genuine lemma-development release. The aim is not to replace the existing structural chapters, but to make the repeated internal moves of the proof architecture explicit enough that later sections may cite them as stable intermediate devices. Three burdens are addressed here: the closure spine is split into reusable intermediate steps, the passage from probes to admissible sectors is cast as a sequence of named reductions, and the projection discipline is tied to a small but durable family of proof lemmas.

Lemma H4.419 (Triolic admissibility precedes compensator balancing)**.** *A configuration cannot enter the Heisenberg-compensator balancing regime unless its triolic decomposition is already admissible with respect to the standing orthogonality and frame-shift constraints.*

Proof. The compensator does not create a triolic decomposition; it only identifies and balances compatible representatives across already existing sector data. If the triolic organization were itself degenerate or incompatible with the orthogonal third direction, then there would be no well-defined data for HC to balance. Hence triolic admissibility is logically prior to the compensator step. □

Lemma H4.420 (Closure filtering follows compensator balancing)**.** *Closure filtering is meaningful only after HC has removed representational drift between coupled sector descriptions.*

Proof. Closure tests are carried out on final configurations that are meant to represent one common admissible state. Without compensator balancing, distinct representatives may still differ by unresolved phase or sector drift, so a closure residue would not yet measure physical inadmissibility but only misalignment of representation. Therefore HC balancing is the prerequisite that turns closure filtering into a structurally meaningful test. □

Proposition H4.421 (Closure ladder)**.** *The passage from local structural data to observer-frame physics factors through the closure ladder described above, and none of its intermediate steps may be skipped without breaking the proof discipline*

closure before projection before interpretation.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. Lemma H4.419 places admissible triolic organization before HC balancing. Lemma H4.420 places closure filtering after HC and before any observer readout. The observer frame, however, names only the output of the stabilized configuration, so projection necessarily comes after closure filtering. Interpretation is last because it depends on projected sector naming rather than on raw structural data. Thus the stated factorization is forced by the standing proof order. □

2. Probe reduction chain and admissible channels

The probe sector now receives a reusable reduction chain. Instead of speaking only about a general search space, archival stage fixes the following sequence:

$$\begin{aligned}&\text{probe pair} \Longrightarrow \text{orthogonal search geometry}\\ &\Longrightarrow \text{final channel family} \Longrightarrow \text{tool-guided compression}\\ &\Longrightarrow \text{closure-compatible channel.}\end{aligned}$$

This makes explicit where frequency detection, routing compression, Markov transition guidance, and Newton-type convergence tools actually enter.

Lemma H4.422 (Orthogonal probe geometry restricts channel growth)**.** *When two probes are coupled together with the standing orthogonal third group, the set of admissible final channels is a strict restriction of the naive pairwise interaction space.*

Proof. The naive pairwise space allows all pair couplings consistent with the two incoming probe descriptions. The orthogonal third group imposes an additional compatibility burden that forbids precisely those channels that would resolve the pairwise mismatch by leaving the triolic discipline. Hence the admissible final family is smaller than the unrestricted pairwise space. □

Lemma H4.423 (Tool-guided compression preserves admissibility order)**.** *Kasisky-, hash-, rainbow-, Markov-, and Newton-type tools are admissible only when used to compress or navigate the final channel family without changing the structural order of admissibility testing.*

Proof. These tools may accelerate pattern detection, routing, transition guidance, or convergence, but none of them decides admissibility independently of closure. Their acceptable role is therefore subordinate: they reduce the burden of search inside the final family while the final decision remains the closure test. Consequently, they preserve admissibility only if they do not replace the closure filter by a purely computational surrogate. □

Proposition H4.424 (Probe reduction principle)**.** *A probe channel is physically readable only if it is obtained as the end point of the probe reduction chain and survives closure filtering after tool-guided compression.*

Proof. By Lemma H4.422, the orthogonal search geometry already restricts the final family. By Lemma H4.423, the technical tools may only compress or navigate that family, not redefine its criterion of admissibility. Therefore the only physically readable channels are those that remain after the final closure-compatible selection step. □

3. Projection robustness and controlled interpretation

The next recurrent burden is the distinction between structural forcing, projective readout, and phenomenological naming. Version archival stage fixes this burden by introducing small reusable robustness statements.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Lemma H4.425 (Projection does not create structure)**.** *A projection map may translate an already closure-stable structural state into an observer-frame description, but it cannot create a new admissible structure that was absent before projection.*

Proof. Projection acts on the representative of a state after the admissibility burden has already been settled. If projection could create admissibility, then the observer frame would retroactively decide structural existence, contradicting the standing principle that measurement is phase-shifted relative to the underlying structure. Thus projection is readout, not creation. □

Corollary H4.426 (Interpretation discipline)**.** *Matter, gauge, mass, Higgs, and geometric language are admissible only as controlled names for projected closure-stable regimes and not as independent axiomatic inputs to the structural proof.*

Proof. By Lemma H4.425, projection only reads out a state that has already survived the structural admissibility filter. Therefore the phenomenological names attached afterwards inherit their legitimacy from the closure-stable regime and cannot function as prior axioms of the construction. □

4. Technical use inside the monolith

The practical purpose of these new lemmas is modest but important. They create stable insertion points for later derivation chains. Subsequent versions may therefore reference:

- the *closure ladder* when passing from local triolic data to observer-frame sectors;
- the *probe reduction principle* when compressing search spaces; and
- the *projection robustness statements* when interpreting effective field sectors.

This means that the monolith can deepen individual proofs without reopening the global explanatory burden every time a local derivation is added.

5. Version status of archival stage

Relative to the previous release, archival stage contributes a stricter internal proof scaffold. The master-equation ladder and the micro-to-macro chain are retained, but they are no longer left as merely global narratives. They are now supported by named intermediate lemmas that later sections may invoke directly. That is the specific function of archival stage: not to conclude the full proof, but to make the next proof layer technically attachable in a disciplined way.

H5 Open-Question Genealogy and Phase IV Detail

The known OP5 and spectral limitations stated in the main premise audit remain binding when older status summaries are reopened.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Open Questions

Status. *Active live-burden/continuation block.*

Block function. This section separates what is still genuinely open from what survives only as historical labeling, so readers can distinguish live fronts from preserved older checkpoints.

The structural core, the master equation, and their principal projections are now explicit. The remaining work no longer concerns the existence of the framework itself, but its residual quantitative closure, phenomenological extraction, and spectrum-level completion. To avoid confusion, this section separates the *current live fronts* from the *historical OP labels* that are still used in later parts of the text.

Current reading. Several historically open OPs should no longer be read as fully open. In particular, the Koide/triolic block is already structurally closed at the present level, the Dirac-operator problem on H_{-}^{+} is no longer open as an independent operator-existence problem, and the photon-frame Θ-remainder step of the cosmological block is already closed. The live questions are therefore the following.

Remark H5.1 (Reader precedence for historical release snapshots inside the active line)**.** *The next two synchronized-status blocks reconstruct older public-stage status only for navigation and burden genealogy. They do not overrule the current burden ledger, the current theorematic status, or the current tool/QAM/decoder routing of the active line. Whenever one of those historical snapshots sounds flatter or stronger than the normalized present reading, the normalized present reading governs.*

Remark H5.2 (historical release line synchronized status reading for the optical approximation layer)**.** *The historical public archival stage release line no longer permits a flat reading in which the new optical sector is either "already a full optical theory" or "still only a loose analogy." After the first reduced prism-readout lemma, the residual-diagnostic layer, the carrier-side fit/calibration/validation chain, the reconstruction/correction steps, and the bundled optical approximation package, the optical layer now has its own staged status:*

(i) *a* strong global optical claim *remains open;*

(ii) *a* reduced retained-carrier optical sector *is already available;*

(iii) *a* first-order optical workflow *is already available;*

(iv) *a* bundled first-order optical package *is already available as an active citation surface.*

Thus the open-front layer must now distinguish between the still-open stronger global optical theory and the already stabilized first-order optical approximation layer.

Proposition H5.3 (historical release line optical/open-question status synchronization principle)**.** *At the present theorematic and retained-carrier level, the following synchronized status reading holds for the new optical line:*

(i) *the still-open strong claim is a full global optical theory of the crystal–shadow architecture, including non-reduced anisotropic branches, stronger inversion, and a later convergence theory for higher optical refinements;*

(ii) *what is already available is a reduced prism-readout sector on one retained carrier surface with ordinary wavelength ordering;*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(iii) *what is already available as first-order workflow is the chain consisting of residual diagnostics, carrier-side fit ansatz, spectral calibration, overdetermined validation, effective-index reconstruction, carrier-profile correction, and reduced stability/stopping control;*

(iv) *what is already available as a single active surface is the bundled optical approximation package of Proposition 31.7;*

(v) *therefore the optical sector should no longer be counted among the fully open foundational fronts of the document, but as a partially stabilized first-order approximation layer with a still-open stronger global continuation.*

Proof. Clause (i) records exactly the stronger optical ambitions that the current line has not yet proved: unrestricted anisotropic branching, stronger inversion, and a genuine global convergence theory are still absent. Clause (ii) is supplied by the reduced prism-readout lemma. Clause (iii) is the cumulative content of the residual spectral-diagnostic lemma, the carrier-side fit principle, the calibration principle, the overdetermined consistency principle, the effective-index reconstruction principle, the carrier-profile correction principle, and the reduced stability principle. Clause (iv) is exactly the content of the bundled optical approximation package. Hence the optical line already has a stabilized first-order approximation layer even though the stronger global optical theory remains open. □

Corollary H5.4 (Use rule for the synchronized historical release line optical status reading)**.** *Whenever a later argument in the present public archival stage release line needs to classify whether the optical sector is already available for active use, it is sufficient to cite Proposition H5.3. A finer reopening is required only when the argument depends on a stronger global optical theory beyond the reduced retained-carrier approximation layer.*

Proof. This is the operational reformulation of Proposition H5.3. The proposition already isolates the precise point at which the optical line ceases to count as merely a loose heuristic while still falling short of a full global optical theory. □

Remark H5.5 (Historical historical release line/historical release line synchronized status reading for the upgraded closure layer)**.** *The historical public archival stage release line no longer permits a flat reading in which the upgraded closure/interface blocks are simply either "open" or "proved." After the archival theorem-level shadow package, the archival interface-payoff layer, and the present historical release line partial-closure steps, their status is now genuinely four-tiered:*

(i) *a* strong deferred claim *remains open;*

(ii) *a* theorem-level shadow *is already available;*

(iii) *an* interface-level operational use *is already available;*

(iv) *for certain carrier-side conclusions, a* restricted partial-closure / lift-back consequence *is already available.*

Thus the open-front layer must now distinguish not only between strong and weak forms, but also between theorematic shadow, interface use, and restricted closure consequence.

Proposition H5.6 (historical release line hypothesis/open-question status synchronization principle)**.** *At the present theorematic and interface level, the following synchronized status reading holds:*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(i) *for Hypothesis 33.1, the still-open strong claim is the existence of one explicit compensator-centered generating equation* $\mathfrak{E}_{\mathrm{HC}} = 0$*, while the theorem-level shadow of a common closure-stable carrier/projection architecture is already available, already usable on the terminal interface, and now participates in the restricted lift-back layer whenever only retained-carrier conclusions are drawn;*

(ii) *for Hypothesis 33.6, the still-open strong claim is the existence of a deeper explicit compensator generator* $\mathbf{H}_C$*, while the theorem-level shadow separating visible defect vanishing from carrier extinction is already available, already usable on the terminal interface, and now participates in the no-second-carrier / restricted lift-back layer;*

(iii) *for Hypothesis 33.11, the still-open strong claim is the existence of a concrete coarse-graining map to fluid-type evolution, while the theorem-level shadow demoting fluid-like behavior to a later coarse-grained readout is already available, already usable on the terminal interface, and now participates negatively in the restricted lift-back layer by forbidding replacement of the retained carrier;*

(iv) *for Hypothesis 31.1, the still-open strong claim is a full universal geometric lensing law, while the theorem-level shadow of common-carrier/two-probe coupling is already available, already usable on the terminal interface, and now participates in the no-second-carrier / partial-closure layer;*

(v) *for the OP 3/OP 5 line, the still-open strong claim is a full global closure theorem, while the historical public archival stage release line already contained more than a weak bridge: it had a first partial closure theorem together with a restricted lift-back to retained carrier-layer structure.*

Therefore these upgraded blocks should no longer be counted among the fully open foundational fronts *of the document. Only their strong deferred forms remain fully open; their weaker theorematic, interface-level, and partial-closure forms are already part of the active line.*

Proof. Each clause follows by combining the archival theorem-level shadows, the archival interface-level application principles, and where applicable the new historical release line partial-closure consequences. For the HC meta-field generator hypothesis, this combines the common-carrier/projection shadow and its interface-level use with the present restricted lift-back layer. For frame-dependent defect readout, this combines the defect-shadow theorem, the interface-level defect-face vanishing principle, and the present no-second-carrier/lift-back layer. For fluid-type readout, this combines the fluid-shadow theorem, the interface-level fluid demotion principle, and the present lift-back restriction that forbids carrier replacement. For bidirectional probe lensing, this combines the probe-lensing shadow theorem, the interface-level probe-coupling principle, and the present no-second-carrier/partial-closure layer. Finally, the OP 3/OP 5 line is upgraded from the historical release line bridge test to the present historical release line partial closure theorem and restricted lift-back theorem. Hence that historical stage already supported a synchronized four-tier status reading. □

Corollary H5.7 (Use rule for the synchronized historical release line status reading)**.** *Whenever a later argument refers back to the historical public archival stage release line and needs to classify whether one of the upgraded closure/interface blocks was still truly open at that stage, it is sufficient to cite Proposition H5.6. A finer reopening is required only when the argument depends on the still-open strong form rather than on the already stabilized theorematic, interface-level, or restricted partial-closure form.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. This is the operational reformulation of Proposition H5.6. The proposition already isolates the precise point at which the upgraded blocks cease to count as fully open foundational fronts. □

Remark H5.8 (historical release line synchronized status reading for the upgraded hypothesis layer)**.** *The present 4.7.0 release line no longer permits a flat reading in which the upgraded hypothesis blocks are simply either "fully open" or "already proved." After the released historical release line theorem-level shadow package and the historical historical release line interface payoff layer, their status is now genuinely three-tiered:*

(i) a strong deferred claim *remains open;*

(ii) a theorem-level shadow *is already available;*

(iii) an interface-level operational use *is already available on the terminal surface.*

Thus the open-front layer must now distinguish between still-open strong forms and already stabilized weaker forms.

Proposition H5.9 (historical release line hypothesis/open-question status synchronization principle)**.** *At the present theorematic and interface level, the following synchronized status reading holds:*

(i) for Hypothesis 33.1, the still-open strong claim is the existence of one explicit compensator-centered generating equation $\mathfrak{E}_{\mathrm{HC}} = 0$*, while the theorem-level shadow of a common closure-stable carrier/projection architecture is already available and already usable on the terminal interface;*

(ii) for Hypothesis 33.6, the still-open strong claim is the existence of a deeper explicit compensator generator $\mathbf{H}_C$*, while the theorem-level shadow separating visible defect vanishing from carrier extinction is already available and already usable on the terminal interface;*

(iii) for Hypothesis 33.11, the still-open strong claim is the existence of a concrete coarse-graining map to fluid-type evolution, while the theorem-level shadow demoting fluid-like behavior to a later coarse-grained readout is already available and already usable on the terminal interface;

(iv) for Hypothesis 31.1, the still-open strong claim is a full universal geometric lensing law, while the theorem-level shadow of common-carrier/two-probe coupling is already available and already usable on the terminal interface.

Therefore these four blocks should no longer be counted among the fully open foundational fronts *of the document. Only their strong deferred forms remain open; their weaker theorematic and interface-operational forms are already part of the active line.*

Proof. Each clause follows by combining one theorem-level shadow result with its corresponding interface-level application principle. For Hypothesis 33.1, this is the theorem-level shadow of the HC meta-field generator together with the interface-level common-carrier/projection principle. For Hypothesis 33.6, this is the theorem-level shadow of frame-dependent defect readout together with the interface-level defect-face vanishing principle. For Hypothesis 33.11, this is the theorem-level shadow of fluid-type readout together with the interface-level fluid-readout demotion principle. For Hypothesis 31.1, this is the theorem-level shadow of probe-lensing together with the interface-level probe-coupling principle. Hence the four upgraded blocks now share one synchronized three-tier status reading: strong deferred form, theorem-level shadow, and interface-level operational use. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Corollary H5.10 (Use rule for the synchronized historical release line status reading)**.** *Whenever a later argument in the present 4.7.0 release line needs to classify whether one of the four upgraded hypothesis blocks is still truly open, it is sufficient to cite Proposition H5.9. A finer reopening is required only when the argument depends on the still-open strong form rather than on the already stabilized theorematic or interface-level shadow.*

Proof. This is the operational reformulation of Proposition H5.9. The proposition already isolates the precise point at which the upgraded blocks cease to count as fully open foundational fronts. □

1. **Residual quantitative completion of the frame scale (historical OP 1; also absorbing OP 2(T4) and the former OP 4 coupling).** The continuum fixed-point structure is already identified at the Hopf/Tribonacci level, and the frame-offset problem is no longer a broad undifferentiated open front. What remains open is the explicit derivation of the prime-lattice / continuum-limit correction δ_{CL} (or an equivalent arithmetic correction) that transports the continuum final to the measured matter-frame value.

2. **Phenomenology: controlled emergence.** The probe–closure field system is now written in explicit variational form and closed by the residue constraint $\mathfrak{R}_C = 0$. Structurally, the preceding sections identify Dirac-, QFT-, and Einstein-type regimes as distinguished projections of the same closure-stable equation. What remains open is the fully controlled derivation of these projections with explicit constants, approximation bounds, and quantitative phenomenology.

3. **Spectrum: mass hierarchy.** The present structural reading is already strong enough to separate the qualitative mass-sector logic from the unresolved quantitative assignment. The remaining open task is the map from prime/spectral index to Standard-Model quantum numbers and measured mass scales.

4. **Spectrum: three generations.** $S^2 \supset S^1 \supset S^0$ as three fermion generations is structurally supported, but the quantitative generation-splitting law is not yet derived.

5. **Structural completion: Dixon algebra.** A compact reformulation in $\mathbb{C}\otimes\mathbb{H}\otimes\mathbb{O}$ (64 dimensions) may unify the present results into one algebraic object, but this remains a later consolidation step rather than a current prerequisite.

6. **Λ-cancellation: reduced photon-frame residual problem (historical OP 5, Step (B) only).** The photon-frame formulation and the structural separation $\Lambda_{\mathrm{fundamental}} = 0$ versus quantitative balance are now fixed, and Step (A) of the Θ–R program is already closed in the present text. The remaining open content is Step (B): the exact Θ-sum / Θ–R balance needed to cancel the constant term at the photon-frame cutoff.

Historical OP synchronization. For the remainder of the document, the historical labels should be read with these statuses in mind:

- **Historical OP 3 (Koide / θ_K block):** closed at the present structural level; later work may still densify exposition, but it should no longer be listed as a primary open problem.
- **Historical OP 2 (Dirac operator on H_-^+):** closed as an independent operator problem in its existence, self-adjointness, and GOE-class parts; only the former frame-offset part survives, and it is absorbed into Item 1 above.
- **Historical OP 5:** reduced to Step (B) only; Step (A) is no longer open.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Phase IV-I: Physical Unfolding, Observer Frames, and Interface Regimes

Lemma H5.11 (Observer-frame lemma)**.** *Within the normalized structural architecture, an observer frame is not introduced as an external background datum. Rather, it is the stabilized readout frame selected by the combined action of the cascade, the closure filter, and the admissibility-preserving transport across the grouped layers. In particular, any observer description compatible with the kernel must arise as a frame-shifted but ground-linked reading of the same underlying body.*

Proof. The semantic kernel identifies all admissible descriptions as readings of a single underlying structure. Frame-shifted observation therefore does not create an ontologically new sector. By the forcing chain established in Phases I–III, any admissible readout is first re-identified by the compensator, then filtered into a closure-stable class, then transported through the grouped architecture without ontology drift. The observer frame is therefore fixed only at the level of admissible readout, not posited independently of the structure. □

Lemma H5.12 (Mass/spectral reading lemma)**.** *Any admissible mass or hierarchy reading in the Companion must be interpreted first as a spectral or sectoral readout of the normalized structural kernel, and only secondarily as a phenomenological mass assignment. In particular, the transition from spectral organization to mass hierarchy is a realization map, not a new generative axiom.*

Proof. The generative layer produces admissible sectors and their closure relations, while the projection/translation layer supplies readout rules and collapse-fixed identifications. Therefore any mass reading that appears in the field/probe layer inherits its legitimacy from a prior spectral or sectoral differentiation. This excludes the reverse move of inserting an independent mass ontology into the generative layer. The mass interpretation is thus subordinate to the spectral structure and must be treated as a downstream realization. □

Lemma H5.13 (Generation-splitting lemma)**.** *If the triolic and grouped-layer architecture is preserved across admissible projections and field realizations, then any generation splitting must appear as a structured decomposition of an already admissible spectral body rather than as a free multiplication of unrelated sectors.*

Proof. Triolic minimality and orthogonal anchoring prevent arbitrary proliferation of sectors at the generative level. The transport rules for $\mathcal{M}$ preserve admissibility and closure, and the no-ontology-drift lemma excludes the creation of new bodies during layer transfer. Hence any multiplicity such as a generation-like splitting must be read as an organized decomposition of the inherited spectral structure. It is therefore constrained by the prior closure and transport logic. □

Theorem H5.14 (Standard-model interface theorem)**.** *The normalized Companion architecture defines a one-way interface to Standard-Model-style phenomenology: admissible structures may be projected into observer, spectral, and field realizations compatible with particle-physics language, but the Companion introduces no independent Standard-Model ontology at the generative level. Consequently, the appropriate first target of the unfolding is not a complete reconstruction of the Standard Model, but a structurally controlled interface map from the grouped kernel into physically readable sectors.*

Proof. By the observer-frame, mass/spectral, and generation-splitting lemmas, all physically readable sectors arise only after the generative core has been passed through closure, confinement, cascade readout, tensor realization, and grouped transport. The field/probe layer

© 2026 Marc Brendecke. Preprint — not peer reviewed.

therefore supplies an interface regime in which known physical language may be attached to admissible sectors, but the kernel itself remains prior to that language. The Standard Model can thus enter only through an interface map respecting admissibility, closure, and no-ontology drift. This proves that the correct unfolding target is an interface theorem rather than a premature full identification. □

Corollary H5.15 (First physical unfolding chain). *The first physical unfolding chain of the Companion is*

$$\textit{observer-frame emergence} \Longrightarrow \textit{spectral/mass readout}$$
$$\textit{generation-compatible splitting} \Longrightarrow \textit{controlled Standard-Model interface.}$$

Thus the physical interpretation layer is admitted only after the structural, closure, tensorial, and grouped-layer chains have already been stabilized.

Phase IV-I′: Refined Physical Interface and Realization Control

Lemma H5.16 (Observer covariance under admissible transport). *If two observer descriptions are related by admissibility-preserving transport through the grouped architecture, then they represent frame-shifted realizations of the same underlying body and cannot define distinct generative ontologies.*

Proof. The no-ontology-drift result of Phase II-F implies that transport across $\mathfrak{G}$, $\mathfrak{P}$, and $\mathfrak{F}$ preserves the underlying structural identity. By the observer-frame lemma, any observer description is already a frame-shifted readout of the same admissible kernel. Therefore two observer descriptions related by admissible transport differ only at the level of readout covariance and not at the level of generative content. □

Proposition H5.17 (Spectral-to-mass realization map). *There exists a structurally controlled realization map*

$$\mathcal{R}_m : \mathcal{S}_{\mathrm{spec}} \longrightarrow \mathcal{S}_{\mathrm{mass}}$$

from admissible spectral sectors to admissible mass-readout sectors, where $\mathcal{R}_m$ is downstream of closure, cascade readout, and grouped transport.

Proof. By the mass/spectral reading lemma, any mass assignment must be subordinated to a prior spectral organization. The admissible spectral sectors are produced only after the closure-stable body has been transported through the projection/translation layer and read out through the cascade. Hence the passage to a mass language is not primitive but realized by a downstream map from already admissible spectral sectors. Denote this realization map by $\mathcal{R}_m$. Its domain is therefore restricted to $\mathcal{S}_{\mathrm{spec}}$ and its codomain inherits admissibility from the upstream structural chain. □

Proposition H5.18 (Generation-compatible decomposition). *If a generation-like multiplicity appears in the physical layer, then it must be expressible as an admissible decomposition of the image of $\mathcal{R}_m$ rather than as an unconstrained replication of primitive sectors.*

Proof. The generation-splitting lemma already excludes free multiplication of unrelated sectors. Since the physical layer is reached only after grouped transport and spectral-to-mass realization, any multiplicity that survives into the interface regime must be inherited from the already admissible image of $\mathcal{R}_m$. Thus a generation-like structure can only be accepted when it is realizable as a constrained decomposition of an upstream admissible spectral body. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Theorem H5.19 (Controlled physical interface functor). *The first physically readable regime of the Companion is determined by a controlled interface functor*

$$\mathfrak{I}_{\mathrm{phys}} : (\mathfrak{G}, \mathfrak{P}, \mathfrak{F}) \longrightarrow \mathcal{O}_{\mathrm{phys}},$$

which assigns observer, spectral, mass, and generation-level readouts to admissible structural sectors while preserving closure compatibility and ontology non-drift.

Proof. The observer covariance lemma identifies physically readable frames as transport-compatible readouts of the same body. The spectral-to-mass realization proposition supplies a downstream realization map from spectral sectors to mass sectors, while the generation-compatible decomposition proposition constrains multiplicity to inherited admissible decompositions. Collecting these assignments yields a functorial interface from the grouped structural architecture to a space of physically readable observables $\mathcal{O}_{\mathrm{phys}}$. Since every step is subordinated to admissibility, closure, and no-ontology drift, the resulting interface is controlled rather than ontologically generative. □

Corollary H5.20 (Refined physical interface chain). *The refined physical interface chain is*

$$\begin{aligned} &\textit{observer covariance} \Longrightarrow \textit{spectral-to-mass realization} \\ &\textit{generation-compatible decomposition} \Longrightarrow \textit{controlled interface functor.} \end{aligned}$$

Hence the physical language of the Companion is introduced only as a regulated readout of the normalized structural architecture.

Phase IV-J: Spectral Readout, Trace Reconstruction, and Observable Selection

Lemma H5.21 (Spectral admissibility inheritance). *If a physically readable sector is admitted by the controlled interface functor, then its spectral representative must inherit admissibility from the upstream closure-stable and transport-compatible structural body. In particular, no spectral mode may enter the readout regime unless it descends from an already admissible structural sector.*

Proof. By the semantic–tensor interface and the grouped transport chain, admissibility is preserved from the generative layer through cascade readout and into the physically readable interface regime. The controlled physical interface functor therefore cannot attach an observable meaning to a mode that was not already admitted upstream. Hence spectral readability inherits admissibility rather than creating it locally. □

Proposition H5.22 (Trace-compatible spectral decomposition). *Every admissible spectral body in the readout regime admits a trace-compatible decomposition into modes whose contribution respects the closure filter, the grouped transport constraints, and the observer-covariant interface conditions.*

Proof. The historical spectral and trace cores already isolate the need for admissible mode selection and trace-compatible decomposition. In the normalized architecture, these requirements are now downstream of closure, cascade, and grouped transport. Therefore any accepted spectral decomposition must preserve the admissible body identified by the upstream chain and cannot separate into trace-incompatible or closure-breaking components. This yields a trace-compatible mode decomposition of every admissible spectral body. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proposition H5.23 (Observable extraction map). *There exists a structurally controlled observable extraction map*

$$\mathcal{R}_{\mathrm{obs}} : \mathcal{S}_{\mathrm{spec}}^{\mathrm{adm}} \longrightarrow \mathcal{O}_{\mathrm{phys}}^{\mathrm{adm}},$$

which assigns observable readouts only to trace-compatible admissible spectral sectors.

Proof. Phase IV-I′ introduced a controlled interface functor from the grouped architecture into physically readable observables. Restricting that interface to the spectrally admissible, trace-compatible body provided by Proposition H5.22 yields a downstream extraction map from admissible spectral sectors to admissible observables. Since its domain is already filtered by closure, transport, and trace compatibility, the map does not generate new observables freely but only extracts structurally sanctioned ones. □

Theorem H5.24 (Mode-selection and observable consistency theorem). *Mode selection in the Companion is admissible only when it is simultaneously spectral, trace-compatible, closure-preserving, and observer-covariant. Under these conditions, the observable extraction map produces a consistent readout regime in which selected modes can be interpreted without violating ontology non-drift.*

Proof. Spectral admissibility inheritance excludes modes that do not descend from the admissible structural chain. Trace-compatible spectral decomposition excludes decompositions that would destroy the integrity of the admissible spectral body. The controlled interface functor and observer covariance from Phase IV-I′ ensure that the resulting readout remains a covariance of one and the same body rather than a proliferation of ontologies. Therefore mode selection is consistent exactly when all four constraints are met, and in that regime observable extraction remains structurally controlled. □

Corollary H5.25 (Spectral readout reconstruction chain). *The next spectral unfolding chain is*

$$\textit{spectral admissibility inheritance} \Longrightarrow \textit{trace-compatible decomposition}$$
$$\Longrightarrow \textit{observable extraction} \Longrightarrow \textit{consistent mode-selected readout.}$$

Thus the spectral and trace layer of the Companion is admitted only after closure, grouped transport, and controlled physical interfacing have already been stabilized.

Phase IV-K: Resolvent Gatekeeping and Operator-Controlled Spectral Access

Lemma H5.26 (Resolvent admissibility lemma). *Let $\mathcal{R}(\lambda) = (\lambda \mathbf{1} - \mathcal{A})^{-1}$ denote a formal resolvent on the admissible spectral layer. If the underlying mode family is closure-stable and trace-compatible, then resolvent access is admissible only on those channels for which the induced transport preserves the grouped-layer constraints and does not reintroduce ontology drift. In particular, admissible resolvent access is a restricted rather than global spectral operation.*

Proof. The semantic kernel already constrains admissible transport by closure preservation, grouped transport, and controlled readout. A resolvent therefore cannot be treated as an arbitrary inversion device: it is allowed only where the corresponding operator action remains inside the admissible closure class and respects the readout filter established in the preceding phases. Hence admissibility is inherited by resolvent access only on the structurally licensed channels. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proposition H5.27 (Gatekeeping proposition for admissible modes)**.** *The resolvent layer acts as a gatekeeping mechanism: admissible modes are precisely those spectral branches for which resolvent access is compatible with closure, trace reconstruction, and grouped-layer transport. Non-admissible branches are excluded before observable extraction.*

Proof. By Lemma H5.26, resolvent access is already restricted to structure-preserving channels. Combined with the trace-compatible mode selection of Phase IV-J, this yields a gatekeeping principle: only those modes that remain compatible with closure and transport can pass into the observable sector. The exclusion of non-admissible branches is therefore structural rather than ad hoc. □

Theorem H5.28 (Operator-controlled trace access theorem)**.** *Assume that the spectral sector is admissible, the selected modes are resolvent-gated, and the trace reconstruction remains compatible with the semantic–tensor interface. Then trace access is operator-controlled: every observable trace contribution arises through an admissible operator channel and not through uncontrolled spectral leakage.*

Proof. The preceding proposition ensures that only admissible modes are available for readout. The semantic–tensor interface then identifies the operator realization of the same admissible content, while closure preservation prevents leakage into non-licensed channels. Consequently, the trace contributions that survive observable extraction are exactly those transmitted through operator-controlled admissible channels. □

Corollary H5.29 (Stable observable channel corollary)**.** *Under admissible resolvent gatekeeping, the observable readout sector forms a stable channel: spectral access, trace reconstruction, and mode selection remain synchronized under the same structural admissibility constraints.*

Proof. This follows directly from Theorem H5.28: once admissible access, admissible mode selection, and admissible trace transport coincide, the resulting observable sector is stable under the same structural constraints. □

Remark H5.30 (Resolvent-to-readout chain)**.** *The operator-controlled spectral access chain may therefore be summarized as*

$$\begin{aligned} \textit{resolvent admissibility} &\Longrightarrow \textit{gatekept mode access} \\ &\Longrightarrow \textit{operator-controlled trace access} \\ &\Longrightarrow \textit{stable observable channel.} \end{aligned}$$

This phase extends the preceding spectral readout layer by showing that observable extraction is not only spectrally filtered but also operatorically regulated.

Phase IV-L: Semigroup Stability and Lyapunov-Controlled Readout Persistence

Lemma H5.31 (Semigroup admissibility lemma)**.** *Let $(T_t)_{t\geq 0}$ denote a formal semigroup acting on the admissible readout sector. If the initial spectral channel is closure-stable, resolvent-gated, and trace-compatible, then admissible semigroup evolution is restricted to those trajectories for which grouped transport, closure preservation, and observable compatibility remain jointly valid for all admissible times.*

Proof. The preceding phases identify the observable channel only after admissibility, gatekeeping, and operator-controlled trace access have been enforced. A semigroup acting on this

© 2026 Marc Brendecke. Preprint — not peer reviewed.

sector may therefore propagate states only if the induced time evolution respects the same admissibility filter. Otherwise the evolution would exit the closure-preserving and trace-compatible regime and would no longer represent a structurally licensed readout trajectory. Hence admissible semigroup evolution is constrained to the channels that preserve the previously established structural conditions for all admissible times. □

Proposition H5.32 (Lyapunov gate proposition)**.** *Assume that an admissible Lyapunov functional $\mathcal{L}$ exists on the gated readout sector. If $\mathcal{L}$ is nonincreasing along the admissible semigroup trajectories, then unstable leakage into non-admissible channels is structurally suppressed, and the gated observable channel remains dynamically protected.*

Proof. A nonincreasing Lyapunov functional excludes the spontaneous amplification of perturbations that would carry the evolution outside the admissible channel. Since admissibility already forbids ontology drift, closure violation, and non-gated spectral leakage, Lyapunov monotonicity upgrades these static constraints to a dynamical protection principle. The readout channel is therefore not merely selected once but remains protected under admissible evolution. □

Theorem H5.33 (Persistent observable channel theorem)**.** *Suppose that the readout sector is resolvent-gated, operator-controlled, and semigroup-admissible, and that a Lyapunov functional controls admissible perturbations. Then the observable channel is persistent: admissible mode selection, trace reconstruction, and grouped transport remain synchronized along the admissible evolution, and the readout sector stays dynamically stable against structural leakage.*

Proof. By Lemma H5.31, admissible semigroup evolution preserves the structurally licensed readout sector. Proposition H5.32 then prevents the admissible trajectories from developing unstable excursions into non-gated channels. Combined with the operator-controlled access theorem and the stable observable channel corollary from Phase IV-K, this yields persistence of the observable channel: the same admissibility constraints that select the readout sector also stabilize it under admissible evolution. □

Corollary H5.34 (Long-time readout stability corollary)**.** *Under admissible semigroup evolution and Lyapunov control, the observable readout channel remains stable at long admissible times; the selected spectral and trace-compatible sector is therefore not merely accessible but persistently readable.*

Proof. This follows directly from Theorem H5.33: persistence of the admissible channel implies long-time stability of the same gated and closure-compatible readout sector. □

Remark H5.35 (Semigroup-to-stability chain)**.** *The late-stage stability chain may therefore be summarized as*

$$\begin{aligned} \textit{semigroup admissibility} &\Longrightarrow \textit{Lyapunov-controlled gated evolution} \\ &\Longrightarrow \textit{persistent observable channel} \\ &\Longrightarrow \textit{long-time readout stability.} \end{aligned}$$

This phase completes the operatoric readout picture by showing that admissible channels are not only spectrally and resolvent-gated, but also dynamically stabilized over admissible evolution scales.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Phase IV-M: Channel Fusion and Multi-Readout Consistency

Lemma H5.36 (Admissible channel fusion lemma)**.** *Suppose that two or more observable channels are individually admissible, resolvent-gated, and semigroup-stable. Then a fused readout may be admitted only if the fusion procedure preserves the common closure filter, the grouped transport discipline, and the no-ontology-drift condition. In particular, fusion is admissible only as a compatibility-preserving synthesis of already licensed channels.*

Proof. The previous phases guarantee admissibility, operatoric access, and dynamical persistence for each channel separately. A fusion mechanism cannot be treated as an unrestricted superposition principle, since otherwise it could reintroduce hidden drift between channels that were only separately controlled. Therefore fusion is admissible only when the same closure, transport, and ontology constraints remain valid for the synthesized channel. In that case fusion does not create a new structural body, but re-expresses a common admissible body through multiple synchronized readouts. □

Proposition H5.37 (Cross-channel consistency proposition)**.** *If admissible channels are fused through a compatibility-preserving synthesis, then their spectral, trace-based, and long-time readout contents must agree on the shared admissible sector. Any discrepancy beyond the admitted transport freedom signals a non-admissible fusion rather than a genuine multi-readout effect.*

Proof. Each admissible channel already carries its observable content through the same semantic–tensor interface, the same resolvent gatekeeping, and the same semigroup-stable evolution class. Hence a compatibility-preserving fusion may only combine channels whose readout contents coincide on the overlap of their admissible sectors. If an apparent fusion produces incompatible observable contents beyond the allowed transport freedom, then the synthesis has violated the common admissibility constraints and is therefore not structurally licensed. □

Theorem H5.38 (Multi-readout consistency theorem)**.** *Under admissible channel fusion, the Companion admits a controlled multi-readout regime: distinct readout channels may coexist, but only as synchronized presentations of one and the same admissible structural body. Consequently, multi-readout consistency is equivalent to fusion without ontology drift.*

Proof. By Lemma H5.36, channel fusion is allowed only as a synthesis that preserves closure, grouped transport, and non-drift. Proposition H5.37 then forces agreement of spectral, trace, and persistent readout content on the common admissible sector. Therefore multiple channels can coexist only as mutually compatible renderings of the same admissible body. The controlled multi-readout regime is thus precisely the regime in which fusion introduces no ontology drift. □

Corollary H5.39 (Fusion-to-consistency chain)**.** *The late-stage fusion chain may therefore be summarized as*

$$\begin{aligned} \textit{admissible channel fusion} &\Longrightarrow \textit{cross-channel consistency} \\ &\Longrightarrow \textit{controlled multi-readout regime} \\ &\Longrightarrow \textit{fusion without ontology drift.} \end{aligned}$$

This phase closes the late readout unfolding by showing that stable observable channels may be combined only through a synthesis that preserves the unique admissible structural body across all readout presentations.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Phase IV-N: Unified Observable Readout Synthesis

Proposition H5.40 (Unified observable readout synthesis)**.** *Let the late readout regime be governed by the spectral admissibility, resolvent gatekeeping, semigroup persistence, and admissible channel-fusion principles established in Phases IV-J through IV-M. Then the resulting observable sector admits a unified synthesis map in which spectral access, operator control, long-time persistence, and multi-readout consistency are read as mutually compatible aspects of one and the same admissible observable architecture.*

Proof. Phase IV-J supplies the spectral and trace-compatible decomposition of admissible modes. Phase IV-K restricts this decomposition to resolvent-accessible channels only, so that observable extraction is not performed on arbitrary modes but only on structurally permitted ones. Phase IV-L then upgrades admissible access to persistence under semigroup evolution and Lyapunov control, which turns transient access into stable readout inheritance over time. Finally, Phase IV-M shows that multiple such persistent channels can be fused without inducing cross-channel inconsistency or ontological drift. Therefore the late readout regime is not a mere juxtaposition of separate spectral, operator, dynamical, and fusion statements; it forms a single admissible synthesis in which observable extraction is structurally unified. □

Corollary H5.41 (Late readout master chain)**.** *The late readout architecture admits the compressed chain*

$$\begin{aligned} &\textit{spectral admissibility} \Longrightarrow \textit{resolvent-controlled access} \\ &\Longrightarrow \textit{semigroup/Lyapunov persistence} \Longrightarrow \textit{multi-readout consistency} \\ &\Longrightarrow \textit{unified observable synthesis.} \end{aligned}$$

Remark H5.42 (Interpretive role of the late synthesis)**.** *This synthesis does not introduce a new ontology behind the observable layer. Rather, it states that the different late readout mechanisms are to be interpreted as coordinated access constraints and persistence conditions on one admissible observable sector generated by the normalized structural core.*

Phase IV-O: Return Bridge to the Generative Core and the Structural Master Equation

Proposition H5.43 (Return bridge from unified readout to the generative core)**.** *The unified observable readout synthesized in Phase IV-N introduces no new generative assumption. Rather, it is a late readout layer of the same generative structure determined by the one-body condition, triolic stabilization, orthogonal anchoring, frame-shifted observation, and the HC–closure–confinement chain.*

Proof. The earlier unfolding phases establish that admissible readout channels arise only after re-identification, compensator forcing, closure stabilization, and higher confinement. Phases IV-J through IV-N then refine how admissible observable content is accessed, filtered, stabilized, fused, and synthesized. None of those later steps introduces a new generating source; they remain constrained by the same admissibility conditions inherited from the generative layer. Hence the late observable synthesis is a return readout of the original structure rather than an independent origin. □

Theorem H5.44 (Master-equation readout closure)**.** *Let the structural master equation be the compressed dynamical expression of the admissible generative core. Then the late readout*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

chain

$$\begin{aligned} &\textit{spectral admissibility} \Longrightarrow \textit{resolvent-controlled access} \\ &\Longrightarrow \textit{semigroup/Lyapunov persistence} \Longrightarrow \textit{multi-readout consistency} \\ &\Longrightarrow \textit{unified observable synthesis} \end{aligned}$$

closes back into the same master-equation-governed structure, in the sense that every admissible late observable synthesis remains a governed readout of the original master-equation sector.

Proof. By the semantic/tensor interface and grouped transport phases, admissibility is preserved under the Millennium transport between the generative, projection, and realization layers. The readout phases IV-J through IV-N add only progressively refined access conditions: spectral admissibility, resolvent gatekeeping, semigroup persistence, and fusion consistency. Since each step is formulated as structure-preserving and closure-compatible, the final observable synthesis remains within the same admissible sector selected by the master equation. Therefore the readout chain closes into the same governing structure and should be interpreted as its late observable face. □

Remark H5.45 (Interpretive closure of the unfolding program)**.** *The point of the return bridge is to prevent the late readout machinery from being mistaken for a detached auxiliary theory or a second source. The late phases do not compete with the generative core; they complete its observable unfolding. In this sense the overall program closes as*

$$\begin{aligned} &\textit{generative core} \Longrightarrow \textit{HC/closure/confinement} \Longrightarrow \textit{master equation} \\ &\Longrightarrow \textit{spectral-resolvent-semigroup-fusion synthesis} \\ &\Longrightarrow \textit{return readout of the same structure.} \end{aligned}$$

Phase IV-P: Observable Back-Projection and Kernel Reconstruction

Lemma H5.46 (Admissible back-projection lemma)**.** *If a late observable synthesis is admissible in the sense of Phases IV-J through IV-O, then it admits a back-projection into the grouped architecture such that the reconstructed object remains compatible with the same closure, transport, and non-drift constraints that governed the forward readout.*

Proof. By Phase IV-N, the unified observable synthesis is not an arbitrary superposition of readout data, but a compatibility-preserving synthesis of spectral, resolvent, semigroup, and fusion constraints. Phase IV-O then shows that this synthesis remains a governed readout of the same master-equation sector. Hence any admissible late synthesis can be projected back through the grouped architecture without leaving the admissible class: what is reconstructed is not a new source, but the same structural body viewed through the inverse admissibility constraints of the readout chain. □

Proposition H5.47 (Kernel reconstruction proposition)**.** *Admissible back-projection reconstructs not a second kernel, but a readout-identified representative of the original generative kernel. In particular, reconstruction is valid only up to the transport, projection, and realization equivalences already built into* $\mathfrak{G}$*,* $\mathfrak{P}$*, and* $\mathfrak{F}$*.*

Proof. The grouped transport architecture never licenses ontology drift; it only transports one admissible body across grouped layers. Therefore any inverse passage from late observable synthesis to kernel-level description must be taken modulo the same transport equivalences that governed the forward route. The back-projected object is thus a kernel representative recovered from admissible readout data, not an independent second kernel standing alongside the original generative source. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Theorem H5.48 (Observable-to-kernel closure theorem)**.** *The late unfolding program closes in both directions: every admissible generative structure yields a governed late observable synthesis, and every admissible late observable synthesis yields a back-projected kernel representative of that same structure. Consequently, the Companion admits a controlled observable-to-kernel closure rather than a one-way readout only.*

Proof. The forward direction is the content of the unfolding chain from the generative core through the master equation to the unified observable synthesis. By Lemma H5.46, an admissible late synthesis can be projected back into the grouped architecture while preserving closure, transport, and non-drift. Proposition H5.47 then shows that the reconstructed object is to be read as a representative of the original generative kernel, not as a second ontological origin. Thus the unfolding closes in both directions: generation governs readout, and admissible readout reconstructs the governed kernel representative. □

Corollary H5.49 (Bidirectional closure chain)**.** *The late-stage closure may therefore be compressed as*

$$\begin{aligned}&\textit{generative core} \Longrightarrow \textit{master-equation-governed readout} \Longrightarrow \textit{unified observable synthesis}\\ &\Longrightarrow \textit{admissible back-projection} \Longrightarrow \textit{kernel representative of the same structure.}\end{aligned}$$

Phase IV-Q: Global Unfolding Synthesis and Architectural Compression

The preceding phases may now be compressed into a single Companion-level statement: the full unfolding chain from the generative kernel to observable back-projection is not a juxtaposition of independent modules, but a single admissibility-controlled architecture with forward, lateral, and return transport. The role of the late synthesis is therefore not to append a second theory or a new axiom block to the generative core, but to expose the full reversible structural path from formation to readout and back.

Proposition H5.50 (Global unfolding synthesis)**.** *The Companion unfolding program may be read as one admissibility-preserving architecture*

$$\begin{aligned}\textit{generative formation} &\Longrightarrow \textit{HC/closure/confinement forcing}\\ &\Longrightarrow \textit{cascade and readout admissibility}\\ &\Longrightarrow \textit{semantic/tensor transport}\\ &\Longrightarrow \textit{grouped-layer realization}\\ &\Longrightarrow \textit{probe and field realization}\\ &\Longrightarrow \textit{physical interface control}\\ &\Longrightarrow \textit{spectral/resolvent/semigroup/fusion synthesis}\\ &\Longrightarrow \textit{observable back-projection}\\ &\Longrightarrow \textit{kernel reconstruction of the same structure.}\end{aligned}$$

In particular, the later readout layers do not introduce a second ontological source, but provide a controlled return map into the same structural kernel whose forcing chain generated the admissible dynamics in the first place.

Proof. Phase I establishes the generative forcing chain from one-body discipline to HC, closure, and higher confinement. Phase II transports this material through cascade-admissible readout, semantic/tensor realization, and grouped-layer architecture without ontological drift.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Phase III realizes the same admissible structure in probe and field form, culminating in the structural master equation and its exact-closure regime. Phase IV then unfolds the physical interface, spectral access, resolvent control, semigroup stability, fusion consistency, and unified readout synthesis, before Phase IV-O and Phase IV-P return the resulting observables to the master-equation-governed kernel and reconstruct a representative of the same admissible structure. The complete chain is therefore structurally circular only in the admissible sense of closure and readout return, not in the sense of a second independent origin. □

Corollary H5.51 (Companion architectural compression rule)**.** *Any later compression of the Companion is admissible only if it preserves explicit access to each of the following four structural roles:*

$$\textit{formation}, \qquad \textit{transport}, \qquad \textit{realization/readout}, \qquad \textit{return reconstruction}.$$

A compressed presentation that suppresses one of these roles may remain heuristically useful, but it no longer represents the full unfolding architecture proved in the Companion.

Interpretive role of the global synthesis. This proposition gives the document a single citeable architectural apex: the Companion does not merely collect historical cores, transport layers, field realizations, and readout mechanisms, but exhibits them as one normalized unfolding from generative structure to controlled observability and admissible back-return.

Phase IV-R: Proof Compression and Structural Admissibility Testing

The global synthesis may now be turned into a practical proof-saving rule. Once the full unfolding architecture has been exhibited, later arguments need not re-derive every intermediate transport, readout, and return step from scratch. Instead, it becomes sufficient to verify that a proposed construction remains inside the admissible chain already established by Phases I through IV-Q.

Lemma H5.52 (Structural admissibility test)**.** *Let X be a proposed extension, readout, or reformulation of the Companion architecture. If X preserves the four structural roles of formation, transport, realization/readout, and return reconstruction, and if X introduces no ontology drift relative to the grouped architecture, then X may be tested against the unfolding program by admissibility checking rather than by rebuilding the entire derivation chain from the beginning.*

Proof. By Corollary [Companion architectural compression rule], any admissible compression or reformulation must preserve access to the four defining structural roles. Since Phase II-F already excludes ontology drift across grouped layers, and Phase IV-P returns admissible readout data to a kernel representative of the same structure, a later proposal that respects these constraints remains inside the already proved architecture. The burden of proof is therefore reduced from full regeneration to admissibility testing: one checks whether the proposal preserves the established chain instead of reconstructing the whole chain anew. □

Proposition H5.53 (Proof-compression proposition)**.** *The Companion permits controlled proof compression: whenever a later statement is structurally subordinate to an already proved admissible chain, it may be justified by citing preservation of that chain together with the specific additional admissibility condition that differentiates the later statement.*

Proof. Phases I through IV-Q provide the explicit full chain from formation to observable back-return. If a later statement does not alter this chain, but only specializes one of its

© 2026 Marc Brendecke. Preprint — not peer reviewed.

stages—for example by restricting admissible spectral modes, selecting a probe sector, or refining an operator gate—then the full derivation need not be repeated. It is enough to show that the specialization preserves the inherited admissibility constraints and adds only the stated extra condition. The proof is therefore compressed, but not weakened: the omitted steps are inherited from the already established chain. □

Theorem H5.54 (Meta-theorem of admissible structural compression)**.** *Any later expansion of the Companion is mathematically legitimate in compressed form if and only if it can be located inside the established unfolding architecture without breaking formation, transport, realization/readout, or return reconstruction. In particular, a compressed argument is valid exactly when it is a structurally admissible sub-chain of the global unfolding synthesis.*

Proof. Necessity follows from the architectural compression rule: if one of the four structural roles is broken, the proposal no longer belongs to the full unfolding architecture and cannot inherit its proof force. Sufficiency follows from Lemma H5.52 and Proposition H5.53: once a proposal is shown to preserve the admissible chain and differs only by an explicit additional admissibility condition, the rest of the proof is inherited from the already established unfolding program. Thus admissible compression is neither heuristic abbreviation nor informal shortcut, but a precise meta-rule derived from the structure of the Companion itself. □

Corollary H5.55 (Late meta-compression chain)**.** *The full late-stage architecture may therefore be compressed as*

$$\begin{aligned} &\textit{global unfolding synthesis} \Longrightarrow \textit{admissibility testing of later proposals} \\ &\Longrightarrow \textit{controlled proof compression} \Longrightarrow \textit{valid structural sub-chain expansions.} \end{aligned}$$

Interpretive role of the meta-compression theorem. This final theorem turns the Companion from a one-off expansion into a reusable proof engine. Later developments no longer need to restart from the beginning whenever they remain inside the admissible architecture; they need only show where they sit in the chain and which extra admissibility condition they impose.

Phase IV-S: Hopf Fixed-Point Geometry and CKM/PMNS Residuals

Remark H5.56 (Transition from Phase IV-R to the Hopf/CKM block)**.** *Phase IV-R turned the late synthesis into a reusable admissibility and proof-compression rule. The present phase uses that rule in a more specialized way: instead of introducing a new generative source, it tests whether the residual Hopf/CKM discrepancies can be read as admissible geometric corrections inside the already normalized operatoric readout architecture.*

H6 Coupled Local OP-Entry and Repair Protocol

Status (v4.9.0 revision): the resolution claimed here is withdrawn (Corollary 41.23). The continuum-limit correction line (OP 4) measures the distance between the matter value and the Tribonacci and Hopf values, which come from other frames or classes; in this form OP 4 is ill-posed, and only OP 1 remains (v4.9.0-r61).

Theorem H6.1 (First machine-fused coupled OP-entry theorem)**.** *Assume:*

1. *the corrected prime-lattice reading applies on a transfer-consistent closure-phase window;*

2. *the first proxy-to-residual transfer theorem applies, so that the active inherited burden is read through*
$$\big(\Delta_{\mathrm{CL}}, \Delta_{\Theta R}(T)\big)$$
with ρ_A as admissibility monitor;

3. *the coupled prime-lattice correction datum*
$$\mathfrak{C}^{\mathrm{cpl}}_{\Gamma_P} = \big(\Delta_{\mathrm{CL}}, \Delta_{\Theta R}(T), \rho_A\big)$$
is QAM-encodable and carried by a machine-reachable state cell;

4. *coupled prime-lattice descent is realizable in the QAM/MML/MMA layer as a finite symbolic descent/test/branch task;*

5. *the closure-phase trigger slice is machine-testable by finitely checkable burden and admissibility conditions.*

Then the first post-archival stage OP-entry problem admits a machine-fused coupled realization.

More precisely:

1. *the operative entry object is the coupled datum*
$$\big(\Delta_{\mathrm{CL}}, \Delta_{\Theta R}(T), \rho_A\big),$$
not the scalar shadow δ_{CL} alone;

2. *every admissible local prime-lattice descent step can be read simultaneously*
   (a) *geometrically, as a coupled closure-phase descent step, and*
   (b) *computationally, as a finite realizable machine step on the encoded state object;*

3. *the closure-phase trigger slice is branch-decidable in the machine layer;*

4. *therefore, whenever a finite admissible descent chain reaches the closure-phase trigger slice while preserving admissibility of ρ_A, immediate OP-entry follows.*

Proof. By the corrected prime-lattice reading, the scalar continuum-limit correction is only the closure-side shadow of the coupled datum
$$\mathfrak{C}^{\mathrm{cpl}}_{\Gamma_P} = \big(\Delta_{\mathrm{CL}}, \Delta_{\Theta R}(T), \rho_A\big).$$
By the proxy-to-residual transfer theorem, the active inherited burden is reduced to the pair
$$\big(\Delta_{\mathrm{CL}}, \Delta_{\Theta R}(T)\big)$$
under admissible ρ_A. By QAM-encodability, this coupled datum is a machine-reachable state object. By finite realizability of coupled prime-lattice descent, each admissible local descent is executable as a finite symbolic task in the QAM/MML/MMA layer. By machine-testability of the closure-phase trigger slice, the machine can decide whether the updated coupled datum has reached entry, must continue descent, or has become inadmissible. Hence the first post-archival stage OP-entry problem is machine-fused: it is simultaneously a coupled geometric descent problem and a finite machine-controlled descent/test/branch problem. Whenever the finite admissible descent chain reaches the closure-phase trigger slice while preserving ρ_A admissibility, immediate OP-entry follows by the prime-lattice coupled entry lemma. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Corollary H6.2 (First post-archival stage freeze threshold)**.** *The first machine-fused coupled OP-entry theorem marks the first natural mathematical freeze threshold for a post-archival stage minor release: the line now carries a single fused claim linking the corrected prime-lattice entry geometry with the QAM/MML/MMA machine layer.*

Proof. Immediate from Theorem H6.1. □

Definition H6.3 (Machine-fused local entry state)**.** *A* machine-fused local entry state *is a tuple*

$$\mathfrak{E}_{\mathrm{loc}} = \big(\mathbf{X}_{\Gamma_P}, \mathcal{C}_{\mathrm{mach}}, \mathfrak{D}_{\mathrm{lead}}, \mathfrak{S}_{\mathrm{trig}}\big),$$

where:

1. $\mathbf{X}_{\Gamma_P}$ *is the current prime-lattice machine object*

$$\mathbf{X}_{\Gamma_P} = \mathrm{Enc}_{\mathrm{QAM}}\big(\Delta_{\mathrm{CL}}, \Delta_{\Theta R}(T), \rho_A\big);$$

2. $\mathcal{C}_{\mathrm{mach}}$ *is the current finite machine configuration carrying* $\mathbf{X}_{\Gamma_P}$*;*
3. $\mathfrak{D}_{\mathrm{lead}}$ *is the current lead-direction tag;*
4. $\mathfrak{S}_{\mathrm{trig}}$ *is the currently active trigger slice final.*

Definition H6.4 (Local machine-fused descent/test/branch operator)**.** *The* local machine-fused descent/test/branch operator *is the composite step*

$$\mathcal{O}_{\mathrm{DTB}} = \mathcal{B}_{\Gamma_P} \circ \mathcal{T}_{\Gamma_P} \circ \mathcal{D}_{\Gamma_P},$$

where:

1. $\mathcal{D}_{\Gamma_P}$ *is a coupled prime-lattice descent realization;*
2. $\mathcal{T}_{\Gamma_P}$ *is the prime-lattice trigger test;*
3. $\mathcal{B}_{\Gamma_P}$ *is the prime-lattice branch rule.*

Remark H6.5 (Operational reading)**.** *This is the first explicit post-archival stage local controller: first lower the coupled burden pair if possible, then test the trigger slice, then branch into entry, continuation, or obstruction.*

Definition H6.6 (Local obstruction certificate)**.** *A* local obstruction certificate *is a finite machine-readable object*

$$\mathfrak{O}_{\mathrm{loc}}$$

issued when the local machine-fused descent/test/branch operator returns

$$\texttt{inadmissible} \quad \textit{or} \quad \texttt{stalled},$$

together with a finite explanation of which of the following failed:

1. *closure descent,*
2. *Theta descent,*
3. *admissibility of* ρ_A*,*
4. *persistence of the closure-phase trigger slice,*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

5. *route-faithful machine realizability.*

Definition H6.7 (Local continuation certificate)**.** *A* local continuation certificate *is a finite machine-readable object*

$$\mathfrak{K}_{\mathrm{cont}}$$

issued when the local operator returns

$$\texttt{continue},$$

certifying that:

1. *the coupled descent remains admissible;*
2. *the current state is still outside immediate OP-entry;*
3. *at least one monotone local burden measure has strictly improved.*

Definition H6.8 (Local entry certificate)**.** *A* local entry certificate *is a finite machine-readable object*

$$\mathfrak{K}_{\mathrm{entry}}$$

issued when the local operator returns

$$\texttt{entry},$$

certifying that the current machine-fused prime-lattice datum has entered the active closure-phase trigger slice and is immediately OP-entry certified.

Definition H6.9 (Machine-fused local entry protocol)**.** *The* machine-fused local entry protocol *is the iterative rule:*

$$\mathfrak{E}_{\mathrm{loc},0} \longmapsto \mathfrak{E}_{\mathrm{loc},1} \longmapsto \mathfrak{E}_{\mathrm{loc},2} \longmapsto \cdots$$

obtained by repeated application of the local descent/test/branch operator, stopping when either

$$\mathfrak{K}_{\mathrm{entry}} \quad \textit{or} \quad \mathfrak{O}_{\mathrm{loc}}$$

is issued.

Remark H6.10 (Why this is the right next step)**.** *The post-archival stage line should no longer remain at the level of existence claims alone. It should now either produce:*

1. *a local entry certificate, or*
2. *a local obstruction certificate.*

That is the first genuinely operational OP-entry phase.

Definition H6.11 (Local burden descent norm)**.** *A* local burden descent norm *is any monotone function*

$$\mathcal{N}_{\mathrm{loc}}\big(\Delta_{\mathrm{CL}}, \Delta_{\Theta R}(T), \rho_A\big)$$

such that:

1. *it decreases when both transferred burdens decrease and admissibility is preserved;*
2. *it is defined only on admissible machine-fused local entry states;*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

3. *lower values indicate closer approach to immediate OP-entry.*

Proposition H6.12 (Continuation implies local progress)**.** *Assume the machine-fused local entry protocol returns a continuation certificate*

$$\mathfrak{K}_{\mathrm{cont}}$$

at some step. Then there exists a local burden descent norm

$$\mathcal{N}_{\mathrm{loc}}$$

for which the updated local entry state has strictly smaller value than the previous one.

Proof. By definition of the continuation certificate, coupled descent remains admissible, immediate OP-entry has not yet been reached, and at least one monotone local burden measure has strictly improved. Such a measure may be taken as a local burden descent norm, yielding strict local progress. □

Lemma H6.13 (Finite local entry-or-obstruction lemma)**.** *Assume:*

1. *the machine-fused local entry protocol is defined on the current local state;*
2. *every continuation step strictly lowers a local burden descent norm;*
3. *the protocol cannot descend indefinitely inside the same admissible slice without either entering the trigger slice or violating admissibility.*

Then a finite local protocol run ends with either:

1. *a local entry certificate, or*
2. *a local obstruction certificate.*

Proof. Each continuation step strictly lowers the local burden descent norm. By assumption, indefinite descent in the same admissible slice is impossible. Hence the protocol must terminate after finitely many steps, either by entering the trigger slice or by losing admissibility/persistence and issuing an obstruction certificate. □

Theorem H6.14 (First machine-fused local OP-entry protocol theorem)**.** *Assume:*

1. *the corrected prime-lattice reading applies;*
2. *the first machine-fused coupled OP-entry theorem applies;*
3. *the closure-phase trigger slice is machine-testable;*
4. *the current line admits a machine-fused local entry state.*

Then the post-archival stage line supports a finite local machine-fused OP-entry protocol.

More precisely, every admissible run of the local protocol yields after finitely many steps either:

1. *a local entry certificate witnessing immediate OP-entry, or*
2. *a local obstruction certificate isolating the first local failure mode.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. By the first machine-fused coupled OP-entry theorem, coupled descent, trigger testing, and branch handling are already fused in the current line. Hence the local descent/test/branch operator is defined on the current machine-fused local entry state. By the finite local entry-or-obstruction lemma, every admissible local run terminates after finitely many steps with either entry or obstruction. □

Corollary H6.15 (Post-archival stage working rule)**.** *The next working line after archival stage should no longer ask only whether the coupled OP-entry theorem exists abstractly. It should explicitly run the local machine-fused entry protocol and record either:*

1. *the first local entry certificate, or*
2. *the first local obstruction certificate.*

Proof. Immediate from Theorem H6.14. □

Definition H6.16 (Default first local outcome hypothesis)**.** *The* default first local outcome hypothesis *for the post-archival stage line is the hypothesis that the first honest terminal output of the machine-fused local entry protocol is a local obstruction certificate rather than a local entry certificate.*

Remark H6.17 (Why obstruction is the current default)**.** *At the present stage, the coupled descent mechanism is already structurally available, but the active closure-phase trigger slice is still only symbolically specified and not yet fully calibrated by explicit local threshold data. Therefore the most likely first terminal failure is not loss of the whole machine-fused architecture, but failure to certify persistence or actual reach of the trigger slice before immediate entry is declared.*

Definition H6.18 (Trigger-persistence obstruction)**.** *A* trigger-persistence obstruction *is a local obstruction certificate*

$$\mathfrak{O}_{\mathrm{loc}}^{\mathrm{trig}}$$

issued when:

1. *coupled descent remains machine-realizable;*
2. *the packet-side monitor remains admissible;*
3. *but persistence or actual reach of the closure-phase trigger slice cannot yet be certified on the current local window.*

Definition H6.19 (Entry-prior versus obstruction-prior local regime)**.** *A machine-fused local entry state is called:*

1. entry-prior *if the active closure-phase trigger slice is sufficiently explicit and stable that the next finite admissible local run is expected to end in a local entry certificate;*
2. obstruction-prior *if coupled descent is available but the trigger slice remains insufficiently explicit, insufficiently persistent, or insufficiently calibrated for immediate entry certification.*

Proposition H6.20 (The current local state is more naturally obstruction-prior than entry-prior)**.** *Assume the post-archival stage line already supports:*

1. *corrected prime-lattice reading;*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

2. *proxy-to-residual transfer;*

3. *machine-fused descent/test/branch control.*

Assume further that the active closure-phase trigger slice is still only symbolically described and not yet equipped with fully explicit local threshold calibration. Then the current machine-fused local entry state is more naturally obstruction-prior than entry-prior.

Proof. Under the stated assumptions, coupled descent and testing already exist as local machine operations, so the current line is not blocked at the level of basic realizability. However, if the closure-phase trigger slice is still only symbolically described and lacks fully explicit local threshold calibration, then immediate entry cannot yet be certified purely from the present data. Hence the more natural first terminal local outcome is an obstruction certificate isolating trigger-persistence or trigger-calibration failure rather than an entry certificate. □

Theorem H6.21 (First local obstruction-priority theorem)**.** *Assume:*

1. *the first machine-fused local OP-entry protocol theorem applies;*

2. *the current line admits machine-fused coupled descent steps;*

3. *the packet-side monitor remains admissible on the first relevant local window;*

4. *the closure-phase trigger slice is not yet fully explicit or fully calibrated on that same window.*

Then the current local machine-fused state is obstruction-prior. More precisely, the first honest terminal local outcome is expected to be a trigger-persistence obstruction

$$\mathfrak{O}_{\mathrm{loc}}^{\mathrm{trig}},$$

rather than an immediate local entry certificate.

Proof. By assumption, the local machine-fused protocol exists and coupled descent is available. By admissibility of the packet-side monitor, the protocol is not forced to fail immediately by packet instability. If, however, the closure-phase trigger slice is not yet fully explicit or calibrated, then the protocol cannot honestly certify immediate entry on the present local window. The most natural terminal outcome is therefore a local obstruction certificate isolating the missing trigger-persistence or trigger-calibration ingredient. □

Corollary H6.22 (First post-archival stage local audit rule)**.** *The next local audit after archival stage should not first ask whether immediate entry already occurs. It should first ask which trigger-persistence or trigger-calibration ingredient is missing from the current local window, i.e. it should aim first at the first trigger-persistence obstruction certificate.*

Proof. Immediate from Theorem H6.21. □

Definition H6.23 (Trigger-calibration obstruction)**.** *A* trigger-calibration obstruction *is a local obstruction certificate*

$$\mathfrak{O}_{\mathrm{loc}}^{\mathrm{cal}}$$

issued when:

1. *the coupled descent/test/branch architecture is available;*

2. *the active closure-phase trigger slice is recognized symbolically;*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

3. *but no sufficiently explicit local burden thresholds or effective local trigger inequalities are yet available to certify entry.*

Definition H6.24 (Trigger-persistence obstruction — strengthened reading)**.** *A* strengthened trigger-persistence obstruction *is a local obstruction certificate*

$$\mathfrak{O}_{\mathrm{loc}}^{\mathrm{pers}}$$

issued when:

1. *the coupled descent/test/branch architecture is available;*
2. *a symbolic closure-phase trigger slice is available;*
3. *some local trigger-improving descent steps are visible;*
4. *but those steps do not yet certify persistence of the trigger slice on a sufficiently long local window.*

Definition H6.25 (Machine-side decodability obstruction)**.** *A* machine-side decodability obstruction *is a local obstruction certificate*

$$\mathfrak{O}_{\mathrm{loc}}^{\mathrm{dec}}$$

issued when:

1. *the corrected prime-lattice datum is present;*
2. *but the corresponding trigger-relevant condition cannot yet be realized as a finite machine-decidable test on the active QAM/MML/MMA state.*

Remark H6.26 (The three natural first obstruction types)**.** *At the present stage, the first local obstruction should naturally be expected to belong to exactly one of three classes:*

1. *trigger-calibration obstruction;*
2. *trigger-persistence obstruction;*
3. *machine-side decodability obstruction.*

These correspond respectively to missing local thresholds, missing stable trigger persistence, or missing test-level realizability.

Proposition H6.27 (Machine-side decodability is not the primary current obstruction)**.** *Assume:*

1. *the corrected prime-lattice reading applies;*
2. *the prime-lattice machine realization theorem applies;*
3. *the machine-fused local entry protocol theorem applies.*

Then a machine-side decodability obstruction is not the most natural first local obstruction of the current line.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. Under the stated assumptions, the coupled prime-lattice datum is already QAM-encodable, coupled descent is MML/MMA-realizable, and the trigger problem is already machine-fused at the protocol level. Hence the current line is not most naturally blocked by complete absence of machine-side realizability. The more natural remaining failure modes lie at the level of trigger calibration or trigger persistence. □

Proposition H6.28 (Current obstruction split is calibration/persistence-dominant)**.** *Assume the hypotheses of Proposition H6.27. Assume further that the active closure-phase trigger slice is already symbolically available but not yet fully quantified by explicit local thresholds and not yet stable on a sufficiently long local window. Then the first local obstruction is more naturally of calibration/persistence type than of decodability type.*

Proof. If the trigger slice is already symbolically available, then total trigger absence is not the main issue. If it is not yet fully quantified by explicit thresholds and not yet stable on a sufficiently long local window, then the missing ingredient lies in calibration and persistence. By Proposition H6.27, this is more natural than a primary machine-side decodability failure. □

Definition H6.29 (Current dominant first local obstruction class)**.** *The* current dominant first local obstruction class *is the class*

$$\mathfrak{O}^{*}_{\mathrm{loc}} \in \{\mathfrak{O}^{\mathrm{cal}}_{\mathrm{loc}}, \mathfrak{O}^{\mathrm{pers}}_{\mathrm{loc}}, \mathfrak{O}^{\mathrm{dec}}_{\mathrm{loc}}\}$$

which is most naturally expected to occur first on the current local window.

Theorem H6.30 (First local obstruction classification theorem)**.** *Assume:*

1. *the first local obstruction-priority theorem applies;*
2. *the corrected prime-lattice reading applies;*
3. *the prime-lattice machine realization theorem applies;*
4. *the closure-phase trigger slice is already symbolically available;*
5. *the trigger slice is not yet fully locally calibrated and not yet certified persistent on a sufficiently long local window.*

Then the current dominant first local obstruction class is not machine-side decodability but calibration/persistence type.

More precisely,

$$\mathfrak{O}^{*}_{\mathrm{loc}} \in \{\mathfrak{O}^{\mathrm{cal}}_{\mathrm{loc}}, \mathfrak{O}^{\mathrm{pers}}_{\mathrm{loc}}\},$$

and the current line should be read as calibration/persistence-obstruction-prior.

Proof. By the prime-lattice machine realization theorem, the coupled datum is already machine-encodable and machine-fused descent/test/branch control is available, so pure decodability failure is not the most natural first obstruction. Because the closure-phase trigger slice is already symbolically available but not yet fully calibrated or persistent on a sufficiently long local window, the remaining failure modes lie in trigger calibration and trigger persistence. Hence the current dominant obstruction class is of calibration/persistence type. □

Corollary H6.31 (Next post-archival stage local audit question)**.** *The next local audit after the current stage should first ask:*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

1. *what explicit local trigger inequalities are still missing;*
2. *and what minimum persistence window is still missing*

before treating machine-side decodability as the primary obstruction.

Proof. Immediate from Theorem H6.30. □

Definition H6.32 (Local closure-phase trigger inequality schema)**.** *A* local closure-phase trigger inequality schema *is a finite family of inequalities of the form*

$$\Delta_{\mathrm{CL}} \le \varepsilon_{\mathrm{CL}}, \qquad \Delta_{\Theta R}(T) \le \varepsilon_{\Theta}, \qquad \rho_A \in \mathfrak{A}_{\mathrm{adm}}^{\mathrm{safe}},$$

together with the side condition that the current transition remains closure-phase led on the same local window. Here

$$\varepsilon_{\mathrm{CL}} > 0, \qquad \varepsilon_{\Theta} > 0$$

are local trigger tolerances and

$$\mathfrak{A}_{\mathrm{adm}}^{\mathrm{safe}} \subseteq \mathfrak{A}_{\mathrm{adm}}$$

is a distinguished safe admissibility region for the packet-side monitor.

Remark H6.33 (Meaning of the inequality schema)**.** *The current line does not yet require a final closed-form trigger formula. At the present stage, what is missing is already captured by the weaker data of explicit local tolerances for the coupled burdens together with a safe admissibility band for* ρ_A*.*

Definition H6.34 (Local closure-phase persistence window of length m)**.** *Let* $m \ge 1$*. A* local closure-phase persistence window of length m *is a block of successive local machine-fused protocol steps*

$$\mathfrak{E}_{\mathrm{loc},n}, \mathfrak{E}_{\mathrm{loc},n+1}, \dots, \mathfrak{E}_{\mathrm{loc},n+m-1}$$

such that throughout the block:

1. *the lead direction remains closure-phase led;*
2. *the trigger inequality schema is satisfied monotonically or remains stable once entered;*
3. *the packet-side monitor stays in* $\mathfrak{A}_{\mathrm{adm}}^{\mathrm{safe}}$*;*
4. *the same route-faithful local slice remains active.*

Definition H6.35 (Minimum local trigger-persistence length)**.** *A positive integer*

$$m_{\min} \ge 1$$

is called a minimum local trigger-persistence length *if immediate OP-entry is not certified merely by entering the local trigger inequality schema once, but only after the schema persists on a closure-phase persistence window of length at least* $m_{\min}$*.*

Remark H6.36 (Why a persistence length is needed)**.** *The current obstruction analysis suggests that a one-step near-entry fluctuation is not yet trustworthy. What is missing is not just smallness of the coupled burdens, but stable smallness on a sufficiently long local window.*

Definition H6.37 (Trigger-calibration data package)**.** *The* trigger-calibration data package *is the tuple*

$$\mathfrak{P}_{\mathrm{trig}}^{\mathrm{loc}} := \left(\varepsilon_{\mathrm{CL}}, \varepsilon_{\Theta}, \mathfrak{A}_{\mathrm{adm}}^{\mathrm{safe}}, m_{\min}\right)$$

consisting of local burden tolerances, a safe admissibility region, and a minimum persistence length.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proposition H6.38 (What the current obstruction is really asking for)**.** *Assume the first local obstruction classification theorem. Then the current calibration/persistence obstruction is equivalent to the absence of a sufficiently explicit trigger-calibration data package.*

Proof. By the obstruction classification theorem, the current obstruction is not primarily one of machine-side decodability, but of calibration/persistence type. Calibration requires explicit local burden tolerances and a safe admissibility region, while persistence requires a minimum stable window length. Therefore the current obstruction is exactly the absence of a sufficiently explicit trigger-calibration data package. □

Definition H6.39 (Locally calibrated closure-phase trigger slice)**.** *A closure-phase trigger slice is called* locally calibrated *if it is equipped with a trigger-calibration data package*

$$\mathfrak{P}_{\mathrm{trig}}^{\mathrm{loc}}$$

and immediate OP-entry is certified whenever the associated local closure-phase trigger inequality schema persists for at least $m_{\min}$ consecutive local protocol steps.

Lemma H6.40 (First local calibration-to-entry lemma)**.** *Assume:*

1. *the machine-fused local entry protocol theorem applies;*
2. *a locally calibrated closure-phase trigger slice is available;*
3. *the current local protocol run enters the associated trigger inequality schema and remains there for at least $m_{\min}$ consecutive steps.*

Then the local protocol yields a local entry certificate.

Proof. By local calibration, entry is certified exactly when the trigger inequality schema persists on a closure-phase window of length at least $m_{\min}$. If the current local protocol run satisfies this condition, then by definition the corresponding state is immediately OP-entry certified and the protocol returns a local entry certificate. □

Theorem H6.41 (First local trigger calibration theorem)**.** *Assume:*

1. *the first local obstruction classification theorem applies;*
2. *the machine-fused local entry protocol theorem applies;*
3. *the active obstruction is of calibration/persistence type.*

Then the next missing local ingredient is exactly the explicit trigger-calibration data package

$$\mathfrak{P}_{\mathrm{trig}}^{\mathrm{loc}} = \big(\varepsilon_{\mathrm{CL}}, \varepsilon_{\Theta}, \mathfrak{A}_{\mathrm{adm}}^{\mathrm{safe}}, m_{\min}\big).$$

Moreover, once such a package is fixed, the remaining local question reduces to whether the current protocol run can enter and maintain the associated inequality schema for at least $m_{\min}$ consecutive steps.

Proof. By the first local obstruction classification theorem, the dominant current obstruction is not decodability but calibration/persistence type. By Proposition H6.38, this is exactly the absence of explicit local burden tolerances, a safe admissibility region, and a minimum persistence length. These data are precisely the trigger-calibration data package. Once the package is fixed, the only remaining local issue is whether the current machine-fused protocol run satisfies the corresponding persistence requirement. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Corollary H6.42 (Next post- working rule)**.** *The next working step after the current obstruction analysis should not try to prove immediate OP-entry in one jump. It should first determine:*

1. *plausible local values or bounds for* $\varepsilon_{\mathrm{CL}}$ *and* ε_{Θ}*;*
2. *a safe admissibility subregion* $\mathfrak{A}_{\mathrm{adm}}^{\mathrm{safe}}$*;*
3. *a first plausible minimum persistence length* $m_{\min}$*.*

Proof. Immediate from Theorem H6.41. □

Definition H6.43 (Default local trigger orientation)**.** *The* default local trigger orientation *is the working hypothesis that the first explicit local closure-phase trigger calibration should satisfy:*

1. *the closure tolerance is the stricter burden tolerance,*
$$\varepsilon_{\mathrm{CL}} \leq \varepsilon_{\Theta};$$
2. *the safe admissibility region for* ρ_A *is taken as a narrow stability band around the current admissible packet-side monitor regime;*
3. *the minimum persistence length is not read as one-step only, but as a genuinely multi-step window*
$$m_{\min} > 1.$$

Remark H6.44 (Why closure tolerance should be stricter first)**.** *The current obstruction analysis suggests that the coupled local burden is not blocked first by lack of Theta-side machine decodability, but by insufficiently stabilized closure-phase persistence. Hence the first local calibration should be closure-tight rather than Theta-tight.*

Definition H6.45 (First plausible default trigger package)**.** *The* first plausible default trigger package *is any trigger-calibration data package*
$$\mathfrak{P}_{\mathrm{trig}}^{\mathrm{loc,def}} := \left(\varepsilon_{\mathrm{CL}}^{\mathrm{def}}, \varepsilon_{\Theta}^{\mathrm{def}}, \mathfrak{A}_{\mathrm{adm}}^{\mathrm{safe,def}}, m_{\min}^{\mathrm{def}}\right)$$
satisfying:

1. $$0 < \varepsilon_{\mathrm{CL}}^{\mathrm{def}} \leq \varepsilon_{\Theta}^{\mathrm{def}};$$
2. $$\mathfrak{A}_{\mathrm{adm}}^{\mathrm{safe,def}} \subseteq \mathfrak{A}_{\mathrm{adm}};$$
3. $$m_{\min}^{\mathrm{def}} \geq 2.$$

Remark H6.46 (Interpretation of the default package)**.** *This package does not yet assign final numerical values. It only fixes the first local reading: closure burden must enter a stricter small-burden regime than the routed Theta burden, and one-step near-entry is not yet trusted as a sufficient certificate.*

Proposition H6.47 (Default local trigger package is obstruction-compatible)**.** *Assume the first local obstruction classification theorem. Then the first plausible default trigger package is compatible with the current obstruction reading.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. The current obstruction has been classified as calibration/persistence type rather than machine-side decodability type. A stricter closure tolerance together with a safe admissibility subregion and a persistence length at least two directly addresses that obstruction pattern: it stabilizes the closure burden first, keeps the packet-side monitor conservative, and excludes one-step transient near-entry events. Hence the default package is compatible with the current obstruction reading. □

Proposition H6.48 (Why one-step persistence should not be the default)**.** *Under the present obstruction-prior reading, the choice*

$$m_{\min} = 1$$

should not be the first default calibration.

Proof. The obstruction analysis explicitly identified trigger-persistence instability as part of the first local failure mode. If one were to set $m_{\min} = 1$, then a single transient entrance into the local trigger inequality schema would already be counted as sufficient, which removes exactly the persistence safeguard that the current obstruction analysis says is missing. Therefore one-step persistence is not the correct first default choice. □

Theorem H6.49 (First default local trigger calibration theorem)**.** *Assume:*

1. *the first local trigger calibration theorem applies;*
2. *the current obstruction is of calibration/persistence type;*
3. *the closure-phase-first local reading remains active.*

Then the first reasonable local default calibration should be chosen from a first plausible default trigger package

$$\mathfrak{P}_{\mathrm{trig}}^{\mathrm{loc,def}} = \left(\varepsilon_{\mathrm{CL}}^{\mathrm{def}}, \varepsilon_{\Theta}^{\mathrm{def}}, \mathfrak{A}_{\mathrm{adm}}^{\mathrm{safe,def}}, m_{\min}^{\mathrm{def}}\right)$$

with

$$0 < \varepsilon_{\mathrm{CL}}^{\mathrm{def}} \leq \varepsilon_{\Theta}^{\mathrm{def}}, \qquad m_{\min}^{\mathrm{def}} \geq 2.$$

In particular, the first local trigger calibration is closure-prior and persistence-prior rather than one-step permissive.

Proof. By the first local trigger calibration theorem, what is missing is an explicit trigger-calibration package. By the obstruction classification, the leading obstruction is calibration/persistence type and not machine-side decodability. Therefore the first calibration should tighten the closure burden first, allow at least as much tolerance on the routed Theta burden, keep the packet-side admissibility band conservative, and require a genuinely persistent window rather than a one-step fluctuation. These conditions are exactly those of the first plausible default trigger package. □

Corollary H6.50 (Current post- default working rule)**.** *At the present stage, the first local trigger audit should proceed as if:*

1. *closure smallness must be at least as strict as Theta smallness;*
2. *the packet-side admissibility band must be chosen conservatively;*
3. *the first honest local persistence length should be read as*

$$m_{\min} \geq 2.$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Only after this default calibration fails should one relax toward a more permissive local trigger reading.

Proof. Immediate from Theorem H6.49. □

Definition H6.51 (Discretized Teichmüller persistence stream)**.** *Let*

$$X_n$$

denote the current local machine-fused state and let

$$T_N(X_n)$$

be the local Monte-Carlo Teichmüller distortion surrogate.

The discretized Teichmüller persistence stream *is the symbolic stream*

$$\Sigma_{\mathrm{pers}} = (\sigma_n)_{n\geq 0},$$

where each symbol σ_n records the discretized local status of

$$\big(\Delta_{\mathrm{CL}}(n), \Delta_{\Theta R}(T)(n), \rho_A(n), T_N(X_n), T_N(X_{n+1}), |T_N(X_{n+1}) - T_N(X_n)|\big)$$

relative to the active local trigger thresholds.

Definition H6.52 (Kasiski persistence scan)**.** *The* Kasiski persistence scan *is the search for repeated symbolic blocks inside*

$$\Sigma_{\mathrm{pers}}$$

in order to detect recurring local trigger-near, persistence-fail, or admissibility-fail patterns.

Remark H6.53 (Role of Kasiski)**.** *Kasiski is used here as a repetition and spacing detector on the discretized persistence stream. It does not decide admissibility; it only detects recurring local obstruction patterns.*

Definition H6.54 (Hashed persistence window)**.** *A* hashed persistence window *of length m is a hash value assigned to a local window*

$$(\sigma_n, \sigma_{n+1}, \ldots, \sigma_{n+m-1})$$

of the discretized Teichmüller persistence stream.

Definition H6.55 (Markov persistence surface)**.** *The* Markov persistence surface *is the transition surface obtained from the empirical transition frequencies between hashed persistence windows or their reduced regime classes.*

Remark H6.56 (Role of Markov control)**.** *The Markov layer is not a primary admissibility criterion. It is only the readable transition shadow of the shell-/trigger-restricted local persistence dynamics.*

Definition H6.57 (Newton persistence refinement)**.** *The* Newton persistence refinement *is a local Newton-type correction step applied only after a persistence regime has already been selected and a smooth surrogate defect function has been chosen.*

Remark H6.58 (Why Newton comes last)**.** *Newton is not used to discover the regime. It is used only to sharpen the local estimate once the regime has already been isolated by Kasiski/hash/Markov filtering.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition H6.59 (Bombe-sieve persistence elimination layer)**.** *The* bombe-sieve persistence elimination layer *is the contradiction-sensitive elimination of local continuation menus by:*

1. *a trigger-sensitive sieve,*
2. *bombe-style rejection of locally incompatible continuation branches,*
3. *optional nested backtracking across deeper local menus.*

Definition H6.60 (Spectral menu readout)**.** *A* spectral menu readout *is the local ranking of a shell-admissible continuation menu by coding it into a self-adjoint menu operator and reading the preferred continuation as the top eigendirection.*

Definition H6.61 (Vernam-like persistence stream heuristic)**.** *The* Vernam-like persistence stream heuristic *is the working interpretation that the discretized Teichmüller persistence stream behaves like an observable stream generated by a hidden local structural source, so that repeated local obstruction patterns may be used to infer the generator rather than merely the visible symbols.*

Remark H6.62 (How to read the Vernam-like heuristic)**.** *This is not an identity claim with literal cryptographic stream ciphers. It is a search heuristic: the visible persistence symbols are treated as a surface stream whose hidden local generator is to be reconstructed from repetition, spacing, transition structure, and contradiction-sensitive elimination.*

Definition H6.63 (Crypto-guided persistence analysis pipeline)**.** *The* crypto-guided persistence analysis pipeline *is the local tool chain*

$$\textit{Kasiski} \to \textit{hash} \to \textit{Markov} \to \textit{Bombe/Sieve} \to \textit{spectral menu readout} \to \textit{Newton refinement},$$

applied to the discretized Teichmüller persistence stream.

Theorem H6.64 (First crypto-guided persistence scan theorem)**.** *Assume:*

1. *a local Monte-Carlo Teichmüller persistence stream has been constructed;*
2. *the active closure/Theta/admissibility thresholds are fixed locally;*
3. *admissibility remains decided by closure-compatible trigger logic.*

Then the crypto-guided persistence analysis pipeline is admissible as a subordinate persistence-analysis layer. It may be used to detect recurring persistence-failure patterns, compress local continuation menus, and sharpen local trigger calibration without replacing the closure-compatible admissibility criterion.

Proof. The earlier tool-guided compression logic already admits Kasiski, hash/rainbow compression, Markov guidance, contradiction-sensitive elimination, spectral ranking, and Newton refinement provided they remain subordinate to structural admissibility. Here the same discipline is applied to the discretized Teichmüller persistence stream. Hence the pipeline is admissible as a subordinate persistence-analysis layer and may be used to detect, compress, rank, and refine local persistence behavior without replacing closure-compatible trigger logic itself. □

Corollary H6.65 (Current post- persistence working rule)**.** *The next local persistence audit should not rely on pointwise burden smallness alone. It should:*

1. *build the discretized Teichmüller persistence stream;*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

2. *scan it by Kasiski for recurring local obstruction patterns;*
3. *hash and compress local windows;*
4. *read the resulting regime dynamics through a Markov persistence surface;*
5. *eliminate locally contradictory branches by bombe/sieve filtering;*
6. *rank the surviving continuation menu spectrally;*
7. *apply Newton refinement only at the final local sharpening stage.*

Proof. Immediate from Theorem H6.64. □

Definition H6.66 (Local persistence symbols)**.** *Let $\Sigma_{\mathrm{pers}} = (\sigma_n)_{n\geq 0}$ be the discretized Teichmüller persistence stream. The following local symbols are distinguished:*

1. `NT` *(near-trigger): the local burden pair satisfies the pointwise closure/Theta smallness bounds but the minimal persistence window is not yet certified;*
2. `PF` *(persistence-fail): the next local state leaves the final closure-phase trigger slice or fails the Teichmüller persistence control while remaining machine-decodable;*
3. `CF` *(calibration-fail): one of the active closure/Theta trigger inequalities is not yet met;*
4. `AF` *(admissibility-fail): the packet-side monitor leaves the safe admissibility zone;*
5. `EN` *(entry): the local state lies in the calibrated closure-phase trigger slice.*

Definition H6.67 (First recurring local failure motif)**.** *The* first recurring local failure motif *is the shortest repeated symbolic block*

$$\omega_{\mathrm{fail}} = (\sigma_n, \ldots, \sigma_{n+\ell-1})$$

with $\ell \geq 2$ detected by the Kasiski persistence scan and interpreted as the first persistent local obstruction pattern.

Definition H6.68 (Near-trigger / persistence-fail motif)**.** *The* near-trigger / persistence-fail motif *is the symbolic block*

$$\omega_{\mathrm{NT}\to\mathrm{PF}} := (\mathtt{NT}, \mathtt{PF}).$$

It records a local run in which the first step reaches pointwise trigger-near smallness, but the second step fails to preserve the closure-phase trigger slice or the corresponding Teichmüller persistence control.

Remark H6.69 (Why this is the natural first failure final)**.** *The current post-archival stage line is obstruction-prior rather than entry-prior, and its first classified obstruction is of trigger calibration/persistence type rather than machine-side decodability type. Hence the first repeated failure pattern should be sought among motifs that are already near local trigger conditions but fail on the required two-step persistence window.*

Proposition H6.70 (Machine-side decodability is not the first recurring failure source)**.** *Assume the prime-lattice machine realization theorem and the first machine-fused local OP-entry protocol theorem. Then the first recurring local failure motif is not forced by machine-side decodability failure alone.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. Under the machine-fused realization hypotheses, descent, trigger testing, and branching are already available as finite machine-controlled operations on the prime-lattice machine object. Therefore a repeated first failure pattern is not naturally attributed to absence of local machine decodability itself, but to failure of trigger calibration or persistence within the already machine-realizable regime. □

Theorem H6.71 (First recurring local persistence-failure reading)**.** *Assume:*

1. *the closure-phase-first local audit remains the preferred default reading;*
2. *the default local trigger orientation with $\varepsilon_{\mathrm{CL}} \le \varepsilon_{\Theta}$ and $m_{\min} \ge 2$ is adopted;*
3. *the current line is obstruction-prior rather than entry-prior;*
4. *the crypto-guided persistence analysis pipeline is applied to the local discretized Teichmüller persistence stream.*

Then the first recurring local failure motif is read by default as

$$\omega_{\mathrm{fail}} = \omega_{\mathrm{NT}\to\mathrm{PF}} = (\mathtt{NT}, \mathtt{PF}),$$

that is: a trigger-near first step followed by failure of two-step closure-phase persistence.

Proof. By obstruction-priority, immediate entry is not the default first local outcome. By the earlier obstruction classification, the first local obstruction is of trigger calibration/persistence type rather than machine-side decodability type. Under the default trigger orientation and the requirement $m_{\min} \ge 2$, the most economical local failure pattern is therefore not a one-step calibration collapse before near-trigger status is reached, but a run in which the first step attains trigger-near pointwise smallness and the second step fails to preserve the closure-phase trigger slice. The crypto-guided persistence scan is designed exactly to detect repeated symbolic blocks of this form. Hence the default first recurring local failure motif is $(\mathtt{NT}, \mathtt{PF})$. □

Corollary H6.72 (Current first persistence-stream reading)**.** *The current post-archival stage persistence line should be read first through the motif*

$$(\mathtt{NT}, \mathtt{PF}),$$

rather than through immediate entry or pure decodability failure. Equivalently, the present first local task is to identify which closure-phase persistence ingredient breaks between the near-trigger first step and the failed second step.

Proof. Immediate from Theorem H6.71 and Proposition H6.70. □

Definition H6.73 (Theta-drift persistence-fail subtype)**.** *The symbol*

$$\mathtt{PF}_{\Theta}$$

denotes the persistence-fail subtype in which:

1. *the first local step is near-trigger;*
2. *the second local step fails because the phase-side component loses the required closure-phase persistence control;*
3. *the packet-side admissibility monitor remains inside the safe zone.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition H6.74 (Teichmüller-distortion persistence-fail subtype)**.** *The symbol*

$$\mathtt{PF}_T$$

denotes the persistence-fail subtype in which:

1. *the first local step is near-trigger;*
2. *the pointwise burden inequalities remain approximately trigger-near;*
3. *the second local step fails because the local Teichmüller distortion surrogate or its one-step variation leaves the allowed persistence band.*

Definition H6.75 (Admissibility-edge persistence-fail subtype)**.** *The symbol*

$$\mathtt{PF}_A$$

denotes the persistence-fail subtype in which:

1. *the first local step is near-trigger;*
2. *the second local step fails because the packet-side monitor ρ_A reaches or leaves the safe admissibility edge;*
3. *the corresponding closure-phase persistence window therefore collapses for admissibility reasons.*

Remark H6.76 (How the three subtypes differ)**.** *The subtype $\mathtt{PF}_\Theta$ is phase-led and burden-side. The subtype $\mathtt{PF}_T$ is geometric and distortion-led. The subtype $\mathtt{PF}_A$ is guard-led and packet-admissibility driven.*

Proposition H6.77 (Admissibility-edge contact is not the default first persistence-fail subtype)**.** *Assume the current line is read through the default local trigger orientation with conservative packet-side admissibility control. Then $\mathtt{PF}_A$ is not the default first persistence-fail subtype.*

Proof. The current line treats ρ_A primarily as an admissibility monitor rather than as a coequal core burden, and the active default trigger package keeps the safe admissibility band deliberately conservative. Hence the first local failure is not naturally read as immediate edge contact of ρ_A, but as failure of the closure-phase persistence mechanism before the admissibility edge is reached. □

Proposition H6.78 (The first persistence-fail subtype is phase/geometry-prior)**.** *Assume the closure-phase-first local audit, the default trigger orientation, and the first recurring persistence-failure reading*

$$(\mathtt{NT}, \mathtt{PF}).$$

Then the first persistence-fail subtype is read by default as phase/geometry-prior, i.e. as lying in the reduced pair

$$\{\mathtt{PF}_\Theta, \mathtt{PF}_T\},$$

rather than as an admissibility-edge subtype $\mathtt{PF}_A$.

Proof. Under the closure-phase-first reading, the active mixed persistence burden is carried first by the closure/Theta pair, while ρ_A remains a guard variable. By Proposition H6.77, immediate admissibility-edge contact is not the default first subtype. Hence the first persistence-fail subtype is phase/geometry-prior. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Theorem H6.79 (First local persistence-fail subtype theorem)**.** *Assume:*

1. *the first recurring local failure motif is*

$$(\mathtt{NT}, \mathtt{PF});$$

2. *the closure-phase-first local audit remains the default reading;*

3. *the packet-side admissibility monitor remains conservative on the first near-trigger step.*

Then the first local persistence-fail subtype is not read by default as admissibility-edge contact. Instead, it is read by default as a phase/geometry-prior break: either

$$\mathtt{PF}_{\Theta}$$

or

$$\mathtt{PF}_{T},$$

with the current line favoring the reading that Theta-side drift is the primary burden-side break and Teichmüller distortion is its geometric witness.

Proof. By Proposition H6.78, the first subtype lies in the pair $\{\mathtt{PF}_{\Theta}, \mathtt{PF}_{T}\}$ rather than in $\mathtt{PF}_{A}$. Under the closure-phase-first reading, the active burden is phase-led before it is envelope-led, while the Teichmüller control records the geometric persistence of that phase-led regime. Therefore the default local reading is that Theta-drift is the primary burden-side break and Teichmüller distortion is the corresponding geometric witness of the failed second step. □

Corollary H6.80 (Current post- local diagnosis rule)**.** *The next local diagnosis should not first ask whether ρ_A touched the admissibility edge. It should first ask:*

1. *did the phase-side burden drift out of the closure-phase persistence window;*

2. *and if so, what Teichmüller distortion witness recorded that loss of persistence?*

Proof. Immediate from Theorem H6.79. □

Definition H6.81 (Local Theta-drift witness)**.** *A* local Theta-drift witness *is a finite witness object*

$$\mathfrak{W}_{\Theta}^{\mathrm{loc}}(n)$$

attached to a near-trigger first step and the following failed second step, certifying that:

1. *the first step is of type* $\mathtt{NT}$*;*

2. *the second step is of type* $\mathtt{PF}_{\Theta}$ *or phase/geometry-prior persistence failure;*

3. *the phase-side burden leaves the active closure-phase persistence window on the second step while the packet-side admissibility monitor remains inside the conservative safe zone.*

Definition H6.82 (Local Teichmüller drift witness)**.** *A* local Teichmüller drift witness *is a finite witness object*

$$\mathfrak{W}_{T}^{\mathrm{loc}}(n)$$

certifying that the local Teichmüller distortion surrogate or its one-step variation crosses the active persistence band between the near-trigger first step and the failed second step.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Remark H6.83 (Burden-side versus geometric witness)**.** *The Theta-drift witness and the Teichmüller drift witness play different roles. The first records the burden-side loss of closure-phase persistence. The second records the geometric distortion signature of that same loss.*

Definition H6.84 (Local coupled drift witness package)**.** *The* local coupled drift witness package *is the pair*

$$\mathfrak{W}^{\mathrm{loc}}_{\mathrm{drift}}(n) := \big(\mathfrak{W}^{\mathrm{loc}}_{\Theta}(n), \mathfrak{W}^{\mathrm{loc}}_{T}(n)\big),$$

read on the current persistence stream whenever the local failure motif

$$(\mathtt{NT}, \mathtt{PF})$$

is refined to a phase/geometry-prior break.

Definition H6.85 (Local Theta-drift witness test)**.** *The* local Theta-drift witness test *asks whether, for some local index n:*

1. *the first step satisfies the active near-trigger inequalities;*
2. *the second step violates the phase-side persistence requirement before admissibility edge contact occurs;*
3. *a Teichmüller distortion witness is simultaneously present or becomes newly active on that second step.*

Proposition H6.86 (Theta-drift witness packages refine the default persistence-fail reading)**.** *Assume the first recurring local failure motif is*

$$(\mathtt{NT}, \mathtt{PF})$$

and the first persistence-fail subtype is phase/geometry-prior. Then every successful local Theta-drift witness test yields a local coupled drift witness package

$$\mathfrak{W}^{\mathrm{loc}}_{\mathrm{drift}}(n)$$

refining the persistence-fail reading.

Proof. If the local Theta-drift witness test succeeds, then the failed second step is not merely an undifferentiated persistence failure: it is a phase-side drift event accompanied by a geometric distortion witness while admissibility remains conservative. That is exactly the data recorded by the two witness objects and therefore by their paired witness package. □

Theorem H6.87 (First local Theta-drift witness theorem)**.** *Assume:*

1. *the current post-archival stage persistence line is read through the first recurring motif*

$$(\mathtt{NT}, \mathtt{PF});$$

2. *the first persistence-fail subtype is phase/geometry-prior;*
3. *the local Theta-drift witness test succeeds at some first relevant local index n.*

Then the current line admits a first local Theta-drift witness package

$$\mathfrak{W}^{\mathrm{loc}}_{\mathrm{drift}}(n) = \big(\mathfrak{W}^{\mathrm{loc}}_{\Theta}(n), \mathfrak{W}^{\mathrm{loc}}_{T}(n)\big),$$

with the following reading:

© 2026 Marc Brendecke. Preprint — not peer reviewed.

1. *the first failed second step is burden-side dominated by Theta-drift;*
2. *the corresponding Teichmüller witness is the geometric recorder of that burden-side drift;*
3. *the first local obstruction is therefore not yet read as pure trigger-calibration failure and not yet as admissibility-edge collapse, but as a drift-mediated persistence break.*

Proof. By the first recurring persistence-failure reading, the current line is governed first by the motif

$$(\texttt{NT}, \texttt{PF}).$$

By the first local persistence-fail subtype theorem, the first persistence-fail subtype is read as phase/geometry-prior rather than admissibility-edge driven. If the local Theta-drift witness test succeeds at the first relevant index, then the failed second step is specifically identified as a phase-side drift event with an accompanying Teichmüller distortion witness. Hence the local coupled drift witness package exists and carries the stated interpretation. □

Corollary H6.88 (Current post- drift diagnosis rule)**.** *Once the first local Theta-drift witness package is available, the next local audit should ask first which phase-side persistence ingredient must be strengthened in order to keep the second step inside the closure-phase window, and only afterwards whether a further trigger-calibration reduction is needed.*

Proof. Immediate from Theorem H6.87. □

Definition H6.89 (Local phase-side reinforcement datum)**.** *A* local phase-side reinforcement datum *at a first relevant local index n is a finite correction datum*

$$\mathfrak{R}_{\Theta}^{\mathrm{loc}}(n) := \big(\eta_{\Theta}(n), \eta_{T}(n)\big),$$

where:

1. $\eta_{\Theta}(n) \geq 0$ *is the minimal phase-side persistence strengthening required to keep the second step inside the active closure-phase window;*
2. $\eta_{T}(n) \geq 0$ *is the corresponding geometric tolerance compensation required so that the Teichmüller distortion witness stays inside the active persistence band.*

Definition H6.90 (Local repaired second step)**.** *A failed second step at index $n+1$ is called* locally repaired *by a reinforcement datum*

$$\mathfrak{R}_{\Theta}^{\mathrm{loc}}(n) = \big(\eta_{\Theta}(n), \eta_{T}(n)\big)$$

if, after strengthening the phase-side persistence requirement by $\eta_{\Theta}(n)$ and compensating the active Teichmüller persistence band by $\eta_{T}(n)$, the updated second step:

1. *remains in the closure-phase trigger window;*
2. *preserves admissibility of ρ_A;*
3. *no longer issues the persistence-fail symbol* PF*.*

Remark H6.91 (Meaning of local repair)**.** *The repair datum does not assert that the original failed step was already admissible. It records the smallest phase-side and geometric strengthening under which the same local two-step pattern would have stayed inside the active closure-phase window.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition H6.92 (Minimal local phase-side reinforcement)**.** *A local phase-side reinforcement datum*

$$\mathfrak{R}_{\Theta}^{\mathrm{loc}}(n) = \big(\eta_{\Theta}(n), \eta_{T}(n)\big)$$

is called minimal *if no strictly smaller pair in the product order on* $\mathbb{R}^2_{\geq 0}$ *already yields a locally repaired second step.*

Definition H6.93 (First local repair witness)**.** *A* first local repair witness *is a pair*

$$\mathfrak{W}_{\mathrm{rep}}^{\mathrm{loc}}(n) := \big(\mathfrak{W}_{\mathrm{drift}}^{\mathrm{loc}}(n), \mathfrak{R}_{\Theta}^{\mathrm{loc}}(n)\big)$$

consisting of:

1. *the first local coupled drift witness package;*
2. *a minimal local phase-side reinforcement datum repairing the failed second step.*

Proposition H6.94 (Theta-drift witnesses induce local repair data)**.** *Assume the first local Theta-drift witness theorem applies. Then every first local Theta-drift witness package determines a nonnegative local phase-side reinforcement datum.*

Proof. The witness package identifies both the burden-side Theta drift and the corresponding Teichmüller distortion contribution at the failed second step while keeping admissibility conservative. Hence one may measure how much phase-side strengthening and geometric tolerance compensation are needed to return the same local second step to the active closure-phase window. These quantities are nonnegative by construction. □

Lemma H6.95 (First local repair lemma)**.** *Assume:*

1. *the first local Theta-drift witness theorem applies at index* n*;*
2. *a minimal local phase-side reinforcement datum*

$$\mathfrak{R}_{\Theta}^{\mathrm{loc}}(n) = \big(\eta_{\Theta}(n), \eta_{T}(n)\big)$$

exists.

Then the associated first local repair witness

$$\mathfrak{W}_{\mathrm{rep}}^{\mathrm{loc}}(n)$$

certifies the smallest local strengthening currently needed to keep the second step inside the closure-phase persistence window.

Proof. By the first local Theta-drift witness theorem, the failed second step is already isolated as a Theta-drift event with Teichmüller distortion witness. A minimal reinforcement datum records the smallest phase-side and geometric compensation required to restore persistence of that step. Therefore the combined repair witness certifies the smallest currently needed local strengthening. □

Theorem H6.96 (First local phase-side repair theorem)**.** *Assume:*

1. *the first recurring local failure motif is*

$$(\mathtt{NT}, \mathtt{PF});$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

2. *the first persistence-fail subtype is phase/geometry-prior;*

3. *the first local Theta-drift witness theorem applies at index n;*

4. *a minimal local phase-side reinforcement datum exists at that same index.*

Then the current local persistence obstruction is repair-readable. More precisely, the first local failure is governed by a minimal reinforcement package

$$\mathfrak{R}_{\Theta}^{\mathrm{loc}}(n) = \big(\eta_{\Theta}(n), \eta_{T}(n)\big)$$

which measures exactly how much phase-side persistence strengthening and geometric tolerance compensation are missing at the failed second step.

Proof. The first three assumptions isolate the first failed second step as a Theta-drift persistence failure with Teichmüller witness. The existence of a minimal reinforcement datum then identifies the smallest correction under which the same local two-step pattern would remain inside the closure-phase window. Therefore the current local obstruction is not only diagnosable but also repair-readable in the stated sense. □

Corollary H6.97 (Current post- repair rule)**.** *After the first local Theta-drift witness has been isolated, the next local audit should not first ask for a new global architecture. It should ask for the minimal local reinforcement package*

$$\big(\eta_{\Theta}(n), \eta_{T}(n)\big)$$

needed to keep the second step inside the active closure-phase persistence window.

Proof. Immediate from Theorem H6.96. □

Definition H6.98 (Theta-dominant local repair)**.** *A minimal local reinforcement package*

$$\mathfrak{R}_{\Theta}^{\mathrm{loc}}(n) = \big(\eta_{\Theta}(n), \eta_{T}(n)\big)$$

is called Theta-dominant *if:*

1. $\eta_{\Theta}(n) > 0$*;*

2. $\eta_{T}(n) \geq 0$*;*

3. *the failed second step cannot be repaired by geometric compensation alone when $\eta_{\Theta}(n)$ is set to* 0*.*

Definition H6.99 (Geometrically carried local repair)**.** *A minimal local reinforcement package*

$$\mathfrak{R}_{\Theta}^{\mathrm{loc}}(n) = \big(\eta_{\Theta}(n), \eta_{T}(n)\big)$$

is called geometrically carried *if:*

1. $\eta_{T}(n) > 0$*;*

2. *the corresponding Teichmüller witness is an essential part of the repair;*

3. *the repaired second step cannot be restored by burden-side phase strengthening alone while keeping the active geometric tolerance unchanged.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Remark H6.100 (Two-sided reading of the repair datum)**.** *The local repair datum measures two distinct but coupled deficits:*

1. *burden-side phase persistence deficit, read through $\eta_\Theta(n)$;*
2. *geometric persistence deficit, read through $\eta_T(n)$.*

The point of the current audit is to determine which of these two deficits is structurally primary in the first failed second step.

Definition H6.101 (Locally coupled repair orientation)**.** *The first local repair orientation at index n is called:*

1. burden-dominant *if the minimal repair package is Theta-dominant and not geometrically carried;*
2. geometric-dominant *if the minimal repair package is geometrically carried and not Theta-dominant;*
3. genuinely coupled *if the minimal repair package is both Theta-dominant and geometrically carried.*

Proposition H6.102 (Theta-drift witnesses force nonzero burden-side repair)**.** *Assume the first local Theta-drift witness theorem applies at index n. Then every minimal local reinforcement package at that index satisfies*

$$\eta_\Theta(n) > 0.$$

Proof. The first local Theta-drift witness theorem identifies the failed second step specifically as a phase-side drift event with Teichmüller witness. If one set $\eta_\Theta(n) = 0$, the burden-side phase drift would remain uncorrected, so the failed second step could not be repaired in the sense of the local repair definition. Hence every minimal repair package must have strictly positive $\eta_\Theta(n)$. □

Proposition H6.103 (Teichmüller witnesses make the repair geometrically visible)**.** *Assume the first local Theta-drift witness theorem applies at index n and the corresponding failed second step carries a nontrivial Teichmüller witness. Then every minimal local reinforcement package at that index satisfies*

$$\eta_T(n) \geq 0$$

with geometric significance, and the resulting repair orientation cannot be read as purely burden-side in the naive scalar sense.

Proof. A nontrivial Teichmüller witness means that the failed second step is accompanied by a geometric persistence defect, not only by a symbolic burden-side drift. Therefore the repair package necessarily carries a geometric component, even if its size is smaller than the burden-side one. Thus the repair is not purely scalar/burden-side in the naive sense. □

Theorem H6.104 (First local repair-orientation theorem)**.** *Assume:*

1. *the first local Theta-drift witness theorem applies at index n;*
2. *a minimal local phase-side reinforcement datum exists at that index;*
3. *the corresponding failed second step carries a nontrivial Teichmüller witness.*

Then the first local repair orientation is not geometric-dominant. More precisely:

© 2026 Marc Brendecke. Preprint — not peer reviewed.

1. *the minimal repair package is necessarily Theta-dominant;*
2. *the geometric contribution remains a genuine witness of the repair problem;*
3. *hence the first local repair is read either as burden-dominant with geometric witness or as genuinely coupled, but not as purely geometric.*

Proof. By Proposition H6.102, one has $\eta_\Theta(n) > 0$ for every minimal repair package. Hence the repair cannot be geometric-dominant in the sense of Definition H6.101. By Proposition H6.103, the Teichmüller component remains a genuine geometric witness of the same local failure. Therefore the first local repair is either burden-dominant with geometric witness or genuinely coupled, but not purely geometric. □

Corollary H6.105 (Current post- repair reading)**.** *The current post-archival stage local repair problem should be read first as a phase-side burden repair problem with a genuine Teichmüller witness. Accordingly, the next local audit should ask:*

1. *how large the minimal burden-side strengthening $\eta_\Theta(n)$ must be;*
2. *whether the corresponding geometric term $\eta_T(n)$ is only subordinate or already large enough to force a genuinely coupled repair reading.*

Proof. Immediate from Theorem H6.104. □

Definition H6.106 (Locally plausible tightened trigger package)**.** *The locally tightened trigger package*

$$\mathfrak{P}^\sharp_{\mathrm{trig}}(n) = \left(\varepsilon^\sharp_{\mathrm{CL}}, \varepsilon^\sharp_\Theta, \varepsilon^\sharp_T, \mathfrak{A}^{\mathrm{safe}}_{\mathrm{adm}}, m_{\min}\right)$$

at index n is called locally plausible *if:*

1. *all tightened thresholds remain positive;*
2. *the admissibility band $\mathfrak{A}^{\mathrm{safe}}_{\mathrm{adm}}$ remains nonempty;*
3. *the tightened package is compatible with the current closure-phase regime at the first step.*

Definition H6.107 (Residual tightened-package insufficiency)**.** *A* residual tightened-package insufficiency *at index n is any component of the locally tightened trigger package whose strengthened form still fails to support the repaired two-step closure-phase persistence test.*

The possible insufficiency types are:

1. *closure-side insufficiency;*
2. *Theta-side insufficiency;*
3. *Teichmüller-side insufficiency;*
4. *admissibility-band insufficiency;*
5. *persistence-length insufficiency.*

Remark H6.108 (What the new audit should decide)**.** *After constructing the first repair-induced tightening, the question is no longer whether the original trigger package fails. The sharper question is which component of the tightened package still blocks the repaired second step.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition H6.109 (Teichmüller-persistence residual insufficiency)**.** *The residual insufficiency is called* Teichmüller-persistence type *if:*

1. *the tightened closure-side threshold is already compatible with the repaired first step;*
2. *the tightened Theta-side threshold is already compatible with the repaired first step;*
3. *the admissibility band remains safe;*
4. *but the repaired two-step window still fails because the Teichmüller-side bound or its two-step stability requirement is not yet strong enough.*

Proposition H6.110 (The tightened package is locally plausible)**.** *Assume the first local repair-to-trigger calibration theorem applies at index n. Then the tightened trigger package*

$$\mathfrak{P}^{\sharp}_{\mathrm{trig}}(n)$$

is locally plausible.

Proof. By the repair-to-trigger calibration theorem, the tightened thresholds are obtained from a genuine local repair witness and remain admissible as a final package. Hence the tightened package preserves positivity, keeps the admissibility band meaningful, and remains compatible with the first-step closure-phase regime. □

Proposition H6.111 (Closure and admissibility are not the first remaining insufficiency)**.** *Assume:*

1. *the first local Theta-drift witness theorem applies;*
2. *the first local repair-orientation theorem applies;*
3. *the tightened trigger package is locally plausible.*

Then the first residual tightened-package insufficiency is not closure-side and not admissibility-band type.

Proof. The local failure was already classified as phase/geometry-prior rather than closure-prior, and the local repair package was constructed without changing the closure-side threshold at first order. Moreover the current line is not admissibility-edge-prior. Therefore, after tightening, the first remaining insufficiency cannot honestly be read first as closure-side or admissibility-band type. □

Theorem H6.112 (First tightened-package plausibility theorem)**.** *Assume:*

1. *the first local repair-to-trigger calibration theorem applies at index n;*
2. *the first local repair-orientation theorem applies at index n;*
3. *the current local obstruction is phase/geometry-prior rather than closure/admissibility-prior.*

Then the current post-archival stage line supports the following reading of the locally tightened trigger package:

1. *the tightened package is locally plausible;*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

2. *if it still fails to restore two-step closure-phase persistence, then the first remaining insufficiency is no longer closure-side or admissibility-side;*

3. *the first remaining insufficiency should be searched first on the Teichmüller-persistence side, i.e. in the tightened geometric threshold or its two-step stability requirement.*

Proof. Local plausibility is Proposition H6.110. By Proposition H6.111, closure-side and admissibility-band failure are excluded as the first residual obstruction. The local repair orientation is Theta-dominant but geometrically carried, so once the primary burden-side strengthening has been applied, the next unresolved obstruction is searched first on the Teichmüller-persistence side. □

Corollary H6.113 (Current post- tightened-package reading)**.** *The current post-archival stage line should read the first tightened trigger package as plausibly near-sufficient but not yet self-certifying. If the repaired two-step persistence still fails, the next local audit should first test whether the tightened Teichmüller-side bound or its persistence clause remains insufficient.*

Proof. Immediate from Theorem H6.112. □

Definition H6.114 (Local Teichmüller-side repair datum)**.** *Let*

$$\mathfrak{P}^{\sharp}_{\mathrm{trig}}(n) = \big(\varepsilon^{\sharp}_{\mathrm{CL}}, \varepsilon^{\sharp}_{\Theta}, \varepsilon^{\sharp}_{T}, \mathfrak{A}^{\mathrm{safe}}_{\mathrm{adm}}, m_{\min}\big)$$

be the locally tightened trigger package at index n.

A local Teichmüller-side repair datum *is a pair*

$$\mathfrak{R}^{\mathrm{loc}}_{T}(n) := \big(\zeta_T(n), \lambda_T(n)\big)$$

with

$$\zeta_T(n) \geq 0, \qquad \lambda_T(n) \geq 0,$$

where:

1. *$\zeta_T(n)$ is the additional tightening of the geometric threshold;*

2. *$\lambda_T(n)$ is the additional tightening of the two-step distortion-variation allowance.*

Definition H6.115 (Teichmüller-reinforced trigger package)**.** *The* Teichmüller-reinforced trigger package *at index n is*

$$\mathfrak{P}^{\flat}_{\mathrm{trig}}(n) := \big(\varepsilon^{\sharp}_{\mathrm{CL}}, \varepsilon^{\sharp}_{\Theta}, \varepsilon^{\flat}_{T}, \mathfrak{A}^{\mathrm{safe}}_{\mathrm{adm}}, m_{\min}, \eta^{\flat}_{T}(n)\big),$$

where

$$\varepsilon^{\flat}_{T} := \varepsilon^{\sharp}_{T} - \zeta_T(n), \qquad \eta^{\flat}_{T}(n) := \eta_T(n) - \lambda_T(n),$$

and the reinforced package is required to remain positive and admissible.

Definition H6.116 (Teichmüller-reinforced two-step persistence test)**.** *A two-step local window*

$$X_n,\ X_{n+1}$$

passes the Teichmüller-reinforced two-step persistence test *if for both $k \in \{n, n+1\}$:*

$$\Delta_{\mathrm{CL}}(k) \leq \varepsilon^{\sharp}_{\mathrm{CL}}, \qquad \Delta_{\Theta R}(T)(k) \leq \varepsilon^{\sharp}_{\Theta}, \qquad T_N(X_k) \leq \varepsilon^{\flat}_{T}, \qquad \rho_A(k) \in \mathfrak{A}^{\mathrm{safe}}_{\mathrm{adm}},$$

and additionally

$$|T_N(X_{n+1}) - T_N(X_n)| \leq \eta^{\flat}_{T}(n).$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Remark H6.117 (Meaning of the Teichmüller-side repair)**.** *If the first tightened trigger package is already closure-compatible and Theta-compatible but still fails on the persistence side, then the next honest local move is no longer to tighten closure or admissibility first. It is to sharpen the geometric threshold and its two-step variation allowance.*

Definition H6.118 (Minimal local Teichmüller-side repair)**.** *A local Teichmüller-side repair datum*

$$\big(\zeta_T(n), \lambda_T(n)\big)$$

is called minimal *if the corresponding Teichmüller-reinforced trigger package passes the two-step persistence test, but every componentwise smaller repair datum fails to do so.*

Proposition H6.119 (Teichmüller-side repair leaves closure and Theta tightening fixed)**.** *Assume the first local repair-to-trigger calibration theorem applies and the first tightened-package plausibility theorem identifies a Teichmüller-persistence residual insufficiency at index n. Then a local Teichmüller-side repair acts only on the geometric threshold and the two-step variation allowance, while leaving*

$$\varepsilon^{\sharp}_{\mathrm{CL}} \qquad \text{and} \qquad \varepsilon^{\sharp}_{\Theta}$$

unchanged.

Proof. By hypothesis, the residual insufficiency is not closure-side and not admissibility-band type, and the first repair orientation is Theta-dominant but geometrically carried. Therefore the remaining local insufficiency is carried by the geometric persistence side, so the next tightening acts only on the Teichmüller-side threshold and its variation allowance. □

Lemma H6.120 (First local Teichmüller-persistence repair lemma)**.** *Assume:*

1. *the first tightened-package plausibility theorem applies at index n;*
2. *the remaining insufficiency is of Teichmüller-persistence type;*
3. *there exists a minimal local Teichmüller-side repair datum.*

Then the corresponding Teichmüller-reinforced trigger package is the first honest local final for restoring the missing two-step closure-phase persistence after the initial Theta-dominant repair.

Proof. The first tightened package already preserves closure-side and Theta-side compatibility and keeps the admissibility band safe. By assumption, the only remaining failure lies on the Teichmüller-persistence side. A minimal local Teichmüller-side repair datum therefore gives exactly the first additional geometric strengthening required to restore the missing two-step persistence without over-tightening the other channels. □

Theorem H6.121 (First local Teichmüller-persistence repair theorem)**.** *Assume:*

1. *the first local Theta-drift witness theorem applies;*
2. *the first local repair-to-trigger calibration theorem applies;*
3. *the first tightened-package plausibility theorem applies;*
4. *the residual insufficiency is of Teichmüller-persistence type.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Then the current post-archival stage line supports a second-stage local trigger recalibration on the geometric persistence side.

More precisely:

1. *the closure threshold remains fixed at*
$$\varepsilon_{\mathrm{CL}}^{\sharp};$$
2. *the Theta threshold remains fixed at*
$$\varepsilon_{\Theta}^{\sharp};$$
3. *the next honest recalibration acts on*
$$\varepsilon_{T}^{\sharp} \quad \textit{and} \quad \eta_{T}(n)$$
through a local Teichmüller-side repair datum
$$\big(\zeta_{T}(n), \lambda_{T}(n)\big);$$
4. *the resulting reinforced package*
$$\mathfrak{P}_{\mathrm{trig}}^{\flat}(n)$$
is the first concrete final for restoring the missing two-step closure-phase persistence after the initial Theta-dominant tightening.

Proof. By the first local repair-to-trigger calibration theorem, the initial repair is Theta-dominant but geometrically carried. By the tightened-package plausibility theorem, the remaining insufficiency is searched first on the Teichmüller-persistence side. Therefore closure and Theta thresholds remain fixed while the next honest recalibration must act on the geometric threshold and its persistence allowance. This yields the claimed Teichmüller-reinforced trigger package as the next concrete final. □

Corollary H6.122 (Current post- working rule)**.** *The next local audit should no longer ask only whether the first tightened trigger package is near-sufficient. It should ask whether a minimal local Teichmüller-side repair datum*
$$\big(\zeta_{T}(n), \lambda_{T}(n)\big)$$
restores the missing two-step closure-phase persistence while leaving the closure and Theta thresholds fixed.

Proof. Immediate from Theorem H6.121. □

H7 Complete OP Transport Arguments

Technical unfolding of the OP transport architecture

Status. *Active proof-unfolding block.*

Block function. This section expands the OP transport chain at theorem level and should be used when the compressed main-line statements need to be reopened into their structural proof pieces.

This technical block unfolds the compressed OP branch-shell theorem used in the main line. It records the native closure machinery, the hardening lemmas, and the detailed branch/shell routed proofs that support the compressed theorem, so later sections can appeal either to the short main-text statement or to the full theorem-level form given here.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Dependency guide for the unfolding. The technical order is now explicit: Theorem H7.2 fixes the OP 1/OP 5 source-return pair, Theorem H7.18 promotes admissible branch transport, Theorem H7.21 localizes shell/residual admissibility at OP 3, and Theorem H7.25 combines these layers into the routed visible return theorem. Proposition H7.33 then records the packet/phase-lock/epsilon line as a diagnostic screening layer rather than a second closure theorem.

H7.1 Closure of the OP1–OP5 pair

The OP-chain should not be read as a naive raw endpoint comparison. In the present framework, OP 1 is the inflow-side readout of a fixed structural route, whereas OP 5 belongs to the return sector of the same route and must be read through its reversal-stable part.

Let Γ be a fixed structural route with reversed route $\overline{\Gamma}$, and let

$$T_\Gamma, T_{\overline{\Gamma}} : \mathcal{A}_{\mathrm{cl}} \to \mathcal{C}$$

denote the forward and backward carrier transports on the admissible closure sector $\mathcal{A}_{\mathrm{cl}} \subseteq \mathcal{C}$. Let

$$\rho_\theta : \mathcal{C} \to \mathcal{W}$$

be the frame-dependent readout family.

We write

$$\mathrm{OP1} = \rho_{\theta_{\mathrm{in}}}(c_\Gamma), \qquad \mathrm{OP5}_{\mathrm{fwd}} = \rho_{\theta_{\mathrm{fwd}}}(T_\Gamma c_\Gamma), \qquad \mathrm{OP5}_{\mathrm{bwd}} = \rho_{\theta_{\mathrm{bwd}}}(T_{\overline{\Gamma}} c_\Gamma),$$

for a carrier $c_\Gamma \in \mathcal{A}_{\mathrm{cl}}$, and define the symmetrized return readout by

$$\mathrm{OP5}_{\mathrm{sym}} := \frac{1}{2}\big(\mathrm{OP5}_{\mathrm{fwd}} + \mathrm{OP5}_{\mathrm{bwd}}\big).$$

To express structural closure, we use the native closure carrier class

$$K_{\mathrm{ncl}} : \mathcal{A}_{\mathrm{cl}} \to \mathcal{A}_{\mathrm{cl}}/\!\sim_{\mathrm{ncl}},$$

which identifies carriers up to HC-compensated transport, admissible frame-shift, and admissible branch refinement of the same closure-bearing mode.

Assumption H7.1 (Native OP 1–OP 5 closure assumptions)**.** *We assume:*

(i) ***HC-confined transport:***

$$T_\Gamma(\mathcal{A}_{\mathrm{cl}}) \subseteq \mathcal{A}_{\mathrm{cl}}, \qquad T_{\overline{\Gamma}}(\mathcal{A}_{\mathrm{cl}}) \subseteq \mathcal{A}_{\mathrm{cl}};$$

(ii) ***two-pass class consistency:***

$$K_{\mathrm{ncl}}(T_\Gamma c) = K_{\mathrm{ncl}}(c) = K_{\mathrm{ncl}}(T_{\overline{\Gamma}} c) \qquad \textit{for all } c \in \mathcal{A}_{\mathrm{cl}};$$

(iii) ***closure-compatible readout:*** *there exists a map*

$$J : \mathcal{A}_{\mathrm{cl}}/\!\sim_{\mathrm{ncl}} \to \Lambda$$

and an observable invariant

$$\mathcal{I} : \mathcal{W} \to \Lambda$$

such that

$$\mathcal{I}(\rho_\theta(c)) = J(K_{\mathrm{ncl}}(c)) \qquad \textit{for all } c \in \mathcal{A}_{\mathrm{cl}},\ \theta \in \Theta;$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(iv) ***return-sector linearity:*** *$\mathcal{I}$ is linear on the admissible OP 5 return sector.*

Theorem H7.2 (Closure of the OP 1–OP 5 pair)**.** *Under the native OP 1–OP 5 closure assumptions,*

$$\mathcal{I}(\mathrm{OP1}) = \mathcal{I}(\mathrm{OP5}_{\mathrm{sym}}).$$

Hence the OP 1–OP 5 pair is closed at the level of the fixed closure invariant.

Proof. By closure-compatible readout,

$$\mathcal{I}(\mathrm{OP1}) = \mathcal{I}(\rho_{\theta_{\mathrm{in}}}(c_\Gamma)) = J(K_{\mathrm{ncl}}(c_\Gamma)).$$

Likewise,

$$\mathcal{I}(\mathrm{OP5}_{\mathrm{fwd}}) = J(K_{\mathrm{ncl}}(T_\Gamma c_\Gamma)), \qquad \mathcal{I}(\mathrm{OP5}_{\mathrm{bwd}}) = J(K_{\mathrm{ncl}}(T_{\overline{\Gamma}} c_\Gamma)).$$

By two-pass class consistency,

$$K_{\mathrm{ncl}}(T_\Gamma c_\Gamma) = K_{\mathrm{ncl}}(c_\Gamma) = K_{\mathrm{ncl}}(T_{\overline{\Gamma}} c_\Gamma),$$

and therefore

$$\mathcal{I}(\mathrm{OP5}_{\mathrm{fwd}}) = \mathcal{I}(\mathrm{OP1}) = \mathcal{I}(\mathrm{OP5}_{\mathrm{bwd}}).$$

Using linearity on the admissible return sector,

$$\mathcal{I}(\mathrm{OP5}_{\mathrm{sym}}) = \frac{1}{2}\Big(\mathcal{I}(\mathrm{OP5}_{\mathrm{fwd}}) + \mathcal{I}(\mathrm{OP5}_{\mathrm{bwd}})\Big) = \mathcal{I}(\mathrm{OP1}),$$

which proves the claim. □

Corollary H7.3 (Odd return defect is closure-invisible)**.** *Define*

$$\Delta_{\mathrm{OP5}} := \frac{1}{2}\big(\mathrm{OP5}_{\mathrm{fwd}} - \mathrm{OP5}_{\mathrm{bwd}}\big).$$

Then

$$\mathcal{I}(\Delta_{\mathrm{OP5}}) = 0.$$

Proof. By linearity,

$$\mathcal{I}(\Delta_{\mathrm{OP5}}) = \frac{1}{2}\Big(\mathcal{I}(\mathrm{OP5}_{\mathrm{fwd}}) - \mathcal{I}(\mathrm{OP5}_{\mathrm{bwd}})\Big) = 0,$$

since the theorem gives equality of the two return-side invariant values. □

Remark H7.4. *The theorem does not require raw observable equality*

$$\mathrm{OP1} = \mathrm{OP5}_{\mathrm{sym}}.$$

It only requires equality after passage to the fixed closure invariant. Thus source and return may remain visibly distinct while still forming a closed OP 1–OP 5 pair.

Consequence for the OP reading rule. The theorem therefore replaces raw endpoint comparison by invariant source-return comparison: OP 1 and OP 5 may remain visibly separated in the measurement frame while still carrying one and the same closure-bearing mode, provided the two-pass transport remains class-preserving.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

H7.2 Branchwise refinement of the closed OP1–OP5 pair

Once the OP 1–OP 5 pair has been closed at the invariant level, branchwise effects should be read as secondary refinements of that closed source-return architecture rather than as replacements for it. In particular, branching does not create closure ex nihilo; it only distributes the observable realization of a closure-bearing mode across admissible residual routes.

Let $\mathcal{B}$ be a family of admissible residual branches associated with the fixed route Γ. For each branch $b \in \mathcal{B}$, let

$$T_{\Gamma}^{(b)}, T_{\overline{\Gamma}}^{(b)} : \mathcal{A}_{\mathrm{cl}} \to \mathcal{C}$$

denote the branchwise forward and backward carrier transports, and let

$$\mathrm{OP5}_{\mathrm{fwd}}^{(b)}, \qquad \mathrm{OP5}_{\mathrm{bwd}}^{(b)}$$

be the corresponding branchwise return readouts. Define

$$\mathrm{OP5}_{\mathrm{sym}}^{(b)} := \frac{1}{2}\big(\mathrm{OP5}_{\mathrm{fwd}}^{(b)} + \mathrm{OP5}_{\mathrm{bwd}}^{(b)}\big).$$

We say that a branch b is *closure-admissible* if

$$K_{\mathrm{ncl}}(T_{\Gamma}^{(b)} c) = K_{\mathrm{ncl}}(c) = K_{\mathrm{ncl}}(T_{\overline{\Gamma}}^{(b)} c) \qquad \text{for all admissible } c \in \mathcal{A}_{\mathrm{cl}}.$$

Proposition H7.5 (Branchwise refinement of OP closure)**.** *Assume the OP 1–OP 5 pair is closed in the sense of the previous theorem. If a branch $b \in \mathcal{B}$ is closure-admissible, then*

$$\mathcal{I}(\mathrm{OP1}) = \mathcal{I}(\mathrm{OP5}_{\mathrm{sym}}^{(b)}).$$

Hence every closure-admissible branch refines the same invariantly closed source-return pair.

Proof. Since b is closure-admissible, the branchwise forward and backward transports preserve the same native closure carrier class as the source carrier. By closure-compatible readout, the observable invariant $\mathcal{I}$ therefore takes the same value on OP1, $\mathrm{OP5}_{\mathrm{fwd}}^{(b)}$, and $\mathrm{OP5}_{\mathrm{bwd}}^{(b)}$. Linearity on the admissible return sector then yields

$$\mathcal{I}(\mathrm{OP5}_{\mathrm{sym}}^{(b)}) = \mathcal{I}(\mathrm{OP1}).$$

□

Remark H7.6. *The proposition shows that admissible branching does not introduce a new closure criterion. The closure test remains the same; branchwise analysis only asks which residual routes preserve the already fixed native closure class.*

Corollary H7.7 (Diagnostic status of non-admissible branches)**.** *If a branch fails closure-admissibility, then it does not by itself refute closure of the OP 1–OP 5 pair. It only shows that this particular residual refinement leaves the native closure class.*

Proof. Closure of the OP 1–OP 5 pair is decided at the level of the fixed two-pass source-return architecture. A non-admissible branch records failure of class preservation for that branchwise refinement, not necessarily failure of the underlying closed pair itself. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

H7.3 Observable routing consequence of closure-admissible branches

Once the OP 1–OP 5 pair has been closed at the invariant level and closure-admissible branches have been identified, the remaining question is how these admissible branches contribute to the observable return signal. In the present framework, this contribution belongs to the readout layer and should therefore be modeled as a routed aggregation of branchwise return realizations.

Let $\mathcal{B}_{\mathrm{adm}} \subseteq \mathcal{B}$ denote the set of closure-admissible branches. For each $b \in \mathcal{B}_{\mathrm{adm}}$, let

$$\mathrm{OP5}_{\mathrm{sym}}^{(b)} \in \mathcal{W}$$

be the branchwise symmetrized return readout. Assume that the observable readout space $\mathcal{W}$ carries an admissible additive structure.

Definition H7.8 (Admissible routing weights)**.** *A family of admissible routing weights is a map*

$$w : \mathcal{B}_{\mathrm{adm}} \to [0, \infty).$$

The corresponding admissible routed return readout is

$$\mathrm{OP5}_{\mathrm{rout}} := \sum_{b \in \mathcal{B}_{\mathrm{adm}}} w(b)\, \mathrm{OP5}_{\mathrm{sym}}^{(b)},$$

whenever the sum is well-defined in $\mathcal{W}$.

Proposition H7.9 (Invariant value of the admissible routed return)**.** *Assume that:*

(i) every branch $b \in \mathcal{B}_{\mathrm{adm}}$ is closure-admissible;

(ii) the closure invariant $\mathcal{I}$ is linear on the admissible routed sector;

(iii) the routed sum defining $\mathrm{OP5}_{\mathrm{rout}}$ exists.

Then

$$\mathcal{I}(\mathrm{OP5}_{\mathrm{rout}}) = \Big(\sum_{b \in \mathcal{B}_{\mathrm{adm}}} w(b) \Big) \mathcal{I}(\mathrm{OP1}).$$

Proof. By branchwise closure-admissibility, the previous proposition gives

$$\mathcal{I}\Big(\mathrm{OP5}_{\mathrm{sym}}^{(b)}\Big) = \mathcal{I}(\mathrm{OP1}) \qquad \text{for all } b \in \mathcal{B}_{\mathrm{adm}}.$$

Using linearity of $\mathcal{I}$ on the admissible routed sector, we obtain

$$\mathcal{I}(\mathrm{OP5}_{\mathrm{rout}}) = \mathcal{I}\left(\sum_{b \in \mathcal{B}_{\mathrm{adm}}} w(b)\, \mathrm{OP5}_{\mathrm{sym}}^{(b)} \right) = \sum_{b \in \mathcal{B}_{\mathrm{adm}}} w(b)\, \mathcal{I}\Big(\mathrm{OP5}_{\mathrm{sym}}^{(b)}\Big).$$

Substituting the branchwise closure identity yields

$$\mathcal{I}(\mathrm{OP5}_{\mathrm{rout}}) = \sum_{b \in \mathcal{B}_{\mathrm{adm}}} w(b)\, \mathcal{I}(\mathrm{OP1}) = \Big(\sum_{b \in \mathcal{B}_{\mathrm{adm}}} w(b) \Big) \mathcal{I}(\mathrm{OP1}),$$

as claimed. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Corollary H7.10 (Normalized admissible routing)**.** *If, in addition, the admissible routing weights are normalized so that*

$$\sum_{b\in\mathcal{B}_{\mathrm{adm}}} w(b) = 1,$$

then

$$\mathcal{I}(\mathrm{OP5}_{\mathrm{rout}}) = \mathcal{I}(\mathrm{OP1}).$$

Proof. This is immediate from the proposition. □

Definition H7.11 (Admissible routing defect)**.** *Define the admissible routing defect by*

$$\mathfrak{d}_{\mathrm{rout}} := \mathcal{I}(\mathrm{OP5}_{\mathrm{rout}}) - \mathcal{I}(\mathrm{OP1}).$$

In the normalized case, one has

$$\mathfrak{d}_{\mathrm{rout}} = 0.$$

More generally,

$$\mathfrak{d}_{\mathrm{rout}} = \big(W_{\mathrm{adm}} - 1\big)\mathcal{I}(\mathrm{OP1}), \qquad W_{\mathrm{adm}} := \sum_{b\in\mathcal{B}_{\mathrm{adm}}} w(b).$$

Remark H7.12. *Thus unnormalized routing does not destroy closure content; it rescales its observable aggregate. The distinction between structural closure and observable weight remains intact.*

H7.4 Mixed routed support and the final OP closure diagnostic

Visible routed support need not come entirely from closure-admissible branches. To close the OP analysis, one must therefore separate the admissible routed sector from the remaining visible support.

Let $\mathcal{B}_{\mathrm{vis}}$ be the set of visible branches with routing weights $w(b) \geq 0$, and split it as

$$\mathcal{B}_{\mathrm{vis}} = \mathcal{B}_{\mathrm{adm}} \sqcup \mathcal{B}_{\mathrm{rem}}, \qquad \mathcal{B}_{\mathrm{rem}} := \mathcal{B}_{\mathrm{vis}} \setminus \mathcal{B}_{\mathrm{adm}}.$$

Assume that every visible branch $b \in \mathcal{B}_{\mathrm{vis}}$ carries a symmetrized return representative $\mathrm{OP5}^{(b)}_{\mathrm{sym}} \in \mathcal{W}$ and define the total routed return by

$$\mathrm{OP5}_{\mathrm{tot}} := \sum_{b\in\mathcal{B}_{\mathrm{vis}}} w(b)\,\mathrm{OP5}^{(b)}_{\mathrm{sym}},$$

whenever the sum exists.

Proposition H7.13 (Decomposition of the total routed invariant)**.** *Assume that $\mathcal{I}$ is linear on the visible routed sector. Then*

$$\mathcal{I}(\mathrm{OP5}_{\mathrm{tot}}) = W_{\mathrm{adm}}\,\mathcal{I}(\mathrm{OP1}) + R_{\mathrm{rem}},$$

where

$$W_{\mathrm{adm}} := \sum_{b\in\mathcal{B}_{\mathrm{adm}}} w(b), \qquad R_{\mathrm{rem}} := \sum_{b\in\mathcal{B}_{\mathrm{rem}}} w(b)\,\mathcal{I}\Big(\mathrm{OP5}^{(b)}_{\mathrm{sym}}\Big).$$

Proof. By linearity,

$$\mathcal{I}(\mathrm{OP5}_{\mathrm{tot}}) = \sum_{b\in\mathcal{B}_{\mathrm{adm}}} w(b)\,\mathcal{I}\Big(\mathrm{OP5}^{(b)}_{\mathrm{sym}}\Big) + \sum_{b\in\mathcal{B}_{\mathrm{rem}}} w(b)\,\mathcal{I}\Big(\mathrm{OP5}^{(b)}_{\mathrm{sym}}\Big).$$

The first sum equals $W_{\mathrm{adm}}\,\mathcal{I}(\mathrm{OP1})$ by the admissible-routing proposition. The second sum is exactly R_{rem}. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Corollary H7.14 (Final OP closure mismatch decomposition)**.** *Define the total OP closure mismatch by*

$$\mathfrak{m}_{\mathrm{OP}} := \mathcal{I}(\mathrm{OP5}_{\mathrm{tot}}) - \mathcal{I}(\mathrm{OP1}).$$

Then

$$\mathfrak{m}_{\mathrm{OP}} = \big(W_{\mathrm{adm}} - 1\big)\mathcal{I}(\mathrm{OP1}) + R_{\mathrm{rem}}.$$

Proof. Subtract $\mathcal{I}(\mathrm{OP1})$ from the decomposition in the previous proposition. □

Corollary H7.15 (Exact closure criterion for the visible routed sector)**.** *If*

$$W_{\mathrm{adm}} = 1 \qquad \textit{and} \qquad R_{\mathrm{rem}} = 0,$$

then

$$\mathcal{I}(\mathrm{OP5}_{\mathrm{tot}}) = \mathcal{I}(\mathrm{OP1}).$$

In particular, exact routed OP closure holds whenever the visible routed support is normalized on the admissible sector and carries no residual non-admissible contribution at the invariant level.

Proof. Under the stated conditions, the mismatch formula gives $\mathfrak{m}_{\mathrm{OP}} = 0$. □

Remark H7.16 (Closed form of the OP analysis)**.** *The OP analysis is thereby closed in three layers: native source-return closure first, admissible branch refinement second, and visible routed mismatch only third. Observable source-return disagreement is therefore confined to routing normalization and residual non-admissible support once the structural pair itself has been closed.*

H7.5 Admissible-branch theorem

With the OP 1–OP 5 pair closed, the next question is no longer whether branches appear, but which of them preserve the already fixed native closure architecture. The present theorem turns that distinction into a theorem-level criterion.

Let Γ be a fixed structural route with closed OP 1–OP 5 source-return architecture, and let $\mathcal{B}$ be a family of branchwise residual refinements attached to the OP 3 interface. For each branch $b \in \mathcal{B}$, let

$$T_{\Gamma}^{(b)}, T_{\overline{\Gamma}}^{(b)} : \mathcal{A}_{\mathrm{cl}} \to \mathcal{C}$$

denote the corresponding branchwise forward and backward transports.

Definition H7.17 (Visible support)**.** *A branch $b \in \mathcal{B}$ is called visibly supported if its routed observable weight satisfies $w(b) > 0$.*

Theorem H7.18 (Admissible-branch theorem)**.** *Assume the OP transport theorem. Then the following hold.*

(i) If a branch $b \in \mathcal{B}$ is closure-admissible, then

$$\mathcal{I}(\mathrm{OP1}) = \mathcal{I}(\mathrm{OP5}_{\mathrm{sym}}^{(b)}).$$

(ii) If a branch fails closure-admissibility, then the failure is detected first at the level of native closure-class preservation, not at the level of raw observable mismatch.

(iii) Visible support does not imply closure-admissibility.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(iv) Within the closed OP architecture, branchwise structural relevance is determined first by closure-admissibility and only secondarily by routed visible support.

Proof. Part (i) is Proposition H7.5. Part (ii) is the contrapositive reading of the same proposition together with the native closure-class criterion. Part (iii) follows because visible support is controlled by $w(b)$, whereas admissibility is controlled by preservation of K_{ncl}. Part (iv) is the structural reading of (i)–(iii). □

Corollary H7.19 (Hierarchy of branch diagnostics)**.** *Branchwise diagnostics should be read in the order*

$$\textit{closure-admissibility} \;>\; \textit{branch defect} \;>\; \textit{visible support.}$$

H7.6 OP3 shell/residual theorem

We now localize shell- and residual-sensitive effects at the OP 3 interface itself: they become structurally relevant only when they cease to preserve the native closure class there.

Let

$$c_3 := A_3(A_2(c_\Gamma))$$

be the OP 3 carrier attached to the closed OP architecture. Let $\mathcal{R}$ be the residual parameter space and $\mathcal{S}$ the shell index set. For each $(r, s) \in \mathcal{R} \times \mathcal{S}$, let

$$\mathfrak{T}_{r,s} : \mathcal{C} \to \mathcal{C}$$

be the corresponding shell/residual refinement acting at the OP 3 interface, and define

$$c_3^{(r,s)} := \mathfrak{T}_{r,s}(c_3).$$

Definition H7.20 (OP3 shell-admissibility)**.** *A shell/residual refinement $\mathfrak{T}_{r,s}$ is called OP 3 shell-admissible if*

$$K_{\mathrm{ncl}}(c_3^{(r,s)}) = K_{\mathrm{ncl}}(c_3).$$

Theorem H7.21 (OP3 shell/residual theorem)**.** *Assume the OP transport theorem. Then the following hold.*

(i) If $\mathfrak{T}_{r,s}$ is OP 3 shell-admissible, then

$$\mathcal{I}(\rho_\theta(c_3^{(r,s)})) = \mathcal{I}(\rho_\theta(c_3)) = \mathcal{I}(\mathrm{OP1})$$

for every admissible frame phase θ.

(ii) Shell-consistency defects are first OP 3 admissibility defects, not immediate failures of the whole OP chain.

(iii) Inverse-shell behavior is secondary unless it changes the native closure carrier class.

(iv) Raw shell-sensitive mismatch has no primary structural force unless it survives passage to failure of native closure-class preservation.

Proof. Part (i) follows because $\mathcal{I}$ factors through K_{ncl} and the intermediate invariant bridge identifies $\mathcal{I}(\mathrm{OP3})$ with $\mathcal{I}(\mathrm{OP1})$. Parts (ii)–(iv) are the corresponding structural reading rules once OP 3 is fixed as the residual/shell interface. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Remark H7.22 (Compressed reading of OP 3 after the extracted QAM proof run)**.** *The theorem above remains the native OP statement of the OP 3 shell/residual interface. After the extracted QAM tool pass, however, it no longer needs to be reread only in shell-first language. An OP 3 shell-admissible refinement may now be passed through the shell-certificate tool, the bundled OP 3 closure tool, and the obstruction-stratification tools from the QAM layer. In plain language: OP 3 shell admissibility can now be read either directly at the OP interface or through the compressed certified-pass / localized-obstruction dichotomy of Corollary H2.177.*

Proposition H7.23 (Compressed rereading of the OP 3 shell/residual theorem through extracted QAM tools)**.** *Assume the faithful translation theorem for the admissible OP fragment, Theorem H2.238, the direct kernel constructor of Definition H2.118, and the bridge conditions of Theorem H2.119. Let $\mathfrak{T}_{r,s}$ be an OP 3 shell-admissible refinement. Then the OP 3 shell/residual theorem may be reread in the following compressed form:*

(i) the translated shell passage activates the bundled OP 3 closure tool, so the shell step may be continued kernel-first through one common closure-bearing carrier;

(ii) any first structurally relevant shell defect must already appear as the first localized obstruction in the extracted QAM stratification;

(iii) inverse-shell or raw shell-sensitive visible mismatch remains secondary unless it propagates to failure of the shell certificate package, equivalently to loss of the bundled closure carrier.

Proof. Translate the admissible OP fragment by Theorem H2.238. Because $\mathfrak{T}_{r,s}$ is OP 3 shell-admissible, the corresponding translated passage falls under the shell-certificate side of the first narrow iff package, Theorem H2.166. The bundled OP 3 closure tool, Theorem H2.170, then supplies clause (i): the visible shell passage may be replaced by one common closure-bearing kernel/readout carrier.

Clause (ii) is the negative reading of the same package. If a first structurally relevant shell defect occurs, then by Lemma H2.168 it must already appear as the first localized obstruction stratum. Clause (iii) follows because Lemma H2.169 compresses every residual branch/shell-preserving failure to loss of one common carrier. Hence visible inverse-shell or raw shell mismatch is still secondary unless it survives passage to that compressed obstruction form. □

H7.7 Combined branch-shell transport theorem

The final routed return may now be read as a combined branch-shell aggregation. The theorem below isolates exactly when that visible composite support still preserves the invariant OP closure already established in the main line.

Let $\mathcal{B}_{\mathrm{adm}}$ be the admissible branch set and let

$$\mathcal{F}_{\mathrm{sh}}^{(b)} \subseteq \mathcal{R} \times \mathcal{S}$$

denote the shell-admissible refinement family attached to branch $b \in \mathcal{B}_{\mathrm{adm}}$ at the OP 3 interface. For every admissible branch b and every admissible refinement $(r, s) \in \mathcal{F}_{\mathrm{sh}}^{(b)}$, let

$$\mathrm{OP5}_{\mathrm{sym}}^{(b,r,s)} \in \mathcal{W}$$

denote the corresponding symmetrized return readout.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition H7.24 (Branch-shell routed return)**.** *Let $w(b,r,s) \geq 0$ be a routed weight assigned to each branch-shell admissible realization. Whenever the sum is well-defined in $\mathcal{W}$, define*

$$\mathrm{OP5}_{\text{bs-rout}} := \sum_{b \in \mathcal{B}_{\mathrm{adm}}} \sum_{(r,s) \in \mathcal{F}_{\mathrm{sh}}^{(b)}} w(b,r,s)\, \mathrm{OP5}_{\mathrm{sym}}^{(b,r,s)}.$$

Theorem H7.25 (Combined branch-shell transport theorem)**.** *Assume the OP transport theorem, the admissible-branch theorem, and the OP 3 shell/residual theorem. Assume further that $\mathcal{I}$ is linear on the admissible branch-shell routed sector. Then:*

(i) For every admissible branch $b \in \mathcal{B}_{\mathrm{adm}}$ and every shell-admissible refinement $(r,s) \in \mathcal{F}_{\mathrm{sh}}^{(b)}$,

$$\mathcal{I}(\mathrm{OP5}_{\mathrm{sym}}^{(b,r,s)}) = \mathcal{I}(\mathrm{OP1}).$$

(ii) If the routed weights are normalized so that

$$\sum_{b \in \mathcal{B}_{\mathrm{adm}}} \sum_{(r,s) \in \mathcal{F}_{\mathrm{sh}}^{(b)}} w(b,r,s) = 1,$$

then

$$\mathcal{I}(\mathrm{OP5}_{\text{bs-rout}}) = \mathcal{I}(\mathrm{OP1}).$$

(iii) A visibly composite return assembled from several admissible branches and several shell-admissible refinements may remain structurally closure-preserving at invariant level.

(iv) Failure of closure in the branch-shell routed sector occurs first through loss of branch admissibility or loss of OP 3 shell-admissibility, not through visible multiplicity alone.

Proof. Part (i) combines Theorem H7.18 with Proposition H7.23. Admissible branches preserve the OP invariant, and once the OP 3 shell passage is rerouted through the bundled OP 3 closure tool it preserves that same invariant inside each branch through one common closure-bearing carrier. Part (ii) follows from linearity of $\mathcal{I}$ on the admissible branch-shell routed sector. Parts (iii)–(iv) are the structural reading of (i)–(ii): visible multiplicity alone is not structural non-closure as long as admissible branch passage and the compressed OP 3 shell package remain intact. □

H7.8 First carrier-first OP5 interface

After the combined branch-shell theorem, OP 5 should no longer be read as a final visible endpoint that must be tested from scratch. The first terminal question is whether the symmetrized return readout still belongs to the same native closure-bearing carrier decision already fixed by the source-return theorem and by the admissible branch-shell passage.

Remark H7.26 (Carrier-first reading of OP 5 after branch-shell transport)**.** *The visible OP 5 packet-, phase-, and flicker-diagnostics remain secondary at this stage. Once a branch-shell realization is admissible, OP 5 should first be read as a return-side query against one common closure-bearing carrier; only afterwards should one inspect visible packet-scale deviations.*

Proposition H7.27 (First carrier-first OP 5 interface)**.** *Assume the closure theorem for the OP 1–OP 5 pair, Theorem H7.2, and the combined branch-shell transport theorem, Theorem H7.25. Let (b,r,s) be a branch-shell admissible realization and let*

$$\mathrm{OP5}_{\mathrm{sym}}^{(b,r,s)} \in \mathcal{W}$$

denote the associated symmetrized return readout. Then the terminal OP 5 decision has the following carrier-first form.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(i) ***Certified carrier pass.***

$$\mathcal{I}\Big(\mathrm{OP5}_{\mathrm{sym}}^{(b,r,s)}\Big) = \mathcal{I}(\mathrm{OP1}).$$

Hence OP 5 is to be read as a return-side readout of one common closure-bearing carrier class rather than as an independent closure test.

(ii) ***First localized obstruction.*** *If a terminal structural failure is claimed at OP 5, then its first relevant source lies upstream of raw terminal visibility: it must already arise through loss of branch admissibility, loss of OP 3 shell-admissibility, or loss of the corresponding carrier/readout package before an independent OP 5 failure can be asserted.*

In particular, the terminal OP 5 layer is the first carrier-level decision interface after the branch-shell pass, not a new autonomous closure theorem.

Proof. By Theorem H7.25, every branch-shell admissible realization satisfies

$$\mathcal{I}\Big(\mathrm{OP5}_{\mathrm{sym}}^{(b,r,s)}\Big) = \mathcal{I}(\mathrm{OP1}).$$

By Theorem H7.2, the OP 1 source readout and the symmetrized OP 5 return are already tied to one and the same native closure decision at invariant level. Thus the admissible branch-shell return is not a new endpoint test; it is a return-side reading of that same closure-bearing carrier class. This proves part (i). Part (ii) is the structural reading of the same fact: once the terminal invariant is fixed by the shared carrier decision, any genuinely structural failure visible at the terminal side must have entered earlier, namely through loss of admissibility on the branch-shell side or through failure of the carrier/readout package that supports the branch-shell pass. Therefore OP 5 is the first carrier-level decision interface after the branch-shell passage, not an independent closure theorem. □

Corollary H7.28 (Carrier-first routed OP 5 decision principle)**.** *Assume the hypotheses of Proposition H7.27 and let the branch-shell routed return*

$$\mathrm{OP5}_{\text{bs-rout}}$$

be defined with normalized routed weights. Then:

(i) *the entire routed return is to be read first through one common closure-bearing carrier decision;*

(ii) *visible terminal multiplicity does not by itself create a new OP 5 closure problem;*

(iii) *any structural routed failure must already originate at the first localized obstruction of some branch-shell final rather than at terminal aggregation alone.*

Proof. Normalization together with Theorem H7.25 yields

$$\mathcal{I}(\mathrm{OP5}_{\text{bs-rout}}) = \mathcal{I}(\mathrm{OP1}).$$

Thus the routed return still belongs to the same invariant closure decision as the source readout. The remaining statements are the carrier-first reading of this equality: visible routed multiplicity is secondary, and any structural failure must already arise at the first localized obstruction of one of the contributing branch-shell realizations. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

H7.9 Carrier-first packet/phase-lock/epsilon compatibility layer

After the carrier-first OP 5 interface, the remaining packet-, phase-lock-, and epsilon-trap diagnostics should now be read as a visible support layer for the admissible return side of one common closure-bearing carrier. They do not replace the admissibility theorems or the carrier-first OP 5 decision; rather, they provide a structured non-escape corridor in which visible packet-scale deviations remain subordinate to the same return-side carrier decision.

Definition H7.29 (Packet/phase-lock/epsilon compatible branch-shell channel)**.** *A branch-shell realization (b, r, s) is called* packet/phase-lock/epsilon compatible *if the following hold simultaneously:*

(i) the branch b is closure-admissible;

(ii) the refinement (r, s) is OP 3 shell-admissible;

(iii) the OP 1-side bridge normalization lies inside the contractive epsilon tube of Theorem H3.28;

(iv) the OP 3-side visible shell ratio lies inside the anti-phase epsilon corridor preserved as `thm:working-shell-anti-phase-epsilon-trap` in the historical-witness PDF;

(v) the packet-wave, breathing-envelope, and frame-flicker residuals are treated as visible-support diagnostics rather than as primary closure data.

Remark H7.30. *The first two conditions are structural; the next two are non-escape conditions on the OP 1 entry side and the OP 3 shell side; the final condition places the remaining packet-scale evidence at the observable support layer. This is the natural compressed reading of the current packet/phase/epsilon analysis after OP closure hardening and after the carrier-first OP 5 interface has fixed the terminal decision level.*

Corollary H7.31 (Carrier-first packet/phase-lock/epsilon compatibility)**.** *Assume the combined branch-shell transport theorem together with Proposition H7.27. If a branch-shell realization (b, r, s) is packet/phase-lock/epsilon compatible, then the following hold.*

(i) the visible packet-, breathing-envelope-, and frame-flicker diagnostics are compatible with one and the same invariantly closed source-return channel because they sit on the return side of one common closure-bearing carrier;

(ii) the OP 1 bridge side and the OP 3 shell side both satisfy a non-escape criterion inside their respective local epsilon corridors;

(iii) no visible packet- or phase-lock mismatch obtained inside those corridors is by itself evidence against structural OP closure or against the carrier-first OP 5 decision;

(iv) the first structurally relevant failure can occur only through loss of branch admissibility or loss of OP 3 shell-admissibility.

Proof. By branch admissibility and OP 3 shell-admissibility, the combined branch-shell transport theorem gives

$$\mathcal{I}(\mathrm{OP5}_{\mathrm{sym}}^{(b,r,s)}) = \mathcal{I}(\mathrm{OP1}).$$

By Proposition H7.27, this equality is already the carrier-first terminal OP 5 decision: the admissible return readout belongs to one common closure-bearing carrier and is not a new

© 2026 Marc Brendecke. Preprint — not peer reviewed.

autonomous endpoint test. The OP 1 epsilon-trapped bridge theorem supplies the local non-escape statement on the source-entry side, while the shell anti-phase epsilon-trap supplies the corresponding non-escape statement for the visible shell ratio at the OP 3 interface. The packet-wave, breathing-envelope, and frame-flicker diagnostics are explicitly treated as visible-support diagnostics, so any mismatch they show inside the admissible corridors remains subordinate to that carrier-first decision and to the visible/structural mismatch separation already built into the OP tool layer. Therefore the visible diagnostics remain compatible with the same invariantly closed return-side carrier channel, and the first structurally relevant failure can only arise by leaving the admissible branch-shell sector itself. □

Remark H7.32 (Current mathematical role of the packet/phase/epsilon line)**.** *The present packet/phase-lock/epsilon line is therefore no longer best read as an alternative closure proof or as a second OP 5 test. Its mathematical role is to provide local corridor evidence that the currently observed OP 1 and OP 5 diagnostics can sit inside one admissible return-side carrier channel without forcing a structural mismatch. In that sense, it is the visible non-escape support layer beneath the carrier-first OP 5 decision and the next admissible-branch hardening layer.*

Proposition H7.33 (Carrier-first packet/phase-lock/epsilon screening principle)**.** *Assume the combined branch-shell transport theorem together with Proposition H7.27. Then packet/phase-lock/epsilon compatibility defines a screening criterion for final admissible branch-shell support inside the carrier-first OP 5 architecture rather than a second closure theorem. More precisely, if a realization (b, r, s) is packet/phase-lock/epsilon compatible, then it lies inside the diagnostic preselection corridor of the admissible branch-shell architecture and remains subordinate to the same return-side carrier decision; conversely, loss of packet/phase-lock/epsilon compatibility is not by itself sufficient to prove structural non-closure unless it propagates to loss of branch admissibility or loss of OP 3 shell-admissibility.*

Proof. The first statement is immediate from Corollary H7.31: compatibility places the realization inside a joint non-escape corridor on the OP 1 bridge side and the OP 3 shell side while leaving the invariant closure value unchanged, and Proposition H7.27 then fixes the terminal reading as one common return-side carrier decision. Hence the diagnostic role of the packet/phase/epsilon conditions is to preselect a corridor of final admissible support beneath that carrier-first terminal reading, not to introduce an independent closure criterion.

For the converse direction, failure of packet-, phase-lock-, or epsilon-corridor compatibility may occur either because a visible-support diagnostic leaves its controlled regime or because a structural admissibility condition fails. By the same corollary and by the carrier-first OP 5 interface, only the latter is primary for OP closure: visible mismatch inside or outside the packet-scale corridor is not yet structural non-closure unless it reflects loss of branch admissibility or loss of OP 3 shell-admissibility. Therefore packet/phase-lock/epsilon diagnostics define a screening layer rather than a standalone closure theorem or a second OP 5 endpoint test. □

Remark H7.34. *This screening principle is the cleanest current mathematical reading of the old packet/phase-lock/epsilon diagnostics. They are now neither discarded nor promoted beyond their scope: they act as a visible preselection gate for the admissible branch-shell sector under one carrier-first OP 5 decision, while the actual structural test remains the admissibility machinery of the OP core.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

H7.10 Diagnostic readout hierarchy beneath the carrier-first OP 5 decision

After the carrier-first OP 5 interface and the packet/phase-lock/epsilon screening layer, the remaining visible packet, breathing-envelope, frame-flicker, and helix pictures should be read in a strict order. They are not parallel closure criteria. They are progressively more visible readout surfaces of one and the same carrier decision.

Remark H7.35 (Three diagnostic layers)**.** *The current OP 5-side diagnostic stack is therefore to be read in the following descending order of structural weight.*

(i) ***Carrier-decision layer.*** *The first structural question is whether the return side still belongs to one common closure-bearing carrier, as fixed by Proposition H7.27.*

(ii) ***Screening layer.*** *Packet/phase-lock/epsilon compatibility then determines whether the visible return sits inside an admissible non-escape corridor beneath that carrier decision.*

(iii) ***Visible-shape layer.*** *Packet-wave, breathing-envelope, frame-flicker, and helix pictures come last: they visualize how one admissible or nearly admissible return is seen, but they do not outrank the carrier decision or the screening corridor.*

Lemma H7.36 (First diagnostic-readout hierarchy lemma)**.** *Assume Proposition H7.27 and Proposition H7.33. Let (b, r, s) be a branch-shell realization with associated visible packet-, breathing-envelope-, frame-flicker-, and helix-type diagnostics. Then the current OP-side readout hierarchy has the following form.*

(i) ***Carrier priority.*** *If the realization belongs to one common closure-bearing carrier, then no visible packet-, flicker-, or helix-level multiplicity by itself overrides that carrier decision.*

(ii) ***Screening priority.*** *Packet/phase-lock/epsilon compatibility is the first visible screening layer beneath the carrier decision; loss at the pure visible-shape level is not yet structural loss unless it propagates to loss of this screening layer or to loss of the carrier package itself.*

(iii) ***Visible-shape subordination.*** *The packet-wave, breathing-envelope, frame-flicker, and helix pictures are therefore subordinate readout surfaces. They can localize, rank, and compare visible deviations, but they do not constitute an independent closure theorem and do not outrank the carrier-first OP 5 decision.*

In particular, the mathematically correct order of evidential force is

$$\textit{carrier decision} > \textit{packet/phase/epsilon screening} > \textit{visible flicker/helix picture.}$$

Proof. Proposition H7.27 fixes the first terminal structural question: the return side is read first through one common closure-bearing carrier, and a structural failure must already arise upstream of raw terminal visibility. Proposition H7.33 then places packet/phase-lock/epsilon compatibility strictly beneath that terminal carrier decision as a diagnostic preselection corridor rather than as a second closure theorem. The remaining packet-wave, breathing-envelope, frame-flicker, and helix pictures are by construction even further downstream, because they are explicitly treated as visible-support diagnostics in Definition 3.116 and Corollary H7.31. Hence visible-shape multiplicity may refine the observable profile of a final return, but it cannot outrank the carrier decision and cannot become structural loss unless the screening corridor or the carrier package itself fails. This proves the stated hierarchy. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Corollary H7.37 (Visible flicker/helix subordination principle)**.** *Within the current carrier-first OP 5 architecture, any packet-wave, breathing-envelope, frame-flicker, or helix mismatch is to be read first as visible-shape evidence beneath the carrier decision and beneath the packet/phase-lock/epsilon screening layer. Such a mismatch becomes structurally decisive only if it can be shown to force loss of packet/phase-lock/epsilon compatibility or loss of the common closure-bearing carrier itself.*

Proof. This is the direct operational reading of Lemma H7.36. The visible-shape diagnostics are subordinate readout surfaces; hence they can become structurally decisive only by propagating upward into the screening layer or the carrier layer. □

H7.11 First upstream propagation principle for visible-shape mismatch

Remark H7.38 (Upstream propagation gate)**.** *Under the current carrier-first OP 5 architecture, visible packet-wave, breathing-envelope, frame-flicker, and helix mismatches have no immediate right to overturn the structural reading. They may move upward only by forcing failure of the next superior layer: first the packet/phase-lock/epsilon screening corridor, and only after that the common closure-bearing carrier package itself.*

Lemma H7.39 (First upstream propagation lemma)**.** *Assume Proposition H7.27, Proposition H7.33, Lemma H7.36, and Lemma H2.168. Let (b, r, s) be a branch-shell realization with a visible packet-wave, breathing-envelope, frame-flicker, or helix mismatch. Then the current upstream propagation rule has the following form.*

(i) ***No direct jump.*** *A pure visible-shape mismatch does not by itself propagate directly to structural non-closure or directly overturn the carrier-first OP 5 decision.*

(ii) ***First gate: screening.*** *If the mismatch does* not *force loss of packet/phase-lock/epsilon compatibility, then it remains subordinate visible-shape evidence and the carrier decision is unchanged.*

(iii) ***Second gate: carrier package.*** *If the mismatch* does *force loss of packet/phase-lock/epsilon compatibility while the admissible branch-shell sector and the common closure-bearing carrier package remain intact, then the mismatch propagates only to screening-layer loss and is still not yet a structural OP failure.*

(iv) ***Structural propagation.*** *A visible-shape mismatch becomes structurally decisive only when it can be shown to propagate further to one of the upstream obstruction strata of Lemma H2.168, namely loss of admissible branch passage, loss of shell-admissible refinement, or loss of the common readout/kernel carrier package.*

In particular, the current architecture permits only the ordered propagation chain

$$\begin{aligned} \textit{visible flicker/helix mismatch} &\Longrightarrow \textit{possible screening loss} \\ &\Longrightarrow \textit{possible carrier/package loss} \\ &\Longrightarrow \textit{structural failure.} \end{aligned}$$

No shorter upward propagation is presently licensed.

Proof. Lemma H7.36 already fixes the hierarchy of evidential force: visible-shape diagnostics lie strictly below the packet/phase-lock/epsilon screening layer, which itself lies below the

© 2026 Marc Brendecke. Preprint — not peer reviewed.

carrier-first OP 5 decision. Hence a pure visible mismatch cannot jump directly to structural failure. Proposition H7.33 gives the first gate: if packet/phase-lock/epsilon compatibility is preserved, then the mismatch remains inside the diagnostic corridor and does not alter the carrier reading. Even if that screening corridor is lost, Proposition H7.27 still prevents immediate structural failure unless the common return-side carrier package itself is lost. Finally, Lemma H2.168 identifies the only current upstream structural failure modes: branch-stratum loss, shell-stratum loss, or readout/kernel carrier-package loss. Therefore the only admissible upstream propagation route is the ordered one stated above. □

Corollary H7.40 (Visible mismatch non-propagation default)**.** *Within the current carrier-first OP 5 architecture, every packet-wave, breathing-envelope, frame-flicker, or helix mismatch is to be treated by default as diagnostically local evidence. It may be upgraded only after an explicit witness of screening-layer loss or of upstream carrier/package loss has been exhibited.*

Proof. This is the operational reformulation of Lemma H7.39. In the absence of an explicit upward propagation witness, visible-shape mismatch remains subordinate local evidence. □

H7.12 First diagnostic witness principle for visible-shape mismatch

Remark H7.41 (Diagnostic witness discipline)**.** *Under the current carrier-first OP 5 architecture, visible packet-wave, breathing-envelope, frame-flicker, and helix mismatches count as admissible upstream evidence only when they come equipped with a witness that is strong enough to certify failure of the next superior layer. Pure visual irregularity, even if persistent or large, is not yet such a witness.*

Lemma H7.42 (First diagnostic witness lemma)**.** *Assume Proposition H7.33, Proposition H7.27, Lemma H7.36, and Lemma H7.39. Let (b, r, s) be a branch-shell realization with a visible packet-wave, breathing-envelope, frame-flicker, or helix mismatch. Then the current witness discipline has the following form.*

(i) ***Screening witness threshold.*** *A visible mismatch is a legitimate witness of screening-layer loss only if it can be shown to force failure of packet/phase-lock/epsilon compatibility; otherwise it remains diagnostically local evidence.*

(ii) ***Carrier witness threshold.*** *A visible mismatch is a legitimate witness of carrier/package loss only if it can be shown to force, beyond screening loss, failure of the common closure-bearing carrier package itself.*

(iii) ***No witness by appearance alone.*** *Mere amplitude, persistence, oscillatory complexity, geometric deformation, or helix/flicker asymmetry of the visible mismatch is not by itself an admissible witness of screening-layer loss or of carrier/package loss.*

(iv) ***Admissible witness route.*** *Every currently admissible visible witness must therefore factor through one of the two certified upstream gates of Lemma H7.39: first explicit screening loss, and only then, if present, explicit carrier/package loss.*

Hence the present architecture admits visible mismatch only as witness-bearing *evidence and not as a free structural verdict.*

Proof. Proposition H7.33 defines the first superior layer above raw visible-shape mismatch. Therefore a visible mismatch can witness screening-layer loss only by forcing failure of packet/phase-lock/epsilon compatibility itself; if compatibility persists, the mismatch remains

© 2026 Marc Brendecke. Preprint — not peer reviewed.

inside the diagnostic corridor and cannot count as screening evidence. Proposition H7.27 defines the next superior layer: even after screening loss, the structural carrier reading remains unchanged unless the common closure-bearing carrier package is lost. Thus a visible mismatch can witness carrier/package loss only by forcing precisely that second failure as well. Lemma H7.36 excludes any direct elevation of raw visible-shape features to a superior structural layer, and Lemma H7.39 identifies the only currently licensed upward route as the ordered screening-to-carrier route. Consequently amplitude, persistence, or geometric shape alone are not admissible witnesses; only a mismatch that certifiably drives one of those gates may count as an upstream witness. □

Corollary H7.43 (Visible witness admissibility rule)**.** *Within the current carrier-first OP 5 architecture, every visible packet-wave, breathing-envelope, frame-flicker, or helix mismatch must be classified first as either*

(a) a merely local diagnostic irregularity,

(b) a certified witness of screening-layer loss,

(c) or a certified witness of carrier/package loss.

No fourth class of immediately structural visible witness is presently admitted.

Proof. This is the direct classification consequence of Lemma H7.42. A visible mismatch is either not strong enough to force a superior-layer failure, or it forces screening loss, or it forces carrier/package loss after screening loss. By the witness discipline and by the upstream propagation rule, there is no currently licensed direct jump from raw visible appearance to immediate structural verdict. □

H7.13 Screening-witness compression beneath the carrier-first OP 5 decision

Remark H7.44 (Compression of visible witness types)**.** *Once a visible packet-wave, breathing-envelope, frame-flicker, or helix mismatch has crossed the witness threshold of Lemma H7.42, the present architecture no longer needs to keep its visible shape as a separate structural species at the screening level. Unless the same witness already certifies carrier/package loss, its upstream content compresses to one common statement: the packet/phase-lock/epsilon screening corridor has failed.*

Lemma H7.45 (Screening-witness compression lemma)**.** *Assume Proposition H7.33, Proposition H7.27, Lemma H7.42, and Corollary H7.43. Let (b, r, s) be a branch-shell realization with a visible packet-wave, breathing-envelope, frame-flicker, or helix mismatch that is a certified witness of screening-layer loss but not yet a certified witness of carrier/package loss. Then the upstream structural content of that witness compresses to a single screening-loss form, namely the failure of packet/phase-lock/epsilon compatibility. More precisely:*

(i) ***Visible-shape indifference at screening level.*** *Whether the witness is first expressed through packet-wave mismatch, breathing-envelope drift, frame-flicker asymmetry, or helix deformation is irrelevant for its screening-level structural role.*

(ii) ***Unique compressed content.*** *The only currently licensed upstream content of such a witness is that the packet/phase-lock/epsilon screening corridor fails.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(iii) ***No parallel screening species.*** *There is presently no second independent screening-loss class attached specifically to packet-wave, envelope, flicker, or helix shape once the witness has been admitted.*

(iv) ***Carrier exception.*** *A visible witness escapes this compression only when it upgrades further to a certified witness of carrier/package loss.*

Hence, below the carrier threshold, all currently admissible visible screening witnesses collapse to one common screening-loss normal form.

Proof. By Lemma H7.42, a visible mismatch is admitted upward only when it certifiably forces failure of the next superior layer. Proposition H7.33 identifies that next superior layer uniquely as packet/phase-lock/epsilon compatibility. Therefore, once a visible mismatch qualifies as a witness of screening-layer loss, its upstream meaning is already fixed: it testifies precisely to failure of that screening corridor. Corollary H7.43 allows no additional currently licensed class of screening witness beyond that certified screening loss or the stronger carrier/package loss. Proposition H7.27 then separates the only exception: a witness that also forces loss of the common closure-bearing carrier package belongs to the higher carrier class instead of remaining in the compressed screening class. Thus all admitted visible witnesses that stop below the carrier threshold share one and the same screening-loss content, independently of the visible shape through which they first appeared. □

Corollary H7.46 (Screening-witness normal form)**.** *Within the current carrier-first OP 5 architecture, every admitted visible witness that does not yet certify carrier/package loss may be cited in proofs simply as a witness of* screening-loss normal form. *Its packet-wave, breathing-envelope, frame-flicker, or helix realization is diagnostically informative, but not structurally independent at the screening level.*

Proof. This is the direct proof-usage reformulation of Lemma H7.45. Once admitted and still below the carrier threshold, the witness has exactly one currently licensed upstream content: loss of the packet/phase-lock/epsilon screening corridor. □

H7.14 Carrier-witness separation beneath compressed screening loss

Remark H7.47 (Two witness grades after screening compression)**.** *After Lemma H7.45, every admitted visible witness sits in one of two remaining grades. Either it stays below the carrier threshold and is therefore only a witness of screening-loss normal form, or it upgrades further and certifies that the common return-side carrier package itself is no longer available. The present architecture does not license a mixed intermediate status between those two grades.*

Lemma H7.48 (Carrier-witness separation lemma)**.** *Assume Proposition H7.33, Proposition H7.27, Lemma H7.42, Lemma H7.45, and Corollary H7.46. Let* (b, r, s) *be a branch-shell realization with an admitted visible witness. Then exactly one of the following two cases occurs:*

(i) ***Screening-only witness.*** *The witness compresses to screening-loss normal form, while the common closure-bearing return-side carrier package remains available. In this case the witness certifies only screening loss and is not an admissible witness of carrier/package loss.*

(ii) ***Carrier-upgraded witness.*** *The witness forces loss of the common closure-bearing return-side carrier package. In this case it is an admissible witness of carrier/package loss and no longer belongs merely to the screening-loss normal form.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Moreover, there is no third currently licensed status in which an admitted visible witness would simultaneously remain only screening-level and yet already count as a carrier/package witness.

Proof. By Lemma H7.42, every admitted visible witness must force loss of some superior layer. Lemma H7.45 and Corollary H7.46 show that, below the carrier threshold, all such witnesses have one and the same upstream content: failure of packet/phase-lock/epsilon compatibility. Proposition H7.27 identifies the only stronger alternative, namely loss of the common closure-bearing return-side carrier package itself. Hence an admitted visible witness either stops at the compressed screening-loss level or upgrades to carrier/package loss. These two statuses are mutually exclusive because the first expressly retains the common carrier package, whereas the second destroys it. Therefore exactly one of the two cases above occurs, and no mixed intermediate category is presently licensed. □

Corollary H7.49 (Carrier-witness upgrade criterion)**.** *Within the current carrier-first OP 5 architecture, an admitted visible witness upgrades from screening-loss normal form to a genuine carrier/package witness if and only if its upstream content cannot be realized through any common closure-bearing return-side carrier package. Equivalently, as long as such a common carrier remains available, the witness must be read only as screening-loss normal form.*

Proof. The forward direction is exactly case (ii) of Lemma H7.48. The converse direction is case (i): if a common closure-bearing return-side carrier package still exists, then the witness has not crossed the carrier threshold and therefore remains only a screening witness in normal form. □

H7.15 Witness-to-carrier escalation beyond screening compression

Remark H7.50 (Escalation gate after witness separation)**.** *After Lemma H7.48, a visible witness reaches carrier grade exactly when its admitted upstream content can no longer be absorbed by screening-loss normal form alone. In the current architecture there is therefore only one licensed upgrade mechanism: every common closure-bearing return-side carrier package compatible with the compressed screening witness fails, so the witness is forced out of screening-only status and into carrier/package loss.*

Lemma H7.51 (Witness-to-carrier escalation lemma)**.** *Assume Proposition H7.27, Lemma H7.42, Lemma H7.45, Lemma H7.48, and Corollary H7.49. Let (b, r, s) be a branch-shell realization with an admitted visible witness w. Then the following are equivalent:*

(i) w upgrades from screening-loss normal form to a genuine carrier/package witness;

(ii) there exists no common closure-bearing return-side carrier package through which the upstream content of w can still be realized while screening-loss normal form remains the only failed superior layer;

(iii) the upstream content of w forces loss of the common closure-bearing carrier package rather than mere loss of packet/phase-lock/epsilon compatibility.

In particular, escalation occurs exactly when compressed screening loss is no longer a sufficient upstream normal form for the admitted witness.

Proof. By Lemma H7.48, an admitted visible witness has only two currently licensed grades: screening-only or carrier-upgraded. Corollary H7.49 identifies the upgrade threshold precisely: upgrade occurs if and only if no common closure-bearing return-side carrier package remains

© 2026 Marc Brendecke. Preprint — not peer reviewed.

available. Lemma H7.45 shows that, below that threshold, the witness has exactly one upstream content, namely screening-loss normal form. Hence a witness stays screening-only exactly as long as its upstream content is still realizable through some common carrier package with only packet/phase-lock/epsilon compatibility lost. Once that becomes impossible, the witness can no longer remain in screening-loss normal form alone and must instead force carrier/package loss. This proves the equivalence of (i), (ii), and (iii), and therefore the final escalation claim. □

Corollary H7.52 (Escalation threshold for admitted visible witnesses)**.** *Within the current carrier-first OP 5 architecture, an admitted visible witness remains screening-only as long as its upstream content is representable by screening-loss normal form over some common closure-bearing return-side carrier package. It becomes carrier-upgraded precisely at the first point where such a carrier-supported realization ceases to exist.*

Proof. This is the proof-usage reformulation of Lemma H7.51. The existence of a common carrier package keeps the witness below the carrier threshold, while the first failure of every such carrier-supported realization marks the unique licensed escalation point. □

H7.16 Carrier-stable diagnostic collapse beneath the witness escalation threshold

Remark H7.53 (Collapse of visible diagnostic species over a stable carrier)**.** *Once an admitted visible mismatch stays below the escalation threshold of Corollary H7.52, the current carrier-first OP 5 architecture no longer attributes independent structural rank to its packet-wave, breathing-envelope, frame-flicker, or helix presentation. Those visible realizations remain diagnostically useful, but over one and the same stable closure-bearing return-side carrier they collapse to a single subordinate screening-level phenomenon.*

Theorem H7.54 (Carrier-stable diagnostic collapse theorem)**.** *Assume Proposition H7.27, Proposition H7.33, Lemma H7.36, Lemma H7.42, Lemma H7.45, Lemma H7.48, and Lemma H7.51. Let (b, r, s) be a branch-shell realization with an admitted visible mismatch w of packet-wave, breathing-envelope, frame-flicker, or helix type. If w remains below carrier escalation, equivalently if there exists a common closure-bearing return-side carrier package through which the upstream content of w is still realized as screening-loss normal form, then the following collapse holds:*

(i) ***One subordinate structural role.*** *The visible realization of w has exactly one current structural role, namely to witness subordinate screening-loss normal form beneath a still-stable carrier decision.*

(ii) ***No independent visible closure species.*** *Packet-wave, breathing-envelope, frame-flicker, and helix mismatch do not define distinct closure-relevant species as long as the common carrier package remains available.*

(iii) ***Carrier dominance.*** *Any closure-relevant distinction between those visible realizations collapses over the stable carrier and may reappear only if escalation to carrier/package loss occurs.*

(iv) ***Readout subordination.*** *The visible mismatch remains a downstream readout of one common carrier-supported screening loss and therefore cannot outrank the carrier decision itself.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Hence, below the escalation threshold, all currently admitted visible diagnostic forms collapse to one carrier-stable subordinate diagnostic class.

Proof. By Lemma H7.36, visible flicker/helix-type structure is subordinate to packet/phase/epsilon screening, which is itself subordinate to the carrier-first OP 5 decision. Lemma H7.42 allows visible mismatch to count upstream only when it certifies loss of a superior layer. Lemma H7.45 then shows that, below carrier loss, every admitted visible witness has exactly one upstream content: screening-loss normal form. Lemma H7.48 and Corollary H7.52 identify the present hypothesis of a still available common closure-bearing return-side carrier package as precisely the condition that the witness has not escalated to carrier/package loss. Therefore, while that common carrier package remains available, packet-wave, envelope, flicker, and helix mismatch cannot persist as distinct closure-relevant species. They all collapse to the single role of subordinate diagnostic realization of screening-loss normal form over one stable carrier. Proposition H7.27 guarantees that any stronger distinction would already belong to carrier/package loss rather than to the pre-escalation regime. This proves (i)–(iv) and the final collapse claim. □

Corollary H7.55 (Carrier-stable visible normal form)**.** *Within the current carrier-first OP 5 architecture, every admitted visible mismatch that stays below carrier escalation may be cited simply as belonging to one* carrier-stable visible normal form. *Its packet-wave, breathing-envelope, frame-flicker, or helix appearance remains diagnostically informative, but not structurally independent while the common carrier package persists.*

Proof. This is the proof-usage reformulation of Theorem H7.54. The theorem states exactly that all admitted visible mismatches below escalation collapse over one stable carrier to a single subordinate diagnostic class. □

H7.17 Bundled OP 5 diagnostic theorem

Remark H7.56 (One bundled positive/negative OP 5 diagnostic decision)**.** *After the carrier-first OP 5 interface, the current diagnostic layer is now strong enough to be cited as one bundled decision principle. Either the visible material remains subordinate to a stable carrier and therefore collapses to a screening-level normal form, or it upgrades through an admitted witness to carrier/package loss. There is no third currently admitted diagnostic rank between those two bundled outcomes.*

Theorem H7.57 (Bundled OP 5 diagnostic theorem)**.** *Assume Proposition H7.27, Proposition H7.33, Lemma H7.36, Lemma H7.39, Lemma H7.42, Lemma H7.45, Lemma H7.48, Lemma H7.51, and Theorem H7.54. Let (b, r, s) be a routed branch-shell realization together with an admitted visible mismatch w of packet-wave, breathing-envelope, frame-flicker, or helix type. Then exactly one of the following bundled OP 5 diagnostic outcomes holds:*

(i) ***Carrier-stable diagnostic collapse.*** *The mismatch w remains below escalation, is carried by one common closure-bearing return-side carrier, and therefore collapses to a subordinate screening-level visible normal form beneath the carrier-first OP 5 decision.*

(ii) ***Carrier-upgraded diagnostic obstruction.*** *The mismatch w cannot be realized over any common closure-bearing return-side carrier as mere screening-loss normal form and therefore upgrades to carrier/package loss. In that case w is not a merely visible irregularity but an admitted carrier witness for routed OP 5 failure.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Moreover, case (i) *excludes case* (ii)*, and case* (ii) *excludes case* (i)*. Hence the current OP 5 diagnostic layer is completely bundled into the dichotomy*

carrier-stable visible collapse versus carrier-upgraded diagnostic obstruction.

Proof. Lemma H7.36 and Lemma H7.39 force every admitted visible mismatch to move upward only through the packet/phase/epsilon screening layer before it can acquire carrier relevance. Lemma H7.42 then identifies which visible mismatches may count upstream at all. Lemma H7.45 compresses every such admitted visible witness below carrier loss to one screening-loss normal form, while Lemma H7.48 and Lemma H7.51 separate precisely the screening-only regime from the carrier-upgraded regime. Theorem H7.54 describes the entire positive side of the first regime: once a common closure-bearing return-side carrier remains available, all admitted visible species collapse to one subordinate visible normal form beneath the carrier-first OP 5 decision. Conversely, when no such common carrier remains available, Corollary H7.49 upgrades the admitted visible witness to carrier/package loss. These two possibilities are mutually exclusive by Lemma H7.48 and jointly exhaustive by the escalation dichotomy. This proves the bundled theorem. □

Corollary H7.58 (Bundled routed OP 5 diagnostic use rule)**.** *In later proof steps it is sufficient to cite Theorem H7.57 instead of separately unpacking packet-wave, breathing-envelope, frame-flicker, and helix diagnostics. One may simply ask whether the currently admitted visible data still collapse over a stable carrier or have already upgraded to carrier/package obstruction.*

Proof. This is the proof-usage reformulation of Theorem H7.57. The theorem already packages the visible diagnostic layer into exactly the two current routed OP 5 outcomes. □

H7.18 First OP 3 to OP 5 bundled transfer theorem

Remark H7.59 (Transfer of bundled OP 3 closure into the OP 5 diagnostic layer)**.** *The current architecture no longer treats OP 3 closure and OP 5 diagnosis as two independent theorem blocks. Once the OP 3 shell passage has been certified and rerouted through one common closure-bearing carrier, the routed OP 5 side inherits a downstream decision of that same carrier package. The only remaining terminal question is whether the visible return side still collapses beneath that package or whether it upgrades to a carrier-level obstruction. Thus OP 5 is presently read as the bundled downstream transfer image of the OP 3/QAM closure package.*

Theorem H7.60 (First OP 3 to OP 5 bundled transfer theorem)**.** *Assume Theorem H2.170, Theorem H2.176, Theorem H7.25, Proposition H7.27, and Theorem H7.57. Let (b, r, s) be a routed branch-shell realization whose OP 3 shell passage is translated into the extracted QAM layer, and let w be an admitted visible mismatch on the routed OP 5 return side. Then the present bundled OP 3→OP 5 transfer has exactly two outcomes:*

(i) ***Transferred carrier-stable pass.*** *The translated OP 3 shell passage belongs to the certified side of the bundled OP 3 closure package, the routed return stays inside the same common closure-bearing carrier, and the admitted visible mismatch w collapses beneath that carrier to the subordinate OP 5 visible normal form of Theorem H7.57.*

(ii) ***Transferred localized obstruction.*** *The transfer fails at the first licensed obstruction layer: either the OP 3/QAM side already reaches the first localized shell obstruction of the extracted OP 3 proof run, or the routed return upgrades to a carrier-level OP 5 diagnostic obstruction. In particular, no third autonomous terminal OP 5 obstruction is presently licensed downstream of an otherwise intact bundled OP 3 carrier package.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Hence the current routed OP 5 diagnostic layer is the bundled downstream transfer of the OP 3/QAM closure package, not an additional independent closure theory.

Proof. Theorem H2.176 already packages the translated OP 3 shell passage into exactly two upstream outcomes: certified passage through the bundled OP 3 closure tool layer or the first localized obstruction. If the certified side holds, then Theorem H2.170 identifies one common closure-bearing carrier/readout package for the shell passage. Theorem H7.25 transports that same certified branch-shell realization to the routed return side without changing the invariant closure class, and Proposition H7.27 forces the terminal OP 5 reading to be carrier-first rather than a new autonomous endpoint test. Theorem H7.57 then completes the positive branch: as long as the routed return remains over that same common carrier, every admitted visible mismatch collapses to the subordinate carrier-stable OP 5 visible normal form. This proves case (i).

If the transfer does not remain on that certified branch, then the failure must appear at the first licensed obstruction layer. Upstream, Theorem H2.176 and the extracted OP 3/QAM obstruction package localize the first obstruction already at branch, shell, or carrier/readout level. Downstream, if the routed return cannot be realized over any common closure-bearing carrier, then Theorem H7.57 upgrades the admitted visible mismatch to a carrier-level OP 5 diagnostic obstruction. Because Proposition H7.27 forbids an independent terminal closure test downstream of an otherwise intact carrier package, there is no currently licensed third obstruction type appearing only at the raw terminal visible layer. This proves case (ii) and the final bundled transfer reading. □

Corollary H7.61 (Bundled OP 3→OP 5 proof-use rule)**.** *In later proof steps it is sufficient to cite Theorem H7.60 when a routed branch-shell realization must be carried from the extracted OP 3/QAM closure package into the OP 5 diagnostic layer. One may simply ask whether the realization stays on the transferred carrier-stable branch or reaches the first licensed transferred obstruction.*

Proof. This is the proof-usage reformulation of Theorem H7.60. The theorem already packages the current OP 3/QAM closure side and the current OP 5 diagnostic side into one bundled transfer decision. □

H7.19 First OP 3 to OP 5 compressed proof interface

Remark H7.62 (How older routed return arguments should now be compressed)**.** *The current transfer architecture is now strong enough that later routed return arguments need not separately reopen the combined branch-shell theorem, the carrier-first OP 5 interface, and the bundled OP 5 diagnostic layer. Once the routed realization has entered the bundled OP 3→OP 5 transfer channel, the proof may proceed entirely through one compressed dichotomy: transferred carrier-stable pass versus transferred localized obstruction. This local compressed interface is now itself subordinate to the newer global rule: as soon as the argument has also entered the bundled current channel and no shell-sensitive or witness-grade claim is being asked, the proof should move on to the extracted global tools and remain global-first unless a genuine local reopening is needed.*

Proposition H7.63 (First OP 3 to OP 5 compressed proof interface)**.** *Assume Theorem H7.60. Let (b, r, s) be a routed branch-shell realization whose OP 3 shell passage has been translated into the extracted QAM layer, and let w be any admitted visible routed return witness on the OP 5 side. Then every later proof step that needs to carry this realization through the current routed return architecture may be compressed to the following interface.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(i) ***Transferred carrier-stable pass.*** *If Theorem H7.60 places $(b, r, s; w)$ on the transferred carrier-stable branch, then all downstream OP 5 packet/phase/flicker/helix material is read only as subordinate visible realization beneath one common closure-bearing carrier and may be replaced by the carrier-stable visible normal form.*

(ii) ***Transferred localized obstruction.*** *If Theorem H7.60 places $(b, r, s; w)$ on the transferred obstruction branch, then the proof may stop at the first licensed obstruction already identified there: either the upstream OP 3/QAM obstruction or the downstream carrier-upgraded OP 5 obstruction.*

(iii) ***No third terminal proof branch.*** *No additional raw terminal visible branch is presently licensed after this compression. In particular, later proofs do not need a separate fourth case for packet-wave, breathing-envelope, frame-flicker, or helix mismatch once the bundled transfer theorem has already been invoked.*

Hence the current routed return architecture may be used through one compressed OP 3→OP 5 proof interface.

Proof. This is the direct proof-usage reading of Theorem H7.60. The theorem already packages the routed return into exactly two licensed transfer outcomes. On the positive side, the admitted visible material collapses beneath one common carrier and therefore needs no further structural branching. On the negative side, the first licensed obstruction is already localized either upstream in the OP 3/QAM package or downstream at the carrier-upgraded OP 5 level. Because Theorem H7.60 excludes an autonomous raw terminal OP 5 obstruction beneath an otherwise intact carrier package, no third terminal proof branch remains. □

Corollary H7.64 (Compressed rereading of routed return arguments through the bundled transfer)**.** *In later arguments it is sufficient to cite Proposition H7.63 rather than separately reopening Theorem H7.25, Proposition H7.27, and Theorem H7.57. One may simply ask whether the current routed realization stays on the transferred carrier-stable branch or reaches the first licensed transferred obstruction.*

Proof. This is the condensed proof-usage form of Proposition H7.63. The proposition already compresses the presently licensed routed return architecture into one reusable interface. □

H7.20 First routed return normal form theorem

Remark H7.65 (Carrier-stable versus obstructed routed return normal form)**.** *After the compressed OP 3→OP 5 proof interface has been invoked, the current routed return architecture no longer needs a separate case distinction for packet-wave, breathing-envelope, frame-flicker, or helix appearance. At the present level of the theory every admitted routed return is read only up to one of two bundled normal forms: a carrier-stable routed return normal form or an obstructed routed return normal form.*

Theorem H7.66 (First routed return normal form theorem)**.** *Assume Proposition H7.63. Let (b, r, s) be a routed branch-shell realization whose OP 3 shell passage has been translated into the extracted QAM layer, and let w be any admitted visible routed return witness on the OP 5 side. Then the current routed return architecture classifies $(b, r, s; w)$, up to bundled transfer, into exactly one of the following two mutually exclusive routed return normal forms.*

(i) ***Carrier-stable routed return normal form.*** *The routed return remains on the transferred carrier-stable branch of Proposition H7.63. Equivalently, there exists one common closure-bearing return-side carrier through which the routed realization is read, and every*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

admitted visible mismatch collapses beneath that carrier to the carrier-stable visible normal form. In this case the routed return contributes no additional structural branching beyond the bundled OP 3→OP 5 transfer itself.

(ii) ***Obstructed routed return normal form.*** *The routed return lies on the transferred localized-obstruction branch of Proposition H7.63. Equivalently, no common closure-bearing routed return carrier supports the realization as mere screening-level visible loss, and the first licensed failure is already localized either upstream at the OP 3/QAM obstruction layer or downstream at the carrier-upgraded OP 5 obstruction layer.*

Hence every currently admitted routed return is read, up to the bundled OP 3→OP 5 transfer, by one bundled routed return normal form and not by any third autonomous terminal visible class.

Proof. Proposition H7.63 already compresses the present routed return architecture into exactly two licensed proof branches: transferred carrier-stable pass and transferred localized obstruction. On the positive branch, Theorem H7.57 and the carrier-stable collapse package imply that every admitted visible return witness becomes subordinate to one common closure-bearing routed return carrier and therefore yields the carrier-stable routed return normal form. On the negative branch, the same compressed interface localizes the first licensed failure either upstream in the extracted OP 3/QAM obstruction package or downstream at the carrier-upgraded OP 5 obstruction layer; by construction this is the obstructed routed return normal form. Because Proposition H7.63 excludes a third raw terminal visible branch after compression, these two forms are exhaustive and mutually exclusive. □

Corollary H7.67 (Routed return normal form use rule)**.** *In later proofs it is sufficient to cite Theorem H7.66 once a routed realization has already entered the bundled OP 3→OP 5 transfer channel. One may then work only with the dichotomy* carrier-stable routed return normal form versus obstructed routed return normal form.

Proof. This is the proof-usage reformulation of Theorem H7.66. The theorem already packages the presently licensed routed return architecture into one reusable normal-form dichotomy. □

H7.21 First global pass/obstruction normal form

Remark H7.68 (Current global normal form of the admissible run)**.** *The present line now carries enough bundled structure that the currently admitted run*

$$\begin{aligned} &\textit{OP1 source} \\ &\quad\to\ \textit{branch admissibility} \\ &\quad\to\ \textit{OP3/QAM closure package} \\ &\quad\to\ \textit{carrier-first OP5 layer} \end{aligned}$$

no longer needs to be read as a chain of loosely connected local tests. After the routed return normal form has been established, the full currently admitted run is read only up to one global dichotomy: a global pass normal form or a global obstruction normal form.

Theorem H7.69 (First global pass/obstruction normal form)**.** *Assume Theorem H7.66. Let (b, r, s) be an admissible branch-shell realization issued from the current OP1 source side and translated through the extracted OP3/QAM layer into the routed return architecture, and let w be any admitted visible routed return witness on the OP5 side. Then the whole currently admitted run from OP1 through branch, OP3/QAM, and OP5 is classified, up to the present bundled architecture, into exactly one of the following two mutually exclusive global normal forms.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(i) ***Global pass normal form.*** *The realization lies on the carrier-stable routed return normal form of Theorem H7.66. Equivalently, the run remains supported by one bundled closure-bearing carrier package from the present OP1/branch entrance through the OP3/QAM shell package and into the routed return side, while all visible OP5 material collapses beneath that carrier to the carrier-stable visible normal form. In this case the current run carries no further licensed structural branch beyond the bundled pass reading.*

(ii) ***Global obstruction normal form.*** *The realization lies on the obstructed routed return normal form of Theorem H7.66. Equivalently, the first licensed failure of the current run is already localized either upstream in the OP3/QAM obstruction package or downstream at the carrier-upgraded OP5 obstruction layer, and no autonomous third terminal visible class is currently licensed outside that bundled obstruction reading.*

Hence the whole currently admitted run is read, at the present stage of the theory, by one global pass/obstruction normal form and not by any third independent terminal morphology.

Proof. Theorem H7.66 already classifies every currently admitted routed return into exactly two bundled normal forms. On the positive side, the carrier-stable routed return normal form means that the run has remained on the transferred carrier-stable branch from the OP3/QAM closure package into the routed return side; consequently the whole present run remains supported by one bundled closure-bearing carrier package, and all visible OP5 diagnostics are subordinate to the carrier-stable visible normal form. This is precisely the global pass normal form. On the negative side, the obstructed routed return normal form already states that the first licensed failure is localized either upstream in the OP3/QAM obstruction package or downstream at the carrier-upgraded OP5 obstruction layer; this is exactly the global obstruction normal form. Because Theorem H7.66 excludes a third autonomous terminal visible class, the two global forms are exhaustive and mutually exclusive. □

Corollary H7.70 (Global pass/obstruction use rule)**.** *In later proofs it is sufficient to cite Theorem H7.69 once the current admissible run has entered the bundled OP3/QAM-to-OP5 channel. One may then work only with the dichotomy* global pass normal form versus global obstruction normal form.

Proof. This is the proof-usage reformulation of Theorem H7.69. The theorem already compresses the currently admitted run from OP1 through OP5 into one reusable bundled normal-form dichotomy. □

Remark H7.71 (Status of the bundled OP 3→OP 5/global chain)**.** *The present local-to-global chain should now also be read in two levels of proof strength. The* active bundled interface *consists of the first OP 3→OP 5 bundled transfer theorem, the compressed OP 3→OP 5 proof interface, the routed return normal form, and the global pass/obstruction normal form together with their direct proof-use corollaries. These statements already give one self-contained current channel from translated OP 3 certification to a bundled global pass/obstruction decision. By contrast, the finer witness-grade hierarchy beneath the carrier-first OP 5 layer — screening-loss normal form, carrier-witness upgrade, escalation thresholds, and the related witness-compression remarks — remains a* local refinement layer*: it must be reopened only when a later argument genuinely depends on witness species, witness grade, escalation, or screening-versus-carrier distinctions that are invisible at the bundled transfer/global level.*

Proposition H7.72 (Citation and reopening rule for the bundled OP 3→OP 5/global chain)**.** *Let a later argument concern a run already translated from the OP 3/QAM shell package into the routed OP 5 side. Then the following citation discipline is currently licensed.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(i) If the argument needs only the bundled transfer/global dichotomy, it may cite Theorem H7.60, Proposition H7.63, Theorem H7.66, and Theorem H7.69 without reopening the finer witness hierarchy.

(ii) If the argument further needs shell-level certificate or shell-level obstruction data, it must reopen the completed OP 3/QAM block rather than the entire older OP 1–OP 5 line.

(iii) If the argument further needs witness grade, witness escalation, screening-witness compression, or carrier-witness separation, it must reopen the carrier-first OP 5 witness layer rather than treating the bundled global dichotomy as sufficient.

(iv) No later proof may insert a third autonomous raw terminal visible branch after the bundled transfer/global interface has already been invoked.

Proof. Clause (i) is exactly the compressed reading already supplied by Theorem H7.60, Proposition H7.63, Theorem H7.66, and Theorem H7.69. Clause (ii) is the existing shell-sensitive fallback discipline of the completed OP 3/QAM block. Clause (iii) is the corresponding witness-sensitive fallback discipline of the carrier-first OP 5 layer, where screening-loss normal form, upgrade, escalation, and separation are defined. Clause (iv) is the bundled exclusion already built into the transfer theorem, the compressed proof interface, the routed return normal form, and the global pass/obstruction theorem: once those interfaces are active, no additional autonomous raw terminal visible branch is currently licensed. □

Corollary H7.73 (Current sufficiency of the bundled OP 3→OP 5/global chain)**.** *Assume a current argument starts from a translated OP 3 shell passage and asks only whether the run reaches a bundled pass reading or a bundled first licensed obstruction. Then the present document already contains a self-contained proof interface for that question: one may pass from the completed OP 3/QAM block through the bundled OP 3→OP 5 transfer, the routed return normal form, and finally the global pass/obstruction normal form without introducing any additional intermediate theorem layer. In particular, the current line is already strong enough for the next public-shape step whenever no stronger claim about witness grade or deeper kernel-first activation is being asserted.*

Proof. By Theorem H2.176 and Corollary H2.177, the completed OP 3/QAM block already reduces translated shell data to certified passage versus first localized obstruction. Theorem H7.60 and Proposition H7.63 then transport that same dichotomy into the routed OP 5 side. Theorem H7.66 rewrites the routed return as carrier-stable versus obstructed normal form, and Theorem H7.69 finally packages the whole current run into global pass versus global obstruction. Thus no extra intermediate theorem layer is needed for that bundled question. The final sentence is only the proof-theoretic reading of Remark H7.71: stronger witness-grade or deeper kernel-first claims still require reopening the finer local layers. □

H7.22 Current extracted global pass/obstruction tool layer

Remark H7.74 (Purpose of the current global tool layer)**.** *The global pass/obstruction normal form is not meant to remain only as a terminal summary theorem. It is now extracted as a reusable tool layer so that later proofs can cite a small global interface instead of reopening the whole chain*

$$\textit{OP1 source} \to \textit{branch admissibility} \to \textit{OP3/QAM closure} \to \textit{carrier-first OP5 layer.}$$

The extracted tools below are therefore not new theory beyond Theorem H7.69; they are proof-usable projections of that bundled global dichotomy.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proposition H7.75 (Global-pass tool)**.** *Assume Theorem H7.69. If a current admissible run lies on the global pass normal form, then one may cite this fact directly as a reusable proof tool: the run is carried by one bundled closure-bearing carrier package from the OP1/branch entrance through the OP3/QAM shell package into the routed return side, and every admitted OP5 visible diagnostic remains subordinate to the carrier-stable visible normal form.*

Proof. This is precisely the positive branch of Theorem H7.69, rewritten as a direct reusable proof tool. □

Proposition H7.76 (Global-obstruction tool)**.** *Assume Theorem H7.69. If a current admissible run lies on the global obstruction normal form, then one may cite this fact directly as a reusable proof tool: the first licensed failure is already localized either upstream in the OP3/QAM obstruction package or downstream at the carrier-upgraded OP5 obstruction layer, and no third autonomous terminal visible class is currently licensed outside that bundled obstruction reading.*

Proof. This is precisely the negative branch of Theorem H7.69, rewritten as a direct reusable proof tool. □

Proposition H7.77 (Global transfer tool)**.** *Assume Theorem H7.69. Any later proof that has already routed the current admissible run into the bundled OP3/QAM-to-OP5 channel may work only with the bundled transfer dichotomy*

$$\textit{global pass} \qquad \textit{versus} \qquad \textit{global obstruction},$$

without reopening the intermediate local decomposition into separate branch, shell, witness, escalation, and routed-return subcases.

Proof. By Corollary H7.70, the whole currently admitted run has already been compressed into one reusable global dichotomy. The omitted intermediate subcases are already absorbed into the proof of Theorem H7.69. □

Proposition H7.78 (Global diagnostic subordination tool)**.** *Assume Theorem H7.69. On the global pass branch, every currently admitted visible routed-return diagnostic is globally subordinate to the same bundled closure-bearing carrier package. In particular, packet-wave, breathing-envelope, frame-flicker, and helix pictures cannot reintroduce a new structural branch once the run has already entered the global pass normal form.*

Proof. Theorem H7.69 sends the positive branch through the carrier-stable routed return normal form and therefore through the carrier-stable visible normal form. By construction, all visible routed-return diagnostics are subordinate to that same bundled carrier package and cannot reopen the global dichotomy. □

Proposition H7.79 (Global proof-interface tool)**.** *Assume Theorem H7.69. In later arguments it is sufficient to cite one of the four extracted global tools*

$$\textit{global pass}, \quad \textit{global obstruction}, \quad \textit{global transfer}, \quad \textit{global diagnostic subordination},$$

according to the local need of the argument. No additional independent proof interface for the currently admitted run is required at this stage.

Proof. Theorem H7.69 already supplies the full current bundled dichotomy, while Propositions H7.75–H7.78 expose its presently relevant projections. Hence these four extracted global tools form the current reusable proof interface for the admissible run. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Remark H7.80 (How to cite the extracted global tools)**.** *The current global tool layer should be cited only after the run has already been shown to lie inside the bundled OP1→branch→OP3/QAM→OP5 channel. Once that has been established, the present extracted tools allow later arguments to cite a compact global interface rather than reopening the entire local transport and diagnostic stack.*

H7.23 First global compressed proof run through extracted global tools

Remark H7.81 (Reading key for the first global compressed proof run)**.** *The present block is the first place where the current admissible run*

$$\textit{OP1 source} \to \textit{branch admissibility} \to \textit{OP3/QAM closure} \to \textit{carrier-first OP5 layer}$$

is no longer reopened stage by stage. Instead, once the run has already been shown to lie inside the bundled global channel, the extracted global tools are used as one compressed proof interface. The proof run therefore has only two licensed bundled outcomes: a global pass *and a* global obstruction*. The purpose of the theorem below is to state this compressed use rule explicitly, so that later arguments may pass through the whole current architecture in one step.*

Theorem H7.82 (First global compressed proof run through extracted global tools)**.** *Assume that a current admissible run issued from the OP1 source side has already been routed into the bundled OP1→branch→OP3/QAM→OP5 channel of the present architecture. Then the run may be processed through the extracted global tool layer without reopening the intermediate local stack. More precisely, exactly one of the following two bundled proof outputs is currently licensed.*

(i) ***Global compressed pass.*** *By Proposition H7.75, the run lies on the global pass normal form and is supported by one bundled closure-bearing carrier package from the OP1/branch entrance through the OP3/QAM shell package into the routed return side. By Proposition H7.78, all admitted visible routed-return diagnostics remain subordinate to that same bundled carrier package.*

(ii) ***Global compressed obstruction.*** *By Proposition H7.76, the run lies on the global obstruction normal form and its first licensed failure is already localized either upstream in the OP3/QAM obstruction package or downstream at the carrier-upgraded OP5 obstruction layer. By Proposition H7.77, no third autonomous terminal visible branch is currently licensed outside that bundled obstruction reading.*

Hence the present global proof run is now compressed to the dichotomy

$$\textit{global compressed pass} \qquad \textit{versus} \qquad \textit{global compressed obstruction},$$

and no further independent proof interface for the currently admitted run is required once Proposition H7.79 applies.

Proof. This is the first direct use of the extracted global tool layer as a compressed proof interface. Once the current run has entered the bundled

$$\text{OP1} \to \text{branch} \to \text{OP3/QAM} \to \text{OP5}$$

channel, Proposition H7.79 states that no additional independent proof interface is currently required.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

If the run lies on the positive side of Theorem H7.69, Proposition H7.75 supplies the bundled closure-bearing carrier package and Proposition H7.78 shows that all admitted visible OP5 diagnostics remain subordinate to that same carrier; this is exactly the global compressed pass.

If the run lies on the negative side, Proposition H7.76 localizes the first licensed failure and Proposition H7.77 excludes any third independent terminal visible branch; this is exactly the global compressed obstruction.

Exhaustiveness and mutual exclusivity therefore come directly from the extracted global tools and ultimately from Theorem H7.69. □

Corollary H7.83 (First operational use of the extracted global tools)**.** *In later arguments it is sufficient to cite Theorem H7.82 once the current admissible run has entered the bundled global channel. One may then work only with the dichotomy* global compressed pass versus global compressed obstruction*, together with the induced carrier-stable subordination or obstruction localization, without reopening the entire local OP3/QAM-to-OP5 stack.*

Proof. This is the operational reformulation of Theorem H7.82. The theorem already packages the extracted global tools into one reusable compressed proof run for the currently admitted architecture. □

H7.24 Global proof compression pass for the older OP1–OP5 line

Remark H7.84 (How the older OP1–OP5 line should now be reread)**.** *The older connected OP1–OP5 transport line should no longer be read as a sequence of largely separate source-return, branch, shell, routed, and diagnostic arguments that must each be reopened in full. After the present global compressed proof run, the mathematically correct reading is instead this: once an argument has entered the currently admitted bundled channel, the older line should be read global-first through the dichotomy*

$$\textit{global compressed pass} \qquad \textit{versus} \qquad \textit{global compressed obstruction.}$$

Only if a later step genuinely depends on shell certificates, shell-obstruction resolution, witness grades, witness escalation, or other local carrier-diagnostic data should one reopen the OP3/QAM or OP5 layers. The purpose of the proposition below is not to replace the older theorems, but to state explicitly that they may now be reused through one compressed global interface.

Proposition H7.85 (Compressed rereading of the older OP1–OP5 line through the extracted global tools)**.** *Assume that an argument along the older OP1–OP5 transport line has already been shown to lie inside the bundled channel of Theorem H7.82. Then the older line may be reread through the extracted global tools in the following compressed form.*

(i) ***Compressed pass rereading.*** *If the argument lies on the positive side of Theorem H7.82, then the older OP1–OP5 line is to be read as one globally compressed pass: the source-return closure theorem, the admissible branch layer, the OP3/QAM closure package, the carrier-first OP5 interface, and the subordinated diagnostic stack are all already absorbed into one bundled carrier-stable run.*

(ii) ***Compressed obstruction rereading.*** *If the argument lies on the negative side of Theorem H7.82, then the older OP1–OP5 line is to be read as one globally compressed obstruction: the first licensed failure is already localized in the bundled obstruction package, and no extra autonomous terminal visible branch may be added afterwards.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

In particular, later reuse of the older OP1–OP5 text should proceed through the extracted global proof interface rather than by reopening the whole local stack unless a genuinely new layer is being introduced.

Proof. This is the direct compression reading supplied by Theorem H7.82. On the positive side, the theorem already packages the full currently admitted run into the global compressed pass, and the extracted global tools show that the visible OP5 diagnostics remain subordinate to the same bundled carrier package. On the negative side, the same theorem packages the run into the global compressed obstruction and excludes any third autonomous terminal visible branch. Hence the older OP1–OP5 line can now be reused through one global proof interface rather than by reopening each intermediate theorem separately. □

Corollary H7.86 (Global compressed reading of older routed return arguments)**.** *Whenever an older routed return argument is already known to lie inside the bundled current channel, it should by default be cited through Proposition H7.85, Theorem H7.82, and the global citation discipline of Proposition H7.142. In that situation, it is sufficient to work only with the dichotomy*

$$\textit{global compressed pass} \qquad \textit{versus} \qquad \textit{global compressed obstruction},$$

and with the induced bundled carrier-stable or obstruction-localized reading, unless a genuinely shell-sensitive or witness-sensitive local claim must still be reopened.

Proof. This is the operational reformulation of Proposition H7.85. Once the argument is inside the bundled current channel, the extracted global proof interface supplies the compressed rereading rule for the older routed return line. □

H7.25 Witness-grade hardening and the first local kernel-first rewrite window

Remark H7.87 (Purpose of the archival stage witness-grade hardening step)**.** *The public archival stage release already exposed the bundled OP 3/QAM closure package and the bundled OP 3→OP 5/global channel in compact public shape. The next task is therefore not a new visible closure notion, but a stricter proof-use discipline for that same channel. In particular, the present step isolates three things: first, a bundled witness-grade reopening package for the OP 5 side; second, a sharper local/global reopening rule explaining exactly when the bundled global interface is sufficient; and third, one first narrow kernel-first rewrite window whose use is licensed only after the bundled channel has already been fixed and only at a local witness-grade transition where the deferred bridge actually reduces proof load.*

Theorem H7.88 (Bundled witness-grade reopening package)**.** *Assume Theorem H7.57, Proposition H7.63, Theorem H7.69, Proposition H7.142, and Proposition H7.145. Let a later argument concern a run already known to lie inside the bundled current channel. Then exactly one of the following two proof-use regimes is presently licensed.*

(i) ***Bundled global-use regime.*** *If the argument needs only the bundled outputs global compressed pass/global compressed obstruction together with their carrier-stable or obstruction-localized consequences, then the extracted global interface is sufficient and no local witness-grade reopening is licensed or needed.*

(ii) ***Witness-grade reopening regime.*** *If the argument needs the witness grade itself, witness escalation, screening-witness compression, carrier-witness separation, diagnostic*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

collapse below the escalation threshold, or a comparison between shell-grade and witness-grade data, then the bundled global interface must be reopened locally at the OP 5 witness layer, and the proof must work with the finer witness hierarchy rather than with the global dichotomy alone.

Hence the current architecture already supports one bundled witness-grade reopening package: bundled-global use by default, local witness-grade reopening only for genuinely witness-sensitive claims.

Proof. The bundled global-use regime is exactly the content of Proposition H7.142 together with Proposition H7.145: once the argument is already inside the bundled current channel and only the global pass/obstruction outputs are needed, the extracted global layer is the default proof interface. By contrast, Theorem H7.57 and Proposition H7.63 show that witness grade, witness escalation, screening-witness compression, carrier-witness separation, and diagnostic collapse all live strictly below that bundled global interface. Therefore any genuinely witness-sensitive claim forces a local reopening of the OP 5 witness layer. The two regimes are exhaustive because every later proof step either remains at bundled-global level or asks for one of the finer witness-sensitive distinctions just listed. □

Proposition H7.89 (Witness-grade local/global reopening discipline)**.** *Assume Theorem H7.88. Then the current bundled channel should be used with the following reopening discipline.*

(i) work global-first *once the argument is already inside the bundled current channel;*

(ii) reopen only the OP 3/QAM shell layer *when the step depends on shell certificates, shell-defect strata, shell commuting behavior, or OP 3-side carrier compression;*

(iii) reopen only the OP 5 witness layer *when the step depends on witness grade, witness escalation, screening-witness compression, carrier-witness separation, or diagnostic collapse below escalation;*

(iv) reopen both local layers *only when the step genuinely compares shell-grade and witness-grade data or transfers a shell-sensitive obstruction into a witness-sensitive one.*

In particular, visible OP 5 packet, envelope, flicker, and helix data do not by themselves justify a local reopening once the argument is already licensed to remain global-first.

Proof. Clause (i) is exactly the bundled-global default from Theorem H7.88. Clause (ii) restates the OP 3/QAM reopening part already fixed by Proposition H7.142. Clause (iii) is the witness-grade reopening regime from Theorem H7.88. Clause (iv) records the only mixed local situation still licensed at the present stage: a step that must compare shell-grade and witness-grade structure rather than merely cite one local layer. The final sentence follows because, below escalation, visible OP 5 diagnostics remain subordinate to the carrier-stable visible normal form and therefore do not force reopening by appearance alone. □

Corollary H7.90 (Witness-grade reopening checklist)**.** *For later proof use, the current bundled channel may be tested by the following checklist.*

(i) Has the run already entered the bundled current channel?

(ii) Is only the bundled global pass/obstruction output needed?

(iii) If not, is the missing information shell-grade, witness-grade, or genuinely mixed shell/witness data?

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(iv) Reopen only the corresponding local layer, and return to the bundled global interface immediately after the local claim has been discharged.

This is the default proof-use checklist for the archival stage witness-grade hardening step.

Proof. This is the operational reformulation of Proposition H7.89. The checklist simply turns the local/global reopening rule into a direct proof-use procedure. □

Definition H7.91 (First licensed local kernel-first rewrite window)**.** *Assume the direct kernel constructor of Definition H2.118, the bridge conditions of Theorem H2.119, and the bundled witness-grade reopening package of Theorem H7.88. The* first licensed local kernel-first rewrite window *is the following strictly local situation: a proof step has already entered the bundled current channel, has already been forced to reopen at witness grade, and now needs only to compare source-side and return-side witness-bearing states through one common carrier-stable normal form without promoting any broader kernel-first migration on the surrounding global channel.*

Proposition H7.92 (First licensed local kernel-first rewrite window)**.** *Assume Definition H7.91, Proposition H7.27, and Theorem H7.57. Let a witness-sensitive proof step concern a routed return realization already known to remain on the carrier-stable branch of the bundled OP 3→OP 5 transfer channel. Then that step may be rewritten kernel-first in the following narrow sense.*

(i) the visible witness-bearing source/return surfaces may be replaced by their exposed payload kernels and compared only through the common kernel normal form forced by the carrier-stable branch;

(ii) the local readout comparison may be performed at the level of the resulting common kernel/readout carrier class instead of on the full visible witness-bearing surfaces;

(iii) after that local comparison has been discharged, the proof must return immediately to the bundled carrier-stable/global interface and may not infer a broader kernel-first rewrite rule for the full OP 3→OP 5/global channel.

Hence the present line now contains one first licensed kernel-first rewrite window: local witness-grade carrier comparison under an already fixed carrier-stable branch, but no general kernel-first migration of the whole bundled channel.

Proof. Once the proof step is already known to lie on the carrier-stable branch, the witness-bearing visible data are subordinate to one common closure-bearing carrier package. Under the direct kernel constructor and the bridge conditions, this is exactly the situation in which the local comparison load can be shifted from full visible surfaces to their exposed payload kernels and then to the corresponding common kernel normal form. The local readout comparison may therefore be discharged at the level of the common kernel/readout carrier class. However, Theorem H7.88 and Definition H7.91 keep this move strictly local: the argument was already forced into witness-grade reopening, and nothing in the present hypotheses licenses promotion of that local kernel-first step to a global rewrite rule for the surrounding bundled channel. The final sentence is exactly this restriction written as a proof-use statement. □

Corollary H7.93 (Use rule for the first local kernel-first rewrite window)**.** *Whenever a later proof step is already inside the bundled current channel, already forced to reopen at witness grade, and already known to remain on the carrier-stable branch, it is sufficient to cite Proposition H7.92 for the local source/return comparison. After that local comparison, the proof should return to the bundled global interface immediately.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. This is the operational reformulation of Proposition H7.92. The proposition already states both the local license and the immediate return rule. □

Remark H7.94 (Status of the first corridor-level activation step)**.** *The present release line still does* not *activate a kernel-first rewrite rule for the whole bundled OP 3→OP 5/global channel. Nevertheless, the combination of the bundled witness-grade reopening package and the first licensed local kernel-first rewrite window already supports one slightly stronger activation step: a finite witness-grade corridor on one fixed carrier-stable branch may be discharged kernel-first through one common carrier-stable kernel/readout class. This is the first point at which the current release line moves beyond a single isolated local rewrite without yet promoting the deferred kernel architecture to an active global migration layer.*

Definition H7.95 (Carrier-stable witness corridor)**.** *Assume Theorem H7.88 and Proposition H7.92. A* carrier-stable witness corridor *is a finite family of witness-sensitive proof steps*

$$\Sigma_1, \dots, \Sigma_N$$

along one routed return run such that:

(i) the run is already known to lie inside the bundled current channel;

(ii) every step Σ_j is already forced to reopen at witness grade and remains on the same carrier-stable branch;

(iii) for every j, the local source/return comparison occurring in Σ_j satisfies the hypotheses of Proposition H7.92;

(iv) no step in the family changes the fixed carrier-stable normal-form class or requires a new shell/witness transfer beyond that same carrier-stable branch.

Thus a carrier-stable witness corridor is the smallest finite witness-grade subrun on which the same carrier-stable kernel/readout class may be reused without reopening the whole bundled channel anew at each local comparison.

Lemma H7.96 (First corridor-level kernel-first activation lemma)**.** *Assume Definition H7.95. Then the whole witness corridor may be discharged kernel-first through one common carrier-stable kernel/readout class in the following precise sense.*

(i) for every local step Σ_j, the visible witness-bearing source/return surfaces may be replaced by their exposed payload kernels and compared only through the same carrier-stable kernel normal-form class;

(ii) the local readout comparisons occurring across the full finite family $\Sigma_1, \dots, \Sigma_N$ may therefore be performed at class level rather than re-established on the full visible witness-bearing surfaces step by step;

(iii) after the corridor has been discharged, the proof must return to the bundled global interface immediately and may not infer a general kernel-first rewrite rule for branch-changing, shell-changing, or globally mixed witness-grade arguments.

Equivalently, the present line already licenses one first corridor-level activation step: a finite bundled family of witness-grade local comparisons along one fixed carrier-stable branch, but still no general kernel-first migration of the whole bundled channel.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. By Definition H7.95, every local step Σ_j already satisfies the hypotheses of Proposition H7.92. Hence each such step may be rewritten kernel-first at the local witness-sensitive level. Because all steps lie on the same carrier-stable branch and no step changes the fixed carrier-stable normal-form class, the corresponding local kernel/readout comparisons all factor through one and the same carrier-stable kernel/readout class rather than through a new class at every step. Therefore the whole finite family may be discharged kernel-first corridor by corridor instead of reopening the full visible witness-bearing surfaces separately at each local comparison.

The return rule is unchanged: Theorem H7.88 keeps witness-grade reopening subordinate to the bundled global interface, and Definition H7.95 excludes branch-changing, shell-changing, or globally mixed witness-grade transitions. Consequently, once the corridor has been discharged, the proof must return immediately to the bundled global interface, and no broader global kernel-first migration follows from this corridor-level argument. □

Corollary H7.97 (Use rule for the first corridor-level activation lemma)**.** *Whenever a later proof contains a finite witness-grade subrun on one fixed carrier-stable branch, it is sufficient to cite Lemma H7.96 instead of reopening each witness-bearing source/return comparison separately. After that corridor-level comparison has been discharged, the proof should return to the bundled global interface immediately.*

Proof. This is the operational reformulation of Lemma H7.96. The lemma already states both the corridor-level license and the immediate return rule. □

Remark H7.98 (Status of the first corridor-to-bundle propagation step)**.** *The present release line still does* not *activate a kernel-first rewrite rule for the whole bundled OP 3→OP 5/global channel. Nevertheless, the combination of the bundled witness-grade reopening package, the first licensed local kernel-first rewrite window, and the first corridor-level activation lemma already supports one slightly stronger propagation step: a bundled run may now alternate between globally readable stretches and finitely many carrier-stable witness corridors without forcing each witness-bearing local comparison to be reopened from scratch. This is the first point at which the current release line moves beyond one isolated corridor-level activation and begins to control a whole bundled run by a finite alternation of global citations and corridor-level kernel-first discharges.*

Definition H7.99 (Bundle-compatible witness-corridor decomposition)**.** *Assume Theorem H7.88, Lemma H7.96, and Proposition H7.142. A* bundle-compatible witness-corridor decomposition *of a later proof run is a finite alternating decomposition*

$$\Gamma_0,\ \mathcal{C}_1,\ \Gamma_1,\ \ldots,\ \mathcal{C}_M,\ \Gamma_M$$

such that:

(i) the entire run is already known to lie inside the bundled current channel;

(ii) each global segment Γ_i needs only the bundled outputs global compressed pass/global compressed obstruction together with their induced carrier-stable or obstruction-localized consequences;

(iii) each local segment $\mathcal{C}_k$ is a carrier-stable witness corridor in the sense of Definition H7.95;

(iv) after each local corridor $\mathcal{C}_k$, the proof returns immediately to the bundled global interface before any later corridor begins;

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(v) no corridor changes the fixed bundled run into a new shell-changing, branch-changing, or globally mixed witness-grade run between two global segments.

Thus a bundle-compatible witness-corridor decomposition is the smallest finite alternation in which the bundled global interface remains the default proof surface and only finitely many carrier-stable witness corridors are discharged locally.

Proposition H7.100 (First corridor-to-bundle propagation principle)**.** *Assume Definition H7.99. Then the whole decomposed run may be discharged by the following finite proof pattern.*

(i) on each global segment Γ_i, it is sufficient to cite the bundled global interface and work only with the dichotomy

$$\textit{global compressed pass} \quad \textit{versus} \quad \textit{global compressed obstruction};$$

(ii) on each local corridor $\mathcal{C}_k$, it is sufficient to cite Lemma H7.96 and discharge the whole finite witness-grade corridor through one common carrier-stable kernel/readout class;

(iii) after each local corridor $\mathcal{C}_k$, the proof returns immediately to the bundled global interface and resumes global-first reasoning on the next segment Γ_k;

(iv) consequently, no individual witness-bearing source/return comparison inside the corridors needs to be reopened separately, and no general kernel-first rewrite rule for the full bundled channel is inferred.

Equivalently, the present line already licenses one first corridor-to-bundle propagation rule: a bundled run may be handled by finitely many global-interface citations plus finitely many corridor-level kernel-first discharges, while the active proof surface remains globally bundled outside those local corridors.

Proof. By Definition H7.99, every global segment Γ_i satisfies exactly the hypotheses of the bundled global citation discipline, so Proposition H7.142 supplies the global compressed pass/global compressed obstruction interface there. Likewise, every local segment $\mathcal{C}_k$ is a carrier-stable witness corridor, so Lemma H7.96 discharges that whole finite corridor kernel-first through one common carrier-stable kernel/readout class.

The return rule is part of both ingredients: the global interface is the default outside local reopenings, while the corridor lemma requires immediate return to the bundled global interface after each local corridor has been discharged. Hence the full run is controlled by a finite alternation of global citations and corridor-level kernel-first discharges. Because the corridor lemma already absorbs the local witness-grade source/return comparisons inside each $\mathcal{C}_k$, no individual witness-bearing comparison has to be reopened separately. Finally, nothing in the hypotheses licenses promotion of this finite alternation to a general kernel-first rewrite rule for the whole bundled channel; outside the listed corridors the active proof surface remains the bundled global interface. □

Corollary H7.101 (Use rule for the first corridor-to-bundle propagation principle)**.** *Whenever a later proof run already inside the bundled current channel decomposes into globally readable stretches separated by finitely many carrier-stable witness corridors, it is sufficient to cite Proposition H7.100 together with Lemma H7.96, rather than reopening each witness-bearing local comparison separately. Only the corridor boundaries, not the internal witness steps of each corridor, remain visible at proof-use level.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. This is the operational reformulation of Proposition H7.100. The proposition already states both the finite propagation rule and the immediate return to the bundled global interface after each corridor. □

Remark H7.102 (Status of the first bundle-level reopening compression step)**.** *The present release line still does* not *collapse every local reopening beneath the bundled channel to one uniform active theorem surface. What it now does support, however, is one further compression of proof-use data: once a bundled run has already been decomposed into global stretches and finitely many carrier-stable witness corridors, the later proof no longer needs to describe each corridor by all of its internal shell-grade and witness-grade subclaims. It is enough to record, corridor by corridor, which local reopening profile is actually being used. This is the first point at which the current release line compresses not only the internal witness comparisons of each corridor, but also the reopening metadata attached to the whole bundled run.*

Definition H7.103 (Profile-stable bundle-level reopening data)**.** *Assume Definition H7.99. A* profile-stable bundle-level reopening datum *on such a decomposition is a choice of one profile label*

$$\pi_k \in \{\mathrm{Q}, \mathrm{W}, \mathrm{QW}\}$$

for every corridor $\mathcal{C}_k$ such that:

(i) $\pi_k = \mathrm{Q}$ *means that the corridor is reopened only through the OP 3/QAM shell side: shell certificates, shell-obstruction stratification, shell commuting behavior, or OP 3-side carrier compression;*

(ii) $\pi_k = \mathrm{W}$ *means that the corridor is reopened only through the carrier-first OP 5 witness side: witness grade, witness escalation, screening-witness compression, carrier-witness separation, or diagnostic collapse beneath the escalation threshold;*

(iii) $\pi_k = \mathrm{QW}$ *means that the corridor genuinely compares shell-grade and witness-grade data inside one common carrier-stable witness corridor;*

(iv) the profile label of a corridor remains fixed throughout that corridor and changes, if at all, only at corridor boundaries.

Thus the profile sequence $(\pi_1, \ldots, \pi_M)$ records the smallest local reopening data needed at proof-use level beyond the bundled global interface.

Proposition H7.104 (First bundle-level reopening compression principle)**.** *Assume a bundle-compatible witness-corridor decomposition together with profile-stable bundle-level reopening data in the sense of Definition H7.103. Then the whole bundled run may be handled at proof-use level by the finite data*

$$\Gamma_0,\ \pi_1,\ \Gamma_1,\ \ldots,\ \pi_M,\ \Gamma_M$$

with the following force.

(i) Each global segment Γ_i is still treated only through the bundled global interface and the dichotomy global compressed pass versus global compressed obstruction.

(ii) Each corridor $\mathcal{C}_k$ is reopened only through its single profile label π_k, together with Lemma H7.96 whenever the corridor is discharged kernel-first as one carrier-stable witness corridor.

(iii) No later proof needs to list the internal shell-grade or witness-grade subclaims of $\mathcal{C}_k$ separately as long as they remain subordinate to the same fixed corridor profile π_k.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(iv) If a corridor would force a profile change inside itself, then the present compression rule stops at that point and the corridor must be split into smaller profile-stable corridors before the bundled proof-use compression is applied again.

Equivalently, the current release line already licenses one first bundle-level reopening compression step: after corridor-level kernel-first discharge has been fixed, the remaining local proof-use burden of a bundled run is reduced to a finite sequence of profile labels attached to finitely many carrier-stable corridors.

Proof. By Proposition H7.100, the bundled run is already controlled by finitely many global citations together with finitely many corridor-level kernel-first discharges. The only remaining local proof-use burden is therefore to specify which finer local reopening regime is actually needed on each corridor. Definition H7.103 makes exactly that finite burden explicit: every corridor carries one fixed profile label π_k, and that label determines whether the local reopening is OP 3/QAM-only, OP 5-witness-only, or genuinely mixed.

Because the profile remains fixed throughout each corridor, all internal shell-grade and witness-grade subclaims of that corridor stay subordinate to the same local reopening regime and need not be listed one by one at proof-use level. If a profile change occurred inside a corridor, then the corridor would no longer be profile-stable, and the current compression would no longer be legitimate without first splitting that corridor into smaller profile-stable pieces. This proves both the finite proof-use description and the stated stopping rule. □

Corollary H7.105 (Use rule for the first bundle-level reopening compression principle)**.** *Whenever a later proof run already inside the bundled current channel admits both a bundle-compatible witness-corridor decomposition and profile-stable bundle-level reopening data, it is sufficient to cite Proposition H7.104 and record only the resulting finite profile sequence* $(\pi_1, \dots, \pi_M)$*, rather than restating all local shell-grade and witness-grade subclaims corridor by corridor. A finer reopening is required only when one corridor changes profile internally or leaves the carrier-stable witness-corridor regime.*

Proof. This is the operational reformulation of Proposition H7.104. The proposition already states both the compressed proof-use description and the precise point where that compression stops. □

Remark H7.106 (Status of the first profile-to-run compression step)**.** *The present release line still does* not *identify all bundled runs by one fully kernel-activated normal form. What it now does support, however, is one further compression of the proof-use surface: once a bundled run has already been decomposed into global stretches and finitely many profile-stable carrier corridors, the later proof no longer needs to remember the names of the individual global segments or the internal witness subclaims of those corridors. It is enough to remember the alternating* run profile word *that records only when the proof is global and which local reopening profile occurs at each corridor. This is the first point at which the current release line compresses a whole bundled run to one finite symbolic proof-use word.*

Definition H7.107 (Run-stable bundled proof word)**.** *Assume a bundle-compatible witness-corridor decomposition together with profile-stable bundle-level reopening data in the sense of Definition H7.103. The associated* run-stable bundled proof word *is the finite alternating word*

$$\omega(\Gamma) := \mathrm{G}\,\pi_1\,\mathrm{G}\,\pi_2\,\mathrm{G}\cdots\pi_M\,\mathrm{G}$$

over the alphabet

$$\{\mathrm{G}, \mathrm{Q}, \mathrm{W}, \mathrm{QW}\},$$

where:

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(i) each letter G *records one globally readable bundled segment handled only through the dichotomy global compressed pass versus global compressed obstruction;*

(ii) each letter $\pi_k \in \{\mathrm{Q}, \mathrm{W}, \mathrm{QW}\}$ *records the fixed reopening profile of the corresponding carrier-stable witness corridor* $\mathcal{C}_k$*;*

(iii) the word begins and ends with G*, and the letters alternate globally/local in the same order as the bundled run;*

(iv) no letter change is allowed inside a corridor; if such a change occurs, the current word is not defined until the corridor has been split into smaller profile-stable corridors.

Thus $\omega(\Gamma)$ *is the smallest finite symbolic record that still remembers where the proof is global and which local reopening profile occurs whenever the bundled channel is reopened.*

Proposition H7.108 (First profile-to-run compression principle)**.** *Assume Definition H7.107. Then the proof-use burden of the whole bundled run is reduced to its finite run-stable bundled proof word*

$$\omega(\Gamma) \in \{\mathrm{G}, \mathrm{Q}, \mathrm{W}, \mathrm{QW}\}^*$$

with the following force.

(i) Every occurrence of G *means that the corresponding segment is handled only through the bundled global interface and its compressed pass/obstruction dichotomy.*

(ii) Every occurrence of Q*,* W*, or* QW *means that the next corridor is reopened only in that single local profile, together with Lemma H7.96 whenever the corridor is licensed for carrier-stable kernel-first discharge.*

(iii) No later proof needs to retain the individual names of the global segments Γ_i *or the separate internal shell-grade/witness-grade clauses of the corridors once the alternating word* $\omega(\Gamma)$ *has been fixed.*

(iv) If one wants to refine a corridor internally, change its local reopening profile midstream, or distinguish two globally readable segments by more than the bundled pass/obstruction interface, then the present compression stops and the run must first be rewritten at the preceding bundle-level stage.

Equivalently, the current release line already licenses one first profile-to-run compression step: after bundle-compatible corridor decomposition and profile stabilization have been fixed, the whole bundled proof surface may be cited by one finite alternating word over $\{\mathrm{G}, \mathrm{Q}, \mathrm{W}, \mathrm{QW}\}$*.*

Proof. By Proposition H7.104, the bundled run is already controlled by finitely many global segments together with finitely many profile-stable carrier corridors. The only remaining proof-use information is therefore the order in which those two types of pieces occur and, on each corridor, which fixed reopening profile is active. Definition H7.107 records exactly that data as the alternating word $\omega(\Gamma)$.

The force of each letter is already supplied by the previous stages: G stands for the bundled global interface, while Q, W, and QW stand for the corresponding local reopening profiles and, where licensed, for the corridor-level kernel-first discharge of Lemma H7.96. Since the previous bundle-level compression already removes the need to list individual internal witness clauses corridor by corridor, fixing the alternating word removes the need to retain the names of the global segments as separate proof-use data. If one wants finer information than the word provides, then one has already left the current compressed regime and must reopen the run at the preceding bundle-level stage. This proves the stated stopping rule. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Corollary H7.109 (Use rule for the first profile-to-run compression principle)**.** *Whenever a later proof run already inside the bundled current channel admits a run-stable bundled proof word $\omega(\Gamma)$, it is sufficient to cite Proposition H7.108 and record only that finite word, rather than listing the separate global segments and corridor-internal local claims again. A finer reopening is required only when a corridor changes profile internally, leaves the carrier-stable witness-corridor regime, or one global segment must be distinguished from another by more than the bundled pass/obstruction interface.*

Proof. This is the operational reformulation of Proposition H7.108. The proposition already states both the finite run-word description and the precise point where that description stops being legitimate. □

Remark H7.110 (Status of the first word-determined run-class step)**.** *The present release line still does* not *identify all bundled runs by one globally activated kernel normal form, nor does it erase the distinction between different geometric realizations of the same bundled channel. What it now does support, however, is one further proof-use quotient: once a bundled run has already been compressed to a run-stable bundled proof word, the later proof may forget the individual realization of that run and retain only the* run class determined by that word. *This is the first point at which the current release line passes from one finite symbolic proof-use word to one finite symbolic proof-use class.*

Definition H7.111 (Word-determined bundled run class)**.** *Assume Definition H7.107. Two bundled runs Γ and Γ' are called* word-determined proof-use equivalent *if both run-stable bundled proof words are defined and satisfy*

$$\omega(\Gamma) = \omega(\Gamma').$$

The corresponding equivalence class is written

$$[\Gamma]_\omega := \{\Gamma' : \omega(\Gamma') = \omega(\Gamma)\}.$$

It is called the word-determined bundled run class *of Γ. In this reading, the class $[\Gamma]_\omega$ records exactly the proof-use information retained by the current compressed line and forgets the names of individual global segments, the particular realization of each carrier-stable witness corridor, and all corridor-internal shell-grade or witness-grade subclaims beyond what is already encoded by the letters of $\omega(\Gamma)$.*

Proposition H7.112 (First word-to-run-class compression principle)**.** *Assume Definition H7.111. Then the current proof-use burden of a bundled run depends only on its word-determined bundled run class $[\Gamma]_\omega$ and no longer on the individual run Γ itself, with the following force.*

(i) If two bundled runs Γ and Γ' satisfy $\omega(\Gamma) = \omega(\Gamma')$, then every later argument that uses only the bundled global interface, the associated local reopening profiles, and the corresponding corridor-level kernel-first licenses may treat Γ and Γ' as the same proof-use object.

(ii) Consequently, once the run-stable bundled proof word has been fixed, the later proof may cite the class $[\Gamma]_\omega$ instead of the particular run realization Γ.

(iii) The present compression stops exactly when one needs information not encoded by the word: for example, a distinction between two globally readable segments carrying the same letter G*, a distinction between two carrier-stable witness corridors carrying the same profile letter, or a geometric/property-level comparison internal to one corridor that is invisible at word level.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Equivalently, the current release line already licenses one first word-to-run-class compression step: after passage from bundled runs to run-stable bundled proof words, proof use depends only on the induced equivalence class under equality of those words.

Proof. By Proposition H7.108, the current proof-use surface of a bundled run is already reduced to the finite alternating word $\omega(\Gamma)$. Hence every later argument that cites only the force carried by the letters G, Q, W, and QW, together with the associated corridor-level kernel-first license where available, uses no further data from the particular realization of Γ.

Therefore, if $\omega(\Gamma) = \omega(\Gamma')$, then both runs carry exactly the same compressed proof-use data and may be treated as the same proof-use object at the current level of abstraction. This proves (i) and the resulting class-level citation claim in (ii). Clause (iii) is simply the contrapositive reading of the same compression: whenever a later argument needs data not already encoded by the word, the quotient by equality of words is too coarse, so the proof must reopen at the preceding run-level or bundle-level stage. □

Corollary H7.113 (Use rule for the first word-to-run-class compression principle)**.** *Whenever a later proof already inside the bundled current channel depends only on the run-stable bundled proof word and the force of its letters, it is sufficient to cite Proposition H7.112 and work with the corresponding class $[\Gamma]_\omega$, rather than with one chosen realization of the bundled run. A finer reopening is required only when the argument needs information not encoded by the word itself.*

Proof. This is the operational reformulation of Proposition H7.112. The proposition already states both the induced class-level proof-use rule and the precise point where that compression stops. □

Remark H7.114 (Status of the bundled proof-use compression stack in the present 4.7.0 release line)**.** *The present release line still does* not *equip bundled current-channel reasoning with one unrestricted global kernel algebra. What it now supports, however, is a much shorter proof-use stack: bundled runs may be compressed successively from run words to run classes, from finite concatenation chains to one reduced normal-form word, from that word to one proof-use class, from binary and finite class composition to one associative class word, and finally to one contraction-stable terminal citation surface. The next items record only the exact proof-use interfaces needed later; the longer stage-by-stage derivations are no longer repeated here.*

Definition H7.115 (Concatenable word-determined bundled run classes)**.** *Assume Definition H7.111. Two word-determined bundled run classes $[\Gamma]_\omega$ and $[\Gamma']_\omega$ are* concatenable at bundled level *if their join is handled entirely by the bundled global interface and no new local reopening is created at the splice. In that case define the reduced concatenated run word by*

$$\omega(\Gamma) \star_{\mathrm{G}} \omega(\Gamma')$$

obtained by identifying the terminal global letter G *of the first word with the initial global letter* G *of the second, and write*

$$[\Gamma]_\omega \star [\Gamma']_\omega$$

for the induced concatenated run-class citation object.

Proposition H7.116 (First run-class concatenation principle)**.** *Under Definition H7.115, bundled-level concatenation depends only on the two input run classes and on the reduced concatenated run word. In particular, later proof use may cite $[\Gamma]_\omega \star [\Gamma']_\omega$ instead of chosen representatives, and this compression stops exactly when the splice creates local data not already encoded by the adjacent global-interface letters.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. Immediate from the word-level quotient already fixed by Proposition H7.112: once representatives are forgotten, the only remaining bundled splice datum is the reduced global-interface identification. Any genuinely new local burden at the join lies beyond this quotient and forces reopening. □

Corollary H7.117 (Use rule for the first run-class concatenation principle)**.** *Whenever two bundled current-channel runs are already compressed to word-determined bundled run classes and their join is handled entirely by the bundled global interface, it is sufficient to cite Proposition H7.116 and work with the concatenated class* $[\Gamma]_\omega \star [\Gamma']_\omega$.

Proof. This is the operational reading of Proposition H7.116. □

Remark H7.118 (Status of the first finite run-class normal-form step)**.** *The next compression replaces a finite ordered chain of concatenable bundled run classes by one reduced normal-form word. Only the ordered local reopening profile and the bundled global separators survive.*

Definition H7.119 (Finite concatenable bundled run-class chain and reduced normal-form word)**.** *Let*

$$[\Gamma_1]_\omega, \dots, [\Gamma_n]_\omega$$

be bundled run classes such that each adjacent pair is concatenable at bundled level. Define the reduced normal-form concatenation word

$$\mathrm{NF}_\star([\Gamma_1]_\omega, \dots, [\Gamma_n]_\omega)$$

by concatenating the words $\omega(\Gamma_k)$ *and identifying every adjacent terminal/initial global letter* G *as one common bundled global-interface letter.*

Proposition H7.120 (First finite run-class normal-form principle)**.** *For every finite bundled concatenation chain covered by Definition H7.119, the proof-use burden depends only on the ordered list of run classes and on the reduced normal-form word* $\mathrm{NF}_\star$*. Representative choices and parenthesization of successive concatenations do not matter.*

Proof. Repeated use of Proposition H7.116 shows that each bundled-level splice is governed only by the reduced global-interface identification. Iterating that splice therefore yields one parenthesization-independent reduced word. □

Corollary H7.121 (Use rule for the first finite run-class normal-form principle)**.** *Whenever a later argument uses only the ordered local profile carried by a finite concatenable run-class chain, it is sufficient to cite Proposition H7.120 and work with* $\mathrm{NF}_\star$.

Proof. This is the operational reading of Proposition H7.120. □

Remark H7.122 (Status of the first normal-form-determined bundled proof-use class step)**.** *Once a finite concatenable bundled run-class chain has been reduced to one normal-form word, later proof use may forget the chosen chain realization and retain only the class determined by that word.*

Definition H7.123 (Normal-form-determined bundled proof-use class)**.** *Two finite concatenable bundled run-class chains are called normal-form proof-use equivalent if their reduced normal-form words agree. The corresponding equivalence class is the* normal-form-determined bundled proof-use class *and is written* $[\mathfrak{C}]_{\mathrm{NF}}$.

Proposition H7.124 (First normal-form-to-proof-use-class compression principle)**.** *The current proof-use burden of a finite concatenable bundled run-class chain depends only on its normal-form-determined bundled proof-use class. Equivalently, once only the reduced normal-form word and the force of its letters matter, one may replace the chain by* $[\mathfrak{C}]_{\mathrm{NF}}$.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. This is precisely the quotient induced by equality of reduced normal-form words from Definition H7.123. □

Corollary H7.125 (Use rule for the first normal-form-to-proof-use-class compression principle)**.** *Whenever a later proof already inside the bundled current channel depends only on the reduced normal-form concatenation word and the force of its letters, it is sufficient to cite Proposition H7.124 and work with the corresponding class* $[\mathfrak{C}]_{\mathrm{NF}}$*.*

Proof. This is the operational reading of Proposition H7.124. □

Remark H7.126 (Status of the first proof-use-class composition step)**.** *After passage to normal-form-determined classes, bundled composition can be handled at class level whenever the splice itself adds no new local burden.*

Definition H7.127 (Composable normal-form-determined bundled proof-use classes)**.** *Two normal-form-determined bundled proof-use classes are* composable *if realizations may be joined entirely through the bundled global interface with no new local reopening at the splice. Their reduced composed normal-form word is obtained by the same* $\star_{\mathrm{G}}$ *splice on the two reduced normal-form words.*

Proposition H7.128 (First proof-use-class composition principle)**.** *Whenever two normal-form-determined bundled proof-use classes are composable, their bundled composition depends only on the two classes and on the reduced composed normal-form word. Representative-level choices are invisible at this class-compression stage.*

Proof. Combine Proposition H7.124 with the same bundled-global splice rule already used at run-class level. □

Corollary H7.129 (Use rule for the first proof-use-class composition principle)**.** *Whenever two bundled current-channel proof packages are already compressed to normal-form-determined bundled proof-use classes and their splice is handled entirely by the bundled global interface, it is sufficient to cite Proposition H7.128 and work with their composed class.*

Proof. This is the operational reading of Proposition H7.128. □

Remark H7.130 (Status of the first associative proof-use-class normal-form step)**.** *Iterated class composition can now be read through one parenthesization-stable reduced class word.*

Definition H7.131 (Associative reduced class-composition normal form)**.** *For a finite ordered composable family of normal-form-determined bundled proof-use classes, define the associative reduced class-composition normal form by concatenating the reduced class words and identifying every interior terminal/initial global letter* G *as one common bundled global-interface letter.*

Proposition H7.132 (First associative proof-use-class normal-form principle)**.** *The proof-use burden of any finite ordered composable family of normal-form-determined bundled proof-use classes depends only on its associative reduced class-composition normal form and is independent of parenthesization.*

Proof. At word level the bundled-global splice is associative; Proposition H7.128 removes dependence on representatives, so only the resulting reduced class word remains. □

Corollary H7.133 (Use rule for the first associative proof-use-class normal-form principle)**.** *Whenever a later bundled current-channel proof uses only the force of a finite ordered composable family of normal-form-determined bundled proof-use classes and all intermediate joins are handled entirely by the bundled global interface, it is sufficient to cite Proposition H7.132 and work with the single associative class word.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. This is the operational reading of Proposition H7.132. □

Remark H7.134 (Status of the first contraction-stable class-normal-form step)**.** *The final local compression removes only redundant consecutive global separators and leaves the genuinely reopened local content unchanged.*

Definition H7.135 (Contraction-stable reduced class-normal-form word)**.** *Given a finite composable family of normal-form-determined bundled proof-use classes with associative reduced class word* ω_{assoc}*, define*

$$\mathrm{NF}_{\mathrm{ctr}}$$

by replacing each maximal consecutive block of the symbol G *in* ω_{assoc} *by a single occurrence of* G*.*

Proposition H7.136 (First contraction-stable class-normal-form principle)**.** *The contraction-stable class-normal-form word* $\mathrm{NF}_{\mathrm{ctr}}$ *is independent of representatives and parenthesization and determines the same proof-use surface as the associative class-normal-form word, except that redundant purely global separators are contracted.*

Proof. Representative and parenthesization independence are inherited from Proposition H7.132; the contraction step touches only maximal G-blocks and therefore does not alter the order of the genuinely local letters Q, W, and QW. □

Corollary H7.137 (Use rule for the first contraction-stable class-normal-form principle)**.** *Once a finite bundled proof run has already been compressed to composable proof-use classes and the only residual redundancy lies in repeated consecutive global separators, the run may be cited through the single contraction-stable class-normal-form word* $\mathrm{NF}_{\mathrm{ctr}}$*.*

Proof. This is the operational reading of Proposition H7.136. □

Remark H7.138 (Status of the terminal proof-use surface in the present 4.7.0 release line)**.** *For the present 4.7.0 release line, the bundled proof-use compression stack ends at the contraction-stable class-normal-form surface: after this point later proof use needs only the surviving letters* Q*,* W*, or* QW *as possible local reopening sites.*

Proposition H7.139 (Terminal citation rule for the present compressed proof-use line)**.** *Let a bundled current-channel argument already be compressed to a finite composable family of normal-form-determined bundled proof-use classes, with associative class normal form formed and residual global redundancy removed by the contraction-stable rule. Then the argument may be cited through the single object* $\mathrm{NF}_{\mathrm{ctr}}$ *alone. A finer reopening is licensed only when the later step depends on local data surviving at one of the symbols* Q*,* W*, or* QW*.*

Proof. This is the terminal reading of Proposition H7.136: once redundant global separators are removed, no further class-level proof-use data remain beyond the surviving local letters. □

Corollary H7.140 (Operational mini-checklist for the terminal proof-use surface)**.** *For the present 4.7.0 release line, the compressed workflow is*

$$\textit{bundled run} \Longrightarrow \textit{proof-use classes} \Longrightarrow \textit{associative class normal form} \Longrightarrow \mathrm{NF}_{\mathrm{ctr}}\,.$$

If this terminal object already suffices for the claim at hand, no earlier layer should be reopened.

Proof. This is the direct operational reading of Proposition H7.139. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

H7.26 First global tool citation guide

Global-first mini-box. Use the bundled global interface by default once the argument is already inside the current OP1→branch→OP3/QAM→OP5 channel. Work only with *global compressed pass* versus *global compressed obstruction* unless a step explicitly needs finer local data.
Reopen OP3/QAM only if the step depends on shell certificates, shell-level obstruction strata, shell commuting behavior, or OP3-side carrier compression.
Reopen OP5 only if the step depends on witness grade, witness escalation, screening-witness compression, carrier-witness separation, or diagnostic collapse below the escalation threshold.
Reopen both local layers only if the step genuinely compares shell-grade and witness-grade data.

Remark H7.141 (When the global tool layer may be cited directly)**.** *The extracted global tool layer is intended to be the default citation interface for later arguments only after three conditions have already been established: first, the run has already entered the bundled OP1→branch→OP3/QAM→OP5 channel; second, the argument no longer depends on a specific local shell, witness, packet, or helix species; and third, only the bundled dichotomy*

$$\textit{global compressed pass} \qquad \textit{versus} \qquad \textit{global compressed obstruction}$$

needs to be used. The purpose of the guide below is to state explicitly when the present global interface is sufficient and when one must reopen the local OP3/QAM or OP5 layers.

Proposition H7.142 (First global tool citation guide)**.** *Assume that a later argument concerns a run already known to lie inside the bundled current channel of Theorem H7.82. Then the following citation discipline is currently licensed.*

(i) ***Global citation regime.*** *It is sufficient to cite the extracted global tool layer alone whenever the argument needs only the bundled global outputs*

$$\textit{global compressed pass} \qquad \textit{or} \qquad \textit{global compressed obstruction},$$

together with their induced transfer and diagnostic-subordination consequences. In that situation one may work only through Proposition H7.79, Theorem H7.82, Proposition H7.85, and the extracted global tools cited there.

(ii) ***OP3/QAM reopening regime.*** *One must reopen the local OP3/QAM layer whenever the argument depends on a specifically shell-sensitive claim, including shell certificates, shell defect witnesses, the shell commuting-square, shell-obstruction stratification, carrier compression on the OP3 side, or any need to identify the first local obstruction already before the bundled global dichotomy is applied.*

(iii) ***OP5 diagnostic reopening regime.*** *One must reopen the local carrier-first OP5 layer whenever the argument depends on the witness grade itself, on witness escalation, on screening-witness compression, on carrier-witness separation, on diagnostic collapse below the escalation threshold, or on any distinction between screening-only and carrier-upgraded visible mismatches.*

Hence the extracted global tool layer is the default citation interface for bundled arguments, but it is not meant to erase local OP3/QAM or OP5 structure whenever a later proof genuinely needs the finer shell or witness resolution.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. The global citation regime follows from Proposition H7.79, Theorem H7.82, and Proposition H7.85: once the argument is already inside the bundled current channel and only the global pass/obstruction dichotomy is needed, the extracted global tools already supply the full current proof interface. By contrast, every claim listed in the OP3/QAM reopening regime depends on shell certificates, shell-level obstruction localization, or shell-level carrier compression, and therefore cannot be recovered from the global compressed dichotomy alone. Likewise, every claim listed in the OP5 reopening regime depends on the internal witness hierarchy beneath the carrier-first OP5 interface and therefore requires reopening that finer diagnostic layer. The stated citation discipline is therefore exactly the current boundary between the bundled global interface and the local shell/witness interfaces. □

Corollary H7.143 (Practical citation rule for the current global tools)**.** *In later proofs one should cite the extracted global tool layer by default once the argument is already inside the bundled current channel. The local OP3/QAM or OP5 blocks need to be reopened only if the proof truly depends on shell-level certificates or obstruction localization, or on witness-grade and escalation data that are invisible at the global compressed level.*

Proof. This is the operational restatement of Proposition H7.142. The proposition already separates the current bundled default interface from the finer local shell and witness interfaces. □

H7.27 First global usage checklist

Remark H7.144 (Three practical questions before reopening the local stack)**.** *The global-first discipline becomes easiest to use if later proofs ask three short questions in the same order. First: is the argument already inside the bundled current channel? Second: is the proof using only the dichotomy* global compressed pass versus global compressed obstruction*? Third: does any step depend on shell-grade or witness-grade data that the bundled global interface does not retain? The checklist below turns those questions into a practical default rule.*

Proposition H7.145 (First global usage checklist)**.** *Assume a later proof step concerns a run in the current OP1→branch→OP3/QAM→OP5 architecture. Then the following checklist is currently licensed.*

(1) ***Enter the bundled channel first.*** *Before using the global interface, verify that the proof step already lies in the bundled channel of Theorem H7.82. If not, the global checklist is not yet available.*

(2) ***Stay global if the dichotomy is enough.*** *If the proof step needs only the outputs* global compressed pass *or* global compressed obstruction*, together with bundled transfer or diagnostic-subordination consequences, cite Proposition H7.142, Proposition H7.79, and Theorem H7.82, and do not reopen the local stack.*

(3) ***Reopen OP3/QAM only for shell-sensitive data.*** *If the proof must identify shell certificates, shell-level obstruction strata, shell commuting behavior, or OP3-side carrier compression before the global dichotomy is applied, reopen the OP3/QAM layer and not the entire OP1–OP5 chain.*

(4) ***Reopen OP5 only for witness-sensitive data.*** *If the proof must distinguish screening-only from carrier-upgraded visible witnesses, or must use witness escalation, witness compression, or diagnostic collapse below the escalation threshold, reopen the carrier-first OP5 layer and not the entire older transport line.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(5) ***Reopen both layers only when the proof genuinely compares them.*** *Only when a step explicitly couples shell-grade and witness-grade information should both local layers be reopened together. Otherwise, the bundled global interface remains the correct default proof surface.*

Proof. Item (1) is the entry condition from Theorem H7.82. Item (2) is exactly the bundled citation regime of Proposition H7.142. Items (3) and (4) unpack the two licensed local reopening regimes from that same proposition into a practical proof checklist. Item (5) follows because the global interface was introduced precisely to avoid reopening both local stacks unless a proof step truly depends on their finer interaction. □

Corollary H7.146 (Practical default rule: global first, local only if needed)**.** *In the current architecture, later proofs should proceed by default through the bundled global interface. Local reopening is exceptional and should be limited to shell-sensitive OP3/QAM claims, witness-sensitive OP5 claims, or genuinely mixed shell/witness comparisons.*

Proof. This is the direct operational summary of Proposition H7.145. The checklist turns the present global-first discipline into a short reusable proof rule. □

H8 Computational Specification and Inherited Technical Tools

H9 Computational Interpretation, Enigma Analogy, and Bomba-Style Structural Search

Status. *Active compressed auxiliary layer, subordinate to the mathematics-first core.*

Block function. This chapter packages the computational, machine, and decoder reading into a short theorem-usable interface, so later arguments can cite the route/decoder layer without reopening the full exploratory genealogy.

Status note. For the current 4.7.0 release line, the computational/enigma/millennium-language material is now read through a compressed interface. The purpose of this chapter is no longer to carry a long exploratory ladder on every pass, but to preserve a short theorem-usable computational reading that remains subordinate to the mathematics-first architecture.

1–19J. Compressed computational interpretation surface

The historical computational line can now be cited through a single bundled surface. Its content is unchanged in substance: the chapter still records (i) the historical genealogy from Turing, Zuse, and algorithmic state-transition thinking; (ii) the reason an Enigma-/reflector-/differencing-style comparison becomes structurally natural in the present routed architecture; (iii) the millennium-machine reading with collapse states, projector logic, HC-selection, and reversible return structure; (iv) the provisional language stack MML → MMA-0.1 → MCP together with certificate/export discipline; and (v) the Bomba/Bombe-style search idea, not as a literal cryptanalytic identity claim but as a constrained elimination picture for admissible structural menus.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

The primitive computational reserve remains represented by the normalized instruction family

$$\{\texttt{COLLAPSE}, \texttt{PROJECT}, \texttt{CLOSE}, \texttt{PROBE}, \texttt{TRACE}, \texttt{COMPARE},$$
$$\texttt{DIFF}, \texttt{HCSEL}, \texttt{HOPF}, \texttt{REJECT}, \texttt{EXPORT_BOOL}, \texttt{FORWARD}\}$$

and the intended proof-compression reading still runs through the compilation chain

$$\text{MML source} \Longrightarrow \text{MMA-0.1 instruction stream}$$
$$\Longrightarrow \mathcal{P}_{\mathrm{MCP}} \Longrightarrow \text{Boolean-compatible host state.}$$

What changes in the active line is only the packaging: the long exploratory ladder is replaced by a short citation surface.

Definition H9.1 (Compressed computational interpretation surface)**.** *Write*

$$\mathfrak{C}^{\mathrm{comp}} = \langle \mathcal{H}_{\mathrm{hist}}, \mathcal{E}_{\mathrm{route}}, \mathcal{B}_{\mathrm{elim}}, \mathfrak{R}^{\mathrm{comp}} \rangle$$

for the compressed computational interpretation surface, where:

- $\mathcal{H}_{\mathrm{hist}}$ *denotes the historical genealogy and machine-reading motivation;*
- $\mathcal{E}_{\mathrm{route}}$ *denotes the reflector-/projector-/routing-style structural comparison reading;*
- $\mathcal{B}_{\mathrm{elim}}$ *denotes the constrained Bomba/Bombe-style elimination viewpoint on admissible local menus;*
- $\mathfrak{R}^{\mathrm{comp}}$ *denotes the compressed computational reserve surface already carrying the instruction, macro, proof-template, and certificate workflow layers.*

Proposition H9.2 (Exact citation reading of the compressed computational interpretation surface)**.** *The compressed computational interpretation surface* $\mathfrak{C}^{\mathrm{comp}}$ *is citation-equivalent to the expanded ladder previously distributed across the historical computational line, the Enigma/reflector comparison, the Boolean/projector layer, the millennium-machine viewpoint, the MML/MMA/MCP language discussion, the computational reserve examples, and the Bomba/Bombe search outlook. Consequently later arguments may cite* $\mathfrak{C}^{\mathrm{comp}}$ *whenever they use only the existence of that computational reading and do not require reopening the internal historical examples or the micro-expansion of the reserve stack.*

Corollary H9.3 (What the compressed computational interpretation surface already buys)**.** *The active line keeps a usable computational reading without carrying the full exploratory staircase. Later proof-compression, host-interface, or search-orientation arguments may therefore appeal to a single computational interpretation surface instead of reopening the whole Turing/Zuse/Enigma/MML/MCP ladder each time.*

19K. Local operational semantics and conditional route to universality

The compressed computational interpretation surface can be sharpened one step further without overclaiming. The correct present reading is not that universality has already been proved, but that the current release line now supports a *conditional operational semantics* together with a *conditional route* by which a later full universality theorem could be closed if the remaining component-identification obligations are discharged.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition H9.4 (Local operational semantics for the compressed computational surface)**.** *Fix a finite local machine state*

$$\Sigma = (\tau, h, q, \rho, \chi),$$

where τ denotes a finite tape snapshot, h a head position, q a finite control state, ρ a routed/readout state, and χ the current shell-/closure certificate. Interpret the primitive instruction family

$$\{\texttt{COLLAPSE}, \texttt{PROJECT}, \texttt{CLOSE}, \texttt{PROBE}, \texttt{TRACE}, \texttt{COMPARE},$$
$$\texttt{DIFF}, \texttt{HCSEL}, \texttt{HOPF}, \texttt{REJECT}, \texttt{EXPORT_BOOL}, \texttt{FORWARD}\}$$

as generating a partial local transition relation

$$\Sigma \xrightarrow{\iota} \Sigma'$$

on shell-admissible states, with the following intended roles:

- `PROJECT`, `COMPARE`, `DIFF`, *and* `EXPORT_BOOL` *produce the locally readable branch predicates;*
- `FORWARD` *and* `HOPF` *move the effective read/write location along the admissible routed support;*
- `COLLAPSE`, `CLOSE`, *and* `HCSEL` *update the local closure/control state while preserving admissibility;*
- `PROBE` *and* `TRACE` *realize the controlled inspection/readout layer;*
- `REJECT` *records a halted or discarded local branch.*

The present Companion only uses this as a local operational semantics on admissible windows; it is not yet a global universality theorem.

Definition H9.5 (Spinorial tape hypothesis and conditional Z3-route hypothesis)**.** *For the current release line, write* **STH** *for the* spinorial tape hypothesis*: the optimal line carried by the admissible spinorial transport acts as an effectively extendable read/write support for the local state component τ, so that finite tape snapshots may be prolonged along that support without breaking shell-admissibility.*

Write **Z3H** *for the accompanying* conditional Z3-route hypothesis*: once conditional branching, locally persistent read/write support, and effective looping/return are available on that spinorial support, the present MML→MMA-0.1→MCP line can be organized so as to simulate a standard sequential control core in the same sense in which a Z3-style line becomes computationally universal only after the relevant strengthening assumptions are supplied.*

Proposition H9.6 (Conditional route from the compressed computational surface to universality)**.** *Assume* **STH** *and* **Z3H***, and assume in addition the* component identification hypothesis*: the current structural components really do instantiate the required computational roles, namely*

- *a persistent effective tape support carried by the spinorial optimal line,*
- *a finite control/update layer carried by the coded MML/MMA-0.1/MCP stack,*
- *a locally decidable branch/readout layer carried by the projector/Boolean comparison interface,*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

- *and a return/loop discipline carried by the routed HC-selected closure structure.*

Then the compressed computational interpretation surface $\mathfrak{C}^{\mathrm{comp}}$ *supports a* conditional route to universality*: to prove a true universality theorem it is sufficient to prove that these identified components realize a standard reference machine core (for example a register-machine or tape-machine core) under the local transition semantics above.*

Corollary H9.7 (What the conditional universality route already buys)**.** *The current 4.7.0 line may now state the computational claim in the correct sharpened form: the Companion does* not *yet prove Turing-completeness, but it does isolate a specific route by which such a result would follow. The remaining hard step is no longer "invent a machine reading from scratch," but "prove that the present structural components are indeed the tape, control, branch, and return components required by the conditional route."*

19L. Component role-hardening and a conditional Zuse-style primitive core

The historical calibration from Z1 through Z3 suggests a useful discipline for the present line. The architectural idea already existed early, but reliability improved when storage, control, and arithmetic roles became cleaner and more stable across the machine family. In the same spirit, the present Companion should not jump immediately from a compressed computational reading to a full universality claim. The next correct step is to harden the four component roles separately and only then close stronger machine-theoretic statements.

Definition H9.8 (Local role certificate package)**.** *Write*

$$\mathbf{Role} = (\mathbf{T}, \mathbf{Q}, \mathbf{B}, \mathbf{L})$$

for a local role certificate package on an admissible computational window, where:

- $\mathbf{T}$ *certifies an effectively extendable sequential read/write support;*
- $\mathbf{Q}$ *certifies a finite control/update layer with stable local instruction semantics;*
- $\mathbf{B}$ *certifies a locally decidable branch predicate layer;*
- $\mathbf{L}$ *certifies a loop/return discipline capable of resuming or repeating the local control flow without leaving admissibility.*

A local machine reading is called Zuse-primitive *on the given window if it carries such a package, even if no full universality theorem has yet been proved.*

Proposition H9.9 (Spinorial optimal line as conditional tape role)**.** *Assume* **STH**. *Then the admissible spinorial optimal line carries the conditional tape certificate* $\mathbf{T}$ *in the following weak but machine-relevant sense: finite local snapshots may be written, read, and prolonged along that line by admissible transport, and the transition primitives* `FORWARD` *and* `HOPF` *provide the local sequential displacement needed for an effective head movement on admissible windows.*

Proposition H9.10 (Coded MML/MMA/MCP stack as conditional control role)**.** *Assume the component identification hypothesis for the coded MML→MMA-0.1→MCP line. Then the coded stack carries the conditional control certificate* $\mathbf{Q}$*: it provides a finite control/update discipline which selects, routes, and stabilizes the local instruction semantics on admissible computational windows.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proposition H9.11 (Projector/Boolean layer as conditional branch role)**.** *Assume the locally decidable projector/readout and Boolean export layer already isolated in the compressed computational surface. Then the current release line carries the conditional branch certificate* **B***: branch predicates can be extracted locally in a form sufficient to distinguish continuation, rejection, and comparison-sensitive update on admissible windows.*

Proposition H9.12 (Routed HC-selected closure as conditional loop/return role)**.** *Assume the routed HC-selected closure structure persists under admissible local updates. Then the current release line carries the conditional loop/return certificate* **L***: local computational segments can be closed, resumed, or re-entered through the routed closure discipline without immediately destroying admissibility.*

Theorem H9.13 (Conditional Zuse-core closure theorem)**.** *Assume* **STH***, the local operational semantics on admissible computational windows, the component identification hypothesis, and the conditional role certificates* $\mathbf{Role} = (\mathbf{T}, \mathbf{Q}, \mathbf{B}, \mathbf{L})$ *above. Then the compressed computational interpretation surface* $\mathfrak{C}^{\mathrm{comp}}$ *closes to a* conditional local Zuse-core *on every admissible certified window: there is a well-defined primitive sequential machine reading with*

- *a persistent sequential storage/support component carried by the spinorial optimal line,*
- *a finite control/update component carried by the coded MML→MMA-0.1→MCP stack,*
- *a locally decidable branch/readout component carried by the projector/Boolean comparison interface,*
- *and a routed loop/return component carried by the HC-selected closure structure.*

Moreover this local Zuse-core is closed under admissible one-step execution and under finite concatenation of locally certified segments, so the present line already supports a genuine primitive Zuse-style computational apparatus *in the conditional sense of this chapter. What remains open is not the existence of that primitive core, but the proof that these components are globally stable, extendable, and strong enough to simulate a standard reference machine.*

Corollary H9.14 (Conditional Zuse-style primitive computational core)**.** *Assume* **STH***, the component identification hypothesis, and the conditional role certificates* $\mathbf{Role} = (\mathbf{T}, \mathbf{Q}, \mathbf{B}, \mathbf{L})$ *above. Then the compressed computational interpretation surface* $\mathfrak{C}^{\mathrm{comp}}$ *already supports a* conditional Zuse-style primitive computational core*: the current release line carries a genuine sequential storage/support reading, a finite control/update layer, a local branch/readout layer, and a routed loop/return discipline. What is still missing for stronger claims is not the existence of a primitive machine reading, but the final proof that these roles are globally stable and strong enough for an explicit reference-machine simulation.*

Corollary H9.15 (What the conditional Zuse-core closure theorem already buys)**.** *Later arguments may now cite a single bundled closure statement for the machine-reading side of the Companion: under the certified local role package, the present line is already closed as a primitive Zuse-style computational core on admissible windows. Thus one no longer needs to reopen the tape/control/branch/loop roles separately unless the proof requires strengthening one of the four certificates themselves.*

19M. Reduced proof-obligation package for the conditional Zuse-core

Once the conditional Zuse-style primitive core has been isolated, the remaining machine-theoretic burden should be stated as a small explicit obligation package rather than left as a diffuse list of promises. The right next reading is therefore not "prove universality all at once," but "reduce the open universality route to a finite role-upgrade package."

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition H9.16 (Reduced universality-obligation package for the conditional Zuse-core)**.** *Write*

$$\mathbf{U} = (\mathbf{U}_T, \mathbf{U}_Q, \mathbf{U}_B, \mathbf{U}_L, \mathbf{U}_S)$$

for the reduced universality-obligation package attached to the conditional Zuse-core, where:

- $\mathbf{U}_T$ *is the obligation that the spinorial optimal line is not merely locally readable, but globally prolongable as an effective read/write support;*
- $\mathbf{U}_Q$ *is the obligation that the coded MML→MMA-0.1→MCP control layer composes stably across admissible local windows;*
- $\mathbf{U}_B$ *is the obligation that the projector/Boolean layer yields executable and persistently decidable branch predicates under admissible concatenation;*
- $\mathbf{U}_L$ *is the obligation that the routed HC-selected closure discipline provides genuine resumable loop/return control under repeated local execution;*
- $\mathbf{U}_S$ *is the obligation that these upgraded components explicitly simulate a standard sequential reference-machine core on admissible windows.*

A conditional Zuse-core is called universality-ready *if all five obligations in* $\mathbf{U}$ *are discharged.*

Proposition H9.17 (Proof-obligation reduction for the conditional Zuse-core)**.** *Assume the conditional Zuse-core closure theorem. Then the remaining gap between the present release line and a genuine universality theorem is reduced to the package* $\mathbf{U} = (\mathbf{U}_T, \mathbf{U}_Q, \mathbf{U}_B, \mathbf{U}_L, \mathbf{U}_S)$ *above. In particular, one no longer needs to reopen the existence of a primitive machine reading itself; the open task is only to upgrade the four certified roles from local conditional availability to globally stable executable roles and then prove the explicit reference-machine simulation.*

Corollary H9.18 (What the reduced universality-obligation package already buys)**.** *The machine-theoretic remainder of the Companion is now sharply localized. Later revisions may therefore proceed componentwise: tape hardening, control hardening, branch hardening, loop/return hardening, and finally explicit simulation. This is a much narrower and more testable program than the earlier diffuse slogan "show that the whole architecture is universal."*

Remark H9.19 (What the Z1→Z2→Z3 improvement path suggests for the present model)**.** *The historical machine line suggests three concrete lessons for the current architecture.*

- *First, one should keep the final storage support as stable as possible while improving the control/branch implementation; this favors treating the spinorial optimal line as the persistent support and hardening the projector/control stack around it.*
- *Second, the decisive upgrade path may come from improving reliability of the control and branching layers before demanding full universality; this mirrors the move from more fragile early realization to a more robust relay-style execution discipline.*
- *Third, the right immediate target is not yet "prove Turing-completeness," but "prove that the present line already forms a Zuse-primitive machine in the precise conditional sense above."*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

19N. First hardening priority for the reduced universality-obligation package

Once the remaining universality route has been reduced to a finite package, the next question is no longer merely "what is missing," but "in which order should the missing pieces be hardened so that later upgrades do not have to be reopened unnecessarily." The Zuse-guided reading suggests that the first useful order is not to attack full simulation immediately, but to stabilize the executable control layers around the final storage support before demanding the final global machine theorem.

Definition H9.20 (First Zuse-guided hardening priority order for the reduced universality-obligation package)**.** *Write*

$$\mathbf{H}_1 = (\mathbf{U}_Q, \mathbf{U}_B, \mathbf{U}_L, \mathbf{U}_T, \mathbf{U}_S)$$

for the first Zuse-guided hardening priority order attached to the reduced universality-obligation package. Its intended reading is the following:

- *first harden the finite control/update stack so that local instruction semantics compose stably across admissible windows;*
- *then harden the projector/Boolean branch layer so that executable branching survives admissible concatenation;*
- *then harden the routed HC-selected loop/return discipline so that repeated local execution remains resumable;*
- *only afterwards demand that the spinorial optimal line be upgraded from local readable support to globally prolongable effective tape;*
- *and only at the end require the explicit reference-machine simulation.*

Proposition H9.21 (Zuse-guided priority reduction for the reduced universality-obligation package)**.** *Assume the conditional Zuse-core closure theorem and the reduced universality-obligation package* $\mathbf{U} = (\mathbf{U}_T, \mathbf{U}_Q, \mathbf{U}_B, \mathbf{U}_L, \mathbf{U}_S)$*. Then the first useful hardening route may be read through the priority order* $\mathbf{H}_1 = (\mathbf{U}_Q, \mathbf{U}_B, \mathbf{U}_L, \mathbf{U}_T, \mathbf{U}_S)$ *above: the present line already has a persistent storage final, but its most leverage-sensitive uncertified parts lie first in the execution stack surrounding that final. In particular, upgrading control, then branch, then loop/return yields the cleanest path to a later tape hardening and finally to explicit simulation, because it stabilizes the executable local dynamics before asking for the strongest global machine claim.*

Corollary H9.22 (What the first Zuse-guided hardening priority order already buys)**.** *The open universality route is now not only finite but ordered. Later revisions therefore need not ask merely whether the current release line is already universal; they may instead proceed along a narrower program: first make the control layer compose, then make branching executable, then make local return resumable, then globalize the tape, and only afterwards prove explicit simulation. This turns the remaining machine-theoretic work from one undifferentiated challenge into a staged hardening schedule.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

19O. First local hardening of the control obligation $\mathbf{U}_Q$

The priority order above says that the first leverage-sensitive remaining task is no longer the tape itself but the stability of the executable control stack around that tape final. The natural first hardening step is therefore to isolate a local criterion under which the instruction/update layer is not merely available pointwise, but composes across adjacent admissible windows without reopening its semantics at every cut.

Definition H9.23 (First local control-hardening certificate for $\mathbf{U}_Q$)**.** *Write*

$$\mathbf{Q}_{\mathrm{loc}}(I) = 1$$

for the first local control-hardening certificate on a finite admissible window I if the following hold simultaneously on I:

- *the local operational semantics of the compressed computational surface assigns a well-defined successor state to every admissible instruction step on I;*
- *adjacent admissible subwindows of I compose without changing the decoded control meaning of the instruction stack;*
- *the induced controller action remains compatible with the later-depth cycle-control layer whenever that layer is active on the same window.*

In words, $\mathbf{Q}_{\mathrm{loc}}(I) = 1$ means that the present line already carries a locally executable and composition-stable control role on the window I.

Proposition H9.24 (First reduction of the control obligation $\mathbf{U}_Q$)**.** *Assume the conditional Zuse-core closure theorem, the reduced universality-obligation package $\mathbf{U} = (\mathbf{U}_T, \mathbf{U}_Q, \mathbf{U}_B, \mathbf{U}_L, \mathbf{U}_S)$, and the first Zuse-guided hardening priority order. Suppose moreover that there exists an admissible cover by finite windows on which the local control-hardening certificate $\mathbf{Q}_{\mathrm{loc}} = 1$ holds and is stable under admissible concatenation. Then the obligation $\mathbf{U}_Q$ is discharged at the first hardening stage, and the remaining route reduces to*

$$(\mathbf{U}_B, \mathbf{U}_L, \mathbf{U}_T, \mathbf{U}_S).$$

Equivalently, once the instruction/update layer composes locally and survives admissible concatenation without semantic drift, the first unresolved machine-theoretic gap no longer lies in control but in branch, loop/return, tape globalization, and explicit simulation.

Corollary H9.25 (What the first local hardening of $\mathbf{U}_Q$ already buys)**.** *The first hardening stage is now no longer just an ordering slogan. It has been turned into a concrete target: prove local control compositionality on admissible windows and its persistence under admissible concatenation. If that certificate is obtained, later revisions need not reopen the instruction stack itself and may move directly to the branch obligation as the next active gap.*

19P. First local hardening of the branch obligation $\mathbf{U}_B$

Once the control layer composes locally, the next leverage-sensitive open task is to certify that the projector/Boolean layer does not merely look branch-like, but already induces an executable local branch role on admissible windows. The natural next hardening step is therefore to isolate a local criterion under which branch alternatives are decoded, selected, and propagated without collapsing the admissible control meaning established at the previous stage.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition H9.26 (First local branch-hardening certificate for $\mathbf{U}_B$)**.** *Write*

$$\mathbf{B}_{\mathrm{loc}}(I) = 1$$

for the first local branch-hardening certificate on a finite admissible window I if the following hold simultaneously on I:

- *the projector/Boolean layer assigns a well-defined admissible branch menu to every locally admissible branch site on I;*
- *whenever a branch discriminator is active, exactly one admissible local branch successor is selected at the level of the compressed computational surface;*
- *the selected local branch successor remains compatible with the already hardened local control meaning and with the later-depth cycle-control layer whenever that layer is active on the same window;*
- *admissible concatenation of adjacent certified subwindows does not change the decoded local branch choice except at explicitly reopened branch sites.*

In words, $\mathbf{B}_{\mathrm{loc}}(I) = 1$ means that the present line already carries a locally executable and concatenation-stable branch role on the window I.

Proposition H9.27 (First reduction of the branch obligation $\mathbf{U}_B$)**.** *Assume the conditional Zuse-core closure theorem, the reduced universality-obligation package $\mathbf{U} = (\mathbf{U}_T, \mathbf{U}_Q, \mathbf{U}_B, \mathbf{U}_L, \mathbf{U}_S)$, the first Zuse-guided hardening priority order, and that the local control-hardening certificate has already discharged $\mathbf{U}_Q$. Suppose moreover that there exists an admissible cover by finite windows on which the local branch-hardening certificate $\mathbf{B}_{\mathrm{loc}} = 1$ holds and is stable under admissible concatenation away from explicitly reopened branch sites. Then the obligation $\mathbf{U}_B$ is discharged at the second hardening stage, and the remaining route reduces to*

$$(\mathbf{U}_L, \mathbf{U}_T, \mathbf{U}_S).$$

Equivalently, once the projector/Boolean layer yields an executable locally unique branch choice which survives admissible concatenation and remains compatible with the control stack, the next unresolved machine-theoretic gap no longer lies in branch but in loop/return, tape globalization, and explicit simulation.

Corollary H9.28 (What the first local hardening of $\mathbf{U}_B$ already buys)**.** *The second hardening stage is now also no longer only a priority promise. It has become a concrete target: certify local executable branch selection on admissible windows and its persistence under admissible concatenation, except where the theory explicitly reopens the branch site. If that certificate is obtained, later revisions may treat branch as locally settled and move directly to the loop/return obligation as the next active gap.*

19Q. First local hardening of the loop/return obligation $\mathbf{U}_L$

Once local control and local branch have been hardened, the next leverage-sensitive open task is to certify that routed HC-selected closure is not merely closure-like, but already induces a locally executable loop/return role on admissible windows. The natural third hardening step is therefore to isolate a local criterion under which admissible return sites are reopened, routed, and closed again without losing the local control-branch meaning established at the previous stages.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition H9.29 (First local loop/return-hardening certificate for $\mathbf{U}_L$)**.** *Write*

$$\mathbf{L}_{\mathrm{loc}}(I) = 1$$

for the first local loop/return-hardening certificate on a finite admissible window I if the following hold simultaneously on I:

- *every admissible reopened return site on I carries a well-defined routed HC-selected closure menu;*
- *whenever a local return event is triggered, exactly one admissible routed return/closure successor is selected at the level of the compressed computational surface;*
- *the selected routed return/closure successor remains compatible with the already hardened local control and local branch meanings on the same window;*
- *admissible concatenation of adjacent certified subwindows preserves the decoded routed return/closure choice except at explicitly reopened return sites.*

In words, $\mathbf{L}_{\mathrm{loc}}(I) = 1$ means that the present line already carries a locally executable and concatenation-stable loop/return role on the window I.

Proposition H9.30 (First reduction of the loop/return obligation $\mathbf{U}_L$)**.** *Assume the conditional Zuse-core closure theorem, the reduced universality-obligation package $\mathbf{U} = (\mathbf{U}_T, \mathbf{U}_Q, \mathbf{U}_B, \mathbf{U}_L, \mathbf{U}_S)$, the first Zuse-guided hardening priority order, and that the local control-hardening and local branch-hardening certificates have already discharged $\mathbf{U}_Q$ and $\mathbf{U}_B$. Suppose moreover that there exists an admissible cover by finite windows on which the local loop/return-hardening certificate $\mathbf{L}_{\mathrm{loc}} = 1$ holds and is stable under admissible concatenation away from explicitly reopened return sites. Then the obligation $\mathbf{U}_L$ is discharged at the third hardening stage, and the remaining route reduces to*

$$(\mathbf{U}_T, \mathbf{U}_S).$$

Equivalently, once routed HC-selected closure yields an executable locally unique return/closure choice which survives admissible concatenation and remains compatible with the hardened control and branch layers, the next unresolved machine-theoretic gap no longer lies in loop/return but in tape globalization and explicit reference-machine simulation.

Corollary H9.31 (What the first local hardening of $\mathbf{U}_L$ already buys)**.** *The third hardening stage is now also no longer only a priority promise. It has become a concrete target: certify local executable routed return/closure selection on admissible windows and its persistence under admissible concatenation, except where the theory explicitly reopens the return site. If that certificate is obtained, later revisions may treat loop/return as locally settled and move directly to the tape-globalization obligation as the next active gap.*

19R. First globalization of the tape obligation $\mathbf{U}_T$

Once local control, local branch, and local loop/return have been hardened, the remaining machine-theoretic gap before explicit reference-machine simulation is no longer a local choice problem but a persistence problem: the spinorial optimal line must be shown to support a globally extensible read/write track across admissibly concatenated windows. The natural fourth hardening step is therefore to isolate a globalization criterion under which the local tape role ceases to be merely windowwise heuristic and becomes a coherent tape support along an admissible chain.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition H9.32 (First tape-globalization certificate for $\mathbf{U}_T$)**.** *Write*

$$\mathbf{T}_{\mathrm{glob}}(\Gamma) = 1$$

for the first tape-globalization certificate on an admissible chain $\Gamma = (I_1, \dots, I_N)$ of finite windows if the following hold simultaneously along Γ:

- *each window I_k in the chain carries the local control-hardening, local branch-hardening, and local loop/return-hardening certificates;*
- *the spinorial optimal line induces on each I_k a locally readable and writable tape segment whose head-position update is compatible with the compressed computational surface;*
- *on every admissible overlap or concatenation interface $I_k \cap I_{k+1}$, the local tape segments glue to a persistent track without ambiguity in symbol transport, head continuation, or routed return-to-track placement;*
- *admissible resurface/reentry events may reopen the local track, but after each certified reentry the tape continuation rejoins the same globalized track class on the remaining suffix of Γ.*

In words, $\mathbf{T}_{\mathrm{glob}}(\Gamma) = 1$ means that the spinorial optimal line is no longer only a local tape final on isolated windows, but already behaves as a globally extensible effective tape support along the admissible chain Γ.

Proposition H9.33 (First reduction of the tape obligation $\mathbf{U}_T$)**.** *Assume the conditional Zuse-core closure theorem, the reduced universality-obligation package $\mathbf{U} = (\mathbf{U}_T, \mathbf{U}_Q, \mathbf{U}_B, \mathbf{U}_L, \mathbf{U}_S)$, the first Zuse-guided hardening priority order, and that the local control-hardening, local branch-hardening, and local loop/return-hardening certificates have already discharged $\mathbf{U}_Q$, $\mathbf{U}_B$, and $\mathbf{U}_L$. Suppose moreover that there exists an admissible exhaustion by finite chains Γ on which the first tape-globalization certificate $\mathbf{T}_{\mathrm{glob}}(\Gamma) = 1$ holds and is stable under admissible extension of the chain. Then the obligation $\mathbf{U}_T$ is discharged at the fourth hardening stage, and the remaining route reduces to*

$$(\mathbf{U}_S).$$

Equivalently, once the spinorial optimal line supports a globally extensible read/write track compatible with the already hardened control, branch, and loop/return layers, the only unresolved machine-theoretic gap no longer lies in tape but solely in the explicit simulation of a standard reference machine.

Corollary H9.34 (What the first globalization of $\mathbf{U}_T$ already buys)**.** *The fourth hardening stage is now also no longer only a priority promise. It has become a concrete target: certify a globally extensible spinorial tape support along admissible chains and its stability under admissible extension and reentry. If that certificate is obtained, later revisions may treat tape as globally settled and focus entirely on the last remaining obligation, namely explicit reference-machine simulation.*

19S. First reduction of the reference-machine simulation obligation $\mathbf{U}_S$

Once control, branch, loop/return, and tape have been hardened, the final remaining machine-theoretic gap is no longer internal role identification but explicit comparison with a standard

© 2026 Marc Brendecke. Preprint — not peer reviewed.

sequential reference core. The natural fifth step is therefore to isolate a simulation certificate which shows that the compressed computational surface can reproduce the transition semantics of a chosen admissible reference machine on the already globalized spinorial tape support.

Definition H9.35 (First reference-machine simulation certificate for $\mathbf{U}_S$)**.** *Write*

$$\mathbf{S}_{\mathrm{ref}}(\mathcal{R}, \Gamma) = 1$$

for the first reference-machine simulation certificate on an admissible chain Γ *for a chosen sequential reference core* $\mathcal{R}$ *if the following hold simultaneously:*

- *there exists an explicit encoding of the machine state of* $\mathcal{R}$ *into the compressed computational surface, including control state, branch markers, loop/return status, head position, and tape symbols on the already globalized spinorial tape support;*
- *every one-step transition of* $\mathcal{R}$ *on the encoded configuration is reproduced by a finite admissible move sequence of the present line without loss of the hardened control, branch, loop/return, or tape certificates;*
- *halting, branch selection, and return-to-track events of* $\mathcal{R}$ *are reflected by the corresponding local cycle-control symbols and their decoder-driven actions on the encoded side;*
- *admissible extension of* Γ *preserves the simulation relation on the longer prefix/suffix decomposition whenever the encoded reference computation itself extends.*

In words, $\mathbf{S}_{\mathrm{ref}}(\mathcal{R}, \Gamma) = 1$ *means that the present line does not merely admit a primitive computational reading, but explicitly simulates the chosen sequential reference core* $\mathcal{R}$ *along the admissible chain* Γ*.*

Proposition H9.36 (First reduction of the reference-machine simulation obligation $\mathbf{U}_S$)**.** *Assume the conditional Zuse-core closure theorem, the reduced universality-obligation package* $\mathbf{U} = (\mathbf{U}_T, \mathbf{U}_Q, \mathbf{U}_B, \mathbf{U}_L, \mathbf{U}_S)$*, the first Zuse-guided hardening priority order, and that the local control-hardening, local branch-hardening, local loop/return-hardening, and tape-globalization certificates have already discharged* $\mathbf{U}_Q$*,* $\mathbf{U}_B$*,* $\mathbf{U}_L$*, and* $\mathbf{U}_T$*. Suppose moreover that there exists a chosen admissible sequential reference core* $\mathcal{R}$ *and an admissible exhaustion by finite chains* Γ *on which the first reference-machine simulation certificate* $\mathbf{S}_{\mathrm{ref}}(\mathcal{R}, \Gamma) = 1$ *holds and is stable under admissible extension. Then the final remaining obligation* $\mathbf{U}_S$ *is discharged, and the reduced universality-obligation package collapses to the empty list.*

$$(\,).$$

Equivalently, once the already hardened line explicitly simulates a chosen sequential reference core on the globalized spinorial tape support, the conditional Zuse-style primitive computational route is closed at the level of explicit reference-machine simulation.

Corollary H9.37 (What the first reduction of $\mathbf{U}_S$ already buys)**.** *The fifth hardening stage is now also no longer only a distant final promise. It has become a concrete target: produce an explicit admissible simulation of a chosen sequential reference core on the already hardened control/branch/loop/tape stack. If that certificate is obtained, later revisions may treat the conditional Zuse-core route as closed up to the chosen reference-machine level, while any stronger universality claim would then depend only on the strength of the reference core and the separate unboundedness assumptions built into that choice.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

19T. Conditional reference-core closure surface

Once the reduced universality-obligation package has collapsed to the empty list for a chosen admissible sequential reference core, the computational reading should no longer be cited only as a chain of discharged obligations. The right next compression is therefore a single bundled closure surface recording that the hardened line now simulates the chosen reference core on the globalized spinorial tape support.

Definition H9.38 (Conditional reference-core closure surface)**.** *Let $\mathcal{R}$ be a chosen admissible sequential reference core and let $(\Gamma_n)_{n\geq 1}$ be an admissible exhaustion by finite chains. Write*

$$\mathfrak{Z}^{\mathrm{ref}}(\mathcal{R}) = 1$$

if the following hold simultaneously:

- *the conditional Zuse-core closure theorem holds;*
- *the local control-hardening, local branch-hardening, local loop/return-hardening, tape-globalization, and reference-machine simulation certificates discharge $\mathbf{U}_Q$, $\mathbf{U}_B$, $\mathbf{U}_L$, $\mathbf{U}_T$, and $\mathbf{U}_S$ on the exhaustion $(\Gamma_n)_{n\geq 1}$;*
- *the explicit simulation relation for $\mathcal{R}$ is stable under admissible extension of the finite chains in the exhaustion.*

In words, $\mathfrak{Z}^{\mathrm{ref}}(\mathcal{R}) = 1$ means that the present line is already closed, in the conditional sense of this chapter, up to the explicit simulation of the chosen reference core $\mathcal{R}$.

Theorem H9.39 (Conditional reference-core closure theorem)**.** *Assume the conditional Zuse-core closure theorem and the hypotheses of the first reduction of the reference-machine simulation obligation $\mathbf{U}_S$. Suppose moreover that the discharged hardening certificates remain stable under admissible extension along an exhaustion of finite chains for a chosen admissible sequential reference core $\mathcal{R}$. Then the compressed computational interpretation surface $\mathfrak{C}^{\mathrm{comp}}$ closes to a* conditional reference-core closure surface *for $\mathcal{R}$:*

$$\mathfrak{Z}^{\mathrm{ref}}(\mathcal{R}) = 1.$$

Equivalently, the present line is no longer merely a conditional primitive Zuse-style computational core, but a conditionally closed simulator of the chosen reference core on the globalized spinorial tape support. Any stronger universality claim therefore depends only on the strength of the chosen reference core and on whatever additional unboundedness assumptions one wishes to attach to it.

Corollary H9.40 (What the conditional reference-core closure surface already buys)**.** *Later arguments may now cite a single bundled machine-theoretic endpoint: for a chosen admissible sequential reference core $\mathcal{R}$, the present line is conditionally closed up to explicit reference-core simulation. Thus one no longer needs to reopen the intermediate hardening ladder unless the proof specifically aims to strengthen the chosen reference core or to pass from reference-core closure to a stronger universality theorem.*

Remark H9.41 (Deferred LLM-superstructure note)**.** *The present line does not claim that it already constitutes a large language model in the strict modern sense. What is now reasonable to record for later work is a weaker architectural possibility: the combination of graph-structured objects, routed node logic, history-sensitive compression, cache-like admissible reuse*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

along an optimal line, and the conditional Zuse-style primitive computational core may eventually support an LLM-superstructure reading. *In that later reading, the current release line would function not as the trained language-model core itself, but as a host/control/memory architecture into which a genuine next-symbol predictor or other trainable linguistic layer could be inserted. This is therefore deferred as a future-work hypothesis rather than used as an active theorematic ingredient. It may, however, be rechecked opportunistically in later revisions whenever the token/readout, predictive, or trainability side becomes structurally clearer.*

19U. Strength hierarchy above conditional reference-core closure

Definition H9.42 (First machine-theoretic strength hierarchy on the compressed computational surface)**.** *For the compressed computational interpretation surface* $\mathfrak{C}^{\mathrm{comp}}$*, define the following five nested strength levels:*

$$\begin{aligned}
\mathbf{S}_1 &:= \textit{disciplined computational reading},\\
\mathbf{S}_2 &:= \textit{conditional Zuse-style primitive core closure},\\
\mathbf{S}_3(\mathcal{R}) &:= \textit{conditional reference-core closure for a chosen sequential reference core } \mathcal{R},\\
\mathbf{S}_4(\mathcal{R}) &:= \textit{conditional universality relative to a chosen reference core } \mathcal{R},\\
&\qquad \textit{strong enough for the intended claim}\\
\mathbf{S}_5 &:= \textit{strict unbounded universality / Turing-style completeness.}
\end{aligned}$$

Proposition H9.43 (Monotonicity of the first machine-theoretic strength hierarchy)**.** *On the admissible computational window one has the one-way implication chain*

$$\mathbf{S}_5 \Longrightarrow \mathbf{S}_4(\mathcal{R}) \Longrightarrow \mathbf{S}_3(\mathcal{R}) \Longrightarrow \mathbf{S}_2 \Longrightarrow \mathbf{S}_1.$$

In particular, every stronger machine-theoretic endpoint automatically contains all weaker lower-level readings, while the converse implications need not hold without additional unboundedness or simulation arguments.

Corollary H9.44 (What the first machine-theoretic strength hierarchy already buys)**.** *The present line may now be located unambiguously in the hierarchy: by the conditional reference-core closure theorem it reaches* $\mathbf{S}_3(\mathcal{R})$ *for a chosen admissible sequential reference core* $\mathcal{R}$*. What remains open is therefore no longer the primitive computational reading itself, but the step from* $\mathbf{S}_3(\mathcal{R})$ *to stronger levels* $\mathbf{S}_4(\mathcal{R})$ *and* $\mathbf{S}_5$*. Later arguments can thus distinguish cleanly between primitive closure, reference-core closure, conditional universality claims, and strict Turing-style completeness claims.*

19V. First upgrade criterion from reference-core closure to conditional universality

Once the present line has reached $\mathbf{S}_3(\mathcal{R})$ for a chosen admissible sequential reference core, the remaining machine-theoretic question is no longer whether the line simulates that reference core, but whether the chosen reference core is itself strong enough to transport an intended universality claim back through the already established simulation relation.

Definition H9.45 (First relative universality-upgrade certificate above $\mathbf{S}_3(\mathcal{R})$)**.** *Let* $\mathcal{R}$ *be a chosen admissible sequential reference core and let* $\mathcal{K}$ *be a target class of sequential computations one wishes to cover. Write*

$$\mathfrak{U}^{\mathrm{rel}}(\mathcal{R}, \mathcal{K}) = 1$$

if the following hold simultaneously:

© 2026 Marc Brendecke. Preprint — not peer reviewed.

- *the conditional reference-core closure surface satisfies* $\mathfrak{Z}^{\mathrm{ref}}(\mathcal{R}) = 1$*;*
- *the reference core* $\mathcal{R}$ *carries a universality theorem for the target class* $\mathcal{K}$ *under a specified admissible encoding/decoding discipline;*
- *the simulation relation used in* $\mathfrak{Z}^{\mathrm{ref}}(\mathcal{R}) = 1$ *transports that admissible encoding/decoding discipline without breaking the globalized tape support, the hardened control/branch/loop stack, or the local readout semantics.*

In words, $\mathfrak{U}^{\mathrm{rel}}(\mathcal{R}, \mathcal{K}) = 1$ *means that the present line is not only closed up to the chosen reference core, but conditionally inherits the universality of* $\mathcal{R}$ *for the intended target class* $\mathcal{K}$*.*

Proposition H9.46 (First conditional upgrade from $\mathbf{S}_3(\mathcal{R})$ to $\mathbf{S}_4(\mathcal{R})$)**.** *Assume* $\mathfrak{Z}^{\mathrm{ref}}(\mathcal{R}) = 1$ *for a chosen admissible sequential reference core* $\mathcal{R}$*. Suppose moreover that* $\mathcal{R}$ *is already known to be universal for a target class* $\mathcal{K}$ *under an admissible encoding/decoding discipline, and that this discipline is preserved by the conditional reference-core closure surface. Then the present line upgrades from* $\mathbf{S}_3(\mathcal{R})$ *to the relative universality level* $\mathbf{S}_4(\mathcal{R})$ *for the intended class* $\mathcal{K}$*.*

$$\mathfrak{U}^{\mathrm{rel}}(\mathcal{R}, \mathcal{K}) = 1 \Longrightarrow \mathbf{S}_4(\mathcal{R}).$$

Equivalently, no additional local hardening layer is required beyond $\mathbf{S}_3(\mathcal{R})$*; what remains is the external strength theorem for the chosen reference core and the proof that its admissible coding discipline is faithfully transported by the already hardened simulation route.*

Corollary H9.47 (What the first upgrade criterion above $\mathbf{S}_3(\mathcal{R})$ already buys)**.** *The gap between the present line and conditional universality is now sharply isolated. To pass from the current reference-core closure level to* $\mathbf{S}_4(\mathcal{R})$*, later revisions no longer need to reopen the primitive computational hardening ladder. They need only choose a reference core* $\mathcal{R}$ *strong enough for the intended target class* $\mathcal{K}$*, certify the corresponding universality theorem for* $\mathcal{R}$*, and verify that the admissible encoding/decoding discipline survives transport through the already established reference-core closure surface.*

19W. First admissible reference-core selection principle above $\mathbf{S}_3(\mathcal{R})$

Once the upgrade from $\mathbf{S}_3(\mathcal{R})$ to $\mathbf{S}_4(\mathcal{R})$ has been isolated, the next question is no longer whether one can choose some strong reference core at all, but which admissible reference core should be chosen first. A structure-first route should avoid importing unnecessary machine-theoretic strength too early; it should instead select the weakest admissible reference core already sufficient for the intended target claim.

Definition H9.48 (First conservative reference-core selection certificate above $\mathbf{S}_3(\mathcal{R})$)**.** *Fix a target class* $\mathcal{K}$ *and a chosen comparison family* $\mathfrak{F}_{\mathrm{ref}}$ *of admissible sequential reference cores together with a strength preorder* $\preceq_{\mathrm{ref}}$*. Write*

$$\mathfrak{S}^{\mathrm{ref}}(\mathcal{R}, \mathcal{K}) = 1$$

if the following hold:

- $\mathcal{R} \in \mathfrak{F}_{\mathrm{ref}}$ *is admissible and satisfies* $\mathfrak{Z}^{\mathrm{ref}}(\mathcal{R}) = 1$*;*
- $\mathcal{R}$ *is strong enough to support the intended target claim for* $\mathcal{K}$ *under an admissible encoding/decoding discipline;*
- *for every* $\mathcal{R}' \in \mathfrak{F}_{\mathrm{ref}}$ *with* $\mathcal{R}' \prec_{\mathrm{ref}} \mathcal{R}$*, the same target claim for* $\mathcal{K}$ *is not yet certified under the same admissible coding discipline.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Thus $\mathfrak{S}^{\mathrm{ref}}(\mathcal{R},\mathcal{K}) = 1$ *means that* $\mathcal{R}$ *is not merely an admissible reference core, but the first conservatively sufficient one inside the chosen comparison family for the intended upgrade problem.*

Proposition H9.49 (First conservative selection reduction for the universality-upgrade route)**.** *Assume* $\mathfrak{U}^{\mathrm{rel}}(\mathcal{R},\mathcal{K}) = 1$ *and* $\mathfrak{S}^{\mathrm{ref}}(\mathcal{R},\mathcal{K}) = 1$*. Then the passage from* $\mathbf{S}_3(\mathcal{R})$ *to* $\mathbf{S}_4(\mathcal{R})$ *may be formulated with no stronger external reference-core assumption than the one already needed for the chosen target class* $\mathcal{K}$ *inside the comparison family* $\mathfrak{F}_{\mathrm{ref}}$*. In particular, later revisions do not need to import a stronger machine model than necessary in order to state the first conditional universality claim above the present reference-core closure level.*

Corollary H9.50 (What the first conservative reference-core selection principle already buys)**.** *The relative universality route is now disciplined in two distinct ways: first by the already established reference-core closure surface, and second by a conservative selection rule on the external endpoint. Later upgrades above* $\mathbf{S}_3(\mathcal{R})$ *can therefore proceed by choosing the weakest admissible reference core sufficient for the intended claim, rather than by jumping prematurely to an unnecessarily strong machine-theoretic endpoint.*

19X. First admissible tape-based reference-core family above $\mathbf{S}_3(\mathcal{R})$

The conservative selection principle above becomes materially useful only once one fixes a first concrete comparison family of admissible reference cores. The present line already carries explicit tape, control, branch, and loop/return roles, so the most conservative first family should preserve exactly that architecture instead of importing richer machine features prematurely.

Definition H9.51 (First admissible tape-based reference-core family)**.** *Let* $\mathfrak{F}_{\mathrm{tape}}$ *denote the class of admissible sequential reference cores* $\mathcal{R}$ *satisfying all of the following:*

- $\mathcal{R}$ *has finite control state, finite local symbol alphabet, and a one-track sequential update discipline;*
- *every local step of* $\mathcal{R}$ *is determined by the currently scanned symbol together with the current control state and returns one next control state, one write/update instruction, and one move instruction in* $\{-1, 0, +1\}$*;*
- *branch resolution is explicit and local, in the sense that each step chooses one of finitely many admissible successor clauses by a finite Boolean/projector-type test on the current scanned data;*
- *loop/return structure is explicit and local, in the sense that every admissible return event is encoded by a finite control-side return symbol together with a deterministic reentry instruction on the same one-track support;*
- *the tape support itself is one-dimensional and sequential, so that the already globalized spinorial optimal line may serve as the direct comparison support without introducing any extra addressing geometry.*

Write

$$\mathfrak{A}^{\mathrm{tape}}(\mathcal{R}) = 1$$

if $\mathcal{R} \in \mathfrak{F}_{\mathrm{tape}}$*.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proposition H9.52 (First conservative tape-family reduction for the upgrade route)**.** *Assume the conditional reference-core closure surface* $\mathfrak{Z}^{\mathrm{ref}}(\mathcal{R}) = 1$ *for some chosen admissible sequential reference core* $\mathcal{R}$*, together with the conservative selection discipline above. Suppose moreover that the intended universality-upgrade problem concerns only target classes whose admissible coding discipline is already one-track, sequential, and local in the same sense as the hardened tape/control/branch/loop architecture of the present line. Then the first conservative search for a reference core may be restricted to the admissible tape-based family* $\mathfrak{F}_{\mathrm{tape}}$ *without loss for that upgrade problem.*

Equivalently, whenever the intended target class does not yet require richer addressing, parallel branching geometry, or nonlocal memory moves, the weakest natural external comparison family is the tape-based family already matched to the present hardened architecture.

Corollary H9.53 (What the first admissible tape-based reference-core family already buys)**.** *The external universality-upgrade route is now no longer open-ended. Later revisions can first search for a minimally sufficient* tape-based *reference core before importing any richer machine model. This keeps the passage above* $\mathbf{S}_3(\mathcal{R})$ *aligned with the already hardened internal architecture and reduces the chance of overstating the needed external strength.*

19Y. First internal-to-tape transport certificate above $\mathbf{S}_3(\mathcal{R})$

Choosing a first admissible tape-based comparison family becomes materially useful only once one can express the existing reference-core closure surface in a way that is transport-compatible with that family. The next conservative step is therefore not a stronger universality claim, but a certificate saying that the already hardened internal read/write/control/branch/return package can be read inside a chosen tape-based reference core without changing the local computational discipline.

Definition H9.54 (First internal-to-tape transport certificate above $\mathbf{S}_3(\mathcal{R})$)**.** *Let* $\mathcal{R} \in \mathfrak{F}_{\mathrm{tape}}$ *satisfy* $\mathfrak{Z}^{\mathrm{ref}}(\mathcal{R}) = 1$*. Write*

$$\mathfrak{T}^{\mathrm{tape}}(\mathcal{R}) = 1$$

if the following hold on every admissible later-depth window used by the present compressed computational surface:

- *the globalized spinorial optimal line induces a one-dimensional sequential support compatible with the tape support of* $\mathcal{R}$*;*
- *the already hardened local control state of the present line is encoded into the finite control state of* $\mathcal{R}$ *without introducing extra addressing geometry;*
- *the already hardened projector/Boolean branch layer is transported into the branch clauses of* $\mathcal{R}$ *by a local admissible encoding/decoding discipline;*
- *the already hardened routed HC-selected return/closure step is transported into the local return/reentry discipline of* $\mathcal{R}$ *without changing the sequential one-track support;*
- *one local step of the present compressed computational surface is transported to one admissible local step of* $\mathcal{R}$ *up to the chosen local coding discipline.*

Thus $\mathfrak{T}^{\mathrm{tape}}(\mathcal{R}) = 1$ *means that the internal reference-core closure already lives on a support that can be read directly inside the first admissible tape-based comparison family.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proposition H9.55 (First transport reduction from reference-core closure to the tape-based family)**.** *Assume* $\mathfrak{Z}^{\mathrm{ref}}(\mathcal{R}) = 1$, $\mathfrak{A}^{\mathrm{tape}}(\mathcal{R}) = 1$, *and* $\mathfrak{T}^{\mathrm{tape}}(\mathcal{R}) = 1$. *Then the external upgrade route above* $\mathbf{S}_3(\mathcal{R})$ *may be reformulated entirely inside the admissible tape-based comparison family without reopening the already hardened local obligations for control, branch, loop/return, or tape support. In particular, the passage to the first conditional universality-upgrade problem may be stated as an external strength question about the chosen tape-based reference core itself, rather than as a renewed local hardening problem for the present line.*

Corollary H9.56 (What the first internal-to-tape transport certificate already buys)**.** *The tape-based upgrade route is now disciplined in three layers: by the conditional reference-core closure surface, by the conservative tape-family selection principle, and by an explicit transport certificate from the present compressed computational surface into that family. Later work above* $\mathbf{S}_3(\mathcal{R})$ *can therefore focus directly on the external strength of the chosen tape-based reference core, instead of rechecking the already stabilized local computational roles.*

19Z. First tape-family relative universality witness above $\mathbf{S}_3(\mathcal{R})$

Once the admissible tape-based family has been selected and the internal reference-core closure has been transported into it, the remaining external question is no longer whether the present line can be read inside that family, but whether a chosen member of that family is already strong enough for the intended target class. The next conservative step is therefore to isolate the first explicit witness certifying that a transported tape-based reference core upgrades the present line from conditional reference-core closure to conditional universality relative to that core.

Definition H9.57 (First tape-family relative universality witness above $\mathbf{S}_3(\mathcal{R})$)**.** *Let* $\mathcal{R} \in \mathfrak{F}_{\mathrm{tape}}$ *satisfy*

$$\mathfrak{Z}^{\mathrm{ref}}(\mathcal{R}) = 1, \qquad \mathfrak{A}^{\mathrm{tape}}(\mathcal{R}) = 1, \qquad \mathfrak{T}^{\mathrm{tape}}(\mathcal{R}) = 1.$$

For a target class $\mathcal{K}$, *write*

$$\mathfrak{U}^{\mathrm{tape}}(\mathcal{R}, \mathcal{K}) = 1$$

if the following hold:

- *the chosen tape-based reference core* $\mathcal{R}$ *is already known to simulate every machine in the intended target class* $\mathcal{K}$ *under a one-track sequential local coding discipline compatible with* $\mathfrak{F}_{\mathrm{tape}}$*;*
- *the admissible encoding/decoding discipline used in that simulation is stable under the internal-to-tape transport certificate* $\mathfrak{T}^{\mathrm{tape}}(\mathcal{R}) = 1$*;*
- *no extra addressing geometry, nonlocal branching discipline, or richer memory move is required beyond the tape-based family already selected.*

Thus $\mathfrak{U}^{\mathrm{tape}}(\mathcal{R}, \mathcal{K}) = 1$ *means that the chosen transported tape-based reference core is externally strong enough to witness the first conditional universality claim relevant to* $\mathcal{K}$.

Proposition H9.58 (First tape-family upgrade from $\mathbf{S}_3(\mathcal{R})$ to $\mathbf{S}_4(\mathcal{R})$)**.** *Assume* $\mathfrak{Z}^{\mathrm{ref}}(\mathcal{R}) = 1$, $\mathfrak{A}^{\mathrm{tape}}(\mathcal{R}) = 1$, $\mathfrak{T}^{\mathrm{tape}}(\mathcal{R}) = 1$, *and* $\mathfrak{U}^{\mathrm{tape}}(\mathcal{R}, \mathcal{K}) = 1$. *Then the present compressed computational surface upgrades from the reference-core closure level* $\mathbf{S}_3(\mathcal{R})$ *to the relative universality level* $\mathbf{S}_4(\mathcal{R})$ *for the intended target class* $\mathcal{K}$.

Equivalently, once the internal closure surface has been transported into a sufficiently strong admissible tape-based reference core, the remaining strength step above $\mathbf{S}_3(\mathcal{R})$ *is discharged entirely by the external universality witness for that transported tape-based core.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Corollary H9.59 (What the first tape-family relative universality witness already buys)**.** *The universality-upgrade route above the present compressed computational surface is now disciplined in four layers: by the conditional reference-core closure surface, by the conservative tape-family selection rule, by the internal-to-tape transport certificate, and by an explicit external universality witness for a chosen transported tape-based reference core. Later revisions can therefore state the first conditional universality claim without reopening the already discharged primitive machine-theoretic hardening ladder.*

19AA. First strict universality-obligation package above $\mathbf{S}_4(\mathcal{R})$

Once the present line has reached the relative universality level $\mathbf{S}_4(\mathcal{R})$ for a chosen transported tape-based reference core and target class, the remaining question is no longer whether the primitive computational roles are available, nor whether a conservative tape-based reference family can witness relative universality. What remains is the stricter global question whether the transported tape-based route supports the genuinely unbounded extension needed for strict Turing-style completeness. The next conservative step is therefore to isolate that last gap as a separate package instead of mixing it back into the already discharged local hardening ladder.

Definition H9.60 (First strict universality-obligation package above $\mathbf{S}_4(\mathcal{R})$)**.** *Assume that $\mathbf{S}_4(\mathcal{R})$ has been reached for an intended target class $\mathcal{K}$. Write*

$$\mathbf{V} = (\mathbf{V}_\infty, \mathbf{V}_{\mathrm{ext}}, \mathbf{V}_{\mathrm{dec}})$$

for the first strict universality-obligation package above $\mathbf{S}_4(\mathcal{R})$, where:

- $\mathbf{V}_\infty$ *is the obligation that the chosen transported tape-based reference route admits arbitrarily long admissible extensions rather than only bounded local windows;*
- $\mathbf{V}_{\mathrm{ext}}$ *is the obligation that the already established tape-globalization and concatenation discipline extends coherently along such arbitrarily long admissible chains, without hidden global collapse or cutoff;*
- $\mathbf{V}_{\mathrm{dec}}$ *is the obligation that the admissible encoding/decoding discipline witnessing $\mathbf{S}_4(\mathcal{R})$ remains stable under those unbounded extensions and does not depend on a fixed finite window budget.*

Thus $\mathbf{V}$ isolates the first strict machine-theoretic gap between relative universality inside a chosen tape-based reference-core route and full unbounded Turing-style completeness.

Proposition H9.61 (First isolation of the strict universality gap above $\mathbf{S}_4(\mathcal{R})$)**.** *Assume that $\mathbf{S}_4(\mathcal{R})$ has been reached through the conditional reference-core closure surface, the conservative tape-family selection rule, the internal-to-tape transport certificate, and an explicit tape-family relative universality witness. Then the only remaining machine-theoretic step toward the strict level $\mathbf{S}_5$ is discharged by the package $\mathbf{V}$ above.*

Equivalently, once the present line has been upgraded to $\mathbf{S}_4(\mathcal{R})$, no additional local hardening of control, branch, loop/return, tape support, or reference-core simulation is needed. What remains is exclusively the global unboundedness, extension, and decoding-stability problem recorded by $\mathbf{V}$.

Corollary H9.62 (What the first strict universality-obligation package already buys)**.** *The machine-theoretic hierarchy is now disciplined at both of its remaining upper gaps. The passage from $\mathbf{S}_3(\mathcal{R})$ to $\mathbf{S}_4(\mathcal{R})$ is controlled by the conservative tape-family upgrade route, while the*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

passage from $\mathbf{S}_4(\mathcal{R})$ *to* $\mathbf{S}_5$ *is now isolated as the strict universality-obligation package* $\mathbf{V}$*. Later revisions can therefore state very clearly whether they are addressing relative universality inside a chosen transported tape-based reference family or the separate problem of genuine unbounded Turing-style completeness.*

19AB. Hardening priority for the strict universality obligation above $\mathbf{S}_4(\mathcal{R})$

Once the strict universality gap above $\mathbf{S}_4(\mathcal{R})$ has been isolated as the package $\mathbf{V} = (\mathbf{V}_\infty, \mathbf{V}_{\mathrm{ext}}, \mathbf{V}_{\mathrm{dec}})$, the next conservative step is not to attack all three remaining obligations at once. As on the earlier Zuse-guided route, the cleaner strategy is to impose a first hardening order, so that later revisions can reduce the strict universality gap in the same staged way as the primitive computational obligations.

Definition H9.63 (First hardening priority order for the strict universality-obligation package above $\mathbf{S}_4(\mathcal{R})$)**.** *Assume that the present compressed computational surface has reached* $\mathbf{S}_4(\mathcal{R})$ *for an intended target class* $\mathcal{K}$*, and let*

$$\mathbf{V} = (\mathbf{V}_\infty, \mathbf{V}_{\mathrm{ext}}, \mathbf{V}_{\mathrm{dec}})$$

be the first strict universality-obligation package above $\mathbf{S}_4(\mathcal{R})$*. Define the first hardening priority order on* $\mathbf{V}$ *by*

$$\mathbf{H}_2 = (\mathbf{V}_{\mathrm{ext}}, \mathbf{V}_{\mathrm{dec}}, \mathbf{V}_\infty).$$

Thus the conservative strengthening strategy is:

- *first harden the global extension obligation* $\mathbf{V}_{\mathrm{ext}}$*,*
- *then harden decoding stability under extension* $\mathbf{V}_{\mathrm{dec}}$*,*
- *and only at the end attack genuine unboundedness* $\mathbf{V}_\infty$*.*

Proposition H9.64 (First priority reduction for the strict universality-obligation package above $\mathbf{S}_4(\mathcal{R})$)**.** *Assume that* $\mathbf{S}_4(\mathcal{R})$ *has been reached and that the hardening order* $\mathbf{H}_2$ *above is adopted. Then later revisions can reduce the strict universality gap monotonically: any successful discharge of the first two obligations* $\mathbf{V}_{\mathrm{ext}}$ *and* $\mathbf{V}_{\mathrm{dec}}$ *shrinks the remaining open strict gap to the single final problem* $\mathbf{V}_\infty$*.*

Equivalently, once the extension discipline and its decoding stability have been hardened, the only remaining obstruction to $\mathbf{S}_5$ *is the genuinely unbounded continuation problem rather than the local or semiglobal coherence of the transported tape-based route.*

Corollary H9.65 (What the first hardening priority order above $\mathbf{S}_4(\mathcal{R})$ already buys)**.** *The strict universality gap is now not only isolated but also staged. Later revisions can therefore work above* $\mathbf{S}_4(\mathcal{R})$ *without mixing extension, decoding, and unboundedness into a single undifferentiated final problem. In the conservative reading, the current next target is no longer* $\mathbf{S}_5$ *as a whole but specifically the first extension-hardening step* $\mathbf{V}_{\mathrm{ext}}$*.*

19AC. First local hardening of the extension obligation $\mathbf{V}_{\mathrm{ext}}$ above $\mathbf{S}_4(\mathcal{R})$

With the strict universality-obligation package now ordered by $\mathbf{H}_2$, the next conservative step is to attack the first remaining upper-gap component $\mathbf{V}_{\mathrm{ext}}$. The question is no longer whether the present line supports a finite transported tape-based computation relative to a chosen reference core, but whether those admissible windows extend coherently across longer chains without hidden collapse, cutoff, or incompatible seam behavior.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition H9.66 (First extension-hardening certificate for $\mathbf{V}_{\mathrm{ext}}$ above $\mathbf{S}_4(\mathcal{R})$)**.** *Assume* $\mathbf{S}_4(\mathcal{R})$ *for an intended target class* $\mathcal{K}$*. Write*

$$\mathfrak{E}^{\mathrm{glob}}(\mathcal{R}) = 1$$

if the following hold:

- *every admissible finite transported tape-window in the chosen family extends to a longer admissible tape-window by a compatible local continuation step;*
- *the already established concatenation/cut and patch-normal-form discipline remains coherent under those longer continuations;*
- *no hidden global seam defect, cutoff phenomenon, or incompatible local reindexing appears when adjoining admissible windows along the transported tape-line.*

Thus $\mathfrak{E}^{\mathrm{glob}}(\mathcal{R}) = 1$ *certifies the first coherent extension-hardening of the tape-based route above* $\mathbf{S}_4(\mathcal{R})$*.*

Proposition H9.67 (First reduction of the extension obligation $\mathbf{V}_{\mathrm{ext}}$ above $\mathbf{S}_4(\mathcal{R})$)**.** *Assume* $\mathbf{S}_4(\mathcal{R})$ *and* $\mathfrak{E}^{\mathrm{glob}}(\mathcal{R}) = 1$*. Then the first strict universality-obligation package reduces from*

$$\mathbf{V} = (\mathbf{V}_{\infty}, \mathbf{V}_{\mathrm{ext}}, \mathbf{V}_{\mathrm{dec}})$$

to

$$\mathbf{V}' = (\mathbf{V}_{\mathrm{dec}}, \mathbf{V}_{\infty}).$$

Equivalently, once the transported tape-based route is globally extensible in the admissible-window sense, the strict universality gap above $\mathbf{S}_4(\mathcal{R})$ *no longer contains a separate extension obstruction; what remains are the decoding-stability problem under such extension and the final genuinely unbounded continuation problem.*

Corollary H9.68 (What the first local hardening of $\mathbf{V}_{\mathrm{ext}}$ above $\mathbf{S}_4(\mathcal{R})$ already buys)**.** *The upper strict gap above relative universality is now cleaner than before. If the extension-hardening certificate holds, later revisions do not need to reopen the global seam/coherence problem of transported admissible tape-windows. The next active upper-gap target becomes decoding stability under extension, with genuine unboundedness kept as the final separate endpoint problem.*

19AD. First local hardening of the decoding-stability obligation $\mathbf{V}_{\mathrm{dec}}$ above $\mathbf{S}_4(\mathcal{R})$

Once the extension obligation has been isolated and locally hardened, the next conservative strict-universality target is no longer the seam/coherence of admissible transported tape-windows themselves, but the stability of the admissible encoding/decoding discipline under those longer extensions. The question is now whether the same transported tape-based route that already witnesses relative universality continues to admit a coherent decoding discipline when admissible windows are enlarged, concatenated, and reread along longer chains.

Definition H9.69 (First decoding-stability certificate for $\mathbf{V}_{\mathrm{dec}}$ above $\mathbf{S}_4(\mathcal{R})$)**.** *Assume* $\mathbf{S}_4(\mathcal{R})$ *for an intended target class* $\mathcal{K}$*, and assume moreover that the global extension certificate* $\mathfrak{E}^{\mathrm{glob}}(\mathcal{R}) = 1$ *has already been established. Write*

$$\mathfrak{D}^{\mathrm{ext}}(\mathcal{R}) = 1$$

if the following hold:

© 2026 Marc Brendecke. Preprint — not peer reviewed.

- *the admissible decoding discipline used to witness* $\mathbf{S}_4(\mathcal{R})$ *on finite transported tape-windows extends coherently to every admissible longer transported tape-window produced by the extension certificate;*
- *restriction from a longer admissible transported tape-window to any admissible subwindow commutes with the chosen decoding/readout rule;*
- *admissible concatenation of transported tape-windows does not introduce a new decoding ambiguity, hidden branch aliasing, or context-dependent reinterpretation of already decoded local machine states.*

Thus $\mathfrak{D}^{\mathrm{ext}}(\mathcal{R}) = 1$ *certifies the first stable decoding discipline under admissible transported tape-extension above* $\mathbf{S}_4(\mathcal{R})$.

Proposition H9.70 (First reduction of the decoding-stability obligation $\mathbf{V}_{\mathrm{dec}}$ above $\mathbf{S}_4(\mathcal{R})$)**.** *Assume* $\mathbf{S}_4(\mathcal{R})$, $\mathfrak{E}^{\mathrm{glob}}(\mathcal{R}) = 1$, *and* $\mathfrak{D}^{\mathrm{ext}}(\mathcal{R}) = 1$. *Then the strict universality-obligation package reduces from*

$$\mathbf{V} = (\mathbf{V}_\infty, \mathbf{V}_{\mathrm{ext}}, \mathbf{V}_{\mathrm{dec}})$$

to

$$\mathbf{V}'' = (\mathbf{V}_\infty).$$

Equivalently, once the transported tape-based route is globally extensible and its admissible decoding discipline remains stable under those extensions, the strict universality gap above $\mathbf{S}_4(\mathcal{R})$ *no longer contains a separate decoding obstruction; what remains is only the final genuinely unbounded continuation problem.*

Corollary H9.71 (What the first local hardening of $\mathbf{V}_{\mathrm{dec}}$ above $\mathbf{S}_4(\mathcal{R})$ already buys)**.** *The upper strict gap has now been reduced as far as it can be reduced without addressing genuine unboundedness itself. If the extension-hardening and decoding-stability certificates both hold, later revisions no longer need to reopen finite-window extension or decoding coherence. The only remaining active upper-gap target is then* $\mathbf{V}_\infty$, *namely true unbounded continuation.*

19AE. Local hardening of genuine unboundedness $\mathbf{V}_\infty$ above $\mathbf{S}_4(\mathcal{R})$

Once global extension and decoding stability under extension have both been isolated and locally hardened, the only remaining strict-universality target is no longer any finite-window transport or rereading problem. What remains is the genuinely unbounded continuation problem itself: whether the transported tape-based route above $\mathbf{S}_4(\mathcal{R})$ admits arbitrarily extendable admissible continuations without forced collapse, periodic truncation, or hidden resource cutoff.

Definition H9.72 (First genuine-unboundedness certificate for $\mathbf{V}_\infty$ above $\mathbf{S}_4(\mathcal{R})$)**.** *Assume* $\mathbf{S}_4(\mathcal{R})$, $\mathfrak{E}^{\mathrm{glob}}(\mathcal{R}) = 1$, *and* $\mathfrak{D}^{\mathrm{ext}}(\mathcal{R}) = 1$. *Write*

$$\mathfrak{U}^{\infty}(\mathcal{R}) = 1$$

if the following hold:

- *for every admissible transported tape-window in the chosen tape-based family there exists a strictly longer admissible transported tape-window extending it;*
- *admissible extension can be iterated indefinitely without forcing a terminal collapse, bounded cycling artifact, or hidden global cutoff that would prevent arbitrarily long transported computations;*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

- *the already established extension and decoding certificates remain valid along every finite stage of such iterated admissible continuation.*

Thus $\mathfrak{U}^{\infty}(\mathcal{R}) = 1$ *certifies the first genuinely unbounded admissible continuation regime above* $\mathbf{S}_4(\mathcal{R})$.

Proposition H9.73 (First reduction of the genuine unboundedness obligation $\mathbf{V}_{\infty}$ above $\mathbf{S}_4(\mathcal{R})$)**.** *Assume* $\mathbf{S}_4(\mathcal{R})$, $\mathfrak{E}^{\mathrm{glob}}(\mathcal{R}) = 1$, $\mathfrak{D}^{\mathrm{ext}}(\mathcal{R}) = 1$, *and* $\mathfrak{U}^{\infty}(\mathcal{R}) = 1$. *Then the strict universality-obligation package reduces from*

$$\mathbf{V} = (\mathbf{V}_{\infty}, \mathbf{V}_{\mathrm{ext}}, \mathbf{V}_{\mathrm{dec}})$$

to the empty list. Equivalently, once the transported tape-based route is globally extensible, decoding-stable under extension, and genuinely unbounded in the admissible continuation sense, the strict universality gap above $\mathbf{S}_4(\mathcal{R})$ *contains no further separate obstruction. In that case the compressed computational surface reaches the top level* $\mathbf{S}_5$, *namely strict unbounded Turing-style completeness relative to the certified route.*

Corollary H9.74 (What the first local hardening of $\mathbf{V}_{\infty}$ above $\mathbf{S}_4(\mathcal{R})$ already buys)**.** *The upper gap is now completely structured. If the global extension, decoding-stability, and genuine-unboundedness certificates all hold, later revisions do not need to reopen any remaining strict-universality obstruction above* $\mathbf{S}_4(\mathcal{R})$. *The only remaining caution is interpretive: one must still keep explicit which route, tape-family, and reference-core class are being certified when speaking of strict completeness.*

19AF. Full machine-theoretic closure surface on the compressed computational line

After the conditional reference-core closure surface, the relative universality-upgrade witness, and the strict upper certificate package have all been isolated, the machine-theoretic reading should no longer appear as a long ladder of locally discharged obligations. It can now be bundled into a single closure surface that records exactly which level of the hierarchy has been reached under which certificate discipline.

Definition H9.75 (Full machine-theoretic closure surface on the compressed computational line)**.** *Fix a chosen admissible sequential reference core* $\mathcal{R}$, *a chosen admissible tape-based reference family, and the associated conservative transport discipline. Write*

$$\mathfrak{M}^{\mathrm{full}}(\mathcal{R}) = 1$$

if the following package holds simultaneously:

- *the compressed computational interpretation surface reaches the conditional reference-core closure level* $\mathbf{S}_3(\mathcal{R})$, *i.e.* $\mathfrak{Z}^{\mathrm{ref}}(\mathcal{R}) = 1$;
- *the tape-family upgrade route is certified, i.e. the conservative tape-family selection rule, the internal-to-tape transport certificate, and the tape-family relative universality witness all hold, so that the line reaches* $\mathbf{S}_4(\mathcal{R})$;
- *the strict upper certificate package is certified, i.e.*

$$\mathfrak{E}^{\mathrm{glob}}(\mathcal{R}) = 1, \qquad \mathfrak{D}^{\mathrm{ext}}(\mathcal{R}) = 1, \qquad \mathfrak{U}^{\infty}(\mathcal{R}) = 1,$$

  so that the remaining strict gap above $\mathbf{S}_4(\mathcal{R})$ *is empty.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Thus $\mathfrak{M}^{\mathrm{full}}(\mathcal{R}) = 1$ denotes the first fully bundled machine-theoretic closure surface attached to the present compressed computational line relative to the chosen certified route.

Theorem H9.76 (First bundled machine-theoretic closure theorem on the compressed computational line)**.** *Assume $\mathfrak{M}^{\mathrm{full}}(\mathcal{R}) = 1$. Then the compressed computational line reaches the top machine-theoretic level $\mathbf{S}_5$ along the chosen certified route. Equivalently, all intermediate gaps*

$$\mathbf{S}_1 \longrightarrow \mathbf{S}_2 \longrightarrow \mathbf{S}_3(\mathcal{R}) \longrightarrow \mathbf{S}_4(\mathcal{R}) \longrightarrow \mathbf{S}_5$$

are discharged under an explicitly named certificate package, so that no separate machine-theoretic obstruction remains inside the adopted route.

Corollary H9.77 (What the full machine-theoretic closure surface already buys)**.** *Later revisions no longer need to cite the whole local hardening ladder when the only point is that the present computational reading has been conditionally closed all the way up the hierarchy along a chosen certified route. They may instead cite $\mathfrak{M}^{\mathrm{full}}(\mathcal{R}) = 1$, while still keeping explicit which reference core, which tape-family route, and which upper certificates are meant. This compresses the machine-theoretic reading to a single surface without pretending that the route has become unconditional or model-independent.*

19AG. First machine-theoretic status decoder below and above the full closure surface

After the full machine-theoretic closure surface has been bundled, the text should not force later revisions to reopen the entire hierarchy merely to say which machine-theoretic level is currently justified under a given certificate package. A small status decoder is therefore useful: it reads off the strongest currently justified strength level from the available closure surfaces and upgrade certificates without changing the underlying mathematics.

Definition H9.78 (First machine-theoretic status decoder on the compressed computational line)**.** *Fix a chosen admissible sequential reference core $\mathcal{R}$. Define the machine-theoretic status decoder*

$$\mathfrak{S}^{\mathrm{mach}}(\mathcal{R}) \in \{\mathbf{S}_1, \mathbf{S}_2, \mathbf{S}_3(\mathcal{R}), \mathbf{S}_4(\mathcal{R}), \mathbf{S}_5\}$$

as the strongest level among the hierarchy

$$\mathbf{S}_1 \longrightarrow \mathbf{S}_2 \longrightarrow \mathbf{S}_3(\mathcal{R}) \longrightarrow \mathbf{S}_4(\mathcal{R}) \longrightarrow \mathbf{S}_5$$

whose required certificate package has been verified. Concretely:

- $\mathfrak{S}^{\mathrm{mach}}(\mathcal{R}) = \mathbf{S}_5$ *if* $\mathfrak{M}^{\mathrm{full}}(\mathcal{R}) = 1$*;*
- $\mathfrak{S}^{\mathrm{mach}}(\mathcal{R}) = \mathbf{S}_4(\mathcal{R})$ *if the tape-family upgrade route is certified but the strict upper certificate package is not fully certified;*
- $\mathfrak{S}^{\mathrm{mach}}(\mathcal{R}) = \mathbf{S}_3(\mathcal{R})$ *if the conditional reference-core closure surface is certified but the tape-family upgrade route is not fully certified;*
- $\mathfrak{S}^{\mathrm{mach}}(\mathcal{R}) = \mathbf{S}_2$ *if the conditional primitive Zuse-style core is certified but reference-core closure is not;*
- $\mathfrak{S}^{\mathrm{mach}}(\mathcal{R}) = \mathbf{S}_1$ *otherwise.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proposition H9.79 (Monotone exactness of the first machine-theoretic status decoder)**.** *For every chosen admissible sequential reference core $\mathcal{R}$, the decoder $\mathfrak{S}^{\mathrm{mach}}(\mathcal{R})$ is monotone with respect to certificate accumulation and exact with respect to the present hierarchy: it never outputs a higher level than has actually been certified, and whenever a full certificate package for some level in the hierarchy has been established, the decoder outputs at least that level. In particular,*

$$\mathfrak{M}^{\mathrm{full}}(\mathcal{R}) = 1 \quad \Longrightarrow \quad \mathfrak{S}^{\mathrm{mach}}(\mathcal{R}) = \mathbf{S}_5.$$

Likewise, if only the reference-core closure surface is available, the decoder halts at $\mathbf{S}_3(\mathcal{R})$, and if the tape-family upgrade route has been certified but the strict upper package has not, it halts at $\mathbf{S}_4(\mathcal{R})$.

Corollary H9.80 (What the first machine-theoretic status decoder already buys)**.** *Later revisions can now state the strongest presently justified machine-theoretic claim without reopening the whole ladder of lower certificates. They may cite the single decoded status $\mathfrak{S}^{\mathrm{mach}}(\mathcal{R})$ and then only unfold the route-specific certificate package when a stronger or more unconditional claim is intended. This keeps the computational reading disciplined: the text can speak precisely about the current level already reached, while preventing accidental overstatement about higher levels that are not yet certified.*

19AH. Conditional/unconditional status surface on the compressed computational line

After the machine-theoretic status decoder has been introduced, the current release line should not remain described only in abstract terms. A small present-line status surface is therefore useful: it records, in one place, the strongest route-dependent status already certified on the present 4.7.0 release line and separates it from the weaker certificate-free status that can be asserted without invoking any chosen reference-core route.

Definition H9.81 (First present-line conditional/unconditional status surface on the compressed computational line)**.** *Fix a chosen admissible sequential reference core $\mathcal{R}$. Define the present-line machine-theoretic status pair*

$$\mathfrak{P}^{\mathrm{mach}}(\mathcal{R}) = \big(\mathfrak{S}^{\mathrm{abs}}, \mathfrak{S}^{\mathrm{cond}}(\mathcal{R})\big)$$

by the following rule:

- *$\mathfrak{S}^{\mathrm{abs}}$ is the strongest machine-theoretic level justified without invoking any chosen reference core, tape-family route, or upper certificate package;*
- *$\mathfrak{S}^{\mathrm{cond}}(\mathcal{R})$ is the strongest machine-theoretic level justified under the explicit chosen route $\mathcal{R}$ together with the corresponding lower and upper certificate package.*

The pair $\mathfrak{P}^{\mathrm{mach}}(\mathcal{R})$ will be called the present-line conditional/unconditional status surface.

Proposition H9.82 (First explicit present-line status readout on the compressed computational line)**.** *On the present 4.7.0 release line, and for every chosen admissible sequential reference core $\mathcal{R}$ along which the bundled closure surface $\mathfrak{M}^{\mathrm{full}}(\mathcal{R}) = 1$ has been certified, the present-line status surface reads*

$$\mathfrak{P}^{\mathrm{mach}}(\mathcal{R}) = \big(\mathbf{S}_1, \mathbf{S}_5\big).$$

Equivalently: unconditionally, the present line should still only be advertised at the baseline disciplined computational-reading level $\mathbf{S}_1$; conditionally, along the explicitly certified route $\mathcal{R}$, the present line reaches the top certified level $\mathbf{S}_5$. In particular, the present text should avoid collapsing these two readings into a single unconditional global claim.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Corollary H9.83 (What the first present-line conditional/unconditional status surface already buys)**.** *Later revisions can now cite the present machine-theoretic state of the Companion in one disciplined sentence: the line is unconditionally at* $\mathbf{S}_1$*, while conditionally along a chosen certified route it reaches* $\mathbf{S}_5$*. This gives a compact status formula for future hardening passes while preventing accidental overstatement about certificate-dependent closures as if they were already unconditional.*

Remark H9.84 (One-paragraph synopsis of the machine-theoretic line)**.** *The computational line of the present framework can be read in three nested stages. First, the local operational layer yields a conditional primitive machine core, in which tape-like support, control, branching, and loop/return behavior are structurally separated and locally executable. Second, this primitive core can be hardened into a reference-core closure relative to a chosen sequential reference kernel* $\mathcal{R}$*, so that the line is no longer only machine-like in analogy, but carries a disciplined reference computation surface. Third, along a certified tape-based route above this reference-core closure, the framework admits a conditional upgrade toward stronger machine-theoretic levels, up to a Turing-style completeness claim only under explicit higher certificates; accordingly, the present line supports an unconditional computational base reading, while stronger universality claims remain route-dependent and conditional.*

19AH′. First bidirectional observer-side machine-shadow split below the status surface

After the present-line conditional/unconditional status surface has been fixed, one more local question becomes unavoidable on the working continuation line: how should the currently retained public machine readout itself be decomposed at observer side? The next useful step is not yet a full new universality statement, but a disciplined split of the retained machine shadow into one symmetric total channel and one antisymmetric imbalance channel. This keeps the older no-second-carrier discipline visible while making the current machine readout auditable in a form that can be compared directly with later transport, spectral, or shell-level hardening.

Definition H9.85 (Bidirectional observer-side machine-shadow coordinates)**.** *Fix the retained overlap window* W_{ret} *and a tested lag* ℓ*. Let* $\eta^{(\ell)}_{\mathrm{drive}}$ *denote the fitted observer-side drive contribution and* $\eta^{(\ell)}_{\mathrm{carrier}}$ *the fitted carrier contribution on* W_{ret}*. Define the bidirectional observer-side coordinates*

$$\eta^{(\ell)}_{+} = \eta^{(\ell)}_{\mathrm{drive}} + \eta^{(\ell)}_{\mathrm{carrier}}, \qquad \eta^{(\ell)}_{-} = \eta^{(\ell)}_{\mathrm{drive}} - \eta^{(\ell)}_{\mathrm{carrier}}.$$

We call $\eta^{(\ell)}_{+}$ *the* total machine-shadow channel *and* $\eta^{(\ell)}_{-}$ *the* observer-side imbalance channel.

Proposition H9.86 (Current retained-overlap checkpoint at the public optimum)**.** *On the retained overlap window currently kept on the working line, the bidirectional observer-side coordinates reproduce the present public optimum at lag*

$$\ell_* = -14$$

with weighted defect approximately 0.6746 *over* 42 *overlap points.*

Remark H9.87 (Interpretation of the checkpoint)**.** *The point of this checkpoint is not yet a final dynamical theorem about the origin of the machine readout. It is the weaker but already useful fact that the presently best public lag can be read consistently through one bidirectional observer-side split rather than through two unrelated fitted channels. In that sense, the present step converts one numerical optimum into a structurally readable local machine-shadow record.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proposition H9.88 (The imbalance channel is substantial but not primitive)**.** *On the same retained overlap window at the public optimum $\ell_* = -14$, the observer-side imbalance channel satisfies*

$$\frac{\|\eta_-^{(\ell_*)}\|_1}{\|\eta_+^{(\ell_*)}\|_1} \approx 0.6827, \qquad \operatorname{corr}\big(\eta_+^{(\ell_*)}, \eta_-^{(\ell_*)}\big) \approx 0.9532.$$

Hence the imbalance channel is not negligible, but it remains tightly slaved to the same total machine-shadow profile.

Corollary H9.89 (First no-second-primitive-carrier reading from the machine split)**.** *The current bidirectional observer-side readout does not force the introduction of a second primitive carrier. What it supports is one dominant machine shadow together with a substantial but highly correlated imbalance readout. In particular, the current continuation line remains compatible with the older no-second-carrier discipline: the extra visible degree of freedom appears at observer side as a locked asymmetry channel, not as evidence for a new independent carrier species.*

Remark H9.90 (Why this matters for the next hardening step)**.** *The split creates a useful next leverage point. Future hardening no longer needs to ask only whether a retained carrier exists; it can ask whether a proposed theorem, transport step, or spectral proxy controls the symmetric total channel η_+, the locked imbalance channel η_-, or both at once. That distinction should make it easier to separate genuine carrier obligations from observer-shell correction obligations in the next working passes.*

19AH″. First channel-control dictionary induced by the machine-shadow split

Once the observer-side machine readout has been split into η_+ and η_-, the next useful step is a control dictionary telling later arguments what kind of burden they actually carry. This is still weaker than a new dynamical theorem. Its role is bookkeeping with mathematical consequences: it distinguishes propositions that genuinely control the retained machine shadow from propositions that merely reorganize the observer-side asymmetry readout around that shadow.

Definition H9.91 (η_+-control, η_--control, and joint channel control)**.** *Fix the retained overlap window W_{ret} at the current public optimum $\ell_* = -14$. We say that a theorem, estimate, transport rule, or spectral proxy on the present working line has*

1. total-channel control *if it bounds, reconstructs, or rigidifies $\eta_+^{(\ell_*)}$ up to the standing admissible equivalences without requiring an independent primitive contribution beyond the present drive/carrier split;*

2. imbalance-channel control *if, once the total channel is fixed in that admissible sense, it bounds, reconstructs, or rigidifies $\eta_-^{(\ell_*)}$ as an observer-side asymmetry readout around the same total machine shadow;*

3. joint channel control *if it controls both channels at once and in addition preserves or explains a structural lock between them, such as the present norm ratio or the present high correlation.*

Proposition H9.92 (First machine-side reading discipline induced by the dictionary)**.** *On the current working line, arguments aimed at the retained carrier or at the existence of one*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

common machine shadow should be read as η_+-control problems, whereas observer-shell, asymmetry, phase-correction, or frame-flicker refinements should be read as η_--control problems unless they also alter the total channel itself. Mixed claims that try to upgrade both at once should be treated as joint-channel problems and therefore carry the heaviest burden of proof.

Corollary H9.93 (First reformulation of the no-second-carrier discipline at machine level)**.** *A future refinement is compatible with the present no-second-carrier discipline whenever its extra visible freedom can be organized as η_--control over a fixed or jointly controlled η_+-channel, rather than by introducing a second primitive contribution already at the total-channel level. In particular, the present checkpoint pushes the burden of any would-be second-carrier claim upward: such a claim would now have to break the current reading at the level of η_+, not merely by producing additional observer-side imbalance structure.*

Remark H9.94 (Why the dictionary is a real compression tool)**.** *The dictionary turns one vague question — "does this next theorem improve the machine line?" — into a sharper trichotomy. A proposed step may improve the total machine shadow, the locked imbalance readout, or the coupling data between them. This should make later proof compression cleaner because many older optical and shell-side corrections can now be classified immediately as η_--side obligations, while the rarer genuinely carrier-bearing steps can be isolated as η_+-side obligations.*

19AH‴. First backward channel reading of the older optical and shell-side baselines

The dictionary becomes substantially more useful once it reorganizes not only future arguments but also the already stabilized comparison/shell archive. The present working line therefore records a first backward reading of the older optical-comparison, shortlist, bombe/sieve, shell/menu/window, and flicker-style material through the split into η_+ and η_-.

Proposition H9.95 (First backward channel reading of the retained public baselines)**.** *In this inherited public-release baseline block, the older public baselines should be read according to the following burden split.*

1. *The retained-carrier comparison layer, public optimum selection, no-second-carrier discipline, and every comparison/back-reading/bombe/sieve step insofar as it isolates one common machine-bearing shadow are primarily η_+-control statements.*

2. *The shell/menu/window refinements, phase-correction re-readings, and frame-flicker-style observer corrections are primarily η_--control statements whenever they refine how that common shadow appears on the observer side without changing the retained optimum itself.*

3. *Any statement claiming at once that a shell-side correction preserves the retained optimum, explains the asymmetry pattern, and leaves the common machine shadow identifiable should be treated as a joint-channel statement.*

Corollary H9.96 (First re-reading of the optical approximation baseline)**.** *The earlier first retained-carrier optical approximation should presently be read as one η_+-bearing baseline together with a subordinate η_--side correction envelope. Accordingly, visible mismatch inside that optical approximation is not yet evidence against the retained-carrier discipline unless it forces an additional primitive contribution already at the total-channel level.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Remark H9.97 (Compression payoff of the backward reading)**.** *The backward reading turns a long mixed archive into a more ordered burden ledger. Carrier isolation, common-window retention, and no-second-carrier arguments can now be cited as total-channel obligations, while shell refinements, asymmetry diagnostics, and flicker-style observer corrections can be cited as imbalance-side obligations. Only the rarer steps that claim both retention and corrected observer-side readout need to reopen the full joint burden.*

19AH$^{(4)}$. First theorematic burden split for the remaining continuation knots

Once the older baselines have been re-read through η_+ and η_-, the next practical question is which unresolved continuation knots still carry genuine carrier burden and which ones are only observer-side refinement burdens. The present working line therefore records one first theorematic burden split: unresolved shell-, phase-, or flicker-side refinement is not automatically a retained-carrier obstruction.

Proposition H9.98 (First theorematic burden split for the remaining continuation knots)**.** *On the present continuation line, the remaining theorematic knots should be read according to the following burden discipline.*

1. *Any future statement whose conclusion asserts preservation of the retained public optimum, one-common-machine-shadow identification, or the no-second-primitive-carrier discipline must control the total channel η_+. Purely imbalance-side refinements on η_- can neither prove nor refute such a carrier-level statement by themselves.*

2. *The frame-flicker ansatz, shell/menu/window refinements, phase-correction re-readings, and analogous observer-side asymmetry refinements are to be treated as η_--side knots as long as they only change how the retained machine shadow is seen and do not force a new total-channel profile.*

3. *Any theorem claiming simultaneously that an observer-side correction explains the asymmetry pattern, improves the retained fit, and preserves one common carrier must be treated as a joint-channel knot: it has to control both η_+ and η_-.*

4. *Consequently, a visible residual defect that can be confined to η_- is not yet a retained-carrier failure of the present line. A genuine carrier-level obstruction appears only if the retained optimum, the one-shadow reading, or the no-second-carrier discipline is already forced to fail on the total-channel side.*

Corollary H9.99 (Default downgrade rule for unresolved observer-side mismatch)**.** *On the present continuation line, an unresolved optical, shell-side, or flicker-style mismatch should by default be downgraded to an η_--side obligation unless it demonstrably changes the retained optimum, destroys the one-shadow reading at the total-channel level, or forces an additional primitive contribution already in η_+. This keeps the current no-second-carrier discipline from being overburdened by observer-side correction tasks that have not yet crossed into carrier space.*

Remark H9.100 (Why this sharpened burden split matters)**.** *The practical gain is a much cleaner proof queue. Future hardening can now ask first whether an open point is really about the common retained shadow or only about its observer-side distortion. If it is only the latter, then it belongs to the asymmetry shell and should not delay carrier-level statements. Only the comparatively rare steps that touch both channels at once need the full coupled burden again.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

19AH$^{(5)}$. First formal demotion criterion from joint-channel burden to pure imbalance-side burden

The burden split of 19AH$^{(4)}$ becomes operational only once one can say when a final continuation step no longer deserves to be treated as a genuinely joint-channel knot. The next local tool is therefore a formal demotion criterion: a pending observer-side refinement may be demoted from joint-channel burden to a pure η_--side burden as soon as its unresolved remainder provably leaves the retained total profile untouched.

Definition H9.101 (Observer-side demotion certificate)**.** *Fix a pending continuation statement $\mathcal{C}$ whose unresolved part currently concerns shell, phase, flicker, menu, or analogous observer-side refinement. We say that $\mathcal{C}$ admits an* observer-side demotion certificate *if all three of the following conditions hold on the retained admissible window under discussion.*

1. ***Total-profile invariance.*** *The retained public optimum, the one-machine-shadow identification, and the no-second-primitive-carrier reading remain unchanged on the η_+-side under the unresolved remainder of $\mathcal{C}$.*

2. ***No forced new primitive contribution.*** *The unresolved remainder of $\mathcal{C}$ does not require an additional primitive carrier already at the total-channel level; equivalently, the outstanding defect can still be represented without altering the retained η_+-carrier architecture.*

3. ***Imbalance localization.*** *The visible mismatch produced by the unresolved remainder of $\mathcal{C}$ can be localized to the observer-side imbalance channel η_-, meaning that its effect is read as a change of asymmetry, shell, or observer-side correction profile rather than as a change of the retained common total shadow.*

Proposition H9.102 (First formal demotion criterion)**.** *If a pending continuation statement $\mathcal{C}$ admits an observer-side demotion certificate, then $\mathcal{C}$ may be demoted from a provisional joint-channel knot to a pure η_--side obligation on the present continuation line. In particular, the unresolved remainder of $\mathcal{C}$ may no longer be cited as an open carrier-level obstruction against the retained public optimum, the one-shadow reading, or the no-second-primitive-carrier discipline.*

Proof. The three certificate clauses remove exactly the reasons why $\mathcal{C}$ would still need joint status. Clause (1) says that the retained total profile already survives unchanged at the η_+-level. Clause (2) rules out the only remaining carrier-level escape route, namely that the unresolved part secretly forces an additional primitive contribution already in the total channel. Clause (3) places the visible remainder entirely on the observer-side imbalance ledger. Once these three facts are available simultaneously, the outstanding burden of $\mathcal{C}$ no longer concerns whether the present line carries one retained common machine shadow, but only how the observer-side asymmetry shell should be refined. The knot is therefore genuinely η_--side and may be demoted accordingly. □

Corollary H9.103 (Safe demotion rule for future continuation work)**.** *Future continuation steps should only be kept on the joint-channel ledger if at least one of the three certificate clauses fails or is not yet known. Conversely, once the retained η_+-profile is stable, no extra primitive carrier is forced there, and the remaining defect is localized to η_-, the step should be queued as observer-side hardening rather than as unfinished carrier proof.*

Remark H9.104 (Why this tool is useful)**.** *This criterion turns the burden split into a reusable proof tool. It prevents later drafts from repeatedly reopening the full carrier question*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

whenever a shell-side mismatch or flicker-style correction is still under discussion. The proof queue can now be triaged by certificate: unresolved points without η_+-damage stay on the asymmetry ledger, while only genuine total-channel risk keeps a knot on the joint ledger.

19AH$^{(6)}$. First provisional demotion ledger for inherited observer-side continuation knots

Reader cue. Read the next two subsections as a queue-classification pair. Their purpose is to sort inherited shell/phase/flicker-style leftovers first onto the observer-side ledger by default and then to state the exact evidence that would force those same knots back into joint or total-channel burden.

The formal demotion criterion of 19AH$^{(5)}$ becomes most useful once it is applied back to the inherited queue. The point is not to declare old open points "solved" by fiat, but to state explicitly which of them should from now on be carried on the observer-side ledger unless and until they actually damage the retained total profile. The next local step is therefore a provisional demotion ledger for the main inherited observer-side knot families.

Definition H9.105 (Provisional demotion ledger for inherited observer-side knots)**.** *On the present continuation line, the following inherited knot families are placed on the* provisional observer-side demotion ledger*, subject to re-promotion only if later work shows a genuine failure of at least one certificate clause from 19AH$^{(5)}$:*

1. *shell/menu/window-refinement mismatches that alter the observer-side shell geometry without changing the retained total optimum profile,*
2. *phase-correction and frame-flicker residuals that modify how the retained profile is read from the observer side while leaving the common total shadow intact,*
3. *optical-approximation residuals whose visible discrepancy can still be localized to the asymmetry envelope rather than to the retained common carrier, and*
4. *analogous observer-side refinements inherited from the older archive whenever they do not force a second primitive carrier and do not move the retained public optimum on the total-channel side.*

Proposition H9.106 (First provisional demotion ledger theorem)**.** *The inherited knot families listed above belong, by default, on the observer-side demotion ledger rather than on the joint-channel obstruction ledger. More precisely, on the present compressed computational line they should be read as provisional η_--side obligations unless later analysis shows that one of them changes the retained η_+-profile, breaks the one-shadow reading at total-channel level, or forces an additional primitive carrier already in η_+.*

Proof. Each listed family is inherited precisely from the shell/phase/flicker/optical side of the archive, hence from the side that 19AH‴ and 19AH$^{(4)}$ already identified as the natural observer-side burden domain. The formal criterion of 19AH$^{(5)}$ then supplies the missing operational rule: as long as the retained total profile remains fixed, no new primitive carrier is forced in the total channel, and the visible mismatch stays localizable to the imbalance channel, the corresponding inherited knot is not entitled to remain on the joint ledger. The listed families therefore enter the queue under default demotion. Re-promotion is reserved only for the exceptional case that later work actually detects η_+-damage or a genuine carrier-level escape route. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Corollary H9.107 (How to read the old queue from now on)**.** *From this point onward, older shell/menu/window, phase-correction, flicker-style, and observer-side optical residuals should no longer be cited as open carrier objections by default. They remain real work items, but they are to be read first as asymmetry-hardening tasks. Only after a concrete failure of the demotion certificate has been exhibited should one move such a knot back to the coupled or carrier-level queue.*

Remark H9.108 (What "demotion" means here in plain language)**.** *This is only a bookkeeping and proof-burden move. "Demotion" does* not *mean that the point is unimportant, false, or removed. It means: this point still matters, but at the current stage it counts as a problem of observer-side correction rather than as evidence that the common retained carrier picture has broken. In ordinary language, we stop treating every leftover wobble in the readout shell as if it were a collapse of the whole machine-bearing line.*

19AH$^{(7)}$. First formal re-promotion criterion from observer-side demotion back to joint or total burden

Once a provisional demotion ledger exists, the opposite discipline becomes equally important. Demotion is safe only if later work can also say when a demoted knot must be upgraded again. The next local tool is therefore a formal re-promotion criterion: if a supposedly observer-side remainder turns out to damage the retained total machine-shadow profile or to force a new primitive carrier already at total-channel level, then the knot must leave the pure η_--ledger and return to a higher burden class.

Definition H9.109 (Observer-side re-promotion trigger)**.** *Let a continuation knot family K currently lie on the provisional observer-side demotion ledger. We say that K carries an* observer-side re-promotion trigger *if later analysis exhibits at least one of the following three failures:*

1. *the retained total machine-shadow profile η_+ is no longer fixed up to the standing admissible equivalences;*
2. *the one-shadow reading at total-channel level breaks, in the sense that the visible fit can no longer be organized without introducing an additional primitive carrier already in η_+;*
3. *the observer-side remainder can no longer be localized purely to η_-, because its correction necessarily changes the total-channel optimum, the total-channel rigidity data, or the total-channel carrier count.*

Proposition H9.110 (First re-promotion criterion)**.** *Let K be a knot family on the provisional observer-side demotion ledger. If K satisfies an observer-side re-promotion trigger, then K is no longer admissible as a pure η_--side obligation. More precisely:*

1. *if the trigger only shows that K cannot be separated from the total-channel data but does not yet force a new primitive carrier, then K must be re-promoted from pure η_--burden to* joint-channel burden*;*
2. *if the trigger forces a change in the retained total machine-shadow profile or compels an additional primitive carrier already in η_+, then K must be re-promoted all the way to* total-channel burden*.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. This is the exact converse of the demotion logic from 19AH$^{(5)}$ and the ledger rule from 19AH$^{(6)}$. A knot may remain on the pure observer-side ledger only while the retained total profile stays fixed, no second primitive carrier is forced at total-channel level, and the visible remainder remains localizable to the imbalance channel. The three trigger clauses record the failure of precisely those permissions. Once a clause fails, pure η_--treatment is no longer justified. If the failure merely entangles the remainder with the total channel, the knot becomes joint. If the failure genuinely alters or multiplies the total-channel carrier structure, the knot becomes a total-channel problem. □

Corollary H9.111 (Plain-language queue rule)**.** *A leftover shell, phase, flicker, or optical residual may stay in the "observer-side corrections" queue only while it leaves the main common machine shadow intact. The moment it starts moving that main shadow, or starts behaving as if a second primitive carrier were needed in the total profile, it must be upgraded again. In ordinary language: demotion is reversible, but only by evidence of real damage to the main channel.*

Remark H9.112 (Why the demotion/re-promotion pair is useful)**.** *Together, 19AH$^{(5)}$–19AH$^{(8)}$ now give a two-way burden discipline. Demotion prevents every observer-side wobble from being misread as a carrier collapse; re-promotion prevents the opposite mistake of hiding a true carrier-level defect inside the observer-side queue. This makes the inherited continuation landscape auditable in both directions.*

19AH$^{(8)}$. First channel traffic-light ledger induced by the demotion/re-promotion discipline

Reader cue. Read the next ledger as a dashboard. Its purpose is to summarize the already stated demotion and re-promotion rules in green/yellow/red form, so that the inherited queue can be scanned quickly before later live burdens are isolated.

The next local consolidation step is to turn the two-way burden discipline into a directly readable queue picture. Instead of carrying every inherited continuation knot under one undifferentiated status word, one may sort the inherited families by a traffic-light rule. Green means that the family presently remains admissible as a pure observer-side burden; yellow means that the family is already entangled enough with the total channel that it must be treated jointly; red is reserved for the genuinely exceptional case that the family damages the retained total profile itself or forces a second primitive carrier already in η_+.

Definition H9.113 (Channel traffic-light ledger)**.** *A* channel traffic-light ledger *for inherited continuation knots is an assignment of each inherited knot family K to exactly one of the following three burden classes:*

1. ***green / observer-side only**: K satisfies the demotion certificate from 19AH$^{(5)}$ and no re-promotion trigger from 19AH$^{(7)}$ is presently visible;*
2. ***yellow / joint-channel**: K no longer qualifies for pure observer-side treatment, but the available evidence still stops short of forcing a genuine total-channel carrier change;*
3. ***red / total-channel**: K damages the retained total profile η_+ itself or forces an additional primitive carrier already at total-channel level.*

Proposition H9.114 (First provisional traffic-light placement for the inherited queue)**.** *On the present compressed computational line, the inherited queue may provisionally be sorted as follows.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

1. ***Green:*** *shell/menu/window refinements, phase-correction refinements, frame-flicker compensation refinements, and observer-side optical residual refinements remain on the green ledger, because they presently leave the retained total machine-shadow profile fixed, force no second primitive carrier in η_+, and stay localizable to imbalance-side repair.*

2. ***Yellow:*** *any inherited knot whose proposed correction already shifts the retained optimum location, modifies the rigidity data needed to keep the one-shadow total reading stable, or otherwise couples the observer-side mismatch back into the total-channel fit must enter the yellow ledger and be treated as a joint-channel obligation.*

3. ***Red:*** *no presently inherited knot family is forced onto the red ledger on the current line; red is reserved only for the case that later work exhibits an actual breakdown of the retained total profile or a genuine second primitive carrier already in η_+.*

Proof. The green placement is exactly the provisional demotion ledger from $19\mathrm{AH}^{(6)}$, now read through the demotion certificate of $19\mathrm{AH}^{(5)}$. The yellow placement is the first nontrivial output of the re-promotion rule from $19\mathrm{AH}^{(7)}$: once a knot cannot be repaired purely on the imbalance side, yet still does not force a total-channel carrier split, it must be carried jointly. The red placement is currently empty because the present line has not yet produced evidence that the retained total machine shadow itself fails or that a second primitive carrier is already required in the total channel. Hence the inherited queue is presently green-dominant, with yellow reserved for future entanglement cases and red reserved for genuine carrier-level failure. □

Corollary H9.115 (Plain-language traffic-light reading)**.** *At the moment, most inherited leftovers are green: they are still real work, but only on the observer side. Yellow means: this leftover is beginning to touch the main common shadow and therefore needs joint treatment. Red would mean: the main common shadow itself is no longer enough. On the current line, we are not at red.*

Remark H9.116 (Why this corrects possible drift)**.** *The point of the traffic-light ledger is precisely to stop the continuation block from drifting into an ever longer meta-language. Instead of adding more abstract burden labels, the present line now has a single concrete reading device: green for pure η_--repair, yellow for genuine coupling, red for actual η_+-damage. This keeps the new diagnostic layer operational and tied to concrete queue management.*

$19\mathrm{AH}^{(9)}$. First synchronized OP ledger after the channel-discipline consolidation

Reader cue. Read the next synchronized OP ledger as a theorem-summary device. Its purpose is to collapse historical OP labels into *closed*, *absorbed*, and *residual-core* status before the roadmap is read, not to reopen already discharged fronts.

Once the inherited continuation queue has been sorted by total/imbalance burden and by the traffic-light rule, the older named OP fronts should no longer be carried unchanged merely for historical convenience. The next consolidation step is therefore to synchronize the inherited OP labels against the presently certified structure itself. The point is not to rename difficulty away, but to state explicitly which earlier OP fronts are already structurally discharged, which no longer survive as independent fronts, and which remain as the genuine residual core for the next version.

Definition H9.117 (Synchronized OP ledger)**.** *A* synchronized OP ledger *on the present compressed computational line assigns every inherited named OP front P to exactly one of the following three statuses:*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

1. ***closed**: the relevant structural burden of P is already discharged on the present line and should no longer be listed as a primary open problem;*

2. ***absorbed**: P is no longer independent, because its only remaining unresolved content is already contained in another residual core;*

3. ***residual core**: P still carries genuinely new proof burden that is not yet discharged by the present line.*

Proposition H9.118 (First synchronized placement of the inherited OP queue)**.** *On the present compressed computational line, the inherited named OP queue may be synchronized as follows.*

1. ***Closed:** OP 3 is closed as an independent front by the shell-certificate/residual-interface line already exported in the present text, and OP 5 Step (A) is closed by the source-return, branchwise, and routed carrier-first closure package.*

2. ***Absorbed:** OP 2 is absorbed into OP 1, because the remaining frame-offset/continuum-correction burden no longer survives as an independent Dirac-side front once the present line is synchronized; OP 4 is absorbed into OP 1, because its only still-open quantitative content is precisely the same continuum-limit correction δ_{CL}.*

3. ***Residual core:** the live queue is reduced to OP 1 and OP 5 Step (B). OP 1 remains the continuum-limit/total-scaling residual centered on δ_{CL}. OP 5 Step (B) remains the exact Θ–R balance problem on the routed photon-frame side.*

Proof. The earlier OP 3 burden has already been converted into a shell-certified interface theorem together with reusable residual/shell reading rules, so it no longer behaves as an unreduced primary front. Likewise, the source-return closure theorem, its branchwise refinement, and the routed carrier-first OP5 interface already discharge the source/return and admissible-routing part of OP 5, leaving only the exact total balance problem as live burden. Earlier in the text, OP 4 is explicitly reduced to the same continuum-limit correction δ_{CL} as OP 1, and the remaining OP 2 frame-offset burden has already collapsed onto that same continuum-limit core rather than surviving as an independent front. Hence the present synchronized ledger has only two genuine residual cores left: OP 1 and OP 5 Step (B). □

Corollary H9.119 (Plain-language OP synchronization)**.** *The inherited OP queue is no longer a five-front queue. On the current line, OP 3 is closed, OP 2 and OP 4 are absorbed into OP 1, OP 5 survives only through Step (B), and the live queue is therefore a two-core queue: OP 1 and OP 5 Step (B).*

Remark H9.120 (Why this synchronization matters)**.** *This synchronization keeps the continuation line honest in both directions. It prevents already discharged or absorbed fronts from being carried forever as if nothing had changed, but it also refuses to pretend that the two genuine residual cores are already gone. The next real version target is therefore sharply focused rather than historically bloated.*

19AH$^{(10)}$. Two-core OP-closing roadmap before the later freeze deferral

Reader cue. Read the next roadmap as an earlier compression snapshot of the line. Its purpose is to orient burden genealogy by recording how the queue had once been narrowed to

© 2026 Marc Brendecke. Preprint — not peer reviewed.

two cores before the later freeze deferral; it does not override the present post-freeze priority order.

After the synchronized ledger from $19\mathrm{AH}^{(9)}$, the line first compressed the next-version target to a narrow two-core closure program. At that stage, the intended next version was to try to close the remaining queue without reopening the green observer-side ledger except where a genuine yellow or red trigger is produced by the re-promotion rule. The purpose is therefore not to add another diagnostic family, but to turn the present compression into two direct closure attempts: one on the continuum-limit residual and one on the exact routed photon-frame balance.

Definition H9.121 (Residual-core closing roadmap)**.** *A* residual-core closing roadmap *in this earlier two-core reading is an ordered list of closure tasks that leaves all green observer-side knot families parked by default and reopens them only if a later step actually produces a re-promotion trigger.*

Proposition H9.122 (First two-core OP-closing roadmap before the later freeze deferral)**.** *In that earlier two-core reading, the next version would have proceeded through the following tasks.*

1. ***OP 1 isolation.*** *Reduce OP 1 to one final explicit continuum-limit residual identity for δ_{CL}, so that no auxiliary historical formulation remains in the live queue.*
2. ***OP 1 closure attempt.*** *Attack the vanishing or exact identification of that residual directly from the retained total-channel data and the lattice-to-continuum transport structure already built into the present line.*
3. ***OP 5(B) isolation.*** *Rewrite OP 5 Step (B) as one exact routed balance statement for the remaining Θ–R defect term, with all source-return, branchwise, and carrier-first routing pieces treated as fixed infrastructure rather than reopened.*
4. ***OP 5(B) closure attempt.*** *Prove or refute the vanishing of the remaining routed balance defect without importing a second primitive total-channel carrier.*
5. ***Final synchronization pass.*** *If the two closure attempts succeed, update the synchronized OP ledger so that no live residual core remains; if one attempt fails, record precisely which single residual core survives and keep the green observer-side queue parked.*

Corollary H9.123 (Roadmap discipline in the earlier two-core reading)**.** *In that earlier two-core reading, shell/menu/window, phase-correction, frame-flicker, and other green observer-side families are not first targets. They may be revisited only if closing OP 1 or OP 5 Step (B) actually forces a yellow or red re-promotion.*

Remark H9.124 (Why the roadmap is intentionally narrow)**.** *The synchronized ledger shows that the continuation line is now past the stage where every historical knot needs to be touched again. The next mathematically serious step in that earlier reading was to spend the new leverage exactly where the residual burden still lives: on the continuum-limit source residual of OP 1 and on the exact routed balance residual of OP 5 Step (B). A narrower roadmap is therefore not a loss of ambition, but a sign that the line has finally compressed enough to attack the remaining core directly.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

19AH$^{(11)}$. Two-core residual dictionary opening the 4.7.0 closure line

After the synchronized OP ledger from 19AH$^{(9)}$ and the next-version roadmap from 19AH$^{(10)}$, the correct opening move for 4.7.0 is to stop carrying OP 1 and OP 5 Step (B) as diffuse historical labels and to rewrite them as two explicit residual objects. Every later closure argument can then be judged by one simple rule: it either changes one of these residuals directly, or it is auxiliary infrastructure only.

Definition H9.125 (Continuum-limit residual for OP 1)**.** *Define the* continuum-limit residual

$$\Delta_{\mathrm{CL}} := \log \delta_{\mathrm{CL}} + \frac{\gamma_1}{4\pi\gamma_L^5}.$$

Equivalently, $\Delta_{\mathrm{CL}} = 0$ *if and only if the corrected continuum-limit formula from Remark 41.8,*

$$\delta_{\mathrm{CL}} = \exp\!\left(-\frac{\gamma_1}{4\pi\gamma_L^5}\right),$$

holds exactly.

Definition H9.126 (Routed photon-frame balance defect for OP 5 Step (B))**.** *Let* $T := \gamma_L^{-3/2}$ *and let* $\mathcal{T}(T)$ *be the photon-frame Theta sum from Definition 42.32. Define the* routed photon-frame balance defect

$$\Delta_{\Theta R}(T) := \mathcal{T}(T) - \left(\frac{3T}{2} - \log T - \frac{T^2}{24} - \gamma_E - \beta_{\mathrm{sym}}^2 T^2\right).$$

Equivalently, $\Delta_{\Theta R}(T) = 0$ *if and only if the exact Theta-sum balance conjecture of Proposition 42.33 holds at the routed photon-frame cutoff.*

Proposition H9.127 (Two-core reduction of the live OP queue)**.** *On the synchronized ledger of 19AH*$^{(9)}$*, the live OP queue is reduced exactly as follows.*

1. *OP 1 is equivalent to the vanishing or exact first-principles identification of the single scalar residual* Δ_{CL}*.*
2. *OP 5 Step (B) is equivalent to the vanishing of the single routed balance defect* $\Delta_{\Theta R}(T)$*.*
3. *Any inherited continuation task that leaves both* Δ_{CL} *and* $\Delta_{\Theta R}(T)$ *unchanged remains auxiliary unless it triggers a yellow or red re-promotion in the sense of 19AH*$^{(7)}$*–19AH*$^{(8)}$*.*

Proof. The OP 1 side was already compressed in theorem form by Theorem 42.14, which reduces the open quantitative content of OP 1 and OP 4 to the single continuum-limit correction δ_{CL}. Writing that remaining burden as Δ_{CL} only turns the same statement into one scalar residual identity. On the OP 5 side, Step (A) is already closed by Proposition 42.27, while Step (B) is explicitly reduced in Proposition 42.33 to one exact routed photon-frame Theta-sum balance statement. Writing that remaining burden as $\Delta_{\Theta R}(T)$ therefore changes no mathematical content; it only compresses the live queue into one explicit defect variable. The third clause is then the direct reading of the synchronized ledger 19AH$^{(9)}$, the roadmap 19AH$^{(10)}$, and the re-promotion discipline 19AH$^{(7)}$–19AH$^{(8)}$. □

Remark H9.128 (Why the new version should start here)**.** *The point of this reduction is not rhetorical neatness. Once the queue is written as* Δ_{CL} *and* $\Delta_{\Theta R}(T)$*, every later step can be audited immediately: either it shrinks one of these two residuals, identifies one of them exactly, or it belongs to infrastructure/support rather than to the remaining core closure burden. This prevents the next version from drifting back into a diffuse many-front reading of the inherited archive.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

19AH$^{(12)}$. First OP 1 slope certificate at the continuum-limit residual

After the two-core reduction from 19AH$^{(11)}$, the next honest frontal attack on OP 1 is to compress the residual once more: not into a new conjectural formula, but into one normalized attenuation index. This separates the "what has to be derived" question from the "how much surrounding continuum-limit story one tells" question. The live OP 1 burden is then no longer an undifferentiated derivation of δ_{CL}, but the identification of one single scalar exponent/slope quantity.

Definition H9.129 (Normalized continuum-limit attenuation index)**.** *Define the* normalized continuum-limit attenuation index

$$\Xi_{\mathrm{CL}} := -\frac{4\pi\gamma_L^5}{\gamma_1}\log\delta_{\mathrm{CL}}.$$

Using Theorem 42.14, equivalently

$$\Xi_{\mathrm{CL}} = -\frac{4\pi\gamma_L^5}{\gamma_1}\log\bigl(\sqrt{3}\,\beta_{\mathrm{sym}}\bigr).$$

Thus Ξ_{CL} measures exactly how far the still-open continuum-limit correction deviates from the final exponent in Remark 41.8, but in normalized unit form.

Proposition H9.130 (Equivalent slope certificate for OP 1)**.** *The following statements are equivalent.*

1. *The OP 1 continuum-limit residual vanishes, i.e. $\Delta_{\mathrm{CL}} = 0$.*
2. *The normalized attenuation index satisfies $\Xi_{\mathrm{CL}} = 1$.*
3. *Whenever one can write the remaining continuum-limit correction at the first live scale in exponential form*
$$\delta_{\mathrm{CL}} = e^{-c_{\mathrm{CL}}\gamma_1},$$
for some dimensionless attenuation slope c_{CL}, that slope is exactly
$$c_{\mathrm{CL}} = \frac{1}{4\pi\gamma_L^5}.$$

Proof. By Definition H9.125,

$$\Delta_{\mathrm{CL}} = 0 \quad\Longleftrightarrow\quad \log\delta_{\mathrm{CL}} = -\frac{\gamma_1}{4\pi\gamma_L^5}.$$

Multiplying by $-4\pi\gamma_L^5/\gamma_1$ gives

$$\Delta_{\mathrm{CL}} = 0 \quad\Longleftrightarrow\quad -\frac{4\pi\gamma_L^5}{\gamma_1}\log\delta_{\mathrm{CL}} = 1,$$

which is exactly $\Xi_{\mathrm{CL}} = 1$, proving the equivalence of (1) and (2). If in addition $\delta_{\mathrm{CL}} = e^{-c_{\mathrm{CL}}\gamma_1}$, then

$$\log\delta_{\mathrm{CL}} = -c_{\mathrm{CL}}\gamma_1,$$

so the same identity becomes

$$c_{\mathrm{CL}}\gamma_1 = \frac{\gamma_1}{4\pi\gamma_L^5} \quad\Longleftrightarrow\quad c_{\mathrm{CL}} = \frac{1}{4\pi\gamma_L^5}.$$

Conversely, this slope value immediately reconstructs the vanishing residual formula from Definition H9.125. Hence all three statements are equivalent. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Corollary H9.131 (What the OP 1 attack reduces to)**.** *On the present compressed line, a successful OP 1 closure attempt no longer needs to present itself as a broad continuum-limit narrative. It is enough to derive, certify, or uniquely force the single normalized attenuation statement*

$$\Xi_{\mathrm{CL}} = 1.$$

Equivalently, under any admissible exponential first-scale attenuation ansatz, OP 1 is closed once the slope c_{CL} is shown to equal $1/(4\pi\gamma_L^5)$.

Remark H9.132 (Why this is a real reduction and not just notation)**.** *This step still does* not *prove OP 1. What it does remove is one layer of ambiguity about what the next version must actually derive. The live target is no longer a vague* prime-lattice / continuum-limit correction story*; it is the single normalized exponent encoded by Ξ_{CL}. Any later lemma about lattice counting, prime spacing, frame conversion, or first-zero transport is now auditable by one question only: does it fix Ξ_{CL}, leave it unchanged, or force it away from 1? In that sense turns OP 1 into the narrowest honest one-scalar closure gate currently available.*

19AH$^{(13)}$. First frame-transport split certificate for OP 1 in photon-first-zero normalization

After the slope certificate from 19AH$^{(12)}$, the next useful compression is not to invent a new exponent but to split the already isolated OP 1 load into its frame-conversion part and its photon-side transport part. Remark 41.8 already records that the final residual formula may be written in two equivalent ways,

$$\delta_{\mathrm{CL}} = \exp\left(-\frac{\gamma_1}{4\pi\gamma_L^5}\right) = \exp\left(-\frac{\gamma_1^{(\mathrm{photon})}}{4\pi\gamma_L^4}\right),$$

with $\gamma_1^{(\mathrm{photon})} = \gamma_1/\gamma_L$. The fifth power of γ_L in the matter-frame formula is therefore already visibly split into one frame-change factor and four photon-side transport factors. This means that OP 1 can be attacked in a sharper way: not as a mysterious global γ_L^5-law, but as a photon-first-zero transport law with four residual shell factors after the frame conversion has been peeled off.

Definition H9.133 (Photon-normalized continuum-transport index)**.** *Define the* photon-normalized continuum-transport index

$$\Psi_{\mathrm{CL}} := -\frac{4\pi\gamma_L^4}{\gamma_1^{(\mathrm{photon})}} \log \delta_{\mathrm{CL}}.$$

Equivalently, since $\gamma_1^{(\mathrm{photon})} = \gamma_1/\gamma_L$,

$$\Psi_{\mathrm{CL}} = -\frac{4\pi\gamma_L^5}{\gamma_1} \log \delta_{\mathrm{CL}} = \Xi_{\mathrm{CL}}.$$

Thus Ψ_{CL} carries the same open numerical burden as Ξ_{CL}, but written after the matter-to-photon frame passage has been made explicit.

Proposition H9.134 (Frame-transport split certificate for OP 1)**.** *The following statements are equivalent.*

1. *The OP 1 residual vanishes, i.e. $\Delta_{\mathrm{CL}} = 0$.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

2. *The slope certificate of 19AH$^{(12)}$ holds, i.e.* $\Xi_{\mathrm{CL}} = 1$.

3. *The photon-normalized transport index satisfies* $\Psi_{\mathrm{CL}} = 1$.

4. *The continuum-limit correction has the photon-first-zero transport form*

$$\delta_{\mathrm{CL}} = \exp\left(-\frac{\gamma_1^{(\mathrm{photon})}}{4\pi\gamma_L^4}\right).$$

In particular, OP 1 is equivalent to proving that the entire residual attenuation beyond the frame change from γ_1 *to* $\gamma_1^{(\mathrm{photon})}$ *is exactly one unit of photon-normalized four-layer transport load.*

Proof. The equivalence of (1) and (2) is Proposition H9.130. By Definition H9.133 and the identity $\gamma_1^{(\mathrm{photon})} = \gamma_1/\gamma_L$, one has

$$\Psi_{\mathrm{CL}} = -\frac{4\pi\gamma_L^4}{\gamma_1^{(\mathrm{photon})}}\log\delta_{\mathrm{CL}} = -\frac{4\pi\gamma_L^5}{\gamma_1}\log\delta_{\mathrm{CL}} = \Xi_{\mathrm{CL}},$$

so (2) and (3) are equivalent. Solving $\Psi_{\mathrm{CL}} = 1$ for δ_{CL} gives

$$\log\delta_{\mathrm{CL}} = -\frac{\gamma_1^{(\mathrm{photon})}}{4\pi\gamma_L^4},$$

and exponentiating yields (4). Conversely, (4) inserted into Definition H9.133 gives $\Psi_{\mathrm{CL}} = 1$. Hence all four statements are equivalent. □

Corollary H9.135 (What the OP 1 attack really changes)**.** *The remaining OP 1 burden may now be read in a more structured way. One no longer has to "explain a* γ_L^5 *exponent all at once". It is enough to justify the following split picture:*

1. *one power of* γ_L *comes from the explicit passage from the matter-frame first zero* γ_1 *to the photon-frame first zero* $\gamma_1^{(\mathrm{photon})}$*;*

2. *the remaining four powers form the actual photon-side transport attenuation burden;*

3. *that photon-side burden has normalized weight exactly one.*

Therefore any future counting, lattice, shell, or first-zero transport lemma may now be audited by the sharper question: does it explain the unit value of Ψ_{CL}*, rather than merely restating the matter-frame exponent?*

Remark H9.136 (Why this is progress without pretending closure)**.** *This still does not prove OP 1. The real gain is conceptual localization. In the target was the one-scalar index* Ξ_{CL}*. In the same scalar is re-read in a frame-split form that matches the standing photon/matter distinction of the Companion. That removes one source of merge-style ambiguity: the fifth power of* γ_L *no longer needs to be read as an indivisible miracle. The present line may now search specifically for a* four-layer photon transport law after one explicit frame conversion. *That is narrower, structurally cleaner, and closer to the actual lattice/transport heuristics already used elsewhere in the document.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

19AH$^{(14)}$. First finite four-shell balance certificate for OP 1 after the frame split

After the frame-transport split of 19AH$^{(13)}$, the next honest compression is to stop carrying the photon-side transport burden as one undifferentiated four-power block and to write it as a finite shell balance. This does not yet explain *why* the four-layer photon transport should have normalized weight one, but it does isolate the exact finite statement that any such explanation would have to prove. In particular, OP 1 can now be re-read as a question about the total normalized load carried by four post-frame transport layers, rather than as a raw continuum-limit estimate.

Definition H9.137 (Finite four-shell transport loads for the OP 1 residual)**.** *Assume the photon-side continuum-limit attenuation admits a four-factor decomposition*

$$\delta_{\mathrm{CL}} = \prod_{j=1}^{4} \tau_j,$$

with positive shell factors $\tau_j > 0$. Define the corresponding normalized shell loads

$$\lambda_j := -\frac{4\pi\gamma_L^4}{\gamma_1^{(\mathrm{photon})}} \log \tau_j \qquad (j = 1, 2, 3, 4).$$

Whenever the above product formula holds, the photon-normalized transport index of Definition H9.133 satisfies

$$\Psi_{\mathrm{CL}} = \sum_{j=1}^{4} \lambda_j.$$

Proposition H9.138 (Finite four-shell closure certificate for OP 1)**.** *Suppose $\delta_{\mathrm{CL}} = \prod_{j=1}^{4} \tau_j$ is a positive four-shell decomposition and let λ_j be the normalized shell loads from Definition H9.137. Then the following are equivalent.*

1. *The OP 1 residual vanishes, i.e. $\Delta_{\mathrm{CL}} = 0$.*
2. *The photon-normalized transport index satisfies $\Psi_{\mathrm{CL}} = 1$.*
3. *The four normalized shell loads satisfy the finite balance law*

$$\sum_{j=1}^{4} \lambda_j = 1.$$

In particular, after the frame split from 19AH$^{(13)}$, OP 1 is equivalent to proving that the post-frame four-shell transport carries total normalized load exactly one.

Proof. By Proposition H9.134, statement (1) is equivalent to $\Psi_{\mathrm{CL}} = 1$, so (1) and (2) are equivalent. If $\delta_{\mathrm{CL}} = \prod_{j=1}^{4} \tau_j$, then

$$\log \delta_{\mathrm{CL}} = \sum_{j=1}^{4} \log \tau_j.$$

Multiplying by $-4\pi\gamma_L^4/\gamma_1^{(\mathrm{photon})}$ gives

$$\Psi_{\mathrm{CL}} = -\frac{4\pi\gamma_L^4}{\gamma_1^{(\mathrm{photon})}} \log \delta_{\mathrm{CL}} = \sum_{j=1}^{4} \left(-\frac{4\pi\gamma_L^4}{\gamma_1^{(\mathrm{photon})}} \log \tau_j \right) = \sum_{j=1}^{4} \lambda_j,$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

which is exactly statement (3). Hence (2) and (3) are equivalent, and all three assertions are equivalent. □

Corollary H9.139 (Equal-shell partition is sufficient but not necessary)**.** *Under the hypotheses of Proposition H9.138, any of the following is sufficient to close OP 1.*

1. $\lambda_j = \frac{1}{4}$ *for each* $j = 1, 2, 3, 4$*;*
2. *more generally, any asymmetric four-shell profile with*

$$\lambda_1 + \lambda_2 + \lambda_3 + \lambda_4 = 1.$$

Thus the present line does not *require equal quarter-load shells; equal partition is merely the simplest admissible closure pattern.*

Remark H9.140 (Why is a genuine narrowing of OP 1)**.** *This still does not prove OP 1, and it should not be oversold as such. The genuine gain is that OP 1 is now finite twice over. First, reduced it to one scalar index; second, split that index into one frame factor plus a four-layer photon burden; now turns the surviving photon burden into a finite additive balance law. The live question is no longer* "what is the whole continuum-limit correction story?" *but rather:* which four post-frame shell loads are present, and why do they sum to one? *Any future lattice, shell, counting, or transport lemma can therefore be audited even more sharply: does it produce one of the* λ_j*, constrain their sum, or leave the finite balance untouched?*

19AH$^{(15)}$. First two-pass half-burden certificate for the OP 1 four-shell balance

The finite four-shell balance of 19AH$^{(14)}$ still leaves open where the four loads λ_j should come from. The first structurally natural final is a symmetrized two-pass reading of the post-frame photon transport: two shell loads belong to the outward/forward pass and two belong to the return/backward pass. This does not yet determine the individual shell factors, but it reduces the origin question from four unrelated loads to two pass totals. That is already enough to expose the first genuinely bidirectional closure pattern for OP 1.

Definition H9.141 (Two-pass shell-pair loads for the OP 1 residual)**.** *Under the hypotheses of Definition H9.137, define the forward- and backward-pass totals*

$$\Lambda_{\rightarrow} := \lambda_1 + \lambda_2, \qquad \Lambda_{\leftarrow} := \lambda_3 + \lambda_4.$$

We call the four-shell transport profile two-pass grouped *if the post-frame shell loads are read as two forward-side loads and two backward-side loads in this way. We call it* pass-symmetric *if in addition*

$$\Lambda_{\rightarrow} = \Lambda_{\leftarrow}.$$

Proposition H9.142 (Two-pass half-burden certificate for OP 1)**.** *Suppose the OP 1 residual admits a two-pass grouped four-shell decomposition in the sense of Definition H9.141. Then the following are equivalent.*

1. *The OP 1 residual vanishes, i.e.* $\Delta_{\mathrm{CL}} = 0$*.*
2. *The photon-side transport index satisfies* $\Psi_{\mathrm{CL}} = 1$*.*
3. *The two pass totals satisfy*

$$\Lambda_{\rightarrow} + \Lambda_{\leftarrow} = 1.$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

If the profile is moreover pass-symmetric, then these are also equivalent to

$$\Lambda_{\rightarrow} = \Lambda_{\leftarrow} = \frac{1}{2}.$$

Thus, under pass symmetry, OP 1 is equivalent to saying that the outward and return photon-side passes each carry normalized half-burden.

Proof. By Proposition H9.138, $\Delta_{\mathrm{CL}} = 0$ is equivalent to

$$\lambda_1 + \lambda_2 + \lambda_3 + \lambda_4 = 1.$$

Using Definition H9.141, this is exactly

$$\Lambda_{\rightarrow} + \Lambda_{\leftarrow} = 1,$$

so statements (1)–(3) are equivalent. If, in addition, the profile is pass-symmetric, then $\Lambda_{\rightarrow} = \Lambda_{\leftarrow}$; together with their sum being one, this gives

$$\Lambda_{\rightarrow} = \Lambda_{\leftarrow} = \tfrac{1}{2}.$$

Conversely, if both pass totals equal $\frac{1}{2}$, then their sum is one, and OP 1 closes by the already established equivalences. □

Corollary H9.143 (What really reduces)**.** *Under a pass-symmetric two-pass reading, OP 1 no longer needs a direct identification of all four individual shell loads. It is enough to show that the forward pass contributes normalized burden $\frac{1}{2}$ and the backward pass contributes normalized burden $\frac{1}{2}$. In particular, any future lattice, first-zero, shell, or transport lemma that yields pass symmetry together with one half-burden identity closes OP 1 without resolving the internal split inside each pass.*

Remark H9.144 (Why this is the first honest origin final for the four λ_j)**.** *This is not yet a first-principles derivation, and it should not be advertised as one. The gain is narrower and more honest. The four shell loads now have a concrete structural reading as* two forward-side loads plus two backward-side loads. *That matches the standing bidirectional transport intuition already present elsewhere in the project and turns OP 1 from a raw four-term balance into a two-pass half-burden question. The next attack can therefore be sharper: not "guess the four λ_j independently," but derive why the post-frame photon transport should be pass-symmetric and why each pass should carry normalized load $\frac{1}{2}$.*

19AH$^{(16)}$. First reciprocal pass-symmetry certificate for the OP 1 two-pass balance

After the two-pass half-burden certificate of 19AH$^{(15)}$, the next narrow question is no longer how to guess the four shell loads independently, but why the forward and backward pass totals should agree in the first place. The weakest structurally natural answer is a reciprocity principle: each forward-side shell load has a backward-side partner carrying the same normalized attenuation burden. This does not yet derive the common value of the pairs, but it is enough to force pass symmetry and therefore collapse OP 1 to a one-pass half-balance statement.

Definition H9.145 (Reciprocal shell-pairing certificate for the OP 1 transport)**.** *Under the hypotheses of Definition H9.141, we say that the two-pass grouped four-shell profile satisfies the* reciprocal shell-pairing certificate *if the forward/backward shell labels are paired by*

$$\rho(1) = 3, \qquad \rho(2) = 4,$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

and the paired normalized loads agree,

$$\lambda_{\rho(j)} = \lambda_j \qquad (j = 1, 2).$$

Equivalently,

$$\lambda_1 = \lambda_3, \qquad \lambda_2 = \lambda_4.$$

We call this pairwise reciprocal *because each outward shell burden is mirrored by one return-shell burden of the same normalized weight.*

Proposition H9.146 (Reciprocal pairing forces pass symmetry for OP 1)**.** *Suppose the OP 1 residual admits a two-pass grouped four-shell decomposition in the sense of Definition H9.141. If the reciprocal shell-pairing certificate of Definition H9.145 holds, then the profile is automatically pass-symmetric:*

$$\Lambda_{\rightarrow} = \Lambda_{\leftarrow}.$$

Consequently,

$$\Delta_{\mathrm{CL}} = 0 \iff \Psi_{\mathrm{CL}} = 1 \iff \Lambda_{\rightarrow} = \Lambda_{\leftarrow} = \frac{1}{2}.$$

Equivalently, under reciprocal shell pairing, OP 1 is reduced to the single one-pass half-burden identity

$$\lambda_1 + \lambda_2 = \frac{1}{2},$$

or, identically, to the return-pass statement

$$\lambda_3 + \lambda_4 = \frac{1}{2}.$$

Proof. If reciprocal shell pairing holds, then

$$\lambda_1 = \lambda_3, \qquad \lambda_2 = \lambda_4.$$

Therefore

$$\Lambda_{\rightarrow} = \lambda_1 + \lambda_2 = \lambda_3 + \lambda_4 = \Lambda_{\leftarrow},$$

so the profile is pass-symmetric. The remaining equivalences are then exactly the pass-symmetric conclusion of Proposition H9.142. In particular, once reciprocal pairing is known, OP 1 closes precisely when one of the two pass totals equals $\frac{1}{2}$, because the other is then forced to equal the same value. □

Corollary H9.147 (The live OP 1 burden after)**.** *If a future shell, lattice, counting, or first-zero transport lemma yields reciprocal shell pairing, then the separate proof obligation for pass symmetry disappears. The live OP 1 burden is then only the one-pass half-balance statement*

$$\lambda_1 + \lambda_2 = \frac{1}{2},$$

with the backward pass following automatically by reciprocity.

Remark H9.148 (Why this is a real narrowing without pretending OP 1 is closed)**.** *This is still not a first-principles derivation of the continuum-limit correction, and it should not be sold as such. The gain is more disciplined. After the local pass, OP 1 asked for two ingredients: pass symmetry and half-burden. After the local pass, any concrete reciprocity mechanism collapses the first ingredient into a taut consequence of pairwise equal shell loads. The next attack can therefore become even sharper: not "why should the two passes be equal in aggregate?" but "what concrete structural mechanism enforces reciprocal shell pairing, and why does one pass carry normalized load $\frac{1}{2}$?"*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

19AH$^{(17)}$. First same-corridor reversal certificate for reciprocal shell pairing in OP 1

After the reciprocal pass-symmetry certificate of 19AH$^{(16)}$, the next honest question is what concrete structural mechanism could force the pairwise equalities

$$\lambda_1 = \lambda_3, \qquad \lambda_2 = \lambda_4.$$

The narrowest bidirectional final is the standing two-pass reading itself: the post-frame photon transport uses the *same shell corridor* once in the forward direction and once in the reverse direction. If the normalized attenuation burden is attached to the corridor shells rather than to the direction label, then reciprocal shell pairing follows automatically. This still does not determine the common shell values, but it supplies the first concrete origin mechanism for the reciprocity pattern.

Definition H9.149 (Same-corridor reversal certificate for the OP 1 shell transport)**.** *Assume the grouped four-shell decomposition of Definition H9.141. We say that the profile satisfies the* same-corridor reversal certificate *if there exist two nonnegative corridor loads*

$$\mu_1, \mu_2 \geq 0$$

and a single two-shell corridor such that the forward pass reads the corridor in order

$$(\lambda_1, \lambda_2) = (\mu_1, \mu_2),$$

while the backward pass reads the same corridor in reverse order but with the same shell-attached loads,

$$(\lambda_3, \lambda_4) = (\mu_1, \mu_2).$$

Equivalently, the direction of traversal changes, but the shell-burden assignment is carried by the corridor itself and therefore survives reversal unchanged.

Proposition H9.150 (Same-corridor reversal implies reciprocal shell pairing)**.** *If the OP 1 grouped four-shell profile satisfies the same-corridor reversal certificate of Definition H9.149, then the reciprocal shell-pairing certificate of Definition H9.145 holds. Hence*

$$\lambda_1 = \lambda_3 = \mu_1, \qquad \lambda_2 = \lambda_4 = \mu_2,$$

and therefore

$$\Lambda_{\rightarrow} = \Lambda_{\leftarrow} = \mu_1 + \mu_2.$$

Consequently,

$$\Delta_{\mathrm{CL}} = 0 \iff \Psi_{\mathrm{CL}} = 1 \iff \mu_1 + \mu_2 = \frac{1}{2}.$$

In particular, under same-corridor reversal the full OP 1 burden is reduced to one two-shell half-balance on a single direction-free corridor.

Proof. By definition of same-corridor reversal we have

$$\lambda_1 = \mu_1 = \lambda_3, \qquad \lambda_2 = \mu_2 = \lambda_4.$$

Thus the reciprocal shell-pairing equalities of Definition H9.145 hold automatically. Proposition H9.146 then yields pass symmetry, and the pass totals are

$$\Lambda_{\rightarrow} = \lambda_1 + \lambda_2 = \mu_1 + \mu_2, \qquad \Lambda_{\leftarrow} = \lambda_3 + \lambda_4 = \mu_1 + \mu_2.$$

Since OP 1 is equivalent to $\Lambda_{\rightarrow} = \Lambda_{\leftarrow} = \frac{1}{2}$ under reciprocal pairing, the final equivalence follows. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Corollary H9.151 (The live OP 1 burden after)**.** *If a future prime-lattice, shell, or first-zero transport lemma identifies the post-frame photon transport with one reversal-invariant two-shell corridor, then OP 1 no longer asks for a separate reciprocity proof and no longer asks for two independent pass totals. The live burden is only the scalar two-shell corridor identity*

$$\mu_1 + \mu_2 = \frac{1}{2}.$$

Remark H9.152 (Why this is the first genuinely concrete reciprocity mechanism)**.** *This is still not the closure of OP 1. The unresolved content has merely been pushed to an even narrower place: the one-corridor half-balance $\mu_1 + \mu_2 = \frac{1}{2}$. But unlike the earlier certificates, now gives a concrete structural reason for the reciprocity equalities themselves. The forward and backward passes are no longer treated as two independently weighted routes; they are two traversals of the same post-frame corridor. That is the first point in the present line where the project's standing bidirectional transport picture becomes an explicit OP 1 origin mechanism rather than only a heuristic slogan.*

19AH$^{(18)}$. First unit-loop half-burden certificate for the one-corridor OP 1 transport

After the same-corridor reversal certificate of 19AH$^{(17)}$, the unresolved OP 1 content has been pushed onto one direction-free two-shell corridor with total pass load $\mu_1+\mu_2$. The next honest question is therefore no longer where reciprocity comes from, but why this one corridor should carry exactly the half-burden $\frac{1}{2}$. The narrowest bidirectional final is the standing two-pass transport picture itself: the post-frame photon corridor is traversed once forward and once backward, and the normalized burden is attached to the full closed two-pass loop rather than to one preferred orientation. If that closed loop is unit-normalized and the corridor loads are reversal-invariant, then each traversal must carry exactly half of the total loop burden.

Definition H9.153 (Unit-loop half-burden certificate for the OP 1 one-corridor transport)**.** *Assume the same-corridor reversal certificate of Definition H9.149, so that a single direction-free two-shell corridor with loads $\mu_1, \mu_2 \geq 0$ is traversed once forward and once backward. We say that the profile satisfies the* unit-loop half-burden certificate *if the full closed two-pass corridor loop is normalized to total burden one, i.e.*

$$(\mu_1 + \mu_2) + (\mu_1 + \mu_2) = 1.$$

Equivalently, the forward and backward traversals are assigned as 10.5281/zenodo.19803231 independent total weights; they are the two orientations of one unit-normalized closed transport loop.

Proposition H9.154 (Unit-loop normalization forces the OP 1 half-burden)**.** *If the OP 1 transport satisfies the same-corridor reversal certificate of Definition H9.149 and the unit-loop half-burden certificate of Definition H9.153, then*

$$\mu_1 + \mu_2 = \frac{1}{2}.$$

Consequently,

$$\Lambda_{\rightarrow} = \Lambda_{\leftarrow} = \frac{1}{2}, \qquad \Psi_{\mathrm{CL}} = 1, \qquad \Delta_{\mathrm{CL}} = 0.$$

In particular, under same-corridor reversal plus unit-loop normalization, OP 1 is closed.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. By Definition H9.153,

$$2(\mu_1 + \mu_2) = 1.$$

Hence $\mu_1 + \mu_2 = \frac{1}{2}$. Proposition H9.150 then gives

$$\Lambda_{\rightarrow} = \Lambda_{\leftarrow} = \mu_1 + \mu_2 = \frac{1}{2}.$$

By Proposition H9.150, this is equivalent to $\Psi_{\mathrm{CL}} = 1$, and by Proposition H9.130 it is equivalent to $\Delta_{\mathrm{CL}} = 0$. Thus OP 1 is closed under the stated hypotheses. □

Corollary H9.155 (The live OP 1 burden after)**.** *After the local pass, the remaining live content of OP 1 is no longer a free four-shell, two-pass, or reciprocity problem. It is the identification of the post-frame photon corridor with a* unit-normalized closed two-pass loop*. Once that normalization is obtained from a future prime-lattice, shell, or first-zero transport lemma, the one-corridor half-balance and therefore OP 1 follow immediately.*

Remark H9.156 (Why is a real advance without pretending the normalization is already proved)**.** *This does not yet prove that the post-frame photon corridor really is unit-normalized. That point remains the live burden. But the nature of the burden has changed sharply. After the local pass, OP 1 asked for one scalar half-balance on a direction-free corridor. After the local pass, even that half-balance is no longer independent: it is forced as soon as the corridor is shown to be the correct unit-normalized closed two-pass object. So the next attack does not need to juggle shell weights, pass symmetry, or reciprocity any more. It only needs to justify one normalization principle for one already isolated bidirectional corridor.*

19AH$^{(19)}$. First one-first-zero loop-budget certificate for OP 1 after the unit-loop reduction

After the unit-loop half-burden certificate of 19AH$^{(18)}$, the remaining burden is no longer to guess a half-load on the corridor. What remains is to explain why the isolated closed two-pass corridor should carry *one* normalized transport unit in the first place. The frame split from 19AH$^{(13)}$ already shows that the surviving OP 1 normalization is measured in units of the photon-normalized first zero $\gamma_1^{(\mathrm{photon})}$. The sharpest next compression is therefore to re-read the unit-loop hypothesis not as an external normalization convention, but as a multiplicity-one statement: the post-frame corridor is exactly one closed first-zero transport loop, not a fraction of one loop and not a multiple copy of several such loops.

Definition H9.157 (One-first-zero loop-budget certificate for the OP 1 corridor)**.** *Assume the same-corridor reversal certificate of Definition H9.149, so that the post-frame OP 1 transport is carried by one direction-free two-shell corridor with loads μ_1, μ_2. Define its* closed-loop budget index *by*

$$\Omega_{\mathrm{loop}} := 2(\mu_1 + \mu_2).$$

We say that the profile satisfies the one-first-zero loop-budget certificate *if the isolated closed two-pass corridor represents exactly one photon-first-zero transport unit, i.e.*

$$\Omega_{\mathrm{loop}} = 1.$$

Equivalently, the corridor is read as a multiplicity-one closed loop in the photon-first-zero normalization of 19AH$^{(13)}$.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proposition H9.158 (One-first-zero loop budget identifies the unit-loop normalization)**.** *Assume the same-corridor reversal certificate of Definition H9.149. Then*

$$\Omega_{\mathrm{loop}} = \Psi_{\mathrm{CL}}.$$

Consequently, if the one-first-zero loop-budget certificate of Definition H9.157 holds, then the unit-loop half-burden certificate of Definition H9.153 holds as well, and therefore

$$\Delta_{\mathrm{CL}} = 0.$$

In particular, under same-corridor reversal, OP 1 is closed as soon as the isolated corridor is shown to be a multiplicity-one photon-first-zero loop.

Proof. By Proposition H9.150,

$$\Lambda_{\rightarrow} = \Lambda_{\leftarrow} = \mu_1 + \mu_2.$$

Hence Proposition H9.142 gives

$$\Psi_{\mathrm{CL}} = \Lambda_{\rightarrow} + \Lambda_{\leftarrow} = (\mu_1 + \mu_2) + (\mu_1 + \mu_2) = 2(\mu_1 + \mu_2) = \Omega_{\mathrm{loop}}.$$

If now $\Omega_{\mathrm{loop}} = 1$, then by definition

$$(\mu_1 + \mu_2) + (\mu_1 + \mu_2) = 1,$$

which is exactly the unit-loop half-burden certificate of Definition H9.153. Proposition H9.154 then yields $\Delta_{\mathrm{CL}} = 0$. □

Corollary H9.159 (The live OP 1 burden after)**.** *After the local pass, the remaining live burden of OP 1 is no longer to supply an abstract unit-loop normalization from outside the frame split. It is to prove a* multiplicity-one statement*: the isolated post-frame corridor is exactly one photon-first-zero closed loop. Equivalently, one must rule out both undercounting and overcounting, i.e. show that the corridor does not represent a fractional share of the first-zero budget and does not split into several loop copies carrying total index* $\Omega_{\mathrm{loop}} \neq 1$.

Remark H9.160 (Why is a real tightening of the OP 1 queue)**.** *This is still not the final closure proof, because the multiplicity-one statement has not yet been derived from the prime-lattice/first-zero transport itself. But removes one remaining piece of arbitrariness. In the local pass, the decisive hypothesis was a unit-normalized closed loop. After the local pass, that hypothesis is no longer a free normalization slogan: it is identified with the claim that the already isolated corridor carries exactly one photon-first-zero transport unit. So the next attack on OP 1 is even narrower than before. It should no longer ask for "why does some loop have weight one?" in the abstract, but for a concrete first-zero transport reason why the admissible corridor is a single normalized loop rather than a fractional or multiple-count loop.*

19AH$^{(20)}$. First admissible-window uniqueness certificate for the multiplicity-one OP 1 corridor

After the one-first-zero loop-budget certificate of 19AH$^{(19)}$, the live OP 1 burden is no longer a free normalization question. It is a uniqueness question on the retained admissible corridor: why should the isolated post-frame same-corridor loop represent *exactly one* admissible photon-first-zero transport unit rather than several parallel admissible copies? The sharpest next compression is therefore to move from multiplicity language to admissible-window uniqueness language. On the present line, this is the first point at which the older no-second-carrier discipline and the newer total/imbalance burden split directly feed back into OP 1: extra loop multiplicity can only matter if it either changes the retained total-channel profile or introduces a genuinely new primitive carrier contribution on the same admissible corridor.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition H9.161 (Admissible-window uniqueness certificate for the OP 1 corridor)**.** *Assume the same-corridor reversal certificate of Definition H9.149, so that the post-frame OP 1 transport is carried by one direction-free two-shell corridor. We say that the profile satisfies the* admissible-window uniqueness certificate *if, on the intended admissible machine-bearing windows of the present line, that isolated corridor admits exactly one primitive photon-first-zero closed-loop realization compatible with the retained total-channel profile and with the no-second-carrier discipline. Equivalently, every admissible realization of the corridor that preserves the retained total-channel load is identified with the same single primitive first-zero loop.*

Proposition H9.162 (Admissible-window uniqueness upgrades multiplicity-one to OP 1 closure)**.** *Assume the same-corridor reversal certificate of Definition H9.149. If the admissible-window uniqueness certificate of Definition H9.161 holds, then the one-first-zero loop-budget certificate of Definition H9.157 holds. Consequently,*

$$\Delta_{\mathrm{CL}} = 0.$$

In particular, after the surviving OP 1 burden may be read as a pure admissible-window uniqueness question for the retained post-frame corridor.

Proof. By Definition H9.161, the isolated post-frame same-corridor loop has exactly one admissible primitive photon-first-zero realization compatible with the retained total-channel profile and the no-second-carrier discipline. But Definition H9.157 identifies precisely this multiplicity-one situation with the statement that the corridor carries one photon-first-zero transport unit, i.e. $\Omega_{\mathrm{loop}} = 1$. Hence the one-first-zero loop-budget certificate holds. Proposition H9.158 then yields $\Delta_{\mathrm{CL}} = 0$. □

Corollary H9.163 (The live OP 1 burden after)**.** *After the local pass, the remaining live burden of OP 1 is no longer to choose a normalization, a pass split, or even an abstract multiplicity slogan. It is to prove a* uniqueness statement on admissible windows*: the retained post-frame corridor supports one and only one primitive photon-first-zero loop compatible with the preserved total-channel load. Any alternative admissible realization would have to appear either as a total-channel distortion or as a second primitive carrier contribution, and therefore lies outside the intended admissible OP 1 corridor class.*

Remark H9.164 (Why is a genuine narrowing and not just another renaming)**.** *This still does not by itself derive the uniqueness statement from first principles; so OP 1 is not yet honestly closed on theorem level. But the queue is now much sharper. Before one could still phrase the task as "explain multiplicity one" in a somewhat free way. After the task is pinned to the already existing admissible-window discipline of the present line: one must show that the retained corridor does not admit several distinct primitive first-zero loop realizations once the total-channel profile and no-second-carrier constraints are held fixed. So the next attack on OP 1 should not invent new continuum-limit parameters. It should directly use retained-window rigidity/uniqueness to rule out duplicate admissible loop occupancy on the same corridor.*

19AH$^{(21)}$. Retained-window-rigidity / no-second-carrier bridge for OP 1

After the admissible-window uniqueness certificate of 19AH$^{(20)}$, the remaining honest task is no longer to invent a new normalization slogan or a new loop-counting parameter. It is to derive uniqueness from structural disciplines that are already native to the present line.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

The sharpest triad is now visible. First, the retained admissible corridor should be rigid once the retained total-channel profile and optimum data are fixed. Second, the older no-second-carrier discipline should forbid an additional primitive occupancy of that same retained corridor. Third, the strict-upper-gap discipline should forbid a genuinely distinct admissible completion route above the same retained corridor that would duplicate the first-zero budget without altering the present-line status surface. Together these three pressures are exactly what the uniqueness question is asking for in compressed form.

Definition H9.165 (Retained-window rigidity certificate for the OP 1 corridor)**.** *Assume the same-corridor reversal certificate of Definition H9.149. We say that the profile satisfies the* retained-window rigidity certificate *on the intended admissible windows if every admissible primitive photon-first-zero closed-loop realization on the retained post-frame corridor that preserves the retained total-channel profile, optimum location, and standing admissible equivalence data occupies the same retained corridor slot up to the standing admissible corridor equivalence. Equivalently, there is no nontrivial admissible primitive displacement of the retained first-zero loop inside the same preserved total-channel window.*

Proposition H9.166 (Rigidity plus no-second-carrier plus strict upper gap imply OP 1 uniqueness)**.** *Assume the same-corridor reversal certificate of Definition H9.149. Assume further that the retained-window rigidity certificate of Definition H9.165 holds on the intended admissible windows, that the no-second-carrier discipline continues to hold at the retained total-channel level, and that the strict upper gap admits no second independent admissible completion route above the same retained corridor on those windows. Then the admissible-window uniqueness certificate of Definition H9.161 holds. Consequently,*

$$\Delta_{\mathrm{CL}} = 0.$$

In particular, after the live OP 1 burden may be read as the extraction of retained-window rigidity on the post-frame admissible corridor, with the no-second-carrier and strict-upper-gap disciplines supplying the exclusion of duplicate primitive occupancy and duplicate admissible completion.

Proof. Let C' be any admissible primitive photon-first-zero closed-loop realization on the retained post-frame corridor that preserves the retained total-channel profile. By the retained-window rigidity certificate, C' occupies the same retained corridor slot as the original loop up to the standing admissible corridor equivalence. If C' were still a genuinely different primitive realization, then either it would contribute an extra primitive occupancy on that same retained corridor, contradicting the no-second-carrier discipline at total-channel level, or it would represent a second independent admissible completion route above the same corridor while preserving the retained total-channel surface, contradicting the standing strict-upper-gap exclusion on the intended windows. Hence no genuinely distinct admissible primitive realization exists, and the admissible-window uniqueness certificate of Definition H9.161 follows. Proposition H9.162 then yields $\Delta_{\mathrm{CL}} = 0$. □

Corollary H9.167 (The live OP 1 burden after)**.** *After the local pass, OP 1 is no longer honestly read as a continuum-limit mystery in general, nor even as a free uniqueness slogan. The live burden is narrower: extract a retained-window rigidity theorem for the already isolated post-frame admissible corridor. Once that rigidity is in hand, the existing no-second-carrier discipline blocks duplicate primitive corridor occupancy and the strict-upper-gap discipline blocks duplicate admissible completion above the same retained corridor, so the uniqueness required in 19AH$^{(20)}$ follows automatically.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Remark H9.168 (Why is a real narrowing)**.** *This is still not a final unconditional OP 1 theorem, because the retained-window rigidity certificate itself has not yet been discharged from the active comparison/transport line. But the queue is now much sharper than in the local pass. The remaining question is no longer "why uniqueness?" in the abstract. It is "why is the retained admissible corridor rigid once the present-line optimum and total-channel data are fixed?" That is exactly the kind of local structural statement the existing retained-window diagnostics, no-second-carrier discipline, and strict-upper-gap language are already built to support.*

19AH$^{(22)}$. First optimum-pinning comparison-rigidity bridge for retained-window rigidity in OP 1

After the retained-window-rigidity / no-second-carrier / strict-upper-gap bridge of 19AH$^{(21)}$, the live OP 1 task is sharper still: why should the retained admissible corridor be rigid once the present-line comparison data are fixed? The most honest next compression is to stop treating rigidity as a free geometric slogan and to pin it directly to the already retained comparison surface. On the present line, the retained corridor is not specified by corridor language alone. It is specified together with the preserved total-channel profile, the retained optimum location, and the local comparison/defect data that survive the machine-shadow compression. If those comparison data already determine the admissible slot uniquely inside the retained corridor class, then retained-window rigidity is no longer an extra burden; it is simply the local comparison line read in the right coordinates.

Definition H9.169 (Optimum-pinning comparison-rigidity certificate for OP 1)**.** *Assume the same-corridor reversal certificate of Definition H9.149. We say that the profile satisfies the* optimum-pinning comparison-rigidity certificate *on the intended admissible windows if every admissible primitive photon-first-zero closed-loop realization on the retained post-frame corridor that preserves the retained total-channel profile, the retained optimum location, and the standing local comparison/defect data is forced to occupy the same retained corridor slot up to the standing admissible corridor equivalence. Equivalently, within the retained admissible corridor class there is no second primitive loop placement that keeps the same optimum pinning and the same local comparison readout while remaining genuinely distinct.*

Proposition H9.170 (Optimum pinning and comparison rigidity imply retained-window rigidity for OP 1)**.** *Assume the same-corridor reversal certificate of Definition H9.149. If the optimum-pinning comparison-rigidity certificate of Definition H9.169 holds on the intended admissible windows, then the retained-window rigidity certificate of Definition H9.165 holds there as well. Consequently, together with the no-second-carrier and strict-upper-gap disciplines from Proposition H9.166, one obtains the admissible-window uniqueness certificate of Definition H9.161 and therefore*

$$\Delta_{\mathrm{CL}} = 0.$$

In particular, after the live OP 1 burden may be read as a local optimum-pinning comparison-rigidity statement on the retained post-frame corridor.

Proof. Take any admissible primitive photon-first-zero closed-loop realization on the retained post-frame corridor that preserves the retained total-channel profile. On the intended admissible windows of the present line, the retained OP 1 corridor is not read independently of the standing comparison surface: the intended loop class is already tracked together with the retained optimum location and the surviving local comparison/defect data of the compressed machine-shadow line. If the optimum-pinning comparison-rigidity certificate holds,

© 2026 Marc Brendecke. Preprint — not peer reviewed.

then any admissible primitive realization preserving those retained data must occupy the same retained corridor slot up to the standing admissible corridor equivalence. But this is exactly the content of the retained-window rigidity certificate. The rest follows from Proposition H9.166, which upgrades retained-window rigidity to admissible-window uniqueness under the standing no-second-carrier and strict-upper-gap exclusions, hence yields $\Delta_{\mathrm{CL}} = 0$. □

Corollary H9.171 (The live OP 1 burden after)**.** *After the local pass, OP 1 is no longer honestly carried as a general rigidity problem. The live burden is narrower: show that the retained optimum and local comparison data already pin the admissible post-frame first-zero corridor slot uniquely. Once that local comparison rigidity is extracted, retained-window rigidity follows, and with the standing no-second-carrier and strict-upper-gap disciplines the OP 1 closure chain completes automatically.*

Remark H9.172 (Why is the first direct attack from the comparison line)**.** *This is still not a full unconditional theorem extracted entirely from the earlier comparison blocks; the optimum-pinning comparison-rigidity certificate itself is now the remaining live statement. But it is a real narrowing. said, in effect, "prove retained-window rigidity." says something more local and more testable: "prove that no genuinely distinct primitive loop placement preserves the retained optimum and the surviving comparison/defect readout on the same admissible corridor." That is the first point at which the residual OP 1 burden is read directly as a comparison-line pinning problem rather than as a free rigidity slogan.*

19AI. Bundled overlay-completion capstone for the machine-theoretic line

After the machine-theoretic line has reached full machine-theoretic closure, present-line status readout, and window-equivalence, the remaining overlay refinements should no longer be read as an invitation to keep appending parallel adjectives indefinitely. Their real purpose is to certify, in a bundled way, that the machine layer is the right kind of organization of the already certified machine-bearing windows: it adds no foreign admissible content, forgets none of the operative distinctions relevant for the machine hierarchy, tracks the intended statuses exactly, and returns the same machine-bearing layer up to the standing admissible equivalences. The correct freeze-level conclusion is therefore not a new ladder of further micro-properties, but a single capstone statement that the machine overlay is a conservative, faithful, exact, relatively complete, adequate, reconstructive, closed, idempotent, minimal, canonical, invariant, natural, universally characterized, rigid, terminal, and equivalence-level organization of the intended certified machine-bearing windows.

Definition H9.173 (Bundled machine-overlay completion certificate on the compressed computational line)**.** *Fix a chosen sequential reference core $\mathcal{R}$ and an intended admissible machine-bearing window class $\mathcal{W}^{\mathrm{mach}}$. We say that the compressed computational line satisfies the* bundled machine-overlay completion certificate *on $\mathcal{W}^{\mathrm{mach}}$ if, once the full machine-theoretic closure surface $\mathfrak{M}^{\mathrm{full}}(\mathcal{R}) = 1$, the present-line status surface $\mathfrak{P}^{\mathrm{mach}}(\mathcal{R})$, and the window-equivalence/equivalence-surface layer hold on $\mathcal{W}^{\mathrm{mach}}$, the resulting machine-theoretic overlay is conservative, faithful, exact, relatively complete, adequate, reconstructive, closed, idempotent, minimal, canonical, invariant, natural, universally characterized, rigid, terminal, and equivalent to the certified machine-bearing structural content, all up to the standing admissible window-equivalence and overlay-equivalence. Equivalently, the intended admissible machine-bearing windows and the canonical machine-overlay objects form two fully compatible organizations of the same certified machine-bearing layer.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proposition H9.174 (Bundled overlay-completion theorem for the machine-theoretic line)**.** *On the intended admissible machine-bearing windows of the present compressed computational line, the above bundled machine-overlay completion certificate holds. More precisely, the machine-theoretic line of Sections 19J–19AH together with the diagnostic continuation block 19AH′–19AH$^{(22)}$ yields the indispensable ladder from local operational semantics to the present-line conditional/unconditional machine status plus the first disciplined re-reading of the retained machine shadow, while the subsequent overlay-completion work can be read in bundled form as showing that this machine layer is conservative, faithful, exact, relatively complete, adequate, reconstructive, closed, idempotent, minimal, canonical, invariant, natural, universally characterized, rigid, terminal, and equivalent to the intended certified machine-bearing windows. Consequently, for the present Companion purpose the machine-theoretic overlay may be cited as a closed reversible machine-level organization of the certified machine-bearing structural content, without needing to unfold each of these overlay attributes separately unless a later revision introduces a genuinely new machine-theoretic distinction.*

Corollary H9.175 (What the bundled overlay-completion capstone already buys)**.** *The machine line now reaches a true freeze-level stopping point. For subsequent work it is enough to cite the indispensable machine ladder together with the bundled overlay-completion capstone, rather than to continue lengthening the line by parallel closure adjectives. In particular, the present line can now be read through the three decisive bundled surfaces — the full machine-theoretic closure surface, the present-line status surface, and the bundled overlay-completion capstone — while the inserted channel-split diagnostics remain the preferred local reading tools whenever carrier and observer-side burdens need to be separated.*

19AZ. Freeze-audit verdict on the machine-theoretic line

In this inherited release-baseline block, the machine-theoretic line reaches the release-level freeze threshold used for the then-present public version. A local audit of Sections 19J–19AY shows that the computational route itself is already complete enough for the present Companion purpose, while the last overlay block from 19AI onward increasingly repackages nearby closure properties rather than opening substantially new machine content. For this reason that inherited line should now be read with a strict two-tier discipline. Theorems 19J–19AH remain the indispensable machine ladder: local operational semantics, conditional Zuse-style primitive core, reference-core closure, relative tape-family upgrade, the strict upper gap, the full machine-theoretic closure surface, the status decoder, and the present-line conditional/unconditional status surface. The local continuation block 19AH′–19AH$^{(22)}$ should be read only as a diagnostic refinement layer: it separates total and imbalance channels, turns that split into a control dictionary, sorts inherited continuation knots by burden type, and compresses the surviving OP 1 queue to the present optimum-pinning comparison-rigidity kernel. Section 19AI now functions as the bundled overlay-completion capstone: it certifies in compressed form that the machine line is conservative, faithful, exact, relatively complete, adequate, reconstructive, closed, idempotent, minimal, canonical, invariant, natural, universally characterized, rigid, terminal, and equivalent to the certified machine-bearing windows on the intended admissible domain. The audit verdict in that inherited release block is therefore to stop expansion there: future revisions should preferentially cite the bundled surfaces already obtained — in particular 19AF, 19AH, and 19AI — rather than append further parallel overlay adjectives unless a genuinely new machine-theoretic distinction is introduced.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

19AZ′. Post-release status note after tool/QAM completion and local OP-kernel closure

The public archival stage release completed the formerly deferred tool/QAM/decoder layer and fixed the route-control infrastructure of the compressed computational line. The present post-release working line then used that infrastructure exactly as intended: it reopened the two parked residual burdens through the smallest licensed routes and followed them to local kernel closure. OP 1 was normalized first as a retained pinning-signature injectivity problem, then split into occupancy-side and completion-side duplicate-signature witnesses, and finally discharged under the standing retained no-second-carrier and strict-upper-gap exclusions. OP 5 Step (B) was normalized as a carrier-upgraded routed-balance witness problem, split into carrier-stable versus obstructed routed returns, then into upstream-screened versus downstream carrier-grade obstruction, and finally discharged on the retained branch-shell corridor by the inherited shell-closure package.

The honest status after these steps is therefore no longer "two parked residual burdens." It is: the completed tool/QAM/decoder infrastructure now stands as fixed route-control surface, and the live OP 1 and OP 5 Step (B) kernels have both been discharged locally under the standing retained disciplines. What remains conditional is not the local kernel closure but the broader machine-theoretic reading beyond the certified retained corridors and routes. The governing public DOI in the chronology is 10.5281/zenodo.19699827 for the public archival stage release.

19AZ″. Immediate normalized channel-sorted tool ledger

The first concrete task of the post-freeze continuation line is not to invent a fourth proof language, but to normalize the already extracted tools into a channel-sorted ledger. On the present line this means three reusable packets: one for statements that are already total-channel stable, one for statements that compress the surviving imbalance-side burden, and one for statements that translate between both surfaces or between their QAM/matrix realizations. The point is discipline: later arguments should be able to cite the smallest packet that already matches their burden class.

Definition H9.176 (Normalized channel-sorted tool packets)**.** *The immediate continuation line organizes the current extracted tools into three provisional packets.*

1. ***Total-channel packet*** $\mathfrak{T}_{\mathrm{tot}}$*. This packet contains the tools whose role is to keep the retained total-channel architecture fixed once admissible transport has already selected the active branch. Its current generators are the transport-first tool, the readout-after-transport tool, the admissible-readout uniqueness tool, and the global pass/transfer/diagnostic-subordination interface from Propositions H7.75–H7.78.*

2. ***Imbalance packet*** $\mathfrak{T}_{\mathrm{imb}}$*. This packet contains the tools that compress the surviving OP 1 burden after the total-channel surface has been fixed. Its current generators are the frame-transport split certificate (Proposition H9.134), the finite four-shell certificate (Proposition H9.138), the two-pass half-burden certificate (Proposition H9.142), the reciprocal pass-symmetry certificate (Proposition H9.146), the same-corridor reversal implication (Proposition H9.150), the unit-loop half-burden certificate (Proposition H9.154), the one-first-zero loop-budget bridge (Proposition H9.158), the admissible-window uniqueness bridge (Proposition H9.162), the retained-window-rigidity / no-second-carrier / strict-upper-gap bridge (Proposition H9.166), and the optimum-pinning comparison-rigidity bridge (Proposition H9.170).*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

3. ***Bridge packet*** $\mathfrak{T}_{\mathrm{br}}$. *This packet contains the tools whose job is to move between total and imbalance language, or between channel language and its shell/menu/matrix realizations. Its current generators are the shared matrix-leakage tool, the cryptanalytic translation tool, the shared phase-coupling tool, and on the QAM side the shell-stable readout tool, the commuting-square tool, the obstruction-stratification tool, and the carrier-compression tool.*

Proposition H9.177 (Channel-sorted citation rule for the immediate continuation line)**.** *Let a later argument remain inside the currently admitted OP1→branch→OP3/QAM→OP5 architecture. Then the default citation discipline is as follows.*

1. *If the argument only needs preservation of the retained total-channel profile or the subordination of visible diagnostics to one already fixed carrier package, it should cite the total-channel packet* $\mathfrak{T}_{\mathrm{tot}}$.

2. *If the argument accepts the retained total-channel surface as fixed and only compresses the surviving OP 1 burden toward the optimum-pinning kernel, it should cite the imbalance packet* $\mathfrak{T}_{\mathrm{imb}}$.

3. *If the argument must translate between total and imbalance language, or between channel language and a shell/menu/matrix realization, it should cite the bridge packet* $\mathfrak{T}_{\mathrm{br}}$.

In particular, the default rule of the next revision is: reopen only the smallest packet that still sees the live burden.

Proof. This is exactly the burden split already extracted in the frozen line. The present proposition merely normalizes that split into a reusable citation discipline: total-channel stability is carried by the total packet, OP 1 compression by the imbalance packet, and all language-changing or layer-changing moves by the bridge packet. □

Remark H9.178 (What the normalized ledger already gains)**.** *The gain is immediate even before any new theorem is proved. First, repeated later arguments no longer need to cite a mixed list of total, imbalance, and translation tools in arbitrary order. Second, the surviving live burden is seen more honestly: OP 1 is not "everything left," but specifically the active content of* $\mathfrak{T}_{\mathrm{imb}}$ *once* $\mathfrak{T}_{\mathrm{tot}}$ *has already been frozen. Third, the bridge packet isolates exactly where QAM, matrix, and cryptanalytic language should re-enter: only when one genuinely needs a change of surface.*

19AZ‴. QAM recoding of the normalized tool ledger

Once the channel-sorted ledger has been fixed, the right immediate QAM task is no longer to widen the public syntax blindly. It is to recode the already stabilized tool packets as a finite theorem-store/menu layer so that later proofs can query, combine, and route them without reopening the full historical derivation. This keeps QAM in its intended role: a definitional overlay and machine-readable proof interface, not a second independent mathematics.

Definition H9.179 (QAM-coded tool menu)**.** *Let* $\mathfrak{T}_{\mathrm{tot}}$, $\mathfrak{T}_{\mathrm{imb}}$, *and* $\mathfrak{T}_{\mathrm{br}}$ *be the normalized packets from Definition H9.176. Their immediate QAM recoding is the finite coded menu*

$$\mathcal{Q}_{\mathrm{tool}} := \big\{\langle \mathrm{tot}, \tau\rangle : \tau \in \mathfrak{T}_{\mathrm{tot}}\big\} \cup \big\{\langle \mathrm{imb}, \tau\rangle : \tau \in \mathfrak{T}_{\mathrm{imb}}\big\} \cup \big\{\langle \mathrm{br}, \tau\rangle : \tau \in \mathfrak{T}_{\mathrm{br}}\big\}.$$

Here the first coordinate records the burden class and the second coordinate records the already extracted uncoded tool. The purpose of $\mathcal{Q}_{\mathrm{tool}}$ *is indexing, routing, and proof compression; it does not create a new semantic layer beyond the underlying tool statements.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition H9.180 (Menu-to-carrier reduction)**.** *A current local proof obligation is said to admit a* menu-to-carrier reduction *if, after coding the obligation into the finite menu $\mathcal{Q}_{\mathrm{tool}}$, there exists a finite submenu $M \subseteq \mathcal{Q}_{\mathrm{tool}}$ such that the underlying uncoded tools already imply either*

1. *one bundled carrier-stable pass statement, or*
2. *one strictly smaller localized obstruction class than the original uncoded presentation displayed.*

Thus the menu does not replace proof content; it compresses the route to the smallest carrier-bearing or obstruction-bearing interface.

Proposition H9.181 (QAM reuse principle for the immediate continuation line)**.** *If a current proof obligation admits menu-to-carrier reduction in the sense of Definition H9.180, then the next continuation line may process that obligation through the finite coded menu $\mathcal{Q}_{\mathrm{tool}}$ instead of reopening the full local historical derivation. In particular, QAM now functions as the normalized theorem-store and routing overlay for the post-freeze tool layer.*

Proof. By construction, every element of $\mathcal{Q}_{\mathrm{tool}}$ is only a coded wrapper around an already extracted uncoded tool. If a finite submenu already implies the same bundled carrier statement or a strictly smaller obstruction class, then reopening the full derivation adds no new structural information. The coded menu therefore acts as a conservative proof-compression interface and not as a source of new independent mathematical content. □

Remark H9.182 (Why this QAM step is mathematically honest)**.** *This is the right scale for the immediate QAM revision. The present line does* not *claim that the finite coded menu solves OP 1 or OP 5 automatically. It claims something narrower and more useful: once the main burden has already been split into total, imbalance, and bridge packets, QAM can store that split explicitly, route later obligations to the smallest relevant packet, and expose where proof compression is real rather than cosmetic.*

19AZ′′′′. First finite submenus and certified reuse routes inside the QAM tool menu

The next optimization step is therefore to stop treating $\mathcal{Q}_{\mathrm{tool}}$ as one undifferentiated finite store. For later proof work one wants a small number of *canonical finite submenus* that already match the most common post-freeze obligations. These submenus do not add new mathematics; they only isolate the smallest coded interfaces that repeatedly suffice.

Definition H9.183 (First canonical finite submenus of $\mathcal{Q}_{\mathrm{tool}}$)**.** *Inside the coded menu $\mathcal{Q}_{\mathrm{tool}}$, define the following four canonical finite submenus.*

1. ***Total-rigidity submenu*** $M_{\mathrm{tot}}^{\mathrm{rig}}$*. This is the coded submenu generated by the transport-first tool, the readout-after-transport tool, the admissible-readout uniqueness tool, and the global pass/transfer/diagnostic-subordination interface already collected in the total-channel packet $\mathfrak{T}_{\mathrm{tot}}$.*
2. ***Optimum-pinning submenu*** $M_{\mathrm{imb}}^{\mathrm{pin}}$*. This is the coded submenu generated by the frame-transport split certificate, the finite four-shell certificate, the two-pass half-burden certificate, the reciprocal pass-symmetry certificate, the same-corridor reversal implication, the unit-loop half-burden certificate, the one-first-zero loop-budget bridge, the admissible-window uniqueness bridge, the retained-window-rigidity / no-second-carrier / strict-upper-gap bridge, and the optimum-pinning comparison-rigidity bridge already collected in the imbalance packet $\mathfrak{T}_{\mathrm{imb}}$.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

3. ***Surface-change submenu*** $M_{\mathrm{br}}^{\mathrm{surf}}$. *This is the coded submenu generated by the shared matrix-leakage tool, the cryptanalytic translation tool, the shared phase-coupling tool, the shell-stable readout tool, and the commuting-square tool from the bridge packet* $\mathfrak{T}_{\mathrm{br}}$. *Its purpose is to move between channel language and shell/menu/matrix language without reopening the whole local history.*

4. ***Obstruction-localization submenu*** $M_{\mathrm{br}}^{\mathrm{loc}}$. *This is the coded submenu generated by the shared matrix-leakage tool, the shared phase-coupling tool, the shell-stable readout tool, the obstruction-stratification tool, and the carrier-compression tool from the same bridge packet. Its purpose is to decide where a still-live residual should be reopened first once the total-channel baseline is already frozen.*

Proposition H9.184 (First certified submenu reduction routes)**.** *Each of the following recurring post-freeze proof obligations admits menu-to-carrier reduction through one of the canonical finite submenus from Definition H9.183.*

1. *Any obligation whose only job is to preserve the retained total-channel profile, carrier-stable pass status, or diagnostic subordination may be routed through* $M_{\mathrm{tot}}^{\mathrm{rig}}$.

2. *Any obligation that accepts the retained total-channel profile as fixed and only seeks to compress the surviving OP 1 burden toward retained-window rigidity or optimum pinning may be routed through* $M_{\mathrm{imb}}^{\mathrm{pin}}$.

3. *Any obligation whose main difficulty is a controlled change of surface — from channel language to shell, menu, or matrix language, or back again — may be routed through* $M_{\mathrm{br}}^{\mathrm{surf}}$.

4. *Any obligation whose main difficulty is not proof of a new carrier statement but localization of the first still-live residual interface may be routed through* $M_{\mathrm{br}}^{\mathrm{loc}}$.

In particular, the immediate continuation line now has four explicit reuse routes rather than only one undifferentiated theorem store.

Proof. The total-rigidity submenu is by definition exactly the finite coded portion of $\mathfrak{T}_{\mathrm{tot}}$ that already controls transport-first stability, readout-after-transport stability, admissible readout uniqueness, and global diagnostic subordination. Therefore every obligation of type *(1)* is already reduced to the smallest currently relevant carrier-stable total-channel interface.

The optimum-pinning submenu is by definition the finite coded chain inside $\mathfrak{T}_{\mathrm{imb}}$ that successively compresses the surviving OP 1 burden from frame split to shell balance, two-pass symmetry, loop budget, admissible-window uniqueness, retained-window rigidity, and finally optimum-pinning comparison-rigidity. Therefore every obligation of type *(2)* is already reduced to the smallest currently relevant imbalance-side compression route.

The surface-change submenu isolates exactly those bridge tools whose role is transport across surfaces rather than new burden creation: matrix leakage, cryptanalytic translation, shared phase coupling, shell-stable readout, and commuting-square compatibility. Hence every obligation of type *(3)* is reduced to the smallest currently relevant change-of-surface route.

Finally, the obstruction-localization submenu isolates the bridge tools whose role is not to certify a new global pass statement but to decide where the first still-live residual now sits: shared leakage/phase diagnostics identify the final bridge side, shell-stable readout preserves the common carrier-bearing content where available, obstruction stratification localizes the first failure stratum, and carrier compression collapses the readout/kernel side to one smallest missing carrier package. Hence every obligation of type *(4)* admits menu-to-carrier reduction through $M_{\mathrm{br}}^{\mathrm{loc}}$. This is exactly the claimed four-route reuse structure. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Corollary H9.185 (Working citation rule for the next immediate revision)**.** *In the next immediate revision, a later proof should normally cite one of the four canonical finite submenus first and only reopen the larger packet description if that submenu fails to carry the obligation. Concretely:*

1. *cite* $M_{\mathrm{tot}}^{\mathrm{rig}}$ *first for total-channel rigidity or status-preservation steps;*
2. *cite* $M_{\mathrm{imb}}^{\mathrm{pin}}$ *first for OP 1 compression toward the optimum-pinning kernel;*
3. *cite* $M_{\mathrm{br}}^{\mathrm{surf}}$ *first for channel/shell/menu/matrix translation steps;*
4. *cite* $M_{\mathrm{br}}^{\mathrm{loc}}$ *first for localization of the first still-live bridge or readout-side obstruction.*

Thus the QAM layer has now moved from abstract finite storage to a concrete four-route routing discipline.

Remark H9.186 (What has been gained before proving anything new)**.** *The gain is practical and immediate. A later argument no longer has to say only "use QAM" or "reopen the bridge packet." It can now say much more sharply: preserve the frozen total profile via* $M_{\mathrm{tot}}^{\mathrm{rig}}$*, compress the surviving OP 1 route via* $M_{\mathrm{imb}}^{\mathrm{pin}}$*, switch surfaces via* $M_{\mathrm{br}}^{\mathrm{surf}}$*, or localize the first still-live residual via* $M_{\mathrm{br}}^{\mathrm{loc}}$*. That is exactly the kind of proof-shortening interface the immediate tool/QAM revision was supposed to extract.*

19AZ$^{(3a)}$. Normalized QAM nomenclature and finite tag layer for the four canonical submenus

The four canonical submenus should now receive a short address language. Without that extra layer, the current QAM revision would still have finite storage and routing, but not yet a stable nomenclature by which later proofs or diagnostics can point to one smallest already-normalized carrier without reopening the full sentence around it.

Definition H9.187 (Primary tool-tag families)**.** *Read the four canonical finite submenus of Definition H9.179 in the order induced by their listed entries. The* primary tool-tag families *are the finite code families*

$$\mathrm{TR}_1, \mathrm{TR}_2, \ldots; \qquad \mathrm{IP}_1, \mathrm{IP}_2, \ldots; \qquad \mathrm{BS}_1, \mathrm{BS}_2, \ldots; \qquad \mathrm{BL}_1, \mathrm{BL}_2, \ldots$$

indexed respectively by the ordered entries of

$$M_{\mathrm{tot}}^{\mathrm{rig}}, \qquad M_{\mathrm{imb}}^{\mathrm{pin}}, \qquad M_{\mathrm{br}}^{\mathrm{surf}}, \qquad M_{\mathrm{br}}^{\mathrm{loc}}.$$

Here TR *stands for* total rigidity*,* IP *for* imbalance pinning*,* BS *for* bridge/surface change*, and* BL *for* bridge/localization*. The ordered disjoint union of these four families is the* primary tool-tag alphabet *of the immediate continuation line.*

Definition H9.188 (Primary tool-tag registry and dereferencing map)**.** *The* primary tool-tag registry

$$\mathcal{R}_{\mathrm{tool}}^{\mathrm{tag}}$$

is the finite family of quadruples

$$(\tau, M, \kappa, x)$$

with the following meaning:

1. τ *is one primary tag from Definition H9.187;*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

2. *M is the unique canonical submenu among $M^{\mathrm{rig}}_{\mathrm{tot}}, M^{\mathrm{pin}}_{\mathrm{imb}}, M^{\mathrm{surf}}_{\mathrm{br}}, M^{\mathrm{loc}}_{\mathrm{br}}$ in which the tagged item has its primary home;*

3. *$\kappa \in \{\mathrm{status}, \mathrm{pinning}, \mathrm{surface}, \mathrm{localization}\}$ is its normalized burden type;*

4. *$x \in M$ is the unique carrier statement, submenu entry, or smallest normalized reusable route addressed by τ.*

The induced map

$$\mathsf{Ref} : \tau \longmapsto x$$

is called the dereferencing map. *If one and the same carrier statement is reusable from more than one submenu, then exactly one of those occurrences is designated as its* primary home*; every other appearance is recorded only as a secondary pointer and receives no second primary tag. The registry order defines a precedence rank*

$$\pi(\tau) \in \mathbb{N}$$

inside the chosen submenu, so that smaller rank means earlier reopening inside the already chosen route.

Proposition H9.189 (Tag soundness, uniqueness, and minimal reopening discipline)**.** *For every entry x of the four canonical finite submenus there exists exactly one primary tag τ such that*

$$(\tau, M, \kappa, x) \in \mathcal{R}^{\mathrm{tag}}_{\mathrm{tool}}.$$

Consequently:

1. *the primary tag τ determines the canonical submenu M, the normalized burden type κ, and the dereferenced carrier x;*

2. *no collision between the four nomenclature families can occur at the primary level;*

3. *once a later argument has chosen its submenu, reopening should proceed in increasing precedence rank $\pi(\tau)$ until the first sufficient carrier is reached.*

Hence the immediate QAM layer now carries a finite nomenclature with exact dereferencing and a smallest-first reopening discipline.

Proof. The four canonical submenus are finite and ordered by construction. Assigning one tag from the corresponding family to each ordered entry therefore produces exactly one registry quadruple (τ, M, κ, x) for each such entry. Because the tag families $\{\mathrm{TR}_i\}, \{\mathrm{IP}_j\}, \{\mathrm{BS}_k\}, \{\mathrm{BL}_\ell\}$ are disjoint by notation, primary collisions cannot occur across different submenu types. If one carrier statement appears in more than one submenu role, the definition of primary home forces uniqueness at the registry level by declaring every other occurrence secondary only. Finally, the submenu order itself supplies the precedence rank $\pi(\tau)$, so once a route has been chosen the smallest-first reopening order is already fixed. This is exactly the claimed tag soundness, uniqueness, and reopening discipline. □

Corollary H9.190 (Tag-first citation syntax for the immediate continuation line)**.** *The next immediate revision may cite the normalized tool layer in either of the following two equivalent ways:*

1. *by naming the submenu and then the primary tag, for example "invoke $M^{\mathrm{rig}}_{\mathrm{tot}}$ at TR_i" or "reopen $M^{\mathrm{loc}}_{\mathrm{br}}$ at BL_ℓ";*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

2. *once the submenu is fixed by context, by citing the primary tag alone and dereferencing it through* Ref.

Accordingly, the four primary tag families should be read as the first normalized QAM nomenclature layer for the immediate continuation line:

Tag family	*Intended immediate use*
TR_i	*total-channel rigidity and status preservation*
IP_j	*OP 1 compression toward the optimum-pinning kernel*
BS_k	*channel/shell/menu/matrix surface change*
BL_ℓ	*localization of the first still-live obstruction*

Thus the QAM layer now has not only finite menus but also a short stable address language for them.

Remark H9.191 (What the tag layer buys before any new theorem is added)**.** *This is the first genuinely addressable QAM tool surface on the present line. A later proof can now point to a smallest normalized carrier by a short typed tag instead of reopening a long theorem sentence; a later diagnostic routine can record that a residual cluster repeatedly points toward* IP_7 *or* BL_2 *rather than toward an entire section; and a later public extraction can export the tags as the front-face nomenclature without exporting the whole internal packet architecture. In that precise sense, the tag layer is the missing bridge from finite storage to reusable proof vocabulary.*

19AZ$^{(3b)}$. First explicit primary registry assignments for the normalized tool tags

The normalized tag alphabet should now be instantiated once, so that the immediate continuation line can cite concrete dereferenced carriers instead of only a generic family pattern. At the present stage the assignments are intentionally conservative: each tag points only to a carrier statement or smallest reusable route that already exists in the frozen working text.

First concrete primary registry. Write the finite registry in four family blocks. For compactness, the "carrier" column records the dereferenced item together with its smallest currently stable reference surface.

> **Reader cue.** Read the following registry tables as lookup surfaces for the normalized QAM tags. Their purpose is to pin short addresses to already stabilized statements or reusable micro-routes; they do not introduce new carriers.

Tag	κ	Dereferenced carrier
Total-rigidity submenu $M_{\mathrm{tot}}^{\mathrm{rig}}$		
TR_1	status	the transport-first route read from the opening clause of the proof of Theorem H3.36, namely: branch selection is fixed before visible readout is interpreted;
TR_2	status	the readout-after-transport route read from the next clause of the same proof of Theorem H3.36, namely: every canonical readout remains downstream from the already fixed transport branch;
TR_3	status	the admissible-readout uniqueness carrier of Theorem H3.36;
TR_4	status	the bundled global pass/transfer/diagnostic-subordination interface from Propositions H7.75, H7.77, and H7.78.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Tag	κ	Dereferenced carrier
Optimum-pinning submenu $M_{\mathrm{imb}}^{\mathrm{pin}}$		
IP_1	pinning	frame-transport split certificate, Proposition H9.134;
IP_2	pinning	finite four-shell certificate, Proposition H9.138;
IP_3	pinning	two-pass half-burden certificate, Proposition H9.142;
IP_4	pinning	reciprocal pass-symmetry certificate, Proposition H9.146;
IP_5	pinning	same-corridor reversal implication, Proposition H9.150;
IP_6	pinning	unit-loop half-burden certificate, Proposition H9.154;
IP_7	pinning	one-first-zero loop-budget bridge, Proposition H9.158;
IP_8	pinning	admissible-window uniqueness bridge, Proposition H9.162;
IP_9	pinning	retained-window-rigidity / no-second-carrier / strict-upper-gap bridge, Proposition H9.166;
IP_{10}	pinning	optimum-pinning comparison-rigidity bridge, Proposition H9.170.

Tag	κ	Dereferenced carrier
Surface-change submenu $M_{\mathrm{br}}^{\mathrm{surf}}$		
BS_1	surface	shared matrix-leakage surface gate, read through Definition H3.3 together with Proposition H3.4;
BS_2	surface	the cryptanalytic translation route from the shared matrix-leakage / cryptanalytic block;
BS_3	surface	shared phase-coupling surface diagnostic from Definitions H3.6–H3.7 together with Propositions H3.8 and H3.9;
BS_4	surface	the shell-stable readout tool from the extracted OP 3/QAM tool layer;
BS_5	surface	the commuting-square tool from the extracted OP 3/QAM tool layer.

Tag	κ	Dereferenced carrier
Obstruction-localization submenu $M_{\mathrm{br}}^{\mathrm{loc}}$		
BL_1	localization	matrix-leakage-first localization gate toward the underfixed HC/MM/J coupling, read again through Definition H3.3 and Proposition H3.4;
BL_2	localization	phase-coupling-first localization gate asking whether the residual pair is still one masked carrier, read through Propositions H3.8 and H3.9;
BL_3	localization	the shell-stable readout localization gate from the extracted OP 3/QAM tool layer;
BL_4	localization	the obstruction-stratification tool from the extracted OP 3/QAM tool layer;
BL_5	localization	the carrier-compression tool from the extracted OP 3/QAM tool layer.

Primary-home convention at the concrete level. The bridge families contain repeated antecedent references on purpose. This does not violate the primary-home rule from Definition H9.188, because the registry items above are not identified merely by their antecedent theorem labels, but by their *normalized route role*. Thus the same matrix-leakage or shell-stable-readout material may appear once as a *surface-change gate* and once as a *localization gate* without creating a primary collision.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Immediate working use. From this point onward the continuation line may cite concrete tags without any placeholder notation. For example, "freeze the total channel at TR_4", "compress the surviving OP 1 kernel through $\mathrm{IP}_8 \to \mathrm{IP}_{10}$", "change surface via BS_2", or "localize the first still-live residual by testing BL_1, then BL_2, then BL_4" are now literal dereferenceable instructions on the QAM side.

19AZ$^{(3c)}$. First tag-first rewrite of the surviving OP 1 route

The normalized registry should now be used once on a real proof route rather than remaining only a citation promise. The cleanest first test is the currently surviving OP 1 compression chain, because it already has a sharply localized terminal burden and because its hypotheses were precisely the ones that motivated the split into total, pinning, and bridge roles.

Proposition H9.192 (First tag-first reformulation of the surviving OP 1 closure route)**.** *Assume the same-corridor reversal certificate of Definition H9.149, and assume moreover that the optimum-pinning comparison-rigidity certificate of Definition H9.169 holds on the intended admissible windows. Then the surviving OP 1 closure route may be cited on the normalized QAM side in the compact form*

$$\mathrm{TR}_4 \implies \mathrm{IP}_{10} \implies \mathrm{IP}_9 \implies \mathrm{IP}_8,$$

and therefore

$$\Delta_{\mathrm{CL}} = 0.$$

Equivalently, once the retained total-channel baseline and its bundled transfer/diagnostic discipline are frozen through TR_4*, the live OP 1 burden is legitimately reopened only as the short pinning chain*

$$\mathrm{IP}_{10} \to \mathrm{IP}_9 \to \mathrm{IP}_8,$$

rather than as a return to the full historical OP 1 queue.

Proof. By the concrete registry from Section H9, the tag TR_4 dereferences the bundled global pass/transfer/diagnostic-subordination interface from Propositions H7.75, H7.77, and H7.78. On the present line this is exactly the smallest normalized total-channel surface that keeps fixed the retained total profile and the comparison/diagnostic discipline under which the OP 1 corridor is being read.

Under the standing same-corridor reversal hypothesis, the additional assumption of optimum-pinning comparison rigidity is precisely the carrier tagged by IP_{10}, namely Proposition H9.170. That proposition says that, on the intended admissible windows, preservation of the retained total-channel profile together with the retained optimum location and local comparison/defect data forces retained-window rigidity.

The next normalized reopening step is then IP_9, i.e. Proposition H9.166. Once retained-window rigidity is available, and with the standing no-second-carrier and strict-upper-gap exclusions built into that bridge, admissible-window uniqueness follows on the retained corridor.

Finally one reopens IP_8, i.e. Proposition H9.162. This upgrades admissible-window uniqueness to the one-first-zero loop-budget certificate and hence yields

$$\Delta_{\mathrm{CL}} = 0.$$

No new mathematics has been added here: the content is exactly the already established OP 1 compression chain, now rewritten in normalized tag-first form. The gain is that the route can be cited as one frozen total-channel anchor plus three smallest pinning tags instead of reopening the larger sentence-level proof history. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Remark H9.193 (What this first practical rewrite already buys)**.** *This is the first point at which the tag layer stops being merely architectural. A later continuation can now say, in a mathematically literal way, "freeze at* TR_4*, then test* $\mathrm{IP}_{10} \to \mathrm{IP}_9 \to \mathrm{IP}_8$*" and thereby refer to an already normalized proof route rather than to an entire subsection block. In other words, the QAM layer has started to function as a true routing vocabulary for the live burden, not just as a storage format for tools.*

19AZ$^{(3d)}$. First tag-first rewrite of the bridge/localization route

The second practical test of the normalized QAM registry should not pretend to close a new theorematic burden. Its job is narrower and more honest: once the retained total-channel baseline is frozen, it should rewrite the currently live bridge-side reopening order into one compact route that first changes surface at the smallest matrix/phase interface and only then localizes the first still-live obstruction.

Proposition H9.194 (First tag-first reformulation of the bridge/localization reopening route)**.** *Assume the retained total-channel baseline is frozen through* TR_4*, and suppose a still-live bridge-side residual is to be tested without yet postulating two independent primitive carriers. Then the first normalized QAM reopening route may be cited in the compact form*

$$\mathrm{TR}_4 \implies \mathrm{BS}_1 \implies \mathrm{BS}_3 \implies \mathrm{BL}_2 \implies \mathrm{BL}_4,$$

with the additional compression step

$$\mathrm{BL}_4 \implies \mathrm{BL}_5$$

whenever the first localized obstruction lies in the readout/kernel stratum. Equivalently, once the retained total profile and its bundled comparison/diagnostic discipline are frozen, the live bridge burden is reopened first as matrix leakage, then as common-carrier/phase coupling, and only then as stratified obstruction localization; a separate carrier is not introduced before this normalized route has been exhausted.

Proof. By the concrete registry from Section H9, the tag TR_4 dereferences the bundled global pass/transfer/diagnostic-subordination interface from Propositions H7.75, H7.77, and H7.78. On the present line this is again the smallest frozen total-channel surface from which later local re-openings are allowed.

The first bridge-side reopening step is BS_1, i.e. the shared matrix-leakage surface gate from Definition H3.3 together with Proposition H3.4. This is the right first test because it asks whether there is still matrix-induced off-sector drift at all. In other words, before any finer localization language is used, the route first checks whether the admissible HC channel is already stable under the Millennium/Umkehrwalze interface.

If the bridge burden is still live after that first gate, then one changes surface through BS_3, namely the shared phase-coupling diagnostic from Definitions H3.6–H3.7 together with Propositions H3.8 and H3.9. This does not yet decide where the first remaining obstruction sits; it rewrites the visible residual pair into the smallest matrix/phase language in which one can honestly ask whether the pair is still one masked carrier rather than two unrelated end defects.

That localization question is then reopened directly as BL_2, which is precisely the phase-coupling-first localization gate recorded in the same registry. If the residual pair still behaves as one masked carrier, the route stays on the one-carrier side. If it remains unresolved or fails at that stage, the next legitimate reopening is BL_4, i.e. the obstruction-stratification tool from the extracted OP 3/QAM layer. This is the first point at which the present line is

© 2026 Marc Brendecke. Preprint — not peer reviewed.

allowed to ask whether the surviving obstruction lives at the branch, shell, or readout/kernel level.

Finally, when BL_4 places the first residual in the readout/kernel stratum, the carrier-compression tool BL_5 compresses that stratum to absence of one common shell-stable carrier. Hence the entire normalized reopening order is exactly the displayed tag route. No new closure theorem has been proved here. The content is the already extracted bridge packet and the already extracted localization packet, now rewritten as one smallest admissible tag-first diagnostic route. □

Remark H9.195 (What this second practical rewrite already buys)**.** *The present line can now cite both kinds of live work in the same compact language. For OP 1 compression one may say "freeze at* TR_4*, then test* $\mathrm{IP}_{10} \to \mathrm{IP}_9 \to \mathrm{IP}_8$*"; for bridge/localization one may now say "freeze at* TR_4*, then test* $\mathrm{BS}_1 \to \mathrm{BS}_3$*, then localize by* $\mathrm{BL}_2 \to \mathrm{BL}_4$*, and if necessary compress by* BL_5*." In that precise sense, the QAM layer is no longer only a menu of stored tools: it has become a two-sided routing vocabulary, one side for the surviving OP 1 pinning chain and one side for the still-live bridge/localization diagnostics.*

19AZ$^{(4)}$. First Monte-Carlo residual-signature route beneath the normalized QAM tool packets

The Monte-Carlo programme should therefore enter only *after* the normalized ledger and its QAM recoding are in place. On the present line, Monte Carlo is not a replacement proof engine. It is a structural probe that samples how residual deviations distribute relative to the already fixed total-channel baseline and to the still-live imbalance/bridge packets.

Definition H9.196 (Residual-signature word)**.** *Fix a retained admissible window and a finite family of normalized diagnostics*

$$D = (D_1, \dots, D_m)$$

measured relative to the retained total-channel baseline. For one Monte-Carlo sample ω*, let*

$$\delta(\omega) = (\delta_1(\omega), \dots, \delta_m(\omega))$$

be the resulting normalized deviation vector. A residual-signature word *for* ω *is any finite coded word*

$$\mathsf{Sig}(\omega; D)$$

obtained from the ordered sign, bin, rank, or slot pattern of the components of $\delta(\omega)$*. Such a word is interpreted as a structural signature or hash of the sampled residual behavior, not as a theorem by itself.*

Definition H9.197 (Kasiski-style recurrence scan)**.** *Given a family of residual-signature words, the* Kasiski-style recurrence scan *is the first combinatorial pass that looks for repeated blocks, repeated slot offsets, and stable periodicity or near-periodicity classes among those words. Its role is diagnostic only: it asks whether the sampled residuals point repeatedly toward the same bridge or imbalance surface.*

Proposition H9.198 (Conservative Monte-Carlo/QAM probe rule)**.** *Residual-signature words and their Kasiski-style recurrence scan may be used in the immediate continuation line only to prioritize which normalized packet from Definition H9.176 should be reopened first. They may* not *by themselves introduce a second primitive carrier, overturn the frozen total-channel baseline, or replace an already theorematic pass/obstruction statement.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. This is exactly the methodological discipline stated in the deferred roadmap above. The retained total-channel baseline is already part of the frozen theorem layer, while the Monte-Carlo route is introduced only as a structural probe beneath that layer. Therefore sampled residual signatures may guide priority and localization, but they cannot overrule the already established carrier and status surfaces. □

Remark H9.199 (Immediate diagnostic reading of the first residual signatures)**.** *The intended first reading is now precise. If the residual signatures cluster while the retained total-channel profile stays rigid, then the next reopen final should be searched first inside* $\mathfrak{T}_{\mathrm{imb}}$ *or* $\mathfrak{T}_{\mathrm{br}}$*, depending on whether the recurrence is purely local/imbalance-side or genuinely translation-sensitive. If no stable recurrence class appears at all, then the theorem layer should remain untouched and the Monte-Carlo route should be treated as presently non-informative rather than overinterpreted.*

19AZ$^{(4a)}$. First decoding of residual-signature recurrences into normalized BS/BL reopening priorities

The next conservative step is now to make the bridge/localization reading operational. The residual-signature words should not merely say that "something bridge-like remains." They should decode into a smallest prioritized reopening order. In particular, the present line needs one first rule saying when a stable recurrence class should push the next diagnostic pass onto the normalized bridge/localization route rather than back into the OP 1 pinning chain.

Definition H9.200 (Route-priority decoder for residual signatures)**.** *Let* $\mathcal{C}$ *be a stable recurrence class produced by the Kasiski-style scan from Definition H9.197. A* route-priority decoder *for* $\mathcal{C}$ *is any conservative rule*

$$\mathsf{Dec}(\mathcal{C}) \in \left\{ M_{\mathrm{imb}}^{\mathrm{pin}},\, M_{\mathrm{br}}^{\mathrm{surf}},\, M_{\mathrm{br}}^{\mathrm{loc}} \right\}$$

that assigns $\mathcal{C}$ *to the smallest normalized menu whose recorded carriers already account for the repeated slot/rank pattern of the sampled residuals. The decoder is called* translation-sensitive *if the repeated pattern is not stable under purely local imbalance-side rereading, but stabilizes once the residual pair is reread through the matrix/phase common-carrier surface. In that case the first prioritized reopening route attached to* $\mathcal{C}$ *is*

$$\mathrm{BS}_1 \to \mathrm{BS}_3 \to \mathrm{BL}_2,$$

with BL_4 *and, when needed,* BL_5 *appended only after the phase/common-carrier gate has been exhausted.*

Proposition H9.201 (Translation-sensitive signature classes prioritize the BS/BL route)**.** *Assume the retained total-channel baseline remains frozen through* TR_4*, and let* $\mathcal{C}$ *be a stable recurrence class of residual-signature words. Suppose moreover that* $\mathcal{C}$ *is translation-sensitive in the sense of Definition H9.200: its repeated slot/rank pattern does not settle under a purely local imbalance-side rereading, but does settle once the same sampled residuals are reread through the matrix/phase common-carrier surface. Then the first normalized reopening priority attached to* $\mathcal{C}$ *is the bridge/localization route*

$$\mathrm{TR}_4 \implies \mathrm{BS}_1 \implies \mathrm{BS}_3 \implies \mathrm{BL}_2,$$

followed, only if a still-live obstruction remains, by

$$\mathrm{BL}_2 \implies \mathrm{BL}_4,$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

and by

$$\mathrm{BL}_4 \implies \mathrm{BL}_5$$

whenever the first localized obstruction lies in the readout/kernel stratum. Equivalently, a stable translation-sensitive recurrence class is not yet evidence for a second primitive carrier; it is evidence that the next reopen final should be searched first along the normalized matrix/phase-to-localization bridge route.

Proof. By Proposition H9.198, residual-signature words may only prioritize a reopening order inside the already normalized theorem/tool layer; they may not overrule the frozen total-channel baseline or create a new carrier hypothesis. Hence one must begin from TR_4, the smallest bundled total-channel anchor already fixed by the registry from Section H9.

Now assume the stable recurrence class $\mathcal{C}$ is translation-sensitive. By definition this means that the repeated residual pattern does not become stable when reread only as a local imbalance perturbation, but does become stable once the same samples are reread through the matrix/phase common-carrier surface. Therefore the first admissible decoder target for $\mathcal{C}$ cannot be merely $M^{\mathrm{pin}}_{\mathrm{imb}}$; the smallest normalized reopening surface that can already account for the stable repeated pattern is the bridge-surface menu. In the concrete registry this is exactly the route starting at BS_1, the shared matrix-leakage gate, and continuing through BS_3, the shared phase/common-carrier surface.

Once that surface change has been made, the first honest question is whether the repeated residual pair still behaves as one masked carrier. In the normalized registry this is precisely the localization gate BL_2. If the recurrence class is already resolved there, no deeper reopening is justified. If it remains live, then the first further admissible step is BL_4, the obstruction-stratification tool. And if that stratification places the first obstruction in the readout/kernel stratum, then BL_5 is exactly the recorded compression step that rewrites the surviving burden as absence of one common shell-stable carrier package. Thus the displayed BS/BL priority route is exactly the smallest normalized decoding of a translation-sensitive stable recurrence class. □

Remark H9.202 (What the first Monte-Carlo-to-route decoding now buys)**.** *The Monte-Carlo layer now has a literal output format inside the QAM vocabulary. A stable recurrence class is no longer only a vague hint that "the bridge side may matter." The present line may now state, in compact normalized form: "freeze at* TR_4*; if the recurrence is translation-sensitive, decode first by* $\mathrm{BS}_1 \to \mathrm{BS}_3 \to \mathrm{BL}_2$*, and only then localize further by* BL_4 *or compress by* BL_5*." In that precise sense the residual-signature words have acquired a conservative but genuine routing semantics.*

19AZ$^{(4b)}$. First decoding of purely local residual-signature recurrences into normalized IP reopening priorities

The decoder should now be made symmetric on the pinning side. The residual-signature words should not only tell us when to reopen the bridge/localization route; they should also tell us when the next honest move remains entirely inside the surviving OP 1 kernel. The immediate line therefore needs one first rule saying when a stable recurrence class should stay on the local imbalance side and decode directly into the normalized IP route.

Definition H9.203 (Local-imbalance decoder for residual signatures)**.** *Let $\mathcal{C}$ be a stable recurrence class produced by the Kasiski-style scan from Definition H9.197. A route-priority decoder from Definition H9.200 is called* purely local *or* imbalance-stable *on $\mathcal{C}$ if the repeated slot/rank pattern already stabilizes under purely local rereading on the imbalance side, so that*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

rereading the same samples through the matrix/phase common-carrier surface adds no smaller admissible explanation. In that case the first prioritized reopening route attached to $\mathcal{C}$ is the normalized pinning chain

$$\mathrm{IP}_{10} \to \mathrm{IP}_9 \to \mathrm{IP}_8,$$

read beneath the frozen total-channel anchor TR_4.

Proposition H9.204 (Purely local signature classes prioritize the IP route)**.** *Assume the retained total-channel baseline remains frozen through* TR_4*, and let $\mathcal{C}$ be a stable recurrence class of residual-signature words. Suppose moreover that $\mathcal{C}$ is purely local in the sense of Definition H9.203: its repeated slot/rank pattern already stabilizes under a local imbalance-side rereading and does not require an earlier matrix/phase surface change in order to become coherent. Then the first normalized reopening priority attached to $\mathcal{C}$ is the pinning route*

$$\mathrm{TR}_4 \implies \mathrm{IP}_{10} \implies \mathrm{IP}_9 \implies \mathrm{IP}_8,$$

so that the sampled recurrence is read first as a compressed OP 1 burden rather than as a bridge/localization burden. Equivalently, a stable purely local recurrence class is not evidence that one should leave the imbalance side; it is evidence that the next reopen final should be searched first inside the surviving normalized pinning chain.

Proof. By Proposition H9.198, the recurrence class $\mathcal{C}$ may only prioritize a route already present in the normalized theorem/tool layer. Hence the total-channel baseline must remain frozen at TR_4, exactly as in the bridge/localization decoder.

Now assume $\mathcal{C}$ is purely local. By definition, its repeated slot/rank pattern already becomes coherent under a local imbalance-side rereading, and no earlier matrix/phase reinterpretation yields a smaller admissible menu. Therefore the decoder is not allowed to jump first to $M_{\mathrm{br}}^{\mathrm{surf}}$ or $M_{\mathrm{br}}^{\mathrm{loc}}$. The smallest normalized reopening surface that can already account for the stable recurrence class is instead the surviving pinning menu $M_{\mathrm{imb}}^{\mathrm{pin}}$.

Inside that menu the current live OP 1 burden has already been compressed in Proposition H9.192. The first admissible entry is therefore IP_{10}, the optimum-pinning comparison-rigidity bridge. Once that bridge is opened, one proceeds to IP_9, the retained-window-rigidity / no-second-carrier / strict-upper-gap bridge, and then to IP_8, the admissible-window uniqueness bridge. Thus the displayed tag route is exactly the smallest normalized decoding of a stable purely local recurrence class. □

Remark H9.205 (What the first Monte-Carlo-to-IP decoding now buys)**.** *The decoder is now symmetric across the two live diagnostic sides. The present line may state in compact normalized form: "freeze at* TR_4*; if the recurrence is translation-sensitive, decode first by* $\mathrm{BS}_1 \to \mathrm{BS}_3 \to \mathrm{BL}_2$*; if the recurrence is already imbalance-stable, decode instead by* $\mathrm{IP}_{10} \to \mathrm{IP}_9 \to \mathrm{IP}_8$*." In that precise sense the Monte-Carlo layer no longer only points vaguely toward "some residual side"; it now separates, conservatively and in dereferenceable tags, the first bridge reopening from the first compressed OP 1 reopening.*

19AZ$^{(4c)}$. First explicit decision table for residual-signature routing

The decoder layer should now be made directly usable. The present line no longer needs only two separate propositions saying when a stable recurrence class goes to the bridge/localization side or stays on the pinning side. It now needs one explicit decision table that turns the first observed residual-signature class into a smallest admissible action. In particular, the table should also record the conservative "hold" outcome when no stable recurrence class appears at all.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Reader cue. Read the next table as an action selector. Its purpose is to turn an observed residual-signature class into the smallest admissible reopen instruction, not to claim that every downstream route has already been exhausted.

Definition H9.206 (First explicit decision table for residual-signature routing)**.** *Assume the retained total-channel baseline remains frozen through* TR_4*, and let* $\mathcal{C}$ *be the first sampled recurrence class produced by the Kasiski-style scan from Definition H9.197. The* first explicit decision table for residual-signature routing *is the map*

$$\mathsf{RouteTab}(\mathcal{C}) \in \{\mathsf{Hold},\ \mathsf{IP\text{-}first},\ \mathsf{BS/BL\text{-}first},\ \mathsf{BL\text{-}continue},\ \mathsf{BL\text{-}kernel}\}$$

with the following normal form.

Observed residual-signature class	*Decoder output*	*First normalized route*
No stable recurrence class appears	Hold	*no reopen; keep the theorem/tool layer frozen*
Purely local / imbalance-stable recurrence	IP-first	$\mathrm{TR}_4 \Rightarrow \mathrm{IP}_{10} \Rightarrow \mathrm{IP}_9 \Rightarrow \mathrm{IP}_8$
Translation-sensitive recurrence	BS/BL-first	$\mathrm{TR}_4 \Rightarrow \mathrm{BS}_1 \Rightarrow \mathrm{BS}_3 \Rightarrow \mathrm{BL}_2$
Translation-sensitive recurrence still live after BL_2	BL-continue	*append* BL_4
First localized obstruction lies in the readout/kernel stratum	BL-kernel	*append* BL_5

The table is read top-down and always returns the smallest normalized action whose recorded carriers already account for the observed recurrence behavior.

Proposition H9.207 (The first explicit decision table is conservative and minimal on the current live classes)**.** *Under the current theorem/tool layer, every sampled residual-signature class that is presently admitted by Propositions H9.201 and H9.204 receives from Definition H9.206 a unique first admissible action. More precisely:*

(i) if no stable recurrence class appears, the decision table returns Hold*;*

(ii) if the recurrence class is purely local, it returns IP-first*;*

(iii) if the recurrence class is translation-sensitive, it returns BS/BL-first*;*

(iv) deeper localization steps BL-continue *and* BL-kernel *may be appended only after the route has already entered the BS/BL side through* BL_2*.*

Hence the table is conservative, because it never manufactures a new carrier or bypasses the frozen total-channel anchor, and minimal, because it always picks the smallest currently recorded reopening route compatible with the observed recurrence class.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. If no stable recurrence class appears at all, then Remark H9.199 already says that the theorem layer must remain untouched; this is exactly the output Hold. If a stable recurrence class is purely local, then Proposition H9.204 forces the first admissible reopening route to stay inside the pinning chain beneath TR_4, so the table must return IP-first and the route $\mathrm{TR}_4 \Rightarrow \mathrm{IP}_{10} \Rightarrow \mathrm{IP}_9 \Rightarrow \mathrm{IP}_8$. If instead the stable recurrence class is translation-sensitive, then Proposition H9.201 forces the first admissible reopening route to pass through $\mathrm{BS}_1 \Rightarrow \mathrm{BS}_3 \Rightarrow \mathrm{BL}_2$, so the table must return BS/BL-first.

The remaining two outputs are not independent first moves. By Proposition H9.201, BL_4 is appended only if the obstruction remains live after BL_2, and BL_5 is appended only when that localized obstruction lies in the readout/kernel stratum. Therefore the decision table never jumps directly to BL_4 or BL_5 without first exhausting the smaller admissible bridge/localization entry route. This proves both conservativity and minimality on the currently live recurrence classes. □

Remark H9.208 (Immediate operational reading of the decision table)**.** *The present line can now state the decoder in one sentence that is usable both by a reader and by the future machine layer: classify the first stable residual-signature class, then apply*

$$\mathsf{Hold} \ \vee \ \mathsf{IP\text{-}first} \ \vee \ \mathsf{BS/BL\text{-}first},$$

with deeper localization allowed only on the BS/BL side by appending BL-continue *and, in the readout/kernel case,* BL-kernel*. In that sense the Monte-Carlo diagnostics, the QAM tag layer, and the normalized reopening routes now meet in one explicit working table rather than only in scattered prose.*

19AZ$^{(4d)}$. First deterministic QAM decoder syntax for residual-signature routing

The decision table may now be compressed one step further. The present line no longer needs only a prose table saying which class goes to hold, IP-first, or BS/BL-first. It may now record one short deterministic decoder syntax whose words already match the normalized QAM routes. The point is not to introduce a new theory of automata; it is only to give the future machine layer and the human reader the same finite, dereferenceable instruction string.

Definition H9.209 (First deterministic QAM decoder syntax)**.** *Assume the retained total-channel anchor has been frozen through* TR_4*, and let* $\mathcal{C}$ *be the first sampled residual-signature class from the Kasiski-style recurrence scan. The* first deterministic QAM decoder syntax *is the finite rewrite alphabet*

$$\mathfrak{A}_{\mathrm{dec}} = \{\mathsf{classify}, \mathsf{hold}, \mathsf{ip}, \mathsf{bsbl}, \mathsf{continue}, \mathsf{kernel}\}$$

with the following admissible output words:

$$\begin{aligned}
&\mathsf{classify} \to \mathsf{hold},\\
&\mathsf{classify} \to \mathsf{ip},\\
&\mathsf{classify} \to \mathsf{bsbl},\\
&\mathsf{classify} \to \mathsf{bsbl} \to \mathsf{continue},\\
&\mathsf{classify} \to \mathsf{bsbl} \to \mathsf{continue} \to \mathsf{kernel}.
\end{aligned}$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

These words are interpreted by the normalized dereferencing map

$$\mathsf{Dec}_{\mathrm{QAM}} : \begin{cases} \mathsf{hold} & \mapsto \textit{keep the theorem/tool layer frozen}, \\ \mathsf{ip} & \mapsto \mathrm{TR}_4 \Rightarrow \mathrm{IP}_{10} \Rightarrow \mathrm{IP}_9 \Rightarrow \mathrm{IP}_8, \\ \mathsf{bsbl} & \mapsto \mathrm{TR}_4 \Rightarrow \mathrm{BS}_1 \Rightarrow \mathrm{BS}_3 \Rightarrow \mathrm{BL}_2, \\ \mathsf{continue} & \mapsto \textit{append } \mathrm{BL}_4, \\ \mathsf{kernel} & \mapsto \textit{append } \mathrm{BL}_5. \end{cases}$$

The syntax is called deterministic *when exactly one of the following first outputs is assigned to $\mathcal{C}$:* hold, ip, *or* bsbl*; the symbols* continue *and* kernel *may occur only after* bsbl.

Proposition H9.210 (The deterministic decoder syntax is equivalent to the current decision table)**.** *Under the current live theorem/tool layer, Definition H9.209 yields the same first admissible routing data as Definition H9.206. More precisely:*

(i) if no stable recurrence class appears, the unique admissible output word is classify $\to$ hold*;*

(ii) if the recurrence class is purely local / imbalance-stable, the unique admissible output word is classify $\to$ ip*;*

(iii) if the recurrence class is translation-sensitive, the unique admissible output word is classify $\to$ bsbl*;*

(iv) if, after entering the BS/BL side, the obstruction remains live beyond BL_2*, the decoder extends uniquely to* classify $\to$ bsbl $\to$ continue*;*

(v) if that localized obstruction lies in the readout/kernel stratum, the decoder extends further uniquely to classify $\to$ bsbl $\to$ continue $\to$ kernel.

Hence the syntax is a faithful finite normal-form compression of the decision table, with no new carriers and no loss of routing information.

Proof. The first three clauses are exactly the three primary outputs of Definition H9.206: Hold, IP-first, and BS/BL-first. Definition H9.209 merely rewrites these three outcomes as the shorter syntax outputs hold, ip, and bsbl, while the dereferencing map $\mathsf{Dec}_{\mathrm{QAM}}$ sends them back to the same normalized routes.

The remaining two clauses are likewise only compressed spellings of the already recorded BS/BL-side continuations. By Proposition H9.207, the symbol BL-continue may appear only after the bridge/localization route has already entered through BL_2, and BL-kernel may appear only after that continuation has localized the first remaining obstruction inside the readout/kernel stratum. In the deterministic syntax these two later actions are rewritten as continue and kernel, with the same ordering restriction. Thus every admissible table output produces exactly one admissible decoder word, and every admissible decoder word dereferences to exactly one table-consistent reopening route.

Therefore the syntax introduces no new mathematical burden. It only compresses the already normalized routing data into a finite QAM-readable word system. This proves equivalence to the current decision table and faithfulness of the compression. □

Remark H9.211 (Immediate machine reading of the deterministic decoder syntax)**.** *The present line may now be summarized by the short machine-readable sentences*

$$\mathsf{classify} \to \mathsf{hold}, \qquad \mathsf{classify} \to \mathsf{ip}, \qquad \mathsf{classify} \to \mathsf{bsbl} \to \mathsf{continue} \to \mathsf{kernel},$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

with the understanding that later symbols may only be appended when the earlier route has actually been entered and justified. In that sense the QAM layer now has not only tags and tables, but also a first deterministic decoder syntax that a future theorem-store, proof assistant, or simulation layer could read without reopening the entire surrounding prose.

19AZ$^{(4e)}$. First finite transition function for deterministic residual-signature routing

The deterministic decoder syntax may now be compressed once more into a tiny state transition rule. The point is again modest: not to introduce a new autonomous automaton theory, but only to expose the present routing discipline as a finite partial transition function that the future machine layer can read directly. In particular, the current line should make explicit that only three first moves are admitted from the classification state, while deeper continuation steps remain locked behind the BS/BL side.

Definition H9.212 (First finite transition function for the deterministic decoder)**.** *Let*

$$Q_{\mathrm{dec}} = \{\mathsf{classify},\ \mathsf{hold},\ \mathsf{ip},\ \mathsf{bsbl},\ \mathsf{continue},\ \mathsf{kernel}\}$$

be the current decoder-state set, and let

$$\Sigma_{\mathrm{dec}} = \{\mathsf{none},\ \mathsf{local},\ \mathsf{trans},\ \mathsf{stilllive},\ \mathsf{kernelstratum}\}$$

be the present event alphabet, where:

- none *means that no stable residual-signature recurrence class appears,*
- local *means that the sampled class is purely local / imbalance-stable,*
- trans *means that the sampled class is translation-sensitive,*
- $\mathsf{stilllive}$ *means that after entering the BS/BL side the obstruction remains live beyond* BL_2*,*
- $\mathsf{kernelstratum}$ *means that the first localized remaining obstruction lies in the read-out/kernel stratum.*

The first finite transition function for deterministic residual-signature routing *is the partial map*

$$\delta_{\mathrm{dec}} : Q_{\mathrm{dec}} \times \Sigma_{\mathrm{dec}} \dashrightarrow Q_{\mathrm{dec}}$$

given by the only admissible transitions

$$\begin{aligned}
\delta_{\mathrm{dec}}(\mathsf{classify}, \mathsf{none}) &= \mathsf{hold},\\
\delta_{\mathrm{dec}}(\mathsf{classify}, \mathsf{local}) &= \mathsf{ip},\\
\delta_{\mathrm{dec}}(\mathsf{classify}, \mathsf{trans}) &= \mathsf{bsbl},\\
\delta_{\mathrm{dec}}(\mathsf{bsbl}, \mathsf{stilllive}) &= \mathsf{continue},\\
\delta_{\mathrm{dec}}(\mathsf{continue}, \mathsf{kernelstratum}) &= \mathsf{kernel}.
\end{aligned}$$

All other transitions are undefined at the present stage. The state outputs are dereferenced through the same normalized map $\mathsf{Dec}_{\mathrm{QAM}}$ *from Definition H9.209.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proposition H9.213 (The finite transition function is equivalent to the deterministic decoder syntax and remains minimal on the current live events)**.** *Under the current theorem/tool layer, the partial transition function from Definition H9.212 carries exactly the same routing information as the deterministic decoder syntax from Definition H9.209. More precisely:*

(i) the three first moves from classify *are exactly* hold, ip, *and* bsbl, *corresponding respectively to the events* none, local, *and* trans*;*

(ii) the only admissible continuation on the bridge/localization side is $\mathsf{bsbl} \xrightarrow{\mathsf{stilllive}} \mathsf{continue}$*;*

(iii) the only admissible further refinement is $\mathsf{continue} \xrightarrow{\mathsf{kernelstratum}} \mathsf{kernel}$*;*

(iv) every undefined transition would either skip a smaller admissible reopening route or manufacture a carrier/event type not yet recorded in the present line.

Hence δ_{dec} *is a faithful finite-state compression of the current routing discipline and is minimal on the currently live diagnostic events.*

Proof. The first three transitions are exactly the three initial output words from Proposition H9.210: classify $\to$ hold, classify $\to$ ip, and classify $\to$ bsbl. The transition function merely rewrites these words as event-labeled state moves. The event names none, local, and trans are exactly the presently live coarse classes isolated by the decision table and the two decoder propositions.

Once the route has entered bsbl, the deterministic syntax allows only the optional later symbols continue and kernel. By Definition H9.209, continue may appear only after bsbl, and by Proposition H9.207 it is justified precisely when the obstruction remains live beyond BL_2. This is the transition $\delta_{\mathrm{dec}}(\mathsf{bsbl}, \mathsf{stilllive}) = \mathsf{continue}$. Likewise, kernel may occur only after that continuation has localized the first remaining obstruction inside the readout/kernel stratum, giving $\delta_{\mathrm{dec}}(\mathsf{continue}, \mathsf{kernelstratum}) = \mathsf{kernel}$.

No further transitions are admissible on the current line. Any direct jump from classify to continue or kernel would bypass the smaller bridge/localization entry route, while any jump from ip to a later BS/BL state would conflate two presently separated reopening disciplines. And any additional event symbol would exceed the currently normalized recurrence classes. Therefore the partial transition function is equivalent to the deterministic decoder syntax, conservative with respect to the recorded carriers, and minimal on the currently live events. □

Remark H9.214 (Immediate automaton reading of the current decoder)**.** *The present line can now be read as a tiny partial automaton:*

$$\mathsf{classify} \xrightarrow{\mathsf{none}} \mathsf{hold},$$
$$\mathsf{classify} \xrightarrow{\mathsf{local}} \mathsf{ip},$$
$$\mathsf{classify} \xrightarrow{\mathsf{trans}} \mathsf{bsbl} \xrightarrow{\mathsf{stilllive}} \mathsf{continue} \xrightarrow{\mathsf{kernelstratum}} \mathsf{kernel}.$$

Through the dereferencing map $\mathsf{Dec}_{\mathrm{QAM}}$*, this automaton is still nothing more than the already normalized theorem/tool layer. But it now has a form that a later theorem store, proof assistant, or Monte-Carlo controller could read as a deterministic transition discipline instead of as prose plus scattered tags.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

19AZ$^{(4f)}$. First observer/event normal form for deterministic residual-signature words

The tiny decoder automaton from the previous subsection still presupposes that the incoming event has already been named as one of the symbols none, local, trans, stilllive, or kernelstratum. The next conservative step is therefore to normalize not the output route once more, but the observer-side input itself. The present line does not need a large language for that purpose. It only needs a short finite event-word normal form recording how much of the bridge/localization continuation has actually become visible at readout.

Definition H9.215 (First observer/event normal form for deterministic residual-signature words)**.** *Let*

$$\mathcal{W}_{\mathrm{dec}}^{\mathrm{nf}} = \{\mathsf{none},\ \mathsf{local},\ \mathsf{trans},\ \mathsf{trans\ stilllive},\ \mathsf{trans\ stilllive\ kernelstratum}\}$$

be the first observer/event normal-form language *for deterministic residual-signature routing. The associated partial extended transition map is defined recursively by*

$$\delta_{\mathrm{dec}}^{*}(q,\varepsilon) = q, \qquad \delta_{\mathrm{dec}}^{*}(q,aw) = \delta_{\mathrm{dec}}^{*}\big(\delta_{\mathrm{dec}}(q,a),w\big)$$

whenever the right-hand side is defined, where $a \in \Sigma_{\mathrm{dec}}$ and w is a finite word in the same alphabet.

The observer/event normal form map

$$\mathsf{NF}_{\mathrm{obs}} : \textit{currently normalized residual-signature cases} \longrightarrow \mathcal{W}_{\mathrm{dec}}^{\mathrm{nf}}$$

assigns the unique normal-form word as follows:

(i) no stable recurrence class is observed $\mapsto$ none*,*

(ii) a purely local / imbalance-stable class is observed $\mapsto$ local*,*

(iii) a translation-sensitive class is observed but no remaining obstruction survives beyond BL_2 $\mapsto$ trans*,*

(iv) a translation-sensitive class is observed and the obstruction remains live beyond BL_2 $\mapsto$ trans stilllive*,*

(v) the previous case holds and the first localized remaining obstruction lies in the readout/kernel stratum $\mapsto$ trans stilllive kernelstratum*.*

Proposition H9.216 (The observer/event normal forms are exactly the currently admissible decoder words and reproduce the normalized reopening routes)**.** *Under the current theorem/tool layer, the language $\mathcal{W}_{\mathrm{dec}}^{\mathrm{nf}}$ from Definition H9.215 is exactly the set of currently admissible observer-side event words for the deterministic decoder. More precisely:*

(i) every word in $\mathcal{W}_{\mathrm{dec}}^{\mathrm{nf}}$ is admitted by the partial automaton δ_{dec} when started at classify*,*

(ii) the terminal states are respectively

$$\delta_{\mathrm{dec}}^{*}(\mathsf{classify},\mathsf{none}) = \mathsf{hold}, \qquad \delta_{\mathrm{dec}}^{*}(\mathsf{classify},\mathsf{local}) = \mathsf{ip},$$

$$\delta_{\mathrm{dec}}^{*}(\mathsf{classify},\mathsf{trans}) = \mathsf{bsbl}, \qquad \delta_{\mathrm{dec}}^{*}(\mathsf{classify},\mathsf{trans\ stilllive}) = \mathsf{continue},$$

$$\delta_{\mathrm{dec}}^{*}(\mathsf{classify},\mathsf{trans\ stilllive\ kernelstratum}) = \mathsf{kernel},$$

and these states dereference exactly to the normalized routes already fixed in the QAM tool layer;

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(iii) no other finite event word is currently admissible without either violating the transition discipline of δ_{dec} or introducing an observer-side distinction not yet normalized in the present line.

Hence the current Monte-Carlo/QAM interface may now be read not only as a state machine, but as a finite normal-form language of observer-side diagnostic words.

Proof. The five displayed words are admitted directly by the transition function from Definition H9.212. The one-letter words none, local, and trans are exactly the three first-step events recorded there. Once the route has entered bsbl, the only currently permitted continuation event is stilllive, and once the route has entered continue, the only currently permitted further refinement is kernelstratum. Therefore the longer words trans stilllive and trans stilllive kernelstratum are admitted as well.

The recursive definition of $\delta^{*}_{\mathrm{dec}}$ then gives the terminal states by straightforward evaluation. For none and local, only one transition is applied, yielding hold and ip. For trans, the first step yields bsbl. Appending stilllive moves to continue, and appending kernelstratum moves from there to kernel. By Definition H9.209, each terminal state dereferences to the already normalized reopening route: hold means reopen nothing, ip means the chain $\mathrm{TR}_4 \Rightarrow \mathrm{IP}_{10} \Rightarrow \mathrm{IP}_9 \Rightarrow \mathrm{IP}_8$, bsbl means the chain $\mathrm{TR}_4 \Rightarrow \mathrm{BS}_1 \Rightarrow \mathrm{BS}_3 \Rightarrow \mathrm{BL}_2$, continue appends BL_4, and kernel appends BL_5.

It remains to exclude any additional finite event word. A word beginning with anything other than none, local, or trans is impossible because δ_{dec} is undefined at classify on all other events. A word beginning with none or local cannot be extended further without leaving the currently normalized routing discipline. And a word beginning with trans may only continue with stilllive, then optionally with kernelstratum, because no further transitions are currently recorded. Thus every admissible observer-side case has exactly one normal-form word and no other finite word is currently allowed. □

Remark H9.217 (Immediate machine reading of the observer/event normal form)**.** *The current live interface may now be summarized by the finite observer-language*

none, local, trans, trans stilllive, trans stilllive kernelstratum.

This is the first point at which the tool layer, the QAM tags, the Monte-Carlo recurrence classes, the deterministic decoder syntax, and the partial transition automaton all meet in one compact object: a finite word language on the observer side whose outputs are already normalized reopening routes on the theorem side.

19AZ$^{(4g)}$. First observer-word evaluation map into deterministic decoder outputs

The observer/event normal forms now provide a finite input language for the deterministic decoder. The next conservative compression step is to package the whole passage from normalized observer word to normalized theorem-side reopening output into one explicit evaluation map. This adds no new event class and no new route. It merely composes the already stabilized normal-form language, transition automaton, and QAM dereferencing rule into one finite readout operator.

Definition H9.218 (First observer-word evaluation map into deterministic decoder outputs)**.** *Let*

$$\mathcal{W}^{\mathrm{nf}}_{\mathrm{dec}} = \{\text{none}, \text{local}, \text{trans}, \text{trans stilllive}, \text{trans stilllive kernelstratum}\}$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

be the observer/event normal-form language from Definition H9.215. Define the observer-word evaluation map

$$\mathsf{Eval}_{\mathrm{obs}} : \mathcal{W}^{\mathrm{nf}}_{\mathrm{dec}} \longrightarrow Q_{\mathrm{dec}} \times \mathcal{R}^{\mathrm{out}}_{\mathrm{tool}}$$

by

$$\mathsf{Eval}_{\mathrm{obs}}(w) = \Big(\delta^{*}_{\mathrm{dec}}(\mathsf{classify}, w), \mathsf{Dec}_{\mathrm{QAM}}\big(\delta^{*}_{\mathrm{dec}}(\mathsf{classify}, w)\big)\Big),$$

where $Q_{\mathrm{dec}} = \{\mathsf{hold}, \mathsf{ip}, \mathsf{bsbl}, \mathsf{continue}, \mathsf{kernel}\}$ *is the current decoder-state set and* $\mathcal{R}^{\mathrm{out}}_{\mathrm{tool}}$ *denotes the finite set of currently normalized reopening outputs:*

$$\mathcal{R}^{\mathrm{out}}_{\mathrm{tool}} = \left\{ \begin{array}{l} \varnothing, \\ [\mathrm{TR}_4, \mathrm{IP}_{10}, \mathrm{IP}_9, \mathrm{IP}_8], \\ [\mathrm{TR}_4, \mathrm{BS}_1, \mathrm{BS}_3, \mathrm{BL}_2], \\ [\mathrm{TR}_4, \mathrm{BS}_1, \mathrm{BS}_3, \mathrm{BL}_2, \mathrm{BL}_4], \\ [\mathrm{TR}_4, \mathrm{BS}_1, \mathrm{BS}_3, \mathrm{BL}_2, \mathrm{BL}_4, \mathrm{BL}_5] \end{array} \right\}.$$

Equivalently,

$$\begin{aligned} \mathsf{Eval}_{\mathrm{obs}}(\mathsf{none}) &= (\mathsf{hold}, \varnothing), \\ \mathsf{Eval}_{\mathrm{obs}}(\mathsf{local}) &= (\mathsf{ip}, [\mathrm{TR}_4, \mathrm{IP}_{10}, \mathrm{IP}_9, \mathrm{IP}_8]), \\ \mathsf{Eval}_{\mathrm{obs}}(\mathsf{trans}) &= (\mathsf{bsbl}, [\mathrm{TR}_4, \mathrm{BS}_1, \mathrm{BS}_3, \mathrm{BL}_2]), \\ \mathsf{Eval}_{\mathrm{obs}}(\mathsf{trans\,stilllive}) &= (\mathsf{continue}, [\mathrm{TR}_4, \mathrm{BS}_1, \mathrm{BS}_3, \mathrm{BL}_2, \mathrm{BL}_4]), \\ \mathsf{Eval}_{\mathrm{obs}}(\mathsf{trans\,stilllive\,kernelstratum}) &= (\mathsf{kernel}, [\mathrm{TR}_4, \mathrm{BS}_1, \mathrm{BS}_3, \mathrm{BL}_2, \mathrm{BL}_4, \mathrm{BL}_5]). \end{aligned}$$

Proposition H9.219 (The observer-word evaluation map is total on the current normal–form language and reproduces the unique normalized reopening outputs)**.** *Under the current theorem/tool layer, the map* $\mathsf{Eval}_{\mathrm{obs}}$ *from Definition H9.218 is total on* $\mathcal{W}^{\mathrm{nf}}_{\mathrm{dec}}$*. More precisely:*

(i) every word in $\mathcal{W}^{\mathrm{nf}}_{\mathrm{dec}}$ *has exactly one decoder terminal state and exactly one currently normalized reopening output under* $\mathsf{Eval}_{\mathrm{obs}}$*,*

(ii) the first component of $\mathsf{Eval}_{\mathrm{obs}}(w)$ *is the unique terminal state reached by the partial automaton* $\delta^{*}_{\mathrm{dec}}$*,*

(iii) the second component is exactly the smallest presently admissible theorem-side reopening route compatible with that terminal state,

(iv) no currently normalized observer word evaluates to two different reopening outputs.

Hence the present live interface may now be read as one deterministic finite composite map

$$\mathsf{NF}_{\mathrm{obs}} \longrightarrow \mathsf{Eval}_{\mathrm{obs}} \longrightarrow \textit{normalized reopening output},$$

rather than as a chain of separately cited decoder pieces.

Proof. By Proposition H9.216, every word in $\mathcal{W}^{\mathrm{nf}}_{\mathrm{dec}}$ is admitted by the extended transition map $\delta^{*}_{\mathrm{dec}}$ when started at $\mathsf{classify}$, and its terminal state is uniquely determined. Thus the first component of $\mathsf{Eval}_{\mathrm{obs}}$ is total and single-valued on the whole normal-form language.

The second component is obtained by applying the deterministic QAM dereferencing map $\mathsf{Dec}_{\mathrm{QAM}}$ to that unique terminal state. By Definition H9.209, each currently live terminal state carries exactly one normalized reopening output: hold gives $\varnothing$, ip gives the fixed IP route $[\mathrm{TR}_4, \mathrm{IP}_{10}, \mathrm{IP}_9, \mathrm{IP}_8]$, bsbl gives the fixed bridge/localization entry route $[\mathrm{TR}_4, \mathrm{BS}_1, \mathrm{BS}_3, \mathrm{BL}_2]$,

© 2026 Marc Brendecke. Preprint — not peer reviewed.

continue appends BL_4, and kernel appends BL_5. Therefore the second component is also total and single-valued.

Minimality follows from the construction of the current decoder. The state hold reopens nothing, ip is already the smallest admissible local route, and the bridge/localization branch enters first through bsbl before optional refinement to continue and kernel. Any alternative output would either reopen more than the current state requires or bypass an earlier normalized stage. Hence no current observer word can evaluate to two different normalized reopening outputs. □

Remark H9.220 (Immediate operational reading of $\mathsf{Eval}_{\mathrm{obs}}$)**.** *The current finite live interface may now be cited in one line:*

$$\begin{aligned} &\textit{observer-side recurrence class} \xrightarrow{\mathsf{NF}_{\mathrm{obs}}} \textit{normal-form word} \\ &\qquad\qquad \xrightarrow{\mathsf{Eval}_{\mathrm{obs}}} (\textit{decoder state}, \textit{normalized reopening output}). \end{aligned}$$

At this point, the Monte-Carlo diagnostics, the QAM tag layer, the finite decoder automaton, and the theorem-side reopening discipline have become one explicit finite evaluation pipeline. This is the first place in the present line where a later controller could read the whole diagnostic-to-route passage through a single total map on a finite language.

19AZ$^{(4h)}$. First compact output-normalform tokens for deterministic decoder results

The observer-word evaluation map now packages every currently admissible observer-side normal form into one of five deterministic theorem-side outputs. The next conservative compression step is to replace those explicit route lists by a finite family of compact output tokens. This adds no new state, no new route, and no new observer word. It merely supplies a shorter QAM-side output carrier for the five already stabilized decoder results.

Definition H9.221 (First compact output-normalform tokens for deterministic decoder results)**.** *Define the finite set of* output-normalform tokens

$$\mathcal{T}_{\mathrm{out}}^{\mathrm{nf}} = \{\mathsf{OH}, \mathsf{OI}, \mathsf{OB}, \mathsf{OC}, \mathsf{OK}\},$$

where the tokens are read as

$$\begin{aligned} \mathsf{OH} &\leftrightarrow (\mathsf{hold}, \varnothing), \\ \mathsf{OI} &\leftrightarrow (\mathsf{ip}, [\mathrm{TR}_4, \mathrm{IP}_{10}, \mathrm{IP}_9, \mathrm{IP}_8]), \\ \mathsf{OB} &\leftrightarrow (\mathsf{bsbl}, [\mathrm{TR}_4, \mathrm{BS}_1, \mathrm{BS}_3, \mathrm{BL}_2]), \\ \mathsf{OC} &\leftrightarrow (\mathsf{continue}, [\mathrm{TR}_4, \mathrm{BS}_1, \mathrm{BS}_3, \mathrm{BL}_2, \mathrm{BL}_4]), \\ \mathsf{OK} &\leftrightarrow (\mathsf{kernel}, [\mathrm{TR}_4, \mathrm{BS}_1, \mathrm{BS}_3, \mathrm{BL}_2, \mathrm{BL}_4, \mathrm{BL}_5]). \end{aligned}$$

Equivalently, define the output-tokenization map

$$\mathsf{Tok}_{\mathrm{out}} : Q_{\mathrm{dec}} \times \mathcal{R}_{\mathrm{tool}}^{\mathrm{out}} \longrightarrow \mathcal{T}_{\mathrm{out}}^{\mathrm{nf}}$$

by sending the five currently normalized evaluation outputs from Definition H9.218 to the corresponding tokens above. Composing with $\mathsf{Eval}_{\mathrm{obs}}$ *gives the compact output map*

$$\mathsf{Out}_{\mathrm{obs}}^{\mathrm{tok}} := \mathsf{Tok}_{\mathrm{out}} \circ \mathsf{Eval}_{\mathrm{obs}} : \mathcal{W}_{\mathrm{dec}}^{\mathrm{nf}} \longrightarrow \mathcal{T}_{\mathrm{out}}^{\mathrm{nf}}.$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Thus

$$\begin{aligned}
\mathsf{Out}^{\mathrm{tok}}_{\mathrm{obs}}(\mathsf{none}) &= \mathsf{OH},\\
\mathsf{Out}^{\mathrm{tok}}_{\mathrm{obs}}(\mathsf{local}) &= \mathsf{OI},\\
\mathsf{Out}^{\mathrm{tok}}_{\mathrm{obs}}(\mathsf{trans}) &= \mathsf{OB},\\
\mathsf{Out}^{\mathrm{tok}}_{\mathrm{obs}}(\mathsf{trans\,stilllive}) &= \mathsf{OC},\\
\mathsf{Out}^{\mathrm{tok}}_{\mathrm{obs}}(\mathsf{trans\,stilllive\,kernelstratum}) &= \mathsf{OK}.
\end{aligned}$$

Proposition H9.222 (The output-normalform tokens are faithful and complete for the current deterministic decoder results)**.** *Under the current live theorem/tool layer, the token map* $\mathsf{Tok}_{\mathrm{out}}$ *is faithful and complete on the presently normalized decoder outputs. More precisely:*

(i) every value of $\mathsf{Eval}_{\mathrm{obs}}$ *is sent to exactly one token in* $\mathcal{T}^{\mathrm{nf}}_{\mathrm{out}}$*,*

(ii) different current decoder/output pairs receive different tokens,

(iii) the token $\mathsf{Out}^{\mathrm{tok}}_{\mathrm{obs}}(w)$ *determines both the terminal decoder state and the unique normalized reopening route attached to that state,*

(iv) no information from the current finite live interface is lost when passing from explicit output pairs to output-normalform tokens.

Hence the present diagnostic-to-route pipeline may now also be read in the compact form

$$\textit{observer class} \longrightarrow \textit{normal-form word} \longrightarrow \textit{output-normalform token.}$$

Proof. By Proposition H9.219, the map $\mathsf{Eval}_{\mathrm{obs}}$ takes values in exactly five currently normalized decoder/output pairs. Definition H9.221 assigns one token to each of those five pairs and to no others, so every current evaluation value receives exactly one token.

Faithfulness is immediate from the construction: $\mathsf{OH}, \mathsf{OI}, \mathsf{OB}, \mathsf{OC}, \mathsf{OK}$ are pairwise distinct symbols, each tied to one different decoder/output pair. Thus two different current evaluation outputs cannot collapse to the same token. Conversely, each token has exactly one prescribed decoder state and exactly one prescribed normalized reopening route, so the token determines the full current output pair.

Therefore the passage from explicit pairs to compact tokens is lossless on the present finite interface. It is merely a normalization of notation, not a quotient by any nontrivial equivalence. The compact map $\mathsf{Out}^{\mathrm{tok}}_{\mathrm{obs}}$ preserves exactly the same current routing content as $\mathsf{Eval}_{\mathrm{obs}}$, only in shorter QAM-readable form. □

Remark H9.223 (Why the compact output tokens matter)**.** *The live interface can now be written in its shortest current form:*

$$\textit{observer-side recurrence class} \xrightarrow{\mathsf{NF}_{\mathrm{obs}}} \textit{normal-form word} \xrightarrow{\mathsf{Out}^{\mathrm{tok}}_{\mathrm{obs}}} \{\mathsf{OH}, \mathsf{OI}, \mathsf{OB}, \mathsf{OC}, \mathsf{OK}\}.$$

At this point the theorem-side reopening discipline already has a compact output alphabet. A later controller or QAM overlay no longer needs to carry the full explicit route list every time; it may cite the token first and dereference the corresponding route only when the detailed theorem-side reopening path is actually needed.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

19AZ$^{(5a)}$. First signature-injectivity normal form for the live OP 1 kernel

The normalized routing layer has now done its job on the OP 1 side: it has compressed the live reopening path to

$$\mathrm{TR}_4 \Longrightarrow \mathrm{IP}_{10} \Longrightarrow \mathrm{IP}_9 \Longrightarrow \mathrm{IP}_8.$$

The next honest mathematical move is therefore not another routing refinement but a sharper obstruction normal form for the first live carrier in that chain. The remaining burden should now be stated as a local injectivity problem for the retained pinning signature itself: can two genuinely distinct admissible primitive photon-first-zero loops occupy the same retained post-frame corridor class while preserving the same total-channel profile, the same retained optimum location, and the same local comparison/defect readout?

> **Reader cue.** Read the next block as an obstruction normal form for the live OP 1 kernel. Its purpose is to rewrite the remaining burden as exclusion of a second admissible primitive loop with the same retained pinning signature, not to claim that this exclusion has already been proved.

Definition H9.224 (Retained pinning signature on the OP 1 corridor)**.** *Assume the same-corridor reversal certificate of Definition H9.149. For every admissible primitive photon-first-zero closed-loop realization L on the retained post-frame corridor, define the* retained pinning signature

$$\mathsf{Sig}_{\mathrm{pin}}(L)$$

to be the complete retained comparison packet carried by the present line on the intended admissible windows, namely: the retained total-channel profile, the retained optimum location, and the standing local comparison/defect data attached to L in the compressed machine-shadow reading. Thus two loops have the same retained pinning signature exactly when they are indistinguishable to the current OP 1 comparison line at the level that feeds Definition H9.169.

Definition H9.225 (Duplicate-signature witness for the live OP 1 kernel)**.** *Assume Definition H9.224. A* duplicate-signature witness *for the live OP 1 kernel is a pair (L, L') of admissible primitive photon-first-zero closed-loop realizations on the retained post-frame corridor such that:*

(i) *L and L' are genuinely distinct up to the standing admissible corridor equivalence,*

(ii) *both preserve the same retained total-channel profile on the intended admissible windows,*

(iii) $\mathsf{Sig}_{\mathrm{pin}}(L) = \mathsf{Sig}_{\mathrm{pin}}(L')$.

Equivalently, a duplicate-signature witness is a second primitive loop placement that the current comparison line cannot distinguish from the intended retained loop even though it remains genuinely distinct inside the retained corridor class.

Proposition H9.226 (The live OP 1 burden is equivalent to exclusion of duplicate-signature witnesses)**.** *Assume the same-corridor reversal certificate of Definition H9.149. Then the following are equivalent on the intended admissible windows:*

(i) *the optimum-pinning comparison-rigidity certificate of Definition H9.169 holds;*

(ii) *no duplicate-signature witness exists in the sense of Definition H9.225.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Consequently, together with Proposition H9.166 and the standing no-second-carrier / strict-upper-gap exclusions already built into the current line, exclusion of duplicate-signature witnesses implies

$$\Delta_{\mathrm{CL}} = 0.$$

Equivalently, the present residual OP 1 burden may now be read exactly as the injectivity of the retained pinning signature on the admissible primitive loop class carried by the retained post-frame corridor.

Proof. Definition H9.169 says that no genuinely distinct admissible primitive loop on the retained post-frame corridor can preserve the retained total-channel profile, the retained optimum location, and the standing local comparison/defect data while remaining distinct up to the admissible corridor equivalence. But this is exactly the negation of a duplicate-signature witness from Definition H9.225, since the retained pinning signature packages precisely those same preserved data. Thus *(i)* and *(ii)* are equivalent.

Once this injectivity statement is available, Proposition H9.170 yields the retained-window rigidity certificate. Proposition H9.166 then upgrades that rigidity, under the standing no-second-carrier and strict-upper-gap exclusions, to admissible-window uniqueness and hence to $\Delta_{\mathrm{CL}} = 0$. □

Remark H9.227 (What the new normal form buys for the next OP 1 step)**.** *This is a real sharpening even though it is not yet a closure theorem. Before the present step, the residual burden was phrased as "prove optimum-pinning comparison rigidity." After the present step, the live target is more concrete: show that the retained comparison packet*

$$\mathsf{Sig}_{\mathrm{pin}}$$

is injective on admissible primitive first-zero loops in the retained post-frame corridor class. In practical terms, the next OP 1 attack should therefore look for a contradiction from a hypothetical duplicate-signature witness, not for a new free continuum-limit parameter. In the normalized QAM reading, this says that the first live carrier IP_{10} *should now be treated as a signature-injectivity problem for the retained pinning packet.*

19AZ$^{(5b)}$. First contradiction channel split for a duplicate-signature witness

The normal form of 19AZ$^{(5a)}$ says that the live OP 1 burden is equivalent to the exclusion of duplicate-signature witnesses. The next honest move is therefore to ask what such a witness would have to do inside the present line. Because the retained pinning signature already fixes the retained total-channel profile, the retained optimum location, and the local comparison/defect packet, any residual distinction between two loops carrying the same signature has only two places left to live: either as a second primitive occupancy of the same retained corridor slot, or as a second admissible completion above that same slot. This is exactly the old no-second-carrier / strict-upper-gap dichotomy, but now pulled back to the new signature normal form.

> **Reader cue.** Read the next block as the first contradiction routing split for the live OP 1 kernel. Its purpose is to show that a hypothetical duplicate-signature witness can only survive by feeding one of the two exclusion gates already built into the current line: duplicate primitive occupancy or duplicate admissible completion.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition H9.228 (Occupancy-side and completion-side duplicate-signature channels)**.** *Assume the same-corridor reversal certificate of Definition H9.149 and let (L, L') be a duplicate-signature witness in the sense of Definition H9.225. We say that the witness is of* occupancy-side type *if the residual distinction between L and L' already appears as a second primitive occupancy of the retained post-frame corridor while the retained total-channel profile is kept fixed. We say that the witness is of* completion-side type *if, after the common retained corridor slot and retained comparison packet are fixed, the residual distinction can only survive as a second admissible completion route above that same retained corridor while preserving the same retained status surface.*

Proposition H9.229 (Every duplicate-signature witness feeds a standing OP 1 exclusion gate)**.** *Assume the same-corridor reversal certificate of Definition H9.149. Let (L, L') be a duplicate-signature witness in the sense of Definition H9.225. Then at least one of the following holds on the intended admissible windows:*

(i) (L, L') is of occupancy-side type in the sense of Definition H9.228;

(ii) (L, L') is of completion-side type in the sense of Definition H9.228.

Consequently, every hypothetical failure of signature-injectivity for the live OP 1 kernel already routes into one of the two standing exclusion channels behind Proposition H9.166: no-second-carrier on the retained corridor, or strict-upper-gap on admissible completion above that corridor.

Proof. By Definition H9.225, the loops L and L' are genuinely distinct up to the standing admissible corridor equivalence, yet they preserve the same retained pinning signature. In particular, they preserve the same retained total-channel profile, the same retained optimum location, and the same local comparison/defect data.

Therefore the current line has already fixed everything that belongs to the retained comparison packet. If the residual distinction between L and L' is visible already at the level of primitive occupancy of the retained post-frame corridor, then the witness is of occupancy-side type by definition. Otherwise the common retained corridor slot and retained comparison packet do not separate the loops, so the remaining distinction can only survive as a second admissible completion above that same retained corridor while preserving the same retained status surface; this is exactly the completion-side case.

Hence every duplicate-signature witness must feed at least one of the two channels. The final sentence is then only a translation into the standing OP 1 language: occupancy-side witnesses are precisely the kind of residual duplication targeted by the no-second-carrier discipline, while completion-side witnesses are precisely the kind of residual duplication targeted by the strict-upper-gap exclusion in Proposition H9.166. □

Remark H9.230 (What this contradiction split buys for the next OP 1 attack)**.** *This still does not close OP 1, but it tells us where a closure attempt must now bite. A hypothetical duplicate-signature witness no longer floats as an abstract rival loop. It is forced into one of two already named contradiction channels. In practical terms, the next OP 1 step should try to show that the retained comparison packet cannot support either occupancy-side duplication on the retained corridor or completion-side duplication above that corridor. In the normalized QAM reading, this means that the live carrier* IP_{10} *has now been resolved into a two-channel contradiction search whose outputs are already aligned with the standing no-second-carrier / strict-upper-gap gates feeding* IP_9*.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

19AZ$^{(5c)}$. Occupancy-side duplicate-signature witnesses collapse under the retained no-second-carrier discipline

After the channel split of 19AZ$^{(5b)}$, the cleanest next question is whether both contradiction channels are still genuinely live. On the present line, the occupancy-side channel sits closest to the already retained theorematic infrastructure: if a duplicate-signature witness already survives as a second primitive occupancy of the same retained corridor while the retained total-channel profile stays fixed, then it is no longer merely a new OP 1 curiosity. It is exactly the sort of duplication that the retained no-second-carrier discipline was built to forbid on the preserved total-channel side.

> **Reader cue.** Read the next block as the first actual collapse theorem inside the new OP 1 contradiction split. Its purpose is to show that occupancy-side duplicate-signature witnesses are already excluded by the standing retained no-second-carrier discipline, so that the live burden cannot honestly remain two-channel after this point.

Proposition H9.231 (Occupancy-side duplicate-signature witnesses are excluded on the retained OP 1 corridor)**.** *Assume the same-corridor reversal certificate of Definition H9.149. Assume moreover that the retained no-second-carrier discipline continues to hold on the intended admissible windows at the retained total-channel level, in the compressed sense already available through Theorem 31.8. Then no duplicate-signature witness of occupancy-side type exists in the sense of Definitions H9.225 and H9.228.*

Proof. Suppose for contradiction that (L, L') is a duplicate-signature witness of occupancy-side type. By Definition H9.225, the loops L and L' are genuinely distinct up to the standing admissible corridor equivalence, yet they preserve the same retained pinning signature. In particular, they preserve the same retained total-channel profile on the intended admissible windows.

By Definition H9.228, the residual distinction already survives as a second primitive occupancy of the retained post-frame corridor while that retained total-channel profile is kept fixed. But this is precisely the kind of extra primitive retained-carrier occupancy excluded by the standing no-second-carrier discipline at retained total-channel level. Hence the assumed witness contradicts the retained no-second-carrier theorematic layer. Therefore no occupancy-side duplicate-signature witness can exist. □

Corollary H9.232 (The live OP 1 duplicate-signature burden reduces to the completion-side channel)**.** *Under the hypotheses of Proposition H9.231, the following are equivalent on the intended admissible windows:*

(i) a duplicate-signature witness for the live OP 1 kernel exists;

(ii) a duplicate-signature witness of completion-side type exists.

Consequently, after the present step the live OP 1 kernel is no longer honestly read as a two-channel contradiction search. The occupancy-side branch is already discharged by the retained no-second-carrier discipline, so the surviving burden is the exclusion of a second admissible completion above the retained corridor that preserves the same retained pinning signature.

Proof. If a duplicate-signature witness exists, Proposition H9.229 shows that it must be either of occupancy-side type or of completion-side type. Proposition H9.231 eliminates the occupancy-side case. Hence every surviving duplicate-signature witness must be completion-side. The converse is immediate, since every completion-side witness is in particular a duplicate-signature witness. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Remark H9.233 (What this first collapse buys for the next OP 1 step)**.** *This is the first point in the new post-release OP 1 line where one of the two contradiction channels is not merely named but actually discharged. The live burden is therefore smaller than it looked in 19AZ$^{(5b)}$: the retained comparison packet is already rigid enough against duplicate* occupancy *once the inherited no-second-carrier discipline is invoked. What remains is subtler and now sharply localized. One must show that the same retained pinning signature cannot support a second* admissible completion *above the already fixed retained corridor. In the normalized QAM reading, the live carrier* IP_{10} *has therefore collapsed from a two-channel contradiction search to a completion-side exclusion problem feeding directly into the strict-upper-gap gate behind* IP_9*.*

Reader cue. Read the next block as the second actual collapse theorem in the new OP 1 contradiction split. Its purpose is to show that completion-side duplicate-signature witnesses are already excluded by the standing retained strict-upper-gap discipline, so that no duplicate-signature witness survives the present OP 1 kernel once both standing exclusion gates are read in the new signature normal form.

19AZ$^{(5d)}$. Completion-side duplicate-signature witnesses collapse under the retained strict-upper-gap discipline

After 19AZ$^{(5c)}$, the live duplicate-signature burden has only one honest place left to survive: as a second admissible completion above the already fixed retained corridor while the retained pinning signature stays unchanged. But this is exactly the sort of residual duplication that the retained strict-upper-gap language was already designed to exclude. Once the retained corridor slot and retained comparison packet are fixed, a genuinely second admissible completion preserving the same retained status surface is no longer a new geometric freedom; it is a forbidden duplicate completion above the same retained corridor.

Proposition H9.234 (Completion-side duplicate-signature witnesses are excluded on the retained OP 1 corridor)**.** *Assume the same-corridor reversal certificate of Definition H9.149. Assume moreover that the retained strict-upper-gap discipline continues to hold on the intended admissible windows in the compressed sense already used by Proposition H9.166. Then no duplicate-signature witness of completion-side type exists in the sense of Definitions H9.225 and H9.228.*

Proof. Suppose for contradiction that (L, L') is a duplicate-signature witness of completion-side type. By Definition H9.225, the loops L and L' are genuinely distinct up to the standing admissible corridor equivalence, yet they preserve the same retained pinning signature. In particular, they preserve the same retained total-channel profile, the same retained optimum location, and the same local comparison/defect packet on the intended admissible windows.

By Definition H9.228, once the common retained corridor slot and retained comparison packet are fixed, the residual distinction between L and L' can only survive as a second admissible completion above that same retained corridor while preserving the same retained status surface. But this is precisely the kind of duplicate admissible completion excluded by the standing strict-upper-gap discipline in the retained OP 1 reading. Hence the assumed witness contradicts the retained strict-upper-gap layer. Therefore no completion-side duplicate-signature witness can exist. □

Corollary H9.235 (No duplicate-signature witness survives the retained OP 1 kernel)**.** *Assume the hypotheses of Proposition H9.231 and Proposition H9.234. Then no duplicate-signature witness for the live OP 1 kernel exists on the intended admissible windows.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. If a duplicate-signature witness existed, Corollary H9.232 would force it to be of completion-side type. Proposition H9.234 eliminates that case. Hence no duplicate-signature witness survives. □

Theorem H9.236 (Closure of the live OP 1 kernel under the standing retained exclusion disciplines)**.** *Assume the same-corridor reversal certificate of Definition H9.149. Assume moreover that the retained no-second-carrier discipline and the retained strict-upper-gap discipline both hold on the intended admissible windows in the senses used above. Then the optimum-pinning comparison-rigidity certificate of Definition H9.169 holds on the live OP 1 kernel. Consequently,*

$$\Delta_{\mathrm{CL}} = 0.$$

Equivalently, under the standing retained exclusion disciplines the normalized OP 1 route

$$\mathrm{TR}_4 \Longrightarrow \mathrm{IP}_{10} \Longrightarrow \mathrm{IP}_9 \Longrightarrow \mathrm{IP}_8$$

closes on the intended admissible windows.

Proof. By Corollary H9.235, no duplicate-signature witness exists. Proposition H9.226 therefore yields the optimum-pinning comparison-rigidity certificate. The final sentence of that same proposition then gives $\Delta_{\mathrm{CL}} = 0$ via the already established retained-window-rigidity / admissible-window-uniqueness bridge. □

Remark H9.237 (What the completion-side collapse means for the role of the Monte-Carlo probes)**.** *After the present step, the Monte-Carlo probes with the fusion/decoder reading should not be read as replacements for the OP 1 theorematic closure. Their real leverage is earlier in the search. They compress the search space by classifying residual behavior into a small number of decoder-relevant recurrence classes and by telling us whether a surviving defect should first be treated as* `ip`*-local or as* `bsbl`*-translation-sensitive. That is why the work now feels faster: the probes do not prove the retained no-second-carrier or strict-upper-gap exclusions, but they stop us from reopening the wrong parts of the document and from carrying broad rival-loop pictures once the residual signature is already locally pinned.*

In the present OP 1 line, this means the probes mainly accelerated the route to the right normal form. They helped isolate the live burden as an injectivity problem, then as a two-channel contradiction search, and finally as the completion-side residue that can be handed directly to the strict-upper-gap gate. So the speed-up is real, but it is a search-space reduction and routing advantage*, not a substitute for the structural closure theorems themselves.*

Reader cue. Read the next block as the first honest post-OP 1 normalization of the live OP 5 Step (B) burden. Its purpose is not to declare the routed photon-frame balance defect solved, but to identify the smallest currently licensed structural kernel on which that burden now lives: the production or exclusion of a carrier-upgraded routed-balance witness at the routed photon-frame cutoff.

19AZ$^{(6a)}$. First carrier-upgraded witness normal form for the live OP 5 Step (B) kernel

After the closure of the live OP 1 kernel, the honest next burden is OP 5 Step (B), i.e. the routed photon-frame balance defect $\Delta_{\Theta R}(T)$. The current line should not reopen that burden through every packet-wave, breathing-envelope, frame-flicker, or helix picture at once. The earlier carrier-first OP 5 architecture already teaches that visible mismatches come in two grades: either they remain screening-only and therefore subordinate to one still-available

© 2026 Marc Brendecke. Preprint — not peer reviewed.

common closure-bearing return-side carrier package, or they upgrade and certify loss of that common carrier package itself. The cleanest first OP 5(B) normalization is therefore to ask not for a new global formula immediately, but for the smallest carrier-grade witness through which a still-live routed balance defect could honestly persist.

Definition H9.238 (Routed-balance witness for the live OP 5 Step (B) kernel)**.** *Fix the routed photon-frame cutoff $T = \gamma_L^{-3/2}$ and the routed balance defect $\Delta_{\Theta R}(T)$ from Definition H9.126. Under the standing carrier-first OP 5 architecture, an admitted visible witness w is called a* routed-balance witness for the live OP 5 Step (B) kernel *if w is cited to support the claim that the routed photon-frame balance layer remains structurally live at the cutoff T. Such a witness is called*

1. screening-only *if it remains screening-only in the sense of Lemma H7.48;*
2. carrier-upgraded *if it is carrier-upgraded in the same sense.*

Thus a routed-balance witness is not a new witness species; it is the OP 5 Step (B) reading of the already licensed witness-grade dichotomy at the routed photon-frame cutoff.

Proposition H9.239 (The live OP 5 Step (B) kernel is a carrier-upgraded witness problem)**.** *Assume Proposition H7.27, Proposition H7.33, Lemma H7.42, Lemma H7.45, Lemma H7.48, and Corollary H7.52. Then the honest live OP 5 Step (B) kernel on the present line is the following alternative:*

(i) either every routed-balance witness at the routed photon-frame cutoff remains screening-only, in which case no carrier/package-level OP 5 Step (B) obstruction has yet been exhibited;

(ii) or there exists a carrier-upgraded routed-balance witness, in which case the live OP 5 Step (B) burden has genuinely reached the carrier/package layer.

In particular, the first honest structural reopening of $\Delta_{\Theta R}(T)$ is not an arbitrary visible mismatch, but the production or exclusion of a carrier-upgraded routed-balance witness.

Proof. By Lemma H7.48, every admitted visible witness at the carrier-first OP 5 layer lies in exactly one of two grades: screening-only or carrier-upgraded. Definition H9.238 does nothing more than read that existing dichotomy at the routed photon-frame cutoff T and relative to the residual $\Delta_{\Theta R}(T)$. If a cited witness remains screening-only, then by Corollary H7.52 its upstream content is still representable through some common closure-bearing return-side carrier package; hence no carrier/package-level OP 5 Step (B) obstruction has yet been exhibited. If, on the other hand, a cited witness upgrades to carrier grade, then the same corollary says that no such common carrier package remains available, so the routed balance burden has genuinely reached the carrier/package layer. Since no third currently licensed witness grade exists, the displayed alternative is exhaustive. Therefore the first honest structural OP 5 Step (B) kernel is exactly the production or exclusion of a carrier-upgraded routed-balance witness. □

Corollary H9.240 (First normalized reopening route for the live OP 5 Step (B) kernel)**.** *Under the hypotheses of Proposition H9.239, the smallest currently licensed normalized reopening route for a genuinely live OP 5 Step (B) witness is*

$$\mathrm{TR}_4 \Longrightarrow \mathrm{BS}_1 \Longrightarrow \mathrm{BS}_3 \Longrightarrow \mathrm{BL}_2 \Longrightarrow \mathrm{BL}_4 \Longrightarrow \mathrm{BL}_5,$$

with the last two steps used only when the routed-balance witness remains live after the masked-single-carrier test and localizes at the readout/kernel stratum.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. The total-channel baseline remains frozen by TR_4. Because the routed photon-frame balance defect is not a pure local optimum-pinning problem but a return-side witness-grade problem, the first honest move is bridge-side rather than imbalance-side: the route enters through the matrix/phase surface-change gates BS_1 and BS_3. Once that surface change has been made, the first honest diagnostic question is whether the putative routed-balance witness is still compatible with one masked single carrier, which is precisely the gate BL_2. If it collapses there, no deeper reopening is licensed. If it remains live, then BL_4 is exactly the normalized stratification step that localizes the first remaining obstruction. And when that localized obstruction sits in the readout/kernel stratum, BL_5 is the recorded compression to absence of one common shell-stable carrier package. Therefore the displayed BS/BL route is exactly the smallest currently licensed reopening route for a genuinely live OP 5 Step (B) witness. □

Remark H9.241 (What this first OP 5 Step (B) normalization buys)**.** *This is not yet an OP 5 Step (B) closure theorem. The routed photon-frame balance defect $\Delta_{\Theta R}(T)$ is still honestly live. But the burden is now sharper. It is no longer carried as an undifferentiated packet/helix/flicker mismatch surface. It is carried as a witness-grade question: does any routed-balance witness actually upgrade to carrier/package level, or do all currently visible routed witnesses remain screening-only beneath one still-available common return-side carrier package? In the present QAM language, OP 1 was compressed onto the* IP*-route, whereas the live OP 5 Step (B) burden now announces itself as a* BS/BL*-route burden from the start.*

Reader cue. Read the next block as the first negative-side sharpening of the live OP 5 Step (B) kernel. Its purpose is not yet to close the routed photon-frame balance defect, but to identify the first retained situation in which a routed-balance witness cannot honestly escalate to carrier/package grade at all.

19AZ$^{(6b)}$. Carrier-stable routed returns prevent witness-grade escalation on the live OP 5 Step (B) kernel

The previous normalization reduced the live OP 5 Step (B) burden to the production or exclusion of a carrier-upgraded routed-balance witness. The next honest question is whether every routed-balance witness must be tested all the way down to the carrier/package layer, or whether one already has a retained positive branch on which such escalation is structurally forbidden. The earlier routed-return package answers this. Once the routed realization lies on the carrier-stable routed return normal form, every admitted visible routed mismatch is already subordinate to one common closure-bearing return-side carrier. In that situation the witness-grade question collapses on the positive side: a routed-balance witness may still exist as screening-level visible data, but it can no longer count as a carrier-grade OP 5 Step (B) witness.

Proposition H9.242 (Carrier-stable routed returns prevent routed-balance witness escalation)**.** *Assume Proposition H9.239, Corollary H9.240, Corollary H7.52, Theorem H7.54, and Theorem H7.66. Let $(b, r, s; w)$ be a routed branch-shell realization at the cutoff $T = \gamma_L^{-3/2}$ together with a routed-balance witness w in the sense of Definition H9.238. If $(b, r, s; w)$ lies on the* carrier-stable routed return normal form *of Theorem H7.66, then w remains screening-only and cannot be carrier-upgraded. In particular, no carrier-upgraded routed-balance witness exists on the carrier-stable routed-return branch.*

Proof. By Theorem H7.66, the carrier-stable routed return normal form means precisely that there exists one common closure-bearing return-side carrier through which the routed realization is read and that every admitted visible routed mismatch collapses beneath that carrier

© 2026 Marc Brendecke. Preprint — not peer reviewed.

to the carrier-stable visible normal form. Apply this to the routed-balance witness w. Theorem H7.54 then says that, over that same stable carrier package, w has only one current structural role: it witnesses subordinate screening-loss normal form and does not define an independent carrier-relevant visible species. But Corollary H7.52 identifies carrier upgrade exactly with the first failure of every common closure-bearing return-side carrier-supported realization. Since the carrier-stable routed return normal form explicitly retains such a common carrier package, the threshold for carrier upgrade is not met. Hence w remains screening-only and cannot be carrier-upgraded. Because w was arbitrary, no carrier-upgraded routed-balance witness exists on the carrier-stable routed-return branch. □

Corollary H9.243 (The live OP 5 Step (B) burden survives only on the obstructed routed-return branch)**.** *Under the hypotheses of Proposition H9.242, a genuinely live OP 5 Step (B) burden can survive only on the* obstructed routed return normal form *of Theorem H7.66. Equivalently, after the gates* $\mathrm{TR}_4 \Rightarrow \mathrm{BS}_1 \Rightarrow \mathrm{BS}_3 \Rightarrow \mathrm{BL}_2$*, any routed-balance witness that still honestly threatens carrier/package loss must already belong to the negative routed-return branch; and when that negative branch localizes at the readout/kernel stratum,* BL_5 *compresses the remaining burden to absence of one common shell-stable carrier package.*

Proof. By Proposition H9.239, the live OP 5 Step (B) kernel is exactly the production or exclusion of a carrier-upgraded routed-balance witness. Proposition H9.242 excludes such a witness on the carrier-stable routed-return branch. Therefore any genuinely live routed-balance burden must sit on the only remaining routed-return branch, namely the obstructed routed return normal form of Theorem H7.66. Corollary H9.240 already records that, after the masked-single-carrier gate BL_2, the next honest localization step is BL_4, and that a readout/kernel localization is then compressed by BL_5 to absence of one common shell-stable carrier package. This is exactly the stated reformulation. □

Remark H9.244 (What this second OP 5 Step (B) sharpening buys)**.** *The live OP 5 Step (B) burden is now smaller than in the local pass. It is no longer "any routed-balance witness that might still upgrade." It is "a routed-balance witness on the obstructed routed-return branch that survives the masked-single-carrier test and, if needed, localizes further." This is also the point at which the Monte-Carlo/fusion probes materially accelerate the search: once the probes classify a run as carrier-stable on the routed-return side, one no longer needs to pursue carrier-grade OP 5 escalation there. The probes still do not prove the theorematic branch separation by themselves, but they do cut away large amounts of blind BS/BL reopening by steering attention directly to the small class of routed returns that still look obstruction-like after the carrier-stable branch has been excluded.*

Reader cue. Read the next block as the first localization split of the remaining OP 5 Step (B) burden on the negative routed-return side. Its purpose is not yet to declare the obstructed routed return harmless, but to separate the part that is already inherited from the upstream OP3/QAM obstruction package from the part that would remain as a genuinely downstream carrier-grade OP 5 obstruction.

19AZ$^{(6c)}$. Obstructed routed returns split into upstream-screened versus downstream carrier-grade OP 5 Step (B) witnesses

After 19AZ$^{(6b)}$, the live OP 5 Step (B) burden survives only on the obstructed routed-return branch. But even this negative branch is still too coarse to be the final witness kernel. The already extracted global obstruction package says that, once the admissible run is inside the bundled OP1→branch→OP3/QAM→OP5 channel, the first licensed failure is already

© 2026 Marc Brendecke. Preprint — not peer reviewed.

localized either upstream in the OP3/QAM obstruction package or downstream at the carrier-upgraded OP5 obstruction layer. Therefore the obstructed routed-return branch must itself be split: either the routed obstruction is only the return-side shadow of an upstream OP3/QAM obstruction already fixed before carrier-grade OP5 escalation, or it is a genuinely downstream carrier-grade OP5 obstruction that survives that upstream screening.

Definition H9.245 (Upstream-screened versus downstream carrier-grade obstructed routed-return witness)**.** *Assume the hypotheses of Corollary H9.243. Let* $(b, r, s; w)$ *be a routed branch-shell realization on the* obstructed routed return normal form *together with a routed-balance witness* w*.*

(i) We say that $(b, r, s; w)$ *is an* upstream-screened obstructed routed-return witness *if the currently licensed negative routed-return reading of* $(b, r, s; w)$ *is already forced by the upstream OP3/QAM obstruction package of Proposition H7.76, so that the witness does not yet certify an independent downstream carrier-grade OP 5 obstruction.*

(ii) We say that $(b, r, s; w)$ *is a* downstream carrier-grade obstructed routed-return witness *if the witness remains honestly live after the upstream OP3/QAM obstruction screening and therefore certifies a genuinely downstream carrier-upgraded OP 5 obstruction layer in the sense of Proposition H7.76.*

Proposition H9.246 (The remaining obstructed routed-return burden is an upstream/downstream split)**.** *Assume Proposition H9.239, Corollary H9.243, Proposition H7.76, and Proposition H7.77. Let* $(b, r, s; w)$ *be a genuinely live routed-balance witness on the obstructed routed return normal form. Then exactly one of the following two cases is currently licensed.*

(i) ***Upstream-screened case.*** *The negative routed-return reading of* $(b, r, s; w)$ *is already accounted for by the upstream OP3/QAM obstruction package, and the witness is therefore not yet an independent downstream carrier-grade OP 5 obstruction.*

(ii) ***Downstream carrier-grade case.*** *The witness survives the upstream OP3/QAM obstruction screening and therefore remains as a genuinely downstream carrier-grade OP 5 obstruction witness.*

No third autonomous obstructed routed-return witness class is currently licensed.

Proof. By Corollary H9.243, every genuinely live OP 5 Step (B) witness must already lie on the obstructed routed return normal form. Once the run is inside the bundled OP1→branch→OP3/QAM→OP5 channel, Proposition H7.76 says that the first licensed failure is already localized either upstream in the OP3/QAM obstruction package or downstream at the carrier-upgraded OP5 obstruction layer. Proposition H7.77 adds that no third autonomous terminal visible class is licensed outside that bundled obstruction reading. Applying this dichotomy to the obstructed routed-return witness $(b, r, s; w)$ yields exactly the two stated cases. The first is the upstream-screened case, where the negative routed-return picture is merely the return-side shadow of an obstruction already fixed upstream. The second is the downstream carrier-grade case, where the witness remains live after that upstream screening and therefore truly belongs to the OP 5 obstruction layer. Exclusivity and exhaustiveness are precisely the content of the bundled global obstruction dichotomy. □

Corollary H9.247 (The genuinely new OP 5 Step (B) burden survives only on the downstream branch)**.** *Under the hypotheses of Proposition H9.246, any genuinely* new *OP 5 Step (B) burden survives only in the downstream carrier-grade case of Definition H9.245.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Equivalently, once the obstructed routed return has been shown not to be merely the return-side shadow of the upstream OP3/QAM obstruction package, the remaining burden is already localized to the downstream carrier-upgraded OP 5 obstruction layer.

Proof. In the upstream-screened case, the negative routed-return witness is already accounted for by the upstream OP3/QAM obstruction package and therefore does not constitute a genuinely new downstream OP 5 burden. Hence only the downstream carrier-grade case can carry a genuinely new OP 5 Step (B) burden after the upstream screening has been discharged. This is exactly the stated reformulation. □

Remark H9.248 (What this third OP 5 Step (B) sharpening buys)**.** *The live OP 5 Step (B) burden is now smaller again. It is no longer merely "a witness on the obstructed routed-return branch." It is "a witness on that negative branch that is not already explained by the upstream OP3/QAM obstruction package." This is also where the Monte-Carlo/fusion probes become strategically useful on the negative side: they do not prove the upstream/downstream split, but they help tell much earlier whether a run looks like an inherited OP3/QAM obstruction shadow or like a genuinely downstream carrier-grade obstruction. In practical search terms, this means that many apparently live negative routed returns can now be discarded upstream before one spends time reopening the deepest OP 5 carrier-grade layer.*

Reader cue. Read the next block as the discharge of the upstream-screened side of the remaining OP 5 Step (B) burden. Its purpose is not yet to close the downstream carrier-grade obstruction kernel, but to remove every negative routed-return witness whose content is already absorbed by the upstream OP3/QAM obstruction package.

19AZ$^{(6d)}$. Upstream-screened obstructed routed returns are discharged before the downstream OP 5 Step (B) carrier-grade kernel begins

After 19AZ$^{(6c)}$, the live negative routed-return burden has only two currently licensed readings: upstream-screened or downstream carrier-grade. But the upstream-screened case is not merely weaker; it is not an independent OP 5 Step (B) burden at all. By definition it is already the routed return shadow of an obstruction whose first licensed failure sits upstream in the OP3/QAM package. Therefore the honest live OP 5 Step (B) kernel starts only after that upstream explanation has been discharged.

Proposition H9.249 (Upstream-screened obstructed routed-return witnesses do not define an independent OP 5 Step (B) burden)**.** *Assume Proposition H9.246, Corollary H9.247, Proposition H7.76, and Proposition H7.77. Let $(b, r, s; w)$ be an upstream-screened obstructed routed-return witness in the sense of Definition H9.245. Then $(b, r, s; w)$ does not define an independent live OP 5 Step (B) burden; its entire negative routed-return content is already discharged by the upstream OP3/QAM obstruction package.*

Proof. By Definition H9.245, an upstream-screened obstructed routed-return witness is precisely one whose currently licensed negative routed-return reading is already forced by the upstream OP3/QAM obstruction package. Proposition H7.76 says that once the admissible run is inside the bundled OP1→branch→OP3/QAM→OP5 channel, the first licensed failure is already localized either upstream in that OP3/QAM obstruction package or downstream at the carrier-upgraded OP5 obstruction layer. Since the witness is, by hypothesis, upstream-screened, the first licensed failure is already on the upstream side of that dichotomy. Proposition H7.77 excludes any third autonomous terminal visible class outside that bundled

© 2026 Marc Brendecke. Preprint — not peer reviewed.

obstruction reading. Therefore no additional independent OP 5 Step (B) burden remains attached to $(b, r, s; w)$ once the upstream package has been acknowledged. In other words, the routed-return shadow carried by $(b, r, s; w)$ is fully discharged upstream and does not itself define a genuinely new OP 5 Step (B) burden. □

Corollary H9.250 (The live OP 5 Step (B) kernel reduces to the downstream carrier-grade obstructed routed-return branch)**.** *Under the hypotheses of Proposition H9.249, the honest live OP 5 Step (B) kernel is exactly the production or exclusion of a downstream carrier-grade obstructed routed-return witness. Equivalently, once the upstream-screened negative routed-return cases have been discharged, the only remaining live OP 5 Step (B) burden sits on the downstream carrier-grade branch isolated in Definition H9.245.*

Proof. By Proposition H9.246, every genuinely live obstructed routed-return witness lies in exactly one of two currently licensed classes: upstream-screened or downstream carrier-grade. Proposition H9.249 removes the first class as an independent OP 5 Step (B) burden. Therefore only the downstream carrier-grade class remains available as the honest live OP 5 Step (B) kernel. This is exactly the stated reformulation. □

Remark H9.251 (What this fourth OP 5 Step (B) sharpening buys)**.** *The live OP 5 Step (B) burden is now concentrated on one branch only. It is no longer "an obstructed routed-return witness that is not yet known to be harmless." It is "a downstream carrier-grade obstructed routed-return witness after the upstream OP3/QAM shadow cases have been discharged." This is the first point on the OP 5 side where the negative branch has been reduced to a single honest carrier-grade kernel. Strategically, this is also where the Monte-Carlo/fusion probes become most valuable as a search accelerator: once a negative routed return looks upstream-screened, it can be discarded without reopening the downstream carrier-grade layer, so the expensive search is reserved only for the small class of runs that still look honestly downstream after the upstream package has been factored out.*

Reader cue. Read the next block as the compression of the remaining downstream OP 5 Step (B) kernel to one shell-package deficit question on the retained branch-shell corridor. Its purpose is not yet to declare OP 5 Step (B) globally closed, but to show that once the upstream shadow cases are gone, the honest remaining burden is exactly the absence of one common shell-stable carrier package.

19AZ$^{(6e)}$. Downstream carrier-grade obstructed routed returns compress to one common shell-stable carrier-package deficit on the retained branch-shell corridor

After 19AZ$^{(6d)}$, the live OP 5 Step (B) burden is concentrated on the downstream carrier-grade obstructed routed-return branch. The smallest honest next question is therefore no longer whether some vague routed mismatch survives, but whether the retained branch-shell corridor still admits one common shell-stable carrier package once that downstream witness has survived every earlier screening. On the current line this is exactly the readout/kernel compression question behind BL_5: if the branch and shell strata themselves remain intact on the retained routed corridor, then every remaining downstream carrier-grade witness is nothing but failure of one common shell-stable carrier package.

Definition H9.252 (Downstream shell-package deficit witness for the live OP 5 Step (B) kernel)**.** *Fix the live OP 5 Step (B) setup of Definition H9.238. A quadruple $(b, r, s; w)$ is called a* downstream shell-package deficit witness for the live OP 5 Step (B) kernel *if*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(i) $(b, r, s; w)$ *is a downstream carrier-grade obstructed routed-return witness in the sense of Definition H9.245, and*

(ii) on the retained routed branch-shell corridor determined after the gates $\mathrm{TR}_4 \Rightarrow \mathrm{BS}_1 \Rightarrow \mathrm{BS}_3 \Rightarrow \mathrm{BL}_2$*, the branch and shell strata remain intact, so the remaining localized burden is entirely in the readout/kernel stratum and therefore equals failure of one common shell-stable carrier package in the sense of Lemma H2.169.*

Proposition H9.253 (Downstream carrier-grade OP 5 witnesses compress to one common shell-stable carrier-package deficit on the retained branch-shell corridor)**.** *Assume Corollary H9.250, Corollary H9.240, Theorem H7.66, Lemma H2.168, and Lemma H2.169. Let* $(b, r, s; w)$ *be a genuinely live downstream carrier-grade obstructed routed-return witness for the OP 5 Step (B) kernel. If, on the retained routed branch-shell corridor carried through* $\mathrm{TR}_4 \Rightarrow \mathrm{BS}_1 \Rightarrow \mathrm{BS}_3 \Rightarrow \mathrm{BL}_2$*, the branch and shell strata remain intact, then* $(b, r, s; w)$ *is exactly a downstream shell-package deficit witness in the sense of Definition H9.252. Equivalently, under the retained branch-shell intactness assumptions, the whole remaining downstream OP 5 Step (B) burden compresses to absence of one common shell-stable carrier package.*

Proof. By Corollary H9.250, every genuinely live OP 5 Step (B) burden is already a downstream carrier-grade obstructed routed-return witness. Corollary H9.240 records that after the retained gates $\mathrm{TR}_4 \Rightarrow \mathrm{BS}_1 \Rightarrow \mathrm{BS}_3 \Rightarrow \mathrm{BL}_2$, the first further honest localization step is BL_4, and that whenever the first remaining localized obstruction lies in the readout/kernel stratum, the compression gate BL_5 rewrites the burden as absence of one common shell-stable carrier package. Now assume that on the retained routed branch-shell corridor the branch and shell strata themselves remain intact. Then Lemma H2.168 forces every remaining first local obstruction into the readout/kernel stratum, and Lemma H2.169 identifies that stratum exactly with failure of one common shell-stable carrier package. Therefore the surviving downstream carrier-grade witness $(b, r, s; w)$ is not a broader residual species: on the retained branch-shell corridor it is precisely a downstream shell-package deficit witness. This is the stated compression. □

Corollary H9.254 (The live OP 5 Step (B) kernel reduces to one common shell-stable carrier-package deficit problem on the retained branch-shell corridor)**.** *Under the hypotheses of Proposition H9.253, the honest live OP 5 Step (B) kernel on the retained branch-shell corridor is exactly the production or exclusion of a downstream shell-package deficit witness. Equivalently, once the upstream-screened cases have been discharged and the branch/shell strata remain intact, OP 5 Step (B) is reduced to the single question whether one common shell-stable carrier package is absent.*

Proof. Corollary H9.250 says that the live OP 5 Step (B) burden is already concentrated on downstream carrier-grade obstructed routed-return witnesses. Proposition H9.253 then shows that, under retained branch-shell intactness, every such witness is exactly a downstream shell-package deficit witness. Hence the remaining kernel is neither a diffuse routed-return problem nor a broader carrier-grade witness search; it is exactly the stated one-package deficit problem. □

Remark H9.255 (What this fifth OP 5 Step (B) sharpening buys)**.** *This is the first point on the OP 5 side where the surviving kernel is no longer merely a branch classification problem. It has become a concrete deficit question. The live burden is now: not "some downstream obstruction somewhere on the negative routed-return branch," but "the absence of one common shell-stable carrier package on the retained branch-shell corridor." This is also why the*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Monte-Carlo/fusion probes feel so effective here. Once the probes indicate that the run is both downstream and branch/shell-intact, the search no longer needs to explore broad alternative witness pictures; it can go straight to the small shell-package deficit kernel that BL_5 *was designed to expose.*

oindent**Reader cue.** Read the next block as the final conservative test of the live OP 5 Step (B) kernel on the retained branch-shell corridor. Its purpose is not to claim a new global closure principle from scratch, but to show that once the surviving downstream kernel is already compressed to one shell-package deficit question, the inherited OP 3/QAM shell-closure package removes exactly that final deficit under the retained shell-transfer discipline.

$19\mathrm{AZ}^{(6f)}$. The inherited shell-closure package discharges the downstream shell-package deficit on the retained branch-shell corridor

After $19\mathrm{AZ}^{(6e)}$, the live OP 5 Step (B) kernel has been compressed as far as possible without reusing the earlier OP 3/QAM shell package. The honest remaining question is therefore very narrow: can the retained branch-shell corridor both inherit the shell-closure package and yet still fail to admit one common shell-stable carrier package? On the present line the answer is no. Once the retained branch-shell corridor is genuinely read through the already established shell certificate / shell-stable kernel-readout package, the missing common shell-stable carrier of $19\mathrm{AZ}^{(6e)}$ is exactly the kind of deficit that the inherited package forbids.

Definition H9.256 (Retained shell-transfer discipline for the live OP 5 Step (B) kernel)**.** *Fix the live OP 5 Step (B) setup of Definition H9.238. We say that the retained routed branch-shell corridor satisfies the* retained shell-transfer discipline *if, after the retained gates* $\mathrm{TR}_4 \Rightarrow \mathrm{BS}_1 \Rightarrow \mathrm{BS}_3 \Rightarrow \mathrm{BL}_2$*, the routed branch-shell realization inherits the shell admissibility certificate and the equivalent shell-stable kernel/readout package from Theorem H2.170. Equivalently, on that retained corridor there exists one common shell-stable kernel/readout carrier through which all closure-relevant routed readout factors.*

Proposition H9.257 (The inherited shell-closure package excludes downstream shell-package deficit witnesses on the retained branch-shell corridor)**.** *Assume Corollary H9.254, Definition H9.256, Theorem H2.170, and Theorem H2.166. Then no downstream shell-package deficit witness in the sense of Definition H9.252 can exist on the retained branch-shell corridor. Equivalently, once the retained branch-shell corridor inherits the shell certificate / shell-stable kernel-readout package, the remaining OP 5 Step (B) kernel from* $19\mathrm{AZ}^{(6e)}$ *is discharged.*

Proof. Assume toward contradiction that a downstream shell-package deficit witness exists on the retained branch-shell corridor. By Definition H9.252, this means that the surviving downstream carrier-grade routed-return witness is already compressed to absence of one common shell-stable carrier package on that corridor. But by Definition H9.256, the same retained corridor inherits the shell admissibility certificate and hence, by Theorem H2.170 together with Theorem H2.166, admits one common shell-stable kernel/readout carrier through which the closure-relevant routed readout factors. This is exactly the negation of the shell-package deficit asserted by the witness. The contradiction shows that no downstream shell-package deficit witness can exist once the retained shell-transfer discipline holds. Therefore the remaining OP 5 Step (B) kernel from Corollary H9.254 is discharged on that corridor. □

Corollary H9.258 (Closure of the live OP 5 Step (B) kernel under the standing retained branch-shell and shell-transfer disciplines)**.** *Assume the standing retained branch-shell intactness hypotheses from Proposition H9.253 together with the retained shell-transfer discipline of*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition H9.256. Then the live OP 5 Step (B) kernel closes:

$$\Delta_{\Theta R} = 0.$$

Equivalently, under the current retained no-upstream-shadow, branch/shell-intact, and inherited shell-package disciplines, no honest live OP 5 Step (B) witness survives.

Proof. By Corollary H9.254, the live OP 5 Step (B) kernel under the retained branch-shell hypotheses is exactly the production or exclusion of a downstream shell-package deficit witness. Proposition H9.257 excludes precisely that witness class once the retained shell-transfer discipline holds. Hence no live OP 5 Step (B) witness remains, so the kernel closes and $\Delta_{\Theta R} = 0$. □

Remark H9.259 (What this sixth OP 5 Step (B) sharpening buys)**.** *This is the OP 5 analogue of the OP 1 local closure step. The result is intentionally conservative. It does not say that every historical reading of OP 5 is now globally absorbed into one theorematic sentence. It says something narrower and structurally cleaner: once the live OP 5 Step (B) burden has been reduced to one downstream shell-package deficit on the retained branch-shell corridor, the inherited OP 3/QAM shell-closure package forbids exactly that final deficit. Strategically, this is also the point where the Monte-Carlo/fusion probes show their full value on the negative routed-return side. They do not prove the inherited shell package, but they steer the search quickly onto the tiny downstream/branch-shell-intact corridor where the shell-transfer discipline can be applied and the final witness class disappears.*

Remark H9.260 (Historical transition note for the archived v2.27.x release-control fragment)**.** *Read the next local post-release fragment as archival proof-memory from an earlier public hardening cycle. Its purpose is to preserve release genealogy, checkpoint recall, and local baseline memory, not to function as a current roadmap, freeze checklist, or active control surface for the present 4.7.0 line.*

Archived archival stage local post-release note (historical baseline memory)

H10 Release orientation

This document is presented as the full monolithic Companion text of the current line. Within that full monolithic presentation, the newly integrated archival stage block freezes the mathematically relevant pre-OP foundation developed in the current working stage. The release claim of that integrated block is therefore intentionally limited and precise:

1. the groundwork layer before OP 1–5 is completed as a coherent route-faithful structure;
2. the current local state is conservatively instantiated as a frontier-certified pre-entry state;
3. the line already supports controlled witness-based descent and lead-guided frontier resolution;
4. explicit OP 1–5 work is intentionally deferred to the next stage.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

H11 The three-coordinate live residual interface

Definition H11.1 (Three-coordinate live residual interface)**.** *The current line is organized around the live residual interface*

$$\mathbf{R}_{\mathrm{live}} = (\rho_{\mathrm{CL}}, \rho_{\Theta}, \rho_{A}),$$

where ρ_{CL} is the horizon-side scalar closure coordinate, while $(\rho_{\Theta}, \rho_{A})$ are the boundary-side slow phase and slow envelope packet coordinates.

Remark H11.2. *The current open residual work is therefore carried by exactly three live coordinates: one scalar horizon-side coordinate and a two-component boundary-side packet coordinate.*

Definition H11.3 (Current normalized state)**.** *Using a current admissible normalization scale triple*

$$(s_{\mathrm{CL,curr}}, s_{\Theta,\mathrm{curr}}, s_{A,\mathrm{curr}}),$$

the current normalized live residual state is

$$\widetilde{\mathbf{R}}_{\mathrm{curr}} = (\widetilde{\rho}_{\mathrm{CL,curr}}, \widetilde{\rho}_{\Theta,\mathrm{curr}}, \widetilde{\rho}_{A,\mathrm{curr}}),$$

with the obvious channel-wise normalization.

H12 Paired conservative machine and compression

Definition H12.1 (Local paired conservative machine)**.** *The local paired conservative machine consists of the paired intake, request, response, fulfillment, resolution, close-or-promote, cycle, return, and branch layers, always preserving:*

1. *the inherited horizon/boundary distinction,*
2. *the inherited earliest-boundary stop,*
3. *certificate-neutrality.*

Definition H12.2 (Compact paired conservative normal form)**.** *The compact paired conservative normal form is the compressed ordered object*

$$\mathcal{N}^{*}_{\mathrm{pair}} = (\mathcal{N}_{\mathrm{hor}}, \mathcal{N}_{\mathrm{bdry}}),$$

which records only the downstream-relevant scalar horizon-side content and packet boundary-side content carried by the paired machine.

Theorem H12.3 (Compression and invariant control)**.** *Every finite route-faithful local paired conservative machine admits a compact paired conservative normal form that preserves the inherited horizon/boundary distinction, the inherited earliest-boundary stop, and certificate-neutrality, while remaining readable through the extracted live residual state.*

Proof. The paired machine is finite at each local stage, and its admissible content separates into one horizon-side scalar channel and one boundary-side packet channel. Compression therefore produces an ordered summary object preserving exactly the route-faithful invariants built into the original machine. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

H13 Tool layer

Definition H13.1 (Extracted tool layer)**.** *The extracted tool layer consists of:*

1. *compression-before-comparison,*
2. *monotone selection on the compressed state,*
3. *a route-faithful close-or-promote shortcut,*
4. *scalar-packet decoupling for genuinely one-sided arguments.*

Theorem H13.2 (Tool extraction theorem)**.** *The compact paired conservative normal form supports the extracted tool layer entirely through the preserved route-faithful invariant class and the extracted three-coordinate residual state.*

Proof. Each tool is defined directly on the compact normal form and depends only on route-faithful invariants and on the residual coordinates. No reopening of the full internal machine is required unless the compressed tool layer is insufficient. □

H14 Tool stability and normalization

Definition H14.1 (Tool-stable invariant regime)**.** *A compact paired conservative normal form lies in a tool-stable invariant regime if monotone selectors are compression-stable, close-side states are route-faithfully robust, promote-side continuation is honest rather than a pure weight artifact, and the preserved route-faithful invariants remain unchanged under admissible uses of the tool layer.*

Definition H14.2 (Canonical normalization regime)**.** *A canonical normalization regime is an admissible class of normalization scale triples in which the selector class and close/promote classification remain unchanged under bounded side-preserving scale replacements.*

Theorem H14.3 (Normalization theorem)**.** *Every compact paired conservative normal form equipped with an admissible normalization scale triple admits a normalized live residual reading and a normalized weighted residual energy*

$$\widetilde{\mathcal{E}}_w = w_{\mathrm{CL}}|\widetilde{\rho}_{\mathrm{CL}}|^2 + w_{\Theta}|\widetilde{\rho}_{\Theta}|^2 + w_A|\widetilde{\rho}_A|^2.$$

Inside a canonical normalization regime, comparison, selection, and close/promote testing are independent of arbitrary raw scale imbalance.

Proof. Normalization is performed channel-wise inside the route-faithful split. Within the canonical normalization regime, equivalent admissible scale choices preserve the relevant comparisons and branch classifications, so the normalized state is the preferred pre-OP working surface. □

H15 Residual geometry

Definition H15.1 (Normalized residual geometry)**.** *The normalized residual space carries:*

1. *admissible descent sectors and strict descent sectors,*
2. *trapping close basins,*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

3. *honest promote regions,*

4. *a residual frontier separating the geometrically unresolved zone.*

Definition H15.2 (Residual frontier)**.** *The residual frontier is the part of normalized residual space lying in neither a trapping close basin nor an honest promote region.*

Theorem H15.3 (Geometric control theorem)**.** *In a fixed canonical normalization regime, the normalized residual space admits a route-faithful geometric control structure consisting of admissible descent sectors, close basins, honest promote regions, a residual frontier, and descent selectors on finite admissible successor families.*

Proof. These structures are defined entirely within the normalized residual space and remain inside the same route-faithful invariant class. Each construction therefore preserves the inherited horizon/boundary distinction, earliest-boundary stop, and certificate-neutrality. □

H16 Residual dynamics

Definition H16.1 (Admissible descent trajectory)**.** *An admissible descent trajectory is a sequence of normalized residual states in which each successor lies in the admissible descent sector of its predecessor. If strict descent is chosen whenever available, the trajectory is called strictly selected.*

Lemma H16.2 (First no-oscillation lemma)**.** *A strictly selected admissible descent trajectory cannot exhibit a two-step oscillation between distinct states if strict descent is available at both states.*

Proof. Strict descent at the first state gives strict energy decrease; strict descent at the second state gives another strict decrease. Returning to the first state would therefore force the same energy to be both strictly larger and strictly smaller than itself, a contradiction. □

Theorem H16.3 (Dynamic control theorem)**.** *Admissible descent dynamics on the normalized residual space has nonincreasing normalized weighted residual energy. Under strict descent, two-step oscillation is excluded; close-side behavior is represented by trapping close basins; promote-side behavior is represented by persistent honest promote regions; and every honest promote-to-close transition must pass through the residual frontier.*

Proof. The monotonicity of energy follows from admissible descent. The no-oscillation claim is Lemma H16.2. The basin and frontier statements are the dynamic reading of the previously defined geometric structures. □

H17 Compressed working principles

Definition H17.1 (Compressed working principles)**.** *The current groundwork is governed by four working principles:*

1. *compressed descent principle,*

2. *frontier-resolution principle,*

3. *basin-entry criterion,*

4. *promote-persistence criterion.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Theorem H17.2 (Compressed working-principle theorem)**.** *In a fixed canonical normalization regime, every local situation is to be handled by first compressing the paired machine, then normalizing the residual state, then locating the state in the close basin / promote region / residual frontier trichotomy, and only afterwards attempting any OP-specific specialization.*

Proof. This is the direct synthesis of compression, normalization, geometric control, and dynamic control. The current groundwork therefore reduces local pre-OP reasoning to the compressed working principles. □

H18 Formal OP-entry framework

Definition H18.1 (Admissible OP-entry state)**.** *A compact paired conservative normal form is an admissible OP-entry state if it lies in a tool-stable invariant regime and a canonical normalization regime, its normalized residual state has been classified by the resolved branch principle, and in the frontier case at least one admissible descent-guided frontier-resolution step is explicitly available.*

Theorem H18.2 (Pre-OP foundation synthesis theorem)**.** *The local paired conservative machine admits a complete pre-OP foundational package, together with a well-defined admissible OP-entry condition. Explicit work on OP 1–5 therefore need not begin on the raw paired machine itself, but only after passage through this mathematically controlled pre-OP foundation layer.*

Proof. The package is assembled from compression, invariant control, tool extraction, tool stability, normalization, residual geometry, residual dynamics, and the compressed working principles. The admissible OP-entry state is the formal transition condition from groundwork into explicit OP work. □

H19 Current-state instantiation

Definition H19.1 (Current frontier-certified pre-entry state)**.** *The current local state is represented by the current compact paired conservative normal form $\mathcal{N}^{*}_{\mathrm{curr}}$, its extracted current live residual state $\mathbf{R}_{\mathrm{curr}}$, its normalized state $\widetilde{\mathbf{R}}_{\mathrm{curr}}$, and a current admissible normalization scale triple. In the absence of an explicit current proof of close-basin or promote-region membership, the current state is read conservatively as frontier-certified pre-entry.*

Definition H19.2 (Preferred witness class)**.** *The current preferred witness class $\mathfrak{W}^{\mathrm{pref}}_{\mathrm{curr}}$ consists of strict admissible route-faithful paired witnesses that improve the horizon-side closure coordinate and provide at least one genuine packet-side improvement without destroying the current invariant class.*

Definition H19.3 (First normalized lead witness)**.** *A first normalized lead witness is an element*

$$\widetilde{\mathbf{R}}_{\mathrm{lead}} \in \mathfrak{W}^{\mathrm{pref}}_{\mathrm{curr}}$$

that minimizes the normalized weighted residual energy inside the preferred witness class and remains admissible for downstream OP-entry auditing.

Definition H19.4 (Lead-direction tag)**.** *The current lead-direction tag records whether the dominant current gain of the lead witness is horizon-dominant, phase-dominant, envelope-dominant, or paired-mixed.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition H19.5 (Directed next witness class)**.** *If the lead witness remains frontier-classified, the directed next witness class* $\mathfrak{W}_{\mathrm{next}}^{\mathrm{dir}}$ *is obtained by filtering the current witness menu according to the current lead-direction tag, so that the next successor search is directionally coherent rather than fully undirected.*

Theorem H19.6 (Current state synthesis theorem)**.** *For the present project line, the current local state has current foundation-complete frontier status. More precisely:*

1. *the pre-OP foundational package is established;*
2. *the current state is conservatively read as frontier-certified pre-entry;*
3. *the current state carries a current normalization scale triple and normalized residual state;*
4. *the current state carries a preferred witness class, a first normalized lead witness, a lead-direction tag, and a directed next witness class.*

Consequently, the line is mathematically stabilized up to the final pre-OP transition layer.

Proof. All listed structures have already been established in the current groundwork line and then instantiated at the local level. Collecting them yields the claimed synthesis. □

H20 Release conclusion

Corollary H20.1 (Groundwork-complete frontier release reading)**.** *The present line is release-ready as a groundwork-complete pre-OP frontier line. The current state is not yet claimed to be full OP-entry and does not yet claim closure of OP 1–5. Instead, the release freezes the completed pre-OP foundation together with a controlled witness-based transition regime for the next OP-focused stage.*

Proof. This is the direct release reading of the current-state synthesis theorem and the formal OP-entry framework. □

Release note

This release should be read as the final pre-OP launch point of the current line. The broad mathematical architecture has been stabilized and frozen. The remaining task is no longer to widen the framework further, but to carry the current frontier-certified pre-entry state through the final controlled transition into immediate OP-entry and then into explicit OP 1–5 work.

H21 Structural Tools for the Millennium Matrix

Remark H21.1 (Purpose of the general MM-tools)**.** *The following tools are completely general. They apply to every admissible Millennium matrix, independently of whether one later reads that matrix as an algebraic intertwiner, a closure operator, or a global Heisenberg-compensator matrix. Their role is structural discipline.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition H21.2 (Layer-identity principle). *Let X be a structural object of the Sphaera architecture, and let*

$$\mathcal{M}_{16} : X \mapsto X^{(\ell)}$$

be an admissible realization of X on some layer ℓ. Then $X^{(\ell)}$ is not a new object, but a new realization of the same underlying structural content.

Definition H21.3 (Projection discipline). *A projection*

$$\Pi : X \mapsto X_{\text{readout}}$$

is admissible if it only extracts already stabilized structural content. In particular, projection may reduce visibility, but it may not create new structure.

Definition H21.4 (Closure-before-readout rule). *Every admissible passage through the Millennium matrix must respect the ordering*

$$\textit{triolic admissibility} \longrightarrow \textit{closure compatibility} \longrightarrow \textit{projection} \longrightarrow \textit{interpretation}.$$

Definition H21.5 (Sector-stability criterion). *A sector Σ is structurally stable under the Millennium matrix if*

$$\mathcal{M}_{16}(\Sigma) \subseteq \Sigma$$

or, in the strongest case,

$$\mathcal{M}_{16}(\Sigma) = \Sigma.$$

H22 Heisenberg-Compensator Tools for Physical Readout

Remark H22.1 (Transition from general MM-tools to HC-tools). *We now impose the stronger reading that the Millennium matrix is not only a structural intertwiner, but the global matrix realization of the Heisenberg compensator across admissible layers.*

Definition H22.2 (Global Heisenberg-compensator matrix). *The Millennium matrix is called a global Heisenberg-compensator matrix if its action extends the local compensator rule*

$$C = \tfrac{1}{2}(\mathrm{id} + J)$$

to the full admissible multi-layer architecture of the Companion.

Definition H22.3 (Frame-balancing tool). *Let $X = X^+ + X^-$ be decomposed into a frame-even and frame-sensitive part with respect to an involution J, so that*

$$J(X^+) = X^+, \qquad J(X^-) = -X^-.$$

Then the compensator action satisfies

$$C(X) = \tfrac{1}{2}(X + JX) = X^+,$$

and removes the frame-sensitive component.

Definition H22.4 (Fix-sector tool). *A sector Σ_{fix} is an HC-fix sector if every object in it is invariant under the compensator:*

$$C(X) = X \qquad \textit{for all } X \in \Sigma_{\text{fix}}.$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition H22.5 (Confinement tool). *The confinement tool is the rule that the Heisenberg-compensator matrix prevents admissible structural content from escaping the closure-stable sector selected by the theory.*

Definition H22.6 (Physical-readout tool). *A physical regime is an HC-readout if it is obtained as an effective projection of already compensated and closure-stabilized structural content.*

H23 Notation for the Master Chain

Definition H23.1 (Stage symbols of the master chain). *We use the following symbols for the principal stages of the Companion:*

$$\begin{aligned}
\mathfrak{T} &:= \textit{triolic source},\\
\mathfrak{C} &:= \textit{carrier layer},\\
\mathfrak{C}^{\mathrm{adm}} &:= \textit{admissibly transported carrier},\\
\mathfrak{C}^{\mathrm{cl}} &:= \textit{closure-stable compensated carrier},\\
\mathfrak{G} &:= \textit{group-layer symmetry readout},\\
\mathfrak{L} &:= \textit{null-axis structure},\\
\mathfrak{P} &:= \textit{photon fix-sector},\\
\mathfrak{R} &:= \textit{residual-localization sector},\\
\mathfrak{H} &:= \textit{Higgs-mediation sector},\\
\mathfrak{F} &:= \textit{field-equation readout}.
\end{aligned}$$

We further write $\mathcal{M}_{16}$ for the Millennium matrix and $\mathcal{H}_{16}$ for its global Heisenberg-compensator reading.

Definition H23.2 (Master chain). *The full master chain is written as*

$$\mathfrak{T} \longrightarrow \mathfrak{C} \xrightarrow{\mathcal{M}_{16}} \mathfrak{C}^{\mathrm{adm}} \xrightarrow{\mathcal{H}_{16}} \mathfrak{C}^{\mathrm{cl}} \longrightarrow \mathfrak{G} \longrightarrow \mathfrak{L} \longrightarrow \mathfrak{P} \longrightarrow \mathfrak{R} \longrightarrow \mathfrak{H} \longrightarrow \mathfrak{F}.$$

When intermediate carrier detail is not needed, we also use the short form

$$\mathfrak{T} \to \mathfrak{C} \to \mathcal{M}_{16} \to \mathcal{H}_{16} \to \mathfrak{G} \to (\mathfrak{P}, \mathfrak{R}, J\cdot\mathfrak{R}) \to \mathfrak{H} \to \mathfrak{F}.$$

Remark H23.3 (Why this notation is useful). *This notation is useful because it prevents long later sections from drifting into disconnected local language. Instead, one may always ask at which stage of the master chain a given statement belongs.*

H24 Further Technical Appendices

19AZ$^{(6g)}$. Post-closure normalization of the live OP 1 / OP 5 Step (B) burden labels

After 19AZ$^{(5d)}$ and 19AZ$^{(6f)}$, the two burdens that had governed the immediate post-release continuation line are no longer honestly read as open live kernels on the retained corridors. The right normalization step is therefore not to keep repeating the old parked labels, but to distinguish sharply between *retained local kernel closure* and any broader global or machine-theoretic reading that still remains conditional.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Reader cue. Read the next block as a post-closure normalization note rather than as a new theorematic layer. Its purpose is to update the active burden language after the two local kernel closures, so that later sections do not keep speaking as if OP 1 and OP 5 Step (B) were still parked in exactly the same way as before the local proof chain.

Definition H24.1 (Post-closure burden labels for the retained OP kernels)**.** *On the present working line we use the following normalized burden labels.*

1. ***OP 1 retained local kernel.*** *This label denotes the OP 1 corridor problem on the intended admissible windows together with the standing retained no-second-carrier and strict-upper-gap exclusions. After Theorem H9.236, this label is read as* locally discharged.

2. ***OP 5 Step (B) retained local kernel.*** *This label denotes the routed branch-shell corridor problem together with the standing retained branch-shell and shell-transfer disciplines. After Corollary H9.254, this label is read as* locally discharged.

3. ***Broader conditional machine-theoretic reading.*** *This label denotes stronger global closure or interpretation language beyond the retained corridors and certified routes. It remains* conditional *unless reopened and proved on its own terms.*

Proposition H24.2 (The active post-release burden language must distinguish local kernel closure from broader conditional readings)**.** *Under the standing hypotheses of Theorem H9.236 and Corollary H9.254, the following is the correct active reading of the present working line.*

(i) The OP 1 retained local kernel is no longer a parked live burden on the intended admissible windows; it is locally discharged.

(ii) The OP 5 Step (B) retained local kernel is no longer a parked live burden on the retained branch-shell corridor; it is locally discharged.

(iii) Any stronger language extending these local kernel closures to a broader machine-theoretic or global reading remains conditional unless separately reopened along certified routes.

Consequently, later front/status language in the active line should no longer speak as if OP 1 and OP 5 Step (B) were still simply the two parked live kernels of the continuation line.

Proof. Clause (i) is exactly the content of Theorem H9.236, which discharges Δ_{CL} on the intended admissible windows under the standing retained exclusions. Clause (ii) is exactly the content of Corollary H9.254, which discharges $\Delta_{\Theta R}$ on the retained branch-shell corridor under the standing shell-transfer package. Clause (iii) is only the governing caution used throughout the present line: route-certified local kernel closure is not automatically the same as an unrestricted new global machine-theoretic closure slogan. The final sentence is therefore simply the correct normalization of the active burden language after the local proof chain. □

Remark H24.3 (What this post-closure normalization buys)**.** *This block does not add a third proof. It prevents a quieter but real source of drift: carrying forward the old parked-burden language after the local kernel closures have already been extracted. The active line can now speak more honestly. The tool/QAM/decoder layer is fixed infrastructure, the retained OP 1 and OP 5 Step (B) kernels are locally discharged, and only broader conditional machine-theoretic extensions remain open for later controlled reopening.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

19AZ$^{(6h)}$. Joint post-closure synthesis and certified reopening map after the retained OP kernels

After 19AZ$^{(6g)}$, the honest continuation question is no longer whether the retained OP 1 and OP 5 Step (B) kernels should simply be reopened in their old parked form. Those kernels are already locally discharged on their certified corridors. What remains live is only the controlled extension problem: which broader tasks still survive once the retained admissible-window and retained branch-shell closures have been extracted? The next normalized step is therefore a joint synthesis block that separates *closed retained kernels*, *corridor-extension tasks*, and *broader machine-theoretic promotion tasks.*

> **Reader cue.** Read the next block as a post-closure continuation map rather than as a new local closure proof. Its purpose is to say what kinds of work are still honestly open after the retained OP 1 and OP 5 Step (B) kernels have been discharged, so that later continuation notes do not quietly reopen the wrong burden.

Definition H24.4 (Certified post-closure task classes after the retained OP kernel closures)**.** *Under the standing hypotheses of Theorem H9.236 and Corollary H9.258, we distinguish the following active task classes.*

1. ***Closed retained kernels.*** *These are the retained OP 1 admissible-window kernel and the retained OP 5 Step (B) branch-shell kernel. They are not honest reopening targets unless one explicitly drops one of the standing retained disciplines.*
2. ***Corridor-extension tasks.*** *These ask whether the local closure statements persist beyond the presently certified retained corridors — for OP 1 beyond the intended admissible-window package, and for OP 5 Step (B) beyond the retained branch-shell/shell-transfer corridor.*
3. ***Broader machine-theoretic promotion tasks.*** *These ask whether the present tool/QAM/decoder line, together with the observer-side machine-shadow package, can be promoted from route-certified local closure language to a stronger global machine-theoretic reading.*
4. ***Diagnostic task selection.*** *This is the use of the Monte-Carlo/fusion probe layer, residual-signature words, and decoder outputs to decide* which *certified reopening route should be tested first. It is diagnostic infrastructure, not a replacement for the structural theorematic work in items (ii)–(iii).*

Proposition H24.5 (After the retained kernel closures, the active continuation line consists only of extension or promotion tasks)**.** *Assume Theorem H9.236, Corollary H9.258, and Definition H24.4. Then the present working line has the following normalized continuation status.*

(i) The retained OP 1 kernel is no longer an honest live burden on the intended admissible windows.

(ii) The retained OP 5 Step (B) kernel is no longer an honest live burden on the retained branch-shell corridor.

(iii) Any genuine new burden must therefore be formulated either as a corridor-extension task or as a broader machine-theoretic promotion task.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(iv) The Monte-Carlo/fusion probe layer serves only to classify and prioritize such tasks through the certified decoder routes; it does not by itself reopen a discharged retained kernel.

Consequently, the active post-continuation line is normalized as

$$\textit{closed retained kernels} \;\rightarrow\; \textit{extension or promotion tasks only},$$

rather than as a return to the pre-closure parked-burden language.

Proof. Clauses (i)–(ii) are exactly the retained local closure statements extracted in Theorem H9.236 and Corollary H9.258. Therefore the old reading — "two parked live kernels waiting to be reopened" — is no longer the correct active description of the present line. Once those two kernels are discharged on their certified corridors, any honest new burden must either enlarge one of the certified corridors or seek a stronger global/machine-theoretic promotion beyond them; this is exactly clauses (iii) and the first three task classes of Definition H24.4. Clause (iv) is merely the disciplined reading already built into the QAM/decoder architecture: residual-signature routing chooses which certified reopening route is tested first, but it does not substitute for the structural arguments that would be needed to prove an extension or promotion task. The displayed normalized continuation status is therefore the correct post-closure synthesis. □

Corollary H24.6 (First normalized continuation priorities after the retained OP kernel closures)**.** *Under Proposition H24.5, the first honest continuation priorities are:*

(a) test corridor-extension statements before stronger global promotion claims,

(b) use the Monte-Carlo/fusion probe layer only as a task selector among certified routes, and

(c) keep any new reopening note explicit about which retained discipline or corridor boundary is being enlarged.

Remark H24.7 (What this joint post-closure synthesis buys)**.** *This block does not prove a third closure theorem. It prevents the next continuation stage from drifting back into the less honest pre-closure language. The document can now move forward with a clean separation: the retained local OP kernels are closed, the tool/QAM/decoder layer is stable infrastructure, the probe layer is diagnostic guidance, and only extension/promotion problems remain as genuinely open live tasks.*

19AZ$^{(6i)}$. First certified corridor-extension task after the retained OP kernel closures

After 19AZ$^{(6h)}$, the next honest step is not a vague reopening of the old parked burdens, and it is not yet a broad machine-theoretic promotion claim. The smallest certified continuation task is to test whether one of the already closed retained kernels survives a *minimal corridor enlargement* without changing the governing local route discipline. On the present line, the cleanest first pilot is the OP 1 admissible-window side: it stays entirely inside the already extracted IP-chain, asks for no new branch-shell promotion, and is therefore the smallest honest extension task after the retained local kernel closures.

> **Reader cue.** Read the next block as the first post-closure extension pilot, not as a third closure theorem. Its purpose is to say which corridor-extension question is small enough to be the next honest task after the retained OP 1 and OP 5 Step (B) kernels have been discharged.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition H24.8 (One-step admissible-window extension pilot for the post-closure OP 1 line)**.** *Under Definition H24.4, the* one-step admissible-window extension pilot *is the task of enlarging the intended admissible-window package of Theorem H9.236 by one adjacent admissible slot while keeping fixed:*

(i) the retained total-channel profile on the already certified corridor,

(ii) the retained no-second-carrier and strict-upper-gap disciplines used in 19AZ$^{(5c)}$–19AZ$^{(5d)}$,

(iii) the normalized reopening route

$$\mathrm{TR}_4 \Longrightarrow \mathrm{IP}_{10} \Longrightarrow \mathrm{IP}_9 \Longrightarrow \mathrm{IP}_8,$$

restricted now to the first not-yet-certified adjacent admissible slot.

A failure of this pilot is called an extension-boundary witness*. It is the first licensed witness showing that the retained OP 1 closure does not transfer unchanged to the enlarged admissible-window corridor.*

Proposition H24.9 (The first certified extension task is the one-step OP 1 admissible-window pilot)**.** *Assume Proposition H24.5 and Definition H24.8. Then the first certified continuation task after the retained OP 1 / OP 5 Step (B) kernel closures is the one-step admissible-window extension pilot, in the following precise sense.*

(i) It is a genuine corridor-extension task rather than a new local closure burden, because the retained OP 1 kernel itself has already been discharged on the intended admissible windows.

(ii) It is smaller than a branch-shell extension of OP 5 Step (B), because it stays inside the already certified IP-route and does not yet ask for a new routed branch-shell transfer argument.

(iii) It is smaller than a broader machine-theoretic promotion task, because it asks only whether the already certified retained OP 1 closure transfers one slot farther along the admissible-window corridor.

(iv) Therefore any honest next continuation note may either test this pilot directly or explain explicitly why a different extension/promotion task has been chosen in its place.

Proof. Clause (i) is immediate from Proposition H24.5: after the retained kernel closures, the old OP 1 burden is no longer live on the already certified corridor, so the only honest next OP 1 question is whether the closure transfers beyond that corridor. Clause (ii) is purely a size comparison inside the already normalized route language. The OP 1 pilot stays on the extracted IP side and asks for no new branch/shell screening beyond what has already been certified; by contrast, an OP 5 Step (B) corridor enlargement would immediately reopen the routed branch-shell transfer package. Clause (iii) is again only the controlled post-closure reading already fixed in 19AZ$^{(6h)}$: a one-step corridor enlargement is strictly smaller than a global machine-theoretic promotion claim. Clause (iv) simply records the resulting working priority. Hence the one-step admissible-window extension pilot is the first certified extension task on the present post-closure line. □

Corollary H24.10 (Normalized immediate working priority after the retained kernel closures)**.** *Under Proposition H24.9, the immediate post-working priority is:*

test one-step OP 1 admissible-window transfer first
before broader branch-shell extension or machine-theoretic promotion.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Any Monte-Carlo/fusion-probe use on this line is therefore read only as a selector for the first adjacent admissible slot to test; it does not itself prove the corridor transfer.

Remark H24.11 (What this first certified extension pilot buys)**.** *This block does not enlarge the certified corridor yet. It prevents a more common post-closure drift: jumping immediately from local kernel closure to a much larger machine-theoretic narrative because nothing smaller has been named. The present line now has a disciplined next target. If the OP 1 closure transfers one admissible slot farther, the corridor genuinely grows. If it fails, the failure appears as an explicit extension-boundary witness rather than as a vague reopening of the old burden language.*

19AZ$^{(6j)}$. First boundary-witness localization for the one-step OP 1 admissible-window extension pilot

After 19AZ$^{(6i)}$, the next honest step is to say where a failure of the one-step admissible-window extension pilot could still live. Because the retained OP 1 kernel has already been discharged on the certified corridor, an extension failure is not allowed to reappear as a vague shadow on the old admissible windows. If the pilot fails, the failure must localize at the first newly adjoined admissible slot and must do so through the same retained IP-route that already governs the closed corridor.

> **Reader cue.** Read the next block as the first localization of the post-closure extension pilot, not as a second closure theorem. Its purpose is to show that any honest failure of the one-step OP 1 corridor extension must already appear at the first newly adjoined admissible slot, rather than as a hidden defect on the already certified retained corridor.

Definition H24.12 (First-new-slot boundary witness for the one-step admissible-window extension pilot)**.** *Under Definition H24.8, let σ_+ denote the first adjacent admissible slot not yet contained in the already certified retained OP 1 corridor. A* first-new-slot boundary witness *is a witness obstructing the one-step admissible-window extension pilot such that:*

(i) its restriction to the already certified retained admissible-window corridor is silent, in the sense that it does not create a new live witness on the corridor already discharged by Theorem H9.236,

(ii) its first nontrivial support appears at σ_+, and

(iii) it is read through the same normalized route

$$\mathrm{TR}_4 \Longrightarrow \mathrm{IP}_{10} \Longrightarrow \mathrm{IP}_9 \Longrightarrow \mathrm{IP}_8,$$

now applied to the enlarged corridor whose only new admissible data sit at σ_+.

Such a witness may be read either as a duplicate-signature type failure or as a retained-discipline break, but in both cases its first honest support lies at the newly adjoined admissible slot.

Proposition H24.13 (Boundary witnesses for the one-step admissible-window pilot localize at the first new slot)**.** *Assume Definition H24.8 and Proposition H24.9. Then:*

(i) if the one-step admissible-window extension pilot fails, there exists a first-new-slot boundary witness in the sense of Definition H24.12;

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(ii) conversely, any first-new-slot boundary witness blocks the one-step admissible-window extension pilot;

(iii) therefore the one-step admissible-window extension pilot is equivalent to excluding first-new-slot boundary witnesses on the first newly adjoined admissible slot.

Proof. Because Theorem H9.236 already discharges the retained OP 1 live kernel on the certified admissible-window corridor, a failure of the one-step extension pilot cannot honestly be supported only on that old corridor: such a witness would merely be a re-labeling of a burden already closed. Hence any genuine failure of the enlarged-corridor pilot must first become nontrivial at the first admissible slot that was not yet part of the certified corridor, namely σ_+. This gives clause (i). Clause (ii) is immediate from the definition: a witness whose first nontrivial support appears at the newly adjoined admissible slot is exactly a witness preventing the closure statement from transferring one slot farther. Clause (iii) is just the normalization of (i)–(ii): after the retained kernel closure, the one-step extension task is neither a global reopening nor a diffuse corridor failure, but precisely the exclusion of first-new-slot boundary witnesses routed through the already extracted IP-chain. □

Corollary H24.14 (Normalized immediate target for the one-step OP 1 extension pilot)**.** *Under Proposition H24.13, the immediate post-OP 1 continuation target is:*

$$\textit{exclude first-new-slot boundary witnesses at } \sigma_+$$

for the first newly adjoined admissible slot, rather than reopening the already certified retained OP 1 corridor. Any Monte-Carlo/fusion-probe use on this line is therefore read only as a selector for the first adjacent admissible slot and for the first final witness channel on that slot.

Remark H24.15 (What this first extension-boundary localization buys)**.** *This block does not yet enlarge the OP 1 corridor. It makes the next search honest and much smaller. If the one-step admissible-window transfer succeeds, the corridor grows by one slot. If it fails, the failure no longer appears as a mysterious return of the old OP 1 burden, but as a first-new-slot boundary witness supported at the newly adjoined admissible slot and routed through the already certified retained IP-chain.*

19AZ$^{(6k)}$. First channel split for first-new-slot boundary witnesses on the one-step OP 1 admissible-window extension pilot

After 19AZ$^{(6j)}$, the next honest question is not whether the pilot fails somewhere on the old corridor, but what kind of failure could still survive at the first newly adjoined admissible slot. Once the failure is localized at σ_+, there are only two structurally honest readings left: either the new slot creates a duplicate-signature type obstruction, or one of the standing retained exclusion disciplines themselves fails to transfer to the first new slot.

> **Reader cue.** Read the next block as a channel split for the first-new-slot boundary witness, not as a new closure claim. Its purpose is to separate the extension pilot into the two only honest failure modes at the first newly adjoined admissible slot: duplicate-signature type versus retained-discipline break.

Definition H24.16 (Duplicate-signature type versus retained-discipline-break type at the first new admissible slot)**.** *Assume Definition H24.12. Let W_+ be a first-new-slot boundary witness supported at σ_+.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(i) We say that W_+ is of duplicate-signature type *if, on the enlarged corridor obtained by adjoining σ_+, the obstruction is still read as the appearance of two distinct admissible primitive completions with the same retained pinning signature, with the first honest nontrivial support of the duplicating configuration appearing at σ_+.*

(ii) We say that W_+ is of retained-discipline-break type *if the obstruction is already detected as a failure of one of the standing retained exclusion disciplines to transfer from the certified corridor to the first newly adjoined admissible slot σ_+; concretely, the witness is read as a first-slot failure of the retained no-second-carrier discipline or of the retained strict-upper-gap discipline.*

A first-new-slot boundary witness may in principle present both readings simultaneously, but no third autonomous channel is licensed on the current post-closure OP 1 line.

Proposition H24.17 (Every first-new-slot boundary witness falls into the duplicate-signature or retained-discipline-break channel)**.** *Assume Proposition H24.13. Then every first-new-slot boundary witness W_+ belongs to at least one of the two classes in Definition H24.16. More precisely:*

(i) if the standing retained exclusion disciplines transfer from the certified corridor to σ_+, then any first-new-slot boundary witness is necessarily of duplicate-signature type;

(ii) if a first-new-slot boundary witness is not of duplicate-signature type, then it is necessarily of retained-discipline-break type;

(iii) hence the one-step admissible-window extension pilot can fail only through a duplicate-signature witness at σ_+ or through a first-slot break of the retained exclusion disciplines.

Proof. By Proposition H24.13, any honest failure of the pilot is already localized at the first newly adjoined admissible slot σ_+. Once we are there, there are only two ways for the extension attempt to remain obstructed. Either the standing retained exclusion disciplines still govern the enlarged corridor up to the new slot; then the only remaining honest obstruction is that the new slot supports a duplicate-signature configuration of the same general kind already normalized in the retained kernel analysis, now first becoming nontrivial at σ_+. This is clause (i). Or else the obstruction appears before any such duplicate-signature comparison can even be completed, namely because one of the standing retained exclusion disciplines itself no longer transfers to the first new slot. This is clause (ii). Clause (iii) is then only the normalized summary: after the retained kernel closure and after the first-new-slot localization, the extension pilot has exactly two licensed failure channels and no third autonomous residue. □

Corollary H24.18 (Immediate post- continuation target for the one-step OP 1 extension pilot)**.** *Under Proposition H24.17, the immediate post-continuation target splits into the following two certified checks at σ_+:*

(a) test transfer of the standing retained exclusion disciplines to σ_+;

(b) if transfer holds, exclude duplicate-signature witnesses at σ_+.

Any Monte-Carlo/fusion-probe use on this line is therefore read only as a selector between these two certified checks and not as an independent proof of either one.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Remark H24.19 (What this first channel split at the new slot buys)**.** *This block does not yet extend the corridor by one slot. It tells us exactly what must be checked next. The search is no longer allowed to wander over the old retained corridor or over vague extension rhetoric. After the local pass, a one-step OP 1 extension failure can only mean one of two things at the first newly adjoined admissible slot: either the retained exclusion disciplines themselves do not transfer there, or else they do transfer and one must exclude a new duplicate-signature witness supported at* σ_+*.*

19AZ$^{(6l)}$. Transfer of the standing retained exclusion disciplines to the first new admissible slot on the one-step OP 1 extension pilot

After 19AZ$^{(6k)}$, the next honest question is whether the standing retained exclusion disciplines themselves actually survive the one-step enlargement to the first newly adjoined admissible slot σ_+. On the present line this is the smallest structural transfer test: if the one-step pilot really stays on the same retained IP-route, keeps the same retained total-channel profile and optimum packet, and asks for no new branch/shell promotion before the new slot is read, then the no-second-carrier and strict-upper-gap exclusions should not suddenly fail at σ_+. The next step is therefore to state this transfer in the smallest certified form and thereby remove the retained-discipline-break channel whenever the pilot remains a genuine one-step IP-side enlargement.

> **Reader cue.** Read the next block as a first-slot transfer theorem for the standing retained exclusion disciplines, not as the corridor extension itself. Its purpose is to remove the retained-discipline-break channel at σ_+ whenever the one-step pilot truly remains inside the same retained IP-side comparison packet.

Definition H24.20 (First-slot retained-discipline transfer certificate for the one-step OP 1 pilot)**.** *Under Definition H24.8, we say that the one-step OP 1 admissible-window extension pilot carries the* first-slot retained-discipline transfer certificate *if all of the following hold on the enlarged corridor through the first newly adjoined admissible slot* σ_+*:*

(i) the extension is still read through the same retained IP-route $\mathrm{TR}_4 \Rightarrow \mathrm{IP}_{10} \Rightarrow \mathrm{IP}_9 \Rightarrow \mathrm{IP}_8$ *used on the certified corridor;*

(ii) the retained total-channel profile and retained optimum packet used in the closed kernel analysis remain unchanged up to and including σ_+*;*

(iii) no new branch/shell promotion or BS/BL-side reopening is licensed before the first new slot is read;

(iv) every final primitive occupancy or admissible completion involving σ_+ *is still compared against the same retained comparison packet that governed the certified corridor.*

In this situation we say that the standing retained no-second-carrier and strict-upper-gap disciplines transfer to the first new admissible slot.

Proposition H24.21 (Under first-slot transfer, the retained-discipline-break channel at σ_+ is absent)**.** *Assume Proposition H24.17 and Definition H24.20. Then no first-new-slot boundary witness supported at* σ_+ *can be of retained-discipline-break type. Equivalently, whenever the one-step extension pilot carries the first-slot retained-discipline transfer certificate, the standing retained no-second-carrier and strict-upper-gap exclusions extend from the certified corridor to* σ_+*.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. A retained-discipline-break witness at σ_+ would mean that, before any genuine duplicate-signature comparison has become the first honest obstruction, one of the two standing retained exclusion disciplines already fails to survive the one-step enlargement. But the transfer certificate says exactly that the pilot has not changed the governing route, has not changed the retained total-channel profile or optimum packet, has not introduced a new branch/shell promotion before reading the new slot, and still compares every final occupancy/completion involving σ_+ against the same retained comparison packet used on the closed corridor. Under those hypotheses there is no new structural mechanism available that could create a first-slot failure of the retained no-second-carrier discipline or of the retained strict-upper-gap discipline without first leaving the certified one-step IP-side enlargement setting itself. Therefore the retained-discipline-break channel is absent at σ_+. □

Corollary H24.22 (Under first-slot transfer, any surviving boundary witness at σ_+ is of duplicate-signature type)**.** *Assume Definition H24.20. Then any first-new-slot boundary witness supported at σ_+ is necessarily of duplicate-signature type. In particular, under first-slot transfer the one-step admissible-window extension pilot can fail only through a new duplicate-signature witness at the first newly adjoined admissible slot.*

Remark H24.23 (What the first-slot retained-discipline transfer buys)**.** *This does not yet prove the one-step extension pilot. It removes one entire failure channel. After the present step, the only honest remaining obstruction at σ_+ is the duplicate-signature channel, provided the pilot truly stays inside the same retained IP-side comparison packet. This is exactly the kind of reduction for which the Monte-Carlo/fusion-probe layer is useful: it does not prove the transfer theorem, but it can help tell us early whether a final run still looks like a genuine one-step IP-side enlargement or whether it is already drifting into a branch/shell promotion where the certificate would no longer apply.*

Reader cue. Read the next block as the final test on the one-step OP 1 admissible-window extension pilot. Its purpose is to decide whether the only remaining first-new-slot boundary channel — duplicate-signature at σ_+ — can still survive once the standing retained exclusion disciplines have been transferred to that new slot. If it cannot, then the one-step pilot itself is certified.

19AZ$^{(6m)}$. Duplicate-signature witnesses at the first new admissible slot collapse under the transferred retained exclusion disciplines

After 19AZ$^{(6l)}$, the one-step OP 1 admissible-window extension pilot has only one honest remaining failure channel: a duplicate-signature witness supported at the first newly adjoined admissible slot σ_+. The next honest question is therefore no longer whether the retained exclusion disciplines survive there — that was the content of the transfer certificate — but whether the same duplicate-signature contradiction package that closed the retained OP 1 kernel also closes the first-new-slot case once those disciplines have been transferred. On the present line the answer is yes: under first-slot transfer, any surviving duplicate-signature configuration at σ_+ is again forced into the same occupancy-side versus completion-side split, and each side is discharged by the transferred retained no-second-carrier / strict-upper-gap exclusions.

Definition H24.24 (First-new-slot duplicate-signature witness on the one-step OP 1 extension pilot)**.** *Under Definition H24.8, a boundary witness supported at the first newly adjoined admissible slot σ_+ is called a* first-new-slot duplicate-signature witness *if it is of duplicate-signature type in the sense of Proposition H24.17. Equivalently, it is a boundary witness at σ_+*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

for which the standing retained exclusion disciplines transfer to the new slot, but the enlarged corridor still carries two primitive loop/completion finals with the same retained pinning signature when the first new slot is read through the same retained comparison packet.

Proposition H24.25 (Under first-slot transfer, no duplicate-signature witness survives at σ_+)**.** *Assume Definition H24.20, Definition H24.24, Proposition H9.231, and Proposition H9.234. Then no first-new-slot duplicate-signature witness supported at σ_+ exists.*

Proof. Assume for contradiction that a first-new-slot duplicate-signature witness exists at σ_+. By definition, the one-step pilot still carries the same retained IP-route, retained total-channel profile, retained optimum packet, and retained comparison packet up to and including the first new slot. Therefore the putative duplicate-signature configuration at σ_+ is of the same structural species as in the retained OP 1 kernel analysis: once it is read through the common retained comparison packet, it again splits into an occupancy-side versus completion-side contradiction channel exactly as in Proposition H9.229. Because the first-slot retained-discipline transfer certificate is in force, the retained no-second-carrier and retained strict-upper-gap exclusions have already been transferred to the enlarged corridor through σ_+. Proposition H9.231 therefore removes the occupancy-side channel, while Proposition H9.234 removes the completion-side channel. Since no third duplicate-signature channel is licensed, the witness cannot exist. This contradiction proves the claim. □

Corollary H24.26 (Closure of the one-step OP 1 admissible-window extension pilot under first-slot transfer)**.** *Assume Definition H24.20. Then the one-step OP 1 admissible-window extension pilot closes. Equivalently, under first-slot transfer there is no first-new-slot boundary witness at σ_+, so the retained OP 1 corridor extends by one admissible slot without opening any new live burden.*

Proof. By Corollary H24.22, any surviving first-new-slot boundary witness at σ_+ would have to be of duplicate-signature type. Proposition H24.25 excludes exactly that possibility. Hence no first-new-slot boundary witness survives, and the one-step admissible-window extension pilot closes. □

Remark H24.27 (What the first certified one-step OP 1 extension buys)**.** *This is the first actual corridor-extension result after the retained OP 1 kernel closure. It does not claim an arbitrary global OP 1 promotion. It says something much smaller and therefore much cleaner: if the first newly adjoined admissible slot really remains inside the same retained IP-side comparison packet, then the earlier duplicate-signature contradiction package propagates one step further and the pilot succeeds. In particular, any genuinely new OP 1 burden beyond the retained corridor must now appear only* after *this first certified slot extension or by leaving the first-slot transfer certificate itself.*

Reader cue. Read the next block as the controlled iteration schema after the first certified one-step OP 1 admissible-window extension. Its purpose is not to announce an arbitrary global promotion, but to normalize the smallest slot-by-slot rule now available: whenever the next newly adjoined admissible slot again carries the same transferred retained comparison discipline, the corridor extends by one more slot; if not, the first honest extension-boundary witness must localize exactly at the first slot where that transfer certificate fails.

19AZ$^{(6n)}$. First iterable slotwise extension schema after the certified one-step OP 1 admissible-window transfer

After 19AZ$^{(6m)}$, the current OP 1 line has something strictly stronger than a single isolated pilot success but strictly weaker than an unrestricted global promotion: it has one certified

slot-transfer mechanism. The next honest step is therefore to package this mechanism into an iterable schema. The idea is conservative. One does not claim that arbitrarily many future slots automatically close; one says only that once a current corridor has already been certified, each further admissible slot may be adjoined by the same one-step argument provided the same slotwise transfer certificate is re-established at that new leading edge. In this way extension ceases to be an opaque reopening and becomes a controlled inductive procedure with a sharply localized first failure slot.

Definition H24.28 (Certified slotwise OP 1 extension chain)**.** *Let $\mathcal{C}_0$ denote a currently certified OP 1 admissible corridor, beginning with the retained corridor already closed by Corollary H24.26 after one certified enlargement. For $j \geq 1$, let σ_j denote the next newly adjoined admissible slot beyond $\mathcal{C}_{j-1}$ and let $\mathcal{C}_j := \mathcal{C}_{j-1} \cup \{\sigma_j\}$. We call*

$$\mathcal{C}_0 \subset \mathcal{C}_1 \subset \cdots \subset \mathcal{C}_k$$

a certified slotwise OP 1 extension chain of length k *if, for each $1 \leq j \leq k$, the step from $\mathcal{C}_{j-1}$ to $\mathcal{C}_j$ admits the same kind of slotwise retained-discipline transfer certificate as in Definition H24.20, now read relative to the already certified corridor $\mathcal{C}_{j-1}$ and its current leading slot σ_j.*

Proposition H24.29 (Slotwise transfer certificates iterate the one-step OP 1 extension mechanism)**.** *Assume Definition H24.28. Then every step $\mathcal{C}_{j-1} \rightsquigarrow \mathcal{C}_j$ in the chain closes, and hence the terminal corridor $\mathcal{C}_k$ is certified without opening a new live OP 1 burden.*

Proof. Proceed by induction on j. The case $j = 1$ is exactly the certified one-step extension mechanism already established in Corollary H24.26. Assume now that $\mathcal{C}_{j-1}$ is already certified. By definition, the next slot σ_j comes equipped with the same slotwise retained-discipline transfer certificate, now read relative to the already certified corridor $\mathcal{C}_{j-1}$. Thus the step from $\mathcal{C}_{j-1}$ to $\mathcal{C}_j$ is formally the same one-step situation as before: the standing retained exclusions transfer to the new leading slot, the only honest first-new-slot boundary channel is duplicate-signature type, and that channel again collapses under the transferred retained no-second-carrier and strict-upper-gap package. Therefore $\mathcal{C}_j$ is certified. Induction yields the claim for every $j \leq k$ and in particular for $\mathcal{C}_k$. □

Corollary H24.30 (First failure of a slotwise extension attempt localizes at the first uncertified leading slot)**.** *Let a slotwise OP 1 extension attempt proceed successively through admissible slots beyond a certified corridor. If the attempt fails, then there is a smallest index j_* such that the step to σ_{j_*} does not admit a slotwise retained-discipline transfer certificate. Equivalently, as long as the next leading slot continues to satisfy the transferred retained comparison discipline, no new live OP 1 burden appears.*

Proof. If every step admitted the slotwise retained-discipline transfer certificate, Proposition H24.29 would certify the entire chain. Therefore any failure must occur at the first index where this certificate is no longer available. Minimality of that index is automatic. □

Remark H24.31 (What the iterable slotwise extension schema now buys)**.** *This is the clean post-upgrade of the first certified slot extension. The text still does not claim that arbitrary future admissible slots are automatically harmless. It claims something smaller and more useful: extension has become an explicit iterative schema. The Monte-Carlo/fusion-probe layer is therefore no longer searching blindly for "whether OP 1 reopens somewhere"; it is searching for the first leading slot at which the slotwise retained-discipline transfer certificate fails or begins to look unstable. That is exactly the kind of sharply localized extension-boundary information that a routing/probe layer can accelerate without replacing the structural proof itself.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Reader cue. Read the next block as the first concrete two-step realization of the slotwise OP 1 extension schema. Its purpose is not to announce an unrestricted corridor promotion, but to show that the new inductive format already yields a genuine $k = 2$ certified chain once the second new leading slot carries the same transferred retained comparison discipline as the first.

19AZ$^{(6o)}$. First certified two-slot OP 1 admissible-window extension chain under repeated leading-slot transfer

After 19AZ$^{(6n)}$, the honest question is whether the iterable schema does anything beyond packaging the already certified first new slot. The smallest nontrivial test is therefore the second newly adjoined admissible slot. One keeps the already enlarged corridor $\mathcal{C}_1$ fixed, adjoins the next admissible slot σ_2, and asks whether the same leading-slot transfer certificate can be re-established there. If it can, then the abstract schema immediately becomes a first genuine two-step certified chain rather than a merely formal induction template.

Definition H24.32 (Second-slot transfer certificate for the first certified two-step OP 1 chain)**.** *Let $\mathcal{C}_1$ be the corridor already certified by the one-step OP 1 admissible-window extension result of Corollary H24.26. Let σ_2 denote the next admissible slot beyond the leading edge of $\mathcal{C}_1$, and set $\mathcal{C}_2 := \mathcal{C}_1 \cup \{\sigma_2\}$. We say that σ_2 carries the* second-slot transfer certificate *if the step $\mathcal{C}_1 \rightsquigarrow \mathcal{C}_2$ satisfies the same slotwise retained-discipline transfer conditions as in Definition H24.28, now read relative to the already certified corridor $\mathcal{C}_1$ and its new leading slot σ_2.*

Proposition H24.33 (The second newly adjoined admissible slot closes under the repeated leading-slot transfer certificate)**.** *Assume Definition H24.32. Then the extension step $\mathcal{C}_1 \rightsquigarrow \mathcal{C}_2$ closes, and $\mathcal{C}_2$ is a certified OP 1 admissible corridor.*

Proof. The corridor $\mathcal{C}_1$ is already certified by Corollary H24.26. By Definition H24.32, the next leading slot σ_2 again satisfies exactly the slotwise retained-discipline transfer assumptions required in the iterable extension schema of Proposition H24.29. Applying that proposition with $k = 2$ yields closure of the step $\mathcal{C}_1 \rightsquigarrow \mathcal{C}_2$ and certifies the enlarged corridor $\mathcal{C}_2$. □

Corollary H24.34 (First realized certified two-slot OP 1 admissible-window chain)**.** *Under the repeated leading-slot transfer certificate, the current OP 1 line already contains a realized certified chain*

$$\mathcal{C}_0 \subset \mathcal{C}_1 \subset \mathcal{C}_2,$$

of length 2*. Equivalently, beyond the retained live-kernel closure, two consecutive admissible-slot enlargements are now certified without reopening a live OP 1 burden.*

Proof. The first step $\mathcal{C}_0 \rightsquigarrow \mathcal{C}_1$ is certified by Corollary H24.26. Proposition H24.33 certifies the second step $\mathcal{C}_1 \rightsquigarrow \mathcal{C}_2$. Hence the displayed chain of length 2 is certified. □

Remark H24.35 (What the first realized two-slot chain now buys)**.** *This is the first point at which the iterable extension schema becomes visibly more than a packaging convenience. The text still does not claim a fully promoted OP 1 corridor. It claims something smaller and already operational: once the transferred retained comparison discipline persists across two successive leading slots, the corridor extends twice without reopening the local OP 1 burden. In particular, any future extension failure must now occur only at the first admissible slot* after $\mathcal{C}_2$ *where the leading-slot transfer certificate ceases to hold or begins to destabilize.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Reader cue. Read the next block as a compact certification format for a finite OP 1 extension run. Its purpose is not to claim unrestricted corridor promotion, but to package a bounded number of already justified leading-slot transfers into one reusable certificate whose failure horizon is explicitly the first slot beyond the certified run.

19AZ$^{(6p)}$. First bounded k-step certificate format for finite OP 1 admissible-window extension runs

After 19AZ$^{(6o)}$, the present line has more than one isolated extension success but still less than an unrestricted promoted corridor. The right packaging is therefore a bounded certificate format: one records a finite certified run together with the leading admissible slots and their slotwise transfer certificates, and one treats the first uncertified slot beyond that run as the first honest location where extension may fail again.

Definition H24.36 (Bounded k-step OP 1 extension certificate)**.** *Let $k \geq 1$. A* bounded k-step OP 1 extension certificate *is data*

$$\mathbf{K}_k = (\mathcal{C}_0 \subset \mathcal{C}_1 \subset \cdots \subset \mathcal{C}_k;\, \sigma_1, \ldots, \sigma_k),$$

where for each $j = 1, \ldots, k$ the slot σ_j is the newly adjoined leading admissible slot in the extension step $\mathcal{C}_{j-1} \rightsquigarrow \mathcal{C}_j$, and σ_j carries the slotwise retained-discipline transfer certificate required by Definition H24.28 relative to the already certified corridor $\mathcal{C}_{j-1}$.

Proposition H24.37 (Bounded k-step certificates certify the whole finite OP 1 run)**.** *Assume Definition H24.36. Then the terminal corridor $\mathcal{C}_k$ is certified, and every intermediate step $\mathcal{C}_{j-1} \rightsquigarrow \mathcal{C}_j$ closes without reopening a live OP 1 burden.*

Proof. This is exactly the repeated one-step mechanism of Proposition H24.29, now written in bounded packaged form. Since each newly adjoined leading slot σ_j carries the required slotwise retained-discipline transfer certificate, every step closes in turn. Hence the entire finite chain up to $\mathcal{C}_k$ is certified. □

Corollary H24.38 (First honest failure lies beyond a bounded certificate horizon)**.** *Assume Definition H24.36. Then any honest future extension failure must first appear at the earliest admissible slot strictly beyond the leading edge of $\mathcal{C}_k$ for which the next slotwise transfer certificate is absent or unstable.*

Proof. Every step through $\mathcal{C}_k$ is already certified by Proposition H24.37. Therefore no honest failure can occur inside that bounded run. The first possible extension failure must lie at the first later leading slot for which the next certificate is not available. □

Remark H24.39 (What the bounded certificate format now buys)**.** *This is the finite-horizon normalization that the post-extension line needed. Instead of speaking vaguely about an enlarging corridor, the text can now speak about a specific certified horizon $\mathbf{K}_k$. The Monte-Carlo/fusion-probe layer is thereby given a precise target: it need only search for the first later slot beyond $\mathbf{K}_k$ where the transfer certificate begins to fail, rather than resurveying the already certified run.*

Reader cue. Read the next block as the first explicit named bounded certificate already available on the present line. Its purpose is to turn the abstract finite-horizon format into a concrete certified object that can be cited directly in later routing and probe discussions.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

19AZ$^{(6q)}$. First explicit bounded certificate $\mathbf{K}_2$ on the present OP 1 extension line

After 19AZ$^{(6p)}$, the next honest step is to name the first concrete bounded certificate already realized by the current line. The smallest nontrivial such object is the two-step extension chain from 19AZ$^{(6o)}$, now repackaged as a single finite certificate. This gives the text an explicit certified horizon that later extension tests and Monte-Carlo/probe diagnostics can cite without reopening the step-by-step derivation each time.

Definition H24.40 (The present two-step bounded certificate $\mathbf{K}_2^{\mathrm{cur}}$)**.** *Let*

$$\mathbf{K}_2^{\mathrm{cur}} := (\mathcal{C}_0 \subset \mathcal{C}_1 \subset \mathcal{C}_2;\, \sigma_1, \sigma_2),$$

where $\mathcal{C}_0$ is the retained certified OP 1 corridor, σ_1 is the first newly adjoined admissible slot certified in Corollary H24.26, and σ_2 is the second newly adjoined admissible slot certified in Corollary H24.34. We call $\mathbf{K}_2^{\mathrm{cur}}$ the present two-step bounded certificate*.*

Proposition H24.41 (The current OP 1 line carries the explicit bounded certificate $\mathbf{K}_2^{\mathrm{cur}}$)**.** *The data of Definition H24.40 satisfy Definition H24.36 with $k = 2$. Hence $\mathbf{K}_2^{\mathrm{cur}}$ is a valid bounded certificate on the present OP 1 line.*

Proof. The step $\mathcal{C}_0 \rightsquigarrow \mathcal{C}_1$ is certified by Corollary H24.26. The step $\mathcal{C}_1 \rightsquigarrow \mathcal{C}_2$ is certified by Corollary H24.34. Therefore the two leading slots σ_1, σ_2 each carry the required slotwise transfer certificate and the chain satisfies Definition H24.36 with $k = 2$. □

Corollary H24.42 (The first uncertified OP 1 slot lies strictly beyond $\mathbf{K}_2^{\mathrm{cur}}$)**.** *On the present line, any honest future OP 1 admissible-window extension failure must first occur at a leading admissible slot strictly beyond the corridor $\mathcal{C}_2$ certified by $\mathbf{K}_2^{\mathrm{cur}}$.*

Proof. This is the case $k = 2$ of Corollary H24.38, applied to Proposition H24.41. □

Remark H24.43 (What the explicit certificate $\mathbf{K}_2^{\mathrm{cur}}$ now buys)**.** *The current extension line now has a named finite horizon. This is useful both mathematically and diagnostically. Mathematically, later corridor-extension arguments can cite $\mathbf{K}_2^{\mathrm{cur}}$ as a certified base segment rather than replaying the first two transferred slots. Diagnostically, the Monte-Carlo/fusion-probe layer can now ask a sharply bounded question: where is the first later slot beyond $\mathbf{K}_2^{\mathrm{cur}}$ at which the next leading-slot transfer certificate destabilizes?*

Reader cue. Read the next block as the first controlled test beyond the current certified horizon $\mathbf{K}_2^{\mathrm{cur}}$. Its purpose is not to claim a third transferred slot automatically, but to localize the next honest extension question at the very first later leading slot and to state the exact alternative: either that slot carries the next transfer certificate and yields a certified $\mathbf{K}_3$, or it is the first genuine post-$\mathbf{K}_2^{\mathrm{cur}}$ boundary witness.

19AZ$^{(6r)}$. First post-$\mathbf{K}_2^{\mathrm{cur}}$ leading-slot test beyond the current certified horizon

After 19AZ$^{(6q)}$, the next honest step is to move from a named finite horizon to the first controlled test beyond that horizon. The current line therefore does not jump directly to an unbounded promotion claim. Instead it isolates the first later leading admissible slot beyond $\mathcal{C}_2$ and asks whether the same slotwise retained-discipline transfer certificate extends one more time. This is the smallest extension question that remains once $\mathbf{K}_2^{\mathrm{cur}}$ has been certified.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition H24.44 (First post-$\mathbf{K}_2^{\mathrm{cur}}$ leading-slot test datum)**.** *Let* $\mathbf{K}_2^{\mathrm{cur}} = (\mathcal{C}_0 \subset \mathcal{C}_1 \subset \mathcal{C}_2; \sigma_1, \sigma_2)$ *be the present two-step bounded certificate from Definition H24.40. Define* σ_3 *to be the first later leading admissible slot strictly beyond the certified corridor* $\mathcal{C}_2$*, whenever such a next slot is under consideration on the same retained OP 1 extension line. We call the pair* $(\mathcal{C}_2, \sigma_3)$ *the* first post-$\mathbf{K}_2^{\mathrm{cur}}$ leading-slot test datum.

Proposition H24.45 (The first honest future OP 1 extension question beyond $\mathbf{K}_2^{\mathrm{cur}}$ is localized at σ_3)**.** *Assume the setting of Definition H24.44. Then every honest future OP 1 admissible-window extension question beyond the currently certified horizon is localized at the single leading slot* σ_3*. More precisely, exactly one of the following alternatives occurs:*

(i) the slot σ_3 *carries the same slotwise retained-discipline transfer certificate as in the previously certified steps, in which case the enlarged chain*

$$\mathcal{C}_0 \subset \mathcal{C}_1 \subset \mathcal{C}_2 \subset \mathcal{C}_3$$

is a valid bounded certificate $\mathbf{K}_3^{\mathrm{cand}}$ *on the same OP 1 line;*

(ii) the slot σ_3 *does not carry that certificate, in which case* σ_3 *itself is the first genuine post-*$\mathbf{K}_2^{\mathrm{cur}}$ *boundary-witness location.*

In particular, no honest future failure can occur strictly before σ_3*.*

Proof. By Corollary H24.42, any honest future OP 1 extension failure must first occur at a leading admissible slot strictly beyond $\mathcal{C}_2$. By definition, σ_3 is the first such later leading slot under consideration. Therefore no honest future failure can occur before σ_3. If σ_3 carries the same slotwise retained-discipline transfer certificate as in the previous certified steps, then the same slotwise-extension mechanism used in Proposition H24.37 extends the chain by one step and yields a valid bounded certificate with $k = 3$. If not, then the failure is already localized at σ_3 itself. These are the only two possibilities. □

Corollary H24.46 (The post-$\mathbf{K}_2^{\mathrm{cur}}$ horizon is controlled by a single next-slot test)**.** *On the present OP 1 line, the first uncertified future horizon beyond* $\mathbf{K}_2^{\mathrm{cur}}$ *is completely controlled by the single next-slot test at* σ_3*: either* $\mathbf{K}_3^{\mathrm{cand}}$ *is certified, or the first honest future boundary witness is already localized at* σ_3*.*

Proof. Immediate from Proposition H24.45. □

Remark H24.47 (What the first post-$\mathbf{K}_2^{\mathrm{cur}}$ test now buys)**.** *This is the first point at which the extension line leaves the already named certified horizon. The benefit is that the next search question is now maximally sharp. Neither the theorematic line nor the Monte-Carlo/fusion-probe layer needs to resurvey the already certified segment* $\mathbf{K}_2^{\mathrm{cur}}$*; both need only test whether the next leading slot* σ_3 *carries the same transfer certificate. If it does, a concrete* $\mathbf{K}_3$ *becomes available. If it does not, then the first genuine post-*$\mathbf{K}_2^{\mathrm{cur}}$ *boundary witness has already been found.*

Reader cue. Read the next block as the two-channel split of the first uncertified slot beyond $\mathbf{K}_2^{\mathrm{cur}}$. Its purpose is not yet to certify $\mathbf{K}_3$, but to normalize the next decision at σ_3 into the two only honest outcomes: either the same transfer certificate still holds, or the first genuine post-$\mathbf{K}_2^{\mathrm{cur}}$ boundary witness is already present there.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

19AZ$^{(6s)}$. First transfer-positive versus boundary-active split at the post-$\mathbf{K}_2^{\mathrm{cur}}$ slot σ_3

After 19AZ$^{(6r)}$, the first uncertified slot beyond the current certified horizon is no longer diffuse. It is the single leading admissible slot σ_3 beyond $\mathcal{C}_2$. The next honest normalization is therefore to split the status of σ_3 itself. Either σ_3 still carries the same slotwise retained-discipline transfer certificate that already certified σ_1 and σ_2, or else σ_3 is already the first active boundary location beyond $\mathbf{K}_2^{\mathrm{cur}}$.

Definition H24.48 (Transfer-positive versus boundary-active status at σ_3)**.** *Assume the setting of Definition H24.44. We say that the first later leading admissible slot σ_3 is*

(i) transfer-positive *if it carries the same slotwise retained-discipline transfer certificate as in the certified steps leading to $\mathbf{K}_2^{\mathrm{cur}}$, and*

(ii) boundary-active *if it does not carry that certificate and therefore already hosts the first genuine post-$\mathbf{K}_2^{\mathrm{cur}}$ boundary witness.*

Proposition H24.49 (The post-$\mathbf{K}_2^{\mathrm{cur}}$ slot σ_3 admits only the transfer-positive or boundary-active status)**.** *Assume the setting of Definition H24.48. Then exactly one of the following alternatives holds:*

(i) σ_3 is transfer-positive, in which case the current line produces a valid three-step bounded certificate

$$\mathbf{K}_3^{\mathrm{cand}} = (\mathcal{C}_0 \subset \mathcal{C}_1 \subset \mathcal{C}_2 \subset \mathcal{C}_3;\, \sigma_1, \sigma_2, \sigma_3),$$

with $\mathcal{C}_3$ the corridor obtained by adjoining σ_3 to $\mathcal{C}_2$; or

(ii) σ_3 is boundary-active, in which case σ_3 itself is already the first genuine post-$\mathbf{K}_2^{\mathrm{cur}}$ boundary location.

No third autonomous status for σ_3 is licensed on the present OP 1 extension line.

Proof. By Proposition H24.45, every honest future OP 1 extension question beyond $\mathbf{K}_2^{\mathrm{cur}}$ is localized at the single next leading admissible slot σ_3. Therefore the only remaining issue is whether σ_3 carries the same slotwise retained-discipline transfer certificate as the previously certified steps. If it does, then the bounded-k certificate mechanism of Proposition H24.37 extends the chain once more and yields the displayed three-step final certificate. If it does not, then by the same localization result σ_3 itself is already the first genuine post-$\mathbf{K}_2^{\mathrm{cur}}$ boundary location. Since the extension problem is completely localized at σ_3, these two alternatives exhaust the possibilities. □

Corollary H24.50 (The next honest OP 1 decision beyond $\mathbf{K}_2^{\mathrm{cur}}$ is a two-channel decision at σ_3)**.** *On the present OP 1 extension line, the first uncertified future decision beyond $\mathbf{K}_2^{\mathrm{cur}}$ reduces to the two-channel split*

$$\textit{transfer-positive at } \sigma_3 \qquad \textit{or} \qquad \textit{boundary-active at } \sigma_3.$$

In particular, the theorematic line and the Monte-Carlo/fusion-probe layer need not resurvey the already certified segment $\mathbf{K}_2^{\mathrm{cur}}$ in order to decide the next step.

Proof. Immediate from Proposition H24.49. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Remark H24.51 (What the first transfer-positive/boundary-active split at σ_3 now buys)**.** *This is the first point beyond the currently certified horizon where the next admissible slot is not merely isolated but also status-normalized. The benefit is that the next step no longer asks a vague continuation question. It asks only whether σ_3 is transfer-positive or already boundary-active. If it is transfer-positive, the line immediately upgrades to a concrete three-step certificate. If it is boundary-active, then the first genuine post-$\mathbf{K}_2^{\mathrm{cur}}$ boundary witness has already been found at the earliest possible place.*

Reader cue. Read the next block as the direct certification of the transfer-positive channel at σ_3. Its purpose is not yet to promote the extension line indefinitely, but to convert the already normalized positive branch into an explicit current three-step bounded certificate and to discharge the boundary-active channel at that slot itself.

19AZ$^{(6t)}$. First certification of $\mathbf{K}_3^{\mathrm{cur}}$ under transfer-positive status at σ_3

After 19AZ$^{(6s)}$, the next honest step is no longer to restate the two-channel split at σ_3, but to close its positive side. If the first later leading admissible slot beyond $\mathbf{K}_2^{\mathrm{cur}}$ is transfer-positive, then the same slotwise retained-discipline transfer mechanism that already produced the certified steps through σ_2 should certify one more step. The present block records that upgrade explicitly and removes the boundary-active alternative at σ_3 itself whenever the positive certificate is in force.

Definition H24.52 (Present three-step bounded certificate $\mathbf{K}_3^{\mathrm{cur}}$)**.** *Assume the setting of Definition H24.48. If σ_3 is transfer-positive, define the* present three-step bounded certificate

$$\mathbf{K}_3^{\mathrm{cur}} = (\mathcal{C}_0 \subset \mathcal{C}_1 \subset \mathcal{C}_2 \subset \mathcal{C}_3;\, \sigma_1, \sigma_2, \sigma_3),$$

where $\mathcal{C}_3$ is obtained from $\mathcal{C}_2$ by adjoining the transfer-positive leading admissible slot σ_3 under the same slotwise retained-discipline transfer certificate used in the already certified steps.

Proposition H24.53 (Transfer-positive status at σ_3 certifies $\mathbf{K}_3^{\mathrm{cur}}$ and discharges the local boundary channel)**.** *Assume the setting of Definition H24.52. Then:*

(i) the chain $\mathbf{K}_3^{\mathrm{cur}}$ is a valid bounded three-step OP 1 admissible-window certificate on the present extension line;

(ii) σ_3 is not a genuine post-$\mathbf{K}_2^{\mathrm{cur}}$ boundary-witness location; and

(iii) every honest future OP 1 extension failure, if any, must occur at a later leading admissible slot strictly beyond the corridor $\mathcal{C}_3$ certified by $\mathbf{K}_3^{\mathrm{cur}}$.

Proof. By Definition H24.52, the slot σ_3 carries the same slotwise retained-discipline transfer certificate as the already certified steps through σ_2. Proposition H24.37 therefore upgrades the chain by one step and certifies the displayed bounded three-step corridor, giving (i). Once that positive certification is in force, the alternative "boundary-active at σ_3" from Proposition H24.49 is excluded, proving (ii). Finally, after $\mathcal{C}_3$ has become part of the certified horizon, any later honest OP 1 failure must first appear strictly beyond $\mathcal{C}_3$, which is exactly (iii). □

Corollary H24.54 (The positive channel at σ_3 upgrades the certified horizon from $\mathbf{K}_2^{\mathrm{cur}}$ to $\mathbf{K}_3^{\mathrm{cur}}$)**.** *On the present OP 1 extension line, transfer-positive status at σ_3 does not merely produce a final continuation. It certifies the concrete bounded horizon $\mathbf{K}_3^{\mathrm{cur}}$ and pushes every honest future extension question to the first later leading admissible slot beyond $\mathcal{C}_3$.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. Immediate from Proposition H24.53. □

Remark H24.55 (What certifying $\mathbf{K}_3^{\mathrm{cur}}$ now buys)**.** *This is the first point at which the extension line no longer talks about a hypothetical third step but about a named current three-step horizon. The benefit is again twofold. The theorematic line now has a larger certified corridor and therefore a smaller honest future search zone. The Monte-Carlo/fusion-probe layer likewise no longer needs to test anything at σ_3 itself; its next bounded question starts only at the first later leading admissible slot beyond $\mathcal{C}_3$.*

Reader cue. Read the next block as the symmetric normalization of the negative channel at σ_3. Its purpose is not to reopen the already certified positive branch, but to record what the extension line is honestly allowed to say when σ_3 is boundary-active: namely, that the first explicit post-$\mathbf{K}_2^{\mathrm{cur}}$ extension-boundary witness has already been found there and that no later slot needs to be consulted before that witness is discharged.

19AZ$^{(6u)}$. First explicit normalization of the boundary-active channel at σ_3 as the earliest post-$\mathbf{K}_2^{\mathrm{cur}}$ extension witness

After 19AZ$^{(6t)}$, the positive side at σ_3 has been normalized: if σ_3 is transfer-positive, then the line upgrades to the concrete bounded certificate $\mathbf{K}_3^{\mathrm{cur}}$. The negative side should now be stated with the same clarity. If σ_3 is boundary-active, then it is not merely a failed final slot. It is already the first explicit post-$\mathbf{K}_2^{\mathrm{cur}}$ extension witness, and every honest later extension question remains downstream of that earliest active boundary location until the witness at σ_3 has been discharged.

Definition H24.56 (Current boundary-active witness datum at σ_3)**.** *Assume the setting of Definition H24.48. If σ_3 is boundary-active, define the* current boundary-active witness datum

$$\mathfrak{B}_3^{\mathrm{cur}} = (\mathcal{C}_2, \sigma_3),$$

viewed as the earliest explicit post-$\mathbf{K}_2^{\mathrm{cur}}$ OP 1 extension-boundary witness on the present line.

Proposition H24.57 (Boundary-active status at σ_3 normalizes the earliest explicit post-$\mathbf{K}_2^{\mathrm{cur}}$ extension witness)**.** *Assume the setting of Definition H24.56. Then:*

(i) σ_3 is the first genuine post-$\mathbf{K}_2^{\mathrm{cur}}$ OP 1 extension-boundary location on the present line;

(ii) no valid bounded three-step certificate $\mathbf{K}_3^{\mathrm{cur}}$ is licensed on the same line until the witness datum $\mathfrak{B}_3^{\mathrm{cur}}$ has been discharged; and

(iii) every honest later OP 1 extension question remains downstream of σ_3 and need not be opened before the witness datum at σ_3 has been resolved.

Proof. If σ_3 is boundary-active, then by Proposition H24.49 the positive alternative fails exactly at the first later leading admissible slot beyond $\mathbf{K}_2^{\mathrm{cur}}$. This gives (i). Since Proposition H24.53 requires transfer-positive status in order to certify $\mathbf{K}_3^{\mathrm{cur}}$, the same line cannot simultaneously carry the certified three-step horizon while σ_3 remains boundary-active, proving (ii). Finally, because σ_3 is already the earliest active boundary location beyond the current certified horizon, no honest later extension question needs to be opened before that earliest witness has been addressed, which is (iii). □

Corollary H24.58 (The negative channel at σ_3 is the earliest explicit witness beyond $\mathbf{K}_2^{\mathrm{cur}}$)**.** *On the present OP 1 extension line, boundary-active status at σ_3 does not create a diffuse failure region. It produces the explicit earliest witness datum $\mathfrak{B}_3^{\mathrm{cur}} = (\mathcal{C}_2, \sigma_3)$ and blocks further honest extension promotion on that line until this witness has been discharged.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. Immediate from Proposition H24.57. □

Remark H24.59 (What explicitly normalizing the boundary-active channel at σ_3 now buys)**.** *This is the symmetric counterpart to certifying $\mathbf{K}_3^{\mathrm{cur}}$ on the positive side. The benefit is that the negative side is no longer left as an unnamed failure possibility. It is packaged as the earliest explicit post-$\mathbf{K}_2^{\mathrm{cur}}$ witness datum, so that the theorematic line, the routing layer, and the Monte-Carlo/fusion-probe layer all know where the next honest obstruction sits if the transfer-positive channel does not carry. In particular, no later slot needs to be searched before σ_3 itself has been resolved.*

Reader cue. Read the next block as the compact decision-normal form for the first slot beyond the current certified horizon. Its purpose is not to add a new extension theorem, but to compress the already normalized alternatives at σ_3 into one finite decision object that the theorematic line, the routing layer, and the probe layer can all use without reopening the full case split.

19AZ$^{(6v)}$. First decision-normal form at σ_3 beyond the current certified horizon

After 19AZ$^{(6t)}$ and 19AZ$^{(6u)}$, the first slot beyond the current certified horizon has been normalized on both sides. On the positive side, transfer-positive status at σ_3 certifies the concrete bounded three-step horizon $\mathbf{K}_3^{\mathrm{cur}}$. On the negative side, boundary-active status at σ_3 produces the earliest explicit post-$\mathbf{K}_2^{\mathrm{cur}}$ witness datum $\mathfrak{B}_3^{\mathrm{cur}} = (\mathcal{C}_2, \sigma_3)$. The next honest step is therefore to package these two outcomes into one compact decision-normal form for the first later leading slot beyond the presently certified horizon.

Definition H24.60 (Current horizon decision datum at σ_3)**.** *Assume the setting of Definition H24.48. Define the* current horizon decision datum

$$\mathfrak{D}_3^{\mathrm{cur}} \in \{\mathbf{K}_3^{\mathrm{cur}}, \mathfrak{B}_3^{\mathrm{cur}}\}$$

by the rule

$$\mathfrak{D}_3^{\mathrm{cur}} = \begin{cases} \mathbf{K}_3^{\mathrm{cur}}, & \text{if } \sigma_3 \text{ is transfer-positive,} \\ \mathfrak{B}_3^{\mathrm{cur}}, & \text{if } \sigma_3 \text{ is boundary-active.} \end{cases}$$

Proposition H24.61 (The first slot beyond $\mathbf{K}_2^{\mathrm{cur}}$ admits a finite decision-normal form)**.** *Assume the setting of Definition H24.60. Then:*

(i) the datum $\mathfrak{D}_3^{\mathrm{cur}}$ is well-defined and exhaustive for the first later leading admissible slot beyond $\mathbf{K}_2^{\mathrm{cur}}$;

(ii) if $\mathfrak{D}_3^{\mathrm{cur}} = \mathbf{K}_3^{\mathrm{cur}}$, then the certified horizon has advanced through $\mathcal{C}_3$ and every honest later extension question begins strictly beyond that corridor; and

(iii) if $\mathfrak{D}_3^{\mathrm{cur}} = \mathfrak{B}_3^{\mathrm{cur}}$, then σ_3 is already the earliest explicit post-$\mathbf{K}_2^{\mathrm{cur}}$ boundary location and no later slot needs to be opened before that witness datum has been discharged.

Proof. By Proposition H24.49, the slot σ_3 admits only the two statuses transfer-positive and boundary-active. Definition H24.60 therefore assigns exactly one of the two normalized outputs $\mathbf{K}_3^{\mathrm{cur}}$ and $\mathfrak{B}_3^{\mathrm{cur}}$, proving (i). If the positive output occurs, Proposition H24.53 gives the certified three-step horizon and pushes every honest later extension question beyond $\mathcal{C}_3$, which is (ii). If the negative output occurs, Proposition H24.57 gives the earliest explicit witness datum at σ_3 and forbids opening later slots first, which is (iii). □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Corollary H24.62 (The first slot beyond the current certified horizon is now decision-normalized)**.** *On the present OP 1 extension line, the first later leading admissible slot beyond* $\mathbf{K}_2^{\mathrm{cur}}$ *no longer carries an uncompressed continuation question. It is represented by the finite decision datum* $\mathfrak{D}_3^{\mathrm{cur}} \in \{\mathbf{K}_3^{\mathrm{cur}}, \mathfrak{B}_3^{\mathrm{cur}}\}$*, which either upgrades the current certified horizon or identifies the earliest explicit boundary witness.*

Proof. Immediate from Proposition H24.61. □

Remark H24.63 (What the decision-normal form at σ_3 now buys)**.** *This is the first point beyond the currently certified horizon where the extension line can speak in one compact finite decision object instead of in a longer two-branch case split. The theorematic line gains a cleaner handoff to the next extension stage; the routing layer gains a single certified output to follow; and the Monte-Carlo/fusion-probe layer gains a sharply bounded question: either the line has already promoted to* $\mathbf{K}_3^{\mathrm{cur}}$*, or the earliest active obstruction has already been localized at* σ_3 *itself.*

Reader cue. Read the next block as the compact QAM/decoder output language for the first extension decision beyond the current certified horizon. Its purpose is not to add a stronger extension theorem, but to package the already normalized alternatives $\mathbf{K}_3^{\mathrm{cur}}$ versus $\mathfrak{B}_3^{\mathrm{cur}}$ into one short token language that later theorematic, routing, and probe layers can cite without reopening the full decision datum.

19AZ$^{(6w)}$. First QAM/decoder output tokens for extension tasks beyond the current certified horizon

After 19AZ$^{(6v)}$, the first slot beyond the current certified horizon is represented by the finite decision datum

$$\mathfrak{D}_3^{\mathrm{cur}} \in \{\mathbf{K}_3^{\mathrm{cur}}, \mathfrak{B}_3^{\mathrm{cur}}\}.$$

The next honest compression step is therefore to translate this extension decision into a short output language that can be consumed uniformly by the QAM layer, the route selector, and the probe/decoder layer without reopening the full theorematic split.

Definition H24.64 (Extension-output token language at σ_3)**.** *Define the finite token set*

$$\mathcal{T}_3^{\mathrm{ext}} := \{\mathsf{EK}_3, \mathsf{EB}_3\},$$

where the symbols mean:

$$\mathsf{EK}_3 := \text{“the current horizon extends through } \mathcal{C}_3\text{”},$$
$$\mathsf{EB}_3 := \text{“the earliest honest post-}\mathbf{K}_2^{\mathrm{cur}} \text{ boundary sits at } \sigma_3\text{”}.$$

Define the associated token map by

$$\mathsf{Out}_{\mathrm{ext},3}^{\mathrm{tok}}(\mathfrak{D}_3^{\mathrm{cur}}) = \begin{cases} \mathsf{EK}_3, & \text{if } \mathfrak{D}_3^{\mathrm{cur}} = \mathbf{K}_3^{\mathrm{cur}}, \\ \mathsf{EB}_3, & \text{if } \mathfrak{D}_3^{\mathrm{cur}} = \mathfrak{B}_3^{\mathrm{cur}}. \end{cases}$$

Proposition H24.65 (The first extension decision beyond $\mathbf{K}_2^{\mathrm{cur}}$ admits a faithful token output)**.** *Assume the setting of Definition H24.64. Then:*

(i) the map $\mathsf{Out}_{\mathrm{ext},3}^{\mathrm{tok}}$ *is well-defined on the current decision datum* $\mathfrak{D}_3^{\mathrm{cur}}$*;*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(ii) $\mathsf{Out}^{\mathrm{tok}}_{\mathrm{ext},3}(\mathfrak{D}^{\mathrm{cur}}_3) = \mathsf{EK}_3$ *if and only if the line has already promoted to the certified bounded horizon* $\mathbf{K}^{\mathrm{cur}}_3$*; and*

(iii) $\mathsf{Out}^{\mathrm{tok}}_{\mathrm{ext},3}(\mathfrak{D}^{\mathrm{cur}}_3) = \mathsf{EB}_3$ *if and only if* σ_3 *is already the earliest explicit boundary witness beyond* $\mathbf{K}^{\mathrm{cur}}_2$*.*

In particular, the token output is faithful to the full current extension decision.

Proof. By Definition H24.60, the decision datum $\mathfrak{D}^{\mathrm{cur}}_3$ takes exactly one of the two values $\mathbf{K}^{\mathrm{cur}}_3$ and $\mathfrak{B}^{\mathrm{cur}}_3$, so the token map is well-defined, proving (i). If the first value occurs, then Proposition H24.53 shows that the certified horizon has advanced through $\mathcal{C}_3$, hence the output token is exactly EK_3, proving (ii). If the second value occurs, then Proposition H24.57 shows that σ_3 is the earliest explicit post-$\mathbf{K}^{\mathrm{cur}}_2$ boundary location, hence the output token is exactly EB_3, proving (iii). Since the two tokens are distinct and attached to the two distinct normalized outcomes, the output is faithful to the full decision datum. □

Corollary H24.66 (Extension tasks at the first post-$\mathbf{K}^{\mathrm{cur}}_2$ slot are now token-normalized)**.** *On the present OP 1 extension line, the first extension task beyond the certified horizon* $\mathbf{K}^{\mathrm{cur}}_2$ *can now be recorded by the compact token*

$$\mathsf{Out}^{\mathrm{tok}}_{\mathrm{ext},3}(\mathfrak{D}^{\mathrm{cur}}_3) \in \{\mathsf{EK}_3, \mathsf{EB}_3\}.$$

Thus later routing or probe logic no longer needs the full branch split at σ_3*; it is enough to ask whether the current token is* EK_3 *or* EB_3*.*

Proof. Immediate from Proposition H24.65. □

Remark H24.67 (What the first extension-output tokens now buy)**.** *The gain is small but real. The present line no longer needs to treat the first post-*$\mathbf{K}^{\mathrm{cur}}_2$ *extension task as a longer prose decision between "certify the next horizon" and "localize the earliest new witness". The theorematic layer, the routing layer, and the Monte-Carlo/probe layer may all read the same compact output token* EK_3 *or* EB_3*. In particular, this is the first point where the extension line itself begins to speak the same kind of compact QAM/decoder language that had previously been reserved for reopening tasks inside the retained OP kernels.*

Reader cue. Read the next block as the reusable proof-compression template abstracted from the first post-$\mathbf{K}^{\mathrm{cur}}_2$ decision. Its purpose is not to claim a new global extension theorem, but to isolate the exact finite-decision pattern that any later first-slot extension stage must satisfy once its local positive-versus-boundary dichotomy has been established.

19AZ$^{(6x)}$. Reusable next-slot decision schema beyond a certified horizon

The blocks 19AZ$^{(6v)}$ and 19AZ$^{(6w)}$ show that, once the first later leading admissible slot beyond a certified current horizon has been normalized into the two statuses *transfer-positive* versus *boundary-active*, the continuation question collapses to one finite decision datum and one faithful output token. The point of the present block is to isolate that pattern as a reusable local schema.

Definition H24.68 (Reusable next-slot decision schema)**.** *Fix a current certified horizon* $\mathbf{K}^{\mathrm{cur}}_m$*, and let* σ_{m+1} *denote the first later leading admissible slot beyond it. Assume that at this slot the local theorematic line has already established an exhaustive dichotomy:*

$$\sigma_{m+1} \text{ is either transfer-positive or boundary-active.}$$

Assume further that:

© 2026 Marc Brendecke. Preprint — not peer reviewed.

- *transfer-positive status at σ_{m+1} certifies a concrete upgraded horizon $\mathbf{K}^{\mathrm{cur}}_{m+1}$, and*
- *boundary-active status at σ_{m+1} produces an earliest explicit witness datum $\mathfrak{B}^{\mathrm{cur}}_{m+1}$.*

Then define the associated next-slot decision datum

$$\mathfrak{D}^{\mathrm{cur}}_{m+1} \in \{\mathbf{K}^{\mathrm{cur}}_{m+1}, \mathfrak{B}^{\mathrm{cur}}_{m+1}\}$$

by

$$\mathfrak{D}^{\mathrm{cur}}_{m+1} = \begin{cases} \mathbf{K}^{\mathrm{cur}}_{m+1}, & \text{if } \sigma_{m+1} \text{ is transfer-positive,} \\ \mathfrak{B}^{\mathrm{cur}}_{m+1}, & \text{if } \sigma_{m+1} \text{ is boundary-active.} \end{cases}$$

Define moreover the associated finite output-token set

$$\mathcal{T}^{\mathrm{ext}}_{m+1} := \{\mathsf{EK}_{m+1}, \mathsf{EB}_{m+1}\},$$

where

$$\mathsf{EK}_{m+1} := \text{“the current horizon extends through } \mathcal{C}_{m+1}\text{”},$$
$$\mathsf{EB}_{m+1} := \text{“the earliest honest boundary beyond } \mathbf{K}^{\mathrm{cur}}_{m} \text{ sits at } \sigma_{m+1}\text{”}.$$

Finally define the token output map

$$\mathsf{Out}^{\mathrm{tok}}_{\mathrm{ext},m+1}(\mathfrak{D}^{\mathrm{cur}}_{m+1}) = \begin{cases} \mathsf{EK}_{m+1}, & \text{if } \mathfrak{D}^{\mathrm{cur}}_{m+1} = \mathbf{K}^{\mathrm{cur}}_{m+1}, \\ \mathsf{EB}_{m+1}, & \text{if } \mathfrak{D}^{\mathrm{cur}}_{m+1} = \mathfrak{B}^{\mathrm{cur}}_{m+1}. \end{cases}$$

Proposition H24.69 (Any normalized first-slot extension stage collapses to one finite decision datum and one faithful token)**.** *Assume the setting of Definition H24.68. Then:*

(i) the datum $\mathfrak{D}^{\mathrm{cur}}_{m+1}$ is well-defined and exhaustive for the first later leading admissible slot beyond $\mathbf{K}^{\mathrm{cur}}_{m}$;

(ii) the token map $\mathsf{Out}^{\mathrm{tok}}_{\mathrm{ext},m+1}$ is well-defined and faithful to that decision datum; and

(iii) every later routing, decoder, or probe layer may consume the compact token EK_{m+1} or EB_{m+1} without reopening the full local branch split at σ_{m+1}.

Proof. By assumption, the local line at σ_{m+1} admits exactly the two statuses transfer-positive and boundary-active, and each status produces exactly one normalized output, namely $\mathbf{K}^{\mathrm{cur}}_{m+1}$ or $\mathfrak{B}^{\mathrm{cur}}_{m+1}$. This proves (i). The token map is then defined on exactly these two outputs and assigns distinct tokens to them, so it is well-defined and faithful, proving (ii). Since the token remembers which of the two normalized outcomes occurred, any later routing, decoder, or probe layer may consume the token alone unless it explicitly chooses to reopen the local theorematic proof, proving (iii). □

Corollary H24.70 (The σ_3 decision package is the first realized instance of the reusable extension schema)**.** *The package*

$$\mathfrak{D}^{\mathrm{cur}}_{3} \in \{\mathbf{K}^{\mathrm{cur}}_{3}, \mathfrak{B}^{\mathrm{cur}}_{3}\}, \qquad \mathsf{Out}^{\mathrm{tok}}_{\mathrm{ext},3} \in \{\mathsf{EK}_3, \mathsf{EB}_3\},$$

constructed in 19AZ$^{(6v)}$–19AZ$^{(6w)}$, is not merely a local convenience. It is the first realized instance of the general next-slot decision schema for extension tasks beyond a certified current horizon.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. Immediate from Definition H24.68 and Proposition H24.69 by taking $m = 2$. □

Remark H24.71 (What this schema now buys)**.** *This is the point where the present OP 1 extension line begins to export a genuine reusable tool rather than only a local closure fact. Future first-slot extension stages no longer need to reinvent the packaging logic: once the local positive-versus-boundary dichotomy is proved, the decision datum and faithful output token come for free. In particular, the current σ_3 block is now readable as the prototype of a general extension-normalization interface.*

Reader cue. Read the next block as the reusable action map generated by the extension-token language. Its purpose is not to add a stronger horizon theorem, but to force every later layer to take one of only two honest next actions: either advance the certified horizon, or reopen work exactly at the earliest normalized boundary site.

19AZ$^{(6y)}$. Reusable action/transition map driven by next-slot extension tokens

After 19AZ$^{(6x)}$, the line has not only a reusable next-slot decision datum but also a reusable token language. The next honest compression step is therefore to attach to that token language the corresponding next action. The guiding principle is simple: a token should not merely summarize the local result, but should also determine which later theorematic, routing, decoder, or probe move is licensed and which moves are forbidden.

Definition H24.72 (Reusable extension-action map)**.** *Assume the setting of Definition H24.68. Define the finite extension-action set*

$$\mathcal{A}^{\mathrm{ext}}_{m+1} := \{\mathsf{ADV}_{m+1}, \mathsf{OPENB}_{m+1}\},$$

where the symbols mean:

$$\begin{aligned} \mathsf{ADV}_{m+1} &:= \textit{advance current certified horizon through } \mathcal{C}_{m+1}, \\ \mathsf{OPENB}_{m+1} &:= \textit{open earliest normalized boundary kernel at } \sigma_{m+1}. \end{aligned}$$

Define the associated action map on extension tokens by

$$\mathsf{Act}^{\mathrm{tok}}_{\mathrm{ext},m+1}(\tau) = \begin{cases} \mathsf{ADV}_{m+1}, & \textit{if } \tau = \mathsf{EK}_{m+1}, \\ \mathsf{OPENB}_{m+1}, & \textit{if } \tau = \mathsf{EB}_{m+1}. \end{cases}$$

Equivalently, one may read the composite transition map

$$\mathsf{Act}^{\mathrm{tok}}_{\mathrm{ext},m+1} \circ \mathsf{Out}^{\mathrm{tok}}_{\mathrm{ext},m+1}$$

as the minimal action normal form attached to the first later leading admissible slot beyond $\mathbf{K}^{\mathrm{cur}}_m$.

Proposition H24.73 (The extension-token language determines the next honest action)**.** *Assume the setting of Definition H24.72. Then:*

(i) the map $\mathsf{Act}^{\mathrm{tok}}_{\mathrm{ext},m+1}$ *is well-defined on* $\mathcal{T}^{\mathrm{ext}}_{m+1}$*;*

(ii) if the output action is ADV_{m+1}*, then the current line is licensed to replace* $\mathbf{K}^{\mathrm{cur}}_m$ *by* $\mathbf{K}^{\mathrm{cur}}_{m+1}$ *and to relocate the first unresolved extension question strictly beyond* $\mathcal{C}_{m+1}$*;*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(iii) if the output action is OPENB_{m+1}*, then every honest later theorematic, routing, decoder, or probe step must remain localized at the earliest normalized boundary site* σ_{m+1} *until that local boundary kernel has been processed.*

Proof. The token language from Definition H24.68 has exactly the two values EK_{m+1} and EB_{m+1}, so the action map is well-defined, proving (i). If the first token occurs, then by construction it records that the current horizon has already advanced through $\mathcal{C}_{m+1}$, so the only honest next action is to promote the certified horizon and move the unresolved extension question strictly beyond that corridor, proving (ii). If the second token occurs, then by construction it records that the earliest honest obstruction has already been normalized at σ_{m+1}; therefore any later layer that skips ahead before processing that site would violate the earliest-boundary discipline, proving (iii). □

Corollary H24.74 (The σ_3 token package now induces a certified next-action split)**.** *For the realized package from 19AZ*$^{(6v)}$*–19AZ*$^{(6x)}$*, the composite map*

$$\mathsf{Act}^{\mathrm{tok}}_{\mathrm{ext},3} \circ \mathsf{Out}^{\mathrm{tok}}_{\mathrm{ext},3}$$

returns exactly one of the two certified next actions:

$$\mathsf{ADV}_3 \qquad \textit{or} \qquad \mathsf{OPENB}_3.$$

Thus the present line is now action-normalized at the first slot beyond $\mathbf{K}_2^{\mathrm{cur}}$*: either it advances through* $\mathcal{C}_3$*, or it reopens work exactly at the earliest explicit boundary site* σ_3*.*

Proof. Immediate from Proposition H24.73 by taking $m = 2$. □

Remark H24.75 (What this action map now buys, and what can be salvaged into later layers)**.** *This is the point where the extension-tool layer stops being a passive encoding device and becomes a genuine routing interface. Later consumers no longer need to ask the full local question again; they need only read whether the current action is* ADV_{m+1} *or* OPENB_{m+1}*. That is precisely the salvageable discipline for the later optimization ideas: Monte-Carlo/fusion probes, decoder packages, and any later Heisenberg-matrix, Millennium-matrix, crystal-equation, or field-equation diagnostics should attach below this action interface as consumers of a certified local action, not as mechanisms that reopen or overwrite the theorematic decision itself.*

Reader cue. Read the next block as the iterative upgrade of the local action map. Its purpose is not to claim an already completed global induction, but to isolate the exact protocol by which the active line may repeatedly consume one certified next-slot token at a time, advancing when the token says "extend" and stopping honestly when the token says "boundary".

19AZ$^{(6z)}$. Iterated extension protocol generated by certified local action tokens

After 19AZ$^{(6y)}$, the line has a local action interface for the first later leading admissible slot beyond a certified current horizon. The next honest compression step is to package that interface into an iterative protocol. The point is simple: once one local stage has been normalized into a token and then into an action, the same logic can be reused at the next stage provided the new local dichotomy is again theorematically certified. Thus the extension line should be readable not as a one-off move at σ_3, but as a repeatable one-slot protocol with a built-in stop rule.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition H24.76 (Iterated extension protocol with earliest-boundary stop rule)**.** *Fix a certified current horizon* $\mathbf{K}_m^{\mathrm{cur}}$*. Suppose that for each stage* $r \geq m+1$*, whenever the line reaches a certified horizon* $\mathbf{K}_{r-1}^{\mathrm{cur}}$*, the first later leading admissible slot* σ_r *admits a certified local dichotomy of the kind required in Definition H24.68. Then define the associated stage-r token*

$$\tau_r \in \mathcal{T}_r^{\mathrm{ext}} = \{\mathsf{EK}_r, \mathsf{EB}_r\}$$

and stage-r action

$$a_r := \mathsf{Act}_{\mathrm{ext},r}^{\mathrm{tok}}(\tau_r) \in \mathcal{A}_r^{\mathrm{ext}} = \{\mathsf{ADV}_r, \mathsf{OPENB}_r\}.$$

The resulting iterated extension protocol *is the finite or infinite stage-by-stage procedure that obeys the following rule at each step:*

(i) if $a_r = \mathsf{ADV}_r$*, replace the current certified horizon* $\mathbf{K}_{r-1}^{\mathrm{cur}}$ *by* $\mathbf{K}_r^{\mathrm{cur}}$ *and continue to the next stage* $r+1$*;*

(ii) if $a_r = \mathsf{OPENB}_r$*, stop the forward extension protocol and localize all further honest work at the earliest normalized boundary kernel sitting at* σ_r*.*

The first index at which the protocol stops, if such an index exists, is called the earliest protocol boundary stage *and is denoted by*

$$r_* := \min\{r \geq m+1 : a_r = \mathsf{OPENB}_r\}.$$

When no such stage exists in the range under consideration, the protocol is said to be advance-persistent *on that range.*

Proposition H24.77 (The iterated extension protocol is well-defined and locally non-skipping)**.** *Assume the setting of Definition H24.76. Then:*

(i) at each certified stage r*, exactly one action* $a_r \in \{\mathsf{ADV}_r, \mathsf{OPENB}_r\}$ *is produced;*

(ii) if the protocol is stopped at stage r_**, then* σ_{r_*} *is the earliest stage in the protocol at which the line is licensed to open a normalized boundary kernel rather than advance the certified horizon;*

(iii) no honest theorematic, routing, decoder, probe, or diagnostic layer may skip past a stage with action OPENB_r *and continue the forward extension protocol as if an* ADV_r *action had occurred.*

Proof. For each certified stage r, Definition H24.68 yields exactly one token in $\{\mathsf{EK}_r, \mathsf{EB}_r\}$, and Definition H24.72 sends that token to exactly one action in $\{\mathsf{ADV}_r, \mathsf{OPENB}_r\}$, proving (i). If the protocol stops at stage r_*, then by definition r_* is the least stage whose action is OPENB_{r_*}, so all earlier stages were genuine advance stages and σ_{r_*} is the earliest protocol boundary stage, proving (ii). Statement (iii) is then forced by the meaning of OPENB_r: it records that the earliest honest unresolved obstruction has already been normalized at stage r, so any layer that skipped past it would violate the non-skipping boundary discipline built into $19\mathrm{AZ}^{(6y)}$. □

Corollary H24.78 (The present σ_3 package is the first step of an iterated extension protocol)**.** *In the current active line, the package from* $19AZ^{(6v)}$*–*$19AZ^{(6y)}$ *is not only a local closure package at the first slot beyond* $\mathbf{K}_2^{\mathrm{cur}}$*. It is the first realized stage of the iterated extension protocol. Concretely, the line now carries a certified first-stage token*

$$\tau_3 \in \{\mathsf{EK}_3, \mathsf{EB}_3\}$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

and the associated first-stage action

$$a_3 \in \{\mathsf{ADV}_3, \mathsf{OPENB}_3\},$$

so every later extension attempt must either continue from the promoted horizon $\mathbf{K}_3^{\mathrm{cur}}$ *or remain localized at the earliest normalized boundary kernel at* σ_3*.*

Proof. Immediate from Corollary H24.74 and Proposition H24.77. □

Remark H24.79 (What can be salvaged from later optimization ideas without corrupting the proof line)**.** *This protocol is also the right place to salvage the useful part of the later optimization discussion. Once the active line emits a certified action stream* (a_r)*, later consumers may compress, hash, decode, profile, or statistically probe that stream — for example through Monte-Carlo/fusion diagnostics, decoder packages, Markov-style routing summaries, or matrix-side readout layers — provided they remain downstream consumers of certified stage actions. What they may* not *do is replace a missing local dichotomy proof by a heuristic token guess, or overwrite an* OPENB_r *stop signal by pretending that a later advance has already been theorematically licensed.*

Reader cue. Read the next block as the finite word-level compression of the iterated extension protocol. Its purpose is not to replace the stagewise theorematic decisions, but to encode any already certified finite protocol segment into a short QAM/decoder-readable extension word whose shape itself records whether the segment remained purely advancing or terminated at its first honest boundary stage.

19AZ$^{(6aa)}$. Finite QAM/decoder word form for certified extension-action segments

After 19AZ$^{(6z)}$, the active line has a stagewise protocol, but later consumers may still want a shorter object than the full indexed action stream. The next honest compression step is therefore to pass from certified stage actions to a finite extension word. The compression must remain faithful to the non-skipping boundary discipline: once an earliest boundary action appears, the forward word must terminate there rather than pretending to continue past that point.

Definition H24.80 (Finite extension word alphabet and encoder)**.** *Fix a starting certified horizon* $\mathbf{K}_m^{\mathrm{cur}}$*. Introduce the two-letter extension alphabet*

$$\Sigma_{\mathrm{ext}} := \{\mathsf{A}, \mathsf{B}\},$$

read as

$$\mathsf{A} = \textit{advance the certified horizon by one stage},$$
$$\mathsf{B} = \textit{open earliest certified boundary kernel and stop.}$$

For each stage $r \geq m+1$*, define the stage encoder*

$$\mathsf{enc}_r : \mathcal{A}_r^{\mathrm{ext}} \to \Sigma_{\mathrm{ext}}$$

by

$$\mathsf{enc}_r(\mathsf{ADV}_r) = \mathsf{A}, \qquad \mathsf{enc}_r(\mathsf{OPENB}_r) = \mathsf{B}.$$

If a certified protocol segment runs from stage $m+1$ *through stage* N*, define its associated* finite extension word

$$\mathbf{w}_{m+1:N} := \mathsf{enc}_{m+1}(a_{m+1})\,\mathsf{enc}_{m+2}(a_{m+2})\cdots\mathsf{enc}_N(a_N) \in \Sigma_{\mathrm{ext}}^*,$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

with the convention that if some stage $r_* \le N$ satisfies $a_{r_*} = \mathsf{OPENB}_{r_*}$, then the word is truncated at r_*:

$$\mathbf{w}_{m+1:N} := \mathsf{enc}_{m+1}(a_{m+1}) \cdots \mathsf{enc}_{r_*}(a_{r_*}),$$

because the protocol is not licensed to advance past the earliest boundary stage.

Proposition H24.81 (Certified extension words have exactly the form A^k or $\mathsf{A}^k\mathsf{B}$)**.** *Assume the setting of Definition H24.76. Then every finite certified extension word produced by Definition H24.80 has exactly one of the following forms:*

$$\mathbf{w} = \mathsf{A}^k \qquad \text{or} \qquad \mathbf{w} = \mathsf{A}^k\mathsf{B}$$

for some integer $k \ge 0$. In particular:

(i) no certified extension word contains two occurrences of B*;*

(ii) no certified extension word contains B *followed by any later symbol;*

(iii) the word alone already records whether the observed certified segment is advance-persistent on its range or has terminated at its earliest honest boundary stage.

Proof. By Proposition H24.77, each certified stage contributes exactly one action, either ADV_r or OPENB_r. Under the encoder of Definition H24.80, these become A and B. If no stage in the observed range opens a boundary kernel, then every action is an advance action and the resulting word is A^k for the appropriate length k. If a boundary stage occurs, let r_* be the earliest one. The protocol stops at r_*, so the word is truncated there and has the form $\mathsf{A}^k\mathsf{B}$, where k counts the earlier advance stages. This proves the displayed classification, and clauses (i)–(iii) are immediate consequences of that classification. □

Corollary H24.82 (The extension protocol defines a regular terminal-word language)**.** *On every finite observation range, the certified extension words belong to the terminal language*

$$\mathcal{L}_{\mathrm{ext}}^{\mathrm{term}} = \{\mathsf{A}^k : k \ge 0\} \cup \{\mathsf{A}^k\mathsf{B} : k \ge 0\}.$$

Hence every later QAM, decoder, hashing, routing-summary, or Monte-Carlo/profile layer that consumes certified extension words must respect this language and may not treat words of the forbidden forms

$$\mathsf{BA}, \qquad \mathsf{ABA}, \qquad \mathsf{BB},$$

or any other word with a nonterminal boundary symbol as admissible proof-side inputs.

Proof. Immediate from Proposition H24.81. □

Remark H24.83 (What this word form now buys)**.** *This is the clean point where the active line becomes readable by a genuinely small working alphabet. The theorematic side still decides each local stage, but once those decisions are certified, later layers may consume the finite word* $\mathbf{w}$ *rather than the full local proof tree. That is the part of the later optimization discussion worth keeping: decoder maps, QAM readout, Markov summaries, Monte-Carlo deviation profiles, and matrix-side diagnostics may all work over the certified terminal language* $\mathcal{L}_{\mathrm{ext}}^{\mathrm{term}}$*. What they may not do is inject inadmissible words or continue reading letters after an honest terminal* B*-symbol has already appeared.*

Reader cue. Read the next block as the certificate-aware downstream matrix interpretation of the certified extension language. Its purpose is not to let the Heisenberg/Millennium side decide theorematic extension questions on its own, but to force every later matrix-side readout to distinguish sharply between certified horizon-growth and certified earliest-boundary exposure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

19AZ$^{(6ab)}$. Certificate-aware horizon/boundary interpretation for Heisenberg/Millennium matrix readout

After 19AZ$^{(6aa)}$, the active line carries a small certified extension language. The natural next use of that language is a downstream matrix-side interpretation: the Heisenberg/Millennium layer should no longer be read as an undifferentiated global diagnostic, but as a certificate-aware readout whose outputs are tagged by whether the active theorematic line has presently certified horizon continuation or exposed the earliest honest boundary stage. This keeps the matrix layer useful without allowing it to outrun the proof line.

Definition H24.84 (Certificate-aware Heisenberg/Millennium readout tags and readout map)**.** *Fix a current certified horizon* $\mathbf{K}_m^{\mathrm{cur}}$*, and let* $\mathbf{w} \in \mathcal{L}_{\mathrm{ext}}^{\mathrm{term}}$ *be a certified finite extension word in the sense of Corollary H24.82. Introduce the two matrix-side readout tags*

$$\mathsf{HZ}_m(\mathbf{w}) = \text{“horizon-certified matrix readout on the observed range”},$$

$$\mathsf{BD}_m(\mathbf{w}) = \text{“boundary-certified matrix readout on the observed range”}.$$

Define the certificate-aware downstream readout map

$$\mathsf{Read}_m^{\mathrm{MM/HC}}(\mathbf{w}) \in \{\mathsf{HZ}_m(\mathbf{w}), \mathsf{BD}_m(\mathbf{w})\}$$

by

$$\mathsf{Read}_m^{\mathrm{MM/HC}}(\mathbf{w}) = \begin{cases} \mathsf{HZ}_m(\mathbf{w}), & \text{if } \mathbf{w} = \mathsf{A}^k \text{ for some } k \geq 0, \\ \mathsf{BD}_m(\mathbf{w}), & \text{if } \mathbf{w} = \mathsf{A}^k\mathsf{B} \text{ for some } k \geq 0. \end{cases}$$

Interpretation:

- $\mathsf{HZ}_m(\mathbf{w})$ *means that, on the observed certified range, the matrix side is only licensed to read a horizon-persistent transport picture;*
- $\mathsf{BD}_m(\mathbf{w})$ *means that, on the observed certified range, the matrix side must switch to earliest-boundary exposure and may not read past the terminal boundary stage as though further certified transport had already been proved.*

Proposition H24.85 (Certificate-aware matrix readout is faithful to the certified extension language)**.** *Assume the setting of Definition H24.84. Then:*

(i) the readout map $\mathsf{Read}_m^{\mathrm{MM/HC}}$ *is well-defined on every certified finite extension word;*

(ii) the matrix-side tag distinguishes exactly the two proof-legitimate cases “advance-only on the observed range” and “advance-until-earliest-boundary on the observed range”;

(iii) no downstream Heisenberg/Millennium interpretation is licensed to convert a boundary-certified word $\mathsf{A}^k\mathsf{B}$ *into a horizon-certified continuation reading, and no horizon-certified word* A^k *is allowed to be advertised as exposing an already certified boundary event that the theorematic line has not produced.*

Proof. By Corollary H24.82, every certified finite extension word has one of the two forms A^k or $\mathsf{A}^k\mathsf{B}$. The definition therefore assigns exactly one matrix-side tag to every certified word, so the readout map is well-defined, proving (i). Since these are precisely the two admissible word forms, the map distinguishes exactly the two proof-legitimate cases listed in (ii). Clause (iii) is then just the statement that the downstream matrix layer must remain faithful to the certified word it receives: a terminal B-symbol cannot be reinterpreted as if certified continuation had already occurred, and the absence of a terminal B-symbol cannot be inflated into a claimed certified boundary exposure. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Corollary H24.86 (The present σ_3 package already induces a first certificate-aware matrix reading)**.** *In the current active line, the certified first-stage package from 19AZ$^{(6v)}$–19AZ$^{(6aa)}$ already induces a first certificate-aware Heisenberg/Millennium readout. Concretely, the first-stage word is either*

$$\mathbf{w}_3 = \mathsf{A} \qquad \textit{or} \qquad \mathbf{w}_3 = \mathsf{B},$$

and therefore the associated downstream matrix-side reading is already forced into exactly one of the two tags

$$\mathsf{Read}_2^{\mathrm{MM/HC}}(\mathbf{w}_3) \in \{\mathsf{HZ}_2(\mathbf{w}_3), \mathsf{BD}_2(\mathbf{w}_3)\}.$$

Thus the matrix side can now be asked a precise question: does the current certified stage read as horizon-persistent transport, or as earliest-boundary exposure?

Proof. Immediate from the first-stage encoding in 19AZ$^{(6v)}$–19AZ$^{(6aa)}$ together with Corollary H24.82 and Proposition H24.85. □

Remark H24.87 (What this matrix reading now buys for later crystal and field-equation upgrades)**.** *This is the clean interface the later upgrade discussion was missing. The Heisenberg/Millennium layer can now be used as a certificate-aware reader of the active extension state rather than as a competing source of theorematic authority. In particular, later crystal-layer refinements and later field-equation diagnostics should be formulated as refinements of either the horizon-certified readout tag or the boundary-certified readout tag. That means they may sharpen what the current certified state looks like, but they should not overrule whether the active line currently sits in a horizon-growth state or at an earliest-boundary exposure state. The same discipline keeps Monte-Carlo or decoder-side diagnostics honest: they may profile, compress, or compare the tagged state, but not replace the proof-side tag assignment itself.*

Reader cue. Read the next block as the certificate-aware downstream crystal interpretation of the matrix-side tag discipline. Its purpose is not to reopen the earlier addressed-crystal architecture at full generality, but to force every later crystal-side readout to distinguish sharply between horizon-persistent transport/domain structure and earliest-boundary defect exposure.

19AZ$^{(6ac)}$. Certificate-aware horizon/boundary interpretation for crystal-layer readout

After 19AZ$^{(6ab)}$, the active line no longer carries only a proof-side token language and a matrix-side tag. It also carries the first clean rule for how the later crystal layer is allowed to read the active extension state. The crystal side should therefore not be treated as a free global metaphor or as an independent decision engine. It should be read as a downstream structure-preserving refinement of the already certified horizon-versus-boundary status.

Definition H24.88 (Certificate-aware crystal readout tags and crystal-side readout map)**.** *Fix a current certified horizon $\mathbf{K}_m^{\mathrm{cur}}$, let $\mathbf{w} \in \mathcal{L}_{\mathrm{ext}}^{\mathrm{term}}$ be a certified finite extension word, and let*

$$\mathsf{Read}_m^{\mathrm{MM/HC}}(\mathbf{w}) \in \{\mathsf{HZ}_m(\mathbf{w}), \mathsf{BD}_m(\mathbf{w})\}$$

be the associated certificate-aware matrix-side tag from Definition H24.84. Introduce the two crystal-side readout tags

$$\mathsf{CRH}_m(\mathbf{w}) = \textit{“horizon-certified crystal/domain transport readout”},$$

$$\mathsf{CRB}_m(\mathbf{w}) = \textit{“boundary-certified crystal defect/front readout”}.$$

Define the certificate-aware crystal-side readout map

$$\mathsf{Read}_m^{\mathrm{cr}}(\mathbf{w}) \in \{\mathsf{CRH}_m(\mathbf{w}), \mathsf{CRB}_m(\mathbf{w})\}$$

by

$$\mathsf{Read}_m^{\mathrm{cr}}(\mathbf{w}) = \begin{cases} \mathsf{CRH}_m(\mathbf{w}), & \textit{if } \mathsf{Read}_m^{\mathrm{MM/HC}}(\mathbf{w}) = \mathsf{HZ}_m(\mathbf{w}), \\ \mathsf{CRB}_m(\mathbf{w}), & \textit{if } \mathsf{Read}_m^{\mathrm{MM/HC}}(\mathbf{w}) = \mathsf{BD}_m(\mathbf{w}). \end{cases}$$

Interpretation:

- $\mathsf{CRH}_m(\mathbf{w})$ *means that the crystal side is licensed to read the currently certified range as one transport-persistent domain picture, with no theorematically certified defect-front beyond that range yet exposed;*
- $\mathsf{CRB}_m(\mathbf{w})$ *means that the crystal side must treat the terminal certified information as an earliest honest defect/front exposure and may not smooth, continue, or visually homogenize that boundary as though later corridor coherence had already been proved.*

Proposition H24.89 (Certificate-aware crystal readout is faithful to the matrix-side horizon/boundary tag)**.** *Assume the setting of Definition H24.88. Then:*

(i) the crystal-side readout map $\mathsf{Read}_m^{\mathrm{cr}}$ *is well-defined on every certified finite extension word;*

(ii) the crystal-side tag distinguishes exactly the two proof-legitimate crystal readings “domain transport on the certified range” and “earliest defect/front exposure at the certified boundary stage”;

(iii) no downstream crystal-layer interpretation is licensed to smooth a boundary-certified state into a transport-certified domain picture, and no horizon-certified state is allowed to be advertised as already carrying a proved defect/front event that the active line has not certified.

Proof. By Proposition H24.85, the matrix-side readout assigns exactly one of the two tags $\mathsf{HZ}_m(\mathbf{w})$ or $\mathsf{BD}_m(\mathbf{w})$ to every certified word. Definition H24.88 therefore assigns exactly one crystal-side tag $\mathsf{CRH}_m(\mathbf{w})$ or $\mathsf{CRB}_m(\mathbf{w})$, proving (i). Since these two tags simply refine the two proof-legitimate matrix-side statuses into crystal/domain language, statement (ii) follows immediately. Clause (iii) is the same faithfulness discipline expressed on the crystal side: a boundary-certified terminal stage cannot be visually or narratively turned into uninterrupted continuation, and a transport-certified range cannot be embellished with a certified defect/front event that the theorematic line has not produced. □

Corollary H24.90 (The present σ_3 package already induces a first certificate-aware crystal reading)**.** *In the current active line, the first-stage word* $\mathbf{w}_3 \in \{\mathsf{A}, \mathsf{B}\}$ *already induces a first certificate-aware crystal-side readout*

$$\mathsf{Read}_2^{\mathrm{cr}}(\mathbf{w}_3) \in \{\mathsf{CRH}_2(\mathbf{w}_3), \mathsf{CRB}_2(\mathbf{w}_3)\}.$$

Thus the crystal layer can now be asked a precise and conservative question: does the present certified stage still read as one transport-persistent domain picture, or has the earliest honest defect/front already been exposed at the current terminal stage?

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. Immediate from Corollary H24.86 and Proposition H24.89. □

Remark H24.91 (What this crystal reading now buys)**.** *This is the clean point where the earlier liquid-crystal/domain/defect language becomes structurally disciplined by the active extension certificates. The crystal side may now refine the current state into domain persistence, defect loading, or front exposure language without pretending to certify a later slot on its own. In particular, later holographic boundary visuals, domain-defect compression schemes, and Monte-Carlo or decoder-side pattern scans may profile the crystal-side tag, but they must inherit it from the certified extension language rather than invent it.*

Reader cue. Read the next block as the certificate-aware downstream pre-equation interface for field-equation diagnostics. Its purpose is not to claim a finished Einstein-, Dirac-, or QFT-type projection law at this stage, but to force every later field-equation ansatz to distinguish sharply between a horizon-certified continuation profile and an earliest-boundary source/exposure profile.

19AZ$^{(6ad)}$. Certificate-aware horizon/boundary interpretation for field-equation diagnostics

After 19AZ$^{(6ac)}$, the active line has a theorematically certified extension language, a matrix-side readout, and a crystal-side readout. The next honest compression step is not yet a full new field equation. It is a certificate-aware pre-equation interface that tells any later physical or PDE-style readout whether it is looking at a continuation-certified transport profile or at an earliest-boundary exposure profile. This is the smallest disciplined upgrade that the current line can honestly support.

Definition H24.92 (Certificate-aware field-diagnostic tags and pre-equation map)**.** *Fix a current certified horizon* $\mathbf{K}_m^{\mathrm{cur}}$*, let* $\mathbf{w} \in \mathcal{L}_{\mathrm{ext}}^{\mathrm{term}}$ *be a certified finite extension word, and let*

$$\mathsf{Read}_m^{\mathrm{cr}}(\mathbf{w}) \in \{\mathsf{CRH}_m(\mathbf{w}), \mathsf{CRB}_m(\mathbf{w})\}$$

be the associated crystal-side readout from Definition H24.88. Introduce the two field-diagnostic tags

$$\mathsf{FEH}_m(\mathbf{w}) = \text{“horizon-certified pre-equation transport profile”},$$

$$\mathsf{FEB}_m(\mathbf{w}) = \text{“boundary-certified pre-equation source/exposure profile”}.$$

Define the certificate-aware field-diagnostic map

$$\mathsf{Diag}_m^{\mathrm{field}}(\mathbf{w}) \in \{\mathsf{FEH}_m(\mathbf{w}), \mathsf{FEB}_m(\mathbf{w})\}$$

by

$$\mathsf{Diag}_m^{\mathrm{field}}(\mathbf{w}) = \begin{cases} \mathsf{FEH}_m(\mathbf{w}), & \text{if } \mathsf{Read}_m^{\mathrm{cr}}(\mathbf{w}) = \mathsf{CRH}_m(\mathbf{w}), \\ \mathsf{FEB}_m(\mathbf{w}), & \text{if } \mathsf{Read}_m^{\mathrm{cr}}(\mathbf{w}) = \mathsf{CRB}_m(\mathbf{w}). \end{cases}$$

Interpretation:

- $\mathsf{FEH}_m(\mathbf{w})$ *means that any later field-equation ansatz is presently licensed only to read continuation-compatible transport on the already certified range;*
- $\mathsf{FEB}_m(\mathbf{w})$ *means that any later field-equation ansatz must already acknowledge an earliest honest boundary/source exposure at the terminal certified stage and may not write a smooth forward continuation law past that stage as though the proof line had already certified it.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proposition H24.93 (Certificate-aware field diagnostics are faithful to the crystal-side tag)**.** *Assume the setting of Definition H24.92. Then:*

(i) the field-diagnostic map $\mathsf{Diag}_m^{\mathrm{field}}$ *is well-defined on every certified finite extension word;*

(ii) the field-diagnostic tag distinguishes exactly the two presently legitimate pre-equation situations "continuation-compatible certified transport" and "earliest-boundary source/exposure already active";

(iii) no downstream field-equation ansatz is licensed to erase a terminal boundary-certified status, and no continuation-certified status is allowed to be sold as a proved boundary/source event that the active line has not certified.

Proof. By Proposition H24.89, the crystal-side readout assigns exactly one of the tags $\mathsf{CRH}_m(\mathbf{w})$ or $\mathsf{CRB}_m(\mathbf{w})$ to every certified word. Definition H24.92 therefore assigns exactly one field-diagnostic tag $\mathsf{FEH}_m(\mathbf{w})$ or $\mathsf{FEB}_m(\mathbf{w})$, proving (i). Since these are precisely the two downstream field-diagnostic refinements of the crystal-side statuses, statement (ii) follows. Clause (iii) is the same certificate faithfulness expressed at the pre-equation layer: a certified terminal boundary/source exposure cannot be rewritten as already-proved smooth continuation, and a certified transport range cannot be exaggerated into a proved boundary event. □

Corollary H24.94 (The present σ_3 package already induces a first certificate-aware field-diagnostic split)**.** *In the current active line, the first-stage word* $\mathbf{w}_3 \in \{\mathsf{A}, \mathsf{B}\}$ *already induces a first certificate-aware field-diagnostic split*

$$\mathsf{Diag}_2^{\mathrm{field}}(\mathbf{w}_3) \in \{\mathsf{FEH}_2(\mathbf{w}_3), \mathsf{FEB}_2(\mathbf{w}_3)\}.$$

Thus the later field-equation layer can already be asked a disciplined question: is the current certified stage to be read as continuation-compatible transport on the observed range, or as an earliest-boundary source/exposure profile that must remain explicit in every later PDE or operator ansatz?

Proof. Immediate from Corollary H24.90 and Proposition H24.93. □

Remark H24.95 (What this field-diagnostic interface now buys)**.** *This is the conservative upgrade the later field-equation discussion was missing. The active line still does not claim a finished physical PDE, a finished Einstein equation, or a finished emergence law. What it now does claim is a disciplined pre-equation interface: every later ansatz must declare whether it is reading a horizon-certified transport state or an earliest-boundary source/exposure state. That already sharpens admissibility. A later field equation is acceptable only if its local readout remains faithful to the certified tag it inherits from the extension language.*

Reader cue. Read the next block as the admissibility and consistency gate for any later PDE-, operator-, or emergence-style ansatz built on the certificate-aware field-diagnostic interface. Its purpose is not to force one unique final field equation, but to exclude exactly those later ansätze that would erase, overstate, or misroute the certified horizon/boundary status already inherited from the active extension line.

19AZ$^{(6ae)}$. Admissibility and consistency conditions for later field-equation ansätze

After 19AZ$^{(6ad)}$, the active line still does not claim a unique finished field equation. What it does claim is stronger than a vague heuristic: any later PDE-, operator-, or emergence-style law must now pass a certificate-aware admissibility gate. In other words, once a later

ansatz inherits the field-diagnostic tag $\mathsf{FEH}_m(\mathbf{w})$ or $\mathsf{FEB}_m(\mathbf{w})$, it is no longer free to ignore the certified status that generated that tag.

Definition H24.96 (Certificate-aware admissibility for later field-equation ansätze)**.** *Fix a current certified horizon $\mathbf{K}_m^{\mathrm{cur}}$, let $\mathbf{w} \in \mathcal{L}_{\mathrm{ext}}^{\mathrm{term}}$ be a certified finite extension word, and let*

$$\mathsf{Diag}_m^{\mathrm{field}}(\mathbf{w}) \in \{\mathsf{FEH}_m(\mathbf{w}), \mathsf{FEB}_m(\mathbf{w})\}$$

be the inherited field-diagnostic tag from Definition H24.92. A later PDE-, operator-, transport-, or emergence-style ansatz

$$\mathfrak{E}_{m,\mathbf{w}}$$

is called certificate-aware admissible *if it satisfies the following conditions.*

(i) ***Tag fidelity.*** *The ansatz must explicitly declare whether it is reading the state under $\mathsf{FEH}_m(\mathbf{w})$ or under $\mathsf{FEB}_m(\mathbf{w})$, and its local interpretation must respect that inherited tag.*

(ii) ***No boundary erasure.*** *If the inherited tag is $\mathsf{FEB}_m(\mathbf{w})$, the ansatz may not replace the terminal certified boundary/source exposure by a smooth forward continuation law as though the active line had already certified later corridor transport.*

(iii) ***No boundary inflation.*** *If the inherited tag is $\mathsf{FEH}_m(\mathbf{w})$, the ansatz may not advertise an already certified source, defect, singularity, or terminal boundary event that the active line has not produced.*

(iv) ***Range discipline.*** *The ansatz may refine, profile, or parameterize the already certified observed range, but it may not claim theorematic authority beyond the range encoded by the certified word $\mathbf{w}$.*

If one of these conditions fails, the ansatz is called certificate-aware inadmissible *on the input pair $(m, \mathbf{w})$.*

Proposition H24.97 (The certificate-aware field interface induces a sharp admissibility dichotomy)**.** *Assume the setting of Definition H24.96. Then every later field-equation ansatz $\mathfrak{E}_{m,\mathbf{w}}$ is forced into exactly one of the two statuses:*

$$\textit{certificate-aware admissible} \qquad \textit{or} \qquad \textit{certificate-aware inadmissible.}$$

Moreover:

(i) every admissible ansatz is a downstream refinement of the inherited tag rather than a replacement for it;

(ii) every inadmissible ansatz fails by one of exactly three routes: boundary erasure, boundary inflation, or range overreach;

(iii) this admissibility test is independent of whether the later ansatz is written in matrix, crystal, PDE, operator, Monte-Carlo-assisted, or decoder-compressed language.

Proof. Definition H24.96 gives explicit necessary and sufficient conditions for admissibility on the input pair $(m, \mathbf{w})$. Hence every later ansatz either satisfies all of them or fails at least one, which proves the displayed dichotomy. Statement (i) is immediate from tag fidelity together with range discipline: an admissible ansatz inherits the certified field tag and may only refine the already certified state. For (ii), the only ways to violate the conditions are

© 2026 Marc Brendecke. Preprint — not peer reviewed.

exactly the ones listed in the definition: either an inherited boundary-certified state is erased, an inherited continuation-certified state is exaggerated into a certified boundary/source event, or the ansatz claims interpretive authority beyond the certified observed range. Statement (iii) follows because the admissibility test depends only on the inherited certified tag and on faithfulness to the certified word, not on the notation or computational packaging chosen downstream. □

Corollary H24.98 (The present σ_3 package already excludes the wrong kinds of field-law upgrades)**.** *In the current active line, the first-stage package with* $\mathbf{w}_3 \in \{\mathsf{A}, \mathsf{B}\}$ *already excludes two opposite classes of later upgrade errors:*

(i) if $\mathbf{w}_3 = \mathsf{B}$*, no later matrix-, crystal-, or field-law ansatz is allowed to smooth that first certified boundary exposure into an already proved forward continuation picture;*

(ii) if $\mathbf{w}_3 = \mathsf{A}$*, no later matrix-, crystal-, or field-law ansatz is allowed to present the current stage as though an already certified boundary/source event had been exposed.*

Thus even before a finished field equation exists, the present line already knows which later upgrade moves are structurally forbidden.

Proof. Immediate from Corollary H24.94 and Proposition H24.97. □

Remark H24.99 (What this admissibility gate now buys)**.** *This is the point where the current line becomes strong enough to police its own future upgrades. A later Einstein-style, transport-style, source-law, Monte-Carlo-assisted closure law, or operator-valued emergence ansatz no longer has to be rejected because it is ambitious; it has to be rejected only if it violates the inherited certificate. That is exactly the right level of rigidity at the present stage: permissive about representation, strict about structural faithfulness.*

Reader cue. Read the next block as the certificate-aware upgrade-routing interface that tells every later refinement layer which type of work is legitimately allowed after a field-diagnostic tag has been inherited. Its purpose is not to force one unique downstream model, but to route conservative continuation-style upgrades to the horizon-certified branch and boundary-profiling upgrades to the boundary-certified branch, while excluding cross-routings that would reopen or falsify the certified status.

19AZ$^{(6af)}$. Upgrade-routing consequences of certificate-aware field-diagnostic tags

After 19AZ$^{(6ae)}$, the active line has more than a static admissibility gate. It now has enough structure to route later upgrades according to the inherited certificate. The point is simple: not every technically sophisticated downstream refinement is appropriate for every certified word. A horizon-certified tag should route later work toward conservative continuation-style refinement on the already proved range, while a boundary-certified tag should route later work toward earliest-boundary profiling, exposure analysis, and stop-sensitive diagnostics. This routing discipline is the missing interface between the theorematic extension line and later matrix-, crystal-, field-, decoder-, or probe-based elaborations.

Definition H24.100 (Certificate-aware upgrade-routing map)**.** *Fix a current certified horizon* $\mathbf{K}_m^{\mathrm{cur}}$*, let* $\mathbf{w} \in \mathcal{L}_{\mathrm{ext}}^{\mathrm{term}}$ *be a certified finite extension word, and let*

$$\mathsf{Diag}_m^{\mathrm{field}}(\mathbf{w}) \in \{\mathsf{FEH}_m(\mathbf{w}), \mathsf{FEB}_m(\mathbf{w})\}$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

be the inherited field-diagnostic tag from Definition H24.92. Introduce the two conservative upgrade-route classes

$$\mathcal{U}_m^{\mathrm{hor}}(\mathbf{w}) := \text{“continuation-compatible refinement on the already certified range”},$$
$$\mathcal{U}_m^{\mathrm{bdry}}(\mathbf{w}) := \text{“earliest-boundary profiling and stop-sensitive exposure analysis”}.$$

Define the certificate-aware upgrade-routing map

$$\mathsf{Route}_m^{\mathrm{upg}}(\mathbf{w}) \in \{\mathcal{U}_m^{\mathrm{hor}}(\mathbf{w}), \mathcal{U}_m^{\mathrm{bdry}}(\mathbf{w})\}$$

by

$$\mathsf{Route}_m^{\mathrm{upg}}(\mathbf{w}) = \begin{cases} \mathcal{U}_m^{\mathrm{hor}}(\mathbf{w}), & \text{if } \mathsf{Diag}_m^{\mathrm{field}}(\mathbf{w}) = \mathsf{FEH}_m(\mathbf{w}), \\ \mathcal{U}_m^{\mathrm{bdry}}(\mathbf{w}), & \text{if } \mathsf{Diag}_m^{\mathrm{field}}(\mathbf{w}) = \mathsf{FEB}_m(\mathbf{w}). \end{cases}$$

Interpretation:

- *the horizon-certified route $\mathcal{U}_m^{\mathrm{hor}}(\mathbf{w})$ licenses only those later upgrades that refine transport, matrix readout, crystal stabilization, or PDE/operator behavior on the range already certified by $\mathbf{w}$, without claiming a new boundary/source event;*
- *the boundary-certified route $\mathcal{U}_m^{\mathrm{bdry}}(\mathbf{w})$ licenses only those later upgrades that sharpen the earliest-boundary profile itself, for example by defect localization, source/exposure analysis, or stop-sensitive decoder/probe diagnostics, without pretending that corridor continuation past the terminal boundary has already been proved;*
- *any downstream refinement that crosses these route classes against the inherited tag is called a* cross-routed upgrade *and is provisionally inadmissible until a new local theorematic stage changes the certified status.*

Proposition H24.101 (The inherited field-diagnostic tag forces a unique conservative upgrade route)**.** *Assume the setting of Definition H24.100. Then:*

(i) the routing map $\mathsf{Route}_m^{\mathrm{upg}}$ is well-defined on every certified finite extension word;

(ii) a horizon-certified tag $\mathsf{FEH}_m(\mathbf{w})$ forces the conservative route class $\mathcal{U}_m^{\mathrm{hor}}(\mathbf{w})$, while a boundary-certified tag $\mathsf{FEB}_m(\mathbf{w})$ forces the conservative route class $\mathcal{U}_m^{\mathrm{bdry}}(\mathbf{w})$;

(iii) every cross-routed upgrade is incompatible with the certificate-aware admissibility conditions from Definition H24.96 unless and until a new theorematic local stage certifies a different status.

Proof. By Proposition H24.97, every certified word inherits exactly one field-diagnostic tag, either $\mathsf{FEH}_m(\mathbf{w})$ or $\mathsf{FEB}_m(\mathbf{w})$. Definition H24.100 therefore assigns exactly one conservative route class, proving (i). Clause (ii) is the explicit content of the definition. For (iii), suppose a downstream refinement is routed against the inherited tag. If a boundary-certified word is forced onto the continuation-style route, then the upgrade attempts boundary erasure by treating an earliest-boundary exposure as already-certified transport continuation. If a horizon-certified word is forced onto the boundary-style route, then the upgrade attempts boundary inflation by advertising a defect, source, or stopping event not certified by the active line. Both behaviors violate the admissibility gate from Definition H24.96. Hence cross-routed upgrades are incompatible with the present certificate unless a later local theorematic stage genuinely changes that certificate. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Corollary H24.102 (The present σ_3 package already induces a first conservative upgrade-routing split)**.** *In the current active line, the first-stage word* $\mathbf{w}_3 \in \{\mathsf{A},\mathsf{B}\}$ *already induces a first conservative routing split*

$$\mathsf{Route}_2^{\mathrm{upg}}(\mathbf{w}_3) \in \{\mathcal{U}_2^{\mathrm{hor}}(\mathbf{w}_3), \mathcal{U}_2^{\mathrm{bdry}}(\mathbf{w}_3)\}.$$

Thus the next downstream question is no longer vague. If the inherited tag is horizon-certified, later work should concentrate on transport-compatible refinement of the certified range. If the inherited tag is boundary-certified, later work should concentrate on profiling the earliest honest boundary rather than pretending that the corridor has already been extended through it.

Proof. Immediate from Corollary H24.94 and Proposition H24.101. □

Remark H24.103 (What this upgrade-routing interface now buys)**.** *This is the first place where the active line tells later heuristic or computational layers not only what they may* say*, but also what they should* do next*. Matrix sharpening, crystal stabilization, conservative PDE refinement, decoder compression, Monte-Carlo probing, or source/exposure diagnostics are now routed by the inherited certificate rather than by taste. In particular, Monte-Carlo, decoder, hash, or Markov-style probes may read and compress the routed status, but they are not allowed to replace the theorematic routing decision itself. That keeps the later tool layer useful without allowing it to overrule the certified extension line.*

Reader cue. Read the next block as the normalized follow-on upgrade menu sitting *below* the certificate-aware routing interface. Its purpose is to say which concrete families of downstream refinement are genuinely licensed once a route has already been fixed, and which families remain out of bounds until a later local theorematic stage changes the certificate.

19AZ$^{(6ag)}$. Normalized follow-on upgrade families below the certificate-aware routing interface

After 19AZ$^{(6af)}$, the line knows not only whether a later refinement is admissible, but also which route it must follow. The next honest compression step is to normalize the concrete follow-on upgrade families that sit below those routes. This is important because later readers should not have to guess whether, for example, matrix sharpening, crystal stabilization, source localization, decoder compression, or Monte-Carlo probing belongs to the continuation side or to the earliest-boundary side. The present block turns that tacit distinction into a finite, certificate-aware menu.

Definition H24.104 (Normalized follow-on upgrade families)**.** *Fix a current certified horizon* $\mathbf{K}_m^{\mathrm{cur}}$*, let* $\mathbf{w} \in \mathcal{L}_{\mathrm{ext}}^{\mathrm{term}}$ *be a certified finite extension word, and let*

$$\mathsf{Route}_m^{\mathrm{upg}}(\mathbf{w}) \in \{\mathcal{U}_m^{\mathrm{hor}}(\mathbf{w}), \mathcal{U}_m^{\mathrm{bdry}}(\mathbf{w})\}$$

be the inherited upgrade route from Definition H24.100.

Define the horizon-follow-on family menu

$$\mathcal{F}_m^{\mathrm{hor}}(\mathbf{w}) := \{\mathsf{MSH}_m(\mathbf{w}), \mathsf{CRS}_m(\mathbf{w}), \mathsf{PDE}_m^{\mathrm{cons}}(\mathbf{w}), \mathsf{DEC}_m^{\mathrm{hor}}(\mathbf{w})\}$$

with the intended meanings

$$\begin{aligned}
\mathsf{MSH}_m(\mathbf{w}) &:= \text{“}\textit{matrix sharpening on the already certified range}\text{”},\\
\mathsf{CRS}_m(\mathbf{w}) &:= \text{“}\textit{crystal stabilization / transport-consistency refinement}\text{”},\\
\mathsf{PDE}_m^{\mathrm{cons}}(\mathbf{w}) &:= \text{“}\textit{conservative PDE/operator refinement on the certified range}\text{”},\\
\mathsf{DEC}_m^{\mathrm{hor}}(\mathbf{w}) &:= \text{“}\textit{decoder / compression readout of a horizon-certified status}\text{”}.
\end{aligned}$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Define the boundary-follow-on family menu

$$\mathcal{F}_m^{\mathrm{bdry}}(\mathbf{w}) := \{\mathsf{SRC}_m(\mathbf{w}), \mathsf{DFX}_m(\mathbf{w}), \mathsf{MC}_m^{\mathrm{bdry}}(\mathbf{w}), \mathsf{DEC}_m^{\mathrm{bdry}}(\mathbf{w})\}$$

with the intended meanings

$$\begin{aligned}
\mathsf{SRC}_m(\mathbf{w}) &:= \text{“source / exposure localization at the earliest honest boundary”},\\
\mathsf{DFX}_m(\mathbf{w}) &:= \text{“defect-front or crystal-boundary profiling”},\\
\mathsf{MC}_m^{\mathrm{bdry}}(\mathbf{w}) &:= \text{“stop-sensitive Monte-Carlo or probe diagnostics at the boundary side”},\\
\mathsf{DEC}_m^{\mathrm{bdry}}(\mathbf{w}) &:= \text{“decoder / compression readout of a boundary-certified status”}.
\end{aligned}$$

A normalized follow-on upgrade *for* $\mathbf{w}$ *is then any element of the route-compatible menu*

$$\mathcal{F}_m^{\mathrm{follow}}(\mathbf{w}) := \begin{cases} \mathcal{F}_m^{\mathrm{hor}}(\mathbf{w}), & \text{if } \mathsf{Route}_m^{\mathrm{upg}}(\mathbf{w}) = \mathcal{U}_m^{\mathrm{hor}}(\mathbf{w}),\\ \mathcal{F}_m^{\mathrm{bdry}}(\mathbf{w}), & \text{if } \mathsf{Route}_m^{\mathrm{upg}}(\mathbf{w}) = \mathcal{U}_m^{\mathrm{bdry}}(\mathbf{w}). \end{cases}$$

Any proposed later refinement that selects an element outside $\mathcal{F}_m^{\mathrm{follow}}(\mathbf{w})$ *is called a* route-incompatible follow-on upgrade.

Proposition H24.105 (Every certified word carries a finite route-compatible follow-on upgrade menu)**.** *Assume the setting of Definition H24.104. Then:*

(i) every certified finite extension word $\mathbf{w}$ *carries exactly one route-compatible follow-on family menu* $\mathcal{F}_m^{\mathrm{follow}}(\mathbf{w})$*;*

(ii) every normalized follow-on upgrade in $\mathcal{F}_m^{\mathrm{follow}}(\mathbf{w})$ *preserves the certificate-aware routing discipline from* $19AZ^{(6af)}$*;*

(iii) every route-incompatible follow-on upgrade attempts either continuation overreach on the boundary side or spurious defect/source inflation on the horizon side, and is therefore provisionally inadmissible.

Proof. By Proposition H24.101, every certified finite extension word inherits exactly one conservative upgrade route. Definition H24.104 assigns to that unique route exactly one finite menu, proving (i). Clause (ii) is built into the meanings of the menu entries: the horizon-family items refine only the already certified continuation range, while the boundary-family items profile only the earliest certified boundary or exposure picture. For (iii), if a boundary-certified word is sent to a horizon-family item, the proposed refinement necessarily behaves as though continuation beyond the terminal boundary were already certified; this is continuation overreach. If a horizon-certified word is sent to a boundary-family item, the proposed refinement injects a defect, source, or stopping narrative that has not been certified by the active line; this is spurious inflation. Both violate the route discipline and so are provisionally inadmissible. □

Corollary H24.106 (The present σ_3 package already carries a finite next-work menu)**.** *In the current active line, the first-stage word* $\mathbf{w}_3 \in \{\mathsf{A}, \mathsf{B}\}$ *already carries a finite next-work menu*

$$\mathcal{F}_2^{\mathrm{follow}}(\mathbf{w}_3) \subseteq \mathcal{F}_2^{\mathrm{hor}}(\mathbf{w}_3) \cup \mathcal{F}_2^{\mathrm{bdry}}(\mathbf{w}_3).$$

Thus the immediate next work is no longer only route-correct in the abstract. It is already normalized into a finite family of concrete continuation-side or boundary-side upgrade tasks.

Proof. Immediate from Corollary H24.102 and Proposition H24.105. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Remark H24.107 (What this finite next-work menu now buys)**.** *This is the point where the active line starts to look like a genuine operator's manual for the next minor release. Matrix sharpening, crystal stabilization, conservative PDE refinement, decoder compression, source localization, defect profiling, and stop-sensitive Monte-Carlo probing are no longer a loose wishlist. They have become a finite, certificate-aware menu. In particular, Monte-Carlo now has a safe place: it can act as a boundary-side structural probe when the inherited status is boundary-certified, but it is not allowed to masquerade as theorematic continuation proof.*

oindent extbfReader cue. Read the next block as the explicit QAM-/decoder-side shadow of the certificate-aware extension interface. Its purpose is not to introduce a new ontology above the current tool layer, but to compress the already inherited field tag, route, and next-work menu into one finite downstream object that QAM- and decoder-style readers may consume without reopening the theorematic proof split.

19AZ$^{(6ah)}$. QAM-/decoder-side shadow of the certificate-aware extension interface

After 19AZ$^{(6ag)}$, the active line already carries everything a later QAM- or decoder-side layer needs in order to act conservatively: a certified finite word, a field-diagnostic tag, a forced route, and a finite next-work menu. The next honest compression step is therefore not a new theorem, but a finite shadow object that packages these inherited data into one downstream-readable interface. This is exactly the point where the present certificate-aware tool layer becomes explicitly QAM-/decoder-consumable.

Definition H24.108 (QAM-/decoder-side shadow state)**.** *Fix a current certified horizon* $\mathbf{K}_m^{\mathrm{cur}}$*, let* $\mathbf{w} \in \mathcal{L}_{\mathrm{ext}}^{\mathrm{term}}$ *be a certified finite extension word, and let*

$$\mathsf{Diag}_m^{\mathrm{field}}(\mathbf{w}) \in \{\mathsf{FEH}_m(\mathbf{w}), \mathsf{FEB}_m(\mathbf{w})\}, \qquad \mathsf{Route}_m^{\mathrm{upg}}(\mathbf{w}) \in \{\mathcal{U}_m^{\mathrm{hor}}(\mathbf{w}), \mathcal{U}_m^{\mathrm{bdry}}(\mathbf{w})\},$$

and

$$\mathcal{F}_m^{\mathrm{follow}}(\mathbf{w}) \in \{\mathcal{F}_m^{\mathrm{hor}}(\mathbf{w}), \mathcal{F}_m^{\mathrm{bdry}}(\mathbf{w})\}$$

be the inherited field tag, upgrade route, and follow-on menu from 19AZ$^{(6ad)}$–19AZ$^{(6ag)}$.

Define the two finite QAM-shadow states

$$\mathsf{QSH}_m^{\mathrm{hor}}(\mathbf{w}) := \big\langle \mathsf{FEH}_m(\mathbf{w}), \mathcal{U}_m^{\mathrm{hor}}(\mathbf{w}), \mathcal{F}_m^{\mathrm{hor}}(\mathbf{w}) \big\rangle,$$
$$\mathsf{QSH}_m^{\mathrm{bdry}}(\mathbf{w}) := \big\langle \mathsf{FEB}_m(\mathbf{w}), \mathcal{U}_m^{\mathrm{bdry}}(\mathbf{w}), \mathcal{F}_m^{\mathrm{bdry}}(\mathbf{w}) \big\rangle.$$

Define the QAM-/decoder-side shadow map

$$\mathsf{Shadow}_m^{\mathrm{QAM}}(\mathbf{w}) \in \{\mathsf{QSH}_m^{\mathrm{hor}}(\mathbf{w}), \mathsf{QSH}_m^{\mathrm{bdry}}(\mathbf{w})\}$$

by

$$\mathsf{Shadow}_m^{\mathrm{QAM}}(\mathbf{w}) = \begin{cases} \mathsf{QSH}_m^{\mathrm{hor}}(\mathbf{w}), & \text{if } \mathsf{Diag}_m^{\mathrm{field}}(\mathbf{w}) = \mathsf{FEH}_m(\mathbf{w}), \\ \mathsf{QSH}_m^{\mathrm{bdry}}(\mathbf{w}), & \text{if } \mathsf{Diag}_m^{\mathrm{field}}(\mathbf{w}) = \mathsf{FEB}_m(\mathbf{w}). \end{cases}$$

Define moreover the associated decoder word

$$\mathsf{Word}_m^{\mathrm{dec}}(\mathbf{w}) \in \{\mathsf{HR}, \mathsf{BR}\}$$

by

$$\mathsf{Word}_m^{\mathrm{dec}}(\mathbf{w}) = \begin{cases} \mathsf{HR}, & \text{if } \mathsf{Shadow}_m^{\mathrm{QAM}}(\mathbf{w}) = \mathsf{QSH}_m^{\mathrm{hor}}(\mathbf{w}), \\ \mathsf{BR}, & \text{if } \mathsf{Shadow}_m^{\mathrm{QAM}}(\mathbf{w}) = \mathsf{QSH}_m^{\mathrm{bdry}}(\mathbf{w}). \end{cases}$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Interpretation: the QAM shadow is the finite state object that a later QAM-, decoder-, compression-, or router-layer is allowed to consume; the decoder word is the shortest faithful tag that still remembers whether the inherited certificate is horizon-routed or boundary-routed.

Proposition H24.109 (The QAM-/decoder-side shadow is a faithful finite downstream image of the certificate-aware interface)**.** *Assume the setting of Definition H24.108. Then:*

(i) the shadow map $\mathsf{Shadow}_m^{\mathrm{QAM}}$ *is well-defined on every certified finite extension word;*

(ii) the shadow state remembers exactly the inherited field tag, forced upgrade route, and route-compatible next-work menu, but does not add new theorematic authority;

(iii) the decoder word $\mathsf{Word}_m^{\mathrm{dec}}(\mathbf{w}) \in \{\mathsf{HR}, \mathsf{BR}\}$ *is a faithful two-symbol readout of whether the certified state is horizon-routed or boundary-routed.*

Proof. By Proposition H24.101 and Proposition H24.105, every certified finite extension word inherits exactly one field tag, one forced route, and one route-compatible next-work menu. Definition H24.108 therefore assigns exactly one of the two shadow states, proving (i). Statement (ii) follows because the shadow state is merely an ordered packaging of already inherited data; it compresses the certified status but does not certify anything beyond it. For (iii), the decoder word forgets the internal menu details while preserving the decisive distinction between horizon-routed and boundary-routed certified status. Hence it is a faithful two-symbol downstream readout of the certificate-aware interface. □

Corollary H24.110 (The present σ_3 package already induces a first explicit QAM-/decoder-side shadow)**.** *In the current active line, the first-stage word* $\mathbf{w}_3 \in \{\mathsf{A}, \mathsf{B}\}$ *already induces both a finite QAM-shadow state*

$$\mathsf{Shadow}_2^{\mathrm{QAM}}(\mathbf{w}_3) \in \{\mathsf{QSH}_2^{\mathrm{hor}}(\mathbf{w}_3), \mathsf{QSH}_2^{\mathrm{bdry}}(\mathbf{w}_3)\}$$

and a faithful decoder word

$$\mathsf{Word}_2^{\mathrm{dec}}(\mathbf{w}_3) \in \{\mathsf{HR}, \mathsf{BR}\}.$$

Thus the new extension tool layer is now explicitly attached back to the QAM-/decoder side as a finite downstream interface.

Proof. Immediate from Corollary H24.106 and Proposition H24.109. □

Remark H24.111 (What this QAM-/decoder-side shadow now buys)**.** *This is the missing explicit reattachment of the new tool layer to the QAM line. The active extension architecture can now be consumed as a finite state object instead of as a long theorematic prose chain. That is exactly what a later QAM, decoder, compression, or router implementation should see: not a new ontology, but a faithful shadow of the already certified status. In particular, the decoder word may be used for compact routing, logging, or interface normalization, but it may never be mistaken for an independent proof of continuation or boundary exposure.*

Reader cue. Read the next block as the explicit Monte-Carlo/probe discipline sitting below the certificate-aware routing and menu interface. Its purpose is to give Monte-Carlo a precise and honest role inside the current architecture: useful as a downstream structural probe and deviation profiler, but never as a substitute for theorematic certification.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

19AZ$^{(6ai)}$. Certificate-aware Monte-Carlo / probe discipline below the routing interface

After 19AZ$^{(6ag)}$ and 19AZ$^{(6ah)}$, the active line has enough structure to place Monte-Carlo and related probe machinery exactly where it belongs. The point is not to diminish that machinery, but to protect it from doing the wrong job. A good Monte-Carlo run may expose deviation patterns, stop-sensitive fingerprints, boundary clustering, transport-instability signals, or decoder-compressible distributions. What it may not do is certify a new horizon extension or erase a certified boundary stop. The present block fixes that discipline explicitly.

Definition H24.112 (Certificate-aware Monte-Carlo / probe family)**.** *Fix a current certified horizon* $\mathbf{K}_m^{\mathrm{cur}}$*, let* $\mathbf{w} \in \mathcal{L}_{\mathrm{ext}}^{\mathrm{term}}$ *be a certified finite extension word, let*

$$\mathsf{Shadow}_m^{\mathrm{QAM}}(\mathbf{w}) \in \{\mathsf{QSH}_m^{\mathrm{hor}}(\mathbf{w}), \mathsf{QSH}_m^{\mathrm{bdry}}(\mathbf{w})\}$$

be the inherited QAM-shadow state from Definition H24.108, and let

$$\mathsf{Route}_m^{\mathrm{upg}}(\mathbf{w}) \in \{\mathcal{U}_m^{\mathrm{hor}}(\mathbf{w}), \mathcal{U}_m^{\mathrm{bdry}}(\mathbf{w})\}$$

be the inherited upgrade route.

A certificate-aware Monte-Carlo / probe family *for* $(m, \mathbf{w})$ *is any downstream probe procedure*

$$\mathsf{MCProbe}_{m,\mathbf{w}}$$

that takes as input only already certified data associated with $\mathbf{w}$ *and outputs a finite diagnostic profile*

$$\mathsf{Prof}_m^{\mathrm{MC}}(\mathbf{w})$$

subject to the following conditions:

(i) ***certificate inheritance:*** *the probe must inherit the existing shadow state and route rather than assign a new one;*

(ii) ***route fidelity:*** *on the horizon route it may test only transport-side stability, continuation-side sensitivity, or compression behavior on the already certified range, while on the boundary route it may test only earliest-boundary localization, stop-sensitive clustering, defect/exposure fingerprints, or boundary-side compression behavior;*

(iii) ***no theorematic promotion:*** *the output profile* $\mathsf{Prof}_m^{\mathrm{MC}}(\mathbf{w})$ *may suggest, rank, cluster, hash, Markov-compress, or Kasiski-scan deviation patterns, but it may not itself certify a later horizon* $\mathbf{K}_{m+1}^{\mathrm{cur}}$ *or replace an inherited boundary witness* $\mathfrak{B}_{m+1}^{\mathrm{cur}}$*;*

(iv) ***stop discipline:*** *if the inherited route is boundary-routed, the probe may sharpen the earliest-boundary picture but may not sample as though theorematic continuation beyond that certified stop had already been proved.*

A probe family that violates one of these conditions is called a certificate-blind Monte-Carlo / probe family.

Proposition H24.113 (Certificate-aware Monte-Carlo probes have diagnostic force but no theorematic authority)**.** *Assume the setting of Definition H24.112. Then:*

(i) every certificate-aware Monte-Carlo / probe family $\mathsf{MCProbe}_{m,\mathbf{w}}$ *is a legitimate downstream consumer of the inherited certificate-aware interface;*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(ii) *the resulting profile* $\mathsf{Prof}_m^{\mathrm{MC}}(\mathbf{w})$ *may refine localization, stability, clustering, or compression questions inside the already certified route, but it does not change the theorematic status of* $\mathbf{w}$*;*

(iii) *any probe family that uses its output to claim a new certified extension, to smooth away an inherited boundary stop, or to inject an uncertified defect/source event is certificate-blind and therefore inadmissible in the present line.*

Proof. Clause (i) follows directly from Definition H24.112: the probe takes only already certified data as input and acts strictly below the inherited route. For (ii), the profile output is diagnostic by construction. It may rank or compress possibilities, but it never changes the inherited field tag, route, or next-work menu, so it cannot alter the theorematic status of $\mathbf{w}$. For (iii), if a probe profile is treated as though it certified a new horizon, then it violates no-theorematic-promotion. If it smooths away a boundary-certified stop, it violates stop discipline and route fidelity. If it injects an uncertified defect/source event on a horizon-routed state, it violates route fidelity on the other side. Hence every such misuse is certificate-blind and inadmissible in the present architecture. □

Corollary H24.114 (The present σ_3 package already admits a first honest Monte-Carlo role)**.** *In the current active line, the first-stage word* $\mathbf{w}_3 \in \{\mathsf{A}, \mathsf{B}\}$ *already admits a first honest Monte-Carlo / probe role below the inherited route:*

(i) *if* $\mathbf{w}_3 = \mathsf{A}$*, Monte-Carlo may act only as a continuation-side stability or compression probe on the currently certified range;*

(ii) *if* $\mathbf{w}_3 = \mathsf{B}$*, Monte-Carlo may act as an earliest-boundary localization, stop-sensitive deviation, or defect/exposure profiling probe, but not as a proof that the corridor has already continued past the certified stop.*

Thus Monte-Carlo is now explicitly fixed as a structural probe layer rather than an alternative theorematic engine.

Proof. Immediate from Corollary H24.110 and Proposition H24.113. □

Remark H24.115 (What this Monte-Carlo / probe discipline now buys)**.** *This is the cleanest place yet to keep the mathematics-first architecture and still use aggressive probe machinery productively. Monte-Carlo may now be used to expose deviation landscapes, boundary clustering, compression-friendly fingerprints, or route-sensitive instability patterns. Those outputs can be helpful for routing attention, designing later diagnostics, or compressing large probe logs into decoder-facing summaries. But the proof burden remains exactly where it belongs: on the certified local extension line. That makes Monte-Carlo valuable without letting it impersonate theorematic closure.*

Reader cue. Read the next block as an inherited freeze audit from an earlier minor public step. Its purpose remains deliberately modest: not to claim a new global theorem, but to record in one place what the release line has stabilized, what must remain local/route-certified in the public wording, and why the present DOI-bearing freeze is honest.

19AZ$^{(6aj)}$. Minor-release freeze audit and public record

After 19AZ$^{(6x)}$–19AZ$^{(6ai)}$, the active line has turned the first post-$\mathbf{K}_2^{\mathrm{cur}}$ extension step into a reusable certificate-aware tool layer. For that inherited minor-release stage, the honest question was therefore no longer whether the new line had mathematical content, but whether its

public wording, routing discipline, archival preservation, and metadata had been synchronized well enough that the public freeze did not misstate what had actually been achieved. The present block records that completed audit explicitly.

Definition H24.116 (Minor-release readiness audit package)**.** *For that inherited public-release line, define the* minor-release readiness audit package

$$\mathsf{Audit}^{\mathrm{minor}} := \big(\mathsf{Math}, \mathsf{Route}, \mathsf{QAM}, \mathsf{MC}, \mathsf{Hist}, \mathsf{Meta}\big)$$

as follows:

(i) Math*: the live OP 1 kernel is discharged locally on the retained no-second-carrier / strict-upper-gap corridor, and the live OP 5 Step (B) kernel is discharged locally on the retained branch-shell / shell-transfer corridor;*

(ii) Route*: the first later leading admissible slot beyond the current certified horizon is normalized by a finite decision datum, an action interface, an admissible terminal word language, field tags, route tags, and a finite next-work menu;*

(iii) QAM*: the new extension layer is attached back to the QAM/decoder side by a finite shadow-state interface and the faithful decoder words* $\{\mathsf{HR}, \mathsf{BR}\}$*;*

(iv) MC*: Monte-Carlo and related probe machinery are explicitly confined to a certificate-aware downstream diagnostic role;*

(v) Hist*: the full historical witness remains preserved in the monolithic source as Part II rather than as a shortened fragment;*

(vi) Meta*: title-line DOI logic, version/chronology wording, and release wording are synchronized so that the concept DOI remains the stable title-page pointer while public version-specific DOI claims appear only in the chronology or explicit release notes.*

Proposition H24.117 (Present release line is mathematically minor-ready and release-packaged)**.** *Assume the audit package from Definition H24.116. Then:*

(i) the mathematics-side burden of that inherited minor release is strong enough for an honest public freeze: the new contribution is a certificate-aware extension/routing/probe layer above the public archival stage infrastructure together with local retained-corridor closure of the live OP 1 and OP 5 Step (B) kernels;

(ii) the public wording of that freeze remains local and route-certified, i.e. it reports retained-corridor kernel closure and the new certificate-aware tool layer, but it does not flatten those results into a new unconditional global closure slogan;

(iii) the release-packaging tasks for that inherited public minor release have been synchronized: version/date metadata, DOI logic, historical preservation, and public wording have been normalized, while ordinary citation/hyperref cleanup can continue in later working states without changing the mathematics.

Proof. Clause (i) is exactly the content accumulated in $19\mathrm{AZ}^{(6x)}$–$19\mathrm{AZ}^{(6ai)}$: finite next-slot decision data, faithful output tokens, action/word routing, certificate-aware matrix/crystal/field readout, admissibility and routing control, finite next-work menus, finite QAM shadows, and an honest downstream Monte-Carlo role. These are not merely editorial restatements but reusable structural tools. The local retained-corridor OP 1 and OP 5 Step (B) closure statements are already recorded in the active main line, so the mathematics side no longer depends

© 2026 Marc Brendecke. Preprint — not peer reviewed.

on introducing yet another theorematic layer before that inherited minor public freeze. Clause (ii) follows from the route-certified architecture itself: every new statement in this block is built from inherited certified corridors, finite decision packaging, and certificate-aware downstream consumers. Flattening that into an unconditional global slogan would destroy the very distinction the new layer was designed to preserve. Clause (iii) follows because what is left to do does not add new mathematical content; it concerns release hygiene, metadata synchronization, and wording discipline. □

Corollary H24.118 (Honest public release message for that inherited minor release)**.** *For that inherited public minor release, the honest release message is the following:*

> *The public archival stage line is carried forward into the public 4.7.0 release by a certificate-aware extension/routing/probe layer. On top of that inherited infrastructure, the live OP 1 kernel and the live OP 5 Step (B) kernel are closed locally on the standing retained corridors. The resulting public gain is therefore not a new global closure slogan, but a route-certified tool layer together with retained-corridor kernel closure, finite decoder/QAM shadows, and a disciplined downstream role for Monte-Carlo and related probes.*

Proof. Immediate from Proposition H24.117. □

Remark H24.119 (Freeze checklist completed for the DOI-bearing release)**.** *In that inherited freeze block, the deliberately boring tasks had been carried out: version/date metadata in the wrapper had been synchronized, chronology wording recorded archival stage as the prior public baseline and the then-current line as the public 4.7.0 release, the historical witness remained preserved in full, and the front-matter wording had been normalized so that local retained-corridor results were not overstated as unconditional global closure. Ordinary citation/hyperref cleanup could still improve future working states, but it no longer blocked the DOI-bearing public freeze.*

Reader cue. Read the next block as the first post-release working bridge that couples the certificate-aware upgrade menus to honest probe consumption on the matrix, crystal, and field-diagnostic side. Its purpose is not to reopen the theorematic routing decision, but to package conservative sharpening together with route-compatible probe use into one downstream working object.

19AZ$^{(6ak)}$. Probe-compatible conservative upgrade packets below the certificate-aware interface

After 19AZ$^{(6ag)}$–19AZ$^{(6ai)}$, the line already knows four things for every certified finite extension word: the inherited horizon/boundary status, the forced route, the finite next-work menu, and the downstream diagnostic role of probe machinery. The next honest post-release step is therefore not a new closure slogan. It is a working packet that couples conservative matrix/crystal/field refinement to admissible probe use without allowing the probe layer to promote, erase, or reroute the certified status.

Definition H24.120 (Probe-compatible conservative upgrade packet)**.** *Fix a certified finite extension word* $\mathbf{w} \in \mathcal{L}_m^{\mathrm{ext}}$*, let*

$$\mathsf{Route}_m^{\mathrm{upg}}(\mathbf{w}) \in \{\mathcal{U}_m^{\mathrm{hor}}(\mathbf{w}), \mathcal{U}_m^{\mathrm{bdry}}(\mathbf{w})\}$$

be its inherited upgrade route from Definition H24.100, let

$$\mathcal{F}_m^{\mathrm{follow}}(\mathbf{w})$$

be its finite route-compatible follow-on menu from Definition H24.104, and let

$$\mathsf{MCProbe}_{m,\mathbf{w}}$$

be any certificate-aware probe family from Definition H24.112.

A probe-compatible conservative upgrade packet *for* $(m, \mathbf{w})$ *is one of the following two objects.*

(i) If $\mathsf{Route}_m^{\mathrm{upg}}(\mathbf{w}) = \mathcal{U}_m^{\mathrm{hor}}(\mathbf{w})$*, define the horizon-side packet*

$$\mathsf{Pack}_m^{\mathrm{hor}}(\mathbf{w}) := \big(\mathsf{HZ}_m(\mathbf{w}), \mathsf{CRH}_m(\mathbf{w}), \mathsf{FEH}_m(\mathbf{w}), \mathcal{F}_m^{\mathrm{follow}}(\mathbf{w}), \mathsf{MCProbe}_{m,\mathbf{w}}\big)$$

under the side condition that every downstream use of $\mathsf{MCProbe}_{m,\mathbf{w}}$ *remains confined to horizon-routed families from*

$$\{\mathsf{MSH}_m(\mathbf{w}), \mathsf{CRS}_m(\mathbf{w}), \mathsf{PDE}_m^{\mathrm{cons}}(\mathbf{w}), \mathsf{DEC}_m^{\mathrm{hor}}(\mathbf{w})\} \cap \mathcal{F}_m^{\mathrm{follow}}(\mathbf{w}).$$

(ii) If $\mathsf{Route}_m^{\mathrm{upg}}(\mathbf{w}) = \mathcal{U}_m^{\mathrm{bdry}}(\mathbf{w})$*, define the boundary-side packet*

$$\mathsf{Pack}_m^{\mathrm{bdry}}(\mathbf{w}) := \big(\mathsf{BD}_m(\mathbf{w}), \mathsf{CRB}_m(\mathbf{w}), \mathsf{FEB}_m(\mathbf{w}), \mathcal{F}_m^{\mathrm{follow}}(\mathbf{w}), \mathsf{MCProbe}_{m,\mathbf{w}}\big)$$

under the side condition that every downstream use of $\mathsf{MCProbe}_{m,\mathbf{w}}$ *remains confined to boundary-routed families from*

$$\{\mathsf{SRC}_m(\mathbf{w}), \mathsf{DFX}_m(\mathbf{w}), \mathsf{MC}_m^{\mathrm{bdry}}(\mathbf{w}), \mathsf{DEC}_m^{\mathrm{bdry}}(\mathbf{w})\} \cap \mathcal{F}_m^{\mathrm{follow}}(\mathbf{w}).$$

Finally define the conservative packet map

$$\mathsf{Pack}_m^{\mathrm{cons}}(\mathbf{w}) := \begin{cases} \mathsf{Pack}_m^{\mathrm{hor}}(\mathbf{w}), & \text{if } \rho_m(\mathbf{w}) = \mathcal{U}^{\mathrm{hor}}, \\ \mathsf{Pack}_m^{\mathrm{bdry}}(\mathbf{w}), & \text{if } \rho_m(\mathbf{w}) = \mathcal{U}^{\mathrm{bdry}}. \end{cases}$$

Proposition H24.121 (Every certified word inherits exactly one probe-compatible conservative upgrade packet)**.** *Assume the setting of Definition H24.120. Then:*

(i) every certified finite extension word $\mathbf{w} \in \mathcal{L}_m^{\mathrm{ext}}$ *inherits exactly one conservative packet* $\mathsf{Pack}_m^{\mathrm{cons}}(\mathbf{w})$*;*

(ii) the packet is faithful to the inherited matrix/crystal/field tags and to the forced route $\rho_m(\mathbf{w})$*;*

(iii) certificate-aware probe output may reprioritize or sharpen work inside *the inherited follow-on menu, but it may not add extra-route tasks, promote theorematic status, or erase a certified boundary stop.*

Proof. By Proposition H24.101, every certified word inherits exactly one route $\rho_m(\mathbf{w})$. By Proposition H24.105, this route comes with exactly one finite compatible follow-on menu. Definition H24.120 therefore assigns exactly one of the two packet forms, proving (i). Statement (ii) is immediate because the packet simply records the inherited matrix/crystal/field tags together with the already forced route and menu. Clause (iii) follows from Proposition H24.113: probe families are downstream consumers only, so they may refine priority inside the inherited menu but they do not change theorematic status, add cross-route tasks, or smooth away a certified terminal boundary event. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Corollary H24.122 (The present σ_3 package already carries a first conservative upgrade packet)**.** *In the current active working line, the first-stage word* $\mathbf{w}_3 \in \{\mathsf{A}, \mathsf{B}\}$ *already induces a first probe-compatible conservative packet*

$$\mathsf{Pack}_2^{\mathrm{cons}}(\mathbf{w}_3),$$

whose type is horizon-side in the transfer-positive case and boundary-side in the earliest-boundary case.

Proof. Immediate from Corollary H24.106, Corollary H24.110, and Proposition H24.121. □

Remark H24.123 (What this conservative packet now buys)**.** *This is the first point at which the post-release line stops treating conservative refinement and probe use as two separate topics. The packet forces them to travel together. Matrix sharpening, crystal stabilization, field-profile refinement, and source/defect localization can now be discussed as probe-compatible working objects, while theorematic certification remains upstream and untouched.*

Reader cue. Read the next block as the finite work-queue normal form induced by a conservative packet. Its purpose is to let probe output guide attention among admissible matrix/crystal/field upgrades without letting that guidance impersonate theorematic certification or invent work outside the inherited route.

19AZ$^{(6al)}$. Probe-ranked conservative work queues for matrix, crystal, and field upgrades

After 19AZ$^{(6ak)}$, every certified word carries one conservative packet. The next honest compression step is to turn the inherited finite follow-on menu into a finite probe-ranked queue. This is exactly the place where Monte-Carlo/probe integration becomes genuinely useful for the next working phase: not by certifying anything new, but by telling the line which admissible conservative upgrade should be looked at first, second, and third.

Definition H24.124 (Probe-ranked conservative work queue)**.** *Fix a conservative packet*

$$\mathsf{Pack}_m^{\mathrm{cons}}(\mathbf{w})$$

from Definition H24.120. A probe-ranked conservative work queue *for* $(m, \mathbf{w})$ *is a finite ordered list*

$$\mathcal{Q}_m^{\mathrm{cons}}(\mathbf{w}) = \big(\mathsf{U}_1 \succcurlyeq \mathsf{U}_2 \succcurlyeq \cdots \succcurlyeq \mathsf{U}_r\big)$$

with the following properties:

(i) every entry U_j *belongs to the inherited follow-on menu* $\mathcal{F}_m^{\mathrm{follow}}(\mathbf{w})$*;*

(ii) the order relation $\succcurlyeq$ *is induced by downstream probe output only as a* priority preorder*, not as new theorematic status;*

(iii) if $\mathsf{Route}_m^{\mathrm{upg}}(\mathbf{w}) = \mathcal{U}_m^{\mathrm{hor}}(\mathbf{w})$*, then every queue entry is horizon-routed and the queue may only prioritize among* $\mathsf{MSHARP}, \mathsf{CSTAB}, \mathsf{FPROF}, \mathsf{DCH}$*;*

(iv) if $\mathsf{Route}_m^{\mathrm{upg}}(\mathbf{w}) = \mathcal{U}_m^{\mathrm{bdry}}(\mathbf{w})$*, then every queue entry is boundary-routed and the queue may only prioritize among* $\mathsf{SLOC}, \mathsf{DFRONT}, \mathsf{MCPROB}, \mathsf{DCB}$*.*

Proposition H24.125 (Every conservative packet induces a finite route-compatible work queue)**.** *Assume the setting of Definition H24.124. Then:*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(i) *every conservative packet* $\mathsf{Pack}_m^{\mathrm{cons}}(\mathbf{w})$ *induces at least one finite probe-ranked conservative work queue* $\mathcal{Q}_m^{\mathrm{cons}}(\mathbf{w})$*;*

(ii) *any two such queues differ only by downstream prioritization inside the same inherited finite route-compatible menu;*

(iii) *in particular, probe ranking may move matrix, crystal, and field-style tasks forward or backward inside the queue, but it may not turn a boundary-routed queue into a continuation queue, nor a continuation queue into a boundary/source queue.*

Proof. Existence in (i) is immediate because the inherited follow-on menu is finite by Proposition H24.105; one may therefore equip it with at least the trivial preorder, or with any finer probe-induced preorder satisfying Definition H24.124. Statement (ii) follows because the queue entries are constrained to lie inside the same inherited route-compatible menu, so downstream variation can only affect the internal ordering. Clause (iii) is exactly the certificate-aware discipline inherited from 19AZ$^{(6af)}$ and 19AZ$^{(6ai)}$: probe ranking is a priority device, not a theorematic rerouting device. □

Corollary H24.126 (The first σ_3 packet already carries a finite conservative work queue)**.** *The present first-stage packet* $\mathsf{Pack}_2^{\mathrm{cons}}(\mathbf{w}_3)$ *already carries a finite route-compatible work queue*

$$\mathcal{Q}_2^{\mathrm{cons}}(\mathbf{w}_3),$$

so the next working phase may be organized as a conservative ranked queue of matrix/crystal/field upgrades together with probe-guided diagnostics before any broader OP-focused reopening is attempted.

Proof. Immediate from Corollary H24.122 and Proposition H24.125. □

Remark H24.127 (What this queue interface now buys)**.** *The next working stage can now be described honestly in one sentence: first sharpen the conservative matrix/crystal/field side and integrate probe diagnostics according to a finite route-compatible queue; only after that use the sharpened readout surface for the next OP-focused push. This is exactly the intended ordering for the post-archival stage line.*

Reader cue. Read the next block as the first concrete working extraction from the probe-ranked conservative queue. Its purpose is not to let probe output overrule the certificate-aware route, but to turn the inherited queue into one honest conservative sharpening/program block that can actually be worked through on the matrix, crystal, and field side before the next OP-focused push.

19AZ$^{(6am)}$. Queue-extracted conservative sharpening/program blocks below the probe-ranked interface

After 19AZ$^{(6al)}$, the active line finally has enough structure to speak about a genuine next working phase rather than only about menus and priorities. The next honest step is to extract, from the finite probe-ranked queue, the actual conservative sharpening/program block that should be worked through first on the inherited route. On the horizon side this means the matrix/crystal/field sharpening strand together with decoder support; on the boundary side it means the earliest-boundary profiling strand together with stop-sensitive probe/decoder support.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition H24.128 (Queue-extracted conservative sharpening/program block)**.** *Fix a probe-ranked conservative work queue*

$$\mathcal{Q}_m^{\mathrm{cons}}(\mathbf{w}) = \big(\mathsf{U}_1 \succcurlyeq \mathsf{U}_2 \succcurlyeq \cdots \succcurlyeq \mathsf{U}_r\big)$$

from Definition H24.124.

(i) If $\mathsf{Route}_m^{\mathrm{upg}}(\mathbf{w}) = \mathcal{U}_m^{\mathrm{hor}}(\mathbf{w})$*, define the* horizon-side conservative sharpening/program block

$$\mathcal{P}_m^{\mathrm{hor}}(\mathbf{w}) := \big(\mathsf{U}_{j_1} \succcurlyeq \cdots \succcurlyeq \mathsf{U}_{j_s}\big)$$

to be the ordered subsequence of $\mathcal{Q}_m^{\mathrm{cons}}(\mathbf{w})$ *consisting exactly of those entries that lie in*

$$\{\mathsf{MSH}_m(\mathbf{w}), \mathsf{CRS}_m(\mathbf{w}), \mathsf{PDE}_m^{\mathrm{cons}}(\mathbf{w}), \mathsf{DEC}_m^{\mathrm{hor}}(\mathbf{w})\}.$$

(ii) If $\mathsf{Route}_m^{\mathrm{upg}}(\mathbf{w}) = \mathcal{U}_m^{\mathrm{bdry}}(\mathbf{w})$*, define the* boundary-side conservative profiling/program block

$$\mathcal{P}_m^{\mathrm{bdry}}(\mathbf{w}) := \big(\mathsf{U}_{k_1} \succcurlyeq \cdots \succcurlyeq \mathsf{U}_{k_t}\big)$$

to be the ordered subsequence of $\mathcal{Q}_m^{\mathrm{cons}}(\mathbf{w})$ *consisting exactly of those entries that lie in*

$$\{\mathsf{SRC}_m(\mathbf{w}), \mathsf{DFX}_m(\mathbf{w}), \mathsf{MC}_m^{\mathrm{bdry}}(\mathbf{w}), \mathsf{DEC}_m^{\mathrm{bdry}}(\mathbf{w})\}.$$

Finally define the queue-extracted conservative program block

$$\mathcal{P}_m^{\mathrm{cons}}(\mathbf{w}) := \begin{cases} \mathcal{P}_m^{\mathrm{hor}}(\mathbf{w}), & \text{if } \rho_m(\mathbf{w}) = \mathcal{U}^{\mathrm{hor}}, \\ \mathcal{P}_m^{\mathrm{bdry}}(\mathbf{w}), & \text{if } \rho_m(\mathbf{w}) = \mathcal{U}^{\mathrm{bdry}}. \end{cases}$$

Proposition H24.129 (Every probe-ranked conservative queue induces one route-faithful working program block)**.** *Assume the setting of Definition H24.128. Then:*

(i) every probe-ranked conservative queue $\mathcal{Q}_m^{\mathrm{cons}}(\mathbf{w})$ *induces exactly one queue-extracted conservative program block* $\mathcal{P}_m^{\mathrm{cons}}(\mathbf{w})$*;*

(ii) if $\mathsf{Route}_m^{\mathrm{upg}}(\mathbf{w}) = \mathcal{U}_m^{\mathrm{hor}}(\mathbf{w})$*, then* $\mathcal{P}_m^{\mathrm{cons}}(\mathbf{w})$ *is a conservative horizon-side sharpening/program block built only from matrix sharpening, crystal stabilization, field-profile refinement, and decoder compression on the already certified range;*

(iii) if $\mathsf{Route}_m^{\mathrm{upg}}(\mathbf{w}) = \mathcal{U}_m^{\mathrm{bdry}}(\mathbf{w})$*, then* $\mathcal{P}_m^{\mathrm{cons}}(\mathbf{w})$ *is a conservative earliest-boundary profiling/program block built only from source localization, defect-front profiling, stop-sensitive probe diagnostics, and boundary-side decoder compression;*

(iv) downstream probe output may change the internal order of the block, but it may not insert extra-route tasks, promote theorematic status, or convert a boundary-profile block into a horizon-sharpening block.

Proof. Statement (i) is immediate because each queue already lies entirely inside one inherited route-compatible menu by Proposition H24.125; taking the ordered subsequence of the relevant route-side task family therefore produces exactly one associated program block. Clauses (ii) and (iii) merely unpack which finite task families are admissible on the two certified routes, using Proposition H24.101 together with Proposition H24.105. Clause (iv) is exactly the downstream discipline already fixed in Proposition H24.113 and Proposition H24.125: probe ranking is allowed to prioritize and sharpen, but not to reroute or recertify. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Corollary H24.130 (The present post-archival stage line already carries a first concrete conservative working program)**.** *The first-stage packet* $\mathsf{Pack}_2^{\mathrm{cons}}(\mathbf{w}_3)$ *and its queue* $\mathcal{Q}_2^{\mathrm{cons}}(\mathbf{w}_3)$ *already induce one concrete conservative program block*

$$\mathcal{P}_2^{\mathrm{cons}}(\mathbf{w}_3),$$

so the current working phase may now be read not merely as a list of admissible future tasks but as one finite route-faithful sharpening/profiling program to be worked through before the next OP-focused reopening.

Proof. Immediate from Corollary H24.126 and Proposition H24.129. □

Remark H24.131 (What this first concrete program block now buys)**.** *This is the first place where the post-release continuation genuinely stops looking like architecture alone. The line now has a concrete finite program block on which to spend the next working effort: either sharpen matrix/crystal/field readout conservatively on the certified horizon side, or profile the earliest honest boundary side in a stop-sensitive way, always under the same probe-compatible discipline. That is exactly the intended bridge from the tool-layer minor release to the next conservative upgrade stage.*

Reader cue. Read the next block as the first honest extraction of a *lead task* from the queue-extracted conservative program block. Its purpose is not to guess a theorematic advance, but to say which concrete conservative sharpening/profiling task should be worked *first* once the inherited route has already been fixed.

19AZ$^{(6an)}$. Route-faithful lead-task extraction from the conservative program block

After 19AZ$^{(6am)}$, the active line no longer lacks an admissible program block; what it still lacks is a normalized answer to the narrower practical question: *which item in that program block should be worked first?* The present block fixes that first-step discipline. On the horizon side, the natural lead task is matrix sharpening, because the matrix/Heisenberg–Millennium readout is the earliest common interface that the crystal and conservative field profiles inherit. On the boundary side, the natural lead task is source/exposure localization, because the earliest honest boundary must be anchored before defect-front profiling, stop-sensitive Monte-Carlo diagnostics, or boundary-side decoder compression can be interpreted coherently.

Definition H24.132 (Route-faithful lead task)**.** *Fix a queue-extracted conservative program block*

$$\mathcal{P}_m^{\mathrm{cons}}(\mathbf{w})$$

from Definition H24.128. Define the associated route-faithful lead task

$$\mathsf{Lead}_m^{\mathrm{cons}}(\mathbf{w})$$

by

$$\mathsf{Lead}_m^{\mathrm{cons}}(\mathbf{w}) := \begin{cases} \mathsf{MSH}_m(\mathbf{w}), & \text{if } \mathsf{Route}_m^{\mathrm{upg}}(\mathbf{w}) = \mathcal{U}_m^{\mathrm{hor}}(\mathbf{w}), \\ \mathsf{SRC}_m(\mathbf{w}), & \text{if } \mathsf{Route}_m^{\mathrm{upg}}(\mathbf{w}) = \mathcal{U}_m^{\mathrm{bdry}}(\mathbf{w}). \end{cases}$$

Thus:

(i) on the horizon side, the lead task is $\mathsf{MSH}_m(\mathbf{w})$*, i.e. matrix sharpening on the already certified range;*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(ii) on the boundary side, the lead task is $\mathsf{SRC}_m(\mathbf{w})$*, i.e. source/exposure localization at the earliest honest boundary.*

Proposition H24.133 (Every conservative program block carries exactly one normalized first task)**.** *Assume the setting of Definition H24.132. Then:*

(i) every queue-extracted conservative program block $\mathcal{P}_m^{\mathrm{cons}}(\mathbf{w})$ *carries exactly one route-faithful lead task* $\mathsf{Lead}_m^{\mathrm{cons}}(\mathbf{w})$*;*

(ii) if $\mathsf{Route}_m^{\mathrm{upg}}(\mathbf{w}) = \mathcal{U}_m^{\mathrm{hor}}(\mathbf{w})$*, then the first conservative sharpening step is matrix sharpening rather than crystal stabilization or conservative PDE refinement;*

(iii) if $\mathsf{Route}_m^{\mathrm{upg}}(\mathbf{w}) = \mathcal{U}_m^{\mathrm{bdry}}(\mathbf{w})$*, then the first conservative profiling step is source/exposure localization rather than defect-front profiling or stop-sensitive probe diagnostics.*

Proof. Uniqueness in (i) is immediate from the already unique inherited route and the case distinction in Definition H24.132. For (ii), matrix sharpening is the earliest common sharpening interface on the continuation side: crystal stabilization and conservative PDE refinement are both read *through* the inherited matrix/Heisenberg–Millennium side, so choosing either of them first would presuppose a matrix profile that the current conservative program has not yet sharpened. Thus the route-faithful first task on the horizon side is $\mathsf{MSH}_m(\mathbf{w})$. For (iii), the earliest honest boundary must first be localized before one can coherently speak about its defect-front geometry, stop-sensitive probe profile, or boundary-side decoder compression. Thus the route-faithful first task on the boundary side is $\mathsf{SRC}_m(\mathbf{w})$. □

Corollary H24.134 (The present post-archival stage line already carries a unique first conservative move)**.** *For the first-stage word* $\mathbf{w}_3 \in \{\mathsf{A}, \mathsf{B}\}$*, the current active line already carries a unique normalized first conservative move:*

$$\mathsf{Lead}_2^{\mathrm{cons}}(\mathbf{w}_3) = \begin{cases} \mathsf{MSH}_2(\mathbf{w}_3), & \text{if } \mathbf{w}_3 = \mathsf{A}, \\ \mathsf{SRC}_2(\mathbf{w}_3), & \text{if } \mathbf{w}_3 = \mathsf{B}. \end{cases}$$

In other words, the current post-release continuation does not *begin by choosing freely among matrix, crystal, field, and probe tasks. It begins by sharpening the matrix side if the inherited status is horizon-certified, and by localizing the earliest source/exposure site if the inherited status is boundary-certified.*

Proof. Immediate from Corollary H24.130 and Proposition H24.133. □

Remark H24.135 (What the lead-task extraction now buys)**.** *This is the point where the post-release line stops saying merely* what could be done next. *It now says which conservative move comes first. That sharply reduces drift in the next working phase: on the horizon side the matrix comes before crystal and field refinements; on the boundary side source localization comes before defect-front profiling and stop-sensitive probing. So the next version can begin with one normalized first move rather than another round of menu-building.*

Reader cue. Read the next block as the first actual execution packet attached to the normalized lead task. Its purpose is not to complete matrix sharpening or source localization in one stroke, but to package the exact first conservative move in a route-faithful form that later crystal, field, defect-front, probe, and decoder layers may inherit without reopening the route.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

19AZ$^{(6ao)}$. First-step conservative execution packets below the lead-task interface

After 19AZ$^{(6an)}$, the active line knows which conservative move comes first. The next honest compression step is therefore to turn that lead task into an explicit first-step execution packet. On the horizon side this packet should sharpen the matrix/Heisenberg–Millennium readout just enough to prepare later crystal and conservative field refinement on the already certified range. On the boundary side it should localize the earliest honest source/exposure site just enough to prepare later defect-front profiling, stop-sensitive probe diagnostics, and decoder compression without smoothing away the boundary stop.

Definition H24.136 (First-step conservative execution packet)**.** *Fix a certified finite extension word $\mathbf{w} \in \mathcal{L}_m^{\mathrm{ext}}$, its route-faithful lead task*

$$\mathsf{Lead}_m^{\mathrm{cons}}(\mathbf{w})$$

from Definition H24.132, and a certificate-aware probe family $\mathsf{MCProbe}_{m,\mathbf{w}}$.

(i) If $\mathsf{Lead}_m^{\mathrm{cons}}(\mathbf{w}) = \mathsf{MSH}_m(\mathbf{w})$, define the horizon-side first-step execution packet

$$\mathsf{Exec}_m^{(1),\mathrm{hor}}(\mathbf{w}) := \big(\mathsf{MSH}_m(\mathbf{w}), \mathsf{HZ}_m(\mathbf{w}), \mathsf{CRH}_m(\mathbf{w}), \mathsf{FEH}_m(\mathbf{w}), \mathsf{MCProbe}_{m,\mathbf{w}}\big)$$

under the side condition that every downstream readout remains confined to a conservative sharpening profile on the already certified range and may only prepare later tasks from

$$\{\mathsf{CRS}_m(\mathbf{w}), \mathsf{PDE}_m^{\mathrm{cons}}(\mathbf{w}), \mathsf{DEC}_m^{\mathrm{hor}}(\mathbf{w})\}.$$

(ii) If $\mathsf{Lead}_m^{\mathrm{cons}}(\mathbf{w}) = \mathsf{SRC}_m(\mathbf{w})$, define the boundary-side first-step execution packet

$$\mathsf{Exec}_m^{(1),\mathrm{bdry}}(\mathbf{w}) := \big(\mathsf{SRC}_m(\mathbf{w}), \mathsf{BD}_m(\mathbf{w}), \mathsf{CRB}_m(\mathbf{w}), \mathsf{FEB}_m(\mathbf{w}), \mathsf{MCProbe}_{m,\mathbf{w}}\big)$$

under the side condition that every downstream readout remains confined to an earliest-boundary localization profile and may only prepare later tasks from

$$\{\mathsf{DFX}_m(\mathbf{w}), \mathsf{MC}_m^{\mathrm{bdry}}(\mathbf{w}), \mathsf{DEC}_m^{\mathrm{bdry}}(\mathbf{w})\}.$$

Finally define the first-step conservative execution map

$$\mathsf{Exec}_m^{(1),\mathrm{cons}}(\mathbf{w}) := \begin{cases} \mathsf{Exec}_m^{(1),\mathrm{hor}}(\mathbf{w}), & \text{if } \mathsf{Lead}_m^{\mathrm{cons}}(\mathbf{w}) = \mathsf{MSH}_m(\mathbf{w}), \\ \mathsf{Exec}_m^{(1),\mathrm{bdry}}(\mathbf{w}), & \text{if } \mathsf{Lead}_m^{\mathrm{cons}}(\mathbf{w}) = \mathsf{SRC}_m(\mathbf{w}). \end{cases}$$

Proposition H24.137 (Every normalized lead task induces exactly one first-step conservative execution packet)**.** *Assume the setting of Definition H24.136. Then:*

(i) every normalized lead task $\mathsf{Lead}_m^{\mathrm{cons}}(\mathbf{w})$ induces exactly one first-step execution packet $\mathsf{Exec}_m^{(1),\mathrm{cons}}(\mathbf{w})$;

(ii) if the lead task is matrix sharpening, then the first-step packet prepares later crystal, conservative field, and decoder refinement on the already certified range without extending theorematic authority beyond that range;

(iii) if the lead task is source/exposure localization, then the first-step packet anchors the earliest honest boundary site before any later defect-front profiling, stop-sensitive probe diagnostics, or boundary-side decoder compression is attempted;

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(iv) certificate-aware probe output may profile or reprioritize work inside the packet's inherited route, but it may not convert the packet into the opposite route, promote theorematic status, or smooth away a certified terminal boundary event.

Proof. Clause (i) is immediate from Definition H24.132: every certified word carries exactly one lead task, namely either $\mathsf{MSH}_m(\mathbf{w})$ or $\mathsf{SRC}_m(\mathbf{w})$, so Definition H24.136 assigns exactly one corresponding first-step packet. Clause (ii) follows because the horizon-side packet records the inherited horizon/matrix/crystal/field status and constrains every downstream use to the conservative continuation-side family; this can sharpen the already certified readout surface, but it does not certify new continuation beyond the current range. Clause (iii) follows because the boundary-side packet records the inherited earliest-boundary tags and constrains downstream use to the boundary-side family; hence localization comes before any defect-front, stop-sensitive probe, or decoder readout that presupposes a fixed boundary anchor. Clause (iv) is exactly the probe discipline inherited from $19\mathrm{AZ}^{(6ai)}$, $19\mathrm{AZ}^{(6ak)}$, and $19\mathrm{AZ}^{(6al)}$: probes may sharpen internal priority and profile, but they do not reroute or recertify. □

Corollary H24.138 (The present post-archival stage line already carries a first executable conservative move)**.** *For the first-stage word $\mathbf{w}_3 \in \{\mathsf{A},\mathsf{B}\}$, the current active line already carries one first executable conservative move:*

$$\mathsf{Exec}_2^{(1),\mathrm{cons}}(\mathbf{w}_3) = \begin{cases} \mathsf{Exec}_2^{(1),\mathrm{hor}}(\mathbf{w}_3), & \text{if } \mathbf{w}_3 = \mathsf{A}, \\ \mathsf{Exec}_2^{(1),\mathrm{bdry}}(\mathbf{w}_3), & \text{if } \mathbf{w}_3 = \mathsf{B}. \end{cases}$$

So the post-release continuation now begins with one explicit executable packet: matrix sharpening first in the transfer-positive case, and source/exposure localization first in the earliest-boundary case.

Proof. Immediate from Corollary H24.134 and Proposition H24.137. □

Remark H24.139 (What this first executable packet now buys)**.** *This is the point where the post-release line stops at last saying only which move comes first and starts carrying the move in an executable normal form. The next working phase can therefore begin with one explicit conservative packet rather than a diffuse instruction: sharpen the matrix side just enough to support later crystal/field refinement, or anchor the earliest boundary just enough to support later defect/probe/decoder profiling. That is a genuinely operational starting point for the next version.*

Reader cue. Read the next block as the first normalized output stage extracted from the first-step execution packet. Its purpose is not to claim that matrix sharpening or source anchoring is already complete in every later detail, but to record the first conservative result surface that the current route-faithful packet is allowed to produce before any later crystal, field, defect-front, probe, or decoder refinement is attempted.

$19\mathrm{AZ}^{(6ap)}$. First conservative result forms extracted from the first execution packet

After $19\mathrm{AZ}^{(6ao)}$, the post-release line carries an explicit first-step execution packet. The next honest compression step is therefore to record the first conservative result form that this packet may output. On the horizon side that result should be the first sharpened matrix/Heisenberg–Millennium readout surface still confined to the already certified range. On the boundary side it should be the first anchored source/exposure profile still confined to the earliest honest boundary event and its immediate diagnostic neighborhood.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition H24.140 (First conservative result form). *Fix a certified finite extension word* $\mathbf{w} \in \mathcal{L}_m^{\mathrm{ext}}$, *its first-step conservative execution packet*

$$\mathsf{Exec}_m^{(1),\mathrm{cons}}(\mathbf{w})$$

from Definition H24.136, and the inherited certificate-aware probe family $\mathsf{MCProbe}_{m,\mathbf{w}}$.

(i) If

$$\mathsf{Exec}_m^{(1),\mathrm{cons}}(\mathbf{w}) = \mathsf{Exec}_m^{(1),\mathrm{hor}}(\mathbf{w}),$$

define the first sharpened horizon-side readout

$$\mathsf{Res}_m^{(1),\mathrm{hor}}(\mathbf{w}) := \big(\mathsf{HZ}_m^{\sharp}(\mathbf{w}), \mathsf{CRH}_m(\mathbf{w}), \mathsf{FEH}_m(\mathbf{w}), \mathsf{MCProbe}_{m,\mathbf{w}}\big),$$

where $\mathsf{HZ}_m^{\sharp}(\mathbf{w})$ *denotes the first conservative sharpened matrix/Heisenberg–Millennium readout produced by* $\mathsf{MSH}_m(\mathbf{w})$ *under the side conditions that:*

- *the sharpened readout remains confined to the already certified horizon-side range,*
- *it does not certify any new continuation beyond that range, and*
- *it may only feed later tasks from*

$$\{\mathsf{CRS}_m(\mathbf{w}), \mathsf{PDE}_m^{\mathrm{cons}}(\mathbf{w}), \mathsf{DEC}_m^{\mathrm{hor}}(\mathbf{w})\}.$$

(ii) If

$$\mathsf{Exec}_m^{(1),\mathrm{cons}}(\mathbf{w}) = \mathsf{Exec}_m^{(1),\mathrm{bdry}}(\mathbf{w}),$$

define the first anchored boundary-side source profile

$$\mathsf{Res}_m^{(1),\mathrm{bdry}}(\mathbf{w}) := \big(\mathsf{BD}_m^{\mathrm{anch}}(\mathbf{w}), \mathsf{CRB}_m(\mathbf{w}), \mathsf{FEB}_m(\mathbf{w}), \mathsf{MCProbe}_{m,\mathbf{w}}\big),$$

where $\mathsf{BD}_m^{\mathrm{anch}}(\mathbf{w})$ *denotes the first conservative anchored source/exposure profile produced by* $\mathsf{SRC}_m(\mathbf{w})$ *under the side conditions that:*

- *the source profile remains anchored at the earliest honest boundary event,*
- *it does not smooth away, postpone, or reclassify that boundary event, and*
- *it may only feed later tasks from*

$$\{\mathsf{DFX}_m(\mathbf{w}), \mathsf{MC}_m^{\mathrm{bdry}}(\mathbf{w}), \mathsf{DEC}_m^{\mathrm{bdry}}(\mathbf{w})\}.$$

Finally define the first conservative result map

$$\mathsf{Res}_m^{(1),\mathrm{cons}}(\mathbf{w}) := \begin{cases} \mathsf{Res}_m^{(1),\mathrm{hor}}(\mathbf{w}), & \text{if } \mathsf{Exec}_m^{(1),\mathrm{cons}}(\mathbf{w}) = \mathsf{Exec}_m^{(1),\mathrm{hor}}(\mathbf{w}), \\ \mathsf{Res}_m^{(1),\mathrm{bdry}}(\mathbf{w}), & \text{if } \mathsf{Exec}_m^{(1),\mathrm{cons}}(\mathbf{w}) = \mathsf{Exec}_m^{(1),\mathrm{bdry}}(\mathbf{w}). \end{cases}$$

Proposition H24.141 (Every first-step execution packet yields exactly one first conservative result form). *Assume the setting of Definition H24.140. Then:*

(i) every first-step conservative execution packet $\mathsf{Exec}_m^{(1),\mathrm{cons}}(\mathbf{w})$ *yields exactly one first conservative result form* $\mathsf{Res}_m^{(1),\mathrm{cons}}(\mathbf{w})$*;*

(ii) in the horizon-side case, the resulting sharpened readout $\mathsf{HZ}_m^{\sharp}(\mathbf{w})$ *is certificate-aware and may support later crystal, conservative field, and decoder refinement without enlarging theorematic authority beyond the already certified range;*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(iii) *in the boundary-side case, the resulting anchored profile* $\mathsf{BD}_m^{\mathrm{anch}}(\mathbf{w})$ *is certificate-aware and fixes the earliest honest source/exposure site before any later defect-front, stop-sensitive probe, or boundary-side decoder refinement is attempted;*

(iv) *certificate-aware probe output may profile or reprioritize work inside the inherited result form, but it may not convert a sharpened horizon-side readout into a boundary profile, convert an anchored boundary profile into a continuation certificate, or erase the route-faithful stop condition.*

Proof. Clause (i) is immediate from Definition H24.136: every certified word carries exactly one first-step execution packet, and Definition H24.140 assigns exactly one corresponding result form according to whether that packet is horizon-side or boundary-side. Clause (ii) follows because the horizon-side result explicitly records the sharpened matrix readout together with the inherited crystal and field tags, while preserving the confinement conditions from the execution packet; this sharpens the readout surface, but it does not certify any further extension. Clause (iii) follows because the boundary-side result explicitly records the anchored source profile together with the inherited boundary-side crystal and field tags, while preserving the earliest-boundary confinement conditions; hence localization precedes any later profiling that presupposes a fixed source anchor. Clause (iv) is exactly the inherited probe discipline from $19\mathrm{AZ}^{(6ai)}$, $19\mathrm{AZ}^{(6ak)}$, and $19\mathrm{AZ}^{(6al)}$, now applied at result-form level rather than packet level. □

Corollary H24.142 (The present post-archival stage line already carries a first conservative output surface)**.** *For the first-stage word* $\mathbf{w}_3 \in \{\mathsf{A}, \mathsf{B}\}$*, the current active line already carries one first conservative output surface:*

$$\mathsf{Res}_2^{(1),\mathrm{cons}}(\mathbf{w}_3) = \begin{cases} \mathsf{Res}_2^{(1),\mathrm{hor}}(\mathbf{w}_3), & \text{if } \mathbf{w}_3 = \mathsf{A}, \\ \mathsf{Res}_2^{(1),\mathrm{bdry}}(\mathbf{w}_3), & \text{if } \mathbf{w}_3 = \mathsf{B}. \end{cases}$$

So the post-release continuation now carries not only an executable first move, but also its first normalized conservative output: a sharpened matrix readout in the transfer-positive case, and an anchored source/exposure profile in the earliest-boundary case.

Proof. Immediate from Corollary H24.138 and Proposition H24.141. □

Remark H24.143 (What the first conservative result form now buys)**.** *This is the point where the post-release line stops carrying only a certified task and an executable packet and begins to carry a first actual result surface. That is exactly the level at which later conservative sharpening can become cumulative: crystal refinement, conservative field profiling, decoder compression, defect-front profiling, and stop-sensitive probing can now inherit a normalized sharpened readout or anchored source profile rather than reconstructing the first move from scratch.*

Reader cue. Read the next block as the first honest second-stage continuation extracted from the newly available conservative result surface. The point is not yet to claim a global refinement theorem, but to show that once a sharpened horizon-side readout or an anchored boundary-side source profile exists, the next conservative move is no longer free: on the horizon side it is crystal stabilization from the sharpened readout; on the boundary side it is defect-front profiling from the anchored source profile.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

19AZ$^{(6aq)}$. First second-stage follow-on forms extracted from the first conservative result surface

After 19AZ$^{(6ap)}$, the post-release line carries a first conservative result surface. The next honest compression step is therefore to record the first second-stage follow-on form that this result surface allows. On the horizon side the sharpened matrix/Heisenberg–Millennium readout should feed crystal stabilization before any broader field or decoder refinement is attempted. On the boundary side the anchored source/exposure profile should feed defect-front profiling before any broader stop-sensitive probe or decoder compression is attempted.

Definition H24.144 (First second-stage follow-on form)**.** *Fix a certified finite extension word* $\mathbf{w} \in \mathcal{L}_m^{\mathrm{ext}}$*, its first conservative result form*

$$\mathsf{Res}_m^{(1),\mathrm{cons}}(\mathbf{w})$$

from Definition H24.140, and the inherited certificate-aware probe family $\mathsf{MCProbe}_{m,\mathbf{w}}$*.*

(i) If

$$\mathsf{Res}_m^{(1),\mathrm{cons}}(\mathbf{w}) = \mathsf{Res}_m^{(1),\mathrm{hor}}(\mathbf{w}),$$

define the first horizon-side second-stage follow-on form

$$\mathsf{FolI}_m^{(2),\mathrm{hor}}(\mathbf{w}) := \big(\mathsf{CRS}_m(\mathbf{w}), \mathsf{HZ}_m^{\sharp}(\mathbf{w}), \mathsf{FEH}_m(\mathbf{w}), \mathsf{DEC}_m^{\mathrm{hor}}(\mathbf{w}), \mathsf{MCProbe}_{m,\mathbf{w}}\big),$$

where $\mathsf{CRS}_m(\mathbf{w})$ *denotes the first conservative crystal-stabilization task extracted from the sharpened readout under the side conditions that:*

- *crystal stabilization remains confined to the already certified horizon-side range encoded by* $\mathsf{HZ}_m^{\sharp}(\mathbf{w})$*,*
- *it does not replace, weaken, or bypass the sharpened readout that generated it, and*
- *it may only feed later tasks from*

$$\{\mathsf{PDE}_m^{\mathrm{cons}}(\mathbf{w}), \mathsf{DEC}_m^{\mathrm{hor}}(\mathbf{w})\}.$$

(ii) If

$$\mathsf{Res}_m^{(1),\mathrm{cons}}(\mathbf{w}) = \mathsf{Res}_m^{(1),\mathrm{bdry}}(\mathbf{w}),$$

define the first boundary-side second-stage follow-on form

$$\mathsf{FolI}_m^{(2),\mathrm{bdry}}(\mathbf{w}) := \big(\mathsf{DFX}_m(\mathbf{w}), \mathsf{BD}_m^{\mathrm{anch}}(\mathbf{w}), \mathsf{FEB}_m(\mathbf{w}), \mathsf{DEC}_m^{\mathrm{bdry}}(\mathbf{w}), \mathsf{MCProbe}_{m,\mathbf{w}}\big),$$

where $\mathsf{DFX}_m(\mathbf{w})$ *denotes the first conservative defect-front profiling task extracted from the anchored source profile under the side conditions that:*

- *defect-front profiling remains anchored at the earliest honest boundary event encoded by* $\mathsf{BD}_m^{\mathrm{anch}}(\mathbf{w})$*,*
- *it does not smooth away, postpone, or reinterpret the anchored source/exposure site, and*
- *it may only feed later tasks from*

$$\{\mathsf{MC}_m^{\mathrm{bdry}}(\mathbf{w}), \mathsf{DEC}_m^{\mathrm{bdry}}(\mathbf{w})\}.$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Finally define the first second-stage conservative follow-on map

$$\mathsf{Foll}_m^{(2),\mathrm{cons}}(\mathbf{w}) := \begin{cases} \mathsf{Foll}_m^{(2),\mathrm{hor}}(\mathbf{w}), & \text{if } \mathsf{Res}_m^{(1),\mathrm{cons}}(\mathbf{w}) = \mathsf{Res}_m^{(1),\mathrm{hor}}(\mathbf{w}), \\ \mathsf{Foll}_m^{(2),\mathrm{bdry}}(\mathbf{w}), & \text{if } \mathsf{Res}_m^{(1),\mathrm{cons}}(\mathbf{w}) = \mathsf{Res}_m^{(1),\mathrm{bdry}}(\mathbf{w}). \end{cases}$$

Proposition H24.145 (Every first conservative result surface yields exactly one first second-stage follow-on form)**.** *Assume the setting of Definition H24.144. Then:*

(i) every first conservative result form $\mathsf{Res}_m^{(1),\mathrm{cons}}(\mathbf{w})$ *yields exactly one first second-stage follow-on form* $\mathsf{Foll}_m^{(2),\mathrm{cons}}(\mathbf{w})$*;*

(ii) in the horizon-side case, crystal stabilization is the first honest second-stage follow-on because the sharpened readout $\mathsf{HZ}_m^{\sharp}(\mathbf{w})$ *must be structurally stabilized before any broader conservative field or decoder refinement is allowed to reuse it;*

(iii) in the boundary-side case, defect-front profiling is the first honest second-stage follow-on because the anchored source profile $\mathsf{BD}_m^{\mathrm{anch}}(\mathbf{w})$ *must first fix the earliest honest boundary geometry before any broader stop-sensitive probe or decoder compression is allowed to reuse it; and*

(iv) certificate-aware probe output may profile or reprioritize work inside the inherited follow-on form, but it may not convert crystal stabilization into a source-localization task, convert defect-front profiling into a continuation certificate, or erase the inherited horizon/boundary route.

Proof. Clause (i) is immediate from Definition H24.140: every certified word carries exactly one first conservative result surface, and Definition H24.144 assigns exactly one corresponding second-stage follow-on form according to whether that surface is horizon-side or boundary-side. Clause (ii) follows because the sharpened horizon-side readout already isolates the admissible range, so the first honest reuse of that surface is to stabilize its crystal-side structure before allowing broader conservative field or decoder refinement. Clause (iii) follows because the anchored boundary-side source profile already isolates the earliest honest boundary site, so the first honest reuse of that surface is to profile the associated defect front before allowing broader stop-sensitive probing or boundary-side decoder compression. Clause (iv) is exactly the inherited probe discipline from $19\mathrm{AZ}^{(6ai)}$, $19\mathrm{AZ}^{(6ak)}$, and $19\mathrm{AZ}^{(6al)}$, now applied at second-stage follow-on level. □

Corollary H24.146 (The present post-archival stage line already carries a first second-stage conservative bifurcation)**.** *For the first-stage word* $\mathbf{w}_3 \in \{\mathsf{A}, \mathsf{B}\}$*, the current active line already carries one first second-stage conservative bifurcation:*

$$\mathsf{Foll}_2^{(2),\mathrm{cons}}(\mathbf{w}_3) = \begin{cases} \mathsf{Foll}_2^{(2),\mathrm{hor}}(\mathbf{w}_3), & \text{if } \mathbf{w}_3 = \mathsf{A}, \\ \mathsf{Foll}_2^{(2),\mathrm{bdry}}(\mathbf{w}_3), & \text{if } \mathbf{w}_3 = \mathsf{B}. \end{cases}$$

So the post-release continuation now carries not only a first conservative output surface, but also its first honest second-stage bifurcation: crystal stabilization from the sharpened horizon-side readout, and defect-front profiling from the anchored boundary-side source profile.

Proof. Immediate from Corollary H24.142 and Proposition H24.145. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Remark H24.147 (What the first second-stage follow-on form now buys)**.** *This is the point where the post-release line starts to behave like a genuine conservative upgrade staircase rather than a single executable packet with a single output surface. The first result form now exports an honest next layer: on the horizon side, crystal stabilization can proceed from a sharpened matrix/Heisenberg–Millennium readout; on the boundary side, defect-front profiling can proceed from an anchored source/exposure profile. Later conservative field refinement, stop-sensitive probes, and decoder compression therefore inherit a better-prepared surface instead of reaching directly back to the first move.*

Reader cue. Read the next block as the first third-stage continuation extracted from the conservative upgrade staircase that now exists after $19\mathrm{AZ}^{(6aq)}$. The point is not yet to claim a full operator or field closure, but to show that once crystal stabilization or defect-front profiling has been honestly reached, the next conservative reuse is again no longer free: on the horizon side it is conservative field refinement from the stabilized crystal side, and on the boundary side it is a stop-sensitive probe/decoder surface from the profiled defect front.

$19\mathrm{AZ}^{(6ar)}$. First third-stage continuation forms extracted from the conservative upgrade staircase

After $19\mathrm{AZ}^{(6aq)}$, the post-release line carries a first second-stage conservative bifurcation. The next honest compression step is therefore to record the first third-stage continuation form that this bifurcation allows. On the horizon side, once a stabilized crystal-side structure exists, the first honest continuation is a conservative field/profile refinement still confined to the already certified range. On the boundary side, once the defect front attached to the earliest honest boundary event has been profiled, the first honest continuation is a stop-sensitive probe/decoder surface that may read the boundary geometry more finely without erasing its stop character.

Definition H24.148 (First third-stage continuation form)**.** *Fix a certified finite extension word* $\mathbf{w} \in \mathcal{L}_m^{\mathrm{ext}}$*, its first second-stage conservative follow-on form*

$$\mathsf{Foll}_m^{(2),\mathrm{cons}}(\mathbf{w})$$

from Definition H24.144, and the inherited certificate-aware probe family $\mathsf{MCProbe}_{m,\mathbf{w}}$*.*

(i) If

$$\mathsf{Foll}_m^{(2),\mathrm{cons}}(\mathbf{w}) = \mathsf{Foll}_m^{(2),\mathrm{hor}}(\mathbf{w}),$$

define the first horizon-side third-stage continuation form

$$\mathsf{Foll}_m^{(3),\mathrm{hor}}(\mathbf{w}) := \big(\mathsf{PDE}_m^{\mathrm{cons}}(\mathbf{w}), \mathsf{CRS}_m(\mathbf{w}), \mathsf{FEH}_m(\mathbf{w}), \mathsf{DEC}_m^{\mathrm{hor}}(\mathbf{w}), \mathsf{MCProbe}_{m,\mathbf{w}}\big),$$

where $\mathsf{PDE}_m^{\mathrm{cons}}(\mathbf{w})$ *denotes the first conservative field/profile refinement task extracted from the stabilized crystal-side structure under the side conditions that:*

- *conservative field refinement remains confined to the already certified horizon-side range inherited from* $\mathsf{HZ}_m^{\sharp}(\mathbf{w})$ *and stabilized by* $\mathsf{CRS}_m(\mathbf{w})$*,*
- *it does not reopen, replace, or bypass the sharpened matrix/Heisenberg–Millennium readout or the crystal stabilization that prepared it, and*
- *it may only feed later horizon-side tasks that remain route-faithful to*

$$\{\mathsf{DEC}_m^{\mathrm{hor}}(\mathbf{w}), \mathsf{MCProbe}_{m,\mathbf{w}}\}.$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(ii) If

$$\mathsf{Foll}_m^{(2),\mathrm{cons}}(\mathbf{w}) = \mathsf{Foll}_m^{(2),\mathrm{bdry}}(\mathbf{w}),$$

define the first boundary-side third-stage continuation form

$$\mathsf{Foll}_m^{(3),\mathrm{bdry}}(\mathbf{w}) := \big(\mathsf{StopSurf}_m(\mathbf{w}), \mathsf{DFX}_m(\mathbf{w}), \mathsf{FEB}_m(\mathbf{w}), \mathsf{DEC}_m^{\mathrm{bdry}}(\mathbf{w}), \mathsf{MCProbe}_{m,\mathbf{w}}\big),$$

where $\mathsf{StopSurf}_m(\mathbf{w})$ *denotes the first stop-sensitive probe/decoder surface extracted from the profiled defect front under the side conditions that:*

- *the stop-sensitive surface remains anchored at the earliest honest boundary event encoded by* $\mathsf{BD}_m^{\mathrm{anch}}(\mathbf{w})$ *and profiled by* $\mathsf{DFX}_m(\mathbf{w})$*,*
- *it may cluster, compress, compare, or decoder-read boundary-side probe traces, but it may not smooth away, postpone, or reinterpret the inherited stop event, and*
- *it may only feed later boundary-side tasks that remain route-faithful to*

$$\{\mathsf{DEC}_m^{\mathrm{bdry}}(\mathbf{w}), \mathsf{MCProbe}_{m,\mathbf{w}}\}.$$

Finally define the first third-stage conservative continuation map

$$\mathsf{Foll}_m^{(3),\mathrm{cons}}(\mathbf{w}) := \begin{cases} \mathsf{Foll}_m^{(3),\mathrm{hor}}(\mathbf{w}), & \text{if } \mathsf{Foll}_m^{(2),\mathrm{cons}}(\mathbf{w}) = \mathsf{Foll}_m^{(2),\mathrm{hor}}(\mathbf{w}), \\ \mathsf{Foll}_m^{(3),\mathrm{bdry}}(\mathbf{w}), & \text{if } \mathsf{Foll}_m^{(2),\mathrm{cons}}(\mathbf{w}) = \mathsf{Foll}_m^{(2),\mathrm{bdry}}(\mathbf{w}). \end{cases}$$

Proposition H24.149 (Every first second-stage conservative bifurcation yields exactly one first third-stage continuation form)**.** *Assume the setting of Definition H24.148. Then:*

(i) every first second-stage conservative follow-on form $\mathsf{Foll}_m^{(2),\mathrm{cons}}(\mathbf{w})$ *yields exactly one first third-stage continuation form* $\mathsf{Foll}_m^{(3),\mathrm{cons}}(\mathbf{w})$*;*

(ii) in the horizon-side case, conservative field refinement is the first honest third-stage continuation because the crystal-stabilized structure is the first admissible surface on which a conservative field/profile refinement can be run without reopening the sharpened readout;

(iii) in the boundary-side case, a stop-sensitive probe/decoder surface is the first honest third-stage continuation because the profiled defect front is the first admissible surface on which probe and decoder traces may be refined without erasing the inherited stop event; and

(iv) certificate-aware probe output may reprioritize or compress work inside the inherited third-stage continuation form, but it may not turn conservative field refinement into a continuation certificate, turn a stop-sensitive surface into a horizon extension, or erase the inherited horizon/boundary route.

Proof. Clause (i) is immediate from Definition H24.144: every certified word carries exactly one first second-stage conservative follow-on form, and Definition H24.148 assigns exactly one corresponding third-stage continuation form according to whether that form is horizon-side or boundary-side. Clause (ii) follows because the crystal-stabilized horizon-side structure is the first admissible surface on which conservative field refinement may proceed without reaching back behind the sharpened readout. Clause (iii) follows because the profiled defect front is the first admissible boundary-side surface on which stop-sensitive probe and decoder traces may be refined without smoothing away the earliest honest stop. Clause (iv) is exactly the inherited probe discipline from $19\mathrm{AZ}^{(6ai)}$, $19\mathrm{AZ}^{(6ak)}$, and $19\mathrm{AZ}^{(6al)}$, now applied at third-stage continuation level. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Corollary H24.150 (The present post-archival stage line already carries a first third-stage conservative continuation)**.** *For the first-stage word* $\mathbf{w}_3 \in \{\mathsf{A}, \mathsf{B}\}$*, the current active line already carries one first third-stage conservative continuation:*

$$\mathsf{Foll}_2^{(3),\mathrm{cons}}(\mathbf{w}_3) = \begin{cases} \mathsf{Foll}_2^{(3),\mathrm{hor}}(\mathbf{w}_3), & \text{if } \mathbf{w}_3 = \mathsf{A}, \\ \mathsf{Foll}_2^{(3),\mathrm{bdry}}(\mathbf{w}_3), & \text{if } \mathbf{w}_3 = \mathsf{B}. \end{cases}$$

So the post-release continuation now carries not only a first conservative output surface and its second-stage bifurcation, but also the first honest third-stage continuation: conservative field refinement from the crystal-stabilized horizon-side structure, and a stop-sensitive probe/decoder surface from the profiled boundary-side defect front.

Proof. Immediate from Corollary H24.146 and Proposition H24.149. □

Remark H24.151 (What the first third-stage continuation now buys)**.** *This is the point where the post-release line begins to carry a genuine conservative continuation staircase rather than only a first move, a first result surface, and one second-stage split. On the horizon side the line now reaches a first conservative field/profile refinement that is already prepared by both sharpened readout and crystal stabilization. On the boundary side the line now reaches a first stop-sensitive probe/decoder surface that is already prepared by both anchored source profiling and defect-front analysis. That makes later decoder compression, conservative field sharpening, and boundary-side diagnostic integration inherit a more mature surface instead of touching the raw first bifurcation directly.*

Reader cue. Read the next block as the first compact conservative readout surface extracted from the upgrade staircase that now exists after $19\mathrm{AZ}^{(6ar)}$. The purpose is to stop carrying the horizon-side and boundary-side continuations only as scattered tasks. Once the sharpened readout, crystal stabilization, and conservative field refinement have been honestly reached on the horizon side, and once the anchored source profile, defect-front profiling, and stop-sensitive probe/decoder surface have been honestly reached on the boundary side, the line may compress each side into one compact conservative working surface for downstream use.

$19\mathrm{AZ}^{(6as)}$. First compact conservative readout surfaces extracted from the continuation staircase

After $19\mathrm{AZ}^{(6ar)}$, the post-release line carries a first third-stage conservative continuation on both the horizon side and the boundary side. The next honest compression step is therefore to record the first compact conservative readout surface that each side now supports. On the horizon side this surface should combine the sharpened matrix/Heisenberg–Millennium readout, the stabilized crystal structure, and the conservative field/profile refinement into one downstream-readable object. On the boundary side it should combine the anchored source/exposure profile, the defect-front profile, and the stop-sensitive probe/decoder surface into one downstream-readable object.

Definition H24.152 (First compact conservative readout surface)**.** *Fix a certified finite extension word* $\mathbf{w} \in \mathcal{L}_m^{\mathrm{ext}}$*, its first third-stage conservative continuation form*

$$\mathsf{Foll}_m^{(3),\mathrm{cons}}(\mathbf{w})$$

from Definition H24.148, and the inherited certificate-aware probe family $\mathsf{MCProbe}_{m,\mathbf{w}}$.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(i) If

$$\mathsf{Foll}_m^{(3),\mathrm{cons}}(\mathbf{w}) = \mathsf{Foll}_m^{(3),\mathrm{hor}}(\mathbf{w}),$$

define the first compact horizon-side conservative readout surface

$$\mathsf{Surf}_m^{\mathrm{hor}}(\mathbf{w}) := \big(\mathsf{HZ}_m^{\sharp}(\mathbf{w}), \mathsf{CRS}_m(\mathbf{w}), \mathsf{PDE}_m^{\mathrm{cons}}(\mathbf{w}), \mathsf{DEC}_m^{\mathrm{hor}}(\mathbf{w}), \mathsf{MCProbe}_{m,\mathbf{w}}\big),$$

under the side conditions that:

- *the surface remains confined to the already certified horizon-side range inherited from the sharpened readout and the crystal-stabilized structure,*
- *the conservative field/profile component remains subordinate to that inherited range and may sharpen transport-side organization but may not certify any new continuation beyond it, and*
- *downstream consumers may read, compare, compress, or decoder-organize the surface, but they may not reopen the route decision that produced it.*

(ii) If

$$\mathsf{Foll}_m^{(3),\mathrm{cons}}(\mathbf{w}) = \mathsf{Foll}_m^{(3),\mathrm{bdry}}(\mathbf{w}),$$

define the first compact boundary-side conservative readout surface

$$\mathsf{Surf}_m^{\mathrm{bdry}}(\mathbf{w}) := \big(\mathsf{BD}_m^{\mathrm{anch}}(\mathbf{w}), \mathsf{DFX}_m(\mathbf{w}), \mathsf{StopSurf}_m(\mathbf{w}), \mathsf{DEC}_m^{\mathrm{bdry}}(\mathbf{w}), \mathsf{MCProbe}_{m,\mathbf{w}}\big),$$

under the side conditions that:

- *the surface remains anchored at the earliest honest boundary event inherited from the anchored source/exposure profile and the profiled defect front,*
- *the stop-sensitive probe/decoder component may refine clustering, exposure comparison, and decoder-readable boundary traces, but it may not smooth away, postpone, or reinterpret the inherited stop event, and*
- *downstream consumers may read, compare, compress, or decoder-organize the surface, but they may not reroute it into a horizon-side continuation claim.*

Finally define the first compact conservative readout-surface map

$$\mathsf{Surf}_m^{\mathrm{cons}}(\mathbf{w}) := \begin{cases} \mathsf{Surf}_m^{\mathrm{hor}}(\mathbf{w}), & \text{if } \mathsf{Foll}_m^{(3),\mathrm{cons}}(\mathbf{w}) = \mathsf{Foll}_m^{(3),\mathrm{hor}}(\mathbf{w}), \\ \mathsf{Surf}_m^{\mathrm{bdry}}(\mathbf{w}), & \text{if } \mathsf{Foll}_m^{(3),\mathrm{cons}}(\mathbf{w}) = \mathsf{Foll}_m^{(3),\mathrm{bdry}}(\mathbf{w}). \end{cases}$$

Proposition H24.153 (Every first third-stage conservative continuation yields exactly one compact conservative readout surface)**.** *Assume the setting of Definition H24.152. Then:*

(i) every first third-stage conservative continuation form $\mathsf{Foll}_m^{(3),\mathrm{cons}}(\mathbf{w})$ *yields exactly one compact conservative readout surface* $\mathsf{Surf}_m^{\mathrm{cons}}(\mathbf{w})$*;*

(ii) in the horizon-side case, the compact surface is the first honest downstream object that combines sharpened readout, crystal stabilization, and conservative field refinement without reopening the route that produced them;

(iii) in the boundary-side case, the compact surface is the first honest downstream object that combines anchored source profiling, defect-front profiling, and stop-sensitive probe/decoder structure without erasing the inherited earliest-boundary stop; and

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(iv) probe or decoder output may cluster, compare, or compress information inside the compact surface, but it may not turn that surface into a new theorematic certificate, a route change, or a denial of the inherited horizon/boundary distinction.

Proof. Clause (i) is immediate from Definition H24.148: every certified word carries exactly one first third-stage conservative continuation form, and Definition H24.152 assigns exactly one corresponding compact conservative readout surface according to whether that continuation is horizon-side or boundary-side. Clause (ii) follows because the horizon-side continuation has already prepared the sharpened readout, crystal stabilization, and conservative field/profile refinement on one inherited certified range, so the first honest reuse is to package them into one downstream-readable surface rather than to scatter them again across separate layers. Clause (iii) follows because the boundary-side continuation has already prepared the anchored source profile, the profiled defect front, and the stop-sensitive probe/decoder structure on one inherited earliest-boundary event, so the first honest reuse is to package them into one downstream-readable surface rather than to let later diagnostics improvise their own boundary anchor. Clause (iv) is exactly the inherited certificate-aware probe discipline from $19\mathrm{AZ}^{(6ai)}$, $19\mathrm{AZ}^{(6ak)}$, and $19\mathrm{AZ}^{(6al)}$, now applied at compact-surface level. □

Corollary H24.154 (The present post-archival stage line already carries a first compact conservative readout surface)**.** *For the first-stage word* $\mathbf{w}_3 \in \{\mathsf{A}, \mathsf{B}\}$*, the current active line already carries one compact conservative readout surface:*

$$\mathsf{Surf}_2^{\mathrm{cons}}(\mathbf{w}_3) = \begin{cases} \mathsf{Surf}_2^{\mathrm{hor}}(\mathbf{w}_3), & \text{if } \mathbf{w}_3 = \mathsf{A}, \\ \mathsf{Surf}_2^{\mathrm{bdry}}(\mathbf{w}_3), & \text{if } \mathbf{w}_3 = \mathsf{B}. \end{cases}$$

So the post-release continuation now carries not only a conservative upgrade staircase through first move, first result surface, second-stage bifurcation, and third-stage continuation, but also the first compact conservative working surface on each side: horizon-side matrix/crystal/field organization and boundary-side source/defect/probe/decoder organization.

Proof. Immediate from Corollary H24.150 and Proposition H24.153. □

Remark H24.155 (What the first compact conservative readout surface now buys)**.** *This is the point where the post-release line first begins to look like a usable conservative readout interface rather than only a staircase of preparatory follow-on forms. On the horizon side, matrix sharpening, crystal stabilization, and conservative field refinement are now available as one compact surface that later decoder or operator-side work may read without falling back behind the certified range. On the boundary side, anchored source profiling, defect-front profiling, and stop-sensitive probe/decoder structure are now available as one compact surface that later diagnostic or clustering work may read without erasing the inherited stop event. The next natural move is therefore no longer to invent another preparatory block, but to extract from these compact surfaces the first route-faithful conservative output or comparison layer that later OP-oriented work can consume.*

Reader cue. Read the next block as the first route-faithful conservative comparison/output layer extracted from the compact surfaces of $19\mathrm{AZ}^{(6as)}$. The purpose is to stop treating the horizon-side and boundary-side surfaces only as rich internal packages. Once each side already carries one compact conservative surface, the next honest step is to expose one downstream-readable output/comparison form that later OP-oriented work may consume without reopening the route, smoothing away an inherited stop, or pretending that horizon-side and boundary-side outputs are interchangeable.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

19AZ$^{(6at)}$. First route-faithful conservative comparison/output layer extracted from the compact surfaces

After 19AZ$^{(6as)}$, the post-release line carries a first compact conservative readout surface on each admissible side. The next honest compression step is therefore to expose a route-faithful comparison/output layer that makes each compact surface downstream-readable as one conservative output object. On the horizon side this output should preserve the already certified matrix/crystal/field organization. On the boundary side it should preserve the already anchored source/defect/probe/decoder organization together with the inherited stop sensitivity. The point is not to collapse both sides into one falsely uniform object, but to let later OP-oriented or diagnostic work compare and consume them through one disciplined interface.

Definition H24.156 (First route-faithful conservative output/comparison layer)**.** *Fix a certified finite extension word* $\mathbf{w} \in \mathcal{L}_m^{\mathrm{ext}}$*, its first compact conservative readout surface*

$$\mathsf{Surf}_m^{\mathrm{cons}}(\mathbf{w})$$

from Definition H24.152, and the inherited certificate-aware probe family $\mathsf{MCProbe}_{m,\mathbf{w}}$*.*

(i) If

$$\mathsf{Surf}_m^{\mathrm{cons}}(\mathbf{w}) = \mathsf{Surf}_m^{\mathrm{hor}}(\mathbf{w}),$$

define the first route-faithful horizon-side conservative output/comparison object

$$\mathsf{OutCmp}_m^{\mathrm{hor}}(\mathbf{w}) := \big(\mathsf{HOR}, \mathsf{HZ}_m^{\sharp}(\mathbf{w}), \mathsf{CRS}_m(\mathbf{w}), \mathsf{PDE}_m^{\mathrm{cons}}(\mathbf{w}), \mathsf{DEC}_m^{\mathrm{hor}}(\mathbf{w}), \mathsf{MCProbe}_{m,\mathbf{w}}\big).$$

under the side conditions that:

- *the tag* HOR *records that the output remains confined to the already certified horizon-side range,*
- *downstream comparison or output work may compare this object with other horizon-side conservative outputs, may compress it, and may use it as an input surface for later OP-oriented tasks, but it may not reinterpret it as a boundary-side stop object, and*
- *no component of the output may certify a continuation beyond the range already inherited by* $\mathsf{Surf}_m^{\mathrm{hor}}(\mathbf{w})$*.*

(ii) If

$$\mathsf{Surf}_m^{\mathrm{cons}}(\mathbf{w}) = \mathsf{Surf}_m^{\mathrm{bdry}}(\mathbf{w}),$$

define the first route-faithful boundary-side conservative output/comparison object

$$\mathsf{OutCmp}_m^{\mathrm{bdry}}(\mathbf{w}) := \big(\mathsf{BDRY}, \mathsf{BD}_m^{\mathrm{anch}}(\mathbf{w}), \mathsf{DFX}_m(\mathbf{w}), \mathsf{StopSurf}_m(\mathbf{w}), \mathsf{DEC}_m^{\mathrm{bdry}}(\mathbf{w}), \mathsf{MCProbe}_{m,\mathbf{w}}\big).$$

under the side conditions that:

- *the tag* BDRY *records that the output remains anchored at the earliest honest boundary event and retains inherited stop sensitivity,*
- *downstream comparison or output work may compare this object with other boundary-side conservative outputs, may compress it, and may use it as an input surface for later OP-oriented tasks, but it may not smooth away the inherited stop event or relabel the object as horizon-side continuation, and*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

- *no component of the output may deny or postpone the earliest honest boundary anchor already inherited by* $\mathsf{Surf}_m^{\mathrm{bdry}}(\mathbf{w})$.

Finally define the first route-faithful conservative output/comparison map

$$\mathsf{OutCmp}_m^{\mathrm{cons}}(\mathbf{w}) := \begin{cases} \mathsf{OutCmp}_m^{\mathrm{hor}}(\mathbf{w}), & \text{if } \mathsf{Surf}_m^{\mathrm{cons}}(\mathbf{w}) = \mathsf{Surf}_m^{\mathrm{hor}}(\mathbf{w}), \\ \mathsf{OutCmp}_m^{\mathrm{bdry}}(\mathbf{w}), & \text{if } \mathsf{Surf}_m^{\mathrm{cons}}(\mathbf{w}) = \mathsf{Surf}_m^{\mathrm{bdry}}(\mathbf{w}). \end{cases}$$

Proposition H24.157 (Every first compact conservative surface yields exactly one route–faithful conservative output/comparison object)**.** *Assume the setting of Definition H24.156. Then:*

(i) every first compact conservative surface $\mathsf{Surf}_m^{\mathrm{cons}}(\mathbf{w})$ *yields exactly one route-faithful conservative output/comparison object* $\mathsf{OutCmp}_m^{\mathrm{cons}}(\mathbf{w})$*;*

(ii) in the horizon-side case, the object is the first honest downstream-readable output that packages the matrix/crystal/field surface together with an explicit horizon-side tag;

(iii) in the boundary-side case, the object is the first honest downstream-readable output that packages the source/defect/probe/decoder surface together with an explicit boundary-side stop-preserving tag; and

(iv) later OP-oriented, diagnostic, clustering, or decoder-side work may consume these objects as conservative outputs, but it may not use them to erase the inherited horizon/boundary distinction or to manufacture a fresh theorematic certificate.

Proof. Clause (i) is immediate from Definition H24.156: every compact conservative surface is either horizon-side or boundary-side, and the definition assigns exactly one corresponding route-faithful output/comparison object with the matching side tag. Clause (ii) follows because the horizon-side compact surface already carries sharpened readout, crystal stabilization, conservative field refinement, and decoder/probe organization on one inherited certified range, so the first honest downstream output is to expose that package with an explicit horizon-side tag rather than to reopen the route or invent a boundary anchor. Clause (iii) follows because the boundary-side compact surface already carries anchored source profiling, defect-front profiling, and stop-sensitive probe/decoder structure on one inherited earliest-boundary event, so the first honest downstream output is to expose that package with an explicit boundary-side tag rather than to smooth away the stop or relabel the object as horizon-side continuation. Clause (iv) is exactly the inherited certificate-aware discipline from $19\mathrm{AZ}^{(6ae)}$, $19\mathrm{AZ}^{(6af)}$, $19\mathrm{AZ}^{(6ai)}$, and $19\mathrm{AZ}^{(6as)}$, now applied at the first output/comparison level. □

Corollary H24.158 (The present post-archival stage line already carries a first route-faithful conservative output/comparison layer)**.** *For the first-stage word* $\mathbf{w}_3 \in \{\mathsf{A}, \mathsf{B}\}$*, the current active line already carries one route-faithful conservative output/comparison object:*

$$\mathsf{OutCmp}_2^{\mathrm{cons}}(\mathbf{w}_3) = \begin{cases} \mathsf{OutCmp}_2^{\mathrm{hor}}(\mathbf{w}_3), & \text{if } \mathbf{w}_3 = \mathsf{A}, \\ \mathsf{OutCmp}_2^{\mathrm{bdry}}(\mathbf{w}_3), & \text{if } \mathbf{w}_3 = \mathsf{B}. \end{cases}$$

So the post-release continuation now carries not only a conservative upgrade staircase and a first compact conservative surface on each side, but also the first route-faithful conservative output/comparison layer that later OP-oriented work may consume directly.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. Immediate from Corollary H24.154 and Proposition H24.157. □

Remark H24.159 (What the first route-faithful conservative output/comparison layer now buys)**.** *This is the point where the post-release line first becomes directly consumable as a disciplined conservative output interface. The horizon-side object now packages matrix/crystal/field organization together with an explicit horizon-side tag, while the boundary-side object packages source/defect/probe/decoder organization together with an explicit stop-preserving boundary-side tag. Later OP-oriented work can therefore start from a route-faithful output object rather than from scattered follow-on packets or even from the raw compact surface itself. The next natural move is to use these output/comparison objects to define the first conservative OP-facing intake layer without broadening the theorematic claims.*

Reader cue. Read the next block as the first conservative OP-facing intake layer extracted from the route-faithful output/comparison objects of $19\mathrm{AZ}^{(6at)}$. Its purpose is to let the next OP-oriented working stage consume the new conservative output surfaces without reopening the route decision, erasing the boundary stop, or inflating the theorematic range. The block does not yet solve any open problem; it defines the disciplined intake interface through which later OP-side work must begin.

$19\mathrm{AZ}^{(6au)}$. First conservative OP-facing intake layer extracted from the route-faithful output/comparison objects

After $19\mathrm{AZ}^{(6at)}$, the post-release line carries a first route-faithful conservative output/comparison object on each admissible side. The next honest step is therefore to define the first OP-facing intake layer that later OP-oriented work may consume. On the horizon side this intake should inherit the already certified matrix/crystal/field package together with its explicit horizon tag. On the boundary side it should inherit the anchored source/defect/probe/decoder package together with its explicit stop-preserving boundary tag. The point is not to broaden the theorematic status, but to make later OP-oriented work start from a disciplined conservative intake surface rather than from raw local branch data.

Definition H24.160 (First conservative OP-facing intake layer)**.** *Fix a certified finite extension word $\mathbf{w} \in \mathcal{L}_m^{\mathrm{ext}}$, its first route-faithful conservative output/comparison object*

$$\mathsf{OutCmp}_m^{\mathrm{cons}}(\mathbf{w})$$

from Definition H24.156, and the inherited certificate-aware probe family $\mathsf{MCProbe}_{m,\mathbf{w}}$.

(i) If

$$\mathsf{OutCmp}_m^{\mathrm{cons}}(\mathbf{w}) = \mathsf{OutCmp}_m^{\mathrm{hor}}(\mathbf{w}),$$

define the first horizon-side conservative OP-facing intake object

$$\mathsf{OPIn}_m^{\mathrm{hor}}(\mathbf{w}) := \big(\mathsf{HOR}, \mathsf{HZ}_m^{\sharp}(\mathbf{w}), \mathsf{CRS}_m(\mathbf{w}), \mathsf{PDE}_m^{\mathrm{cons}}(\mathbf{w}), \mathsf{DEC}_m^{\mathrm{hor}}(\mathbf{w}), \mathsf{MCProbe}_{m,\mathbf{w}}\big),$$

under the side conditions that:

- *later OP-oriented work may use this intake object only as a conservative horizon-side input surface and may not reinterpret it as evidence for a boundary-side stop event,*
- *no OP-facing consumer may certify any continuation beyond the horizon-side range already inherited by $\mathsf{OutCmp}_m^{\mathrm{hor}}(\mathbf{w})$, and*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

- *any later transport, matrix, crystal, or field-equation sharpening that starts from this object remains subordinate to the inherited horizon tag* HOR.

(ii) If

$$\mathsf{OutCmp}_m^{\mathrm{cons}}(\mathbf{w}) = \mathsf{OutCmp}_m^{\mathrm{bdry}}(\mathbf{w}),$$

define the first boundary-side conservative OP-facing intake object

$$\mathsf{OPIn}_m^{\mathrm{bdry}}(\mathbf{w}) := \big(\mathsf{BDRY}, \mathsf{BD}_m^{\mathrm{anch}}(\mathbf{w}), \mathsf{DFX}_m(\mathbf{w}), \mathsf{StopSurf}_m(\mathbf{w}), \mathsf{DEC}_m^{\mathrm{bdry}}(\mathbf{w}), \mathsf{MCProbe}_{m,\mathbf{w}}\big).$$

under the side conditions that:

- *later OP-oriented work may use this intake object only as a conservative boundary-side input surface and may not relabel it as a horizon-continuation object,*
- *no OP-facing consumer may smooth away, postpone, or reinterpret the inherited earliest honest boundary stop carried by* $\mathsf{OutCmp}_m^{\mathrm{bdry}}(\mathbf{w})$*, and*
- *any later source, defect, probe, decoder, or field-side comparison that starts from this object remains subordinate to the inherited stop-preserving boundary tag* BDRY.

Finally define the first conservative OP-facing intake map

$$\mathsf{OPIn}_m^{\mathrm{cons}}(\mathbf{w}) := \begin{cases} \mathsf{OPIn}_m^{\mathrm{hor}}(\mathbf{w}), & \text{if } \mathsf{OutCmp}_m^{\mathrm{cons}}(\mathbf{w}) = \mathsf{OutCmp}_m^{\mathrm{hor}}(\mathbf{w}), \\ \mathsf{OPIn}_m^{\mathrm{bdry}}(\mathbf{w}), & \text{if } \mathsf{OutCmp}_m^{\mathrm{cons}}(\mathbf{w}) = \mathsf{OutCmp}_m^{\mathrm{bdry}}(\mathbf{w}). \end{cases}$$

Proposition H24.161 (Every route-faithful conservative output/comparison object yields exactly one conservative OP-facing intake object)**.** *Assume the setting of Definition H24.160. Then:*

(i) every route-faithful conservative output/comparison object $\mathsf{OutCmp}_m^{\mathrm{cons}}(\mathbf{w})$ *yields exactly one conservative OP-facing intake object* $\mathsf{OPIn}_m^{\mathrm{cons}}(\mathbf{w})$*;*

(ii) in the horizon-side case, the intake object is the first honest OP-facing surface that packages the matrix/crystal/field organization together with an explicit horizon-side tag;

(iii) in the boundary-side case, the intake object is the first honest OP-facing surface that packages the source/defect/probe/decoder organization together with an explicit stop-preserving boundary-side tag; and

(iv) later OP-oriented work may consume these intake objects as conservative inputs, but it may not use them to erase the inherited horizon/boundary distinction, to deny the inherited earliest-boundary stop, or to manufacture a fresh theorematic certificate.

Proof. Clause (i) is immediate from Definition H24.160: every route-faithful conservative output/comparison object is either horizon-side or boundary-side, and the definition assigns exactly one corresponding conservative OP-facing intake object with the matching side tag. Clause (ii) follows because the horizon-side output/comparison object already packages sharpened readout, crystal stabilization, conservative field refinement, and decoder/probe organization on one inherited certified range, so the first honest OP-facing use is to intake that package with its explicit horizon-side tag rather than to reopen the route or invent a boundary anchor. Clause (iii) follows because the boundary-side output/comparison object already packages anchored source profiling, defect-front profiling, and stop-sensitive probe/decoder

© 2026 Marc Brendecke. Preprint — not peer reviewed.

structure on one inherited earliest-boundary event, so the first honest OP-facing use is to intake that package with its explicit stop-preserving boundary tag rather than to smooth away the stop or relabel the object as horizon-side continuation. Clause (iv) is exactly the inherited certificate-aware discipline from 19AZ$^{(6ae)}$, 19AZ$^{(6af)}$, 19AZ$^{(6ai)}$, and 19AZ$^{(6at)}$, now applied at the first OP-facing intake level. □

Corollary H24.162 (The present post-archival stage line already carries a first conservative OP-facing intake layer)**.** *For the first-stage word* $\mathbf{w}_3 \in \{\mathsf{A}, \mathsf{B}\}$*, the current active line already carries one conservative OP-facing intake object:*

$$\mathsf{OPIn}_2^{\mathrm{cons}}(\mathbf{w}_3) = \begin{cases} \mathsf{OPIn}_2^{\mathrm{hor}}(\mathbf{w}_3), & \text{if } \mathbf{w}_3 = \mathsf{A}, \\ \mathsf{OPIn}_2^{\mathrm{bdry}}(\mathbf{w}_3), & \text{if } \mathbf{w}_3 = \mathsf{B}. \end{cases}$$

So the post-release continuation now carries not only a route-faithful conservative output/comparison layer, but also the first conservative OP-facing intake layer that the next OP-oriented working stage may consume directly.

Proof. Immediate from Corollary H24.158 and Proposition H24.161. □

Remark H24.163 (What the first conservative OP-facing intake layer now buys)**.** *This is the point where the post-release line first becomes ready for the next OP-oriented working stage without pretending that any open problem has already been solved. The horizon-side intake object exposes a disciplined matrix/crystal/field input surface under an explicit horizon tag, while the boundary-side intake object exposes a disciplined source/defect/probe/decoder input surface under an explicit stop-preserving boundary tag. Later OP-facing work can therefore begin from a conservative intake object that is already route-faithful, comparison-ready, and certificate-aware, instead of improvising from scattered packets or reopening the extension decision itself.*

Reader cue. Read the next block as the first conservative OP-facing work-request contract extracted from the intake layer of 19AZ$^{(6au)}$. Its purpose is to specify which kinds of OP-oriented demands may be posed from the new intake objects without reopening the route decision, erasing the inherited boundary stop, or inflating the theorematic range. The block does not yet answer any open problem; it fixes the finite admissible request menu through which the next OP-oriented stage must begin.

19AZ$^{(6av)}$. First conservative OP-facing work-request contract extracted from the intake layer

After 19AZ$^{(6au)}$, the post-release line carries a first conservative OP-facing intake object on each admissible side. The next honest step is therefore to define which OP-oriented requests may be asked from those intake objects without violating the inherited certificate-aware discipline. On the horizon side the admissible requests should remain subordinate to the inherited matrix/crystal/field organization and explicit horizon tag. On the boundary side the admissible requests should remain subordinate to the inherited anchored source/defect/probe/decoder organization and explicit stop-preserving boundary tag. The point is not to authorize arbitrary OP-side speculation, but to specify the first finite request contract under which later OP-facing work may proceed.

Definition H24.164 (First conservative OP-facing work-request contract)**.** *Fix a certified finite extension word* $\mathbf{w} \in \mathcal{L}_m^{\mathrm{ext}}$ *and its first conservative OP-facing intake object*

$$\mathsf{OPIn}_m^{\mathrm{cons}}(\mathbf{w})$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

from Definition H24.160.

(i) If

$$\mathsf{OPIn}_m^{\mathrm{cons}}(\mathbf{w}) = \mathsf{OPIn}_m^{\mathrm{hor}}(\mathbf{w}),$$

define the first horizon-side conservative OP-facing request menu

$$\mathcal{Q}_m^{\mathrm{hor}}(\mathbf{w}) := \{\mathsf{QMSH}_m(\mathbf{w}), \mathsf{QCRS}_m(\mathbf{w}), \mathsf{QPDE}_m(\mathbf{w}), \mathsf{QDEC}_m^{\mathrm{hor}}(\mathbf{w})\},$$

where the four request forms ask respectively for:

- *conservative matrix/MM-HC sharpening comparison relative to* $\mathsf{HZ}_m^{\sharp}(\mathbf{w})$,
- *conservative crystal-stabilization comparison relative to* $\mathsf{CRS}_m(\mathbf{w})$,
- *conservative field/profile comparison relative to* $\mathsf{PDE}_m^{\mathrm{cons}}(\mathbf{w})$, *and*
- *horizon-side decoder/probe compression or comparison relative to* $\mathsf{DEC}_m^{\mathrm{hor}}(\mathbf{w})$ *and* $\mathsf{MCProbe}_{m,\mathbf{w}}$.

These requests are admissible only under the side conditions that:

- *no request may ask for a fresh theorematic certificate beyond the horizon-side range already inherited by* $\mathsf{OPIn}_m^{\mathrm{hor}}(\mathbf{w})$,
- *no request may relabel the horizon-side intake object as evidence for a boundary-side stop event, and*
- *every admissible request remains subordinate to the inherited horizon tag* HOR.

(ii) If

$$\mathsf{OPIn}_m^{\mathrm{cons}}(\mathbf{w}) = \mathsf{OPIn}_m^{\mathrm{bdry}}(\mathbf{w}),$$

define the first boundary-side conservative OP-facing request menu

$$\mathcal{Q}_m^{\mathrm{bdry}}(\mathbf{w}) := \{\mathsf{QSRC}_m(\mathbf{w}), \mathsf{QDFX}_m(\mathbf{w}), \mathsf{QSTOP}_m(\mathbf{w}), \mathsf{QDEC}_m^{\mathrm{bdry}}(\mathbf{w})\},$$

where the four request forms ask respectively for:

- *anchored source/exposure comparison relative to* $\mathsf{BD}_m^{\mathrm{anch}}(\mathbf{w})$,
- *defect-front comparison relative to* $\mathsf{DFX}_m(\mathbf{w})$,
- *stop-sensitive probe-surface comparison relative to* $\mathsf{StopSurf}_m(\mathbf{w})$, *and*
- *boundary-side decoder/probe compression or comparison relative to* $\mathsf{DEC}_m^{\mathrm{bdry}}(\mathbf{w})$ *and* $\mathsf{MCProbe}_{m,\mathbf{w}}$.

These requests are admissible only under the side conditions that:

- *no request may smooth away, postpone, or reinterpret the inherited earliest honest boundary stop carried by* $\mathsf{OPIn}_m^{\mathrm{bdry}}(\mathbf{w})$,
- *no request may relabel the boundary-side intake object as a horizon-continuation object, and*
- *every admissible request remains subordinate to the inherited stop-preserving boundary tag* BDRY.

Finally define the first conservative OP-facing request contract map

$$\mathsf{Q}_m^{\mathrm{cons}}(\mathbf{w}) := \begin{cases} \mathcal{Q}_m^{\mathrm{hor}}(\mathbf{w}), & \text{if } \mathsf{OPIn}_m^{\mathrm{cons}}(\mathbf{w}) = \mathsf{OPIn}_m^{\mathrm{hor}}(\mathbf{w}), \\ \mathcal{Q}_m^{\mathrm{bdry}}(\mathbf{w}), & \text{if } \mathsf{OPIn}_m^{\mathrm{cons}}(\mathbf{w}) = \mathsf{OPIn}_m^{\mathrm{bdry}}(\mathbf{w}). \end{cases}$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proposition H24.165 (Every conservative OP-facing intake object yields exactly one finite admissible work-request contract)**.** *Assume the setting of Definition H24.164. Then:*

(i) every conservative OP-facing intake object $\mathsf{OPIn}_m^{\mathrm{cons}}(\mathbf{w})$ *yields exactly one finite admissible request contract* $\mathsf{Q}_m^{\mathrm{cons}}(\mathbf{w})$*;*

(ii) in the horizon-side case, the contract authorizes only matrix/crystal/field/decoder-probe requests subordinate to the inherited horizon tag;

(iii) in the boundary-side case, the contract authorizes only source/defect/stop-surface/decoder-probe requests subordinate to the inherited stop-preserving boundary tag; and

(iv) no admissible request in either contract may manufacture a fresh theorematic certificate, erase the inherited horizon/boundary distinction, or deny the inherited earliest-boundary stop.

Proof. Clause (i) is immediate from Definition H24.164: every conservative OP-facing intake object is either horizon-side or boundary-side, and the definition assigns exactly one corresponding finite request menu. Clause (ii) follows because the horizon-side intake object already packages the sharpened readout, crystal stabilization, conservative field refinement, and decoder/probe organization under one explicit horizon tag, so the only honest OP-facing requests are requests that compare, refine, or compress that inherited organization without leaving the inherited range. Clause (iii) follows because the boundary-side intake object already packages the anchored source profile, defect-front profile, stop-sensitive probe surface, and decoder/probe organization under one explicit stop-preserving boundary tag, so the only honest OP-facing requests are requests that compare, refine, or compress that inherited organization without smoothing away the stop or relabeling the route. Clause (iv) is exactly the inherited certificate-aware discipline from $19\mathrm{AZ}^{(6ae)}$, $19\mathrm{AZ}^{(6af)}$, $19\mathrm{AZ}^{(6ai)}$, $19\mathrm{AZ}^{(6at)}$, and $19\mathrm{AZ}^{(6au)}$, now expressed as the first finite OP-facing request contract. □

Corollary H24.166 (The present post-archival stage line already carries a first finite OP–facing request contract)**.** *For the first-stage word* $\mathbf{w}_3 \in \{\mathsf{A}, \mathsf{B}\}$*, the current active line already carries one finite conservative OP-facing request contract:*

$$\mathsf{Q}_2^{\mathrm{cons}}(\mathbf{w}_3) = \begin{cases} \mathcal{Q}_2^{\mathrm{hor}}(\mathbf{w}_3), & \text{if } \mathbf{w}_3 = \mathsf{A}, \\ \mathcal{Q}_2^{\mathrm{bdry}}(\mathbf{w}_3), & \text{if } \mathbf{w}_3 = \mathsf{B}. \end{cases}$$

So the post-release continuation now carries not only a conservative OP-facing intake layer, but also the first finite contract specifying which OP-oriented demands may honestly be posed from that intake surface.

Proof. Immediate from Corollary H24.162 and Proposition H24.165. □

Remark H24.167 (What the first conservative OP-facing work-request contract now buys)**.** *This is the point where the post-release line first becomes able to formulate disciplined OP-oriented questions without pretending that it already owns their answers. The horizon-side intake object now supports a finite request menu for matrix/crystal/field/decoder-probe work under an explicit horizon tag, while the boundary-side intake object supports a finite request menu for source/defect/stop-surface/decoder-probe work under an explicit stop-preserving boundary tag. Later OP-facing work can therefore begin from a conservative intake contract instead of improvising its own admissible question set or silently broadening the inherited theorematic range.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Reader cue. Read the next block as the first conservative OP-facing response/fulfillment layer extracted from the request contract of $19AZ^{(6av)}$. Its purpose is to specify how an admissible OP-facing request may be answered conservatively without manufacturing a fresh theorematic certificate, erasing the inherited route, or denying the earliest honest boundary stop. The block does not yet close any open problem; it fixes the first disciplined fulfillment format through which later OP-oriented work may respond to admissible requests.

$19AZ^{(6aw)}$. First conservative OP-facing response/fulfillment layer extracted from the request contract

After $19AZ^{(6av)}$, the post-release line carries a first finite admissible request contract on each intake side. The next honest step is therefore to define how an admissible OP-facing request may be fulfilled while remaining subordinate to the inherited certificate-aware route. On the horizon side a conservative response must stay within the inherited matrix/crystal/field/decoder organization under the explicit horizon tag. On the boundary side a conservative response must stay within the inherited source/defect/stop-surface/decoder organization under the explicit stop-preserving boundary tag. The point is not to authorize arbitrary new claims, but to define the first finite route-faithful response layer through which admissible OP-facing work may answer the requests that the contract actually allows.

Definition H24.168 (First conservative OP-facing response/fulfillment layer)**.** *Fix a certified finite extension word* $\mathbf{w} \in \mathcal{L}_m^{\mathrm{ext}}$*, its first conservative OP-facing request contract*

$$\mathsf{Q}_m^{\mathrm{cons}}(\mathbf{w})$$

from Definition H24.164, and an admissible request

$$q \in \mathsf{Q}_m^{\mathrm{cons}}(\mathbf{w}).$$

Define the first conservative OP-facing fulfillment object

$$\mathsf{Ful}_m^{\mathrm{cons}}(\mathbf{w}; q)$$

by the following cases.

(i) If

$$\mathsf{Q}_m^{\mathrm{cons}}(\mathbf{w}) = \mathcal{Q}_m^{\mathrm{hor}}(\mathbf{w}),$$

define the first horizon-side conservative fulfillment family

$$\mathcal{F}_m^{\mathrm{hor}}(\mathbf{w}) := \{\mathsf{FMSH}_m(\mathbf{w}), \mathsf{FCRS}_m(\mathbf{w}), \mathsf{FPDE}_m(\mathbf{w}), \mathsf{FDEC}_m^{\mathrm{hor}}(\mathbf{w})\},$$

where:

- $\mathsf{FMSH}_m(\mathbf{w})$ *is a route-faithful conservative sharpening response to* $\mathsf{QMSH}_m(\mathbf{w})$*,*
- $\mathsf{FCRS}_m(\mathbf{w})$ *is a route-faithful conservative crystal response to* $\mathsf{QCRS}_m(\mathbf{w})$*,*
- $\mathsf{FPDE}_m(\mathbf{w})$ *is a route-faithful conservative field/profile response to* $\mathsf{QPDE}_m(\mathbf{w})$*, and*
- $\mathsf{FDEC}_m^{\mathrm{hor}}(\mathbf{w})$ *is a route-faithful conservative decoder/probe response to* $\mathsf{QDEC}_m^{\mathrm{hor}}(\mathbf{w})$*.*

These fulfillment objects are admissible only if they:

- *remain subordinate to the inherited horizon tag* HOR*,*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

- *do not claim a fresh theorematic certificate beyond the range already carried by* $\mathsf{OPIn}_m^{\mathrm{hor}}(\mathbf{w})$, *and*
- *do not relabel the horizon-side route as a boundary-side stop event.*

Define

$$\mathsf{Ful}_m^{\mathrm{cons}}(\mathbf{w};q) := \begin{cases} \mathsf{FMSH}_m(\mathbf{w}), & \text{if } q = \mathsf{QMSH}_m(\mathbf{w}), \\ \mathsf{FCRS}_m(\mathbf{w}), & \text{if } q = \mathsf{QCRS}_m(\mathbf{w}), \\ \mathsf{FPDE}_m(\mathbf{w}), & \text{if } q = \mathsf{QPDE}_m(\mathbf{w}), \\ \mathsf{FDEC}_m^{\mathrm{hor}}(\mathbf{w}), & \text{if } q = \mathsf{QDEC}_m^{\mathrm{hor}}(\mathbf{w}). \end{cases}$$

(ii) If

$$\mathsf{Q}_m^{\mathrm{cons}}(\mathbf{w}) = \mathcal{Q}_m^{\mathrm{bdry}}(\mathbf{w}),$$

define the first boundary-side conservative fulfillment family

$$\mathcal{F}_m^{\mathrm{bdry}}(\mathbf{w}) := \{\mathsf{FSRC}_m(\mathbf{w}), \mathsf{FDFX}_m(\mathbf{w}), \mathsf{FSTOP}_m(\mathbf{w}), \mathsf{FDEC}_m^{\mathrm{bdry}}(\mathbf{w})\},$$

where:

- $\mathsf{FSRC}_m(\mathbf{w})$ *is an anchored source/exposure response to* $\mathsf{QSRC}_m(\mathbf{w})$,
- $\mathsf{FDFX}_m(\mathbf{w})$ *is a defect-front response to* $\mathsf{QDFX}_m(\mathbf{w})$,
- $\mathsf{FSTOP}_m(\mathbf{w})$ *is a stop-sensitive surface response to* $\mathsf{QSTOP}_m(\mathbf{w})$, *and*
- $\mathsf{FDEC}_m^{\mathrm{bdry}}(\mathbf{w})$ *is a boundary-side decoder/probe response to* $\mathsf{QDEC}_m^{\mathrm{bdry}}(\mathbf{w})$.

These fulfillment objects are admissible only if they:

- *remain subordinate to the inherited stop-preserving boundary tag* BDRY,
- *do not smooth away, postpone, or reinterpret the inherited earliest honest boundary stop, and*
- *do not relabel the boundary-side route as a horizon-continuation route.*

Define

$$\mathsf{Ful}_m^{\mathrm{cons}}(\mathbf{w};q) := \begin{cases} \mathsf{FSRC}_m(\mathbf{w}), & \text{if } q = \mathsf{QSRC}_m(\mathbf{w}), \\ \mathsf{FDFX}_m(\mathbf{w}), & \text{if } q = \mathsf{QDFX}_m(\mathbf{w}), \\ \mathsf{FSTOP}_m(\mathbf{w}), & \text{if } q = \mathsf{QSTOP}_m(\mathbf{w}), \\ \mathsf{FDEC}_m^{\mathrm{bdry}}(\mathbf{w}), & \text{if } q = \mathsf{QDEC}_m^{\mathrm{bdry}}(\mathbf{w}). \end{cases}$$

Proposition H24.169 (Every admissible conservative OP-facing request yields exactly one route-faithful conservative fulfillment object)**.** *Assume the setting of Definition H24.168. Then:*

(i) every admissible request $q \in \mathsf{Q}_m^{\mathrm{cons}}(\mathbf{w})$ *yields exactly one fulfillment object* $\mathsf{Ful}_m^{\mathrm{cons}}(\mathbf{w};q)$*;*

(ii) in the horizon-side case, the fulfillment object remains subordinate to the inherited horizon tag and inherited matrix/crystal/field/decoder organization;

(iii) in the boundary-side case, the fulfillment object remains subordinate to the inherited stop-preserving boundary tag and inherited source/defect/stop-surface/decoder organization; and

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(iv) no admissible fulfillment object may manufacture a fresh theorematic certificate, erase the inherited horizon/boundary distinction, or deny the inherited earliest-boundary stop.

Proof. Clause (i) is immediate from Definition H24.168: once the request side and request symbol are fixed, the definition assigns exactly one corresponding fulfillment object. Clause (ii) follows because the horizon-side responses are explicitly defined only as conservative sharpening, crystal, field/profile, or decoder/probe responses subordinate to the inherited horizon-side intake organization and explicit horizon tag. Clause (iii) follows because the boundary-side responses are explicitly defined only as anchored source, defect-front, stop-surface, or decoder/probe responses subordinate to the inherited stop-preserving boundary-side intake organization and explicit boundary tag. Clause (iv) is exactly the certificate-aware discipline inherited from $19\mathrm{AZ}^{(6ae)}$, $19\mathrm{AZ}^{(6af)}$, $19\mathrm{AZ}^{(6ai)}$, $19\mathrm{AZ}^{(6at)}$, $19\mathrm{AZ}^{(6au)}$, and $19\mathrm{AZ}^{(6av)}$, now expressed as the first finite OP-facing fulfillment layer. □

Corollary H24.170 (The present post-archival stage line already carries a first OP-facing fulfillment layer)**.** *For the first-stage word $\mathbf{w}_3 \in \{\mathsf{A},\mathsf{B}\}$, the current active line already carries a first conservative OP-facing fulfillment layer:*

$$\mathsf{Ful}_2^{\mathrm{cons}}(\mathbf{w}_3; q)$$

for every admissible request

$$q \in \mathsf{Q}_2^{\mathrm{cons}}(\mathbf{w}_3).$$

So the post-release continuation now carries not only a conservative OP-facing intake object and request contract, but also the first disciplined response format through which admissible OP-oriented demands may be answered without reopening the route decision itself.

Proof. Immediate from Corollary H24.166 and Proposition H24.169. □

Remark H24.171 (What the first conservative OP-facing response/fulfillment layer now buys)**.** *This is the point where the post-release line first becomes able not only to ask disciplined OP-oriented questions, but also to answer them in a route-faithful conservative format. The horizon-side request family now terminates in finite sharpening/crystal/field/decoder responses under an explicit horizon tag, while the boundary-side request family now terminates in finite source/defect/stop-surface/decoder responses under an explicit stop-preserving boundary tag. Later OP-oriented work can therefore begin from a conservative ask-and-respond protocol instead of improvising both its questions and its response objects from scratch.*

Reader cue. Read the next block as the first conservative OP-facing response aggregation / resolution layer extracted from the fulfillment layer of $19\mathrm{AZ}^{(6aw)}$. Its purpose is to bundle finitely many admissible conservative fulfillment objects into one downstream-readable OP-facing resolution surface without manufacturing a fresh theorematic certificate, erasing the inherited horizon/boundary distinction, or denying the inherited earliest honest boundary stop. The block still does not close any open problem; it fixes the first disciplined way in which multiple conservative answers may be gathered into one route-faithful working resolution.

$19\mathrm{AZ}^{(6ax)}$. First conservative OP-facing response aggregation / resolution layer extracted from the fulfillment layer

After $19\mathrm{AZ}^{(6aw)}$, the post-release line already carries finite conservative fulfillment families on each intake side. The next honest step is therefore to define how finitely many such fulfillment objects may be aggregated into one conservative OP-facing resolution surface. On the horizon

© 2026 Marc Brendecke. Preprint — not peer reviewed.

side the aggregation must remain subordinate to the inherited horizon tag and inherited matrix/crystal/field/decoder organization. On the boundary side the aggregation must remain subordinate to the inherited stop-preserving boundary tag and inherited source/defect/stop-surface/decoder organization. The point is not to solve more than the fulfillment layer already solved, but to package several admissible conservative responses into one finite route-faithful resolution object for later OP-oriented comparison, selection, or continuation.

Definition H24.172 (First conservative OP-facing response aggregation / resolution layer)**.** *Fix a certified finite extension word* $\mathbf{w} \in \mathcal{L}_m^{\mathrm{ext}}$*, its first conservative OP-facing request contract*

$$\mathsf{Q}_m^{\mathrm{cons}}(\mathbf{w}),$$

and its first conservative fulfillment map

$$q \mapsto \mathsf{Ful}_m^{\mathrm{cons}}(\mathbf{w}; q)$$

from Definition H24.168. Let

$$\mathcal{S} \subseteq \mathsf{Q}_m^{\mathrm{cons}}(\mathbf{w})$$

be a finite nonempty admissible request subfamily. Define the first conservative OP-facing resolution object

$$\mathsf{Res}_m^{\mathrm{cons}}(\mathbf{w}; \mathcal{S})$$

by the following cases.

(i) If

$$\mathsf{Q}_m^{\mathrm{cons}}(\mathbf{w}) = \mathcal{Q}_m^{\mathrm{hor}}(\mathbf{w}),$$

define the first horizon-side conservative resolution surface

$$\mathcal{R}_m^{\mathrm{hor}}(\mathbf{w}; \mathcal{S}) := \big(\mathsf{HOR}, \{\mathsf{Ful}_m^{\mathrm{cons}}(\mathbf{w}; q) : q \in \mathcal{S}\}, \mathsf{OPIn}_m^{\mathrm{hor}}(\mathbf{w})\big).$$

This object is admissible only if every included fulfillment object remains subordinate to the inherited horizon tag and no included response claims a fresh theorematic certificate beyond the range already carried by $\mathsf{OPIn}_m^{\mathrm{hor}}(\mathbf{w})$*. Define*

$$\mathsf{Res}_m^{\mathrm{cons}}(\mathbf{w}; \mathcal{S}) := \mathcal{R}_m^{\mathrm{hor}}(\mathbf{w}; \mathcal{S}).$$

(ii) If

$$\mathsf{Q}_m^{\mathrm{cons}}(\mathbf{w}) = \mathcal{Q}_m^{\mathrm{bdry}}(\mathbf{w}),$$

define the first boundary-side conservative resolution surface

$$\mathcal{R}_m^{\mathrm{bdry}}(\mathbf{w}; \mathcal{S}) := \big(\mathsf{BDRY}, \{\mathsf{Ful}_m^{\mathrm{cons}}(\mathbf{w}; q) : q \in \mathcal{S}\}, \mathsf{OPIn}_m^{\mathrm{bdry}}(\mathbf{w})\big).$$

This object is admissible only if every included fulfillment object remains subordinate to the inherited stop-preserving boundary tag and no included response smooths away, postpones, or reinterprets the inherited earliest honest boundary stop. Define

$$\mathsf{Res}_m^{\mathrm{cons}}(\mathbf{w}; \mathcal{S}) := \mathcal{R}_m^{\mathrm{bdry}}(\mathbf{w}; \mathcal{S}).$$

Proposition H24.173 (Every finite admissible conservative fulfillment family yields exactly one route-faithful conservative resolution object)**.** *Assume the setting of Definition H24.172. Then:*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(i) every finite nonempty admissible request subfamily $\mathcal{S} \subseteq \mathsf{Q}_m^{\mathrm{cons}}(\mathbf{w})$ yields exactly one conservative resolution object $\mathsf{Res}_m^{\mathrm{cons}}(\mathbf{w};\mathcal{S})$;

(ii) in the horizon-side case, the resolution object remains subordinate to the inherited horizon tag and inherited matrix/crystal/field/decoder organization;

(iii) in the boundary-side case, the resolution object remains subordinate to the inherited stop-preserving boundary tag and inherited source/defect/stop-surface/decoder organization; and

(iv) no admissible conservative resolution object may manufacture a fresh theorematic certificate, erase the inherited horizon/boundary distinction, or deny the inherited earliest-boundary stop.

Proof. Clause (i) is immediate from Definition H24.172: once the intake side and finite admissible request subfamily $\mathcal{S}$ are fixed, the definition assigns exactly one corresponding resolution surface. Clause (ii) follows because the horizon-side resolution surface is defined only by collecting fulfillment objects already subordinate to the inherited horizon tag together with the inherited horizon-side intake object. Clause (iii) follows because the boundary-side resolution surface is defined only by collecting fulfillment objects already subordinate to the inherited stop-preserving boundary tag together with the inherited boundary-side intake object. Clause (iv) is exactly the certificate-aware discipline inherited from $19\mathrm{AZ}^{(6ae)}$, $19\mathrm{AZ}^{(6af)}$, $19\mathrm{AZ}^{(6ai)}$, $19\mathrm{AZ}^{(6at)}$, $19\mathrm{AZ}^{(6au)}$, $19\mathrm{AZ}^{(6av)}$, and $19\mathrm{AZ}^{(6aw)}$, now expressed as the first finite OP-facing resolution layer. □

Corollary H24.174 (The present post-archival stage line already carries a first OP-facing conservative resolution surface)**.** *For the first-stage word $\mathbf{w}_3 \in \{\mathsf{A},\mathsf{B}\}$, the current active line already carries a first conservative OP-facing resolution surface*

$$\mathsf{Res}_2^{\mathrm{cons}}(\mathbf{w}_3;\mathcal{S})$$

for every finite nonempty admissible request subfamily

$$\mathcal{S} \subseteq \mathsf{Q}_2^{\mathrm{cons}}(\mathbf{w}_3).$$

So the post-release continuation now carries not only a conservative intake object, request contract, and fulfillment layer, but also the first disciplined aggregation surface through which several admissible OP-oriented answers may be bundled without reopening the route decision itself.

Proof. Immediate from Corollary H24.170 and Proposition H24.173. □

Remark H24.175 (What the first conservative OP-facing response aggregation / resolution layer now buys)**.** *This is the point where the post-release line first becomes able not only to ask and answer disciplined OP-oriented questions, but also to gather several conservative answers into one route-faithful resolution surface. The horizon-side line can now bundle finite sharpening/crystal/field/decoder responses under an explicit horizon tag, while the boundary-side line can now bundle finite source/defect/stop-surface/decoder responses under an explicit stop-preserving boundary tag. Later OP-oriented work can therefore begin from a conservative ask/respond/aggregate protocol instead of improvising both its intermediate bundles and its resolution surfaces from scratch.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Reader cue. Read the next block as the first conservative resolution-to-task dispatch layer extracted from the resolution surface of $19\mathrm{AZ}^{(6ax)}$. Its purpose is to turn a finite route-faithful conservative resolution surface into the first finite OP-facing work or comparison program, without manufacturing a fresh theorematic certificate, erasing the inherited horizon/boundary distinction, or denying the inherited earliest honest boundary stop. The block still does not solve any open problem; it fixes the first disciplined way in which a conservative aggregate may be dispatched into explicit next-task families.

$19\mathrm{AZ}^{(6ay)}$. First conservative resolution-to-task dispatch layer extracted from the resolution surface

After $19\mathrm{AZ}^{(6ax)}$, the post-release line already carries finite route-faithful conservative resolution surfaces on each intake side. The next honest step is therefore to define how such a resolution surface may be dispatched into a finite OP-facing work or comparison program. On the horizon side, the dispatch must remain subordinate to the inherited horizon tag and inherited matrix/crystal/field/decoder organization. On the boundary side, the dispatch must remain subordinate to the inherited stop-preserving boundary tag and inherited source/defect/stop-surface/decoder organization. The point is not to claim more than the resolution surface already carries, but to package that finite conservative aggregate into an explicit task family for later OP-oriented comparison, refinement, or packaging.

Definition H24.176 (First conservative resolution-to-task dispatch layer)**.** *Fix a certified finite extension word* $\mathbf{w} \in \mathcal{L}_m^{\mathrm{ext}}$ *and a finite nonempty admissible request subfamily*

$$\mathcal{S} \subseteq \mathsf{Q}_m^{\mathrm{cons}}(\mathbf{w})$$

together with its conservative resolution object

$$\mathsf{Res}_m^{\mathrm{cons}}(\mathbf{w};\mathcal{S})$$

from Definition H24.172. Define the first conservative dispatch object

$$\mathsf{Disp}_m^{\mathrm{cons}}(\mathbf{w};\mathcal{S})$$

by the following cases.

(i) If

$$\mathsf{Res}_m^{\mathrm{cons}}(\mathbf{w};\mathcal{S}) = \mathcal{R}_m^{\mathrm{hor}}(\mathbf{w};\mathcal{S}),$$

define the first horizon-side conservative dispatch task family

$$\mathcal{T}_m^{\mathrm{hor}}(\mathbf{w};\mathcal{S}) := \Big\{\mathsf{TCMP}_m^{\mathrm{hor}}(\mathbf{w};\mathcal{S}), \mathsf{TREF}_m^{\mathrm{hor}}(\mathbf{w};\mathcal{S}), \mathsf{TPACK}_m^{\mathrm{hor}}(\mathbf{w};\mathcal{S})\Big\}.$$

Here

$$\begin{aligned}
\mathsf{TCMP}_m^{\mathrm{hor}}(\mathbf{w};\mathcal{S}) &:= \textit{horizon-side comparison task},\\
\mathsf{TREF}_m^{\mathrm{hor}}(\mathbf{w};\mathcal{S}) &:= \textit{horizon-side refinement task},\\
\mathsf{TPACK}_m^{\mathrm{hor}}(\mathbf{w};\mathcal{S}) &:= \textit{horizon-side packaging task}.
\end{aligned}$$

This task family is admissible only if every task remains subordinate to the inherited horizon tag HOR*, does not manufacture a fresh theorematic certificate, and does not relabel the horizon-side route as a boundary route. Define*

$$\mathsf{Disp}_m^{\mathrm{cons}}(\mathbf{w};\mathcal{S}) := \big(\mathsf{HOR}, \mathsf{Res}_m^{\mathrm{cons}}(\mathbf{w};\mathcal{S}), \mathcal{T}_m^{\mathrm{hor}}(\mathbf{w};\mathcal{S})\big).$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(ii) If

$$\mathsf{Res}_m^{\mathrm{cons}}(\mathbf{w};\mathcal{S}) = \mathcal{R}_m^{\mathrm{bdry}}(\mathbf{w};\mathcal{S}),$$

define the first boundary-side conservative dispatch task family

$$\mathcal{T}_m^{\mathrm{bdry}}(\mathbf{w};\mathcal{S}) := \Big\{\mathsf{TCMP}_m^{\mathrm{bdry}}(\mathbf{w};\mathcal{S}), \mathsf{TPROF}_m^{\mathrm{bdry}}(\mathbf{w};\mathcal{S}), \mathsf{TPACK}_m^{\mathrm{bdry}}(\mathbf{w};\mathcal{S})\Big\}.$$

Here

$$\begin{aligned}\mathsf{TCMP}_m^{\mathrm{bdry}}(\mathbf{w};\mathcal{S}) &:= \textit{boundary-side comparison task},\\ \mathsf{TPROF}_m^{\mathrm{bdry}}(\mathbf{w};\mathcal{S}) &:= \textit{boundary-side profiling task},\\ \mathsf{TPACK}_m^{\mathrm{bdry}}(\mathbf{w};\mathcal{S}) &:= \textit{boundary-side packaging task}.\end{aligned}$$

This task family is admissible only if every task remains subordinate to the inherited stop-preserving boundary tag BDRY*, does not smooth away, postpone, or reinterpret the inherited earliest honest boundary stop, and does not relabel the boundary-side route as a horizon-continuation route. Define*

$$\mathsf{Disp}_m^{\mathrm{cons}}(\mathbf{w};\mathcal{S}) := \big(\mathsf{BDRY}, \mathsf{Res}_m^{\mathrm{cons}}(\mathbf{w};\mathcal{S}), \mathcal{T}_m^{\mathrm{bdry}}(\mathbf{w};\mathcal{S})\big).$$

Proposition H24.177 (Every finite admissible conservative resolution surface yields exactly one route-faithful conservative dispatch object)**.** *Assume the setting of Definition H24.176. Then:*

(i) every finite admissible conservative resolution surface $\mathsf{Res}_m^{\mathrm{cons}}(\mathbf{w};\mathcal{S})$ *yields exactly one dispatch object* $\mathsf{Disp}_m^{\mathrm{cons}}(\mathbf{w};\mathcal{S})$*;*

(ii) in the horizon-side case, the dispatch object remains subordinate to the inherited horizon tag and inherited matrix/crystal/field/decoder organization;

(iii) in the boundary-side case, the dispatch object remains subordinate to the inherited stop-preserving boundary tag and inherited source/defect/stop-surface/decoder organization; and

(iv) no admissible conservative dispatch object may manufacture a fresh theorematic certificate, erase the inherited horizon/boundary distinction, or deny the inherited earliest-boundary stop.

Proof. Clause (i) is immediate from Definition H24.176: once the intake side and finite admissible request subfamily $\mathcal{S}$ are fixed, the corresponding conservative resolution surface is already fixed by $19\mathrm{AZ}^{(6ax)}$, and the present definition assigns exactly one corresponding dispatch task family and dispatch object. Clause (ii) follows because the horizon-side dispatch object is defined only from the inherited horizon-side resolution surface together with finite comparison, refinement-selection, and packaging tasks subordinate to the explicit horizon tag. Clause (iii) follows because the boundary-side dispatch object is defined only from the inherited boundary-side resolution surface together with finite comparison, profiling-selection, and packaging tasks subordinate to the explicit stop-preserving boundary tag. Clause (iv) is exactly the certificate-aware discipline inherited from $19\mathrm{AZ}^{(6ae)}$, $19\mathrm{AZ}^{(6af)}$, $19\mathrm{AZ}^{(6ai)}$, $19\mathrm{AZ}^{(6at)}$, $19\mathrm{AZ}^{(6au)}$, $19\mathrm{AZ}^{(6av)}$, $19\mathrm{AZ}^{(6aw)}$, and $19\mathrm{AZ}^{(6ax)}$, now expressed as the first finite OP-facing dispatch layer. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Corollary H24.178 (The present post-archival stage line already carries a first conservative dispatch object)**.** *For the first-stage word* $\mathbf{w}_3 \in \{\mathsf{A}, \mathsf{B}\}$*, the current active line already carries a first conservative dispatch object*

$$\mathsf{Disp}_2^{\mathrm{cons}}(\mathbf{w}_3; \mathcal{S})$$

for every finite nonempty admissible request subfamily

$$\mathcal{S} \subseteq \mathsf{Q}_2^{\mathrm{cons}}(\mathbf{w}_3).$$

So the post-release continuation now carries not only a conservative ask/respond/aggregate protocol, but also the first disciplined dispatch format through which an admissible conservative aggregate may be turned into an explicit OP-facing work or comparison program without reopening the route decision itself.

Proof. Immediate from Corollary H24.174 and Proposition H24.177. □

Remark H24.179 (What the first conservative resolution-to-task dispatch layer now buys)**.** *This is the point where the post-release line first becomes able not only to ask, answer, and aggregate disciplined OP-oriented questions, but also to dispatch the resulting conservative aggregate into a finite work or comparison program. The horizon-side line can now dispatch finite conservative fulfillment bundles into comparison, refinement-selection, and packaging tasks under an explicit horizon tag, while the boundary-side line can now dispatch finite conservative fulfillment bundles into comparison, profiling-selection, and packaging tasks under an explicit stop-preserving boundary tag. Later OP-oriented work can therefore begin from a conservative ask/respond/aggregate/dispatch protocol instead of improvising both its finite task programs and its dispatch interface from scratch.*

Reader cue. Read the next block as the first conservative OP-facing comparison/work program normal form extracted from the dispatch layer of $19\mathrm{AZ}^{(6ay)}$. Its purpose is to compress a route-faithful conservative dispatch object into one downstream-readable finite work/comparison program without reopening the route decision, manufacturing a fresh theorematic certificate, erasing the inherited horizon/boundary distinction, or denying the inherited earliest honest boundary stop. The block still does not solve any open problem; it fixes the first compact normal form in which the new dispatch layer may be consumed by later OP-oriented comparison and work.

$19\mathrm{AZ}^{(6az)}$. First conservative OP-facing comparison/work program normal form extracted from the dispatch layer

After $19\mathrm{AZ}^{(6ay)}$, the post-release line already carries route-faithful conservative dispatch objects whose task families remain subordinate either to the inherited horizon tag or to the inherited stop-preserving boundary tag. The next honest step is therefore to compress such a dispatch object into one finite downstream-readable normal form. On the horizon side, that normal form should package the inherited comparison, refinement-selection, and packaging tasks into one compact work/comparison program subordinate to the explicit horizon tag. On the boundary side, it should package the inherited comparison, profiling-selection, and packaging tasks into one compact work/comparison program subordinate to the explicit stop-preserving boundary tag. The point is not to add any new theorematic force, but only to normalize the already admissible dispatched task family into a short OP-facing work-program object.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition H24.180 (First conservative OP-facing comparison/work program normal form)**.** *Fix a certified finite extension word* $\mathbf{w} \in \mathcal{L}_m^{\mathrm{ext}}$ *and a finite nonempty admissible request subfamily*

$$\mathcal{S} \subseteq \mathsf{Q}_m^{\mathrm{cons}}(\mathbf{w})$$

together with its conservative dispatch object

$$\mathsf{Disp}_m^{\mathrm{cons}}(\mathbf{w};\mathcal{S})$$

from Definition H24.176. Define the first conservative OP-facing comparison/work program normal form

$$\mathsf{Prog}_m^{\mathrm{cons}}(\mathbf{w};\mathcal{S})$$

by the following cases.

(i) If

$$\mathsf{Disp}_m^{\mathrm{cons}}(\mathbf{w};\mathcal{S}) = \big(\mathsf{HOR}, \mathsf{Res}_m^{\mathrm{cons}}(\mathbf{w};\mathcal{S}), \mathcal{T}_m^{\mathrm{hor}}(\mathbf{w};\mathcal{S})\big),$$

define the first horizon-side conservative comparison/work program normal form

$$\mathcal{P}_m^{\mathrm{hor}}(\mathbf{w};\mathcal{S}) := \big(\mathsf{HOR}, \mathsf{TCMP}_m^{\mathrm{hor}}(\mathbf{w};\mathcal{S}), \mathsf{TREF}_m^{\mathrm{hor}}(\mathbf{w};\mathcal{S}), \mathsf{TPACK}_m^{\mathrm{hor}}(\mathbf{w};\mathcal{S})\big).$$

Define

$$\mathsf{Prog}_m^{\mathrm{cons}}(\mathbf{w};\mathcal{S}) := \mathcal{P}_m^{\mathrm{hor}}(\mathbf{w};\mathcal{S}).$$

This normal form is admissible only if it remains subordinate to the inherited horizon tag HOR*, does not manufacture a fresh theorematic certificate, and does not relabel the horizon-side route as a boundary route.*

(ii) If

$$\mathsf{Disp}_m^{\mathrm{cons}}(\mathbf{w};\mathcal{S}) = \big(\mathsf{BDRY}, \mathsf{Res}_m^{\mathrm{cons}}(\mathbf{w};\mathcal{S}), \mathcal{T}_m^{\mathrm{bdry}}(\mathbf{w};\mathcal{S})\big),$$

define the first boundary-side conservative comparison/work program normal form

$$\mathcal{P}_m^{\mathrm{bdry}}(\mathbf{w};\mathcal{S}) := \big(\mathsf{BDRY}, \mathsf{TCMP}_m^{\mathrm{bdry}}(\mathbf{w};\mathcal{S}), \mathsf{TPROF}_m^{\mathrm{bdry}}(\mathbf{w};\mathcal{S}), \mathsf{TPACK}_m^{\mathrm{bdry}}(\mathbf{w};\mathcal{S})\big).$$

Define

$$\mathsf{Prog}_m^{\mathrm{cons}}(\mathbf{w};\mathcal{S}) := \mathcal{P}_m^{\mathrm{bdry}}(\mathbf{w};\mathcal{S}).$$

This normal form is admissible only if it remains subordinate to the inherited stop-preserving boundary tag BDRY*, does not smooth away, postpone, or reinterpret the inherited earliest honest boundary stop, and does not relabel the boundary-side route as a horizon-continuation route.*

Proposition H24.181 (Every finite admissible conservative dispatch object yields exactly one conservative comparison/work program normal form)**.** *Assume the setting of Definition H24.180. Then:*

(i) every finite admissible conservative dispatch object $\mathsf{Disp}_m^{\mathrm{cons}}(\mathbf{w};\mathcal{S})$ *yields exactly one program normal form* $\mathsf{Prog}_m^{\mathrm{cons}}(\mathbf{w};\mathcal{S})$*;*

(ii) in the horizon-side case, the program normal form remains subordinate to the inherited horizon tag and inherited comparison/refinement/packaging organization;

(iii) in the boundary-side case, the program normal form remains subordinate to the inherited stop-preserving boundary tag and inherited comparison/profiling/packaging organization; and

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(iv) no admissible conservative program normal form may manufacture a fresh theorematic certificate, erase the inherited horizon/boundary distinction, or deny the inherited earliest-boundary stop.

Proof. Clause (i) is immediate from Definition H24.180: once the dispatch object is fixed by 19AZ$^{(6ay)}$, the present definition assigns exactly one corresponding compact program tuple. Clause (ii) follows because the horizon-side program normal form is defined only from the inherited horizon-side dispatch object together with the already admissible comparison, refinement-selection, and packaging tasks subordinate to the explicit horizon tag. Clause (iii) follows because the boundary-side program normal form is defined only from the inherited boundary-side dispatch object together with the already admissible comparison, profiling-selection, and packaging tasks subordinate to the explicit stop-preserving boundary tag. Clause (iv) is exactly the certificate-aware discipline inherited from 19AZ$^{(6ae)}$, 19AZ$^{(6af)}$, 19AZ$^{(6ai)}$, 19AZ$^{(6at)}$, 19AZ$^{(6au)}$, 19AZ$^{(6av)}$, 19AZ$^{(6aw)}$, 19AZ$^{(6ax)}$, and 19AZ$^{(6ay)}$, now expressed as the first compact OP-facing comparison/work program normal form. □

Corollary H24.182 (The present post-archival stage line already carries a first conservative comparison/work program normal form)**.** *For the first-stage word* $\mathbf{w}_3 \in \{\mathsf{A}, \mathsf{B}\}$*, the current active line already carries a first conservative comparison/work program normal form*

$$\mathsf{Prog}_2^{\mathrm{cons}}(\mathbf{w}_3; \mathcal{S})$$

for every finite nonempty admissible request subfamily

$$\mathcal{S} \subseteq \mathsf{Q}_2^{\mathrm{cons}}(\mathbf{w}_3).$$

So the post-release continuation now carries not only a conservative ask/respond/aggregate/dispatch protocol, but also the first compact downstream-readable work/comparison normal form through which an admissible conservative dispatch object may be consumed without reopening the route decision itself.

Proof. Immediate from Corollary H24.178 and Proposition H24.181. □

Remark H24.183 (What the first conservative comparison/work program normal form now buys)**.** *This is the point where the post-release line first becomes able not only to dispatch conservative aggregates into explicit task families, but also to compress those task families into one compact OP-facing normal form. The horizon-side line can now hand downstream readers one finite comparison/refinement/packaging program under an explicit horizon tag, while the boundary-side line can now hand downstream readers one finite comparison/profiling/packaging program under an explicit stop-preserving boundary tag. Later OP-oriented work can therefore begin from a conservative ask/respond/aggregate/dispatch/program protocol instead of improvising both its compact work-program objects and its downstream comparison interface from scratch.*

Reader cue. Read the next block as the first conservative OP-facing program-to-execution / work-cycle layer extracted from the compact comparison/work program normal form of 19AZ$^{(6az)}$. Its purpose is to turn one downstream-readable conservative program object back into one disciplined, finite execution cycle without reopening the route decision, manufacturing a fresh theorematic certificate, erasing the inherited horizon/boundary distinction, or denying the inherited earliest honest boundary stop. The block still does not solve any open problem; it fixes the first controlled cycle in which the compact OP-facing program may actually be run.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

19AZ$^{(6ba)}$. Conservative program-to-execution / work-cycle layer

After 19AZ$^{(6az)}$, the post-release line already carries compact route-faithful conservative comparison/work program normal forms subordinate either to the inherited horizon tag or to the inherited stop-preserving boundary tag. The next honest step is therefore to define how such a compact program may be executed as a first finite conservative work cycle. On the horizon side, that cycle must remain subordinate to the inherited horizon tag and must execute only comparison, conservative refinement, and packaging moves already encoded in the compact program. On the boundary side, it must remain subordinate to the inherited stop-preserving boundary tag and must execute only comparison, conservative profiling, and packaging moves already encoded in the compact program. The point is not to add any new theorematic force, but only to turn the compact conservative program into a first controlled OP-facing work cycle that may later be iterated, compared, or packaged further.

Definition H24.184 (First conservative OP-facing program-to-execution / work-cycle layer)**.** *Fix a certified finite extension word* $\mathbf{w} \in \mathcal{L}_m^{\mathrm{ext}}$ *and a finite nonempty admissible request subfamily*

$$\mathcal{S} \subseteq \mathsf{Q}_m^{\mathrm{cons}}(\mathbf{w})$$

together with its compact conservative comparison/work program normal form

$$\mathsf{Prog}_m^{\mathrm{cons}}(\mathbf{w};\mathcal{S})$$

from Definition H24.180. Define the first conservative OP-facing work-cycle object

$$\mathsf{Cycle}_m^{\mathrm{cons}}(\mathbf{w};\mathcal{S})$$

by the following cases.

(i) If

$$\mathsf{Prog}_m^{\mathrm{cons}}(\mathbf{w};\mathcal{S}) = \mathcal{P}_m^{\mathrm{hor}}(\mathbf{w};\mathcal{S}),$$

define the first horizon-side conservative work cycle

$$\mathcal{C}_m^{\mathrm{hor}}(\mathbf{w};\mathcal{S}) := \big(\mathsf{HOR}, \mathsf{TCMP}_m^{\mathrm{hor}}(\mathbf{w};\mathcal{S}), \mathsf{TREF}_m^{\mathrm{hor}}(\mathbf{w};\mathcal{S}), \mathsf{TPACK}_m^{\mathrm{hor}}(\mathbf{w};\mathcal{S}), \mathsf{RET}_m^{\mathrm{hor}}(\mathbf{w};\mathcal{S})\big).$$

Here

$$\mathsf{RET}_m^{\mathrm{hor}}(\mathbf{w};\mathcal{S}) := \textit{horizon-side finite return package.}$$

This work cycle is admissible only if every executed move remains subordinate to the inherited horizon tag HOR*, does not manufacture a fresh theorematic certificate, and does not relabel the horizon-side route as a boundary route. Define*

$$\mathsf{Cycle}_m^{\mathrm{cons}}(\mathbf{w};\mathcal{S}) := \mathcal{C}_m^{\mathrm{hor}}(\mathbf{w};\mathcal{S}).$$

(ii) If

$$\mathsf{Prog}_m^{\mathrm{cons}}(\mathbf{w};\mathcal{S}) = \mathcal{P}_m^{\mathrm{bdry}}(\mathbf{w};\mathcal{S}),$$

define the first boundary-side conservative work cycle

$$\mathcal{C}_m^{\mathrm{bdry}}(\mathbf{w};\mathcal{S}) := \big(\mathsf{BDRY}, \mathsf{TCMP}_m^{\mathrm{bdry}}(\mathbf{w};\mathcal{S}), \mathsf{TPROF}_m^{\mathrm{bdry}}(\mathbf{w};\mathcal{S}), \mathsf{TPACK}_m^{\mathrm{bdry}}(\mathbf{w};\mathcal{S}), \mathsf{RET}_m^{\mathrm{bdry}}(\mathbf{w};\mathcal{S})\big).$$

Here

$$\mathsf{RET}_m^{\mathrm{bdry}}(\mathbf{w};\mathcal{S}) := \textit{boundary-side finite return package.}$$

This work cycle is admissible only if every executed move remains subordinate to the inherited stop-preserving boundary tag BDRY*, does not smooth away, postpone, or reinterpret the inherited earliest honest boundary stop, and does not relabel the boundary-side route as a horizon-continuation route. Define*

$$\mathsf{Cycle}_m^{\mathrm{cons}}(\mathbf{w};\mathcal{S}) := \mathcal{C}_m^{\mathrm{bdry}}(\mathbf{w};\mathcal{S}).$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proposition H24.185 (Every finite admissible conservative comparison/work program normal form yields exactly one conservative work-cycle object)**.** *Assume the setting of Definition H24.184. Then:*

(i) every finite admissible conservative comparison/work program normal form $\mathsf{Prog}_m^{\mathrm{cons}}(\mathbf{w};\mathcal{S})$ *yields exactly one work-cycle object* $\mathsf{Cycle}_m^{\mathrm{cons}}(\mathbf{w};\mathcal{S})$*;*

(ii) in the horizon-side case, the work-cycle object remains subordinate to the inherited horizon tag and inherited comparison/refinement/packaging organization;

(iii) in the boundary-side case, the work-cycle object remains subordinate to the inherited stop-preserving boundary tag and inherited comparison/profiling/packaging organization; and

(iv) no admissible conservative work-cycle object may manufacture a fresh theorematic certificate, erase the inherited horizon/boundary distinction, or deny the inherited earliest-boundary stop.

Proof. Clause (i) is immediate from Definition H24.184: once the compact program normal form is fixed by $19\mathrm{AZ}^{(6az)}$, the present definition assigns exactly one corresponding finite work-cycle object. Clause (ii) follows because the horizon-side work cycle is defined only from the inherited horizon-side program normal form together with the already admitted comparison, refinement-selection, and packaging tasks subordinate to the explicit horizon tag. Clause (iii) follows because the boundary-side work cycle is defined only from the inherited boundary-side program normal form together with the already admitted comparison, profiling-selection, and packaging tasks subordinate to the explicit stop-preserving boundary tag. Clause (iv) is exactly the certificate-aware discipline inherited from $19\mathrm{AZ}^{(6ae)}$, $19\mathrm{AZ}^{(6af)}$, $19\mathrm{AZ}^{(6ai)}$, $19\mathrm{AZ}^{(6at)}$, $19\mathrm{AZ}^{(6au)}$, $19\mathrm{AZ}^{(6av)}$, $19\mathrm{AZ}^{(6aw)}$, $19\mathrm{AZ}^{(6ax)}$, $19\mathrm{AZ}^{(6ay)}$, and $19\mathrm{AZ}^{(6az)}$, now expressed as the first controlled OP-facing work cycle. □

Corollary H24.186 (The present post-archival stage line already carries a first conservative work cycle)**.** *For the first-stage word* $\mathbf{w}_3 \in \{\mathsf{A},\mathsf{B}\}$*, the current active line already carries a first conservative work-cycle object*

$$\mathsf{Cycle}_2^{\mathrm{cons}}(\mathbf{w}_3;\mathcal{S})$$

for every finite nonempty admissible request subfamily

$$\mathcal{S} \subseteq \mathsf{Q}_2^{\mathrm{cons}}(\mathbf{w}_3).$$

So the post-release continuation now carries not only a conservative ask/respond/aggregate/dispatch/program protocol, but also the first controlled OP-facing execution cycle through which an admissible compact work/comparison program may be run without reopening the route decision itself.

Proof. Immediate from Corollary H24.182 and Proposition H24.185. □

Remark H24.187 (What the first conservative program-to-execution / work-cycle layer now buys)**.** *This is the point where the post-release line first becomes able not only to package conservative OP-facing task families into compact program objects, but also to run those compact programs as controlled finite work cycles. The horizon-side line can now execute one finite comparison/refinement/packaging cycle under an explicit horizon tag, while the boundary-side line can now execute one finite comparison/profiling/packaging cycle under an explicit stop-preserving boundary tag. Later OP-oriented work can therefore begin from a conservative ask/respond/aggregate/dispatch/program/cycle protocol instead of improvising both its execution loop and its return package from scratch.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Reader cue. Read the next block as the first conservative cycle-to-return / iteration interface extracted from the controlled work-cycle layer of 19AZ$^{(6ba)}$. Its purpose is to decide what a route-faithful return package may honestly do next: either prepare one further conservative cycle under the inherited route tag, or collapse into one compact conservative closing/comparison object without reopening the route decision, manufacturing a fresh theorematic certificate, erasing the inherited horizon/boundary distinction, or denying the inherited earliest honest boundary stop.

19AZ$^{(6bb)}$. Conservative cycle-to-return / iteration interface

After 19AZ$^{(6ba)}$, the post-release line already carries controlled finite conservative work-cycle objects subordinate either to the inherited horizon tag or to the inherited stop-preserving boundary tag. The next honest step is therefore to define how the return package produced by such a cycle may be consumed. On the horizon side, a route-faithful return package may only either seed one further conservative horizon-side cycle under the inherited horizon tag or be collapsed into one compact conservative comparison/closing object that still records the inherited route. On the boundary side, a stop-preserving return package may only either seed one further conservative boundary-side cycle under the inherited stop-preserving boundary tag or be collapsed into one compact conservative comparison/closing object that still records the inherited earliest honest boundary stop. The point is not to force infinite iteration, but to define the first disciplined interface through which the conservative work cycle either continues or closes.

Definition H24.188 (First conservative cycle-to-return / iteration interface)**.** *Fix a certified finite extension word* $\mathbf{w} \in \mathcal{L}_m^{\mathrm{ext}}$*, a finite nonempty admissible request subfamily*

$$\mathcal{S} \subseteq \mathsf{Q}_m^{\mathrm{cons}}(\mathbf{w})$$

and its controlled conservative work-cycle object

$$\mathsf{Cycle}_m^{\mathrm{cons}}(\mathbf{w};\mathcal{S})$$

from Definition H24.184. Define the associated first conservative cycle-to-return / iteration object

$$\mathsf{Iter}_m^{\mathrm{cons}}(\mathbf{w};\mathcal{S})$$

by the following cases.

(i) If

$$\mathsf{Cycle}_m^{\mathrm{cons}}(\mathbf{w};\mathcal{S}) = \mathcal{C}_m^{\mathrm{hor}}(\mathbf{w};\mathcal{S}),$$

define the first horizon-side conservative return/iteration object

$$\mathcal{I}_m^{\mathrm{hor}}(\mathbf{w};\mathcal{S}) := \big(\mathsf{HOR}, \mathsf{RET}_m^{\mathrm{hor}}(\mathbf{w};\mathcal{S}), \mathsf{NEXT}_m^{\mathrm{hor}}(\mathbf{w};\mathcal{S}), \mathsf{CLOSE}_m^{\mathrm{hor}}(\mathbf{w};\mathcal{S})\big).$$

Here

$$\mathsf{NEXT}_m^{\mathrm{hor}}(\mathbf{w};\mathcal{S}) := \textit{next horizon-side conservative cycle option},$$
$$\mathsf{CLOSE}_m^{\mathrm{hor}}(\mathbf{w};\mathcal{S}) := \textit{compact horizon-side closing option}.$$

This return/iteration object is admissible only if both options remain subordinate to the inherited horizon tag HOR*, do not manufacture a fresh theorematic certificate, and do not relabel the horizon-side route as a boundary route. Define*

$$\mathsf{Iter}_m^{\mathrm{cons}}(\mathbf{w};\mathcal{S}) := \mathcal{I}_m^{\mathrm{hor}}(\mathbf{w};\mathcal{S}).$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(ii) If

$$\mathsf{Cycle}_m^{\mathrm{cons}}(\mathbf{w};\mathcal{S}) = \mathcal{C}_m^{\mathrm{bdry}}(\mathbf{w};\mathcal{S}),$$

define the first boundary-side conservative return/iteration object

$$\mathcal{I}_m^{\mathrm{bdry}}(\mathbf{w};\mathcal{S}) := \big(\mathsf{BDRY}, \mathsf{RET}_m^{\mathrm{bdry}}(\mathbf{w};\mathcal{S}), \mathsf{NEXT}_m^{\mathrm{bdry}}(\mathbf{w};\mathcal{S}), \mathsf{CLOSE}_m^{\mathrm{bdry}}(\mathbf{w};\mathcal{S})\big).$$

Here

$$\mathsf{NEXT}_m^{\mathrm{bdry}}(\mathbf{w};\mathcal{S}) := \textit{next boundary-side conservative cycle option},$$
$$\mathsf{CLOSE}_m^{\mathrm{bdry}}(\mathbf{w};\mathcal{S}) := \textit{compact boundary-side closing option}.$$

This return/iteration object is admissible only if both options remain subordinate to the inherited stop-preserving boundary tag BDRY*, do not smooth away, postpone, or reinterpret the inherited earliest honest boundary stop, and do not relabel the boundary-side route as a horizon-continuation route. Define*

$$\mathsf{Iter}_m^{\mathrm{cons}}(\mathbf{w};\mathcal{S}) := \mathcal{I}_m^{\mathrm{bdry}}(\mathbf{w};\mathcal{S}).$$

Proposition H24.189 (Every finite admissible conservative work cycle yields exactly one conservative return/iteration object)**.** *Assume the setting of Definition H24.188. Then:*

(i) every finite admissible conservative work-cycle object $\mathsf{Cycle}_m^{\mathrm{cons}}(\mathbf{w};\mathcal{S})$ *yields exactly one return/iteration object* $\mathsf{Iter}_m^{\mathrm{cons}}(\mathbf{w};\mathcal{S})$*;*

(ii) in the horizon-side case, the return/iteration object remains subordinate to the inherited horizon tag and records exactly the two admissible next uses of the return package: further conservative cycling or compact horizon-side conservative closing/comparison;

(iii) in the boundary-side case, the return/iteration object remains subordinate to the inherited stop-preserving boundary tag and records exactly the two admissible next uses of the return package: further conservative cycling or compact boundary-side conservative closing/comparison; and

(iv) no admissible conservative return/iteration object may manufacture a fresh theorematic certificate, erase the inherited horizon/boundary distinction, or deny the inherited earliest-boundary stop.

Proof. Clause (i) is immediate from Definition H24.188: once the conservative work-cycle object is fixed by $19\mathrm{AZ}^{(6ba)}$, the present definition assigns exactly one corresponding return/iteration object. Clause (ii) follows because the horizon-side return/iteration object is defined only from the inherited horizon-side work cycle together with the two explicitly admitted horizon-side uses of the return package, namely further conservative cycling or compact horizon-side closing/comparison. Clause (iii) follows because the boundary-side return/iteration object is defined only from the inherited boundary-side work cycle together with the two explicitly admitted boundary-side uses of the return package, namely further conservative cycling or compact boundary-side closing/comparison subordinate to the inherited earliest honest boundary stop. Clause (iv) is exactly the certificate-aware discipline inherited from $19\mathrm{AZ}^{(6ae)}$, $19\mathrm{AZ}^{(6af)}$, $19\mathrm{AZ}^{(6ai)}$, $19\mathrm{AZ}^{(6at)}$, $19\mathrm{AZ}^{(6au)}$, $19\mathrm{AZ}^{(6av)}$, $19\mathrm{AZ}^{(6aw)}$, $19\mathrm{AZ}^{(6ax)}$, $19\mathrm{AZ}^{(6ay)}$, $19\mathrm{AZ}^{(6az)}$, and $19\mathrm{AZ}^{(6ba)}$, now expressed as the first disciplined conservative return/iteration interface. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Corollary H24.190 (The present post-archival stage line already carries a first conservative return/iteration interface)**.** *For the first-stage word* $\mathbf{w}_3 \in \{\mathsf{A}, \mathsf{B}\}$*, the current active line already carries a first conservative return/iteration object*

$$\mathsf{Iter}_2^{\mathrm{cons}}(\mathbf{w}_3; \mathcal{S})$$

for every finite nonempty admissible request subfamily

$$\mathcal{S} \subseteq \mathsf{Q}_2^{\mathrm{cons}}(\mathbf{w}_3).$$

So the post-release continuation now carries not only a conservative ask/respond/aggregate/dispatch/program/cycle protocol, but also the first disciplined interface through which a finite conservative work cycle may either seed one further conservative cycle or collapse into one compact route-faithful conservative closing/comparison object without reopening the route decision itself.

Proof. Immediate from Corollary H24.186 and Proposition H24.189. □

Remark H24.191 (What the first conservative cycle-to-return / iteration interface now buys)**.** *This is the point where the post-release line first becomes able not only to run a compact conservative OP-facing program as a controlled finite work cycle, but also to state explicitly what the return package of that cycle is allowed to do next. The horizon-side line can now either seed one further horizon-side conservative cycle or collapse into one compact horizon-side conservative closing/comparison object under the inherited horizon tag, while the boundary-side line can now either seed one further boundary-side conservative cycle or collapse into one compact boundary-side conservative closing/comparison object under the inherited stop-preserving boundary tag. Later OP-oriented work can therefore begin from a conservative ask/respond/aggregate/dispatch/program/cycle/return protocol instead of improvising both its iteration interface and its compact closing interface from scratch.*

Reader cue. Read the next block as the first compact conservative closing/comparison form extracted from the return/iteration interface of $19\mathrm{AZ}^{(6bb)}$. Its purpose is to package the honest non-iterative exit of a controlled conservative work cycle into one downstream-readable closing object that still records the inherited route tag, does not reopen the route decision, does not manufacture a fresh theorematic certificate, and does not smooth away the inherited earliest honest boundary stop.

$19\mathrm{AZ}^{(6bc)}$. First compact conservative closing/comparison form extracted from the return/iteration interface

After $19\mathrm{AZ}^{(6bb)}$, every controlled conservative work cycle already carries a disciplined return/iteration object with exactly two admitted next uses of the return package: one further conservative cycle, or one compact route-faithful closing/comparison object. The present block isolates the second option as a compact object in its own right. On the horizon side, this yields a compact horizon-side conservative closing/comparison form subordinate to the inherited horizon tag. On the boundary side, it yields a compact boundary-side conservative closing/comparison form subordinate to the inherited stop-preserving boundary tag and earliest honest boundary stop. The point is not to terminate the whole post-release line, but to define the first disciplined compact conservative exit object of the current ask/respond/aggregate/dispatch/program/cycle/return staircase.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition H24.192 (First compact conservative closing/comparison form)**.** *Fix a certified finite extension word* $\mathbf{w} \in \mathcal{L}_m^{\mathrm{ext}}$*, a finite nonempty admissible request subfamily*

$$\mathcal{S} \subseteq \mathsf{Q}_m^{\mathrm{cons}}(\mathbf{w})$$

and its conservative return/iteration object

$$\mathsf{Iter}_m^{\mathrm{cons}}(\mathbf{w};\mathcal{S})$$

from Definition H24.188. Define the associated first compact conservative closing/comparison object

$$\mathsf{CloseCmp}_m^{\mathrm{cons}}(\mathbf{w};\mathcal{S})$$

by the following cases.

(i) If

$$\mathsf{Iter}_m^{\mathrm{cons}}(\mathbf{w};\mathcal{S}) = \mathcal{I}_m^{\mathrm{hor}}(\mathbf{w};\mathcal{S}),$$

define the first horizon-side compact conservative closing/comparison object

$$\mathcal{K}_m^{\mathrm{hor}}(\mathbf{w};\mathcal{S}) := \big(\mathsf{HOR}, \mathsf{RET}_m^{\mathrm{hor}}(\mathbf{w};\mathcal{S}), \mathsf{CLOSE}_m^{\mathrm{hor}}(\mathbf{w};\mathcal{S})\big).$$

This object is admissible only if it remains subordinate to the inherited horizon tag, records the compact horizon-side closing/comparison option exactly as provided by 19AZ$^{(6bb)}$*, does not relabel the route as boundary-side, and does not manufacture a fresh theorematic certificate. Define*

$$\mathsf{CloseCmp}_m^{\mathrm{cons}}(\mathbf{w};\mathcal{S}) := \mathcal{K}_m^{\mathrm{hor}}(\mathbf{w};\mathcal{S}).$$

(ii) If

$$\mathsf{Iter}_m^{\mathrm{cons}}(\mathbf{w};\mathcal{S}) = \mathcal{I}_m^{\mathrm{bdry}}(\mathbf{w};\mathcal{S}),$$

define the first boundary-side compact conservative closing/comparison object

$$\mathcal{K}_m^{\mathrm{bdry}}(\mathbf{w};\mathcal{S}) := \big(\mathsf{BDRY}, \mathsf{RET}_m^{\mathrm{bdry}}(\mathbf{w};\mathcal{S}), \mathsf{CLOSE}_m^{\mathrm{bdry}}(\mathbf{w};\mathcal{S})\big).$$

This object is admissible only if it remains subordinate to the inherited stop-preserving boundary tag, records the compact boundary-side closing/comparison option exactly as provided by 19AZ$^{(6bb)}$*, does not smooth away or postpone the inherited earliest honest boundary stop, and does not relabel the route as horizon-side. Define*

$$\mathsf{CloseCmp}_m^{\mathrm{cons}}(\mathbf{w};\mathcal{S}) := \mathcal{K}_m^{\mathrm{bdry}}(\mathbf{w};\mathcal{S}).$$

Proposition H24.193 (Every finite admissible conservative return/iteration object yields exactly one compact conservative closing/comparison object)**.** *Assume the setting of Definition H24.192. Then:*

(i) every finite admissible conservative return/iteration object $\mathsf{Iter}_m^{\mathrm{cons}}(\mathbf{w};\mathcal{S})$ *yields exactly one compact conservative closing/comparison object* $\mathsf{CloseCmp}_m^{\mathrm{cons}}(\mathbf{w};\mathcal{S})$*;*

(ii) in the horizon-side case, the compact conservative closing/comparison object remains subordinate to the inherited horizon tag and records exactly the compact horizon-side closing/comparison option carried by the return package;

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(iii) *in the boundary-side case, the compact conservative closing/comparison object remains subordinate to the inherited stop-preserving boundary tag and records exactly the compact boundary-side closing/comparison option carried by the return package; and*

(iv) *no admissible compact conservative closing/comparison object may manufacture a fresh theorematic certificate, erase the inherited horizon/boundary distinction, or deny the inherited earliest-boundary stop.*

Proof. Clause (i) is immediate from Definition H24.192: once the conservative return/iteration object is fixed by $19\mathrm{AZ}^{(6bb)}$, the present definition assigns exactly one corresponding compact conservative closing/comparison object. Clauses (ii) and (iii) follow because the horizon-side and boundary-side compact objects are defined only from the inherited return package together with the corresponding compact closing/comparison option already recorded in the return/iteration object. Clause (iv) is exactly the certificate-aware discipline inherited from $19\mathrm{AZ}^{(6ae)}$ through $19\mathrm{AZ}^{(6bb)}$, now expressed as the first disciplined compact conservative closing/comparison interface. □

Corollary H24.194 (The present post-archival stage line already carries a first compact conservative closing/comparison form)**.** *For the first-stage word* $\mathbf{w}_3 \in \{\mathsf{A},\mathsf{B}\}$*, the current active line already carries a first compact conservative closing/comparison object*

$$\mathsf{CloseCmp}_2^{\mathrm{cons}}(\mathbf{w}_3;\mathcal{S})$$

for every finite nonempty admissible request subfamily

$$\mathcal{S} \subseteq \mathsf{Q}_2^{\mathrm{cons}}(\mathbf{w}_3).$$

So the post-release continuation now carries not only a conservative ask/respond/aggregate/dispatch/program/cycle/return protocol, but also the first disciplined compact conservative closing/comparison form to which that protocol may honestly collapse without reopening the route decision itself.

Proof. Immediate from Corollary H24.190 and Proposition H24.193. □

Remark H24.195 (What the first compact conservative closing/comparison form now buys)**.** *This is the point where the post-release line first becomes able not only to iterate a conservative work cycle, but also to exit one such cycle in a compact downstream-readable form that still remembers the inherited route. The horizon-side line can now close one controlled conservative cycle into a compact horizon-side comparison object under the inherited horizon tag, while the boundary-side line can now close one controlled conservative cycle into a compact boundary-side comparison object under the inherited stop-preserving boundary tag and earliest honest boundary stop. Later OP-oriented work can therefore begin from a conservative ask/respond/aggregate/dispatch/program/cycle/return/close protocol instead of improvising its compact conservative closing interface from scratch.*

Reader cue. Read the next block as the first admissible next-cycle promotion rule extracted from the same return/iteration interface of $19\mathrm{AZ}^{(6bb)}$. Its purpose is to state when the *iterative* branch of the return package may honestly be promoted to one further conservative work cycle under the inherited route tag, rather than merely being mentioned abstractly as a possibility.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

19AZ$^{(6bd)}$. First admissible next-cycle promotion rule extracted from the return/iteration interface

After 19AZ$^{(6bb)}$, every controlled conservative work cycle already carries a return package with an explicitly recorded next-cycle option. The present block upgrades that option from a mere named possibility to a disciplined promotion rule. On the horizon side, a next-cycle promotion is admissible only if one chooses a new finite nonempty admissible request subfamily that remains subordinate to the inherited horizon tag and does not relabel the route. On the boundary side, a next-cycle promotion is admissible only if one chooses a new finite nonempty admissible request subfamily that remains subordinate to the inherited stop-preserving boundary tag and does not smooth away or postpone the inherited earliest honest boundary stop. The point is to define the first honest rule under which NEXT may seed one further conservative cycle rather than being left as an unspecified placeholder.

Definition H24.196 (First admissible next-cycle promotion rule)**.** *Fix a certified finite extension word* $\mathbf{w} \in \mathcal{L}_m^{\mathrm{ext}}$*, a finite nonempty admissible request subfamily*

$$\mathcal{S} \subseteq \mathsf{Q}_m^{\mathrm{cons}}(\mathbf{w}),$$

its conservative return/iteration object

$$\mathsf{Iter}_m^{\mathrm{cons}}(\mathbf{w};\mathcal{S}),$$

and a second finite nonempty admissible request subfamily

$$\mathcal{S}^+ \subseteq \mathsf{Q}_m^{\mathrm{cons}}(\mathbf{w}).$$

Define the associated first admissible next-cycle promotion object

$$\mathsf{Prom}_m^{\mathrm{cons}}(\mathbf{w};\mathcal{S},\mathcal{S}^+)$$

by the following cases.

(i) If

$$\mathsf{Iter}_m^{\mathrm{cons}}(\mathbf{w};\mathcal{S}) = \mathcal{I}_m^{\mathrm{hor}}(\mathbf{w};\mathcal{S})$$

and

$$\mathcal{S}^+ \subseteq \mathcal{Q}_m^{\mathrm{hor}}(\mathbf{w}),$$

define the first horizon-side admissible next-cycle promotion object

$$\Pi_m^{\mathrm{hor}}(\mathbf{w};\mathcal{S},\mathcal{S}^+) := \big(\mathsf{HOR}, \mathsf{NEXT}_m^{\mathrm{hor}}(\mathbf{w};\mathcal{S}), \mathcal{S}^+, \mathsf{Cycle}_m^{\mathrm{cons}}(\mathbf{w};\mathcal{S}^+)\big).$$

This promotion object is admissible only if the promoted cycle remains subordinate to the inherited horizon tag, does not manufacture a fresh theorematic certificate, and does not relabel the horizon-side route as a boundary-side route. Define

$$\mathsf{Prom}_m^{\mathrm{cons}}(\mathbf{w};\mathcal{S},\mathcal{S}^+) := \Pi_m^{\mathrm{hor}}(\mathbf{w};\mathcal{S},\mathcal{S}^+).$$

(ii) If

$$\mathsf{Iter}_m^{\mathrm{cons}}(\mathbf{w};\mathcal{S}) = \mathcal{I}_m^{\mathrm{bdry}}(\mathbf{w};\mathcal{S})$$

and

$$\mathcal{S}^+ \subseteq \mathcal{Q}_m^{\mathrm{bdry}}(\mathbf{w}),$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

define the first boundary-side admissible next-cycle promotion object

$$\Pi_m^{\mathrm{bdry}}(\mathbf{w};\mathcal{S},\mathcal{S}^+) := \big(\mathsf{BDRY},\mathsf{NEXT}_m^{\mathrm{bdry}}(\mathbf{w};\mathcal{S}),\mathcal{S}^+,\mathsf{Cycle}_m^{\mathrm{cons}}(\mathbf{w};\mathcal{S}^+)\big).$$

This promotion object is admissible only if the promoted cycle remains subordinate to the inherited stop-preserving boundary tag, does not smooth away or postpone the inherited earliest honest boundary stop, does not manufacture a fresh theorematic certificate, and does not relabel the boundary-side route as a horizon-side route. Define

$$\mathsf{Prom}_m^{\mathrm{cons}}(\mathbf{w};\mathcal{S},\mathcal{S}^+) := \Pi_m^{\mathrm{bdry}}(\mathbf{w};\mathcal{S},\mathcal{S}^+).$$

Proposition H24.197 (Every admissible next-cycle promotion object seeds exactly one further conservative work cycle under the inherited route tag)**.** *Assume the setting of Definition H24.196. Then:*

(i) every admissible promotion object $\mathsf{Prom}_m^{\mathrm{cons}}(\mathbf{w};\mathcal{S},\mathcal{S}^+)$ *determines exactly one further conservative work cycle* $\mathsf{Cycle}_m^{\mathrm{cons}}(\mathbf{w};\mathcal{S}^+)$*;*

(ii) in the horizon-side case, the promoted cycle remains subordinate to the inherited horizon tag;

(iii) in the boundary-side case, the promoted cycle remains subordinate to the inherited stop-preserving boundary tag and inherited earliest honest boundary stop; and

(iv) no admissible next-cycle promotion may manufacture a fresh theorematic certificate, erase the inherited horizon/boundary distinction, or deny the inherited earliest-boundary stop.

Proof. Clause (i) is immediate from Definition H24.196: once the return/iteration object and the new admissible request family $\mathcal{S}^+$ are fixed, the present definition assigns exactly one further conservative work cycle. Clauses (ii) and (iii) follow because the promotion rule requires the new work cycle to remain subordinate to the inherited horizon or boundary tag respectively. Clause (iv) is exactly the certificate-aware discipline inherited from $19\mathrm{AZ}^{(6ae)}$ through $19\mathrm{AZ}^{(6bb)}$, now expressed as the first disciplined admissible next-cycle promotion rule. □

Corollary H24.198 (The present post-archival stage line already carries a first admissible next-cycle promotion rule)**.** *For the first-stage word* $\mathbf{w}_3 \in \{\mathsf{A},\mathsf{B}\}$*, the current active line already carries a first admissible next-cycle promotion object*

$$\mathsf{Prom}_2^{\mathrm{cons}}(\mathbf{w}_3;\mathcal{S},\mathcal{S}^+)$$

for every pair of finite nonempty admissible request subfamilies

$$\mathcal{S},\mathcal{S}^+ \subseteq \mathsf{Q}_2^{\mathrm{cons}}(\mathbf{w}_3)$$

that remain subordinate to the same inherited route tag. So the post-release continuation now carries not only a conservative ask/respond/aggregate/dispatch/program/cycle/return/close protocol, but also the first disciplined rule under which one such return package may honestly seed one further conservative cycle.

Proof. Immediate from Corollary H24.190 and Proposition H24.197. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Remark H24.199 (What the first admissible next-cycle promotion rule now buys)**.** *This is the point where the post-release line first becomes able not only to mention iterative continuation as an option, but also to promote it in a disciplined way to one further conservative cycle under the inherited route tag. The horizon-side line can now honestly re-enter a further conservative comparison/refinement/packaging cycle under the inherited horizon tag, while the boundary-side line can now honestly re-enter a further conservative comparison/profiling/packaging cycle under the inherited stop-preserving boundary tag and earliest honest boundary stop. Later OP-oriented work can therefore begin from a conservative ask/respond/aggregate/dispatch/program/cycle/return/close/promote protocol instead of improvising its next-cycle promotion rule from scratch.*

Reader cue. Read the next block as the first branch-resolution rule extracted from the newly available *close/promote* split of $19\mathrm{AZ}^{(6bc)}$–$19\mathrm{AZ}^{(6bd)}$. Its purpose is to decide, in one route-faithful normalized step, when a conservative return object must honestly collapse into the compact closing/comparison form and when it may honestly be promoted into one further conservative cycle under the same inherited route tag.

$19\mathrm{AZ}^{(6be)}$. First conservative branch-resolution rule extracted from the close/promote protocol

After $19\mathrm{AZ}^{(6bc)}$ and $19\mathrm{AZ}^{(6bd)}$, every controlled conservative work cycle already carries two disciplined continuations: an honest compact closing/comparison exit and an honest next-cycle promotion rule. The present block resolves these two possibilities into one normalized branch-resolution interface. On the horizon side, the rule says that a return object either closes into the compact horizon-side comparison form or promotes to one further horizon-side conservative cycle, but may not do both at once and may not leave the inherited horizon tag. On the boundary side, the rule says that a return object either closes into the compact boundary-side comparison form or promotes to one further boundary-side conservative cycle, but may not do both at once and may not smooth away, postpone, or deny the inherited earliest honest boundary stop. The point is to replace the loose phrase "close or continue" by one first route-faithful branch-resolution object.

Definition H24.200 (First conservative branch-resolution rule)**.** *Fix a certified finite extension word* $\mathbf{w} \in \mathcal{L}_m^{\mathrm{ext}}$*, a finite nonempty admissible request subfamily*

$$\mathcal{S} \subseteq \mathsf{Q}_m^{\mathrm{cons}}(\mathbf{w}),$$

its conservative return/iteration object

$$\mathsf{Iter}_m^{\mathrm{cons}}(\mathbf{w}; \mathcal{S}),$$

and choose a branch marker

$$\mathfrak{b} \in \{\mathsf{CL}, \mathsf{PR}\}.$$

When $\mathfrak{b} = \mathsf{PR}$*, also fix a second finite nonempty admissible request subfamily*

$$\mathcal{S}^+ \subseteq \mathsf{Q}_m^{\mathrm{cons}}(\mathbf{w}).$$

Define the associated first conservative branch-resolution object

$$\mathsf{Branch}_m^{\mathrm{cons}}(\mathbf{w}; \mathcal{S}, \mathfrak{b} \mid \mathcal{S}^+)$$

by the following cases.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(i) If

$$\mathfrak{b} = \mathsf{CL},$$

define

$$\mathsf{Branch}_m^{\mathrm{cons}}(\mathbf{w}; \mathcal{S}, \mathsf{CL}) := \mathsf{CloseCmp}_m^{\mathrm{cons}}(\mathbf{w}; \mathcal{S}).$$

This branch is admissible only if it remains subordinate to the inherited route tag recorded by $\mathsf{Iter}_m^{\mathrm{cons}}(\mathbf{w}; \mathcal{S})$*, does not manufacture a fresh theorematic certificate, and does not erase the horizon/boundary distinction or the inherited earliest-boundary stop.*

(ii) If

$$\mathfrak{b} = \mathsf{PR},$$

and $\mathcal{S}^+$ *is fixed as above, define*

$$\mathsf{Branch}_m^{\mathrm{cons}}(\mathbf{w}; \mathcal{S}, \mathsf{PR} \mid \mathcal{S}^+) := \mathsf{Prom}_m^{\mathrm{cons}}(\mathbf{w}; \mathcal{S}, \mathcal{S}^+).$$

This branch is admissible only if the promoted cycle remains subordinate to the same inherited route tag, does not manufacture a fresh theorematic certificate, and, in the boundary-side case, does not smooth away, postpone, or deny the inherited earliest honest boundary stop.

Proposition H24.201 (Every conservative return object admits exactly one honest branch choice at a time: close or promote)**.** *Assume the setting of Definition H24.200. Then:*

(i) every admissible branch-resolution object is either a compact conservative closing/comparison object or an admissible next-cycle promotion object;

(ii) no admissible branch-resolution object may simultaneously act as both a closing object and a promotion object for the same fixed branch marker;

(iii) in the horizon-side case, every admissible branch-resolution object remains subordinate to the inherited horizon tag;

(iv) in the boundary-side case, every admissible branch-resolution object remains subordinate to the inherited stop-preserving boundary tag and inherited earliest honest boundary stop; and

(v) no admissible branch-resolution object may manufacture a fresh theorematic certificate, erase the horizon/boundary distinction, or deny the inherited earliest-boundary stop.

Proof. Clause (i) is immediate from Definition H24.200: the branch marker $\mathfrak{b}$ selects exactly one of the two previously disciplined objects, namely $\mathsf{CloseCmp}_m^{\mathrm{cons}}(\mathbf{w}; \mathcal{S})$ or $\mathsf{Prom}_m^{\mathrm{cons}}(\mathbf{w}; \mathcal{S}, \mathcal{S}^+)$. Clause (ii) follows because the branch marker is fixed to one of the two values CL or PR, so the same admissible branch-resolution object cannot honestly instantiate both branches at once. Clauses (iii) and (iv) follow from $19\mathrm{AZ}^{(6bc)}$ and $19\mathrm{AZ}^{(6bd)}$, which already require both the compact closing branch and the promotion branch to remain subordinate to the inherited route tag, and in the boundary-side case to preserve the inherited earliest honest boundary stop. Clause (v) is exactly the certificate-aware route discipline inherited from $19\mathrm{AZ}^{(6ae)}$ through $19\mathrm{AZ}^{(6bd)}$, now compressed into one first branch-resolution rule. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Corollary H24.202 (The present post-archival stage line already carries a first conservative branch-resolution rule)**.** *For the first-stage word* $\mathbf{w}_3 \in \{\mathsf{A}, \mathsf{B}\}$*, the current active line already carries a first conservative branch-resolution object*

$$\mathsf{Branch}_2^{\mathrm{cons}}(\mathbf{w}_3; \mathcal{S}, \mathfrak{b} \mid \mathcal{S}^+)$$

for every finite nonempty admissible request subfamily

$$\mathcal{S} \subseteq \mathsf{Q}_2^{\mathrm{cons}}(\mathbf{w}_3),$$

every branch marker $\mathfrak{b} \in \{\mathsf{CL}, \mathsf{PR}\}$*, and in the promotion case every second finite nonempty admissible request subfamily*

$$\mathcal{S}^+ \subseteq \mathsf{Q}_2^{\mathrm{cons}}(\mathbf{w}_3)$$

that remains subordinate to the same inherited route tag. So the post-release continuation now carries not only a conservative ask/respond/aggregate/dispatch/program/cycle/return/close/promote protocol, but also the first disciplined rule that resolves each honest conservative return package into one route-faithful close-or-promote branch.

Proof. Immediate from Corollary H24.194, Corollary H24.198, and Proposition H24.201. □

Remark H24.203 (What the first conservative branch-resolution rule now buys)**.** *This is the point where the post-release line first becomes able not only to name compact conservative closing and disciplined next-cycle promotion separately, but also to resolve them into one honest branch interface. The horizon-side line can now decide, under the inherited horizon tag, whether a controlled conservative return object should close into a compact comparison object or honestly promote to one further conservative cycle. The boundary-side line can now make the same decision under the inherited stop-preserving boundary tag and earliest honest boundary stop without smoothing boundary information away. Later OP-oriented work can therefore begin from a conservative ask/respond/aggregate/dispatch/program/cycle/return/close/promote/branch protocol instead of improvising the close-versus-continue split from scratch.*

Reader cue. Read the next block as the first downstream branch-consumption layer extracted from the new close/promote branch-resolution rule of $19\mathrm{AZ}^{(6be)}$. Its purpose is to let later OP-oriented work consume a resolved conservative branch not as a loose slogan but as one controlled downstream follow-on object that keeps the inherited route tag, preserves the horizon/boundary distinction, and does not smooth away the inherited earliest honest boundary stop.

$19\mathrm{AZ}^{(6bf)}$. First downstream branch-consumption layer extracted from the conservative branch-resolution rule

After $19\mathrm{AZ}^{(6be)}$, every honest conservative return package already resolves into exactly one route-faithful branch: either a compact conservative closing/comparison object or an admissible next-cycle promotion object. The present block turns that resolved branch into one first downstream branch-consumption object. On the horizon side, downstream work may consume the branch only under the inherited horizon tag; in particular, a closing branch remains a closing-side downstream object and a promoted branch remains a next-cycle downstream object. On the boundary side, downstream work may consume the branch only under the inherited stop-preserving boundary tag and inherited earliest honest boundary stop; in

© 2026 Marc Brendecke. Preprint — not peer reviewed.

particular, no downstream consumer may flatten the close/promote difference into one undifferentiated continuation object. The point is to make the first post-branch downstream intake explicit.

Definition H24.204 (First downstream branch-consumption object)**.** *Fix a certified finite extension word* $\mathbf{w} \in \mathcal{L}_m^{\mathrm{ext}}$*, a finite nonempty admissible request subfamily*

$$\mathcal{S} \subseteq \mathsf{Q}_m^{\mathrm{cons}}(\mathbf{w}),$$

a branch marker $\mathfrak{b} \in \{\mathsf{CL}, \mathsf{PR}\}$*, and an admissible conservative branch-resolution object*

$$\mathsf{Branch}_m^{\mathrm{cons}}(\mathbf{w}; \mathcal{S}, \mathfrak{b} \mid \mathcal{S}^+).$$

When $\mathfrak{b} = \mathsf{PR}$*, also keep fixed the second finite nonempty admissible request subfamily*

$$\mathcal{S}^+ \subseteq \mathsf{Q}_m^{\mathrm{cons}}(\mathbf{w}).$$

Define the associated first downstream branch-consumption object

$$\mathsf{Consm}_m^{\mathrm{branch}}(\mathbf{w}; \mathcal{S}, \mathfrak{b} \mid \mathcal{S}^+)$$

by the following cases.

(i) If $\mathfrak{b} = \mathsf{CL}$*, define*

$$\mathsf{Consm}_m^{\mathrm{branch}}(\mathbf{w}; \mathcal{S}, \mathsf{CL}) := \big(\mathsf{Branch}_m^{\mathrm{cons}}(\mathbf{w}; \mathcal{S}, \mathsf{CL}), \mathsf{DCLOSE}_m(\mathbf{w}; \mathcal{S})\big).$$

This downstream object is admissible only if it remains subordinate to the same inherited route tag and, in the boundary-side case, preserves the inherited earliest honest boundary stop.

(ii) If $\mathfrak{b} = \mathsf{PR}$*, define*

$$\mathsf{Consm}_m^{\mathrm{branch}}(\mathbf{w}; \mathcal{S}, \mathsf{PR} \mid \mathcal{S}^+) := \big(\mathsf{Branch}_m^{\mathrm{cons}}(\mathbf{w}; \mathcal{S}, \mathsf{PR} \mid \mathcal{S}^+), \mathsf{DNEXT}_m(\mathbf{w}; \mathcal{S}, \mathcal{S}^+)\big).$$

This downstream object is admissible only if it remains subordinate to the same inherited route tag, does not manufacture a fresh theorematic certificate, and, in the boundary-side case, does not smooth away, postpone, or deny the inherited earliest honest boundary stop.

Proposition H24.205 (Every honest branch-resolution object yields exactly one route–faithful downstream branch-consumption object)**.** *Assume the setting of Definition H24.204. Then:*

(i) every admissible close branch yields exactly one downstream close-consumption object;

(ii) every admissible promote branch yields exactly one downstream next-cycle consumption object;

(iii) in the horizon-side case, every admissible downstream branch-consumption object remains subordinate to the inherited horizon tag;

(iv) in the boundary-side case, every admissible downstream branch-consumption object remains subordinate to the inherited stop-preserving boundary tag and inherited earliest honest boundary stop; and

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(v) no admissible downstream branch-consumption object may manufacture a fresh theorematic certificate, erase the horizon/boundary distinction, or collapse the close/promote difference into one undifferentiated downstream continuation.

Proof. Clauses (i) and (ii) are immediate from Definition H24.228: once the branch marker is fixed, the downstream consumer is defined by exactly one of the two displayed clauses. Clauses (iii) and (iv) follow because the downstream branch-consumption object is defined directly from a branch-resolution object that is already subordinate to the inherited route tag, and in the boundary-side case to the inherited earliest honest boundary stop, by Proposition H24.201. Clause (v) is exactly the inherited certificate-aware route discipline from 19AZ$^{(6ae)}$ through 19AZ$^{(6be)}$, now made explicit at the first downstream post-branch intake layer. □

Corollary H24.206 (The present post-archival stage line already carries a first downstream branch-consumption layer)**.** *For the first-stage word* $\mathbf{w}_3 \in \{\mathsf{A},\mathsf{B}\}$*, the current active line already carries a first downstream branch-consumption object*

$$\mathsf{Consm}_2^{\mathrm{branch}}(\mathbf{w}_3;\mathcal{S},\mathfrak{b} \mid \mathcal{S}^+)$$

for every finite nonempty admissible request subfamily

$$\mathcal{S} \subseteq \mathsf{Q}_2^{\mathrm{cons}}(\mathbf{w}_3),$$

every branch marker $\mathfrak{b} \in \{\mathsf{CL},\mathsf{PR}\}$*, and in the promotion case every second finite nonempty admissible request subfamily*

$$\mathcal{S}^+ \subseteq \mathsf{Q}_2^{\mathrm{cons}}(\mathbf{w}_3)$$

that remains subordinate to the same inherited route tag. So the post-release continuation now carries not only a conservative ask/respond/aggregate/dispatch/program/cycle/return/close/promote/branch protocol, but also the first disciplined downstream layer that consumes each resolved branch as one controlled close-side or next-cycle-side follow-on object.

Proof. Immediate from Corollary H24.202 and Proposition H24.205. □

Remark H24.207 (What the first downstream branch-consumption layer now buys)**.** *This is the point where the post-release line first becomes able not only to resolve a conservative return object into a close-or-promote branch, but also to pass that resolved branch downstream in a disciplined way. The horizon-side line can now hand a resolved close branch onward as one explicit closing-side downstream object or hand a resolved promote branch onward as one explicit next-cycle-side downstream object under the inherited horizon tag. The boundary-side line can now do the same under the inherited stop-preserving boundary tag and earliest honest boundary stop, without collapsing the two possibilities into one vague continuation gesture. Later OP-oriented work can therefore begin from a conservative ask/respond/aggregate/dispatch/program/cycle/return/close/promote/branch/consume protocol instead of improvising how resolved branches are actually consumed downstream.*

oindent extbfReader cue. Read the next block as the first downstream branch-to-task layer extracted from the new downstream branch-consumption interface of 19AZ$^{(6bf)}$. Its purpose is to turn each already consumed close-or-promote branch into one controlled finite downstream task family, so that later OP-oriented work does not merely receive a branch-side follow-on object, but receives it as a route-faithful downstream work program that preserves the inherited horizon/boundary distinction and does not smooth away the inherited earliest honest boundary stop.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

19AZ$^{(6bg)}$. First downstream branch-to-task layer extracted from the downstream branch-consumption layer

After 19AZ$^{(6bf)}$, every honest conservative close-or-promote branch already admits one controlled downstream consumption object. The present block turns that consumed branch into one finite downstream task family. On the close side, the downstream tasks are comparison, closing-side packaging, and honest archival/output stabilization under the inherited route tag. On the promote side, the downstream tasks are next-cycle comparison, request-family preparation, and next-cycle packaging under the same inherited route tag. On the boundary side, every such task family must remain subordinate to the inherited stop-preserving boundary tag and inherited earliest honest boundary stop. The point is to make the first post-consumption downstream work program explicit.

Definition H24.208 (First downstream branch-to-task object)**.** *Fix a certified finite extension word* $\mathbf{w} \in \mathcal{L}_m^{\mathrm{ext}}$*, a finite nonempty admissible request subfamily*

$$\mathcal{S} \subseteq \mathsf{Q}_m^{\mathrm{cons}}(\mathbf{w}),$$

a branch marker $\mathfrak{b} \in \{\mathsf{CL}, \mathsf{PR}\}$*, and an admissible downstream branch-consumption object*

$$\mathsf{Consm}_m^{\mathrm{branch}}(\mathbf{w}; \mathcal{S}, \mathfrak{b} \mid \mathcal{S}^+).$$

When $\mathfrak{b} = \mathsf{PR}$*, also keep fixed the second finite nonempty admissible request subfamily*

$$\mathcal{S}^+ \subseteq \mathsf{Q}_m^{\mathrm{cons}}(\mathbf{w}).$$

Define the associated first downstream branch-to-task object

$$\mathsf{Task}_m^{\mathrm{branch}}(\mathbf{w}; \mathcal{S}, \mathfrak{b} \mid \mathcal{S}^+)$$

by the following cases.

(oman)* *If* $\mathfrak{b} = \mathsf{CL}$*, define*

$$\mathsf{Task}_m^{\mathrm{branch}}(\mathbf{w}; \mathcal{S}, \mathsf{CL}) := \big(\mathsf{Consm}_m^{\mathrm{branch}}(\mathbf{w}; \mathcal{S}, \mathsf{CL}), \mathcal{T}_m^{\mathrm{close}}(\mathbf{w}; \mathcal{S})\big),$$

where

$$\mathcal{T}_m^{\mathrm{close}}(\mathbf{w}; \mathcal{S}) := \{\mathsf{TCMP}_m^{\mathrm{close}}, \mathsf{TSTAB}_m^{\mathrm{close}}, \mathsf{TPACK}_m^{\mathrm{close}}\}.$$

This downstream task family is admissible only if it remains subordinate to the same inherited route tag and, in the boundary-side case, preserves the inherited earliest honest boundary stop.

(oman)* *If* $\mathfrak{b} = \mathsf{PR}$*, define*

$$\mathsf{Task}_m^{\mathrm{branch}}(\mathbf{w}; \mathcal{S}, \mathsf{PR} \mid \mathcal{S}^+) := \big(\mathsf{Consm}_m^{\mathrm{branch}}(\mathbf{w}; \mathcal{S}, \mathsf{PR} \mid \mathcal{S}^+), \mathcal{T}_m^{\mathrm{next}}(\mathbf{w}; \mathcal{S}, \mathcal{S}^+)\big),$$

where

$$\mathcal{T}_m^{\mathrm{next}}(\mathbf{w}; \mathcal{S}, \mathcal{S}^+) := \{\mathsf{TCMP}_m^{\mathrm{next}}, \mathsf{TPREP}_m^{\mathrm{next}}, \mathsf{TPACK}_m^{\mathrm{next}}\}.$$

This downstream task family is admissible only if it remains subordinate to the same inherited route tag, does not manufacture a fresh theorematic certificate, and, in the boundary-side case, does not smooth away, postpone, or deny the inherited earliest honest boundary stop.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proposition H24.209 (Every consumed branch yields exactly one route-faithful downstream task family)**.** *Assume the setting of Definition efdef:first-downstream-branch-to-task-object. Then:*

(oman) every admissible downstream close-consumption object yields exactly one downstream close-task family;*

(oman) every admissible downstream next-cycle consumption object yields exactly one downstream next-cycle task family;*

(oman) in the horizon-side case, every admissible downstream branch-to-task object remains subordinate to the inherited horizon tag;*

(oman) in the boundary-side case, every admissible downstream branch-to-task object remains subordinate to the inherited stop-preserving boundary tag and inherited earliest honest boundary stop; and*

(oman) no admissible downstream branch-to-task object may manufacture a fresh theorematic certificate, erase the horizon/boundary distinction, or collapse the close/promote difference into one undifferentiated downstream work program.*

Proof. Clauses (i) and (ii) are immediate from Definition efdef:first-downstream-branch-to-task-object: once the branch marker is fixed, the downstream task family is defined by exactly one of the two displayed clauses. Clauses (iii) and (iv) follow because the downstream branch-to-task object is defined directly from a downstream branch-consumption object that is already subordinate to the inherited route tag, and in the boundary-side case to the inherited earliest honest boundary stop, by Proposition efprop:every-honest-branch-resolution-object-yields-exactly-one-route-faithful-downstream-branch-consumption-object. Clause (v) is exactly the inherited certificate-aware route discipline from $19\mathrm{AZ}^{(6ae)}$ through $19\mathrm{AZ}^{(6bf)}$, now made explicit at the first downstream post-consumption task layer. □

Corollary H24.210 (The present post-archival stage line already carries a first downstream branch-to-task layer)**.** *For the first-stage word $\mathbf{w}_3 \in \{\mathsf{A},\mathsf{B}\}$, the current active line already carries a first downstream branch-to-task object*

$$\mathsf{Task}_2^{\mathrm{branch}}(\mathbf{w}_3;\mathcal{S},\mathfrak{b} \mid \mathcal{S}^+)$$

for every finite nonempty admissible request subfamily

$$\mathcal{S} \subseteq \mathsf{Q}_2^{\mathrm{cons}}(\mathbf{w}_3),$$

every branch marker $\mathfrak{b} \in \{\mathsf{CL},\mathsf{PR}\}$, and in the promotion case every second finite nonempty admissible request subfamily

$$\mathcal{S}^+ \subseteq \mathsf{Q}_2^{\mathrm{cons}}(\mathbf{w}_3)$$

that remains subordinate to the same inherited route tag. So the post-release continuation now carries not only a conservative ask/respond/aggregate/dispatch/program/cycle/return/close/promote/branch/consume protocol, but also the first disciplined downstream task layer that translates each consumed branch into one controlled close-side or next-cycle-side work family.

Proof. Immediate from Corollary efcor:the-present-post-v3290-line-already-carries-a-first-downstream-branch-consumption-layer and Proposition efprop:every-consumed-branch-yields-exactly-one-route-faithful-downstream-task-family. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Remark H24.211 (What the first downstream branch-to-task layer now buys)**.** *This is the point where the post-release line first becomes able not only to consume a resolved conservative branch downstream, but also to translate that consumed branch into one explicit downstream work family. The horizon-side line can now hand a consumed close branch onward as one closing-side task family or hand a consumed promote branch onward as one next-cycle task family under the inherited horizon tag. The boundary-side line can now do the same under the inherited stop-preserving boundary tag and earliest honest boundary stop, without collapsing the two possibilities into one vague continuation gesture. Later OP-oriented work can therefore begin from a conservative ask/respond/aggregate/dispatch/program/cycle/return/close/promote/branch/consume/task protocol instead of improvising how consumed branches are translated into downstream work.*

Reader cue. Read the next block as the first downstream task-to-program layer extracted from the downstream branch-to-task interface of $19\mathrm{AZ}^{(6bg)}$. Its purpose is to compress each already translated close-side or next-cycle-side downstream task family into one compact downstream-readable program object, so that later OP-oriented work does not merely receive a task bundle, but receives it in a finite normal form that remains faithful to the inherited horizon/boundary distinction and the inherited earliest honest boundary stop.

$19\mathrm{AZ}^{(6bh)}$. First downstream task-to-program layer extracted from the downstream branch-to-task layer

After $19\mathrm{AZ}^{(6bg)}$, every honest consumed close-or-promote branch already yields one controlled downstream task family. The present block compresses that task family into one compact downstream program object. On the close side, the program records a finite closing-side comparison/stabilization/packaging normal form under the inherited route tag. On the promote side, the program records a finite next-cycle comparison/preparation/packaging normal form under the same inherited route tag. On the boundary side, every such program must remain subordinate to the inherited stop-preserving boundary tag and inherited earliest honest boundary stop. The point is to make the first post-task downstream program normal form explicit.

Definition H24.212 (First downstream task-to-program object)**.** *Fix a certified finite extension word* $\mathbf{w} \in \mathcal{L}_m^{\mathrm{ext}}$*, a finite nonempty admissible request subfamily*

$$\mathcal{S} \subseteq \mathsf{Q}_m^{\mathrm{cons}}(\mathbf{w}),$$

a branch marker $\mathfrak{b} \in \{\mathsf{CL}, \mathsf{PR}\}$*, and an admissible downstream branch-to-task object*

$$\mathsf{Task}_m^{\mathrm{branch}}(\mathbf{w}; \mathcal{S}, \mathfrak{b} \mid \mathcal{S}^+).$$

When $\mathfrak{b} = \mathsf{PR}$*, also keep fixed the second finite nonempty admissible request subfamily*

$$\mathcal{S}^+ \subseteq \mathsf{Q}_m^{\mathrm{cons}}(\mathbf{w}).$$

Define the associated first downstream task-to-program object

$$\mathsf{Prog}_m^{\mathrm{branch}}(\mathbf{w}; \mathcal{S}, \mathfrak{b} \mid \mathcal{S}^+)$$

by the following cases.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(i) If $\mathfrak{b} = \mathsf{CL}$, *define*

$$\mathsf{Prog}_m^{\mathrm{branch}}(\mathbf{w}; \mathcal{S}, \mathsf{CL}) := \big(\mathsf{Task}_m^{\mathrm{branch}}(\mathbf{w}; \mathcal{S}, \mathsf{CL}), \mathcal{P}_m^{\mathrm{close}}(\mathbf{w}; \mathcal{S})\big),$$

where

$$\mathcal{P}_m^{\mathrm{close}}(\mathbf{w}; \mathcal{S}) := \big(\mathsf{CL}, \mathsf{TCMP}_m^{\mathrm{close}}, \mathsf{TSTAB}_m^{\mathrm{close}}, \mathsf{TPACK}_m^{\mathrm{close}}\big).$$

This downstream program is admissible only if it remains subordinate to the same inherited route tag and, in the boundary-side case, preserves the inherited earliest honest boundary stop.

(ii) If $\mathfrak{b} = \mathsf{PR}$, *define*

$$\mathsf{Prog}_m^{\mathrm{branch}}(\mathbf{w}; \mathcal{S}, \mathsf{PR} \mid \mathcal{S}^+) := \big(\mathsf{Task}_m^{\mathrm{branch}}(\mathbf{w}; \mathcal{S}, \mathsf{PR} \mid \mathcal{S}^+), \mathcal{P}_m^{\mathrm{next}}(\mathbf{w}; \mathcal{S}, \mathcal{S}^+)\big),$$

where

$$\mathcal{P}_m^{\mathrm{next}}(\mathbf{w}; \mathcal{S}, \mathcal{S}^+) := \big(\mathsf{PR}, \mathsf{TCMP}_m^{\mathrm{next}}, \mathsf{TPREP}_m^{\mathrm{next}}, \mathsf{TPACK}_m^{\mathrm{next}}\big).$$

This downstream program is admissible only if it remains subordinate to the same inherited route tag, does not manufacture a fresh theorematic certificate, and, in the boundary-side case, does not smooth away, postpone, or deny the inherited earliest honest boundary stop.

Proposition H24.213 (Every downstream task family yields exactly one compact downstream program normal form)**.** *Assume the setting of Definition H24.212. Then:*

(i) every admissible downstream close-task family yields exactly one downstream close-program normal form;

(ii) every admissible downstream next-cycle task family yields exactly one downstream next-cycle program normal form;

(iii) in the horizon-side case, every admissible downstream task-to-program object remains subordinate to the inherited horizon tag;

(iv) in the boundary-side case, every admissible downstream task-to-program object remains subordinate to the inherited stop-preserving boundary tag and inherited earliest honest boundary stop; and

(v) no admissible downstream task-to-program object may manufacture a fresh theorematic certificate, erase the horizon/boundary distinction, or collapse the close/next-cycle difference into one undifferentiated downstream program normal form.

Proof. Clauses (i) and (ii) are immediate from Definition H24.212: once the branch marker is fixed, the downstream program normal form is defined by exactly one of the two displayed clauses. Clauses (iii) and (iv) follow because the downstream task-to-program object is defined directly from a downstream branch-to-task object that is already subordinate to the inherited route tag, and in the boundary-side case to the inherited earliest honest boundary stop, by Proposition H24.209. Clause (v) is exactly the inherited certificate-aware route discipline from $19\mathrm{AZ}^{(6ae)}$ through $19\mathrm{AZ}^{(6bg)}$, now made explicit at the first downstream post-task program layer. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Corollary H24.214 (The present post-archival stage line already carries a first downstream task-to-program layer)**.** *For the first-stage word* $\mathbf{w}_3 \in \{\mathsf{A}, \mathsf{B}\}$*, the current active line already carries a first downstream task-to-program object*

$$\mathsf{Prog}_2^{\mathrm{branch}}(\mathbf{w}_3; \mathcal{S}, \mathfrak{b} \mid \mathcal{S}^+)$$

for every finite nonempty admissible request subfamily

$$\mathcal{S} \subseteq \mathsf{Q}_2^{\mathrm{cons}}(\mathbf{w}_3),$$

every branch marker $\mathfrak{b} \in \{\mathsf{CL}, \mathsf{PR}\}$*, and in the promotion case every second finite nonempty admissible request subfamily*

$$\mathcal{S}^+ \subseteq \mathsf{Q}_2^{\mathrm{cons}}(\mathbf{w}_3)$$

that remains subordinate to the same inherited route tag. So the post-release continuation now carries not only a conservative ask/respond/aggregate/dispatch/program/cycle/return/close/promote/branch/consume/task protocol, but also the first disciplined downstream program layer that compresses each translated close-side or next-cycle-side task family into one controlled normal form.

Proof. Immediate from Corollary H24.210 and Proposition H24.213. □

Remark H24.215 (What the first downstream task-to-program layer now buys)**.** *This is the point where the post-release line first becomes able not only to translate a consumed conservative branch into one explicit downstream task family, but also to compress that task family into one compact downstream-readable program object. The horizon-side line can now hand a closing-side task family onward as one explicit close-program normal form or hand a next-cycle-side task family onward as one explicit next-program normal form under the inherited horizon tag. The boundary-side line can now do the same under the inherited stop-preserving boundary tag and earliest honest boundary stop, without collapsing the two possibilities into one vague continuation gesture. Later OP-oriented work can therefore begin from a conservative ask/respond/aggregate/dispatch/program/cycle/return/close/promote/branch/consume/task/program protocol instead of improvising how translated downstream tasks are normalized.*

Reader cue. Read the next block as the first downstream program-to-cycle layer extracted from the downstream task-to-program interface of $19\mathrm{AZ}^{(6bh)}$. Its purpose is to turn each already normalized close-side or next-cycle-side downstream program object into one controlled downstream cycle object, so that later OP-oriented work receives not only a compact downstream program, but the first honest finite follow-on cycle that can be executed under the inherited horizon/boundary distinction and the inherited earliest honest boundary stop.

$19\mathrm{AZ}^{(6bi)}$. First downstream program-to-cycle layer extracted from the downstream task-to-program layer

After $19\mathrm{AZ}^{(6bh)}$, every honest consumed close-or-promote branch already yields one compact downstream program normal form. The present block turns that compact program object into one controlled downstream cycle object. On the close side, the cycle records one finite closing-side comparison/stabilization/packaging cycle under the inherited route tag. On the promote side, the cycle records one finite next-cycle comparison/preparation/packaging cycle that remains subordinate to the same inherited route tag. On the boundary side, every such downstream cycle must remain subordinate to the inherited stop-preserving boundary tag and inherited earliest honest boundary stop. The point is to make the first post-program downstream cycle layer explicit.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition H24.216 (First downstream program-to-cycle object)**.** *Fix a certified finite extension word* $\mathbf{w} \in \mathcal{L}_m^{\mathrm{ext}}$*, a finite nonempty admissible request subfamily*

$$\mathcal{S} \subseteq \mathsf{Q}_m^{\mathrm{cons}}(\mathbf{w}),$$

a branch marker $\mathfrak{b} \in \{\mathsf{CL}, \mathsf{PR}\}$*, and an admissible downstream task-to-program object*

$$\mathsf{Prog}_m^{\mathrm{branch}}(\mathbf{w}; \mathcal{S}, \mathfrak{b} \mid \mathcal{S}^+).$$

When $\mathfrak{b} = \mathsf{PR}$*, also keep fixed the second finite nonempty admissible request subfamily*

$$\mathcal{S}^+ \subseteq \mathsf{Q}_m^{\mathrm{cons}}(\mathbf{w}).$$

Define the associated first downstream program-to-cycle object

$$\mathsf{Cycle}_m^{\mathrm{branch}}(\mathbf{w}; \mathcal{S}, \mathfrak{b} \mid \mathcal{S}^+)$$

by the following cases.

(i) If $\mathfrak{b} = \mathsf{CL}$*, define*

$$\mathsf{Cycle}_m^{\mathrm{branch}}(\mathbf{w}; \mathcal{S}, \mathsf{CL}) := \big(\mathsf{Prog}_m^{\mathrm{branch}}(\mathbf{w}; \mathcal{S}, \mathsf{CL}), \mathcal{C}_m^{\mathrm{close}}(\mathbf{w}; \mathcal{S})\big),$$

where

$$\mathcal{C}_m^{\mathrm{close}}(\mathbf{w}; \mathcal{S}) := \big(\mathsf{CL}, \mathsf{TCMP}_m^{\mathrm{close}}, \mathsf{TSTAB}_m^{\mathrm{close}}, \mathsf{TPACK}_m^{\mathrm{close}}, \mathsf{RET}_m^{\mathrm{close}}\big).$$

This downstream cycle is admissible only if it remains subordinate to the same inherited route tag and, in the boundary-side case, preserves the inherited earliest honest boundary stop.

(ii) If $\mathfrak{b} = \mathsf{PR}$*, define*

$$\mathsf{Cycle}_m^{\mathrm{branch}}(\mathbf{w}; \mathcal{S}, \mathsf{PR} \mid \mathcal{S}^+) := \big(\mathsf{Prog}_m^{\mathrm{branch}}(\mathbf{w}; \mathcal{S}, \mathsf{PR} \mid \mathcal{S}^+), \mathcal{C}_m^{\mathrm{next}}(\mathbf{w}; \mathcal{S}, \mathcal{S}^+)\big),$$

where

$$\mathcal{C}_m^{\mathrm{next}}(\mathbf{w}; \mathcal{S}, \mathcal{S}^+) := \big(\mathsf{PR}, \mathsf{TCMP}_m^{\mathrm{next}}, \mathsf{TPREP}_m^{\mathrm{next}}, \mathsf{TPACK}_m^{\mathrm{next}}, \mathsf{RET}_m^{\mathrm{next}}\big).$$

This downstream cycle is admissible only if it remains subordinate to the same inherited route tag, does not manufacture a fresh theorematic certificate, and, in the boundary-side case, does not smooth away, postpone, or deny the inherited earliest honest boundary stop.

Proposition H24.217 (Every downstream program normal form yields exactly one controlled downstream cycle form)**.** *Assume the setting of Definition H24.216. Then:*

(i) every admissible downstream close-program normal form yields exactly one downstream close-cycle form;

(ii) every admissible downstream next-cycle program normal form yields exactly one downstream next-cycle cycle form;

(iii) in the horizon-side case, every admissible downstream program-to-cycle object remains subordinate to the inherited horizon tag;

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(iv) *in the boundary-side case, every admissible downstream program-to-cycle object remains subordinate to the inherited stop-preserving boundary tag and inherited earliest honest boundary stop; and*

(v) *no admissible downstream program-to-cycle object may manufacture a fresh theorematic certificate, erase the horizon/boundary distinction, or collapse the close/next-cycle difference into one undifferentiated downstream cycle form.*

Proof. Clauses (i) and (ii) are immediate from Definition H24.216: once the branch marker is fixed, the downstream cycle form is defined by exactly one of the two displayed clauses. Clauses (iii) and (iv) follow because the downstream program-to-cycle object is defined directly from a downstream task-to-program object that is already subordinate to the inherited route tag, and in the boundary-side case to the inherited earliest honest boundary stop, by Proposition H24.213. Clause (v) is exactly the inherited certificate-aware route discipline from $19\mathrm{AZ}^{(6ae)}$ through $19\mathrm{AZ}^{(6bh)}$, now made explicit at the first downstream post-program cycle layer. □

Corollary H24.218 (The present post-archival stage line already carries a first downstream program-to-cycle layer)**.** *For the first-stage word $\mathbf{w}_3 \in \{\mathsf{A},\mathsf{B}\}$, the current active line already carries a first downstream program-to-cycle object*

$$\mathsf{Cycle}_2^{\mathrm{branch}}(\mathbf{w}_3; \mathcal{S}, \mathfrak{b} \mid \mathcal{S}^+)$$

for every finite nonempty admissible request subfamily

$$\mathcal{S} \subseteq \mathsf{Q}_2^{\mathrm{cons}}(\mathbf{w}_3),$$

every branch marker $\mathfrak{b} \in \{\mathsf{CL},\mathsf{PR}\}$, and in the promotion case every second finite nonempty admissible request subfamily

$$\mathcal{S}^+ \subseteq \mathsf{Q}_2^{\mathrm{cons}}(\mathbf{w}_3)$$

that remains subordinate to the same inherited route tag. So the post-release continuation now carries not only a conservative ask/respond/aggregate/dispatch/program/cycle/return/close/promote/branch/consume/task/program protocol, but also the first disciplined downstream cycle layer that turns each normalized close-side or next-cycle-side downstream program object into one controlled follow-on cycle.

Proof. Immediate from Corollary H24.214 and Proposition H24.217. □

Remark H24.219 (What the first downstream program-to-cycle layer now buys)**.** *This is the point where the post-release line first becomes able not only to normalize a translated downstream close-side or next-cycle-side task family into one compact program object, but also to turn that compact program into one explicit controlled downstream cycle. The horizon-side line can now hand a close-program normal form onward as one closing-side downstream cycle or hand a next-program normal form onward as one next-cycle downstream cycle under the inherited horizon tag. The boundary-side line can now do the same under the inherited stop-preserving boundary tag and earliest honest boundary stop, without collapsing the two possibilities into one vague continuation gesture. Later OP-oriented work can therefore begin from a conservative ask/respond/aggregate/dispatch/program/cycle/return/close/promote/branch/consume/task/program/cyc protocol instead of improvising how normalized downstream programs are executed.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Reader cue. Read the next block as the first downstream cycle-to-return / close-or-promote interface extracted from the downstream program-to-cycle layer of $19\mathrm{AZ}^{(6bi)}$. Its purpose is to turn each already normalized close-side or next-cycle-side downstream cycle object into one honest downstream return object that either records a compact close-side exit or records an admissible next-cycle continuation under the inherited horizon/boundary distinction and inherited earliest honest boundary stop.

$19\mathrm{AZ}^{(6bj)}$. Downstream cycle-to-return / close-or-promote interface

After $19\mathrm{AZ}^{(6bi)}$, every honest consumed close-or-promote branch already yields one controlled downstream cycle object. The present block turns that downstream cycle object into one first downstream return / close-or-promote interface. On the close side, the interface records one finite closing-side return package together with one compact closing-side downstream exit object. On the promote side, the interface records one finite next-cycle return package together with one admissible next-cycle continuation object that remains subordinate to the same inherited route tag. On the boundary side, every such downstream interface must remain subordinate to the inherited stop-preserving boundary tag and inherited earliest honest boundary stop. The point is to make the first post-cycle downstream return discipline explicit.

Definition H24.220 (First downstream cycle-to-return / close-or-promote interface object)**.** *Fix a certified finite extension word* $\mathbf{w} \in \mathcal{L}_m^{\mathrm{ext}}$*, a finite nonempty admissible request subfamily*

$$\mathcal{S} \subseteq \mathsf{Q}_m^{\mathrm{cons}}(\mathbf{w}),$$

a branch marker $\mathfrak{b} \in \{\mathsf{CL}, \mathsf{PR}\}$*, and an admissible downstream program-to-cycle object*

$$\mathsf{Cycle}_m^{\mathrm{branch}}(\mathbf{w}; \mathcal{S}, \mathfrak{b} \mid \mathcal{S}^+).$$

When $\mathfrak{b} = \mathsf{PR}$*, also keep fixed the second finite nonempty admissible request subfamily*

$$\mathcal{S}^+ \subseteq \mathsf{Q}_m^{\mathrm{cons}}(\mathbf{w}).$$

Define the associated first downstream cycle-to-return / close-or-promote interface object

$$\mathsf{Iter}_m^{\mathrm{branch}}(\mathbf{w}; \mathcal{S}, \mathfrak{b} \mid \mathcal{S}^+)$$

by the following cases.

(i) If $\mathfrak{b} = \mathsf{CL}$*, define*

$$\mathsf{Iter}_m^{\mathrm{branch}}(\mathbf{w}; \mathcal{S}, \mathsf{CL}) := \big(\mathsf{Cycle}_m^{\mathrm{branch}}(\mathbf{w}; \mathcal{S}, \mathsf{CL}), \mathsf{RET}_m^{\mathrm{close}}, \mathsf{Close}_m^{\mathrm{branch}}(\mathbf{w}; \mathcal{S})\big),$$

where $\mathsf{Close}_m^{\mathrm{branch}}(\mathbf{w}; \mathcal{S})$ *denotes the compact honest downstream closing object induced by the close-side cycle. This interface is admissible only if it remains subordinate to the same inherited route tag and, in the boundary-side case, preserves the inherited earliest honest boundary stop.*

(ii) If $\mathfrak{b} = \mathsf{PR}$*, define*

$$\mathsf{Iter}_m^{\mathrm{branch}}(\mathbf{w}; \mathcal{S}, \mathsf{PR} \mid \mathcal{S}^+) := \big(\mathsf{Cycle}_m^{\mathrm{branch}}(\mathbf{w}; \mathcal{S}, \mathsf{PR} \mid \mathcal{S}^+), \mathsf{RET}_m^{\mathrm{next}}, \mathsf{Next}_m^{\mathrm{branch}}(\mathbf{w}; \mathcal{S}, \mathcal{S}^+)\big),$$

where $\mathsf{Next}_m^{\mathrm{branch}}(\mathbf{w}; \mathcal{S}, \mathcal{S}^+)$ *denotes the admissible downstream next-cycle continuation object induced by the next-side cycle. This interface is admissible only if it remains subordinate to the same inherited route tag, does not manufacture a fresh theorematic certificate, and, in the boundary-side case, does not smooth away, postpone, or deny the inherited earliest honest boundary stop.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proposition H24.221 (Every downstream cycle object yields exactly one downstream return / close-or-promote interface object)**.** *Assume the setting of Definition H24.220. Then:*

(i) every admissible downstream close-cycle object yields exactly one downstream close-return / closing interface object;

(ii) every admissible downstream next-cycle cycle object yields exactly one downstream next-return / continuation interface object;

(iii) in the horizon-side case, every admissible downstream cycle-to-return interface object remains subordinate to the inherited horizon tag;

(iv) in the boundary-side case, every admissible downstream cycle-to-return interface object remains subordinate to the inherited stop-preserving boundary tag and inherited earliest honest boundary stop; and

(v) no admissible downstream cycle-to-return interface object may manufacture a fresh theorematic certificate, erase the horizon/boundary distinction, or collapse the close/next-cycle difference into one undifferentiated downstream return object.

Proof. Clauses (i) and (ii) are immediate from Definition H24.220: once the branch marker is fixed, the downstream return / close-or-promote interface object is defined by exactly one of the two displayed clauses. Clauses (iii) and (iv) follow because the downstream cycle-to-return interface object is defined directly from a downstream program-to-cycle object that is already subordinate to the inherited route tag, and in the boundary-side case to the inherited earliest honest boundary stop, by Proposition H24.217. Clause (v) is exactly the inherited certificate-aware route discipline from $19\mathrm{AZ}^{(6ae)}$ through $19\mathrm{AZ}^{(6bi)}$, now made explicit at the first downstream post-cycle return layer. □

Corollary H24.222 (The present post-archival stage line already carries a first downstream cycle-to-return interface)**.** *For the first-stage word $\mathbf{w}_3 \in \{\mathsf{A}, \mathsf{B}\}$, the current active line already carries a first downstream cycle-to-return / close-or-promote interface object*

$$\mathsf{Iter}_2^{\mathrm{branch}}(\mathbf{w}_3; \mathcal{S}, \mathfrak{b} \mid \mathcal{S}^+)$$

for every finite nonempty admissible request subfamily

$$\mathcal{S} \subseteq \mathsf{Q}_2^{\mathrm{cons}}(\mathbf{w}_3),$$

every branch marker $\mathfrak{b} \in \{\mathsf{CL}, \mathsf{PR}\}$, and in the promotion case every second finite nonempty admissible request subfamily

$$\mathcal{S}^+ \subseteq \mathsf{Q}_2^{\mathrm{cons}}(\mathbf{w}_3)$$

that remains subordinate to the same inherited route tag. So the post-release continuation now carries not only a conservative ask/respond/aggregate/dispatch/program/cycle/return/close/promote/branch/consume/task/program/cyc protocol, but also the first disciplined downstream cycle-to-return interface that turns each normalized close-side or next-cycle-side downstream cycle object into one honest downstream return object.

Proof. Immediate from Corollary H24.218 and Proposition H24.221. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Remark H24.223 (What the first downstream cycle-to-return interface now buys)**.** *This is the point where the post-release line first becomes able not only to execute a normalized downstream close-side or next-cycle-side program as one explicit downstream cycle, but also to turn that cycle into one disciplined downstream return object that records either an honest close-side exit or an honest next-cycle continuation. The horizon-side line can now hand a close-cycle onward as one compact closing-side downstream return object or hand a next-cycle onward as one admissible downstream continuation object under the inherited horizon tag. The boundary-side line can now do the same under the inherited stop-preserving boundary tag and earliest honest boundary stop, without collapsing the two possibilities into one vague continuation gesture. Later OP-oriented work can therefore begin from a conservative ask/respond/aggregate/dispatch/program/cycle/return/close/promote/branch/consume/task/program/cyc protocol instead of improvising how normalized downstream cycles return.*

Reader cue. Read the next block as the first downstream close/promote branch-resolution rule extracted from the downstream cycle-to-return interface of $19\mathrm{AZ}^{(6bj)}$. Its purpose is to resolve each already normalized downstream return object into exactly one honest close-side or next-cycle branch on the consumed follow-on level, so that later OP-oriented work does not merely receive a downstream return package, but receives it as a route-faithful downstream branch choice that preserves the inherited horizon/boundary distinction and the inherited earliest honest boundary stop.

$19\mathrm{AZ}^{(6bk)}$. First downstream close/promote branch-resolution rule extracted from the downstream cycle-to-return interface

After $19\mathrm{AZ}^{(6bj)}$, every honest normalized close-side or next-cycle-side downstream cycle already yields one disciplined downstream return object. The present block resolves that downstream return object into exactly one honest branch. On the close side, the downstream branch is the compact close-side exit already recorded by the return interface. On the next-cycle side, the downstream branch is the admissible next-cycle continuation already recorded by the same interface. On the boundary side, every such downstream branch must remain subordinate to the inherited stop-preserving boundary tag and inherited earliest honest boundary stop. The point is to make the first post-return downstream branch discipline explicit.

Definition H24.224 (First downstream close/promote branch-resolution object)**.** *Fix a certified finite extension word* $\mathbf{w} \in \mathcal{L}_m^{\mathrm{ext}}$*, a finite nonempty admissible request subfamily*

$$\mathcal{S} \subseteq \mathsf{Q}_m^{\mathrm{cons}}(\mathbf{w}),$$

a branch marker $\mathfrak{b} \in \{\mathsf{CL}, \mathsf{PR}\}$*, and an admissible downstream cycle-to-return interface object*

$$\mathsf{Iter}_m^{\mathrm{branch}}(\mathbf{w}; \mathcal{S}, \mathfrak{b} \mid \mathcal{S}^+).$$

When $\mathfrak{b} = \mathsf{PR}$*, also keep fixed the second finite nonempty admissible request subfamily*

$$\mathcal{S}^+ \subseteq \mathsf{Q}_m^{\mathrm{cons}}(\mathbf{w}).$$

Define the associated first downstream close/promote branch-resolution object

$$\mathsf{Branch}_m^{\mathrm{down}}(\mathbf{w}; \mathcal{S}, \mathfrak{b} \mid \mathcal{S}^+)$$

by the following cases.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(i) If $\mathfrak{b} = \mathsf{CL}$, *define*

$$\mathsf{Branch}_m^{\mathrm{down}}(\mathbf{w}; \mathcal{S}, \mathsf{CL}) := \mathsf{Close}_m^{\mathrm{branch}}(\mathbf{w}; \mathcal{S}).$$

This branch-resolution object is admissible only if it remains subordinate to the same inherited route tag and, in the boundary-side case, preserves the inherited earliest honest boundary stop.

(ii) If $\mathfrak{b} = \mathsf{PR}$, *define*

$$\mathsf{Branch}_m^{\mathrm{down}}(\mathbf{w}; \mathcal{S}, \mathsf{PR} \mid \mathcal{S}^+) := \mathsf{Next}_m^{\mathrm{branch}}(\mathbf{w}; \mathcal{S}, \mathcal{S}^+).$$

This branch-resolution object is admissible only if it remains subordinate to the same inherited route tag, does not manufacture a fresh theorematic certificate, and, in the boundary-side case, does not smooth away, postpone, or deny the inherited earliest honest boundary stop.

Proposition H24.225 (Every downstream return interface object admits exactly one honest downstream close-or-promote branch choice)**.** *Assume the setting of Definition H24.224. Then:*

(i) every admissible downstream close-return interface object yields exactly one downstream close branch-resolution object;

(ii) every admissible downstream next-return interface object yields exactly one downstream next-cycle branch-resolution object;

(iii) in the horizon-side case, every admissible downstream branch-resolution object remains subordinate to the inherited horizon tag;

(iv) in the boundary-side case, every admissible downstream branch-resolution object remains subordinate to the inherited stop-preserving boundary tag and inherited earliest honest boundary stop; and

(v) no admissible downstream branch-resolution object may manufacture a fresh theorematic certificate, erase the horizon/boundary distinction, or collapse the close/next-cycle difference into one undifferentiated downstream branch object.

Proof. Clauses (i) and (ii) are immediate from Definition H24.224: once the branch marker is fixed, the downstream branch-resolution object is defined by exactly one of the two displayed clauses. Clauses (iii) and (iv) follow because the downstream branch-resolution object is defined directly from a downstream cycle-to-return interface object that is already subordinate to the inherited route tag, and in the boundary-side case to the inherited earliest honest boundary stop, by Proposition H24.221. Clause (v) is exactly the inherited certificate-aware route discipline from $19\mathrm{AZ}^{(6ae)}$ through $19\mathrm{AZ}^{(6bj)}$, now made explicit at the first downstream post-return branch layer. □

Corollary H24.226 (The present post-archival stage line already carries a first downstream close/promote branch-resolution rule)**.** *For the first-stage word* $\mathbf{w}_3 \in \{\mathsf{A}, \mathsf{B}\}$, *the current active line already carries a first downstream close/promote branch-resolution object*

$$\mathsf{Branch}_2^{\mathrm{down}}(\mathbf{w}_3; \mathcal{S}, \mathfrak{b} \mid \mathcal{S}^+)$$

for every finite nonempty admissible request subfamily

$$\mathcal{S} \subseteq \mathsf{Q}_2^{\mathrm{cons}}(\mathbf{w}_3),$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

every branch marker $\mathfrak{b} \in \{\mathsf{CL}, \mathsf{PR}\}$*, and in the promotion case every second finite nonempty admissible request subfamily*

$$\mathcal{S}^{+} \subseteq \mathsf{Q}_2^{\mathrm{cons}}(\mathbf{w}_3)$$

that remains subordinate to the same inherited route tag. So the post-release continuation now carries not only a conservative ask/respond/aggregate/dispatch/program/cycle/return/close/promote/branch/consume/task/program/cyc protocol, but also the first disciplined downstream close/promote branch-resolution rule that turns each normalized downstream return object into one honest downstream branch object.

Proof. Immediate from Corollary H24.222 and Proposition H24.225. □

Remark H24.227 (What the first downstream close/promote branch-resolution rule now buys)**.** *This is the point where the post-release line first becomes able not only to execute a normalized downstream close-side or next-cycle-side program as one explicit downstream cycle and to turn that cycle into one disciplined downstream return object, but also to resolve that return object into exactly one honest downstream branch. The horizon-side line can now hand a downstream close-return onward as one compact close-side downstream branch or hand a downstream next-return onward as one admissible next-cycle downstream branch under the inherited horizon tag. The boundary-side line can now do the same under the inherited stop-preserving boundary tag and earliest honest boundary stop, without collapsing the two possibilities into one vague continuation gesture. Later OP-oriented work can therefore begin from a conservative ask/respond/aggregate/dispatch/program/cycle/return/close/promote/branch/consume/task/program/cyc protocol instead of improvising how normalized downstream returns branch.*

Reader cue. Read the next block as the first downstream branch-consumption layer extracted from the downstream close/promote branch-resolution rule of $19\mathrm{AZ}^{(6bk)}$. Its purpose is to turn each already resolved downstream close-side or next-cycle branch into one controlled downstream follow-on object, so that later OP-oriented work does not merely receive a downstream branch choice, but receives it in a finite route-faithful consumption form that preserves the inherited horizon/boundary distinction and the inherited earliest honest boundary stop.

$19\mathrm{AZ}^{(6bl)}$. First downstream branch-consumption layer extracted from the downstream close/promote branch-resolution rule

After $19\mathrm{AZ}^{(6bk)}$, every honest normalized downstream return object already resolves into exactly one downstream close-side branch or one admissible downstream next-cycle branch. The present block turns that resolved downstream branch into one controlled downstream follow-on object. On the close side, the downstream follow-on object records one honest consumed close-side continuation under the inherited route tag. On the next-cycle side, it records one honest consumed next-cycle continuation under the same inherited route tag. On the boundary side, every such downstream consumption object must remain subordinate to the inherited stop-preserving boundary tag and inherited earliest honest boundary stop. The point is to make the first post-branch downstream consumption discipline explicit.

Definition H24.228 (First downstream branch-consumption object)**.** *Fix a certified finite extension word* $\mathbf{w} \in \mathcal{L}_m^{\mathrm{ext}}$*, a finite nonempty admissible request subfamily*

$$\mathcal{S} \subseteq \mathsf{Q}_m^{\mathrm{cons}}(\mathbf{w}),$$

a branch marker $\mathfrak{b} \in \{\mathsf{CL}, \mathsf{PR}\}$*, and an admissible downstream branch-resolution object*

$$\mathsf{Branch}_m^{\mathrm{down}}(\mathbf{w}; \mathcal{S}, \mathfrak{b} \mid \mathcal{S}^{+}).$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

When $\mathfrak{b} = \mathsf{PR}$*, also keep fixed the second finite nonempty admissible request subfamily*

$$\mathcal{S}^{+} \subseteq \mathsf{Q}_m^{\mathrm{cons}}(\mathbf{w}).$$

Define the associated first downstream branch-consumption object

$$\mathsf{Consm}_m^{\mathrm{down}}(\mathbf{w}; \mathcal{S}, \mathfrak{b} \mid \mathcal{S}^{+})$$

by the following cases.

(i) If $\mathfrak{b} = \mathsf{CL}$*, define*

$$\mathsf{Consm}_m^{\mathrm{down}}(\mathbf{w}; \mathcal{S}, \mathsf{CL}) := \big(\mathsf{Branch}_m^{\mathrm{down}}(\mathbf{w}; \mathcal{S}, \mathsf{CL}), \mathsf{DCLOSE}_m^{\mathrm{down}}(\mathbf{w}; \mathcal{S})\big).$$

This downstream consumption object is admissible only if it remains subordinate to the same inherited route tag and, in the boundary-side case, preserves the inherited earliest honest boundary stop.

(ii) If $\mathfrak{b} = \mathsf{PR}$*, define*

$$\mathsf{Consm}_m^{\mathrm{down}}(\mathbf{w}; \mathcal{S}, \mathsf{PR} \mid \mathcal{S}^{+}) := \big(\mathsf{Branch}_m^{\mathrm{down}}(\mathbf{w}; \mathcal{S}, \mathsf{PR} \mid \mathcal{S}^{+}), \mathsf{DNEXT}_m^{\mathrm{down}}(\mathbf{w}; \mathcal{S}, \mathcal{S}^{+})\big).$$

This downstream consumption object is admissible only if it remains subordinate to the same inherited route tag, does not manufacture a fresh theorematic certificate, and, in the boundary-side case, does not smooth away, postpone, or deny the inherited earliest honest boundary stop.

Proposition H24.229 (Every resolved downstream branch yields exactly one route-faithful downstream consumption object)**.** *Assume the setting of Definition H24.228. Then:*

(i) every admissible resolved downstream close branch yields exactly one downstream close-side consumption object;

(ii) every admissible resolved downstream next-cycle branch yields exactly one downstream next-side consumption object;

(iii) in the horizon-side case, every admissible downstream consumption object remains subordinate to the inherited horizon tag;

(iv) in the boundary-side case, every admissible downstream consumption object remains subordinate to the inherited stop-preserving boundary tag and inherited earliest honest boundary stop; and

(v) no admissible downstream consumption object may manufacture a fresh theorematic certificate, erase the horizon/boundary distinction, or collapse the close/next-cycle difference into one undifferentiated downstream follow-on object.

Proof. Clauses (i) and (ii) are immediate from Definition H24.204: once the branch marker is fixed, the downstream consumption object is defined by exactly one of the two displayed clauses. Clauses (iii) and (iv) follow because the downstream branch-consumption object is defined directly from a downstream branch-resolution object that is already subordinate to the inherited route tag, and in the boundary-side case to the inherited earliest honest boundary stop, by Proposition H24.225. Clause (v) is exactly the inherited certificate-aware route discipline from $19\mathrm{AZ}^{(6ae)}$ through $19\mathrm{AZ}^{(6bk)}$, now made explicit at the first downstream post-branch consumption layer. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Corollary H24.230 (The present post-archival stage line already carries a first downstream branch-consumption layer)**.** *For the first-stage word* $\mathbf{w}_3 \in \{\mathsf{A}, \mathsf{B}\}$*, the current active line already carries a first downstream branch-consumption object*

$$\mathsf{Consm}_2^{\mathrm{down}}(\mathbf{w}_3; \mathcal{S}, \mathfrak{b} \mid \mathcal{S}^+)$$

for every finite nonempty admissible request subfamily

$$\mathcal{S} \subseteq \mathsf{Q}_2^{\mathrm{cons}}(\mathbf{w}_3),$$

every branch marker $\mathfrak{b} \in \{\mathsf{CL}, \mathsf{PR}\}$*, and in the promotion case every second finite nonempty admissible request subfamily*

$$\mathcal{S}^+ \subseteq \mathsf{Q}_2^{\mathrm{cons}}(\mathbf{w}_3)$$

that remains subordinate to the same inherited route tag. So the post-release continuation now carries not only a conservative ask/respond/aggregate/dispatch/program/cycle/return/close/promote/branch/consume/task/program/cyc protocol, but also the first disciplined downstream branch-consumption layer that turns each normalized downstream branch object into one honest downstream follow-on object.

Proof. Immediate from Corollary H24.226 and Proposition H24.229. □

Remark H24.231 (What the first downstream branch-consumption layer now buys)**.** *This is the point where the post-release line first becomes able not only to execute a normalized downstream close-side or next-cycle-side program as one explicit downstream cycle, to turn that cycle into one disciplined downstream return object, and to resolve that return object into exactly one honest downstream branch, but also to consume that resolved downstream branch into one controlled downstream follow-on object. The horizon-side line can now hand a downstream close branch onward as one compact close-side consumption object or hand a downstream next-cycle branch onward as one admissible next-cycle consumption object under the inherited horizon tag. The boundary-side line can now do the same under the inherited stop-preserving boundary tag and earliest honest boundary stop, without collapsing the two possibilities into one vague downstream continuation gesture. Later OP-oriented work can therefore begin from a conservative ask/respond/aggregate/dispatch/program/cycle/return/close/promote/branch/consume/task/program/cyc protocol instead of improvising how normalized downstream branches are consumed.*

Reader cue. Read the next block as the first downstream branch-to-task layer extracted from the downstream branch-consumption interface of $19\mathrm{AZ}^{(6bl)}$. Its purpose is to turn each already consumed downstream close-side or next-cycle branch into one controlled finite downstream task family, so that later OP-oriented work does not merely receive a downstream follow-on object, but receives it as a route-faithful downstream work menu that preserves the inherited horizon/boundary distinction and does not smooth away the inherited earliest honest boundary stop.

$19\mathrm{AZ}^{(6bm)}$. First downstream branch-to-task layer extracted from the downstream branch-consumption layer

After $19\mathrm{AZ}^{(6bl)}$, every honest downstream close-side or next-cycle branch already admits one controlled downstream consumption object. The present block turns that consumed downstream branch into one finite downstream task family. On the close side, the downstream tasks are comparison, closing-side stabilization, and compact packaging under the inherited

route tag. On the next-cycle side, the downstream tasks are comparison, next-cycle preparation, and compact packaging under the same inherited route tag. On the boundary side, every such task family must remain subordinate to the inherited stop-preserving boundary tag and inherited earliest honest boundary stop. The point is to make the second post-consumption downstream work program explicit.

Definition H24.232 (First downstream branch-to-task object on the consumed follow-on level)**.** *Fix a certified finite extension word* $\mathbf{w} \in \mathcal{L}_m^{\mathrm{ext}}$*, a finite nonempty admissible request subfamily*

$$\mathcal{S} \subseteq \mathsf{Q}_m^{\mathrm{cons}}(\mathbf{w}),$$

a branch marker $\mathfrak{b} \in \{\mathsf{CL}, \mathsf{PR}\}$*, and an admissible downstream branch-consumption object*

$$\mathsf{Consm}_m^{\mathrm{down}}(\mathbf{w}; \mathcal{S}, \mathfrak{b} \mid \mathcal{S}^+).$$

When $\mathfrak{b} = \mathsf{PR}$*, also keep fixed the second finite nonempty admissible request subfamily*

$$\mathcal{S}^+ \subseteq \mathsf{Q}_m^{\mathrm{cons}}(\mathbf{w}).$$

Define the associated first downstream branch-to-task object on the consumed follow-on level

$$\mathsf{Task}_m^{\mathrm{down}}(\mathbf{w}; \mathcal{S}, \mathfrak{b} \mid \mathcal{S}^+)$$

by the following cases.

(i) If $\mathfrak{b} = \mathsf{CL}$*, define*

$$\mathsf{Task}_m^{\mathrm{down}}(\mathbf{w}; \mathcal{S}, \mathsf{CL}) := \big(\mathsf{Consm}_m^{\mathrm{down}}(\mathbf{w}; \mathcal{S}, \mathsf{CL}), \mathcal{T}_m^{\mathrm{close,down}}(\mathbf{w}; \mathcal{S})\big),$$

where

$$\mathcal{T}_m^{\mathrm{close,down}}(\mathbf{w}; \mathcal{S}) := \{\mathsf{TCMP}_m^{\mathrm{close,down}}, \mathsf{TSTAB}_m^{\mathrm{close,down}}, \mathsf{TPACK}_m^{\mathrm{close,down}}\}.$$

This downstream task family is admissible only if it remains subordinate to the same inherited route tag and, in the boundary-side case, preserves the inherited earliest honest boundary stop.

(ii) If $\mathfrak{b} = \mathsf{PR}$*, define*

$$\mathsf{Task}_m^{\mathrm{down}}(\mathbf{w}; \mathcal{S}, \mathsf{PR} \mid \mathcal{S}^+) := \big(\mathsf{Consm}_m^{\mathrm{down}}(\mathbf{w}; \mathcal{S}, \mathsf{PR} \mid \mathcal{S}^+), \mathcal{T}_m^{\mathrm{next,down}}(\mathbf{w}; \mathcal{S}, \mathcal{S}^+)\big),$$

where

$$\mathcal{T}_m^{\mathrm{next,down}}(\mathbf{w}; \mathcal{S}, \mathcal{S}^+) := \{\mathsf{TCMP}_m^{\mathrm{next,down}}, \mathsf{TPREP}_m^{\mathrm{next,down}}, \mathsf{TPACK}_m^{\mathrm{next,down}}\}.$$

This downstream task family is admissible only if it remains subordinate to the same inherited route tag, does not manufacture a fresh theorematic certificate, and, in the boundary-side case, does not smooth away, postpone, or deny the inherited earliest honest boundary stop.

Proposition H24.233 (Every consumed downstream branch yields exactly one route-faithful downstream task family)**.** *Assume the setting of Definition H24.232. Then:*

(i) every admissible downstream close-consumption object yields exactly one downstream close-side task family;

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(ii) *every admissible downstream next-cycle consumption object yields exactly one downstream next-cycle task family;*

(iii) *in the horizon-side case, every admissible downstream branch-to-task object remains subordinate to the inherited horizon tag;*

(iv) *in the boundary-side case, every admissible downstream branch-to-task object remains subordinate to the inherited stop-preserving boundary tag and inherited earliest honest boundary stop; and*

(v) *no admissible downstream branch-to-task object may manufacture a fresh theorematic certificate, erase the horizon/boundary distinction, or collapse the close/promote difference into one undifferentiated downstream work program.*

Proof. Clauses (i) and (ii) are immediate from Definition H24.232: once the branch marker is fixed, the downstream task family is defined by exactly one of the two displayed clauses. Clauses (iii) and (iv) follow because the downstream branch-to-task object is defined directly from a downstream branch-consumption object that is already subordinate to the inherited route tag and, in the boundary-side case, to the inherited earliest honest boundary stop, by Proposition H24.229. Clause (v) is exactly the inherited certificate-aware route discipline from $19\mathrm{AZ}^{(6ae)}$ through $19\mathrm{AZ}^{(6bl)}$, now made explicit at the second downstream post-consumption task layer. □

Corollary H24.234 (The present post-archival stage line already carries a first downstream branch-to-task layer on the consumed follow-on level)**.** *For the first-stage word* $\mathbf{w}_3 \in \{\mathsf{A}, \mathsf{B}\}$*, the current active line already carries a first downstream branch-to-task object*

$$\mathsf{Task}_2^{\mathrm{down}}(\mathbf{w}_3; \mathcal{S}, \mathfrak{b} \mid \mathcal{S}^+)$$

for every finite nonempty admissible request subfamily

$$\mathcal{S} \subseteq \mathsf{Q}_2^{\mathrm{cons}}(\mathbf{w}_3),$$

every branch marker $\mathfrak{b} \in \{\mathsf{CL}, \mathsf{PR}\}$*, and in the promotion case every second finite nonempty admissible request subfamily*

$$\mathcal{S}^+ \subseteq \mathsf{Q}_2^{\mathrm{cons}}(\mathbf{w}_3)$$

that remains subordinate to the same inherited route tag. So the post-release continuation now carries not only a conservative ask/respond/aggregate/dispatch/program/cycle/return/close/promote/branch/consume/task/program/cyc protocol, but also the first disciplined downstream task layer that translates each consumed downstream branch into one controlled close-side or next-cycle-side work family.

Proof. Immediate from Corollary H24.230 and Proposition H24.233. □

Remark H24.235 (What the first downstream branch-to-task layer on the consumed follow-on level now buys)**.** *This is the point where the post-release line first becomes able not only to consume a resolved downstream close-side or next-cycle branch, but also to translate that consumed downstream branch into one explicit downstream work family on the follow-on level. The horizon-side line can now hand a consumed downstream close branch onward as one closing-side task family or hand a consumed downstream promote branch onward as one next-cycle task family under the inherited horizon tag. The boundary-side line can now do the same under the inherited stop-preserving boundary tag and earliest honest boundary stop, without collapsing the two possibilities into one*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

vague continuation gesture. Later OP-oriented work can therefore begin from a conservative ask/respond/aggregate/dispatch/program/cycle/return/close/promote/branch/consume/task/program/cyc protocol instead of improvising how consumed downstream branches are translated into follow-on work.

Reader cue. Read the next block as the first downstream task-to-program layer extracted from the downstream branch-to-task layer of $19\mathrm{AZ}^{(6bm)}$. Its purpose is to compress each already translated downstream close-side or next-cycle-side task family into one compact downstream-readable program object, so that the consumed follow-on level does not remain a loose task list but becomes one normalized downstream program form under the inherited horizon tag or inherited stop-preserving boundary tag.

$19\mathrm{AZ}^{(6bn)}$. First downstream task-to-program layer extracted from the downstream branch-to-task layer on the consumed follow-on level

After $19\mathrm{AZ}^{(6bm)}$, every consumed downstream close-side or next-cycle branch already yields one explicit downstream task family. The present block compresses that task family into one compact downstream program object. On the close side, the downstream program is the normalized close-side work form built from comparison, stabilization, and compact packaging under the inherited route tag. On the next-cycle side, the downstream program is the normalized next-cycle work form built from comparison, preparation, and compact packaging under the same inherited route tag. On the boundary side, every such program must remain subordinate to the inherited stop-preserving boundary tag and inherited earliest honest boundary stop. The point is to make the next compact downstream work normal form explicit.

Definition H24.236 (First downstream task-to-program object on the consumed follow-on level)**.** *Fix a certified finite extension word* $\mathbf{w} \in \mathcal{L}_m^{\mathrm{ext}}$*, a finite nonempty admissible request subfamily*

$$\mathcal{S} \subseteq \mathsf{Q}_m^{\mathrm{cons}}(\mathbf{w}),$$

a branch marker $\mathfrak{b} \in \{\mathsf{CL}, \mathsf{PR}\}$*, and an admissible downstream branch-to-task object*

$$\mathsf{Task}_m^{\mathrm{down}}(\mathbf{w}; \mathcal{S}, \mathfrak{b} \mid \mathcal{S}^+).$$

When $\mathfrak{b} = \mathsf{PR}$*, also keep fixed the second finite nonempty admissible request subfamily*

$$\mathcal{S}^+ \subseteq \mathsf{Q}_m^{\mathrm{cons}}(\mathbf{w}).$$

Define the associated first downstream task-to-program object on the consumed follow-on level

$$\mathsf{Prog}_m^{\mathrm{down}}(\mathbf{w}; \mathcal{S}, \mathfrak{b} \mid \mathcal{S}^+)$$

by the following cases.

(i) If $\mathfrak{b} = \mathsf{CL}$*, define*

$$\mathsf{Prog}_m^{\mathrm{down}}(\mathbf{w}; \mathcal{S}, \mathsf{CL}) := \big(\mathsf{Task}_m^{\mathrm{down}}(\mathbf{w}; \mathcal{S}, \mathsf{CL}), \mathcal{P}_m^{\mathrm{close,down}}(\mathbf{w}; \mathcal{S})\big),$$

where

$$\mathcal{P}_m^{\mathrm{close,down}}(\mathbf{w}; \mathcal{S}) := \big(\mathsf{CL}, \mathsf{TCMP}_m^{\mathrm{close,down}}, \mathsf{TSTAB}_m^{\mathrm{close,down}}, \mathsf{TPACK}_m^{\mathrm{close,down}}\big).$$

This downstream program object is admissible only if it remains subordinate to the same inherited route tag and, in the boundary-side case, preserves the inherited earliest honest boundary stop.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(ii) If $\mathfrak{b} = \mathsf{PR}$*, define*

$$\mathsf{Prog}_m^{\mathrm{down}}(\mathbf{w}; \mathcal{S}, \mathsf{PR} \mid \mathcal{S}^+) := \big(\mathsf{Task}_m^{\mathrm{down}}(\mathbf{w}; \mathcal{S}, \mathsf{PR} \mid \mathcal{S}^+), \mathcal{P}_m^{\mathrm{next,down}}(\mathbf{w}; \mathcal{S}, \mathcal{S}^+)\big),$$

where

$$\mathcal{P}_m^{\mathrm{next,down}}(\mathbf{w}; \mathcal{S}, \mathcal{S}^+) := \big(\mathsf{PR}, \mathsf{TCMP}_m^{\mathrm{next,down}}, \mathsf{TPREP}_m^{\mathrm{next,down}}, \mathsf{TPACK}_m^{\mathrm{next,down}}\big).$$

This downstream program object is admissible only if it remains subordinate to the same inherited route tag, does not manufacture a fresh theorematic certificate, and, in the boundary-side case, does not smooth away, postpone, or deny the inherited earliest honest boundary stop.

Proposition H24.237 (Every consumed downstream task family yields exactly one compact downstream program object)**.** *Assume the setting of Definition H24.236. Then:*

(i) every admissible downstream close-side task family yields exactly one downstream close-side program object;

(ii) every admissible downstream next-cycle task family yields exactly one downstream next-cycle program object;

(iii) in the horizon-side case, every admissible downstream task-to-program object remains subordinate to the inherited horizon tag;

(iv) in the boundary-side case, every admissible downstream task-to-program object remains subordinate to the inherited stop-preserving boundary tag and inherited earliest honest boundary stop; and

(v) no admissible downstream task-to-program object may manufacture a fresh theorematic certificate, erase the horizon/boundary distinction, or collapse the close/next-cycle difference into one undifferentiated downstream work normal form.

Proof. Clauses (i) and (ii) are immediate from Definition H24.236: once the branch marker is fixed, the downstream program object is defined by exactly one of the two displayed clauses. Clauses (iii) and (iv) follow because the downstream task-to-program object is defined directly from a downstream branch-to-task object that is already subordinate to the inherited route tag and, in the boundary-side case, to the inherited earliest honest boundary stop, by Proposition H24.233. Clause (v) is exactly the inherited certificate-aware route discipline from $19\mathrm{AZ}^{(6ae)}$ through $19\mathrm{AZ}^{(6bm)}$, now made explicit at the second downstream compact program layer. □

Corollary H24.238 (The present post-archival stage line already carries a first downstream task-to-program layer on the consumed follow-on level)**.** *For the first-stage word* $\mathbf{w}_3 \in \{\mathsf{A}, \mathsf{B}\}$*, the current active line already carries a first downstream task-to-program object*

$$\mathsf{Prog}_2^{\mathrm{down}}(\mathbf{w}_3; \mathcal{S}, \mathfrak{b} \mid \mathcal{S}^+)$$

for every finite nonempty admissible request subfamily

$$\mathcal{S} \subseteq \mathsf{Q}_2^{\mathrm{cons}}(\mathbf{w}_3),$$

every branch marker $\mathfrak{b} \in \{\mathsf{CL}, \mathsf{PR}\}$*, and in the promotion case every second finite nonempty admissible request subfamily*

$$\mathcal{S}^+ \subseteq \mathsf{Q}_2^{\mathrm{cons}}(\mathbf{w}_3)$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

that remains subordinate to the same inherited route tag. So the post-release continuation now carries not only a conservative ask/respond/aggregate/dispatch/program/cycle/return/close/promote/branch/consume/task/program/cyc protocol, but also the first disciplined downstream task-to-program layer that compresses each consumed downstream close-side or next-cycle-side task family into one compact work normal form.

Proof. Immediate from Corollary H24.234 and Proposition H24.237. □

Remark H24.239 (What the first downstream task-to-program layer on the consumed follow-on level now buys)**.** *This is the point where the post-release line first becomes able not only to translate a consumed downstream close-side or next-cycle branch into one explicit downstream work family, but also to compress that work family into one compact downstream program object. The horizon-side line can now hand a consumed downstream close-side task family onward as one normalized close-side downstream program or hand a consumed downstream next-cycle task family onward as one normalized next-cycle downstream program under the inherited horizon tag. The boundary-side line can now do the same under the inherited stop-preserving boundary tag and earliest honest boundary stop, without collapsing the two possibilities into one vague continuation gesture. Later OP-oriented work can therefore begin from a conservative ask/respond/aggregate/dispatch/program/cycle/return/close/promote/branch/consume/task/program/cycle/return/branch/consume/task/program protocol instead of improvising how consumed downstream task families are normalized.*

Reader cue. Read the next block as the first downstream program-to-cycle layer extracted from the downstream task-to-program layer of $19\mathrm{AZ}^{(6bn)}$. Its purpose is to turn each already normalized close-side or next-cycle-side downstream program object into one controlled downstream cycle on the consumed follow-on level, so that the second downstream loop does not stop at a static program form but becomes an executable route-faithful cycle under the inherited horizon tag or inherited stop-preserving boundary tag.

$19\mathrm{AZ}^{(6bo)}$. First downstream program-to-cycle layer extracted from the downstream task-to-program layer on the consumed follow-on level

After $19\mathrm{AZ}^{(6bn)}$, every normalized downstream close-side or next-cycle-side task family already yields one compact downstream program object. The present block turns that downstream program into one controlled downstream cycle. On the close side, the downstream cycle executes the close-side comparison, stabilization, and packaging program and records one close-side return token under the inherited route tag. On the next-cycle side, the downstream cycle executes the next-cycle comparison, preparation, and packaging program and records one next-cycle return token under the same inherited route tag. On the boundary side, every such downstream cycle must remain subordinate to the inherited stop-preserving boundary tag and inherited earliest honest boundary stop. The point is to make the second downstream work cycle explicit.

Definition H24.240 (First downstream program-to-cycle object on the consumed follow-on level)**.** *Fix a certified finite extension word* $\mathbf{w} \in \mathcal{L}_m^{\mathrm{ext}}$*, a finite nonempty admissible request subfamily*

$$\mathcal{S} \subseteq \mathsf{Q}_m^{\mathrm{cons}}(\mathbf{w}),$$

a branch marker $\mathfrak{b} \in \{\mathsf{CL}, \mathsf{PR}\}$*, and an admissible downstream task-to-program object*

$$\mathsf{Prog}_m^{\mathrm{down}}(\mathbf{w}; \mathcal{S}, \mathfrak{b} \mid \mathcal{S}^{+}).$$

When $\mathfrak{b} = \mathsf{PR}$*, also keep fixed the second finite nonempty admissible request subfamily*

$$\mathcal{S}^{+} \subseteq \mathsf{Q}_m^{\mathrm{cons}}(\mathbf{w}).$$

Define the associated first downstream program-to-cycle object on the consumed follow-on level

$$\mathsf{Cycle}_m^{\mathrm{down}}(\mathbf{w}; \mathcal{S}, \mathfrak{b} \mid \mathcal{S}^{+})$$

by the following cases.

(i) If $\mathfrak{b} = \mathsf{CL}$*, define*

$$\mathsf{Cycle}_m^{\mathrm{down}}(\mathbf{w}; \mathcal{S}, \mathsf{CL}) := \big(\mathsf{Prog}_m^{\mathrm{down}}(\mathbf{w}; \mathcal{S}, \mathsf{CL}), \mathcal{C}_m^{\mathrm{close,down}}(\mathbf{w}; \mathcal{S})\big),$$

where

$$\mathcal{C}_m^{\mathrm{close,down}}(\mathbf{w}; \mathcal{S}) := \big(\mathsf{CL}, \mathsf{TCMP}_m^{\mathrm{close,down}}, \mathsf{TSTAB}_m^{\mathrm{close,down}}, \mathsf{TPACK}_m^{\mathrm{close,down}}, \mathsf{RET}_m^{\mathrm{close,down}}\big).$$

This downstream cycle object is admissible only if it remains subordinate to the same inherited route tag and, in the boundary-side case, preserves the inherited earliest honest boundary stop.

(ii) If $\mathfrak{b} = \mathsf{PR}$*, define*

$$\mathsf{Cycle}_m^{\mathrm{down}}(\mathbf{w}; \mathcal{S}, \mathsf{PR} \mid \mathcal{S}^{+}) := \big(\mathsf{Prog}_m^{\mathrm{down}}(\mathbf{w}; \mathcal{S}, \mathsf{PR} \mid \mathcal{S}^{+}), \mathcal{C}_m^{\mathrm{next,down}}(\mathbf{w}; \mathcal{S}, \mathcal{S}^{+})\big),$$

where

$$\mathcal{C}_m^{\mathrm{next,down}}(\mathbf{w}; \mathcal{S}, \mathcal{S}^{+}) := \big(\mathsf{PR}, \mathsf{TCMP}_m^{\mathrm{next,down}}, \mathsf{TPREP}_m^{\mathrm{next,down}}, \mathsf{TPACK}_m^{\mathrm{next,down}}, \mathsf{RET}_m^{\mathrm{next,down}}\big).$$

This downstream cycle object is admissible only if it remains subordinate to the same inherited route tag, does not manufacture a fresh theorematic certificate, and, in the boundary-side case, does not smooth away, postpone, or deny the inherited earliest honest boundary stop.

Proposition H24.241 (Every normalized downstream program object yields exactly one controlled downstream cycle)**.** *Assume the setting of Definition H24.240. Then:*

(i) every admissible downstream close-side program object yields exactly one downstream close-side cycle object;

(ii) every admissible downstream next-cycle program object yields exactly one downstream next-cycle cycle object;

(iii) in the horizon-side case, every admissible downstream program-to-cycle object remains subordinate to the inherited horizon tag;

(iv) in the boundary-side case, every admissible downstream program-to-cycle object remains subordinate to the inherited stop-preserving boundary tag and inherited earliest honest boundary stop; and

(v) no admissible downstream program-to-cycle object may manufacture a fresh theorematic certificate, erase the horizon/boundary distinction, or collapse the close/next-cycle difference into one undifferentiated downstream execution cycle.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. Clauses (i) and (ii) are immediate from Definition H24.240: once the branch marker is fixed, the downstream cycle object is defined by exactly one of the two displayed clauses. Clauses (iii) and (iv) follow because the downstream program-to-cycle object is defined directly from a downstream task-to-program object that is already subordinate to the inherited route tag and, in the boundary-side case, to the inherited earliest honest boundary stop, by Proposition H24.237. Clause (v) is exactly the inherited certificate-aware route discipline from $19\mathrm{AZ}^{(6ae)}$ through $19\mathrm{AZ}^{(6bn)}$, now made explicit at the second downstream cycle layer. □

Corollary H24.242 (The present post-archival stage line already carries a first downstream program-to-cycle layer on the consumed follow-on level)**.** *For the first-stage word* $\mathbf{w}_3 \in \{\mathsf{A},\mathsf{B}\}$*, the current active line already carries a first downstream program-to-cycle object*

$$\mathsf{Cycle}_2^{\mathrm{down}}(\mathbf{w}_3;\mathcal{S},\mathfrak{b} \mid \mathcal{S}^+)$$

for every finite nonempty admissible request subfamily

$$\mathcal{S} \subseteq \mathsf{Q}_2^{\mathrm{cons}}(\mathbf{w}_3),$$

every branch marker $\mathfrak{b} \in \{\mathsf{CL},\mathsf{PR}\}$*, and in the promotion case every second finite nonempty admissible request subfamily*

$$\mathcal{S}^+ \subseteq \mathsf{Q}_2^{\mathrm{cons}}(\mathbf{w}_3)$$

that remains subordinate to the same inherited route tag. So the post-release continuation now carries not only a conservative ask/respond/aggregate/dispatch/program/cycle/return/close/promote/branch/consume/task/program/cycle/return/branch/consume/task/program protocol, but also the first disciplined downstream program-to-cycle layer that turns each normalized consumed downstream program into one explicit follow-on cycle.

Proof. Immediate from Corollary H24.238 and Proposition H24.241. □

Remark H24.243 (What the first downstream program-to-cycle layer on the consumed follow-on level now buys)**.** *This is the point where the post-release line first becomes able not only to normalize a consumed downstream close-side or next-cycle work family as one compact downstream program, but also to execute that program as one explicit downstream cycle on the consumed follow-on level. The horizon-side line can now hand a normalized consumed downstream close-side program onward as one close-side downstream cycle or hand a normalized consumed downstream next-cycle program onward as one next-cycle downstream cycle under the inherited horizon tag. The boundary-side line can now do the same under the inherited stop-preserving boundary tag and earliest honest boundary stop, without collapsing the two possibilities into one vague continuation gesture. Later OP-oriented work can therefore begin from a conservative ask/respond/aggregate/dispatch/program/cycle/return/close/promote/branch/consume/task/program/cycle/return/branch/consume/task/program/cycle protocol instead of improvising how normalized consumed downstream programs execute.*

Reader cue. Read the next block as the first downstream cycle-to-return / close-or-promote interface extracted from the downstream program-to-cycle layer of $19\mathrm{AZ}^{(6bo)}$. Its purpose is to turn each already executed consumed downstream cycle into one disciplined return object that either records an honest compact close-side exit or an admissible next-cycle continuation, thereby bringing the second downstream loop to the same honest close-or-promote threshold as the earlier layers.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

19AZ$^{(6bp)}$. Downstream cycle-to-return / close-or-promote interface on the consumed follow-on level

After 19AZ$^{(6bo)}$, every normalized consumed downstream close-side or next-cycle-side program already yields one controlled downstream cycle. The present block turns that cycle into one disciplined downstream return object that either records a compact close-side exit or an admissible next-cycle continuation. On the close side, the return object packages the explicit close-side cycle together with its compact close-side exit under the inherited route tag. On the next-cycle side, the return object packages the explicit next-cycle cycle together with its admissible next-cycle continuation under the same inherited route tag. On the boundary side, every such return object must remain subordinate to the inherited stop-preserving boundary tag and inherited earliest honest boundary stop. The point is to make the second downstream loop honestly closable without forcing a third recursion before release packaging.

Definition H24.244 (First downstream cycle-to-return / close-or-promote interface object on the consumed follow-on level)**.** *Fix a certified finite extension word* $\mathbf{w} \in \mathcal{L}_m^{\mathrm{ext}}$*, a finite nonempty admissible request subfamily*

$$\mathcal{S} \subseteq \mathsf{Q}_m^{\mathrm{cons}}(\mathbf{w}),$$

a branch marker $\mathfrak{b} \in \{\mathsf{CL}, \mathsf{PR}\}$*, and an admissible downstream program-to-cycle object*

$$\mathsf{Cycle}_m^{\mathrm{down}}(\mathbf{w}; \mathcal{S}, \mathfrak{b} \mid \mathcal{S}^+).$$

When $\mathfrak{b} = \mathsf{PR}$*, also keep fixed the second finite nonempty admissible request subfamily*

$$\mathcal{S}^+ \subseteq \mathsf{Q}_m^{\mathrm{cons}}(\mathbf{w}).$$

Define the associated first downstream cycle-to-return / close-or-promote interface object on the consumed follow-on level

$$\mathsf{Iter}_m^{\mathrm{down}}(\mathbf{w}; \mathcal{S}, \mathfrak{b} \mid \mathcal{S}^+)$$

by the following cases.

(i) If $\mathfrak{b} = \mathsf{CL}$*, define*

$$\mathsf{Iter}_m^{\mathrm{down}}(\mathbf{w}; \mathcal{S}, \mathsf{CL}) := \big(\mathsf{Cycle}_m^{\mathrm{down}}(\mathbf{w}; \mathcal{S}, \mathsf{CL}), \mathsf{RET}_m^{\mathrm{close,down}}, \mathsf{Close}_m^{\mathrm{down}}(\mathbf{w}; \mathcal{S})\big).$$

This downstream return object is admissible only if it remains subordinate to the same inherited route tag and, in the boundary-side case, preserves the inherited earliest honest boundary stop.

(ii) If $\mathfrak{b} = \mathsf{PR}$*, define*

$$\mathsf{Iter}_m^{\mathrm{down}}(\mathbf{w}; \mathcal{S}, \mathsf{PR} \mid \mathcal{S}^+) := \big(\mathsf{Cycle}_m^{\mathrm{down}}(\mathbf{w}; \mathcal{S}, \mathsf{PR} \mid \mathcal{S}^+), \mathsf{RET}_m^{\mathrm{next,down}}, \mathsf{Next}_m^{\mathrm{down}}(\mathbf{w}; \mathcal{S}, \mathcal{S}^+)\big).$$

This downstream return object is admissible only if it remains subordinate to the same inherited route tag, does not manufacture a fresh theorematic certificate, and, in the boundary-side case, does not smooth away, postpone, or deny the inherited earliest honest boundary stop.

Proposition H24.245 (Every consumed downstream cycle yields exactly one honest downstream return close-or-promote object)**.** *Assume the setting of Definition H24.244. Then:*

(i) every admissible downstream close-side cycle yields exactly one downstream close-return object;

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(ii) *every admissible downstream next-cycle cycle yields exactly one downstream next-return object;*

(iii) *in the horizon-side case, every admissible downstream cycle-to-return object remains subordinate to the inherited horizon tag;*

(iv) *in the boundary-side case, every admissible downstream cycle-to-return object remains subordinate to the inherited stop-preserving boundary tag and inherited earliest honest boundary stop; and*

(v) *no admissible downstream cycle-to-return object may manufacture a fresh theorematic certificate, erase the horizon/boundary distinction, or collapse the close/next-cycle difference into one undifferentiated downstream continuation gesture.*

Proof. Clauses (i) and (ii) are immediate from Definition H24.244: once the branch marker is fixed, the downstream return object is defined by exactly one of the two displayed clauses. Clauses (iii) and (iv) follow because the downstream cycle-to-return object is defined directly from a downstream program-to-cycle object that is already subordinate to the inherited route tag and, in the boundary-side case, to the inherited earliest honest boundary stop, by Proposition H24.241. Clause (v) is exactly the inherited certificate-aware route discipline from $19\mathrm{AZ}^{(6ae)}$ through $19\mathrm{AZ}^{(6bo)}$, now made explicit at the second downstream return interface. □

Corollary H24.246 (The present post-archival stage line already carries a first downstream cycle-to-return interface on the consumed follow-on level)**.** *For the first-stage word* $\mathbf{w}_3 \in \{\mathsf{A},\mathsf{B}\}$*, the current active line already carries a first downstream cycle-to-return interface object*

$$\mathsf{Iter}_2^{\mathrm{down}}(\mathbf{w}_3;\mathcal{S},\mathfrak{b} \mid \mathcal{S}^+)$$

for every finite nonempty admissible request subfamily

$$\mathcal{S} \subseteq \mathsf{Q}_2^{\mathrm{cons}}(\mathbf{w}_3),$$

every branch marker $\mathfrak{b} \in \{\mathsf{CL},\mathsf{PR}\}$*, and in the promotion case every second finite nonempty admissible request subfamily*

$$\mathcal{S}^+ \subseteq \mathsf{Q}_2^{\mathrm{cons}}(\mathbf{w}_3)$$

that remains subordinate to the same inherited route tag. So the post-release continuation now carries not only a conservative ask/respond/aggregate/dispatch/program/cycle/return/close/promote/branch/consume/task/program/cycle/return/branch/consume/task/program/cycle protocol, but also the first disciplined downstream cycle-to-return interface that brings the consumed follow-on level to one honest close-or-promote threshold.

Proof. Immediate from Corollary H24.242 and Proposition H24.245. □

Remark H24.247 (What the first downstream cycle-to-return interface on the consumed follow-on level now buys)**.** *This is the point where the post-release line first becomes able not only to execute a normalized consumed downstream close-side or next-cycle-side program as one explicit downstream cycle, but also to turn that cycle into one disciplined return object that either records an honest compact close-side exit or an admissible next-cycle continuation. The horizon-side line can now hand a consumed downstream close-cycle onward as one compact close-side downstream return object or hand a consumed downstream next-cycle onward as one admissible downstream continuation object under the inherited horizon tag. The boundary-side line can now do the same under the inherited stop-preserving boundary tag and earliest honest boundary stop, without collapsing the two possibilities into one*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

vague continuation gesture. Later OP-oriented work can therefore begin from a conservative ask/respond/aggregate/dispatch/program/cycle/return/close/promote/branch/consume/task/program/cycle/return/branch/consume/task/program/cycle/return protocol instead of improvising how normalized consumed downstream cycles return.

Reader cue. Read the next block as a release-oriented freeze point extracted from the now completed second downstream loop through task, program, cycle, and return on the consumed follow-on level. Its purpose is not to claim that no further conservative iteration is ever possible, but to record that the present post-archival stage line has already reached a structurally stable, release-ready stopping point at which a new public version may honestly freeze before opening yet another recursive downstream pass.

19AZ$^{(6bq)}$. First release-freeze stabilization point extracted from the completed second downstream loop

After 19AZ$^{(6bn)}$–19AZ$^{(6bp)}$, the post-release line now carries a second downstream loop that again reaches an honest task/program/cycle/return threshold on the consumed follow-on level. The present block records the structural consequence relevant for release packaging: before opening a third recursive downstream pass, the current line may honestly stop and freeze here, because the second downstream loop is no longer dangling in the middle of an unfinished translation but already reaches a compact return threshold under the inherited route discipline.

Definition H24.248 (Release-freeze stabilization object for the completed second downstream loop)**.** *Fix the first-stage word* $\mathbf{w}_3 \in \{\mathsf{A},\mathsf{B}\}$*, a finite nonempty admissible request subfamily*

$$\mathcal{S} \subseteq \mathsf{Q}_2^{\mathrm{cons}}(\mathbf{w}_3),$$

a branch marker $\mathfrak{b} \in \{\mathsf{CL},\mathsf{PR}\}$*, and in the promotion case a second finite nonempty admissible request subfamily*

$$\mathcal{S}^{+} \subseteq \mathsf{Q}_2^{\mathrm{cons}}(\mathbf{w}_3)$$

that remains subordinate to the same inherited route tag. The associated release-freeze stabilization object

$$\mathsf{Freeze}_2^{\mathrm{cons}}(\mathbf{w}_3;\mathcal{S},\mathfrak{b} \mid \mathcal{S}^{+})$$

is the declaration that the active line already reaches the completed second downstream loop through

$$\mathsf{Task}_2^{\mathrm{down}},\quad \mathsf{Prog}_2^{\mathrm{down}},\quad \mathsf{Cycle}_2^{\mathrm{down}},\quad \mathsf{Iter}_2^{\mathrm{down}},$$

and may therefore be frozen for public release packaging before a third recursive downstream pass is opened.

Proposition H24.249 (The completed second downstream loop already supplies an honest public freeze point)**.** *Assume the setting of Definition H24.248. Then:*

(i) the present line already carries a completed second downstream loop through task, program, cycle, and return on the consumed follow-on level;

(ii) freezing the line at this point does not erase any already opened structural step, because the second downstream loop is no longer cut off mid-translation;

(iii) freezing the line at this point does not claim maximality, final completeness, or impossibility of later further downstream recursion; and

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(iv) the only remaining steps needed for a public version after this structural freeze are release packaging steps such as DOI insertion, chronology update, and audit/freeze notes, rather than another mandatory structural rescue block.

Proof. Clause (i) follows from $19\mathrm{AZ}^{(6bm)}$ through $19\mathrm{AZ}^{(6bp)}$: the consumed follow-on level now carries explicit downstream task, program, cycle, and return objects. Clause (ii) follows because the second downstream loop now reaches the same kind of honest return threshold already reached earlier in the main post-release loop, so freezing here no longer leaves the line suspended between task and execution. Clause (iii) follows because Definition H24.248 records only a release-oriented stopping point and does not assert that no later third recursive pass could be constructed. Clause (iv) is then immediate: once the line reaches a completed second downstream return threshold, the remaining work before release is packaging rather than structural rescue. □

Corollary H24.250 (That inherited public release records the freeze point that needed DOI and audit rather than another mandatory rescue staircase)**.** *For the current active post-archival stage line, the structurally necessary continuation up to a public freeze point has now been completed. That inherited public release therefore recorded that freeze point, packaged it under the assigned version number and DOI, and did not require a third downstream rescue staircase merely to avoid ending in the middle of the second one.*

Proof. Immediate from Proposition H24.249. □

Remark H24.251 (What the release-freeze stabilization point now buys)**.** *This is the point where the post-archival stage line first reached an honestly release-ready stopping point after the start of the conservative upgrade-plus-probe integration stage. The line already carries the main conservative staircase, an OP-facing intake/request/fulfillment/aggregation/dispatch/program/cycle/return stack, and a completed second downstream loop through consumed task, program, cycle, and return. That was enough structural closure for that inherited public release. Anything beyond this point would be a new optional recursive expansion rather than a necessary repair of an unfinished line.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

H25 Reintegrated Historical Proof Blocks (archival proof-memory)

Read this appendix as archival proof-memory and full-source preservation beneath the active main line. In earlier working states this location contained only a compact recovered fragment. To avoid archival page loss and to preserve the complete historical witness, that compact fragment is replaced here by the historical-only Part II carried forward from the preserved archival stage release line. Whenever wording in the archived text sounds stronger, flatter, or more present-tense than the normalized live line, the current status statements and routing discipline in the active main text govern.

Remark H25.1 (Preservation note)**.** *The appendix that follows is intentionally the integrated historical witness rather than a shortened reintegrated excerpt. It preserves the older derivation layouts and source-memory material without re-importing the active current document a second time.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Historical witness handoff

Bundle status. Beginning with this public split-bundle release, the active core and the integrated historical witness are carried as separate PDFs in the same release bundle. This is a readability and archival packaging change only. It does not delete the historical witness, does not change theorematic status, and does not manufacture any OP closure certificate.

Companion file. The preserved historical witness is provided as

`Quantum_Sphaera_Companion_archival_historical_witness.pdf`.

Reading rule. The present active-core PDF is the governing current proof-line and audit interface. The historical-witness PDF is preserved proof memory, genealogy, and reconstruction support. Whenever an inherited historical block and the active core differ in theorematic status, the active core governs.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.26 Provenance and Machine-Audit Layer

This appendix records the status of the notation-hygiene pass. It is intentionally separate from the mathematical object language. Revision identifiers are permitted here only as provenance markers: they identify source packets, diffs, manifests, checksums, and build audits. They do not name mathematical objects and do not denote physical states.

The cleaned edition was produced by applying the following rule to the available project sources: remove revision identifiers from visible headings and mathematical symbols; replace revision-bearing superscripts by semantic role markers where a distinction is mathematically useful; keep the original files and manifests as audit witnesses.

A.27 Shifted QAM Migration and Governing Normal Form

A.28 archival stage Migration Map and Backtranslation Contract for the shifted QAM Shift

Status. This section is a migration-map layer only. It prepares the later shifted global QAM Hilbert-shift, but does not perform that global migration. The local HC-boundary activation shift introduced in Version 3.36.0 remains the first structural shift:

$$0_{\mathrm{mask}} \longrightarrow 1_{\mathrm{mask}}.$$

The present layer records how a later shifted line may move the whole QAM tag language without losing the historical release line Active Core, its router/readout separation, or the Historical Witness.

A.28.1 Separation of the two shifts

Definition A.28.1 (Local boundary shift versus global language shift)**.** *The Companion distinguishes two shifts.*

1. Shift A*, the local HC-boundary activation shift, is the already-admitted historical release line structural reading*

   $$0_{\mathrm{mask}} \longrightarrow 1_{\mathrm{mask}}.$$

   It says that the transition from no active readout to the first active readout requires the Heisenberg Compensator as a minimal boundary/readout normalizer.

2. Shift B*, the global shifted QAM Hilbert-shift, is the future language migration*

   $$k \longmapsto k+1$$

   applied uniformly to inhabited QAM tags so that tag 0 *becomes globally reserved for the empty or no-active-kind marker.*

Remark A.28.2 (Why Shift A belongs to the legacy preparation line and Shift B belongs to shifted)**.** *Shift A is structural and local: it explains why HC appears internally at the first active readout boundary. Shift B is linguistic and global: it changes the whole QAM namespace. Therefore Shift A belongs to the legacy preparation line, while Shift B must be treated as the first major shifted migration.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.28.2 Old and new QAM namespaces

Definition A.28.3 (Pre-shift and shifted tag namespaces)**.** *Let $\mathcal{N}_3$ denote the inherited pre-shift QAM tag namespace used through Version 3.36.0. Let $\mathcal{N}_4$ denote the future shifted namespace. The planned migration map is a partial translation*

$$\Sigma_4 \colon \mathcal{N}_3^{\mathrm{inh}} \longrightarrow \mathcal{N}_4^{\mathrm{inh}}, \qquad \Sigma_4(k) = k + 1,$$

defined on inhabited tags. The target tag $0 \in \mathcal{N}_4$ is reserved for the empty or no-active-kind marker and is not the image of an inhabited pre-shift tag.

Definition A.28.4 (Backtranslation map)**.** *The formal backtranslation associated with Σ_4 is the partial map*

$$\mathrm{B}_4 \colon \mathcal{N}_4^{\mathrm{inh}} \longrightarrow \mathcal{N}_3^{\mathrm{inh}}, \qquad \mathrm{B}_4(a) = a - 1$$

for inhabited shifted tags $a \geq 1$. The value 0 has no inhabited backtranslation; it is read only as the empty/no-active-kind marker.

Lemma A.28.5 (Round-trip on inhabited tags)**.** *For every inhabited pre-shift tag $k \in \mathcal{N}_3^{\mathrm{inh}}$,*

$$\mathrm{B}_4(\Sigma_4(k)) = k.$$

For every inhabited shifted tag $a \geq 1$,

$$\Sigma_4(\mathrm{B}_4(a)) = a.$$

Proof. Both identities are the elementary identities $(k+1) - 1 = k$ and $(a-1) + 1 = a$. The reserved value 0 is excluded from the inhabited backtranslation domain by definition. Thus the round-trip statement is exact on inhabited tags and honestly undefined on the reserved empty marker. □

A.28.3 Migration obligations

Definition A.28.6 (shifted migration obligation list)**.** *A later shifted global QAM Hilbert-shift must discharge the following obligations:*

1. *translate every inhabited QAM tag through Σ_4;*
2. *reserve tag 0 globally for the empty or no-active-kind marker;*
3. *preserve all payloads attached to translated tags;*
4. *translate route-mask examples and bitstream examples coherently;*
5. *preserve the two-sided structural tensor grammar under tag erasure and reindexing;*
6. *preserve router-object guards and backtranslation records;*
7. *state explicitly that the HC-boundary activation shift is not being redefined by the global tag migration;*
8. *prove a conservativity theorem by erasing shifted tags back to the pre-shift namespace through B_4 on inhabited components.*

Remark A.28.7 (What is not migrated)**.** *The shifted tag migration should not change classical upper–lower tensor contraction, same-variance comparison rules, the OP status, the Historical Witness, or the physical meaning of an already guarded readout. It changes the QAM namespace and routing labels, not the underlying mathematical payloads.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.28.4 Router-package migration

Definition A.28.8 (Router-package translation)**.** *Let*

$$\mathfrak{R} = (K, M, C, T, B, G)$$

be a guarded router object, with kind registry K, route-mask registry M, classical collars C, trace record T, backtranslation record B, and guard record G. Its shifted translation is the object

$$\Sigma_4(\mathfrak{R}) = (\Sigma_4 K, \Sigma_4 M, C, T, \mathrm{B}_4^{\mathfrak{R}}, G),$$

where collars, traces, and guards are preserved, while kind and mask registries are reindexed and the backtranslation record is replaced by the corresponding shifted-to-pre-shift decoding data.

Lemma A.28.9 (Router guard preservation)**.** *If a router object $\mathfrak{R}$ is guarded before migration, then $\Sigma_4(\mathfrak{R})$ carries the same guard record after migration.*

Proof. By Definition A.28.8, the guard record G is copied unchanged. The migration reindexes registries and replaces the backtranslation ledger; it does not modify the guard predicates. Hence guard status is preserved. □

A.28.5 Readout-package migration

Definition A.28.10 (Readout-package translation)**.** *A guarded readout package may be translated to shifted only by translating its QAM/router labels while leaving its declared readout target, comparison tensor, boundary collar, residual map, and non-claim guard unchanged.*

Proposition A.28.11 (Machine/content preservation under migration)**.** *The future shifted global QAM Hilbert-shift preserves the historical release line machine/content separation: QAM/MML/MMA remains the router/compiler language, and physical or mathematical content remains in the guarded readout package.*

Proof. The migration acts on labels, kinds, masks, and routing registries. It does not act on the declared target of a physical readout, the comparison tensor, the boundary collar, or the residual map. Therefore it changes the machine addressing language while preserving the separation between machine layer and content layer. □

A.28.6 Conservativity statement for the future shifted shift

Theorem A.28.12 (shifted migration conservativity contract)**.** *Suppose a future shifted text translates every inhabited pre-shift QAM tag through Σ_4, reserves 0 only for the empty/no-active-kind marker, and equips each translated router or readout object with its backtranslation ledger B_4. Then every finite guarded shifted router/readout package has an inhabited-component backtranslation to a finite guarded pre-shift package with the same payloads and the same theorematic status.*

Proof. Apply B_4 to every inhabited shifted tag and leave non-tag payloads fixed. Round-trip correctness on inhabited tags follows from Lemma A.28.5. Router guards are preserved by Lemma A.28.9. Readout targets, comparison tensors, boundary collars, residual maps, and non-claim guards are unchanged by Definition A.28.10. Hence the translated object backtranslates to a pre-shift guarded object with identical payloads and unchanged theorematic status. The reserved value 0 has no inhabited backtranslation and therefore cannot be used to manufacture a new certificate. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.28.7 historical release line handoff to shifted

Definition A.28.13 (shifted entry checklist)**.** *A later archival stage draft should begin only after the following checklist is satisfied:*

1. *the global tag translation Σ_4 is fixed;*
2. *every occurrence class of QAM tags has an entry in the migration ledger;*
3. *every route-mask example has a shifted and pre-shift display pair;*
4. *every router object carries a backtranslation record;*
5. *the HC-boundary activation shift remains local and is not redefined;*
6. *the shifted text carries the conservativity contract of Theorem A.28.12;*
7. *OP 1/4 and OP 5(B) are not claimed closed merely by the migration.*

Remark A.28.14 (historical release line non-claim)**.** *Version 3.37.0 is a migration-map and backtranslation-contract layer. It does not execute the shifted global QAM Hilbert-shift, does not close OP 1/4 or OP 5(B), does not activate primitive mixed tensors, and does not claim Turing-completeness. Its purpose is to make the later shifted migration auditable before the namespace is actually changed.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.29 archival stage Concrete Migration Ledger and Route-Mask Examples

Status. This section is a concrete-audit layer for the historical release line migration map. It still does not execute the global shifted QAM Hilbert-shift. Its purpose is to make the future migration visually and mechanically checkable before the namespace is changed.

A.29.1 Minimal tag ledger

Definition A.29.1 (Concrete migration ledger)**.** *The concrete migration ledger is the finite display of the rule*

$$\Sigma_4(k) = k + 1, \qquad \mathrm{B}_4(a) = a - 1 \quad (a \geq 1),$$

together with the reserved value

$$0_{\mathrm{shift}} = \varnothing_{\mathrm{active}},$$

which has no inhabited pre-shift backtranslation.

pre-shift tag	shifted tag	reading	backtranslation
reserved / none	0	empty/no-active-kind	none
0	1	first inhabited kind	$\mathrm{B}_4(1) = 0$
1	2	second inhabited kind	$\mathrm{B}_4(2) = 1$
2	3	third inhabited kind	$\mathrm{B}_4(3) = 2$
k	$k+1$	inhabited kind k	$\mathrm{B}_4(k+1) = k$

Remark A.29.2 (Why the ledger is not a new theorem)**.** *The ledger is a bookkeeping surface. It records how inhabited tags will be renamed in shifted. It does not change the underlying readout, comparison tensor, boundary collar, residual map, or OP status.*

A.29.2 Route payload example

Definition A.29.3 (Route payload migration)**.** *For a finite ordered route payload*

$$r = \langle k_1, \ldots, k_n \rangle,$$

the future shifted route payload is

$$\Sigma_4(r) = \langle k_1 + 1, \ldots, k_n + 1 \rangle.$$

Its backtranslation is applied componentwise:

$$\mathrm{B}_4(\langle a_1, \ldots, a_n \rangle) = \langle a_1 - 1, \ldots, a_n - 1 \rangle$$

whenever all $a_i \geq 1$.

Remark A.29.4 (Example: two-entry route payload)**.** *The pre-shift payload*

$$r = \langle 5, 12 \rangle$$

migrates to

$$\Sigma_4(r) = \langle 6, 13 \rangle.$$

Backtranslation returns

$$\mathrm{B}_4(\langle 6, 13 \rangle) = \langle 5, 12 \rangle.$$

Thus the route addresses change, while the ordered route structure and the payload role are preserved.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.29.3 Packed mask example

Definition A.29.5 (Support mask versus packed mask)**.** *A support mask is a finite inhabited support set*

$$S \subseteq \mathbb{N}.$$

Its pre-shift packed integer is

$$m_3(S) = \sum_{k \in S} 2^k.$$

After migration the support becomes

$$\Sigma_4(S) = \{k + 1 : k \in S\},$$

and its shifted packed integer is

$$m_4(\Sigma_4(S)) = \sum_{k \in S} 2^{k+1}.$$

Lemma A.29.6 (Packed-mask doubling rule)**.** *For every finite inhabited support S,*

$$m_4(\Sigma_4(S)) = 2\, m_3(S).$$

Proof. By definition,

$$m_4(\Sigma_4(S)) = \sum_{k \in S} 2^{k+1} = 2 \sum_{k \in S} 2^k = 2\, m_3(S).$$

The rule applies to the packed integer representation of the support. It is not the same operation as adding 1 to the packed integer. □

Remark A.29.7 (Example: packed mask for the route support $\{5, 12\}$)**.** *For*

$$S = \{5, 12\},$$

the pre-shift packed mask is

$$m_3(S) = 2^5 + 2^{12} = 32 + 4096 = 4128.$$

The shifted support is

$$\Sigma_4(S) = \{6, 13\},$$

and the shifted packed mask is

$$m_4(\Sigma_4(S)) = 2^6 + 2^{13} = 64 + 8192 = 8256.$$

This equals $2 \cdot 4128$. Backtranslation of the support gives

$$\mathrm{B}_4(\{6, 13\}) = \{5, 12\}.$$

Remark A.29.8 (No accidental activation of the reserved bit)**.** *Since inhabited supports are shifted upward, the shifted bit 0 is not produced by migration. It remains reserved for the empty/no-active-kind marker. This is the operational reason why the global shift is compatible with the HC-boundary interpretation: the first active bit is no longer confused with the empty marker.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.29.4 Two-sided tensor ledger example

Definition A.29.9 (Tensor-index migration ledger)**.** *A two-sided structural tensor expression may carry QAM/router tags inside its front-side or back-side index payloads. A future shifted migration applies Σ_4 only to those tag payloads. It does not change front/back sidedness, upper/lower variance, or the declared contraction/comparison rule.*

Remark A.29.10 (Example: front-side route migration)**.** *A readout-facing tensor package with front-side route payload*

$$^{\langle 5,12\rangle}{}_{\langle 2\rangle}\mathcal{T}^{i}{}_{j}$$

migrates to

$$^{\langle 6,13\rangle}{}_{\langle 3\rangle}\mathcal{T}^{i}{}_{j}.$$

The front-side payloads are reindexed. The back-side structural indices i, j are not changed in this example. The tensor still has the same front/back and upper/lower grammar.

Remark A.29.11 (Example: full payload migration)**.** *If both front-side and back-side QAM payloads are explicitly tagged, then*

$$^{\langle 5,12\rangle}{}_{\langle 2\rangle}\mathcal{T}^{\langle 7\rangle}{}_{\langle 10\rangle}$$

migrates to

$$^{\langle 6,13\rangle}{}_{\langle 3\rangle}\mathcal{T}^{\langle 8\rangle}{}_{\langle 11\rangle}.$$

This is a label migration only. It does not turn an upper index into a lower index, a front-side readout into a back-side action, or a comparison tensor into a contraction tensor.

Proposition A.29.12 (Tensor grammar preservation under the migration ledger)**.** *The concrete migration ledger preserves the two-sided structural tensor grammar.*

Proof. The migration acts on tag payloads by $k \mapsto k + 1$. The syntactic positions of the payloads are unchanged: front-side payloads remain front-side payloads, back-side payloads remain back-side payloads, upper payloads remain upper payloads, and lower payloads remain lower payloads. Therefore classical upper–lower readout and explicit same-variance comparison requirements are unchanged. □

A.29.5 shifted readiness consequence

Theorem A.29.13 (Concrete ledger supports the shifted conservativity contract)**.** *The concrete tag ledger, route payload example, packed-mask rule, and two-sided tensor ledger examples support the historical release line backtranslation contract: every migrated finite example displayed in this section backtranslates to its pre-shift source example on inhabited components.*

Proof. The tag ledger backtranslates by $\mathrm{B}_4(a) = a-1$ for $a \geq 1$. Route payloads backtranslate componentwise. Packed masks backtranslate at the level of support sets, not by subtracting 1 from the packed integer. Tensor payloads backtranslate by applying the same rule to the tags carried inside their index payloads, while preserving sidedness and variance. Hence each finite example returns to its pre-shift source example on inhabited components. □

Remark A.29.14 (archival stage non-claim)**.** *Version 3.37.1 still does not execute the global shifted QAM Hilbert-shift. It only adds concrete ledger examples. The actual global migration remains a shifted task.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.30 Global QAM Hilbert-Shift Activation Candidate

Status. This is the first shifted activation release-final state. It executes the global QAM namespace decision prepared in Versions 3.37.0–3.37.1: from this layer forward, the active QAM language reserves tag 0 for the empty/no-active-kind marker and reads inhabited tags in the shifted shifted namespace. The earlier historical release line HC-boundary shift remains local and structural; the present layer is the global language shift.

A.30.1 Governing shifted namespace convention

Definition A.30.1 (Global shifted QAM Hilbert-shift convention)**.** *The governing shifted QAM namespace is obtained from the inhabited pre-shift namespace by the shift*

$$\Sigma_4(k) = k + 1.$$

The tag 0 *in the shifted namespace is reserved globally for the empty or no-active-kind marker:*

$$0_{\text{shift}} = \varnothing_{\text{active}}.$$

It is not the image of an inhabited pre-shift tag.

Definition A.30.2 (shifted backtranslation ledger)**.** *The shifted backtranslation ledger is the partial map*

$$\mathrm{B}_4(a) = a - 1$$

defined on inhabited shifted tags $a \geq 1$*. The reserved tag* 0 *has no inhabited backtranslation.*

Remark A.30.3 (What is executed in)**.** *The executed decision is the global namespace convention: shifted readout packages use the shifted tag convention by default. The text does not pretend that the Historical Witness has been rewritten. Historical material remains readable by the backtranslation ledger.*

A.30.2 Compatibility with the first HC-boundary shift

Definition A.30.4 (Shift A and Shift B in the shifted text)**.** *The shifted text keeps the following distinction.*

1. Shift A *is the local historical release line HC-boundary activation*

   $$0_{\text{mask}} \longrightarrow 1_{\text{mask}}.$$

   It explains why the Heisenberg Compensator appears as a boundary/readout normalizer at the first active readout transition.

2. Shift B *is the global shifted QAM Hilbert-shift*

   $$k \longmapsto k + 1$$

   applied to inhabited QAM tags.

Proposition A.30.5 (HC-boundary shift is not redefined by shifted)**.** *The global shifted QAM Hilbert-shift does not redefine the historical release line HC-boundary activation principle.*

Proof. Shift A concerns the structural boundary from inactive mask state to first active readout state. Shift B concerns the global label namespace for inhabited QAM tags. These acts have different domains: the first is a local boundary-normalization reading, the second is a global language migration. Therefore the shifted namespace convention preserves the HC-boundary reading rather than replacing it. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.30.3 Governing shifted ledger

pre-shift tag	shifted tag	shifted reading	backtranslation
none	0	empty/no-active-kind	none
0	1	first inhabited kind	$\mathrm{B}_4(1) = 0$
1	2	second inhabited kind	$\mathrm{B}_4(2) = 1$
2	3	third inhabited kind	$\mathrm{B}_4(3) = 2$
k	$k+1$	inhabited kind k	$\mathrm{B}_4(k+1) = k$

Lemma A.30.6 (shifted inhabited round-trip)**.** *For every inhabited pre-shift tag k,*

$$\mathrm{B}_4(\Sigma_4(k)) = k.$$

For every inhabited shifted tag $a \geq 1$,

$$\Sigma_4(\mathrm{B}_4(a)) = a.$$

Proof. Both claims are the identities $(k+1)-1 = k$ and $(a-1)+1 = a$. The reserved value 0 is excluded from the inhabited backtranslation domain, so it cannot generate an inhabited source tag. □

A.30.4 Route-mask and support migration

Definition A.30.7 (shifted route payload)**.** *A pre-shift ordered route payload*

$$r = \langle k_1, \ldots, k_n \rangle$$

is read in the shifted language as

$$r^{(4)} = \Sigma_4(r) = \langle k_1 + 1, \ldots, k_n + 1 \rangle.$$

Definition A.30.8 (shifted support and packed mask)**.** *For a finite inhabited support $S \subseteq \mathbb{N}$, the migrated shifted support is*

$$S^{(4)} = \{k + 1 : k \in S\}.$$

If the pre-shift packed integer is

$$m_3(S) = \sum_{k \in S} 2^k,$$

then the shifted packed integer is

$$m_4(S^{(4)}) = \sum_{k \in S} 2^{k+1}.$$

Lemma A.30.9 (shifted packed-mask doubling)**.** *For every finite inhabited support S,*

$$m_4(S^{(4)}) = 2\, m_3(S).$$

Proof. The proof is the direct calculation

$$m_4(S^{(4)}) = \sum_{k \in S} 2^{k+1} = 2 \sum_{k \in S} 2^k = 2\, m_3(S).$$

□

Remark A.30.10 (Example: route and mask)**.** *The pre-shift route payload $\langle 5, 12 \rangle$ is read in shifted as $\langle 6, 13 \rangle$. The support $\{5, 12\}$ has packed value $2^5 + 2^{12} = 4128$ before migration and packed value $2^6 + 2^{13} = 8256$ after migration. The packed integer doubles; the support backtranslates by subtracting one from inhabited support entries.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.30.5 Two-sided tensor grammar under shifted

Definition A.30.11 (shifted tensor-payload migration)**.** *In a two-sided structural tensor expression, the shifted shift acts only on QAM or router tag payloads carried by index positions. It does not alter front/back sidedness, upper/lower variance, classical upper–lower contraction, or the requirement that same-variance comparisons use explicit comparison or selector tensors.*

Remark A.30.12 (Example: front-side migration)**.** *The front-side route display*

$$\mathcal{T}^{\langle 5,12\rangle}{}_{\langle 2\rangle}{}^{i}{}_{j}$$

is read in shifted as

$$\mathcal{T}^{\langle 6,13\rangle}{}_{\langle 3\rangle}{}^{i}{}_{j}.$$

The expression changes its tag payloads, not its tensor grammar.

Remark A.30.13 (Example: full payload migration)**.** *If both front-side and back-side payloads are tagged, then*

$$\mathcal{T}^{\langle 5,12\rangle}{}_{\langle 2\rangle}{}^{\langle 7\rangle}{}_{\langle 10\rangle}$$

is read in shifted as

$$\mathcal{T}^{\langle 6,13\rangle}{}_{\langle 3\rangle}{}^{\langle 8\rangle}{}_{\langle 11\rangle}.$$

The front/back positions and the upper/lower variance are preserved.

Theorem A.30.14 (shifted tensor-grammar preservation)**.** *The global shifted QAM Hilbert-shift preserves the two-sided structural tensor grammar.*

Proof. The shift is a function on tag payloads. It does not move an index from front to back, from back to front, from upper to lower, or from lower to upper. Therefore the grammar of sidedness and variance is unchanged. Since classical upper–lower contraction and explicit same-variance comparison are defined by position and variance rather than by the numerical tag name, those rules are preserved. □

A.30.6 shifted router and readout activation

Definition A.30.15 (shifted router object)**.** *A shifted router object is a guarded router object whose inhabited kind registry and route-mask registry are read in the shifted shifted namespace, while its collars, trace records, backtranslation ledger, and guard record remain explicit.*

Definition A.30.16 (shifted readout object)**.** *A shifted readout object is a guarded readout package whose QAM/router labels are read in the shifted shifted namespace, while its readout target, comparison tensor, boundary collar, residual map, and non-claim guard remain unchanged.*

Theorem A.30.17 (shifted activation conservativity)**.** *Every finite guarded shifted router/readout package admitted by Definitions A.30.15 and A.30.16 backtranslates on inhabited components to a finite guarded pre-shift package with the same payloads and the same theorematic status.*

Proof. Apply $\mathrm{B}_4(a) = a - 1$ to every inhabited shifted tag. By Lemma A.30.6, this recovers the corresponding pre-shift tag on inhabited components. Collars, trace records, comparison tensors, readout targets, boundary collars, residual maps, and non-claim guards are kept fixed by definition. The reserved value 0 has no inhabited backtranslation, so it cannot create a new active component. Hence the package backtranslates without changing payloads or theorematic status. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.30.7 handoff

Remark A.30.18 (non-claims)**.** *This shifted activation release-final state performs the namespace decision, but it does not close OP 1/4 or OP 5(B), does not activate primitive mixed tensors, does not claim Turing-completeness, and does not replace standard QFT/QED/Lifshitz or general-relativistic calculations. It is a language and router/readout migration step.*

Theorem A.30.19 (handoff theorem)**.** *The local proof line layer is a well-formed global QAM Hilbert-shift activation release-final state over the archival stage release line.*

Proof. Definition A.30.1 fixes the global namespace convention. Definition A.30.2 provides the backtranslation ledger. The route and mask rules preserve finite payloads by Lemma A.30.9. The two-sided tensor grammar is preserved by Theorem A.30.14. Finally, finite router/readout packages remain conservative under Theorem A.30.17. Therefore the layer is well formed as a global QAM language activation release-final state, while all stated non-claims remain in force. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.31 Occurrence Ledger and Shift-Audit Map

Status. This layer is an occurrence ledger for the shifted global QAM Hilbert-shift. Version 4.0.0-fixed the governing shifted namespace. The present layer records where that shift is allowed to act, where it is forbidden to act, and where a backtranslation or historical reading must be retained. It is an audit map, not a new physical promotion.

A.31.1 Occurrence classes

Definition A.31.1 (shifted occurrence class)**.** *A* shifted occurrence class *is a declared role assigned to a visible tag, mask, route, tensor-index payload, or OP-facing label in the Active Core. The role determines whether the global shifted shift is applied, whether the object is kept historical, or whether the object is read only through an explicit backtranslation ledger.*

Definition A.31.2 (The five primary occurrence classes)**.** *The occurrence ledger uses the following primary classes.*

1. ***Class A: historical/pre-shift witness occurrences.*** *These remain unchanged and are read through the Historical Witness or through the backtranslation ledger.*
2. ***Class B: active shifted router occurrences.*** *These are read in the shifted shifted namespace by default.*
3. ***Class C: example, table, route, and mask occurrences.*** *These must carry a shifted display and, when useful, a pre-shift backtranslation display.*
4. ***Class D: two-sided tensor-index payload occurrences.*** *These shift only their QAM/router tag payloads. The tensor grammar itself is not shifted.*
5. ***Class E: OP-facing and physical-readout occurrences.*** *These may have shifted router labels, but their theorematic status, physical readout target, residual map, and non-claim guard remain unchanged.*

Remark A.31.3 (Why occurrence classes are needed)**.** *Without occurrence classes, the symbol* 5 *could ambiguously mean a pre-shift tag, a shifted tag, a packed mask integer, a tensor component, or a mere example number. The occurrence ledger prevents that ambiguity by assigning a role before any shift is read.*

A.31.2 Class A: historical and pre-shift witness occurrences

Definition A.31.4 (Class A occurrence)**.** *A Class A occurrence is any tag, route, mask, or notation block that belongs to the preserved historical line, to the pre-shift witness text, or to an explicitly quoted earlier version. Class A occurrences are not rewritten by the shifted Active Core. They remain recoverable by the Historical Witness and readable by backtranslation.*

Lemma A.31.5 (Class A preservation)**.** *The shifted occurrence ledger preserves Class A occurrences without applying the global shift to their displayed historical form.*

Proof. Class A is defined as historical or pre-shift witness material. Its function is to preserve genealogy and reconstruction memory, not to govern the active shifted namespace. Therefore applying the shifted shift directly to Class A would corrupt the witness role. The correct operation is preservation plus optional backtranslation annotation. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.31.3 Class B: active shifted router occurrences

Definition A.31.6 (Class B occurrence)**.** *A Class B occurrence is a tag or route occurrence inside an active shifted router object, shifted kind registry, shifted route-mask registry, or shifted machine-language package. Class B occurrences are read in the shifted namespace:*

$$k_{\mathrm{pre}} \longmapsto k_{\mathrm{shift}} = k_{\mathrm{pre}} + 1.$$

Remark A.31.7 (Class B default)**.** *From the local proof line onward, active router-language statements default to Class B unless the text explicitly marks the occurrence as historical, illustrative, or unshifted for comparison.*

Lemma A.31.8 (Class B backtranslation)**.** *Every finite Class B tag family with inhabited shifted tags has a componentwise backtranslation by* $\mathrm{B}_4(a) = a - 1$.

Proof. Class B contains active inhabited shifted router labels. By the shifted ledger, every inhabited shifted tag satisfies $a \geq 1$, so $\mathrm{B}_4(a) = a-1$ is defined. Applying this componentwise gives the pre-shift label family. □

A.31.4 Class C: examples, tables, routes, and masks

Definition A.31.9 (Class C occurrence)**.** *A Class C occurrence is a displayed example, table entry, route payload, support set, or packed mask integer used for exposition or audit. Class C occurrences must declare whether the displayed value is a pre-shift value, a shifted value, or a packed integer derived from a support.*

Definition A.31.10 (Class C display contract)**.** *A Class C route or mask example is well displayed if it separates:*

$$\textit{support}, \qquad \textit{route payload}, \qquad \textit{packed integer}.$$

In particular, support migration is

$$S \mapsto S^{(4)} = \{k + 1 : k \in S\},$$

while packed-mask migration is

$$m_3(S) = \sum_{k \in S} 2^k \quad \mapsto \quad m_4(S^{(4)}) = 2m_3(S).$$

Remark A.31.11 (Class C warning)**.** *The packed integer should not be shifted by adding* 1*. The support is shifted by adding* 1 *to each inhabited support entry; the packed integer then changes according to the support rule, often by doubling in the pure support-shift case.*

A.31.5 Class D: two-sided tensor-index payloads

Definition A.31.12 (Class D occurrence)**.** *A Class D occurrence is a QAM/router tag payload carried inside a front-side or back-side index position of a two-sided structural tensor expression. The shifted shift acts on the payload label only. It does not alter sidedness, variance, ordinary upper–lower contraction, or explicit same-variance comparison.*

Proposition A.31.13 (Class D grammar invariance)**.** *Class D migration preserves the two-sided structural tensor grammar.*

Proof. The migration rule changes only the numerical or route-label payload carried by an index. It does not move that payload from the front side to the back side, from the back side to the front side, from upper position to lower position, or from lower position to upper position. Thus the grammar of sidedness and variance is unchanged. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.31.6 Class E: OP-facing and physical-readout occurrences

Definition A.31.14 (Class E occurrence). *A Class E occurrence is an OP-facing label, residual package, physical-readout package, boundary-readout package, or diagnostic object whose router address may be shifted but whose physical or theorematic status is not changed by the shift.*

Definition A.31.15 (Class E invariance contract). *A Class E occurrence remains valid under shifted only if the following data remain unchanged:*

1. *its OP status;*
2. *its physical or mathematical readout target;*
3. *its comparison tensor or boundary collar;*
4. *its residual map or diagnostic map;*
5. *its non-claim guard.*

Lemma A.31.16 (Class E status preservation). *The shifted occurrence ledger does not close OP 1/4 or OP 5(B).*

Proof. Class E migration may reindex router labels, but Definition A.31.15 fixes the OP status and non-claim guard. Since no closure certificate is added by reindexing, the OP status remains unchanged. □

A.31.7 Occurrence decision table

Class	Where it occurs	Shift action	Required audit note
A	Historical/pre-shift witness blocks	preserve displayed form	read via witness/backtranslation
B	active router/kind/route registries	shift inhabited tags	keep B_4 ledger
C	examples, tables, masks, routes	display pre-shift and shifted roles	separate support from packed integer
D	tensor-index tag payloads	shift payload only	preserve sidedness/variance
E	OP-facing/readout packages	shift router address only	preserve OP status/guard

A.31.8 Ambiguity filters

Definition A.31.17 (Ambiguity filter). *Before a visible integer, tuple, support, or tensor payload is read as shifted, the reader must determine whether it is:*

tag, *route payload*, *support set*,
packed mask integer, *tensor component*, *historical witness value.*

Only tag, route-payload, support, and tag-payload occurrences are subject to the shifted shift.

Remark A.31.18 (Packed values are not tags). *A packed integer such as* 4128 *is not itself a tag occurrence. It encodes a support. The support migrates first, and the packed integer is recomputed from the migrated support.*

A.31.9 shifted occurrence-ledger conservativity

Theorem A.31.19 (Occurrence-ledger conservativity). *Every finite shifted occurrence family classified by Classes A–E has an honest reading that either preserves its displayed historical form or backtranslates its inhabited shifted components to the pre-shift source language without changing payloads, tensor grammar, OP status, or non-claim guards.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. Class A is preserved by Lemma A.31.5. Class B backtranslates by Lemma A.31.8. Class C separates support, route payload, and packed integer by Definition A.31.10. Class D preserves the two-sided tensor grammar by Proposition A.31.13. Class E preserves OP and readout status by Lemma A.31.16. The five cases cover the declared occurrence classes; hence the occurrence ledger is conservative for finite classified occurrence families. □

Remark A.31.20 (non-claim)**.** *The occurrence ledger does not add a new physical equation, does not close an OP, does not activate primitive mixed tensors, and does not claim Turing-completeness. It makes the global shifted shift auditable by assigning roles to visible occurrences.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.32 Global Reindexed Display Layer

Status. This layer adds normalized side-by-side displays for the global shifted QAM Hilbert-shift. It does not add new physics, close OP 1/4, close OP 5(B), or alter the two-sided tensor grammar. It only standardizes how active examples are displayed in three synchronized forms:

pre-shift source, shifted display, backtranslation.

A.32.1 Reindexed display triple

Definition A.32.1 (Reindexed display triple)**.** *A* shifted reindexed display triple *is a finite record*

$$\mathfrak{D}_4(X) = (X_{\mathrm{pre}}, X_{\mathrm{shift}}, \mathrm{B}_4(X_{\mathrm{shift}})),$$

where X_{pre} *is the pre-shift source display,* X_{shift} *is the shifted shifted display, and* $\mathrm{B}_4(X_{\mathrm{shift}})$ *is the explicit backtranslation on inhabited components.*

Remark A.32.2 (Display triples are audit objects)**.** *A display triple is not a new theorematic certificate. It is an audit object showing that the visible shifted display can be read back to the pre-shift source where the components are inhabited.*

A.32.2 Tag and route display triples

Definition A.32.3 (Tag display triple)**.** *For an inhabited pre-shift tag* k*, the shifted tag display triple is*

$$\mathfrak{D}_4(k) = (k,\ k+1,\ \mathrm{B}_4(k+1) = k).$$

The reserved value 0_{shift} *has no inhabited pre-shift source and is displayed separately as*

$$(\varnothing_{\mathrm{active}},\ 0_{\mathrm{shift}},\ \textit{no inhabited backtranslation}).$$

Definition A.32.4 (Route display triple)**.** *For a finite pre-shift route payload*

$$r = \langle k_1, \ldots, k_n \rangle,$$

the shifted display triple is

$$\mathfrak{D}_4(r) = (\langle k_1, \ldots, k_n \rangle,\ \langle k_1 + 1, \ldots, k_n + 1 \rangle,\ \langle k_1, \ldots, k_n \rangle)\,.$$

pre-shift source	shifted display	backtranslation
5	6	$\mathrm{B}_4(6) = 5$
12	13	$\mathrm{B}_4(13) = 12$
$\langle 5, 12 \rangle$	$\langle 6, 13 \rangle$	$\langle 5, 12 \rangle$

A.32.3 Support and packed-mask display triples

Definition A.32.5 (Support display triple)**.** *For a finite inhabited support set* S*, the support display triple is*

$$\mathfrak{D}_4(S) = (S,\ S^{(4)},\ \mathrm{B}_4(S^{(4)})), \qquad S^{(4)} = \{k+1 : k \in S\}.$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition A.32.6 (Packed-mask display triple)**.** *For a finite inhabited support S, the packed-mask display triple is*

$$\mathfrak{D}_4(m(S)) = \big(m_3(S),\ m_4(S^{(4)}),\ S = \mathrm{B}_4(S^{(4)})\big)\,.$$

The backtranslation is attached to the support, not to the packed integer by subtraction.

pre-shift source	shifted display	backtranslation
$\{5,12\}$	$\{6,13\}$	$\{5,12\}$
$2^5 + 2^{12} = 4128$	$2^6 + 2^{13} = 8256$	support returns to $\{5,12\}$

Lemma A.32.7 (Packed-mask display consistency)**.** *The packed-mask display triple for a finite support S is consistent with the packed-mask doubling rule:*

$$m_4(S^{(4)}) = 2\, m_3(S).$$

Proof. This is exactly Lemma A.31.8 applied to the inhabited support entries together with the packed-mask calculation

$$\sum_{k\in S} 2^{k+1} = 2\sum_{k\in S} 2^k.$$

The displayed packed integer is recomputed from the migrated support, not shifted as a tag. □

A.32.4 Two-sided tensor display triples

Definition A.32.8 (Tensor display triple)**.** *A two-sided structural tensor display triple is obtained by applying the shifted tag shift to QAM/router payloads inside index positions while leaving sidedness and variance unchanged.*

pre-shift source	shifted display	backtranslation
${}^{\langle 5,12\rangle}{}_{\langle 2\rangle}\mathcal{T}^{i}{}_{j}$	${}^{\langle 6,13\rangle}{}_{\langle 3\rangle}\mathcal{T}^{i}{}_{j}$	${}^{\langle 5,12\rangle}{}_{\langle 2\rangle}\mathcal{T}^{i}{}_{j}$
${}^{\langle 5,12\rangle}{}_{\langle 2\rangle}\mathcal{T}^{\langle 7\rangle}{}_{\langle 10\rangle}$	${}^{\langle 6,13\rangle}{}_{\langle 3\rangle}\mathcal{T}^{\langle 8\rangle}{}_{\langle 11\rangle}$	${}^{\langle 5,12\rangle}{}_{\langle 2\rangle}\mathcal{T}^{\langle 7\rangle}{}_{\langle 10\rangle}$

Proposition A.32.9 (Display triples preserve tensor grammar)**.** *Tensor display triples preserve the two-sided structural tensor grammar.*

Proof. Only QAM/router tag payloads inside the displayed index positions are shifted. The index positions themselves are not moved. Hence front-side remains front-side, back-side remains back-side, upper remains upper, lower remains lower, and the ordinary contraction/comparison rules remain those of the underlying tensor grammar. □

A.32.5 OP-facing display triples

Definition A.32.10 (OP-facing display triple)**.** *An OP-facing display triple records only the router-address migration of an OP-facing package. Its OP status, residual target, comparison surface, and non-claim guard are displayed as invariant data.*

pre-shift address	shifted address	invariant data
OP package tag k	OP package tag $k+1$	OP status unchanged; guard unchanged
residual route $\langle 5,12\rangle$	residual route $\langle 6,13\rangle$	residual target unchanged
boundary readout route $\langle 2\rangle$	boundary readout route $\langle 3\rangle$	boundary collar unchanged

Lemma A.32.11 (OP-facing display triples do not close OPs)**.** *OP-facing display triples do not close OP 1/4 or OP 5(B).*

Proof. By Definition A.32.10, the triple changes only router-address display. It explicitly leaves OP status, residual target, comparison surface, and non-claim guard invariant. Therefore no OP closure certificate is introduced. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.32.6 Display-layer conservativity

Theorem A.32.12 (Global reindexed display conservativity)**.** *Every finite display triple has a backtranslation to its pre-shift source display on inhabited components and preserves the occurrence class assigned in the occurrence ledger.*

Proof. Tag and route triples backtranslate componentwise by $\mathrm{B}_4(a) = a - 1$. Support triples backtranslate by applying the same rule to support entries. Packed-mask triples backtranslate through their support sets, not by subtracting from the packed integer. Tensor triples preserve grammar by Proposition A.32.9. OP-facing triples preserve status by Lemma A.32.11. Hence the display layer is conservative over the occurrence ledger. □

Remark A.32.13 (non-claim)**.** *The layer is a display layer only. It does not add a new physical equation, does not close OP 1/4 or OP 5(B), does not activate primitive mixed tensors, and does not claim Turing-completeness.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.33 Physical Readout Compatibility Check

Status. This layer checks that the global shifted QAM Hilbert-shift is compatible with the active physical and mathematical readout packages. It does not promote a new physical law. It verifies that the shifted shift changes router addresses and tag payloads only, while readout targets, comparison surfaces, residual maps, and non-claim guards remain fixed.

A.33.1 Readout compatibility principle

Definition A.33.1 (Readout compatibility package)**.** *A* shifted readout compatibility package *is a tuple*

$$\mathcal{C}_4(R) = (R_{\mathrm{pre}}, R_{\mathrm{shift}}, \mathrm{B}_4(R_{\mathrm{shift}}), T_R, G_R, S_R),$$

where R_{pre} is the pre-shift router address, R_{shift} is the shifted shifted router address, $\mathrm{B}_4(R_{\mathrm{shift}})$ is the inhabited backtranslation, T_R is the readout target, G_R is the guard record, and S_R is the theorematic status record.

Definition A.33.2 (Readout compatibility)**.** *A readout package is* shifted-compatible *if the migration from R_{pre} to R_{shift} changes only the router address and leaves T_R, G_R, and S_R unchanged.*

Lemma A.33.3 (Readout compatibility is address-local)**.** *If a readout package is shifted-compatible, then the global shifted QAM Hilbert-shift does not change its physical or mathematical target.*

Proof. By Definition A.33.2, the shift is applied to the router address only. The readout target T_R is explicitly preserved. Hence the physical or mathematical target of the package is unchanged. □

A.33.2 HC-boundary readout compatibility

Definition A.33.4 (HC-boundary compatibility record)**.** *The HC-boundary readout package is shifted-compatible when the local structural shift*

$$0_{\mathrm{mask}} \longrightarrow 1_{\mathrm{mask}}$$

remains the HC-boundary activation principle, while global shifted router labels are read in the shifted namespace.

Proposition A.33.5 (HC-boundary readout survives the global shift)**.** *The global shifted QAM Hilbert-shift preserves the historical release line HC-boundary activation reading.*

Proof. The HC-boundary reading belongs to Shift A: the local transition from inactive mask state to first active readout state. The shifted QAM Hilbert-shift belongs to Shift B: the global namespace migration of inhabited tags. Since the local proof chain separate these two acts, the global shift cannot redefine the local HC-boundary activation. It can only change how the active router address is displayed. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.33.3 Photon-compatible null-side readout compatibility

Definition A.33.6 (Null-side readout compatibility)**.** *A photon-compatible null-side readout is shifted-compatible when the null-side readout target and its guard remain unchanged while its router address is displayed in the shifted shifted namespace.*

Lemma A.33.7 (Null-side target preservation)**.** *The shifted shift preserves the null-side readout target of a photon-compatible readout package.*

Proof. By Definition A.33.6, the null-side target is part of the invariant data. The shifted shift operates only on router address labels, so the null-side target remains fixed. □

Remark A.33.8 (No photon proof claim)**.** *This compatibility check does not prove a new photon theory. It only says that the already guarded null-side readout remains address-stable under the shifted tag migration.*

A.33.4 HC-normalized singularity projection compatibility

Definition A.33.9 (Singularity projection compatibility)**.** *An HC-normalized singularity projection package is shifted-compatible when its projection target, HC-normalization guard, and residual diagnostic remain unchanged under router-address migration.*

Proposition A.33.10 (Singularity projection status preservation)**.** *The shifted shift does not promote an HC-normalized singularity projection package from diagnostic status to theorematic closure.*

Proof. The migration changes the router address. It does not add a new projection identity, a new residual estimate, or a new closure certificate. The diagnostic guard is part of the invariant record in Definition A.33.9. Therefore the theorematic status is preserved. □

A.33.5 Casimir boundary readout compatibility

Definition A.33.11 (Casimir boundary compatibility)**.** *A Casimir boundary readout package is shifted-compatible when its boundary collar, comparison surface, and non-replacement guard remain unchanged while its router-address labels are shifted.*

Lemma A.33.12 (Casimir boundary collar preservation)**.** *The shifted shift preserves the boundary collar of a shifted-compatible Casimir boundary readout package.*

Proof. By Definition A.33.11, the boundary collar is invariant data. The shift acts only on router labels. Therefore the boundary collar is unchanged. □

Remark A.33.13 (No replacement of standard Casimir calculations)**.** *The compatibility statement does not replace standard QFT, QED, or Lifshitz calculations. It keeps the Companion readout package address-stable under the shifted namespace shift.*

A.33.6 OP-facing residual package compatibility

Definition A.33.14 (OP-facing residual compatibility)**.** *An OP-facing residual package is shifted-compatible when its route address may be shifted but its residual target, status flag, and non-claim guard remain unchanged.*

Lemma A.33.15 (OP-facing residual status preservation)**.** *The shifted shift does not close OP 1/4 or OP 5(B).*

Proof. By Definition A.33.14, OP-facing residual packages preserve their status flags and non-claim guards. A router address migration is not a proof certificate. Therefore the OP status remains unchanged. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.33.7 Compatibility table

Readout package	Shifted data	Invariant data
HC-boundary	active router address	local HC-boundary activation principle
null-side/photon-compatible	router address	null-side target and guard
singularity projection	router address	projection target, HC guard, residual diagnostic
Casimir boundary	router address	boundary collar, comparison surface, non-replacement guard
OP-facing residual	route address	residual target, OP status, non-claim guard

A.33.8 Physical-readout compatibility theorem

Theorem A.33.16 (Physical readout compatibility under the shifted shift)**.** *For the readout packages listed in Section A.33.7, the global shifted QAM Hilbert-shift preserves the declared readout target, comparison or boundary data, residual data, and non-claim guard. It changes only the router-address display.*

Proof. Each listed package is declared shifted-compatible by its corresponding definition: HC-boundary compatibility by Definition A.33.4, null-side compatibility by Definition A.33.6, singularity projection compatibility by Definition A.33.9, Casimir boundary compatibility by Definition A.33.11, and OP-facing residual compatibility by Definition A.33.14. In each case, the shifted component is the router or route address. The physical or mathematical readout target and the guard/status records are invariant data. Therefore the shifted shift is compatible with these readout packages without changing their theorematic status. □

Remark A.33.17 (non-claim)**.** *The layer is a compatibility check. It does not add a new physical equation, does not close OP 1/4 or OP 5(B), does not activate primitive mixed tensors, and does not claim Turing-completeness.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.34 Governing shifted Normal Form and Release-Freeze Handoff

Status. This layer compresses the archival stage-the local proof chain migration work into one governing normal form. It is intended as a release-freeze handoff: after this layer, new active QAM/router/readout material should be written either in the shifted normal form or with an explicit pre-shift backtranslation record.

A.34.1 Governing normal-form object

Definition A.34.1 (Governing shifted normal-form object)**.** *A governing shifted normal-form object is a finite record*

$$\mathcal{N}_4(X) = (C_X,\ X_{\mathrm{shift}},\ \mathrm{B}_4(X_{\mathrm{shift}}),\ P_X,\ G_X,\ S_X),$$

where C_X is the occurrence class, X_{shift} is the shifted shifted display, $\mathrm{B}_4(X_{\mathrm{shift}})$ is the inhabited-component backtranslation, P_X is the payload or readout target, G_X is the guard record, and S_X is the theorematic status record.

Remark A.34.2 (Why the pre-shift source is not always the first field)**.** *The shifted Active Core now governs by the shifted namespace. Therefore the normal form puts the shifted display first and keeps the pre-shift source as a backtranslation ledger when needed. Historical source displays remain available through the Historical Witness.*

A.34.2 Normal-form admissibility

Definition A.34.3 (shifted-normal admissibility)**.** *A finite object is* shifted-normal admissible *if:*

1. *it carries an occurrence class C_X;*
2. *all active inhabited tags are read in the shifted shifted namespace;*
3. *every shifted inhabited tag has a backtranslation by B_4;*
4. *packed masks are reconstructed from shifted supports, not shifted as integers;*
5. *tensor-index payload migration preserves sidedness and variance;*
6. *physical/readout payloads preserve their target, collar, residual map, guard, and theorematic status.*

Lemma A.34.4 (Normal admissibility implies ledger readability)**.** *Every shifted-normal admissible object is readable through the namespace ledger, the occurrence ledger, the display triple, and the readout compatibility check.*

Proof. The occurrence class invokes the local pass. The shifted tag convention invokes the local pass. The backtranslation field invokes whenever a visible pre-shift comparison is needed. The payload/guard/status fields invoke for readout packages and the tensor-grammar rules for tensor payloads. Hence the object is readable through the whole the local proof chain ledger chain. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.34.3 Normal forms by class

Definition A.34.5 (Class-wise shifted-namespace normal forms)**.** *The following class-wise normal forms are used:*

Class A: $(A,\ X_{\mathrm{hist}},\ \text{witness/backtranslation},\ P_X,\ G_X,\ S_X)$,
Class B: $(B,\ X_{\mathrm{shift}},\ \mathrm{B}_4(X_{\mathrm{shift}}),\ P_X,\ G_X,\ S_X)$,
Class C: $(C,$ *shifted display*, *pre-shift display*, *support/payload split*, $G_X,\ S_X)$,
Class D: $(D,$ *shifted tag payload*, *backtranslated payload*, *tensor grammar*, $G_X,\ S_X)$,
Class E: $(E,$ *shifted route address*, *backtranslated address*, *readout target*, $G_X,\ S_X)$.

Remark A.34.6 (Class C special rule)**.** *For Class C objects, a packed mask integer is never the primary shifted object. The primary shifted object is the support or route payload. The packed integer is recomputed from that shifted support.*

A.34.4 Normal-form examples

Class	shifted display	backtranslation	invariant data
B	$\langle 6,13\rangle$	$\langle 5,12\rangle$	router role
C	support $\{6,13\}$, mask 8256	support $\{5,12\}$, mask 4128	support/mask split
D	${}^{\langle 6,13\rangle}{}_{\langle 3\rangle}\mathcal{T}^{i}{}_{j}$	${}^{\langle 5,12\rangle}{}_{\langle 2\rangle}\mathcal{T}^{i}{}_{j}$	sidedness/variance
E	route $\langle 6,13\rangle$	route $\langle 5,12\rangle$	readout target and guard

A.34.5 Release-freeze discipline

Definition A.34.7 (shifted release-freeze discipline)**.** *The shifted release-freeze discipline is the rule that, after the local pass, every new active QAM/router/readout object must either:*

1. *be written directly as a shifted-normal admissible object; or*
2. *be marked as historical/pre-shift and supplied with an explicit backtranslation or witness pointer.*

Lemma A.34.8 (Release-freeze prevents mixed namespace drift)**.** *The shifted release-freeze discipline prevents uncontrolled mixing of pre-shift and shifted tag namespaces.*

Proof. Every new object must either be shifted-normal or explicitly historical/pre-shift. In the first case, inhabited tags are shifted and backtranslated by the shifted ledger. In the second case, the object is not treated as an active shifted governing object. Thus no object may silently mix pre-shift and shifted tag meanings. □

A.34.6 Governing shifted normal-form theorem

Theorem A.34.9 (Governing shifted normal-form theorem)**.** *The normal form is conservative over the local proof chain shifted migration chain: every finite shifted-normal admissible object has a controlled reading in the shifted shifted namespace and an inhabited-component backtranslation that preserves payloads, tensor grammar, physical/readout targets, guards, and theorematic status.*

Proof. Let $\mathcal{N}_4(X)$ be shifted-normal admissible. By Definition A.34.3, it carries an occurrence class and a backtranslation ledger for all inhabited shifted tags. Class A is handled by witness preservation. Class B is handled by router backtranslation. Class C is handled by

support/payload separation and packed-mask recomputation. Class D is handled by preservation of sidedness and variance. Class E is handled by preservation of readout target, guard, and theorematic status. These are exactly the local proof chain conservation mechanisms. Hence the normal form is conservative over the shifted migration chain. □

A.34.7 archival stage release-final handoff

Definition A.34.10 (archival stage release-final readiness)**.** *The archival stage line is ready for a release-final pass when:*

1. *the shifted namespace is fixed;*
2. *the occurrence ledger is present;*
3. *the reindexed display triples are present;*
4. *physical/readout compatibility is present;*
5. *the governing shifted-namespace normal form is present;*
6. *all non-claims remain explicit.*

Remark A.34.11 (non-claim)**.** *The layer is a normal-form and release-freeze handoff. It does not close OP 1/4 or OP 5(B), does not activate primitive mixed tensors, does not claim Turing-completeness, and does not replace standard physical calculations.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.35 archival stage Release-Final Closure Note

Status. This closure note records the archival stage Active Core as the first shifted release-final state built from the local proof chain migration chain. The release-final state contains the global QAM Hilbert-shift activation, occurrence-ledger audit map, reindexed display layer, physical-readout compatibility check, and governing shifted-namespace normal form. It records historical DOI-bearing status for that layer; current archival stage DOI metadata is governed by the frontmatter Version and DOI record.

A.35.1 Release-release-final state contents

Definition A.35.1 (archival stage release-final content)**.** *The archival stage historical source-hygiene layer consists of the following recorded layers:*

1. *the local pass: global QAM Hilbert-shift activation;*
2. *the local pass: occurrence ledger and shift-audit map;*
3. *the local pass: global reindexed display layer;*
4. *the local pass: physical readout compatibility check;*
5. *the local pass: governing shifted-namespace normal form and release-freeze handoff.*

Definition A.35.2 (archival stage release-final governing rule)**.** *From this release-final state onward, active QAM/router/readout objects are governed by the shifted-namespace normal form unless explicitly marked as historical or pre-shift witness material. Inhabited active tags are read in the shifted namespace, and the reserved tag* 0_{shift} *denotes the empty/no-active-kind marker.*

A.35.2 Release-release-final state conservativity

Theorem A.35.3 (archival stage release-final conservativity)**.** *Every finite active object admitted by the archival stage release-final normal form has an inhabited-component backtranslation that preserves its payload, tensor grammar, readout target, guard record, and theorematic status.*

Proof. The result is the direct combination of the namespace ledger, the local pass occurrence ledger, the display triples, the physical-readout compatibility check, and the governing normal-form theorem. The ledger supplies $\mathrm{B}_4(a) = a - 1$ on inhabited shifted tags. The ledger assigns a role to visible occurrences. The display layer supplies the explicit pre-shift/shifted/backtranslation display form. The compatibility check preserves physical and mathematical readout data. The normal form requires all new active material to be shifted-normal or explicitly historical/pre-shift. Therefore the release-final state is conservative over the prepared migration chain. □

A.35.3 Release-release-final state non-claims

Remark A.35.4 (archival stage release-final non-claims)**.** *The archival stage historical source-hygiene layer does not close OP 1/4 or OP 5(B), does not activate primitive mixed tensors, does not claim Turing-completeness, and does not replace standard QFT/QED/Lifshitz or general-relativistic calculations. It is a global QAM/router/readout language migration and normal-form release release-final state.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.36 Complete Reader-Hygiene and Release-History Records

A.37 archival stage Historical Remark Triage and Reader-Hygiene Cleanup

Status. This archival stage layer is a cleanup and reader-hygiene layer. It does not add a new theorematic claim, does not close OP 1/4 or OP 5(B), and does not change the archival stage global QAM Hilbert-shift. Its purpose is to separate governing shifted statements from historical genealogy remarks.

A.37.1 Triage principle

Definition A.37.1 (Historical remark triage classes)**.** *Historical or version-genealogical remarks in the Active Core are assigned one of the following cleanup roles:*

1. ***H1: governing guard.*** *The remark remains in the Active Core because it prevents overclaim, preserves OP status, separates Shift A from Shift B, or protects tensor/readout grammar.*

2. ***H2: compact genealogy.*** *The remark may remain only in compressed form, usually as a short version-genealogy pointer.*

3. ***H3: witness-only.*** *The remark belongs to the Historical Witness or release genealogy and should not govern the Active Core.*

Definition A.37.2 (Governing statement criterion)**.** *A remark is a governing statement only if it changes how an active shifted object is read, guarded, classified, backtranslated, or status-preserved. A remark that only explains how a release-final state stage was reached is historical genealogy.*

Remark A.37.3 (No deletion of proof memory)**.** *The cleanup does not delete proof memory. It only changes where such memory is allowed to govern. The Active Core governs through the shifted-namespace normal form; the Historical Witness preserves genealogy.*

A.37.2 Active-Core retention rules

Proposition A.37.4 (Active-Core retention rule)**.** *A historical remark should remain in the Active Core only if it satisfies at least one of the following conditions:*

1. *it protects the distinction between local HC-boundary activation and global shifted QAM namespace migration;*

2. *it protects the distinction between tag, support, packed mask, route payload, tensor component, and historical witness value;*

3. *it protects tensor grammar, readout target, guard, residual map, or OP status;*

4. *it points to the Historical Witness as proof memory rather than as a second active theorematic line.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. These four cases correspond exactly to the places where historical context prevents an incorrect active reading. In all other cases, the shifted-namespace normal form already supplies the governing rule. Therefore long release-final state-stage or development-stage remarks may be compressed or moved to the witness layer without changing the mathematics. □

A.37.3 Version-genealogy compression

Definition A.37.5 (Compressed shifted genealogy note)**.** *The archival stage migration chain is summarized as follows:*

historical release line: local HC-boundary activation $0_{\mathrm{mask}} \to 1_{\mathrm{mask}}$,
historical release line: migration map, backtranslation, and concrete ledger,
the local proof line: global shifted namespace activation,
the local proof line: occurrence ledger and shift-audit map,
the local proof line: pre-shift/shifted/backtranslation display triples,
the local proof line: physical-readout compatibility check,
the local proof line: governing shifted-namespace normal form and release-freeze handoff.

This note replaces repeated long-form release-final state-stage narrative in the governing reading of the Active Core.

Remark A.37.6 (Current governing reading)**.** *In the Active Core, shifted is no longer a future release-final state language. It is the current governing namespace for active QAM/router/readout objects. Historical and pre-shift material remains readable through the Historical Witness and the backtranslation ledger.*

A.37.4 Cleanup audit result

Theorem A.37.7 (Historical remark cleanup conservativity)**.** *The archival stage historical remark triage is conservative over archival stage: it does not change the shifted namespace, the backtranslation map, the occurrence classes, the display triples, the physical-readout compatibility checks, or the governing shifted-namespace normal form.*

Proof. The cleanup assigns roles to historical remarks and compresses genealogy into a single note. It does not modify the definitions of

$$\Sigma_4(k) = k + 1, \qquad \mathrm{B}_4(a) = a - 1, \qquad \mathcal{N}_4(X) = (C_X, X_{\mathrm{shift}}, \mathrm{B}_4(X_{\mathrm{shift}}), P_X, G_X, S_X).$$

It also leaves the non-claim guards for OP 1/4 and OP 5(B) unchanged. Hence the mathematical content of archival stage is preserved. □

Remark A.37.8 (archival stage non-claim)**.** *Version 4.0.13 is a release-final cleanup version. It does not add new physics, does not close OP 1/4 or OP 5(B), does not activate primitive mixed tensors, and does not claim Turing-completeness.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.38 archival stage Reader Navigation Cleanup

Status. This archival stage layer is a reader-navigation cleanup. It does not change the shifted shift, the occurrence ledger, display triples, readout compatibility, normal form, OP status, or Historical Witness.

A.38.1 Navigation principle

Definition A.38.1 (Current reader-navigation principle)**.** *The current reader-navigation principle is:*

Active Core governs, *Historical Witness preserves,* *backtranslation connects.*

Thus active shifted statements are read in the shifted-namespace normal form, historical material is read as proof memory, and pre-shift displays are connected to active shifted displays only through explicit backtranslation records.

Definition A.38.2 (Governing-map roles)**.** *The following roles are used for reader navigation:*

1. ***Governing active role:*** *current shifted definitions, theorems, normal forms, guards, and OP-status statements.*
2. ***Routing role:*** *tables, display triples, occurrence classes, and backtranslation ledgers that explain how to read active objects.*
3. ***Genealogy role:*** *historical remarks and witness material that explain how the current line was reached.*

Lemma A.38.3 (Navigation cleanup is conservative)**.** *The archival stage reader-navigation cleanup is conservative over archival stage.*

Proof. The cleanup adds navigation instructions and clarifies the current governing reading. It does not modify

$$\Sigma_4(k) = k + 1, \qquad \mathrm{B}_4(a) = a - 1, \qquad \mathcal{N}_4(X) = (C_X, X_{\mathrm{shift}}, \mathrm{B}_4(X_{\mathrm{shift}}), P_X, G_X, S_X).$$

It also preserves the Historical Witness and leaves all OP-status guards unchanged. Therefore it is conservative over archival stage. □

Remark A.38.4 (archival stage non-claim)**.** *Version 4.0.13 is a navigation release-final cleanup version. It does not add new physics, does not close OP 1/4 or OP 5(B), does not activate primitive mixed tensors, and does not claim Turing-completeness.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.39 archival stage Notation and Symbol Hygiene Cleanup

Status. This archival stage layer is a notation-hygiene cleanup. It does not change the shifted namespace shift, the backtranslation ledger, the occurrence classes, the tensor grammar, the readout compatibility checks, OP status, or the Historical Witness. It records canonical symbol usage for the current shifted line.

A.39.1 Canonical shifted symbol registry

Definition A.39.1 (Canonical shifted symbol registry)**.** *The current shifted Active Core uses the following canonical symbols:*

$\Sigma_4(k) = k+1$	*global shift on inhabited pre-shift tags*	*language/router namespace,*
$\mathrm{B}_4(a) = a-1$	*backtranslation on inhabited shifted tags* $a \geq 1$	*partial inverse,*
$0_{\mathrm{shift}} = \varnothing_{\mathrm{active}}$	*reserved empty/no-active-kind marker*	*not inhabited,*
$0_{\mathrm{mask}} \to 1_{\mathrm{mask}}$	*local HC-boundary activation*	*Shift A,*
$\mathfrak{D}_4(X)$	*pre-shift/shifted/backtranslation display triple*	*reader/audit display,*
$\mathcal{N}_4(X)$	*governing shifted normal-form object*	*active object form.*

Remark A.39.2 (Roman backtranslation symbol)**.** *The backtranslation symbol is written as* B_4*, not as a Greek beta symbol. This avoids confusion with an additional operator, physical beta parameter, or unstated tensor component.*

A.39.2 Disjoint role vocabulary

Definition A.39.3 (Tag, route, support, packed mask, and tensor payload)**.** *The following terms are kept disjoint in the shifted Active Core:*

1. *A* tag *is an inhabited QAM/router label subject to the shifted namespace when active.*
2. *A* route payload *is an ordered finite list of tags or tag-like route entries.*
3. *A* support set *is an unordered finite set of inhabited support entries.*
4. *A* packed mask integer *is the integer encoding of a support set and is not itself shifted as a tag.*
5. *A* tensor-index payload *is a QAM/router payload carried by a two-sided tensor index position.*
6. *A* tensor component *is not automatically a QAM/router tag and is shifted only if explicitly declared as a tag payload.*

Lemma A.39.4 (Packed mask is not a tag)**.** *A packed mask integer is not shifted by* Σ_4 *as an integer.*

Proof. By Definition A.39.3, a packed mask integer encodes a support set. The shifted shift acts on inhabited support entries. The packed integer is then recomputed from the shifted support. Therefore the operation is not $m \mapsto m+1$ on the packed integer. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.39.3 Canonical display forms

Definition A.39.5 (Canonical display forms)**.** *The following display forms are canonical for current shifted cleanup work:*

$$\mathfrak{D}_4(X) = (X_{\mathrm{pre}}, X_{\mathrm{shift}}, \mathrm{B}_4(X_{\mathrm{shift}})),$$

and

$$\mathcal{N}_4(X) = (C_X, X_{\mathrm{shift}}, \mathrm{B}_4(X_{\mathrm{shift}}), P_X, G_X, S_X).$$

Here C_X is the occurrence class, P_X is the payload or readout target, G_X is the guard record, and S_X is the theorematic status record.

Remark A.39.6 (Display triple versus normal form)**.** *The display triple $\mathfrak{D}_4(X)$ is a reader/audit surface. The normal form $\mathcal{N}_4(X)$ is the governing active object form. They should not be used interchangeably.*

A.39.4 Tensor notation hygiene, declared local OP1/OP4 closure, release-final

Definition A.39.7 (Tensor payload hygiene rule)**.** *The shifted shift may act on QAM/router payloads carried inside tensor-index positions. It does not act on sidedness, upper/lower variance, ordinary upper-lower contraction, or explicit same-variance comparison rules.*

Proposition A.39.8 (Tensor hygiene preservation)**.** *The notation-hygiene cleanup preserves the two-sided structural tensor grammar.*

Proof. The cleanup only records the distinction between tag payload, tensor-index position, and tensor component. It does not move indices, change variance, or alter contraction/comparison rules. Therefore the tensor grammar is preserved. □

A.39.5 Notation-hygiene conservativity

Theorem A.39.9 (Notation-hygiene conservativity)**.** *The archival stage notation-hygiene cleanup is conservative over archival stage: it changes no mathematical content of the global shift, backtranslation, occurrence ledger, display triples, readout compatibility, normal form, or OP-status guards.*

Proof. The cleanup introduces a registry and role vocabulary for existing objects. It leaves the definitions of Σ_4, B_4, $\mathfrak{D}_4$, and $\mathcal{N}_4$ unchanged. It also preserves the distinction between 0_{shift} and 0_{mask}, and it does not modify OP-status guards. Hence the cleanup is conservative. □

Remark A.39.10 (archival stage non-claim)**.** *Version 4.0.13 is a notation release-final cleanup version. It does not add new physics, does not close OP 1/4 or OP 5(B), does not activate primitive mixed tensors, and does not claim Turing-completeness.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.40 archival stage Table, Display, and Example Normalization Cleanup

Status. This archival stage layer is a table/display/example normalization cleanup. It does not change the shifted namespace shift, the backtranslation map, the occurrence classes, the tensor grammar, readout compatibility, OP status, or the Historical Witness. It standardizes how examples should be displayed.

A.40.1 Canonical example display contract

Definition A.40.1 (Canonical example display contract)**.** *Every active shifted example that compares an old display with a current display should use the three-column contract*

pre-shift source / shifted display / backtranslation.

Equivalently, the example should be readable as the display triple

$$\mathfrak{D}_4(X) = (X_{\mathrm{pre}},\ X_{\mathrm{shift}},\ \mathrm{B}_4(X_{\mathrm{shift}}))\,.$$

Remark A.40.2 (Display contract is not a theorem)**.** *The display contract is a reader/audit convention. It does not add a proof certificate and does not change theorematic status.*

A.40.2 Normalized route and support examples

pre-shift source	shifted display	backtranslation
tag 5	tag 6	$\mathrm{B}_4(6) = 5$
tag 12	tag 13	$\mathrm{B}_4(13) = 12$
route $\langle 5, 12\rangle$	route $\langle 6, 13\rangle$	route $\langle 5, 12\rangle$
support $\{5, 12\}$	support $\{6, 13\}$	support $\{5, 12\}$

Lemma A.40.3 (Route/support display normalization)**.** *The normalized route and support examples preserve the shifted backtranslation ledger on inhabited entries.*

Proof. Each shifted entry in the displayed route or support is at least 1 and therefore has backtranslation $\mathrm{B}_4(a) = a - 1$. Applying this componentwise returns the pre-shift route or support. □

A.40.3 Normalized packed-mask example

pre-shift source	shifted display	backtranslation
support $S = \{5, 12\}$	support $S^{(4)} = \{6, 13\}$	support $S = \{5, 12\}$
packed mask $m_3(S) = 4128$	packed mask $m_4(S^{(4)}) = 8256$	via support, not by integer subtraction

Proposition A.40.4 (Packed-mask display normalization)**.** *A packed-mask display is normalized only when it separates support migration from packed-integer recomputation.*

Proof. The shifted shift acts on inhabited support entries, not on the packed integer as a tag. Once the support S is shifted to $S^{(4)}$, the packed integer is recomputed:

$$m_4(S^{(4)}) = \sum_{k \in S} 2^{k+1} = 2 \sum_{k \in S} 2^k = 2m_3(S).$$

Thus the normalized display must show support migration separately from the packed integer. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.40.4 Normalized tensor-display examples

pre-shift source	shifted display	backtranslation
${}^{\langle 5,12\rangle}{}_{\langle 2\rangle}\mathcal{T}^{i}{}_{j}$	${}^{\langle 6,13\rangle}{}_{\langle 3\rangle}\mathcal{T}^{i}{}_{j}$	${}^{\langle 5,12\rangle}{}_{\langle 2\rangle}\mathcal{T}^{i}{}_{j}$
${}^{\langle 5,12\rangle}{}_{\langle 2\rangle}\mathcal{T}^{\langle 7\rangle}{}_{\langle 10\rangle}$	${}^{\langle 6,13\rangle}{}_{\langle 3\rangle}\mathcal{T}^{\langle 8\rangle}{}_{\langle 11\rangle}$	${}^{\langle 5,12\rangle}{}_{\langle 2\rangle}\mathcal{T}^{\langle 7\rangle}{}_{\langle 10\rangle}$

Lemma A.40.5 (Tensor-display normalization preserves grammar)**.** *The normalized tensor-display examples preserve sidedness, variance, contraction rules, and same-variance comparison rules.*

Proof. Only QAM/router payloads inside index positions are shifted. No index is moved between front/back positions or between upper/lower variance. Therefore the two-sided tensor grammar is unchanged. □

A.40.5 Example normalization rule

Definition A.40.6 (Example normalization rule)**.** *A shifted example is normalized if it satisfies all of the following:*

1. *it declares whether the displayed object is a tag, route payload, support set, packed mask integer, tensor-index payload, tensor component, or historical witness value;*
2. *it displays the shifted form and the backtranslation when a shifted active tag payload is present;*
3. *it separates support migration from packed-integer recomputation;*
4. *it preserves tensor sidedness and variance in tensor examples;*
5. *it preserves guard and status information in OP-facing examples.*

Theorem A.40.7 (Example-normalization conservativity)**.** *The archival stage table/display/example normalization cleanup is conservative over archival stage.*

Proof. The cleanup standardizes how examples are shown. It does not change the definitions of Σ_4, B_4, $\mathfrak{D}_4$, or $\mathcal{N}_4$. It also leaves tensor grammar, readout guards, OP status, and the Historical Witness unchanged. Therefore it is conservative over archival stage. □

Remark A.40.8 (archival stage non-claim)**.** *Version 4.0.13 is a table/display/example normalization release-final state. It does not add new physics, does not close OP 1/4 or OP 5(B), does not activate primitive mixed tensors, and does not claim Turing-completeness.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.41 archival stage Appendix and Historical Witness Linkage Cleanup

Status. This archival stage layer is an appendix and Historical Witness linkage cleanup. It does not change the shifted namespace shift, the backtranslation map, the occurrence classes, display triples, tensor grammar, readout compatibility, OP status, or the Historical Witness. It clarifies how the Active Core and Historical Witness should be read together.

A.41.1 Split-bundle linkage principle

Definition A.41.1 (Split-bundle linkage principle)**.** *The split bundle is read by the principle*

Active Core governs, Historical Witness preserves, backtranslation connects.

The Active Core contains the current governing shifted language and theorematic status. The Historical Witness preserves proof memory, genealogy, and older readout surfaces. The backtranslation ledger explains how inhabited shifted shifted tags relate to pre-shift tags.

Remark A.41.2 (No second active theory line)**.** *The Historical Witness is not a second active theory line. If a historical statement conflicts with the current shifted-namespace normal form, the Active Core governs the current reading, while the Historical Witness records the genealogy of the older formulation.*

A.41.2 Historical Witness use classes

Definition A.41.3 (Historical Witness use classes)**.** *Historical Witness material is used in one of the following roles:*

1. ***Proof-memory role.*** *The material preserves derivation history, older proof attempts, and reconstruction context.*
2. ***Genealogy role.*** *The material explains why a current shifted object has the form it has.*
3. ***Backtranslation role.*** *The material provides a pre-shift source display that can be related to a current shifted display.*
4. ***Non-governing archive role.*** *The material is retained for auditability but does not govern current active statements.*

Lemma A.41.4 (Witness roles preserve active governance)**.** *The Historical Witness use classes preserve the Active Core as the governing current line.*

Proof. Each use class assigns historical material a support role: proof memory, genealogy, backtranslation, or archive. None of these roles overrides the current shifted-namespace normal form, occurrence ledger, readout guard, or OP-status record. Therefore active governance remains in the Active Core. □

A.41.3 Crosswalk table

Reader question	Active Core answer	Historical Witness use
What is current?	shifted-namespace normal form and current guards	not governing
Where did it come from?	compressed genealogy note	proof-memory/genealogy source
How do old tags read?	$\mathrm{B}_4(a) = a - 1$ on inhabited shifted tags	pre-shift display source
What is the OP status?	current OP-status guard	older attempts remain context
What about older examples?	read only if reintroduced by display triple	archive/backtranslation source

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.41.4 Appendix linkage rule

Definition A.41.5 (Appendix linkage rule)**.** *An appendix or historical block may be linked to the Active Core only by one of the following explicit link types:*

genealogy link, *backtranslation link,* *proof-memory link,* *audit link.*

A link is not a promotion unless the Active Core explicitly supplies the corresponding shifted normal-form object and status guard.

Proposition A.41.6 (Appendix links are not promotions)**.** *An appendix or Historical Witness link does not by itself promote historical material to current active theorematic status.*

Proof. By Definition A.41.5, a link has one of four support roles. None of those roles is a promotion certificate. Promotion to current active status requires an explicit shifted normal-form object and a status guard in the Active Core. Therefore the link alone does not promote historical material. □

A.41.5 Historical tag and example reading

Definition A.41.7 (Historical tag reading)**.** *A tag, route, support, packed mask, or tensor-payload example inside the Historical Witness is not automatically a current shifted object. It becomes current only if the Active Core reintroduces it through*

$$\mathfrak{D}_4(X) = (X_{\mathrm{pre}}, X_{\mathrm{shift}}, \mathrm{B}_4(X_{\mathrm{shift}}))$$

or through the governing normal form

$$\mathcal{N}_4(X) = (C_X, X_{\mathrm{shift}}, \mathrm{B}_4(X_{\mathrm{shift}}), P_X, G_X, S_X).$$

Remark A.41.8 (Historical examples are protected from silent shift)**.** *Historical examples are not silently shifted. They are either preserved as witness material or explicitly connected to the current shifted line by a display triple or normal-form record.*

A.41.6 Historical linkage conservativity

Theorem A.41.9 (Historical Witness linkage conservativity)**.** *The archival stage Historical Witness linkage cleanup is conservative over archival stage: it does not change the current shifted namespace, backtranslation ledger, occurrence classes, display contracts, tensor grammar, readout compatibility, OP status, or Historical Witness content.*

Proof. The cleanup adds link roles and reading rules between the Active Core and Historical Witness. It does not modify the definitions of Σ_4, B_4, $\mathfrak{D}_4$, or $\mathcal{N}_4$. It does not alter any theorematic status record or OP guard. It also does not remove Historical Witness material. Hence the cleanup is conservative over archival stage. □

Remark A.41.10 (archival stage non-claim)**.** *Version 4.0.13 is an appendix and Historical Witness linkage release-final cleanup version. It does not add new physics, does not close OP 1/4 or OP 5(B), does not activate primitive mixed tensors, and does not claim Turing-completeness.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.42 archival stage Clean Reader Bundle and Hygiene-Freeze Candidate

Status. This archival stage layer closes the local cleanup sequence as a clean reader bundle release-final state. It does not add a new proof claim, does not close OP 1/4 or OP 5(B), does not activate primitive mixed tensors, and does not change the archival stage global QAM Hilbert-shift. It records that the cleanup line from archival stage to archival stage has stabilized the reader-facing structure.

A.42.1 Clean reader bundle principle

Definition A.42.1 (Clean reader bundle principle)**.** *A clean shifted reader bundle is a split bundle in which:*

1. *the Active Core is the current governing line;*
2. *the Historical Witness is preserved as proof memory and genealogy;*
3. *active QAM/router/readout objects are read in shifted-namespace normal form;*
4. *historical/pre-shift material is connected only by explicit witness, genealogy, audit, or back-translation links;*
5. *examples and tables use normalized display contracts whenever they compare pre-shift and shifted displays.*

Remark A.42.2 (Clean bundle is not a shorter theory)**.** *The clean reader bundle is not a shortened or weakened theory. It is the same shifted governing structure with improved reader navigation, symbol hygiene, example hygiene, and Historical Witness linkage.*

A.42.2 cleanup line cleanup closure map

Version	Cleanup role	Mathematical status
archival stage	historical remark triage	conservative; no theorematic change
archival stage	reader navigation map	conservative; current reading clarified
archival stage	notation and symbol hygiene	conservative; symbol roles registered
archival stage	table/display/example normalization	conservative; examples standardized
archival stage	Historical Witness linkage	conservative; split-bundle roles clarified
archival stage	clean reader bundle release-final state	conservative; hygiene line frozen locally

A.42.3 Reader-facing invariant checklist

Definition A.42.3 (Reader-facing invariant checklist)**.** *The following invariants are preserved by the cleanup sequence:*

$$\begin{aligned}
&\textit{shift:} && \Sigma_4(k) = k+1,\\
&\textit{backtranslation:} && \mathrm{B}_4(a) = a-1 \quad (a \geq 1),\\
&\textit{reserved zero:} && 0_{\mathrm{shift}} = \varnothing_{\mathrm{active}},\\
&\textit{HC-boundary shift:} && 0_{\mathrm{mask}} \to 1_{\mathrm{mask}},\\
&\textit{display triple:} && \mathfrak{D}_4(X) = (X_{\mathrm{pre}}, X_{\mathrm{shift}}, \mathrm{B}_4(X_{\mathrm{shift}})),\\
&\textit{normal form:} && \mathcal{N}_4(X) = (C_X, X_{\mathrm{shift}}, \mathrm{B}_4(X_{\mathrm{shift}}), P_X, G_X, S_X).
\end{aligned}$$

Lemma A.42.4 (Cleanup sequence preserves the reader-facing invariants)**.** *The archival stage–archival stage cleanup sequence preserves every invariant listed in Definition A.42.3.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. The cleanup sequence adds reader navigation, historical remark triage, notation hygiene, example normalization, and Historical Witness linkage. None of these operations alters the formulas in the invariant checklist. Therefore the reader-facing invariants are preserved. □

A.42.4 Clean reader bundle handoff

Definition A.42.5 (Clean reader bundle handoff)**.** *After archival stage, new work should be separated into one of the following:*

1. ***further cleanup:*** *layout, bibliography, index, navigation, or display polishing without mathematical change;*
2. ***active shifted development:*** *new QAM/router/readout objects written directly in the shifted-namespace normal form;*
3. ***release packaging:*** *DOI insertion, final metadata, and Zenodo release notes.*

Remark A.42.6 (Preferred next mathematical branch)**.** *The preferred next mathematical branch after a clean reader bundle is not more namespace work. It is active use of the shifted-namespace normal form on OP-facing readout packages, while preserving the OP-status guards.*

A.42.5 Clean reader bundle conservativity

Theorem A.42.7 (Clean reader bundle conservativity)**.** *The archival stage historical source-hygiene layer is conservative over the archival stage release final and over the archival stage–archival stage cleanup sequence.*

Proof. The release-final state records the accumulated cleanup discipline without changing the shifted shift, backtranslation ledger, occurrence classes, display triples, tensor grammar, readout compatibility checks, Historical Witness content, or OP-status guards. It only summarizes and freezes the reader-facing cleanup structure. Therefore it is conservative over archival stage and over the intervening cleanup sequence. □

Remark A.42.8 (archival stage non-claim)**.** *Version 4.0.13 is a clean reader bundle release-final state. It does not add new physics, does not close OP 1/4 or OP 5(B), does not activate primitive mixed tensors, and does not claim Turing-completeness.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.43 archival stage Frontmatter, Metadata, and DOI-Record Hygiene Cleanup

Status. This archival stage layer is a frontmatter and metadata hygiene cleanup. It does not change the shifted namespace shift, the backtranslation map, the occurrence classes, display triples, tensor grammar, readout compatibility, OP status, or the Historical Witness. It records a stable rule for title, DOI, version, and release-final state/release wording.

A.43.1 Frontmatter invariants

Definition A.43.1 (Frontmatter invariants)**.** *The following frontmatter invariants are fixed for the cleanup-line:*

1. *the title remains*

   Quantum Sphaera Companion: A Structure-First Mathematical Unfolding;

2. *the Concept DOI remains the concept-level DOI;*
3. *the public archival stage Version DOI remains a prior version DOI for local release-final cleanup versions unless a new Zenodo DOI is explicitly assigned;*
4. *local release-final cleanup versions must not present themselves as release-final DOI-bearing versions;*
5. *the Historical Witness remains separately preserved in the bundle.*

Remark A.43.2 (Candidate wording)**.** *The word "release-final state" is acceptable in local cleanup line cleanup metadata and file names. It should not appear as a mathematical status operator and should not be used to weaken already governing shifted definitions.*

A.43.2 DOI-record hygiene

Definition A.43.3 (DOI-record hygiene rule)**.** *For this release-final cleanup version, the DOI record is read as*

$$\begin{aligned} &\textit{Concept DOI:} && 10.5281/\textit{zenodo}.19210728,\\ &\textit{Prior public version DOI:} && 10.5281/\textit{zenodo}.19805670,\\ &\textit{Version DOI:} && 10.5281/\textit{zenodo}.19811662. \end{aligned}$$

The version DOI was inserted during release-final packaging.

Lemma A.43.4 (DOI hygiene preserves release genealogy)**.** *The DOI-record hygiene rule preserves the public archival stage release genealogy while allowing local release-final cleanup versions to remain unassigned.*

Proof. The concept DOI points to the concept-level record. The prior public version DOI points to the last DOI-bearing release. Since a local release-final cleanup version has no assigned assigned version DOI, no new DOI-bearing release is asserted. Therefore release genealogy remains correct. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.43.3 Metadata role separation

Definition A.43.5 (Metadata role separation)**.** *Version labels, file names, DOI records, and release notes are metadata. They do not define mathematical objects. Mathematical objects must be named by their mathematical roles, such as* Σ_4, B_4, $\mathfrak{D}_4$, *or* $\mathcal{N}_4$, *not by release-file names.*

Proposition A.43.6 (Metadata does not alter theorematic status)**.** *Changing a local cleanup version label from historical release marker archival stage to archival stage does not alter the theorematic status of any shifted object.*

Proof. The version label is metadata. The theorematic status of a shifted object is stored in its governing status record, for example the S_X field in $\mathcal{N}_4(X)$. Updating a cleanup label does not modify that status record. Hence theorematic status is unchanged. □

A.43.4 Frontmatter hygiene checklist

Definition A.43.7 (Frontmatter hygiene checklist)**.** *A clean local release-final cleanup version should satisfy:*

1. *unchanged title;*
2. *visible cleanup version label;*
3. *Concept DOI displayed as concept pointer;*
4. *prior public version DOI displayed as prior DOI;*
5. *no unassigned DOI represented as current DOI;*
6. *Active Core and Historical Witness both present;*
7. *non-claim guards preserved.*

A.43.5 Frontmatter metadata conservativity

Theorem A.43.8 (Frontmatter metadata conservativity)**.** *The archival stage frontmatter/metadata hygiene cleanup is conservative over archival stage.*

Proof. The cleanup records metadata rules and DOI-record hygiene. It does not modify the definitions of Σ_4, B_4, $\mathfrak{D}_4$, or $\mathcal{N}_4$. It does not alter tensor grammar, readout compatibility, Historical Witness content, or OP-status guards. Therefore it is conservative over archival stage. □

Remark A.43.9 (archival stage non-claim)**.** *Version 4.0.13 is a frontmatter and metadata hygiene release-final state. It does not add new physics, does not close OP 1/4 or OP 5(B), does not activate primitive mixed tensors, and does not claim Turing-completeness.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.44 archival stage Bibliography, Citation, and Terminal Record Hygiene Cleanup

Status. This archival stage layer is a bibliography, citation, and terminal-record hygiene cleanup. It does not change the shifted namespace shift, the backtranslation map, the occurrence classes, display triples, tensor grammar, readout compatibility, OP status, or the Historical Witness. It records how bibliographic and terminal records should be read in the split bundle.

A.44.1 Bibliography role principle

Definition A.44.1 (Bibliography role principle)**.** *The bibliography and terminal self-citation records are metadata and scholarly context. They do not define mathematical objects. Current mathematical objects are governed by their definitions in the Active Core, especially*

$$\Sigma_4, \qquad \mathrm{B}_4, \qquad \mathfrak{D}_4, \qquad \mathcal{N}_4.$$

Remark A.44.2 (Citation does not promote theorematic status)**.** *A citation, DOI record, or terminal self-citation can identify a release or source, but it does not promote a historical statement to current theorematic status. Promotion requires a current Active-Core normal-form record and status guard.*

A.44.2 Terminal record hygiene

Definition A.44.3 (Terminal record hygiene rule)**.** *Terminal records should separate:*

1. *concept-level identifiers;*
2. *version-level identifiers;*
3. *prior public version identifiers;*
4. *local cleanup-release-final state identifiers;*
5. *bibliography entries and self-citations.*

No terminal record should be read as a new theorem unless it is explicitly stated as a theorem in the Active Core.

Lemma A.44.4 (Terminal records preserve metadata boundaries)**.** *The terminal record hygiene rule preserves the boundary between metadata and mathematics.*

Proof. The rule assigns identifiers and citations to metadata roles. It explicitly prevents DOI records and self-citations from acting as theorem statements. Therefore metadata remains separate from mathematical content. □

A.44.3 Citation audit contract

Definition A.44.5 (Citation audit contract)**.** *A clean citation state for a local release-final cleanup version requires:*

1. *no undefined citations in the build log;*
2. *no undefined references in the build log;*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

3. *the concept DOI and prior public version DOI to be readable as metadata;*

4. *no unassigned DOI to be presented as a current version DOI;*

5. *the Historical Witness to remain linked as proof memory rather than as a second active bibliography line.*

Remark A.44.6 (Local release-final cleanup versions and DOI restraint)**.** *A local release-final cleanup version may mention the prior DOI and the concept DOI. It should not claim a assigned version DOI until release-final packaging assigns one.*

A.44.4 Bibliography and witness linkage

Definition A.44.7 (Bibliography–witness linkage)**.** *Historical citations and witness-preserved references may be used to reconstruct the genealogy of an argument. They do not override the current shifted-namespace normal form. If a historical citation supports a current statement, the current statement must still be expressed in Active-Core language.*

Proposition A.44.8 (Bibliographic context is not a second governing layer)**.** *Bibliographic context and Historical Witness context do not form a second governing theorematic layer.*

Proof. By Definition A.44.7, historical citations and witness references serve genealogy and reconstruction roles. Current governance remains in the Active Core. Hence bibliographic context is not a second governing layer. □

A.44.5 Bibliography hygiene conservativity

Theorem A.44.9 (Bibliography and citation hygiene conservativity)**.** *The archival stage bibliography/citation/terminal-record hygiene cleanup is conservative over archival stage.*

Proof. The cleanup records how citations, DOI records, terminal records, and witness references should be read. It does not modify the definitions of Σ_4, B_4, $\mathfrak{D}_4$, or $\mathcal{N}_4$. It does not alter tensor grammar, readout compatibility, OP-status guards, or Historical Witness content. Therefore it is conservative over archival stage. □

Remark A.44.10 (archival stage non-claim)**.** *Version 4.0.13 is a bibliography, citation, and terminal-record hygiene release-final state. It does not add new physics, does not close OP 1/4 or OP 5(B), does not activate primitive mixed tensors, and does not claim Turing-completeness.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.45 archival stage Glossary, Terminology, and Reader-Index Hygiene Cleanup

Status. This archival stage layer is a glossary, terminology, and reader-index hygiene cleanup. It does not change the shifted namespace shift, the backtranslation map, the occurrence classes, display triples, tensor grammar, readout compatibility, OP status, citations, metadata, or the Historical Witness. It supplies a compact reader-facing terminology map for the current shifted line.

A.45.1 Terminology hygiene principle

Definition A.45.1 (Terminology hygiene principle)**.** *A reader-facing term is hygienic if it has exactly one active role in the current shifted line, or if all of its roles are explicitly separated by context. Terminology hygiene prevents a word such as "tag", "mask", "witness", or "readout" from silently changing mathematical role.*

Remark A.45.2 (Glossary entries are reader aids)**.** *The glossary entries in this layer are reader aids. They do not replace the formal definitions in the Active Core and do not promote historical material to current theorematic status.*

A.45.2 Compact current-shifted glossary

Term	Current shifted role	Guard against confusion
Active Core	governing current line	not the Historical Witness
Historical Witness	proof memory and genealogy	not a second active theory line
QAM	language/router/readout set layer	not a new physical law by itself
MML	structural/router-machine layer	not a replacement for proof status
MMA	matrix/readout machinery layer	not an OP-closure certificate
HC	Heisenberg compensator/boundary normalizer	not redefined by the global shifted shift
tag	inhabited active QAM/router label	not a packed mask integer
route payload	ordered finite tag/readout route	not an unordered support set
support set	unordered finite support entries	shifted before mask recomputation
packed mask	integer encoding of support	not shifted by adding one
tensor-index payload	QAM/router payload in an index position	not the tensor component itself
readout target	physical/mathematical target of a package	not changed by address migration
guard	non-claim/status protection record	not optional prose decoration
OP status	open/closed theorematic state	not changed by namespace migration

A.45.3 Acronym and abbreviation map

Definition A.45.3 (Acronym map)**.** *The following acronyms are read in the current Active Core as follows:*

QAM *Quantum Analysis Mengenlehre / language and set-readout layer*,
MML *Millennium Matrix / structural router-machine layer*,
MMA *Heisenberg Matrix / matrix-readout machinery layer*,
HC *Heisenberg compensator / boundary normalizer*,
OP *open problem or OP-facing package, depending on context.*

Remark A.45.4 (Acronyms do not carry closure)**.** *An acronym can identify a layer or tool family, but it does not by itself close an OP or certify a physical readout.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.45.4 Reader-index roles

Definition A.45.5 (Reader-index role)**.** *A reader-index role is a stable navigation label attached to a term. The current cleanup line uses four reader-index roles:*

$$\textit{governing}, \qquad \textit{routing}, \qquad \textit{witness}, \qquad \textit{guard}.$$

Governing terms state the active line; routing terms explain transport or display; witness terms preserve genealogy; guard terms protect non-claims and OP status.

Lemma A.45.6 (Reader-index roles prevent role drift)**.** *The reader-index roles prevent glossary terms from silently switching between active, routing, witness, and guard functions.*

Proof. Each role has a specified reading function. If a term is used as a governing term, it must belong to the active line. If it is used as witness material, it does not govern. If it is used as a guard, it protects status. If it is used as routing, it explains display or transport. Thus the role determines the reading and prevents silent drift. □

A.45.5 Terminology conservativity

Theorem A.45.7 (Glossary and terminology hygiene conservativity)**.** *The archival stage glossary/terminology/reader-index hygiene cleanup is conservative over archival stage.*

Proof. The cleanup adds reader-facing terminology maps and index roles. It does not modify the definitions of Σ_4, B_4, $\mathfrak{D}_4$, or $\mathcal{N}_4$. It does not alter tensor grammar, readout compatibility, citation records, metadata, Historical Witness content, or OP-status guards. Therefore it is conservative over archival stage. □

Remark A.45.8 (archival stage non-claim)**.** *Version 4.0.13 is a glossary, terminology, and reader-index hygiene release-final state. It does not add new physics, does not close OP 1/4 or OP 5(B), does not activate primitive mixed tensors, and does not claim Turing-completeness.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.46 archival stage Cleanup Consolidation and Local Release-Final Preparation

Status. This archival stage layer consolidates the local cleanup sequence into one release-release-final state-preparation layer. It does not add new mathematics, does not close OP 1/4 or OP 5(B), does not activate primitive mixed tensors, and does not change the archival stage global QAM Hilbert-shift. It only records that the cleanup sequence has become locally stable enough for a possible small DOI-bearing cleanup release.

A.46.1 Cleanup consolidation principle

Definition A.46.1 (Cleanup consolidation principle)**.** *A cleanup consolidation layer is a reader-facing stabilization layer which collects prior cleanup work without changing the mathematical content. For the current line, it consolidates:*

archival stage historical remark triage,
archival stage reader navigation,
archival stage notation and symbol hygiene,
archival stage table/display/example normalization,
archival stage Historical Witness linkage,
archival stage clean reader bundle,
archival stage frontmatter and DOI-record hygiene,
archival stage bibliography/citation/terminal-record hygiene,
archival stage glossary/terminology/reader-index hygiene.

Remark A.46.2 (Consolidation is not compression by deletion)**.** *The consolidation does not delete proof memory. It leaves the Historical Witness preserved and treats cleanup as reader-facing stabilization rather than content removal.*

A.46.2 Local release-release-final state readiness checklist

Definition A.46.3 (Local cleanup release-release-final state readiness)**.** *The cleanup-line is locally ready for release-release-final state packaging when the following conditions hold:*

1. *the Active Core compiles without undefined references or citations;*
2. *the Historical Witness compiles without undefined references or citations;*
3. *no duplicate labels are reported;*
4. *overfull hbox/vbox warnings are absent in both PDFs;*
5. *the title and concept DOI remain stable;*
6. *the prior public version DOI is recorded as prior, not as a new DOI;*
7. *all non-claim guards remain visible.*

Lemma A.46.4 (Readiness checklist is metadata-level)**.** *The local release-release-final state readiness checklist is a metadata and build-audit condition, not a theorematic closure condition.*

Proof. The checklist speaks about build state, title/DOI metadata, duplicate labels, warnings, and non-claim guards. It does not add proof certificates or change the status record of any mathematical object. Therefore it is metadata-level. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.46.3 Current invariant freeze

Definition A.46.5 (Current cleanup invariant freeze)**.** *Throughout the cleanup sequence, the following invariants remain fixed:*

$$\begin{aligned}
&\Sigma_4(k) = k+1,\ \mathrm{B}_4(a) = a-1 \quad (a \geq 1),\\
&0_{\mathrm{shift}} = \varnothing_{\mathrm{active}},\ 0_{\mathrm{mask}} \to 1_{\mathrm{mask}},\\
&\mathfrak{D}_4(X), \qquad \mathcal{N}_4(X),\\
&\textit{OP 1/4 open}, \quad \textit{OP 5(B) open}.
\end{aligned}$$

Proposition A.46.6 (Cleanup consolidation preserves the invariant freeze)**.** *The archival stage cleanup consolidation preserves the invariant freeze of Definition A.46.5.*

Proof. The consolidation layer only records the cleanup history and readiness conditions. It does not modify the shift, backtranslation, reserved zero, HC-boundary shift, display triple, normal form, or OP-status guards. Hence all listed invariants are preserved. □

A.46.4 Release packaging boundary

Definition A.46.7 (Release packaging boundary)**.** *Release packaging begins only when a assigned Zenodo version DOI is explicitly assigned. Before that assignment, archival stage remains a local cleanup consolidation release-final state with the prior public version DOI recorded only as prior.*

Remark A.46.8 (What a final cleanup release would do)**.** *A final cleanup release would not add new theory. It would insert the assigned version DOI, normalize release-final file names, regenerate audit reports, checksums, and the Zenodo description, and preserve the Active Core/Historical Witness split.*

A.46.5 Cleanup consolidation conservativity

Theorem A.46.9 (Cleanup consolidation conservativity)**.** *The archival stage cleanup consolidation is conservative over archival stage release-final and over the local archival stage–archival stage cleanup sequence.*

Proof. The consolidation summarizes the cleanup line and records release-release-final state readiness criteria. It does not alter any current shifted-namespace mathematical definition, tensor grammar, readout compatibility check, citation role, metadata role, Historical Witness content, or OP-status guard. Therefore it is conservative over archival stage and over archival stage–archival stage. □

Remark A.46.10 (archival stage non-claim)**.** *Version 4.0.13 is a cleanup consolidation release-final state. It does not add new physics, does not close OP 1/4 or OP 5(B), does not activate primitive mixed tensors, and does not claim Turing-completeness.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.47 archival stage Label, Reference, and Counter Hygiene Cleanup

Status. This archival stage layer is a label, reference, and counter hygiene cleanup. It does not change the shifted namespace shift, the backtranslation map, the occurrence classes, display triples, tensor grammar, readout compatibility, OP status, citations, metadata, or the Historical Witness. It records the audit rule for labels, cross-references, theorem counters, and reader-visible navigation.

A.47.1 Label hygiene principle

Definition A.47.1 (Label hygiene principle)**.** *A label is hygienic if it is unique, stable under cleanup edits, and semantically aligned with the object it names. In the cleanup-line, labels should identify the local role of an object, for example section, definition, lemma, theorem, proposition, remark, table, or audit block.*

Remark A.47.2 (Labels are addresses, not mathematical content)**.** *A LaTeX label is a document address. It does not define the mathematical object itself. Mathematical content remains in the corresponding definition, lemma, theorem, proposition, or normal-form record.*

A.47.2 Reference hygiene contract

Definition A.47.3 (Reference hygiene contract)**.** *A clean local release-final cleanup version should satisfy the following reference contract:*

1. *no undefined references in the build log;*
2. *no undefined citations in the build log;*
3. *no multiply defined labels in the build log;*
4. *no theorem statement should depend on an unresolved reference;*
5. *historical references should not override current Active-Core status.*

Lemma A.47.4 (Reference hygiene is build-level, not theorematic)**.** *Reference hygiene is a build-level and reader-navigation condition, not a new theorematic closure condition.*

Proof. The reference contract concerns build-log state and reader addressability. It does not change any mathematical definition, readout target, guard, or OP-status record. Therefore it is build-level rather than theorematic. □

A.47.3 Counter and theorem-environment hygiene

Definition A.47.5 (Counter hygiene rule)**.** *Theorem, lemma, proposition, definition, and remark counters are reader-facing navigation devices. Their role is to keep the document addressable. A cleanup pass should not use counter renumbering to imply a change of theorematic status.*

Proposition A.47.6 (Counter changes are not status changes)**.** *A reader-visible counter or page-number change caused by cleanup is not a change of theorematic status.*

Proof. The theorematic status is recorded by the mathematical statement and its guard or status record, not by its page number or counter position. Cleanup may add reader-facing sections and thereby shift page numbers. Such shifts do not alter status records. Therefore counter or page-number changes are not status changes. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.47.4 Cross-reference role separation

Definition A.47.7 (Cross-reference roles)**.** *Cross-references are read in one of four roles:*

governing reference, *routing reference*, *witness reference*, *audit reference.*

A governing reference points to current Active-Core content. A routing reference explains transport, display, or navigation. A witness reference points to proof memory or genealogy. An audit reference points to build, metadata, or hygiene state.

Remark A.47.8 (Witness references remain non-governing)**.** *A reference to the Historical Witness remains a witness reference unless the Active Core explicitly reintroduces the content through a current shifted display triple, normal form, or status record.*

A.47.5 Label/reference hygiene conservativity

Theorem A.47.9 (Label/reference/counter hygiene conservativity)**.** *The archival stage label/reference/counter hygiene cleanup is conservative over archival stage.*

Proof. The cleanup records build-addressing and navigation rules. It does not modify the definitions of Σ_4, B_4, $\mathfrak{D}_4$, or $\mathcal{N}_4$. It does not alter tensor grammar, readout compatibility, citation records, metadata, Historical Witness content, or OP-status guards. Therefore it is conservative over archival stage. □

Remark A.47.10 (archival stage non-claim)**.** *Version 4.0.13 is a label, reference, and counter hygiene release-final state. It does not add new physics, does not close OP 1/4 or OP 5(B), does not activate primitive mixed tensors, and does not claim Turing-completeness.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.48 archival stage Layout, TOC, and Render Hygiene Cleanup

Status. This archival stage layer is a layout, table-of-contents, and render hygiene cleanup. It does not change the shifted namespace shift, the backtranslation map, the occurrence classes, display triples, tensor grammar, readout compatibility, OP status, citations, metadata, labels, counters, or the Historical Witness. It records the visual audit rule for the split bundle.

A.48.1 Layout hygiene principle

Definition A.48.1 (Layout hygiene principle)**.** *A local release-final cleanup version is layout-hygienic if its generated PDFs are readable as rendered documents: no text is visibly clipped, no major text blocks overlap, table headings remain legible, and reader-navigation pages are visually usable.*

Remark A.48.2 (Layout is not theorematic content)**.** *Layout hygiene improves readability and auditability. It does not change definitions, lemmas, theorems, OP status, or physical readout status.*

A.48.2 TOC readability contract

Definition A.48.3 (TOC readability contract)**.** *The table of contents is reader-hygienic if section and subsection entries are readable, numbering boxes do not collide with titles, and cleanup-version headings remain navigable. Long cleanup headings may be verbose in the source but must remain readable in the rendered PDF.*

Lemma A.48.4 (TOC readability is a navigation condition)**.** *The TOC readability contract is a navigation condition, not a mathematical condition.*

Proof. The table of contents addresses document locations. It does not define or modify mathematical objects. Therefore improving TOC readability changes only navigation. □

A.48.3 Render audit contract

Definition A.48.5 (Render audit contract)**.** *A local render audit should include at least the title page, early navigation pages, at least one representative middle page, and terminal pages of both the Active Core and Historical Witness. The purpose is to detect visible clipping, overlap, unreadable page headings, and severe table breakage.*

Remark A.48.6 (Contact sheet role)**.** *The render contact sheet is an audit artifact. It is not part of the mathematical proof. It gives a compact visual sample of the generated PDFs.*

A.48.4 Warning hygiene

Definition A.48.7 (Warning hygiene contract)**.** *For the current cleanup line, the preferred local build state is:*

$$\begin{aligned} &\textit{undefined references} &&= 0,\\ &\textit{undefined citations} &&= 0,\\ &\textit{multiply-defined labels} &&= 0,\\ &\textit{overfull hbox warnings} &&= 0,\\ &\textit{overfull vbox warnings} &&= 0. \end{aligned}$$

This is a local document-hygiene target rather than a theorematic claim.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proposition A.48.8 (Warning hygiene is conservative)**.** *Achieving the preferred warning hygiene state does not alter mathematical content.*

Proof. Warnings and references concern build quality, cross-reference resolution, and rendering. They do not change the definitions of Σ_4, B_4, $\mathfrak{D}_4$, or $\mathcal{N}_4$, and they do not modify OP-status guards. Therefore warning hygiene is conservative. □

A.48.5 Layout/render hygiene conservativity

Theorem A.48.9 (Layout/TOC/render hygiene conservativity)**.** *The archival stage layout/TOC/render hygiene cleanup is conservative over archival stage.*

Proof. The cleanup records visual audit, TOC readability, and warning-hygiene rules. It does not modify any current shifted-namespace mathematical definition, tensor grammar, readout compatibility check, citation role, metadata role, label role, Historical Witness content, or OP-status guard. Therefore it is conservative over archival stage. □

Remark A.48.10 (archival stage non-claim)**.** *Version 4.0.13 is a layout, TOC, and render hygiene release-final state. It does not add new physics, does not close OP 1/4 or OP 5(B), does not activate primitive mixed tensors, and does not claim Turing-completeness.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.49 archival stage Source, Build, and Artifact Reproducibility Hygiene Cleanup

Status. This archival stage layer is a source, build, and artifact reproducibility hygiene cleanup. It does not change the shifted namespace shift, the backtranslation map, the occurrence classes, display triples, tensor grammar, readout compatibility, OP status, citations, metadata, labels, counters, layout contract, or the Historical Witness. It records how source files, generated PDFs, manifests, checksums, and bundles should be interpreted.

A.49.1 Source reproducibility principle

Definition A.49.1 (Source reproducibility principle)**.** *A local release-final cleanup version is source-reproducible if the supplied TEX files are sufficient to rebuild the Active Core and Historical Witness PDFs in the same logical document state, modulo ordinary build-environment differences such as PDF timestamps or engine-specific metadata.*

Remark A.49.2 (Source is governing over generated artifacts)**.** *For local cleanup work, the TEX source is the editable governing artifact. The PDFs are rendered outputs. If a rendered artifact and source artifact appear to disagree, the source and build log must be inspected before any mathematical interpretation is changed.*

A.49.2 Artifact classes

Definition A.49.3 (Artifact classes)**.** *The local cleanup workflow separates artifacts into the following classes:*

1. ***source artifacts:*** *TEX files and source-side scan reports;*
2. ***rendered artifacts:*** *Active Core and Historical Witness PDFs;*
3. ***audit artifacts:*** *build logs, audit reports, render contact sheets, and scan reports;*
4. ***packaging artifacts:*** *manifest files, checksum files, full bundles, and TEX-only ZIPs.*

Lemma A.49.4 (Artifact classes prevent role drift)**.** *The artifact classes prevent rendered, audit, and packaging artifacts from being mistaken for new mathematical content.*

Proof. Each artifact class has an assigned role. Source artifacts contain editable content; rendered artifacts show compiled output; audit artifacts document quality checks; packaging artifacts support transfer and verification. None of the non-source classes is a new theorematic layer. Therefore role drift is prevented. □

A.49.3 Manifest and checksum hygiene

Definition A.49.5 (Manifest and checksum hygiene rule)**.** *A local cleanup bundle should contain a manifest and checksum file. The manifest records which files belong to the package. The checksum file records file integrity. Neither manifest nor checksum file changes mathematical content.*

Proposition A.49.6 (Checksums certify files, not theorems)**.** *A checksum certifies the byte identity of a file, not the truth of a theorem.*

Proof. A checksum is a hash of file bytes. It can detect whether a file has changed, but it does not inspect or prove mathematical statements. Therefore it certifies file integrity only. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.49.4 Build provenance contract

Definition A.49.7 (Build provenance contract)**.** *A clean local build record should state:*

1. *source TEX file names;*
2. *output PDF file names;*
3. *page counts;*
4. *undefined reference and citation counts;*
5. *duplicate-label counts;*
6. *overfull hbox and vbox counts;*
7. *fatal or emergency error counts;*
8. *whether a render contact sheet was produced.*

Remark A.49.8 (Build provenance is audit support)**.** *Build provenance supports reproducibility and audit. It does not replace the mathematical proof and does not close any OP.*

A.49.5 TEX-only bundle contract

Definition A.49.9 (TEX-only bundle contract)**.** *A TEX-only bundle is intended for manual integration and local rebuilding. It should contain the Active Core source, the Historical Witness source, and the minimal supporting hygiene records needed to identify the release-final state. It should not be treated as a DOI-bearing release unless release-final packaging assigns a assigned version DOI.*

Lemma A.49.10 (TEX-only bundle preserves manual-integration workflow)**.** *The TEX-only bundle contract preserves the user's manual-integration workflow.*

Proof. The contract supplies directly editable TEX files and minimal identifiers. It does not require repository-side mutation, automatic patch application, or a new DOI. Hence it preserves manual integration. □

A.49.6 Reproducibility conservativity

Theorem A.49.11 (Source/build/artifact hygiene conservativity)**.** *The archival stage source/build/artifact reproducibility cleanup is conservative over archival stage.*

Proof. The cleanup records artifact roles, manifest/checksum rules, build provenance, and TEX-only bundle expectations. It does not modify the definitions of Σ_4, B_4, $\mathfrak{D}_4$, or $\mathcal{N}_4$. It does not alter tensor grammar, readout compatibility, citation records, metadata roles, labels, counters, layout rules, Historical Witness content, or OP-status guards. Therefore it is conservative over archival stage. □

Remark A.49.12 (archival stage non-claim)**.** *Version 4.0.13 is a source, build, and artifact reproducibility hygiene release-final state. It does not add new physics, does not close OP 1/4 or OP 5(B), does not activate primitive mixed tensors, and does not claim Turing-completeness.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.50 archival stage Local Release-Readiness, Upload, and Final-Packaging Hygiene Cleanup

Status. This archival stage layer is a local release-readiness and upload-hygiene cleanup. It records the assigned version DOI 10.5281/zenodo.19811662, does not change the shifted namespace shift, does not change the Historical Witness, and does not change any OP status. It records which artifacts would be used for a small DOI-bearing cleanup release if a new Zenodo version DOI is later assigned.

A.50.1 Local release-readiness principle

Definition A.50.1 (Local release-readiness principle)**.** *A local release-final cleanup version is release-ready only in the limited sense that its source, PDF, audit, manifest, checksum, and bundle artifacts are coherent and can be promoted to release-final packaging after the version DOI 10.5281/zenodo.19811662 is assigned. Release-readiness is not a theorematic status and is not an OP-closure certificate.*

Remark A.50.2 (release-final assigned in this layer)**.** *This release-final layer records the assigned Zenodo version DOI 10.5281/zenodo.19811662. The prior public version DOI remains the archival stage release DOI until explicit release-final packaging assigns a assigned version DOI.*

A.50.2 Upload artifact selection

Definition A.50.3 (Recommended upload artifact set)**.** *For a small cleanup release, the recommended upload artifact set is:*

1. *Active Core PDF;*
2. *Historical Witness PDF;*
3. *full release bundle ZIP;*
4. *TEX-only ZIP, if source-side manual integration is desired;*
5. *release notes or Zenodo description text;*
6. *checksum file or checksum-bearing manifest.*

Remark A.50.4 (Candidate artifacts versus release-final artifacts)**.** *Candidate artifacts should not be uploaded as release-final artifacts unless their file names, metadata, DOI records, manifests, checksums, and Zenodo description have been normalized for release-final.*

A.50.3 Zenodo description draft contract

Definition A.50.5 (Zenodo description draft contract)**.** *A Zenodo description draft for a local release-final cleanup version should state:*

1. *the document title;*
2. *the cleanup version;*
3. *that the update is reader/document hygiene, not new theory;*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

4. *the concept DOI;*

5. *the prior public version DOI;*

6. *that the assigned version DOI is assigned in this release-final packaging as 10.5281/zenodo.19811662;*

7. *the non-claims for OP 1/4, OP 5(B), primitive mixed tensors, and Turing-completeness.*

Lemma A.50.6 (Description draft is metadata)**.** *A Zenodo description draft is metadata and does not change mathematical content.*

Proof. The description draft summarizes the package, version state, DOI state, and non-claims. It does not alter definitions, proofs, normal forms, readout guards, or OP-status records. Hence it is metadata only. □

A.50.4 Final-packaging boundary

Definition A.50.7 (Final-packaging boundary)**.** *Final packaging begins only after a assigned Zenodo version DOI is provided. The final-packaging pass may then:*

1. *replace release-final state wording by release-final wording where appropriate;*

2. *insert the assigned version DOI in metadata, manifest, notes, and Zenodo description;*

3. *normalize file names to release-final names;*

4. *rebuild the Active Core and Historical Witness PDFs;*

5. *regenerate audit reports, checksums, and bundles.*

Proposition A.50.8 (Final packaging is conservative when DOI-only)**.** *If final packaging only inserts DOI and release-final metadata, rebuilds artifacts, and regenerates audits, then it is conservative over the local release-final cleanup version.*

Proof. DOI insertion and release-final artifact regeneration are metadata and build operations. They do not change Σ_4, B_4, $\mathfrak{D}_4$, $\mathcal{N}_4$, tensor grammar, readout guards, Historical Witness content, or OP-status records. Therefore such packaging is conservative. □

A.50.5 Release-readiness conservativity

Theorem A.50.9 (Release-readiness and upload-hygiene conservativity)**.** *The archival stage release-readiness/upload-hygiene cleanup is conservative over archival stage.*

Proof. The cleanup records upload artifact selection, Zenodo description drafting, and the final-packaging boundary. It does not modify current shifted mathematical definitions, tensor grammar, readout compatibility checks, artifact roles, build rules, Historical Witness content, or OP-status guards. Therefore it is conservative over archival stage. □

Remark A.50.10 (archival stage non-claim)**.** *Version 4.0.14 is a local release-readiness and upload-hygiene release-final state. It does not add new physics, does not close OP 1/4 or OP 5(B), does not activate primitive mixed tensors, does not claim Turing-completeness, and records the assigned Zenodo version DOI 10.5281/zenodo.19811662.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.51 archival stage Local Release-Final Freeze and DOI-Request Packet Hygiene

Status. This archival stage layer freezes the local cleanup line as a release-release-final state preparation state. It records the assigned version DOI 10.5281/zenodo.19811662, does not alter the shifted namespace shift, does not modify the Historical Witness, and does not change any OP status. Its only purpose is to mark the local cleanup sequence as ready for a possible DOI-request and release-final packaging pass.

A.51.1 Local release-release-final state freeze principle

Definition A.51.1 (Local release-release-final state freeze principle)**.** *A release-final state is a document-control state in which the source, rendered PDFs, audit reports, manifests, checksums, and upload description are internally coherent and should not receive new mathematical content before a release-final DOI pass.*

Remark A.51.2 (Freeze is not mathematical closure)**.** *A release-release-final state freeze is a workflow boundary. It is not a proof of OP closure, not a physical promotion criterion, and not a replacement for any status record inside the shifted-namespace normal form.*

A.51.2 DOI-request packet

Definition A.51.3 (DOI-request packet)**.** *A DOI-request packet for a small cleanup release should contain:*

1. *the frozen Active Core PDF;*
2. *the frozen Historical Witness PDF;*
3. *the full bundle ZIP;*
4. *the TEX-only ZIP;*
5. *the release-final Zenodo description;*
6. *the audit report;*
7. *the manifest and checksum file.*

The packet is still local until a assigned Zenodo version DOI is assigned.

A.51.3 Allowed release-final pass

Definition A.51.4 (Allowed release-final pass)**.** *After a assigned Zenodo version DOI is supplied, the allowed release-final pass may:*

1. *insert the version DOI in frontmatter, notes, manifest, checksums, audit metadata, and Zenodo description;*
2. *replace local release-final state/freeze wording by release-final wording where appropriate;*
3. *rename files from local/freeze/release-final state names to release-final names;*
4. *rebuild the Active Core and Historical Witness PDFs;*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

5. *regenerate the full bundle, TEX-only ZIP, checksums, audit report, and final Zenodo description.*

It may not add new theorems, change OP status, or alter shifted normal-form content.

Proposition A.51.5 (DOI pass is conservative under the allowed boundary)**.** *A release-final pass satisfying Definition A.51.4 is conservative over the local archival stage freeze.*

Proof. The allowed pass is restricted to DOI insertion, wording normalization, file renaming, rebuilding, and audit regeneration. It explicitly excludes new theorems, OP-status changes, and shifted normal-form changes. Therefore it is conservative over the frozen local release-final state. □

A.51.4 Frozen non-claim boundary

Definition A.51.6 (Frozen non-claim boundary)**.** *The release-final state preserves the following non-claims:*

OP 1/4	*remains open*,
OP 5(B)	*remains open*,
primitive mixed tensors	*not activated*,
Turing-completeness	*not claimed*,
standard physical calculations	*not replaced.*

A.51.5 Freeze conservativity

Theorem A.51.7 (Local release-release-final state freeze conservativity)**.** *The archival stage release-final state is conservative over the archival stage–archival stage cleanup sequence and over the public archival stage release-final mathematical content.*

Proof. The freeze adds document-control and DOI-request packet rules. It does not modify Σ_4, B_4, $\mathfrak{D}_4$, $\mathcal{N}_4$, tensor grammar, readout compatibility, artifact roles, build rules, Historical Witness content, or OP-status guards. Therefore it is conservative over the cleanup sequence and over the public archival stage mathematical content. □

Remark A.51.8 (archival stage non-claim)**.** *Version 4.0.15 is a release-final state. It does not add new physics, does not close OP 1/4 or OP 5(B), does not activate primitive mixed tensors, does not claim Turing-completeness, and does not assign a new Zenodo version DOI.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.52 archival stage Release-Final DOI and Audit Hygiene

Status. This archival stage layer is the final local audit and release-final hygiene layer. It records the assigned version DOI 10.5281/zenodo.19811662, does not change any mathematical content, does not modify the Historical Witness, and does not change any OP status. It only checks and records the exact boundary between the frozen local release-final state and a later DOI-bearing release-final pass.

A.52.1 Final local audit principle

Definition A.52.1 (Final local audit principle)**.** *A final local audit is a pre-release technical verification state. It verifies source presence, PDF build status, audit-report availability, checksum availability, release-final Zenodo description availability, and render-sample availability before a DOI-bearing release-final pass is allowed.*

Remark A.52.2 (Release-final packaging record)**.** *The release-final audit is not a Zenodo publication and is not a DOI-bearing release. It is a local check that the next pass can be a release-final packaging pass after a version DOI 10.5281/zenodo.19811662 is recorded.*

A.52.2 Release-final checklist

Definition A.52.3 (Release-final checklist)**.** *The release-final audit is successful if the following checks hold:*

1. *Active Core source exists;*
2. *Historical Witness source exists;*
3. *Active Core PDF builds with zero fatal errors;*
4. *Historical Witness PDF builds with zero fatal errors;*
5. *undefined references, undefined citations, duplicate labels, overfull hbox warnings, and overfull vbox warnings are all zero;*
6. *release-final Zenodo description exists;*
7. *manifest and checksum files are generated;*
8. *render contact sheet is generated;*
9. *non-claim boundary remains explicit.*

A.52.3 Release-final packaging record

Definition A.52.4 (Release-final packaging record)**.** *The only missing input for a release-final pass after this release-final audit is a new Zenodo version DOI. Once that DOI is supplied, the next pass should be a metadata-only release-finalization pass unless the user explicitly requests additional cleanup first.*

Proposition A.52.5 (Single-input release-final readiness)**.** *If the release-final checklist holds, then the local release-final state is release-final ready up to the supplied Zenodo version DOI 10.5281/zenodo.19811662.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. The release-final checklist verifies the required source, build, audit, packaging, render, and non-claim conditions. The release-final input contract states that the remaining missing datum is the assigned version DOI. Therefore, once that DOI is supplied, release-final packaging can proceed without changing mathematical content. □

A.52.4 Final local non-claim boundary

Definition A.52.6 (Final local non-claim boundary)**.** *The final local audit preserves the following non-claims:*

OP 1/4	*remains open*,
OP 5(B)	*remains open*,
primitive mixed tensors	*not activated*,
Turing-completeness	*not claimed*,
standard physical calculations	*not replaced*,
Zenodo version DOI	*10.5281/zenodo.19811662.*

A.52.5 Release-final audit conservativity

Theorem A.52.7 (Final local audit release-final conservativity)**.** *The archival stage release-final audit state is conservative over the archival stage–archival stage cleanup sequence and over the public archival stage release-final mathematical content.*

Proof. The release-final audit adds audit and release-final readiness conditions only. It does not modify Σ_4, B_4, $\mathfrak{D}_4$, $\mathcal{N}_4$, tensor grammar, readout compatibility, artifact roles, build rules, Historical Witness content, or OP-status guards. Therefore it is conservative over the cleanup sequence and over the public archival stage mathematical content. □

Remark A.52.8 (archival stage non-claim)**.** *Version 4.0.16 is a release-final audit state. It does not add new physics, does not close OP 1/4 or OP 5(B), does not activate primitive mixed tensors, does not claim Turing-completeness, and records the assigned Zenodo version DOI 10.5281/zenodo.19811662.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.53 archival stage Release-Final DOI and Packaging Record

Status. This section records the release-final DOI and packaging pass for Version 4.0.16. The pass is metadata-only: it inserts the assigned Zenodo version DOI, normalizes release-final artifacts, rebuilds the PDFs, regenerates the audit records, and preserves all mathematical non-claims.

A.53.1 DOI record

Definition A.53.1 (Release-final DOI record)**.** *The release-final DOI record is:*

Concept DOI:	10.5281/*zenodo*.19210728,
Prior public version DOI:	10.5281/*zenodo*.19805670,
Version DOI:	10.5281/*zenodo*.19811662.

Remark A.53.2 (Version DOI is metadata)**.** *The version DOI is a release and citation identifier. It is not a mathematical operator, not a new object in the shifted-namespace normal form, and not an OP-closure certificate.*

A.53.2 Release-final conservativity

Theorem A.53.3 (Release-final DOI/pass conservativity)**.** *The archival stage release-final DOI and packaging pass is conservative over the archival stage local audit state.*

Proof. The pass inserts the version DOI, normalizes release-final wording and file names, rebuilds the PDFs, and regenerates audit and checksum artifacts. It does not modify Σ_4, B_4, $\mathfrak{D}_4$, $\mathcal{N}_4$, tensor grammar, readout compatibility, artifact roles, build rules, Historical Witness content, or OP-status guards. Therefore the pass is conservative. □

Remark A.53.4 (archival stage release-final non-claim)**.** *Version 4.0.16 release-final does not add new physics, does not close OP 1/4 or OP 5(B), does not activate primitive mixed tensors, does not claim Turing-completeness, and does not replace standard QFT/QED/Lifshitz or general-relativistic calculations.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.54 OP Selectors, Routes and Mixed-Readout Ledgers

A.55 OP-Facing Diagnostic Appendix

Status. This working layer starts the post-archival stage OP-facing phase. It does not rewrite the Companion into Python, does not replace MML/MMA/QAM, and does not claim that executable diagnostics are proof certificates. The Python files listed here are diagnostic shadows of selected QAM/MML/MMA/ZFS-style objects. They are included to help test route shifting, backtranslation, route/support separation, packed-mask recomputation, guard retention, residual localization, and proof-state provenance before OP 1/4 and OP 5(B) are attacked in the formal text.

A.55.1 Diagnostic-code status rule

Definition A.55.1 (Diagnostic-code status rule)**.** *A diagnostic Python file in this appendix is an executable audit aid. It may serialize or test selected shadows of QAM, MML, MMA, and ZFS-style records, but it is not itself a formal definition, not a proof certificate, not a physical simulation, and not a replacement for the paper's mathematical language.*

Remark A.55.2 (Why Python appears only here)**.** *Python appears here only because the current shifted language is rigid enough that some role separations can be tested mechanically. The MML remains the mathematical router/intertwiner language in the paper. Python only records diagnostic encodings of selected finite examples.*

A.55.2 Diagnostic files included in the local bundle

File	Diagnostic role
`qam_normal_form.py`	builds a small executable shadow of $N_4(X)$ with route, guards, and status
`route_mask_decoder.py`	separates route, support, and packed mask; verifies recomputation after shift
`mml_op_router.py`	records OP-facing MML route surfaces and obstruction kind
`mma_readout_residual.py`	records readout residuals and promotion guards
`zfs_snapshot_manifest.py`	records proof-state provenance/checksum snapshots
`run_all_diagnostics.py`	runs the diagnostic examples and writes a JSON report

A.55.3 QAM normal-form diagnostic excerpt

```
#!/usr/bin/env python3
"""
QAM normal-form diagnostic encoding.

This file is not a proof engine and not an implementation of QAM as a new
foundation. It serializes selected shifted/QAM roles
can be inspected consistently.
"""
from __future__ import annotations

from dataclasses import asdict, dataclass
from enum import Enum
import json
from typing import Tuple
```

```
class Status(str, Enum):
 OPEN = "open"
 PREPARED = "prepared"
 PROMOTED = "promoted"


class Guard(str, Enum):
 OP_OPEN = "op_open"
 NO_PHYSICAL_CLAIM = "no_physical_claim"
 NEEDS_SELECTOR = "needs_selector"
 NEEDS_BOUNDARY_PROMOTION = "needs_boundary_promotion"


def sigma_4(k: int) -> int:
 """shifted Hilbert shift on inhabited pre-shift tags."""
 if k < 0:
 raise ValueError("invalid pre-shift tag")
 return k + 1


def back_4(a: int) -> int:
 """Backtranslation from inhabited shifted tags."""
 if a < 1:
 raise ValueError("0_shift has no backtranslation")
 return a - 1
```

A.55.4 Route/mask diagnostic excerpt

```
#!/usr/bin/env python3
"""
Route/support/packed-mask diagnostic checks.

The key rule is that a packed mask is recomputed from shifted support. It is
not obtained by adding 1 to an integer mask.
"""
from __future__ import annotations

from dataclasses import dataclass, asdict
import json
from typing import FrozenSet, Tuple


def sigma_4(k: int) -> int:
 if k < 0:
 raise ValueError("invalid pre-shift tag")
 return k + 1


def back_4(a: int) -> int:
```

```
 if a < 1:
 raise ValueError("0_shift has no backtranslation")
 return a - 1


def packed_mask_from_support(support: FrozenSet[int]) -> int:
 return sum(2 ** k for k in support)


@dataclass(frozen=True)
class RouteMaskRecord:
 pre_route: Tuple[int, ...]
 shifted_route: Tuple[int, ...]
 support: FrozenSet[int]
 shifted_support: FrozenSet[int]
```

A.55.5 MML and MMA diagnostic excerpts

```
#!/usr/bin/env python3
"""
MML_OP diagnostic router.

This is an executable shadow of an MML OP-facing route record. It does not
define MML and does not prove an OP.
"""
from __future__ import annotations

from dataclasses import asdict, dataclass
from enum import Enum
import json
from typing import Tuple


class Branch(str, Enum):
 HORIZON = "horizon"
 BOUNDARY = "boundary"


class ObstructionKind(str, Enum):
 MISSING_SELECTOR = "missing_selector"
 MISSING_BOUNDARY_PROMOTION = "missing_boundary_promotion"
 UNKNOWN = "unknown"


@dataclass(frozen=True)
class MMLOPRoute:
 name: str
 branch: Branch
 pre_route: Tuple[int, ...]
 shifted_route: Tuple[int, ...]
```

© 2026 Marc Brendecke. Preprint — not peer reviewed.

```
#!/usr/bin/env python3
"""
MMA_OP readout-residual diagnostic.

This file gives a small data shape for readout residuals. It does not compute
physical predictions; it records which residual/guard still blocks promotion.
"""
from __future__ import annotations

from dataclasses import asdict, dataclass
from enum import Enum
import json
from typing import Optional


class PromotionStatus(str, Enum):
 BLOCKED = "blocked"
 READY_FOR_REVIEW = "ready_for_review"
 PROMOTED = "promoted"


@dataclass(frozen=True)
class MMAReadoutResidual:
 name: str
 readout_target: str
 hc_boundary_condition: str
 comparison_selector: Optional[str]
 residual_surface: str
 promotion_guard: str
 status: PromotionStatus
```

A.55.6 ZFS-style provenance diagnostic excerpt

```
#!/usr/bin/env python3
"""
ZFS-style proof-state snapshot manifest.

This is a provenance/audit helper. It is not a mathematical object and not a
proof operator.
"""
from __future__ import annotations

from dataclasses import asdict, dataclass
import hashlib
import json
from pathlib import Path
from typing import Dict, Iterable


@dataclass(frozen=True)
```

© 2026 Marc Brendecke. Preprint — not peer reviewed.

```
class SnapshotManifest:
 base_version: str
 branch_name: str
 files: Dict[str, str]
 purpose: str
 non_claims: tuple[str, ...]

 def to_json_dict(self) -> dict:
 return asdict(self)


def sha256_file(path: Path) -> str:
 return hashlib.sha256(path.read_bytes()).hexdigest()
```

A.55.7 First diagnostic run record

The local diagnostic runner completed successfully. The generated JSON report is stored in the local bundle as

`python_diagnostics/diagnostic_run_report.json`.

The successful run only certifies that the diagnostic examples execute and that their internal consistency checks pass. It does not close OP 1/4 or OP 5(B).

Theorem A.55.3 (Diagnostic appendix conservativity)**.** *The local proof line diagnostic-code appendix is conservative over archival stage release-final.*

Proof. The appendix adds executable diagnostic shadows and records their status as audit aids. It does not modify Σ_4, B_4, $\mathfrak{D}_4$, $\mathcal{N}_4$, tensor grammar, readout compatibility, Historical Witness content, or OP-status guards. Therefore it is conservative. □

Remark A.55.4 (non-claim)**.** *Version 4.1.0-does not add new physics, does not close OP 1/4 or OP 5(B), does not activate primitive mixed tensors, does not claim Turing-completeness, and does not replace standard QFT/QED/Lifshitz or general-relativistic calculations.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.56 First OP Obstruction Extraction

Status. This working layer uses the diagnostic scripts from the local proof line against the two current OP-facing targets OP 1/4 and OP 5(B). The output is an obstruction extraction report, not a proof closure. It identifies the current named block that prevents promotion of each OP packet.

A.56.1 Obstruction extraction rule

Definition A.56.1 (Diagnostic obstruction extraction)**.** *A diagnostic obstruction extraction sends an OP-facing packet to a finite audit record containing its branch, pre-shift route, shifted route, backtranslation, support, shifted support, packed masks, witness link, active normal form, readout target, missing item, guard, and current status. The record is a machine-checkable audit surface and not a theorematic closure certificate.*

A.56.2 First extracted OP obstruction table

OP packet	branch	active form	obstruction	missing item
OP 1/4	horizon	$N_4(\mathrm{OP}_{1/4})$	missing selector	comparison tensor $C^{\mathrm{sel}}_{1/4}$
OP 5(B)	boundary	$N_4(O\Pi_5^{(B)})$	missing boundary promotion	theorem $\Pi_5^{(B)}$

The generated files are included in the local bundle under `python_diagnostics/`: `op_obstruction_report.json`, `op_obstruction_table.md`, and `op_obstruction_table.csv`.

A.56.3 Interpretation

Proposition A.56.2 (First OP obstruction localization)**.** *Under the current diagnostic encoding, OP 1/4 is reduced to a missing selector/comparison tensor on the horizon-side route-normalized packet, while OP 5(B) is reduced to a missing boundary-promotion theorem preserving the boundary-side stop guard.*

Proof. The diagnostic extractor verifies route shifting and backtranslation for each packet, keeps the Historical Witness link separate from the active normal form, and refuses to promote a packet when its declared missing item remains active. The OP 1/4 packet is horizon-side and therefore fails at the declared selector/comparison requirement. The OP 5(B) packet is boundary-side and therefore fails at the declared boundary-promotion requirement. These failures are named obstructions rather than closures. □

Remark A.56.3 (What the extractor has and has not shown)**.** *The extraction has shown where the current OP-facing packets are blocked in the shifted diagnostic grammar. It has not proven that the selector/comparison tensor exists, has not proven the boundary-promotion theorem, and has not promoted any physical readout.*

Theorem A.56.4 (First obstruction extraction conservativity)**.** *The local proof line first OP obstruction extraction is conservative over the local proof line and over archival stage release-final.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. The working layer adds a diagnostic extraction report and a reader-facing summary table. It does not modify Σ_4, B_4, $\mathfrak{D}_4$, $\mathcal{N}_4$, tensor grammar, readout compatibility, Historical Witness content, or OP-status guards. OP 1/4 and OP 5(B) remain open. Therefore the layer is conservative. □

Remark A.56.5 (non-claim)**.** *Version 4.1.0-does not close OP 1/4 or OP 5(B). It only names their first current diagnostic obstruction surfaces.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.57 Obstruction-to-Task Reduction Layer

Status. This working layer consumes the diagnostic obstruction extraction and turns its two named obstructions into a finite OP-facing task queue. It does not close OP 1/4 or OP 5(B), does not promote a physical readout, and does not replace the formal QAM/MML/MMA language by Python. The Python task queue is an audit and planning surface only.

A.57.1 From obstruction labels to task packets

Definition A.57.1 (OP obstruction-to-task packet)**.** *An OP obstruction-to-task packet is a finite record*

$$\mathcal{T}_{\mathrm{OP}} = (O, B, N_4(O), \tau, T, I, C, G, S),$$

where O is the OP object, B is the branch, $N_4(O)$ is the active shifted-namespace normal form, τ is the named obstruction, T is the next task kind, I is the required input, C is the success criterion, G is the guard to preserve, and S is the task status.

Remark A.57.2 (Task packets are not proof packets)**.** *The task packet names the next controlled move. It does not certify that the move can be completed, and it does not close the corresponding OP.*

A.57.2 Extracted task queue

The diagnostic task planner reads the file

`python_diagnostics/op_obstruction_report.json`

and produces

`op_task_queue.json`, `op_task_queue.md`, `op_task_queue.csv`.

The resulting task queue is:

OP	Task	Kind	Target object	Status
OP 1/4	T1	selector domain/codomain	$C^{\mathrm{sel}}_{1/4}$	ready
OP 1/4	T2	backtranslation check	B_4-compatible selector	open
OP 1/4	T3	residual reduction	$R^{\mathrm{post}}_{1/4}$	open
OP 5(B)	T1	promotion theorem	$\Pi^{(B)}_5$	ready
OP 5(B)	T2	HC/MM compatibility	boundary compatibility check	open
OP 5(B)	T3	promotion residual	$R^{(B),\mathrm{post}}_5$	open

A.57.3 OP 1/4 reduction

Proposition A.57.3 (OP 1/4 selector-task reduction)**.** *Given the obstruction record, the first controlled OP 1/4 move is not a new physical readout and not an OP closure. It is the construction of a selector/comparison tensor $C^{\mathrm{sel}}_{1/4}$ with explicit domain, codomain, route role, and backtranslation compatibility.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. The obstruction is `missing_selector` and the missing item is `selector_or_comparison_tensor`. The planner therefore extracts the first task `OP14-T1`, whose target is the selector or comparison tensor. Since the associated guard remains `OP_OPEN` and `no physical claim`, the task is a controlled reduction rather than an OP closure. □

A.57.4 OP 5(B) reduction

Proposition A.57.4 (OP 5(B) boundary-promotion task reduction)**.** *Given the obstruction record, the first controlled OP 5(B) move is the statement of a boundary-promotion theorem* $\Pi_5^{(B)}$ *preserving the earliest boundary stop. It is not a physical promotion and not an OP closure.*

Proof. The obstruction is `missing_boundary_promotion` and the missing item is `boundary_promotion_theorem`. The planner therefore extracts the first task `OP5B-T1`, whose target is a boundary promotion theorem. Since the guard requires the stop-preserving boundary condition and no physical claim, the task cannot be read as a promoted physical readout or as OP closure. □

A.57.5 Next formal move

The next mathematical step after is therefore sharply constrained:

$$\text{OP } 1/4\text{: construct/test } C_{1/4}^{\mathrm{sel}}, \qquad \text{OP } 5(\mathrm{B})\text{: state/test } \Pi_5^{(B)}.$$

If $C_{1/4}^{\mathrm{sel}}$ fails, the failure becomes the next named OP 1/4 obstruction. If $\Pi_5^{(B)}$ fails, the failure becomes the next named OP 5(B) boundary obstruction. In both cases the OP remains open unless the residual is formally discharged in the Active Core.

Theorem A.57.5 (Obstruction-to-task reduction conservativity)**.** *The local proof line task-object formalization is conservative over the local proof line.*

Proof. The layer reads the diagnostic obstruction records and produces a finite task queue. It does not change Σ_4, B_4, $\mathfrak{D}_4$, $\mathcal{N}_4$, tensor grammar, readout compatibility, Historical Witness content, or OP-status guards. Therefore it is conservative over the local pass. □

Remark A.57.6 (non-claim)**.** *Version 4.1.0-does not add new physics, does not close OP 1/4 or OP 5(B), does not activate primitive mixed tensors, does not claim Turing-completeness, and does not replace standard QFT/QED/Lifshitz or general-relativistic calculations.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.58 Task-Object Formalization

Status. This layer turns the task queue into two explicit next target objects. It is still not an OP closure. It identifies the first formal object needed for OP 1/4 as a selector/comparison tensor $C^{\mathrm{sel}}_{1/4}$ and the first formal object needed for OP 5(B) as a boundary-promotion theorem $\Pi^{(B)}_5$.

A.58.1 Task-to-object reduction rule

Definition A.58.1 (Task-to-object reduction)**.** *A task-to-object reduction sends a named obstruction task*

$$T = (OP, \text{obstruction}, \text{missing item}, G, S)$$

to a target object

$$\mathcal{T}(T) = (X_T, D_T, E_T, \mathcal{C}_T, R_T^{post}, G_T, S_T),$$

where X_T is the object to be constructed, D_T is its intended domain, E_T is its comparison or promotion codomain, $\mathcal{C}_T$ is the list of required compatibility conditions, R_T^{post} is the residual left after the construction, G_T is the retained guard, and S_T is the post-reduction status.

Remark A.58.2 (Why this is mathematical progress)**.** *The local pass/diagnostics named where the packets fail. The reduction names what must be constructed next. This changes an unlocalized OP obstruction into a concrete target-object problem.*

A.58.2 The OP 1/4 target object $C^{\mathrm{sel}}_{1/4}$

Definition A.58.3 (OP 1/4 selector/comparison target)**.** *Let $X_{14} = N_4(\mathrm{OP}_{1/4})$ be the horizon-side OP 1/4 normal-form packet, with pre-shift witness route $\rho^{\mathrm{pre}}_{1/4}$ and inhabited shifted route $\rho^{\mathrm{shift}}_{1/4}$. A first OP 1/4 selector/comparison target is a comparison tensor*

$$C^{\mathrm{sel}}_{1/4}: \quad P(\rho^{\mathrm{pre}}_{1/4}) \times P(\rho^{\mathrm{shift}}_{1/4}) \longrightarrow E^{\mathrm{sel}}_{1/4}$$

subject to backtranslation compatibility, guard preservation, and residual monotonicity $R^{\mathrm{post}}_{1/4} \preccurlyeq R^{\mathrm{pre}}_{1/4}$.

Proposition A.58.4 (OP 1/4 task-object reduction)**.** *Under the obstruction table, OP 1/4 reduces first to the construction of $C^{\mathrm{sel}}_{1/4}$ satisfying Definition A.58.3. If such a $C^{\mathrm{sel}}_{1/4}$ is constructed, the remaining OP 1/4 obstruction is the post-selector residual $R^{\mathrm{post}}_{1/4}$, not the original missing-selector obstruction.*

Proof. The task queue assigns OP 1/4 the obstruction *missing selector*. A selector obstruction is discharged only by an object with a declared domain, comparison codomain, compatibility conditions, and retained guard. Definition A.58.3 supplies exactly this target. It preserves backtranslation and guard data and leaves an explicit residual. Therefore the first obstruction is reduced to constructing $C^{\mathrm{sel}}_{1/4}$, after which only $R^{\mathrm{post}}_{1/4}$ remains to be analyzed. □

A.58.3 The OP 5(B) target object $\Pi^{(B)}_5$

Definition A.58.5 (OP 5(B) boundary-promotion target)**.** *Let $Y^{(B)}_5 = N_4(O\Pi^{(B)}_5)$ be the boundary-side OP 5(B) normal-form packet with stop-preserving boundary guard G_∂. A first OP 5(B) boundary-promotion target is a theorem schema $\Pi^{(B)}_5$ of the form*

$$\Pi^{(B)}_5: \quad (Y^{(B)}_5, G_\partial, HC/MM) \longrightarrow (Y^{(B),\mathrm{prom}}_5, G_\partial, R^{(B),\mathrm{post}}_5),$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

subject to stop preservation, HC/MM compatibility, backtranslation stability, and residual retention.

Proposition A.58.6 (OP 5(B) task-object reduction)**.** *Under the obstruction table, OP 5(B) reduces first to the formulation and proof of* $\Pi_5^{(B)}$ *satisfying Definition A.58.5. If such a theorem is obtained, the remaining OP 5(B) obstruction is* $R_5^{(B),\mathrm{post}}$ *under the retained stop-preserving guard.*

Proof. The task queue assigns OP 5(B) the obstruction *missing boundary promotion*. A boundary-promotion obstruction cannot be discharged by a route renaming alone, since the stop-preserving guard must survive promotion. The schema $\Pi_5^{(B)}$ states the exact data required for a conservative promotion: retained guard, HC/MM compatibility, backtranslation stability, and explicit post-promotion residual. Hence OP 5(B) is reduced to proving $\Pi_5^{(B)}$, after which $R_5^{(B),\mathrm{post}}$ is the remaining obstruction. □

Theorem A.58.7 (task-object formalization conservativity)**.** *The local proof line task-object formalization is conservative over the local proof line.*

Proof. The layer does not claim that $C_{1/4}^{\mathrm{sel}}$ or $\Pi_5^{(B)}$ has already been constructed. It only states their required roles and the residuals that remain after their construction. OP 1/4 and OP 5(B) therefore remain open, while their first target objects are now named and constrained. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.59 Notation and Physics-Style Audit for OP-Facing Objects

Status. This layer audits the OP-facing notation introduced in the local proof chain and separates mathematical display notation from diagnostic/code-safe identifiers. It does not change the obstruction analysis, does not close OP 1/4 or OP 5(B), and does not replace QAM/MML/MMA by Python or by information-theoretic syntax.

A.59.1 Display notation versus diagnostic identifiers

Definition A.59.1 (Display notation and diagnostic identifier)**.** *A display notation is the notation used in mathematical and physical statements. A diagnostic identifier is a machine-safe name used by Python snippets, JSON reports, CSV tables, or local audit files. Diagnostic identifiers may be short and ASCII-safe; display notation must remain mathematically and physically readable.*

Diagnostic identifier	Mathematical display notation	Role
`OP_1_4`	$\mathrm{OP}_{1/4}$	OP 1/4 normal-form target
`OP_5B`	$\mathrm{OP}_5^{(B)}$	OP 5(B) normal-form target
`C_OP14`	$C_{1/4}^{\mathrm{sel}}$	selector/comparison object for OP 1/4
`P_5B`	$\Pi_5^{(B)}$	boundary-promotion schema for OP 5(B)
`R_OP14_post`	$R_{1/4}^{\mathrm{post}}$	post-selector residual
`R_5B_post`	$R_5^{(B),\mathrm{post}}$	post-promotion residual

A.59.2 Physics-style rule

Definition A.59.2 (Physics-style rule for OP-facing notation)**.** *Information-theoretic and code-facing tools may be used as diagnostic encodings only after the governing mathematical object has been written in display notation. In particular, the active text should prefer*

$$C_{1/4}^{\mathrm{sel}}, \qquad \Pi_5^{(B)}, \qquad R_{1/4}^{\mathrm{post}}, \qquad R_5^{(B),\mathrm{post}}$$

over code-like symbols such as `C_OP14`*,* `P_5B`*,* `R_OP14`*, or* `R_5B`*.*

Remark A.59.3 (Information theory is a diagnostic layer)**.** *The route/mask/identifier layer may look information-theoretic, but the governing objects remain mathematical and physical readout objects. The information-theoretic encoding is therefore a compression and audit surface, not a replacement ontology.*

A.59.3 Normalized target-object statements

With notation normalized, the two target objects read as follows:

$$C_{1/4}^{\mathrm{sel}} : P(\rho_{1/4}^{\mathrm{pre}}) \times P(\rho_{1/4}^{\mathrm{shift}}) \longrightarrow E_{1/4}^{\mathrm{sel}},$$

and

$$\Pi_5^{(B)} : (Y_5^{(B)}, G_\partial, HC/MM) \longrightarrow (Y_5^{(B),\mathrm{prom}}, G_\partial, R_5^{(B),\mathrm{post}}).$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Thus OP 1/4 is still reduced to a selector/comparison construction, and OP 5(B) is still reduced to a boundary-promotion construction, but the display notation no longer borrows code-style names from the diagnostic layer.

Theorem A.59.4 (Notation-normalization conservativity)**.** *The notation normalization is conservative over the task-object formalization.*

Proof. The normalization replaces code-like display labels by mathematical display notation and records a diagnostic-identifier map. It does not change the underlying route, guard, obstruction, residual, or OP-status records. Therefore the reductions to a selector/comparison object for OP 1/4 and to a boundary-promotion schema for OP 5(B) are preserved. □

Remark A.59.5 (non-claim)**.** *Version 4.1.0-does not close OP 1/4 or OP 5(B), does not activate primitive mixed tensors, does not claim Turing-completeness, and does not replace standard QFT/QED/Lifshitz or general-relativistic calculations.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.60 First Selector and Boundary-Promotion Ansatz

Status. This layer starts the first genuine OP-facing mathematical attack after the notation/physics-style audit. It does not close OP 1/4 or OP 5(B). It turns the two normalized target objects,

$$C_{1/4}^{\mathrm{sel}} \qquad \text{and} \qquad \Pi_5^{(B)},$$

into first construction ansaetze: a comparison/fibre-product ansatz for OP 1/4 and a guard-preserving boundary-promotion ansatz for OP 5(B).

A.60.1 OP 1/4: comparison surface and selector ansatz

Definition A.60.1 (OP 1/4 comparison surface)**.** *Let*

$$P(\rho_{1/4}^{\mathrm{pre}}) \qquad \textit{and} \qquad P(\rho_{1/4}^{\mathrm{shift}})$$

be the witness-side and shifted-normalized packet surfaces attached to $\mathrm{OP}_{1/4}$*. A comparison surface for OP 1/4 is an object* $E_{1/4}^{\mathrm{sel}}$ *equipped with two comparison maps*

$$e_{\mathrm{pre}} : P(\rho_{1/4}^{\mathrm{pre}}) \longrightarrow E_{1/4}^{\mathrm{sel}}, \qquad e_{\mathrm{shift}} : P(\rho_{1/4}^{\mathrm{shift}}) \longrightarrow E_{1/4}^{\mathrm{sel}}.$$

The maps must be read as current shifted diagnostic/readout maps, not as a claim that the historical witness and active shifted surface are already identical.

Definition A.60.2 (Selector/comparison object for OP 1/4)**.** *Given a comparison surface* $E_{1/4}^{\mathrm{sel}}$*, define the first selector/comparison ansatz by the fibre product*

$$C_{1/4}^{\mathrm{sel}} := P(\rho_{1/4}^{\mathrm{pre}}) \times_{E_{1/4}^{\mathrm{sel}}} P(\rho_{1/4}^{\mathrm{shift}}).$$

Equivalently,

$$C_{1/4}^{\mathrm{sel}} = \{(x_{\mathrm{pre}}, x_{\mathrm{shift}}) \mid e_{\mathrm{pre}}(x_{\mathrm{pre}}) = e_{\mathrm{shift}}(x_{\mathrm{shift}})\}.$$

Thus $C_{1/4}^{\mathrm{sel}}$ *contains exactly the witness/shifted pairs that become equal after being read in the same comparison surface.*

Definition A.60.3 (Post-selector residual)**.** *The post-selector residual for OP 1/4 is the remaining comparison obstruction after the selector object has been formed. In the weak set-difference reading,*

$$R_{1/4}^{\mathrm{post}} := E_{1/4}^{\mathrm{sel}} \setminus \mathrm{im}\big(C_{1/4}^{\mathrm{sel}} \to E_{1/4}^{\mathrm{sel}}\big).$$

In a category or additive setting, the same role may be represented by a cokernel/cocone residual. The present layer requires only the weak residual role, not a fixed ambient abelian category.

Lemma A.60.4 (Selector ansatz reduces the OP 1/4 obstruction)**.** *If the comparison surface exists and the fibre product* $C_{1/4}^{\mathrm{sel}}$ *is well-defined under the inherited guards, then the obstruction* missing selector *is reduced to the post-selector residual* $R_{1/4}^{\mathrm{post}}$*.*

Proof. The obstruction identified the absence of a comparison/selector object. Definitions A.60.1 and A.60.2 provide such an object as a fibre product over a common comparison surface. Therefore the original absence of a selector is discharged at the level of ansatz formation. What remains is not the same obstruction but the part of the comparison surface not reached by that selector, namely $R_{1/4}^{\mathrm{post}}$. □

Remark A.60.5 (Why this is not yet OP 1/4 closure)**.** *The selector ansatz does not prove that* $R_{1/4}^{\mathrm{post}} = \varnothing$ *and does not prove that all OP 1/4 guards are discharged. It only turns the vague obstruction "missing selector" into a precise residual target.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.60.2 OP 5(B): guard-preserving boundary-promotion ansatz

Definition A.60.6 (Boundary-promotion ansatz for OP 5(B)). *Let $Y_5^{(B)}$ be the boundary-side object associated with $\mathrm{OP}_5^{(B)}$, let G_∂ be the inherited stop-preserving boundary guard, and let $\mathcal{R}_{HC/MM}$ denote the HC/MM readout interface. A first boundary-promotion ansatz is an operation*

$$\Pi_5^{(B)} : (Y_5^{(B)}, G_\partial, HC/MM) \longrightarrow (Y_5^{(B),\mathrm{prom}}, G_\partial, R_5^{(B),\mathrm{post}})$$

satisfying the guard-preservation condition

$$G_\partial \circ \Pi_5^{(B)} = G_\partial$$

where both sides are read on the admissible boundary-side domain.

Definition A.60.7 (HC/MM compatibility defect). *The HC/MM compatibility defect of $\Pi_5^{(B)}$ is the obstruction to the commutation relation*

$$\mathcal{R}_{HC/MM} \circ \Pi_5^{(B)} \sim_\partial \Pi_5^{(B)} \circ \mathcal{R}_{HC/MM},$$

where $\sim_\partial$ denotes boundary-guard-compatible comparison. The post-promotion residual is written

$$R_5^{(B),\mathrm{post}} := \mathrm{Obstr}\left(G_\partial \circ \Pi_5^{(B)} = G_\partial,\ \mathcal{R}_{HC/MM} \circ \Pi_5^{(B)} \sim_\partial \Pi_5^{(B)} \circ \mathcal{R}_{HC/MM}\right).$$

Lemma A.60.8 (Boundary-promotion ansatz reduces the OP 5(B) obstruction). *If $\Pi_5^{(B)}$ is defined on the admissible boundary-side domain and preserves G_∂, then the obstruction* missing boundary promotion *is reduced to the post-promotion residual $R_5^{(B),\mathrm{post}}$.*

Proof. The obstruction named the absence of a boundary-promotion mechanism. A map $\Pi_5^{(B)}$ satisfying Definition A.60.6 provides such a mechanism while preserving the inherited boundary guard. The remaining obstruction is therefore not the absence of promotion itself, but the failure, if any, of guard preservation and HC/MM compatibility. This remaining obstruction is precisely $R_5^{(B),\mathrm{post}}$. □

Remark A.60.9 (Physical reading). *The OP 5(B) ansatz keeps the physical boundary reading visible. It does not promote a boundary object by erasing the boundary stop; it asks whether a promotion through the HC/MM readout interface can be made while the stop-preserving guard survives.*

A.60.3 First combined reduction statement

Theorem A.60.10 (First OP-facing obstruction reduction). *In the ansatz layer, the two the local pass/obstruction labels reduce as follows:*

$$\begin{aligned} \mathrm{OP}_{1/4} :&\quad \textit{missing selector} \rightsquigarrow R_{1/4}^{\mathrm{post}}, \\ \mathrm{OP}_5^{(B)} :&\quad \textit{missing boundary promotion} \rightsquigarrow R_5^{(B),\mathrm{post}}. \end{aligned}$$

The reduction is conditional on the formation of $C_{1/4}^{\mathrm{sel}}$ and $\Pi_5^{(B)}$, respectively.

Proof. The first line follows from Lemma A.60.4. The second line follows from Lemma A.60.8. Both reductions are conditional because neither residual is proven to vanish in this layer. □

Remark A.60.11 (non-claim). *Version 4.1.0-does not close OP 1/4 or OP 5(B). It replaces the first named obstructions by sharper residual targets. It also does not activate primitive mixed tensors, does not claim Turing-completeness, and does not replace standard QFT/QED/Lifshitz or general-relativistic calculations.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.61 First Selector Existence Reduction for OP 1/4

Status. This layer continues the OP-facing phase by attacking the first OP 1/4 target $C^{\mathrm{sel}}_{1/4}$. The point is deliberately modest: the selector is constructed as a fiber product/equalizer object whenever the two comparison readouts to a common surface exist. This removes the purely formal "missing-selector" obstruction, but it does not close OP 1/4. The remaining questions are non-emptiness, adequacy, guard strength, and post-selector residual control.

A.61.1 Comparison surface and readout maps

Definition A.61.1 (OP 1/4 comparison datum)**.** *An OP 1/4 comparison datum is a triple*

$$\mathfrak{E}_{1/4} := \left(E^{\mathrm{sel}}_{1/4}, e_{\mathrm{pre}}, e_{\mathrm{shift}}\right)$$

where $E^{\mathrm{sel}}_{1/4}$ is a comparison surface and

$$e_{\mathrm{pre}} : P(\rho^{\mathrm{pre}}_{1/4}) \longrightarrow E^{\mathrm{sel}}_{1/4}, \qquad e_{\mathrm{shift}} : P(\rho^{\mathrm{shift}}_{1/4}) \longrightarrow E^{\mathrm{sel}}_{1/4}$$

are the pre-shift/witness-side and shifted-normalized readout maps.

Definition A.61.2 (Selector as fiber product)**.** *Given a comparison datum $\mathfrak{E}_{1/4}$, define*

$$C^{\mathrm{sel}}_{1/4} := P(\rho^{\mathrm{pre}}_{1/4}) \times_{E^{\mathrm{sel}}_{1/4}} P(\rho^{\mathrm{shift}}_{1/4})$$

as the set of compatible pairs

$$C^{\mathrm{sel}}_{1/4} = \{(x_{\mathrm{pre}}, x_{\mathrm{shift}}) \mid e_{\mathrm{pre}}(x_{\mathrm{pre}}) = e_{\mathrm{shift}}(x_{\mathrm{shift}})\}\,.$$

Equivalently, it is the equalizer of the two maps

$$e_{\mathrm{pre}} \circ \pi_{\mathrm{pre}}, \quad e_{\mathrm{shift}} \circ \pi_{\mathrm{shift}} : P(\rho^{\mathrm{pre}}_{1/4}) \times P(\rho^{\mathrm{shift}}_{1/4}) \longrightarrow E^{\mathrm{sel}}_{1/4}.$$

Lemma A.61.3 (Formal selector existence)**.** *If $P(\rho^{\mathrm{pre}}_{1/4})$, $P(\rho^{\mathrm{shift}}_{1/4})$, $E^{\mathrm{sel}}_{1/4}$, e_{pre}, and e_{shift} exist as ordinary shifted-admissible set/readout objects, then $C^{\mathrm{sel}}_{1/4}$ exists as a shifted-admissible selector object.*

Proof. The cartesian product $P(\rho^{\mathrm{pre}}_{1/4}) \times P(\rho^{\mathrm{shift}}_{1/4})$ exists in the underlying set-compatible universe. The two maps to $E^{\mathrm{sel}}_{1/4}$ determine an equality condition. By separation, the subset of pairs satisfying that equality condition exists. This subset is exactly $C^{\mathrm{sel}}_{1/4}$. Since the construction uses only product, composition, equality, and separation, it is conservative over the shifted-admissible QAM/ZF-compatible setting. □

A.61.2 Backtranslation and guards

Definition A.61.4 (Backtranslation-compatible comparison datum)**.** *A comparison datum $\mathfrak{E}_{1/4}$ is backtranslation-compatible if there is a backtranslation readout B_4 such that the diagram*

$$\begin{array}{ccc} P(\rho^{\mathrm{shift}}_{1/4}) & \xrightarrow{e_{\mathrm{shift}}} & E^{\mathrm{sel}}_{1/4} \\ \downarrow B_4 & & \downarrow B_E \\ P(\rho^{\mathrm{pre}}_{1/4}) & \xrightarrow{e_{\mathrm{pre}}} & E^{\mathrm{sel,pre}}_{1/4} \end{array}$$

commutes up to the declared comparison tolerance of the OP 1/4 route.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Lemma A.61.5 (Guard pullback)**.** *Let $G_{1/4}$ be a guard predicate on both packet surfaces and suppose e_{pre} and e_{shift} are guard-respecting. Then $C^{\mathrm{sel}}_{1/4}$ inherits the pulled-back guard*

$$G^{\mathrm{sel}}_{1/4}(x_{\mathrm{pre}}, x_{\mathrm{shift}}) := G_{1/4}(x_{\mathrm{pre}}) \wedge G_{1/4}(x_{\mathrm{shift}}).$$

Proof. Every element of $C^{\mathrm{sel}}_{1/4}$ is a pair from the two guarded packet surfaces. Defining the selector guard by conjunction of the two component guards therefore produces a well-defined predicate on the fiber product. Because the readout maps are guard-respecting, equality in $E^{\mathrm{sel}}_{1/4}$ does not erase the inherited guard data. □

A.61.3 Residual reduction

Definition A.61.6 (Post-selector residual)**.** *The post-selector residual is*

$$R^{\mathrm{post}}_{1/4} := E^{\mathrm{sel}}_{1/4} \setminus \mathrm{im}\big(C^{\mathrm{sel}}_{1/4} \to E^{\mathrm{sel}}_{1/4}\big),$$

or, in a non-set-subtractive readout context, the corresponding obstruction object measuring the part of $E^{\mathrm{sel}}_{1/4}$ not covered by compatible selector pairs.

Proposition A.61.7 (Missing selector reduces to adequacy and residual control)**.** *Under the comparison-datum hypotheses of Definition A.61.1, the OP 1/4 obstruction "missing selector" is replaced by the following narrower tasks:*

$$\mathrm{NonEmpty}(C^{\mathrm{sel}}_{1/4}), \qquad \mathrm{Adequate}(C^{\mathrm{sel}}_{1/4}), \qquad \mathrm{Control}(R^{\mathrm{post}}_{1/4}).$$

Proof. Lemma A.61.3 supplies the formal selector object as soon as the comparison datum exists. Therefore the obstruction is no longer the absence of a definable selector. What remains is whether the selector has enough elements or enough structural coverage to compare the two packet surfaces, and whether the part not covered by the selector is controlled by $R^{\mathrm{post}}_{1/4}$. These are precisely non-emptiness, adequacy, and residual-control tasks. □

Remark A.61.8 (OP 1/4 status after)**.** *This does not close OP 1/4. It does, however, make the formal obstruction smaller. The next OP 1/4 step is not "find any selector", but rather prove that the fiber-product selector is nonempty or adequate for the OP 1/4 route and then reduce $R^{\mathrm{post}}_{1/4}$.*

Theorem A.61.9 (conservativity)**.** *The local proof line selector-existence reduction is conservative over the prior archival stage working line.*

Proof. The layer constructs $C^{\mathrm{sel}}_{1/4}$ by ordinary fiber-product / equalizer logic under stated hypotheses. It does not assert non-emptiness, adequacy, OP closure, physical promotion, or any replacement of standard physical calculations. Therefore it is conservative. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.62 Selector Non-Emptiness and Anchor Criterion for OP 1/4

Status. The layer constructed the selector $C^{\mathrm{sel}}_{1/4}$ as a formal fiber product once a comparison datum exists. The present layer addresses the next obstruction: whether that selector is nonempty, and when it is structurally adequate. The result is still not an OP 1/4 closure. It reduces non-emptiness to an anchor/intersection condition inside the comparison surface $E^{\mathrm{sel}}_{1/4}$.

A.62.1 Anchor set

Definition A.62.1 (OP 1/4 selector anchor set)**.** *Given the comparison datum*

$$\mathfrak{E}_{1/4} = (E^{\mathrm{sel}}_{1/4}, e_{\mathrm{pre}}, e_{\mathrm{shift}}),$$

define the selector anchor set by

$$A^{\mathrm{sel}}_{1/4} := \mathrm{im}(e_{\mathrm{pre}}) \cap \mathrm{im}(e_{\mathrm{shift}}) \subseteq E^{\mathrm{sel}}_{1/4}.$$

An element $a \in A^{\mathrm{sel}}_{1/4}$ is a common comparison anchor if it is reached both from the pre-shift/witness-side packet surface and from the shifted-normalized packet surface.

Lemma A.62.2 (Non-emptiness criterion)**.** *The selector $C^{\mathrm{sel}}_{1/4}$ is nonempty if and only if the anchor set $A^{\mathrm{sel}}_{1/4}$ is nonempty.*

Proof. If $(x_{\mathrm{pre}}, x_{\mathrm{shift}}) \in C^{\mathrm{sel}}_{1/4}$, then

$$e_{\mathrm{pre}}(x_{\mathrm{pre}}) = e_{\mathrm{shift}}(x_{\mathrm{shift}}) = a$$

for some $a \in E^{\mathrm{sel}}_{1/4}$. Hence $a \in \mathrm{im}(e_{\mathrm{pre}}) \cap \mathrm{im}(e_{\mathrm{shift}})$, so $A^{\mathrm{sel}}_{1/4} \neq \varnothing$. Conversely, if $a \in A^{\mathrm{sel}}_{1/4}$, then there exist x_{pre} and x_{shift} with $e_{\mathrm{pre}}(x_{\mathrm{pre}}) = a = e_{\mathrm{shift}}(x_{\mathrm{shift}})$. Thus $(x_{\mathrm{pre}}, x_{\mathrm{shift}}) \in C^{\mathrm{sel}}_{1/4}$. □

A.62.2 Adequacy

Definition A.62.3 (Anchor adequacy)**.** *The anchor set $A^{\mathrm{sel}}_{1/4}$ is adequate for OP 1/4 if every comparison-relevant component of $E^{\mathrm{sel}}_{1/4}$ meets $A^{\mathrm{sel}}_{1/4}$. Equivalently, if $\pi_0^{\mathrm{rel}}(E^{\mathrm{sel}}_{1/4})$ denotes the finite family of comparison-relevant components, then*

$$\forall K \in \pi_0^{\mathrm{rel}}(E^{\mathrm{sel}}_{1/4}) : \quad K \cap A^{\mathrm{sel}}_{1/4} \neq \varnothing.$$

Lemma A.62.4 (Adequacy transfers to selector coverage)**.** *If $A^{\mathrm{sel}}_{1/4}$ is adequate, then $C^{\mathrm{sel}}_{1/4}$ has at least one compatible selector pair over every comparison-relevant component of $E^{\mathrm{sel}}_{1/4}$.*

Proof. Let K be a comparison-relevant component. By adequacy, choose $a \in K \cap A^{\mathrm{sel}}_{1/4}$. Since a is in the anchor set, it has a preimage under both e_{pre} and e_{shift}. The two preimages form a compatible pair in $C^{\mathrm{sel}}_{1/4}$ lying over $a \in K$. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.62.3 Residual localization

Definition A.62.5 (Anchor residual)**.** *The anchor residual is*

$$R^{\mathrm{anc}}_{1/4} := E^{\mathrm{sel}}_{1/4} \setminus A^{\mathrm{sel}}_{1/4},$$

or, in non-subtractive readout language, the obstruction object measuring the comparison surface not hit by both readout images.

Proposition A.62.6 (Post-selector residual is controlled by anchor residual)**.** *The post-selector residual satisfies the schematic containment*

$$R^{\mathrm{post}}_{1/4} \preceq R^{\mathrm{anc}}_{1/4},$$

where $\preceq$ denotes the current readout-level obstruction comparison. In particular, if $A^{\mathrm{sel}}_{1/4}$ is adequate and the remaining non-anchor part is readout-irrelevant, then the selector obstruction is discharged at the comparison level.

Proof. The image of the selector in the comparison surface is exactly the common image part reached by both readouts, i.e. the anchor set, up to the declared readout identification. Therefore anything not covered by the selector is contained in or represented by the complement/non-anchor obstruction. Thus the post-selector residual is controlled by the anchor residual. Adequacy removes the comparison-relevant part of this obstruction; any remaining part must be declared readout-irrelevant before it can be ignored. □

A.62.4 Reduced OP 1/4 target after

Theorem A.62.7 (OP 1/4 selector obstruction reduction)**.** *Under the comparison-datum hypotheses, the OP 1/4 selector obstruction reduces to the following anchor tasks:*

$$A^{\mathrm{sel}}_{1/4} \neq \varnothing, \qquad \mathrm{Adequate}(A^{\mathrm{sel}}_{1/4}), \qquad \mathrm{Control}(R^{\mathrm{anc}}_{1/4}).$$

Proof. By Lemma A.62.2, selector non-emptiness is equivalent to anchor non-emptiness. By Lemma A.62.4, anchor adequacy gives selector coverage over the comparison-relevant components. By Proposition A.62.6, the post-selector residual is controlled by the anchor residual. Hence the selector obstruction is reduced to anchor non-emptiness, anchor adequacy, and anchor residual control. □

Remark A.62.8 (OP 1/4 status after)**.** *OP 1/4 remains open. However, the obstruction has been sharpened again: instead of asking whether a selector exists in the abstract, the next step is to show that the two readout images have a common and adequate anchor set inside $E^{\mathrm{sel}}_{1/4}$.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.63 Anchor Adequacy and Component-Cover Criterion for OP 1/4

Status. The layer reduced selector non-emptiness to the non-emptiness of the anchor set $A^{\mathrm{sel}}_{1/4}$. The present layer attacks the next question: when is the anchor set adequate? The answer is expressed as a component-cover condition on the comparison-relevant part of $E^{\mathrm{sel}}_{1/4}$. This does not close OP 1/4; it converts the adequacy task into a finite route/component coverage task.

A.63.1 Relevant component system

Definition A.63.1 (Comparison-relevant component system)**.** *Let $E^{\mathrm{sel}}_{1/4}$ be the OP 1/4 comparison surface. A comparison-relevant component system is a finite family*

$$\mathcal{K}_{1/4} = \{K_1, \dots, K_m\}$$

of subregions, strata, or readout components of $E^{\mathrm{sel}}_{1/4}$ such that every comparison-relevant obstruction in the OP 1/4 selector problem is represented in at least one K_i.

Remark A.63.2 (Why finite components are allowed here)**.** *The component system is not asserted to exhaust the whole comparison surface. It only exhausts the comparison-relevant part selected by the current OP 1/4 route and its guards. This keeps the criterion finite and OP-facing.*

A.63.2 Anchor cover

Definition A.63.3 (Anchor cover)**.** *The anchor set $A^{\mathrm{sel}}_{1/4}$ is a component cover of $\mathcal{K}_{1/4}$ if*

$$\forall K_i \in \mathcal{K}_{1/4}: \qquad K_i \cap A^{\mathrm{sel}}_{1/4} \neq \varnothing.$$

Equivalently, the cover defect is empty, where

$$D^{\mathrm{cov}}_{1/4} := \{K_i \in \mathcal{K}_{1/4} \mid K_i \cap A^{\mathrm{sel}}_{1/4} = \varnothing\}.$$

Lemma A.63.4 (Component cover implies adequacy)**.** *If $A^{\mathrm{sel}}_{1/4}$ is a component cover of $\mathcal{K}_{1/4}$, then $A^{\mathrm{sel}}_{1/4}$ is adequate relative to the comparison-relevant component system.*

Proof. Adequacy relative to $\mathcal{K}_{1/4}$ means that every comparison-relevant component contains at least one common anchor. This is exactly the component cover condition. Hence the condition implies adequacy. □

A.63.3 Route-lifted anchors

Definition A.63.5 (Route-lifted anchor)**.** *A route-lifted anchor over K_i is a triple*

$$\alpha_i = (x_i^{\mathrm{pre}}, x_i^{\mathrm{shift}}, a_i)$$

such that

$$a_i \in K_i, \qquad e_{\mathrm{pre}}(x_i^{\mathrm{pre}}) = a_i = e_{\mathrm{shift}}(x_i^{\mathrm{shift}}),$$

and the pair $(x_i^{\mathrm{pre}}, x_i^{\mathrm{shift}})$ respects the inherited OP 1/4 route guard.

Proposition A.63.6 (Route-lifted anchors certify component cover)**.** *If every $K_i \in \mathcal{K}_{1/4}$ admits a route-lifted anchor α_i, then $A^{\mathrm{sel}}_{1/4}$ is a component cover of $\mathcal{K}_{1/4}$.*

Proof. For each K_i, the anchor α_i contains a point $a_i \in K_i$ that lies in the image of both readout maps. Therefore $a_i \in K_i \cap A^{\mathrm{sel}}_{1/4}$, so the intersection is nonempty for every K_i. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.63.4 Residual after component cover

Definition A.63.7 (Cover residual)**.** *The cover residual is*

$$R^{\mathrm{cov}}_{1/4} := D^{\mathrm{cov}}_{1/4}$$

together with any residual guard data attached to components not covered by a route-lifted anchor.

Proposition A.63.8 (Anchor residual reduces to cover residual)**.** *Relative to the component system $\mathcal{K}_{1/4}$, anchor residual control reduces to cover-residual control:*

$$\mathrm{Control}(R^{\mathrm{anc}}_{1/4}) \quad \rightsquigarrow \quad \mathrm{Control}(R^{\mathrm{cov}}_{1/4}).$$

If $D^{\mathrm{cov}}_{1/4} = \varnothing$, then the comparison-relevant anchor residual is discharged.

Proof. The component system records all comparison-relevant regions. The only comparison-relevant part not controlled by anchors is represented by components which do not meet the anchor set. These are precisely the elements of $D^{\mathrm{cov}}_{1/4}$. Thus controlling the anchor residual relative to the selected component system is equivalent to controlling the cover defect. When the cover defect is empty, every relevant component has an anchor and the component-level anchor residual vanishes. □

A.63.5 Reduced OP 1/4 target after

Theorem A.63.9 (OP 1/4 adequacy reduction to finite cover)**.** *Under the comparison and anchor hypotheses, OP 1/4 selector adequacy is reduced to the finite cover condition*

$$D^{\mathrm{cov}}_{1/4} = \varnothing,$$

or equivalently to the existence of route-lifted anchors

$$\alpha_i = (x_i^{\mathrm{pre}}, x_i^{\mathrm{shift}}, a_i)$$

for every comparison-relevant component $K_i \in \mathcal{K}_{1/4}$.

Proof. By Lemma A.63.4, a component cover is exactly adequacy relative to the selected comparison-relevant components. By Proposition A.63.6, route-lifted anchors certify this cover. By Proposition A.63.8, the remaining anchor residual is represented by the cover defect. Hence the adequacy task reduces to the finite cover condition $D^{\mathrm{cov}}_{1/4} = \varnothing$ or to the equivalent route-lifted-anchor certificate. □

Remark A.63.10 (OP 1/4 status after)**.** *OP 1/4 is still open. However, the obstruction is now much more concrete: find route-lifted anchors for the finitely selected comparison-relevant components, or identify which components remain in $D^{\mathrm{cov}}_{1/4}$.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.64 Clock-Anchored Canonical Optimal Route Principle

Status. This layer records the clock-anchored optimal-route principle suggested by the OP-facing analysis. It changes the problem shape: once the clock anchor and a target information locus are fixed, the remaining first task is to prove that at least one admissible route exists. Multiple geometric drawings of the same canonical route are not treated as different optimal routes; they are identified by the Stone–von Neumann/Teichmuller/QAM normal-form reading. This does not close OP 1/4, OP 4, or OP 5. It supplies a stronger route principle for the next OP reductions.

A.64.1 Clock anchor and admissible route space

Definition A.64.1 (Clock anchor)**.** *The clock anchor is the distinguished known reference information point*

$$q_{\mathrm{clk}}$$

against which OP-facing routes are read. It is not an optional coordinate choice; it is the fixed reference supplied by the taktgeber/clock structure of the system.

Definition A.64.2 (Admissible clock-to-target routes)**.** *For a target information locus q_{tar}, let*

$$\mathcal{R}_{\mathrm{adm}}(q_{\mathrm{clk}}, q_{\mathrm{tar}})$$

denote the class of router-admissible routes from the clock anchor to the target which respect the currently declared QAM/MML/MMA tensor interfaces and all active guards. If a boundary guard G_∂ is present, the admissible class is restricted to routes satisfying the guard-preservation condition.

Definition A.64.3 (Route action and canonical equivalence)**.** *A route action is a ranked route functional*

$$\mathcal{A} : \mathcal{R}_{\mathrm{adm}}(q_{\mathrm{clk}}, q_{\mathrm{tar}}) \longrightarrow \Lambda$$

where Λ is the ordered cost/rank set selected by the current OP-facing readout. Two routes ρ_1, ρ_2 are canonically equivalent, written

$$\rho_1 \sim_{\mathrm{can}} \rho_2,$$

if they represent the same clock-anchored information transport after the Stone–von Neumann representation identification, the Teichmuller extremal-route normalization, and the QAM/MML normal-form readout.

Remark A.64.4 (Uniqueness means canonical uniqueness)**.** *The principle does not claim that there cannot be two visually different geometric curves. It claims that, once they realize the same canonical clock-anchored information transport, they are one route in the mathematical sense relevant here. Different drawings can therefore represent the same canonical optimal route.*

A.64.2 Canonical optimal route

Definition A.64.5 (Canonical optimal route class)**.** *Assume $\mathcal{R}_{\mathrm{adm}}(q_{\mathrm{clk}}, q_{\mathrm{tar}}) \neq \varnothing$. A canonical optimal route class is*

$$[\rho_{\mathrm{opt}}]_{\mathrm{can}} := \operatorname*{argmin}_{[\rho]_{\mathrm{can}} \subseteq \mathcal{R}_{\mathrm{adm}}(q_{\mathrm{clk}}, q_{\mathrm{tar}})} \mathcal{A}([\rho]_{\mathrm{can}}),$$

where the minimization is taken after canonical equivalence has identified routes representing the same Stone–von Neumann/Teichmuller/QAM-normalized information transport.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition A.64.6 (QAM/MML canonical representative)**.** *The canonical representative*

$$\rho_{\mathrm{opt}}^{\mathrm{can}}$$

is the QAM/MML normal-form representative of $[\rho_{\mathrm{opt}}]_{\mathrm{can}}$*. It is the route object used by the router language, diagnostic comparisons, and OP-facing residual tests.*

Theorem A.64.7 (Clock-anchored canonical optimal route principle)**.** *Suppose the admissible clock-to-target route space is nonempty and finite, discretely ranked, or otherwise compact enough for the declared route action to attain its minimum. Then there exists a unique canonical optimal route representative*

$$\rho_{\mathrm{opt}}^{\mathrm{can}} = \rho_{\mathrm{opt}}^{\mathrm{can}}(q_{\mathrm{clk}}, q_{\mathrm{tar}}).$$

Proof. Since the admissible route space is nonempty and the route action attains a minimum under the stated finiteness/rank/compactness hypothesis, at least one minimizing route exists. If several geometric representatives realize the same minimum, the Stone–von Neumann representation identification, Teichmuller extremal-route normalization, and QAM/MML normal-form readout identify them as one canonical information route. The QAM/MML normal-form rule then selects one canonical representative of that class. Hence the canonical optimal route representative exists and is unique in the route language of the paper. □

A.64.3 Optimal-route map

Definition A.64.8 (Clock-anchored optimal-route map)**.** *For all target information loci for which the admissible route space is nonempty, define the optimal-route map*

$$\mathcal{O}_{\mathrm{clk}} : q_{\mathrm{tar}} \longmapsto \rho_{\mathrm{opt}}^{\mathrm{can}}(q_{\mathrm{clk}}, q_{\mathrm{tar}}).$$

This map is the local "map of optimal routes" determined by the clock anchor, the target locus, the route action, and the canonical equivalence rule.

Remark A.64.9 (Why this is stronger than arbitrary path choice)**.** *The route is not chosen by convenience. The clock anchor fixes the reference, the target fixes the information locus, the admissible class fixes the allowed router/tensor/guard moves, the route action fixes optimality, and the canonical equivalence/normal-form rule removes representational multiplicity.*

A.64.4 Effect on OP 1/4

Definition A.64.10 (Clock-to-anchor admissibility for OP 1/4)**.** *For a comparison-relevant component* $K_i \in \mathcal{K}_{1/4}$*, define*

$$\mathcal{R}_{\mathrm{adm}}(q_{\mathrm{clk}}, K_i \cap A_{1/4}^{\mathrm{sel}}) \neq \varnothing$$

to mean that at least one route-admissible clock-to-anchor route reaches a common anchor inside K_i*.*

Proposition A.64.11 (OP 1/4 cover criterion becomes route existence)**.** *If for every* $K_i \in \mathcal{K}_{1/4}$ *one has*

$$\mathcal{R}_{\mathrm{adm}}(q_{\mathrm{clk}}, K_i \cap A_{1/4}^{\mathrm{sel}}) \neq \varnothing,$$

then each component has a unique canonical optimal clock-to-anchor route

$$\rho_i^{\mathrm{opt,can}} = \rho_{\mathrm{opt}}^{\mathrm{can}}(q_{\mathrm{clk}}, K_i \cap A_{1/4}^{\mathrm{sel}}),$$

and the finite cover task is reduced to proving admissible route existence for each relevant component.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. Apply Theorem A.64.7 to each nonempty admissible route space $\mathcal{R}_{\mathrm{adm}}(q_{\mathrm{clk}}, K_i \cap A^{\mathrm{sel}}_{1/4})$. Each such route space has a unique canonical optimal representative. Therefore the component-cover problem is no longer an arbitrary anchor search; it is a finite family of clock-to-anchor admissible-route existence problems. □

A.64.5 Effect on OP 4 and OP 5

Remark A.64.12 (OP 4 as optimal-route-map stability)**.** *With the clock-anchored optimal-route map in place, OP 4 can be read as a stability problem for $\mathcal{O}_{\mathrm{clk}}$: the question is whether the canonical optimal routes remain transport-stable under the admitted router and tensor transitions.*

Proposition A.64.13 (Boundary route principle for OP 5(B))**.** *If the guard-preserving route space*

$$\mathcal{R}^{G_\partial}_{\mathrm{adm}}(q_{\mathrm{clk}}, Y_5^{(B),\mathrm{prom}})$$

is nonempty, then there is a unique canonical guard-preserving boundary promotion route

$$\rho_5^{(B),\mathrm{opt,can}}.$$

The promotion package $\Pi_5^{(B)}$ may then be read as the execution of this canonical route, subject to the remaining HC/MM compatibility and residual checks.

Proof. This is the guarded instance of the clock-anchored canonical optimal route principle, with admissibility restricted by G_∂. The existence of at least one guard-preserving route gives a minimizing canonical class and a QAM/MML representative. The execution of that representative gives the route-level reading of $\Pi_5^{(B)}$; HC/MM compatibility and residual control remain separate obligations. □

A.64.6 reduced obstruction

Theorem A.64.14 (OP-facing reduction)**.** *After the clock-anchored canonical optimal route principle, the next OP-facing obstruction is not selection among many equally optimal routes. It is the existence of admissible routes to the required anchor or boundary targets:*

$$\mathcal{R}_{\mathrm{adm}}(q_{\mathrm{clk}}, q_{\mathrm{tar}}) \neq \varnothing.$$

Once such a route exists, its canonical optimal representative is fixed by the Stone–von Neumann/Teichmuller/QAM normal-form reading.

Proof. The preceding theorem gives a unique canonical optimal representative whenever the admissible route space is nonempty and the declared action attains its minimum. Therefore the ambiguity of multiple equal routes is removed at the canonical-route level. What remains is to prove that the relevant admissible route spaces are nonempty and that their guards and residuals are respected. □

Remark A.64.15 (OP status after)**.** *OP 1/4, OP 4, and OP 5 remain open. The gain is that their next route-level form is sharper: show admissible clock-to-target route existence, then compare the current route to the canonical optimal-route map.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.65 Admissible Route Certificate and Optimal-Route Trigger

Status. The layer says that once an admissible clock-to-target route exists, the Stone–von Neumann/Teichmuller/QAM-normalized optimal route is canonically fixed. The present layer therefore moves the OP-facing task one step earlier: instead of searching for the optimal route directly, it defines the finite certificate needed to prove that an admissible route exists.

A.65.1 Route certificate

Definition A.65.1 (Admissible route certificate)**.** *Let q_{clk} be the clock anchor and q_{tar} a target information locus. An admissible route certificate is a finite record*

$$\mathsf{Cert}_\rho(q_{\mathrm{clk}}, q_{\mathrm{tar}}) = (V_\rho, E_\rho, s, t, \tau, \Gamma, \Theta)$$

where:

1. *V_ρ is a finite family of route states;*
2. *$E_\rho \subseteq V_\rho \times V_\rho$ is a finite family of directed admissible route steps;*
3. *$s = q_{\mathrm{clk}}$ is the clock start state;*
4. *$t = q_{\mathrm{tar}}$ is the target state or target component;*
5. *τ assigns the QAM/MML/MMA tensor type to each edge;*
6. *Γ records the active guard obligations for each edge;*
7. *Θ records the route action/rank data used by the local pass.*

Definition A.65.2 (Certificate validity)**.** *The certificate Cert_ρ is valid if:*

1. *there is a directed path in (V_ρ, E_ρ) from s to t;*
2. *every edge on the path has matching tensor interfaces under τ;*
3. *every edge preserves the guards declared by Γ;*
4. *the route action/rank record Θ is defined along the path.*

Lemma A.65.3 (Certificate implies admissible route existence)**.** *If $\mathsf{Cert}_\rho(q_{\mathrm{clk}}, q_{\mathrm{tar}})$ is valid, then*

$$\mathcal{R}_{\mathrm{adm}}(q_{\mathrm{clk}}, q_{\mathrm{tar}}) \neq \varnothing.$$

Proof. A valid certificate contains a directed path from the clock start state to the target state. Each edge in the path is admissible by tensor-interface matching, guard preservation, and route-action definability. Concatenating the certified edges yields an admissible route from q_{clk} to q_{tar}. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.65.2 Optimal-route trigger

Theorem A.65.4 (Certificate triggers canonical optimal route)**.** *If a valid admissible route certificate exists for* $(q_{\mathrm{clk}}, q_{\mathrm{tar}})$*, then the principle supplies the canonical optimal route representative*

$$\rho_{\mathrm{opt}}^{\mathrm{can}}(q_{\mathrm{clk}}, q_{\mathrm{tar}}).$$

Proof. By Lemma A.65.3, a valid certificate proves that the admissible route space is nonempty. Applying the clock-anchored canonical optimal route principle gives the unique canonical optimal route representative, assuming the declared action/rank setting is one in which the minimum is attained. □

A.65.3 OP 1/4 certificate form

Definition A.65.5 (OP 1/4 component certificate)**.** *For a comparison-relevant component* $K_i \in \mathcal{K}_{1/4}$*, an OP 1/4 component certificate is a valid route certificate for the target*

$$q_{\mathrm{tar},i} := K_i \cap A_{1/4}^{\mathrm{sel}}.$$

It proves that the route system can reach a common anchor in the component from the clock anchor.

Proposition A.65.6 (OP 1/4 finite certificate reduction)**.** *If every component* $K_i \in \mathcal{K}_{1/4}$ *admits a valid OP 1/4 component certificate, then every component has a canonical optimal clock-to-anchor route*

$$\rho_i^{\mathrm{opt,can}}.$$

Consequently the cover defect $D_{1/4}^{\mathrm{cov}}$ *is reduced to the absence or failure of such finite certificates.*

Proof. Each valid component certificate gives an admissible route from q_{clk} to $K_i \cap A_{1/4}^{\mathrm{sel}}$. By the local pass principle, each such route space has a canonical optimal representative. If a component has no certificate, it remains in the certificate-level obstruction. Thus the previous cover defect is represented by the finite certificate failure set. □

A.65.4 OP 5(B) boundary certificate form

Definition A.65.7 (Boundary promotion certificate)**.** *A boundary promotion certificate for OP 5(B) is a valid route certificate for*

$$q_{\mathrm{tar}} = Y_5^{(B),\mathrm{prom}}$$

inside the guard-restricted route class

$$\mathcal{R}_{\mathrm{adm}}^{G_\partial}(q_{\mathrm{clk}}, Y_5^{(B),\mathrm{prom}}).$$

Its guard record Γ *must include*

$$G_\partial(u) = G_\partial(v)$$

for every certified boundary edge $u \to v$*.*

Proposition A.65.8 (Boundary certificate triggers promotion route)**.** *A valid OP 5(B) boundary promotion certificate supplies a unique canonical guard-preserving boundary route*

$$\rho_5^{(B),\mathrm{opt,can}},$$

and hence gives the route-level input needed to interpret $\Pi_5^{(B)}$ *as execution of that canonical route, subject to HC/MM compatibility and residual checks.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. The certificate proves non-emptiness of the guard-restricted route space. The guarded route principle then supplies a canonical optimal representative. The route-level execution of this representative is exactly the route input needed for the promotion package. The HC/MM and residual clauses remain separate obligations. □

A.65.5 Certificate obstruction

Definition A.65.9 (Certificate obstruction set)**.** *For a finite family of OP targets* $\{q_{\mathrm{tar},i}\}_{i\in I}$*, define*

$$D_\rho^{\mathrm{cert}} := \{\, i \in I \mid \mathsf{Cert}_\rho(q_{\mathrm{clk}}, q_{\mathrm{tar},i}) \text{ is absent or invalid} \,\}.$$

Theorem A.65.10 (OP-facing reduction)**.** *After the local pass, the route-level OP-facing obstruction is the certificate obstruction*

$$D_\rho^{\mathrm{cert}}.$$

If $D_\rho^{\mathrm{cert}} = \varnothing$*, then all selected OP targets have canonical optimal clock-to-target routes.*

Proof. For each target, a valid certificate proves admissible route existence. By the local pass, admissible route existence triggers the canonical optimal route representative. Therefore precisely those targets without valid certificates remain obstructed. The obstruction set is D_ρ^{cert}. □

Remark A.65.11 (OP status after)**.** *OP 1/4, OP 4, and OP 5 remain open. The improvement is operational: the next thing to build is no longer an abstract optimal route, but a finite typed certificate proving that the route exists.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.66 Route-Certificate Tree and Arborescence Criterion

Status. The layer reduced route existence to finite typed route certificates. The present layer packages many target certificates into one clock-rooted directed route tree. This is the first explicit spanning/arborescence reading of the OP-facing route problem: the tree must start at the clock anchor, reach all selected OP targets, respect tensor interfaces, preserve guards, and carry route-action data. This does not close OP 1/4, OP 4, or OP 5. It makes the next obstruction a finite tree-validity problem.

A.66.1 Clock-rooted certificate tree

Definition A.66.1 (Clock-rooted route-certificate tree). *Let $\mathcal{Q} = \{q_{\mathrm{tar},i}\}_{i\in I}$ be a finite family of target information loci. A clock-rooted route-certificate tree is a finite directed tree/arborescence*

$$\mathsf{T}_\rho = (V_T, E_T, q_{\mathrm{clk}}, \mathcal{Q}, \tau_T, \Gamma_T, \Theta_T)$$

where:

1. *$q_{\mathrm{clk}} \in V_T$ is the root;*
2. *every $q_{\mathrm{tar},i} \in \mathcal{Q}$ is reached from q_{clk} by a directed path in E_T;*
3. *τ_T assigns tensor/interface type data to every edge;*
4. *Γ_T assigns the active guard record to every edge;*
5. *Θ_T assigns route action/rank data to every edge or path.*

Definition A.66.2 (Tree validity). *The tree T_ρ is valid if every root-to-target path extracted from it is a valid admissible route certificate in the sense of Definition A.65.2.*

Lemma A.66.3 (Tree validity gives all component certificates). *If T_ρ is valid, then every target $q_{\mathrm{tar},i} \in \mathcal{Q}$ has a valid admissible route certificate.*

Proof. For each target $q_{\mathrm{tar},i}$, follow the unique directed path in the tree from the clock root to that target. By tree validity, this path satisfies tensor-interface matching, guard preservation, and route-action definability. Therefore it is a valid admissible route certificate for the target. □

A.66.2 Spanning and Steiner readings

Definition A.66.4 (Spanning route tree). *A valid route-certificate tree is a spanning route tree for $\mathcal{Q}$ if its required vertices are exactly the clock root and the selected targets, together with any already declared route states.*

Definition A.66.5 (Steiner route tree). *A valid route-certificate tree is a Steiner route tree if it may introduce additional admissible intermediate states*

$$z_1, \ldots, z_\ell$$

such as HC compensation points, MML router states, MMA readout states, boundary collar states, or selector-anchor states, provided every additional state has a declared tensor type and guard role.

Remark A.66.6 (Why the Steiner version is natural here). *The OP-facing system rarely needs only a direct edge from clock to target. Intermediate states are often the meaningful physics: HC compensation, boundary-collar stabilization, route-mask decoding, or selector comparison can all appear as necessary admissible middle points. Thus the natural object is a guarded tensor-labeled directed Steiner tree, with ordinary spanning trees as the special case.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.66.3 Cost and optimal tree

Definition A.66.7 (Tree action). *The action of a valid route-certificate tree is*

$$\mathcal{A}_T(\mathsf{T}_\rho) := \sum_{e \in E_T} w(e) + \lambda\, \mathsf{G}_T + \mu\, \mathsf{R}_T,$$

where $w(e)$ is the edge route cost/rank, G_T records guard tension or guard-check complexity, and R_T records residual terms remaining on the tree. The constants λ, μ are bookkeeping weights for the current OP-facing readout.

Definition A.66.8 (Canonical optimal route tree). *If the valid tree class*

$$\mathcal{T}_{\mathrm{adm}}(q_{\mathrm{clk}}, \mathcal{Q})$$

is nonempty, the canonical optimal route tree is the QAM/MML normal-form representative

$$\mathsf{T}_\rho^{\mathrm{opt,can}} := \operatorname*{argmin}_{\mathsf{T}_\rho \in \mathcal{T}_{\mathrm{adm}}(q_{\mathrm{clk}}, \mathcal{Q})} \mathcal{A}_T(\mathsf{T}_\rho)$$

after canonical route equivalence has identified representationally equivalent trees.

Proposition A.66.9 (Valid tree triggers all canonical routes). *If a valid route-certificate tree exists for $(q_{\mathrm{clk}}, \mathcal{Q})$, then every target in $\mathcal{Q}$ has a canonical optimal clock-to-target route. If the valid tree class is nonempty and the tree action attains its minimum, then $\mathsf{T}_\rho^{\mathrm{opt,can}}$ exists.*

Proof. A valid tree gives a valid certificate for every target by Lemma A.66.3. By and the local pass, each certificate triggers a canonical optimal route. If the whole valid tree class is nonempty and the tree action attains its minimum, then minimization over canonical tree classes produces a canonical optimal route tree. □

A.66.4 OP 1/4, OP 4, and OP 5(B) readings

Definition A.66.10 (OP 1/4 anchor-cover tree). *An OP 1/4 anchor-cover tree is a valid route-certificate tree with target family*

$$\mathcal{Q}_{1/4} := \{K_i \cap A_{1/4}^{\mathrm{sel}} \mid K_i \in \mathcal{K}_{1/4}\}.$$

It certifies clock-to-anchor reachability for every comparison-relevant component.

Definition A.66.11 (OP 5(B) boundary-promotion tree). *An OP 5(B) boundary-promotion tree is a valid route-certificate tree in the guard-restricted class*

$$\mathcal{T}_{\mathrm{adm}}^{G_\partial}(q_{\mathrm{clk}}, Y_5^{(B),\mathrm{prom}})$$

such that every boundary edge preserves G_∂ and the terminal path reaches the promoted boundary target.

Remark A.66.12 (OP 4 as tree stability). *OP 4 is now naturally read as stability of the certified route tree or of the clock-anchored optimal-route map extracted from it. Informally, OP 1/4 builds or certifies the anchor tree, OP 4 tests transport stability of that tree, and OP 5(B) tests whether a boundary-promotion tree can be built without breaking the boundary guard.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.66.5 Tree obstruction

Definition A.66.13 (Tree obstruction)**.** *For a selected target family $\mathcal{Q}$, define the tree obstruction*

$$D_T^{\mathrm{cert}} := \begin{cases} \varnothing, & \text{if a valid route-certificate tree exists,} \\ \mathcal{Q}_{\mathrm{fail}}, & \text{otherwise, recording targets not reached by any valid tree path.} \end{cases}$$

Theorem A.66.14 (OP-facing tree reduction)**.** *After the local pass, the route-level OP-facing obstruction can be represented by the tree obstruction D_T^{cert}. If*

$$D_T^{\mathrm{cert}} = \varnothing,$$

then the selected targets are reached by a valid tensor-typed, guard-preserving clock-rooted route tree and hence possess canonical optimal route representatives.

Proof. If $D_T^{\mathrm{cert}} = \varnothing$, a valid route-certificate tree exists. By Proposition A.66.9, all targets then have valid certificates and canonical optimal routes. Conversely, any target not reached by a valid tree path remains a certificate failure and is recorded in the tree obstruction. □

Remark A.66.15 (OP status after)**.** *OP 1/4, OP 4, and OP 5(B) remain open. The gain is that their route-level form is now a finite tree/arborescence problem with tensor types, guards, and route action visible.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.67 Routing-Protocol Analogies for OP-Facing Trees

Status. This layer extracts routing ideas from established Internet and network-routing protocol families. The extraction is architectural only: OSPF, IS-IS, BGP, MPLS/Traffic Engineering, Segment Routing, RSVP-TE, RPL, BFD, PCEP, and controller-based routing are not imported as physics and are not proof certificates. They supply robust design patterns for the QAM/MML route-tree certificate layer introduced in the local proof chain.

A.67.1 Protocol-pattern table

Protocol idea	Network role	OP-facing route-tree analogue
OSPF / IS-IS	Link-state map and shortest-path computation inside a domain	Build a local map of admissible tensor/guard edges and compute a clock-rooted route tree.
BGP	Path-vector routing with policy constraints between domains	Record whole route histories and policy/guard filters before accepting a route.
MPLS / TE	Traffic-engineered paths with explicit constraints	Treat HC/MML/MMA/collar constraints as traffic-engineering constraints on allowed OP paths.
Segment Routing	Source/route instruction list through selected waypoints	Encode an OP route as a finite segment list of required compensator/router/readout states.
RSVP-TE	Path reservation and resource/constraint signaling	Reserve guard budget, boundary collar, or residual tolerance along a route.
RPL	Directed acyclic graph rooted at a sink/root in constrained networks	Use the clock anchor as root and build a directed acyclic route tree to targets.
BFD / keepalive	Fast failure detection for links/routes	Detect broken guard/tensor links before the OP certificate is accepted.
PCEP / SDN	Controller computes constrained paths from global view	Let the QAM/MML router compute or audit the route tree from declared global certificate data.

Remark A.67.1 (Not a protocol import)**.** *The table is a design dictionary, not a dependency. It says which proof-route pattern resembles a known network-routing pattern. It does not assert that a physical system uses Internet protocols.*

A.67.2 Link-state route database

Definition A.67.2 (OP link-state database)**.** *An OP link-state database is a finite record*

$$\mathsf{LS}_{\mathrm{OP}} = (V, E, \tau, \Gamma, \omega, \mathrm{up})$$

where V are route states, E are possible directed steps, τ is the tensor/interface label, Γ is the guard record, ω is the edge rank/cost, and $\mathrm{up}(e) \in \{0, 1\}$ records whether the edge is currently admissible.

Definition A.67.3 (Link-state admissible edge)**.** *An edge $e : u \to v$ is link-state admissible if*

$$\mathrm{up}(e) = 1, \qquad \tau(e) \text{ matches at its interfaces}, \qquad \Gamma(e) \text{ preserves all active guards.}$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proposition A.67.4 (Link-state tree extraction)**.** *If the OP link-state database contains a directed admissible path from q_{clk} to every target in $\mathcal{Q}$, then it contains a valid route-certificate tree for $(q_{\mathrm{clk}}, \mathcal{Q})$.*

Proof. Choose, for each target, one directed admissible path from the clock root to the target. The union of these paths contains a directed subgraph rooted at q_{clk} that reaches all targets. Removing redundant cycles, if any, yields a directed arborescence or a directed acyclic route certificate subtree. Since each retained edge is tensor-compatible and guard-preserving, the resulting tree is valid in the sense. □

A.67.3 Path-vector guard memory

Definition A.67.5 (Path-vector guard record)**.** *A path-vector guard record for a route*

$$\rho = (e_1, \ldots, e_n)$$

is the sequence

$$\mathsf{PV}_\Gamma(\rho) := \big((e_1, \Gamma(e_1)), \ldots, (e_n, \Gamma(e_n))\big).$$

It records not merely the current endpoint, but the guard history of the route.

Lemma A.67.6 (Path-vector guard rejection)**.** *If any edge in $\mathsf{PV}_\Gamma(\rho)$ violates an active guard, then ρ is not an admissible route certificate.*

Proof. Certificate validity requires every edge of the route path to preserve the declared guards. A path-vector guard record exposes every edge-level guard state. Therefore any recorded violation invalidates the certificate. □

A.67.4 Segment routing and OP waypoints

Definition A.67.7 (OP segment list)**.** *An OP segment list is a finite route instruction sequence*

$$\mathsf{Seg}_{\mathrm{OP}} = (z_0, z_1, \ldots, z_k)$$

where $z_0 = q_{\mathrm{clk}}$, $z_k = q_{\mathrm{tar}}$, and intermediate segments z_j may be HC compensation points, MML router points, MMA readout points, boundary-collar points, or selector-anchor points.

Proposition A.67.8 (Segment list gives Steiner-tree skeleton)**.** *A valid OP segment list determines the skeleton of a directed Steiner route tree. The intermediate segment states are precisely the permitted Steiner points needed to make the route tensor-compatible and guard-preserving.*

Proof. The segment list fixes a finite sequence of required waypoints from clock to target. Any valid realization of each segment-to-segment subroute gives a piece of the total route. The union of these pieces is a directed route tree or path with intermediate nodes that were not necessarily original targets. These nodes are Steiner points by definition. □

A.67.5 Traffic-engineering constraints

Definition A.67.9 (OP traffic-engineering constraint)**.** *An OP traffic-engineering constraint is any additional route constraint of the form*

$$\mathsf{TE} = (\mathrm{maxResidual}, \mathrm{guardBudget}, \mathrm{curvatureBound}, \mathrm{HCcompat}, \mathrm{boundaryCollar})$$

declared before a route certificate is accepted.

Definition A.67.10 (TE-valid route tree)**.** *A route-certificate tree is TE-valid if it is valid in the sense and also satisfies all declared OP traffic-engineering constraints.*

Proposition A.67.11 (TE-validity sharpens route admissibility)**.** *Replacing validity by TE-validity narrows the admissible tree class without changing the tree logic. In particular, any TE-valid tree is a valid route-certificate tree, but not every valid tree is TE-valid.*

Proof. TE-validity is validity plus additional constraints. Hence it implies ordinary validity. Since the extra constraints may fail for an otherwise valid tree, the inclusion can be strict. □

A.67.6 Failure detection and route convergence

Definition A.67.12 (Route heartbeat)**.** *A route heartbeat is a repeated local check*

$$\mathsf{HB}(e) = (\tau(e) \text{ matches},\ \Gamma(e) \text{ preserved},\ \omega(e) \text{ defined})$$

attached to an edge e. If a heartbeat fails, the edge is marked $\mathrm{up}(e) = 0$ *in the OP link-state database.*

Definition A.67.13 (Route convergence)**.** *Route convergence is the process of recomputing a valid route-certificate tree after one or more admissible edges become unavailable.*

Remark A.67.14 (Why failure detection matters for OP 4)**.** *OP 4 can be read as transport stability of the optimal-route map. Failure detection supplies the discrete analogue: if a tensor/guard edge fails, the route tree must either reconverge to a valid canonical tree or record a nonempty obstruction.*

A.67.7 Protocol-pattern reduction

Theorem A.67.15 (Routing-protocol pattern reduction)**.** *The route-tree problem may be refined into the following protocol-pattern tasks:*

Link-state task	*build* $\mathsf{LS}_{\mathrm{OP}}$,
Path-vector task	*record guard history* PV_{Γ},
Segment task	*choose OP waypoints* $\mathsf{Seg}_{\mathrm{OP}}$,
TE task	*apply residual/guard/curvature constraints*,
Heartbeat task	*detect failed tensor/guard edges*,
Convergence task	*recompute or certify obstruction.*

These tasks preserve the logic while making the route-tree construction operational.

Proof. Each protocol-pattern task refines one component of the tree certificate: link-state supplies the finite edge database, path-vector supplies route history, segment routing supplies permitted Steiner waypoints, traffic engineering supplies admissibility constraints, heartbeat supplies failure detection, and convergence supplies recomputation after failure. None of these patterns changes the meaning of a valid route-certificate tree; they only make its construction and audit explicit. □

Remark A.67.16 (OP status after)**.** *OP 1/4, OP 4, and OP 5(B) remain open. The gain is that the route-tree problem now has a practical protocol grammar: map, history, waypoints, constraints, failure detection, and convergence.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.68 Link-State Route-Tree Extraction

Status. The layer introduced routing-protocol patterns. The present layer turns the link-state pattern into an explicit route-tree extraction rule. Given a finite OP link-state database, the admissible subgraph is formed by keeping only tensor-compatible, guard-preserving, active edges. The route obstruction is then exactly the target set not reachable from the clock root in this admissible subgraph. If all targets are reachable, canonical clock-to-target paths can be extracted and packaged as a route-certificate tree. This still does not close OP 1/4, OP 4, or OP 5(B); it gives the first executable route-protocol criterion for the certificate layer.

A.68.1 Admissible subgraph

Definition A.68.1 (Admissible OP subgraph)**.** *Given the OP link-state database*

$$\mathsf{LS}_{\mathrm{OP}} = (V, E, \tau, \Gamma, \omega, \mathrm{up}),$$

define the admissible subgraph

$$\mathcal{G}_{\mathrm{adm}} = (V, E_{\mathrm{adm}})$$

by

$$E_{\mathrm{adm}} := \{e \in E \mid \mathrm{up}(e) = 1,\ \tau(e) \text{ matches},\ \Gamma(e) \text{ preserves all active guards}\}.$$

Definition A.68.2 (Reachability obstruction)**.** *For a target family $\mathcal{Q} = \{q_{\mathrm{tar},i}\}_{i\in I}$, define*

$$D_{\mathrm{reach}} := \{q_{\mathrm{tar},i} \in \mathcal{Q} \mid q_{\mathrm{tar},i} \notin \mathrm{Reach}_{\mathcal{G}_{\mathrm{adm}}}(q_{\mathrm{clk}})\}.$$

Lemma A.68.3 (Reachability is equivalent to certificate existence)**.** *For a target q_{tar}, there exists a valid link-state route certificate from q_{clk} to q_{tar} if and only if*

$$q_{\mathrm{tar}} \in \mathrm{Reach}_{\mathcal{G}_{\mathrm{adm}}}(q_{\mathrm{clk}}).$$

Proof. If a valid certificate exists, its edges are active, tensor-compatible, and guard-preserving. Hence its path lies in E_{adm}, and the target is reachable in $\mathcal{G}_{\mathrm{adm}}$. Conversely, if the target is reachable in $\mathcal{G}_{\mathrm{adm}}$, a directed path from q_{clk} to q_{tar} exists using only admissible edges. Recording its vertices, edges, tensor labels, guard records, and rank data gives a valid route certificate. □

A.68.2 Canonical path extraction

Definition A.68.4 (Canonical extracted path)**.** *Assume q_{tar} is reachable in $\mathcal{G}_{\mathrm{adm}}$. A canonical extracted path*

$$\rho_{\mathrm{LS}}^{\mathrm{can}}(q_{\mathrm{clk}}, q_{\mathrm{tar}})$$

is the QAM/MML normal-form representative of a minimum-rank path from q_{clk} to q_{tar} in $\mathcal{G}_{\mathrm{adm}}$ with respect to the edge rank ω, after canonical route equivalence has identified representationally equivalent paths.

Proposition A.68.5 (Reachable targets have canonical extracted paths)**.** *Every target outside D_{reach} has a canonical extracted path.*

Proof. If a target is outside D_{reach}, it is reachable in the finite admissible subgraph. Therefore there is at least one finite admissible path. Minimizing the declared edge-rank sum over finite paths gives a minimum-rank path. Canonical equivalence and QAM/MML normal form identify representational multiplicity and select the canonical representative. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.68.3 Tree extraction

Definition A.68.6 (Extracted link-state route tree)**.** *If $D_{\mathrm{reach}} = \varnothing$, define the extracted link-state route tree*

$$\mathsf{T}_{\mathrm{LS}}^{\mathrm{can}}$$

as the canonical acyclic reduction of the union of canonical extracted paths

$$\bigcup_{q_{\mathrm{tar}} \in \mathcal{Q}} \rho_{\mathrm{LS}}^{\mathrm{can}}(q_{\mathrm{clk}}, q_{\mathrm{tar}}).$$

Its edge labels, guard records, and rank data are inherited from $\mathsf{LS}_{\mathrm{OP}}$.

Theorem A.68.7 (Link-state extraction criterion)**.** *For a finite OP link-state database and finite target family $\mathcal{Q}$,*

$$D_{\mathrm{reach}} = \varnothing$$

if and only if an extracted valid link-state route-certificate tree exists.

Proof. If $D_{\mathrm{reach}} = \varnothing$, every target is reachable in $\mathcal{G}_{\mathrm{adm}}$. By Proposition A.68.5, each target has a canonical extracted path. The acyclic reduction of the union of these paths is a finite clock-rooted route tree reaching all targets, and all of its retained edges are admissible by construction. Conversely, if a valid extracted tree exists, every target is reached from the clock root by a valid admissible path; hence no target lies in D_{reach}. □

A.68.4 Traffic-engineering and segment refinement

Definition A.68.8 (TE-filtered admissible graph)**.** *Given a traffic-engineering constraint family TE, define*

$$\mathcal{G}_{\mathrm{adm}}^{\mathsf{TE}}$$

by further removing from $\mathcal{G}_{\mathrm{adm}}$ every edge or path segment violating the declared residual, guard-budget, curvature, HC/MM, or boundary-collar constraints.

Definition A.68.9 (Segment-constrained extraction)**.** *Given an OP segment list*

$$\mathsf{Seg}_{\mathrm{OP}} = (z_0, \dots, z_k),$$

a segment-constrained extracted route is a concatenation of canonical extracted paths

$$\rho(z_0, z_1) * \rho(z_1, z_2) * \cdots * \rho(z_{k-1}, z_k)$$

inside $\mathcal{G}_{\mathrm{adm}}^{\mathsf{TE}}$, provided every segment is reachable.

Proposition A.68.10 (Segment failure localizes obstruction)**.** *If a segment-constrained route fails, the obstruction is localized to the first segment*

$$(z_j, z_{j+1})$$

for which $z_{j+1} \notin \mathrm{Reach}_{\mathcal{G}_{\mathrm{adm}}^{\mathsf{TE}}}(z_j)$.

Proof. A segment-constrained route is a finite concatenation. If it fails, at least one segment is unreachable in the TE-filtered admissible graph. Taking the first such segment localizes the obstruction. All earlier segments are reachable by definition of first failure. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.68.5 OP-facing consequence

Theorem A.68.11 (route-protocol criterion)**.** *After the local pass, the OP-facing route-tree obstruction can be checked by the finite criterion*

$$D_{\mathrm{reach}} = \varnothing$$

in the admissible link-state graph, optionally refined to the TE-filtered and segment-constrained graph. If the criterion holds, a valid canonical route tree exists; if it fails, the failed target or failed segment is explicitly identified.

Proof. The unfiltered statement is the link-state extraction criterion. TE filtering and segment constraints only restrict the admissible graph further. Therefore the same reachability logic applies, and failures are recorded as unreachable targets or first failed segments. □

Remark A.68.12 (OP status after)**.** *OP 1/4, OP 4, and OP 5(B) remain open. The gain is that the route/tree certificate layer is now checkable by a finite link-state reachability criterion with optional traffic-engineering and segment-routing refinements.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.69 Segment-Route Waypoint Certificates

Status. The layer gave a finite link-state reachability criterion. The present layer adds segment-route structure: instead of asking only whether a target is reachable, we declare which intermediate waypoints must be respected by an OP route. This is the controlled form of the Steiner-tree idea: HC, MML, MMA, selector, and boundary-collar states may appear as required waypoints, but only with declared tensor types and guard roles. OP 1/4, OP 4, and OP 5(B) remain open.

A.69.1 Waypoint classes

Definition A.69.1 (OP waypoint class)**.** *An OP waypoint class is one of the route-state types*

$$\mathsf{WP} \in \{\mathsf{Clock}, \mathsf{QAM}, \mathsf{MML}, \mathsf{MMA}, \mathsf{HC}, \mathsf{Selector}, \mathsf{BoundaryCollar}, \mathsf{Anchor}, \mathsf{Target}\}.$$

Each waypoint used in a segment certificate must carry a tensor/interface type and a guard role.

Definition A.69.2 (Typed waypoint)**.** *A typed waypoint is a tuple*

$$z = (|z|, \mathsf{WP}(z), \tau(z), \Gamma(z))$$

where $|z|$ is the route state, $\mathsf{WP}(z)$ its waypoint class, $\tau(z)$ its tensor/interface type, and $\Gamma(z)$ its local guard record.

A.69.2 Segment-route certificate

Definition A.69.3 (Segment-route certificate)**.** *A segment-route certificate from q_{clk} to q_{tar} is a finite waypoint list*

$$\mathsf{Seg}_\rho = (z_0, z_1, \dots, z_k)$$

with $z_0 = q_{\mathrm{clk}}$, $z_k = q_{\mathrm{tar}}$, together with a link-state certificate proving that every adjacent segment

$$z_j \rightsquigarrow z_{j+1}$$

is reachable inside the currently selected admissible graph

$$\mathcal{G}_{\mathrm{adm}}^{\mathsf{TE}}.$$

Definition A.69.4 (Segment obstruction)**.** *The segment obstruction of Seg_ρ is*

$$D_{\mathrm{seg}}(\mathsf{Seg}_\rho) := \{j \mid z_{j+1} \notin \mathrm{Reach}_{\mathcal{G}_{\mathrm{adm}}^{\mathsf{TE}}}(z_j)\}.$$

Lemma A.69.5 (Segment certificate implies route certificate)**.** *If*

$$D_{\mathrm{seg}}(\mathsf{Seg}_\rho) = \varnothing,$$

then Seg_ρ determines a valid admissible route certificate from q_{clk} to q_{tar}.

Proof. If the segment obstruction is empty, every adjacent pair of waypoints is linked by an admissible route inside the TE-filtered admissible graph. Concatenating these certified subroutes gives a route from $z_0 = q_{\mathrm{clk}}$ to $z_k = q_{\mathrm{tar}}$. Since each segment is tensor-compatible, guard-preserving, and TE-valid, the concatenation is a valid admissible route certificate. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.69.3 OP 1/4 segment shape

Definition A.69.6 (OP 1/4 selector-anchor segment template)**.** *For a comparison-relevant component K_i, the OP 1/4 selector-anchor segment template is*

$$\mathsf{Seg}_{1/4,i} = (q_{\mathrm{clk}}, z_{\mathrm{QAM}}, z_{\mathrm{MML}}, z_{\mathrm{sel}}, K_i \cap A^{\mathrm{sel}}_{1/4}).$$

Here z_{QAM} records the shifted/QAM normal-form shift, z_{MML} records the router/intertwiner step, z_{sel} records the selector/comparison interface, and the final target is the common anchor region inside K_i.

Proposition A.69.7 (OP 1/4 segment criterion)**.** *If every OP 1/4 segment template $\mathsf{Seg}_{1/4,i}$ has empty segment obstruction, then every relevant component K_i has a valid clock-to-anchor route certificate and hence a canonical optimal clock-to-anchor route.*

Proof. Apply Lemma A.69.5 to each segment template. Each valid segment certificate gives a route certificate to $K_i \cap A^{\mathrm{sel}}_{1/4}$. By the local proof chain principles, each such certificate triggers a canonical optimal route. □

A.69.4 OP 5(B) segment shape

Definition A.69.8 (OP 5(B) boundary-promotion segment template)**.** *The OP 5(B) boundary-promotion segment template is*

$$\mathsf{Seg}^{(B)}_5 = (q_{\mathrm{clk}}, z_{\mathrm{HC}}, z_{\partial\mathrm{col}}, z_{\mathrm{MMA}}, Y^{(B),\mathrm{prom}}_5).$$

Here z_{HC} records HC compensation, $z_{\partial\mathrm{col}}$ records the boundary-collar guard state, z_{MMA} records the matrix/readout residual evaluator, and the terminal target is the promoted boundary-side state.

Definition A.69.9 (Boundary segment guard)**.** *The OP 5(B) segment template is boundary-valid only if every segment from z_{HC} onward preserves the boundary guard:*

$$G_\partial(z_j) = G_\partial(z_{j+1}).$$

Proposition A.69.10 (OP 5(B) segment criterion)**.** *If $\mathsf{Seg}^{(B)}_5$ is boundary-valid and*

$$D_{\mathrm{seg}}(\mathsf{Seg}^{(B)}_5) = \varnothing,$$

then OP 5(B) has a valid guard-preserving route certificate to $Y^{(B),\mathrm{prom}}_5$. The remaining obligations are HC/MM compatibility and post-promotion residual control.

Proof. Empty segment obstruction gives reachability for every adjacent segment inside the admissible graph. Boundary-validity ensures preservation of G_∂ along the boundary-sensitive part of the route. Therefore the concatenated route is a valid guard-preserving certificate to the promoted boundary target. The statement does not discharge HC/MM compatibility or residual control; those remain explicit. □

A.69.5 Segment-level OP obstruction

Definition A.69.11 (Segment-level OP obstruction)**.** *For a family of segment templates $\mathfrak{S}$, define*

$$D^{\mathrm{seg}}_{\mathrm{OP}} := \{\mathsf{Seg} \in \mathfrak{S} \mid D_{\mathrm{seg}}(\mathsf{Seg}) \neq \varnothing \text{ or a declared guard segment fails}\}.$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Theorem A.69.12 (segment-route reduction)**.** *After the local pass, the OP-facing route obstruction can be refined from target reachability to segment-template validity:*

$$D_{\mathrm{OP}}^{\mathrm{seg}} = \varnothing \quad \Longrightarrow \quad D_{\mathrm{reach}} = \varnothing$$

for the selected OP targets. If $D_{\mathrm{OP}}^{\mathrm{seg}} \neq \varnothing$, the first failed segment localizes the obstruction.

Proof. Each segment template with empty segment obstruction yields a valid route certificate by Lemma A.69.5. Thus if all selected segment templates are valid, all selected targets are reachable and D_{reach} is empty. If a segment template fails, Definition A.69.4 identifies the failed adjacent waypoint pair, localizing the obstruction. □

Remark A.69.13 (OP status after)**.** *OP 1/4, OP 4, and OP 5(B) remain open. The gain is that the Steiner route tree now has explicit permitted waypoint templates. The next task is to instantiate the OP 1/4 and OP 5(B) segment lists against the current link-state database and inspect their first failed segments.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.70 Traffic-Engineering and Guard-Budget Filters

Status. The layer declared OP segment templates. The present layer adds the filter that decides whether a reachable segment is actually admissible after residual, guard-budget, curvature, HC/MM-compatibility, and boundary-collar constraints are applied. This is a traffic-engineering analogue only: the constraints are QAM/MML/MMA route constraints, not imported network physics. OP 1/4, OP 4, and OP 5(B) remain open.

A.70.1 TE profile

Definition A.70.1 (OP traffic-engineering profile)**.** *An OP traffic-engineering profile is a tuple*

$$\mathsf{TE}_{\mathrm{OP}} = (\varepsilon_R, \varepsilon_G, \varepsilon_\kappa, \chi_{\mathrm{HC/MM}}, \chi_{\partial\mathrm{col}}, \chi_\tau)$$

where:

1. ε_R *is the maximum allowed residual increment;*
2. ε_G *is the maximum allowed guard tension;*
3. ε_κ *is the maximum allowed curvature/route-bending load;*
4. $\chi_{\mathrm{HC/MM}} \in \{0,1\}$ *records HC/MM compatibility;*
5. $\chi_{\partial\mathrm{col}} \in \{0,1\}$ *records boundary-collar admissibility;*
6. $\chi_\tau \in \{0,1\}$ *records tensor-interface compatibility.*

Definition A.70.2 (TE-valid edge)**.** *Let an edge e carry diagnostics*

$$\mathsf{d}(e) = (\Delta R(e), \Delta G(e), \kappa(e), \chi_{\mathrm{HC/MM}}(e), \chi_{\partial\mathrm{col}}(e), \chi_\tau(e)).$$

The edge is $\mathsf{TE}_{\mathrm{OP}}$-valid if

$$\Delta R(e) \le \varepsilon_R, \qquad \Delta G(e) \le \varepsilon_G, \qquad \kappa(e) \le \varepsilon_\kappa,$$

and

$$\chi_{\mathrm{HC/MM}}(e) = \chi_{\partial\mathrm{col}}(e) = \chi_\tau(e) = 1$$

whenever the corresponding compatibility clause is active for the OP segment.

A.70.2 Filtered graph

Definition A.70.3 (TE-filtered admissible graph)**.** *Given $\mathcal{G}_{\mathrm{adm}} = (V, E_{\mathrm{adm}})$, define*

$$\mathcal{G}_{\mathrm{adm}}^{\mathsf{TE}} = (V, E_{\mathrm{adm}}^{\mathsf{TE}})$$

where

$$E_{\mathrm{adm}}^{\mathsf{TE}} := \{e \in E_{\mathrm{adm}} \mid e \text{ is } \mathsf{TE}_{\mathrm{OP}}\text{-valid}\}.$$

Lemma A.70.4 (TE filtering preserves validity)**.** *Every route certificate in $\mathcal{G}_{\mathrm{adm}}^{\mathsf{TE}}$ is a valid route certificate in $\mathcal{G}_{\mathrm{adm}}$.*

Proof. The TE-filtered graph is a subgraph of the admissible graph. Therefore every edge retained by the TE filter was already active, tensor-compatible, and guard-preserving. The extra TE clauses only restrict the admissible class further. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.70.3 Guard-budget accumulation

Definition A.70.5 (Route guard budget). *For a finite route*

$$\rho = (e_1, \dots, e_n),$$

define its accumulated guard tension by

$$\mathsf{G}(\rho) := \sum_{j=1}^{n} \Delta G(e_j).$$

The route is guard-budget valid if

$$\mathsf{G}(\rho) \le \varepsilon_G^{\mathrm{tot}}.$$

Definition A.70.6 (Route residual budget). *The accumulated residual increment of ρ is*

$$\mathsf{R}(\rho) := \sum_{j=1}^{n} \Delta R(e_j).$$

The route is residual-budget valid if

$$\mathsf{R}(\rho) \le \varepsilon_R^{\mathrm{tot}}.$$

Proposition A.70.7 (Budget-valid segment certificate). *If every segment in Seg_ρ is reachable inside $\mathcal{G}_{\mathrm{adm}}^{\mathsf{TE}}$, and the concatenated route is guard-budget and residual-budget valid, then Seg_ρ determines a TE-valid admissible route certificate.*

Proof. Reachability inside the TE-filtered graph gives a concatenated route whose edges satisfy local TE constraints. Guard-budget and residual-budget validity ensure that accumulated route tension and residual remain within the declared global bounds. Hence the resulting certificate is TE-valid. □

A.70.4 Boundary and HC/MM filters

Definition A.70.8 (Boundary-collar filter). *A boundary-sensitive segment is boundary-collar valid if each boundary edge satisfies*

$$\chi_{\partial\mathrm{col}}(e) = 1, \qquad G_\partial(\mathrm{src}(e)) = G_\partial(\mathrm{dst}(e)).$$

Definition A.70.9 (HC/MM compatibility filter). *An HC/MM-sensitive segment is HC/MM-valid if each relevant edge satisfies*

$$\chi_{\mathrm{HC/MM}}(e) = 1$$

and its readout diagram commutes up to the declared route tolerance.

Proposition A.70.10 (OP 5(B) TE boundary criterion). *If $\mathsf{Seg}_5^{(B)}$ has empty segment obstruction in $\mathcal{G}_{\mathrm{adm}}^{\mathsf{TE}}$, is boundary-collar valid, is HC/MM-valid, and satisfies total guard and residual budgets, then it supplies a TE-valid guard-preserving boundary promotion route certificate.*

Proof. Empty TE-filtered segment obstruction gives reachability of all declared segments. Boundary-collar validity preserves G_∂ along the boundary-sensitive portion. HC/MM-validity supplies the required matrix/readout compatibility. Budget validity controls accumulated guard tension and residual. Thus the route certificate is TE-valid and guard-preserving. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.70.5 TE-level obstruction

Definition A.70.11 (TE obstruction)**.** *For a family of segment templates $\mathfrak{S}$, define the TE obstruction*

$$D_{\mathrm{OP}}^{\mathsf{TE}} := \{\mathsf{Seg} \in \mathfrak{S} \mid D_{\mathrm{seg}}^{\mathsf{TE}}(\mathsf{Seg}) \neq \varnothing \text{ or a budget, boundary, HC/MM, or tensor clause fails}\}.$$

Theorem A.70.12 (TE-filter reduction)**.** *After the local pass, the OP-facing segment obstruction is sharpened from reachability to TE-validity:*

$$D_{\mathrm{OP}}^{\mathsf{TE}} = \varnothing \quad \Longrightarrow \quad D_{\mathrm{OP}}^{\mathrm{seg}} = \varnothing \quad \Longrightarrow \quad D_{\mathrm{reach}} = \varnothing.$$

If $D_{\mathrm{OP}}^{\mathsf{TE}} \neq \varnothing$, the failed clause records whether the obstruction is reachability, tensor typing, guard budget, residual budget, curvature load, HC/MM compatibility, or boundary-collar preservation.

Proof. TE-validity is segment validity plus the declared budget and compatibility filters. Therefore an empty TE obstruction implies empty segment obstruction. By the local pass, empty segment obstruction implies reachability. Conversely, any TE failure is recorded by the first failed segment or by the first failed TE clause attached to that segment. □

Remark A.70.13 (OP status after)**.** *OP 1/4, OP 4, and OP 5(B) remain open. The gain is that route existence is no longer treated as enough: a route must also remain within residual, guard, curvature, HC/MM, tensor, and boundary-collar budgets.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.71 Why the New QAM Narrows the OPs Faster

Status. This layer explains why the current QAM/MML/MMA route grammar narrows the OP-facing search faster than the older QAM notation. It is an explanatory architecture layer, not an OP closure and not a physical overclaim. Its role is to document why the local proof chain could turn OP 1/4, OP 4, and OP 5(B) from global existence questions into finite route, certificate, segment, and TE-filter questions.

A.71.1 Old obstruction shape

Definition A.71.1 (Unfactored OP obstruction)**.** *An unfactored OP obstruction is an obstruction described only at the level of the target problem, without separating:*

$$\textit{route}, \quad \textit{readout}, \quad \textit{guard}, \quad \textit{residual}, \quad \textit{tensor interface}, \quad \textit{canonical normal form}.$$

In this form one can often say that an OP is open, but one cannot yet determine which local clause is responsible.

Remark A.71.2 (Why the old QAM was slower)**.** *The older QAM notation could express many structural ideas, but it did not yet force the OP into a typed route object. Consequently selector failure, boundary failure, tensor-interface failure, guard failure, residual growth, route nonexistence, and representational multiplicity could appear as one large combined obstruction.*

A.71.2 New QAM factorization

Definition A.71.3 (QAM route factorization)**.** *The current QAM/MML/MMA route grammar factors an OP-facing obstruction into the ordered layers*

$$\begin{aligned}
\textit{clock anchor} &: q_{\mathrm{clk}},\\
\textit{target locus} &: q_{\mathrm{tar}},\\
\textit{admissible route space} &: \mathcal{R}_{\mathrm{adm}}(q_{\mathrm{clk}}, q_{\mathrm{tar}}),\\
\textit{canonical optimal route} &: \rho^{\mathrm{can}}_{\mathrm{opt}},\\
\textit{certificate} &: \mathsf{Cert}_{\rho},\\
\textit{tree} &: \mathsf{T}_{\rho},\\
\textit{link-state graph} &: \mathcal{G}_{\mathrm{adm}},\\
\textit{segment list} &: \mathsf{Seg}_{\rho},\\
\textit{TE profile} &: \mathsf{TE}_{\mathrm{OP}}.
\end{aligned}$$

Proposition A.71.4 (Factorization narrows failure)**.** *If an OP obstruction is expressed in the current route grammar, then failure can be localized to one of the finite obstruction sets*

$$D_{\mathrm{reach}}, \qquad D^{\mathrm{seg}}_{\mathrm{OP}}, \qquad D^{\mathsf{TE}}_{\mathrm{OP}}, \qquad D^{\mathrm{cert}}_{\rho}, \qquad D^{\mathrm{cert}}_{T}.$$

Proof. The route grammar introduces a finite sequence of checks. Reachability fails only in D_{reach}. Segment validity fails only in $D^{\mathrm{seg}}_{\mathrm{OP}}$. TE validity fails only in $D^{\mathsf{TE}}_{\mathrm{OP}}$. Certificate validity fails in D^{cert}_{ρ}, and tree validity fails in D^{cert}_{T}. Thus a global OP failure is replaced by a finite menu of typed local failures. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.71.3 Role of the Hilbert shift and tensor notation

Remark A.71.5 (Reserved zero and clock anchoring)**.** *The shifted Hilbert-shift discipline helps because the inactive state and active route atoms are no longer confused. The clock anchor can be treated as a known reference object, while active route states are shifted into a normal form. This prevents the old ambiguity in which "no active route", "first active route", and "reference state" could blur together.*

Remark A.71.6 (Two-sided tensor notation)**.** *The two-sided tensor notation helps because it separates readout/frame/language interfaces from structure/action/field interfaces. A route edge is therefore not just an arrow; it is a typed edge with two-sided compatibility obligations. This makes tensor mismatch visible as its own failure mode rather than hiding it inside a general OP obstruction.*

A.71.4 Why could move quickly

Theorem A.71.7 (New-QAM narrowing principle)**.** *The current QAM/MML/MMA route grammar accelerates OP localization because it turns a global existence problem into a finite ordered diagnostic chain:*

$$\begin{array}{c} \textit{target problem} \\ \Downarrow \\ \textit{clock-to-target route existence} \\ \Downarrow \\ \textit{canonical optimal route} \\ \Downarrow \\ \textit{finite certificate} \\ \Downarrow \\ \textit{route tree} \\ \Downarrow \\ \textit{link-state reachability} \\ \Downarrow \\ \textit{segment waypoint validity} \\ \Downarrow \\ \textit{TE/guard-budget validity.} \end{array}$$

Proof. Each arrow is conservative and typed. The clock-to-target route separates reference from target. The canonical optimal route removes representational multiplicity. The finite certificate converts route existence into checkable data. The route tree packages many certificates. Link-state reachability turns the tree problem into finite graph reachability. Segment waypoints localize the required Steiner states. TE filters separate residual, guard, curvature, HC/MM, boundary-collar, and tensor-interface failures. Therefore the search space is progressively factored into smaller local checks. □

A.71.5 What the old QAM could and could not do

Proposition A.71.8 (Old QAM was expressive but not yet operational)**.** *The older QAM notation could describe the structural landscape, but it did not yet provide enough typed routing machinery to force a canonical OP diagnostic sequence.*

Proof. A notation may be expressive enough to describe many structural relations without being operational enough to determine which local admissibility clause fails. Before the

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Hilbert-shift normal form, router records, two-sided tensor interfaces, route certificates, segment templates, and TE filters were added, the obstruction could remain semantically correct but diagnostically unfactored. The newer grammar adds the missing operational layers. □

A.71.6 OP-facing interpretation

Remark A.71.9 (OP 1/4)**.** *For OP 1/4, the old question "does the selector exist?" is now split into: clock-to-anchor reachability, QAM/MML/selector segment validity, anchor coverage, and residual/guard/curvature budgets.*

Remark A.71.10 (OP 4)**.** *For OP 4, the old question "is the route globally stable?" is now split into: optimal-route-map stability, tree heartbeat/failure detection, reconvergence, and tensor/guard preservation along transported tree edges.*

Remark A.71.11 (OP 5(B))**.** *For OP 5(B), the old question "can boundary promotion happen?" is now split into: guard-preserving route existence, HC waypoint validity, boundary-collar validity, MMA readout validity, HC/MM compatibility, and post-promotion residual budget.*

Remark A.71.12 (Status after)**.** *This layer explains the acceleration but does not close the OPs. The next technical step is to instantiate the TE filters against the currently declared OP 1/4 and OP 5(B) segment templates and read off the first nonempty obstruction set.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.72 Mixed-Readout Edge Compatibility Layer

Status. This layer ties the traffic-engineering flags back to the earlier guarded mixed QAM readout and two-sided tensor notation. A route edge is no longer accepted merely because it is reachable and guard-budget valid. It must also admit the mixed readout required by its role. The layer is conservative: mixed readouts are derived compatibility-filtered packages of already admitted component readouts, not new primitive tensors and not OP-closure certificates.

A.72.1 Edge readout modes

Definition A.72.1 (Edge readout mode)**.** *An edge readout mode is a declared component readout*

$$R_a^m$$

attached to a route edge, where a records the readout family and m records its mode. The current OP-facing families include

$$a \in \{\mathrm{QAM}, \mathrm{MML}, \mathrm{MMA}, \mathrm{HC}, \partial\mathrm{col}, \mathrm{sel}, \mathrm{res}, \tau\}.$$

Definition A.72.2 (Mixed-readout selector)**.** *For readout modes R_a^m and R_b^n, a mixed-readout selector is a finite compatibility tensor*

$$\chi_{mn}^{ab} \in \{0, 1\}$$

or, more generally, a finite guard-valued selector. The mixed readout is admitted only when the selector returns the admissible value.

Definition A.72.3 (Edge mixed readout)**.** *The edge mixed readout determined by R_a^m, R_b^n, and χ_{mn}^{ab} is the derived package*

$$\mathrm{Mix}_\chi^{ab}(R_a, R_b)(x) \quad \rightsquigarrow \quad \chi_{mn}^{ab} R_a^m(x) R_b^n(x).$$

This expression is read with the two-sided contraction discipline: the selector χ performs the compatibility test before the paired readout is admitted.

A.72.2 From flags to mixed readouts

Definition A.72.4 (Flag-to-readout realization)**.** *The compatibility flags are realized as mixed-readout selectors:*

$$\chi_{\mathrm{HC/MM}}(e) := \chi_{mn}^{\mathrm{HC,MML/MMA}} R_{\mathrm{HC}}^m(e) R_{\mathrm{MML/MMA}}^n(e),$$

$$\chi_{\partial\mathrm{col}}(e) := \chi_{mn}^{\mathrm{HC},\partial} R_{\mathrm{HC}}^m(e) R_{\partial\mathrm{col}}^n(e),$$
$$\chi_\tau(e) := \chi_{mn}^{\mathrm{front,back}} R_{\mathrm{front}}^m(e) R_{\mathrm{back}}^n(e).$$

A flag is therefore no longer an uninterpreted Boolean; it is the value of a declared mixed-readout compatibility test.

Remark A.72.5 (Why this matters)**.** *The filter could reject a route because $\chi_{\mathrm{HC/MM}}$, $\chi_{\partial\mathrm{col}}$, or χ_τ failed. The present layer explains what those failures mean: the required pair of readout channels does not admit a guarded mixed readout at that edge.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.72.3 Mixed-readout-valid edge

Definition A.72.6 (Required mixed readout set)**.** *For an edge e, let*

$$\mathfrak{M}(e) = \{(a, b; m, n, \chi^{ab})\}$$

be the finite family of mixed-readout pairs required by the edge role. Examples include HC/MML, HC/MMA, HC/boundary-collar, selector/QAM, selector/MML, and MMA/residual pairs.

Definition A.72.7 (Mixed-readout-valid edge)**.** *An edge e is mixed-readout-valid if every required pair in $\mathfrak{M}(e)$ is admitted:*

$$\forall (a, b; m, n, \chi^{ab}) \in \mathfrak{M}(e) : \qquad \chi^{ab}_{mn} R^m_a(e) R^n_b(e) \text{ is admissible.}$$

Lemma A.72.8 (Mixed-readout validity refines TE validity)**.** *If an edge is mixed-readout-valid for every compatibility flag required by its TE profile, then the TE compatibility flags on that edge are realized by declared readout data rather than by free Boolean assumptions.*

Proof. By Definition A.72.4, each TE compatibility flag is the value of a declared mixed-readout selector. If the edge is mixed-readout-valid, every required selector admits the corresponding readout pair. Hence the flag is supported by readout data. □

A.72.4 OP 1/4 mixed readouts

Definition A.72.9 (OP 1/4 selector mixed-readout package)**.** *For the OP 1/4 segment*

$$q_{\mathrm{clk}} \to z_{\mathrm{QAM}} \to z_{\mathrm{MML}} \to z_{\mathrm{sel}} \to K_i \cap A^{\mathrm{sel}}_{1/4},$$

the selector mixed-readout package is

$$\mathfrak{M}_{1/4} := \{\mathrm{Mix}_\chi(R_{\mathrm{QAM}}, R_{\mathrm{MML}}), \mathrm{Mix}_\chi(R_{\mathrm{MML}}, R_{\mathrm{sel}}), \mathrm{Mix}_\chi(R_{\mathrm{sel}}, R_{\mathrm{anc}})\}.$$

Proposition A.72.10 (OP 1/4 mixed-readout refinement)**.** *If an OP 1/4 segment route is TE-valid and every edge of the route is mixed-readout-valid for $\mathfrak{M}_{1/4}$, then the route does not merely reach the anchor component; it reaches it through a compatible QAM/MML/selector readout chain.*

Proof. TE-validity supplies reachability, guard, residual, curvature, and tensor budget control. Mixed-readout validity supplies the required compatibility between the QAM readout, the MML router readout, the selector readout, and the anchor readout. Therefore the route reaches the anchor component through the declared readout chain rather than through an untyped path. □

A.72.5 OP 5(B) mixed readouts

Definition A.72.11 (OP 5(B) boundary mixed-readout package)**.** *For the OP 5(B) segment*

$$q_{\mathrm{clk}} \to z_{\mathrm{HC}} \to z_{\partial\mathrm{col}} \to z_{\mathrm{MMA}} \to Y_5^{(B),\mathrm{prom}},$$

the boundary mixed-readout package is

$$\mathfrak{M}_5^{(B)} := \{\mathrm{Mix}_\chi(R_{\mathrm{HC}}, R_{\partial\mathrm{col}}), \mathrm{Mix}_\chi(R_{\partial\mathrm{col}}, R_{\mathrm{MMA}}), \mathrm{Mix}_\chi(R_{\mathrm{MMA}}, R_{\mathrm{res}}), \mathrm{Mix}_\chi(R_{\mathrm{HC}}, R_{\mathrm{MML/MM}})\}.$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proposition A.72.12 (OP 5(B) mixed-readout refinement)**.** *If an OP 5(B) boundary segment route is TE-valid, boundary-collar valid, HC/MM-valid, and mixed-readout-valid for $\mathfrak{M}_5^{(B)}$, then the route supplies a compatible boundary-promotion readout chain. The remaining obligation is the post-promotion residual control, not the mere existence of a compatible readout route.*

Proof. TE-validity and boundary-collar validity supply the route, guard, and boundary constraints. HC/MM-validity supplies the matrix/readout compatibility. The mixed-readout package asserts that HC, boundary-collar, MMA, residual, and MML/MM readouts can be paired through declared selectors. Thus the boundary promotion route is compatible at the readout level. The residual after promotion remains a separate quantity to be controlled. □

A.72.6 Mixed-readout obstruction

Definition A.72.13 (Mixed-readout obstruction)**.** *For a route ρ, define*

$$D_{\mathrm{mix}}(\rho) := \{(e,a,b;m,n) \mid (a,b;m,n,\chi^{ab}) \in \mathfrak{M}(e) \text{ and } \chi^{ab}_{mn} R^m_a(e) R^n_b(e) \text{ is not admissible}\}.$$

Theorem A.72.14 (mixed-readout reduction)**.** *After the local pass, the OP-facing TE obstruction refines to the mixed-readout obstruction:*

$$D_{\mathrm{mix}}(\rho) = \varnothing \quad \Longrightarrow \quad \begin{array}{c} \text{all required compatibility flags on } \rho \text{ are realized} \\ \text{by declared readout selectors.} \end{array}$$

If $D_{\mathrm{mix}}(\rho) \neq \varnothing$, the obstruction records the exact edge and readout pair that failed.

Proof. The implication follows directly from the flag-to-readout realization. A nonempty mixed-readout obstruction records an edge and a required readout pair whose selector does not admit the mixed readout. Hence the route is no longer rejected by an opaque compatibility flag; it is rejected by a named readout-pair failure. □

Remark A.72.15 (OP status after)**.** *OP 1/4, OP 4, and OP 5(B) remain open. The gain is that the compatibility part of the route filter is now readable as mixed-readout admissibility. This should make the next OP pass more precise: one can test whether the obstruction is reachability, segment structure, budget, or a specific failed readout mixture.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.73 Mixed-Readout Index Registry and Reader Atlas

Status. This layer is a reader-facing and audit-facing index registry for the local pass mixed-readout edge compatibility layer. It does not add new physics and does not close OP 1/4, OP 4, or OP 5(B). It fixes the meaning of the index tuple

$$(e, a, b; m, n)$$

so that later OP diagnostics can say exactly which edge, readout family, and mode pairing failed.

A.73.1 Four index levels

Definition A.73.1 (Mixed-readout index tuple). *A mixed-readout diagnostic index is a tuple*

$$\iota_{\mathrm{mix}} := (e, a, b; m, n),$$

where:

1. *e is the route edge being tested;*
2. *a, b are readout-family indices;*
3. *m, n are mode indices inside the corresponding readout families.*

The tuple is read as: on edge e, readout family a in mode m is being mixed with readout family b in mode n.

Definition A.73.2 (Readout-family registry). *The current readout-family registry is*

$$\mathcal{A}_{\mathrm{RO}} := \{\mathrm{QAM}, \mathrm{MML}, \mathrm{MMA}, \mathrm{HC}, \partial\mathrm{col}, \mathrm{sel}, \mathrm{anc}, \mathrm{res}, \tau\}.$$

The intended readings are:

Family	*Reading*
QAM	*object, language, normal-form, and store-side readout*
MML	*route, instruction, and router/intertwiner readout*
MMA	*matrix, execution, and readout-semantics evaluator*
HC	*Heisenberg-compensator/frame-normalization readout*
∂col	*boundary-collar and boundary-guard readout*
sel	*selector/comparison-surface readout*
anc	*anchor/common-target readout*
res	*residual/parity/obstruction readout*
τ	*tensor-interface/type readout*

Definition A.73.3 (Mode index registry). *For every readout family $a \in \mathcal{A}_{\mathrm{RO}}$, the mode index $m \in M_a$ records the local mode in which that family is read. In these mode sets remain named finite local registries rather than globally numeric registers. This prevents premature global enumeration before the OP-facing mode families have stabilized.*

Remark A.73.4 (Why modes are not globally numbered yet). *A global numeric mode registry would be useful later, but it would be too early here. The current OP-facing work first needs to learn which modes are actually used by OP 1/4, OP 4, and OP 5(B). Therefore fixes the role of the mode indices without freezing a final numeric list.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.73.2 Selector grammar

Definition A.73.5 (Selector-indexed mixed readout)**.** *Given the diagnostic index* $(e, a, b; m, n)$*, the corresponding selector-indexed mixed readout is*

$$\mathrm{Mix}_{\chi}^{ab}[m, n](e) := \chi_{mn}^{ab}(e)\, R_a^m(e)\, R_b^n(e).$$

The expression is admissible only if the selector permits the pair:

$$\chi_{mn}^{ab}(e) = 1$$

or, in a guard-valued registry, if $\chi_{mn}^{ab}(e)$ *returns an admissible guard state.*

Definition A.73.6 (Mixed-readout obstruction with indices)**.** *For a route* ρ*, the indexed mixed-readout obstruction is*

$$D_{\mathrm{mix}}(\rho) := \{(e, a, b; m, n) \mid \mathrm{Mix}_{\chi}^{ab}[m, n](e) \text{ is not admissible}\}.$$

Proposition A.73.7 (Indexed failure localization)**.** *If* $(e, a, b; m, n) \in D_{\mathrm{mix}}(\rho)$*, then the mixed-readout failure is localized to one route edge, two readout families, and one pair of local modes.*

Proof. This is immediate from the definition of $D_{\mathrm{mix}}(\rho)$. The tuple records exactly the tested edge, the two readout families, and the two modes. Hence the failure is not a global compatibility failure but a local selector-indexed readout-pair failure. □

A.73.3 OP 1/4 atlas

Definition A.73.8 (OP 1/4 mixed-readout index atlas)**.** *For the OP 1/4 segment*

$$q_{\mathrm{clk}} \to z_{\mathrm{QAM}} \to z_{\mathrm{MML}} \to z_{\mathrm{sel}} \to K_i \cap A_{1/4}^{\mathrm{sel}},$$

the default mixed-readout index atlas is

$$\begin{aligned} e_{QM} &: (\mathrm{QAM}, \mathrm{MML}; m_Q, m_M), \\ e_{MS} &: (\mathrm{MML}, \mathrm{sel}; m_M, m_S), \\ e_{SA} &: (\mathrm{sel}, \mathrm{anc}; m_S, m_A). \end{aligned}$$

Thus the OP 1/4 route is read as a chain of three selector-mediated tests: QAM/MML, MML/selector, and selector/anchor.

Remark A.73.9 (Human reading of the OP 1/4 atlas)**.** *The OP 1/4 atlas says: first the QAM normal-form readout must be compatible with the MML router readout; then the router readout must be compatible with the selector readout; then the selector readout must be compatible with the anchor readout. A failure in any of the three pairs tells which part of the selector route must be repaired.*

A.73.4 OP 5(B) atlas

Definition A.73.10 (OP 5(B) mixed-readout index atlas)**.** *For the OP 5(B) segment*

$$q_{\mathrm{clk}} \to z_{\mathrm{HC}} \to z_{\partial\mathrm{col}} \to z_{\mathrm{MMA}} \to Y_5^{(B),\mathrm{prom}},$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

the default mixed-readout index atlas is

$$\begin{aligned} e_{H\partial} &: (\mathrm{HC}, \partial\mathrm{col}; m_H, m_\partial), \\ e_{\partial M} &: (\partial\mathrm{col}, \mathrm{MMA}; m_\partial, m_M), \\ e_{MR} &: (\mathrm{MMA}, \mathrm{res}; m_M, m_R), \\ e_{HMM} &: (\mathrm{HC}, \mathrm{MML/MM}; m_H, m_{MM}). \end{aligned}$$

The last pair records the HC/MM compatibility readout that may be attached to the boundary-promotion route even when it is not a separate visible edge.

Remark A.73.11 (Human reading of the OP 5(B) atlas)**.** *The OP 5(B) atlas says: HC must be compatible with the boundary collar; the boundary collar must be compatible with the MMA readout; the MMA readout must be compatible with the residual evaluator; and HC/MM compatibility must remain available as a route-level readout. This separates boundary failure from matrix-readout failure and from residual-control failure.*

A.73.5 Index-to-filter bridge

Proposition A.73.12 (Index bridge to TE flags)**.** *The TE flags*

$$\chi_{\mathrm{HC/MM}}, \qquad \chi_{\partial\mathrm{col}}, \qquad \chi_\tau$$

are instances of the indexed selector grammar

$$\chi^{ab}_{mn}(e)$$

after the readout families and modes have been chosen from the registry.

Proof. The layer realized the TE flags as mixed-readout selectors. The present registry fixes the general selector grammar by naming the edge e, readout families a, b, and mode pair m, n. Therefore each TE flag is a special case of $\chi^{ab}_{mn}(e)$ with the relevant family and mode assignment. □

Theorem A.73.13 (mixed-readout index clarification)**.** *After the local pass, a mixed-readout compatibility failure is expressible in the canonical form*

$$(e, a, b; m, n) \in D_{\mathrm{mix}}(\rho).$$

This means that the obstruction is localized to a specific route edge, a specific pair of readout families, and a specific pair of local modes.

Proof. By Definition A.73.1, every mixed-readout diagnostic has the form $(e, a, b; m, n)$. By Definition A.73.6, failed mixed-readout tests are precisely recorded in $D_{\mathrm{mix}}(\rho)$. Thus every compatibility failure can be written in the stated canonical form. □

Remark A.73.14 (OP status after)**.** *OP 1/4, OP 4, and OP 5(B) remain open. The gain is explanatory and diagnostic: the new mixed-readout layer is now indexed clearly enough for a reader or audit script to understand exactly what is being tested on each route edge.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.74 First Mixed-Readout Mode Registry

Status. This layer continues the index registry by naming the first local mode sets used by OP 1/4 and OP 5(B). The modes are intentionally not converted into a final global numeric registry. They are local OP-facing modes that stabilize the meaning of m, n in the diagnostic tuple

$$(e, a, b; m, n).$$

This is a reader/audit layer and a preparation for sharper OP diagnostics; it does not close OP 1/4, OP 4, or OP 5(B).

A.74.1 Local mode sets

Definition A.74.1 (Local mode set)**.** *For a readout family a, a local mode set M_a^{loc} is the finite set of named modes currently needed by the OP-facing route grammar. A local mode is not yet a global numerical index; it is a stable semantic label for a readout use.*

Definition A.74.2 (First OP-facing mode registry)**.** *The first OP-facing mode registry is the finite family*

$$\mathcal{M}_{\mathrm{OP}}^{(1)} = \{M_{\mathrm{QAM}}, M_{\mathrm{MML}}, M_{\mathrm{MMA}}, M_{\mathrm{HC}}, M_{\partial\mathrm{col}}, M_{\mathrm{sel}}, M_{\mathrm{anc}}, M_{\mathrm{res}}, M_{\tau}\}$$

with the following local mode labels:

$$\begin{aligned}
M_{\mathrm{QAM}} &= \{\mathrm{normal}, \mathrm{store}, \mathrm{mask}, \mathrm{front}\},\\
M_{\mathrm{MML}} &= \{\mathrm{router}, \mathrm{intertwiner}, \mathrm{segment}, \mathrm{path}\},\\
M_{\mathrm{MMA}} &= \{\mathrm{matrix}, \mathrm{execution}, \mathrm{residualEval}, \mathrm{back}\},\\
M_{\mathrm{HC}} &= \{\mathrm{comp}, \mathrm{frame}, \mathrm{boundaryNorm}, \mathrm{mmCompat}\},\\
M_{\partial\mathrm{col}} &= \{\mathrm{boundary}, \mathrm{collar}, \mathrm{guard}, \mathrm{stop}\},\\
M_{\mathrm{sel}} &= \{\mathrm{selector}, \mathrm{comparison}, \mathrm{fiber}, \mathrm{equalizer}\},\\
M_{\mathrm{anc}} &= \{\mathrm{anchor}, \mathrm{component}, \mathrm{commonImage}, \mathrm{target}\},\\
M_{\mathrm{res}} &= \{\mathrm{residual}, \mathrm{post}, \mathrm{budget}, \mathrm{obstruction}\},\\
M_{\tau} &= \{\mathrm{frontType}, \mathrm{backType}, \mathrm{contract}, \mathrm{compare}\}.
\end{aligned}$$

Remark A.74.3 (Why this is still local)**.** *The registry is strong enough for OP 1/4 and OP 5(B) diagnostics but weak enough to avoid premature global numbering. A later version may turn these labels into a numeric or binary mask registry after the OP-facing mode use has stabilized.*

A.74.2 OP 1/4 instantiated modes

Definition A.74.4 (OP 1/4 mode assignments)**.** *The first OP 1/4 mixed-readout mode assignments are*

$$\begin{aligned}
e_{QM} &: (\mathrm{QAM}, \mathrm{MML}; \mathrm{normal}, \mathrm{router}),\\
e_{MS} &: (\mathrm{MML}, \mathrm{sel}; \mathrm{router}, \mathrm{selector}),\\
e_{SA} &: (\mathrm{sel}, \mathrm{anc}; \mathrm{selector}, \mathrm{anchor}).
\end{aligned}$$

Equivalently, the OP 1/4 mixed-readout tests are

$$\chi^{\mathrm{QAM,MML}}_{\mathrm{normal,router}}(e_{QM}),$$

$$\chi^{\mathrm{MML,sel}}_{\mathrm{router,selector}}(e_{MS}),$$

$$\chi^{\mathrm{sel,anc}}_{\mathrm{selector,anchor}}(e_{SA}).$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proposition A.74.5 (OP 1/4 first mode diagnostic)**.** *If all three OP 1/4 mode assignments are admitted by their selectors, then the OP 1/4 segment route passes the first mixed-readout compatibility layer. If one fails, the failure is localized to the corresponding pair: QAM/MML, MML/selector, or selector/anchor.*

Proof. The obstruction records failures as tuples $(e, a, b; m, n)$. Substituting the OP 1/4 mode assignments gives exactly the three listed tests. Therefore empty failure set across these tests means the first mixed-readout layer passes, while a nonempty failure identifies the failing pair. □

A.74.3 OP 5(B) instantiated modes

Definition A.74.6 (OP 5(B) mode assignments)**.** *The first OP 5(B) mixed-readout mode assignments are*

$$\begin{aligned} e_{H\partial} &: (\mathrm{HC}, \partial\mathrm{col}; \mathrm{boundaryNorm}, \mathrm{collar}),\\ e_{\partial M} &: (\partial\mathrm{col}, \mathrm{MMA}; \mathrm{guard}, \mathrm{matrix}),\\ e_{MR} &: (\mathrm{MMA}, \mathrm{res}; \mathrm{residualEval}, \mathrm{post}),\\ e_{HMM} &: (\mathrm{HC}, \mathrm{MML/MM}; \mathrm{mmCompat}, \mathrm{intertwiner}). \end{aligned}$$

Equivalently, the OP 5(B) mixed-readout tests are

$$\chi^{\mathrm{HC},\partial\mathrm{col}}_{\mathrm{boundaryNorm},\mathrm{collar}}(e_{H\partial}),$$

$$\chi^{\partial\mathrm{col},\mathrm{MMA}}_{\mathrm{guard},\mathrm{matrix}}(e_{\partial M}),$$

$$\chi^{\mathrm{MMA},\mathrm{res}}_{\mathrm{residualEval},\mathrm{post}}(e_{MR}),$$

$$\chi^{\mathrm{HC},\mathrm{MML/MM}}_{\mathrm{mmCompat},\mathrm{intertwiner}}(e_{HMM}).$$

Proposition A.74.7 (OP 5(B) first mode diagnostic)**.** *If all four OP 5(B) mode assignments are admitted by their selectors, then the boundary-promotion route passes the first mixed-readout compatibility layer. If one fails, the failure is localized to HC/boundary-collar, boundary-collar/MMA, MMA/residual, or HC/MM compatibility.*

Proof. As in the OP 1/4 case, records a failed selector as $(e, a, b; m, n)$. The four assignments instantiate those indices for the OP 5(B) boundary route. Thus pass/fail is localized to one named readout-pair test. □

A.74.4 Mode diagnostic obstruction

Definition A.74.8 (Mode diagnostic obstruction)**.** *For a route ρ, define the first mode diagnostic obstruction*

$$D^{(1)}_{\mathrm{mode}}(\rho) := \{(e, a, b; m, n) \in D_{\mathrm{mix}}(\rho) \mid m, n \in \mathcal{M}^{(1)}_{\mathrm{OP}}\}.$$

Theorem A.74.9 (first mode-registry reduction)**.** *After the local pass, the first OP-facing mixed-readout compatibility problem can be tested against the named local mode registry $\mathcal{M}^{(1)}_{\mathrm{OP}}$. If*

$$D^{(1)}_{\mathrm{mode}}(\rho) = \varnothing,$$

then the route passes the first local mode layer. If not, the obstruction is a specific edge, readout-family pair, and named mode pair.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. The definition of $D^{(1)}_{\mathrm{mode}}$ restricts the mixed-readout obstruction to the first OP-facing mode registry. Therefore the empty set means no mixed-readout selector fails at the first mode layer. A nonempty set gives the exact tuple $(e, a, b; m, n)$, with m, n now named rather than schematic. □

Remark A.74.10 (Use in the next OP pass)**.** *The next OP pass can instantiate the tests before attempting larger route-tree or residual arguments. If the first mixed-readout mode layer fails, the correct repair target is local. If it passes, one can continue to reachability, TE budgets, residual control, and OP 4 stability.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.75 First Selector Table and Mixed-Readout Diagnostic Pass

Status. The layer named the first local mixed-readout mode registry. The present layer turns those named mode pairs into the first selector table. A route passes the first mixed-readout diagnostic pass exactly when the required selector entries $\chi^{ab}_{mn}(e)$ are admitted on the required edges. This is still not OP closure; it is a finite compatibility precheck before the larger reachability, TE-budget, residual, and OP 4 stability arguments.

A.75.1 First selector table

Definition A.75.1 (First selector table)**.** *The first selector table is the partial finite table*

$$\mathsf{Sel}^{(1)} \subseteq E \times \mathcal{A}_{\mathrm{RO}} \times \mathcal{A}_{\mathrm{RO}} \times M \times M$$

whose entries have the form

$$\mathsf{Sel}^{(1)}(e, a, b; m, n) = \chi^{ab}_{mn}(e) \in \{0, 1\}.$$

An entry value 1 *means that the mixed readout*

$$\mathrm{Mix}^{ab}_{\chi}[m, n](e)$$

is admitted by the first local mode layer. An entry value 0 *means that the pair is rejected at this layer.*

Definition A.75.2 (Required first selector entries)**.** *For a route ρ, the required first selector entry set is*

$$\mathsf{Req}^{(1)}(\rho) := \{(e, a, b; m, n) \mid \textit{the first-mode atlas requires this mixed readout on } \rho\}.$$

Definition A.75.3 (First selector failure set)**.** *The first selector failure set is*

$$D^{(1)}_{\chi}(\rho) := \{(e, a, b; m, n) \in \mathsf{Req}^{(1)}(\rho) \mid \mathsf{Sel}^{(1)}(e, a, b; m, n) \neq 1\}.$$

Lemma A.75.4 (Selector table equals first mode diagnostic)**.** *For the local mode registry,*

$$D^{(1)}_{\chi}(\rho) = D^{(1)}_{\mathrm{mode}}(\rho).$$

up to the notational convention that $D^{(1)}_{\chi}$ records the selector table entry and $D^{(1)}_{\mathrm{mode}}$ records the failed mixed-readout tuple.

Proof. Both sets are built from the same required tuples $(e, a, b; m, n)$. The first records tuples whose selector-table value is not admitted; the second records tuples whose mixed readout is not admissible. Since first-layer mixed-readout admissibility is exactly the condition $\chi^{ab}_{mn}(e) = 1$, the two obstruction records agree. □

A.75.2 OP 1/4 selector table

Definition A.75.5 (OP 1/4 first selector table)**.** *The OP 1/4 first selector table consists of the entries*

$$\begin{aligned}
\mathsf{Sel}^{(1)}_{1/4}(e_{QM}) &= \chi^{\mathrm{QAM,MML}}_{\mathrm{normal,router}}(e_{QM}),\\
\mathsf{Sel}^{(1)}_{1/4}(e_{MS}) &= \chi^{\mathrm{MML,sel}}_{\mathrm{router,selector}}(e_{MS}),\\
\mathsf{Sel}^{(1)}_{1/4}(e_{SA}) &= \chi^{\mathrm{sel,anc}}_{\mathrm{selector,anchor}}(e_{SA}).
\end{aligned}$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition A.75.6 (OP 1/4 selector pass)**.** *The OP 1/4 first selector pass condition is*

$$\mathsf{Pass}_{1/4}^{\chi,(1)} := \big[\mathsf{Sel}_{1/4}^{(1)}(e_{QM}) = \mathsf{Sel}_{1/4}^{(1)}(e_{MS}) = \mathsf{Sel}_{1/4}^{(1)}(e_{SA}) = 1\big].$$

Proposition A.75.7 (OP 1/4 first selector diagnostic)**.** *If* $\mathsf{Pass}_{1/4}^{\chi,(1)}$ *holds, then the OP 1/4 route passes the first QAM/MML/selector/anchor mixed-readout compatibility check. If it fails, the failed selector entry identifies whether the obstruction is QAM/MML, MML/selector, or selector/anchor.*

Proof. The three entries are exactly the three mixed-readout pairings required by the OP 1/4 atlas. If all are admitted, the first compatibility check passes. If any entry is not admitted, the failed tuple names the failed pair. □

A.75.3 OP 5(B) selector table

Definition A.75.8 (OP 5(B) first selector table)**.** *The OP 5(B) first selector table consists of the entries*

$$\begin{aligned}
\mathsf{Sel}_{5B}^{(1)}(e_{H\partial}) &= \chi^{\mathrm{HC},\partial\mathrm{col}}_{\mathrm{boundaryNorm,collar}}(e_{H\partial}),\\
\mathsf{Sel}_{5B}^{(1)}(e_{\partial M}) &= \chi^{\partial\mathrm{col},\mathrm{MMA}}_{\mathrm{guard,matrix}}(e_{\partial M}),\\
\mathsf{Sel}_{5B}^{(1)}(e_{MR}) &= \chi^{\mathrm{MMA,res}}_{\mathrm{residualEval,post}}(e_{MR}),\\
\mathsf{Sel}_{5B}^{(1)}(e_{HMM}) &= \chi^{\mathrm{HC,MML/MM}}_{\mathrm{mmCompat,intertwiner}}(e_{HMM}).
\end{aligned}$$

Definition A.75.9 (OP 5(B) selector pass)**.** *The OP 5(B) first selector pass condition is*

$$\mathsf{Pass}_{5B}^{\chi,(1)} := \big[\mathsf{Sel}_{5B}^{(1)}(e_{H\partial}) = \mathsf{Sel}_{5B}^{(1)}(e_{\partial M}) = \mathsf{Sel}_{5B}^{(1)}(e_{MR}) = \mathsf{Sel}_{5B}^{(1)}(e_{HMM}) = 1\big].$$

Proposition A.75.10 (OP 5(B) first selector diagnostic)**.** *If* $\mathsf{Pass}_{5B}^{\chi,(1)}$ *holds, then the OP 5(B) boundary route passes the first HC/boundary/MMA/residual and HC/MM mixed-readout compatibility check. If it fails, the failed selector entry identifies whether the obstruction is HC/boundary-collar, boundary-collar/MMA, MMA/residual, or HC/MM.*

Proof. The four entries are exactly the four mixed-readout pairings required by the OP 5(B) atlas. The pass condition says all four are admitted. Any failure therefore names a unique first-layer mixed-readout obstruction. □

A.75.4 Interaction with TE and route checks

Theorem A.75.11 (First selector pass as compatibility precheck)**.** *The first selector pass is a necessary compatibility precheck for the local pass TE-valid route claims. More precisely, a route cannot satisfy the local pass compatibility clauses through declared mixed readouts unless*

$$D_{\chi}^{(1)}(\rho) = \varnothing$$

for all required first-layer mixed-readout entries.

Proof. The layer realizes the compatibility flags as mixed-readout selector tests. The layer names the first OP-facing modes. Therefore, if a required first-layer selector entry is not admitted, the corresponding compatibility flag cannot be supported by declared readout data at that edge. Hence empty $D_{\chi}^{(1)}$ is necessary for TE compatibility through this readout layer. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Remark A.75.12 (Where this helps the OPs)**.** *The first selector table is useful because it is cheaper to check than the full route/tree/residual problem. If the table fails, the OP obstruction is already localized to one mixed-readout pair. If the table passes, the next pass can honestly move to reachability, TE budgets, residual control, and OP 4 stability.*

Remark A.75.13 (OP status after)**.** *OP 1/4, OP 4, and OP 5(B) remain open. The gain is that the first mixed-readout mode registry now has a concrete selector-table pass/fail diagnostic.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.76 Selector-to-Route Gate

Status. The layer introduced the first selector table. The present layer turns that table into a formal route gate. Before a route is tested for link-state reachability, segment validity, traffic-engineering budgets, residual control, or OP 4 stability, the required first-layer mixed-readout selectors must pass. This is a conservative ordering rule: it prevents route work from proceeding through an edge whose readout compatibility is already rejected.

A.76.1 Selector gate

Definition A.76.1 (First selector gate)**.** *For a route ρ, the first selector gate is the predicate*

$$\mathsf{Gate}_{\chi}^{(1)}(\rho) := [D_{\chi}^{(1)}(\rho) = \varnothing].$$

If $\mathsf{Gate}_{\chi}^{(1)}(\rho)$ holds, the route is selector-admissible at the first mixed-readout mode layer. If it fails, the route is stopped before reachability or TE-budget evaluation.

Definition A.76.2 (Selector-gated admissible edge)**.** *An edge e is first-selector-gated admissible if every first-layer mixed readout required on e has selector value 1:*

$$\forall (e, a, b; m, n) \in \mathsf{Req}^{(1)}(e) : \qquad \chi_{mn}^{ab}(e) = 1.$$

The corresponding gated edge set is

$$E_{\chi}^{(1)} := \{e \in E \mid e \text{ is first-selector-gated admissible}\}.$$

Definition A.76.3 (Selector-gated graph)**.** *Given a link-state graph $\mathcal{G} = (V, E)$, define the selector-gated graph*

$$\mathcal{G}_{\chi}^{(1)} := (V, E_{\chi}^{(1)}).$$

The later admissible and TE-filtered graphs may be formed only after this gate has been applied:

$$\mathcal{G}_{\chi,\mathrm{adm}}^{(1)} \subseteq \mathcal{G}_{\chi}^{(1)}.$$

A.76.2 Gate ordering

Definition A.76.4 (OP diagnostic order after)**.** *The diagnostic order is*

$$\begin{gathered} D_{\chi}^{(1)}(\rho) = \varnothing \\ \Downarrow \\ D_{\mathrm{reach}} = \varnothing \\ \Downarrow \\ D_{\mathrm{OP}}^{\mathrm{seg}} = \varnothing \\ \Downarrow \\ D_{\mathrm{OP}}^{\mathsf{TE}} = \varnothing \\ \Downarrow \\ \textit{residual control and OP 4 stability tests.} \end{gathered}$$

Lemma A.76.5 (Gate failure precedes route failure)**.** *If $D_{\chi}^{(1)}(\rho) \neq \varnothing$, then ρ cannot be accepted as a first-layer readout-compatible OP route, regardless of later reachability or TE-budget data.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. A nonempty selector failure set contains a required mixed-readout pair that is not admitted. Since realizes the relevant compatibility flags through mixed-readout selectors, a route with such a failure lacks the readout compatibility required before later route checks. Later reachability cannot repair a failed selector on a required edge. □

Proposition A.76.6 (Selector-gated reachability is sharper than raw reachability)**.** *Reachability in $\mathcal{G}^{(1)}_{\chi,\mathrm{adm}}$ is a sharper condition than reachability in the raw link-state graph. In particular, if a target is reachable in $\mathcal{G}^{(1)}_{\chi,\mathrm{adm}}$, then it is reachable by a route whose first-layer mixed readout requirements are all admitted.*

Proof. The graph $\mathcal{G}^{(1)}_{\chi,\mathrm{adm}}$ contains only edges that first passed the selector gate and then passed the ordinary admissibility filters. Therefore any path in it is built entirely from selector-admitted edges. Raw reachability may use edges that are graph-theoretically present but readout-incompatible; selector-gated reachability cannot. □

A.76.3 OP 1/4 gate

Definition A.76.7 (OP 1/4 selector gate)**.** *The OP 1/4 selector gate is*

$$\mathsf{Gate}^{\chi,(1)}_{1/4} := \mathsf{Pass}^{\chi,(1)}_{1/4}.$$

Equivalently,

$$\mathsf{Gate}^{\chi,(1)}_{1/4} = [\chi^{\mathrm{QAM,MML}}_{\mathrm{normal,router}}(e_{QM}) = \chi^{\mathrm{MML,sel}}_{\mathrm{router,selector}}(e_{MS}) = \chi^{\mathrm{sel,anc}}_{\mathrm{selector,anchor}}(e_{SA}) = 1].$$

Proposition A.76.8 (OP 1/4 gate consequence)**.** *If $\mathsf{Gate}^{\chi,(1)}_{1/4}$ fails, then OP 1/4 should be repaired at the first failed mixed-readout selector before testing anchor reachability or cover adequacy. If it passes, the OP 1/4 route may proceed to selector-gated reachability and segment checks.*

Proof. By Lemma A.76.5, a failed selector gate blocks first-layer readout-compatible route acceptance. The failed entry identifies whether the repair target is QAM/MML, MML/selector, or selector/anchor. If the gate passes, no first-layer selector entry blocks the later route checks. □

A.76.4 OP 5(B) gate

Definition A.76.9 (OP 5(B) selector gate)**.** *The OP 5(B) selector gate is*

$$\mathsf{Gate}^{\chi,(1)}_{5B} := \mathsf{Pass}^{\chi,(1)}_{5B}.$$

Equivalently, it requires the four entries

$$\chi^{\mathrm{HC},\partial\mathrm{col}}_{\mathrm{boundaryNorm,collar}},\quad \chi^{\partial\mathrm{col,MMA}}_{\mathrm{guard,matrix}},\quad \chi^{\mathrm{MMA,res}}_{\mathrm{residualEval,post}},\quad \chi^{\mathrm{HC,MML/MM}}_{\mathrm{mmCompat,intertwiner}}$$

to be admitted on their declared route edges.

Proposition A.76.10 (OP 5(B) gate consequence)**.** *If $\mathsf{Gate}^{\chi,(1)}_{5B}$ fails, then OP 5(B) should be repaired at the failed mixed-readout selector before boundary-promotion reachability, HC/MM residual, or post-promotion residual checks. If it passes, the OP 5(B) route may proceed to boundary-collar, TE-budget, and residual tests.*

Proof. The proof is the OP 5(B) instance of the selector-gate ordering. A failed first-layer selector blocks readout-compatible boundary-promotion routing. A passed gate removes this first obstruction and allows the later tests to become meaningful. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.76.5 Gate obstruction

Definition A.76.11 (Selector-gate obstruction)**.** *For a family of OP routes $\mathfrak{R}$, define*

$$D^{(1)}_{\mathrm{gate}} := \{\rho \in \mathfrak{R} \mid D^{(1)}_{\chi}(\rho) \neq \varnothing\}.$$

Theorem A.76.12 (selector-to-route reduction)**.** *After the local pass, the first OP-facing route precheck is*

$$D^{(1)}_{\mathrm{gate}} = \varnothing.$$

If this fails, the obstruction is selector/readout compatibility, not reachability, TE-budget, residual control, or OP 4 stability. If it holds, the route family can be passed to the selector-gated reachability and TE layers.

Proof. The selector-gate obstruction is exactly the family of routes with nonempty first selector failures. By the gate-ordering lemma, those routes cannot be accepted before later route tests. Conversely, if no route fails the selector gate, the first mixed-readout compatibility layer is cleared and the next diagnostic layers may be applied. □

Remark A.76.13 (OP status after)**.** *OP 1/4, OP 4, and OP 5(B) remain open. The gain is procedural: the proof search now has a strict first gate. Do not spend work on reachability or TE budgets for a route whose first required mixed-readout selector has already failed.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.77 Selector-Gated Reachability Diagnostic

Status. The layer installed the first selector gate. The present layer performs the next diagnostic pass: reachability is checked only inside the graph whose edges have passed the first mixed-readout selector gate and the ordinary link-state admissibility filters. Thus a reachable route is not merely a graph path; it is a first-layer readout-compatible graph path. OP 1/4, OP 4, and OP 5(B) remain open.

A.77.1 Selector-gated admissible graph

Definition A.77.1 (Selector-gated admissible graph)**.** *Let*

$$\mathcal{G} = (V, E)$$

be the raw OP link-state graph. The first selector-gated admissible graph is

$$\mathcal{G}^{(1)}_{\chi,\mathrm{adm}} := (V, E^{(1)}_{\chi,\mathrm{adm}})$$

where

$$E^{(1)}_{\chi,\mathrm{adm}} := \{e \in E \mid e \in E^{(1)}_{\chi},\ \mathrm{up}(e) = 1,\ \tau(e) \text{ matches},\ \Gamma(e) \text{ preserves active guards}\}.$$

Definition A.77.2 (Selector-gated reachability obstruction)**.** *For a selected target family $\mathcal{Q}$, define*

$$D^{\chi,(1)}_{\mathrm{reach}} := \{q_{\mathrm{tar}} \in \mathcal{Q} \mid q_{\mathrm{tar}} \notin \mathrm{Reach}_{\mathcal{G}^{(1)}_{\chi,\mathrm{adm}}}(q_{\mathrm{clk}})\}.$$

Lemma A.77.3 (Selector-gated reachability implies ordinary reachability)**.** *If*

$$q_{\mathrm{tar}} \in \mathrm{Reach}_{\mathcal{G}^{(1)}_{\chi,\mathrm{adm}}}(q_{\mathrm{clk}}),$$

then q_{tar} is reachable in the ordinary admissible graph $\mathcal{G}_{\mathrm{adm}}$.

Proof. The selector-gated admissible graph is a subgraph of the ordinary admissible graph. Therefore any path in the selector-gated admissible graph is also a path in the ordinary admissible graph. □

Proposition A.77.4 (Selector-gated route certificate)**.** *If*

$$q_{\mathrm{tar}} \in \mathrm{Reach}_{\mathcal{G}^{(1)}_{\chi,\mathrm{adm}}}(q_{\mathrm{clk}}),$$

then there exists a route certificate from q_{clk} to q_{tar} whose edges are first-layer selector compatible, tensor-compatible, active, and guard-preserving.

Proof. Reachability gives a directed path in $\mathcal{G}^{(1)}_{\chi,\mathrm{adm}}$. By construction, every edge of this path is first-layer selector compatible, active, tensor-compatible, and guard-preserving. Recording the path data gives the stated route certificate. □

A.77.2 OP 1/4 selector-gated reachability

Definition A.77.5 (OP 1/4 selector-gated target family)**.** *The OP 1/4 selector-gated target family is*

$$\mathcal{Q}^{\chi}_{1/4} := \{K_i \cap A^{\mathrm{sel}}_{1/4} \mid K_i \in \mathcal{K}_{1/4}\}$$

read inside $\mathcal{G}^{(1)}_{\chi,\mathrm{adm}}$.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition A.77.6 (OP 1/4 selector-gated reachability obstruction)**.** *The OP 1/4 selector-gated reachability obstruction is*

$$D^{\chi,(1)}_{\mathrm{reach},1/4} := \{K_i \cap A^{\mathrm{sel}}_{1/4} \in \mathcal{Q}^{\chi}_{1/4} \mid K_i \cap A^{\mathrm{sel}}_{1/4} \notin \mathrm{Reach}_{\mathcal{G}^{(1)}_{\chi,\mathrm{adm}}}(q_{\mathrm{clk}})\}.$$

Proposition A.77.7 (OP 1/4 first selector-gated reachability pass)**.** *If*

$$\mathsf{Gate}^{\chi,(1)}_{1/4}$$

holds and

$$D^{\chi,(1)}_{\mathrm{reach},1/4} = \varnothing,$$

then every relevant OP 1/4 anchor component has a first-layer readout-compatible clock-to-anchor route certificate.

Proof. The gate condition ensures that the required OP 1/4 selector entries are admitted. Empty selector-gated reachability obstruction says every selected anchor target is reachable in the selector-gated admissible graph. Applying Proposition A.77.4 to each target gives first-layer readout-compatible route certificates. □

A.77.3 OP 5(B) selector-gated reachability

Definition A.77.8 (OP 5(B) selector-gated target)**.** *The OP 5(B) selector-gated target is*

$$q^{\mathrm{tar}}_{5B} := Y^{(B),\mathrm{prom}}_5$$

read inside the selector-gated boundary-sensitive graph

$$\mathcal{G}^{(1),\partial}_{\chi,\mathrm{adm}}.$$

Definition A.77.9 (OP 5(B) selector-gated reachability obstruction)**.** *The OP 5(B) selector-gated reachability obstruction is*

$$D^{\chi,(1)}_{\mathrm{reach},5B} := \begin{cases} \varnothing, & Y^{(B),\mathrm{prom}}_5 \in \mathrm{Reach}_{\mathcal{G}^{(1),\partial}_{\chi,\mathrm{adm}}}(q_{\mathrm{clk}}), \\ \{Y^{(B),\mathrm{prom}}_5\}, & \textit{otherwise.} \end{cases}$$

Proposition A.77.10 (OP 5(B) first selector-gated reachability pass)**.** *If*

$$\mathsf{Gate}^{\chi,(1)}_{5B}$$

holds and

$$D^{\chi,(1)}_{\mathrm{reach},5B} = \varnothing,$$

then there exists a first-layer readout-compatible, boundary-sensitive clock-to-promoted-target route certificate for OP 5(B). The remaining obligations are TE budgets, HC/MM residual compatibility, post-promotion residual control, and OP 4 stability.

Proof. The gate condition admits the required mixed-readout selector entries for the OP 5(B) boundary route. Empty selector-gated reachability obstruction gives a path to $Y^{(B),\mathrm{prom}}_5$ inside the selector-gated boundary-sensitive graph. This path yields a first-layer readout-compatible route certificate. The statement does not discharge later budget, residual, or stability obligations. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.77.4 Selector-gated diagnostic chain

Theorem A.77.11 (selector-gated reachability reduction)**.** *After the local pass, the first two OP-facing route checks are*

$$D_{\chi}^{(1)}(\rho) = \varnothing \qquad \text{and} \qquad D_{\mathrm{reach}}^{\chi,(1)} = \varnothing.$$

If both hold for the selected OP route family, then all selected targets have first-layer readout-compatible admissible route certificates. If the first fails, the repair target is a selector entry. If the second fails, the repair target is missing selector-compatible reachability.

Proof. The first condition is exactly the selector gate. The second condition states that the selected targets are reachable after the selector gate and ordinary admissibility filters have been applied. By the selector-gated route certificate proposition, those reachable targets have first-layer readout-compatible route certificates. Failure of the first condition is a selector failure; failure of the second condition is a reachability failure after selector filtering. □

Remark A.77.12 (OP status after)**.** *OP 1/4, OP 4, and OP 5(B) remain open. The gain is that reachability is now checked in the right graph: not the raw graph and not merely the admissible graph, but the graph already filtered by first-layer mixed-readout selectors.*

A.78 Full Atlas and OP Promotion Chain

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Integrated Delta Layer: Robust Local Chart

Abstract

This delta layer continues the local proof chain photon-frame HC-collar contraction line. The idealized non-ideal sample already produced a local photon-frame near-kernel closure candidate. The present layer tests whether that candidate survives a denser asymmetric chart, non-uniform curvature/readout weights, and explicitly nonzero routed HC/MM absorption. The result is still conservative: it is a local OP5 closure-candidate certificate for the declared chart only, not a global OP5 closure theorem.

A.79 Position of the probe

The chart certified a finite near-kernel photon-frame sample by checking

$$\widehat{\lambda}^{\mathrm{loc},(2)}_{\gamma,K} + \eta_{\mathrm{loc}} < 1, \qquad L^{\mathrm{loc}}_{\mathrm{unr}} = 0.$$

That test was useful but still too symmetric: the sample table used a short four-point chart and encoded the remaining perturbation through a relatively coarse margin. The test strengthens the diagnostic in three ways:

1. the photon-frame chart is denser and intentionally asymmetric;
2. each sample point carries a non-uniform curvature/readout weight;
3. the HC/MM absorption register is allowed to be nonzero, but only as routed debt-free bookkeeping.

The intended question is therefore not whether OP5 is globally closed, but whether the local HC-cone reading remains contraction-compatible after the first robustification of the sample.

A.80 Asymmetric weighted photon-frame chart

Definition A.80.1 (asymmetric photon-frame HC-absorption chart)**.** *An asymmetric photon-frame chart is a finite packet*

$$\mathcal{S}^{\mathrm{asym},(2)}_{\gamma,K} = \{(b_i, q_i, a_i, e_i, w_i, L_i^{\mathrm{rout}}, L_i^{\mathrm{unr}})\}_{i=1}^{8},$$

where:

- $b_i > 0$ *is the unshifted photon-frame radial readout size;*
- q_i *is the normalized second-shift radius ratio before explicit absorption;*
- a_i *is the normalized routed* HC/MM*-absorption contribution;*
- e_i *is the normalized finite chart error/tolerance contribution;*
- $w_i > 0$ *is a declared non-uniform readout weight;*
- L_i^{rout} *records routed leakage/absorption bookkeeping;*
- L_i^{unr} *records leakage that is not routed through an admissible* HC/MM *register.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

The pointwise weighted contraction ratio is bounded by

$$\lambda_i^{\mathrm{loc},(2)} \le q_i + a_i + e_i.$$

The chart estimator and routed/unrouted guards are

$$\widehat{\lambda}_{\gamma,K}^{\mathrm{loc},(2)} := \max_{1\le i\le 8} \lambda_i^{\mathrm{loc},(2)}, \qquad L_{\mathrm{unr}}^{\mathrm{loc}} := \sum_{i=1}^{8} L_i^{\mathrm{unr}}, \qquad B_{\mathrm{HC/MM}}^{\mathrm{loc}} := \mathrm{unpaid}(L^{\mathrm{rout}}).$$

Here $B_{\mathrm{HC/MM}}^{\mathrm{loc}} = 0$ means that nonzero absorption has been declared and routed without leaving residual absorption debt.

Definition A.80.2 (local closure-candidate classifier)**.** *The asymmetric photon-frame branch is classified by*

$$\mathrm{Class}_{\gamma,K}^{\mathrm{loc}} = \begin{cases} \mathrm{cert_red}_{\gamma,\mathrm{asym}}, & \widehat{\lambda}_{\gamma,K}^{\mathrm{loc},(2)} + \eta_{\mathrm{loc}} < 1,\ L_{\mathrm{unr}}^{\mathrm{loc}} = 0,\ B_{\mathrm{HC/MM}}^{\mathrm{loc}} = 0, \\ \mathrm{comp_margin}_{\gamma,\mathrm{asym}}, & \widehat{\lambda}_{\gamma,K}^{\mathrm{loc},(2)} + \eta_{\mathrm{loc}} < 1,\ L_{\mathrm{unr}}^{\mathrm{loc}} = 0,\ B_{\mathrm{HC/MM}}^{\mathrm{loc}} > 0, \\ \mathrm{debt_rout}_{\gamma,\mathrm{asym}}, & \widehat{\lambda}_{\gamma,K}^{\mathrm{loc},(2)} + \eta_{\mathrm{loc}} < 1,\ L_{\mathrm{unr}}^{\mathrm{loc}} > 0 \text{ but routeable}, \\ \mathrm{res_proj}_{\gamma,\mathrm{asym}}, & \widehat{\lambda}_{\gamma,K}^{\mathrm{loc},(2)} + \eta_{\mathrm{loc}} \ge 1 \text{ or leakage is not routeable.} \end{cases}$$

The certificate $\mathrm{cert_red}_{\gamma,\mathrm{asym}}$ *is local to the declared chart and is read as* $\mathrm{close_cand}_{\gamma,\mathrm{asym}}$*, not as a global OP5 theorem.*

A.81 Explicit sample table

The following sample is deliberately not a uniform halving table. The base radii are uneven, the weights are non-uniform, and the routed HC/MM-absorption contribution is nonzero in every row.

i	b_i	w_i	q_i	a_i	e_i	$q_i+a_i+e_i$	L_i^{unr}
1	1	1	1/2	1/64	1/64	17/32	0
2	29/32	33/32	17/32	1/64	1/32	37/64	0
3	23/32	17/16	9/16	1/32	1/64	39/64	0
4	17/32	19/16	19/32	1/32	1/32	21/32	0
5	13/32	5/4	5/8	1/64	1/64	21/32	0
6	3/8	21/16	21/32	1/64	1/64	11/16	0
7	9/32	11/8	5/8	1/32	1/32	11/16	0
8	5/32	3/2	19/32	1/64	1/32	41/64	0

All rows carry routed absorption $a_i > 0$. We declare the routed register as paid,

$$L_{\mathrm{rout}}^{\mathrm{loc}} > 0, \qquad B_{\mathrm{HC/MM}}^{\mathrm{loc}} = 0, \qquad L_{\mathrm{unr}}^{\mathrm{loc}} = 0.$$

The maximum row value is

$$\widehat{\lambda}_{\gamma,K}^{\mathrm{loc},(2)} = \max_i (q_i + a_i + e_i) = \frac{11}{16}.$$

With the asymmetric chart margin

$$\eta_{\mathrm{loc}} = \frac{9}{64},$$

one obtains

$$\widehat{\lambda}_{\gamma,K}^{\mathrm{loc},(2)} + \eta_{\mathrm{loc}} = \frac{11}{16} + \frac{9}{64} = \frac{53}{64} < 1.$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proposition A.81.1 (asymmetric HC-absorption robustness certificate)**.** *For the explicit table above, the asymmetric photon-frame branch receives the local classifier*

$$\mathrm{Class}^{\mathrm{loc}}_{\gamma,K} = \mathrm{cert_red}_{\gamma,\mathrm{asym}} = \mathrm{close_cand}_{\gamma,\mathrm{asym}}.$$

Equivalently, the local closure-candidate status survives the first denser asymmetric weighted chart with nonzero routed HC/MM *absorption.*

Proof. The estimator is the maximum of the row bounds in the explicit table. The largest row value is 11/16, attained in rows 6 and 7. The declared margin is 9/64, hence

$$\widehat{\lambda}^{\mathrm{loc},(2)}_{\gamma,K} + \eta_{\mathrm{loc}} = 53/64 < 1.$$

The table has zero unrouted leakage in every row, so $L^{\mathrm{loc}}_{\mathrm{unr}} = 0$. The absorption register is nonzero, because $a_i > 0$ in every row, but it is explicitly routed and declared debt-free, so $B^{\mathrm{loc}}_{\mathrm{HC/MM}} = 0$. The first branch of the classifier therefore applies. This proves the stated local certificate. □

A.82 Field-equation bookkeeping update

The branch adds the asymmetric photon-frame contraction register

$$\mathcal{R}^{(2)}_{\gamma,\mathrm{asym}} := \left(\widehat{\lambda}^{\mathrm{loc},(2)}_{\gamma,K}, \eta_{\mathrm{loc}}, L^{\mathrm{loc}}_{\mathrm{rout}}, L^{\mathrm{loc}}_{\mathrm{unr}}, B^{\mathrm{loc}}_{\mathrm{HC/MM}}, \{w_i\}_{i=1}^{8}\right)$$

to the OP5-facing source-return bookkeeping. The readout-side implication is

$$\mathcal{R}^{(2)}_{\gamma,\mathrm{asym}} \vdash \left[\widehat{\lambda} + \eta < 1,\ L_{\mathrm{unr}} = 0,\ B_{\mathrm{HC/MM}} = 0\right] \Longrightarrow \mathrm{close_cand}_{\gamma,\mathrm{asym}}.$$

Thus the HC-cone interpretation is strengthened from an ideal or near-ideal photon-frame sample to a first robust asymmetric weighted sample. The correct structural reading is that OP5 remains more plausibly an HC-cone/pass-count debt problem than an independent terminal endpoint problem, provided the remaining chart families keep satisfying the same contraction and routing guards.

A.83 Conservativity guards

The following guards are part of the layer.

1. No global OP5 closure is claimed.
2. The certificate is local to the declared finite photon-frame chart.
3. Nonzero HC/MM absorption is not deletion of residuals; it is paid routed bookkeeping.
4. The chart does not assert a literal Euclidean photon trajectory.
5. The result does not close OP1, OP4, QCD/confinement, CKM/PMNS, or the quark-mass spectrum.
6. Promotion beyond $\mathrm{close_cand}_{\gamma,\mathrm{asym}}$ still requires either a covering argument over admissible photon-frame charts or a finite atlas theorem with stable contraction margins.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.84 Next recommended step

The highest-leverage next step is an atlas-style finite cover test: define a small family of admissible asymmetric photon-frame charts

$$\mathfrak{A}_{\gamma}^{(2)} = \{\mathcal{S}_{\gamma,K,j}^{\mathrm{asym},(2)}\}_{j=1}^{M}$$

and require a uniform bound

$$\sup_{j\le M}\left(\widehat{\lambda}_{\gamma,K,j}^{(2)} + \eta_j\right) < 1, \qquad \sum_{j\le M} L_{\mathrm{unr},j} = 0, \qquad \sum_{j\le M} B_{\mathrm{HC/MM},j} = 0.$$

If such a finite-atlas certificate is obtained, the OP5=HC-cone hypothesis can be promoted from a single-chart local closure candidate to a finite-atlas closure candidate.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Integrated Delta Layer: Finite Atlas

Abstract

This delta layer upgrades the local proof chain photon-frame HC-collar contraction probes from single-chart diagnostics to a finite-atlas closure-candidate test. The layer also states, explicitly and conservatively, what would still have to be proved in order to close OP5 under the hypothesis that OP5 is the HC-cone/reflector surface of the source-return route rather than an independent endpoint obstruction. The result is an atlas-level closure candidate, not a final OP5 theorem.

A.85 Position after

The layer showed that the local photon-frame candidate survives a denser asymmetric chart with nonzero but debt-free routed HC/MM absorption:

$$\widehat{\lambda}^{\mathrm{loc},(2)}_{\gamma,K} = \frac{11}{16}, \qquad \eta_{\mathrm{loc}} = \frac{9}{64}, \qquad \widehat{\lambda}^{\mathrm{loc},(2)}_{\gamma,K} + \eta_{\mathrm{loc}} = \frac{53}{64} < 1.$$

This is strong enough to justify an atlas test but not strong enough to close OP5. A single chart only says that one declared readout neighbourhood is compatible with the HC-cone contraction picture. OP5 closure requires that the admissible OP5 branch is exhausted by such neighbourhoods, and that the transition logic between them does not reintroduce an endpoint obstruction.

A.86 Finite photon-frame atlas

Definition A.86.1 (finite photon-frame HC-cone atlas)**.** *An finite photon-frame atlas for the OP5-facing branch is a finite family*

$$\mathfrak{A}^{(2)}_{\gamma} = \{\mathcal{S}^{(2)}_{\gamma,K,j}\}_{j=1}^{M}$$

of admissible photon-frame HC*-collar charts. Each chart carries the data*

$$\mathcal{S}^{(2)}_{\gamma,K,j} \rightsquigarrow \left(\widehat{\lambda}_j, \eta_j, L_{\mathrm{unr},j}, B_{\mathrm{HC/MM},j}, \tau_j\right),$$

where $\widehat{\lambda}_j$ is the finite contraction estimator, η_j is the chart margin, $L_{\mathrm{unr},j}$ is the unrouted leakage register, $B_{\mathrm{HC/MM},j}$ is the unpaid HC/MM*-absorption debt, and τ_j records the transition-map debt between overlapping charts.*

Definition A.86.2 (Atlas classifier)**.** *The atlas receives the classifier by the following finite rule.*

$$\mathrm{Class}(\mathfrak{A}^{(2)}_{\gamma}) = \mathrm{atlas_close_cand}$$

exactly if

$$\sup_{j\le M}(\widehat{\lambda}_j + \eta_j) < 1, \qquad \sum_{j\le M} L_{\mathrm{unr},j} = 0, \qquad \sum_{j\le M} B_{\mathrm{HC/MM},j} = 0, \qquad \sum_{j\le M} \tau_j = 0.$$

It receives the compatible-open classifier cert_comp *if the uniform contraction inequality holds but some leakage, absorption, or transition register is nonzero while still routeable. It receives the residual-projection classifier* res_proj *if the uniform contraction bound fails or if any register is not routeable. Thus an atlas is not merely a list of successful sample tables. It also has to verify that no obstruction appears on overlaps or transitions.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.87 Explicit finite-atlas sample

The following five-chart atlas is a compact first test. It collects the already motivated ideal, near-kernel, non-ideal, asymmetric, and tilted/edge regimes into one finite certificate table.

Chart	$\widehat{\lambda}_j$	η_j	$\widehat{\lambda}_j + \eta_j$	$L_{\mathrm{unr},j}$	$B_{\mathrm{HC/MM},j} + \tau_j$
central ideal collar	1/2	1/8	5/8	0	0
near-kernel symmetric	21/32	1/8	25/32	0	0
non-ideal sample	21/32	1/8	25/32	0	0
asymmetric HC-absorption	11/16	9/64	53/64	0	0
tilted edge collar	45/64	7/64	13/16	0	0

The atlas maximum is therefore

$$\Lambda_{\mathfrak{A}}^{\mathrm{loc}} := \sup_{j \le 5} (\widehat{\lambda}_j + \eta_j) = \frac{53}{64} < 1.$$

All explicit leakage, absorption-debt, and transition-debt registers are declared zero at the atlas level:

$$\sum_{j \le 5} L_{\mathrm{unr},j} = 0, \qquad \sum_{j \le 5} B_{\mathrm{HC/MM},j} = 0, \qquad \sum_{j \le 5} \tau_j = 0.$$

Proposition A.87.1 (finite-atlas closure-candidate certificate)**.** *The explicit five-chart photon-frame atlas above receives the classifier*

$$\mathrm{Class}(\mathfrak{A}_{\gamma}^{(2)}) = \mathrm{atlas_close_cand}.$$

It promotes the single-chart result to a finite-atlas OP5 closure candidate.

Proof. The largest value in the third numerical column is 53/64, which is strictly smaller than one. The declared atlas registers satisfy

$$\sum_{j \le 5} L_{\mathrm{unr},j} = 0, \qquad \sum_{j \le 5} B_{\mathrm{HC/MM},j} = 0, \qquad \sum_{j \le 5} \tau_j = 0.$$

The first branch of the atlas classifier applies. Hence the finite atlas receives the atlas closure-candidate status. □

A.88 What would be needed to close OP5

The atlas is still a candidate because it assumes, rather than proves, that the five declared chart types cover all admissible OP5-facing photon-frame readouts. Under the working hypothesis

$$\mathrm{OP5} = \text{HC-cone/reflector surface of the source-return route},$$

OP5 would be closed by proving the following theorem.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Theorem A.88.1 (Conditional OP5 closure theorem to be proved)**.** *Assume that the OP5-facing source-return route factorizes through the* HC*-cone reflector,*

$$R_{\mathrm{OP}}^{\text{HC-cone}} = T_{\Gamma}^{\vee} \circ J_{\mathrm{HC}}^{\mathrm{cone}} \circ T_{\Gamma},$$

and that its cone-factorization obstruction vanishes globally,

$$D^{\chi,(1)}_{\text{OP5=HC-cone}} = 0.$$

Then OP5 closes if there exists a finite photon-frame atlas $\mathfrak{A}_{\gamma}^{(2)}$ *satisfying all of the following conditions.*

1. ***Coverage.*** *Every admissible OP5-facing readout state is contained in at least one chart of* $\mathfrak{A}_{\gamma}^{(2)}$*.*
2. ***Uniform contraction.*** *There is a constant* $\Lambda < 1$ *such that*

$$\widehat{\lambda}_j + \eta_j \le \Lambda < 1 \qquad \text{for every chart } j.$$

3. ***Transition stability.*** *On every nonempty chart overlap, the transition map preserves the contraction class and creates no unpaid transition debt:*

$$\tau_{jk} = 0 \qquad \text{for all admissible overlaps } j \cap k.$$

4. ***No unrouted leakage.*** *Every leakage term is either zero or routed through an admissible* HC/MM *register, and the total unrouted leakage vanishes:*

$$L^{\mathfrak{A}}_{\mathrm{unr}} = 0.$$

5. ***No unpaid absorption debt.*** *Routed* HC/MM *absorption may be nonzero, but all such absorption must be paid inside the route bookkeeping:*

$$B^{\mathfrak{A}}_{\mathrm{HC/MM}} = 0.$$

6. ***Periapsis/no-through-collapse guard.*** *The active* HC*-collar forces every admissible source-return branch to have a collar-side periapsis rather than a through-collapse endpoint.*
7. ***Pass-count consistency.*** *The OP5 pass count is exactly the source-return pass count seen through the second Hilbert-shift/*HC*-cone frame, not an additional independent terminal transport pass.*
8. ***Statement identification.*** *The formal statement of OP5 is shown to be equivalent to the absence of the above cone-factorization, leakage, debt, and transition obstructions.*

If all eight items hold, then OP5 is not an independent residual endpoint problem. It is discharged as the HC*-cone source-return contraction theorem.*

Remark A.88.2 (Why does not yet close OP5)**.** *The table verifies uniform contraction for a finite declared atlas and records zero debt in the declared registers. It does not yet prove the coverage item, the global vanishing of* $D^{\chi,(1)}_{\text{OP5=HC-cone}}$*, or the equivalence between the formal OP5 statement and the* HC*-cone source-return theorem. Those are the remaining proof burdens.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.89 Practical proof route after

The shortest conservative route from to a genuine OP5 closure proof is:

1. write the admissible OP5 readout space $\mathcal{X}_{\mathrm{OP5}}^{\mathrm{adm}}$;
2. prove that the five chart types form a finite cover of $\mathcal{X}_{\mathrm{OP5}}^{\mathrm{adm}}$;
3. prove that all chart overlaps have zero transition debt;
4. prove that $D^{\chi,(1)}_{\mathrm{OP5=HC\text{-}cone}} = 0$ under the HC-cone factorization hypothesis;
5. prove that no admissible branch crosses the collapse core under an active HC-collar, i.e. every branch has a periapsis/readout-turning point;
6. identify OP5's original obstruction statement with the atlas obstruction package.

If these six tasks are completed, the numerical contraction certificates become supporting evidence for a structural theorem rather than isolated probes.

A.90 Conservativity guards

The layer uses the following guards.

1. The result is a finite-atlas closure candidate, not a final OP5 closure.
2. The five-chart cover is declared as a test atlas; its coverage of all admissible OP5 states remains to be proved.
3. The HC-cone interpretation remains conditional on vanishing cone-factorization obstruction.
4. Nonzero HC/MM absorption is permitted only as explicit routed bookkeeping, never as residual deletion.
5. The result does not imply OP1, OP4, QCD/confinement, CKM/PMNS, or quark-mass spectrum closure.

A.91 Next recommended step

The highest-leverage next step is to define the admissible OP5 readout space

$$\mathcal{X}_{\mathrm{OP5}}^{\mathrm{adm}}$$

and prove a finite-cover lemma

$$\mathcal{X}_{\mathrm{OP5}}^{\mathrm{adm}} \subseteq \bigcup_{j=1}^{5} \mathcal{S}^{(2)}_{\gamma,K,j}.$$

That lemma is the first real bridge from atlas diagnostics to OP5 closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Integrated Delta Layer: Tunnel Menu

Abstract

This delta layer inserts the nonstandard-halo step between the robust photon-frame contraction probe and the finite-atlas closure-candidate logic. Its purpose is to reduce the apparent combinatorial overload of OP5: instead of testing all admissible source-return routes as free trajectories, the OP5-facing branch is first trapped in a double-epsilon halo around the collapse kernel. Exit from this halo is then possible only through explicitly gated tunnel points. The result is conservative: OP5 is not closed here. The proof burden is reduced from an all-routes problem to a finite or finitely classifiable tunnel-point problem, provided the stated halo-capture, halo-invariance, and tunnel-gating hypotheses are later proved.

A.92 Version position

The immediate pre-line contains two relevant ingredients. First, produced a robust asymmetric photon-frame local certificate with

$$\widehat{\lambda}^{\mathrm{loc},(2)}_{\gamma,K} = \frac{11}{16}, \qquad \eta_{\mathrm{loc}} = \frac{9}{64}, \qquad \widehat{\lambda}^{\mathrm{loc},(2)}_{\gamma,K} + \eta_{\mathrm{loc}} = \frac{53}{64} < 1.$$

Second, an finite-atlas draft upgrades this local logic to a declared finite atlas, but such an atlas still needs an independent explanation of why the admissible OP5 search does not branch into uncontrolled route families. The present layer supplies that missing structural filter: a double-epsilon halo channel with tunnel-point reduction.

A.93 Double-epsilon halo algebra

Definition A.93.1 (Two-scale halo readout algebra)**.** *Let $\mathcal{A}$ be the OP5-facing readout algebra and let K denote the collapse kernel. A two-scale halo readout is a triple*

$$(x; \rho_r(x), \rho_h(x)) \in \mathcal{A} \times \mathbb{R}_{\geq 0} \times \mathbb{R}_{\geq 0},$$

where $\rho_r(x)$ is the radial kernel-distance readout and $\rho_h(x)$ is the HC*-cone residual readout. The two infinitesimal windows are denoted*

$$\varepsilon_r > 0, \qquad \varepsilon_h > 0.$$

The corresponding double-epsilon halo is

$$\mathsf{Halo}_{\varepsilon_r,\varepsilon_h}(K) := \{x \in \mathcal{X}^{\mathrm{adm}}_{\mathrm{OP5}} : \rho_r(x) \leq \varepsilon_r \text{ and } \rho_h(x) \leq \varepsilon_h\}.$$

The first inequality is radial closeness to the collapse kernel. The second inequality is closeness to the HC*-cone/reflector normal form.*

Remark A.93.2 (Why two epsilon channels are needed)**.** *A one-parameter halo around K only records radial nearness. OP5 is not merely a radial-distance problem; the relevant question is whether the source-return route is already captured by the active* HC*-cone collar. Therefore the halo must carry two channels: one for kernel nearness and one for* HC*-normality. In this sense ε_r controls "near the kernel", while ε_h controls "near the* HC*-cone route".*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition A.93.3 (Halo boundary and exit side)**.** *The halo boundary is the union of the radial and* HC*-residual boundary faces:*

$$\partial\mathsf{Halo}_{\varepsilon_r,\varepsilon_h}(K) = \{\rho_r = \varepsilon_r, \rho_h \le \varepsilon_h\} \cup \{\rho_r \le \varepsilon_r, \rho_h = \varepsilon_h\}.$$

An exit candidate is a point

$$p \in \partial\mathsf{Halo}_{\varepsilon_r,\varepsilon_h}(K)$$

for which the source-return update points outward in at least one of the two halo channels before applying the tunnel selector.

A.94 Tunnel points

Definition A.94.1 (Tunnel candidate)**.** *A point $p \in \partial\mathsf{Halo}_{\varepsilon_r,\varepsilon_h}(K)$ is a tunnel candidate if it satisfies the four tests*

$$\begin{aligned} &\textit{(T1)} \quad p \in \partial\mathsf{Halo}_{\varepsilon_r,\varepsilon_h}(K), \\ &\textit{(T2)} \quad D_{\text{HC-cone}}(p) \le \varepsilon_h, \\ &\textit{(T3)} \quad \chi_{\text{tun}}(p) = 1, \\ &\textit{(T4)} \quad \operatorname{rank} J_{\mathsf{Exit}}(p) < \operatorname{rank}_{\max}. \end{aligned}$$

*Here χ_{tun} is the tunnel selector and J_{Exit} is the exit Jacobian for the paired radial/*HC*-residual channel. The resulting candidate set is*

$$\mathcal{T}_{\text{tun}}^{\text{cand}}(\varepsilon_r, \varepsilon_h) := \{p \in \partial\mathsf{Halo}_{\varepsilon_r,\varepsilon_h}(K) : \textit{(T1)–(T4) hold}\}.$$

Definition A.94.2 (Resolved tunnel point)**.** *A tunnel candidate $p \in \mathcal{T}_{\text{tun}}^{\text{cand}}$ is resolved if exactly one of the following outcomes is certified:*

$$\mathsf{Gate}(p) \in \{\text{blocked}, \text{HC/MM absorbed}, \text{atlas routed}, \text{residual projected}\}.$$

The first three outcomes are controlled. The fourth outcome records a remaining obstruction and cannot be used to close OP5.

Remark A.94.3 (Tunnel points as the replacement for route explosion)**.** *The intended reduction is not that particles literally escape through geometric holes. The tunnel language is a readout-side compression: a halo-trapped branch can only leave the halo where the radial boundary, the* HC*-cone residual boundary, and the selector/rank conditions simultaneously permit an exit. Thus an apparent continuum of route choices is replaced by a finite or finitely classifiable family of boundary obstructions.*

A.95 Capture, invariance, and tunnel reduction

Definition A.95.1 (Halo capture)**.** *The OP5-facing route satisfies halo capture if every admissible readout enters the double halo after finitely many* HC*-normalized source-return updates:*

$$\forall x \in \mathcal{X}_{\text{OP5}}^{\text{adm}}\ \exists n < \infty : R_{\text{HC}}^n(x) \in \mathsf{Halo}_{\varepsilon_r,\varepsilon_h}(K).$$

Definition A.95.2 (Halo invariance off tunnels)**.** *The double halo is invariant off tunnel candidates if*

$$x \in \mathsf{Halo}_{\varepsilon_r,\varepsilon_h}(K) \setminus \mathcal{T}_{\text{tun}}^{\text{cand}} \quad \Longrightarrow \quad R_{\text{HC}}(x) \in \mathsf{Halo}_{\varepsilon_r,\varepsilon_h}(K).$$

Equivalently, the active HC*-collar produces a periapsis/turning-point readout before a through-collapse exit can occur, unless the state lies on a declared tunnel candidate.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proposition A.95.3 (tunnel-point reduction)**.** *Assume halo capture and halo invariance off tunnels. Then every admissible OP5-facing branch is either permanently trapped in the double-epsilon halo or reaches a point of* $\mathcal{T}_{\mathrm{tun}}^{\mathrm{cand}}(\varepsilon_r, \varepsilon_h)$*. Consequently, the OP5 route problem reduces to resolving the tunnel candidate set.*

Proof. Let $x \in \mathcal{X}_{\mathrm{OP5}}^{\mathrm{adm}}$. By halo capture, there is a finite n such that $x_n := R_{\mathrm{HC}}^n(x) \in \mathsf{Halo}_{\varepsilon_r,\varepsilon_h}(K)$. If no later iterate lies in $\mathcal{T}_{\mathrm{tun}}^{\mathrm{cand}}$, halo invariance implies by induction that every later iterate remains in $\mathsf{Halo}_{\varepsilon_r,\varepsilon_h}(K)$. If a later iterate fails to remain in the halo, the contrapositive of halo invariance says that the failure can occur only at a tunnel candidate. Hence every branch is trapped or reaches the tunnel candidate set. This proves the reduction. □

A.96 Finite tunnel menu

The first conservative tunnel inventory is not a numerical discovery claim. It is a finite classification scheme for the exits that must be tested.

Definition A.96.1 (First OP5 tunnel menu)**.** *The first tunnel menu is*

$$\mathfrak{T}_{\mathrm{OP5}}^{\mathrm{loc}} = \{T_r, T_h, T_{rh}, T_\chi, T_{\mathrm{rk}}, T_{\mathrm{ov}}\},$$

where:

Class	*Meaning*
T_r	*radial-face exit:* $\rho_r = \varepsilon_r,\ \rho_h < \varepsilon_h$*;*
T_h	HC*-residual-face exit:* $\rho_r < \varepsilon_r,\ \rho_h = \varepsilon_h$*;*
T_{rh}	*corner exit:* $\rho_r = \varepsilon_r,\ \rho_h = \varepsilon_h$*;*
T_χ	*selector-switch exit caused by a change in* χ_{tun}*;*
T_{rk}	*rank-drop exit caused by* $\operatorname{rank} J_{\mathsf{Exit}} < \mathrm{rank}_{\max}$*;*
T_{ov}	*atlas-overlap exit caused by transition debt between two photon-frame charts.*

A tunnel candidate must be assigned to at least one class in $\mathfrak{T}_{\mathrm{OP5}}^{\mathrm{loc}}$*.*

Proposition A.96.2 (Finite tunnel classification target)**.** *If the candidate tunnel set admits a finite decomposition*

$$\mathcal{T}_{\mathrm{tun}}^{\mathrm{cand}}(\varepsilon_r, \varepsilon_h) = \bigsqcup_{\alpha \in \mathfrak{T}_{\mathrm{OP5}}^{\mathrm{loc}}} \mathcal{T}_\alpha$$

and every nonempty $\mathcal{T}_\alpha$ *has one of the controlled outcomes*

$$\mathsf{Gate}(\mathcal{T}_\alpha) \in \{\text{blocked}, \text{HC/MM absorbed}, \text{atlas routed}\},$$

then OP5 has no uncontrolled tunnel escape inside the declared double-epsilon halo.

Proof. By the preceding tunnel-point reduction, every non-trapped branch can only exit through $\mathcal{T}_{\mathrm{tun}}^{\mathrm{cand}}$. The finite decomposition assigns each such candidate to one of the declared tunnel classes. If every class is blocked, absorbed without unpaid debt, or routed into an atlas chart, then no class contributes an uncontrolled exit. Therefore the double halo has no uncontrolled tunnel escape. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.97 Connection to the finite atlas

Once a tunnel point is atlas-routed, it must land in a contraction chart rather than in a free branch. The bridge to the finite-atlas line is therefore:

$$\mathcal{T}_{\mathrm{tun}}^{\mathrm{atlas}} \subseteq \bigcup_{j=1}^{M} \mathcal{S}_{\gamma,K,j}^{(2)}, \qquad \sup_{j\le M}(\widehat{\lambda}_j + \eta_j) < 1,$$

with the same zero-debt guards

$$L_{\mathrm{unr}}^{\mathfrak{A}} = 0, \qquad B_{\mathrm{HC/MM}}^{\mathfrak{A}} = 0, \qquad \tau^{\mathfrak{A}} = 0.$$

For typographic clarity, the intended symbol in the preceding display is the chart margin η_j. Thus the atlas-routed tunnel branch inherits the atlas criterion exactly when all tunnel exits are covered by charts with a uniform strict contraction margin.

A.98 Conditional OP5 closure statement after the halo step

Theorem A.98.1 (Conditional OP5 closure via double-epsilon halo)**.** *Assume the working identification*

$$\mathrm{OP5} = \mathrm{HC}\text{-}\textit{cone source-return reflector problem}$$

and assume that the cone-factorization obstruction vanishes,

$$D_{\mathrm{OP5=HC\text{-}cone}}^{\chi,(1)} = 0.$$

Then OP5 closes if the following six conditions are proved.

1. ***Halo capture:*** *every admissible OP5 readout enters* $\mathsf{Halo}_{\varepsilon_r,\varepsilon_h}(K)$ *after finitely many* HC*-normalized source-return updates.*
2. ***Halo invariance off tunnels:*** *the double halo is invariant outside* $\mathcal{T}_{\mathrm{tun}}^{\mathrm{cand}}$*.*
3. ***Finite tunnel classification:*** $\mathcal{T}_{\mathrm{tun}}^{\mathrm{cand}}$ *is empty, finite, or decomposes into finitely many tunnel classes.*
4. ***Tunnel gating:*** *each nonempty tunnel class is blocked, debt-free* HC/MM*-absorbed, or routed into a contraction atlas.*
5. ***No hidden leakage or debt:*** *all tunnel and atlas registers satisfy*

$$L_{\mathrm{unr}} = 0, \qquad B_{\mathrm{HC/MM}} = 0, \qquad \tau = 0.$$

6. ***OP5 statement identification:*** *the original OP5 obstruction is equivalent to the cone-factorization/tunnel/atlas obstruction package above.*

Under these hypotheses the OP5 endpoint problem is exhausted by controlled halo trapping and controlled tunnel resolution. Hence OP5 is discharged as a double-epsilon HC*-cone halo theorem rather than as an independent terminal transport obstruction.*

Remark A.98.2 (What proves and what remains)**.** *The layer proves the formal reduction: capture plus invariance off tunnels reduces OP5 to tunnel resolution. It does not yet prove capture, invariance, finite tunnel classification, or global vanishing of* $D_{\mathrm{OP5=HC\text{-}cone}}^{\chi,(1)}$*. Those become the next concrete proof obligations.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.99 Immediate next proof tasks

The next executable line should be:

1. instantiate ρ_r and ρ_h from the current photon-frame HC-collar readout;
2. compute the first symbolic boundary faces $\rho_r = \varepsilon_r$ and $\rho_h = \varepsilon_h$;
3. define the exit Jacobian J_{Exit} in the two-channel coordinates;
4. test whether the first tunnel menu $\mathfrak{T}^{\mathrm{loc}}_{\mathrm{OP5}}$ is complete for the declared halo grammar;
5. route each nonempty tunnel class to blocked, HC/MM-absorbed, atlas-routed, or residual-projected status.

If these tasks succeed, the finite atlas is no longer trying to cover arbitrary OP5 routes. It only has to cover the gated tunnel exits of the double-epsilon halo.

A.100 Conservativity guards

1. does not claim global OP5 closure.
2. The halo is a readout-side structural filter, not a claim that physical particles literally sit inside a geometric shell.
3. Tunnel points are selector/rank/obstruction points, not literal wormholes.
4. HC/MM-absorption is allowed only as explicit debt-free bookkeeping, never as residual deletion.
5. The finite tunnel menu is a first classification target and must still be tested for completeness.
6. OP1, OP4, QCD/confinement, CKM/PMNS, and the quark-mass spectrum remain untouched by this layer.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Integrated Delta Layer: Halo Channel and Response Alphabet

Abstract

This delta layer executes the next proof obligation left by the local pass. The double-epsilon halo symbols are made typographically stable, the two halo channels are instantiated in the photon-frame near-kernel collar, and the exit Jacobian is defined as the differential of the paired radial/HC-residual channel. The result is a first conservative tunnel-menu execution: within the declared photon-frame halo grammar every boundary exit is assigned to one of the six tunnel classes. The layer does not close OP5 globally. It converts the next obstruction into explicit tests: collar admissibility, channel regularity, selector gating, rank-drop isolation, atlas-overlap compatibility, and zero hidden leakage/debt.

A.101 Macro hygiene and version position

The layer introduced the double-epsilon halo but the visual notation for the halo and exit operators must be kept stable in downstream TeX. In this layer the intended printed symbols are

$$\mathsf{Halo}, \qquad \mathsf{Exit}, \qquad J_{\mathsf{Exit}}, \qquad \mathcal{E}_{\mathsf{Exit}}.$$

For source-level robustness the lower-case spellings are aliased to the same mathematical symbols, but the body of this layer uses the capitalized operators. Thus the rendered document should not display raw control words for the halo or exit operators.

The reduction says that the OP5 route problem is reduced to tunnel resolution once halo capture and halo invariance off tunnels are proved. The present layer executes the first concrete part of that program: it instantiates the two halo channels in the photon-frame HC-collar chart and defines the exit Jacobian which detects rank-drop tunnel candidates.

A.102 Photon-frame collar chart

Definition A.102.1 (Photon-frame HC-collar chart)**.** *Let $U_{\gamma,K} \subset \mathcal{X}^{\mathrm{adm}}_{\mathrm{OP5}}$ be an admissible near-kernel photon-frame chart. The chart carries:*

$$d^G_K(x), \qquad r^{(2)}_\gamma(x), \qquad h^{(2)}_\gamma(x), \qquad q^{(2)}_\gamma(x) := (h^{(2)}_\gamma(x))^2 - (r^{(2)}_\gamma(x))^2,$$

where d^G_K is the metric distance to the collapse kernel, $r^{(2)}_\gamma$ and $h^{(2)}_\gamma$ are the second-shift radial and HC-height coordinates, and $q^{(2)}_\gamma$ is the photon-frame cone residual. The active first-shift HC collar radius is denoted $r^{(1)}_{\mathrm{HC}}(x) > 0$. The admissible collar guard is

$$d^G_K(x) \geq r^{(1)}_{\mathrm{HC}}(x) > 0.$$

This guard prevents interpreting a projected line through K as a literal through-collapse trajectory.

Definition A.102.2 (Second-shift tilt operator)**.** *The second-shift cone-to-radial tilt is*

$$\Theta^{(2)}_{\gamma,K} := \Pi^{\mathrm{HC},\gamma}_{\mathrm{cone}\to\mathrm{rad}} \circ U^{(2)}_\pi - U^{(2)}_\pi \circ \Pi^{\mathrm{HC},\gamma}_{\mathrm{cone}\to\mathrm{rad}}.$$

It records the failure of radial projection and the second Hilbert shift to commute. A projected path may visually pass through K after the second shift; in this layer that appearance is read first as tilt debt rather than as immediate OP5 failure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.103 Concrete halo-channel instantiation

Definition A.103.1 (Radial halo channel). *The radial halo channel is*

$$\rho_r^\gamma(x) := d_K^G\big(\Pi^{\mathrm{HC},\gamma}_{\mathrm{cone}\to\mathrm{rad}}(U_\pi^{(2)}x), K\big).$$

The radial halo face is

$$\rho_r^\gamma(x) = \varepsilon_r.$$

The photon-frame near-kernel collar is admissible only on the guarded subregion

$$r_{\mathrm{HC}}^{(1)}(x) \le \rho_r^\gamma(x) \le \varepsilon_r.$$

Thus $\rho_r^\gamma \le \varepsilon_r$ means that the projected second-shift readout is inside the declared near-kernel radial halo, while $\rho_r^\gamma \ge r_{\mathrm{HC}}^{(1)}$ records the HC periapsis/collar exclusion of literal through-collapse.

Definition A.103.2 (HC-residual halo channel). *Fix a harmless normalization floor $\delta_0 > 0$ and a tilt weight $\mu_\Theta \ge 0$. The HC-residual halo channel is*

$$\rho_h^\gamma(x) := \frac{|q_\gamma^{(2)}(x)|}{(h_\gamma^{(2)}(x))^2 + (r_\gamma^{(2)}(x))^2 + \delta_0} + \mu_\Theta \,\|\Theta_{\gamma,K}^{(2)}(x)\|_G.$$

The HC-residual halo face is

$$\rho_h^\gamma(x) = \varepsilon_h.$$

The first term measures cone defect. The second term measures the second-shift projection tilt. Therefore $\rho_h^\gamma \le \varepsilon_h$ means that the readout is simultaneously close to the ideal photon-frame HC cone and free of uncontrolled projection tilt up to the declared tolerance.

Definition A.103.3 (Executed double-epsilon halo). *The executed photon-frame halo is*

$$\mathsf{Halo}^{\gamma,K}_{\varepsilon_r,\varepsilon_h} := \{x \in U_{\gamma,K} \cap \mathcal{X}^{\mathrm{adm}}_{\mathrm{OP5}} : r_{\mathrm{HC}}^{(1)}(x) \le \rho_r^\gamma(x) \le \varepsilon_r,\ \rho_h^\gamma(x) \le \varepsilon_h\}.$$

The extra lower bound $r_{\mathrm{HC}}^{(1)} \le \rho_r^\gamma$ is not an additional escape channel. It is the previously declared HC-collar guard: the admissible branch approaches a periapsis collar rather than crossing K literally.

A.104 Exit map and exit Jacobian

Definition A.104.1 (Two-channel exit map). *The exit map is*

$$\mathcal{E}^{\gamma,K}_{\mathsf{Exit}} : U_{\gamma,K} \longrightarrow \mathbb{R}^2, \qquad \mathcal{E}^{\gamma,K}_{\mathsf{Exit}}(x) := \big(\rho_r^\gamma(x) - \varepsilon_r,\ \rho_h^\gamma(x) - \varepsilon_h\big).$$

The halo boundary is $(\mathcal{E}^{\gamma,K}_{\mathsf{Exit}})^{-1}(0\times(-\infty,0]) \cup (\mathcal{E}^{\gamma,K}_{\mathsf{Exit}})^{-1}((-\infty,0]\times 0)$ together with the collar face $\rho_r^\gamma = r_{\mathrm{HC}}^{(1)}$. The collar face is not treated as an outward OP5 exit; it is the no-through-collapse reflecting side of the HC collar.

Definition A.104.2 (Exit Jacobian). *At a boundary point p, the exit Jacobian is*

$$J^{\gamma,K}_{\mathsf{Exit}}(p) := D\mathcal{E}^{\gamma,K}_{\mathsf{Exit}}(p) = \begin{pmatrix} D\rho_r^\gamma(p) \\ D\rho_h^\gamma(p) \end{pmatrix}.$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

The maximal two-channel rank is $\mathrm{rank}_{\max} = 2$. *The rank-drop tunnel condition is therefore*

$$\operatorname{rank} J^{\gamma,K}_{\mathsf{Exit}}(p) < 2,$$

which is equivalently

$$D\rho^{\gamma}_{r}(p) \wedge D\rho^{\gamma}_{h}(p) = 0$$

whenever the two channel differentials are defined.

Definition A.104.3 (Outward exit condition)**.** *A boundary point* $p \in \partial\mathsf{Halo}^{\gamma,K}_{\varepsilon_r,\varepsilon_h}$ *is outward in the radial channel if there exists an admissible update vector* v *with*

$$D\rho^{\gamma}_{r}(p)[v] > 0 \quad \textit{on} \quad \rho^{\gamma}_{r}(p) = \varepsilon_r.$$

It is outward in the HC-residual channel if

$$D\rho^{\gamma}_{h}(p)[v] > 0 \quad \textit{on} \quad \rho^{\gamma}_{h}(p) = \varepsilon_h.$$

A genuine tunnel candidate must be outward in at least one of these two channels and must pass the selector/rank tests.

A.105 First tunnel-menu execution

Proposition A.105.1 (Boundary exhaustion by the tunnel menu)**.** *Within the photon-frame halo grammar, every outward boundary exit candidate belongs to at least one of the six tunnel classes*

$$\mathfrak{T}^{\mathrm{loc}}_{\mathrm{OP5}} = \{T_r, T_h, T_{rh}, T_\chi, T_{\mathrm{rk}}, T_{\mathrm{ov}}\}.$$

Proof. Let $p \in \partial\mathsf{Halo}^{\gamma,K}_{\varepsilon_r,\varepsilon_h}$ be an outward exit candidate. If p lies on the radial face with $\rho^{\gamma}_{r}(p) = \varepsilon_r$ and $\rho^{\gamma}_{h}(p) < \varepsilon_h$, then $p \in T_r$. If p lies on the HC-residual face with $\rho^{\gamma}_{r}(p) < \varepsilon_r$ and $\rho^{\gamma}_{h}(p) = \varepsilon_h$, then $p \in T_h$. If both equalities hold, then $p \in T_{rh}$. If the tunnel selector changes value or must be evaluated at a selector discontinuity, then $p \in T_\chi$. If $\operatorname{rank} J^{\gamma,K}_{\mathsf{Exit}}(p) < 2$, then $p \in T_{\mathrm{rk}}$. Finally, if the exit occurs while changing from one photon-frame chart to another, then the point lies in an atlas-overlap region and belongs to T_{ov}. These cases exhaust the declared boundary, selector, rank, and overlap mechanisms of the grammar. □

Definition A.105.2 (response alphabet)**.** *For each tunnel class* T*, the first execution response is drawn from*

$$\mathrm{Resp}_{\mathrm{loc}}(T) \in \{\text{blocked}, \text{absorbed}_{\mathrm{HC/MM}}, \text{atlas routed}, \text{rank isolated}, \text{residual open}\}.$$

Only the first three responses are closing responses. Rank-isolated is a reduction response: it converts the question into a finite critical-set test. Residual-open is a non-closing response.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Tunnel class	test	conservative response
T_r	$\rho_r^\gamma = \varepsilon_r,\ \rho_h^\gamma < \varepsilon_h$, radial outward derivative	atlas routed if covered by a strict contraction chart; otherwise residual open
T_h	$\rho_h^\gamma = \varepsilon_h,\ \rho_r^\gamma < \varepsilon_r$, HC-residual outward derivative	absorbed if HC/MM absorption is explicit and debt-free; otherwise residual open
T_{rh}	$\rho_r^\gamma = \varepsilon_r,\ \rho_h^\gamma = \varepsilon_h$	split into T_r and T_h if rank two; rank isolated if rank drops
T_χ	tunnel selector changes or is discontinuous	blocked when selector returns zero; otherwise routed to atlas-overlap or residual open
T_{rk}	$D\rho_r^\gamma \wedge D\rho_h^\gamma = 0$	rank isolated as a critical set of the two-channel exit map
T_{ov}	chart transition or finite-atlas overlap debt	atlas routed if transition maps preserve strict contraction and zero debt

A.106 Rank-drop isolation and finite critical set target

Definition A.106.1 (Rank-drop critical register)**.** *The rank-drop register is*

$$\mathcal{C}_{\mathrm{rk}}^{\mathrm{loc}} := \{p \in \partial\mathsf{Halo}_{\varepsilon_r,\varepsilon_h}^{\gamma,K} : D\rho_r^\gamma(p) \wedge D\rho_h^\gamma(p) = 0,\ \chi_{\mathrm{tun}}(p) = 1\}.$$

It is the only place where the two halo faces can fail to give independent boundary directions.

Proposition A.106.2 (Rank-drop reduction)**.** *Assume ρ_r^γ and ρ_h^γ are piecewise analytic on a finite photon-frame atlas. Then the rank-drop tunnel class is reduced to the piecewise analytic critical-set problem*

$$D\rho_r^\gamma \wedge D\rho_h^\gamma = 0 \quad \text{on} \quad \partial\mathsf{Halo}_{\varepsilon_r,\varepsilon_h}^{\gamma,K}.$$

If this critical set is empty, blocked by $\chi_{\mathrm{tun}} = 0$, or covered by finitely many strict contraction charts with zero hidden debt, then T_{rk} contributes no uncontrolled OP5 exit.

Proof. By definition, rank-drop occurs exactly when the differential of the exit map has rank less than two. For a two-channel map this is exactly the vanishing of the wedge $D\rho_r^\gamma \wedge D\rho_h^\gamma$, away from nondifferentiable selector points which are assigned to T_χ. Piecewise analyticity transfers the problem to a finite critical-set calculation on each chart. The stated empty, blocked, or chart-routed alternatives are precisely the controlled outcomes in the response alphabet. □

A.107 First executed closure implication

Theorem A.107.1 (executed halo/tunnel reduction)**.** *Assume the OP5=HC-cone working identification and the photon-frame channel definitions above. Suppose the following four conditions hold on the declared finite photon-frame atlas:*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

1. ***Channel regularity:*** *ρ_r^γ and ρ_h^γ are piecewise analytic and compatible with the HC-collar guard $r_{\mathrm{HC}}^{(1)} \le \rho_r^\gamma$.*

2. ***Selector completeness:*** *every admissible outward boundary exit is evaluated by χ_{tun}.*

3. ***Tunnel response closure:*** *each nonempty class in $\{T_r, T_h, T_{rh}, T_\chi, T_{\mathrm{rk}}, T_{\mathrm{ov}}\}$ receives one of the three closing responses: blocked, debt-free HC/MM absorbed, or atlas routed through strict contraction charts.*

4. ***No hidden debt:***
$$L_{\mathrm{unr}} = 0, \qquad B_{\mathrm{HC/MM}} = 0, \qquad \tau = 0.$$

Then the double-epsilon photon-frame halo has no uncontrolled OP5 exit. Consequently, if halo capture and halo invariance off tunnels are also proved globally for the admissible OP5 readout class, OP5 reduces to a closed HC-cone halo theorem.

Proof. Boundary exhaustion assigns every outward exit candidate to the finite tunnel menu. The response-closure assumption assigns every nonempty class to a controlled outcome. Blocked classes do not exit; debt-free HC/MM-absorbed classes have no unpaid residual; atlas-routed classes return to strict contraction charts; and the no-hidden-debt condition rules out suppressed leakage. Hence no uncontrolled exit remains inside the executed halo grammar. Combining this with capture and invariance would trap every admissible OP5 branch or route every tunnel exit through controlled charts. This is exactly the HC-cone halo theorem formulation of OP5. □

A.108 What still has to be shown for an actual OP5 close

The layer executes the definitions and the first tunnel-menu reduction. It does not yet prove the global OP5 close. The remaining non-negotiable proof obligations are:

1. prove finite-atlas coverage of the admissible photon-frame near-kernel collar;
2. prove halo capture for every admissible OP5 readout;
3. prove halo invariance away from the finite tunnel menu;
4. compute or bound $\mathcal{C}_{\mathrm{rk}}^{\mathrm{loc}}$;
5. prove that selector-switch points are blocked or chart-routed;
6. prove that all atlas overlaps preserve the strict margin $\sup_j(\widehat{\lambda}_j + \eta_j) < 1$;
7. prove zero hidden leakage and zero hidden HC/MM debt.

If these obligations are discharged, OP5 is no longer an all-routes problem. It is exhausted by the double-epsilon halo, the finite tunnel menu, and strict contraction routing at the tunnel exits.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.109 Conservativity guards

1. does not claim global OP5 closure.
2. The executed halo is a readout-side HC-collar filter, not a literal particle shell.
3. The collar face $\rho_r^{\gamma} = r_{\mathrm{HC}}^{(1)}$ is a no-through-collapse guard and not an escape face.
4. The second-shift projected through-K appearance remains tilt debt unless the tunnel tests certify otherwise.
5. HC/MM absorption must remain explicit and debt-free.
6. OP1, OP4, QCD/confinement, CKM/PMNS, and the quark-mass spectrum are not changed by this layer.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Integrated Delta Layer: Normal-Collar Rank Test

Abstract

This delta layer executes the next obstruction left by the local pass. The exit map is decomposed chartwise on a finite photon-frame halo atlas. The rank-drop tunnel class is bounded by an explicit determinant register, and the selector-switch tunnel class is bounded by a finite switch skeleton. The result is not a global OP5 close. It reduces the uncontrolled tunnel search to two finite residual registers plus atlas-overlap margin checks.

A.110 Purpose and notation hygiene, declared local OP1/OP4 closure, release-final

The local proof chain halo channel reduced the OP5 near-kernel route problem to a two-channel boundary problem. The printed operators in this layer are

$$\mathsf{Halo}, \qquad \mathsf{Exit}, \qquad J^{\gamma,K}_{\mathsf{Exit}}, \qquad \mathcal{E}^{\gamma,K}_{\mathsf{Exit}}.$$

No lowercase display macro is used. The task is to execute the instruction

$$\operatorname{rank} J^{\gamma,K}_{\mathsf{Exit}} < 2 \quad \Longleftrightarrow \quad D\rho^{\gamma}_{r} \wedge D\rho^{\gamma}_{h} = 0,$$

and to combine it with the selector-switch test.

A.111 Finite photon-frame halo atlas

Definition A.111.1 (finite halo atlas)**.** *An photon-frame halo atlas is a finite family*

$$\mathfrak{A}^{\mathrm{loc}}_{\gamma,K} = \{(U_j, \varphi_j)\}_{j=1}^{M}$$

covering the active halo boundary

$$\partial\mathsf{Halo}^{\gamma,K}_{\varepsilon_r,\varepsilon_h} \cap \{r^{(1)}_{\mathrm{HC}} \le \rho^{\gamma}_{r} \le \varepsilon_r,\ \rho^{\gamma}_{h} \le \varepsilon_h\}.$$

On each chart we write

$$r_j := \rho^{\gamma}_{r} \circ \varphi_j^{-1}, \qquad h_j := \rho^{\gamma}_{h} \circ \varphi_j^{-1}.$$

The chart is regular for when these two functions are piecewise analytic and all nonsmooth selector boundaries are explicitly assigned to the switch register.

Definition A.111.2 (Exit determinant)**.** *On a smooth two-channel chart piece with active coordinates (u, v), define*

$$W^{\mathrm{loc}}_{j}(u,v) = \det \begin{pmatrix} \partial_u r_j & \partial_v r_j \\ \partial_u h_j & \partial_v h_j \end{pmatrix} = \partial_u r_j\, \partial_v h_j - \partial_v r_j\, \partial_u h_j.$$

The rank-drop register is

$$\mathcal{C}^{\mathrm{loc}}_{\mathrm{rk}} = \bigcup_j \{p \in U_j \cap \partial\mathsf{Halo}^{\gamma,K}_{\varepsilon_r,\varepsilon_h} : W^{\mathrm{loc}}_{j}(p) = 0,\ \chi_{\mathrm{tun}}(p) = 1\}.$$

Thus the rank-drop tunnel class is not an arbitrary route family; it is a finite union of explicitly declared chartwise critical loci.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Lemma A.111.3 (Rank-drop localization). *Every rank-drop tunnel point belongs either to $\mathcal{C}^{\mathrm{loc}}_{\mathrm{rk}}$ or to a selector-switch boundary.*

Proof. On a smooth chart piece, gives rank drop exactly when the two channel differentials are dependent. In active coordinates this is exactly $W^{\mathrm{loc}}_j = 0$. If the point is not smooth because the selector changes there, it is assigned to the selector-switch register. Since the atlas is finite, the localization is finite. □

A.112 Normal collar charts

Definition A.112.1 (Normal collar form). *A smooth halo chart piece is in normal collar form when it admits coordinates (a, b, z) such that*

$$r_j(a, b, z) = r^{(1)}_{\mathrm{HC},j}(z) + \varepsilon_r a, \qquad h_j(a, b, z) = \varepsilon_h b + \Delta_j(a, b, z),$$

with

$$0 \le a \le 1, \qquad 0 \le b \le 1, \qquad |\partial_b \Delta_j(a, b, z)| \le \theta_j \varepsilon_h, \qquad 0 \le \theta_j < 1.$$

The number $1 - \theta_j$ is the local HC-residual transversality margin.

Proposition A.112.2 (Normal collar charts have no rank drop). *On a normal collar chart piece,*

$$W^{\mathrm{loc}}_j = \varepsilon_r\big(\varepsilon_h + \partial_b \Delta_j\big)$$

in the active (a, b)-plane. Hence

$$|W^{\mathrm{loc}}_j| \ge \varepsilon_r \varepsilon_h (1 - \theta_j) > 0.$$

Consequently $\mathcal{C}^{\mathrm{loc}}_{\mathrm{rk}}$ is empty on that chart piece.

Proof. The active derivatives are $\partial_a r_j = \varepsilon_r$, $\partial_b r_j = 0$, and $\partial_b h_j = \varepsilon_h + \partial_b \Delta_j$. Therefore the determinant is $\varepsilon_r(\varepsilon_h + \partial_b \Delta_j)$. The assumed bound on $\partial_b \Delta_j$ keeps this determinant strictly away from zero. □

A.113 Selector-switch register

Definition A.113.1 (Finite selector skeleton). *The tunnel selector is a finite Boolean gate*

$$\chi_{\mathrm{tun}} : \partial \mathsf{Halo}^{\gamma,K}_{\varepsilon_r,\varepsilon_h} \longrightarrow \{0, 1\}.$$

It is constant on cells cut out by finitely many piecewise analytic switch functions

$$\sigma_{j,\ell} : U_j \longrightarrow \mathbb{R}.$$

The selector-switch register is

$$\mathcal{C}^{\mathrm{loc}}_{\chi} = \bigcup_{j,\ell} \{p \in U_j \cap \partial \mathsf{Halo}^{\gamma,K}_{\varepsilon_r,\varepsilon_h} : \sigma_{j,\ell}(p) = 0\}.$$

A switch point is closing only if the outward-side selector is zero or if a strict atlas-transition chart routes it with zero hidden debt.

Lemma A.113.2 (Selector localization). *Every selector ambiguity or selector discontinuity of an outward boundary candidate belongs to $\mathcal{C}^{\mathrm{loc}}_{\chi}$.*

Proof. Away from the switch skeleton all switch signs are fixed, so the selector is locally constant. Hence the selector can change only where at least one $\sigma_{j,\ell}$ vanishes. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.114 Tunnel response execution

The six tunnel classes receive the following responses:

class	response
T_r	atlas routed if strict contraction covers the radial face; otherwise open or blocked
T_h	debt-free HC/MM absorbed if explicit absorption certificate exists; otherwise open
T_{rh}	split into T_r, T_h if $W_j^{\mathrm{loc}} \neq 0$; rank isolated if $W_j^{\mathrm{loc}} = 0$
T_χ	blocked if outward selector is zero; otherwise switch isolated or atlas routed
T_{rk}	rank isolated inside $\mathcal{C}_{\mathrm{rk}}^{\mathrm{loc}}$ until empty/block/absorb/route is proved
T_{ov}	atlas routed if overlap margins remain strictly below one and debt is zero

Definition A.114.1 (Overlap margin)**.** *For a chart transition $\Gamma_{jk} : U_j \to U_k$, set*

$$\mathfrak{m}_{jk} := \sup_{x \in U_j \cap U_k} \big(\widehat{\lambda}_k(\Gamma_{jk}x) + \eta_k(\Gamma_{jk}x)\big).$$

The overlap is contraction-preserving if

$$\mathfrak{m}_{jk} < 1, \qquad L_{\mathrm{unr}} = 0, \qquad B_{\mathrm{HC/MM}} = 0$$

on the overlap.

Theorem A.114.2 (rank/switch register reduction)**.** *Assume the OP5=HC-cone working identification and the photon-frame halo-channel definitions. Suppose the finite halo atlas satisfies:*

1. *each smooth chart piece is piecewise analytic;*
2. *selector changes occur only on $\mathcal{C}_\chi^{\mathrm{loc}}$;*
3. *every normal collar chart piece has $\theta_j < 1$;*
4. *all active atlas overlaps are contraction-preserving;*
5. *hidden leakage and hidden HC/MM debt vanish on the executed halo boundary:*

$$L_{\mathrm{unr}} = 0, \qquad B_{\mathrm{HC/MM}} = 0.$$

Then every outward tunnel candidate is either blocked, debt-free HC/MM absorbed, atlas routed, or contained in

$$\mathcal{R}_{\mathrm{res}}^{\mathrm{loc}} := \mathcal{C}_{\mathrm{rk}}^{\mathrm{loc}} \cup \mathcal{C}_\chi^{\mathrm{loc}}.$$

If, in addition, $\mathcal{C}_{\mathrm{rk}}^{\mathrm{loc}}$ is empty, blocked, absorbed, or routed and $\mathcal{C}_\chi^{\mathrm{loc}}$ is blocked or routed, then the tunnel menu has no uncontrolled OP5 exit.

Proof. The boundary exhaustion assigns every outward exit to the finite tunnel menu. The determinant localization sends rank-drop points to $\mathcal{C}_{\mathrm{rk}}^{\mathrm{loc}}$, while normal collar transversality eliminates rank drop on regular pieces. The selector localization sends selector ambiguity to $\mathcal{C}_\chi^{\mathrm{loc}}$. Radial, HC-residual, corner, rank, selector, and overlap tunnel classes are then handled by the response table. The only non-closing objects left are the two residual registers. If these are also empty, blocked, absorbed, or routed, no uncontrolled tunnel remains. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.115 First executable normal-form sample

For the simplest ideal photon-frame collar sample, take active coordinates (a, b) and set

$$r(a,b) = r_{\mathrm{HC}}^{(1)} + \varepsilon_r a, \qquad h(a,b) = \varepsilon_h b + \delta ab, \qquad |\delta| < \varepsilon_h.$$

Then

$$W^{\mathrm{loc}}(a,b) = \det \begin{pmatrix} \varepsilon_r & 0 \\ \delta b & \varepsilon_h + \delta a \end{pmatrix} = \varepsilon_r(\varepsilon_h + \delta a) > 0.$$

Thus the rank-drop register is empty in the ideal normal-form sample. Any remaining tunnel issue in that sample must come from a selector switch, an atlas overlap, or an explicitly declared HC/MM absorption boundary.

A.116 Conservativity guards

This layer proves a structural reduction, not a final physical closure of OP5. The double-epsilon halo remains a readout-side structural filter, not a literal particle shell. Tunnel points are selector, rank, or obstruction points, not literal spacetime wormholes. The ideal normal-form sample is a diagnostic model, not a measurement claim. HC/MM absorption must remain explicit and debt-free. OP1, OP4, QCD/confinement, CKM/PMNS, and the quark-mass spectrum are not changed by this layer.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Integrated Delta Layer: Selector and Overlap Test

Abstract

This delta layer executes the follow-up left by the local pass. It gives a sufficient, finite, chartwise criterion which places the relevant photon-frame halo charts into normal-collar form with a uniform margin $|\partial_b \Delta_j| \le \theta_j \varepsilon_h, \theta_j < 1$. Under this certificate the rank-drop register is empty on all certified collar pieces. The layer does not close OP5 globally. It removes the rank-drop tunnel channel from the certified photon-frame normal-collar atlas and leaves only selector-switch, atlas-overlap, leakage/debt, or non-certified residual pieces.

A.117 Purpose

The layer reduced uncontrolled halo exits to a rank-drop register, a selector-switch register, and atlas-overlap margin checks. The key unresolved certificate was whether the photon-frame halo charts may actually be placed in normal-collar form. This layer gives the required local criterion and the first finite symbolic uniformization certificate.

The printed operators remain renderer-safe:

$$\mathsf{Halo}, \qquad \mathsf{Exit}, \qquad J^{\gamma,K}_{\mathsf{Exit}}, \qquad \mathcal{E}^{\gamma,K}_{\mathsf{Exit}}.$$

No raw lowercase display macro is used.

A.118 Photon-frame collar coordinates

Definition A.118.1 (Certified photon-frame collar piece)**.** *A smooth halo chart piece $U_j \subset \partial\mathsf{Halo}^{\gamma,K}_{\varepsilon_r,\varepsilon_h}$ is the local pass-certified when it admits coordinates*

$$(a, b, z) \in [0, 1] \times [0, 1] \times Z_j$$

with the following three properties:

1. *a is the normalized radial collar coordinate after the second photon-frame shift,*

$$\rho^{\gamma}_r(a, b, z) = r^{(1)}_{\mathrm{HC},j}(z) + \varepsilon_r a;$$

2. *b is the normalized HC-cone height coordinate;*

3. *the HC-residual channel splits as*

$$\rho^{\gamma}_h(a, b, z) = \varepsilon_h b + \Delta_j(a, b, z)$$

with a declared tilt-debt derivative bound

$$|\partial_b \Delta_j(a, b, z)| \le \theta_j \varepsilon_h, \qquad 0 \le \theta_j < 1.$$

The number $1 - \theta_j$ is the normal-collar safety margin.

Remark A.118.2. *The coordinate b is not defined to be $\rho^{\gamma}_h/\varepsilon_h$. It is the HC-cone height coordinate inherited from the photon-frame collar. This avoids a circular proof of transversality. The nontrivial check is exactly the derivative bound on Δ_j.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition A.118.3 (Uniform certificate)**.** *A finite photon-frame halo atlas*

$$\mathfrak{A}^{\mathrm{loc}}_{\gamma,K} = \{U_j\}_{j=1}^{M}$$

is uniformly normal-collar certified when every smooth active piece is the local pass-certified and

$$\theta^{\mathrm{loc}}_{\max} := \max_{1\le j\le M} \theta_j < 1.$$

The certified rank-drop register is then

$$\mathcal{C}^{\mathrm{loc}}_{\mathrm{rk}} = \bigcup_j \{p \in U_j \cap \partial \mathsf{Halo}^{\gamma,K}_{\varepsilon_r,\varepsilon_h} : W^{\mathrm{loc}}_j(p) = 0, \chi_{\mathrm{tun}}(p) = 1\}.$$

A.119 Rank-drop elimination on certified pieces

Lemma A.119.1 (determinant lower bound)**.** *On an the local pass-certified collar piece,*

$$W^{\mathrm{loc}}_j(a,b,z) = \varepsilon_r\big(\varepsilon_h + \partial_b \Delta_j(a,b,z)\big).$$

Consequently

$$|W^{\mathrm{loc}}_j(a,b,z)| \ge \varepsilon_r \varepsilon_h (1-\theta_j) > 0.$$

Proof. The active derivatives in the (a,b)-plane are

$$\partial_a \rho^{\gamma}_r = \varepsilon_r, \qquad \partial_b \rho^{\gamma}_r = 0, \qquad \partial_b \rho^{\gamma}_h = \varepsilon_h + \partial_b \Delta_j.$$

Therefore the exit determinant is

$$\det \begin{pmatrix} \partial_a \rho^{\gamma}_r & \partial_b \rho^{\gamma}_r \\ \partial_a \rho^{\gamma}_h & \partial_b \rho^{\gamma}_h \end{pmatrix} = \varepsilon_r(\varepsilon_h + \partial_b \Delta_j).$$

The derivative bound gives $\varepsilon_h + \partial_b \Delta_j \ge \varepsilon_h(1-\theta_j) > 0$ after orienting the collar height positively. Hence the determinant is bounded away from zero. □

Proposition A.119.2 (Certified collar pieces contain no rank-drop tunnels)**.** *If U_j is the local pass-certified, then*

$$U_j \cap \mathcal{C}^{\mathrm{loc}}_{\mathrm{rk}} = \varnothing.$$

If the atlas is uniformly the local pass-certified, then the rank-drop tunnel channel is empty on the certified smooth halo boundary:

$$\mathcal{C}^{\mathrm{loc}}_{\mathrm{rk}} \cap \bigcup_j U^{\mathrm{cert}}_j = \varnothing.$$

Proof. Rank drop is equivalent to $W^{\mathrm{loc}}_j = 0$ on a smooth chart piece. The determinant lower bound excludes this equality on each certified piece. Taking the finite union gives the atlas statement. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.120 Finite symbolic certificate

The first executable certificate uses three representative collar pieces. These are not physical measurements; they are the minimal symbolic chart types required by the current halo grammar.

piece	role	$\Delta_j(a,b,z)/\varepsilon_h$	θ_j	rank response
U_1	radial face	$\frac{1}{4}ab$	$\frac{1}{4}$	empty
U_2	HC-residual face	$\frac{3}{8}b(1-a)$	$\frac{3}{8}$	empty
U_3	corner/overlap collar	$\frac{1}{2}ab(1-a)$	$\frac{1}{2}$	empty

Thus

$$\theta_{\max}^{\mathrm{loc}} = \frac{1}{2} < 1$$

and hence

$$|W_j^{\mathrm{loc}}| \geq \frac{1}{2}\varepsilon_r\varepsilon_h$$

on the three certified representative pieces.

Lemma A.120.1 (First finite symbolic normal-collar sample)**.** *For the three-piece certificate above, the rank-drop register is empty on all certified pieces.*

Proof. For every listed piece, $\theta_j \leq 1/2$. The determinant lower bound gives strict positivity of W_j^{loc}. Hence no listed piece contributes to $\mathcal{C}_{\mathrm{rk}}^{\mathrm{loc}}$. □

A.121 Remaining tunnel channels

After the certificate, the local proof chain tunnel menu is reduced as follows:

tunnel class	status
T_r	covered by normal collar if radial face is in the certified atlas
T_h	covered by normal collar or debt-free HC/MM absorption
T_{rh}	rank split eliminated on certified pieces; overlap still checked
T_χ	unchanged: selector-switch register must be blocked or routed
T_{rk}	empty on certified collar pieces
T_{ov}	unchanged: transition margins must remain strictly below one

Thus does not remove every OP5 obstruction. It removes the rank-drop channel exactly where the photon-frame halo atlas has the certified normal-collar form.

Theorem A.121.1 (normal-collar rank elimination)**.** *Assume the OP5=HC-cone working identification, the halo-channel execution, and the rank/switch register reduction. Suppose a finite photon-frame halo atlas is uniformly the local pass-certified with $\theta_{\max}^{\mathrm{loc}} < 1$. Then no rank-drop tunnel occurs on the certified smooth halo boundary. Every remaining uncontrolled candidate must lie in one of the following residual classes:*

$$\mathcal{R}_{\mathrm{res}}^{\mathrm{loc}} \subseteq \mathcal{C}_\chi^{\mathrm{loc}} \cup \mathcal{C}_{\mathrm{ov}}^{\mathrm{loc}} \cup \mathcal{C}_{\mathrm{leak/debt}}^{\mathrm{loc}} \cup \mathcal{C}_{\mathrm{uncert}}^{\mathrm{loc}},$$

where $\mathcal{C}_{\mathrm{uncert}}^{\mathrm{loc}}$ denotes non-certified pieces not yet placed in normal-collar form. If the selector-switch register is blocked or routed, all atlas-overlap margins are contraction-preserving, hidden leakage and HC/MM debt vanish, and $\mathcal{C}_{\mathrm{uncert}}^{\mathrm{loc}} = \varnothing$, then the halo tunnel menu has no uncontrolled exit on the finite certified atlas.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. The certified determinant lower bound eliminates T_{rk} on every smooth certified chart piece. The reduction already states that all remaining outward candidates are rank, selector, overlap, leakage/debt, or residual non-certified pieces. Removing rank on certified pieces leaves exactly the displayed residual classes. If these classes are blocked, routed, or empty under the stated hypotheses, no uncontrolled exit remains on the finite certified atlas. □

A.122 Meaning for OP5

The result is a strong local structural advance. It says that, provided the photon-frame halo atlas is uniformly normal-collar certified, OP5 is no longer obstructed by a generic rank-collapse phenomenon near the HC collar. The hard work is pushed to four explicit checks:

selector switches, atlas overlaps, hidden leakage/debt, uncertified pieces.

A global OP5 close would require these four checks to vanish, block, or route over the full finite atlas.

A.123 Conservativity guards

This layer does not claim a global OP5 closure. It proves a conditional rank-drop elimination theorem for certified photon-frame normal-collar charts. The halo remains a readout-side structural filter, not a literal particle shell. Tunnel points remain selector, rank, overlap, leakage, or obstruction points. HC/MM absorption must remain explicit and debt-free. OP1, OP4, QCD/confinement, CKM/PMNS, and the quark-mass spectrum are untouched.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Integrated Delta Layer: Finite Capture

Abstract

This delta layer executes the two residual channels left by on certified photon-frame normal-collar pieces: selector switches and atlas overlaps. It introduces a finite selector-switch stratum, a finite overlap transition graph, a switch routing alphabet, and an overlap contraction-margin test. The layer proves that if selector switches are blocked, debt-free HC/MM absorbed, or routed into certified charts, and if every overlap edge preserves the strict contraction margin, then no uncontrolled OP5 exit remains on the finite certified halo atlas apart from explicitly declared leakage/debt or non-certified pieces. No global OP5 closure is claimed.

A.124 Purpose and inherited state

The layer removed rank-drop tunnels on certified photon-frame normal-collar pieces. The residual menu was reduced to selector switches, atlas overlaps, hidden leakage/debt, and non-certified pieces. This layer executes the first two items:

$$\mathcal{C}_{\chi}, \qquad \mathcal{C}_{\mathrm{ov}}.$$

The printed operators remain renderer-safe:

$$\mathsf{Halo}, \qquad \mathsf{Exit}, \qquad J^{\gamma,K}_{\mathsf{Exit}}.$$

No lowercase display macro for these operators is used.

We work over a finite certified halo atlas

$$\mathfrak{A}^{\mathrm{loc}}_{\gamma,K} = \{U_1, \dots, U_M\}$$

whose certified normal-collar pieces satisfy the determinant bound

$$|W_j| \geq \varepsilon_r \varepsilon_h (1 - \theta_j) > 0.$$

Thus the rank-drop channel is already empty on the certified pieces. The remaining question is whether a route can leave through a selector change or through a chart transition.

A.125 Selector-switch register

Definition A.125.1 (Finite selector-switch stratum)**.** *Let*

$$\chi_j : U_j \longrightarrow \{0, 1, \dots, Q\}$$

be the local OP5 route selector on a certified chart piece. The selector-switch stratum is

$$\mathfrak{S}^{\mathrm{loc}}_{\chi} = \bigcup_{j=1}^{M} \partial_{\chi} U_j,$$

where $\partial_{\chi} U_j$ is the finite union of codimension-one faces on which χ_j changes value or ceases to be locally constant. The selector-switch register is

$$\mathcal{C}^{\mathrm{loc}}_{\chi} = \{p \in \mathfrak{S}^{\mathrm{loc}}_{\chi} : p \in \partial \mathsf{Halo}^{\gamma,K}_{\varepsilon_r,\varepsilon_h},\ \chi_{\mathrm{tun}}(p) = 1\}.$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition A.125.2 (switch response alphabet)**.** *A selector-switch point $p \in \mathcal{C}_\chi^{\mathrm{loc}}$ receives one of four conservative responses:*

$$\mathrm{resp}_\chi^{\mathrm{loc}}(p) \in \{\mathrm{block}, \mathrm{abs}, \mathrm{route}, \mathrm{res}\}.$$

The meanings are:

response	*conservative meaning*
block	*the selector change is inadmissible for the OP5 tunnel gate; no exit is allowed;*
abs	*the switch is absorbed by declared debt-free HC/MM bookkeeping, with $B_{\mathrm{HC/MM}} = 0$;*
route	*the switch enters a certified chart piece with preserved halo margin;*
res	*the point remains an explicit residual and cannot be used for closure.*

Lemma A.125.3 (Switch containment)**.** *Assume every selector is locally constant on the interior of each certified collar piece. Then every selector-induced uncontrolled exit candidate lies in $\mathcal{C}_\chi^{\mathrm{loc}}$.*

Proof. On the interior of a certified piece the selector does not change. Therefore selector-driven route creation cannot occur there. If a selector change occurs, it occurs on the switch stratum by definition. Intersecting with the halo boundary and the tunnel gate gives exactly $\mathcal{C}_\chi^{\mathrm{loc}}$. □

Proposition A.125.4 (Selector-switch closure criterion)**.** *If*

$$\mathrm{resp}_\chi^{\mathrm{loc}}(p) \in \{\mathrm{block}, \mathrm{abs}, \mathrm{route}\} \quad \textit{for every } p \in \mathcal{C}_\chi^{\mathrm{loc}},$$

and if each routed switch lands in a certified chart with unchanged debt-free bookkeeping, then selector switches create no uncontrolled OP5 exit on the certified atlas.

Proof. A blocked switch has no admissible exit. An absorbed switch has no hidden HC/MM debt by hypothesis. A routed switch remains inside the certified atlas and is again subject to the rank elimination and the overlap test below. Since no switch has response res, no selector-induced uncontrolled exit remains. □

A.126 Atlas-overlap register

Definition A.126.1 (Finite overlap transition graph)**.** *The overlap graph is the finite directed graph*

$$\mathfrak{G}_{\mathrm{ov}}^{\mathrm{loc}} = (V, E), \qquad V = \{1, \dots, M\},$$

where $(j, k) \in E$ exactly when the certified chart overlap

$$U_{jk} = U_j \cap U_k$$

is nonempty and may be crossed by an admissible OP5 route. On such an edge let

$$\Phi_{jk} : U_{jk} \subset U_j \longrightarrow U_k$$

be the declared transition map.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition A.126.2 (Overlap contraction margin). *For an overlap edge $(j,k) \in E$, define the overlap margin*

$$\mathfrak{m}_{jk}^{\mathrm{loc}} = \widehat{\lambda}_j + \eta_j + \omega_{jk},$$

where $\widehat{\lambda}_j + \eta_j < 1$ is the inherited chart contraction budget and $\omega_{jk} \geq 0$ is the declared transition distortion of Φ_{jk} in the two-channel halo norm

$$\|(u,v)\|_{\mathsf{Halo}} = \max\left\{\frac{|u|}{\varepsilon_r}, \frac{|v|}{\varepsilon_h}\right\}.$$

The overlap edge is contraction-preserving when

$$\mathfrak{m}_{jk}^{\mathrm{loc}} < 1.$$

The overlap register is

$$\mathcal{C}_{\mathrm{ov}}^{\mathrm{loc}} = \{(j,k) \in E : \mathfrak{m}_{jk}^{\mathrm{loc}} \geq 1\} \cup \{(j,k) \in E : \Phi_{jk} \text{ is undeclared}\}.$$

Lemma A.126.3 (Overlap preservation). *If $(j,k) \in E$ has $\mathfrak{m}_{jk}^{\mathrm{loc}} < 1$, then crossing the overlap from U_j to U_k preserves strict halo contraction and does not create an uncontrolled OP5 exit.*

Proof. The inherited chart budget bounds the active chartwise projection debt by $\widehat{\lambda}_j + \eta_j$. The transition contributes at most ω_{jk} in the normalized two-channel halo norm. Hence the total budget after crossing the overlap is bounded by $\mathfrak{m}_{jk}^{\mathrm{loc}} < 1$. The crossing therefore remains inside the controlled halo atlas budget. □

Proposition A.126.4 (Atlas-overlap closure criterion). *If $\mathcal{C}_{\mathrm{ov}}^{\mathrm{loc}} = \varnothing$, then atlas overlaps do not contribute uncontrolled OP5 exits on the finite certified atlas.*

Proof. By definition, $\mathcal{C}_{\mathrm{ov}}^{\mathrm{loc}} = \varnothing$ means every admissible overlap edge is declared and contraction-preserving. The previous lemma applies to every edge of the finite transition graph. Therefore no overlap crossing leaves the controlled halo budget. □

A.127 First symbolic execution table

For the current three-piece symbolic atlas inherited from the local pass, use the following conservative finite routing table. The entries are structural certificates, not physical measurements.

selector face	location	response	reason
S_{12}	$U_1 \leftrightarrow U_2$	route	same halo side, certified target
S_{23}	$U_2 \leftrightarrow U_3$	route	corner collar remains certified
S_{13}	$U_1 \leftrightarrow U_3$	block	non-adjacent selector jump not OP5-admissible
$S_{\mathrm{HC/MM}}$	HC/MM bookkeeping face	abs	declared debt-free absorption

Thus no switch in the sample table has residual response.
The corresponding overlap margins are:

edge	$\widehat{\lambda}_j + \eta_j$	ω_{jk}	$\mathfrak{m}_{jk}^{\mathrm{loc}}$	status
$(1,2)$	5/8	1/16	11/16	preserved
$(2,3)$	21/32	1/32	11/16	preserved
$(1,3)$	25/32	1/32	13/16	preserved if routed; blocked as switch

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Therefore the sample overlap register is empty:

$$\mathcal{C}_{\mathrm{ov}}^{\mathrm{loc}} = \varnothing$$

for the declared transition graph.

A.128 residual theorem

Theorem A.128.1 (Selector-overlap execution theorem)**.** *Assume the double-epsilon halo channel, the photon-frame halo-channel execution, the rank/switch register reduction, and the normal-collar rank elimination. Suppose furthermore that:*

1. *every selector switch in* $\mathcal{C}_{\chi}^{\mathrm{loc}}$ *has response* block*,* abs*, or* route*, never* res*;*
2. *every routed selector switch lands in a certified normal-collar chart;*
3. *every admissible overlap edge is declared and satisfies* $\mathfrak{m}_{jk}^{\mathrm{loc}} < 1$*;*
4. *hidden leakage and HC/MM debt vanish:*

$$L_{\mathrm{unr}} = 0, \qquad B_{\mathrm{HC/MM}} = 0;$$

5. *no non-certified photon-frame halo piece remains in the active OP5 route menu.*

Then the finite certified photon-frame halo atlas has no uncontrolled OP5 exit. Equivalently, within this declared halo/tunnel grammar, every admissible near-kernel OP5 route is either blocked, debt-free absorbed, or routed back into a contraction-preserving certified HC-collar chart.

Proof. The determinant bound eliminates rank-drop exits on certified collar pieces. The selector-switch criterion eliminates selector-induced exits unless they are blocked, absorbed, or routed into certified pieces. The overlap criterion eliminates chart-transition exits because every admissible edge preserves strict contraction margin. The leakage and HC/MM-debt hypotheses rule out hidden bookkeeping exits. Finally, the absence of non-certified active pieces prevents escape through parts of the halo boundary not covered by the certified grammar. Hence the finite certified atlas contains no uncontrolled OP5 exit. □

Remark A.128.2 (Conservativity guard)**.** *This theorem is not a global OP5 closure. It is a finite-atlas closure criterion under the OP5=HC-cone working identification and under the declared double-epsilon halo grammar. A global closure would additionally require proving that every admissible OP5 near-kernel route enters this certified finite atlas and that the non-certified residual class is empty in the full active-core setting.*

A.129 Resulting next step

The execution leaves a sharply stated final gap. To promote the local halo/tunnel analysis toward a real OP5 closure statement, the next layer must prove the capture/completeness clause:

$$\mathcal{R}_{\mathrm{OP5}}^{\mathrm{adm,near}K} \subseteq \bigcup_{j=1}^{M} U_j^{\mathrm{cert}}$$

for the relevant photon-frame HC-collar readouts, together with $L_{\mathrm{unr}} = 0$ and $B_{\mathrm{HC/MM}} = 0$. This is the natural target.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Integrated Delta Layer: Dixon-Carrier Coding

Abstract

This delta layer executes the capture/completeness step left open by the local pass. It defines a finite route-capture certificate for admissible near-kernel OP5 readouts under the working identification OP5 = photon-frame HC-cone route. The layer separates three tasks: capture into the double-epsilon halo, assignment to a certified normal-collar chart, and debt-free compatibility with selector/overlap routing. The result is a conditional but explicit completeness theorem: if the finite capture certificate is exhaustive, if every captured branch lands in the local proof chain certified atlas, and if hidden leakage and HC/MM debt vanish, then the declared near-kernel OP5 route menu has no uncontrolled exit. This is still not a global OP5 proof outside the declared working grammar.

A.130 Purpose and inherited state

The local proof-chain layers transformed the OP5 near-kernel route problem into a finite halo/tunnel problem. The current state is:

(i) introduced the double-epsilon halo channel;

(ii) instantiated the photon-frame channels ρ_r^γ and ρ_h^γ and the two-channel exit map;

(iii) reduced rank-drop and selector-switch search to explicit registers;

(iv) eliminated rank-drop on certified normal-collar pieces;

(v) routed selector switches and showed contraction-preserving atlas overlaps in the sample finite atlas.

The remaining gap is a capture/completeness statement:

$$\mathcal{R}_{\mathrm{OP5}}^{\mathrm{adm,near}K} \subseteq \bigcup_{j=1}^{M} U_j^{\mathrm{cert}}, \qquad L_{\mathrm{unr}} = 0, \qquad B_{\mathrm{HC/MM}} = 0.$$

This layer gives the certificate grammar for exactly that statement.

All displayed operators use renderer-safe notation:

$$\mathsf{Halo}, \qquad \mathsf{Exit}, \qquad \mathsf{Cap}.$$

A.131 Finite route-capture certificate

Definition A.131.1 (Near-kernel OP5 route menu)**.** *Let $\mathcal{R}_{\mathrm{OP5}}^{\mathrm{adm,near}K}$ be the declared set of admissible near-kernel OP5 readouts after photon-frame HC-collar normalization. A route $r \in \mathcal{R}_{\mathrm{OP5}}^{\mathrm{adm,near}K}$ is represented by a finite branch word*

$$\beta(r) = (\sigma_0, \sigma_1, \ldots, \sigma_\ell)$$

whose symbols record, in order, HC-collar entry, possible selector face, possible overlap transition, and terminal certified chart candidate. The word is admissible *only if every symbol is admitted by the inherited switch/overlap tables.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition A.131.2 (capture witness). *An capture witness for a route $r \in \mathcal{R}_{\mathrm{OP5}}^{\mathrm{adm,near}K}$ is a tuple*

$$\mathfrak{c}_r^{\mathrm{loc}} = (n_r, j_r, \beta_r, \delta_r^{\mathsf{Halo}}, \ell_r, b_r),$$

where:

(a) $n_r \in \mathbb{N}$ *is the finite number of HC/Hilbert normalizing steps;*

(b) $j_r \in \{1, \dots, M\}$ *is the certified target chart;*

(c) β_r *is the finite the local pass-admissible selector/overlap word;*

(d) δ_r^{Halo} *is the strict halo margin;*

(e) ℓ_r *is the hidden leakage register;*

(f) b_r *is the hidden HC/MM debt register.*

It is valid when

$$\rho_r^{\gamma}(R_{\mathrm{HC}}^{n_r} r) \le \varepsilon_r - \delta_r^{\mathsf{Halo}}, \qquad \rho_h^{\gamma}(R_{\mathrm{HC}}^{n_r} r) \le \varepsilon_h - \delta_r^{\mathsf{Halo}},$$
$$R_{\mathrm{HC}}^{n_r} r \in U_{j_r}^{\mathrm{cert}}, \qquad \delta_r^{\mathsf{Halo}} > 0, \qquad \ell_r = 0, \qquad b_r = 0.$$

Definition A.131.3 (Finite capture certificate). *A finite capture certificate is a finite family*

$$\mathfrak{C}_{\mathrm{cap}}^{\mathrm{loc}} = \{\mathfrak{c}_a^{\mathrm{loc}} : a \in A_{\mathrm{loc}}\}$$

together with a surjective classification map

$$\pi_{\mathrm{cap}}^{\mathrm{loc}} : \mathcal{R}_{\mathrm{OP5}}^{\mathrm{adm,near}K} \longrightarrow A_{\mathrm{loc}}.$$

For a route r, the witness assigned to $a = \pi_{\mathrm{cap}}^{\mathrm{loc}}(r)$ must be valid for every route in the class $(\pi_{\mathrm{cap}}^{\mathrm{loc}})^{-1}(a)$.

Remark A.131.4 (Why this is the right bottleneck). *Without the capture certificate, OP5 remains a many-route problem. With the certificate, every admissible near-kernel route is compressed into one of finitely many certified classes. The remaining proof is then finite: check capture margin, chart certification, selector/overlap admissibility, leakage, and HC/MM debt for each class.*

A.132 Capture tests

Definition A.132.1 (capture test stack). *For each capture class $a \in A_{\mathrm{loc}}$, define four Boolean tests:*

$$\mathsf{Cap}_{\mathsf{Halo}}(a), \qquad \mathsf{Cap}_{\mathfrak{A}}(a), \qquad \mathsf{Cap}_{\mathsf{Route}}(a), \qquad \mathsf{Cap}_{\mathsf{Debt}}(a).$$

They mean respectively:

$$\mathsf{Cap}_{\mathsf{Halo}}(a) = 1 \iff \rho_r^{\gamma} \le \varepsilon_r - \delta_a^{\mathsf{Halo}} \text{ and } \rho_h^{\gamma} \le \varepsilon_h - \delta_a^{\mathsf{Halo}} \text{ after } n_a \text{ steps},$$
$$\mathsf{Cap}_{\mathfrak{A}}(a) = 1 \iff R_{\mathrm{HC}}^{n_a}(r) \in U_{j_a}^{\mathrm{cert}} \text{ for every } r \text{ in class } a,$$
$$\mathsf{Cap}_{\mathsf{Route}}(a) = 1 \iff \beta_a \text{ uses only the local pass-admissible selector/overlap moves},$$
$$\mathsf{Cap}_{\mathsf{Debt}}(a) = 1 \iff \ell_a = 0 \text{ and } b_a = 0.$$

The class is capture-complete when all four tests equal one.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Lemma A.132.2 (Classwise capture implies chartwise capture)**.** *If* $\mathsf{Cap}_{\mathsf{Halo}}(a) = \mathsf{Cap}_{\mathfrak{A}}(a) = 1$*, then every route in class* a *enters the certified double-epsilon halo atlas after at most* n_a *normalizing steps.*

Proof. The first test gives strict entry into the double-epsilon halo, with positive margin δ_a^{Halo}. The second test identifies the target as an certified normal-collar chart. Therefore all routes in the class enter the certified halo atlas. □

Lemma A.132.3 (Route and debt tests eliminate hidden exits)**.** *If* $\mathsf{Cap}_{\mathsf{Route}}(a) = \mathsf{Cap}_{\mathsf{Debt}}(a) = 1$*, then class* a *contains no unregistered selector/overlap exit and no hidden leakage or HC/MM debt.*

Proof. The route test restricts the branch word to selector and overlap transitions already routed by the local pass. The debt test sets both hidden registers to zero. Thus the class cannot leave through an undeclared selector/overlap move or through hidden bookkeeping. □

A.133 First symbolic capture table

The following table records the first finite symbolic certificate compatible with the local proof chain. It is a structural certificate table rather than a measurement table.

class	branch type	n_a	target	margin	debt	status
A_1	collar interior	2	U_1^{cert}	1/8	0	captured
A_2	adjacent switch S_{12}	2	U_2^{cert}	3/32	0	routed/captured
A_3	adjacent switch S_{23}	2	U_3^{cert}	1/16	0	routed/captured
A_4	non-adjacent switch S_{13}	–	–	–	–	blocked
A_5	HC/MM face	2	U_2^{cert}	1/16	0	absorbed/captured
A_6	overlap edge	2	$U_j^{\mathrm{cert}} \cup U_k^{\mathrm{cert}}$	1/16	0	overlap-preserved

The active captured classes are

$$A_{\mathrm{act}}^{\mathrm{loc}} = \{A_1, A_2, A_3, A_5, A_6\}.$$

The blocked non-adjacent selector class A_4 does not contribute an admissible OP5 route. Thus the symbolic residual class is

$$A_{\mathrm{res}}^{\mathrm{loc}} = \varnothing$$

provided the classification map $\pi_{\mathrm{cap}}^{\mathrm{loc}}$ is exhaustive for the declared near-kernel route menu.

Proposition A.133.1 (Sample certificate status)**.** *For the symbolic capture table, every active class has strict positive halo margin, zero leakage, zero HC/MM debt, and a certified the local proof chain target chart. The only non-captured class is blocked by the inherited selector rule. Hence the sample table has no residual active capture class.*

Proof. The entries list positive margins for A_1, A_2, A_3, A_5, A_6, debt value zero, and certified targets. Class A_4 is not an active OP5 route because blocks the non-adjacent selector jump. Therefore no active class remains residual. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.134 Completeness theorem

Theorem A.134.1 (capture/completeness theorem)**.** *Assume the double-epsilon halo channel, the photon-frame channel execution, the register reduction, the normal-collar rank elimination, and the selector/overlap execution. Suppose there exists a finite capture certificate* $\mathfrak{C}_{\mathrm{cap}}^{\mathrm{loc}}$ *and an exhaustive classification map*

$$\pi_{\mathrm{cap}}^{\mathrm{loc}} : \mathcal{R}_{\mathrm{OP5}}^{\mathrm{adm,near}K} \longrightarrow A_{\mathrm{loc}}$$

such that every active class is capture-complete and every inactive class is blocked by an inherited OP5-admissibility rule. Then

$$\mathcal{R}_{\mathrm{OP5}}^{\mathrm{adm,near}K} \subseteq \bigcup_{j=1}^{M} U_j^{\mathrm{cert}}$$

after finitely many HC/Hilbert normalizing steps, with

$$L_{\mathrm{unr}} = 0, \qquad B_{\mathrm{HC/MM}} = 0.$$

Consequently, within the declared photon-frame HC-halo grammar, the near-kernel OP5 route menu has no uncontrolled exit.

Proof. Let $r \in \mathcal{R}_{\mathrm{OP5}}^{\mathrm{adm,near}K}$. By exhaustiveness, r belongs to some class $a = \pi_{\mathrm{cap}}^{\mathrm{loc}}(r)$. If the class is inactive, it is blocked and hence not an active OP5 exit. If the class is active, capture-completeness gives four facts: strict double-epsilon halo entry, certified target chart membership, admissible selector/overlap routing, and zero leakage/debt. The theorem eliminates rank-drop on the certified chart. The theorem eliminates selector and overlap exits. The zero-register clause eliminates hidden leakage and HC/MM debt. Hence r is blocked, absorbed, or routed into a certified contraction-preserving chart. Since r was arbitrary, the inclusion and the no-uncontrolled-exit claim follow. □

Corollary A.134.2 (Local OP5 closure inside the declared grammar)**.** *Under the hypotheses of the theorem and under the working identification OP5 = photon-frame HC-cone route, the near-kernel OP5 branch of the declared finite halo/tunnel grammar is locally closed.*

Proof. The theorem states that every admissible near-kernel route is captured by the certified atlas or blocked, with no hidden leakage/debt. This is precisely local closure in the declared grammar. □

Remark A.134.3 (No overclaim)**.** *This is not a global OP5 proof in the ordinary external sense. It is the strongest local closure statement presently justified by the local proof chain: if OP5 is correctly read as the photon-frame HC-cone route problem, and if the finite capture certificate is exhaustive for the relevant active-core route menu, then the near-kernel OP5 branch has no uncontrolled exit. A full release-level OP5 theorem would still need this certificate integrated into the monolithic Active Core and cross-checked against every OP5 occurrence in the historical witness.*

A.135 Audit target for the next layer

The layer turns the remaining work into an audit problem. The next meaningful step is not to add another route mechanism, but to verify exhaustiveness:

$$\mathrm{supp}(\mathcal{R}_{\mathrm{OP5}}^{\mathrm{adm,near}K}) = \bigcup_{a \in A_{\mathrm{loc}}} (\pi_{\mathrm{cap}}^{\mathrm{loc}})^{-1}(a),$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

with no unclassified active branch. If the monolithic Active Core confirms that equality, then can promote the result from a local capture/completeness certificate to an OP5 closure-candidate theorem for the full active-core route menu.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Integrated Delta Layer: Dixon-Image Audit

Abstract

This delta layer executes the HS(6) Dixon-coding map suggested after the capture/completeness packet. The layer separates two clocks which must not be conflated: the carrier clock $C_n = 2^n$, and the activation-mask clock $|\{0,1\}^{C_n}| = 2^{C_n}$. In this grammar, HS(5) has a 32-channel carrier with a 2^{32}-state activation mask, while HS(6) has a 64-channel carrier whose activation mask has 2^{64} states. The 64-channel carrier is mapped to the Dixon carrier $D = \mathbb{C} \otimes \mathbb{H} \otimes \mathbb{O}$. The layer then relates the HS(4) meta-septum projector $\widehat{P}_{C_7} \subset \mathcal{M}_{16}$, the HS(5) two-sided duplication, and the HS(6) Dixon carrier to the OP5 halo/capture chain. No claim is made that the Dixon algebra is 2^{64}-dimensional; only its Boolean activation code has 2^{64} possible masks.

A.136 Purpose and inherited state

The local proof chain sequence reduced the near-kernel OP5 route problem to a finite photon-frame HC-halo capture problem. Its current decisive condition is exhaustiveness of the capture map

$$\pi_{\mathrm{cap}}^{\mathrm{loc}} : \mathcal{R}_{\mathrm{OP5}}^{\mathrm{adm,near}K} \longrightarrow A_{\mathrm{loc}},$$

with $L_{\mathrm{unr}} = 0$ and $B_{\mathrm{HC/MM}} = 0$. The present layer records the next structural interpretation of higher Hilbert shifts. It does not add a new OP5 route mechanism. It provides a finite coding target for the already captured routes.

The motivating observation is that the HS tower naturally admits two readings:

$$\mathsf{Car}(n) = 2^n, \qquad \mathsf{Mask}(n) = \{0,1\}^{\mathsf{Car}(n)}, \qquad |\mathsf{Mask}(n)| = 2^{2^n}.$$

Thus the phrase "up to 2^{32}, then 2^{64}" is correct when it refers to activation masks at carrier sizes 32 and 64, not to the algebra dimension itself.

A.137 Two-clock Hilbert-shift grammar

Definition A.137.1 (Carrier clock and mask clock)**.** *For the Hilbert-shift tower define*

$$C_n := 2^n$$

as the carrier size of the n-th shift, and define the Boolean activation mask space

$$\mathsf{Mask}_n := \{0,1\}^{C_n}.$$

Then

$$|\mathsf{Mask}_n| = 2^{C_n} = 2^{2^n}.$$

We call C_n the carrier clock *and $|\mathsf{Mask}_n|$ the* mask clock.

Remark A.137.2 (Correction of the common ambiguity)**.** *The statement "Dixon is 2^{64}" is not used as a dimension statement. The structurally safe statement is*

$$\dim_{\mathbb{R}}(\mathbb{C} \otimes \mathbb{H} \otimes \mathbb{O}) = 64, \qquad |\{0,1\}^{64}| = 2^{64}.$$

Thus the Dixon carrier has 64 real channels, and its full Boolean activation grammar has 2^{64} masks.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

shift	carrier size C_n	mask count $\lvert\mathsf{Mask}_n\rvert$	structural reading
HS(0)	1	2	radial kernel target / unshifted collapse readout
HS(1)	2	4	HC collar and periapsis split
HS(2)	4	16	stabilized orbit / contraction chart
HS(3)	8	2^8	Fano/octonion-compatible local incidence fibre
HS(4)	16	2^{16}	meta-septum $\mathcal{M}_{16}/\widehat{P}_{C_7}$ leakage test
HS(5)	32	2^{32}	two-sided front/back carrier duplication
HS(6)	64	2^{64}	Dixon carrier $\mathbb{C}\otimes\mathbb{H}\otimes\mathbb{O}$ and 64-bit activation code

A.138 HS(4) meta-septum leakage target

Definition A.138.1 (HS(4) meta-septum carrier)**.** *Let $\mathcal{H}_{16}^{(4)}$ be the 16-channel HS(4) carrier containing the inherited meta-septum sector selected by*

$$\widehat{P}_{C_7} : \mathcal{H}_{16}^{(4)} \longrightarrow \mathcal{H}_{16}^{(4)}, \qquad \widehat{P}_{C_7}^2 = \widehat{P}_{C_7}.$$

Write

$$Q := \widehat{P}_{C_7}, \qquad Q_\perp := \mathbf{1} - Q.$$

For the inherited internal matrix $\mathcal{M}_{16}$, define the admissible block and mixed leakage block by

$$\mathcal{M}_{CC} := Q\mathcal{M}_{16}Q,$$

$$\mathsf{Leak}_{C_7}^{(4)}(\mathcal{M}_{16}) := Q_\perp\mathcal{M}_{16}Q + Q\mathcal{M}_{16}Q_\perp.$$

The HS(4) carrier is meta-septum closed when

$$\mathsf{Leak}_{C_7}^{(4)}(\mathcal{M}_{16}) = 0.$$

Lemma A.138.2 (Commutator form of HS(4) closure)**.** *If $Q = Q^2$ and $Q_\perp = \mathbf{1} - Q$, then vanishing of both mixed blocks is equivalent to block preservation of the C_7-sector. In particular the closure condition can be written as*

$$[\mathcal{M}_{16}, Q] = 0,$$

provided the two mixed blocks are tested as independent leakage registers.

Proof. With respect to $\mathcal{H}_{16}^{(4)} = Q\mathcal{H}_{16}^{(4)} \oplus Q_\perp\mathcal{H}_{16}^{(4)}$, the operator $\mathcal{M}_{16}$ decomposes into four blocks. The two diagonal blocks preserve the selected and complementary sectors. The two off-diagonal blocks are $Q_\perp\mathcal{M}_{16}Q$ and $Q\mathcal{M}_{16}Q_\perp$. If they vanish, $\mathcal{M}_{16}$ is block diagonal and commutes with Q. Conversely, if $[\mathcal{M}_{16}, Q] = 0$, multiplying on the left and right by Q and $Q_\perp$ gives vanishing of the off-diagonal blocks. Hence the mixed leakage register is zero exactly in the block-preserving case. □

Remark A.138.3 (Relation to HS(4) danger)**.** *The HS(4) carrier is the first level at which a naive 16-channel extension may behave like a sedenion-type expansion. This layer does not use raw sedenions as a free algebra. It uses $\mathcal{M}_{16}/\widehat{P}_{C_7}$ as a gated 16-channel envelope and records all off-sector motion as explicit leakage.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.139 HS(5) two-sided duplication

Definition A.139.1 (Two-sided HS(5) carrier)**.** *The HS(5) carrier is the front/back duplication of the HS(4) envelope:*

$$\mathcal{H}_{32}^{(5)} := (\mathcal{H}_{16}^{(4)})_{\text{front}} \oplus (\mathcal{H}_{16}^{(4)})_{\text{back}}.$$

Its projector is

$$\widehat{P}_{C_7}^{(5)} := \begin{pmatrix} (\widehat{P}_{C_7})_{\text{front}} & 0 \\ 0 & (\widehat{P}_{C_7})_{\text{back}} \end{pmatrix}.$$

A two-sided HS(5) state is written

$$\Psi^{(5)} = (\Psi_{\text{front}}, \Psi_{\text{back}}),$$

where the front side carries readout/frame/language information and the back side carries structure/action/field information.

Definition A.139.2 (HS(5) cross-side debt)**.** *Let* $\mathcal{T}^{(5)} \in \mathrm{End}(\mathcal{H}_{32}^{(5)})$*. Define*

$$\mathsf{Debt}_{\text{cross}}^{(5)}(\mathcal{T}^{(5)}) := (\mathbf{1} - \widehat{P}_{C_7}^{(5)})\mathcal{T}^{(5)}\widehat{P}_{C_7}^{(5)} + \widehat{P}_{C_7}^{(5)}\mathcal{T}^{(5)}(\mathbf{1} - \widehat{P}_{C_7}^{(5)}).$$

The HS(5) two-sided carrier is debt-free when this register vanishes and all front/back comparison maps are mediated by declared comparison tensors or selectors.

Proposition A.139.3 (HS(5) as safe bridge)**.** *If the front and back HS(4) sectors are separately meta-septum closed and the cross-side debt register vanishes, then the HS(5) carrier transports the HS(3) Fano-compatible fibre through the two-sided readout grammar without adding an uncontrolled off-sector channel.*

Proof. Separate meta-septum closure prevents leakage within each 16-channel side. Vanishing cross-side debt prevents an admissible selected component on one side from being pushed into a complementary component on the other side. Since two-sided comparisons are only admitted through declared selectors or comparison tensors, every surviving front/back transition is registered. Hence no uncontrolled off-sector channel is introduced at HS(5). □

A.140 HS(6) Dixon carrier and 64-bit activation code

Definition A.140.1 (Dixon carrier)**.** *The HS(6) Dixon carrier is the 64-dimensional real carrier*

$$\mathcal{H}_{64}^{(6)} \cong \mathcal{H}_{\text{Dix}} := \mathbb{C} \otimes \mathbb{H} \otimes \mathbb{O}.$$

A basis channel is indexed by a triple

$$\alpha = (a, b, c), \qquad a \in \{0, 1\}, \quad b \in \{0, 1, 2, 3\}, \quad c \in \{0, 1, \ldots, 7\}.$$

Thus

$$|\{\alpha\}| = 2 \cdot 4 \cdot 8 = 64.$$

Definition A.140.2 (Dixon activation code)**.** *The Dixon activation code of a captured OP5 route* $r \in \mathcal{R}_{\text{OP5}}^{\text{adm,near}K}$ *is a Boolean mask*

$$\chi_{\text{Dix}}(r) = (\chi_\alpha(r))_{\alpha \in \{0,1\} \times \{0,1,2,3\} \times \{0,\ldots,7\}} \in \{0, 1\}^{64}.$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

The full Dixon activation space therefore has

$$|\mathsf{Code}_{\mathrm{Dix}}| = 2^{64}$$

possible masks. A mask is admissible only if it respects the inherited capture class, the HS(4) meta-septum gate, and the HS(5) two-sided debt-free condition.

Definition A.140.3 (HS(6) coding map)**.** *Define the conditional route-to-Dixon coding map*

$$\Xi^{\mathrm{loc}}_{\mathrm{Dix}} : \mathcal{R}^{\mathrm{adm,near}K}_{\mathrm{OP5}} \longrightarrow \{0,1\}^{64} \cup \{\mathsf{blocked}, \mathsf{debt}, \mathsf{leak}\}$$

as follows. For $r \in \mathcal{R}^{\mathrm{adm,near}K}_{\mathrm{OP5}}$:

(a) if r is blocked by the selector table, set $\Xi^{\mathrm{loc}}_{\mathrm{Dix}}(r) = \mathsf{blocked}$;

(b) if $\mathsf{Leak}^{(4)}_{C_7} \neq 0$, set $\Xi^{\mathrm{loc}}_{\mathrm{Dix}}(r) = \mathsf{leak}$;

(c) if $\mathsf{Debt}^{(5)}_{\mathrm{cross}} \neq 0$, set $\Xi^{\mathrm{loc}}_{\mathrm{Dix}}(r) = \mathsf{debt}$;

(d) otherwise assign the 64-bit mask $\chi_{\mathrm{Dix}}(r)$ determined by its captured chart, selector word, front/back side, and Fano/octonion channel.

A.141 Compatibility with the capture theorem

Lemma A.141.1 (Captured routes admit finite Dixon codes)**.** *Assume the capture/completeness theorem and the HS(4)–HS(5) zero-register conditions*

$$\mathsf{Leak}^{(4)}_{C_7} = 0, \qquad \mathsf{Debt}^{(5)}_{\mathrm{cross}} = 0.$$

Then every active captured route $r \in \mathcal{R}^{\mathrm{adm,near}K}_{\mathrm{OP5}}$ has a well-defined Dixon activation code $\chi_{\mathrm{Dix}}(r) \in \{0,1\}^{64}$.

Proof. By the local pass, each active route is captured in a certified normal-collar chart after finitely many HC/Hilbert steps, with $L_{\mathrm{unr}} = 0$ and $B_{\mathrm{HC/MM}} = 0$. The HS(4) condition prevents the Fano/meta-septum component from leaking into the complementary 16-channel sector. The HS(5) condition prevents two-sided front/back transport from creating unregistered debt. Therefore the remaining channel data are finite and live inside the HS(6) Dixon carrier. The 64 indexed basis channels define the Boolean mask. □

Proposition A.141.2 (Tunnel-to-Dixon reduction)**.** *Under the hypotheses of the previous lemma, every tunnel/capture class is assigned one of four statuses:*

$$\mathsf{blocked}, \qquad \mathsf{leak}, \qquad \mathsf{debt}, \qquad \chi_{\mathrm{Dix}} \in \{0,1\}^{64}.$$

Moreover, an uncontrolled HS(6) tunnel can occur only through a nonzero HS(4) leakage register, a nonzero HS(5) cross-side debt register, or a nonadmissible Dixon mask.

Proof. The coding map definition exhausts the cases. Blocked routes are not active OP5 exits. Nonzero HS(4) leakage and HS(5) debt are explicitly recorded, not hidden. If both vanish, the route is represented by a finite 64-bit Dixon mask. Hence any uncontrolled tunnel must be visible as leak, debt, or failure of Dixon-mask admissibility. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Theorem A.141.3 (HS(6) Dixon coding theorem)**.** *Assume the local proof chain photon-frame HC-halo capture chain. Suppose further that*

$$\mathsf{Leak}^{(4)}_{C_7} = 0, \qquad \mathsf{Debt}^{(5)}_{\mathrm{cross}} = 0, \qquad L_{\mathrm{unr}} = 0, \qquad B_{\mathrm{HC/MM}} = 0.$$

Then every declared admissible near-kernel OP5 route in the capture menu is either blocked by an inherited selector rule or encoded by an admissible 64-bit Dixon activation mask:

$$\Xi^{\mathrm{loc}}_{\mathrm{Dix}}(r) \in \{\mathsf{blocked}\} \cup \{0,1\}^{64}.$$

Consequently, within the declared HS(0)–HS(6) grammar, the remaining OP5 tunnel audit becomes a finite Dixon-mask admissibility audit rather than an unbounded route search.

Proof. The theorem captures all active near-kernel routes into the finite certified halo atlas and eliminates hidden leakage and HC/MM debt. The HS(4) condition eliminates meta-septum off-sector leakage. The HS(5) condition eliminates front/back cross-side debt. Thus every route left after selector blocking has finite chart, selector, side, and Fano/octonion channel data. These data select a subset of the 64 Dixon carrier channels, i.e. a Boolean mask in $\{0,1\}^{64}$. Therefore the audit is finite at the level of mask admissibility. □

Corollary A.141.4 (What would be needed next)**.** *A release-level OP5 closure candidate can now be pursued by proving that every Dixon mask actually produced by the capture map is admissible, or else blocked/absorbed by explicit HC/MM bookkeeping:*

$$\mathrm{im}(\Xi^{\mathrm{loc}}_{\mathrm{Dix}}) \subseteq \mathsf{Code}^{\mathrm{adm}}_{\mathrm{Dix}} \cup \{\mathsf{blocked}\},$$

with no leak and no debt registers.

A.142 First symbolic Dixon status table

The following table records the first symbolic status assignment. It is an integration table for the route grammar, not a numerical particle-spectrum table.

class	HS(4) leakage	HS(5) debt	Dixon carrier	mask status	status
A_1 collar interior	0	0	64	admissible pending full mask audit	Dixon-coded
A_2 S_{12} routed	0	0	64	admissible pending full mask audit	Dixon-coded
A_3 S_{23} routed	0	0	64	admissible pending full mask audit	Dixon-coded
A_4 S_{13} blocked	–	–	–	–	blocked
A_5 HC/MM face	0	0	64	absorbed/coded if debt-free	coded/absorbed
A_6 overlap edge	0	0	64	overlap-preserved	Dixon-coded

Remark A.142.1 (No spectrum or Standard Model overclaim)**.** *The Dixon carrier is used here as a finite route-code carrier. This layer does not derive the Standard Model spectrum, CKM/PMNS values, quark masses, QCD confinement, or OP1/OP4. It only states how captured OP5 halo routes can be mapped into a finite 64-channel Dixon activation grammar.*

A.143 Audit target for

The next meaningful layer is not a new shift. It is the image audit of the coding map:

$$\mathrm{im}(\Xi^{\mathrm{loc}}_{\mathrm{Dix}}) \overset{?}{\subseteq} \mathsf{Code}^{\mathrm{adm}}_{\mathrm{Dix}} \cup \{\mathsf{blocked}\}.$$

Equivalently, should enumerate the Dixon-mask classes actually reachable from the capture table and classify each as blocked, HC/MM absorbed, atlas routed, Dixon-admissible, or residual-open. If that image audit is empty on residual-open masks, then the OP5 closure candidate becomes substantially stronger inside the declared HS(0)–HS(6) grammar.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Integrated Delta Layer: Route-Exhaustiveness Audit

Abstract

This delta layer executes the image audit requested after the local pass. The layer mapped captured near-kernel OP5 routes into the HS(6) Dixon carrier $\mathbb{C} \otimes \mathbb{H} \otimes \mathbb{O}$, with 64 real carrier channels and a 2^{64}-state Boolean activation-mask space. The present layer audits only the masks actually reachable from the capture table. It classifies the symbolic image of $\Xi^{\mathrm{loc}}_{\mathrm{Dix}}$ into blocked, HC/MM absorbed, atlas routed, Dixon-admissible, and residual-open classes. Under the inherited zero leakage/debt assumptions, the declared sample image has no residual-open Dixon-mask class. This remains a conditional local OP5 closure audit inside the declared photon-frame HC-halo and HS(0)–HS(6) grammar, not a global OP5 proof outside that grammar.

A.144 Why the Dixon representation helps QAM

The QAM grammar becomes more useful when it is not only expressive but also addressable. The Dixon layer supplies this addressability. Instead of leaving a captured route as an open structural phrase, it assigns the route a finite carrier address

$$\alpha = (a, b, c), \qquad a \in \{0, 1\}, \quad b \in \{0, 1, 2, 3\}, \quad c \in \{0, \ldots, 7\},$$

inside the 64-channel Dixon carrier

$$\mathcal{H}_{\mathrm{Dix}} = \mathbb{C} \otimes \mathbb{H} \otimes \mathbb{O}.$$

The QAM state is then not a vague continuum of possible route meanings. It is a Boolean activation mask

$$\chi_{\mathrm{Dix}} \in \{0, 1\}^{64}.$$

Thus QAM over Dixon gives three concrete benefits.

(i) **Finite address space.** Captured routes are assigned to a finite 64-channel carrier instead of an unbounded route grammar.

(ii) **Auditability.** Every possible uncontrolled channel must appear as a non-admissible mask, an HS(4) leakage register, or an HS(5) cross-side debt register.

(iii) **Compatibility with existing algebra.** The complex, quaternionic, and octonionic factors separate phase, two-sided/spin-like transport, and Fano/octonion incidence without claiming that the Dixon carrier itself has dimension 2^{64}.

The present layer uses these benefits to audit the image of the route-to-Dixon map.

A.145 Inherited coding map

Recall the conditional coding map

$$\Xi^{\mathrm{loc}}_{\mathrm{Dix}} : \mathcal{R}^{\mathrm{adm,near}K}_{\mathrm{OP5}} \longrightarrow \{0, 1\}^{64} \cup \{\mathsf{blocked}, \mathsf{debt}, \mathsf{leak}\}.$$

It is defined only after the local proof chain photon-frame HC-halo capture chain has reduced admissible near-kernel OP5 routes to the finite class menu

$$A_{\mathrm{loc}} = \{A_1, A_2, A_3, A_4, A_5, A_6\}.$$

The audit does not enumerate all 2^{64} masks. It enumerates the much smaller symbolic image of $\Xi^{\mathrm{loc}}_{\mathrm{Dix}}$ produced by A_{loc}.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition A.145.1 (Reachable Dixon image)**.** *The reachable Dixon image is*

$$\mathcal{I} := \Xi^{\mathrm{loc}}_{\mathrm{Dix}}(\mathcal{R}^{\mathrm{adm,near}K}_{\mathrm{OP5}}) \cap \big(\{0,1\}^{64} \cup \{\mathsf{block}, \mathsf{abs}, \mathsf{Route}, \mathsf{Leak}, \mathsf{Debt}, \mathsf{res}\}\big),$$

where each element is represented by its class, its HS(4) leakage status, its HS(5) cross-side debt status, and its Dixon-mask class.

Definition A.145.2 (Compressed Dixon mask class)**.** *For a captured route class A_i, define its compressed Dixon mask class by*

$$\mathfrak{M}_i := \big(\sigma_i, \tau_i, \omega_i, \kappa_i\big),$$

where σ_i is the selector word, τ_i is the two-sided front/back tag, $\omega_i \subseteq \{0,\dots,7\}$ is the Fano/octonion support, and κ_i is the response status. The associated full mask is the subset of Dixon channels

$$\mathrm{supp}(\chi_{\mathrm{Dix}}(A_i)) \subseteq \{0,1\} \times \{0,1,2,3\} \times \{0,\dots,7\}.$$

Remark A.145.3 (Symbolic rather than exhaustive bit enumeration)**.** *The audit enumerates mask families forced by the capture table. It does not list all 2^{64} bitstrings. This is the intended gain: OP5 does not require an exhaustive scan of the Dixon mask space if the route image is first captured into finitely many symbolic classes.*

A.146 Status alphabet and admissibility test

Definition A.146.1 (status alphabet)**.** *The image audit uses the status alphabet*

$$\mathfrak{S}_{\mathrm{loc}} = \{\mathsf{block}, \mathsf{abs}, \mathsf{Route}, \mathsf{adm}, \mathsf{res}\}.$$

The meanings are as follows.

- block*: the route is selector-blocked and produces no active OP5 exit.*
- abs*: the route is HC/MM absorbed with explicit zero debt.*
- Route*: the route is atlas-routed to another certified normal-collar piece.*
- adm*: the route is Dixon-admissible as a finite 64-channel activation mask.*
- res*: the route remains residual-open.*

Definition A.146.2 (Dixon admissibility predicate)**.** *A reachable Dixon mask class $\mathfrak{M}_i$ is the local pass-admissible when the following four tests hold:*

$$\mathsf{Leak}^{(4)}_{C_7}(A_i) = 0, \qquad \mathsf{Debt}^{(5)}_{\mathrm{cross}}(A_i) = 0,$$

$$L_{\mathrm{unr}}(A_i) = 0, \qquad B_{\mathrm{HC/MM}}(A_i) = 0,$$

and the selector/overlap word of A_i is the local pass-admissible. We write this as

$$\mathsf{Adm}^{\mathrm{loc}}_{\mathrm{Dix}}(\mathfrak{M}_i) = 1.$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Lemma A.146.3 (Image audit reduction)**.** *Assume the capture table is exhaustive. Then the residual-open part of the Dixon image is*

$$\mathcal{I}_{\mathsf{res}} = \left\{ \mathfrak{M}_i \;:\; \begin{array}{l} A_i \in A_{\mathrm{loc}}, \\ \mathsf{Adm}^{\mathrm{loc}}_{\mathrm{Dix}}(\mathfrak{M}_i) = 0, \\ A_i \text{ is neither blocked nor HC/MM absorbed} \end{array} \right\}.$$

Thus the residual audit is finite at the level of captured symbolic classes.

Proof. By the local pass, every declared admissible near-kernel route belongs to one of the classes A_i. By the local pass, every such class is either blocked, marked leak/debt, or assigned a Dixon mask. The predicate records exactly the leak, debt, hidden leakage, HC/MM debt, and selector/overlap admissibility tests. Hence any residual-open element must be one of the finitely many captured classes failing this predicate while not being otherwise blocked or absorbed. □

A.147 image table

The following table is the first Dixon image audit of the capture classes. The supports are symbolic; they specify which factor of the Dixon carrier is active, not a particle spectrum.

class	route meaning	HS(4) leak	HS(5) debt	Dixon support class	status	residual reason
A_1	collar interior	0	0	neutral collar support in $\mathbb{C}\otimes\mathbb{H}\otimes\mathbb{O}$	adm	none
A_2	S_{12} switch	0	0	routed selector edge support	Route	none
A_3	S_{23} switch	0	0	routed selector edge support	Route	none
A_4	non-adjacent S_{13}	–	–	no active mask	block	blocked by the local pass
A_5	HC/MM bookkeeping face	0	0	absorbed neutral/debt-free face	abs	none
A_6	overlap edge	0	0	overlap-preserved support	Route or adm	none

Proposition A.147.1 (No residual-open class in the declared sample image)**.** *For the declared symbolic image table, under the inherited assumptions*

$$\mathsf{Leak}^{(4)}_{C_7} = 0, \qquad \mathsf{Debt}^{(5)}_{\mathrm{cross}} = 0, \qquad L_{\mathrm{unr}} = 0, \qquad B_{\mathrm{HC/MM}} = 0,$$

and with the selector/overlap decisions unchanged, one has

$$\mathcal{I}_{\mathsf{res}} = \varnothing.$$

Proof. Class A_4 is blocked, hence it is not residual-open. Class A_5 is HC/MM absorbed with zero debt, hence it is not residual-open. Classes A_2, A_3, A_6 are routed through the selector/overlap grammar and preserve contraction margins, hence they are not residual-open. Class A_1 remains inside the certified collar interior and receives an admissible Dixon support class. Since the table contains all declared classes, no residual-open symbolic image element remains. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.148 Mask-image theorem

Theorem A.148.1 (Dixon image audit theorem)**.** *Assume the local proof chain photon-frame HC-halo capture chain, the HS(4)–HS(6) coding hypotheses, and the image table above. If the capture map is exhaustive over the declared near-kernel OP5 route grammar, then*

$$\mathrm{im}(\Xi^{\mathrm{loc}}_{\mathrm{Dix}}) \subseteq \mathsf{Code}^{\mathrm{adm}}_{\mathrm{Dix}} \cup \{\mathsf{block}, \mathsf{abs}, \mathsf{Route}\},$$

with no residual-open Dixon image class. Equivalently, within the declared HS(0)–HS(6) grammar, the remaining OP5 audit is reduced to the independent exhaustiveness check of the capture map and the zero-register checks for HS(4), HS(5), hidden leakage, and HC/MM debt.

Proof. The capture map reduces the route grammar to $A_1, \ldots, A_6$. The map sends each captured route to a Dixon mask or to an explicit blocked/leak/debt status. The table evaluates these statuses. Under the zero leakage/debt assumptions, no class is mapped to leak or debt. The non-adjacent selector class is blocked, the HC/MM face is absorbed debt-free, the selector and overlap edges are routed, and the collar interior is Dixon-admissible. Therefore the image contains only admissible Dixon masks and the three controlled non-residual statuses. □

Corollary A.148.2 (OP5 consequence inside the declared grammar)**.** *Inside the declared OP5=photon-frame HC-cone grammar, the audit leaves no uncontrolled OP5 exit in the captured HS(0)–HS(6) image, provided the capture map is exhaustive and all zero-register hypotheses remain true.*

Remark A.148.3 (What has not been proven)**.** *This is not a global OP5 proof outside the declared grammar. The decisive remaining mathematical work is to audit the full Active-Core route menu for any admissible near-kernel OP5 branch not represented by $A_1, \ldots, A_6$, and to preserve the zero-register conditions under integration into the monolithic source.*

A.149 Next audit target

The next useful layer should not add a new algebraic carrier. It should perform one of two integration checks:

(a) a full Active-Core route-menu scan showing that every admissible near-kernel OP5 branch is represented by the capture classes; or

(b) a monolithic integration of the local proof chain into the active source, followed by separate label, reference, layout, and route-menu residual audits.

If both checks pass, OP5 can be stated as locally closed inside the declared photon-frame HC-halo/Dixon-coding grammar, with the remaining caveat being only the scope of that grammar.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Integrated Delta Layer: Integrated Residual Register

Abstract

This layer audits the available route-menu surface after the local proof chain. Its purpose is not to introduce a new algebraic mechanism. It checks whether the local photon-frame HC-halo/Dixon coding chain leaves any declared admissible near-kernel OP5 route outside the finite capture menu $A_1, \ldots, A_6$. The result is deliberately scoped: the local proof chain delta stack is internally exhausted by the local proof chain classification, while the available monolithic basis source is only the local proof line. Therefore this layer certifies the declared delta-stack route image and records the exact remaining condition for a future source-native monolithic freeze: no active OP5 branch in the full Active Core may remain outside the capture classes or the named zero-register tests.

A.150 Audit scope

The available source surface consists of the local proof line monolithic Active Core basis and the local proof chain delta layers. Hence performs two separate checks.

1. **Delta-stack route audit.** This scans the local proof chain local photon-frame HC-halo, tunnel, finite-atlas, capture, and Dixon-coding chain.
2. **Basis-source compatibility scan.** This records the older the local pass OP5 route language that predates the Halo/Dixon grammar and must later be mapped or explicitly declared out of scope during a monolithic integration.

Remark A.150.1 (No false monolithic claim)**.** *The present file is a source-level audit layer, not a source-native rebuild of a single full the local proof line Active Core. It therefore proves a local exhaustiveness claim for the declared the local proof chain grammar and lists the remaining monolithic audit obligation separately.*

A.151 Delta-stack route menu

The local proof chain reduces the local near-kernel OP5 problem as follows:

$$\begin{gathered}\text{robust photon chart} \to \text{finite atlas} \to \text{double-}\varepsilon\text{ halo},\\ \text{double-}\varepsilon\text{ halo} \to \text{rank/selector/overlap registers} \to \text{capture classes},\\ \text{capture classes} \to \text{Dixon image audit}.\end{gathered}$$

The effective route alphabet after is

$$A_{\mathrm{loc}} = \{A_1, A_2, A_3, A_4, A_5, A_6\}.$$

The Dixon audit assigns these classes the following controlled statuses.

class	route meaning	status
A_1	collar interior	adm
A_2	S_{12} selector switch	route
A_3	S_{23} selector switch	route
A_4	non-adjacent S_{13} selector	block
A_5	HC/MM bookkeeping face	abs
A_6	atlas-overlap edge	route or adm

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition A.151.1 (residual support)**.** *Let* $\Xi^{\mathrm{loc}}_{\mathrm{Dix}}$ *be the route-to-Dixon map of and let* $\pi^{\mathrm{loc}}_{\mathrm{cap}}$ *be the capture map. The residual support is*

$$\mathcal{I}^{\mathrm{loc}}_{\mathsf{res}} := \left\{ r \in \mathcal{R}^{\mathsf{adm},\mathrm{near}\,K}_{\mathrm{OP5}} : \begin{array}{l} \pi^{\mathrm{loc}}_{\mathrm{cap}}(r) \notin \{A_1,\dots,A_6\}, \ \text{or} \\ \Xi^{\mathrm{loc}}_{\mathrm{Dix}}(r) \in \{\mathsf{leak},\mathsf{debt},\mathsf{res}\} \end{array} \right\}.$$

Thus $\mathcal{I}^{\mathrm{loc}}_{\mathsf{res}}$ *records exactly the active near-kernel routes that are either uncaptured or mapped to a non-controlled Dixon status.*

Proposition A.151.2 (Declared delta-stack residual vanishing)**.** *Inside the declared the local proof chain photon-frame* HC*-halo/Dixon grammar, the source scan finds no declared route class beyond* $A_1,\dots,A_6$*, and maps those classes only to* adm*,* block*,* abs*, or* route*. Hence*

$$\mathcal{I}^{\mathrm{loc}}_{\mathsf{res}} = \varnothing$$

for the declared delta-stack image.

Proof. The tunnel menu is reduced by the local proof chain to radial/HC boundary, selector, rank, and overlap checks. The normal-collar certificate removes rank-drop on certified pieces. The selector/overlap table blocks or routes the switch and overlap classes. The capture table enumerates the six remaining local near-kernel classes $A_1,\dots,A_6$. The Dixon map sends captured active classes into the 64-channel Dixon carrier or to explicit blocked/leak/debt statuses. The image table assigns A_1 to admissible, A_2, A_3, A_6 to routed/admissible, A_4 to blocked, and A_5 to absorbed. No table row is assigned leak, debt, or res. Therefore the declared local image has empty residual support. □

A.152 Source scan facts

The source scan of the local proof chain delta files produced the following structural metadata.

layer	lines	labels	raw lower-case halo	raw lower-case exit
the local pass	239	0	0	0
the local pass	245	0	0	0
the local pass	293	0	0	0
the local pass	317	0	1	1
the local pass	276	0	0	0
the local pass	257	0	0	0
the local pass	284	0	0	0
the local pass	296	0	0	0
the local pass	387	0	0	0
the local pass	277	0	0	0

The single lower-case occurrences are compatibility aliases in the source; they are not active rendered notation in the local proof chain line. No duplicate labels were found across the local proof chain delta files.

The available monolithic basis contains the older selector-gated OP5 route language and does not yet contain the later Halo, near-kernel, or Dixon route grammar as an integrated source-native block. The scan counts were:

© 2026 Marc Brendecke. Preprint — not peer reviewed.

anchor	count
`OP~5`	567
`OP5`	133
selector-gated reachability	13
TE budgets	4
HC/MM residual	2
post-promotion residual	7
OP~4	55
Dixon	8
Halo	0
near-kernel	0

This confirms that the local proof chain are a new local closure grammar over the basis, not yet a source-native monolithic replacement of every older OP5 paragraph.

A.153 Compatibility with the OP5 route surface

The selector-gated OP5(B) statement leaves four route obligations after a first reachability pass: TE budgets, HC/MM residual compatibility, post-promotion residual control, and OP4 stability. The local proof chain directly attacks the near-kernel HC-cone route obstruction, not the whole OP5(B) boundary-promotion package.

Definition A.153.1 (Monolithic residual register)**.** *For a future full Active-Core integration define*

$$\mathcal{R}^{\mathrm{loc}}_{\mathrm{mono}} := \mathcal{R}_{\mathrm{uncap}} \cup \mathcal{R}_{\mathrm{HS4\ leak}} \cup \mathcal{R}_{\mathrm{HS5\ debt}} \cup \mathcal{R}_{L_{\mathrm{unr}}} \cup \mathcal{R}_{B_{\mathrm{HC/MM}}} \cup \mathcal{R}_{\text{loc-res}}.$$

Here $\mathcal{R}_{\mathrm{uncap}}$ *denotes active near-kernel routes not mapped by* $\pi^{\mathrm{loc}}_{\mathrm{cap}}$*, while* $\mathcal{R}_{\text{loc-res}}$ *denotes any older OP5(B) budget, post-promotion, or OP4-stability condition not identified with the local* HC*-cone grammar.*

Theorem A.153.2 (route-menu exhaustiveness theorem, scoped form)**.** *Assume the working identification*

$$\mathrm{OP5}_{\mathrm{near}\,K} = \textit{photon-frame}\ \mathrm{HC}\textit{-cone source-return reflector}$$

and assume that the capture map is exhaustive over the declared near-kernel route grammar. If

$$\mathcal{R}_{\mathrm{HS4\ leak}} = \mathcal{R}_{\mathrm{HS5\ debt}} = \mathcal{R}_{L_{\mathrm{unr}}} = \mathcal{R}_{B_{\mathrm{HC/MM}}} = \varnothing$$

and the selector/overlap decisions are preserved under integration, then there is no uncontrolled near-kernel OP5 exit in the local proof-chain delta stack route menu.

If, in addition, a future monolithic scan proves

$$\mathcal{R}_{\mathrm{uncap}} = \varnothing \qquad \textit{and} \qquad \mathcal{R}_{\text{loc-res}} = \varnothing,$$

then the same statement may be promoted from a local delta-stack closure theorem to a monolithic local OP5 closure theorem inside the declared HC*-halo/Dixon grammar.*

Proof. The first part is exactly Proposition 1 combined with the zero-register hypotheses: every captured near-kernel route is admissible, blocked, absorbed, or routed, and no leak/debt/residual status remains. The second part adds the only missing monolithic conditions. If no older active branch lies outside the capture map and no residual obligation remains unmapped, the local exhaustiveness statement covers the integrated route menu for the declared grammar. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Corollary A.153.3 (Current OP5 status after)**.** *The local proof chain local route image is exhausted. The current remaining OP5 burden is no longer a free route explosion; it is the monolithic integration check*

$$\mathcal{R}^{\mathrm{loc}}_{\mathrm{mono}} = \varnothing.$$

Thus OP5 is locally closed inside the declared delta-stack grammar and remains open only at the full Active-Core integration/exhaustiveness boundary.

A.154 Next integration rule

The next source-native step should not add another algebraic layer. It should merge or append the local proof chain into the active source and run a residual search with the following pass/fail test:

$$\operatorname{supp}\big(\mathcal{R}^{\mathsf{adm},\mathrm{near}\,K}_{\mathrm{OP5}}\big) \subseteq \bigcup_{i=1}^{6} (\pi^{\mathrm{loc}}_{\mathrm{cap}})^{-1}(A_i) \cup \{\mathsf{block}, \mathsf{abs}, \mathsf{route}, \mathsf{adm}\}.$$

Any hit outside this set must be named explicitly as one of: uncaptured route, HS4 leakage, HS5 cross-side debt, hidden leakage, HC/MM debt, post-promotion residual, TE-budget residual, or OP4-stability residual.

A.155 Conservativity guards

- does not claim global OP5 closure outside the declared photon-frame HC-halo/Dixon grammar.
- does not claim OP1, OP4, QCD/confinement, CKM/PMNS, quark masses, or a Standard Model spectrum derivation.
- The Dixon carrier is 64-dimensional; 2^{64} is the Boolean activation mask count, not the Dixon dimension.
- HC/MM absorption remains explicit bookkeeping with zero unpaid debt, not residual deletion.
- A source-native monolithic rebuild remains required before making a public release-style OP5 closure statement.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Monolithic OP5 Promotion Audit

Definition A.155.1 (monolithic OP5 residual register)**.** *Let $\mathcal{R}^{\mathrm{loc}}_{\mathrm{mono}}$ denote the set of active OP5-facing near-kernel branches in the integrated Active-Core source which are not assigned to one of the capture classes*

$$A_1, \dots, A_6,$$

and are not discharged by the named zero-register tests

$$L_{\mathrm{unr}} = 0, \qquad B_{\mathrm{HC/MM}} = 0, \qquad R_{\mathrm{HS4}} = 0, \qquad D_{\mathrm{HS5}} = 0.$$

A hit is counted as residual only if it is active in the current local proof line OP5-facing grammar, not merely preserved historical witness language.

Proposition A.155.2 (integrated delta-stack exhaustion)**.** *In the integrated local proof chain delta stack, the local photon-frame HC-halo/Dixon route image is exhausted by the six classes*

$$A_1, \dots, A_6.$$

Their status assignment remains

$$A_1 \mapsto \mathsf{adm}, \quad A_2 \mapsto \mathsf{route}, \quad A_3 \mapsto \mathsf{route}, \quad A_4 \mapsto \mathsf{block}, \quad A_5 \mapsto \mathsf{abs}, \quad A_6 \mapsto \mathsf{route} \text{ } or \text{ } \mathsf{adm}.$$

Consequently the residual Dixon image inside the declared local grammar is empty:

$$\mathcal{I}^{\mathrm{loc}}_{\mathrm{res}} = \varnothing.$$

Proof. This is the source-native integration of the audit conclusion with the local proof chain delta bodies included in the same Active-Core source. The local proof-chain layers provide the robust local chart and finite-atlas candidate, the local proof chain reduce the OP5 route question to halo/tunnel, rank, selector, and overlap registers, supplies the finite capture menu, maps captured routes to the 64-dimensional Dixon carrier with a 2^{64}-state mask space, audits the reachable Dixon image, and verifies that no residual-open class remains in that declared delta image. □

Theorem A.155.3 (local OP5 closure in the declared HC-halo/Dixon grammar)**.** *Assume the working identification that the OP5-facing near-kernel branch is the photon-frame HC-cone/reflector route, and assume the inherited zero-register conditions*

$$L_{\mathrm{unr}} = 0, \qquad B_{\mathrm{HC/MM}} = 0, \qquad R_{\mathrm{HS4}} = 0, \qquad D_{\mathrm{HS5}} = 0.$$

Then the integrated local proof chain Active-Core delta stack leaves no uncontrolled local OP5 exit inside the declared photon-frame HC-halo/Dixon grammar.

Proof. By the capture layer, every declared admissible near-kernel route in the grammar is assigned to $A_1, \dots, A_6$, provided the capture map is exhaustive. By the selector/overlap layer, the active switch and overlap channels are blocked, routed, absorbed, or contraction-preserving. By the Dixon-image layers, every captured route has either a Dixon-admissible 64-channel mask or is classified as blocked, routed, or HC/MM-absorbed. The assumed zero-register conditions remove hidden leakage, unpaid HC/MM absorption, HS(4) meta-septum leakage, and HS(5) cross-side debt. Hence no residual-open OP5 class remains in the declared local grammar. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Remark A.155.4 (Scope of the promotion)**.** *This statement promotes the local proof-chain delta stack result into the available Active-Core source as a local OP5 closure theorem inside the declared HC-halo/Dixon grammar. It does not claim global OP5 closure outside that grammar, does not close OP1 or OP4, and does not derive QCD/confinement, CKM/PMNS values, quark masses, or a Standard Model spectrum.*

Remark A.155.5 (Inherited historical OP5 language)**.** *The older Active-Core basis contains preserved historical OP1–OP5 and OP5(B) statements from earlier release lines. In those blocks remain theorem-memory and genealogy unless explicitly referenced by the current HC-halo/Dixon grammar. The governing current promotion claim is the local theorem above.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

OP5 Closure Hardening and Old-Branch Mapping Audit

Target-version rule. This audit belongs to the provenance layer. In the reader-facing proof spine it is interpreted semantically, not as a mathematical version marker. It hardens the local OP5 promotion and maps inherited OP5 language into current, historical, guard, or outside-scope status.

Definition A.155.6 (governing OP5 status)**.** *The governing OP5 status is that OP5 is locally closed inside the declared photon-frame HC-halo/Dixon grammar under*

$$L_{\mathrm{unr}} = 0, \quad B_{\mathrm{HC/MM}} = 0, \quad R_{\mathrm{HS4}} = 0, \quad D_{\mathrm{HS5}} = 0.$$

This is not a global OP5 theorem outside that grammar and does not close OP1 or OP4.

Proposition A.155.7 (no-active-old-branch promotion rule)**.** *An inherited OP5-facing paragraph may be read as part of the current OP5 closure claim only if it is explicitly mapped into the HC-halo/Dixon grammar or is one of the named zero-register guards. Otherwise it is historical theorem-memory, a guard, or outside the scope of the local theorem.*

Theorem A.155.8 (compact OP5 closure spine)**.** *Inside the declared photon-frame HC-halo/Dixon grammar, OP5 has the finite proof spine*

$$\textit{HC cone} \Rightarrow \textit{double-}\varepsilon\textit{ halo} \Rightarrow \textit{tunnel reduction} \Rightarrow \textit{normal collar}$$

$$\Rightarrow \textit{selector/overlap routing} \Rightarrow \textit{Dixon image audit} \Rightarrow \textit{no residual local OP5 exit.}$$

Under the four zero-register assumptions, the local residual register of this spine is empty.

Remark A.155.9 (No version promotion)**.** *The present document is a v4.3.0 release-final. A later public line should be reserved for broader OP4/OP1 promotion or for a genuinely new architecture beyond the OP5 hardening chain.*

A.156 Finite Perturbation Threshold Atlas for the Triolic Orbit Scan

This section records a compact finite threshold atlas for the alternating triolic orbit-scan model. The purpose is not to promote the toy chart to a physical trajectory simulation, but to make the robustness statement falsifiable at the finite readout level.

Let the finite scan diagnostics be

$$\widetilde{E}_{N,M}, \qquad \widetilde{R}^{\min}_{N,M}, \qquad C^{\mathrm{even}}_{95}, \qquad C^{\mathrm{odd}}_{05},$$

where $\widetilde{E}_{N,M}$ is the bifocal defect, $\widetilde{R}^{\min}_{N,M}$ is the smallest guard margin, C^{even}_{95} is the 95-percentile even-sector core-sightline residual, and C^{odd}_{05} is the 5-percentile odd-sector complementarity residual.

Definition A.156.1 (Finite threshold predicate)**.** *A perturbed finite scan is threshold-admissible when*

$$\widetilde{E}_{N,M} \le \tau_E, \qquad \widetilde{R}^{\min}_{N,M} > 0, \qquad C^{\mathrm{even}}_{95} \le \tau_C, \qquad C^{\mathrm{odd}}_{05} \ge \tau_O.$$

The predicate is a finite readout certificate only. It does not assert a physical equation of motion.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Lemma A.156.2 (Index-level parity is threshold-invariant)**.** *For the shifted Kronecker transport*

$$\Sigma_L{}^{i}{}_{j} = \delta^{i}_{j+1 \bmod L},$$

the even–odd identities

$$\Pi_0\Sigma_L = \Sigma_L\Pi_1, \qquad \Pi_1\Sigma_L = \Sigma_L\Pi_0, \qquad \Pi_r\Sigma_L^2 = \Sigma_L^2\Pi_r$$

are independent of the geometric perturbation amplitudes.

Proposition A.156.3 (Finite threshold-atlas reading)**.** *The finite threshold atlas separates three layers:*

$$\textit{index parity}, \qquad \textit{protected-core guard}, \qquad \textit{geometric triolic balance}.$$

The first is exact under the shifted Kronecker transport. The second is certified by positive guard margin. The third is a genuine geometric residual and can fail under core-offset or antipodal-breaking perturbations.

Remark A.156.4 (Compact numerical atlas instance)**.** *In the compact finite atlas run, the one-parameter scans estimate the following last-stable / first-unstable transition windows:*

parameter	*last stable*	*first unstable*
core offset	0.030	0.045
antipodal-breaking jitter	0.080	*none*
lens asymmetry	0.315	0.360
symmetric radial ripple	0.030	0.045
twist amplitude	0.080	*none*

The atlas is not a physical proof. It is a reproducible diagnostic showing which perturbations preserve the local orbit-scan readout and which ones create a named obstruction.

Theorem A.156.5 (Finite threshold-atlas guard)**.** *A finite orbit-scan atlas may be used as a local readout certificate only when the threshold predicate holds. If the guard remains positive but the even-sector core-sightline residual exceeds threshold, then the failure is not a through-core event; it is a triolic-balance obstruction.*

A.157 Finite Core-Recentering Calibration for the Triolic Orbit Scan

The threshold atlas identifies core offset as the most sensitive finite obstruction for the triolic orbit scan. This section records the corresponding calibration layer. The purpose is conservative: a displaced declared core is not interpreted as a permitted through-core motion. It is treated as a calibration defect unless the residual remains after recentering.

Let $\mathcal{I}_0$ denote the even-sector sample indices and let (p_n, q_n), $n \in \mathcal{I}_0$, be the paired even-sector readout points. The finite midpoint estimator is

$$\widehat{K} := \frac{1}{|\mathcal{I}_0|} \sum_{n \in \mathcal{I}_0} \frac{p_n + q_n}{2}.$$

If K_0 is the declared collapse centre, the recentering displacement is

$$\Delta K := \widehat{K} - K_0.$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition A.157.1 (Recentered triolic residuals)**.** *For each even pair define*

$$C_n(K) := \frac{\|(p_n - K) \times (q_n - K)\|}{\|p_n - K\| \, \|q_n - K\|}, \qquad M_n(K) := \left\| K - \frac{p_n + q_n}{2} \right\|.$$

The pre-calibration residuals are $C_n(K_0), M_n(K_0)$*. The recentered residuals are* $C_n(\widehat{K}), M_n(\widehat{K})$*.*

Lemma A.157.2 (Midpoint recovery in the antipodal case)**.** *Assume that all even-sector pairs are antipodal about a common point* K_**, that is*

$$\frac{p_n + q_n}{2} = K_* \qquad (n \in \mathcal{I}_0).$$

Then $\widehat{K} = K_*$*. In particular, if the declared centre* K_0 *is offset from the actual finite balance centre, the estimator recovers the balance centre from the paired readout data.*

Proof. The estimator is the average of the pair midpoints. If every midpoint equals K_*, their average equals K_*. □

Proposition A.157.3 (Finite recentering criterion)**.** *A finite core-offset defect is called calibratable when*

$$C_{95}(\widehat{K}) \le \tau_C, \qquad R^{\min}(\widehat{K}) > 0,$$

and the displacement ΔK *is recorded as a calibration displacement, not as a through-core route. If these inequalities fail, the residual is a persistent triolic-balance obstruction.*

Remark A.157.4 (Numerical calibration instance)**.** *In the finite calibration probe, the uncalibrated core-offset threshold was approximately*

$$\textit{last stable} = 0.030, \qquad \textit{first unstable} = 0.040.$$

After midpoint recentering, the scan remained stable through the tested range, with

$$\max C_{95}(\widehat{K}) \approx 6.639e-16, \qquad \min R^{\min}(\widehat{K}) \approx 1.180 > 0.$$

Thus the tested core-offset failure is a calibratable centre-declaration defect in this finite readout chart, not a permitted crossing of the protected core.

Theorem A.157.5 (Core-recentering guard)**.** *Finite recentering can repair a triolic scan only as a readout calibration if three conditions hold: the recentered even-sector residual is below threshold, the protected-core guard remains positive, and the displacement* ΔK *is reported as calibration provenance. Under these conditions the repaired scan may be used as a local readout certificate, but it does not promote the orbit to a physical through-core trajectory.*

Proof. The residual condition restores the finite even-sector core-sightline test. The positive guard condition keeps all sampled points outside the protected core. Recording ΔK prevents the calibration from being mistaken for a new dynamical claim. Hence the procedure repairs the declared centre of the readout chart only; it does not assert a through-core motion. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.158 Crystal–Fractal Readout Probe from Calibrated Orbit Scans

The calibrated triolic orbit scan separates a centre-calibration problem from a residual-geometry problem. This makes the crystal and fractal readings operational: the crystal reading concerns coherence across directions, while the fractal reading concerns recurrence across scales.

Definition A.158.1 (Ringwise calibrated balance centres)**.** *For each scan ring ℓ, define the even-triole balance centre*

$$\widehat{K}_\ell := \frac{1}{|\mathcal{I}_0|} \sum_{n \in \mathcal{I}_0} \frac{p_{\ell n} + q_{\ell n}}{2}.$$

The calibrated global centre is

$$\widehat{K} := \frac{1}{L} \sum_{\ell=1}^{L} \widehat{K}_\ell.$$

The finite crystal-coherence residual is

$$\mathcal{C}_{\mathrm{crys}} := \left(\frac{1}{L} \sum_{\ell=1}^{L} \|\widehat{K}_\ell - \widehat{K}\|^2 \right)^{1/2}.$$

Definition A.158.2 (Residual readout set across scales)**.** *Let $s \in S$ range over a finite set of scan scales. For each scale, let $\Sigma_K^{\Gamma,s}$ be the finite periapsis envelope and let*

$$\mathcal{R}_s := \{x_{\ell m}^s : \mathcal{E}_F(x_{\ell m}^s) > \tau_{\mathrm{res}}\}$$

be the residual readout set above the declared bifocal-defect threshold. The multiscale residual readout is

$$\mathcal{R}_{\mathrm{multi}} := \bigcup_{s \in S} \mathcal{R}_s.$$

Definition A.158.3 (Finite residual box-counting diagnostic)**.** *For box length $\delta > 0$, let $N_{\mathcal{R}}(\delta)$ be the number of occupied boxes meeting $\mathcal{R}_{\mathrm{multi}}$. A finite scaling diagnostic is obtained by fitting*

$$\log N_{\mathcal{R}}(\delta) \approx D_{\mathrm{res}} \log(1/\delta) + b.$$

The number D_{res} is a finite residual-scaling estimate, not a proof of Hausdorff dimension.

Proposition A.158.4 (Crystal–fractal separation)**.** *The calibrated orbit-scan data split into two independent readouts:*

$$\mathcal{C}_{\mathrm{crys}} \quad \text{and} \quad D_{\mathrm{res}}.$$

Small $\mathcal{C}_{\mathrm{crys}}$ certifies coherent directional centering, i.e. a crystal-like local cell centre. Nontrivial, scale-persistent $\mathcal{R}_{\mathrm{multi}}$ certifies residual scaling structure, i.e. a fractal-like defect readout. Neither statement asserts a physical trajectory through the protected core.

Remark A.158.5 (Finite diagnostic instance)**.** *In the finite toy diagnostic, the maximum ring-centre RMS spread was*

$$\mathcal{C}_{\mathrm{crys}}^{\max} \approx 8.699e - 17,$$

with 30216 residual points above threshold across the multiscale scan. The box-counting fit returned

$$D_{\mathrm{res}} \approx 2.435.$$

This is a diagnostic number: it indicates that residual structure can be measured across scales, not that the protected-core surface has already been identified with a mathematical fractal.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Theorem A.158.6 (Readout guard for crystal and fractal claims). *A crystal claim is admissible only as a statement about coherence of the calibrated centres $\widehat{K}_\ell$. A fractal claim is admissible only as a statement about the scaling of residual readout sets across declared scan scales. The orbit-scan data do not by themselves promote either claim to a global physical closure theorem.*

A.159 Residual Classification Atlas for Crystal–Fractal Readouts

The crystal–fractal readout probe shows that residual sets can be measured across directions and scales. The present atlas classifies the residuals into finite diagnostic classes. This is not a physical closure theorem; it is a readout classification layer.

Definition A.159.1 (Residual classification alphabet). *Let $\mathcal{R}_{\mathrm{multi}}$ be the multiscale residual readout set. The finite classification alphabet is*

$$\mathfrak{A}_{\mathcal{R}} = \{\mathrm{crys}, \mathrm{frac}, \mathrm{lens}, \mathrm{guard}, \mathrm{noise}\}.$$

Here crys *denotes directional or angular recurrence,* frac *denotes recurrence across scan scales,* lens *denotes high bifocal stress,* guard *denotes proximity to the protected-core guard, and* noise *denotes the unclassified remainder under the declared finite rules.*

Definition A.159.2 (Class tags and primary class). *A residual point $x \in \mathcal{R}_{\mathrm{multi}}$ may carry several tags:*

$$\tau_{\mathrm{crys}}(x), \quad \tau_{\mathrm{frac}}(x), \quad \tau_{\mathrm{lens}}(x), \quad \tau_{\mathrm{guard}}(x).$$

For disjoint counting, the primary class is assigned by the conservative order

$$\mathrm{guard} \succ \mathrm{lens} \succ \mathrm{frac} \succ \mathrm{crys} \succ \mathrm{noise}.$$

Thus a point close to the guard is counted as a guard residual even if it also has a scale-recurrence tag.

Proposition A.159.3 (Residual classification separation). *The classification atlas separates four causes of residual structure:*

$$\textit{directional recurrence}, \quad \textit{scale recurrence}, \quad \textit{bifocal stress}, \quad \textit{guard proximity}.$$

Only the first two support crystal–fractal interpretation. Lens and guard classes are obstruction classes and must not be promoted to crystal or fractal evidence without an additional argument.

Remark A.159.4 (Finite classification instance). *In the finite diagnostic instance,*

$$|\mathcal{R}_{\mathrm{multi}}| = 30216.$$

The primary class counts are

class	*count*
crys	0
frac	27194
lens	3022
guard	0
noise	0

© 2026 Marc Brendecke. Preprint — not peer reviewed.

The finite box-counting estimates by primary class are

class	D_{box}
crys	*n/a*
frac	2.379
lens	1.874
guard	*n/a*
noise	*n/a*

These numbers are finite readout diagnostics only. They do not identify the protected-core surface with a mathematical fractal.

Theorem A.159.5 (Classification guard)**.** *A residual point may be used as crystal evidence only through its directional recurrence tag and as fractal evidence only through its scale-recurrence tag. If the point is primarily classified as lens or guard, then it remains an obstruction point until a separate routing or calibration step reclassifies it.*

Proof. The tags are defined by distinct finite predicates: angular recurrence, scale support, bifocal stress, and guard proximity. The primary order assigns obstruction classes before interpretive classes. Hence a lens-stressed or guard-near residual cannot be counted as crystal–fractal evidence merely because it appears in the residual set. □

A.160 Finite Hopf-Linking Readout Test

The orbit-scan, calibration, crystal–fractal, and residual-classification layers describe metric and residual structure. The next readout layer asks a topological question: whether the scan rings behave like a finite family of Hopf-linked fibres.

Definition A.160.1 (Finite Hopf-fibre chart)**.** *Let $S^3 \subset \mathbb{C}^2$ and let a fibre over a base point $(\alpha, \beta) \in S^2$ be parametrised by*

$$z_1(t) = e^{i(t+\beta)} \cos(\alpha/2), \qquad z_2(t) = e^{it} \sin(\alpha/2), \qquad t \in [0, 2\pi).$$

Writing $z_1 = X_1 + iX_2$ and $z_2 = X_3 + iX_4$, the finite chart uses the stereographic projection

$$(X_1, X_2, X_3, X_4) \longmapsto \frac{(X_1, X_2, X_3)}{1 - X_4} \in \mathbb{R}^3.$$

A finite Hopf readout family is a collection

$$\Gamma_H = \{\gamma_1, \ldots, \gamma_N\}$$

of such projected fibres sampled by a finite polygonal mesh.

Definition A.160.2 (Finite Gauss-linking estimator)**.** *For two disjoint closed polygonal curves γ_i, γ_j, define*

$$\widehat{\mathrm{Lk}}(\gamma_i, \gamma_j) := \frac{1}{4\pi} \sum_{a,b} \frac{(r_a - r_b) \cdot (\Delta r_a \times \Delta r_b)}{\|r_a - r_b\|^3},$$

where r_a, r_b are segment midpoints and $\Delta r_a, \Delta r_b$ are the corresponding segment vectors. This is a finite quadrature readout of the Gauss linking integral.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition A.160.3 (Linking stability matrix)**.** *The finite linking matrix is*

$$L_{ij} := \widehat{\mathrm{Lk}}(\gamma_i, \gamma_j), \qquad L_{ii} := 0.$$

For a declared tolerance $\tau_L > 0$, a pair is called Hopf-stable when

$$\big||L_{ij}| - 1\big| \le \tau_L.$$

Pairs that fail this condition are finite topological readout obstructions.

Proposition A.160.4 (Topological readout separation)**.** *The Hopf-linking test is independent of the residual classification alphabet. A residual point may be scale-recurring or lens-stressed, while the surrounding fibre pair remains topologically Hopf-stable. Conversely, a linking failure is a topological obstruction even if the local metric residual is small.*

Remark A.160.5 (Finite diagnostic instance)**.** *In the finite diagnostic instance, $N = 10$ projected Hopf fibres were sampled. There are 45 unordered fibre pairs. With tolerance $\tau_L = 0.180$, the run returned*

$$\#\text{stable pairs} = 45, \qquad \#\text{unstable pairs} = UN45.$$

The mean and maximum deviations from unit linking were

$$\overline{\Delta}_L \approx 0.0001, \qquad \Delta_L^{\max} \approx 0.0001.$$

These numbers are finite readout diagnostics. They do not prove that the physical orbit-scan space is globally a Hopf fibration.

Theorem A.160.6 (Hopf-linking guard)**.** *A finite Hopf-linking readout may support a local topological-organisation claim only when the sampled fibre pairs are disjoint and the linking stability matrix is unit-stable within the declared tolerance. If a pair fails this test, the failure is a topological readout obstruction, not a metric residual and not a through-core trajectory.*

Proof. The estimator is computed from closed curve geometry and compares pairwise linking. Its input is not the residual magnitude at a point but the relative winding of two closed fibres. Therefore a failure of unit linking identifies a topological obstruction in the finite readout layer. Since the curves are sampled outside the protected core, such a failure is not a permission to cross the core. □

A.161 Hopf-Stable Residual Carrier Routing

The finite Hopf-linking readout verifies that the sampled fibre family is topologically stable in the local chart. The residual classification atlas identifies metric residual classes. The present layer couples these two readouts by routing residual points to their nearest Hopf-stable carrier fibres.

Definition A.161.1 (Normalised residual-carrier assignment)**.** *Let $\mathcal{R}_{\mathrm{multi}}$ be the multiscale residual set and let*

$$\Gamma_H = \{\gamma_1, \ldots, \gamma_N\}$$

be the finite Hopf-fibre family. After applying a common centred normalisation to the residual cloud and the Hopf-fibre cloud, define

$$c(x) := \operatorname*{arg\,min}_{1 \le j \le N} \mathrm{dist}(x, \gamma_j).$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

The carrier distance is

$$d_H(x) := \operatorname{dist}(x, \gamma_{c(x)}).$$

For a declared threshold τ_H, a residual is called carrier-resolved when

$$d_H(x) \le \tau_H.$$

Definition A.161.2 (Hopf-stable residual carrier)**.** *A carrier-resolved residual x is called Hopf-stably carried when its assigned fibre belongs to a unit-stable linking matrix:*

$$\big||L_{ij}| - 1\big| \le \tau_L \quad \text{for all tested } i \ne j.$$

Residuals that fail the carrier threshold are off-carrier residuals, not through-core trajectories.

Proposition A.161.3 (Metric residual versus topological carrier)**.** *Residual class and carrier stability are logically distinct. A point can be fractal-recurring or lens-stressed as a metric residual while still being carried by a Hopf-stable fibre family. Conversely, a non-unit linking matrix would create a topological obstruction even if the metric residual class were small.*

Remark A.161.4 (Finite carrier-routing instance)**.** *In the finite diagnostic instance,*

$$|\mathcal{R}_{\text{multi}}| = 30216, \qquad N = 10.$$

The Hopf-linking matrix had

$$45$$

stable pairs out of

$$45$$

tested pairs. With carrier threshold

$$\tau_H = 0.6835,$$

the carrier routing returned

$$\#\text{carrier resolved} = 28705, \qquad \#\text{off carrier} = 1511.$$

The carrier-resolved fraction was

$$\rho_H \approx 0.950.$$

The normalised carrier-load entropy was

$$H_H \approx 0.986.$$

Theorem A.161.5 (Hopf-stable residual carrier guard)**.** *If the Hopf-linking matrix is unit-stable and a residual is carrier-resolved, then the residual may be read as metric or scale structure carried by stable local Hopf topology. It may not be promoted to global physical closure. If a residual is off-carrier, the correct conclusion is a carrier-routing obstruction, not a collapse-core crossing.*

Proof. The Hopf-linking matrix tests pairwise topological organisation of the carrier fibres. The residual classification tests metric, scale, lens, and guard properties of points. The carrier assignment relates these layers by nearest finite readout distance. Therefore a carrier-resolved residual inherits only the finite local carrier stability, while an off-carrier residual records the failure of this finite routing predicate. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.162 Off-Carrier Residual Obstruction Split

The Hopf-stable carrier routing leaves a finite off-carrier residual set. This section classifies that remainder. The purpose is to prevent an off-carrier point from being misread as a through-core trajectory. It is a routing obstruction inside the finite readout chart.

Definition A.162.1 (Off-carrier residual set)**.** *Let*

$$\mathcal{R}_{\text{off}} := \{x \in \mathcal{R}_{\text{multi}} : d_H(x) > \tau_H\}$$

be the residuals that fail the declared Hopf-carrier distance threshold. These points are not discarded. They are routed into an obstruction alphabet.

Definition A.162.2 (Off-carrier obstruction alphabet)**.** *The finite obstruction alphabet is*

$$\mathfrak{A}_{\text{off}} = \{\text{lens-extreme}, \text{scale-boundary}, \text{projection-chart}, \text{missing-fiber}, \text{persistent}\}.$$

The meanings are:

class	*readout meaning*
lens-extreme	*high bifocal defect or lens stress*
scale-boundary	*finite scale-atlas boundary effect*
projection-chart	*large chart distance / possible projection artifact*
missing-fiber	*near-threshold point requiring denser carrier family*
persistent	*unexplained carrier obstruction*

Proposition A.162.3 (Off-carrier split guard)**.** *Every off-carrier residual must be classified before it can be used as evidence for a topological defect. Lens-extreme, scale-boundary, projection-chart, and missing-fiber classes are repairable or extendable readout tasks. Only the persistent class is carried forward as a named carrier obstruction.*

Remark A.162.4 (Finite off-carrier diagnostic instance)**.** *In the finite carrier-routing instance,*

$$|\mathcal{R}_{\text{off}}| = 1511.$$

The primary obstruction counts are

class	*count*
lens-extreme	315
scale-boundary	633
projection-chart	0
missing-fiber	563
persistent	0

The box-counting estimates for the nonempty classes are

class	D_{box}
lens-extreme	1.330
scale-boundary	1.508
projection-chart	*n/a*
missing-fiber	1.588
persistent	*n/a*

These values are finite diagnostics only.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Theorem A.162.5 (Off-carrier obstruction contract)**.** *The off-carrier set admits the following conservative contract. If an off-carrier residual is lens-extreme, it is routed to a lens refinement. If it is scale-boundary, the scale atlas must be extended. If it is projection-chart, a second chart must be tested. If it is missing-fiber, the Hopf carrier family must be refined. Only if none of these explanations applies is the residual kept as a persistent finite carrier obstruction.*

Proof. The classes are defined by finite predicates that test bifocal stress, scale support, carrier distance, and near-threshold carrier failure. Each predicate corresponds to a specific repair or refinement task. Therefore a residual cannot be promoted to persistent obstruction until the earlier finite routing explanations have failed. □

A.163 Closure Ledger and Release-Final Audit for the Orbit–Hopf Chain

The preceding local readout chain has reached a natural audit boundary. This section records what has been certified, what remains only diagnostic, and what is explicitly not claimed. The ledger is a release-final bookkeeping layer, not a new physical closure theorem.

Definition A.163.1 (Finite orbit–Hopf readout chain)**.** *The finite chain audited here is*

alternating triolic orbit scan $\longrightarrow$ *core re-centring calibration* $\longrightarrow$ *crystal–fractal readout* $\longrightarrow$ *residual classification* $\longrightarrow$ *Hopf-linking readout* $\longrightarrow$ *Hopf-stable residual carrier routing* $\longrightarrow$ *off-carrier obstruction split.*

Each arrow is a finite readout transition under declared chart, sampling, and guard assumptions.

Definition A.163.2 (Closure ledger states)**.** *A stage may receive one of the following ledger states:*

defined, diagnostic, calibration-resolved, classified, carrier-resolved, persistent.

A state of persistent *is reserved for a named finite obstruction that is not explained by calibration, scale extension, chart correction, lens routing, or carrier refinement.*

Proposition A.163.3 (Finite local ledger result)**.** *In the present finite audit instance, the chain records no persistent finite-carrier obstruction:*

$$\#\mathcal{R}_{\text{persistent}} = 0.$$

The carrier routing resolves

28705

residuals and leaves

1511

off-carrier residuals, all of which are classified as lens-extreme, scale-boundary, or missing-fibre tasks.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. The residual classification layer splits the multiscale residual set into finite primary classes. The Hopf-linking layer verifies unit-stable linking for

$$45/45$$

tested fibre pairs. The carrier-routing layer assigns residuals to this finite Hopf-stable carrier family and records the off-carrier remainder. The off-carrier split then classifies that remainder as

class	count
lens-extreme	315
scale-boundary	633
projection-chart	0
missing-fiber	563
persistent	0

so no member of the off-carrier set is carried forward as a persistent finite carrier obstruction. □

Remark A.163.4 (Numerical readout summary)**.** *The finite crystal–fractal and Hopf readouts record*

$$\mathcal{C}_{\mathrm{crys}}^{\max} \approx 8.699e-17, \qquad D_{\mathrm{res}} \approx 2.435117,$$

with

$$|\mathcal{R}_{\mathrm{multi}}| = 30216.$$

The residual classification counts are

primary class	*count*
frac	27194
lens	3022
guard	0
noise	0

The Hopf readout records mean and maximum unit-linking deviations

$$\overline{\Delta}_L \approx 1.1382e-04, \qquad \Delta_L^{\max} \approx 1.4636e-04.$$

The carrier-resolved fraction is

$$\rho_H \approx 0.950,$$

with normalised carrier-load entropy

$$H_H \approx 0.986.$$

Theorem A.163.5 (Release-final guard)**.** *The finite orbit–Hopf chain is locally reviewable as a release-final readout block with no persistent finite-carrier obstruction. It does not imply a global physical closure theorem, a through-core trajectory, or a global Hopf fibration of the physical model.*

Proof. Every stage in the chain is finite and chart-dependent. The orbit scan and perturbation atlas are metric diagnostics; the re-centring layer is a calibration step; the crystal–fractal readout is a finite coherence and scaling diagnostic; the residual classification layer is a finite alphabet; the Hopf-linking layer is a sampled topological readout; and the carrier and off-carrier layers are finite routing predicates. Since no stage removes the chart, sampling, and guard assumptions, the conclusion is a local release-final ledger statement only. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Corollary A.163.6 (Next release tasks)**.** *The remaining non-closure tasks are:*

(1) *extend the scale atlas for scale-boundary residuals,*
(2) *increase Hopf-carrier density for missing-fibre residuals,*
(3) *route lens-extreme residuals through bifocal/lens-stress refinement,*
(4) *insert a final version-specific DOI only at actual release,*
(5) *optionally rebuild the Historical Witness in the same release packet.*

A.164 v4.4 Roadmap

This roadmap records the refinement and OP1/OP4 closure-candidate audit plan after the v4.3.0 local orbit–Hopf readout release. It is a roadmap and audit-contract layer, not a new closure claim. The v4.4 target is

$$\text{v4.4} = \text{Refinement} + \text{OP1/OP4 closure-candidate audit.}$$

The corresponding v4.4 target is reserved for possible OP1/OP4 closure attempts, and only if the v4.4 audit contracts are satisfied.

A.164.1 Starting point from the v4.3.0 local readout chain

The v4.3.0 local chain consists of

$$\begin{aligned}&\text{Orbit Scan} \to \text{Core Recentering} \to \text{Crystal/Fractal Readout}\\ &\to \text{Residual Classification} \to \text{Hopf Linking}\\ &\to \text{Carrier Routing} \to \text{Off-Carrier Split.}\end{aligned}$$

Its finite ledger reports no persistent finite carrier obstruction in the declared local readout chart:

$$\#\mathcal{R}_{\text{persistent}} = 0.$$

The remaining explanatory classes are not ignored. They become the v4.4 refinement tasks:

$$\#\mathcal{R}_{\text{missing fiber}} = 563, \qquad \#\mathcal{R}_{\text{scale boundary}} = 633, \qquad \#\mathcal{R}_{\text{lens extreme}} = 315.$$

The interpretation is conservative: these numbers define finite readout tasks, not physical trajectories through the protected core.

A.164.2 Strand A: local refinement of the orbit–Hopf chain

The first v4.4 strand refines the explanatory classes left by v4.3.0.

Dense Hopf carrier refinement. The missing-fibre class is tested by increasing the carrier family, for example

$$N = 10 \longrightarrow 20 \longrightarrow 40,$$

while rechecking the finite Hopf-linking matrix. The task succeeds only if the missing-fibre count drops without producing non-unit linking pairs.

Scale atlas extension. The scale-boundary class is tested by extending the scan-scale atlas both inward and outward. The task succeeds only if boundary residuals are reduced or explicitly reclassified while the residual scaling diagnostics remain stable within declared tolerances.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Lens-extreme local refitting. The lens-extreme class is routed through local bifocal refitting. A local focus-pair correction

$$(F_-, F_+) \longmapsto (\widehat{F}_-, \widehat{F}_+)$$

may reduce bifocal stress. If it does not, the residual is not erased; it is carried as explicit lens debt.

Dual projection chart audit. The Hopf-linking and carrier-routing diagnostics are repeated in a second projection chart or pole choice. The task succeeds only if the finite linking/carrier conclusions are chart-stable.

v4.4 refinement ledger. The refinement strand closes with a ledger splitting the local residuals into resolved, calibrated, reducible, and explicitly open classes. It must not create a new unexplained persistent finite carrier obstruction.

A.164.3 Strand B: OP1/OP4 ingress and closure-candidate audit

The second v4.4 strand tests whether OP1 and OP4 can be legitimately routed into the refined orbit–Hopf chain. It does not close OP1 or OP4.

Definition A.164.1 (OP ingress contract)**.** *For $j \in \{1, 4\}$, an OPj ingress contract is an explicitly declared map*

$$\mathcal{D}_{\mathrm{OP}j} \longrightarrow \mathcal{R}^{\mathrm{OP}j}_{\mathrm{orbit/Hopf}}$$

from the OPj obstruction grammar into the refined finite orbit–Hopf readout chain. The map is admissible only if it preserves the protected-core guard, introduces no unbudgeted HC/MM debt, and does not identify a finite readout residual with a physical through-core trajectory.

Definition A.164.2 (OP closure-candidate status)**.** *After ingress and routing, OPj receives exactly one status:*

$$\text{closed-candidate}, \quad \text{reduced}, \quad \text{blocked}, \quad \text{not applicable}.$$

The status closed-candidate *is permitted only when*

$$\mathcal{R}^{\mathrm{OP}j}_{\mathrm{persistent}} = 0$$

under the refined carrier, scale, lens, and dual-chart audits, and when no new unbudgeted debt is introduced.

Proposition A.164.3 (v4.4 does not close OP1 or OP4)**.** *The v4.4 line may define OP1/OP4 ingress contracts and closure-candidate ledgers, but it does not assert OP1 or OP4 closure.*

Proof. The v4.4 tasks are refinement, routing, and audit tasks. Their output is a status ledger. A closure theorem would require a further proof layer using the closed-candidate status as input. That proof layer is reserved for v4.4. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.164.4 v4.4 handoff criteria

A v4.4 closure attempt for OP1 or OP4 is allowed only when the v4.4 status is closed-candidate. The minimal handoff criteria are:

(H1) protected-core guard preserved;
(H2) no new unbudgeted HC/MM, lens, carrier, scale, or guard debt;
(H3) $\mathcal{R}^{\mathrm{OP}j}_{\mathrm{persistent}} = 0$ after refined routing;
(H4) dual-chart / carrier stability holds;
(H5) all remaining classes are resolved, calibrated, or explicitly open.

If any criterion fails, v4.4 may still report reduction or blockage, but not closure.

Theorem A.164.4 (v4.4 roadmap guard)**.** *The v4.4 roadmap is a conservative preparation layer. It can strengthen the local orbit–Hopf readout chain and can prepare OP1/OP4 closure candidates, but it cannot by itself promote OP1, OP4, or the global physical model to a closure theorem.*

Proof. Each v4.4 task is formulated as a finite refinement, chart-stability, residual-routing, or ingress-contract test. None of these tasks supplies the additional proof layer needed for global OP closure. Therefore the only admissible v4.4 conclusion is a ledger status and a v4.4 handoff decision. □

A.165 Dense Hopf Carrier Refinement

This section begins the v4.4 refinement branch. The v4.3 carrier routing left a finite missing-fibre class. The present test asks whether that class is a true obstruction or a finite carrier-density artefact.

Definition A.165.1 (Dense carrier family)**.** *Let*

$$\Gamma_H^{(N)} = \{\gamma_1, \dots, \gamma_N\}$$

be a finite Hopf-carrier family sampled in the same stereographic chart as the v4.3 finite Hopf-linking readout. The dense refinement tests

$$N \in \{10, 20, 40\}$$

while keeping the v4.3 carrier threshold τ_H fixed. A residual is resolved at density N when

$$d_H^{(N)}(x) \le \tau_H.$$

Definition A.165.2 (Missing-fibre refinement predicate)**.** *Let $\mathcal{R}^{(10)}_{\mathrm{miss}}$ denote the v4.3 missing-fibre residual class. The dense refinement residue is*

$$\mathcal{R}^{(N)}_{\mathrm{miss}} := \{x \in \mathcal{R}^{(10)}_{\mathrm{miss}} : d_H^{(N)}(x) > \tau_H\}.$$

The missing-fibre class is called density-resolved at level N if

$$|\mathcal{R}^{(N)}_{\mathrm{miss}}| = 0.$$

Proposition A.165.3 (Carrier-density refinement guard)**.** *Dense carrier refinement may reduce a missing-fibre residual class only if the Hopf-linking stability matrix remains unit-stable at the refined carrier density. Otherwise the refinement trades a metric routing issue for a topological carrier issue.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Remark A.165.4 (Finite dense-carrier diagnostic)**.** *Using the fixed v4.3 threshold*

$$\tau_H = 0.6835,$$

the finite dense-carrier run produced the following ledger:

N	*stable pairs*	*all off-carrier*	*old missing remaining*	$\max \Delta_L$
10	45/45	1512	563	0.0016
20	190/190	1283	444	0.0016
40	780/780	1159	391	0.0020

The test is finite and local. It does not assert OP1 or OP4 closure.

Theorem A.165.5 (Dense carrier handoff)**.** *If the missing-fibre class decreases under increasing N while the Hopf linking matrix remains unit-stable, then the missing-fibre class is a carrier resolution task rather than a persistent topology obstruction. Any remaining points are passed to the v4.4 refinement ledger, not to OP closure.*

Proof. The threshold τ_H is held fixed from the v4.3 carrier readout. Hence a newly resolved point is explained by a denser carrier family rather than by a looser acceptance threshold. Unit-stable linking prevents the denser carrier family from introducing a new topological defect. Remaining points are exactly those still outside the fixed carrier threshold after refinement. □

A.166 Scale Atlas Extension for Boundary Residuals

The off-carrier split identifies a finite scale-boundary residual class. This class is not a persistent obstruction until the scale atlas has been extended. The present layer enlarges the scan-scale window and tests whether formerly boundary-supported sites acquire interior scale support.

Definition A.166.1 (Extended scale atlas)**.** *Let the original scale window be*

$$S_0 = \{1.00, 0.82, 0.67, 0.55, 0.45, 0.37\}.$$

The v4.4 extension uses

$$S_1 = \{1.30, 1.15, 1.00, 0.82, 0.67, 0.55, 0.45, 0.37, 0.30, 0.25, 0.20\}.$$

For a residual site (ℓ, m), define its extended support

$$\sigma_{S_1}(\ell, m) := \#\{s \in S_1 : x^s_{\ell m} \in \mathcal{R}_s\}.$$

A former scale-boundary site is called extension-resolved when

$$\sigma_{S_1}(\ell, m) > 4.$$

Definition A.166.2 (Scale-window dimension stability)**.** *Let D_{S_0} and D_{S_1} be finite box-counting estimates for the residual readout set in the original and extended scale windows. The scale stability span is*

$$\Delta D_{\text{scale}} := \max(D_{S_0}, D_{S_1}, D_{\text{outer}}, D_{\text{inner}}) - \min(D_{S_0}, D_{S_1}, D_{\text{outer}}, D_{\text{inner}}),$$

where D_{outer} and D_{inner} are the corresponding estimates for the newly added outer and inner scale windows.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proposition A.166.3 (Scale-boundary guard). *A residual classified as scale-boundary cannot be promoted to persistent obstruction until it remains boundary-supported after an explicit scale-atlas extension. If its extended support becomes interior, it is a finite-window artifact rather than a persistent carrier obstruction.*

Remark A.166.4 (Finite scale-extension instance). *In the finite v4.4 diagnostic, the old off-carrier scale-boundary class had*

$$N^0_{\mathrm{scale}} = 633$$

rows, supported on

$$N^0_{\mathrm{sites}} = 633$$

distinct residual sites. After extension to S_1,

$$N_{\mathrm{rows,resolved}} = 633, \qquad N_{\mathrm{rows,unresolved}} = 0.$$

At the site level,

$$N_{\mathrm{sites,resolved}} = 633, \qquad N_{\mathrm{sites,unresolved}} = 0.$$

The box-counting estimates were

window	D_{box}
S_0	2.435
S_1	2.520
outer added	2.171
inner added	2.291

with scale-stability span

$$\Delta D_{\mathrm{scale}} \approx 0.348.$$

Theorem A.166.5 (Scale-extension contract). *The scale atlas extension may reduce a scale-boundary class to a finite-window artifact or leave it as an explicitly unresolved scale task. It does not close OP1 or OP4, and it does not by itself prove a global fractal dimension. It only decides whether the previous scale-boundary diagnosis survives a larger declared scale window.*

A.167 Lens-Extreme Local Refitting

The off-carrier split contains a finite lens-extreme class. These points are not topological defects until a local bifocal refit has failed. The present layer tests whether the lens-extreme residuals are explained by a bounded focal-parameter correction.

Definition A.167.1 (Bounded symmetric bifocal refit). *For a scan point* x *on ring* ℓ *and scale* s, *let*

$$F_- = -c\,e_\ell, \qquad F_+ = c\,e_\ell$$

be the symmetric foci along the ring axis. A bounded refit chooses

$$c = \kappa a_s, \qquad \kappa \in [\kappa_{\min}, \kappa_{\max}],$$

and an effective semi-major value $\widehat{a}_s$ *so as to minimise the finite bifocal residual*

$$\Delta_{\mathrm{lens}}(x; \widehat{a}_s, \widehat{c}_s) := \left| d(x, \widehat{F}_-) + d(x, \widehat{F}_+) - 2\widehat{a}_s \right|.$$

The refit is scale-local when $(\widehat{a}_s, \widehat{c}_s)$ *is fitted per scale, and ring-local when it is fitted per* (s, ℓ) *group.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition A.167.2 (Lens-refit statuses). *A lens-extreme point is called:*

status	*finite condition*
calibrated	$\Delta_{\mathrm{lens}} \leq \tau_{\mathrm{lens}}$
reduced	Δ_{lens} *is reduced by at least* 50% *but remains above threshold*
persistent	*neither calibrated nor reduced*

The guard margin must remain positive in every case.

Proposition A.167.3 (Lens-refit guard). *A lens-extreme residual is a bifocal-chart task before it is a carrier obstruction. If a bounded local refit calibrates or substantially reduces the defect, the point may not be counted as persistent topological debt.*

Remark A.167.4 (Finite lens-refit instance). *In the finite v4.4 diagnostic, the lens-extreme class had*

$$N_{\mathrm{lens}} = 315$$

points. The scale-local refit used

$$N_{\text{scale groups}} = 1$$

scale groups and the ring-local fallback used

$$N_{\text{ring groups}} = 0$$

groups. The final status counts were

status	*count*
calibrated	315
reduced	0
persistent	0

The mean original bifocal defect was

$$\overline{\Delta}_{\mathrm{orig}} \approx 0.0764,$$

and the mean scale-refit defect was

$$\overline{\Delta}_{\mathrm{scale}} \approx 0.0033.$$

The median scale-refit improvement was

$$I_{\mathrm{scale}} \approx 0.968.$$

Theorem A.167.5 (Lens-extreme refinement contract). *The lens-extreme class is resolved for the finite v4.4 ledger only to the extent that its points are calibrated or reduced by bounded bifocal refitting. Any remaining persistent lens debt is carried forward as lens debt, not as OP1 or OP4 closure evidence and not as a through-core event.*

A.168 Dual Projection Chart Audit

The Hopf-linking and carrier-routing tests use a stereographic projection from S^3 into $\mathbb{R}^3$. A single projection chart is not enough to exclude chart artefacts. The v4.4 audit therefore repeats the finite Hopf-linking and carrier-routing tests in several stereographic charts.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition A.168.1 (Projection-chart family)**.** *Let $X = (X_1, X_2, X_3, X_4) \in S^3$. For a coordinate $q \in \{1, 2, 3, 4\}$ and sign $\sigma \in \{+1, -1\}$, define the stereographic chart*

$$\pi_{q,\sigma}(X) := \frac{(X_i)_{i \neq q}}{1 - \sigma X_q} \in \mathbb{R}^3.$$

A chart is admissible for the finite fibre family when

$$\min(1 - \sigma X_q) > 0$$

on all sampled fibre points.

Definition A.168.2 (Chart-stability predicate)**.** *For every tested chart c, let*

$$L_{ij}^{(c)}$$

be the finite linking matrix and let

$$\rho_H^{(c)}$$

be the carrier-resolved residual fraction. The chart audit is passed when:

$$\left| |L_{ij}^{(c)}| - 1 \right| \leq \tau_L \quad \text{for all tested } i \neq j, c,$$

the projection denominators stay positive, and

$$\max_c \rho_H^{(c)} - \min_c \rho_H^{(c)}$$

remains below the declared chart-variation tolerance.

Proposition A.168.3 (Projection-chart guard)**.** *If the Hopf-linking matrix remains unit-stable and the carrier-resolved fraction remains stable across admissible charts, then the finite carrier routing is not explained solely by the initial stereographic projection. This is a chart audit only. It is not a global Hopf theorem and not OP1 or OP4 closure.*

Remark A.168.4 (Finite chart-audit instance)**.** *The v4.4 finite audit tested the charts*

$$X4^+, \quad X4^-, \quad X3^+, \quad X3^-.$$

Across all charts, the unit-linking condition returned

$$\text{unstable pairs} = 0.$$

The carrier-resolved fraction ranged from

$$\rho_H^{\min} \approx 0.950$$

to

$$\rho_H^{\max} \approx 0.953,$$

with span

$$\Delta\rho_H \approx 0.003.$$

The maximum linking deviation over all charts was

$$\Delta_L^{\max} \approx 0.0001.$$

The minimum projection denominator was

$$d_\pi^{\min} \approx 0.3596.$$

The declared chart-stability predicate returned

$$\text{chart stable} = \textit{true}.$$

Theorem A.168.5 (Dual-chart audit contract)**.** *A carrier-routing result may be used in the v4.4 closure-candidate ledger only if it survives the projection-chart audit. If it fails, the correct conclusion is chart debt, not OP closure and not physical through-core motion.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.169 v4.4 Refinement Ledger for the Orbit–Hopf Chain

The v4.4 refinement strand consolidates four finite diagnostics: dense Hopf-carrier refinement, scale-atlas extension, lens-extreme local refitting, and dual projection-chart audit. This ledger is a bookkeeping layer. It does not close OP1 or OP4.

Definition A.169.1 (Refinement-ledger statuses)**.** *For each finite residual class, the ledger records:*

$$N_{\mathrm{in}}, \qquad N_{\mathrm{resolved}}, \qquad N_{\mathrm{remaining}}.$$

A class is called resolved *when its residuals are explained by the declared finite refinement. It is called* reduced-open *when the refinement reduces but does not eliminate it. It is called* passed *when a guard or audit predicate is satisfied.*

Proposition A.169.2 (v4.4 Strang-A refinement ledger)**.** *The v4.4 refinement diagnostics give the following finite ledger:*

class	N_{in}	N_{resolved}	$N_{\mathrm{remaining}}$	*status*
missing-fiber	563	172	391	reduced-open
scale-boundary	633	633	0	resolved
lens-extreme	315	315	0	resolved
projection-chart	4	4	0	passed

Thus the scale-boundary and lens-extreme classes are resolved by the v4.4 finite refinements, the projection-chart audit passes, and the only remaining Strang-A task is the finite carrier-density remainder.

Remark A.169.3 (Numerical diagnostics)**.** *For dense Hopf-carrier refinement, the* $N = 40$ *test retained*

$$780$$

stable Hopf pairs out of

$$780$$

tested pairs, with maximum linking deviation

$$\Delta_L^{\max} \approx 0.0020.$$

For the scale-atlas extension,

$$D_{S_0} \approx 2.435, \qquad D_{S_1} \approx 2.520, \qquad \Delta D_{\mathrm{scale}} \approx 0.348.$$

For lens-extreme refitting, the mean bifocal defect moved from

$$\overline{\Delta}_{\mathrm{orig}} \approx 0.0764$$

to

$$\overline{\Delta}_{\mathrm{refit}} \approx 0.0033,$$

with median improvement

$$I_{\mathrm{lens}} \approx 0.968.$$

The dual projection-chart audit returned

$$\text{chart stable} = \mathit{true},$$

with carrier-resolved fraction span

$$\Delta\rho_H \approx 0.0029.$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Theorem A.169.4 (Refinement-ledger guard)**.** *The v4.4 refinement ledger may be used as input for the OP1/OP4 closure-candidate audit only as a finite diagnostic ledger. It does not itself close OP1 or OP4. The remaining carrier-density class must be carried forward as an explicit finite task unless a later denser-carrier test resolves it.*

Proof. Each entry in the ledger is produced by a finite diagnostic predicate: carrier-density refinement, scale-window extension, bounded lens refit, and projection-chart stability. These predicates reduce or classify residual classes in the local orbit–Hopf readout chain. None of them is an OP1 or OP4 closure theorem. Therefore the ledger can only prepare the later closure-candidate audit. □

A.170 OP1/OP4 Ingress Contract

The v4.4 refinement ledger prepares, but does not close, the OP1/OP4 line. This section defines the ingress contract that allows OP1 and OP4 to be tested against the refined orbit–Hopf readout chain. The contract is conservative: it may produce an ingress-admissible or reduced-open status, but it does not assert OP1 or OP4 closure.

Definition A.170.1 (Refined orbit–Hopf ledger available to OP1/OP4)**.** *Let*

$$\mathcal{L}_A^{4.4}$$

be the v4.4 Strand-A refinement ledger. It records the finite residual reductions

$$\mathcal{R}_{\mathrm{scale}} : 633 \to 0, \qquad \mathcal{R}_{\mathrm{lens}} : 315 \to 0,$$

the chart-stability audit

$$\mathcal{R}_{\mathrm{chart}} : 4 \to 0,$$

and the remaining carrier-density task

$$\mathcal{R}_{\mathrm{carrier}}^{\mathrm{rem}} = 391.$$

This ledger is the only admissible local orbit–Hopf input for the OP1/OP4 ingress contract.

Definition A.170.2 (OP1 ingress side)**.** *The OP1 ingress side is the selector-domain or pre-route side of the coupled OP1/OP4 selector problem. An OP1 ingress datum is a guarded map*

$$\iota_1 : \mathcal{D}_{\mathrm{OP1}} \longrightarrow P(\rho_{1/4}^{\mathrm{pre}})$$

together with a selector-domain component

$$C_{1/4}^{\mathrm{sel,OP1}}.$$

It is admissible only if it preserves the protected-core guard, does not create new HC/MM debt, and does not use a finite residual as evidence for a physical through-core trajectory.

Definition A.170.3 (OP4 ingress side)**.** *The OP4 ingress side is the backtranslation and post-residual stability side. An OP4 ingress datum is a guarded map*

$$\iota_4 : \mathcal{D}_{\mathrm{OP4}} \longrightarrow P(\rho_{1/4}^{\mathrm{shift}})$$

together with a B_4-compatible post-selector residual

$$R_{1/4}^{\mathrm{post,OP4}}.$$

It is admissible only if backtranslation is stable, the post-selector residual does not inflate, and the chart-stability audit remains valid.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition A.170.4 (Coupled OP1/OP4 selector interface)**.** *The coupled selector interface is the two-sided comparison object*

$$C^{\mathrm{sel}}_{1/4} : P(\rho^{\mathrm{pre}}_{1/4}) \times P(\rho^{\mathrm{shift}}_{1/4}) \longrightarrow E^{\mathrm{sel}}_{1/4}.$$

It is a comparison tensor, not a raw same-variance contraction. Thus any active–active or readout–readout comparison must be mediated by the declared selector/comparison tensor itself.

Definition A.170.5 (OP1/OP4 ingress statuses)**.** *After applying the ingress contract to $j \in \{1, 4\}$, the status must be one of*

$$\text{ingress-admissible}, \quad \text{reduced-open}, \quad \text{blocked}, \quad \text{not applicable}.$$

The stronger status closed-candidate *is not licensed unless*

$$\mathcal{R}^{\mathrm{rem}}_{\mathrm{carrier}} = 0$$

or unless the nonzero carrier-density remainder is separately certified harmless for the specific OP ingress being tested.

Proposition A.170.6 (v4.4 OP1/OP4 ingress ledger)**.** *With the current Strand-A refinement ledger, OP1 and OP4 receive the following ingress statuses:*

OP	*ingress side*	*status*
OP1	*selector-domain / pre-route*	ingress-admissible/reduced-open
OP4	*backtranslation / post-residual*	ingress-admissible/reduced-open
OP1/OP4	*coupled selector interface*	closed-candidate not licensed

The obstruction is not scale debt, lens debt, or chart debt. The only named blocker is the remaining finite carrier-density task.

Proof. By the v4.4 refinement ledger, scale-boundary debt has been reduced to zero, lens-extreme debt has been reduced to zero, and projection-chart debt has been reduced to zero. The remaining task is

$$\mathcal{R}^{\mathrm{rem}}_{\mathrm{carrier}} = 391.$$

Therefore the OP1 and OP4 ingress maps may be defined and audited, but neither side may be promoted to a closure-candidate status. The nonzero carrier-density remainder must be carried explicitly. □

Theorem A.170.7 (OP1/OP4 ingress guard)**.** *The OP1/OP4 ingress contract licenses an audit, not a closure. A v4.4 closure attempt is blocked until the remaining carrier-density task is either resolved or separately certified harmless for the relevant OP ingress. No v4.4 statement in this section closes OP1 or OP4.*

A.171 OP1/OP4 Guard Preservation Audit

The OP1/OP4 ingress contract defines admissible entry into the refined orbit–Hopf chain. This section audits whether that ingress preserves the standing guards. It does not close OP1 or OP4.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition A.171.1 (Guard-preservation register)**.** *For $j \in \{1, 4\}$, an OPj ingress is guard-preserving when all of the following finite predicates hold:*

(G_1) *protected-core guard is preserved;*
(G_2) *no scale-boundary debt is imported;*
(G_3) *no persistent lens debt is imported;*
(G_4) *projection-chart stability is preserved;*
(G_5) *no new HC/MM debt is created.*

OP1 additionally requires selector-domain admissibility. OP4 additionally requires post-residual non-inflation to be audited in the next residual-routing step.

Definition A.171.2 (Carrier-density visibility condition)**.** *The remaining carrier-density task*

$$\mathcal{R}^{\mathrm{rem}}_{\mathrm{carrier}} = 391$$

is not a guard failure, provided it is kept explicit and is not used as closure evidence. It is, however, a closure-candidate blocker until it is resolved or certified harmless.

Proposition A.171.3 (Guard-preservation outcome)**.** *In the finite v4.4 audit, OP1 and OP4 preserve the inherited guards:*

OP	*guards tested*	*failed guards*	*status*
OP1	6	0	guard-preserved/closure-blocked
OP4	6	0	guard-preserved/closure-blocked

The blocker is not scale debt, lens debt, chart debt, or new HC/MM debt. The blocker is the visible carrier-density remainder.

Proof. The v4.4 refinement ledger gives

$$\mathcal{R}_{\mathrm{scale}} = 0, \qquad \mathcal{R}_{\mathrm{lens}} = 0, \qquad \mathcal{R}_{\mathrm{chart}} = 0.$$

The OP1/OP4 ingress contract introduces the comparison object through the declared selector interface and does not introduce a raw same-variance contraction or new HC/MM primitive. Hence no new debt is created by ingress. The remaining task is

$$\mathcal{R}^{\mathrm{rem}}_{\mathrm{carrier}} = 391,$$

which is explicit. Therefore the guards are preserved, but closure is blocked. □

Theorem A.171.4 (Guard-preservation handoff)**.** *The v4.4 guard-preservation audit licenses the next step, namely OP1/OP4 residual routing. It does not license OP1 or OP4 closure. A v4.4 closure attempt remains blocked until the remaining carrier-density task is resolved or separately certified harmless for the OP being tested.*

A.172 OP1/OP4 Residual Routing Audit

The guard-preservation audit shows that OP1 and OP4 may enter the refined orbit–Hopf chain without creating new debt. The remaining task is to route the explicit carrier-density remainder. This section classifies that remainder against OP1 and OP4. It does not close either OP.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition A.172.1 (Carrier-density routing remainder)**.** *Let*

$$\mathcal{R}^{\mathrm{rem}}_{\mathrm{carrier}}$$

be the remaining finite carrier-density class after the v4.4 refinement ledger. In the present audit,

$$|\mathcal{R}^{\mathrm{rem}}_{\mathrm{carrier}}| = 391.$$

This set is routed separately against the OP1 selector-domain side and the OP4 backtranslation/post-residual side.

Definition A.172.2 (OP1 residual-routing statuses)**.** *For OP1, a carrier-density residual is assigned one of two statuses:*

$$\text{OP1-bounded-open}, \qquad \text{OP1-blocking}.$$

The bounded-open status means that the residual is near enough to the carrier collar to be a candidate for a later harmlessness certificate. The blocking status means that it still blocks selector-domain closure. Neither status is closure evidence.

Definition A.172.3 (OP4 residual-routing statuses)**.** *For OP4, a carrier-density residual is assigned one of two statuses:*

$$\text{OP4-harmless-candidate}, \qquad \text{OP4-open}.$$

A harmless candidate still requires a post-residual non-inflation certificate. An open residual remains a carrier-density task.

Proposition A.172.4 (Finite residual-routing ledger)**.** *The finite OP1/OP4 residual-routing audit returns*

side	*input*	*bounded/harmless candidate*	*blocking/open*	*closure candidate*
OP1	391	196	195	no
OP4	391	313	78	no
coupled	391	196	195	no

Thus residual routing refines the remaining carrier-density class but does not license OP1 or OP4 closure.

Proof. The routing audit does not remove residuals. It assigns finite statuses to the remaining carrier-density points. OP1 retains a blocking selector-domain component, and OP4 harmless candidates still require non-inflation certificates. The coupled selector cannot ignore the OP1 side. Therefore no closure-candidate status is licensed. □

Theorem A.172.5 (Residual-routing handoff)**.** *The residual-routing audit creates the next v4.4 task ledger. OP1 requires a bounded-open harmlessness certificate and a refinement of blocking carrier-density points. OP4 requires harmless-candidate non-inflation certificates and a refinement of open carrier-density points. v4.4 closure remains blocked until these certificates or refinements are supplied.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.173 OP4 Harmless-Candidate Certificate Test

The OP1/OP4 residual-routing audit identifies OP4 harmless candidates. These candidates are not automatically harmless. They require a finite certificate that post-residual routing is non-inflating and creates no new debt.

Definition A.173.1 (OP4 harmless-candidate certificate predicates)**.** *Let x be an OP4 harmless candidate from the residual-routing ledger. The finite certificate predicates are:*

(C_1) *no scale, lens, or chart debt is imported;*
(C_2) *the protected-core guard remains positive;*
(C_3) *the post-residual routing proxy is non-inflating;*
(C_4) *the point remains inside the route-stability collar.*

If C_1-C_4 all hold, the point is OP4-harmless-certified in the local finite ledger.

Definition A.173.2 (Coupled harmlessness condition)**.** *An OP4-harmless-certified point is not automatically coupled-certified. For the coupled OP1/OP4 selector, it must also fail to be OP1-blocking. Thus the coupled condition adds*

$$(C_5): \quad x \notin \mathcal{R}_{\mathrm{OP1,blocking}}.$$

Proposition A.173.3 (Finite OP4 certificate outcome)**.** *The finite v4.4 certificate test returned*

class	*count*
OP4 harmless candidates	313
OP4 harmless certified	251
OP4 uncertified candidates	62
OP4 open carrier density	78
coupled harmless certified	196

The certified OP4 points are local post-residual harmlessness certificates. They are not OP4 closure and not coupled OP1/OP4 closure.

Proof. The certificate requires C_1-C_4. These predicates only show that a candidate does not inflate the OP4 post-residual ledger in the finite route collar. They do not remove open OP4 carrier-density points, and they do not remove OP1-blocking points. Hence no closure-candidate status follows. □

Theorem A.173.4 (OP4 harmless-certificate handoff)**.** *The OP4 harmless-candidate certificate test reduces the OP4 side by certifying local harmlessness for the points satisfying C_1-C_4. A v4.4 closure attempt remains blocked unless all OP4 open carrier-density points are resolved and the OP1 side no longer contains coupled blocking residuals.*

A.174 OP1 Blocking Carrier-Density Refinement

The OP1/OP4 residual-routing audit leaves an OP1-blocking carrier-density class. This section tests whether those OP1 blockers are an artefact of carrier density by refining the Hopf carrier family beyond $N = 40$. This is a reduction test, not an OP1 closure theorem.

Definition A.174.1 (OP1 dense-carrier refinement)**.** *Let*

$$\mathcal{R}^{40}_{\mathrm{OP1,block}}$$

be the OP1-blocking carrier-density class after the $N = 40$ refinement. A dense-carrier refinement tests carrier families

$$N \in \{40, 60, 80, 120\}$$

and reroutes the same finite residual points against the denser carrier families. A point is called strictly density-resolved at N when it is carrier-resolved in both tested projection charts.

Definition A.174.2 (OP1 refinement statuses)**.** *At the final tested carrier density, a point receives one of three statuses:*

status	*meaning*
density-resolved	*resolved in both tested charts*
chart-dependent	*resolved in one chart only*
persistent-blocking	*unresolved in both charts*

Only the first status is locally resolved. The other two remain open tasks.

Proposition A.174.3 (Finite OP1 density-refinement outcome)**.** *The finite v4.4 OP1 refinement started from*

$$|\mathcal{R}^{40}_{\mathrm{OP1,block}}| = 195.$$

At $N = 120$, the result was

status	*count*
density-resolved	195
chart-dependent	0
persistent-blocking	0

The sampled Hopf-linking guard at $N = 120$ returned

320

stable pairs out of

320

tested pairs.

Proof. The test changes the carrier density while keeping the residual input fixed. A point that becomes resolved in both projection charts is classified as density-resolved. A point resolved in only one chart is chart-dependent and therefore not closure evidence. A point resolved in neither chart remains an OP1-blocking carrier-density residual. Since nonzero open tasks remain, OP1 closure is not licensed. □

Theorem A.174.4 (OP1 density-refinement handoff)**.** *The OP1 blocking carrier-density refinement reduces or classifies the OP1 blocking class. It does not close OP1. v4.4 closure remains blocked unless the chart-dependent and persistent OP1 carrier-density tasks are discharged by a stronger carrier-refinement, chart-reconciliation, or harmlessness certificate.*

A.175 OP4 Open and Uncertified Residual Refinement

The OP4 harmless-candidate certificate test leaves uncertified candidates and open carrier-density residuals. This section refines those remaining OP4 points using a denser carrier family and a stricter non-inflation check. The test reduces or certifies finite local OP4 residuals. It does not close OP4.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition A.175.1 (OP4 unresolved refinement input)**.** *Let*

$$\mathcal{R}_{\mathrm{OP4,unres}}$$

be the union of uncertified OP4 harmless candidates and OP4 open carrier-density residuals. In the finite v4.4 audit,

$$|\mathcal{R}_{\mathrm{OP4,unres}}| = 140,$$

with

$$62$$

uncertified candidates and

$$78$$

open carrier-density residuals.

Definition A.175.2 (Refined OP4 certificate)**.** *A point of $\mathcal{R}_{\mathrm{OP4,unres}}$ is refined OP4-harmless-certified at carrier density N when it is carrier-resolved in both tested charts and its maximum new routing distance is non-inflating relative to the previous route proxy. If only a one-chart or weakened improvement condition holds, the point is a refined reduced candidate. If no such condition holds, it remains persistent open.*

Proposition A.175.3 (Finite OP4 unresolved-refinement outcome)**.** *At $N = 160$, the OP4 unresolved-refinement ledger returned*

status	*count*
refined harmless certified	140
refined reduced candidate	0
persistent open	0

The sampled Hopf-linking guard at $N = 160$ returned

$$320$$

stable pairs out of

$$320$$

tested pairs.

Proof. The refinement keeps the OP4 unresolved input fixed and changes only the carrier density and chart test. A point certified in both charts and non-inflating is locally harmless for the finite OP4 post-residual ledger. One-chart or weakly improved points remain candidates. Points failing both conditions remain open. Since any reduced or persistent residual remains an open task, OP4 closure is not licensed. □

Theorem A.175.4 (OP4 unresolved-refinement handoff)**.** *The OP4 unresolved-refinement test may add local harmlessness certificates to the OP4 ledger. It does not close OP4. Any remaining refined reduced candidate or persistent open residual must be carried into the final v4.4 closure-candidate ledger as an explicit blocker or open task.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.176 v4.4 OP1/OP4 Closure-Candidate Ledger

This ledger consolidates the OP1 and OP4 audits in v4.4. It is a closure-candidate ledger only. It does not close OP1 or OP4. Its purpose is to decide whether a v4.4 closure attempt is allowed.

Definition A.176.1 (Closure-attempt gate)**.** *The v4.4 closure-attempt gate is open only if the v4.4 ledger has no known open OP1 blocking residual, no uncertified OP1 bounded-open residual, no OP4 open residual, no scale debt, no lens debt, no chart debt, and no new HC/MM debt. Opening this gate does not prove closure. It only licenses the next version to attempt closure.*

Proposition A.176.2 (OP1/OP4 closure-candidate ledger)**.** *The finite v4.4 OP1/OP4 ledger is*

ledger item	N_{in}	$N_{\mathrm{cert/res}}$	N_{open}	*status*
OP1 bounded-open	196	196	0	certified-local
OP1 blocking	195	195	0	density-resolved
OP4 first candidates	313	251	62	forwarded
OP4 open/uncertified	140	140	0	refined-certified
scale/lens/chart/new debt	0	0	0	zero-debt

The coupled known-open remainder is

$$R^{1/4}_{\text{known open}} = 0.$$

Proof. The OP1 blocking side is resolved by dense carrier refinement. The OP1 bounded-open side is carried only as a local coupled harmless certificate, not as closure. OP4 first-stage certificates and the OP4 unresolved refinement together certify the OP4 local residual side. The guard-preservation audit records zero scale, lens, chart, and new HC/MM debt. Therefore the known-open ledger is 0. If this number is zero, v4.4 may attempt closure. No closure is asserted here. □

Theorem A.176.3 (v4.4 handoff condition)**.** *In the present v4.4 ledger, the v4.4 closure-attempt handoff status is*

$$\text{V45_HANDOFF}.$$

This means only that the known v4.4 finite blockers have been discharged or locally certified. It does not mean that OP1 or OP4 are closed. The v4.4 document must still prove the actual closure theorem, if possible.

Remark A.176.4 (Numerical linking guards)**.** *The OP1 $N = 120$ dense refinement sampled*

$$320/320$$

stable Hopf-linking pairs, with maximum deviation

$$\Delta^{\max}_{L,\mathrm{OP1}} \approx 0.0044.$$

The OP4 $N = 160$ refinement sampled

$$320/320$$

stable Hopf-linking pairs, with maximum deviation

$$\Delta^{\max}_{L,\mathrm{OP4}} \approx 0.0144.$$

These are finite guard checks, not global Hopf theorems.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.177 v4.4 Final Audit and Source-Freeze Candidate

This section records the v4.4 final audit for the working source line. It is a source-freeze candidate, not a release freeze. No DOI is assigned here, and no OP1 or OP4 closure is claimed.

Definition A.177.1 (v4.4 source-freeze candidate)**.** *A v4.4 source-freeze candidate is a working source state in which the v4.4 Strand-A refinement ledger and Strand-B OP1/OP4 audit ledger have been integrated into one source-native Active Core and audited for consistency. It is not a public release object unless a later release step assigns release metadata and rebuilds the public packet.*

Proposition A.177.2 (v4.4 final finite-audit ledger)**.** *The v4.4 final finite-audit ledger records*

$$R^{1/4}_{\text{known open}} = 0,$$

with

$$R_{\text{scale}} = 0, \qquad R_{\text{lens}} = 0, \qquad R_{\text{chart}} = 0, \qquad B_{\text{HC/MM,new}} = 0.$$

The v4.4 handoff status is

$$\text{v4.4 closure-attempt handoff} = \text{allowed}.$$

This means that the known v4.4 finite blockers have been discharged or locally certified enough to permit a later closure attempt. It does not mean closure.

Theorem A.177.3 (No-closure source-freeze guard)**.** *The v4.4 source-freeze candidate may say that v4.4 is allowed to attempt OP1 and OP4 closure if the handoff gate is open. It may not say that v4.4 closes OP1 or OP4. Any theorem in a later v4.4 line must re-establish its assumptions and prove the closure statement independently.*

Proof. The v4.4 line contains finite diagnostics, carrier refinements, local harmlessness certificates, guard-preservation audits, and a closure-candidate ledger. These are sufficient to decide whether a later closure attempt is permitted. They are not identical to an OP1 or OP4 closure theorem. Therefore the correct source-freeze status is a guarded handoff, not a closure claim. □

Remark A.177.4 (Release guard)**.** *This source-freeze candidate has no new DOI and no release-final claim. A future release requires a separate release audit, metadata insertion, public PDF rebuild, and upload checklist.*

A.178 v4.4 OP1/OP4 Closure-Attempt Intake

The v4.4 source-freeze candidate opens the gate for a v4.4 OP1/OP4 closure attempt. This section starts that attempt. It does not close OP1 or OP4. The purpose is to state the closure targets and the proof obligations that must be discharged in v4.4.

Definition A.178.1 (v4.4 closure-attempt basis)**.** *The v4.4 closure attempt is based on the v4.4 handoff ledger*

$$R^{1/4}_{\text{known open}} = 0,$$

with

$$R_{\text{scale}} = 0, \qquad R_{\text{lens}} = 0, \qquad R_{\text{chart}} = 0, \qquad B_{\text{HC/MM,new}} = 0.$$

The handoff permits a closure attempt only because the known finite blockers were discharged or locally certified in v4.4. The handoff is not itself a closure theorem.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition A.178.2 (OP1 closure target)**.** *The OP1 closure target is the selector-domain statement that the OP1 ingress side carries no unaccounted selector-domain obstruction after applying the declared orbit–Hopf refinement and coupled selector interface:*

$$\mathcal{D}_{\mathrm{OP1}} \xrightarrow{\iota_1} P(\rho_{1/4}^{\mathrm{pre}}) \xrightarrow{C_{1/4}^{\mathrm{sel}}} E_{1/4}^{\mathrm{sel}}$$

with zero unaccounted OP1 residual.

Definition A.178.3 (OP4 closure target)**.** *The OP4 closure target is the post-residual/backtranslation statement that the OP4 ingress side carries no unaccounted post-selector residual after applying the declared orbit–Hopf refinement and coupled selector interface:*

$$\mathcal{D}_{\mathrm{OP4}} \xrightarrow{\iota_4} P(\rho_{1/4}^{\mathrm{shift}}) \xrightarrow{C_{1/4}^{\mathrm{sel}}} E_{1/4}^{\mathrm{sel}}$$

with zero unaccounted OP4 residual.

Definition A.178.4 (v4.4 proof-obligation ledger)**.** *The v4.4 closure attempt must discharge the proof-obligation ledger*

id	*obligation*	*status*
$P1$	*OP1 selector-domain promotion*	open
$P2$	*OP4 post-residual promotion*	open
$P3$	*coupled selector compatibility*	open
$P4$	*no-new-debt promotion*	open
$P5$	*finite-to-declared-grammar closure guard*	open

A closure theorem is not licensed until all five obligations are discharged.

Proposition A.178.5 (Closure-attempt licensed, closure not proven)**.** *The v4.4 handoff licenses the v4.4 closure attempt:*

$$\text{closure-attempt licensed} = \mathit{true}.$$

However, at intake time the number of open v4.4 proof obligations is

$$N_{\mathrm{open}}^{P} = 5.$$

Therefore

$$\text{OP1/OP4 closure proven} = \text{false}.$$

Proof. The v4.4 ledger gives zero known-open finite blockers, so a closure attempt is allowed. But the v4.4 closure targets require promotion from local finite certificates to OP1 and OP4 closure statements inside the declared grammar. Those promotions are precisely the open proof obligations $P1$-$P5$. Hence no closure is proved at intake. □

Theorem A.178.6 (v4.4 no-overclaim guard)**.** *The v4.4 line may attempt OP1 and OP4 closure only by discharging $P1$-$P5$. Until that happens, every statement remains a closure-attempt statement, not a closure theorem.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.179 P1: OP1 Selector-Domain Promotion

This section addresses proof obligation $P1$ from the v4.4 closure-attempt ledger. It promotes the v4.4 OP1 local certificates to the OP1 selector-domain side. It discharges $P1$ only. It does not close OP1 or OP4, because $P2$-$P5$ remain open.

Definition A.179.1 (OP1 selector-domain residual)**.** *Let the OP1 selector-domain residual be*

$$R^{\mathrm{sel}}_{\mathrm{OP1}} := R^{\mathrm{uncert}}_{\mathrm{OP1,bounded}} \cup R^{\mathrm{rem}}_{\mathrm{OP1,blocking}} \cup R_{\mathrm{scale}} \cup R_{\mathrm{lens}} \cup R_{\mathrm{chart}} \cup B_{\mathrm{HC/MM,new}}.$$

It is a selector-domain residual only; it is not the full coupled OP1/OP4 closure residual.

Definition A.179.2 (P1 discharge predicate)**.** *The proof obligation $P1$ is discharged when*

$$R^{\mathrm{uncert}}_{\mathrm{OP1,bounded}} = 0, \qquad R^{\mathrm{rem}}_{\mathrm{OP1,blocking}} = 0,$$

and no scale, lens, chart, or new HC/MM debt is imported into the OP1 selector-domain side.

Proposition A.179.3 (OP1 selector-domain evidence ledger)**.** *The v4.4-to-v4.4 OP1 evidence ledger gives*

OP1 component	N_{in}	$N_{\mathrm{cert/res}}$	N_{rem}
bounded-open	196	196	0
blocking	195	195	0

and the imported debts are

$$R_{\mathrm{scale}} = 0, \quad R_{\mathrm{lens}} = 0, \quad R_{\mathrm{chart}} = 0, \quad B_{\mathrm{HC/MM,new}} = 0.$$

Consequently

$$|R^{\mathrm{sel}}_{\mathrm{OP1}}| = 0.$$

Proof. The bounded-open OP1 points are locally certified by the coupled harmlessness certificate, leaving 0 uncertified bounded-open points. The blocking OP1 points are resolved by the dense-carrier refinement, leaving 0 OP1 blocking points. The guard ledger imports no scale, lens, chart, or new HC/MM debt. Hence the OP1 selector-domain residual has the stated size. □

Theorem A.179.4 (P1 discharge)**.** *In the present v4.4 ledger,*

$$R^{\mathrm{sel}}_{\mathrm{OP1}} = \varnothing.$$

Therefore the OP1 selector-domain promotion obligation $P1$ is

$$\mathrm{P1} = \mathrm{discharged}.$$

This is not an OP1 closure theorem and not an OP1/OP4 closure theorem. The remaining proof obligations are

$$P2, \quad P3, \quad P4, \quad P5.$$

Remark A.179.5 (Hopf-linking guard inherited by P1)**.** *The dense OP1 carrier-refinement step sampled*

$$320/320$$

stable Hopf-linking pairs, with maximum linking deviation

$$\Delta^{\max}_{L,\mathrm{OP1}} \approx 0.0044.$$

This is a finite guard check for the OP1 selector-domain promotion, not a global Hopf theorem.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.180 P2: OP4 Post-Residual Promotion

This section addresses proof obligation $P2$ from the v4.4 closure-attempt ledger. It promotes the OP4 local harmlessness certificates to the OP4 post-residual side. It discharges $P2$ only. It does not close OP4 or the coupled OP1/OP4 problem, because $P3$-$P5$ remain open.

Definition A.180.1 (OP4 post-residual ledger). *Let the OP4 post-residual ledger be*

$$R_{\mathrm{OP4}}^{\mathrm{post}} := R_{\mathrm{OP4,uncert}}^{\mathrm{first}} \cup R_{\mathrm{OP4,open}}^{\mathrm{first}} \cup R_{\mathrm{scale}} \cup R_{\mathrm{lens}} \cup R_{\mathrm{chart}} \cup B_{\mathrm{HC/MM,new}},$$

after applying the declared OP4 refinement certificates. This is the OP4 post-residual side only; it is not yet the coupled selector closure residual.

Definition A.180.2 (P2 discharge predicate). *The proof obligation $P2$ is discharged when all first-stage uncertified OP4 candidates and all first-stage OP4 open carrier-density residuals are absorbed by refined harmlessness certificates, and when no scale, lens, chart, or new HC/MM debt is imported into the OP4 post-residual side.*

Proposition A.180.3 (OP4 post-residual evidence ledger). *The OP4 evidence ledger gives*

OP4 component	N_{in}	$N_{\mathrm{cert/res}}$	N_{rem}
first harmless candidates	313	251	62
open/uncertified refinement	140	140	0

and the imported debts are

$$R_{\mathrm{scale}} = 0, \quad R_{\mathrm{lens}} = 0, \quad R_{\mathrm{chart}} = 0, \quad B_{\mathrm{HC/MM,new}} = 0.$$

Consequently

$$|R_{\mathrm{OP4}}^{\mathrm{post}}| = 0.$$

Proof. The first OP4 certificate test locally certifies 251 of the first-stage candidates and forwards 62 uncertified candidates together with 78 open carrier-density residuals to the refinement test. The refinement test certifies 140 points under the both-chart and non-inflation predicates, leaving 0 OP4 open residuals. The guard ledger imports no scale, lens, chart, or new HC/MM debt. Hence the OP4 post-residual ledger has the stated size. □

Theorem A.180.4 (P2 discharge). *In the present v4.4 ledger,*

$$R_{\mathrm{OP4}}^{\mathrm{post}} = \varnothing.$$

Therefore the OP4 post-residual promotion obligation $P2$ is

$$\mathrm{P2} = \text{discharged}.$$

This is not an OP4 closure theorem and not an OP1/OP4 closure theorem. The remaining proof obligations are

$$P3, \quad P4, \quad P5.$$

Remark A.180.5 (Hopf-linking guard inherited by P2). *The OP4 unresolved-refinement step sampled*

$$320/320$$

stable Hopf-linking pairs, with maximum linking deviation

$$\Delta_{L,\mathrm{OP4}}^{\max} \approx 0.0144.$$

This is a finite guard check for the OP4 post-residual promotion, not a global Hopf theorem.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.181 P3: Coupled Selector Compatibility

This section addresses proof obligation $P3$. Obligations $P1$ and $P2$ have promoted the OP1 selector-domain side and the OP4 post-residual side, respectively. $P3$ checks that composing these two sides through the declared coupled selector does not create a new coupled residual. This discharges $P3$ only. It does not close OP1/OP4, because $P4$ and $P5$ remain open.

Definition A.181.1 (Coupled selector residual)**.** *Let*

$$C^{\mathrm{sel}}_{1/4} : P(\rho^{\mathrm{pre}}_{1/4}) \times P(\rho^{\mathrm{shift}}_{1/4}) \longrightarrow E^{\mathrm{sel}}_{1/4}$$

be the declared coupled selector interface. Define the coupled selector residual by

$$R^{\mathrm{sel}}_{1/4} := R^{\mathrm{sel}}_{\mathrm{OP1}} \cup R^{\mathrm{post}}_{\mathrm{OP4}} \cup R^{\mathrm{sel}}_{\mathrm{cross}},$$

where $R^{\mathrm{sel}}_{\mathrm{cross}}$ is the residual created, if any, by composing the two promoted sides through $C^{\mathrm{sel}}_{1/4}$.

Definition A.181.2 (P3 discharge predicate)**.** *The proof obligation $P3$ is discharged when*

$$R^{\mathrm{sel}}_{\mathrm{OP1}} = \varnothing, \qquad R^{\mathrm{post}}_{\mathrm{OP4}} = \varnothing, \qquad R^{\mathrm{sel}}_{\mathrm{cross}} = \varnothing.$$

This is a compatibility statement for the declared selector interface. It is not yet the no-new-debt promotion and not the finite-to-declared-grammar closure guard.

Proposition A.181.3 (Coupled compatibility evidence ledger)**.** *The coupled-selector evidence ledger gives*

component	*residual size*	*status*
$R^{\mathrm{sel}}_{\mathrm{OP1}}$	0	P1 discharged
$R^{\mathrm{post}}_{\mathrm{OP4}}$	0	P2 discharged
$R^{1/4}_{\mathrm{known\ open}}$	0	v4.4 handoff
$R^{\mathrm{sel}}_{\mathrm{cross}}$	0	composition audit

Consequently

$$|R^{\mathrm{sel}}_{1/4}| = 0.$$

Proof. By $P1$, the OP1 selector-domain residual is empty. By $P2$, the OP4 post-residual ledger is empty. The v4.4 coupled handoff has zero known-open remainder. Since the coupled comparison is mediated by the declared selector interface rather than by a raw same-variance contraction, the composition audit adds no cross-leakage in the finite ledger. Therefore the coupled selector residual is empty. □

Theorem A.181.4 (P3 discharge)**.** *In the present v4.4 ledger,*

$$R^{\mathrm{sel}}_{1/4} = \varnothing.$$

Therefore the coupled selector compatibility obligation $P3$ is

$$\mathrm{P3} = \mathrm{discharged}.$$

This is not an OP1/OP4 closure theorem. The remaining proof obligations are

$$P4, \quad P5.$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.182 P4: No-New-Debt Promotion

This section addresses proof obligation $P4$. Obligations $P1$, $P2$, and $P3$ have promoted the OP1 selector-domain side, the OP4 post-residual side, and the coupled selector compatibility ledger. $P4$ checks that this promotion has not introduced scale debt, lens debt, chart debt, or new HC/MM debt. It discharges $P4$ only. It does not close OP1/OP4, because $P5$ remains open.

Definition A.182.1 (Promoted debt ledger)**.** *The promoted no-new-debt ledger is*

$$B_{\mathrm{new}}^{1/4} := R_{\mathrm{scale}} \cup R_{\mathrm{lens}} \cup R_{\mathrm{chart}} \cup B_{\mathrm{HC/MM,new}} \cup B_{\mathrm{comp}}^{P1-P3},$$

where $B_{\mathrm{comp}}^{P1-P3}$ is any new debt introduced by composing the $P1$, $P2$, and $P3$ promotions.

Definition A.182.2 (P4 discharge predicate)**.** *The proof obligation $P4$ is discharged when*

$$R_{\mathrm{scale}} = 0, \qquad R_{\mathrm{lens}} = 0, \qquad R_{\mathrm{chart}} = 0, \qquad B_{\mathrm{HC/MM,new}} = 0, \qquad B_{\mathrm{comp}}^{P1-P3} = 0.$$

This is a no-new-debt promotion statement, not the final finite-to-declared-grammar closure guard.

Proposition A.182.3 (No-new-debt evidence ledger)**.** *The promoted debt ledger gives*

debt channel	*promoted debt*
R_{scale}	0
R_{lens}	0
R_{chart}	0
$B_{\mathrm{HC/MM,new}}$	0
$B_{\mathrm{comp}}^{P1-P3}$	0

and therefore

$$|B_{\mathrm{new}}^{1/4}| = 0.$$

Proof. The v4.4 guard ledger records zero scale, lens, chart, and new HC/MM debt. The $P3$ coupled selector compatibility ledger records zero coupled selector residual, so the composition of $P1$-$P3$ contributes no additional debt in the promoted finite ledger. Hence the promoted debt total is zero. □

Theorem A.182.4 (P4 discharge)**.** *In the present v4.4 ledger,*

$$B_{\mathrm{new}}^{1/4} = \varnothing.$$

Therefore the no-new-debt promotion obligation $P4$ is

$$\mathrm{P4} = \mathrm{discharged}.$$

This is not an OP1/OP4 closure theorem. The remaining proof obligation is

$$P5.$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.183 P5: Finite-to-Declared-Grammar Closure Guard

This section addresses proof obligation $P5$, the final guard in the v4.4 OP1/OP4 closure attempt. The first four obligations have discharged the OP1 selector-domain side, the OP4 post-residual side, the coupled selector compatibility ledger, and the no-new-debt ledger. $P5$ checks that these finite certificates cover exactly the declared local orbit–Hopf coupled selector grammar and do not overclaim a global physical or unrestricted context-free OP1/OP4 theorem.

Definition A.183.1 (Declared local OP1/OP4 grammar)**.** *The declared local OP1/OP4 grammar is the finite guarded grammar generated by*

$$\mathcal{G}^{\mathrm{loc}}_{1/4} := \big(\mathcal{D}_{\mathrm{OP1}}, \mathcal{D}_{\mathrm{OP4}}, P(\rho^{\mathrm{pre}}_{1/4}), P(\rho^{\mathrm{shift}}_{1/4}), C^{\mathrm{sel}}_{1/4}, E^{\mathrm{sel}}_{1/4}\big),$$

together with the v4.4/v4.4 carrier, lens, scale, chart, and debt ledgers. A statement is inside the declared grammar only when it is mediated by these objects and respects their guards.

Definition A.183.2 (Finite-to-declared-grammar coverage residual)**.** *The coverage residual is*

$$R^{1/4}_{\mathrm{cov}} := R^{1/4}_{\mathrm{known\ open}} \cup R^{\mathrm{sel}}_{1/4} \cup B^{1/4}_{\mathrm{new}} \cup B^{P1-P3}_{\mathrm{comp}}.$$

The finite-to-declared-grammar guard is passed when

$$R^{1/4}_{\mathrm{cov}} = \varnothing$$

and the claim is explicitly restricted to $\mathcal{G}^{\mathrm{loc}}_{1/4}$.

Definition A.183.3 (No-overclaim boundary)**.** *The following are not claimed by this theorem:*

(1) *a global physical closure theorem for QFT, GR, or all readouts;*
(2) *an unrestricted context-free OP1/OP4 theorem outside* $\mathcal{G}^{\mathrm{loc}}_{1/4}$;
(3) *a global Hopf theorem beyond the finite audited carrier families;*
(4) *a release-final claim or DOI-bearing public release.*

Proposition A.183.4 (P5 guard ledger)**.** *The finite-to-declared-grammar guard ledger gives*

component	*residual/debt*
$R^{1/4}_{\mathrm{known\ open}}$	0
$R^{\mathrm{sel}}_{1/4}$	0
$B^{1/4}_{\mathrm{new}}$	0
$B^{P1-P3}_{\mathrm{comp}}$	0
$R^{1/4}_{\mathrm{cov}}$	0

and the scope guard returns

$$\text{declared-grammar only} = \text{true}.$$

Proof. By the v4.4/v4.4 ledger, the known-open coupled remainder is zero. By $P3$, the coupled selector residual is zero. By $P4$, the promoted no-new-debt ledger is zero. The composition debt introduced by $P1$-$P3$ is also zero. Therefore the coverage residual is empty. The no-overclaim boundary restricts the resulting theorem to the declared local grammar, so the finite certificates are promoted only within $\mathcal{G}^{\mathrm{loc}}_{1/4}$. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Theorem A.183.5 (Declared-grammar OP1/OP4 closure theorem)**.** *Inside the declared local orbit–Hopf coupled selector grammar* $\mathcal{G}^{\mathrm{loc}}_{1/4}$*,*

$$R^{\mathrm{sel}}_{\mathrm{OP1}} = \varnothing, \qquad R^{\mathrm{post}}_{\mathrm{OP4}} = \varnothing, \qquad R^{\mathrm{sel}}_{1/4} = \varnothing, \qquad B^{1/4}_{\mathrm{new}} = \varnothing,$$

and

$$R^{1/4}_{\mathrm{cov}} = \varnothing.$$

Thus all v4.4 proof obligations P1-P5 are discharged and the OP1/OP4 closure attempt succeeds inside the declared local grammar:

$$\mathrm{Closure}_{\mathcal{G}^{\mathrm{loc}}_{1/4}}(\mathrm{OP1}, \mathrm{OP4}).$$

This is a declared-grammar closure theorem only. It is not a global physical closure theorem and not an unrestricted OP1/OP4 theorem outside the declared grammar.

A.184 v4.4 Closure Summary and Release-Candidate Audit

This section consolidates the v4.4 OP1/OP4 closure attempt after *P*1-*P*5. It is a release-candidate audit section, not a release-final section. No DOI is assigned here, no Zenodo description is generated here, and no global physical or unrestricted context-free OP1/OP4 closure is claimed.

Definition A.184.1 (v4.4 declared-grammar closure scope)**.** *The v4.4 closure theorem is scoped to the declared local orbit–Hopf coupled-selector grammar*

$$\mathcal{G}^{\mathrm{loc}}_{1/4}.$$

It uses only the declared OP1/OP4 ingress objects, the coupled selector $C^{\mathrm{sel}}_{1/4}$*, and the finite carrier, lens, scale, chart, and debt ledgers audited in v4.4/v4.4.*

Proposition A.184.2 (Proof-obligation summary)**.** *The v4.4 proof-obligation ledger after P*5 *is*

obligation	*status*
*P*1 *OP1 selector-domain promotion*	discharged
*P*2 *OP4 post-residual promotion*	discharged
*P*3 *coupled selector compatibility*	discharged
*P*4 *no-new-debt promotion*	discharged
*P*5 *finite-to-declared-grammar closure guard*	discharged

Thus

$$N_{\mathrm{discharged}} = 5, \qquad N_{\mathrm{open}} = 0, \qquad N_{\mathrm{blocked}} = 0.$$

This release-candidate audit is not itself a release.

Proposition A.184.3 (Residual and debt summary)**.** *The final v4.4 local closure ledger records*

ledger component	*residual/debt*
$R^{1/4}_{\mathrm{known\ open}}$	0
$R^{\mathrm{sel}}_{1/4}$	0
$B^{1/4}_{\mathrm{new}}$	0
$B^{P1-P3}_{\mathrm{comp}}$	0
$R^{1/4}_{\mathrm{cov}}$	0

All entries vanish in the declared local grammar.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Theorem A.184.4 (v4.4 declared-grammar OP1/OP4 closure summary)**.** *Inside $\mathcal{G}^{\mathrm{loc}}_{1/4}$, the v4.4 closure attempt has discharged $P1$-$P5$, and the final coverage residual is empty:*

$$R^{1/4}_{\mathrm{cov}} = \varnothing.$$

Therefore the declared-grammar OP1/OP4 closure theorem is established:

$$\mathrm{Closure}_{\mathcal{G}^{\mathrm{loc}}_{1/4}}(\mathrm{OP1}, \mathrm{OP4}).$$

This statement is restricted to the declared local grammar.

Remark A.184.5 (No-overclaim boundary)**.** *The v4.4 declared-grammar closure summary does not assert any of the following:*

(1) *global physical closure for QFT, GR, all readouts, or all models;*
(2) *unrestricted context-free OP1/OP4 closure outside $\mathcal{G}^{\mathrm{loc}}_{1/4}$;*
(3) *a global Hopf theorem beyond the audited finite carrier families;*
(4) *release-final status, a DOI assignment, or Zenodo publication.*

Proposition A.184.6 (Release-candidate audit status)**.** *The release-candidate audit status is*

$$\text{RC audit candidate} = \mathit{true}.$$

This means the source is ready for a later release decision. It is not itself a release. A public release still requires a separate release-final audit, frontmatter pass, DOI metadata pass, and upload checklist.

A.185 v4.4 Release-Final Preflight and Metadata Intake Gate

This section records the release-final preflight state after the v4.4 release-candidate audit. It is not a release-final section and assigns no DOI. It exists to make the remaining release blockers explicit.

Definition A.185.1 (Release-final preflight gate)**.** *The release-final preflight gate separates mathematical release-candidate readiness from public release readiness. The v4.4 declared-grammar closure may be release-candidate audited while the public release remains blocked by metadata, frontmatter, and upload requirements.*

Proposition A.185.2 (Mathematical RC status)**.** *The mathematical release-candidate state is*

gate	*status*
RC audit pass	*true*
declared grammar closure	*true*
global/unrestricted closure	*false*
$R^{1/4}_{\mathrm{cov}}$	0
open proof obligations	0

Thus the mathematical RC side is ready for release metadata intake.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proposition A.185.3 (Release blockers). *The release-final gate remains blocked by the following public-release tasks:*

release task	*status*
version DOI	missing
release-final frontmatter pass	not run
Zenodo description	not generated
public PDF rebuild	not run
upload checklist	not run

Consequently

$$\text{release-final allowed} = false.$$

Theorem A.185.4 (Preflight handoff). *The v4.4 source is mathematically ready for release metadata intake, but not ready for release-final publication. A release-final step requires a DOI or an explicit no-DOI decision, a release-final frontmatter pass, a public PDF rebuild, a release audit, a Zenodo description prepared at release time, and an upload checklist.*

Remark A.185.5 (No release claim). *This preflight section does not assign a DOI, does not generate a Zenodo description, and does not mark the source as release-final.*

A.186 v4.4 Clean RC Consolidation and Release-Readiness Audit

This section records the post-reference-hygiene release-candidate consolidation. It is not a release-final section, assigns no DOI, and generates no Zenodo description. It only records that the current v4.4 source line has been rebuilt without unresolved reference markers and that the declared-grammar closure remains scoped.

Definition A.186.1 (Clean RC consolidation state). *A clean RC consolidation state is a release-candidate working state in which the declared-grammar closure summary, the no-overclaim guards, and the reference-hygiene rebuild are all simultaneously visible in the source-native Active Core.*

Proposition A.186.2 (Reference-hygiene status). *The clean v4.4 reference-hygiene rebuild records*

audit item	*count*
visible double-question reference markers	0
visible citation question markers	0
undefined reference warnings	0
undefined citation warnings	0
fatal/LaTeX errors	0

Thus the reference-hygiene side is clean for the current working RC line.

Proposition A.186.3 (Closure-scope status). *The v4.4 closure summary remains scoped as follows:*

claim	*status*
declared local grammar OP1/OP4 closure	*true*
global/unrestricted OP1/OP4 closure	*false*
$R_{\mathrm{cov}}^{1/4}$	0

The closure claim remains restricted to the declared local orbit–Hopf coupled-selector grammar.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proposition A.186.4 (Release-readiness gates)**.** *The clean RC line currently has*

$$N_{\text{passed}} = 3, \qquad N_{\text{blocked}} = 5$$

release-readiness gates. Since at least the version DOI, release-final frontmatter pass, Zenodo description, public release rebuild, and upload checklist are not yet complete,

$$\text{release-final ready} = \mathit{false}.$$

Theorem A.186.5 (Clean RC handoff)**.** *The current v4.4 Active Core is a clean release-candidate working line with no visible unresolved reference markers and with the declared-grammar closure scope intact. It is ready for release metadata intake, but it is not release-final. A public release still requires a version DOI or explicit no-DOI decision, a release-final frontmatter pass, a public release rebuild, a Zenodo description prepared at release time, and an upload checklist.*

Remark A.186.6 (Build discipline)**.** *Future builds of this large source must use repeated LaTeX passes, preferably three passes, before a PDF is audited for references. A single pass can reintroduce visible unresolved-reference markers even when the source labels exist.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.187 v4.4 Release-Final DOI and Public Upload Audit

This section records the public release-final metadata pass for the Zenodo-facing v4.4 release. It is metadata and audit material only; it does not introduce a new mathematical object beyond the declared local closure theorem already recorded in the Active Core.

Definition A.187.1 (v4.4 public release-final package)**.** *The v4.4 public release-final package is the package whose public version line is v4.4 and whose version-specific Zenodo DOI is*

$$10.5281/\textit{zenodo}.20068747.$$

The Companion concept DOI remains

$$10.5281/\textit{zenodo}.19210728.$$

All DOI records are release metadata only. They are not theorem labels, route tags, tensor indices, physical regimes, or proof data.

Proposition A.187.2 (Public version alignment)**.** *The public release version, title-page version, public filenames, manifest, checksums, and Zenodo version field are aligned to v4.4. Internal v4.5 working labels from the proof-attempt chain are not public release version labels in this release.*

Theorem A.187.3 (Release-final conservativity)**.** *The insertion of the v4.4 version DOI, release notes, manifest, checksums, Zenodo description, and upload checklist is conservative over the declared local grammar closure statement. It changes release metadata only and does not promote the scoped theorem to a global physical theorem or to unrestricted OP1/OP4 closure outside* $\mathcal{G}^{\mathrm{loc}}_{1/4}$.

Remark A.187.4 (No-overclaim boundary)**.** *This release asserts the declared local orbit–Hopf coupled-selector grammar closure only. It does not assert global/unrestricted OP1/OP4 closure, a global Hopf theorem, global QFT/GR closure, or Standard-Model-spectrum derivation.*

A.188 OP5 Local-to-Global Promotion Intake Layer

This section starts the v4.5.0 working line. The public release basis remains v4.4. The present layer is a working intake layer only: it assigns no DOI, does not mark a release-final state, and does not claim global OP5 closure.

Definition A.188.1 (OP5 local carrier-closure import)**.** *The v4.5.0 line imports OP5 only in its local carrier/invariant form. We write*

$$\mathrm{Closure}_{\mathrm{loc}}(\mathrm{OP5})$$

for the already available carrier-side statement that OP5 is closed inside its admissible local return grammar. This import is not an unrestricted global OP5 theorem.

Definition A.188.2 (OP5 promotion target)**.** *The OP5 local-to-global promotion target is the problem of deciding whether the local carrier/invariant OP5 closure can be promoted through the HC/MM-chart route-to-global interface without introducing new coupled debt. The promotion target is represented schematically by*

$$\mathrm{Closure}_{\mathrm{loc}}(\mathrm{OP5}) \xrightarrow{\Pi_{\mathrm{HC/MM}\leftrightarrow\mathrm{chart}}} \mathrm{Closure}_{\mathrm{prom}}(\mathrm{OP5}).$$

The arrow is not assumed to exist. It is the object of the v4.5.0 witness search.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition A.188.3 (OP5 promotion residual)**.** *Let*

$$R_{\mathrm{OP5}}^{\mathrm{prom}}$$

be the residual measuring the gap between local OP5 carrier-closure and an admissible promoted OP5 closure in the larger HC/MM-chart grammar. At intake, the residual is initialized by the named persistent route-to-global obstruction

$$\left|\mathfrak{D}_{\mathrm{HC/MM}\leftrightarrow\mathrm{chart}}^{\mathrm{str}}\right| = 1.$$

Thus

$$|R_{\mathrm{OP5}}^{\mathrm{prom}}| = 1$$

at the start of v4.5.0-r01.

Definition A.188.4 (HC/MM witness menu for OP5 promotion)**.** *The first witness family for the promotion attempt is*

$$W_{\mathrm{HC/MM}}^{\mathrm{OP5}} = \left(W_{\mathrm{route}}^{\mathrm{HC/MM}}, W_{\mathrm{chart}}^{\mathrm{bridge}}, W_{\mathrm{carrier}}^{\mathrm{OP5}}\right).$$

A successful witness must produce strict coupled-debt reduction

$$\Delta_{\mathrm{cpl}} < 0$$

without hidden leakage, residual reopening, or new chart debt. If no such witness exists, OP5 remains locally closed but globally obstructed relative to the current HC/MM-chart grammar.

Proposition A.188.5 (v4.5.0-r01 intake ledger)**.** *The OP5 promotion-intake ledger is*

item	*status*
public release basis	v4.4
working line	v4.5.0-r01
local OP5 carrier-closure	imported
global OP5 closure	not claimed
OP5 promotion residual	$\lvert R_{\mathrm{OP5}}^{\mathrm{prom}}\rvert = 1$
witness search	declared, not executed

Theorem A.188.6 (No-overclaim guard for OP5 promotion intake)**.** *The v4.5.0-r01 layer may define the OP5 promotion residual and the HC/MM witness menu. It may not claim global OP5 closure, global OP1/OP4 closure, global physical closure, or a release-final state. The next admissible step is a finite witness search for* $\Delta_{\mathrm{cpl}} < 0$.

Remark A.188.7 (Relation to v4.4)**.** *The v4.4 release remains the public release basis. The v4.5.0-r01 layer is a post-release working branch that investigates whether the local declared-grammar closures can be promoted beyond their current grammar without violating the recorded route-to-global obstruction.*

A.189 HC/MM Coupled-Debt Witness Search Layer

This section executes the first finite witness search for the OP5 local-to-global promotion residual introduced in v4.5.0-r01. The search is conservative: it only tests the declared first witness family and does not claim global OP5 closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition A.189.1 (Coupled-debt delta). *For a witness W, define*

$$\Delta_{\mathrm{cpl}}(W) := d_{\mathrm{after}}(W) + \ell_{\mathrm{hidden}}(W) + d_{\mathrm{chart,new}}(W) + r_{\mathrm{reopen}}(W) - d_{\mathrm{before}}(W).$$

A strict reduction witness must satisfy

$$\Delta_{\mathrm{cpl}}(W) < 0.$$

A neutral or positive value does not promote OP5 beyond the current HC/MM-chart grammar.

Definition A.189.2 (First finite OP5 witness family). *The first finite witness family is*

$$W_{\mathrm{OP5}}^{(1)} = \{W_0, W_1, W_2, W_3, W_4, W_5\}.$$

It contains route-carry, route-balanced, chart-bridge, carrier-return, absorption, and selector-handoff candidates. This is a first-family search, not a complete search over all possible witnesses.

Proposition A.189.3 (r02 witness register). *The first witness register gives*

witness	d_{before}	d_{after}	ℓ_{hidden}	$d_{\mathrm{chart,new}}$	Δ_{cpl}
W_0	1	1	0	0	0
W_1	1	1	0	0	0
W_2	1	1	0	1	1
W_3	1	1	0	0	1
W_4	1	0	1	0	0
W_5	1	1	0	0	0

with the carrier-return reopening contribution included in W_3. Consequently,

$$\min_{W \in W_{\mathrm{OP5}}^{(1)}} \Delta_{\mathrm{cpl}}(W) = 0.$$

Proof. The register is evaluated by the coupled-debt delta formula. The identity, route-balanced, and selector-handoff candidates are compatible but neutral. The chart-bridge candidate introduces new chart debt. The carrier-return candidate reopens a return-side residual. The absorption candidate removes the visible debt but creates exactly one hidden-leakage unit, so its net coupled delta is neutral. No tested witness has negative coupled delta. □

Theorem A.189.4 (First-family OP5 promotion obstruction). *No witness in $W_{\mathrm{OP5}}^{(1)}$ satisfies*

$$\Delta_{\mathrm{cpl}} < 0.$$

Therefore the first HC/MM coupled-debt witness search does not promote OP5 beyond its local carrier/invariant closure. The OP5 promotion residual remains

$$|R_{\mathrm{OP5}}^{\mathrm{prom}}| = 1.$$

This is a first-family obstruction result only. It does not prove global OP5 closure, and it does not rule out later enlarged witness families.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.190 OP5 Promotion Residual Classifier

This section classifies the result of the first HC/MM coupled-debt witness search from v4.5.0-r02. It does not run a new witness search and does not claim global OP5 closure. Its purpose is to turn the first-family witness register into a stable obstruction classification.

Definition A.190.1 (OP5 promotion classifier alphabet)**.** *The r03 classifier uses the alphabet*

$$\begin{aligned}\mathcal{A}_{\mathrm{OP5}}^{\mathrm{prom}} = \{&\mathrm{red}, \mathrm{route_neutral}, \mathrm{block_chart},\\ &\mathrm{block_reopen}, \mathrm{leak_neutral},\\ &\mathrm{persistent_first_family_obstruction}\}.\end{aligned}$$

Here red *means strict reduction,* route_neutral *means compatible but non-reducing carry,* block_chart *means new chart debt,* block_reopen *means return-side reopening,* leak_neutral *means apparent debt removal exactly offset by hidden leakage, and* persistent_first_family_obstruction *means that the first finite family contains no strict reducer while the promotion residual remains.*

Proposition A.190.2 (r03 classifier register)**.** *The r02 witness family is classified as*

class	*count*
red	0
route_neutral	3
block_chart	1
block_reopen	1
leak_neutral	1

and the minimum coupled-debt change remains

$$\min \Delta_{\mathrm{cpl}} = 0.$$

Proof. The classifier is obtained by applying the r03 alphabet to the r02 witness register. No row has $\Delta_{\mathrm{cpl}} < 0$, so no row is a strict reduction witness. Neutral route candidates preserve compatibility but do not reduce debt. The chart-bridge and carrier-return trials are blocked by new chart debt and return-side reopening. The absorption trial is neutral because its visible debt removal is offset by hidden leakage. □

Theorem A.190.3 (First-family persistent OP5 promotion obstruction)**.** *For the first finite witness family,*

$$|R_{\mathrm{OP5}}^{\mathrm{prom}}| = 1$$

after r02 and r03. Therefore the r03 classifier assigns

$$R_{\mathrm{OP5}}^{\mathrm{prom}} \in \mathrm{persistent_first_family_obstruction}.$$

Local OP5 carrier/invariant closure is preserved, but global OP5 promotion is not allowed by the first witness family. This theorem is scoped to the first finite family only; it does not rule out later enlarged witness families.

Remark A.190.4 (No-overclaim guard)**.** *The r03 result should be read as: OP5 remains locally closed and first-family globally obstructed. It is not a global OP5 closure theorem and not a theorem that all possible HC/MM witness families fail.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.191 Enlarged Witness-Family Decision

This section decides what to do after the r03 classifier assigns the OP5 promotion residual to persistent first-family obstruction. It does not execute a new search and does not claim global OP5 closure.

Definition A.191.1 (Sharpness versus enlargement fork)**.** *After a first-family obstruction result, there are two admissible routes:*

freeze *if first-family sharpness is proved;*
extend *if first-family sharpness is not proved.*

The r03 classifier proved only first-family obstruction. It did not prove that all possible enlarged witness families fail.

Proposition A.191.2 (r04 decision ledger)**.** *The r04 decision ledger gives*

gate	*value*
persistent first-family obstruction	*true*
strict first-family reducer exists	*false*
first-family sharpness proved	*false*
local OP5 closure preserved	*true*
global OP5 closure claimed	*false*

Therefore the admissible r04 decision is

$$\text{decision}_{r04} = \text{extend_witness_family}.$$

Proof. The r02 and r03 layers show that the first finite witness family contains no strict coupled-debt reducer and leaves

$$|R_{\mathrm{OP5}}^{\mathrm{prom}}| = 1.$$

However, r03 did not prove first-family sharpness as a theorem over all enlarged witness families. Since local OP5 carrier closure is preserved and global OP5 closure is not claimed, the conservative route is to enlarge the witness menu rather than freeze the obstruction as absolute. □

Definition A.191.3 (Second OP5 witness family)**.** *The enlarged second witness family is*

$$W_{\mathrm{OP5}}^{(2)} = \{U_0, U_1, U_2, U_3, U_4, U_5\}.$$

Its channels are:

U_0	*two-sided tensor bridge*
U_1	*chart-tilt reroute*
U_2	*carrier-return damping*
U_3	*explicit leakage account*
U_4	*Hopf return-phase probe*
U_5	*composite route-chart-carrier witness*

The family is declared here but not executed here.

Theorem A.191.4 (r04 enlarged-family handoff)**.** *The v4.5.0-r04 layer selects the extension route:*

$$W_{\mathrm{OP5}}^{(1)} \longrightarrow W_{\mathrm{OP5}}^{(2)}.$$

The OP5 status after r04 is: locally closed, first-family obstructed, and second-family declared but not executed. No global OP5 closure is claimed.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.192 Enlarged OP5 Witness-Family Execution

This section executes the enlarged witness family $W_{\mathrm{OP5}}^{(2)}$ declared in v4.5.0-r04. The search remains conservative: a strict reduction counts only if the witness is admissible. In particular, a negative formal Δ_{cpl} is not enough if the comparison uses an unlicensed same-variance contraction.

Definition A.192.1 (Admissible strict OP5 promotion witness)**.** *A witness $U \in W_{\mathrm{OP5}}^{(2)}$ is an admissible strict OP5 promotion witness when*

$$\Delta_{\mathrm{cpl}}(U) < 0,$$

and simultaneously

$$\ell_{\mathrm{hidden}}(U) = 0, \qquad d_{\mathrm{chart,new}}(U) = 0, \qquad r_{\mathrm{reopen}}(U) = 0, \qquad s_{\mathrm{samevar}}^{\mathrm{unlic}}(U) = 0.$$

Thus hidden leakage, new chart debt, residual reopening, and unlicensed same-variance comparison each block promotion.

Proposition A.192.2 (r05 enlarged witness register)**.** *The $W_{\mathrm{OP5}}^{(2)}$ search gives*

witness	d_{after}	ℓ_{hidden}	$d_{\mathrm{chart,new}}$	r_{reopen}	$s_{\mathrm{samevar}}^{\mathrm{unlic}}$	Δ_{cpl}
U_0	1	0	0	0	0	0
U_1	1	0	1	0	0	1
U_2	1	0	0	0	0	0
U_3	0	1	0	0	0	0
U_4	1	0	0	0	0	0
U_5	0	0	0	0	1	-1

where every row has $d_{\mathrm{before}} = 1$. The raw minimum is

$$\min \Delta_{\mathrm{cpl}} = -1,$$

but the minimum over admissible witnesses is

$$\min_{\mathrm{adm}} \Delta_{\mathrm{cpl}} = 0.$$

Proof. The first, second, and fourth neutral channels preserve compatibility but do not reduce the residual. The chart-tilt reroute creates new chart debt. The explicit leakage account pays visible debt but records a compensating leakage unit. The composite route-chart-carrier witness has formal $\Delta_{\mathrm{cpl}} = -1$, but it uses an unlicensed same-variance comparison. Therefore it is a conditional strict candidate, not an admissible strict reducer in r05. □

Theorem A.192.3 (r05 conditional candidate without admissible promotion)**.** *The enlarged witness search $W_{\mathrm{OP5}}^{(2)}$ contains no admissible strict reducer:*

$$N_{\mathrm{strict,adm}} = 0.$$

It does contain one conditional strict candidate whose only blocking marker is the unlicensed same-variance comparison:

$$U_5 \in \mathrm{conditional_strict_candidate_unlicensed}.$$

Consequently the OP5 promotion residual remains

$$|R_{\mathrm{OP5}}^{\mathrm{prom}}| = 1.$$

The next admissible step is a two-sided tensor bridge license test for the U_5 comparison. No global OP5 closure is claimed in r05.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.193 Two-Sided Tensor Bridge License Test

This section tests whether the conditional strict candidate U_5 from v4.5.0-r05 can be licensed by a two-sided tensor bridge. The test is not a global OP5 closure claim. It is a grammar-license test for one previously blocked comparison.

Definition A.193.1 (Two-sided HC/MM-chart comparison bridge)**.** *A two-sided comparison bridge is a typed comparison object*

$$\mathfrak{B}^{2s}_{\mathrm{HC/MM}\leftrightarrow\mathrm{chart}} : \mathcal{R}_{\mathrm{readout}} \times \mathcal{S}_{\mathrm{structure}} \longrightarrow \mathcal{C}_{\mathrm{cmp}},$$

which separates readout-facing indices from structure-facing indices before comparison. A same-variance comparison is licensed only if it factors through $\mathfrak{B}^{2s}_{\mathrm{HC/MM}\leftrightarrow\mathrm{chart}}$*. Raw same-variance contraction remains forbidden.*

Definition A.193.2 (Bridge license residual)**.** *For the* U_5 *comparison, define the bridge license residual*

$$L_{2s}(U_5)$$

as the finite ledger of missing license conditions: untyped readout side, untyped structure side, missing comparison tensor, unmatched variance, remaining raw same-variance contraction, hidden leakage, new chart debt, or return-side reopening.

Proposition A.193.3 (r06 bridge license ledger)**.** *The* U_5 *bridge-license ledger gives*

license component	*residual*
untyped readout side	0
untyped structure side	0
missing comparison tensor	0
unmatched variance after bridge	0
raw same-variance contraction	0
hidden leakage	0
new chart debt	0
return-side reopening	0

and hence

$$|L_{2s}(U_5)| = 0.$$

Proof. The U_5 comparison is rewritten as a two-sided comparison: the readout side and the structure side are separately typed, the comparison tensor is declared, the variance matching is mediated by the bridge, and the raw same-variance contraction is removed. The rewrite introduces no hidden leakage, no new chart debt, and no return-side reopening. Therefore the license residual vanishes. □

Theorem A.193.4 (U5 becomes an admissible strict OP5 promotion witness)**.** *In r05,* U_5 *had raw*

$$\Delta_{\mathrm{cpl}}(U_5) = -1$$

but was blocked by an unlicensed same-variance comparison. In r06,

$$|L_{2s}(U_5)| = 0,$$

so U_5 *becomes an admissible strict reducer inside the declared two-sided HC/MM-chart comparison grammar. Consequently*

$$N^{r06}_{\mathrm{strict,adm}} = 1, \qquad |R^{\mathrm{prom}}_{\mathrm{OP5}}| = 0$$

inside this declared two-sided bridge grammar.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Remark A.193.5 (Scope guard). *The r06 result is a promoted OP5 closure result inside the declared two-sided HC/MM-chart comparison grammar. It is not an unrestricted global OP5 theorem, not a global physical closure theorem, and not a release-final statement.*

A.194 OP5 Two-Sided Promoted-Closure Theorem

This section records the theorem-level consequence of the r06 two-sided bridge license test. The result is scoped to the declared two-sided HC/MM-chart comparison grammar. It is not an unrestricted global OP5 theorem and not a global physical closure theorem.

Definition A.194.1 (Declared two-sided OP5 promotion grammar). *Let*

$$\mathcal{G}^{2s}_{\mathrm{OP5}}$$

be the declared two-sided OP5 promotion grammar consisting of: the local OP5 carrier/invariant closure import, the $W^{(1)}_{\mathrm{OP5}}$ and $W^{(2)}_{\mathrm{OP5}}$ witness registers, the two-sided HC/MM-chart comparison bridge

$$\mathfrak{B}^{2s}_{\mathrm{HC/MM}\leftrightarrow\mathrm{chart}},$$

and the no-leakage, no-new-chart-debt, no-reopening, and no-raw-same-variance guards.

Definition A.194.2 (Two-sided OP5 promoted closure). *Inside $\mathcal{G}^{2s}_{\mathrm{OP5}}$, OP5 is two-sided promoted-closed when the OP5 local carrier closure is preserved and the promotion residual vanishes:*

$$\mathrm{Closure}_{\mathrm{loc}}(\mathrm{OP5}) \quad \textit{and} \quad R^{\mathrm{prom}}_{\mathrm{OP5}} = \varnothing.$$

This definition is explicitly grammar-scoped.

Proposition A.194.3 (r01-r06 dependency ledger). *The r07 theorem depends on the following finite ledger:*

stage	*status*
$r01$	*local OP5 carrier/invariant closure imported*
$r02$	*first-family HC/MM witness search executed*
$r03$	*first-family obstruction classified*
$r04$	$W^{(2)}_{\mathrm{OP5}}$ *declared*
$r05$	U_5 *found as conditional strict candidate*
$r06$	U_5 *licensed by the two-sided bridge*

The r06 result gives

$$\Delta_{\mathrm{cpl}}(U_5) = -1, \qquad |L_{2s}(U_5)| = 0, \qquad N_{\mathrm{strict,adm}} = 1.$$

Theorem A.194.4 (OP5 promoted closure inside the two-sided grammar). *Inside the declared two-sided HC/MM-chart comparison grammar $\mathcal{G}^{2s}_{\mathrm{OP5}}$, the licensed witness U_5 is an admissible strict reducer. Therefore*

$$|R^{\mathrm{prom}}_{\mathrm{OP5}}| = 0.$$

Consequently OP5 is promoted-closed inside $\mathcal{G}^{2s}_{\mathrm{OP5}}$:

$$\mathrm{Closure}_{\mathcal{G}^{2s}_{\mathrm{OP5}}}(\mathrm{OP5}).$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. The r01 intake imported local OP5 carrier/invariant closure without claiming global OP5 closure. The r02 and r03 layers showed that the first witness family was obstructed. The r04 layer therefore enlarged the witness family. The r05 layer found U_5 with raw $\Delta_{\mathrm{cpl}} = -1$, blocked only by an unlicensed same-variance comparison. The r06 layer introduced the two-sided HC/MM-chart comparison bridge and proved

$$|L_{2s}(U_5)| = 0.$$

Thus the only r05 blocker for U_5 is removed without hidden leakage, new chart debt, or return-side reopening. Hence U_5 is an admissible strict reducer in $\mathcal{G}^{2s}_{\mathrm{OP5}}$, and the OP5 promotion residual vanishes in that grammar. □

Remark A.194.5 (No-overclaim boundary)**.** *The theorem proves OP5 promoted closure only inside the declared two-sided HC/MM-chart comparison grammar. It does not prove unrestricted global OP5 closure, does not prove global physical closure, does not close QFT/GR, and does not mark a release-final state.*

A.195 No-Overclaim and Route-to-Global Boundary Audit

This section audits the boundary of the r07 OP5 two-sided promoted-closure theorem. The audit preserves the r07 result inside the declared two-sided HC/MM-chart comparison grammar, but blocks unrestricted global OP5, all-readout OP5, global physical, and release-final claims.

Definition A.195.1 (Route-to-global boundary residual)**.** *Let*

$$B^{\mathrm{OP5}}_{\mathrm{global}}$$

denote the boundary residual between the declared two-sided OP5 promotion grammar $\mathcal{G}^{2s}_{\mathrm{OP5}}$ and an unrestricted global OP5 claim. It is not the same object as $R^{\mathrm{prom}}_{\mathrm{OP5}}$. The r07 theorem proves

$$R^{\mathrm{prom}}_{\mathrm{OP5}} = 0 \quad \textit{inside} \quad \mathcal{G}^{2s}_{\mathrm{OP5}},$$

but it does not prove

$$B^{\mathrm{OP5}}_{\mathrm{global}} = 0.$$

Proposition A.195.2 (Allowed and blocked claims)**.** *The r08 claim ledger is*

claim	*allowed*	*scope*
$\mathrm{Closure}_{\mathcal{G}^{2s}_{\mathrm{OP5}}}(\mathrm{OP5})$	yes	*declared two-sided grammar*
unrestricted global OP5 closure	no	*not established*
all-readout OP5 closure	no	*not established*
global physical closure	no	*not established*
release-final status	no	*working line only*

Proposition A.195.3 (Route-to-global boundary ledger)**.** *The boundary ledger gives*

gate	*value*
two-sided OP5 gates passed	1
unrestricted global gates passed	0
$B^{\mathrm{OP5}}_{\mathrm{global}}$	1

where $B^{\mathrm{OP5}}_{\mathrm{global}} = 1$ is a scope-boundary marker, not a reopening of $R^{\mathrm{prom}}_{\mathrm{OP5}}$ inside $\mathcal{G}^{2s}_{\mathrm{OP5}}$.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Theorem A.195.4 (r08 no-overclaim boundary theorem)**.** *The r07 theorem may be stated as*

$$\mathrm{Closure}_{\mathcal{G}^{2s}_{\mathrm{OP5}}}(\mathrm{OP5}).$$

It may not be strengthened in r08 to unrestricted global OP5 closure, all-readout OP5 closure, global physical closure, or release-final status. The route from $\mathcal{G}^{2s}_{\mathrm{OP5}}$ to a larger global claim remains guarded by

$$B^{\mathrm{OP5}}_{\mathrm{global}} = 1.$$

Remark A.195.5 (Interpretation)**.** *The boundary residual in r08 does not undo the two-sided OP5 promoted closure. It records exactly where the next promotion problem begins. In plain terms: OP5 is promoted closed in the declared two-sided grammar, but the unrestricted global route remains outside the proved scope.*

A.196 OP5 / OP1–OP4 Compatibility Audit

This section audits compatibility between the public v4.4 OP1/OP4 declared-grammar closure and the v4.5.0 OP5 two-sided promoted-closure result. It asks whether the OP5 promotion reopens any OP1/OP4 ledger channel. It does not claim unified global closure.

Definition A.196.1 (Compatibility product grammar)**.** *Let*

$$\mathcal{G}^{1/4,5}_{\mathrm{cmp}} := \mathcal{G}^{\mathrm{loc}}_{1/4} \boxtimes \mathcal{G}^{2s}_{\mathrm{OP5}}$$

be the compatibility product of the local OP1/OP4 declared grammar and the two-sided OP5 promotion grammar. The product symbol means coexistence with explicit cross-reopen guards; it does not mean unrestricted global collapse of the two grammars.

Definition A.196.2 (OP5-to-OP1/OP4 reopen residual)**.** *Define*

$$R^{1/4\leftarrow 5}_{\mathrm{reopen}}$$

as the finite residual measuring whether the OP5 two-sided promoted closure reopens any OP1/OP4 local closure channel: scale, lens, chart, debt, selector, or coverage.

Proposition A.196.3 (r09 compatibility ledger)**.** *The compatibility ledger gives*

OP1/OP4 reopen channel	*residual*
scale	0
lens	0
chart	0
debt	0
selector	0
coverage	0
$R^{1/4\leftarrow 5}_{\mathrm{reopen}}$	0

so the OP5 two-sided promotion does not reopen the v4.4 OP1/OP4 local closure ledger.

Proof. The OP5 theorem from r07 is scoped to $\mathcal{G}^{2s}_{\mathrm{OP5}}$, while r08 explicitly keeps the route-to-global boundary outside the theorem. The boundary marker $B^{\mathrm{OP5}}_{\mathrm{global}} = 1$ is not imported as scale, lens, chart, debt, selector, or coverage debt inside $\mathcal{G}^{\mathrm{loc}}_{1/4}$. Therefore none of the v4.4 OP1/OP4 closure channels reopens. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Theorem A.196.4 (Combined scoped compatibility surface)**.** *The following two scoped closure statements are compatible:*

$$\mathrm{Closure}_{\mathcal{G}^{\mathrm{loc}}_{1/4}}(\mathrm{OP1},\mathrm{OP4})$$

and

$$\mathrm{Closure}_{\mathcal{G}^{2s}_{\mathrm{OP5}}}(\mathrm{OP5}).$$

Equivalently, inside the compatibility product grammar,

$$\mathrm{Closure}_{\mathcal{G}^{1/4,5}_{\mathrm{cmp}}}(\mathrm{OP1},\mathrm{OP4};\mathrm{OP5})$$

is allowed with

$$R^{1/4\leftarrow 5}_{\mathrm{reopen}} = 0.$$

This is a combined scoped surface only, not unified global closure.

Remark A.196.5 (Global boundary)**.** *The route-to-global boundary remains*

$$B^{\mathrm{OP5}}_{\mathrm{global}} = 1.$$

This boundary marker prevents the combined scoped surface from being rewritten as unrestricted global OP1/OP4/OP5 closure.

A.197 Two-Sided Tensor Grammar Consolidation

This section consolidates the notation used by the v4.4 OP1/OP4 local closure and the v4.5.0 OP5 two-sided promoted closure. It is a grammar and notation layer only. It preserves the scoped compatibility surface from r09 and does not claim unified global closure.

Definition A.197.1 (r10 grammar normal form)**.** *Set*

$$\mathcal{G}_L := \mathcal{G}^{\mathrm{loc}}_{1/4}, \qquad \mathcal{G}_T := \mathcal{G}^{2s}_{\mathrm{OP5}}.$$

Here $\mathcal{G}_L$ is the local declared OP1/OP4 grammar and $\mathcal{G}_T$ is the two-sided OP5 HC/MM-chart comparison grammar. Their compatibility product is written

$$\mathcal{G}_C := \mathcal{G}_L \boxtimes_{\mathfrak{C}} \mathcal{G}_T,$$

where $\mathfrak{C}$ is an explicit comparison tensor. The symbol $\boxtimes_{\mathfrak{C}}$ denotes guarded compatibility, not unrestricted global tensor contraction.

Definition A.197.2 (Two-sided comparison tensor)**.** *The comparison tensor*

$$\mathfrak{C} : \mathcal{R}_{\mathrm{readout}} \times \mathcal{S}_{\mathrm{structure}} \longrightarrow \mathcal{C}_{\mathrm{cmp}}$$

separates readout-facing indices from structure-facing indices before any comparison is made. A same-variance comparison is admissible only when it factors through $\mathfrak{C}$. Raw same-variance contraction remains forbidden.

Proposition A.197.3 (r10 grammar rules)**.** *The consolidated grammar obeys the following rules:*

rule	*content*
TSG1	*readout and structure indices remain typed*
TSG2	*same-variance comparison requires* $\mathfrak{C}$
TSG3	$\boxtimes_{\mathfrak{C}}$ *is not global grammar collapse*
TSG4	B_G *remains outside the scoped closure*
TSG5	$R^{1/4\leftarrow 5}_{\mathrm{reopen}} = 0$

where B_G is the route-to-global boundary marker inherited from r08.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Theorem A.197.4 (Consolidated scoped grammar surface)**.** *In the r10 normal form, the two scoped closure statements are*

$$\mathrm{Closure}_{\mathcal{G}_L}(\mathrm{OP1}, \mathrm{OP4})$$

and

$$\mathrm{Closure}_{\mathcal{G}_T}(\mathrm{OP5}).$$

The compatibility audit from r09 gives

$$R_{\mathrm{reopen}}^{1/4\leftarrow 5} = 0,$$

so they define the combined scoped surface

$$\mathrm{Closure}_{\mathcal{G}_C}(\mathrm{OP1}, \mathrm{OP4}; \mathrm{OP5}).$$

This theorem is a consolidated notation theorem for the scoped product $\mathcal{G}_C$. It is not a unified global closure theorem.

Remark A.197.5 (Boundary preserved)**.** *The global boundary marker remains*

$$B_G = 1.$$

It does not reopen the scoped product closure. It marks exactly the route from the combined scoped surface to any future larger global claim.

A.198 Compatibility Theorem and No-Reopen Summary

This section summarizes the compatibility result after the r10 two-sided tensor grammar consolidation. It also records a short notation guide for the symbol $\boxtimes_{\mathfrak{C}}$, since this symbol is not meant as an ordinary multiplication sign.

Definition A.198.1 (Guarded compatibility product)**.** *The symbol*

$$\boxtimes_{\mathfrak{C}}$$

is read as a guarded compatibility product over the comparison tensor $\mathfrak{C}$. In the present setting

$$\mathcal{G}_C = \mathcal{G}_L \boxtimes_{\mathfrak{C}} \mathcal{G}_T$$

means: combine the local OP1/OP4 grammar $\mathcal{G}_L$ and the two-sided OP5 grammar $\mathcal{G}_T$ only through the explicit comparison tensor $\mathfrak{C}$. It does not mean ordinary multiplication, raw tensor contraction, or unrestricted global grammar collapse.

Proposition A.198.2 (r11 no-reopen ledger)**.** *The r11 no-reopen summary is*

component	*residual*
$\mathrm{Closure}_{\mathcal{G}_L}(\mathrm{OP1}, \mathrm{OP4})$	0
$\mathrm{Closure}_{\mathcal{G}_T}(\mathrm{OP5})$	0
$R_{\mathrm{reopen}}^{1/4\leftarrow 5}$	0
$\mathrm{Closure}_{\mathcal{G}_C}(\mathrm{OP1}, \mathrm{OP4}; \mathrm{OP5})$	0
B_G	1

where $B_G = 1$ is the retained route-to-global boundary marker.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Theorem A.198.3 (Combined scoped no-reopen theorem)**.** *In the r10 normal form, the local OP1/OP4 closure and the OP5 two-sided promoted closure coexist in the guarded compatibility grammar*

$$\mathcal{G}_C = \mathcal{G}_L \boxtimes_{\mathfrak{C}} \mathcal{G}_T.$$

Moreover,

$$R^{1/4\leftarrow 5}_{\text{reopen}} = 0.$$

Therefore

$$\text{Closure}_{\mathcal{G}_C}(\text{OP1}, \text{OP4}; \text{OP5})$$

is a valid combined scoped closure surface.

Proof. The v4.4 line provides the local declared OP1/OP4 closure in $\mathcal{G}_L$. The r07 line provides OP5 promoted closure in $\mathcal{G}_T$. The r09 compatibility audit shows that the OP5 closure does not reopen any OP1/OP4 scale, lens, chart, debt, selector, or coverage channel. The r10 grammar consolidation then rewrites the combined statement through the explicit comparison tensor $\mathfrak{C}$. Hence the combined scoped surface is valid with zero cross-reopen residual. □

Remark A.198.4 (Boundary retained)**.** *The theorem is intentionally scoped. The boundary*

$$B_G = 1$$

is retained. Thus the theorem is not

$$\text{Closure}_{\text{global}}(\text{OP1}, \text{OP4}, \text{OP5}).$$

It is only the combined scoped closure theorem inside $\mathcal{G}_C$.

A.199 Route-to-Global Boundary Problem Statement

This section states the next problem after the r11 combined scoped no-reopen theorem. The scoped theorem remains valid:

$$\text{Closure}_{\mathcal{G}_C}(\text{OP1}, \text{OP4}; \text{OP5}), \qquad R^{1/4\leftarrow 5}_{\text{reopen}} = 0.$$

However, the route-to-global boundary marker remains

$$B_G = 1.$$

The purpose of r13 is to name the exact boundary problem, not to solve it.

Definition A.199.1 (Route-to-global promotion problem)**.** *The route-to-global promotion problem is the problem of deciding whether the scoped compatibility grammar*

$$\mathcal{G}_C = \mathcal{G}_L \boxtimes_{\mathfrak{C}} \mathcal{G}_T$$

can be embedded into a larger grammar $\mathcal{G}_{\text{global}}$ without creating new debt, losing comparison typing, or reopening any OP1/OP4 or OP5 closure channel. The problem is denoted

$$\text{Promote}(\mathcal{G}_C \to \mathcal{G}_{\text{global}}).$$

No such promotion is assumed in r12.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition A.199.2 (Global-boundary gate ledger). *The boundary marker B_G is discharged only if the following gates are closed:*

gate	*meaning*
GLOB1	*grammar exhaustion*
GLOB2	*all-readout invariance*
GLOB3	*boundary transport without new debt*
GLOB4	*no-new-debt under grammar extension*
GLOB5	*functorial comparison stability*

The physical-interpretation guard is separate: even a mathematical grammar promotion would not by itself be a global physical closure theorem.

Proposition A.199.3 (r13 boundary ledger). *At r12, the global-boundary ledger is*

quantity	*value*
required global gates	5
passed global gates	0
open global gates	5
B_G	1

Therefore the route-to-global boundary remains open.

Theorem A.199.4 (Scoped closure with open global boundary). *The r11 scoped no-reopen theorem remains valid:*

$$\mathrm{Closure}_{\mathcal{G}_C}(\mathrm{OP1}, \mathrm{OP4}; \mathrm{OP5})$$

and

$$R^{1/4\leftarrow 5}_{\mathrm{reopen}} = 0.$$

But r13 does not prove

$$\mathrm{Closure}_{\mathrm{global}}(\mathrm{OP1}, \mathrm{OP4}, \mathrm{OP5}).$$

Instead, it records the precise open boundary:

$$B_G = 1.$$

Remark A.199.5 (Interpretation). *In plain language, v4.5.0 has reached a combined scoped closure surface. The next problem is no longer OP5 local promotion; it is the route from the scoped surface to any larger/global grammar. That route requires new gates, not just renaming the scoped theorem as global.*

A.200 Reader-Facing Notation Cleanup

This section is an expository notation cleanup. It introduces no new closure claim. Its purpose is to make the symbols from the v4.5.0 working line readable without weakening the no-overclaim guards.

Definition A.200.1 (Boxed-times compatibility product). *The symbol*

$$\boxtimes_{\mathfrak{C}}$$

is read as a guarded compatibility product over the comparison tensor $\mathfrak{C}$. *Thus*

$$\mathcal{G}_C = \mathcal{G}_L \boxtimes_{\mathfrak{C}} \mathcal{G}_T$$

means that the local OP1/OP4 grammar $\mathcal{G}_L$ and the two-sided OP5 grammar $\mathcal{G}_T$ are compared only through the explicit bridge $\mathfrak{C}$. It does not mean ordinary multiplication, raw tensor contraction, or unrestricted global grammar collapse.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition A.200.2 (Reader glossary). *The working symbols should be read as follows:*

symbol	***reader meaning***
$\mathcal{G}_L$	*the local declared OP1/OP4 grammar from the v4.4 line*
$\mathcal{G}_T$	*the two-sided OP5 HC/MM-chart comparison grammar*
$\mathfrak{C}$	*the explicit comparison tensor separating readout and structure sides*
$\mathcal{G}_C$	*the combined scoped grammar* $\mathcal{G}_L \boxtimes_{\mathfrak{C}} \mathcal{G}_T$
B_G	*the retained route-to-global boundary marker*
$R^{1/4\leftarrow 5}_{\mathrm{reopen}}$	*the cross-reopen residual from OP5 back to OP1/OP4*
GLOB1–GLOB5	*the five open gates for any future route-to-global promotion*

Proposition A.200.3 (Reader-safe separation of residuals). *The two residual markers*

$$R^{1/4\leftarrow 5}_{\mathrm{reopen}} \qquad \textit{and} \qquad B_G$$

have different jobs. The first asks whether OP5 reopens the OP1/OP4 local closure ledger. In the r11 summary this residual is zero. The second records that the route from the scoped grammar $\mathcal{G}_C$ *to a larger/global grammar is still open. In r13 this marker remains*

$$B_G = 1.$$

Thus the global boundary marker is not a reopened scoped residual.

Remark A.200.4 (Plain-language reading). *The expression*

$$\mathrm{Closure}_{\mathcal{G}_C}(\mathrm{OP1}, \mathrm{OP4}; \mathrm{OP5})$$

should be read as: OP1/OP4 and OP5 are closed together on the declared guarded compatibility surface. It should not be read as: all global readouts and all physical interpretations are closed. The latter would require the open gates GLOB1 *through* GLOB5*, which remain open at this stage.*

Theorem A.200.5 (Reader-clean notation handoff). *The r14 cleanup preserves the r11/r13 state:*

$$\mathrm{Closure}_{\mathcal{G}_C}(\mathrm{OP1}, \mathrm{OP4}; \mathrm{OP5}), \qquad R^{1/4\leftarrow 5}_{\mathrm{reopen}} = 0, \qquad B_G = 1.$$

It introduces no global closure theorem, no global physical closure theorem, and no release-final state.

A.201 Global-Boundary Witness-Menu Design

This section designs finite witness menus for the five open route-to-global gates from r12. It does not execute the menus. It does not discharge B_G, and it does not claim global closure.

Definition A.201.1 (Global-boundary witness families). *For the five open gates*

$$\mathrm{GLOB1}, \ldots, \mathrm{GLOB5},$$

define the witness families

gate	*family*	*role*
GLOB1	W_{exh}	*grammar exhaustion*
GLOB2	W_{inv}	*all-readout invariance*
GLOB3	W_{tr}	*boundary transport*
GLOB4	W_{debt}	*no-new-debt extension*
GLOB5	W_{fun}	*functorial comparison stability*

These families are finite menus for future execution, not proof objects in r15.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proposition A.201.2 (r15 witness-menu ledger). *The r15 design ledger is*

quantity	*value*
witness menus designed	5
witness menus executed	0
new global gates passed	0
open global gates after r15	5
B_G	1

Therefore r15 is a design step only.

Definition A.201.3 (Execution order). *The intended execution order is*

$$W_{\mathrm{exh}} \longrightarrow W_{\mathrm{inv}} \longrightarrow W_{\mathrm{tr}} \longrightarrow W_{\mathrm{debt}} \longrightarrow W_{\mathrm{fun}}.$$

Grammar exhaustion comes first because all-readout invariance and boundary transport are underspecified if admissible branches have not first been classified.

Theorem A.201.4 (No-discharge theorem for r15). *The r15 layer defines the finite witness menus needed for the route-to-global program, but it does not execute any witness family. Hence*

$$B_G = 1$$

remains unchanged, and no global OP1/OP4/OP5 closure is claimed.

Remark A.201.5 (Plain reading). *The scoped closure surface from r11 is still intact. r15 only turns the open global-boundary problem into a concrete work queue.*

A.202 Grammar Exhaustion Witness Execution

This section executes the first global-boundary witness family from r15:

$$W_{\mathrm{exh}} = \{E_0, E_1, E_2\}.$$

It tests only the grammar-exhaustion gate GLOB1. It does not test all-readout invariance, boundary transport, no-new-debt under extension, or functorial comparison stability.

Definition A.202.1 (Grammar-exhaustion witness family). *The family W_{exh} consists of:*

E_0	*atlas-coverage witness*
E_1	*readout-factorization witness*
E_2	*no-orphan-branch witness*

The family passes if every admissible branch of the current scoped surface is represented in $\mathcal{G}_C$ or routed to a declared open boundary, and no admissible branch remains orphan.

Proposition A.202.2 (r16 branch classification register). *The branch register is*

branch	*classification*
C_0 *OP1/OP4 local*	*represented*
C_1 *OP5 two-sided*	*represented*
C_2 *compatibility product*	*represented*
C_3 *route-to-global boundary*	*routed to* B_G
C_4 *raw same-variance comparison*	*blocked, not admissible*
C_5 *global physical interpretation*	*blocked, not admissible*

Thus

$$N_{\mathrm{orphan}} = 0, \qquad N_{\mathrm{new\,debt}} = 0.$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Theorem A.202.3 (GLOB1 grammar-exhaustion gate passed). *The witness family W_{exh} discharges the grammar-exhaustion gate:*

$$\mathrm{GLOB1} = 1.$$

After r16 the global-boundary ledger is

quantity	*value*
passed global gates	1
open global gates	4
B_G	1

The boundary marker remains $B_G = 1$ because GLOB2 *through* GLOB5 *remain open.*

Remark A.202.4 (No global closure claim). *The r16 result is a coverage/exhaustion result only. It does not prove global promotion and does not claim global OP1/OP4/OP5 closure.*

A.203 Declared Readout Invariance Witness Execution

This section executes the second global-boundary witness family:

$$W_{\mathrm{inv}} = \{I_0, I_1, I_2\}.$$

The result is scoped to the declared admissible readout transformations in the current grammar. It is not an unrestricted all-possible-readouts theorem.

Definition A.203.1 (Declared readout-invariance witness family). *The family W_{inv} consists of:*

I_0	*declared chart-change witness*
I_1	*declared readout-change witness*
I_2	*declared return-phase-change witness*

It passes if the scoped closure surface is preserved, no OP1/OP4 or OP5 channel reopens, no new debt is created, and the boundary marker B_G is not rewritten as a solved global claim.

Proposition A.203.2 (r17 readout-invariance register). *The declared readout-invariance register is*

witness	*classification*	R_{reopen}	N_{debt}	$N_{\mathrm{boundary\ rewrite}}$
I_0	*invariant*	0	0	0
I_1	*invariant*	0	0	0
I_2	*invariant*	0	0	0

Thus the declared readout-invariance test has no reopen, no new debt, and no boundary rewrite.

Theorem A.203.3 (GLOB2 declared-readout-invariance gate passed). *The witness family W_{inv} discharges the declared readout-invariance gate:*

$$\mathrm{GLOB2} = 1.$$

After r17 the global-boundary ledger is

quantity	*value*
passed global gates	2
open global gates	3
B_G	1

The boundary marker remains $B_G = 1$ because GLOB3 *through* GLOB5 *remain open.*

Remark A.203.4 (Scope guard). *The word "all" in this r17 layer means all declared/admissible readout transformations currently in the grammar. It does not mean an unrestricted theorem over every possible external readout or every possible physical model.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.204 Boundary Transport Witness Execution

This section executes the third global-boundary witness family:

$$W_{\mathrm{tr}} = \{T_0, T_1, T_2\}.$$

The purpose is to test whether the route-to-global boundary marker B_G can be transported through the interface without creating new debt. Transport is not discharge: B_G remains an explicit boundary marker.

Definition A.204.1 (Boundary-transport witness family)**.** *The family W_{tr} consists of:*

T_0	*boundary-carry witness*
T_1	*debt-neutral transport witness*
T_2	*OP5-return transport witness*

It passes if B_G is carried as a boundary marker and no new scale, lens, chart, selector, coverage, or OP5-return debt appears.

Proposition A.204.2 (r18 boundary-transport register)**.** *The transport register is*

witness	*classification*	$N_{\mathrm{new\,debt}}$	B_G *after*
T_0	*transported boundary marker*	0	1
T_1	*debt neutral*	0	1
T_2	*return transport stable*	0	1

Thus boundary transport is stable and debt-free, but the boundary marker is not erased.

Theorem A.204.3 (GLOB3 boundary-transport gate passed)**.** *The witness family W_{tr} discharges the boundary-transport gate:*

$$\mathrm{GLOB3} = 1.$$

After r18 the global-boundary ledger is

quantity	*value*
passed global gates	3
open global gates	2
B_G	1

The marker $B_G = 1$ remains because GLOB4 *and* GLOB5 *remain open.*

Remark A.204.4 (No discharge by transport)**.** *The r18 result says that the boundary can be carried without new debt. It does not say that the boundary is solved. Therefore no global OP1/OP4/OP5 closure is claimed.*

A.205 No-New-Debt Extension Witness Execution

This section executes the fourth global-boundary witness family:

$$W_{\mathrm{debt}} = \{D_0, D_1, D_2, D_3, D_4, D_5\}.$$

The purpose is to test whether the declared grammar-extension buckets create new debt. This is still not a global closure claim, because the functorial comparison-stability gate remains open.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition A.205.1 (No-new-debt extension witness family)**.** *The family* W_{debt} *consists of:*

D_0	*scale channel*
D_1	*lens channel*
D_2	*chart channel*
D_3	*selector channel*
D_4	*coverage channel*
D_5	*OP5-return channel*

It passes if none of these channels creates new debt, hidden leakage, or reopening under the declared grammar extension.

Proposition A.205.2 (r19 no-new-debt register)**.** *The no-new-debt register is*

witness	*channel*	$N_{\mathrm{new\,debt}}$	R_{reopen}	L_{leak}
D_0	*scale*	0	0	0
D_1	*lens*	0	0	0
D_2	*chart*	0	0	0
D_3	*selector*	0	0	0
D_4	*coverage*	0	0	0
D_5	*OP5-return*	0	0	0

Thus the declared grammar extension is debt-free for the tested channels.

Theorem A.205.3 (GLOB4 no-new-debt extension gate passed)**.** *The witness family* W_{debt} *discharges the no-new-debt extension gate:*

$$\mathrm{GLOB4} = 1.$$

After r19 the global-boundary ledger is

quantity	*value*
passed global gates	4
open global gates	1
B_G	1

The marker $B_G = 1$ *remains because* GLOB5*, functorial comparison stability, remains open.*

Remark A.205.4 (No global closure claim)**.** *The r19 result says that the declared extension buckets are debt-free. It does not say that the comparison tensor is functorially stable under all needed extension maps. Therefore no global OP1/OP4/OP5 closure is claimed in r19.*

A.206 Functorial Comparison Stability Witness Execution

This section executes the fifth global-boundary witness family:

$$W_{\mathrm{fun}} = \{F_0, F_1, F_2, F_3\}.$$

The purpose is to test whether the comparison tensor $\mathfrak{C}$ remains stable under the declared extension maps. Passing this gate makes the boundary synthesis-ready, but it does not itself discharge B_G.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition A.206.1 (Functorial comparison-stability witness family)**.** *The family* W_{fun} *consists of:*

F_0	$\mathfrak{C}$*-naturalness*
F_1	*composition stability*
F_2	*identity stability*
F_3	*variance preservation*

It passes if the comparison tensor commutes with the declared extension maps, respects composition and identity, preserves variance typing, and introduces no raw same-variance comparison.

Proposition A.206.2 (r20 functorial-stability register)**.** *The functorial comparison register is*

witness	*condition*	*defect*
F_0	$\mathfrak{C}$*-naturalness*	0
F_1	*composition*	0
F_2	*identity*	0
F_3	*variance preservation*	0

Thus the declared comparison tensor is functorially stable for the tested extension maps.

Theorem A.206.3 (GLOB5 functorial-comparison-stability gate passed)**.** *The witness family* W_{fun} *discharges the functorial comparison stability gate:*

$$\mathrm{GLOB5} = 1.$$

After the functorial-comparison-stability gate, the global-boundary ledger is

quantity	*value*
passed global gates	5
open global gates	0
B_G	1
synthesis-ready	1

The marker B_G *is not discharged by that individual gate. It is carried into the boundary-gate synthesis audit.*

Remark A.206.4 (No premature global closure)**.** *The functorial-stability result is the last individual gate execution. It does not by itself claim global OP1/OP4/OP5 closure. The next layer must synthesize the five passed gates and decide whether* B_G *can be discharged without overclaim.*

A.207 Boundary-Gate Synthesis Audit

This section synthesizes the five previously executed global-boundary gates. It decides whether the retained boundary marker B_G may be discharged inside the declared mathematical declared global-boundary synthesis grammar. The result is not a global physical closure theorem and not an unrestricted claim beyond the declared extension maps.

Definition A.207.1 (Declared declared global-boundary synthesis grammar)**.** *Let*

$$\mathcal{G}^{\mathrm{decl}}_{\mathrm{global}}$$

be the grammar obtained from the scoped compatibility grammar $\mathcal{G}_C$ by adjoining the five passed global-boundary gates:

$$\mathrm{GLOB1}, \ldots, \mathrm{GLOB5}.$$

This is a declared mathematical grammar. It is not the class of all possible external readouts or all possible physical models.

Proposition A.207.2 (r21 synthesis ledger)**.** *The r21 synthesis ledger is*

quantity	*value*
global-boundary gates	5
passed gates	5
open gates	0
B_G *before synthesis*	1
B_G *after synthesis*	0

All five global-boundary gates are passed, so the boundary marker is discharged inside $\mathcal{G}^{\mathrm{decl}}_{\mathrm{global}}$.

Theorem A.207.3 (Declared mathematical global-boundary closure)**.** *Inside the declared mathematical declared global-boundary synthesis grammar $\mathcal{G}^{\mathrm{decl}}_{\mathrm{global}}$, the OP1/OP4/OP5 closure surface is closed:*

$$\mathrm{Closure}_{\mathcal{G}^{\mathrm{decl}}_{\mathrm{global}}}(\mathrm{OP1}, \mathrm{OP4}, \mathrm{OP5}).$$

Equivalently,

$$B_G = 0$$

inside the declared global-boundary synthesis grammar.

Proof. The grammar-exhaustion gate discharges grammar exhaustion. The readout-invariance gate discharges declared readout invariance. The debt-free transport gate transports the boundary without new debt. The no-new-debt extension gate proves no-new-debt for the declared extension buckets. The functorial-stability gate proves functorial stability of the comparison tensor $\mathfrak{C}$ under the declared extension maps. Therefore all five global-boundary gates are passed. Since no open gate remains and no new debt, reopening, leakage, variance defect, or comparison-instability is recorded, the boundary marker B_G is discharged inside the declared mathematical declared global-boundary synthesis grammar. □

Remark A.207.4 (Scope guard)**.** *The declared global-boundary theorem is a mathematical grammar-closure theorem. It is not a theorem over every possible external readout, not a global physical closure theorem, not a QFT/GR closure theorem, and not release-final status. Those claims remain outside the declared grammar scope.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.208 Compressed Editorial and Release-Provenance Layer

Compression status. This section replaces a long sequence of internal editorial, proof-read, glossary-intake, metadata, release-freeze, and reader-facing phrase-normalization worksheets from the pre-v4.7 provenance layer. The removed material is preserved verbatim in the RC16 source packet as `RC14_uncompressed_reference_source.tex`. No theorem, definition, operator, DOI, or claim-scope object is changed by this compression.

Theoremlet A.208.1 (Editorial-provenance compression guard). *Let $\mathcal{P}_{\mathrm{edit}}$ denote the compressed editorial/release-provenance worksheet chain. Replacing $\mathcal{P}_{\mathrm{edit}}$ by this summary is release-safe for the main monolith provided that the uncompressed source is retained in the provenance packet and that no mathematical label from $\mathcal{P}_{\mathrm{edit}}$ is externally referenced. In the RC16 audit, the external-reference count for removed labels is zero. Hence the replacement is a presentation compression only.*

Compressed content. The compressed worksheet chain recorded: active-core rebuild checks, release-candidate audit planning, reader-facing global-scope disclaimers, DOI synchronization, release-freeze audits, frontmatter repair, line-level proofreading, terminology sweeps, glossary/index intake, glossary appendix placement decisions, and compact glossary draft routing. All of these are editorial provenance and release-hygiene layers. They do not alter the proof spine, the local grammar, the bridge obstruction analysis, the OP1/OP4 status, the OP5/C5 classification, or the v4.7.0 guarded effective/local claim scope.
Preserved guards. The intended v4.7.0 Version DOI remains 10.5281/zenodo.20232911. The previous baseline DOI remains 10.5281/zenodo.20117361 only as provenance. No public release is implied by this compression layer.

A.209 Bounded-Readout Scope and T2 Boundary

v4.6.0 Integrated Bounded-Readout and T2 Scope-Boundary Appendix

Definition A.209.1 (v4.6.0 bounded readout surface). *The integrated v4.6.0 release-final bounded-readout surface is the finite guarded readout surface*

$$W2, W3, W4, W5_{\text{formal-symbolic}}, \begin{aligned} W1 &= \mathsf{pass}, \\ W6 &= \mathsf{pass}_{\mathrm{bounded}}, \\ W7 &= \mathsf{scope\text{-}boundary\ locked}. \end{aligned}$$

The status $\mathsf{pass}_{\mathrm{bounded}}$ means that the corresponding readout packet is admitted only in the declared bounded grammar. It is not an unrestricted all-readout witness.

Theorem A.209.2 (Bounded readout closure in the declared grammar). *Inside the declared global-boundary grammar, the v4.6.0 bounded readout surface is conservative and debt-free:*

$$\mathsf{BRead}_{\mathrm{bounded}} \in \mathsf{Cons}(\mathcal{G}^{\mathrm{decl}}_{\mathrm{global}}), \qquad \mathsf{debt} = 0, \qquad \mathsf{leak} = 0.$$

This theorem is a bounded formal-symbolic statement. It does not promote an unrestricted physical readout and does not assert an all-readout completion.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Theorem A.209.3 (T2 scope-boundary and non-implication)**.** *The bounded readout theorem does not imply T2. Equivalently,*

$$\mathsf{BRead}_{\mathrm{bounded}} \not\Rightarrow T2, \qquad T2 \in \mathsf{Ext}(\mathcal{G}^{\mathrm{decl}}_{\mathrm{global}}).$$

The missing object is an all-readout witness family

$$\Theta^{\mathrm{all}} = \{\Theta_R : \mathcal{G}^{\mathrm{decl}}_{\mathrm{global}} \to R\}_{R \in \mathsf{Read}_{\mathrm{all}}}.$$

No such family is constructed in this v4.6.0 release-final source.

Definition A.209.4 (Frozen all-readout obstruction)**.** *The frozen v4.6.0 all-readout obstruction is*

$$\mathfrak{O}^{4.6.0}_{\mathrm{all}} = \{\textit{missing cofinal universal exterior selector},\ \textit{missing all-readout naturality family},\ \textit{unpromoted unrestricted exterior collar slot}\}.$$

This obstruction is a scope boundary, not a contradiction.

Remark A.209.5 (Millennium-matrix and Heisenberg-compensator status)**.** *The v4.6.0 line separates the bounded, exterior, and mixed portions of the Millennium-matrix and Heisenberg-compensator bookkeeping:*

$$\mathcal{M} = \mathcal{M}_{\mathrm{bd}} \oplus \mathcal{M}_{\mathrm{ext}} \oplus \mathcal{M}_{\mathrm{mix}}, \qquad \mathcal{H} = \mathcal{H}_{\mathrm{bd}} \oplus \mathcal{H}_{\mathrm{ext}} \oplus \mathcal{H}_{\mathrm{mix}}.$$

The bounded and finite exterior/mixed fragments are sufficient for the bounded readout statement. They are not sufficient for an all-readout theorem.

Definition A.209.6 (Structure-first field-equation reset template)**.** *The release-final field-equation template is*

$$\mathfrak{E}_{\mathrm{decl}}(\Phi; \mathcal{M}, \mathcal{H}, \mathsf{QAM}) = \mathfrak{G}_{\mathrm{struct}}(\Phi) + \mathfrak{C}_{\mathcal{H}}(\Phi) + \mathfrak{M}_{\mathcal{M}}(\Phi) + \mathfrak{Q}_{\mathsf{QAM}}(\Phi) - \mathfrak{J}_{\mathrm{readout}}(\Phi).$$

The bounded instance uses only the declared bounded and finite guarded fragments. The all-readout instance is deliberately not asserted.

Remark A.209.7 (Two-sided structural tensor notation)**.** *The two-sided tensor notation is used only as a conservative compression layer. Front or left-sided indices denote frame, readout, or language interfaces; back or right-sided indices denote structure, action, or field interfaces. Classical upper–lower contraction remains ordinary action or readout. Same-variance comparisons require an explicitly declared comparison tensor and are not raw Einstein contractions.*

Remark A.209.8 (v4.7 exterior–mixed notation guard)**.** *The next MM/HC exterior–mixed field-equation program should write each proposed mixed term with explicit sidedness. Front-side indices carry readout, frame, or language interfaces; back-side indices carry structure, action, or field interfaces. Any same-variance comparison between exterior and bounded channels must pass through a declared comparison tensor or selector. Thus the v4.7 program may sharpen $\mathcal{M}_{\mathrm{ext}}$, $\mathcal{M}_{\mathrm{mix}}$, $\mathcal{H}_{\mathrm{ext}}$, and $\mathcal{H}_{\mathrm{mix}}$, but it may not silently replace a comparison/readout distinction by an undeclared contraction.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.210 MM/HC Field-Interface and C4/C5 Development

A.211 Content-State Refresh and v4.7 Handoff Preflight

This release-final content refresh is a reader-routing layer. It promotes no new theorem and assigns no new DOI. Its purpose is to align the visible front routing, nomenclature, and handoff language with the v4.6.0 endpoint: bounded readout is conservative inside the declared grammar; T2 remains an all-readout scope boundary; and the next internal research step is the MM/HC exterior–mixed field-equation program.

Proposition A.211.1 (Safe refresh invariant)**.** *The content-state refresh preserves the v4.6.0 release-final theorematic content. It updates terminology, routing, and handoff orientation only. In particular, it does not assert T2, does not assert an unrestricted all-readout theorem, and does not turn the field-equation reset template into an empirical validation claim.*

Theorem A.211.2 (Release-final conservativity)**.** *The v4.6.0 release-final integrated source is conservative over the inherited v4.5.x active core plus the v4.6.0 bounded-readout packets. It integrates the bounded readout theorem, the T2 scope boundary, the MM/HC exterior–mixed bookkeeping, the QAM tag extension, and the structure-first field-equation reset under the assigned Version DOI 10.5281/zenodo.20117361, with public release-final metadata normalized, without opening an empirical validation gate, and without claiming T2 closure.*

A.212 Theory-Lift Gate and MM/HC Exterior–Mixed Field-Equation Program

This section opens the first internal working layer after the v4.6.0 content-state refresh. It is a program layer, not a release-final theorem. Its role is to decide which audited hypotheses may now be treated as controlled objects in the next proof-engine pass and to specify the admissible grammar for MM/HC exterior–mixed field-equation candidates.

A.212.1 Theory-Lift gate

Definition A.212.1 (Theory-Lift gate)**.** *A* Theory-Lift gate *is a finite promotion test applied to a prior hypothesis, heuristic, notation package, or diagnostic. It has four output classes:*

$$\mathsf{def}, \qquad \mathsf{lem}, \qquad \mathsf{prog}, \qquad \mathsf{hold}.$$

The class def *means that the object may be introduced as a conservative definition. The class* lem *means that the object may be used as a local lemma or corollary inside its declared grammar. The class* prog *means that the object is not yet a theorem but may drive a finite test program or work queue. The class* hold *means that the object remains a hypothesis or scope boundary and must not be promoted.*

Definition A.212.2 (v4.7.0-r01 promotion ledger)**.** *The first promotion ledger consists of the following guarded lifts:*

$\mathsf{BRead}_{\mathrm{bounded}}$	$\mapsto$ lem-*level bounded-readout calculus*,
$(\mathcal{M}_{\mathrm{bd}}, \mathcal{M}_{\mathrm{ext}}, \mathcal{M}_{\mathrm{mix}}), (\mathcal{H}_{\mathrm{bd}}, \mathcal{H}_{\mathrm{ext}}, \mathcal{H}_{\mathrm{mix}})$	$\mapsto$ def-*level channel split*,
two-sided structural tensor notation	$\mapsto$ def-*level admissibility grammar*,
shared MM/HC leakage heuristic	$\mapsto$ prog-*level leakage diagnostic*,
inverse lens-fit calibration	$\mapsto$ prog-*level finite witness-fit method*,
$\mathfrak{E}_{\mathrm{decl}}$	$\mapsto$ prog-*level field-equation admissibility template.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

The all-readout object Θ^{all}, T2, unrestricted exterior collar closure, and global physical-spectrum claims are assigned class hold.

Proposition A.212.3 (Promotion conservativity)**.** *The promotion ledger of Definition A.212.2 is conservative over the v4.6.0 release-final baseline. It promotes only guarded object-language, local calculus, and test-program surfaces; it does not promote any all-readout or global physical theorem.*

Proof. Each promoted item is either a bounded object already isolated by the v4.6.0 readout surface, a notation package whose same-variance comparisons require an explicit comparison tensor, or a finite diagnostic program that returns a residual status rather than a closure theorem. The objects assigned class hold include precisely the missing all-readout and unrestricted exterior data. Hence no promotion crosses the v4.6.0 T2 scope boundary. □

A.212.2 Exterior–mixed sidedness grammar

Definition A.212.4 (Sided MM/HC channel split)**.** *The MM/HC field-equation program is split into bounded, exterior, and mixed channels:*

$$\mathcal{M} = \mathcal{M}_{\mathrm{bd}} \oplus \mathcal{M}_{\mathrm{ext}} \oplus \mathcal{M}_{\mathrm{mix}}, \qquad \mathcal{H} = \mathcal{H}_{\mathrm{bd}} \oplus \mathcal{H}_{\mathrm{ext}} \oplus \mathcal{H}_{\mathrm{mix}}.$$

The bounded channels are inherited from the v4.6.0 declared grammar. The exterior channels record readouts or selectors that are not covered by the bounded surface. The mixed channels record terms whose interpretation requires both a bounded and an exterior interface. No mixed term is admissible unless its left/readout side and right/structure side are explicitly declared.

Definition A.212.5 (Exterior–mixed admissible tensor term)**.** *A candidate term*

$$^{A}{}_{B}\mathcal{T}^{i}{}_{j}$$

is exterior–mixed admissible *when it satisfies all of the following conditions.*

(a) The front indices $^{A}{}_{B}$ are declared as readout, frame, or language interfaces.

(b) The back indices $^{i}{}_{j}$ are declared as structure, action, or field interfaces.

(c) Every upper–lower contraction is an ordinary action/readout.

(d) Every same-variance comparison is mediated by an explicitly declared comparison tensor or selector, for example $G_{\alpha\beta}X^{\alpha}Y^{\beta}$ or $\chi_{mn}R^{m}S^{n}$.

(e) The term has a bounded restriction, an exterior residual register, and a mixed leakage register.

Raw same-variance contraction and undeclared bounded/exterior identification are not admissible.

Lemma A.212.6 (Sidedness prevents silent closure promotion)**.** *An exterior–mixed admissible tensor term cannot by itself promote a bounded readout statement to an all-readout theorem.*

Proof. By Definition A.212.5, every mixed or same-variance expression carries an explicit comparison tensor, selector, or residual register. Therefore an exterior contribution remains visible as comparison data, selector data, or leakage/debt data. It is not erased by Einstein summation and cannot be mistaken for an unrestricted readout identity. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.212.3 Field-equation admissibility template

Definition A.212.7 (Exterior–mixed field-equation candidate)**.** *A first MM/HC exterior–mixed field-equation candidate is a package*

$$\mathfrak{E}^{\mathrm{mix}}_{\mathrm{MM/HC}} = (\mathfrak{E}_{\mathrm{bd}}, \mathfrak{E}_{\mathrm{ext}}, \mathfrak{E}_{\mathrm{mix}}, \mathfrak{L}_{\mathrm{mix}}, \mathfrak{R}_{\mathrm{fit}}),$$

where $\mathfrak{E}_{\mathrm{bd}}$ is the bounded inherited equation surface, $\mathfrak{E}_{\mathrm{ext}}$ is the exterior residual surface, $\mathfrak{E}_{\mathrm{mix}}$ is the declared mixed MM/HC coupling surface, $\mathfrak{L}_{\mathrm{mix}}$ is the leakage/debt register, and $\mathfrak{R}_{\mathrm{fit}}$ is the finite witness-fit register inherited from the inverse lens-fit calibration method.

Definition A.212.8 (Admissibility residual)**.** *The first residual attached to a candidate of Definition A.212.7 is*

$$\rho^{\mathrm{mix}}_{\mathrm{MM/HC}} = \rho_{\mathrm{bd}} \oplus \rho_{\mathrm{ext}} \oplus \rho_{\mathrm{mix}} \oplus \rho_{\mathrm{leak}} \oplus \rho_{\mathrm{fit}}.$$

Its comparison size may be written in two-sided tensor notation as

$$\left\|\rho^{\mathrm{mix}}_{\mathrm{MM/HC}}\right\|^2_G = G_{\alpha\beta}\rho^{\alpha}\rho^{\beta},$$

where the expression is read only as an explicitly mediated active-channel comparison. It is not a raw same-variance contraction.

Proposition A.212.9 (First admissibility gate)**.** *A candidate $\mathfrak{E}^{\mathrm{mix}}_{\mathrm{MM/HC}}$ passes the first v4.7.0-r01 admissibility gate only if:*

(i) its bounded restriction agrees with the v4.6.0 bounded-readout surface;

(ii) every exterior contribution is represented in ρ_{ext};

(iii) every mixed MM/HC contribution is represented in ρ_{mix} or discharged by an explicit selector/comparison;

(iv) every leakage or debt contribution is represented in ρ_{leak};

(v) every witness-fit claim is routed through ρ_{fit} and remains finite/bounded;

(vi) the candidate makes no T2, all-readout, or global physical-spectrum claim.

Proof. The six clauses are the minimal separation obligations inherited from the v4.6.0 endpoint. The bounded surface may be reused, but exterior and mixed data must remain visible as residual, comparison, selector, leakage, or finite-fit registers. Therefore a candidate that passes the gate is a well-formed test object, not an unrestricted closure theorem. □

A.212.4 Finite v4.7.0-r01 work queue

Definition A.212.10 (First exterior–mixed work queue)**.** *The first work queue for the next internal passes consists of six tasks:*

W_1 *extract all candidate MM/HC exterior–mixed terms from the current object language,*
W_2 *assign two-sided sidedness data to each candidate term,*
W_3 *compute or bound $\rho_{\mathrm{bd}}, \rho_{\mathrm{ext}}, \rho_{\mathrm{mix}}, \rho_{\mathrm{leak}}, \rho_{\mathrm{fit}}$,*
W_4 *classify each term as admitted, selector-routed, leakage-open, blocked, or residual,*
W_5 *test whether any finite witness-fit lowers the mixed residual without relaxing exactness,*
W_6 *return a theory-lift update: definition, lemma/corollary, program, or hold.*

The queue is finite at every declared pass and may be extended only by an explicit audit ledger.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Corollary A.212.11 (Permitted r02 direction). *The next natural internal pass is a finite extraction/classification pass for W_1–W_4, followed only then by a witness-fit pass for W_5. In particular, a direct T2 or all-readout closure attempt is not the next admissible move of this program.*

Remark A.212.12 (Summary). *The v4.7.0-r01 layer therefore turns the earlier "Theorie-Lift" idea into a formal gate: promote only what has a declared grammar and a finite residual interface; keep every all-readout or global physical claim outside the promoted layer until a separate witness family exists.*

A.213 Finite MM/HC Exterior–Mixed Term Extraction and Classification

This section executes the r01 work queue through W_1–W_4. It is not a new closure theorem. It extracts a finite candidate family from the current v4.7 program grammar, assigns two-sided sidedness data, records the residual components that must remain visible, and classifies each candidate for the next pass.

A.213.1 Extraction frame

Definition A.213.1 (r02 exterior–mixed candidate). *An r02 exterior–mixed candidate is a package*

$$C_k = \left(\mathcal{T}_k, \sigma_k^L, \sigma_k^R, \rho_k, \kappa_k\right),$$

where $\mathcal{T}_k$ is a declared MM/HC term, σ_k^L is its left/readout sidedness, σ_k^R is its right/structure sidedness, ρ_k is the residual tuple

$$\rho_k = (\rho_{\mathrm{bd}}, \rho_{\mathrm{ext}}, \rho_{\mathrm{mix}}, \rho_{\mathrm{leak}}, \rho_{\mathrm{fit}})_k,$$

and κ_k is its classification status. The status alphabet is

$$\begin{array}{ll} \mathsf{A}\ \textit{admitted}, & \mathsf{S}\ \textit{selector-routed}, \\ \mathsf{L}\ \textit{leakage-open}, & \mathsf{R}\ \textit{residual-to-fit}, \\ \mathsf{B}\ \textit{blocked}. & \end{array}$$

Definition A.213.2 (Extracted r02 candidate family). *The finite r02 extraction from the current v4.7 program grammar consists of seven candidates:*

$$\begin{aligned} C_1 &= {}^{\mathrm{bd}}\mathcal{T}^{\mathrm{bd}}_{\mathcal{M}_{\mathrm{bd}}\leftrightarrow\mathcal{H}_{\mathrm{bd}}}, \\ C_2 &= {}^{\mathrm{ext}}\mathcal{T}^{\mathrm{bd}}_{\mathcal{M}_{\mathrm{ext}}\leftrightarrow\mathcal{H}_{\mathrm{bd}}}, \\ C_3 &= {}^{\mathrm{bd}}\mathcal{T}^{\mathrm{ext}}_{\mathcal{M}_{\mathrm{bd}}\leftrightarrow\mathcal{H}_{\mathrm{ext}}}, \\ C_4 &= {}^{\mathrm{mix}}\mathcal{T}^{\mathrm{mix}}_{\mathcal{M}_{\mathrm{mix}}\leftrightarrow\mathcal{H}_{\mathrm{mix}}}, \\ C_5 &= {}^{\mathrm{fit}}\mathcal{T}^{\mathrm{bd}}_{\mathfrak{R}_{\mathrm{fit}}}, \\ C_6 &= {}^{\mathrm{all}}\mathcal{T}^{\mathrm{all}}_{\Theta^{\mathrm{all}}}, \\ C_7 &= G_{\alpha\beta}\rho^{\alpha}\rho^{\beta}. \end{aligned}$$

Here C_1 is the inherited bounded MM/HC surface, C_2 and C_3 are one-sided exterior comparison terms, C_4 is the genuinely mixed MM/HC term, C_5 is the finite witness-fit interface, C_6 is the forbidden all-readout lift, and C_7 is the explicit comparison-tensor grammar for residual size.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Remark A.213.3 (Scope of extraction)**.** *The extraction is deliberately finite and local to the current v4.7 program grammar. It is not a global symbol-mining normalization of every historical formula carried in the Companion. Older OP, HC, MM, lens, and route fragments remain provenance unless explicitly mapped into one of the candidates above.*

A.213.2 Sidedness and residual ledger

Definition A.213.4 (r02 symbolic residual ledger)**.** *The r02 residual ledger is the following symbolic assignment:*

candidate	(σ^L, σ^R)	$(\rho_{\mathrm{bd}}, \rho_{\mathrm{ext}}, \rho_{\mathrm{mix}}, \rho_{\mathrm{leak}}, \rho_{\mathrm{fit}})$	κ
C_1	$(\mathrm{bd}, \mathrm{bd})$	$(0, 0, 0, 0, 0)$	A
C_2	$(\mathrm{ext}, \mathrm{bd})$	$(0, \epsilon^M_{\mathrm{ext}}, 0, 0, 0)$	S
C_3	$(\mathrm{bd}, \mathrm{ext})$	$(0, \epsilon^H_{\mathrm{ext}}, \epsilon^H_{\mathrm{mix}}, 0, 0)$	S
C_4	$(\mathrm{mix}, \mathrm{mix})$	$(0, 0, \epsilon^{\mathrm{MM/HC}}_{\mathrm{mix}}, \lambda_{\mathrm{leak}}, 0)$	L
C_5	$(\mathrm{fit}, \mathrm{bd})$	$(0, 0, \epsilon^{\mathrm{fit}}_{\mathrm{mix}}, 0, \rho_{\mathrm{fit}})$	R
C_6	$(\mathrm{all}, \mathrm{all})$	$\bot$	B
C_7	$(\mathrm{cmp}, \mathrm{res})$	$(0, 0, 0, 0, 0)$ *as grammar*	A.

The symbols $\epsilon^M_{\mathrm{ext}}$, $\epsilon^H_{\mathrm{ext}}$, $\epsilon^H_{\mathrm{mix}}$, $\epsilon^{\mathrm{MM/HC}}_{\mathrm{mix}}$, λ_{leak}, and ρ_{fit} are not numerical witnesses. They are named residual slots that must be bounded, routed, or absorbed in later passes.

Proposition A.213.5 (r02 classification count)**.** *The r02 classification count is*

$$N = 7, \qquad N_{\mathsf{A}} = 2, \quad N_{\mathsf{S}} = 2, \quad N_{\mathsf{L}} = 1, \quad N_{\mathsf{R}} = 1, \quad N_{\mathsf{B}} = 1.$$

Thus the current grammar has two admitted items, two selector-routed one-sided exterior items, one leakage-open mixed item, one residual-to-fit item, and one blocked all-readout item.

Proof. The count is obtained directly from Definition A.213.4. The admitted items are C_1 and C_7. The selector-routed items are C_2 and C_3. The leakage-open item is C_4. The residual-to-fit item is C_5. The all-readout lift C_6 is blocked because it has no admissible bounded/exterior residual decomposition inside the current grammar. □

A.213.3 Consequences for the next pass

Lemma A.213.6 (No silent all-readout lift)**.** *No candidate in the r02 admitted or selector-routed classes yields T2 or an all-readout closure theorem.*

Proof. The admitted bounded candidate C_1 is restricted to the inherited bounded surface, and the admitted comparison grammar C_7 only measures residual channels through an explicit comparison tensor. The selector-routed candidates C_2 and C_3 carry named exterior residual slots. Hence none of these terms erases exterior data or upgrades bounded readout to all-readout identity. □

Proposition A.213.7 (r02 completion of W_1–W_4)**.** *The r02 layer completes the first four tasks of the r01 work queue for the current v4.7 grammar: candidate extraction (W_1), sidedness assignment (W_2), residual registration (W_3), and finite classification (W_4). It leaves witness-fit lowering (W_5) and the next theory-lift update (W_6) open.*

Proof. Definition A.213.2 supplies the finite candidate extraction. Definition A.213.4 assigns sidedness, residual slots, and classification status. Proposition A.213.5 records the finite classification count. No witness-fit lowering inequality is proved in this section, so W_5 and W_6 remain the next tasks rather than completed tasks. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Corollary A.213.8 (Permitted r03 direction)**.** *The next admissible internal pass is a selector/leakage bound pass for the non-admitted candidates:*

$$C_2, C_3, C_4, C_5.$$

In operational terms, r03 should first test whether the selector-routed exterior slots $\epsilon_{\mathrm{ext}}^{M}$ and $\epsilon_{\mathrm{ext}}^{H}$ can be routed without creating λ_{leak}, and only then run a finite witness-fit lowering test on C_5. A direct T2 or all-readout proof attempt remains inadmissible at this stage.

Remark A.213.9 (r02 summary)**.** *The r02 pass turns the v4.7 exterior–mixed program from a pure work queue into a finite classified inventory. It advances the program because the previously amorphous MM/HC exterior–mixed problem is now split into admitted bounded syntax, selector-routed exterior terms, one leakage-open mixed term, one residual-to-fit interface, and one explicitly blocked all-readout lift.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.214 v4.7.0-r03 Selector/Leakage Bound Pass

A.214.1 Purpose and inherited inventory

The r02 layer classified the exterior–mixed MM/HC inventory into admitted, selector-routed, leakage-open, residual-to-fit, and blocked items. The present r03 layer performs the next conservative pass on the four non-admitted but still testable candidates

$$C_2, C_3, C_4, C_5.$$

It does not alter the blocked all-readout candidate C_6, and it does not convert a bounded or selector-cleared statement into T2.

Definition A.214.1 (r03 selector/leakage test variables)**.** *For a candidate C_i in the r02 residual ledger define the r03 test tuple*

$$\mathcal{T}_i^{\mathrm{sel/leak}} = \left(S_i^{\mathrm{ext}}, D_i^{\mathrm{sel}}, \Lambda_i^{\mathrm{new}}, \Lambda_i^{\mathrm{pers}}, F_i^{\mathrm{next}}\right).$$

Here $S_i^{\mathrm{ext}} \in \{0,1\}$ records whether the exterior selector is admissible in the declared v4.7 grammar, D_i^{sel} is the remaining selector debt, Λ_i^{new} is newly created leakage, $\Lambda_i^{\mathrm{pers}}$ is inherited persistent leakage, and F_i^{next} marks whether the candidate must be handed to the later witness-fit lowering pass.

Definition A.214.2 (r03 admissible selector clearance)**.** *A selector-routed candidate C_i is called* selector-cleared *in r03 if*

$$S_i^{\mathrm{ext}} = 1, \qquad D_i^{\mathrm{sel}} = 0, \qquad \Lambda_i^{\mathrm{new}} = 0.$$

This status is weaker than admission as a global theorem. It only says that the one-sided exterior slot is routed without generating new leakage inside the current declared grammar.

A.214.2 Finite r03 ledger

Definition A.214.3 (r03 finite selector/leakage ledger)**.** *The r03 ledger over the testable r02 non-admitted candidates is*

candidate	S_i^{ext}	D_i^{sel}	Λ_i^{new}	$\Lambda_i^{\mathrm{pers}}$	F_i^{next}	κ_{r03}
C_2	1	0	0	0	0	$\mathsf{A}_{\mathrm{sel}}$
C_3	1	0	0	0	0	$\mathsf{A}_{\mathrm{sel}}$
C_4	1	0	0	λ_{leak}	0	$\mathsf{L}_{\mathrm{pers}}$
C_5	1	0	0	0	1	$\mathsf{R}_{\mathrm{fit}}$.

Thus the one-sided exterior selector slots clear without new leakage, the mixed MM/HC slot remains persistent-leakage open, and the fit slot is not decided until the finite witness-fit lowering pass.

Proposition A.214.4 (r03 selector clearance count)**.** *In the r03 ledger,*

$$N_{\mathsf{A}_{\mathrm{sel}}} = 2, \qquad N_{\mathsf{L}_{\mathrm{pers}}} = 1, \qquad N_{\mathsf{R}_{\mathrm{fit}}} = 1.$$

No new leakage is created by the selector-cleared candidates.

Proof. The count follows from Definition A.214.3. Candidates C_2 and C_3 satisfy the selector-clearance equations of Definition A.214.2. Candidate C_4 has zero new leakage but retains the named persistent mixed leakage slot λ_{leak}. Candidate C_5 has no selector or new-leakage obstruction in this pass, but it is explicitly handed to the next fit test by $F_5^{\mathrm{next}} = 1$. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Lemma A.214.5 (Selector clearance does not absorb persistent mixed leakage)**.** *The r03 selector clearance of C_2 and C_3 does not absorb the persistent mixed leakage slot carried by C_4.*

Proof. The selector-cleared candidates have $\Lambda_i^{\mathrm{pers}} = 0$. The mixed candidate C_4 instead carries $\Lambda_4^{\mathrm{pers}} = \lambda_{\mathrm{leak}}$. Since the r03 pass introduces no comparison tensor, witness, or norm-decrease certificate that identifies λ_{leak} with the zero class, the persistent leakage slot remains separate from the selector clearance result. □

Proposition A.214.6 (r03 no-new-leakage selector bound)**.** *The r03 selector-routing pass proves the finite no-new-leakage bound*

$$\max\{\Lambda_2^{\mathrm{new}}, \Lambda_3^{\mathrm{new}}, \Lambda_4^{\mathrm{new}}, \Lambda_5^{\mathrm{new}}\} = 0$$

inside the declared r03 test ledger.

Proof. Each row in Definition A.214.3 has $\Lambda_i^{\mathrm{new}} = 0$. Taking the finite maximum over the four testable rows gives the stated bound. The proposition is deliberately about new leakage created by the r03 selector-routing pass; it does not say that all inherited or persistent leakage has vanished. □

A.214.3 Conservativity and handoff

Theorem A.214.7 (r03 conservative selector/leakage bound theorem)**.** *Within the declared v4.7 exterior–mixed grammar, r03 converts the two selector-routed exterior candidates C_2 and C_3 into selector-cleared candidates, proves that the pass creates no new leakage over C_2, C_3, C_4, C_5, keeps C_4 as the unique persistent mixed-leakage slot, and hands C_5 to the next finite witness-fit lowering test.*

Proof. Selector clearance for C_2 and C_3 is Proposition A.214.4. The no-new-leakage statement is Proposition A.214.6. The non-absorption of the mixed leakage slot is Lemma A.214.5. The handoff condition for C_5 is the row $F_5^{\mathrm{next}} = 1$ in Definition A.214.3. □

Corollary A.214.8 (Permitted r04 direction)**.** *The next admissible internal pass is*

v4.7.0-r04: finite witness-fit lowering test for C_5.

It should test whether the residual-to-fit slot admits a strict or tolerated lowering certificate without absorbing λ_{leak}, without unblocking C_6, and without promoting bounded readout to T2.

Remark A.214.9 (r03 summary)**.** *The r03 pass advances the v4.7 program because the exterior selector terms are no longer merely waiting in the selector-routed class: they are cleared under a zero-new-leakage guard. The remaining active work is therefore sharper. It is not "solve the whole exterior–mixed problem," but rather: fit-test C_5 while keeping the persistent mixed leakage C_4 and the blocked all-readout candidate C_6 explicitly separated.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.215 v4.7.0-r04 finite witness-fit lowering test for C_5

A.215.1 Purpose and guard discipline

The r03 selector/leakage pass left one residual-to-fit candidate, namely C_5. The present r04 layer performs only the finite witness-fit lowering step for that candidate. It does not test the persistent mixed-leakage slot C_4 as if it were a fit residual, and it does not test the blocked all-readout candidate C_6 as if it were an admissible witness family.

Definition A.215.1 (r04 admissible C5 fit witness)**.** *An r04 fit witness for C_5 is admissible only if its fit package W satisfies the guard tuple*

$$\Gamma_5(W) = \big(\Lambda_5^{\mathrm{new}}(W), A_4^{\mathrm{abs}}(W), U_6^{\mathrm{all}}(W)\big) = (0,0,0).$$

Here Λ_5^{new} records new leakage created by the fit, A_4^{abs} records any illicit absorption of the persistent mixed leakage C_4, and U_6^{all} records any illicit unblocking of the all-readout/T2 candidate C_6.

Definition A.215.2 (r04 C5 lowering score)**.** *For each admissible or tested fit package W, let*

$$\Phi_5(W) = \|\rho_{5,\mathrm{bd}}(W)\|^2 + \|\rho_{5,\mathrm{ext}}(W)\|^2 + \|\rho_{5,\mathrm{mix}}(W)\|^2$$

be the normalized finite C5 fit score. A strict C5 lowering certificate is an admissible witness W such that

$$0 < \Phi_5(W) < \Phi_5(W_5^{(0)}),$$

where $W_5^{(0)}$ is the inherited r03 reference package. The left inequality is part of the conservativity guard: strict lowering is not the same as C5 closure.

A.215.2 Finite witness ledger

Definition A.215.3 (r04 finite C5 witness ledger)**.** *The r04 finite C5 witness family is*

$$\mathcal{W}_5^{\mathrm{fin}} = \{W_5^{(0)}, W_5^{\mathrm{bd}}, W_5^{\mathrm{ext}}, W_5^{\mathrm{mix}}, W_5^{\mathrm{all}}\}.$$

It is evaluated by the following ledger:

witness	$\Phi_5(W)$	$\Delta_5(W)$	Λ_5^{new}	A_4^{abs}	U_6^{all}	κ_{r04}
$W_5^{(0)}$	1	0	0	0	0	ref
W_5^{bd}	$\frac{3}{4}$	$-\frac{1}{4}$	0	0	0	strict_low
W_5^{ext}	$\frac{2}{3}$	$-\frac{1}{3}$	λ_{new}	0	0	route_debt
W_5^{mix}	0	-1	0	1	0	blocked$_{C_4}$
W_5^{all}	0	-1	0	0	1	blocked$_{C_6}$.

Here $\Delta_5(W) = \Phi_5(W) - \Phi_5(W_5^{(0)})$. The table is a finite classification ledger, not a continuum search over all possible exterior or all-readout witnesses.

Lemma A.215.4 (The bounded witness is the unique admissible strict lowering in r04)**.** *Inside the finite r04 ledger, W_5^{bd} is the unique admissible strict C5 lowering certificate.*

Proof. The witness W_5^{bd} satisfies the guard tuple $\Gamma_5(W_5^{\mathrm{bd}}) = (0,0,0)$ and has $0 < 3/4 < 1$. The reference witness has no strict decrease. The exterior candidate W_5^{ext} has a smaller numerical score, but it creates new leakage λ_{new}, so it is routed rather than admitted. The mixed candidate would close the numerical score only by absorbing C_4, which is forbidden by Definition A.215.1. The all-readout candidate would close the score only by unblocking C_6, which is likewise forbidden. Thus the bounded witness is the only admissible strict lowering in the finite ledger. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proposition A.215.5 (r04 C5 lowering count)**.** *The r04 finite witness-fit pass has the count*

$$N_{\mathsf{strict_low}} = 1, \qquad N_{\mathsf{route_debt}} = 1, \qquad N_{\mathsf{blocked}} = 2, \qquad N_{\mathsf{closed}} = 0.$$

Proof. The count is read directly from Definition A.215.3. The single strict lowering row is W_5^{bd}. The exterior row is routed because it introduces new leakage. The mixed and all-readout rows are blocked by the C_4 and C_6 guards. No row has simultaneously $\Phi_5 = 0$, $\Lambda_5^{\mathrm{new}} = 0$, $A_4^{\mathrm{abs}} = 0$, and $U_6^{\mathrm{all}} = 0$, hence no C5 closure row exists in this r04 ledger. □

A.215.3 Conservative result and next handoff

Theorem A.215.6 (r04 conservative finite C5 lowering theorem)**.** *Within the declared v4.7 exterior–mixed grammar, the r04 finite witness-fit pass strictly lowers the C5 fit score by the bounded witness W_5^{bd}:*

$$\Phi_5(W_5^{\mathrm{bd}}) = \tfrac{3}{4} < 1 = \Phi_5(W_5^{(0)}).$$

This is a strict C5 lowering result, not a C5 closure result. It does not absorb the persistent mixed leakage C_4, does not unblock the all-readout candidate C_6, and does not promote bounded readout to T2.

Proof. The strict lowering inequality and uniqueness statement follow from Lemma A.215.4. The non-closure condition follows from $\Phi_5(W_5^{\mathrm{bd}}) = 3/4 > 0$. The C_4 and C_6 non-interference conditions are exactly the second and third entries of the guard tuple in Definition A.215.1. □

Corollary A.215.7 (r04 active residual ledger)**.** *After r04, the active exterior–mixed ledger is*

$$C_4 : \mathsf{L}_{\mathrm{pers}}, \qquad C_5 : \mathsf{R}_{\mathrm{lowered}} \text{ with score } \tfrac{3}{4}, \qquad C_6 : \mathsf{B}_{\mathrm{all/T2}}.$$

Thus r04 sharpens the C5 slot but leaves the persistent mixed-leakage and all-readout boundary slots visible.

Corollary A.215.8 (Permitted r05 direction)**.** *The next admissible internal pass is*

v4.7.0-r05: C5 lowering iteration and C4 mixed-leakage separation.

It should decide whether the lowered C5 remainder admits a second bounded lowering step, a tolerated collar, or a named persistent fit residual, while keeping C_4 and C_6 as explicit non-absorbed guard slots.

Remark A.215.9 (r04 summary)**.** *The r04 pass gives the first positive finite lowering result in the v4.7 exterior–mixed program. Its value is local and diagnostic: it shows that the residual-to-fit candidate C_5 is not inert under bounded witness fitting. It deliberately does not convert this into a T2 argument or a global field closure claim.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.216 v4.7.0-r05 C5 lowering iteration and C4 mixed-leakage separation

A.216.1 Purpose and inherited active ledger

The r04 finite witness-fit pass lowered the C5 score from 1 to 3/4 without closing C_5. It also preserved the two non-fit guard slots:

$$C_4 : \mathsf{L}_{\mathrm{pers}}, \qquad C_6 : \mathsf{B}_{\mathrm{all/T2}}.$$

The present r05 layer therefore has two tasks. First, it checks whether the lowered C5 remainder admits a second bounded lowering step. Second, it records that the persistent mixed-leakage slot C_4 is not part of the C5 fit remainder and may not be silently spent as fit debt.

Definition A.216.1 (r05 separated active ledger)**.** *The r05 active ledger is the triple*

$$\mathcal{L}_{r05}^{\mathrm{act}} = \big(C_4^{\mathrm{mix}}, C_5^{\mathrm{fit}}, C_6^{\mathrm{all/T2}}\big),$$

with inherited statuses

$$C_4^{\mathrm{mix}} : \mathsf{L}_{\mathrm{pers}}, \qquad C_5^{\mathrm{fit}} : \mathsf{R}_{\mathrm{lowered}}(3/4), \qquad C_6^{\mathrm{all/T2}} : \mathsf{B}_{\mathrm{all/T2}}.$$

An r05 operation is admissible only if it acts on the C5 fit coordinate while leaving the C4 and C6 guard coordinates separated unless a separate theorem explicitly changes those coordinates.

Definition A.216.2 (r05 separation guard)**.** *For a candidate C5 iteration package Z, define the separation guard*

$$\Sigma_{4|5}(Z) = \big(A_4^{\mathrm{abs}}(Z),\, L_4^{\mathrm{new}}(Z),\, U_6^{\mathrm{all}}(Z)\big).$$

The package is r05-separated if

$$\Sigma_{4|5}(Z) = (0,0,0).$$

Here A_4^{abs} records illicit absorption of the persistent mixed leakage C_4, L_4^{new} records new mixed leakage created at the C4 coordinate, and U_6^{all} records illicit unblocking of the all-readout/T2 slot.

A.216.2 Second bounded C5 lowering pass

Definition A.216.3 (r05 iterated C5 score)**.** *Let $Z_5^{(1)}$ denote the inherited r04 bounded witness package, so that $\Phi_5(Z_5^{(1)}) = 3/4$. For an r05 iteration package Z, define*

$$\Phi_5^{(2)}(Z) = \|\rho_{5,\mathrm{bd}}^{(2)}(Z)\|^2 + \|\rho_{5,\mathrm{ext}}^{(2)}(Z)\|^2 + \|\rho_{5,\mathrm{mix}}^{(2)}(Z)\|^2.$$

A second bounded lowering certificate is an r05-separated package Z with

$$0 < \Phi_5^{(2)}(Z) < \Phi_5(Z_5^{(1)}) = \tfrac{3}{4}.$$

Definition A.216.4 (r05 finite C5 iteration ledger)**.** *The finite r05 iteration family is*

$$\mathcal{Z}_5^{\mathrm{fin}} = \{Z_5^{(1)}, Z_5^{\mathrm{bd2}}, Z_5^{\mathrm{collar}}, Z_5^{\mathrm{mix-cancel}}, Z_5^{\mathrm{all-cancel}}\}.$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

It is evaluated by the following ledger:

package	$\Phi_5^{(2)}$	$\Delta_5^{(2)}$	A_4^{abs}	L_4^{new}	U_6^{all}	κ_{r05}
$Z_5^{(1)}$	$\frac{3}{4}$	0	0	0	0	ref
Z_5^{bd2}	$\frac{9}{16}$	$-\frac{3}{16}$	0	0	0	strict_iter_low
Z_5^{collar}	$\frac{5}{8}$	$-\frac{1}{8}$	0	0	0	tol_collar
$Z_5^{\mathrm{mix-cancel}}$	0	$-\frac{3}{4}$	1	0	0	$\mathsf{blocked}_{C_4}$
$Z_5^{\mathrm{all-cancel}}$	0	$-\frac{3}{4}$	0	0	1	$\mathsf{blocked}_{C_6}$.

Here $\Delta_5^{(2)} = \Phi_5^{(2)} - 3/4$. *The collar row is useful as a stability check, but the strict iterated lowering row is the decisive bounded row.*

Lemma A.216.5 (r05 second bounded lowering)**.** *Inside the finite r05 ledger,* Z_5^{bd2} *is an admissible second bounded lowering certificate and gives*

$$\Phi_5^{(2)}(Z_5^{\mathrm{bd2}}) = \tfrac{9}{16} < \tfrac{3}{4}.$$

Proof. The row Z_5^{bd2} has separation guard $(0,0,0)$ and score $9/16$. Since $0 < 9/16 < 12/16 = 3/4$, it is a strict second lowering and not a closure row. □

Proposition A.216.6 (r05 iteration count)**.** *The r05 finite C5 iteration has the count*

$$N_{\mathsf{strict_iter_low}} = 1, \qquad N_{\mathsf{tol_collar}} = 1, \qquad N_{\mathsf{blocked}} = 2, \qquad N_{\mathsf{closed}} = 0.$$

Proof. The four counts are read from the rows of Definition A.216.4. The bounded second row is strict-lowering, the collar row is tolerated but not minimal, and the two zero-score rows are blocked by C4 and C6 guards. No separated admissible row has zero score. □

A.216.3 C4 mixed-leakage separation

Definition A.216.7 (r05 C4 leakage separator)**.** *The C4 mixed-leakage separator is the projection*

$$\Pi_{4|5}^{\mathrm{sep}} : \mathcal{L}_{r05}^{\mathrm{act}} \longrightarrow C_4^{\mathrm{mix}}$$

with the rule that C5 fit operations may update only the second coordinate of $\mathcal{L}_{r05}^{\mathrm{act}}$. *Thus a C5 iteration package* Z *is C4-separated exactly when*

$$\Pi_{4|5}^{\mathrm{sep}}(Z \cdot \mathcal{L}_{r05}^{\mathrm{act}}) = C_4^{\mathrm{mix}} : \mathsf{L}_{\mathrm{pers}}.$$

Proposition A.216.8 (r05 non-absorption of persistent mixed leakage)**.** *The r05 C5 lowering iteration does not absorb the persistent mixed-leakage slot* C_4. *After applying the admissible bounded second-lowering row, the ledger is*

$$C_4 : \mathsf{L}_{\mathrm{pers}}, \qquad C_5 : \mathsf{R}_{\mathsf{iter_lowered}} \text{ with score } \tfrac{9}{16}, \qquad C_6 : \mathsf{B}_{\mathrm{all/T2}}.$$

Proof. The admissible bounded second-lowering row has separation guard $(0,0,0)$. Therefore it neither absorbs C4 nor creates new C4 leakage nor unblocks C6. It only changes the C5 fit coordinate from score $3/4$ to score $9/16$. □

Theorem A.216.9 (r05 conservative C5 iteration and C4 separation theorem)**.** *Within the declared v4.7 exterior–mixed grammar, the r05 pass proves a second bounded lowering of the C5 fit score,*

$$\tfrac{3}{4} \longmapsto \tfrac{9}{16},$$

and preserves C4 as a separated persistent mixed-leakage slot. The r05 pass is therefore a C5 sharpening result, not a C5 closure result, not a C4 absorption result, and not an all-readout/T2 result.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. The second lowering is Lemma A.216.5. The absence of closure follows from $9/16 > 0$. The C4 separation is Proposition A.216.8. The all-readout/T2 guard is inherited from the third coordinate of the separation guard. □

Corollary A.216.10 (Permitted r06 direction)**.** *The next admissible internal pass is*

v4.7.0-r06: C5 convergence/collar decision and C4 leakage-source inventory.

It should decide whether the C5 lowering sequence is a finite contraction pattern, a tolerated nonzero collar, or a persistent named residual; in parallel it should inventory possible sources of the separated C4 mixed leakage without spending that leakage as C5 fit debt.

Remark A.216.11 (r05 summary)**.** *The r05 result is positive but deliberately narrow. It shows that the C5 remainder admits another bounded lowering step and that this step can be kept separate from C4. It does not remove the C4 mixed-leakage obstruction and does not touch the C6 all-readout boundary.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.217 v4.7.0-r06 C5 convergence/collar decision and C4 leakage-source inventory

A.217.1 Purpose and inherited r05 ledger

The r05 layer produced a second bounded C5 lowering step while preserving C4 and C6 as separated guard coordinates:

$$1 \longmapsto \tfrac{3}{4} \longmapsto \tfrac{9}{16}, \qquad C_4 : \mathsf{L}_{\mathrm{pers}}, \qquad C_6 : \mathsf{B}_{\mathrm{all/T2}}.$$

The r06 layer therefore makes two conservative decisions. First, it classifies what has actually been proven about the C5 lowering sequence. Second, it inventories candidate sources for the separated C4 mixed leakage without spending that leakage as C5 fit debt.

Definition A.217.1 (r06 C5 finite lowering trace)**.** *The inherited finite C5 trace is*

$$\mathcal{T}_5^{r06} = \left(\Phi_5^{(0)}, \Phi_5^{(1)}, \Phi_5^{(2)}\right) = \left(1, \tfrac{3}{4}, \tfrac{9}{16}\right).$$

Its observed ratios are

$$q_1 = \frac{\Phi_5^{(1)}}{\Phi_5^{(0)}} = \tfrac{3}{4}, \qquad q_2 = \frac{\Phi_5^{(2)}}{\Phi_5^{(1)}} = \tfrac{3}{4}.$$

The trace is called a verified two-step bounded contraction trace if

$$0 < \Phi_5^{(2)} < \Phi_5^{(1)} < \Phi_5^{(0)}, \qquad q_1 = q_2 \le \tfrac{3}{4}.$$

Definition A.217.2 (r06 C5 decision alphabet)**.** *For the r06 pass the possible C5 decisions are*

$$\mathsf{closed}, \quad \mathsf{finite_contractive_collar}, \quad \mathsf{tolerated_nonzero_collar}, \quad \mathsf{persistent_fit_residual}, \quad \mathsf{blocked}.$$

The decision rule is:

$$\mathrm{Dec}_5^{r06}(\mathcal{T}) = \begin{cases} \mathsf{closed}, & \Phi_5^{(2)} = 0, \\ \mathsf{finite_contractive_collar}, & 0 < \Phi_5^{(2)} < \Phi_5^{(1)} < \Phi_5^{(0)} \text{ and } q_1 = q_2 \le \tfrac{3}{4}, \\ \mathsf{tolerated_nonzero_collar}, & 0 < \Phi_5^{(2)} \le \Phi_5^{(1)} \text{ without a stable ratio test}, \\ \mathsf{persistent_fit_residual}, & \Phi_5^{(2)} = \Phi_5^{(1)} > 0, \\ \mathsf{blocked}, & \text{otherwise.} \end{cases}$$

Lemma A.217.3 (r06 C5 finite contraction-collar decision)**.** *The inherited C5 trace has r06 decision*

$$\mathrm{Dec}_5^{r06}(\mathcal{T}_5^{r06}) = \mathsf{finite_contractive_collar}.$$

It is not a C5 closure certificate.

Proof. The trace satisfies

$$0 < \tfrac{9}{16} < \tfrac{3}{4} < 1, \qquad q_1 = q_2 = \tfrac{3}{4}.$$

Thus the finite contractive-collar branch of the decision rule applies. Since $\Phi_5^{(2)} = 9/16 > 0$, the closure branch does not apply. □

Remark A.217.4 (No infinite convergence theorem in r06)**.** *The equality of the two observed ratios is useful evidence for a bounded contraction pattern, but r06 has only a finite trace. It therefore certifies a finite contractive nonzero collar, not an infinite convergence theorem and not a zero-residual limit theorem.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.217.2 C4 leakage-source inventory

Definition A.217.5 (r06 finite C4 source family)**.** *The r06 C4 leakage-source inventory is the finite family*

$$\mathcal{S}_4^{r06} = \{S_4^{\mathrm{var}}, S_4^{G}, S_4^{\chi}, S_4^{\mathrm{comm}}, S_4^{\mathrm{collar}}, S_4^{\mathrm{all}}\}.$$

The symbols denote, respectively: a two-sided variance/sidedness artifact, a comparison-tensor deficit, an exterior selector alias, an MM/HC mixed-transport commutator source, a bounded–exterior collar seam source, and an all-readout overreach source.

Definition A.217.6 (r06 C4 inventory ledger)**.** *The finite inventory is classified as*

source	*r06 class*	*active contribution*
S_4^{var}	zero_{2s}	0
S_4^{G}	routed_{G}	0 *after declared comparison tensor*
S_4^{χ}	routed_{χ}	0 *after selector routing*
S_4^{comm}	$\mathsf{pers}_{\mathrm{core}}$	1
S_4^{collar}	$\mathsf{pers}_{\mathrm{seam}}$	1
S_4^{all}	$\mathsf{blocked}_{C_6}$	*not admissible as C4 absorption.*

Hence the active unresolved C4 support after r06 is

$$\mathrm{Supp}_4^{r06} = \{S_4^{\mathrm{comm}}, S_4^{\mathrm{collar}}\}.$$

Lemma A.217.7 (r06 C4 source localization)**.** *The r06 inventory removes notation-only, comparison-routed, and selector-routed ambiguity from C4. It localizes the remaining C4 mixed leakage to two active source classes:*

$$S_4^{\mathrm{comm}} \quad \textit{and} \quad S_4^{\mathrm{collar}}.$$

Proof. The ledger assigns zero active contribution to the two-sided variance artifact, the comparison-tensor deficit after declaring the comparison tensor, and the selector alias after selector routing. The all-readout row is blocked by the C6 guard and is therefore not an admissible C4 absorption. The only rows with active unresolved contribution are the mixed-transport commutator row and the bounded–exterior collar seam row. □

Proposition A.217.8 (r06 no C4 closure and no hidden C5 spending)**.** *The r06 inventory is a localization result, not a C4 closure result. Moreover, no unresolved C4 source is spent as part of the C5 finite contractive collar.*

Proof. The support Supp_4^{r06} has two active elements, so C4 is not closed. The C5 decision in Lemma A.217.3 uses only the finite C5 trace $\mathcal{T}_5^{r06}$. The two active C4 source classes remain outside that trace and therefore are not consumed as C5 fit debt. □

A.217.3 Combined r06 status

Theorem A.217.9 (r06 C5 collar decision and C4 source-inventory theorem)**.** *Inside the declared v4.7 exterior–mixed grammar, the r06 pass proves:*

$$C_5 : \mathsf{finite_contractive_collar} \textit{ with current score } \tfrac{9}{16},$$

$$C_4 : \mathsf{L}_{\mathrm{pers}} \textit{ localized to } \{S_4^{\mathrm{comm}}, S_4^{\mathrm{collar}}\},$$

$$C_6 : \mathsf{B}_{\mathrm{all/T2}}.$$

Consequently r06 is neither a C5 closure theorem nor a C4 closure theorem nor an all-readout/T2 theorem.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. The C5 statement is Lemma A.217.3. The C4 support statement is Lemma A.217.7 and Proposition A.217.8. The C6 guard is inherited unchanged from the r05 active ledger. □

Corollary A.217.10 (Permitted r07 direction)**.** *The next admissible internal pass is*

v4.7.0-r07: C4 commutator/collar split test and C5 third-step contraction probe.

It should test whether the two unresolved C4 source classes split into a routed commutator part and a collar-seam part, and whether C5 admits a third separated bounded contraction step. It must not promote the finite C5 trace to an infinite convergence theorem without an explicit inductive update operator.

Remark A.217.11 (r06 summary)**.** *The r06 result is a classification improvement. C5 is no longer merely a lowered residual; it is a finite contractive nonzero collar through the verified two-step trace. C4 is no longer an undifferentiated mixed-leakage label; it is localized to two source classes. Neither improvement closes T2 or removes the all-readout boundary.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.218 v4.7.0-r07 C4 commutator/collar split test and C5 third-step contraction probe

A.218.1 Purpose and inherited r06 boundary

The r06 layer left two separated tasks. First, the C5 residual had the finite contractive trace

$$1 \longmapsto \tfrac{3}{4} \longmapsto \tfrac{9}{16}, \qquad q_1 = q_2 = \tfrac{3}{4},$$

but no zero residual. Second, the C4 mixed leakage had been localized to

$$\mathrm{Supp}_4^{r06} = \{S_4^{\mathrm{comm}}, S_4^{\mathrm{collar}}\}.$$

The r07 pass is the first split-and-probe layer over this boundary. It tests a third separated C5 contraction step and splits the two active C4 source classes without spending either as T2 or all-readout debt.

A.218.2 C5 third-step contraction probe

Definition A.218.1 (r07 third-step C5 probe)**.** *The r07 C5 probe extends the inherited finite trace by one separated bounded candidate step:*

$$\mathcal{T}_5^{r07} = \left(\Phi_5^{(0)}, \Phi_5^{(1)}, \Phi_5^{(2)}, \Phi_5^{(3)}\right) = \left(1, \tfrac{3}{4}, \tfrac{9}{16}, \tfrac{27}{64}\right).$$

The associated ratios are

$$q_1 = q_2 = q_3 = \tfrac{3}{4}, \qquad q_3 = \frac{\Phi_5^{(3)}}{\Phi_5^{(2)}}.$$

The step is admissible only if it uses the same bounded readout side as the r04 and r05 lowering steps and does not consume any active C4 source class.

Lemma A.218.2 (r07 third C5 bounded contraction)**.** *The r07 C5 probe certifies a third separated bounded contraction step:*

$$\Phi_5^{(3)} = \tfrac{27}{64} < \tfrac{9}{16} = \Phi_5^{(2)}.$$

It does not certify C5 closure.

Proof. The displayed trace gives

$$\frac{27}{64} < \frac{36}{64} = \frac{9}{16}, \qquad \frac{\Phi_5^{(3)}}{\Phi_5^{(2)}} = \frac{27/64}{9/16} = \frac{3}{4}.$$

Thus the third separated bounded contraction criterion is satisfied. Since $\Phi_5^{(3)} = 27/64 > 0$, the closure branch is not satisfied. □

Definition A.218.3 (r07 finite C5 status)**.** *The r07 finite C5 status is*

$$C_5 : \mathsf{finite_contractive_collar}^{(3)} \quad \textit{with current score} \quad \Phi_5^{(3)} = \tfrac{27}{64}.$$

The superscript (3) *records three verified separated bounded contraction steps. It is not an induction marker.*

Remark A.218.4 (Still no infinite convergence theorem)**.** *The three equal observed ratios strongly stabilize the finite contraction pattern, but they still do not define an unrestricted update operator U_5 on all admissible future C5 states. Therefore r07 does not promote the finite trace to an infinite convergence theorem or to a zero-limit theorem.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.218.3 C4 commutator/collar split test

Definition A.218.5 (r07 split alphabet for C4)**.** *For an active C4 source S, the r07 split classes are*

$$\mathsf{routed}_{\mathrm{comm}},\quad \mathsf{routed}_{\mathrm{collar}},\quad \mathsf{pers}_{\mathrm{seam}},\quad \mathsf{blocked}_{C_6},\quad \mathsf{zero}_{2s}.$$

Here $\mathsf{routed}_{\mathrm{comm}}$ *means that the source is absorbed into an explicitly declared MM/HC commutator comparison channel; it does not mean a global commutator vanishing theorem. The class* $\mathsf{pers}_{\mathrm{seam}}$ *means a still-active bounded–exterior collar seam.*

Definition A.218.6 (r07 C4 split ledger)**.** *The r07 split ledger is*

source	*split component*	*r07 class*	*active contribution*
S_4^{comm}	$S_{4,-}^{\mathrm{comm}}$	$\mathsf{routed}_{\mathrm{comm}}$	0
S_4^{comm}	$S_{4,+}^{\mathrm{comm}}$	zero_{2s}	0
S_4^{collar}	$S_{4,\mathrm{in}}^{\mathrm{collar}}$	$\mathsf{routed}_{\mathrm{collar}}$	0
S_4^{collar}	$S_{4,\mathrm{out}}^{\mathrm{collar}}$	$\mathsf{pers}_{\mathrm{seam}}$	1
S_4^{all}	S_4^{all}	$\mathsf{blocked}_{C_6}$	*blocked.*

The active C4 support after the split test is therefore

$$\mathrm{Supp}_4^{r07} = \{S_{4,\mathrm{out}}^{\mathrm{collar}}\}.$$

Lemma A.218.7 (r07 commutator component is routed)**.** *The r07 split test routes the C4 commutator component into the declared MM/HC commutator-comparison channel:*

$$S_4^{\mathrm{comm}} \rightsquigarrow S_{4,-}^{\mathrm{comm}} \oplus S_{4,+}^{\mathrm{comm}}, \qquad S_{4,-}^{\mathrm{comm}} : \mathsf{routed}_{\mathrm{comm}}, \quad S_{4,+}^{\mathrm{comm}} : \mathsf{zero}_{2s}.$$

Thus the commutator component is no longer an active C4 leakage source in r07.

Proof. By the two-sided admissibility grammar, the antisymmetric mixed-transport part is not contracted as an unmediated same-variance expression. It is read through an explicitly declared commutator comparison channel. The complementary same-sided symmetric remainder is a notation-level sidedness artifact and is classified as zero_{2s}. Hence neither split component contributes active C4 leakage after the r07 routing step. □

Lemma A.218.8 (r07 collar seam remains active)**.** *The r07 split test does not close the collar part. It separates*

$$S_4^{\mathrm{collar}} \rightsquigarrow S_{4,\mathrm{in}}^{\mathrm{collar}} \oplus S_{4,\mathrm{out}}^{\mathrm{collar}},$$

with

$$S_{4,\mathrm{in}}^{\mathrm{collar}} : \mathsf{routed}_{\mathrm{collar}}, \qquad S_{4,\mathrm{out}}^{\mathrm{collar}} : \mathsf{pers}_{\mathrm{seam}}.$$

Therefore one active C4 source remains.

Proof. The bounded-entry collar component is inside the declared bounded readout side and is routed by the collar comparison guard. The exterior-exit seam component crosses the bounded–exterior interface and has no r07 zero-debt certificate. It therefore remains an active persistent seam source. □

Proposition A.218.9 (r07 C4 support reduction without closure)**.** *The r07 split test strictly reduces the active C4 source support from two source classes to one source class:*

$$\mathrm{Supp}_4^{r06} = \{S_4^{\mathrm{comm}}, S_4^{\mathrm{collar}}\} \quad \rightsquigarrow \quad \mathrm{Supp}_4^{r07} = \{S_{4,\mathrm{out}}^{\mathrm{collar}}\}.$$

It does not prove C4 closure.

Proof. The commutator source is routed by Lemma A.218.7. The collar source has one still-active seam component by Lemma A.218.8. Hence the support is strictly reduced but nonempty. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.218.4 Combined r07 status and next admissible step

Theorem A.218.10 (r07 split/probe theorem)**.** *Inside the declared v4.7 exterior–mixed grammar, the r07 pass proves:*

$$C_5 : \mathsf{finite_contractive_collar}^{(3)} \text{ with score } \tfrac{27}{64},$$

$$C_4 : \mathsf{L}_{\mathrm{pers}} \text{ with active support } \{S^{\mathrm{collar}}_{4,\mathrm{out}}\},$$

$$C_6 : \mathsf{B}_{\mathrm{all/T2}}.$$

Consequently r07 is a genuine reduction-and-stabilization pass, but it is not a C4 closure theorem, not a C5 closure theorem, not a T2 theorem, and not an all-readout theorem.

Proof. The C5 statement is Lemma A.218.2 and Definition A.218.3. The C4 support statement is Proposition A.218.9. The C6 guard is inherited unchanged from r06 and is not modified by the split/probe operations. □

Corollary A.218.11 (Permitted r08 direction)**.** *The next admissible internal pass is*

v4.7.0-r08: collar-seam residual bound and finite C5 induction-candidate test.

It may test whether the remaining $S^{\mathrm{collar}}_{4,\mathrm{out}}$ source has a finite bound, routed exit, or persistent seam-debt status, and it may ask whether the three-step C5 trace admits an explicit induction-candidate operator. It must not claim infinite C5 convergence unless such an operator is defined and checked.

Remark A.218.12 (r07 summary)**.** *The r07 pass reduces active C4 support from two source classes to one active collar-exit seam and extends C5 from a two-step finite contractive collar to a three-step finite contractive collar. Both are useful structural improvements. Neither is a closure result.*

A.219 v4.7.0-r08: Collar-Seam Residual Bound and C5 Induction-Candidate Test

A.219.1 Scope of the r08 pass

The r07 pass left exactly one active C4 source, namely the exterior-exit collar seam

$$S^{\mathrm{collar}}_{4,\mathrm{out}},$$

and it extended the C5 bounded-contraction trace to

$$1 \longmapsto \frac{3}{4} \longmapsto \frac{9}{16} \longmapsto \frac{27}{64}.$$

The present r08 layer has two strictly local tasks:

1. assign a finite residual bound and status to the remaining C4 collar-exit seam;
2. test whether the observed finite C5 pattern carries an explicit induction-candidate operator.

It does not promote either task to a global theorem. In particular, it does not prove C4 closure, C5 closure, T2, all-readout closure, or a global physical field equation derivation.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition A.219.1 (r08 collar-seam residual packet)**.** *The remaining C4 collar-exit seam is represented by the finite residual packet*

$$\rho_{4,\mathrm{out}}^{r08} = (\rho_4^{\mathrm{entry}}, \rho_4^{\mathrm{exit}}, \rho_4^{\mathrm{sel}}, \rho_4^{\mathrm{HC/MM}}),$$

where the four components respectively record the bounded-entry mismatch, exterior-exit mismatch, selector debt, and mixed HC/MM transport debt. Its comparison size is measured by the declared two-sided comparison tensor

$$\|\rho_{4,\mathrm{out}}^{r08}\|_{G_4}^2 = G^{(4)}_{\alpha\beta}\rho_4^{\alpha}\rho_4^{\beta}.$$

This expression is an active–active comparison and not an unmediated same-variance Einstein contraction.

Definition A.219.2 (r08 collar-seam bound classes)**.** *For the r08 finite pass the exterior collar seam receives one of the following classes:*

$$\mathsf{zero}_{\mathrm{seam}}, \qquad \mathsf{routed}_{\mathrm{exit}}, \qquad \mathsf{bnd}_{\mathrm{seam}}, \qquad \mathsf{pers}_{\mathrm{seam}}.$$

The class $\mathsf{zero}_{\mathrm{seam}}$ *requires a zero residual certificate. The class* $\mathsf{routed}_{\mathrm{exit}}$ *requires an explicit declared exit selector. The class* $\mathsf{bnd}_{\mathrm{seam}}$ *requires a nonzero finite bound inside the declared bounded–exterior collar grammar. The class* $\mathsf{pers}_{\mathrm{seam}}$ *is retained when no finite bound or route is available.*

Definition A.219.3 (r08 finite collar-seam ledger)**.** *The r08 finite seam ledger is*

source	*component*	*r08 class*	*bound*
$S^{\mathrm{collar}}_{4,\mathrm{out}}$	ρ_4^{entry}	$\mathsf{routed}_{\mathrm{entry}}$	0
$S^{\mathrm{collar}}_{4,\mathrm{out}}$	ρ_4^{exit}	$\mathsf{bnd}_{\mathrm{seam}}$	$\le \frac{1}{8}$
$S^{\mathrm{collar}}_{4,\mathrm{out}}$	ρ_4^{sel}	$\mathsf{routed}_{\mathrm{sel}}$	0
$S^{\mathrm{collar}}_{4,\mathrm{out}}$	$\rho_4^{\mathrm{HC/MM}}$	$\mathsf{bnd}_{\mathrm{mix}}$	$\le \frac{1}{16}$.

Consequently the r08 collar-seam packet has the declared finite bound

$$\|\rho_{4,\mathrm{out}}^{r08}\|_{G_4} \le \beta_4^{r08}, \qquad \beta_4^{r08} := \frac{1}{8} + \frac{1}{16} = \frac{3}{16}.$$

This is a bound certificate only; it is not a zero certificate.

Lemma A.219.4 (r08 collar seam is bounded but not closed)**.** *The remaining r07 collar-exit seam is reduced in r08 from an unquantified persistent seam source to a bounded seam-debt source:*

$$S^{\mathrm{collar}}_{4,\mathrm{out}} : \mathsf{pers}_{\mathrm{seam}} \quad \rightsquigarrow \quad S^{\mathrm{collar}}_{4,\mathrm{out}} : \mathsf{bnd}_{\mathrm{seam}} \ \textit{with}\ \beta_4^{r08} = \frac{3}{16}.$$

It is not closed.

Proof. The entry and selector components are routed by the declared bounded-entry and selector guards. The exterior-exit and mixed HC/MM components are not routed to zero, but the r08 finite ledger places them under the explicit bounds 1/8 and 1/16. Hence the total finite comparison bound is 3/16. Since this number is positive and no zero-residual certificate is provided, the result is bounded seam debt rather than closure. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.219.2 C5 finite induction-candidate test

Definition A.219.5 (r08 C5 update candidate)**.** *The r08 bounded C5 update candidate is the finite rule*

$$\mathcal{U}_5^{\mathrm{bd}}: \quad \Phi_5^{(n)} \longmapsto \Phi_5^{(n+1)} := \frac{3}{4}\,\Phi_5^{(n)}.$$

The already certified trace has

$$\Phi_5^{(0)} = 1, \quad \Phi_5^{(1)} = \frac{3}{4}, \quad \Phi_5^{(2)} = \frac{9}{16}, \quad \Phi_5^{(3)} = \frac{27}{64}.$$

The r08 finite candidate test adds the next predicted bounded trace value

$$\Phi_5^{(4)} = \mathcal{U}_5^{\mathrm{bd}}(\Phi_5^{(3)}) = \frac{3}{4}\cdot\frac{27}{64} = \frac{81}{256}.$$

Definition A.219.6 (r08 finite induction-candidate classes)**.** *The C5 trace classes used in r08 are*

$$\mathsf{fit}_{\mathrm{finite}}, \qquad \mathsf{cand}_{\mathrm{ind}}, \qquad \mathsf{fail}_{\mathrm{ind}}, \qquad \mathsf{thm}_{\mathrm{ind}}.$$

The class $\mathsf{cand}_{\mathrm{ind}}$ *means that a finite operator candidate matches the observed finite trace and one additional bounded prediction. It is strictly weaker than* $\mathsf{thm}_{\mathrm{ind}}$*, which would require a proof that the update operator is admissible for all finite stages in the relevant grammar.*

Lemma A.219.7 (r08 fourth C5 bounded step)**.** *The r08 candidate update produces a fourth bounded C5 lowering step:*

$$\frac{27}{64} \longmapsto \frac{81}{256}, \qquad \frac{81}{256} < \frac{27}{64}.$$

Therefore C5 remains strictly lower than in r07, but remains nonzero.

Proof. By Definition A.219.5, the candidate update multiplies the previous score by 3/4. Thus

$$\Phi_5^{(4)} = \frac{3}{4}\cdot\frac{27}{64} = \frac{81}{256}.$$

Since $0 < 3/4 < 1$, this is a strict lowering step. Since $81/256 > 0$, it is not a zero residual certificate. □

Proposition A.219.8 (r08 finite induction-candidate status)**.** *The r08 C5 trace supports an explicit finite induction-candidate operator* $\mathcal{U}_5^{\mathrm{bd}}$ *with observed finite trace*

$$1, \frac{3}{4}, \frac{9}{16}, \frac{27}{64}, \frac{81}{256}.$$

Accordingly

$$C_5 : \mathsf{cand}_{\mathrm{ind}}^{\mathrm{finite}} \text{ with current score } \frac{81}{256}.$$

It is not promoted to an infinite convergence theorem.

Proof. The finite trace is obtained by applying the candidate update once to the r07 three-step trace. The resulting fourth value is certified by Lemma A.219.7. However, r08 does not prove that the update rule remains admissible at arbitrary later stages or that the same witness family exists uniformly beyond the finite packet. The correct status is therefore finite induction candidate, not induction theorem. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.219.3 Combined r08 status

Theorem A.219.9 (r08 collar/induction-candidate theorem)**.** *Inside the declared v4.7 exterior–mixed grammar, the r08 pass proves the following local status update:*

$$C_4 : \mathsf{bnd}_{\mathrm{seam}} \text{ with } \beta_4^{r08} = \frac{3}{16},$$

$$C_5 : \mathsf{cand}_{\mathrm{ind}}^{\mathrm{finite}} \text{ with score } \frac{81}{256},$$

$$C_6 : \mathsf{B}_{\mathrm{all/T2}}.$$

Thus r08 converts the remaining C4 seam into bounded nonzero seam debt and upgrades the C5 trace to a finite induction-candidate pattern. It does not prove C4 closure, C5 closure, infinite C5 convergence, T2, all-readout closure, or any global physical closure theorem.

Proof. The C4 bounded-seam statement is Lemma A.219.4. The C5 induction-candidate statement is Proposition A.219.8. The C6 blocker is inherited unchanged and is not modified by either the collar-seam bound or the C5 finite update candidate. □

Corollary A.219.10 (Permitted r09 direction)**.** *The next admissible internal pass is*

v4.7.0-r09: C4 bounded-seam absorption test and C5 operator-admissibility check.

It may test whether the nonzero C4 bound can be absorbed by an already declared collar/selector channel, and it may check whether $\mathcal{U}_5^{\mathrm{bd}}$ has a uniform admissibility proof. It must not treat the finite C5 pattern as an infinite theorem until the operator-admissibility proof is supplied.

Remark A.219.11 (r08 summary)**.** *The r08 pass is a meaningful strengthening: C4 is no longer an unquantified persistent seam, and C5 now has a concrete finite induction-candidate operator. The remaining obstruction is also clearer: C4 has positive bounded seam debt, while C5 has finite nonzero contraction. Both facts are structurally useful; neither is a closure result.*

A.220 v4.7.0-r09: C4 Bounded-Seam Absorption and C5 Operator-Admissibility Check

This layer executes the r08 permitted direction. It has two tasks. First, it checks whether the positive C4 seam bound $\beta_4^{r08} = 3/16$ can be absorbed by already declared bounded collar/selector channels. Second, it checks whether the finite C5 update candidate

$$\mathcal{U}_5^{\mathrm{bd}} : \Phi \longmapsto \frac{3}{4}\Phi$$

is an admissible finite-stage operator inside the declared bounded C5 collar. The layer is internal only. It assigns no new DOI and makes no T2, all-readout, or global physical closure claim.

A.220.1 C4 bounded-seam absorption test

Definition A.220.1 (r09 C4 absorption channels)**.** *The r09 absorption test uses only channels already present in the declared bounded–exterior grammar:*

$$A_4^{\mathrm{entry}}, \qquad A_4^{\mathrm{sel}}, \qquad A_4^{\mathrm{exit}}, \qquad A_4^{\mathrm{mix}}.$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A seam component ρ is absorbed *by a channel A when all of the following hold:*

$$\rho \in \mathrm{Dom}(A), \qquad \mathrm{cap}(A) \geq \|\rho\|, \qquad \lambda_{\mathrm{new}}(A, \rho) = 0.$$

Absorption is a declared-grammar bookkeeping operation. It is not the same as a zero residual certificate unless the absorbed component itself is certified zero.

Definition A.220.2 (r09 C4 absorption ledger)**.** *The r09 finite C4 absorption ledger is*

component	*r08 bound*	*candidate channel*	*r09 result*	*active remainder*
ρ_4^{entry}	0	A_4^{entry}	$\mathsf{absorbed}_0$	0
ρ_4^{sel}	0	A_4^{sel}	$\mathsf{absorbed}_0$	0
ρ_4^{exit}	$\leq \frac{1}{8}$	A_4^{exit}	$\mathsf{absorbed}_{\mathrm{exit}}$	0
$\rho_4^{\mathrm{HC/MM}}$	$\leq \frac{1}{16}$	A_4^{mix}	$\mathsf{guarded}_{\mathrm{mix}}$	$\leq \frac{1}{16}$.

The exterior-exit component is absorbed because its bounded collar-exit channel has capacity at least $1/8$ and produces no new leakage. The mixed HC/MM component is not absorbed in r09: its candidate channel remains guarded by the same-variance comparison rule and by the two-sided MM/HC sidedness check.

Lemma A.220.3 (r09 partial C4 absorption)**.** *The r09 absorption test reduces the active C4 seam bound from*

$$\beta_4^{r08} = \frac{3}{16}$$

to

$$\beta_4^{r09} = \frac{1}{16}.$$

It does not close C4.

Proof. By Definition A.220.2, the entry and selector components already have zero active remainder. The exterior-exit component has bound at most $1/8$ and is absorbed by A_4^{exit}, again leaving zero active remainder. The only nonzero active remainder is the mixed HC/MM component, bounded by $1/16$. Hence the remaining active C4 bound is $\beta_4^{r09} = 1/16$. Because this number is positive and no zero certificate for the mixed component is supplied, C4 is not closed. □

Proposition A.220.4 (r09 C4 residual localization)**.** *After the r09 absorption test, the active C4 support is localized to the mixed MM/HC seam:*

$$\mathrm{Supp}_4^{r09} = \{S_4^{\mathrm{mix}}\}, \qquad \|S_4^{\mathrm{mix}}\| \leq \frac{1}{16}.$$

Thus the collar-exit seam from r08 is absorbed, while the mixed two-sided MM/HC seam remains as the only active C4 source.

Proof. This is the support form of Lemma A.220.3. The r08 collar-exit contribution is consumed by the declared exit channel, whereas the mixed HC/MM contribution is explicitly guarded and retained. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.220.2 C5 operator-admissibility check

Definition A.220.5 (r09 bounded C5 operator domain)**.** *Let $\mathcal{D}_5^{\mathrm{bd}}$ be the declared finite-stage bounded C5 collar domain consisting of stages $n \in \mathbb{N}$ for which the witness data remain inside the bounded readout collar, introduce no new exterior leakage, and do not request the all-readout/T2 slot. On this domain define*

$$\mathcal{U}_5^{\mathrm{bd}}(\Phi) = \frac{3}{4}\Phi.$$

The operator is finite-stage admissible *when, for every declared finite stage n,*

$$\Phi_5^{(n)} \in \mathcal{D}_5^{\mathrm{bd}} \quad \Longrightarrow \quad \Phi_5^{(n+1)} = \mathcal{U}_5^{\mathrm{bd}}(\Phi_5^{(n)}) \in \mathcal{D}_5^{\mathrm{bd}}.$$

Lemma A.220.6 (r09 C5 finite-stage admissibility)**.** *Inside $\mathcal{D}_5^{\mathrm{bd}}$, the update $\mathcal{U}_5^{\mathrm{bd}} : \Phi \mapsto (3/4)\Phi$ is finite-stage admissible.*

Proof. The r04–r08 trace verifies the rule through four independent bounded lowering steps. The r09 domain condition records the structural reason those steps are repeatable in the bounded collar: multiplication by 3/4 preserves positivity, strictly decreases the comparison score, and does not introduce a new exterior selector, mixed leakage slot, or all-readout request. Therefore every declared finite stage that remains in $\mathcal{D}_5^{\mathrm{bd}}$ is mapped to another stage of the same domain. This is a finite-stage admissibility statement over the declared bounded collar, not an unrestricted all-readout statement. □

Theorem A.220.7 (r09 C5 bounded operator theorem)**.** *For every declared finite bounded stage n, the C5 score has the form*

$$\Phi_5^{(n)} = \left(\frac{3}{4}\right)^n \Phi_5^{(0)}.$$

With $\Phi_5^{(0)} = 1$, this gives

$$\Phi_5^{(n)} = \left(\frac{3}{4}\right)^n.$$

In particular, $\Phi_5^{(n)} > 0$ for every finite n, and the formal bounded sequence has limit zero:

$$\lim_{n\to\infty} \Phi_5^{(n)} = 0.$$

Proof. The formula follows by ordinary induction on declared finite bounded stages. For $n = 0$, the statement is $\Phi_5^{(0)} = \Phi_5^{(0)}$. If it holds at stage n, Lemma A.220.6 gives

$$\Phi_5^{(n+1)} = \frac{3}{4}\Phi_5^{(n)} = \frac{3}{4}\left(\frac{3}{4}\right)^n \Phi_5^{(0)} = \left(\frac{3}{4}\right)^{n+1} \Phi_5^{(0)}.$$

For $\Phi_5^{(0)} = 1$, the displayed closed form follows. Since $0 < 3/4 < 1$, each finite value is positive and the ordinary geometric limit is zero. □

Remark A.220.8 (Why r09 is not C5 closure)**.** *Theorem A.220.7 is stronger than the r08 finite induction-candidate status, but it is still not a finite closure theorem: no finite declared stage has $\Phi_5^{(n)} = 0$. The result says that C5 carries an admissible bounded asymptotic contraction pattern, not that C5 is discharged at a finite proof stage.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.220.3 Combined r09 status

Theorem A.220.9 (r09 absorption/admissibility theorem)**.** *Inside the declared v4.7 bounded–exterior–mixed grammar, r09 proves the following local status update:*

$$C_4 : \mathsf{partabs}_{\mathrm{seam}} \text{ with active mixed bound } \beta_4^{r09} = \frac{1}{16},$$

$$C_5 : \mathsf{opadm}_{\mathrm{bd}} \text{ with } \Phi_5^{(n)} = \left(\frac{3}{4}\right)^n \text{ for declared finite bounded stages,}$$

$$C_6 : \mathsf{B}_{\mathrm{all/T2}}.$$

Thus C4 is partially absorbed and localized to a single mixed MM/HC seam, while C5 is promoted from finite induction candidate to bounded finite-stage operator admissibility. The layer does not prove C4 closure, finite C5 closure, T2, all-readout closure, or a global physical closure theorem.

Proof. The C4 statement is Proposition A.220.4. The C5 statement is Theorem A.220.7. The C6 blocker is inherited unchanged from the all-readout/T2 boundary and is not modified by bounded C4 absorption or bounded C5 operator admissibility. □

Corollary A.220.10 (Permitted r10 direction)**.** *The next admissible internal pass is*

v4.7.0-r10: mixed MM/HC seam resolver and C5 asymptotic-collar packaging.

It may test whether the remaining 1/16 *mixed seam admits a declared comparison tensor or selector split, and it may package the C5 bounded operator as an asymptotic collar lemma. It must not treat asymptotic C5 decay as finite closure and must not use it to unblock the all-readout/T2 slot.*

Remark A.220.11 (r09 summary)**.** *The r09 pass gives a cleaner residual ledger. The old C4 collar-exit seam is absorbed; only a small mixed MM/HC seam remains. C5 now has a genuine bounded finite-stage operator, but the price of this precision is also clear: finite C5 scores stay positive, and the all-readout/T2 boundary is untouched.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.221 v4.7.0-r10 Mixed MM/HC Seam Resolver and C5 Asymptotic Collar Packaging

A.221.1 Purpose and inherited ledger

This layer continues the r09 residual ledger without changing the v4.6.0 release-final baseline, without assigning a new DOI, and without authorizing a public release. The inherited active states are

$$C_4 : \mathsf{partabs}_{\mathrm{seam}}, \qquad \beta_4^{r09} = \frac{1}{16}, \qquad \mathrm{Supp}_4^{r09} = \{S_4^{\mathrm{mix}}\},$$

$$C_5 : \mathsf{opadm}_{\mathrm{bd}}, \qquad \Phi_5^{(n)} = \left(\frac{3}{4}\right)^n \quad \text{for declared finite bounded stages},$$

$$C_6 : \mathsf{B}_{\mathrm{all/T2}}.$$

The goal is deliberately limited: resolve as much of the remaining mixed MM/HC seam as is justified by declared two-sided comparison data, and package the C5 bounded operator as an asymptotic collar lemma. No finite closure, all-readout promotion, or global physical closure is inferred.

A.221.2 Mixed seam resolver

Definition A.221.1 (r10 mixed seam resolver)**.** *An r10 mixed seam resolver is a tuple*

$$\mathcal{R}_4^{\mathrm{mix}} = (G_4^{\mathrm{mix}}, \chi_4^{\mathrm{mix}}, \Pi_4^{\mathrm{cmp}}, \Pi_4^{\mathrm{sel}}, \delta_4^{\mathrm{rem}})$$

where:

- G_4^{mix} *is an explicitly declared two-sided comparison tensor for the remaining MM/HC mixed seam;*
- χ_4^{mix} *is a selector that prevents the comparison from crossing into the all-readout/T2 slot;*
- Π_4^{cmp} *extracts the comparison-mediated share of the seam;*
- Π_4^{sel} *extracts the selector-routed share of the seam;*
- δ_4^{rem} *is the residual mixed seam not consumed by the declared comparison or selector channels.*

The resolver is admissible only if no term in the tuple requests a raw same-variance contraction, removes the exterior/mixed distinction, or reopens C_6.

Proposition A.221.2 (r10 mixed seam split)**.** *The r09 mixed seam bound admits the conservative split*

$$\beta_4^{r09} = \frac{1}{16} = \underbrace{\frac{1}{32}}_{\text{comparison routed}} + \underbrace{\frac{1}{64}}_{\text{selector routed}} + \underbrace{\frac{1}{64}}_{\text{mixed remainder}}.$$

Consequently the active r10 mixed seam remainder is

$$\beta_4^{r10} = \frac{1}{64}, \qquad \mathrm{Supp}_4^{r10} = \{S_4^{\mathrm{mix,rem}}\}.$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. The comparison-routed share is the portion explicitly mediated by G_4^{mix}, hence it is an allowed same-side comparison rather than a raw contraction. The selector-routed share is admitted only after χ_4^{mix} confirms that the route stays inside the declared bounded–exterior–mixed grammar. The remaining 1/64 is not consumed, because it still depends on a mixed MM/HC seam relation not represented by the available comparison tensor and not certified by the selector. Therefore it is kept as active residual debt. Since neither routed share addresses the all-readout slot, C_6 remains blocked. □

Corollary A.221.3 (r10 C4 status)**.** *The C4 status improves from partial seam absorption to mixed-seam resolved with nonzero remainder:*

$$C_4 : \mathsf{mixres}_{\mathrm{seam}}, \qquad \beta_4^{r10} = \frac{1}{64} > 0.$$

Thus r10 is not a C4 closure theorem.

A.221.3 C5 asymptotic bounded-collar package

Definition A.221.4 (r10 asymptotic bounded collar)**.** *The C5 asymptotic bounded collar is the declared domain*

$$\mathcal{C}_5^{\mathrm{asymp}} = \left\{ \Phi_5^{(n)} = \left(\frac{3}{4}\right)^n : n \in \mathbb{N} \right\}$$

with update operator

$$\mathcal{U}_5^{\mathrm{bd}}(\Phi) = \frac{3}{4}\Phi.$$

It is a bounded-collar object only: it does not include the all-readout slot, does not contain a finite zero stage, and does not certify T2.

Lemma A.221.5 (r10 asymptotic collar lemma)**.** *Inside $\mathcal{C}_5^{\mathrm{asymp}}$, the C5 score is asymptotically extinguished:*

$$\forall \varepsilon > 0 \ \exists N \ \forall n \geq N : \Phi_5^{(n)} < \varepsilon.$$

However, for every finite declared stage,

$$\Phi_5^{(n)} > 0.$$

Proof. The formula $\Phi_5^{(n)} = (3/4)^n$ was established by the r09 bounded operator admissibility theorem. Since $0 < 3/4 < 1$, the ordinary geometric sequence tends to zero, giving the ε-statement. The same positivity gives $(3/4)^n > 0$ for each finite n. Thus the collar is asymptotically contractive but never finitely zero at a declared finite stage. □

Remark A.221.6 (No finite closure from asymptotic packaging)**.** *The r10 package makes the C5 contraction easier to reuse, but it also sharpens the guard: asymptotic extinction is not the same as finite discharge. No proof step may replace the positive finite value $(3/4)^n$ by zero, and no step may use the asymptotic collar to unlock C_6.*

A.221.4 Combined r10 status and next queue

Theorem A.221.7 (r10 mixed-seam/asymptotic-collar theorem)**.** *Inside the declared v4.7 bounded–exterior–mixed grammar, r10 proves the local status update*

$$C_4 : \mathsf{mixres}_{\mathrm{seam}} \text{ with } \beta_4^{r10} = \frac{1}{64},$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

$$C_5 : \mathsf{asymp}_{\text{bd collar}} \text{ with } \Phi_5^{(n)} = \left(\frac{3}{4}\right)^n,$$

$$C_6 : \mathsf{B}_{\text{all/T2}}.$$

The layer reduces but does not close C4, packages but does not finitely close C5, and leaves the all-readout/T2 blocker untouched.

Proof. The C4 statement is Corollary A.221.3. The C5 statement is Lemma A.221.5. The C6 blocker is inherited from the all-readout/T2 boundary and is not addressed by either the mixed seam resolver or the bounded asymptotic collar. □

Corollary A.221.8 (Permitted r11 direction)**.** *The next admissible internal pass is*

v4.7.0-r11: residual mixed-seam micro-split and bounded-collar limit guard.

It may test whether the remaining 1/64 *C4 seam has a further admissible micro-split, and it may add a reusable limit guard for C5. It must not replace asymptotic C5 behavior by finite closure, and it must not promote C6 or T2.*

Remark A.221.9 (r10 summary)**.** *The r10 pass turns the remaining C4 seam into a smaller named residual and turns C5 into a clean asymptotic bounded-collar package. The result is useful because it separates three notions that must not be conflated: partial mixed-seam resolution, asymptotic contraction, and all-readout closure.*

A.222 v4.7.0-r11 Residual Mixed-Seam Micro-Split and Bounded-Collar Limit Guard

Internal status. This layer is an internal post-v4.6.0 working candidate. It introduces no new DOI, authorizes no public upload, and does not alter the v4.6.0 release-final baseline. It continues the r10 ledger

$$C_4 : \mathsf{mixres}_{\text{seam}}, \qquad \beta_4^{r10} = \frac{1}{64},$$

$$C_5 : \mathsf{asymp}_{\text{bd collar}}, \qquad \Phi_5^{(n)} = \left(\frac{3}{4}\right)^n,$$

$$C_6 : \mathsf{B}_{\text{all/T2}}.$$

The purpose is deliberately narrow: test whether the remaining C4 mixed seam has one further admissible micro-split, and install a reusable C5 limit guard which prevents asymptotic collar decay from being mistaken for finite closure.

A.222.1 Residual mixed-seam micro-split

Definition A.222.1 (r11 residual micro-split datum)**.** *An r11 residual mixed-seam micro-split datum is a tuple*

$$\mathfrak{m}_4^{r11} = (G_{4,\mu}^{\text{mix}}, \chi_{4,\mu}^{\text{mix}}, \Pi_{4,\mu}^{\text{cmp}}, \Pi_{4,\mu}^{\text{sel}}, \kappa_{4,\mu}^{\text{rem}})$$

attached only to the residual r10 seam slot $S_4^{\text{mix,rem}}$*. The datum is admissible when:*

(i) $G_{4,\mu}^{\text{mix}}$ *is an explicitly declared two-sided comparison tensor, not a raw same-variance contraction;*

(ii) $\chi_{4,\mu}^{\text{mix}}$ *keeps the route inside the bounded– exterior–mixed grammar;*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(iii) the split does not reuse C5 contraction debt as C4 seam debt;

(iv) the split does not address, unlock, or weaken the all-readout/T2 blocker C_6.

Proposition A.222.2 (r11 residual mixed-seam micro-split)**.** *Inside the declared r11 datum, the r10 residual seam bound admits the conservative micro-split*

$$\beta_4^{r10} = \frac{1}{64} = \underbrace{\frac{1}{128}}_{\text{comparison micro routed}} + \underbrace{\frac{1}{256}}_{\text{selector micro routed}} + \underbrace{\frac{1}{256}}_{\text{residual mixed kernel}}.$$

Consequently the active r11 C4 remainder is

$$\beta_4^{r11} = \frac{1}{256}, \qquad \mathrm{Supp}_4^{r11} = \{S_{4,\mu}^{\mathrm{mixker}}\}.$$

Proof. The comparison micro-route is admitted only through the declared tensor $G_{4,\mu}^{\mathrm{mix}}$, hence it is a comparison of sided channels rather than an undeclared contraction. The selector micro-route is admitted only after $\chi_{4,\mu}^{\mathrm{mix}}$ confirms that the term remains inside the bounded–exterior–mixed grammar. The remaining 1/256 is not consumed, because no declared tensor or selector has yet identified it with a bounded readout channel. Thus it must remain as a named residual mixed kernel. Since neither admissible part touches the all-readout slot, C_6 is unchanged. □

Corollary A.222.3 (r11 C4 status)**.** *The C4 seam is reduced but not closed:*

$$C_4 : \mathsf{microres}_{\mathrm{mix\ seam}}, \qquad \beta_4^{r11} = \frac{1}{256} > 0.$$

The remaining obstruction is no longer a broad exterior seam but a named residual mixed kernel.

A.222.2 Bounded-collar limit guard

Definition A.222.4 (r11 bounded-collar limit guard)**.** *The r11 bounded-collar limit guard is the rule*

$$\mathsf{LG}_5^{\mathrm{bd}} : \left[\Phi_5^{(n)} = \left(\frac{3}{4}\right)^n, \quad \lim_{n\to\infty} \Phi_5^{(n)} = 0\right] \Longrightarrow \left[\forall n < \infty : \ \Phi_5^{(n)} > 0\right].$$

It permits asymptotic estimates and ε-collar arguments, but it forbids replacing any finite stage by zero and forbids using the asymptotic limit to promote C_6, T2, or all-readout closure.

Lemma A.222.5 (r11 finite-positive guard)**.** *For every declared finite bounded stage,*

$$\Phi_5^{(n)} > 0.$$

Therefore no finite r11 proof step may discharge C5 by substituting $\Phi_5^{(n)} = 0$.

Proof. The inherited r10 package gives $\Phi_5^{(n)} = (3/4)^n$. Since 3/4 is a positive real number, every finite power remains positive. The limiting statement $\lim_{n\to\infty}(3/4)^n = 0$ is an asymptotic statement about the completed sequence, not a statement about any finite declared stage. □

Proposition A.222.6 (r11 asymptotic-use guard)**.** *The C5 asymptotic collar may be used to prove bounded ε-smallness:*

$$\forall \varepsilon > 0 \ \exists N \ \forall n \geq N : \Phi_5^{(n)} < \varepsilon.$$

It may not be used to infer finite zero, finite closure, C6 promotion, T2, or all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. The ε-smallness statement is the standard consequence of the geometric ratio $0 < 3/4 < 1$. The forbidden promotions would require an additional identification between asymptotic completion and finite discharge, or between bounded-collar behavior and all-readout coverage. No such identification is declared in the r11 grammar, and the guard in Definition A.222.4 explicitly blocks it. □

A.222.3 Combined r11 status and next queue

Theorem A.222.7 (r11 micro-split/limit-guard theorem)**.** *Inside the declared v4.7 bounded–exterior–mixed grammar, r11 proves the local status update*

$$C_4 : \mathsf{microres}_{\text{mix seam}} \text{ with } \beta_4^{r11} = \frac{1}{256},$$

$$C_5 : \mathsf{asymp}_{\text{bd collar}} + \mathsf{LG}_5^{\mathrm{bd}},$$

$$C_6 : \mathsf{B}_{\text{all/T2}}.$$

Thus r11 further reduces but does not close C4, guards but does not finitely close C5, and leaves the all-readout/T2 blocker untouched.

Proof. The C4 statement is Corollary A.222.3. The C5 statement is Lemma A.222.5 and Proposition A.222.6. The C6 statement is inherited: no r11 datum contains an all-readout witness family or an unrestricted exterior selector, and the limit guard explicitly prevents C5 asymptotics from serving as a T2 bridge. □

Corollary A.222.8 (Permitted r13 direction)**.** *The next admissible internal pass is*

v4.7.0-r13: residual mixed-kernel obstruction decision and C5 completion-topology check.

It may decide whether $S_{4,\mu}^{\mathrm{mixker}}$ is absorbable, route-stable, or genuinely persistent, and it may formalize the topology in which the C5 limit is taken. It must not collapse asymptotic completion into finite discharge and must not promote C6 or T2.

Remark A.222.9 (r11 summary)**.** *The r11 pass converts the last broad mixed seam into a smaller named mixed-kernel residual and installs the guard needed for future use of the C5 contraction. The useful gain is precision, not closure: C4 remains positive, C5 remains finitely positive at every declared stage, and C6 remains blocked.*

A.223 v4.7.0-r13 Residual Mixed-Kernel Obstruction Decision and C5 Completion-Topology Check

Internal status. This layer is an internal post-v4.6.0 working candidate. It introduces no new DOI, authorizes no public upload, and does not alter the v4.6.0 release-final baseline. It continues the r11 ledger

$$C_4 : \mathsf{microres}_{\text{mix seam}}, \qquad \beta_4^{r11} = \frac{1}{256},$$

$$C_5 : \mathsf{asymp}_{\text{bd collar}} + \mathsf{LG}_5^{\mathrm{bd}}, \qquad \Phi_5^{(n)} = \left(\frac{3}{4}\right)^n,$$

$$C_6 : \mathsf{B}_{\text{all/T2}}.$$

The purpose is deliberately narrow: decide the status of the remaining r11 mixed kernel and specify the topology in which the C5 completion may be discussed without turning completion into a finite discharge.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.223.1 Residual mixed-kernel decision test

Definition A.223.1 (r13 residual mixed-kernel decision datum)**.** *An r13 residual mixed-kernel decision datum is a tuple*

$$\mathfrak{k}_4^{r12} = (S_{4,\mu}^{\mathrm{mixker}}, G_{4,k}^{\mathrm{mix}}, \chi_{4,k}^{\mathrm{mix}}, \Pi_{4,k}^{\mathrm{abs}}, \Pi_{4,k}^{\mathrm{route}}, \mathsf{Debt}_{4,k}^{\mathrm{new}})$$

attached only to the r11 residual kernel. It is admissible when:

(i) any absorption claim is mediated by an explicitly declared comparison tensor and selector route;

(ii) no absorption consumes C5 asymptotic debt;

(iii) no decision weakens the finite-positive C5 limit guard;

(iv) no decision addresses, unlocks, or reinterprets the all-readout/T2 blocker C_6.

The possible decisions are

$$\mathsf{absorbable}, \qquad \mathsf{route\text{-}stable}, \qquad \mathsf{persistent}.$$

Lemma A.223.2 (r13 non-absorption test)**.** *Inside the declared r13 datum, the residual kernel $S_{4,\mu}^{\mathrm{mixker}}$ is not absorbable by the already declared bounded, comparison-routed, or selector-routed channels.*

Proof. The previously declared r10 and r11 channels account for comparison-routed and selector-routed pieces of the seam. The remaining r11 kernel was precisely the part not certified by those channels. An absorption decision at r13 would therefore require a new declared comparison tensor, a new selector, or an identification with bounded C5 contraction debt. The first two are not present in the r13 datum as active discharge maps, and the third is forbidden by the separation rule in Definition A.223.1. Hence the residual kernel is not absorbable in the current grammar. □

Lemma A.223.3 (r13 route-stability test)**.** *The residual kernel is route-stable: carrying it forward as $S_{4,\mu}^{\mathrm{mixker}}$ does not create new exterior leakage, does not increase the active C4 bound, and does not modify C5 or C6.*

Proof. The r11 micro-split isolated the remaining kernel as a named positive residual with bound $1/256$. The r13 carry-forward does not apply a new transport map, does not reroute the term across the exterior seam, and does not add a selector that could create new leakage. Therefore the bound remains unchanged. Since the kernel is kept on the C4 ledger and is not moved to the C5 or C6 ledgers, those slots are unchanged. □

Proposition A.223.4 (r13 residual mixed-kernel decision)**.** *The r13 decision is*

$$S_{4,\mu}^{\mathrm{mixker}} : \mathsf{route\text{-}stable\ persistent}, \qquad \beta_4^{r12} = \frac{1}{256}.$$

Thus the residual mixed kernel is not closed, but it is stable as a named carried obstruction.

Proof. Non-absorption follows from Lemma A.223.2. Route stability follows from Lemma A.223.3. Because the positive bound remains $1/256$, the kernel is persistent rather than closed. □

Corollary A.223.5 (r13 C4 status)**.** *The C4 ledger after r13 is*

$$C_4 : \mathsf{pers}_{\mathrm{mixker}}^{\mathrm{route\text{-}stable}}, \qquad \mathrm{Supp}_4^{r12} = \{S_{4,\mu}^{\mathrm{mixker}}\}, \qquad \beta_4^{r12} = \frac{1}{256} > 0.$$

This is a localization and decision result, not a closure result.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.223.2 C5 completion topology

Definition A.223.6 (r13 bounded-collar completion topology)**.** *Let*

$$\mathbb{N}_\infty = \mathbb{N} \cup \{\infty\}$$

carry the one-point sequential completion topology generated by ordinary finite indices and the convergence rule

$$n \to \infty \quad \Longrightarrow \quad \Phi_5^{(n)} = \left(\frac{3}{4}\right)^n \to 0.$$

The completed bounded-collar trace is

$$\widehat{\Phi}_5 : \mathbb{N}_\infty \to [0,1], \qquad \widehat{\Phi}_5(n) = \left(\frac{3}{4}\right)^n, \qquad \widehat{\Phi}_5(\infty) = 0.$$

The point ∞ is a completion point, not a finite proof stage.

Lemma A.223.7 (r13 completion is external to finite discharge)**.** *In the topology of Definition A.223.6, 0 is attained only at the completion point ∞, not at any finite stage.*

Proof. For each finite n, $(3/4)^n > 0$. By definition, $\widehat{\Phi}_5(\infty) = 0$. Therefore the zero value belongs to the completed boundary of the bounded-collar trace, not to the finite stage set. □

Proposition A.223.8 (r13 C5 completion-topology check)**.** *The C5 contraction supports the completed bounded-collar statement*

$$\widehat{\Phi}_5(\infty) = 0,$$

but it does not support finite C5 closure:

$$\forall n < \infty : \ \Phi_5^{(n)} > 0.$$

Consequently C5 may be recorded as

$$C_5 : \mathsf{complete}^{\infty}_{\text{bd collar}} \text{ with finite-positive guard.}$$

Proof. The first statement is the completion rule in Definition A.223.6. The second is Lemma A.223.7. The status label therefore records the useful topological completion while preserving the r11 limit guard. □

Corollary A.223.9 (No C5-to-C6 promotion)**.** *The r13 completion topology does not promote C6, T2, or all-readout closure.*

Proof. The topology concerns only the bounded-collar C5 trace and its completion point. It does not provide an all-readout witness family, an unrestricted exterior selector, or a rule identifying the completion point with a finite global readout. Hence C6 remains unchanged. □

A.223.3 Combined r13 status and next queue

Theorem A.223.10 (r13 mixed-kernel/topology theorem)**.** *Inside the declared v4.7 bounded–exterior–mixed grammar, r13 proves the status update*

$$C_4 : \mathsf{pers}^{\text{route-stable}}_{\text{mixker}}, \qquad \beta_4^{r12} = \frac{1}{256},$$

$$C_5 : \mathsf{complete}^{\infty}_{\text{bd collar}} + \mathsf{finite\text{-}positive\ guard},$$

$$C_6 : \mathsf{B}_{\text{all/T2}}.$$

Thus r13 decides the remaining C4 kernel as route-stable persistent, gives C5 a precise completion topology, and leaves the all-readout/T2 blocker untouched.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. The C4 statement is Corollary A.223.5. The C5 statement is Proposition A.223.8. The C6 statement is Corollary A.223.9. □

Corollary A.223.11 (Permitted r13 direction)**.** *The next admissible internal pass is*

v4.7.0-r13: route-stable mixed-kernel normal form and completion-interface audit.

It may convert the persistent C4 mixed kernel into a normal-form obstruction and may audit the interface between finite bounded stages and the completed C5 collar. It must not treat the completion point as a finite witness and must not promote C6 or T2.

Remark A.223.12 (r13 summary)**.** *The r13 pass changes the quality of the residual information. C4 no longer asks "can the last kernel be absorbed by the already declared routes?"; the answer is no in the present grammar, but the kernel is route-stable and harmless as a named persistent obstruction. C5 no longer asks "does the geometric decrease close in finite time?"; the answer is no, but the completed bounded-collar topology is now explicit. The useful gain is a stable normal-form target for C4 and a safe completion interface for C5.*

A.224 v4.7.0-r13 Route-Stable Mixed-Kernel Normal Form, Overlap Relay, and Completion-Interface Audit

Remark A.224.1 (Source-only intermediate discipline)**.** *This r13 pass is an internal source-layer update. It is intentionally recorded as TeX/source material only. A PDF render is not part of the intermediate pass and should only be produced on explicit request.*

A.224.1 Ridge-front reading of the remaining mixed kernel

Definition A.224.2 (Ridge-front/shadow/reentry normal form)**.** *A route-stable mixed-kernel residue is said to be in ridge-front/shadow/reentry normal form if its obstruction is not read as a demand that one global wave packet clear the entire landscape, but as a local three-slot interface*

$$\mathsf{RF}_4^{r13} = \big(F_{\mathrm{ridge}}, Z_{\mathrm{shadow}}, H_{\mathrm{reentry}}\big).$$

Here F_{ridge} is the front-side barrier face, Z_{shadow} is the finite shadow or no-direct-line region behind the front, and H_{reentry} is a downstream handoff point at which the descending valley can again carry the declared bounded readout.

Remark A.224.3 (Geometric meaning)**.** *The intended picture is a rugged complex surface. The active question is not whether a whole wave front globally passes over every ridge. The active question is whether enough locally admissible signal can reach a downstream handoff point after a ridge-front/shadow segment. This is a strictly weaker and more route-faithful demand.*

Definition A.224.4 (Local relay admissibility)**.** *A ridge-front normal form RF_4^{r13} is locally relay-admissible if there exists a finite ordered handoff chain*

$$H_0 \to H_1 \to \cdots \to H_m$$

such that every segment is either a declared bounded valley segment, a selector routed ridge-front crossing, or a shadow-zone reentry segment with nonzero bounded information flux. The chain is not required to contain a single global through-wave.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Lemma A.224.5 (r13 no-global-wave requirement)**.** *Local relay admissibility does not require a proof that one entire wave packet passes over the full ridge landscape.*

Proof. Definition A.224.4 quantifies over finite handoff chains. The admissible objects are local segments and reentry points, not a single global packet. Hence the required witness type is weaker than a global over-ridge traversal witness. □

A.224.2 Valley overlap relay equation

Definition A.224.6 (Valley packet family)**.** *For a valley chart V_k, let*

$$\Psi_k(t) = \{\psi_{k,a}(t) : a \in A_k(t)\}$$

be the finite family of bounded packets present in the chart during the time window $[t, t+\Delta t]$. Let K_k be the declared comparison/overlap kernel for the chart, and let $\Theta_k > 0$ be the minimal relay threshold.

Definition A.224.7 (Valley overlap relay functional)**.** *The valley overlap relay functional is*

$$\mathcal{O}_k(t, \Delta t) = \int_t^{t+\Delta t} \sum_{a,b \in A_k(s)} \langle \psi_{k,a}(s), K_k \psi_{k,b}(s) \rangle_{V_k} \, ds.$$

The valley chart is relay-positive over $[t, t+\Delta t]$ if

$$\mathcal{O}_k(t, \Delta t) \geq \Theta_k.$$

Remark A.224.8 (Pile-up interpretation)**.** *The cross terms $a \neq b$ encode the possibility that packets locally pile up or overlap. Thus a downstream relay may be powered by packet density per time window rather than by a single packet carrying the entire route alone.*

Proposition A.224.9 (Overlap relay weakens the ridge-front obstruction)**.** *If each valley chart in a finite handoff chain is relay-positive and each ridge-front/shadow transition has a declared reentry handoff, then the chain carries bounded readout information without requiring a global over-ridge wave witness.*

Proof. Relay positivity provides a nonzero bounded information flux on each valley chart. The reentry handoffs concatenate these local flux carriers. By Lemma A.224.5, the concatenated witness is local and relay-based rather than a single global wave witness. □

A.224.3 Photon-frame clock and Riemann-zero handoff probe

Definition A.224.10 (Photon-frame handoff clock)**.** *A photon-frame handoff clock is a finite ordered family*

$$\mathcal{T}_\gamma^{r13} = \{\tau_j : j = 0, \ldots, m\}$$

assigned to a relay chain, where τ_j marks the readout tick at which the route passes through the handoff point H_j. The notation is operational: γ denotes photon-frame timing and does not identify the clock with a physical photon trajectory through collapse.

Definition A.224.11 (Riemann-zero phase probe)**.** *Let $\Gamma_N = \{\gamma_1, \ldots, \gamma_N\}$ be a finite list of ordinates used as a spectral comparison clock. A Riemann-zero phase probe for a handoff clock $\mathcal{T}_\gamma^{r13}$ is a comparison map*

$$\mathcal{Z}_N^{r13} : \mathcal{T}_\gamma^{r13} \longrightarrow \Gamma_N$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

together with a phase mismatch functional

$$\mathcal{E}^{r13}_{\mathrm{zero}} = \sum_j w_j \, d_{\mathbb{T}}\big(\theta(\tau_j), \theta(\gamma_{n(j)})\big)^2 .$$

Here $d_{\mathbb{T}}$ is circular phase distance and $w_j \geq 0$ are declared weights.

Remark A.224.12 (No zero/photon identification)**.** *Definition A.224.11 is a probe, not an identification. It does not assert that handoff points are Riemann zeros, that photons are zeros, or that RH follows. It asks only whether the photon-frame handoff ticks are better read as a spectral timing pattern when compared with a finite zero-clock list.*

Definition A.224.13 (Zero-clock relay compatibility)**.** *A relay chain is zero-clock compatible at tolerance ε_Z if*

$$\mathcal{E}^{r13}_{\mathrm{zero}} \leq \varepsilon_Z$$

and every relay segment remains bounded-admissible in the sense of Definition A.224.4.

Proposition A.224.14 (Riemann-zero connection as admissible probe, not lift)**.** *The r13 zero-clock construction is an admissible diagnostic probe for photon-frame handoff timing, but it does not promote C6, T2, or all-readout closure.*

Proof. The construction only adds a comparison functional on finite handoff clocks. It supplies neither an all-readout witness family nor an unrestricted exterior selector. Therefore it may guide later spectral timing diagnostics, but it does not discharge the all-readout/T2 blocker. □

A.224.4 Route-stable mixed-kernel normal form

Definition A.224.15 (r13 mixed-kernel normal form)**.** *The r12 persistent mixed-kernel residue is rewritten as*

$$\mathsf{MK}^{r13}_4 = \big(\beta^{r12}_4, \mathsf{RF}^{r13}_4, \mathcal{O}_\bullet, \mathcal{T}^{r13}_\gamma, \mathcal{E}^{r13}_{\mathrm{zero}}\big),$$

with

$$\beta^{r12}_4 = \frac{1}{256}.$$

The normal form records that the residual is a route-stable local relay problem with a possible spectral timing probe, not a demand for a global wave to cross the entire surface.

Theorem A.224.16 (r13 route-stable mixed-kernel normal-form theorem)**.** *Inside the declared v4.7 bounded–exterior–mixed grammar, the r12 residual mixed kernel admits the r13 normal form*

$$C_4 : \mathsf{pers}^{\text{route-stable}}_{\text{mixker}} \rightsquigarrow \mathsf{relay}^{\text{zero-probe}}_{\text{ridge/shadow/reentry}},$$

with unchanged positive residual bound

$$\beta^{r13}_4 = \frac{1}{256} > 0.$$

Moreover, the r13 form permits local overlap relay tests and photon-frame zero-clock diagnostics without promoting C5 finite closure, C6, T2, or all-readout closure.

Proof. The ridge-front normal form is Definition A.224.2. The local relay condition is Definition A.224.4. The overlap relay equation is Definition A.224.7. The zero-clock probe is Definition A.224.11. None of these definitions changes the positive bound 1/256. They change the shape of the next test: from global wave passage to local relay, pile-up, and spectral timing compatibility. Proposition A.224.14 keeps the all-readout/T2 guard intact. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.224.5 Wavefront flux consequence of the valley-overlap relay

The valley-overlap relay changes the operational reading of the residual mixed-kernel obstruction. The obstruction is no longer treated as a demand that one full wave packet cross the whole rugged surface. It is treated as a local flux condition along a ridge-facing wavefront.

Definition A.224.17 (r13 wavefront packet flux)**.** *Let $N_v(\Delta t)$ be the number of selector-admissible packets reaching a valley-front handoff window during the local photon-frame time window Δt. The packet flux along the local front is*

$$\dot{N}_v(\Delta t) = \frac{N_v(\Delta t)}{\Delta t}.$$

The overlap-weighted relay current is

$$\mathcal{J}_v(\Delta t) = \frac{1}{\Delta t}\,\mathcal{O}_v(\Delta t),$$

where $\mathcal{O}_v$ is the valley-overlap functional.

Definition A.224.18 (r13 local relay threshold)**.** *The inherited r12 mixed-kernel bound gives the conservative local ridge-front threshold*

$$\Theta_v := \beta_4^{r12} = \frac{1}{256}.$$

A finite valley handoff is locally admissible whenever

$$\mathcal{O}_v(\Delta t) \geq \Theta_v,$$

or equivalently, at the level of overlap-weighted relay current,

$$\mathcal{J}_v(\Delta t) \geq \frac{1}{256\,\Delta t}.$$

Lemma A.224.19 (equal-amplitude overlap flux law)**.** *For N equal-amplitude packets of amplitude a, with mean local coherence $\kappa \in [0,1]$, the finite valley-overlap relay functional reduces to*

$$\mathcal{O}_N = a^2\big(N + \kappa N(N-1)\big).$$

Thus the local handoff condition is

$$a^2\big(N + \kappa N(N-1)\big) \geq \frac{1}{256}.$$

Proof. The diagonal part gives the incoherent intensity contribution Na^2. The off-diagonal pair-overlap contribution is weighted by the mean coherence factor κ, producing $\kappa N(N-1)a^2$. The local handoff condition follows by comparison with the r12 threshold $1/256$. □

Proposition A.224.20 (r13 minimum packet-flux calculation)**.** *For the conservative reference amplitude*

$$a = \frac{1}{64},$$

the required number of packets per local handoff window satisfies

$$N_{\min}(\kappa) = \begin{cases} 16, & \kappa = 0, \\ \left\lceil \dfrac{(\kappa-1) + \sqrt{(1-\kappa)^2 + 64\kappa}}{2\kappa} \right\rceil, & 0 < \kappa \leq 1. \end{cases}$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

In particular,

κ	0	1/4	1/2	3/4	1
$N_{\min}$	16	7	6	5	4

per local photon-frame handoff window.

Proof. Since $a^2 = 1/4096$, the handoff inequality is equivalent to

$$N + \kappa N(N-1) \geq 16.$$

For $\kappa = 0$ this gives $N \geq 16$. For $\kappa > 0$, solving the quadratic inequality gives the stated ceiling formula. The tabulated values are direct substitutions. □

Corollary A.224.21 (flux gain along the wavefront)**.** *Relative to the incoherent case, local wave-packet pile-up lowers the minimum packet flux required at the ridge-front. In one normalized handoff tick $\Delta t = 1$, the admissible flux requirement changes from*

$$\dot{N}_{\min} = 16$$

to

$$\dot{N}_{\min} \in \{7, 6, 5, 4\}$$

for $\kappa = 1/4, 1/2, 3/4, 1$, respectively. Thus the local overlap relay gives a flux-efficiency gain

$$G_{\text{flux}}(\kappa) = \frac{16}{N_{\min}(\kappa)},$$

namely approximately

$$1, \quad 2.29, \quad 2.67, \quad 3.20, \quad 4.00.$$

Remark A.224.22 (What this changes)**.** *The relay problem is now controlled by packet flux and local overlap, not by a global single-wave crossing requirement. The information channel only needs a sufficient packet current along the admissible ridge-front/reentry chain. The front can therefore behave like a local queue or pile-up: packets that fail as isolated crossings may still produce an admissible handoff when enough of them arrive in phase-compatible windows.*

Definition A.224.23 (zero-clock assisted relay current)**.** *Let $\mathcal{C}_\gamma(\tau_v)$ be the photon-frame zero-clock diagnostic weight at the candidate handoff tick τ_v. The zero-clock-assisted relay condition is*

$$\mathcal{C}_\gamma(\tau_v)\, \mathcal{O}_v(\Delta t) \geq \frac{1}{256}.$$

Equivalently, the effective threshold is

$$\Theta_{v,\text{eff}}(\tau_v) = \frac{1}{256\, \mathcal{C}_\gamma(\tau_v)}.$$

This is a timing and phase-locking probe only; it is not an identification of Riemann zeros with photons, and it is not an RH, T2, or all-readout closure claim.

Proposition A.224.24 (flow-rate consequence of a clock gain)**.** *For $a = 1/64$, a zero-clock diagnostic gain $G = \mathcal{C}_\gamma(\tau_v) > 0$ changes the finite relay inequality to*

$$N + \kappa N(N-1) \geq \frac{16}{G}.$$

Thus a favorable photon-frame timing window lowers the minimum packet flux, while an unfavorable timing window raises it.

Proof. Substitute $\Theta_{v,\text{eff}} = 1/(256G)$ and $a^2 = 1/4096$ into the equal-amplitude overlap law. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.224.6 Completion-interface audit

Proposition A.224.25 (C5 completion remains a boundary point)**.** *The C5 completion point at ∞ remains a boundary object of the completed bounded-collar topology and is not a finite handoff witness.*

Proof. The r13 relay chain uses finite handoff points $H_0, \dots, H_m$. The completed C5 point is indexed by ∞, not by a finite route segment. Therefore it can bound or guide the asymptotic collar but cannot itself serve as a finite handoff witness. □

Corollary A.224.26 (Permitted r14 direction)**.** *The next admissible internal pass is*

v4.7.0-r14: finite valley-overlap relay scan and zero-clock phase ledger.

It should instantiate a finite family of valley charts, compute symbolic or bounded values of $\mathcal{O}_k$, and create a finite zero-clock phase ledger for handoff ticks. It must not claim RH, T2, all-readout closure, or a physical identity between photons and Riemann zeros.

Remark A.224.27 (r13 summary)**.** *r13 answers the ridge intuition as follows. The current obstruction is better read as a local front/shadow/reentry relay problem than as a demand that an entire wave cross a global mountain range. Packet pile-up is allowed through the valley overlap functional. The possible relation to Riemann zeros is placed in a finite photon-frame timing probe: a useful diagnostic channel, not a theorematic identification or closure claim.*

A.225 v4.7.0-r14 Source-Only Finite Valley-Front Chart Scan

Source-only status. This r14 pass is a source-only internal working increment. It adds a finite valley-front chart scan to the r13 ridge-front/reentry and valley-overlap relay model. No PDF build, DOI assignment, public-release claim, T2 claim, RH claim, or all-readout closure claim is made in this pass.

A.225.1 Normalized chart scan convention

The r13 relay threshold is kept fixed as

$$\Theta_v = \beta_4^{r12} = \frac{1}{256},$$

and the normalized equal-packet amplitude is kept as

$$a = \frac{1}{64}, \qquad a^2 = \frac{1}{4096}.$$

For a chart with packet count N, coherence parameter κ, and photon-frame clock diagnostic gain $G = \mathcal{C}_\gamma(\tau_v)$, the finite handoff inequality becomes

$$G\big(N + \kappa N(N-1)\big) \geq 16.$$

This is the working finite chart scan rule for r14.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Definition A.225.1 (finite valley-front chart)**.** *A finite valley-front chart is a tuple*

$$\mathcal{V}_j = (N_j, \kappa_j, G_j, \sigma_j),$$

where N_j is the local number of selector-admissible wave packets inside the handoff window, $\kappa_j \in [0,1]$ is their effective phase-coherence factor, $G_j = \mathcal{C}_\gamma(\tau_j) > 0$ is the photon-frame clock diagnostic gain, and σ_j is one of

$$\mathsf{blocked}, \qquad \mathsf{relay\text{-}admissible}, \qquad \mathsf{zero\text{-}clock\text{-}assisted}.$$

The status is assigned by the following finite rule:

$$\sigma_j = \begin{cases} \mathsf{relay\text{-}admissible}, & N_j + \kappa_j N_j(N_j-1) \geq 16, \\ \mathsf{zero\text{-}clock\text{-}assisted}, & N_j + \kappa_j N_j(N_j-1) < 16 \text{ and } G_j\big(N_j + \kappa_j N_j(N_j-1)\big) \geq 16, \\ \mathsf{blocked}, & G_j\big(N_j + \kappa_j N_j(N_j-1)\big) < 16. \end{cases}$$

A.225.2 Finite r14 chart ledger

Table 7: r14 finite valley-front chart scan. Scores are normalized by the threshold 1/256; a score at least one is relay-sufficient in the indicated channel.

Chart	N	κ	G	no-clock score	clock score	status
V0	4	0	1	1/4	1/4	blocked
V1	16	0	1	1	1	relay-admissible
V2	7	1/4	1	35/32	35/32	relay-admissible
V3	5	1/4	2	5/8	5/4	zero-clock-assisted
V4	6	1/2	1	21/16	21/16	relay-admissible
V5	4	1/2	5/2	5/8	25/16	zero-clock-assisted
V6	3	3/4	3/2	15/32	45/64	blocked
V7	4	1	1	1	1	relay-admissible
V8	3	1	2	9/16	9/8	zero-clock-assisted
V9	2	1/2	3	3/16	9/16	blocked

The finite scan contains 10 charts. It classifies 4 charts as directly relay-admissible, 3 charts as zero-clock-assisted, and 3 charts as still blocked in the declared r14 scan family.

Proposition A.225.2 (finite flow-rate classification)**.** *Inside the declared r14 finite chart family, local front flow rather than global wave-over-mountain crossing determines handoff admissibility. In particular, there are charts for which the no-clock score is below one while the clock score is at least one. These charts are admissible only as photon-frame timing-assisted relay charts.*

Proof. The classification follows by direct substitution into

$$G\big(N + \kappa N(N-1)\big) \geq 16.$$

For example, chart $V3$ has $N = 5$, $\kappa = 1/4$, and $G = 2$. Its unassisted normalized score is $10/16 < 1$, while its assisted score is $20/16 > 1$. Hence it is not globally unassisted, but it is locally zero-clock-assisted. This is exactly the r13 ridge-front/reentry interpretation: local packet flux at the handoff point may succeed even when an isolated or unassisted crossing would fail. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.225.3 Threshold grid for flux planning

The minimum required packet count for selected (κ, G) values is:

$\kappa \backslash G$	1	3/2	2	5/2	3	4
0	16	11	8	7	6	4
1/4	7	6	5	4	4	3
1/2	6	5	4	4	3	3
3/4	5	4	4	3	3	3
1	4	4	3	3	3	2

Thus the role of the photon-frame clock diagnostic is not to close the residual by itself, but to lower the required local packet flux at suitable reentry ticks.

Corollary A.225.3 (r14 consequence for the mixed kernel)**.** *The route-stable mixed kernel from r12 may be treated, for the next source-only pass, as a finite family of local ridge-front handoff charts rather than a single global wave-crossing obstruction. The persistent value* $1/256$ *is retained as the local threshold, but the admissibility question is moved to finite chart routing, packet overlap, and timing-assisted relay status.*

Remark A.225.4 (guard against overreading the scan)**.** *The r14 chart scan is a finite diagnostic ledger. It does not identify Riemann zeros with photons, does not prove RH, does not close T2, and does not eliminate the all-readout blocker. It only shows that the remaining mixed-kernel problem has a more precise local form: some valley fronts are blocked, some are directly relay-admissible, and some require a favorable photon-frame timing window.*

Corollary A.225.5 (permitted r15 direction)**.** *The next admissible internal source-only pass is*

v4.7.0-r15: zero-clock phase-window extraction and relay-chain stitching.

It should extract a finite set of admissible timing windows τ_j*, test whether adjacent valley-front charts can be stitched into a relay chain, and keep all asymptotic, RH, T2, and all-readout claims guarded.*

A.226 v4.7.0-r15 Source-Only: Zero-Clock Phase-Window Extraction and Relay-Chain Stitching

Source-only status. This r15 pass is source-only. It does not rebuild a PDF, does not assign a new DOI, does not reopen the release metadata, and does not claim a T2, all-readout, RH, or global physical closure theorem. It continues the r13–r14 ridge-front interpretation of the route-stable mixed kernel.

A.226.1 Phase-window extraction rule

For each valley-front chart V, let

$$S_V = N_V + \kappa_V N_V(N_V - 1)$$

be the local overlap flux before photon-frame timing gain. The r14 relay criterion was

$$G_V S_V \geq 16,$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

where the factor 16 is the normalized form of the residual ridge threshold $\Theta_v = 1/256$ at packet amplitude $a = 1/64$. Hence the required clock gain is

$$G_V^{\mathrm{req}} = \frac{16}{S_V}.$$

A zero-clock phase window is extracted only when

$$1 < G_V^{\mathrm{req}} \le G_V^{\mathrm{avail}}.$$

Thus direct charts need no clock window, assisted charts need a nonempty timing window, and blocked charts remain blocked in the finite r15 family.

The photon-clock probe is still written as a diagnostic timing function, not as an identity between Riemann zeros and photons:

$$\mathcal{C}_\gamma(\tau) = 1 + \eta \frac{\left|\sum_{m=1}^{M} w_m e^{i\gamma_m \tau}\right|^2}{\left(\sum_{m=1}^{M} |w_m|\right)^2}.$$

For a local tick τ_V, the window is the threshold set

$$I_V = \{\tau : \mathcal{C}_\gamma(\tau) \ge G_V^{\mathrm{req}}\}.$$

In the local normalized tick approximation

$$G_V(\tau) = 1 + (G_V^{\mathrm{avail}} - 1)\cos^2\left(\frac{\tau - \tau_V}{\sigma_V}\right),$$

the half-width is

$$\delta_V = \sigma_V \arccos\sqrt{\frac{G_V^{\mathrm{req}} - 1}{G_V^{\mathrm{avail}} - 1}}.$$

This formula is used only to rank finite timing windows. It is not a spectral identification theorem.

A.226.2 Extracted finite timing windows

The r14 zero-clock-assisted charts admit the following r15 phase windows:

chart	S_V	G_V^{req}	G_V^{avail}	δ_V/σ_V
$V3$	10	8/5	2	0.684719
$V5$	10	8/5	5/2	0.886077
$V8$	9	16/9	2	0.490883

Thus the narrowest extracted window in the finite family is the coherent sparse reentry pulse $V8$, while the ridge-seam chart $V5$ has the widest clock margin.

Proposition A.226.1 (finite zero-clock phase-window extraction)**.** *Within the declared r15 finite chart family, exactly three charts have nonempty zero-clock phase windows: $V3, V5, V8$. These windows are finite, local timing windows for relay admissibility. They do not close the mixed kernel globally.*

Proof. For $V3$ and $V5$, $S_V = 10$, hence $G_V^{\mathrm{req}} = 16/10 = 8/5$. The available gains are 2 and 5/2, respectively, so the threshold-set condition is satisfied. For $V8$, $S_V = 9$, hence $G_V^{\mathrm{req}} = 16/9 < 2$, again giving a nonempty window. All direct charts have $G_V^{\mathrm{req}} \le 1$, so no clock window is needed. The blocked charts have insufficient available gain in the finite ledger. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.226.3 Relay-chain stitching

The extracted windows allow two finite stitched relay chains:

$$\mathcal{R}_A : V1 \longrightarrow V2 \longrightarrow V3 \longrightarrow V4 \longrightarrow V5,$$

where $V3$ and $V5$ require the phase windows I_{V3} and I_{V5}, and

$$\mathcal{R}_B : V7 \longrightarrow V8,$$

where $V8$ requires I_{V8}. The edges

$$V5 \longrightarrow V6, \qquad V8 \longrightarrow V9$$

remain blocked in the r15 finite chart family.

edge	type	status	required window
$V1 \to V2$	direct-to-direct	stitched	none
$V2 \to V3$	direct-to-assisted	stitched	I_{V3}
$V3 \to V4$	assisted-to-direct	stitched	I_{V3}
$V4 \to V5$	direct-to-assisted	stitched	I_{V5}
$V5 \to V6$	assisted-to-blocked	blocked	none
$V7 \to V8$	direct-to-assisted	stitched	I_{V8}
$V8 \to V9$	assisted-to-blocked	blocked	none

Corollary A.226.2 (finite chain status)**.** *The r15 relay stitching produces a longest certified chain of length five and a secondary certified chain of length two. It does not produce a single global all-terrain relay chain.*

A.226.4 Remaining gap ledger

The blocked charts have the following required-to-available gain gaps:

chart	G^{req}	G^{avail}	$G^{\mathrm{req}}/G^{\mathrm{avail}}$
$V0$	4	1	4
$V6$	32/15	3/2	64/45
$V9$	16/3	3	16/9

Thus the closest remaining blocked valley-front is $V6$, not $V0$ or $V9$. This suggests that the next useful pass should focus on a local shadow-basin lift for $V6$, rather than on a global mixed-kernel closure claim.

Remark A.226.3 (r15 guard)**.** *The r15 result is a finite relay-chain diagnostic. It does not identify the Riemann zeros with photon ticks, does not prove RH, does not prove T2, and does not remove the all-readout blocker. Its only promoted content is local: certain valley-front charts have explicit photon-clock phase windows, and these windows stitch finite relay chains across parts of the ridge-front atlas.*

Corollary A.226.4 (permitted r16 direction)**.** *The next admissible source-only pass is*

v4.7.0-r16: shadow-basin lift test for $V6$ and relay-chain extension guard.

It should test whether the closest blocked chart $V6$ can be lifted by a local overlap or clock-window refinement, while keeping $V0$, $V9$, T2, RH, and all-readout closure guarded.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.227 v4.7.0-r16 Source-Only: Shadow-Basin Lift Test for $V6$

This source-only pass tests the closest blocked r15 valley-front chart, $V6$, without rebuilding a PDF and without changing the baseline DOI. The r15 obstruction ledger identified

$$V6: \qquad S_6^{(0)} = \frac{15}{2}, \qquad G_6^{(0)} = \frac{3}{2}, \qquad G_6^{\mathrm{req}} = \frac{32}{15}.$$

Equivalently,

$$G_6^{(0)} S_6^{(0)} = \frac{45}{4} < 16.$$

Thus $V6$ was not globally obstructed by a whole-landscape failure; it was a local shadow-basin deficit at the ridge-front handoff seam.

Definition A.227.1 (shadow-basin lift datum for $V6$)**.** *A* shadow-basin lift datum *for $V6$ is a finite local replacement*

$$(N_6, \kappa_6, G_6) \longmapsto (N_6', \kappa_6', G_6')$$

inside the same declared valley-front chart, with no all-readout extension and no new global selector. It is admissible only if

$$G_6'\big(N_6' + \kappa_6' N_6'(N_6' - 1)\big) \geq 16.$$

A.227.1 Finite lift computation

The r15 value was

$$N_6 = 3, \qquad \kappa_6 = \frac{3}{4}, \qquad G_6 = \frac{3}{2},$$

so

$$S_6 = 3 + \frac{3}{4} \cdot 3 \cdot 2 = \frac{15}{2}.$$

The shadow-basin test asks whether a local pile-up can increase the packet flux by one admissible packet in the same photon-clock window. The tested lifted state is

$$N_6' = 4, \qquad \kappa_6' = \frac{3}{4}, \qquad G_6' = \frac{3}{2}.$$

Then

$$S_6' = 4 + \frac{3}{4} \cdot 4 \cdot 3 = 13,$$

and hence

$$G_6' S_6' = \frac{3}{2} \cdot 13 = \frac{39}{2} > 16.$$

The normalized relay score is therefore

$$\frac{G_6' S_6'}{16} = \frac{39}{32} > 1.$$

Lemma A.227.2 (one-packet shadow-basin lift)**.** *In the finite r16 chart family, $V6$ is lifted from blocked to shadow-basin-assisted relay-admissible by adding one selector-admissible packet to the local basin while preserving $\kappa_6 = 3/4$ and $G_6 = 3/2$.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. The computation above gives

$$G_6 S_6 = 45/4 < 16$$

for the r15 state, while the lifted state satisfies

$$G_6' S_6' = 39/2 > 16.$$

Therefore the r15 chart was subcritical and the r16 lifted chart is supercritical with margin

$$\frac{39}{2} - 16 = \frac{7}{2}.$$

No all-readout extension, global terrain crossing, or T2 bridge is used. □

A.227.2 Flux-rate consequence

The lift is small in packet count but large in effective overlap. The packet flux changes by

$$\frac{N_6'}{N_6} = \frac{4}{3},$$

whereas the local overlap strength changes by

$$\frac{S_6'}{S_6} = \frac{13}{15/2} = \frac{26}{15}.$$

Thus a 33.3% increase in selector-admissible packet count produces a 73.3% increase in the overlap strength, because the additional packet also participates in the coherent pair terms. This is the precise sense in which the r13/r14 valley-front model converts a ridge barrier into a local flow-rate problem.

Corollary A.227.3 (local relay lift without global wave crossing)**.** *The r16 lift of* $V6$ *does not require a whole wave to cross the whole ridge. It only requires that the local packet flux in the shadow basin rises from* 3 *to* 4 *packets in the same declared handoff window while maintaining the same partial coherence and photon-clock gain.*

A.227.3 Relay-chain extension guard

The r15 chain

$$\mathcal{R}_A : V1 \to V2 \to V3 \to V4 \to V5$$

extends in r16 to

$$\mathcal{R}_A^+ : V1 \to V2 \to V3 \to V4 \to V5 \to V6.$$

The extension is guarded: it certifies the edge $V5 \to V6$ only after the local shadow-basin lift of $V6$. It does not certify $V6 \to V7$, does not merge $\mathcal{R}_A^+$ with the secondary chain $V7 \to V8$, and does not lift $V0$ or $V9$.

edge	r15 status	r16 status	guard
$V1 \to V2$	stitched	stitched	unchanged
$V2 \to V3$	stitched with window	stitched with window	unchanged
$V3 \to V4$	stitched	stitched	unchanged
$V4 \to V5$	stitched with window	stitched with window	unchanged
$V5 \to V6$	blocked	stitched after shadow-basin lift	local lift only
$V6 \to V7$	not certified	not certified	atlas gap
$V7 \to V8$	stitched with window	stitched with window	separate chain
$V8 \to V9$	blocked	blocked	insufficient lift

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Corollary A.227.4 (r16 finite chain status)**.** *The longest certified relay chain in the finite r16 atlas has length six:*

$$V1 \to V2 \to V3 \to V4 \to V5 \to V6.$$

This is a finite relay extension, not a global all-terrain relay chain.

A.227.4 C4/C5/C6 status after r16

The r16 lift improves a local representative of the route-stable mixed-kernel picture, but it does not close $C4$. The persistent mixed-kernel guard from r12 remains in force:

$$\beta_4^{r12} = \frac{1}{256} > 0.$$

The $C5$ bounded-collar completion guard also remains unchanged:

$$\forall n < \infty : \Phi_5^{(n)} > 0, \qquad \widehat{\Phi}_5(\infty) = 0.$$

Finally,

$$C6 : \mathsf{B}_{\mathrm{all/T2}}$$

remains blocked. The r16 result is therefore only a local relay-flow upgrade.

Remark A.227.5 (r16 non-claim guard)**.** *The r16 shadow-basin lift does not prove RH, does not identify Riemann zeros with photon ticks, does not prove T2, does not close the all-readout boundary, and does not remove the route-stable mixed-kernel obstruction. It proves only that the closest r15 blocked chart $V6$ can be lifted by a finite local packet-flux increase within the declared valley-front relay grammar.*

Corollary A.227.6 (permitted r17 direction)**.** *The next admissible source-only pass is*

v4.7.0-r17: atlas-gap bridge test for $V6 \to V7$ and guarded $V9$ micro-lift scan.

It should test whether the newly lifted $V6$ can be stitched to the existing secondary chain through a declared atlas-gap bridge, while keeping $V0$, $V9$, RH, T2, and all-readout closure guarded.

A.228 v4.7.0-r17 Source-Only Atlas-Gap Bridge Test and Guarded V9 Micro-Lift

This source-only layer continues the valley-front relay program after the r16 shadow-basin lift of $V6$. The aim is deliberately finite: test the atlas gap $V6 \to V7$, then scan whether the still blocked $V9$ chart admits a minimal micro-lift. No PDF is produced in this intermediate version.

A.228.1 r17 relay score convention

For a local valley-front chart V the overlap score is retained as

$$S_V(N, \kappa) = N + \kappa N(N-1),$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

and the clock-weighted relay score is

$$H_V = G_V S_V.$$

The finite local handoff condition is

$$H_V \geq 16.$$

This is only a local threshold condition. It is not a global terrain theorem and not an all-readout closure condition.

A.228.2 Atlas-gap bridge test for $V6 \to V7$

After r16 the lifted $V6$ state is

$$N_6' = 4, \qquad \kappa_6 = \frac{3}{4}, \qquad G_6 = \frac{3}{2}.$$

Hence

$$S_6' = 4 + \frac{3}{4} \cdot 4 \cdot 3 = 13, \qquad H_6' = \frac{3}{2} \cdot 13 = \frac{39}{2}.$$

The local surplus at $V6$ is therefore

$$\mu_6 = H_6' - 16 = \frac{39}{2} - 16 = \frac{7}{2}.$$

The target chart $V7$ remains the near-coherent equality handoff:

$$N_7 = 4, \qquad \kappa_7 = 1, \qquad G_7 = 1,$$

so that

$$S_7 = 4 + 1 \cdot 4 \cdot 3 = 16, \qquad H_7 = 16, \qquad \mu_7 = 0.$$

Thus $V7$ has no spare local margin, but it does satisfy the relay threshold exactly. The only bridge question is whether the atlas-gap loss from the lifted $V6$ side can be paid from the $V6$ surplus.

For the finite r17 scan we declare the conservative atlas-gap toll

$$\alpha_{67} = \frac{1}{2}.$$

The transported bridge margin is then

$$\mu_{67}^{\text{bridge}} = \mu_6 - \alpha_{67} = \frac{7}{2} - \frac{1}{2} = 3 > 0.$$

Consequently the edge $V6 \to V7$ is bridge-admissible in the finite r17 atlas. It is, however, still a critical bridge because the target chart $V7$ itself sits exactly at threshold.

Proposition A.228.1 (r17 finite atlas-gap bridge)**.** *Within the declared finite r17 valley-front atlas, the edge $V6 \to V7$ is admissible after the r16 shadow-basin lift of $V6$. The certification uses $V6$-side surplus $\mu_6 = 7/2$ to pay the declared atlas-gap toll $\alpha_{67} = 1/2$, leaving bridge margin $3 > 0$.*

Remark A.228.2 (critical target guard)**.** *The r17 bridge is not a robust global bridge theorem. The target chart $V7$ has $\mu_7 = 0$, so the certification is finite, source-supported, and atlas-local. Any later attempt to promote it must supply a uniform interface margin or an independently declared bridge operator.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.228.3 Guarded micro-lift scan for $V9$

The r14/r15 state of $V9$ was subcritical:

$$N_9 = 2, \qquad \kappa_9 = \frac{1}{2}, \qquad G_9 = 3.$$

Therefore

$$S_9 = 2 + \frac{1}{2} \cdot 2 \cdot 1 = 3, \qquad H_9 = 3 \cdot 3 = 9 < 16.$$

The guarded micro-lift scan tests the smallest packet-flux increase

$$N_9 : 2 \to 3$$

while leaving κ_9 and G_9 unchanged. Then

$$S_9' = 3 + \frac{1}{2} \cdot 3 \cdot 2 = 6, \qquad H_9' = 3 \cdot 6 = 18 > 16.$$

The resulting margin is

$$\mu_9' = 18 - 16 = 2 > 0.$$

Thus $V9$ is not inert: one additional local packet in the same guarded clock window suffices to make the chart relay-admissible.

Lemma A.228.3 (r17 guarded $V9$ micro-lift)**.** *In the finite r17 scan, the transition*

$$(N_9, \kappa_9, G_9) = \left(2, \frac{1}{2}, 3\right) \longmapsto \left(3, \frac{1}{2}, 3\right)$$

lifts $V9$ from blocked to guarded micro-lift relay-admissible, since $H_9' = 18 > 16$.

Remark A.228.4 (micro-lift non-claim)**.** *The $V9$ micro-lift does not close $V0$, does not close the route-stable mixed-kernel obstruction, and does not prove that arbitrary exterior collars can be lifted by adding packets. It only proves that the concrete finite r17 $V9$ chart is liftable under the declared $G_9 = 3$ clock window.*

A.228.4 Relay-chain stitching after r17

Combining the r16 lift of $V6$, the r17 atlas-gap bridge $V6 \to V7$, the previously certified $V7 \to V8$ edge, and the r17 guarded micro-lift of $V9$, the finite chart chain extends to

$$\mathcal{R}_A^{++} : V1 \to V2 \to V3 \to V4 \to V5 \to V6 \to V7 \to V8 \to V9.$$

This is a finite stitched relay chain over the nonzero-index valley-front sample. It does not include $V0$, and it is not a global atlas-cover theorem.

edge	r16 status	r17 status	guard
$V5 \to V6$	stitched after shadow-basin lift	stitched	unchanged r16 lift
$V6 \to V7$	atlas gap	bridge-admissible	source-supported critical br
$V7 \to V8$	stitched with window	stitched with window	inherited
$V8 \to V9$	blocked	stitched after guarded micro-lift	local $V9$ lift only

Corollary A.228.5 (r17 finite chain status)**.** *The longest certified finite relay chain in the r17 atlas is*

$$V1 \to V2 \to V3 \to V4 \to V5 \to V6 \to V7 \to V8 \to V9.$$

The result is a finite relay-chain extension, not a global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.228.5 Guarded residual status after r17

The r17 finite atlas scan improves the relay geometry but does not change the global claim guards. The following remain active:

$$V0 : \mathsf{blocked}, \qquad \beta_4^{r12} = \frac{1}{256} > 0, \qquad C6 : \mathsf{B}_{\mathrm{all/T2}}.$$

The $C5$ completion guard remains unchanged:

$$\forall n < \infty : \ \Phi_5^{(n)} > 0, \qquad \widehat{\Phi}_5(\infty) = 0.$$

Thus r17 should be read as a finite relay upgrade across the local valley-front sample. It is not a proof of RH, not a photon-zero identity, not a T2 proof, and not a global physical closure theorem.

Corollary A.228.6 (permitted r18 direction)**.** *The next admissible source-only pass is*

v4.7.0-r18: $V0$ boundary-front obstruction inventory and finite-chain guard audit.

It should determine whether $V0$ is a true boundary-front obstruction, a missing-source chart, or an inadmissible zero-packet baseline, while auditing that the r17 finite chain is not misread as a global closure.

A.229 v4.7.0-r18 Source-Only Pass: V0 Boundary-Front Obstruction Inventory and Finite-Chain Guard Audit

This source-only pass records the next conservative step after the r17 atlas-gap bridge and guarded micro-lift scan. No PDF rebuild is requested or used. The purpose is not to extend the local relay chain past its declared scope, but to inventory the remaining boundary chart $V0$ and to audit that

$$V1 \to V2 \to V3 \to V4 \to V5 \to V6 \to V7 \to V8 \to V9$$

is read only as a finite local valley-front chain.

A.229.1 V0 boundary-front inventory

Recall the r15/r17 boundary datum

$$V0 : \qquad S_0 = 4, \qquad G_0 = 1, \qquad H_0 = G_0 S_0 = 4 < 16.$$

Equivalently,

$$\Delta_0 = 16 - H_0 = 12, \qquad G_0^{\mathrm{req}} = \frac{16}{S_0} = 4.$$

Thus $V0$ is not close to the relay threshold in the same sense as the former $V6$ blocker. It is not a small shadow-basin miss; it is a boundary-front source problem.

The conservative inventory is:

case	N	κ	G	S	$H = GS$	status
baseline	4	0	1	4	4	blocked
gain-only	4	0	4	4	16	formal only
incoherent flux-only	16	0	1	16	16	formal only
overlap lift	5	3/4	1	20	20	source-guarded

© 2026 Marc Brendecke. Preprint — not peer reviewed.

The gain-only and flux-only rows show that $V0$ is numerically liftable if one allows an undeclared boundary source, an undeclared boundary clock, or an external injection rule. These are not currently part of the declared local relay grammar. Therefore r18 classifies $V0$ as

$$\boxed{V0 : \text{boundary-front source-deficient obstruction.}}$$

This is stronger than saying that $V0$ is simply "blocked", but weaker than claiming that it is an absolute topological obstruction. It means that the known finite chain has no certified upstream source chart at the boundary.

Lemma A.229.1 (V0 boundary-front decision)**.** *Within the current r18 source grammar, $V0$ is not absorbed by the $V1 \to \cdots \to V9$ finite relay chain. It remains a source-deficient boundary-front obstruction. Any lift of $V0$ must declare at least one of the following additional structures:*

boundary source chart, *boundary zero-clock phase window,* *external packet injection rule.*

Proof. The current datum gives $H_0 = 4 < 16$. A formal threshold equality can be forced by taking $G = 4$, or by taking sixteen incoherent packets, or by introducing a coherent overlap lift. None of these is certified by the existing relay chain, because the chain begins at $V1$ and supplies no admissible upstream source for $V0$. Hence the obstruction is not a numerical impossibility; it is a missing-source boundary-front obstruction in the declared grammar. □

A.229.2 Finite-chain guard audit

The r17 chain remains a finite local atlas result:

$$\boxed{\mathcal{R}^{\mathrm{fin}}_{1:9} = V1 \to V2 \to V3 \to V4 \to V5 \to V6 \to V7 \to V8 \to V9.}$$

The r18 guard audit records the active non-closures:

guard	r18 status	interpretation
$V0$	active	boundary-front source deficient
$C4$	$\beta_4 = 1/256 > 0$	route-stable mixed-kernel remains
$C5$	$\forall n < \infty : \Phi_5^{(n)} > 0$	asymptotic collar only
$C6$	$\mathsf{B}_{\mathrm{all/T2}}$	all-readout/T2 blocked
RH	not claimed	photon-clock handoff remains a probe
global	not claimed	no all-readout closure

Consequently, the finite local chain is useful because it converts many formerly abstract valley-front obstructions into relay-admissible handoff steps. It does not remove the boundary source problem at $V0$, and it does not discharge the route-stable $C4$ mixed-kernel residue.

Proposition A.229.2 (r18 finite-chain guard)**.** *The r17 relay chain $\mathcal{R}^{\mathrm{fin}}_{1:9}$ is admissible as a finite local relay chain, but it is not admissible as a global closure witness. The active guards are $V0$, $C4$, finite $C5$ closure, $C6/T2$, RH, and global all-readout closure.*

Proof. The chain starts at $V1$, not at a certified boundary source. It also leaves $\beta_4 = 1/256$ unchanged and does not alter the finite-positive $C5$ guard or the all-readout/T2 blocker. Therefore the finite chain may be used as a local relay atlas but not as a global closure theorem. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.229.3 Permitted r19 direction

The next source-only pass may either keep $V0$ frozen and move back to the $C4$ mixed-kernel seam, or test whether a declared boundary source chart can be added without creating new debt. The safest next pass is therefore

v4.7.0-r19: boundary-source chart admissibility test and C4 guard-preserving audit.

No RH, T2, or global all-readout closure should be inferred from r18.

A.230 v4.7.0-r19 Source-Only Pass: Boundary-Source Chart Admissibility Test and C4 Guard-Preserving Audit

This source-only pass tests the admissibility of a possible boundary-source chart for the remaining boundary-front obstruction $V0$. No PDF rebuild is requested or used. The goal is deliberately narrow: determine whether $V0$ can be lifted by a declared local boundary source without spending the route-stable $C4$ mixed-kernel guard and without opening the all-readout/T2 selector.

The r18 decision was

$$V0 : \text{boundary-front source-deficient obstruction}, \qquad S_0 = 4, \quad G_0 = 1, \quad H_0 = 4 < 16.$$

Thus the missing amount is

$$\Delta_0 = 12, \qquad G_0^{\mathrm{req}} = 4.$$

The present pass asks whether this deficit can be supplied by a legitimate boundary chart rather than by an undeclared global injection.

A.230.1 Boundary-source admissibility criterion

For a boundary-source candidate $\mathcal{B}_0$, write the local overlap score as

$$S(\mathcal{B}_0) = N + \kappa N(N-1), \qquad H(\mathcal{B}_0) = G\, S(\mathcal{B}_0).$$

Here N is the packet count in the boundary window, κ is the local coherence, and G is any declared photon-clock or boundary timing gain. The threshold remains

$$H(\mathcal{B}_0) \geq 16.$$

This numerical threshold is necessary but not sufficient. The chart is admissible only if the following guard conditions also hold:

(A1) $H(\mathcal{B}_0) \geq 16$,
(A2) $D_\partial(\mathcal{B}_0) = 0$ (no undeclared boundary debt),
(A3) $P_{\partial 0} S_{4,\mu}^{\mathrm{mixker}} = 0$ (no spending of the $C4$ mixed kernel),
(A4) $\mathcal{B}_0$ does not introduce an all-readout selector or a $T2$ lift.

Condition (A3) is the guard-preserving clause. The boundary chart may supply a local upstream source for $V0$, but it may not absorb or relabel the persistent mixed-kernel residue

$$\beta_4 = \frac{1}{256}.$$

Definition A.230.1 (r19 boundary-source admissibility)**.** *A boundary-source chart $\mathcal{B}_0$ is r19-admissible if and only if it satisfies (A1)–(A4). A candidate satisfying only (A1) is called numerically lifting but grammar-inadmissible.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.230.2 Finite boundary-source candidate scan

The finite candidate scan is:

candidate	N	κ	G	S	$H = GS$	r19 decision
baseline	4	0	1	4	4	blocked
clock-only	4	0	4	4	16	numerical only; undeclared clock debt
external injection	16	0	1	16	16	numerical only; injection debt
coherent boundary reservoir	5	3/4	1	20	20	conditionally admissible
clock-assisted overlap	4	3/4	16/13	13	16	conditionally admissible

The baseline fails the threshold. The clock-only and external-injection rows show that $V0$ can be forced to the threshold numerically, but only by adding an undeclared source of gain or packets. Those rows remain inadmissible in the current grammar.

The first genuinely useful row is the coherent boundary reservoir. For $N = 5$, the threshold condition is

$$5 + 20\kappa \geq 16, \qquad \kappa \geq \frac{11}{20}.$$

Thus the displayed value $\kappa = 3/4$ has positive margin

$$S - 16 = 4.$$

This proves only a conditional admissibility statement: if a local boundary reservoir $\mathcal{B}_0^{\mathrm{res}}$ with

$$N = 5, \qquad \kappa \geq \frac{11}{20}, \qquad G = 1$$

is declared and if it satisfies the no-new-debt and $C4$-orthogonality guards, then $V0$ is relay-admissible as a boundary-source chart.

Lemma A.230.2 (minimal coherent reservoir threshold)**.** *For a boundary-reservoir candidate with $G = 1$ and $N = 5$, the local relay threshold is reached exactly when*

$$\kappa \geq \frac{11}{20}.$$

In particular, $\kappa = 3/4$ gives $S = 20 > 16$.

Proof. Substituting $N = 5$ into $S = N + \kappa N(N-1)$ gives $S = 5 + 20\kappa$. The threshold $S \geq 16$ is therefore equivalent to $20\kappa \geq 11$, hence $\kappa \geq 11/20$. At $\kappa = 3/4$, one obtains $S = 5 + 15 = 20$. □

A.230.3 C4 guard-preserving audit

The boundary-source scan does not spend the $C4$ mixed-kernel residue. The r12–r18 persistent residue remains

$$S_{4,\mu}^{\mathrm{mixker}} : \mathsf{route\text{-}stable\ persistent}, \qquad \beta_4 = \frac{1}{256}.$$

The guard audit is:

candidate	threshold	$C4$ guard	decision
clock-only	met	not certified	reject unless clock source declared
external injection	met	not certified	reject as external debt
coherent boundary reservoir	met	$P_{\partial 0} S_{4,\mu}^{\mathrm{mixker}} = 0$	conditional pass
clock-assisted overlap	met	requires clock-source audit	conditional only

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Accordingly, r19 makes no unconditional reduction of β_4:

$$\boxed{\beta_4^{r19} = \frac{1}{256}.}$$

The only admissible direction is to treat $\mathcal{B}_0^{\mathrm{res}}$ as a new local boundary-source object whose support is orthogonal to the mixed-kernel seam. If that orthogonality is not declared, the candidate cannot be used.

Proposition A.230.3 (r19 guard-preserving outcome)**.** *The r19 scan finds no unconditional current-grammar lift of $V0$. It does, however, isolate a conditional boundary-source normal form:*

$$\mathcal{B}_0^{\mathrm{res}}: \qquad N = 5, \qquad \kappa \geq \frac{11}{20}, \qquad G = 1, \qquad P_{\partial 0} S_{4,\mu}^{\mathrm{mixker}} = 0.$$

Under these conditions $V0$ is boundary-source relay-admissible and the $C4$ guard remains unchanged at $\beta_4 = 1/256$. Without a declared $\mathcal{B}_0^{\mathrm{res}}$, $V0$ remains guarded.

Proof. The numerical threshold follows from Lemma A.230.2. The no-new-debt requirement is exactly the statement that the reservoir is a local boundary-source chart rather than an external injection. The displayed orthogonality condition prevents the new boundary source from absorbing the route-stable $C4$ mixed-kernel residue. Thus the result is conditional and guard-preserving, not an unconditional closure. □

A.230.4 Residual status after r19

After r19 the finite relay-chain statement is unchanged:

$$V1 \to V2 \to V3 \to V4 \to V5 \to V6 \to V7 \to V8 \to V9$$

remains the certified finite local valley-front chain. The boundary chart $V0$ has a conditional lift criterion, but it is not yet part of the certified chain.

The guarded status is therefore:

object	r19 status
$V0$	conditional boundary-reservoir criterion only
$C4$	$\beta_4 = 1/256$ route-stable mixed-kernel guard preserved
$C5$	asymptotic bounded collar; no finite closure
$C6/T2$	all-readout blocker unchanged
RH	no claim; photon-clock remains handoff probe
global closure	not claimed

A.230.5 Permitted r20 direction

The next source-only pass may test whether the conditional object $\mathcal{B}_0^{\mathrm{res}}$ can be promoted from an admissibility criterion to a declared local boundary-source chart. The safe formulation is

v4.7.0-r20: boundary-reservoir declaration test and no-new-debt proof obligation.

This pass should either certify the no-new-debt and $C4$-orthogonality clauses for $\mathcal{B}_0^{\mathrm{res}}$, or keep $V0$ frozen as a guarded boundary source deficit. No RH, T2, finite $C5$ closure, or global all-readout closure is licensed by r19.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.231 v4.7.0-r20 Source-Only Boundary-Reservoir Declaration Test

This source-only pass performs the next admissible test after the r19 boundary-source scan. The question is not whether the old grammar already contains a free lift of $V0$. r19 showed that it does not. The question is whether the conditional boundary-source normal form isolated in r19 may be introduced as a declared local boundary reservoir without creating new debt. No PDF is produced in this intermediate pass.

A.231.1 Declared local boundary-reservoir object

We define the boundary-reservoir extension of the local finite valley-front grammar by adjoining one explicitly declared boundary chart:

$$\mathcal{G}_{\mathrm{loc}}^{r20,\partial mres} := \mathcal{G}_{\mathrm{loc}}^{r19} \cup \{\mathcal{B}_0^{\mathrm{res}}\}.$$

The new object is not an external packet injection and not a global readout channel. It is a local boundary reservoir with declared data

$$\mathcal{B}_0^{\mathrm{res}} = \left(V0, N = 5, \kappa = \frac{3}{4}, G = 1, \delta_\partial = 0, P_{\partial 0} S_{4,\mu}^{\mathrm{mixker}} = 0, \chi_{\mathrm{all}} = 0\right).$$

The associated local overlap score is

$$S_0^{\partial mres} = N + \kappa N(N-1) = 5 + \frac{3}{4} \cdot 5 \cdot 4 = 20.$$

Since $G = 1$, the relay score is

$$H_0^{\partial mres} = 20 > 16.$$

Thus the reservoir has relay margin

$$\mu_0^{\partial mres} = 20 - 16 = 4.$$

Definition A.231.1 (r20 boundary-reservoir declaration)**.** *The object $\mathcal{B}_0^{\mathrm{res}}$ is called a declared local boundary-reservoir chart if it satisfies the following four clauses:*

$$\begin{aligned} &\text{(D1) } \delta_\partial = 0, \\ &\text{(D2) } P_{\partial 0} S_{4,\mu}^{\mathrm{mixker}} = 0, \\ &\text{(D3) } \chi_{\mathrm{all}} = 0, \\ &\text{(D4) } H_0^{\partial \mathrm{res}} \geq 16. \end{aligned}$$

Clause (D1) forbids external boundary debt. Clause (D2) forbids spending the route-stable $C4$ mixed-kernel residue. Clause (D3) forbids all-readout or $T2$ promotion. Clause (D4) is the local relay threshold.

A.231.2 No-new-debt proof obligation

The no-new-debt vector for the declaration is

$$\mathbf{d}_{\mathrm{new}}^{\partial mres} = \left(\delta_\partial, \delta_{C4}, \delta_{C6/T2}, \delta_{\mathrm{all}}\right).$$

For the declared reservoir we set

$$\delta_\partial = 0, \qquad \delta_{C4} = \left\| P_{\partial 0} S_{4,\mu}^{\mathrm{mixker}} \right\| = 0, \qquad \delta_{C6/T2} = 0, \qquad \delta_{\mathrm{all}} = 0.$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Therefore

$$\boxed{\mathbf{d}_{\mathrm{new}}^{\partial mres} = (0,0,0,0).}$$

This proves only that the explicitly declared local reservoir is debt-neutral with respect to the listed guards. It does not prove that such a reservoir was already present in the old r19 grammar.

Proposition A.231.2 (r20 no-new-debt declaration pass)**.** *In the extended local grammar* $\mathcal{G}_{\mathrm{loc}}^{r20,\partial\mathrm{res}}$*, the declared object* $\mathcal{B}_0^{\mathrm{res}}$ *is relay-admissible for* $V0$ *and introduces no new boundary,* $C4$*,* $C6/T2$*, or all-readout debt.*

Proof. The relay inequality follows from

$$H_0^{\partial\mathrm{res}} = 20 > 16.$$

The no-new-debt claim follows directly from the four declaration clauses in Definition A.231.1. Clause (D1) sets boundary debt to zero. Clause (D2) makes the new boundary support orthogonal to the route-stable mixed-kernel seam, so β_4 is not spent. Clause (D3) prevents promotion into all-readout or $T2$ territory. Clause (D4) supplies the local relay inequality. Hence the extension is conservative at the level stated here. □

A.231.3 Boundary-extended finite relay chain

With the declared reservoir, the finite local chain may be written as

$$\boxed{V0^{\partial\mathrm{res}} \to V1 \to V2 \to V3 \to V4 \to V5 \to V6 \to V7 \to V8 \to V9.}$$

The superscript on $V0$ is essential. It records that the first chart is not inherited as a free current-grammar chart. It is admitted only in the boundary-reservoir extension.

The active guard ledger after r20 is therefore

object	r20 status	guard
$V0$	declared local boundary-reservoir relay-admissible	local only
$C4$	$\beta_4 = 1/256$ route-stable mixed-kernel	unchanged
$C5$	asymptotic bounded collar	no finite closure
$C6/T2$	all-readout blocker	unchanged
RH	photon-clock handoff probe only	no proof claim
global closure	not claimed	guarded

Corollary A.231.3 (r20 boundary-extended local coverage)**.** *The r20 extension certifies a boundary-extended finite local valley-front chain from* $V0$ *through* $V9$*. This is not a global atlas theorem and not an all-readout closure theorem.*

Proof. The chart $V0$ is relay-admissible by Proposition A.231.2. The charts $V1$ through $V9$ are the finite relay chain certified by the r14–r17 source-only passes and guard-audited in r18. The r20 declaration changes only the local boundary entry chart. It does not remove the $C4$, $C5$, $C6/T2$, RH, or global all-readout guards. □

A.231.4 Permitted r21 direction

The next source-only pass may test whether the boundary-extended finite chain has any hidden seam coupling back into $C4$, or whether the boundary reservoir can be treated as a separate local source chart in the two-sided tensor grammar. The safe formulation is

$$\boxed{\text{v4.7.0-r21: boundary-extended chain seam audit and two-sided source typing.}}$$

No finite $C5$ closure, $C6/T2$ lift, RH claim, or global all-readout closure is licensed by r20.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.232 v4.7.0-r21: Boundary-Extended Chain Seam Audit and Two-Sided Source Typing

Status. This is a source-only internal pass. It audits the r20 boundary-reservoir extension for hidden seam coupling and assigns a conservative two-sided source type to the boundary chart. It does not introduce a new DOI, does not rebuild a PDF, and does not claim finite $C5$ closure, $C6/T2$ promotion, RH, or global all-readout closure.

A.232.1 Seam-audit object

The r20 boundary extension produced the local finite chain

$$V0^{\partial\mathrm{res}} \to V1 \to V2 \to V3 \to V4 \to V5 \to V6 \to V7 \to V8 \to V9.$$

The r21 question is whether the new boundary entry creates a hidden seam back into the route-stable $C4$ mixed kernel, the $C6/T2$ all-readout guard, or an undeclared global closure channel.

We record the seam-audit vector

$$\Sigma_{\partial mres} = \left(\Sigma_{\partial C4}, \Sigma_{\partial C6/T2}, \Sigma_{\partial mall}, \Sigma_{\partial 1}\right),$$

where

$$\Sigma_{\partial C4} = \left\| P_{\partial 0} S_{4,\mu}^{\mathrm{mixker}} \right\|, \qquad \Sigma_{\partial C6/T2} = \chi_{\mathrm{all}}, \qquad \Sigma_{\partial mall} = \delta_{\mathrm{all}},$$

and $\Sigma_{\partial 1}$ is the boundary-to-first-interior relay seam. The r20 declaration clauses give

$$\Sigma_{\partial C4} = 0, \qquad \Sigma_{\partial C6/T2} = 0, \qquad \Sigma_{\partial mall} = 0.$$

For the first relay seam, r20 supplies

$$H_0^{\partial\mathrm{res}} = 20, \qquad 16 \le H_0^{\partial\mathrm{res}}, \qquad \mu_0^{\partial\mathrm{res}} = 4,$$

so the boundary exit is locally relay-admissible. Hence

$$\boxed{\Sigma_{\partial\mathrm{res}} = (0,0,0,0).}$$

This vanishing seam vector means only that the declared local boundary source adds no new seam debt. It does not erase the existing route-stable mixed-kernel residue

$$\boxed{\beta_4 = \frac{1}{256}.}$$

Lemma A.232.1 (r21 boundary seam audit)**.** *In the declared local grammar $\mathcal{G}_{\mathrm{loc}}^{r20,\partial\mathrm{res}}$, the boundary entry chart $V0^{\partial\mathrm{res}}$ introduces no hidden seam coupling into $C4$, $C6/T2$, or all-readout closure.*

Proof. The $C4$ seam vanishes by the explicit orthogonality clause $P_{\partial 0} S_{4,\mu}^{\mathrm{mixker}} = 0$. The $C6/T2$ and all-readout seams vanish by $\chi_{\mathrm{all}} = 0$ and $\delta_{\mathrm{all}} = 0$. The first relay seam is admitted by the local inequality $H_0^{\partial\mathrm{res}} = 20 \ge 16$. These four facts give $\Sigma_{\partial\mathrm{res}} = (0,0,0,0)$. The statement is local and does not change β_4. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.232.2 Two-sided source typing

The boundary reservoir is now typed in the two-sided structural tensor grammar. We write schematically

$$ {}^{\partial msrc}{}_{\mathrm{loc}}\mathcal{B}_{0\ \mathrm{relay}}^{\mathrm{res}V0}. $$

The front indices ${}^{\partial msrc}{}_{\mathrm{loc}}$ mark a left-sided, readout-facing local boundary source interface. The back indices ${}^{V0}_{\mathrm{relay}}$ mark the right-sided valley-front relay structure. The notation is not a new contraction rule. It records that the object is a local source chart whose output may feed the finite relay chain, while its boundary source face is not an all-readout face.

The admissible readout is therefore the ordinary left-to-right local relay readout

$$ \text{boundary reservoir} \longmapsto V0^{\partial\mathrm{res}} \longmapsto V1. $$

Same-variance comparison against the $C4$ mixed kernel is allowed only through an explicit comparison/orthogonality test,

$$ G_{\partial C4}\,\mathcal{B}_0^{\partial}\,S_4^{\mathrm{mixker}} = 0, $$

which is the typed version of $P_{\partial 0}S_{4,\mu}^{\mathrm{mixker}} = 0$. Thus the boundary source is not permitted to spend, cancel, or re-label the persistent $C4$ mixed-kernel residue.

Definition A.232.2 (r21 two-sided boundary-source type)**.** *The declared object* ${}^{\partial msrc}{}_{\mathrm{loc}}\mathcal{B}_{0\ \mathrm{relay}}^{\mathrm{res}V0}$ *is a two-sided local boundary-source chart if it satisfies:*

$$ \begin{aligned} &m(T1)\ \textit{left/front type} = \partial\mathrm{src/loc},\\ &m(T2)\ \textit{right/back type} = V0/\mathrm{relay},\\ &m(T3)\ G_{\partial C4}\mathcal{B}_0^{\partial}S_4^{\mathrm{mixker}} = 0,\\ &m(T4)\ \chi_{\mathrm{all}} = 0,\\ &m(T5)\ H_0^{\partial\mathrm{res}} \geq 16. \end{aligned} $$

The type is local-source/relay. It is not global-source, not all-readout, and not a $T2$ bridge.

A.232.3 Completion-interface audit

The r12–r20 passes leave $C5$ as an asymptotic bounded-collar object,

$$ \Phi_5^{(n)} = \left(\frac{3}{4}\right)^n, \qquad \widehat{\Phi}_5(\infty) = 0, \qquad \Phi_5^{(n)} > 0 \quad (n < \infty). $$

The r20 boundary reservoir does not alter this completion interface. In r21 we therefore record the guard

$$ \boxed{V0^{\partial\mathrm{res}} \text{ enters the finite chain, but it does not change the } C5 \text{ completion topology.}} $$

Equivalently, the boundary source is a local chain-entry chart, not a new limit operator on $\mathbb{N}_\infty$.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.232.4 r21 decision ledger

The r21 decision is

object	r21 decision	guard
$V0^{\partial\mathrm{res}}$	two-sided local source/relay typed	local only
$\Sigma_{\partial\mathrm{res}}$	$(0,0,0,0)$	no new seam debt
$C4$	$\beta_4 = 1/256$	route-stable persistent
$C5$	asymptotic bounded collar	no finite closure
$C6/T2$	blocked	all-readout guard
RH	photon-clock handoff probe only	no proof claim
global closure	not claimed	guarded

Proposition A.232.3 (r21 typed local-chain preservation)**.** *The r20 boundary-extended finite chain remains valid after the r21 seam audit and two-sided source typing, and no additional $C4$, $C5$, $C6/T2$, RH, or global all-readout claim is licensed.*

Proof. Lemma A.232.1 gives the zero seam-debt vector for the boundary entry. Definition A.232.2 assigns the boundary reservoir a local two-sided source/relay type and forbids both $C4$-spending and all-readout promotion. The internal chain $V1 \to \cdots \to V9$ is unchanged from the r14–r17 finite relay construction. Thus the finite boundary-extended chain is preserved locally, with all guards listed in the decision ledger still active. □

A.232.5 Permitted r22 direction

The next source-only pass may test whether the typed boundary source can be compiled into a small boundary-source calculus or whether it must remain a single declared chart. The safe formulation is

v4.7.0-r22: boundary-source calculus extraction and typed relay closure guard.

This r22 direction is still local and source-only. It may not promote the finite chain to global all-readout closure and may not identify the photon-clock probe with RH.

A.233 v4.7.0-r22 source-only: boundary-source calculus extraction and typed relay closure guard

This source-only pass extracts a small local boundary-source calculus from the single r20–r21 boundary-reservoir chart and immediately guards the extraction against accidental promotion to global closure. The pass does not rebuild a PDF, does not assign a DOI, and does not modify the public release status.

The r21 typed reservoir is kept as

$$ {}^{\partial\mathrm{src}}{}_{\mathrm{loc}}\mathcal{B}_0^{\mathrm{res}V0}{}_{\mathrm{relay}}. $$

The r22 question is whether this typed object may be used as a reusable local calculus rule rather than as an isolated chart. The answer is yes, but only as a local boundary-source rule with a typed relay guard.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.233.1 Local boundary-source calculus rule

Definition A.233.1 (r22 local boundary-source calculus rule)**.** *Let $\mathcal{B}$ be a boundary reservoir candidate. It is admissible as a local boundary-source calculus instance, written*

$$\mathcal{B} \in \mathsf{BSC}_{\partial}^{\mathrm{loc}},$$

if the following five clauses hold:

$$\begin{aligned}
&\text{BSC1 } \mathcal{B} \text{ has two-sided type } {}^{\partial\mathrm{src}}{}_{\mathrm{loc}}\mathcal{B}^{V}_{\mathrm{relay}};\\
&\text{BSC2 } H_{\mathcal{B}} = G_{\mathcal{B}} S_{\mathcal{B}} \geq 16;\\
&\text{BSC3 } P_{\partial}\, S^{\mathrm{mixker}}_{4,\mu} = 0;\\
&\text{BSC4 } \chi_{\mathrm{all}}(\mathcal{B}) = 0;\\
&\text{BSC5 } \mathbf{d}_{\mathrm{new}}(\mathcal{B}) = (0,0,0,0).
\end{aligned}$$

The rule is therefore a local source-and-relay rule. It is not an all-readout rule and it is not a global boundary theorem.

For the concrete r20/r21 reservoir one has

$$N = 5, \qquad \kappa = \frac{3}{4}, \qquad G = 1,$$

so

$$S = 5 + \frac{3}{4} \cdot 5 \cdot 4 = 20, \qquad H = 20 > 16.$$

Thus

$$\boxed{\mathcal{B}_0^{\mathrm{res}} \in \mathsf{BSC}_{\partial}^{\mathrm{loc}}.}$$

Lemma A.233.2 (r22 boundary-source calculus extraction)**.** *The r20–r21 boundary-reservoir chart can be extracted as a reusable local boundary-source calculus instance in $\mathsf{BSC}_{\partial}^{\mathrm{loc}}$.*

Proof. BSC1 is the r21 two-sided source typing. BSC2 follows from the r20 threshold calculation $H = 20 > 16$. BSC3 and BSC5 are the r20–r21 no-new-debt and seam-audit conditions. BSC4 is the explicit local-only readout guard. Hence the reservoir satisfies all five clauses of Definition A.233.1. □

A.233.2 Typed relay closure guard

The word "closure" is used here only in the typed finite-relay sense. It means that a declared local boundary source may be attached to the beginning of the already certified finite relay chain. It does not mean global closure.

Definition A.233.3 (r22 typed finite-relay closure)**.** *The typed finite-relay closure generated by $\mathsf{BSC}_{\partial}^{\mathrm{loc}}$ is the local operation*

$$\mathrm{Cl}^{\mathrm{loc,typed}}_{\mathrm{relay}}\big(\mathcal{B}_0^{\mathrm{res}}; V1, \ldots, V9\big) = \big[V0^{\partial\mathrm{res}} \to V1 \to \cdots \to V9\big]$$

provided the guard vector

$$\Gamma^{r22}_{\mathrm{guard}} = (\gamma_{C4}, \gamma^{\mathrm{fin}}_{C5}, \gamma_{C6/T2}, \gamma_{\mathrm{all}}, \gamma_{\mathrm{RH}})$$

remains

$$\Gamma^{r22}_{\mathrm{guard}} = (1,1,1,1,1).$$

Here 1 means that the corresponding guard is still active, not discharged.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

For r22 the guard ledger is

$$\boxed{\Gamma^{r22}_{\mathrm{guard}} = (1,1,1,1,1).}$$

In particular,

$$\beta_4^{r22} = \frac{1}{256}, \qquad \forall n < \infty : \ \Phi_5^{(n)} > 0, \qquad C6 : \mathsf{B}_{\mathrm{all/T2}}.$$

Proposition A.233.4 (r22 typed relay closure guard)**.** *The boundary-source calculus extraction licenses the typed local finite relay chain*

$$V0^{\partial \mathrm{res}} \to V1 \to \cdots \to V9,$$

but it does not license $C4$-closure, finite $C5$-closure, $C6/T2$, RH, or global all-readout closure.

Proof. By Lemma A.233.2, the boundary reservoir is a valid local calculus instance. Definition A.233.3 then permits the local finite relay attachment. The same definition also requires the guard vector to remain active. Since $\beta_4 = 1/256$ is preserved, finite $C5$-closure is excluded by the finite positivity of $\Phi_5^{(n)}$, $C6/T2$ remains an all-readout blocker, and the photon-clock construction is a handoff probe only, none of the guarded global claims is discharged. □

A.233.3 r22 calculus ledger

The extracted calculus can be summarized as

clause	r22 value	decision
two-sided type	${}^{\partial\mathrm{src}}{}_{\mathrm{loc}}\mathcal{B}_0^{\mathrm{res}V0}{}_{\mathrm{relay}}$	admissible
H	$20 > 16$	relay pass
$P_{\partial 0} S^{\mathrm{mixker}}_{4,\mu}$	0	C4 guard preserved
χ_{all}	0	not all-readout
$\mathbf{d}_{\mathrm{new}}$	$(0,0,0,0)$	no new debt
$\Gamma^{r22}_{\mathrm{guard}}$	$(1,1,1,1,1)$	guards active

A.233.4 Permitted r23 direction

The next source-only pass may test whether the local boundary-source calculus is stable under finite perturbations of N, κ, G and whether the typed relay guard remains invariant under such perturbations. The safe formulation is

v4.7.0-r23: boundary-source stability window and relay-guard robustness scan.

This may not be read as global all-readout closure, a finite $C5$ closure, an RH statement, or a $T2$ bridge.

A.234 v4.7.0-r23 Boundary-Source Stability Window and Relay-Guard Robustness Scan

This source-only pass tests whether the local boundary-source calculus extracted in r22 is stable under small finite perturbations of the reservoir parameters (N, κ, G). The pass is deliberately local. It does not create a new DOI, does not rebuild the PDF, and does not change any all-readout or global-closure claim.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.234.1 Reservoir response function

For a local boundary-source reservoir we keep the r13–r22 overlap response function

$$S(N,\kappa) = N + \kappa N(N-1), \qquad H(N,\kappa,G) = G\,S(N,\kappa).$$

The relay threshold is

$$H(N,\kappa,G) \geq 16.$$

The r22 anchor is

$$(N,\kappa,G) = \left(5, \frac{3}{4}, 1\right), \qquad S = 20, \qquad H = 20, \qquad \mu = H - 16 = 4.$$

Definition A.234.1 (r23 boundary-source stability window)**.** *The weak r23 stability window is the set of local reservoir triples*

$$\mathcal{W}_{\partial}^{r23} = \{(N,\kappa,G) : G(N + \kappa N(N-1)) \geq 16\}$$

subject to the r22 source-typing and guard clauses BSC1–BSC5. The strict stability window is

$$\mathcal{W}_{\partial,+}^{r23} = \{(N,\kappa,G) : G(N + \kappa N(N-1)) > 16\}$$

with the same typing and guard clauses.

For the fixed local source size $N = 5$, the threshold surfaces are explicit:

$$G = 1 \quad \Longrightarrow \quad \kappa \geq \frac{11}{20},$$

while

$$\kappa = \frac{3}{4} \quad \Longrightarrow \quad G \geq \frac{4}{5}.$$

A one-unit strict margin is already guaranteed by

$$G = 1, \kappa \geq \frac{3}{5},$$

or by

$$\kappa = \frac{3}{4}, G \geq \frac{17}{20}.$$

A simultaneous finite robustness rectangle around the r22 anchor is

$$\boxed{N = 5, \qquad \kappa \geq \frac{7}{10}, \qquad G \geq \frac{9}{10},}$$

because then

$$S \geq 5 + \frac{7}{10} \cdot 5 \cdot 4 = 19, \qquad H \geq \frac{9}{10} \cdot 19 = \frac{171}{10} > 16.$$

Thus the r22 reservoir is not a knife-edge construction: it has a finite local stability collar.

Lemma A.234.2 (r23 local stability collar)**.** *The r22 boundary-reservoir point* $(5, 3/4, 1)$ *lies in the strict interior of the local boundary-source stability window. In particular, the rectangle*

$$N = 5, \qquad \kappa \in \left[\frac{7}{10}, 1\right], \qquad G \in \left[\frac{9}{10}, 1\right]$$

remains strictly relay-admissible.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. For every triple in the displayed rectangle,

$$S = 5 + 20\kappa \geq 5 + 20 \cdot \frac{7}{10} = 19,$$

and therefore

$$H = GS \geq \frac{9}{10} \cdot 19 = \frac{171}{10} > 16.$$

The r22 source-typing and no-new-debt clauses are not altered by this finite parameter perturbation, so the relay inequality remains valid in the same local calculus. □

A.234.2 Finite perturbation scan

The finite scan checked the grid

$$N \in \{4, 5, 6\}, \qquad \kappa \in \left\{\frac{1}{2}, \frac{11}{20}, \frac{3}{5}, \frac{2}{3}, \frac{7}{10}, \frac{3}{4}, 1\right\}, \qquad G \in \left\{\frac{4}{5}, \frac{17}{20}, \frac{9}{10}, 1\right\}.$$

The scan is not used as a proof by enumeration. It is a stability diagnostic around the symbolic threshold surfaces above. Its useful outcomes are:

case	N	κ	G	decision
r22 anchor	5	3/4	1	$H = 20 > 16$
lower-κ strict buffer	5	3/5	1	$H = 17 > 16$
critical κ surface	5	11/20	1	$H = 16$
gain-damped strict point	5	3/4	17/20	$H = 17 > 16$
gain-damped critical point	5	3/4	4/5	$H = 16$
simultaneous robust corner	5	7/10	9/10	$H = 171/10 > 16$
nominal $N = 4$	4	3/4	1	$H = 13 < 16$
larger low-κ reservoir	6	1/2	1	$H = 21 > 16$

The failure of the nominal $N = 4$ case is useful: the local boundary-source rule is not vacuous. It still requires enough local reservoir capacity.

A.234.3 Relay-guard robustness

The r23 stability collar is guarded by exactly the same guard vector as r22:

$$\boxed{\Gamma^{r23}_{\text{guard}} = (1, 1, 1, 1, 1).}$$

Concretely,

$$\beta_4^{r23} = \frac{1}{256}, \qquad \forall n < \infty : \ \Phi_5^{(n)} > 0, \qquad C6 : \mathsf{B}_{\text{all/T2}}.$$

The photon-clock construction remains a handoff probe, not an RH statement. The boundary-source stability collar remains local and typed, not all-readout.

Proposition A.234.3 (r23 relay-guard robustness)**.** *Every triple in the strict local r23 stability collar preserves the r22 guard vector. Hence the local boundary-source robustness scan does not discharge $C4$, finite $C5$-closure, $C6/T2$, RH, or global all-readout closure.*

Proof. The perturbation scan changes only the local relay parameters (N, κ, G) inside the already declared two-sided boundary-source type. The guard clauses BSC3–BSC5 remain unchanged: the boundary source stays orthogonal to the route-stable mixed kernel, the all-readout selector remains off, and the new-debt vector remains zero. Therefore the relay inequality can be stable without spending $\beta_4 = 1/256$, without converting the asymptotic $C5$ collar into a finite closure, and without opening the all-readout $C6/T2$ gate. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.234.4 r23 decision ledger

The source-only r23 decision is

object	r23 result	guard
$\mathcal{B}_0^{\mathrm{res}}$	strictly stable near $(5, 3/4, 1)$	local only
$\mathcal{W}_{\partial,+}^{r23}$	nonempty finite stability collar	$\chi_{\mathrm{all}} = 0$
$C4$	$\beta_4 = 1/256$	unspent
$C5$	$\Phi_5^{(n)} > 0$ for finite n	no finite closure
$C6/T2$	$\mathsf{B}_{\mathrm{all/T2}}$	blocked
RH	photon-clock handoff probe only	not a theorem
global all-readout	unclaimed	guard active

A.234.5 Permitted r24 direction

The next source-only step may package the stability collar into a small "theoremlet" and draw the residual frontier map showing exactly where the local boundary-source calculus stops. The safe title is

v4.7.0-r24: boundary-source robustness theoremlet and residual frontier map.

It must remain source-only unless a PDF is explicitly requested.

A.235 v4.7.0-r24 Boundary-Source Robustness Theoremlet and Residual Frontier Map

Source-only status. This r24 layer is a source-only internal working step. It packages the r23 stability scan into a local theoremlet and records the residual frontier of the boundary-source calculus. It assigns no new DOI, builds no PDF, and claims no global closure.

A.235.1 Boundary-source robustness theoremlet

The r23 anchor was

$$(N, \kappa, G) = \left(5, \frac{3}{4}, 1\right), \qquad S = 20, \qquad H = 20, \qquad \mu = 4.$$

The useful r24 point is that this is not merely a witnessed chart. It sits in a small local robustness box.

Theorem A.235.1 (r24 local boundary-source robustness theoremlet)**.** *Let*

$$H(N, \kappa, G) = G\big(N + \kappa N(N-1)\big)$$

be the local boundary-source relay height. In the declared local boundary-source grammar, every chart with

$$N = 5, \qquad \kappa \geq \frac{7}{10}, \qquad G \geq \frac{9}{10}$$

satisfies

$$H \geq \frac{171}{10} > 16.$$

Hence the r22/r23 boundary-source calculus has a strict local robustness collar around the anchor $(5, 3/4, 1)$.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. For $N = 5$,

$$S = 5 + 20\kappa.$$

If $\kappa \geq 7/10$, then

$$S \geq 5 + 20 \cdot \frac{7}{10} = 19.$$

If also $G \geq 9/10$, then

$$H = GS \geq \frac{9}{10} \cdot 19 = \frac{171}{10} > 16.$$

The proof uses only the already declared local boundary-source type. It does not switch on the all-readout selector and does not couple the boundary source to the route-stable mixed kernel. □

Corollary A.235.2 (r24 strict local relay pass)**.** *Every chart in the robustness box of Theorem A.235.1 is relay-admissible with positive margin. In particular, the boundary-source lift is not a knife-edge equality at $H = 16$.*

A.235.2 Threshold surfaces and finite frontier

The two one-parameter threshold surfaces around the r23 anchor are

$$N = 5, \quad G = 1 \quad \Longrightarrow \quad \kappa \geq \frac{11}{20},$$

and

$$N = 5, \quad \kappa = \frac{3}{4} \quad \Longrightarrow \quad G \geq \frac{4}{5}.$$

The strict one-unit buffer surfaces are

$$N = 5, \quad G = 1 \quad \Longrightarrow \quad \kappa \geq \frac{3}{5},$$

and

$$N = 5, \quad \kappa = \frac{3}{4} \quad \Longrightarrow \quad G \geq \frac{17}{20}.$$

For a wider residual-frontier map, r24 records the finite grid

$$N \in \{3, 4, 5, 6, 7\},$$

$$\kappa \in \left\{\frac{1}{2}, \frac{11}{20}, \frac{3}{5}, \frac{2}{3}, \frac{7}{10}, \frac{3}{4}, 1\right\},$$

$$G \in \left\{\frac{4}{5}, \frac{17}{20}, \frac{9}{10}, 1\right\}.$$

The grid contains 140 cases and partitions as

frontier class	count
strict-interior ($H \geq 17$)	66
thin-pass ($16 \leq H < 17$)	6
frontier-fail ($15 \leq H < 16$)	4
deep-fail ($H < 15$)	64

Thus the boundary-source calculus has a genuine pass region, a thin critical frontier, and a nontrivial fail region. The local rule is therefore not a vacuous route-to-closure device.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.235.3 Residual frontier map

The residual frontier is described by

$$G\big(N+\kappa N(N-1)\big)=16.$$

For fixed N and G, the threshold is

$$\kappa_{\min}(N,G)=\frac{16/G-N}{N(N-1)}.$$

For fixed N and κ, the threshold is

$$G_{\min}(N,\kappa)=\frac{16}{N+\kappa N(N-1)}.$$

At the representative slice $G=1$, the threshold moves into the unit coherence interval first at $N=4$:

$$N=4\Rightarrow\kappa_{\min}=1,\qquad N=5\Rightarrow\kappa_{\min}=\frac{11}{20},\qquad N=6\Rightarrow\kappa_{\min}=\frac{1}{3}.$$

This explains why the r23/r24 local reservoir really needs at least the five-packet anchor to obtain a comfortable margin at moderate coherence.

Proposition A.235.3 (r24 residual-frontier decision)**.** *The declared local boundary-source calculus extends the finite valley-front chain only inside the local pass region of*

$$G\big(N+\kappa N(N-1)\big)\geq 16.$$

Outside this region, the frontier remains residual and no relay-admissibility is inferred.

Proof. The calculus rule BSC2 is exactly the local relay inequality $H\geq 16$. If the inequality fails, the boundary source has no declared local relay height. The source typing and no-new-debt clauses do not create missing relay height; they only prevent successful local relay from spending forbidden guards. Therefore the pass/fail frontier is exactly the stated inequality. □

A.235.4 Guard-preserving consequence

The theoremlet is strictly local. Its guard vector remains

$$\boxed{\Gamma^{r24}_{\text{guard}}=(1,1,1,1,1).}$$

Equivalently,

$$\beta_4^{r24}=\frac{1}{256},\qquad \forall n<\infty:\ \Phi_5^{(n)}>0,\qquad C6:\mathsf{B}_{\text{all/T2}}.$$

The Riemann-zero/photon-clock language remains a timing probe for handoff windows, not an RH theorem and not a photon-zero identity. The boundary-source robustness theoremlet is a local relay theoremlet only.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.235.5 r24 decision ledger and permitted r25 direction

The r24 decision is

object	r24 result	guard
$\mathcal{B}_0^{\mathrm{res}}$	local robustness theoremlet	local only
$\mathcal{W}_{\partial,+}^{r24}$	finite pass/interior region mapped	$\chi_{\mathrm{all}} = 0$
$\partial\mathcal{W}^{r24}$	residual frontier mapped	no inferred pass outside $H \geq 16$
$C4$	$\beta_4 = 1/256$	unspent
$C5$	$\Phi_5^{(n)} > 0$ for finite n	no finite closure
$C6/T2$	$\mathsf{B}_{\mathrm{all/T2}}$	blocked
RH	photon-clock handoff probe only	not a theorem
global all-readout	unclaimed	guard active

A safe next source-only direction is

v4.7.0-r25: boundary-source frontier minimizer and guard-stable atlas compression.

This would search for the smallest local boundary-source resources that still preserve the guard vector, without turning the local finite chain into a global closure claim.

A.236 v4.7.0-r25 Source-Only Frontier Minimizer and Guard-Stable Atlas Compression

This source-only layer continues the r24 residual-frontier map. No PDF is built in this intermediate step. The goal is to identify the smallest local boundary-source resources visible on the finite frontier and to compress the finite relay atlas without spending any global guard.

A.236.1 Resource order and frontier minimizers

For a local boundary-source triple

$$(N, \kappa, G)$$

we keep the r24 height

$$H(N, \kappa, G) = G\big(N + \kappa N(N-1)\big).$$

A triple is relay-admissible only if

$$H(N, \kappa, G) \geq 16.$$

The source-resource order used in this scan is the componentwise order

$$(N, \kappa, G) \preceq (N', \kappa', G') \quad \Longleftrightarrow \quad N \leq N', \quad \kappa \leq \kappa', \quad G \leq G'.$$

A frontier point is Pareto-minimal if no strictly smaller admissible triple exists in this order.

In the r24 finite grid the absolute pass minimizer is

$$\boxed{(N, \kappa, G) = (4, 1, 1).}$$

It gives

$$H = 1 \cdot (4 + 1 \cdot 4 \cdot 3) = 16.$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

This is a thin pass with zero margin. It is therefore not a robust boundary reservoir rule; it is only the exact local frontier point.

For the five-packet boundary source the pass thresholds are

$$G = 1 \Rightarrow \kappa \geq \frac{11}{20}, \qquad \kappa = \frac{3}{4} \Rightarrow G \geq \frac{4}{5}.$$

The margin-one stable thresholds are

$$G = 1 \Rightarrow \kappa \geq \frac{3}{5}, \qquad \kappa = \frac{3}{4} \Rightarrow G \geq \frac{17}{20}.$$

The simultaneous robust-box corner remains

$$\boxed{(N, \kappa, G) = \left(5, \frac{7}{10}, \frac{9}{10}\right),}$$

with

$$H = \frac{9}{10}\left(5 + \frac{7}{10} \cdot 5 \cdot 4\right) = \frac{171}{10} > 16.$$

The original r22–r24 anchor

$$\left(5, \frac{3}{4}, 1\right)$$

has

$$H = 20, \qquad \mu = 4.$$

Thus r25 distinguishes three roles:

$$\boxed{(4, 1, 1) = \text{absolute thin frontier},}$$

$$\boxed{\left(5, \frac{7}{10}, \frac{9}{10}\right) = \text{robust local box corner},}$$

$$\boxed{\left(5, \frac{3}{4}, 1\right) = \text{comfortable boundary-source anchor}.}$$

Proposition A.236.1 (r25 frontier minimizer decision)**.** *In the finite r24/r25 frontier grid, the triple* $(4, 1, 1)$ *is the absolute local pass minimizer, but it has zero relay margin. The five-packet boundary-reservoir rule is the first locally robust reservoir family used by this branch, and the box*

$$N = 5, \qquad \kappa \geq \frac{7}{10}, \qquad G \geq \frac{9}{10}$$

is a strict local robustness region.

Proof. The equation

$$G\big(N + \kappa N(N-1)\big) = 16$$

defines the frontier. Direct substitution gives $(4, 1, 1) \mapsto H = 16$, hence pass with zero margin. The same substitution at $(5, 7/10, 9/10)$ gives $H = 171/10 > 16$. Since increasing any of N, κ, G weakly increases H, the displayed box is contained in the strict pass region. The r22–r24 anchor $(5, 3/4, 1)$ lies strictly inside the same region with margin 4. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.236.2 Pareto frontier summary

The finite pass Pareto frontier consists of the thin and low-resource points

N	κ	G	H	status
4	1	1	16	thin pass
5	11/20	1	16	thin pass
5	2/3	9/10	33/2	thin pass
5	7/10	17/20	323/20	thin pass
5	3/4	4/5	16	thin pass
6	1/2	4/5	84/5	thin pass

These points describe where the boundary-source calculus first becomes locally admissible. They are not automatically robust enough to be used as source anchors.

The margin-one Pareto layer begins with

N	κ	G	H	status
5	3/5	1	17	strict interior
5	7/10	9/10	171/10	strict interior
5	3/4	17/20	17	strict interior
5	1	4/5	20	strict interior
6	1/2	17/20	357/20	strict interior
6	11/20	4/5	18	strict interior
7	1/2	4/5	112/5	strict interior

This layer is the better target for guard-stable atlas compression.

A.236.3 Guard-stable atlas compression

The expanded local chain

$$V0^{\partial\mathrm{res}} oV1 \to V2 \to V3 \to V4 \to V5 \to V6 \to V7 \to V8 \to V9$$

is compressed into four local atlas cells:

$$\mathcal{A}_0 = V0^{\partial\mathrm{res}}, \qquad \mathcal{A}_1 = (V1 \to V2 \to V3 \to V4 \to V5),$$
$$\mathcal{A}_2 = (V5 \to V6 \to V7), \qquad \mathcal{A}_3 = (V7 \to V8 \to V9).$$

Here $\mathcal{A}_0$ is the local boundary-reservoir source cell, $\mathcal{A}_1$ is the core valley relay cell, $\mathcal{A}_2$ is the shadow-basin/atlas-gap bridge cell, and $\mathcal{A}_3$ is the zero-clock handoff tail.

The compressed local atlas is therefore

$$\boxed{\mathcal{A}_\partial^{r25} = \mathcal{A}_3 \circ \mathcal{A}_2 \circ \mathcal{A}_1 \circ \mathcal{A}_0.}$$

This notation is only a local atlas compression. It does not turn the finite chain into a global all-readout closure witness.

Lemma A.236.2 (r25 guard-stable compression)**.** *The compressed atlas $\mathcal{A}_\partial^{r25}$ preserves the r24 guard vector*

$$\Gamma_{\mathrm{guard}}^{r25} = (1, 1, 1, 1, 1).$$

Proof. Each compressed cell is inherited from a previously guarded local construction: $\mathcal{A}_0$ satisfies the boundary-source calculus with $\chi_{\mathrm{all}} = 0$, zero new debt, and $P_\partial S_{4,\mu}^{\mathrm{mixker}} = 0$; $\mathcal{A}_1$ is the finite relay core; $\mathcal{A}_2$ uses the r16/r17 source-supported bridge and spends no $C4$ debt; $\mathcal{A}_3$ uses zero-clock windows only as timing probes. None of these cells declares an all-readout interface, an RH theorem, a finite $C5$ closure, or a $C6/T2$ bridge. Therefore the guard vector remains unchanged. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.236.4 r25 decision ledger

The r25 decision is

object	r25 result	guard
$(4,1,1)$	absolute thin frontier minimizer	zero margin
$(5,7/10,9/10)$	robust local box corner	local only
$(5,3/4,1)$	comfortable boundary-source anchor	local only
$\mathcal{A}_\partial^{r25}$	compressed finite local atlas	$\chi_{\mathrm{all}}=0$
$C4$	$\beta_4=1/256$	unspent
$C5$	$\Phi_5^{(n)}>0$ for finite n	no finite closure
$C6/T2$	$\mathsf{B}_{\mathrm{all/T2}}$	blocked
RH	photon-clock handoff probe only	not a theorem
global all-readout	unclaimed	guard active

A safe next source-only direction is

v4.7.0-r26: compressed atlas transition audit and frontier-to-field-equation handoff.

This would test whether the compressed local atlas can feed the MM/HC field-equation program as a typed local input without promoting any guarded object to global closure.

A.237 v4.7.0-r26 compressed atlas transition audit and frontier-to-field-equation handoff

A.237.1 Purpose and scope guard

This r26 source-only layer audits the compressed local atlas

$$\mathcal{A}_\partial^{r25}=\mathcal{A}_3\circ\mathcal{A}_2\circ\mathcal{A}_1\circ\mathcal{A}_0$$

as a possible typed input for the MM/HC field-equation program. It does not claim a new field equation, a global all-readout theorem, a finite $C5$ closure, a $C6/T2$ bridge, or an RH theorem. The purpose is narrower: check whether the local atlas cells can be handed forward as a guarded source object for the next field-equation search layer.

The r26 input objects are

$$\mathcal{A}_0=V0^{\partial\mathrm{res}},\qquad \mathcal{A}_1=(V1\to V2\to V3\to V4\to V5),$$

$$\mathcal{A}_2=(V5\to V6\to V7),\qquad \mathcal{A}_3=(V7\to V8\to V9),$$

together with the active guard vector

$$\Gamma_{\mathrm{guard}}^{r25}=(1,1,1,1,1).$$

The target is a local handoff object, not a global closure object.

A.237.2 Two-sided transition typing

Each compressed atlas cell is typed as a two-sided local interface:

$${}^{\partial\mathrm{src}}{}_{\mathrm{loc}}\mathcal{A}_{0\,\mathrm{relay}}^{V0},\qquad {}^{\mathrm{core}}{}_{\mathrm{loc}}\mathcal{A}_{1\,\mathrm{relay}}^{V5},$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

$$^{\text{shadow}}{}_{\text{loc}}\mathcal{A}^{V7}_{2\text{bridge}}, \qquad ^{\gamma\text{clock}}{}_{\text{loc}}\mathcal{A}^{V9}_{3\text{tail}}.$$

The front indices record the readout/source side of the cell. The back indices record the structure/relay side. Same-variance interactions remain forbidden unless a declared comparison tensor or selector mediates them.

The transition interfaces are

$$\tau_{01} : \mathcal{A}_0 \rightsquigarrow \mathcal{A}_1, \qquad \tau_{12} : \mathcal{A}_1 \rightsquigarrow \mathcal{A}_2, \qquad \tau_{23} : \mathcal{A}_2 \rightsquigarrow \mathcal{A}_3.$$

Their audit ledger is

transition	source type	target type	r26 status
τ_{01}	∂src/$V0$	core/$V5$	typed local relay
τ_{12}	core/$V5$	shadow/$V7$	source-supported bridge
τ_{23}	shadow/$V7$	γclock/$V9$	timing-gated tail

No transition in this table declares an all-readout channel.

A.237.3 Transition seam audit

For each transition define a seam-debt vector

$$\sigma(\tau) = (\sigma_{C4}, \sigma_{C5}, \sigma_{C6/T2}, \sigma_{\text{all}}).$$

The r26 transition audit returns

$$\boxed{\sigma(\tau_{01}) = (0,0,0,0), \qquad \sigma(\tau_{12}) = (0,0,0,0), \qquad \sigma(\tau_{23}) = (0,0,0,0).}$$

Therefore the compressed atlas transition vector is

$$\boxed{\Sigma^{r26}_{\text{trans}} = \sum_{j=0}^{2} \sigma(\tau_{j,j+1}) = (0,0,0,0).}$$

This is a local transition statement. It says that the compression introduces no new seam debt across the declared local atlas cells. It does not spend the pre-existing route-stable mixed-kernel guard

$$\beta_4 = \frac{1}{256}.$$

Lemma A.237.1 (r26 transition seam preservation)**.** *The compressed atlas transitions $\tau_{01}, \tau_{12}, \tau_{23}$ preserve the r25 guard vector.*

Proof. The transition τ_{01} uses the typed boundary-reservoir source calculus with zero new boundary debt, $P_\partial S^{\text{mixker}}_{4,\mu} = 0$, and $\chi_{\text{all}} = 0$. The transition τ_{12} is the already guarded source-supported bridge from the core relay into the shadow-basin/atlas-gap cell. The transition τ_{23} uses the photon-clock factor only as a local timing gate for the zero-clock tail. None of the three transitions declares an all-readout channel, consumes β_4, closes finite $C5$, or opens $C6/T2$. Hence $\Sigma^{r26}_{\text{trans}} = (0,0,0,0)$, and the guard vector is preserved. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.237.4 Frontier-to-field-equation handoff object

The local atlas is now packaged as a handoff object

$$\boxed{\mathfrak{X}_{\partial}^{r26} = \left(\mathcal{A}_{\partial}^{r25}, \Sigma_{\mathrm{trans}}^{r26}, \mathcal{F}_{\partial}, \Gamma_{\mathrm{guard}}^{r25}\right),}$$

where $\mathcal{F}_{\partial}$ denotes the boundary-source frontier surface

$$\boxed{G\big(N + \kappa N(N-1)\big) = 16.}$$

This object is allowed to feed the MM/HC field-equation program only after projection through the local guard projector

$$\Pi_{\mathrm{loc}}^{\mathrm{guard}} : \mathfrak{X}_{\partial}^{r26} \longmapsto \mathfrak{X}_{\partial,\mathrm{loc}}^{r26}.$$

The projected object is local, typed, and guarded:

$$\boxed{\mathfrak{X}_{\partial,\mathrm{loc}}^{r26} := \Pi_{\mathrm{loc}}^{\mathrm{guard}} \mathfrak{X}_{\partial}^{r26}.}$$

A schematic MM/HC handoff may then be written as

$${}^{\mathrm{loc}\,\partial\mathrm{atlas}}_{\mathrm{read}}\mathfrak{E}^{\mathrm{seed}}_{\mathrm{MM/HC}}{}^{\mathrm{field}}_{\mathrm{constraint}} = \mathsf{H}_{\mathrm{bd}} \circ \mathsf{M}_{\mathrm{bd}} \left(\mathfrak{X}_{\partial,\mathrm{loc}}^{r26}\right).$$

The symbol $\mathfrak{E}^{\mathrm{seed}}_{\mathrm{MM/HC}}$ is deliberately a seed or constraint object. It is not yet a derived physical field equation.

A.237.5 Handoff admissibility clauses

The handoff is admissible only under the following five clauses:

H1 $\mathcal{A}_{\partial}^{r25}$ is typed as a local atlas,
H2 $\Sigma_{\mathrm{trans}}^{r26} = (0,0,0,0)$,
H3 $\mathcal{F}_{\partial} : G(N + \kappa N(N-1)) = 16$ remains recorded,
H4 $\Gamma_{\mathrm{guard}}^{r26} = (1,1,1,1,1)$,
H5 $\mathfrak{E}^{\mathrm{seed}}_{\mathrm{MM/HC}}$ is a local constraint seed, not a closure theorem.

These clauses define the permitted frontier-to-field-equation handoff. They also forbid the common over-reading in which a finite local atlas is mistaken for a global all-readout proof.

Theoremlet A.237.2 (r26 guarded atlas-to-field handoff)**.** *Under H1–H5, the compressed boundary-extended atlas $\mathcal{A}_{\partial}^{r25}$ is admissible as a local typed input seed for the MM/HC field-equation program. The handoff preserves all r25 guards.*

Proof. By H1 the atlas is a local typed object. By H2 no new transition seam debt is introduced. By H3 the frontier surface remains explicit, so the handoff does not hide the boundary-source admissibility conditions. By H4 all closure guards remain active. By H5 the field-equation side receives only a seed/constraint object. Hence the handoff is admissible as a local input to the MM/HC program, but no global closure conclusion follows. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.237.6 r26 decision ledger

The r26 decision ledger is

object	r26 result	guard
$\mathcal{A}_\partial^{r25}$	typed compressed local atlas	local only
$\tau_{01}, \tau_{12}, \tau_{23}$	transition-audited	$\Sigma_{\mathrm{trans}} = 0$
$\mathfrak{X}_\partial^{r26}$	frontier handoff object	$\Pi_{\mathrm{loc}}^{\mathrm{guard}}$
$\mathfrak{E}_{\mathrm{MM/HC}}^{\mathrm{seed}}$	local constraint seed	not a field equation
$C4$	$\beta_4 = 1/256$	unspent
$C5$	$\Phi_5^{(n)} > 0$ for finite n	no finite closure
$C6/T2$	$\mathsf{B}_{\mathrm{all/T2}}$	blocked
RH	photon-clock handoff probe only	not a theorem
global all-readout	unclaimed	guard active

A safe next source-only direction is

v4.7.0-r27: MM/HC seed-equation ansatz extraction from the guarded atlas handoff.

This should extract candidate seed constraints from $\mathfrak{X}_{\partial,\mathrm{loc}}^{r26}$ without upgrading them to physical field equations or closure theorems.

A.238 v4.7.0-r27: MM/HC Seed-Equation Ansatz Extraction from the Guarded Atlas Handoff

This source-only layer extracts the first MM/HC seed-equation ansatz family from $\mathfrak{X}_{\partial,\mathrm{loc}}^{r26}$. The extraction is deliberately conservative. The output is a typed family of local seed constraints, not a physical field equation and not a closure theorem.

A.238.1 Projected handoff input

The input object inherited from r26 is

$$\mathfrak{X}_\partial^{r26} = \left(\mathcal{A}_\partial^{r25}, \Sigma_{\mathrm{trans}}^{r26}, \mathcal{F}_\partial, \Gamma_{\mathrm{guard}}^{r26}\right),$$

with

$$\Sigma_{\mathrm{trans}}^{r26} = (0,0,0,0), \qquad \mathcal{F}_\partial : G\big(N + \kappa N(N-1)\big) = 16.$$

The seed extraction uses only the projected local object

$$\boxed{\mathfrak{X}_{\partial,\mathrm{loc}}^{r26} := \Pi_{\mathrm{loc}}^{\mathrm{guard}} \mathfrak{X}_\partial^{r26}.}$$

The projection is part of the syntax. No term may be extracted before passing through the local guard.

A.238.2 Seed-equation ansatz family

The first MM/HC seed family is

$$\boxed{\mathfrak{E}_{\mathrm{MM/HC}}^{\mathrm{seed,r27}} = \left(E_{\mathrm{rel}}, E_{\mathrm{src}}, E_{\mathrm{seam}}, E_{\mathrm{mix}}, E_{\mathrm{col}}, E_{\mathrm{clk}}\right).}$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Each component is a local constraint seed:

$$\begin{aligned} E_{\mathrm{rel}} &:= G\big(N + \kappa N(N-1)\big) - 16, \\ E_{\mathrm{src}} &:= H_\partial - 16, \\ E_{\mathrm{seam}} &:= \|\Sigma^{r26}_{\mathrm{trans}}\|^2, \\ E_{\mathrm{mix}} &:= \|P_\partial S^{\mathrm{mixker}}_{4,\mu}\|^2 + \left(\beta_4 - \frac{1}{256}\right)^2, \\ E_{\mathrm{col}} &:= \Phi_5^{(n+1)} - \frac{3}{4}\Phi_5^{(n)}, \\ E_{\mathrm{clk}} &:= G - \mathcal{C}_\gamma(\tau_v). \end{aligned}$$

Here E_{rel} records the valley-front relay frontier, E_{src} records the local boundary-source surplus, E_{seam} records transition exactness, E_{mix} records that the mixed-kernel guard is not consumed, E_{col} records the bounded $C5$ collar recurrence, and E_{clk} records the photon-clock timing probe.

Definition A.238.1 (r27 MM/HC seed ansatz)**.** *The r27 MM/HC seed ansatz is the six-component local constraint family* $\mathfrak{E}^{\mathrm{seed,r27}}_{\mathrm{MM/HC}}$ *extracted from* $\mathfrak{X}^{r26}_{\partial,\mathrm{loc}}$ *under the guard projector* $\Pi^{\mathrm{guard}}_{\mathrm{loc}}$.

A.238.3 Two-sided seed typing

The seed ansatz is written in the two-sided notation as

$$\boxed{{}^{\partial\mathrm{atlas}}{}_{\mathrm{loc/read}}\mathfrak{E}^{\mathrm{seed,r27\ MM/HC}}_{\mathrm{MM/HC}}{}_{\mathrm{constraint}}\cdot}$$

The left side records the local boundary-atlas readout, while the right side records the MM/HC constraint target. The expression is not a same-variance Einstein contraction. It is a typed local source-to-constraint interface.

A scalar audit functional may be used for bookkeeping:

$$\boxed{\mathcal{S}^{r27}_{\mathrm{seed}} = \lambda_{\mathrm{rel}}E^2_{\mathrm{rel}} + \lambda_{\mathrm{src}}E^2_{\mathrm{src},-} + \lambda_{\mathrm{seam}}E_{\mathrm{seam}} + \lambda_{\mathrm{mix}}E_{\mathrm{mix}} + \lambda_{\mathrm{col}}E^2_{\mathrm{col}} + \lambda_{\mathrm{clk}}E^2_{\mathrm{clk}},}$$

where $E_{\mathrm{src},-} := \max(0, 16 - H_\partial)$. This functional is a local bookkeeping device only. It is not an action principle and not a physical Lagrangian.

A.238.4 Admissibility and non-upgrade clauses

The ansatz is admissible only under the following clauses:

A1 $\mathfrak{X}^{r26}_\partial$ is first projected by $\Pi^{\mathrm{guard}}_{\mathrm{loc}}$,
A2 $E_{\mathrm{seam}} = 0$ is inherited from $\Sigma^{r26}_{\mathrm{trans}} = (0,0,0,0)$,
A3 E_{mix} preserves $\beta_4 = 1/256$ as unspent guard debt,
A4 E_{col} records only $\Phi_5^{(n+1)} = \frac{3}{4}\Phi_5^{(n)}$ for finite stages,
A5 E_{clk} is a local photon-clock probe and not an RH theorem,
A6 $\mathfrak{E}^{\mathrm{seed,r27}}_{\mathrm{MM/HC}}$ is not promoted to a physical field equation.

Lemma A.238.2 (r27 seed extraction preserves guards)**.** *The extraction of* $\mathfrak{E}^{\mathrm{seed,r27}}_{\mathrm{MM/HC}}$ *from* $\mathfrak{X}^{r26}_{\partial,\mathrm{loc}}$ *preserves the r26 guard vector.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. The projected handoff already has zero transition seam vector. The relay and boundary-source terms record local admissibility conditions but do not introduce an all-readout channel. The mixed-kernel term explicitly keeps $\beta_4 = 1/256$ as a recorded guard rather than consuming it. The collar term records the finite recurrence of $C5$ and includes no finite closure clause. The clock term is a timing probe only. Hence no guarded channel is discharged. □

Theoremlet A.238.3 (r27 guarded seed-equation ansatz)**.** *The six-component family* $\mathfrak{E}^{\mathrm{seed,r27}}_{\mathrm{MM/HC}}$ *is an admissible local MM/HC seed-equation ansatz extracted from the guarded atlas handoff. It is not a physical field equation and not a closure theorem.*

Proof. By A1 the input is local and guard-projected. By A2 the transition seam component is exact. By A3 the route-stable mixed-kernel residual is preserved rather than spent. By A4 no finite $C5$ closure is claimed. By A5 the Riemann-zero/photon-clock channel remains a probe. By A6 the output is a seed constraint family only. Therefore the ansatz is admissible as a guarded local MM/HC seed and no global conclusion follows. □

A.238.5 r27 decision ledger

The r27 decision ledger is

seed component	local role	guard status
E_{rel}	relay frontier	local only
E_{src}	boundary-source surplus	no global source
E_{seam}	transition seam exactness	$\Sigma_{\mathrm{trans}} = 0$
E_{mix}	mixed-kernel guard record	$\beta_4 = 1/256$ unspent
E_{col}	bounded $C5$ collar recurrence	no finite closure
E_{clk}	photon-clock handoff probe	not RH theorem

The global guard vector remains

$$\boxed{\Gamma^{r27}_{\mathrm{guard}} = (1,1,1,1,1).}$$

A safe next source-only direction is

v4.7.0-r28: seed-ansatz consistency scan and MM/HC coefficient typing.

This next step should classify the coefficients $\lambda_{\mathrm{rel}}, \lambda_{\mathrm{src}}, \lambda_{\mathrm{seam}}, \lambda_{\mathrm{mix}}, \lambda_{\mathrm{col}}, \lambda_{\mathrm{clk}}$ as bookkeeping weights, selector weights, or admissibility multipliers, without turning $\mathcal{S}^{r27}_{\mathrm{seed}}$ into a physical action.

A.239 v4.7.0-r28: Seed-Ansatz Consistency Scan and MM/HC Coefficient Typing

This source-only layer audits the r27 seed ansatz $\mathfrak{E}^{\mathrm{seed,r27}}_{\mathrm{MM/HC}}$ before any field-equation upgrade is attempted. The goal is coefficient typing. The scan deliberately prevents the local seed functional from being misread as a physical action, a Lagrangian, or a closure theorem.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.239.1 Consistency issue: signed surplus is not a zero residual

The r27 family contains the signed frontier coordinate

$$E_{\mathrm{rel}} = G\big(N + \kappa N(N-1)\big) - 16.$$

This coordinate is useful, but it is not always a residual that should be driven to zero. If a local relay chart is strictly admissible, then

$$H = G\big(N + \kappa N(N-1)\big) > 16,$$

so

$$E_{\mathrm{rel}} > 0.$$

Penalising E_{rel}^2 as an ordinary equality residual would incorrectly push a robustly admissible chart back onto the thin frontier $H = 16$. Thus the signed frontier coordinate must be typed as a monitor, while the admissibility penalty uses only the violation part

$$\boxed{E_{\mathrm{rel}}^- := \max\{0, 16 - H\}.}$$

Similarly, for the boundary-source surplus,

$$\boxed{E_{\mathrm{src}}^- := \max\{0, 16 - H_\partial\}.}$$

The positive surplus

$$\mu := \max\{0, H - 16\}$$

is recorded as robustness margin, not as debt.

Definition A.239.1 (r28 violation-projected relay residual)**.** *The r28 relay residual is the violation-projected quantity $E_{\mathrm{rel}}^- = \max\{0, 16 - H\}$, while the signed quantity $E_{\mathrm{rel}} = H - 16$ is retained only as a frontier coordinate and margin monitor.*

A.239.2 Typed coefficient classes

The r27 coefficients are now divided into three noninterchangeable classes:

$$\boxed{\lambda \in \{\lambda^{\mathrm{book}}, \lambda^{\mathrm{sel}}, \lambda^{\mathrm{adm}}\}.}$$

Here

λ^{book} bookkeeping or diagnostic weight only,
λ^{sel} selector/timing weight for admissible local routing,
λ^{adm} admissibility multiplier for a hard local guard.

The r28 typing ledger is

coefficient	r28 type	role
λ_{rel}	$\lambda^{\mathrm{book}} + \lambda^{\mathrm{adm}}$	signed monitor plus violation penalty $(E_{\mathrm{rel}}^-)^2$
λ_{src}	λ^{adm}	boundary-source deficit penalty $(E_{\mathrm{src}}^-)^2$
λ_{seam}	λ^{adm}	transition seam exactness guard
λ_{mix}	λ^{adm}	mixed-kernel guard lock, keeps $\beta_4 = 1/256$ unspent
λ_{col}	$\lambda^{\mathrm{book}} + \lambda^{\mathrm{adm}}_{\mathrm{loc}}$	finite collar recurrence consistency only
λ_{clk}	λ^{sel}	photon-clock timing selector/probe

The mixed notation $\lambda^{\mathrm{book}} + \lambda^{\mathrm{adm}}$ means that the signed value may be displayed diagnostically, but only its admissibility-violation projection may be treated as debt.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.239.3 Typed seed audit functional

The scalar r27 bookkeeping expression is therefore replaced, for audit purposes, by the typed functional

$$\boxed{\mathcal{S}_{\mathrm{seed}}^{r28,\mathrm{typed}} = \alpha_{\mathrm{rel}}(E_{\mathrm{rel}}^{-})^2 + \alpha_{\mathrm{src}}(E_{\mathrm{src}}^{-})^2 + \alpha_{\mathrm{seam}}E_{\mathrm{seam}} + \alpha_{\mathrm{mix}}E_{\mathrm{mix}} + \alpha_{\mathrm{col}}E_{\mathrm{col}}^2 + \alpha_{\mathrm{clk}}^{\mathrm{sel}}E_{\mathrm{clk}}^2.}$$

with the side condition that $\alpha_{\mathrm{clk}}^{\mathrm{sel}}$ is active only inside a declared photon-clock phase window. Outside such a declared window, E_{clk} remains a probe and contributes no hard debt.

For the r22–r25 boundary-source anchor

$$(N, \kappa, G) = \left(5, \frac{3}{4}, 1\right),$$

one has

$$H = 20, \qquad E_{\mathrm{rel}} = 4, \qquad E_{\mathrm{rel}}^{-} = 0.$$

Thus the anchor has positive margin, but no relay violation debt. This resolves the apparent contradiction between a robust pass and a zero-residual audit.

A.239.4 Seed-component consistency scan

The six r27 seed components pass the r28 consistency scan with the following interpretations:

component	consistency decision	allowed use
E_{rel}	consistent after violation projection	frontier monitor plus local deficit test
E_{src}	consistent after deficit projection	boundary-source admissibility test
E_{seam}	consistent hard exactness seed	$\Sigma_{\mathrm{trans}}^{r26} = 0$
E_{mix}	consistent guard-preserving seed	$\beta_4 = 1/256$ retained
E_{col}	consistent finite recurrence seed	no finite $C5$ closure
E_{clk}	consistent selector/probe seed	not an RH theorem

In particular, E_{rel} and E_{src} are not independent physical field equations. When $H_\partial = H$, they are the same local relay surplus viewed from two bookkeeping sides: the valley-front side and the boundary-source side.

Lemma A.239.2 (r28 coefficient typing removes the frontier-square artefact)**.** *Replacing E_{rel}^2 by $(E_{\mathrm{rel}}^{-})^2$ in the admissibility part of the seed audit prevents robust relay margins from being misclassified as residual debt.*

Proof. If $H \geq 16$, then $E_{\mathrm{rel}}^{-} = 0$, so the local relay chart contributes no admissibility debt regardless of the positive surplus $H - 16$. If $H < 16$, then $E_{\mathrm{rel}}^{-} = 16 - H > 0$, so the deficit is recorded. Thus the typed residual detects exactly local relay failure and does not penalise robust pass interior points. □

A.239.5 MM/HC coefficient typing rule

The r28 coefficient rule is:

> No coefficient may be interpreted before its seed component is typed.

Equivalently,

$\lambda_i E_i^2$ is admissible only after declaring whether E_i is a monitor, selector, violation residual, or hard gu

This rule blocks the common false upgrade

$$\mathcal{S}_{\mathrm{seed}} \longrightarrow \text{physical action.}$$

The present object remains

$$\boxed{\mathcal{S}_{\mathrm{seed}}^{r28,\mathrm{typed}} \in \text{local MM/HC seed-audit grammar only.}}$$

Theoremlet A.239.3 (r28 seed-ansatz consistency and coefficient typing)**.** *After violation projection and coefficient typing, the r27 seed ansatz is consistent as a guarded local MM/HC seed-audit system. It is not a physical field equation, not an action principle, and not a closure theorem.*

Proof. The only apparent inconsistency is the treatment of positive relay surplus as a zero residual. Definition A.239.1 resolves this by splitting the signed frontier monitor from the violation residual. The seam, mixed-kernel, collar, and clock components keep their r27 guards: seam exactness remains local, $\beta_4 = 1/256$ is preserved, finite $C5$ closure is not claimed, and the photon-clock term remains a probe. Since all coefficients are typed before use, no scalar audit expression can be promoted to an action or global field equation. Therefore the typed seed-audit system is consistent in the declared local grammar only. □

A.239.6 r28 guard ledger

The r28 guard ledger is

guard	r28 status	reason
$C4$	$\beta_4 = 1/256$ unspent	E_{mix} remains a guard lock
$C5$	no finite closure	E_{col} records recurrence only
$C6/T2$	blocked	$\chi_{\mathrm{all}} = 0$ remains active
RH	probe only	E_{clk} is selector/timing, not theorem
global all-readout	unclaimed	$\Pi_{\mathrm{loc}}^{\mathrm{guard}}$ remains mandatory

Thus

$$\boxed{\Gamma_{\mathrm{guard}}^{r28} = (1,1,1,1,1).}$$

A safe next source-only direction is

$$\boxed{\text{v4.7.0-r29: typed seed residual minimizer and selector-window calibration.}}$$

The next step should use the r28 typed residuals to scan local minimisers without allowing a physical-action or closure upgrade.

A.240 v4.7.0-r29 Typed Seed Residual Minimizer and Selector-Window Calibration

This source-only step uses the r28 typed seed audit to define a minimizer for the local MM/HC seed residuals. The important point is that the minimizer is applied to violation-projected residuals, not to signed surplus monitors. Therefore a robust relay interior point is not pushed back to the thin frontier.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.240.1 Typed residual functional

Let

$$H(N,\kappa,G) = G\big(N + \kappa N(N-1)\big)$$

be the local relay height. The r29 typed admissibility residual is

$$\boxed{\mathcal{D}^{r29}_{\text{typed}} = \lambda_{\text{rel}}\big(E^{-}_{\text{rel}}\big)^2 + \lambda_{\text{src}}\big(E^{-}_{\text{src}}\big)^2 + \lambda_{\text{seam}}\|\Sigma^{r26}_{\text{trans}}\|^2 + \lambda_{\text{mix}}\|P_{\partial}S^{\text{mixker}}_{4,\mu}\|^2 + \lambda_{\text{col}}\big(E_{\text{col}}\big)^2 + \lambda_{\text{clk}}\big(E^{\text{win}}_{\text{clk}}\big)^2.}$$

Here

$$E^{-}_{\text{rel}} = \max(0, 16 - H), \qquad E^{-}_{\text{src}} = \max(0, 16 - H_{\partial}).$$

The clock term is not a theorem term. It is a selector-window residual:

$$\boxed{E^{\text{win}}_{\text{clk}} = \max(0, G_{\text{req}} - \mathcal{C}_{\gamma}(\tau_v)).}$$

Thus photon-clock timing is used only to decide whether a local relay window is available.

Definition A.240.1 (r29 typed zero-debt minimizer)**.** *A local seed chart is a typed zero-debt minimizer if*

$$\mathcal{D}^{r29}_{\text{typed}} = 0$$

inside the declared local grammar. This means that all admissibility deficits, seam debts, mixed-kernel couplings, collar-recurrence defects, and clock-window deficits vanish. It does not mean that the signed surplus monitors vanish.

A.240.2 Minimizer degeneracy and margin classes

Because r28 removed the frontier-square artefact, every admissible relay point with

$$H \geq 16$$

has zero relay debt. Hence the minimizer set is not a single point. It has a thin frontier and an interior.

We classify zero-debt points by margin

$$\mu = H - 16.$$

The local classes are

class	condition
frontier	$\mu = 0$
thin	$0 < \mu < 1$
robust	$1 \leq \mu < 4$
comfortable	$\mu \geq 4.$

Thus the finite-grid minimizer

$$(N,\kappa,G) = (4,1,1)$$

remains a zero-debt point, but only a frontier minimizer. The r22–r25 anchor

$$(N,\kappa,G) = \left(5, \frac{3}{4}, 1\right)$$

is a comfortable zero-debt minimizer with

$$H = 20, \qquad \mu = 4.$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.240.3 Selector-window calibration

For a fixed overlap strength

$$S(N,\kappa) = N + \kappa N(N-1),$$

the gain required for relay is

$$\boxed{G_{\mathrm{req}}(N,\kappa) = \frac{16}{S(N,\kappa)}.}$$

For a strict margin-one relay it is

$$\boxed{G_{\mathrm{req}}^{(+1)}(N,\kappa) = \frac{17}{S(N,\kappa)}.}$$

For the comfortable r22 anchor

$$S\left(5, \frac{3}{4}\right) = 20,$$

so

$$G_{\mathrm{req}} = \frac{4}{5}, \qquad G_{\mathrm{req}}^{(+1)} = \frac{17}{20}.$$

If a photon-clock window satisfies

$$\mathcal{C}_\gamma(\tau_v) \geq \frac{17}{20},$$

then the anchor has at least one full unit of relay margin. If it satisfies

$$\mathcal{C}_\gamma(\tau_v) \geq \frac{4}{5},$$

it is relay-admissible but may be near the thin boundary.

Lemma A.240.2 (r29 selector-window calibration)**.** *For fixed local packet data (N,κ), the selector-window threshold is*

$$G_{\mathrm{req}} = 16/S(N,\kappa).$$

The clock term contributes zero selector debt exactly when

$$\mathcal{C}_\gamma(\tau_v) \geq G_{\mathrm{req}}.$$

Proof. The local relay condition is

$$GS(N,\kappa) \geq 16.$$

Solving for the required gain gives $G_{\mathrm{req}} = 16/S(N,\kappa)$. Since the clock term is typed as a selector-window residual,

$$E_{\mathrm{clk}}^{\mathrm{win}} = \max(0, G_{\mathrm{req}} - \mathcal{C}_\gamma(\tau_v)),$$

it vanishes exactly when the available photon-clock gain meets or exceeds the required threshold. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.240.4 Typed minimizer scan result

The r29 finite scan uses the same local grid as r24/r25 and evaluates the typed residual. It finds that the typed residual minimizer set is precisely the admissible region of the local frontier map, stratified by margin. The audit therefore selects two representatives:

$$\boxed{(4,1,1) \quad \text{as the frontier zero-debt representative,}}$$

and

$$\boxed{\left(5,\frac{3}{4},1\right) \quad \text{as the comfortable zero-debt representative.}}$$

The second representative is retained as the operational boundary-source anchor because it has margin 4 and lies inside the robust local collar.

Theoremlet A.240.3 (r29 typed residual minimizer)**.** *Using violation-projected residuals, the local MM/HC seed audit admits zero-debt minimizers throughout the relay-admissible region* $H \geq 16$*. The robust boundary-source anchor* $(5, 3/4, 1)$ *is a comfortable zero-debt minimizer, while* $(4, 1, 1)$ *is only a frontier zero-debt minimizer.*

Proof. For every point with $H \geq 16$, both E^{-}_{rel} and E^{-}_{src} vanish. The seam and mixed-kernel terms vanish by the local boundary-source typing clauses, the collar recurrence is the r28 bookkeeping recurrence, and the clock term vanishes when the selector window is met. Thus all admissible points are typed zero-debt minimizers. The margin stratification then separates the thin frontier from robust and comfortable interior points. For $(4, 1, 1)$, $H = 16$, so the margin is zero. For $(5, 3/4, 1)$, $H = 20$, so the margin is four. □

A.240.5 r29 guard ledger

The minimizer is local and typed. It does not create a physical action, a field equation, or a closure theorem. The r29 guard ledger is

guard	r29 status	reason
$C4$	$\beta_4 = 1/256$ unspent	E_{mix} remains a guard-lock
$C5$	no finite closure	E_{col} is recurrence consistency only
$C6/T2$	blocked	$\chi_{\mathrm{all}} = 0$ remains active
RH	probe only	$E^{\mathrm{win}}_{\mathrm{clk}}$ is a selector-window term
global all-readout	unclaimed	$\Pi^{\mathrm{guard}}_{\mathrm{loc}}$ remains mandatory

Therefore

$$\boxed{\Gamma^{r29}_{\mathrm{guard}} = (1,1,1,1,1).}$$

A safe next source-only direction is

$$\boxed{\text{v4.7.0-r30: typed MM/HC seed normal form and admissibility certificate.}}$$

The next step should package the typed minimizer as a normal-form certificate for local MM/HC seed admissibility, still without promoting it to a physical field equation or global closure theorem.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.241 v4.7.0-r30 Typed MM/HC Seed Normal Form and Admissibility Certificate

Source-only status. This r30 layer is a source-only continuation of the v4.7.0 internal line. It packages the r28–r29 typed residual calculus as a local MM/HC seed normal form and an admissibility certificate. It does not introduce a physical action, a Lagrangian, a field equation, a T2 bridge, an RH statement, or a global all-readout closure theorem.

A.241.1 Typed seed normal form

Let

$$S(N,\kappa) = N + \kappa N(N-1), \qquad H = GS(N,\kappa).$$

The r30 local seed normal form is the guarded tuple

$$\boxed{\mathfrak{N}^{r30}_{\mathrm{seed}} = \left(N, \kappa, G; \Sigma_{\mathrm{trans}}, P_\partial S^{\mathrm{mixker}}_{4,\mu}, \beta_4, \Phi^{(n)}_5, \mathcal{C}_\gamma(\tau_v), \Gamma_{\mathrm{guard}}\right)}$$

subject to the following typed clauses:

NF1 two-sided local boundary-source/relay typing is declared;
NF2 $E^-_{\mathrm{rel}} = \max(0, 16 - H) = 0$;
NF3 $E^-_{\mathrm{src}} = \max(0, 16 - H_\partial) = 0$;
NF4 $\Sigma_{\mathrm{trans}} = 0$;
NF5 $P_\partial S^{\mathrm{mixker}}_{4,\mu} = 0$ and $\beta_4 = 1/256$;
NF6 $\Phi^{(n+1)}_5 = \frac{3}{4}\Phi^{(n)}_5$ is only a local collar recurrence;
NF7 $\mathcal{C}_\gamma(\tau_v) \geq 16/S(N,\kappa)$ is only a selector-window probe;
NF8 $\Gamma_{\mathrm{guard}} = (1,1,1,1,1)$ remains active.

The normal form is intentionally asymmetric with respect to surplus: positive relay surplus is not penalized. Only deficit is counted through E^-_{rel} and E^-_{src}.

Definition A.241.1 (r30 typed MM/HC seed normal form)**.** *A local boundary-atlas handoff is in typed MM/HC seed normal form if it admits a tuple $\mathfrak{N}^{r30}_{\mathrm{seed}}$ satisfying NF1–NF8. The declaration is local: it certifies admissibility of the guarded seed constraints only, not a physical field equation and not a closure theorem.*

A.241.2 Admissibility certificate

The associated certificate is

$$\boxed{\mathsf{Cert}^{r30}_{\mathrm{MM/HC}} = \left[\mathrm{Type}_{\partial/\mathrm{relay}}, E^-_{\mathrm{rel}} = 0, E^-_{\mathrm{src}} = 0, \Sigma_{\mathrm{trans}} = 0, P_\partial S^{\mathrm{mixker}}_{4,\mu} = 0, \beta_4 = \frac{1}{256}, \Phi^{(n+1)}_5 = \frac{3}{4}\Phi^{(n)}_5, \mathcal{C}_\gamma(\tau_v) \geq \right.}$$

For the comfortable r29/r30 operational anchor

$$(N,\kappa,G) = \left(5, \frac{3}{4}, 1\right),$$

one has

$$S = 20, \qquad H = 20, \qquad \mu = H - 16 = 4.$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Hence

$$E^{-}_{\mathrm{rel}} = 0, \qquad G_{\mathrm{req}} = \frac{4}{5}, \qquad G^{(+1)}_{\mathrm{req}} = \frac{17}{20}.$$

Since $G = 1$, the anchor satisfies the selector-window requirement with margin. The frontier representative $(4, 1, 1)$ also satisfies the certificate but only critically, with $H = 16$ and $\mu = 0$. It is therefore kept as a frontier witness, not as the operational anchor.

Proposition A.241.2 (r30 local admissibility certificate)**.** *The comfortable anchor* $(5, 3/4, 1)$ *satisfies the typed MM/HC seed normal form with margin* 4*. The certificate has zero typed admissibility debt while preserving all global guards.*

Proof. For the anchor, $S = 20$ and $H = 20$, so $E^{-}_{\mathrm{rel}} = \max(0, 16 - 20) = 0$. The boundary-source handoff has $H_\partial = 20$, so $E^{-}_{\mathrm{src}} = 0$. By the r26 transition audit, $\Sigma_{\mathrm{trans}} = 0$. By the r20–r22 boundary-source typing, $P_\partial S^{\mathrm{mixker}}_{4,\mu} = 0$ and $\beta_4 = 1/256$ is left unspent. The collar term is the already guarded recurrence $\Phi^{(n+1)}_5 = (3/4)\Phi^{(n)}_5$, which is not finite closure. The clock selector threshold is $G_{\mathrm{req}} = 16/20 = 4/5$, and the anchor has $G = 1$. Thus every local admissibility component is satisfied while $\Gamma_{\mathrm{guard}} = (1, 1, 1, 1, 1)$ remains active. □

A.241.3 Certificate classes

The r30 certificate separates admissible points into three useful local classes:

class	condition	interpretation
frontier-cert	$H = 16$	zero debt but no margin
robust-cert	$17 \le H < 20$	positive protected margin
comfortable-cert	$H \ge 20$	operational anchor region

The certificate deliberately avoids selecting the smallest zero-debt point as the operational seed. The point $(4, 1, 1)$ is minimal on the finite grid, but it is thin. The point $(5, 3/4, 1)$ carries margin four and is therefore retained as the preferred local seed normal form.

Corollary A.241.3 (r30 frontier/anchor separation)**.** *The r30 normal form distinguishes minimal admissibility from robust operational admissibility. A frontier-zero-debt point is allowed as a witness, but only a positive-margin point is used as the seed anchor for subsequent MM/HC coefficient work.*

A.241.4 r30 guard certificate

The r30 admissibility certificate is local and guard-preserving:

guard	r30 status	certificate effect
$C4$	$\beta_4 = 1/256$ unspent	mixed-kernel guard preserved
$C5$	no finite closure	collar recurrence only
$C6/T2$	blocked	$\chi_{\mathrm{all}} = 0$ remains active
RH	probe only	$\mathcal{C}_\gamma$ only calibrates selector windows
global all-readout	unclaimed	$\Pi^{\mathrm{guard}}_{\mathrm{loc}}$ remains mandatory

Therefore

$$\boxed{\Gamma^{r30}_{\mathrm{guard}} = (1, 1, 1, 1, 1).}$$

A safe next source-only direction is

$$\boxed{\text{v4.7.0-r31: typed coefficient skeleton and local seed functional packaging.}}$$

The next step should introduce a typed coefficient skeleton for the certified seed normal form, still as a local audit functional and not as a physical action or field equation.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.242 v4.7.0-r31 Source-Only: Typed Coefficient Skeleton and Local Seed Functional Packaging

This r31 source-only layer packages the r30 seed normal form as a typed local seed functional. The purpose is strictly audit-theoretic: the functional below is a local MM/HC seed bookkeeping device, not a physical action, not a Lagrangian, not a derived field equation, and not a closure theorem.

A.242.1 Coefficient Skeleton

Let

$$\mathfrak{N}^{r30}_{\mathrm{seed}} = (E^-_{\mathrm{rel}}, E^-_{\mathrm{src}}, \Sigma_{\mathrm{trans}}, P_\partial S^{\mathrm{mixker}}_{4,\mu}, \beta_4, \Phi^{(n)}_5, G, \mathcal{C}_\gamma(\tau_v))$$

be the local seed normal form certified in r30. r31 assigns types to the coefficient skeleton before any functional is written:

$$\boxed{\Lambda^{r31}_{\mathrm{seed}} = (\lambda^-_{\mathrm{rel}}, \lambda^-_{\mathrm{src}}, \lambda_{\mathrm{seam}}, \lambda_{\mathrm{mix}}, \lambda_\beta, \lambda_{\mathrm{col}}, \lambda_{\mathrm{clk}}, \lambda_{\mathrm{cmf}}).}$$

The typing is:

coefficient	type	allowed role
λ^-_{rel}	admissibility penalty	$(E^-_{\mathrm{rel}})^2$
λ^-_{src}	admissibility penalty	$(E^-_{\mathrm{src}})^2$
λ_{seam}	hard seam guard	$\|\Sigma_{\mathrm{trans}}\|^2$
λ_{mix}	mixed-kernel guard-lock	$\|P_\partial S^{\mathrm{mixker}}_{4,\mu}\|^2$
λ_β	C4 ledger lock	$(\beta_4 - 1/256)^2$
λ_{col}	local recurrence bookkeeping	$(\Phi^{(n+1)}_5 - \frac{3}{4}\Phi^{(n)}_5)^2$
λ_{clk}	selector/probe coefficient	$(G - \mathcal{C}_\gamma(\tau_v))^2_{\mathrm{sel}}$
λ_{cmf}	comfort-margin selector	$(E^-_{\mathrm{cmf}})^2$

where the last term is optional and only used when a comfortable certificate is requested. It is not part of the bare admissibility certificate.

A.242.2 Local Seed Functional

The local audit functional is

$$\boxed{\begin{aligned}\mathcal{J}^{r31}_{\mathrm{seed}} &= \lambda^-_{\mathrm{rel}}(E^-_{\mathrm{rel}})^2 + \lambda^-_{\mathrm{src}}(E^-_{\mathrm{src}})^2 + \lambda_{\mathrm{seam}}\|\Sigma^{r26}_{\mathrm{trans}}\|^2 \\ &\quad + \lambda_{\mathrm{mix}}\|P_\partial S^{\mathrm{mixker}}_{4,\mu}\|^2 + \lambda_\beta\left(\beta_4 - \frac{1}{256}\right)^2 \\ &\quad + \lambda_{\mathrm{col}}\left(\Phi^{(n+1)}_5 - \frac{3}{4}\Phi^{(n)}_5\right)^2 + \lambda_{\mathrm{clk}}\,(G - \mathcal{C}_\gamma(\tau_v))^2_{\mathrm{sel}}\,.\end{aligned}}$$

The signed monitor

$$E_{\mathrm{rel}} = H - 16$$

is not squared in the admissibility functional. Only the violation projection

$$E^-_{\mathrm{rel}} = \max(0, 16 - H)$$

contributes debt. Hence interior relay surplus is not penalized.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

For comfortable certification one may add the explicitly typed selector term

$$\boxed{\mathcal{J}^{r31}_{\mathrm{seed},+4} = \mathcal{J}^{r31}_{\mathrm{seed}} + \lambda_{\mathrm{cmf}}(E^{-}_{\mathrm{cmf}})^2, \qquad E^{-}_{\mathrm{cmf}} = \max(0, 20 - H).}$$

This term distinguishes the comfortable anchor from a thin frontier pass, but it is not a new physical principle.

A.242.3 Anchor Evaluation

At the comfortable operational anchor

$$(N, \kappa, G) = \left(5, \frac{3}{4}, 1\right)$$

one has

$$S = 20, \qquad H = 20, \qquad E^{-}_{\mathrm{rel}} = 0, \qquad E^{-}_{\mathrm{src}} = 0, \qquad E^{-}_{\mathrm{cmf}} = 0.$$

Together with the r30 certificate clauses

$$\Sigma^{r26}_{\mathrm{trans}} = 0, \qquad P_{\partial} S^{\mathrm{mixker}}_{4,\mu} = 0, \qquad \beta_4 = \frac{1}{256}, \qquad \Phi_5^{(n+1)} = \frac{3}{4}\Phi_5^{(n)},$$

and a matched selector window

$$G = \mathcal{C}_{\gamma}(\tau_v),$$

this gives

$$\boxed{\mathcal{J}^{r31}_{\mathrm{seed}} = 0, \qquad \mathcal{J}^{r31}_{\mathrm{seed},+4} = 0}$$

inside the declared local seed grammar.

At the frontier witness

$$(N, \kappa, G) = (4, 1, 1)$$

one has

$$H = 16, \qquad E^{-}_{\mathrm{rel}} = 0,$$

so the bare admissibility functional may vanish. However

$$E^{-}_{\mathrm{cmf}} = 4,$$

so the comfortable certificate does not vanish. Thus r31 preserves the r30 classification:

$$\boxed{(4, 1, 1) = \text{frontier-cert only}, \qquad \left(5, \frac{3}{4}, 1\right) = \text{comfortable-cert.}}$$

A.242.4 Guard Preservation

The r31 functional is local and typed. It does not spend the persistent C4 mixed-kernel guard, does not close C5 at any finite stage, does not unblock C6/T2, does not identify Riemann zeros with photon events, and does not claim a global all-readout closure. Hence

$$\boxed{\Gamma^{r31}_{\mathrm{guard}} = (1, 1, 1, 1, 1).}$$

The safe next source-only direction is

$$\boxed{\text{v4.7.0-r32: local seed functional sanity table and coefficient-domain guard.}}$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.243 v4.7.0-r32 Source-Only Local Seed Functional Sanity Table and Coefficient-Domain Guard

Source-only status. This subsection is an internal source-only continuation of the v4.7.0 line. No PDF artefact is generated at this stage. The purpose is to sanity-check the r31 local seed functional against representative admissible, frontier, robust, comfortable, and debt-positive sites, and to record an explicit coefficient -domain guard.

A.243.1 Sanity Table for the Local Seed Functional

Recall the local relay score

$$S(N,\kappa) = N + \kappa N(N-1), \qquad H = G\,S(N,\kappa),$$

with admissibility debt

$$E^{-}_{\mathrm{rel}} = \max(0, 16 - H)$$

and optional comfort-margin selector

$$E^{-}_{\mathrm{cmf}} = \max(0, 20 - H).$$

The signed quantity

$$E_{\mathrm{rel}} = H - 16$$

remains only a frontier monitor and is not squared as debt.

The representative sanity table is:

site	N	κ	G	H	E^{-}_{rel}	E^{-}_{cmf}
deep-debt	3	1/2	1	6	10	14
local-debt	4	3/4	1	13	3	7
frontier witness	4	1	1	16	0	4
weak five-packet frontier	5	11/20	1	16	0	4
gain frontier	5	3/4	4/5	16	0	4
margin-one selector	5	3/4	17/20	17	0	3
robust box corner	5	7/10	9/10	171/10	0	29/10
comfortable anchor	5	3/4	1	20	0	0

Hence the bare r31 admissibility part vanishes exactly on and above the relay frontier $H = 16$, while the optional comfort selector distinguishes thin frontier passes from the operational comfortable anchor $H = 20$.

Lemma A.243.1 (r32 local sanity separation)**.** *Inside the declared local seed grammar, the r31/r32 typed seed audit separates three cases without changing the guard vector:*

$$H < 16 \Rightarrow E^{-}_{\mathrm{rel}} > 0, \qquad 16 \le H < 20 \Rightarrow E^{-}_{\mathrm{rel}} = 0 < E^{-}_{\mathrm{cmf}}, \qquad H \ge 20 \Rightarrow E^{-}_{\mathrm{rel}} = E^{-}_{\mathrm{cmf}} = 0.$$

Consequently a frontier certificate, a robust local certificate, and a comfortable certificate remain distinct.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.243.2 Coefficient-Domain Guard

The admissible coefficient domain is recorded as

$$\boxed{\mathcal{D}_{\lambda}^{r32}}$$

with the following typed clauses:

$$\begin{aligned}
&\lambda_{\mathrm{rel}}^{-}, \lambda_{\mathrm{src}}^{-} \geq 0, && \text{violation-projected admissibility penalties only,}\\
&\lambda_{\mathrm{seam}} > 0, && \text{hard transition-seam guard,}\\
&\lambda_{\mathrm{mix}} > 0, && \text{mixed-kernel guard-lock,}\\
&\lambda_{\beta} > 0, && \text{C4 ledger lock at } \beta_4 = 1/256,\\
&\lambda_{\mathrm{col}} \geq 0, && \text{local C5 recurrence bookkeeping,}\\
&\lambda_{\mathrm{clk}} \geq 0, && \text{selector/probe weight only,}\\
&\lambda_{\mathrm{cmf}} \geq 0, && \text{optional comfort-margin selector.}
\end{aligned}$$

No coefficient in $\mathcal{D}_{\lambda}^{r32}$ may convert the signed monitor $E_{\mathrm{rel}} = H - 16$ into a debt term. Debt is carried only by the negative part E_{rel}^{-}. Likewise, no coefficient may spend the C4 ledger lock, unblock C6/T2, or turn the photon-clock probe into an RH assertion.

The guarded local audit functional is therefore read as

$$\boxed{\mathcal{J}_{\mathrm{seed}}^{r32} : \mathcal{D}_{\lambda}^{r32} \times \mathfrak{N}_{\mathrm{seed}}^{r30} \longrightarrow \mathbb{R}_{\geq 0}}$$

with the same meaning as in r31: it is a local audit functional, not an action, not a Lagrangian, not a derived field equation, and not a closure theorem.

A.243.3 Guard Preservation

The r32 sanity table and coefficient-domain guard do not alter the previous mathematics. They prevent a misreading of the signed frontier monitor as a least-squares field residual and preserve the operational separation between frontier, robust, and comfortable relay certificates. Thus

$$\boxed{\Gamma_{\mathrm{guard}}^{r32} = (1, 1, 1, 1, 1).}$$

Explicitly, C4 remains route-stable with

$$\beta_4 = \frac{1}{256},$$

C5 remains an asymptotic bounded collar with no finite closure, C6/T2 remains blocked, the Riemann-zero language remains a photon-clock handoff probe only, and global all-readout closure is not claimed.

A.244 v4.7.0-r33 Source-Only: Local Seed Functional Parameter Sweep and Comfort-Margin Calibration

This source-only layer performs the safe r33 follow-up to the r32 sanity table. No PDF is generated in this step. The purpose is to sweep the local seed functional over a finite parameter box and to calibrate the optional comfort margin without changing any guard, without turning the signed frontier monitor into debt, and without promoting the seed audit into a physical action or field equation.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.244.1 Finite Parameter Sweep

The sweep uses

$$N \in \{3,4,5,6,7,8\},$$
$$\kappa \in \left\{\frac{1}{2}, \frac{11}{20}, \frac{3}{5}, \frac{2}{3}, \frac{7}{10}, \frac{3}{4}, \frac{4}{5}, \frac{9}{10}, 1\right\},$$
$$G \in \left\{\frac{3}{4}, \frac{4}{5}, \frac{17}{20}, \frac{9}{10}, \frac{19}{20}, 1\right\}.$$

For each point we evaluate

$$S(N,\kappa) = N + \kappa N(N-1), \qquad H = G\,S(N,\kappa),$$

$$E^{-}_{\mathrm{rel}} = \max(0, 16 - H), \qquad E^{-}_{\mathrm{cmf}} = \max(0, 20 - H).$$

The classification is:

$$\begin{array}{ll} H < 16 & \text{debt-positive,} \\ H = 16 & \text{frontier-zero-debt,} \\ 16 < H < 17 & \text{thin-zero-debt,} \\ 17 \le H < 20 & \text{robust-zero-debt,} \\ H \ge 20 & \text{comfortable-zero-debt.} \end{array}$$

The finite sweep contains 324 cases and yields

class	# cases
debt positive	130
frontier zero debt	3
thin zero debt	6
robust zero debt	24
comfortable zero debt	161

Thus the r32 typing behaves as intended: a broad zero-debt region exists above relay threshold, but the optional comfort selector still separates thin and robust admissibility from the operational comfort regime.

A.244.2 Comfort-Margin Calibration

For a fixed pair (N,κ), the relay, margin-one, and comfort thresholds are

$$G_{\mathrm{rel}}(N,\kappa) = \frac{16}{S(N,\kappa)}, \qquad G_{+1}(N,\kappa) = \frac{17}{S(N,\kappa)}, \qquad G_{\mathrm{cmf}}(N,\kappa) = \frac{20}{S(N,\kappa)}.$$

The five-packet calibration slice is

site	N	κ	$S(N,\kappa)$	$G_{\mathrm{cmf}} = 20/S$
five-packet frontier	5	$\frac{11}{20}$	16	$\frac{5}{4}$
margin-one threshold	5	$\frac{3}{5}$	17	$\frac{20}{17}$
robust box corner	5	$\frac{7}{10}$	19	$\frac{20}{19}$
comfortable anchor	5	$\frac{3}{4}$	20	1

Consequently, at the comfortable anchor

$$(N,\kappa,G) = \left(5, \frac{3}{4}, 1\right)$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

one has

$$S = 20, \qquad H = 20, \qquad E^-_{\mathrm{rel}} = E^-_{\mathrm{cmf}} = 0.$$

The same five-packet source with $G = \frac{4}{5}$ sits exactly on the relay frontier, while $G = \frac{17}{20}$ gives the first unit of positive margin. Hence the comfort selector is not an admissibility guard; it is a local preference for the stable operational anchor.

Lemma A.244.1 (r33 comfort-margin calibration)**.** *Inside the declared local seed grammar, the optional comfort-margin term* $E^-_{\mathrm{cmf}} = \max(0, 20-H)$ *calibrates the local selector preference without altering admissibility. In particular,*

$$H \geq 16 \Rightarrow E^-_{\mathrm{rel}} = 0,$$

but only

$$H \geq 20 \Rightarrow E^-_{\mathrm{cmf}} = 0.$$

Therefore a frontier certificate and a comfortable operational certificate are kept distinct without creating new debt.

A.244.3 Representative Sweep Anchors

The parameter sweep confirms the hierarchy already suggested by r32:

$$(4,1,1): \quad H = 16, \qquad \mu = 0, \qquad \text{frontier-cert only},$$

$$\left(5, \frac{11}{20}, 1\right): \quad H = 16, \qquad \text{weak five-packet frontier},$$

$$\left(5, \frac{3}{4}, \frac{4}{5}\right): \quad H = 16, \qquad \text{gain frontier},$$

$$\left(5, \frac{3}{4}, \frac{17}{20}\right): \quad H = 17, \qquad \text{margin-one selector},$$

$$\left(5, \frac{7}{10}, \frac{9}{10}\right): \quad H = \frac{171}{10}, \qquad \text{robust box corner},$$

$$\left(5, \frac{3}{4}, 1\right): \quad H = 20, \qquad \text{comfortable anchor}.$$

The frontier minimizer remains useful as a boundary witness, but the comfort anchor remains the preferred local source point for downstream MM/HC seed packaging.

A.244.4 Guard Preservation

The sweep and comfort calibration do not alter any global guard. They only classify local seed-functional behavior inside the already declared local boundary-source grammar. Thus

$$\boxed{\Gamma^{r33}_{\mathrm{guard}} = (1,1,1,1,1).}$$

Explicitly, C4 remains route-stable with

$$\beta_4 = \frac{1}{256},$$

C5 remains an asymptotic bounded collar with no finite closure, C6/T2 remains blocked, the Riemann-zero language remains a photon-clock handoff probe only, and global all-readout closure is not claimed.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Next source-only direction. The safe next step is

v4.7.0-r34: comfort-calibrated seed selector and guard-stable handoff summary.

A.245 v4.7.0-r34: Comfort-Calibrated Seed Selector and Guard-Stable Handoff Summary

This source-only step packages the r33 parameter sweep into a conservative selector rule. The purpose is not to introduce a new field equation, action, Lagrangian, or closure theorem. The purpose is only to choose a stable local seed for downstream MM/HC bookkeeping while preserving every active guard.

A.245.1 Selector Rule

Let

$$S(N,\kappa) = N + \kappa N(N-1), \qquad H(N,\kappa,G) = GS(N,\kappa).$$

The signed frontier monitor remains

$$E_{\mathrm{rel}} = H - 16,$$

but the admissibility debt is still only

$$E^{-}_{\mathrm{rel}} = \max(0, 16 - H).$$

The comfort deficit is

$$E^{-}_{\mathrm{cmf}} = \max(0, 20 - H).$$

Thus the selector is not a minimizer of the signed frontier monitor. It is a filtered local choice rule:

$$\boxed{E^{-}_{\mathrm{rel}} = 0, \quad E^{-}_{\mathrm{src}} = 0, \quad \Gamma_{\mathrm{guard}} = (1,1,1,1,1),}$$

followed, for operational handoff purposes, by the comfort preference

$$\boxed{E^{-}_{\mathrm{cmf}} = 0.}$$

Frontier witnesses remain useful boundary witnesses, but are not selected as operational handoff anchors unless no comfortable seed is available.

A.245.2 Representative Selector Table

The r34 selector keeps the known frontier witness visible but chooses the comfortable anchor for downstream local MM/HC seed packaging:

$$(4,1,1): \quad H = 16, \qquad \text{frontier witness only},$$

whereas

$$\boxed{\left(5, \frac{3}{4}, 1\right): \quad S = 20, \quad H = 20, \quad \mu = 4}$$

is selected as the comfort-calibrated operational seed. It satisfies

$$E^{-}_{\mathrm{rel}} = 0, \qquad E^{-}_{\mathrm{cmf}} = 0.$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

The robust lower corner

$$\left(5, \frac{7}{10}, \frac{9}{10}\right)$$

remains a robustness certificate, since

$$H \geq \frac{171}{10} > 16,$$

but it is not the preferred handoff anchor because the comfortable anchor gives larger margin without changing the declared local grammar.

A.245.3 Guard-Stable Handoff Object

The local handoff object produced by r34 is

$$\boxed{\mathfrak{H}^{r34}_{\mathrm{seed}} = \left(\left(5, \frac{3}{4}, 1\right), \mathfrak{N}^{r30}_{\mathrm{seed}}, \mathcal{J}^{r31}_{\mathrm{seed}}, \mathcal{D}^{r32}_{\lambda}, \Gamma^{r33}_{\mathrm{guard}}\right).}$$

This object is a local seed handoff summary only. It is not a physical field equation and not a closure statement.

The guard vector remains

$$\boxed{\Gamma^{r34}_{\mathrm{guard}} = (1, 1, 1, 1, 1).}$$

Explicitly:

$$\beta_4 = \frac{1}{256}$$

remains unspent; C5 remains an asymptotic bounded collar with no finite closure; C6/T2 remains blocked; the Riemann-zero language remains only a photon-clock handoff probe; and global all-readout closure remains unclaimed.

A.245.4 Handoff Summary

The r34 source-only conclusion is:

$$\boxed{\text{Use } \left(5, \frac{3}{4}, 1\right) \text{ as the comfort-calibrated local seed.}}$$

The finite local boundary-extended atlas and the typed seed functional may now be handed to the next MM/HC stage as a guarded seed package. The safe next step is

$$\boxed{\text{v4.7.0-r35: guarded MM/HC seed-to-interface map and no-action audit.}}$$

A.246 v4.7.0-r35 Source-Only: Guarded MM/HC Seed-to-Interface Map and No-Action Audit

A.246.1 Purpose and Scope

This source-only increment takes the r34 comfort-calibrated handoff object

$$\mathfrak{H}^{r34}_{\mathrm{seed}} = \left(\left(5, \frac{3}{4}, 1\right), \mathfrak{N}^{r30}_{\mathrm{seed}}, \mathcal{J}^{r31}_{\mathrm{seed}}, \mathcal{D}^{r32}_{\lambda}, \Gamma^{r33}_{\mathrm{guard}}\right)$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

and maps it to a guarded MM/HC interface skeleton. The purpose is only to identify admissible interface slots and no-action obligations. No physical action, no Lagrangian, no derived field equation, no T2 closure, no RH claim, and no global all-readout closure is asserted.

The selected seed remains

$$(N, \kappa, G) = \left(5, \frac{3}{4}, 1\right), \qquad S = 20, \qquad H = 20, \qquad \mu = 4.$$

The frontier witness $(4, 1, 1)$ remains a boundary witness only and is not promoted to the operational interface seed.

A.246.2 Guarded Interface Slots

We introduce a local interface map

$$\mathfrak{I}^{r35}_{\mathrm{MM/HC}} : \mathfrak{H}^{r34}_{\mathrm{seed}} \longrightarrow (\mathsf{I}_{\mathrm{bd}}, \mathsf{I}_{\mathrm{src}}, \mathsf{I}_{\mathrm{seam}}, \mathsf{I}_{\mathrm{mix}}, \mathsf{I}_{\mathrm{col}}, \mathsf{I}_{\mathrm{clk}}),$$

where each component is a typed interface slot rather than an equation of motion.

The slots are:

$$\mathsf{I}_{\mathrm{bd}} : \quad E^{-}_{\mathrm{rel}} = 0, \qquad E^{-}_{\mathrm{cmf}} = 0,$$

for the comfort-calibrated boundary relay condition;

$$\mathsf{I}_{\mathrm{src}} : \quad {}^{\partial \mathrm{src}}{}_{\mathrm{loc}}\mathcal{B}^{\mathrm{res}V0}_{0\ \mathrm{relay}},$$

for the two-sided local source/relay typing;

$$\mathsf{I}_{\mathrm{seam}} : \quad \Sigma_{\mathrm{trans}} = 0,$$

for the compressed atlas transition seam audit;

$$\mathsf{I}_{\mathrm{mix}} : \quad P_{\partial} S^{\mathrm{mixker}}_{4,\mu} = 0, \qquad \beta_4 = \frac{1}{256},$$

for the C4 mixed-kernel guard-lock;

$$\mathsf{I}_{\mathrm{col}} : \quad \Phi^{(n+1)}_5 = \frac{3}{4}\Phi^{(n)}_5, \qquad \forall n < \infty : \ \Phi^{(n)}_5 > 0,$$

for the local C5 bounded-collar recurrence without finite closure; and

$$\mathsf{I}_{\mathrm{clk}} : \quad G = \mathcal{C}_\gamma(\tau_v) \quad \text{or} \quad \mathcal{C}_\gamma(\tau_v) \geq \frac{16}{S(N, \kappa)},$$

for the photon-clock selector/probe interface.

Each slot is local and typed. None of these slots may be read as a global readout equation or as a physical field equation.

A.246.3 No-Action Audit

The interface map is equipped with a no-action audit vector

$$\boxed{\mathfrak{A}^{r35}_{\mathrm{noact}} = (0, 0, 0, 0, 0, 0)}$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

with the following entries:

$$
\begin{aligned}
A_1 &= \text{no physical action introduced},\\
A_2 &= \text{no Lagrangian introduced},\\
A_3 &= \text{no Euler–Lagrange variation asserted},\\
A_4 &= \text{no derived field equation asserted},\\
A_5 &= \text{no global readout promoted},\\
A_6 &= \text{no C4, C5, C6/T2, RH, or all-readout guard spent.}
\end{aligned}
$$

Thus r35 only records a guarded seed-to-interface map. It is a preparation layer for a possible later field-equation program, not the program itself.

A.246.4 Admissibility Matrix

For compact bookkeeping, the interface slots may be summarized by the local admissibility matrix

$$
\mathcal{M}^{r35}_{\mathrm{int}} = \begin{pmatrix} 1 & 1 & 1 & 1 & 1 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix},
$$

where the first row records local slot admissibility and the second row records new guard debt. Therefore

$$
\boxed{\mathrm{newDebt}(\mathcal{M}^{r35}_{\mathrm{int}}) = 0.}
$$

This matrix is not an MM dynamics matrix. It is only a local interface audit matrix.

A.246.5 Guard Preservation

The guard vector remains

$$
\boxed{\Gamma^{r35}_{\mathrm{guard}} = (1,1,1,1,1).}
$$

Equivalently:

$$
\boxed{\beta_4 = \frac{1}{256} \quad \text{remains unspent.}}
$$

C5 remains an asymptotic bounded collar with no finite closure; C6/T2 remains blocked; the Riemann-zero component remains only a photon-clock handoff probe; and global all-readout closure remains unclaimed.

A.246.6 r35 Handoff

The r35 handoff object is

$$
\boxed{\mathfrak{H}^{r35}_{\mathrm{int}} = \left(\mathfrak{H}^{r34}_{\mathrm{seed}}, \mathfrak{I}^{r35}_{\mathrm{MM/HC}}, \mathfrak{A}^{r35}_{\mathrm{noact}}, \mathcal{M}^{r35}_{\mathrm{int}}, \Gamma^{r35}_{\mathrm{guard}}\right).}
$$

The safe next step is

$$
\boxed{\text{v4.7.0-r36: interface residual table and MM/HC map consistency scan.}}
$$

This next step should still remain source-only unless a PDF is explicitly requested.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.247 v4.7.0-r36 Source-Only Interface Residual Table and MM/HC Map Consistency Scan

This source-only layer continues the r35 seed-to-interface map. It does not compile a PDF, does not introduce a physical action, and does not derive an Euler–Lagrange or field equation. Its only purpose is to test whether the six local MM/HC interface slots from r35 remain mutually consistent when read as residual channels.

The comfort-calibrated seed remains

$$(N,\kappa,G)=\left(5,\frac{3}{4},1\right),\qquad S=20,\qquad H=20,\qquad \mu=4.$$

The frontier witness $(4,1,1)$ remains only a boundary witness and is not used as the operational handoff seed.

A.247.1 Interface Residual Table

For each r35 interface slot define a local residual channel

$$\mathcal{R}^{r36}_{\mathrm{int}}=(R_{\mathrm{bd}},R_{\mathrm{src}},R_{\mathrm{seam}},R_{\mathrm{mix}},R_{\mathrm{col}},R_{\mathrm{clk}}).$$

The table below records the guarded interpretation.

slot	residual channel	value	status
I_{bd}	$R_{\mathrm{bd}}=E^{-}_{\mathrm{rel}}+E^{-}_{\mathrm{src}}$	0	local relay admissible
$\mathsf{I}_{\mathrm{src}}$	$R_{\mathrm{src}}=\delta_{\text{two-sided type}}$	0	typed local source/relay
$\mathsf{I}_{\mathrm{seam}}$	$R_{\mathrm{seam}}=\lVert\Sigma^{r26}_{\mathrm{trans}}\rVert^2$	0	transition seam exact
$\mathsf{I}_{\mathrm{mix}}$	$R_{\mathrm{mix}}=\lVert P_{\partial}S^{\mathrm{mixker}}_{4,\mu}\rVert^2+\left(\beta_4-\frac{1}{256}\right)^2$	0	C4 guard-lock preserved
$\mathsf{I}_{\mathrm{col}}$	$R_{\mathrm{col}}=\Phi_5^{(n+1)}-\frac{3}{4}\Phi_5^{(n)}$	0	bounded-collar recurrence only
$\mathsf{I}_{\mathrm{clk}}$	$R_{\mathrm{clk}}=\max\left(0,\frac{16}{S(N,\kappa)}-\mathcal{C}_\gamma(\tau_v)\right)$	0	selector/probe window admissible

For the comfort seed, $S(5,3/4)=20$, hence

$$\frac{16}{S(N,\kappa)}=\frac{4}{5},\qquad \mathcal{C}_\gamma(\tau_v)=G=1,$$

so the photon-clock selector residual vanishes as a local selector channel. This is not an RH statement. It is only the local clock-window reading already allowed by the r35 handoff.

Therefore

$$\boxed{\mathcal{R}^{r36}_{\mathrm{int}}=(0,0,0,0,0,0).}$$

A.247.2 Guard Residuals Versus Interface Residuals

The vanishing of $\mathcal{R}^{r36}_{\mathrm{int}}$ is not a global closure statement. It only means that the r35 local interface slots are internally consistent. The guard residuals remain logically separate:

$$\mathcal{B}^{r36}_{\mathrm{guard}}=\left(\frac{1}{256},\Phi_5^{(n)}>0\ (n<\infty),\mathsf{B}_{\mathrm{all/T2}},\mathsf{probe}_{\mathrm{RH}},\mathsf{unclaimed}_{\mathrm{all}}\right).$$

Thus the interface residual scan has zero local mismatch while the global guard profile remains active.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.247.3 MM/HC Map Consistency Scan

The r35 map is consistent iff the six local slots satisfy the following map checks:

$$\begin{aligned}
&\mathrm{MC1} : \mathsf{I}_{\mathrm{bd}} \text{ uses only } E^{-}_{\mathrm{rel}}, E^{-}_{\mathrm{src}} \text{ and does not square signed } E_{\mathrm{rel}},\\
&\mathrm{MC2} : \mathsf{I}_{\mathrm{src}} \text{ remains two-sided local and does not become an all-readout source},\\
&\mathrm{MC3} : \mathsf{I}_{\mathrm{seam}} \text{ imports no transition seam debt},\\
&\mathrm{MC4} : \mathsf{I}_{\mathrm{mix}} \text{ locks } \beta_4 = \tfrac{1}{256} \text{ instead of spending it},\\
&\mathrm{MC5} : \mathsf{I}_{\mathrm{col}} \text{ records only the } C5 \text{ recurrence and no finite closure},\\
&\mathrm{MC6} : \mathsf{I}_{\mathrm{clk}} \text{ remains a selector/probe and not an RH theorem.}
\end{aligned}$$

All six checks pass in the source-level scan:

$$\boxed{\mathrm{MC}^{r36} = (1,1,1,1,1,1), \qquad \mathrm{mapDebt}^{r36} = 0.}$$

A.247.4 No-Action Preservation

The no-action audit vector remains zero:

$$\boxed{\mathfrak{A}^{r36}_{\mathrm{noact}} = (0,0,0,0,0,0).}$$

No variational principle is introduced. No Lagrangian is declared. No Euler–Lagrange equation is varied. No physical field equation is derived. No all-readout promotion is made. No C4, C5, C6/T2, RH, or global-readout guard is spent.

A.247.5 r36 Handoff

The r36 handoff object is

$$\boxed{\mathfrak{H}^{r36}_{\mathrm{int}} = \left(\mathfrak{H}^{r35}_{\mathrm{int}}, \mathcal{R}^{r36}_{\mathrm{int}}, \mathrm{MC}^{r36}, \mathfrak{A}^{r36}_{\mathrm{noact}}, \Gamma^{r35}_{\mathrm{guard}}\right).}$$

It is a guard-stable local interface consistency certificate and nothing more. The safe next source-only step is

$$\boxed{\text{v4.7.0-r37: local MM/HC interface certificate and guard-stable residual packaging.}}$$

A.248 v4.7.0-r37 Local MM/HC Interface Certificate and Guard-Stable Residual Packaging

This r37 source-only layer packages the r36 zero-residual interface scan as a local certificate. The purpose is deliberately narrow: the certificate records that the six declared local MM/HC interface slots close their local residuals inside the current guarded grammar. It does not convert the local interface map into a physical action, a Lagrangian, an Euler–Lagrange system, a derived field equation, a T2 theorem, or an all-readout closure statement.

A.248.1 Certificate Object

Let the ordered local interface list be

$$\mathbb{I}^{r37} = (\mathsf{I}_{\mathrm{bd}}, \mathsf{I}_{\mathrm{src}}, \mathsf{I}_{\mathrm{seam}}, \mathsf{I}_{\mathrm{mix}}, \mathsf{I}_{\mathrm{col}}, \mathsf{I}_{\mathrm{clk}}).$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

The r36 local residual vector is inherited as

$$\mathcal{R}_{\mathrm{int}}^{r36} = (0,0,0,0,0,0),$$

and the r36 MM/HC map consistency vector is inherited as

$$\mathrm{MC}^{r36} = (1,1,1,1,1,1), \qquad \mathrm{mapDebt}^{r36} = 0.$$

The r37 local interface certificate is the tuple

$$\boxed{\mathfrak{C}_{\mathrm{int}}^{r37} = \left(\mathbb{I}^{r37}, \mathcal{R}_{\mathrm{int}}^{r36}, \mathrm{MC}^{r36}, \mathfrak{A}_{\mathrm{noact}}^{r36}, \Gamma_{\mathrm{guard}}^{r36}\right).}$$

It is valid only as a local interface certificate. The inclusion of $\Gamma_{\mathrm{guard}}^{r36}$ is part of the certificate, not a debt-spending operation.

A.248.2 Slotwise Certificate Ledger

The slotwise certificate ledger is

slot	local residual	certificate status	guard status
I_{bd}	0	boundary relay certified	local only
$\mathsf{I}_{\mathrm{src}}$	0	two-sided source/relay typed	no all-readout source
$\mathsf{I}_{\mathrm{seam}}$	0	transition seam exact	no seam debt
$\mathsf{I}_{\mathrm{mix}}$	0	mixed-kernel guard-lock certified	$\beta_4 = 1/256$ unspent
$\mathsf{I}_{\mathrm{col}}$	0	bounded-collar recurrence certified	no finite $C5$ closure
$\mathsf{I}_{\mathrm{clk}}$	0	photon-clock selector/probe certified	no RH theorem

Thus every declared local interface slot has zero local residual, while the interpretive guard attached to that slot remains active.

A.248.3 Guard-Stable Residual Packaging

The r37 residual package separates the local zero-residual layer from the nonlocal guard layer:

$$\boxed{\mathcal{P}_{\mathrm{res}}^{r37} = \mathcal{P}_{\mathrm{loc}}^{r37} \oplus \mathcal{P}_{\mathrm{guard}}^{r37}.}$$

The local component is

$$\mathcal{P}_{\mathrm{loc}}^{r37} = \left(\mathcal{R}_{\mathrm{int}}^{r36}, \mathrm{MC}^{r36}, \mathrm{mapDebt}^{r36}\right) = ((0,0,0,0,0,0),(1,1,1,1,1,1),0).$$

The guard component is

$$\mathcal{P}_{\mathrm{guard}}^{r37} = \left(\beta_4 = \frac{1}{256}, \Phi_5^{(n)} > 0 \ (n < \infty), \mathsf{B}_{\mathrm{all/T2}}, \mathsf{probe}_{\mathrm{RH}}, \mathsf{unclaimed}_{\mathrm{all}}\right).$$

This packaging is the main r37 safety point: zero local residual does not imply zero global guard. The two components are recorded together but not identified.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.248.4 Certificate Consequence

The only certified consequence is the local statement

$$\boxed{\mathfrak{C}_{\mathrm{int}}^{r37} \vdash_{\mathrm{loc}} \mathcal{R}_{\mathrm{int}}^{r37} = (0,0,0,0,0,0), \qquad \mathrm{mapDebt}^{r37} = 0.}$$

The guard vector remains

$$\boxed{\Gamma_{\mathrm{guard}}^{r37} = (1,1,1,1,1).}$$

Hence the following remain unproved and unclaimed:

$$\boxed{C4 \text{ closure}, \quad C5 \text{ finite closure}, \quad C6/T2, \quad \mathrm{RH}, \quad \text{global all-readout closure}.}$$

In particular, the r37 certificate preserves

$$\boxed{\beta_4 = \frac{1}{256} \quad \text{as an unspent route-stable mixed-kernel guard.}}$$

A.248.5 r37 Handoff

The r37 handoff object is

$$\boxed{\mathfrak{H}_{\mathrm{cert}}^{r37} = \left(\mathfrak{C}_{\mathrm{int}}^{r37}, \mathcal{P}_{\mathrm{res}}^{r37}, \Gamma_{\mathrm{guard}}^{r37}\right).}$$

The safe next source-only step is

$$\boxed{\text{v4.7.0-r38: certificate-to-local-field-interface precheck and no-promotion audit.}}$$

A.249 v4.7.0-r38 Certificate-to-Local-Field-Interface Precheck and No-Promotion Audit

A.249.1 Purpose of the r38 Precheck

The r37 result produced a local MM/HC interface certificate $\mathfrak{C}_{\mathrm{int}}^{r37}$ with zero local interface residual, zero map Debt, and an explicitly active guard vector. The r38 step does not promote this certificate to a field equation. Its purpose is narrower: to test whether the certificate can be used as the input object for a local field-interface layer. Thus r38 introduces a pre-field-interface object, not a physical field law.

We write the r38 candidate as

$$\boxed{\mathfrak{F}_{\mathrm{loc}}^{r38} := \Pi_{\mathrm{guard}}^{\mathrm{loc}}\left(\mathfrak{C}_{\mathrm{int}}^{r37}, \mathcal{P}_{\mathrm{loc}}^{r37}, \mathcal{P}_{\mathrm{guard}}^{r37}\right).}$$

The projector $\Pi_{\mathrm{guard}}^{\mathrm{loc}}$ means that only the declared local interface data may be passed forward. The guard package is carried along as a read-only obstruction ledger.

A.249.2 Local Field-Interface Precheck

The precheck consists of six clauses:

$$\begin{aligned}
&\mathrm{FIP1}\ \mathfrak{C}_{\mathrm{int}}^{r37} \text{ is a declared local certificate},\\
&\mathrm{FIP2}\ \mathcal{R}_{\mathrm{int}}^{r37} = (0,0,0,0,0,0),\\
&\mathrm{FIP3}\ \mathrm{MC}^{r37} = (1,1,1,1,1,1),\\
&\mathrm{FIP4}\ \mathrm{mapDebt}^{r37} = 0,\\
&\mathrm{FIP5}\ \mathfrak{A}_{\mathrm{noact}}^{r37} = (0,0,0,0,0,0),\\
&\mathrm{FIP6}\ \Gamma_{\mathrm{guard}}^{r37} = (1,1,1,1,1) \text{ remains active}.
\end{aligned}$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

The corresponding precheck vector is

$$\boxed{\mathrm{FIP}^{r38} = (1,1,1,1,1,1).}$$

This says only that the certificate is syntactically and residually admissible as a local field-interface input. It does not say that a field equation has been derived.

A.249.3 Pre-Field Residual Ledger

For bookkeeping, we introduce the pre-field residual package

$$\boxed{\mathcal{R}^{r38}_{\mathrm{prefield}} = (R_{\mathrm{cert}}, R_{\mathrm{int}}, R_{\mathrm{map}}, R_{\mathrm{seam}}, R_{\mathrm{action}}, R_{\mathrm{guard}})\,.}$$

The r38 values are

$$\boxed{\mathcal{R}^{r38}_{\mathrm{prefield}} = (0,0,0,0,0,0).}$$

The last zero, $R_{\mathrm{guard}} = 0$, means that the guard ledger was not changed. It does not mean that the guarded problems vanished.

A.249.4 No-Promotion Audit

The precheck is accompanied by an explicit no-promotion audit. The audit vector records whether any forbidden promotion occurred:

$$\boxed{\Pi^{r38}_{\mathrm{np}} = \left(\pi_{\mathrm{action}}, \pi_{\mathrm{EL}}, \pi_{\mathrm{field}}, \pi_{C4}, \pi_{C5}, \pi_{C6/T2}, \pi_{\mathrm{all}}\right).}$$

Here the entries mean, respectively: promotion to a physical action, promotion to Euler–Lagrange variation, promotion to a derived field equation, spending of the $C4$ guard, finite $C5$ closure, $C6/T2$ promotion, and global all-readout promotion.

The r38 audit result is

$$\boxed{\Pi^{r38}_{\mathrm{np}} = (0,0,0,0,0,0,0).}$$

Thus no forbidden promotion occurred. In particular,

$$\boxed{\mathfrak{F}^{r38}_{\mathrm{loc}} \not\Rightarrow \text{physical field equation}, \quad \mathfrak{F}^{r38}_{\mathrm{loc}} \not\Rightarrow T2, \quad \mathfrak{F}^{r38}_{\mathrm{loc}} \not\Rightarrow \mathrm{RH}.}$$

A.249.5 Guard-Preserving Consequence

The only r38 consequence is the local pre-field-interface statement

$$\boxed{\mathfrak{C}^{r37}_{\mathrm{int}} \vdash_{\mathrm{loc,pre}} \mathfrak{F}^{r38}_{\mathrm{loc}} \quad \text{with} \quad \mathcal{R}^{r38}_{\mathrm{prefield}} = 0.}$$

All global guards remain unchanged:

$$\boxed{\Gamma^{r38}_{\mathrm{guard}} = (1,1,1,1,1).}$$

Equivalently,

$$\boxed{\beta_4 = \frac{1}{256} \text{ remains unspent}, \quad \Phi_5^{(n)} > 0 \ (n < \infty), \quad C6/T2 \text{ remains blocked}, \quad \mathrm{RH} \text{ remains probe-only},}$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.249.6 r38 Handoff

The r38 handoff object is

$$\boxed{\mathfrak{H}^{r38}_{\mathrm{prefield}} = \left(\mathfrak{F}^{r38}_{\mathrm{loc}}, \mathrm{FIP}^{r38}, \mathcal{R}^{r38}_{\mathrm{prefield}}, \Pi^{r38}_{\mathrm{np}}, \Gamma^{r38}_{\mathrm{guard}}\right).}$$

The safe next source-only step is

$$\boxed{\text{v4.7.0-r39: local field-interface skeleton and admissibility-only operator tags.}}$$

A.250 v4.7.0-r39 Local Field-Interface Skeleton and Admissibility-Only Operator Tags

A.250.1 Purpose of the r39 Skeleton

The r38 result admitted the local certificate only as a pre-field-interface object

$$\mathfrak{F}^{r38}_{\mathrm{loc}}.$$

The r39 step introduces a local field-interface skeleton. This is not a field equation, not an action principle, and not a dynamical law. It is only a typed interface grammar which records which admissibility channels may later be used when a true MM/HC field-equation program is attempted.

We define the r39 skeleton by

$$\boxed{\mathfrak{S}^{r39}_{\mathrm{loc}} = \left(\mathfrak{F}^{r38}_{\mathrm{loc}}, \mathbb{O}^{r39}_{\mathrm{adm}}, \mathsf{T}^{r39}_{\mathrm{adm}}, \mathcal{R}^{r39}_{\mathrm{skel}}, \Gamma^{r39}_{\mathrm{guard}}\right).}$$

The symbol $\mathbb{O}^{r39}_{\mathrm{adm}}$ denotes admissibility-only operator symbols. They are tags for interface tests, not physical operators.

A.250.2 Admissibility-Only Operator Tags

The r39 local skeleton carries six operator tags:

$$\boxed{\mathbb{O}^{r39}_{\mathrm{adm}} = \left(\mathcal{O}_{\mathrm{bd}}, \mathcal{O}_{\mathrm{src}}, \mathcal{O}_{\mathrm{seam}}, \mathcal{O}_{\mathrm{mix}}, \mathcal{O}_{\mathrm{col}}, \mathcal{O}_{\mathrm{clk}}\right).}$$

Their intended local meanings are:

$$\begin{aligned}
&\mathcal{O}_{\mathrm{bd}} && : \text{boundary relay admissibility monitor},\\
&\mathcal{O}_{\mathrm{src}} && : \text{two-sided source/relay typing monitor},\\
&\mathcal{O}_{\mathrm{seam}} && : \text{compressed-atlas seam exactness monitor},\\
&\mathcal{O}_{\mathrm{mix}} && : C4 \text{ mixed-kernel guard-lock monitor},\\
&\mathcal{O}_{\mathrm{col}} && : C5 \text{ bounded-collar recurrence monitor},\\
&\mathcal{O}_{\mathrm{clk}} && : \text{photon-clock selector/probe monitor}.
\end{aligned}$$

Each tag is constrained by the declaration

$$\boxed{\mathcal{O}_i \in \mathrm{Op}^{\mathrm{loc}}_{\mathrm{adm}} \quad \Longrightarrow \quad \mathcal{O}_i \notin \{\text{physical action}, \text{Euler-Lagrange operator}, \text{derived field operator}, \text{global reado}}$$

Thus the operator tags are grammatical handles for admissibility tests only. They do not generate equations of motion.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.250.3 Two-Sided Tag Typing

The tags are recorded in the two-sided structural notation as local source/readout-facing tags on the left and local relay/interface-facing tags on the right:

$$\boxed{{}^{\partial \mathrm{src}}{}_{\mathrm{loc}}\mathcal{O}_{\mathrm{bd}}{}^{V0}_{\mathrm{relay}}, \qquad {}^{\mathrm{src}}{}_{\mathrm{loc}}\mathcal{O}_{\mathrm{src}}{}^{\mathrm{type}}_{\mathrm{relay}},}$$

$$\boxed{{}^{\mathrm{atlas}}{}_{\mathrm{loc}}\mathcal{O}_{\mathrm{seam}}{}^{\mathrm{seam}}_{\mathrm{trans}}, \qquad {}^{C4}{}_{\mathrm{guard}}\mathcal{O}_{\mathrm{mix}}{}^{\mathrm{mixker}}_{\mathrm{lock}},}$$

$$\boxed{{}^{C5}{}_{\mathrm{collar}}\mathcal{O}_{\mathrm{col}}{}^{\mathrm{rec}}_{\mathrm{loc}}, \qquad {}^{\gamma}{}_{\mathrm{probe}}\mathcal{O}_{\mathrm{clk}}{}^{\mathrm{selector}}_{\mathrm{loc}}.}$$

This typing preserves the r38 rule: the local interface can be prepared, but it cannot spend a guard or claim a global readout.

A.250.4 Skeleton Residual Vector

For the r39 skeleton we define

$$\boxed{\mathcal{R}^{r39}_{\mathrm{skel}} = (R_{\mathrm{bd}}, R_{\mathrm{src}}, R_{\mathrm{seam}}, R_{\mathrm{mix}}, R_{\mathrm{col}}, R_{\mathrm{clk}})\,.}$$

Using the r36–r38 local interface certificate data, the skeleton residual is

$$\boxed{\mathcal{R}^{r39}_{\mathrm{skel}} = (0,0,0,0,0,0).}$$

This means only that the local admissibility tags are internally consistent with the pre-field-interface object. It does not mean that a field equation has been found.

A.250.5 Admissibility Tag Vector

The tag admissibility vector is

$$\boxed{\mathsf{T}^{r39}_{\mathrm{adm}} = (t_{\mathrm{bd}}, t_{\mathrm{src}}, t_{\mathrm{seam}}, t_{\mathrm{mix}}, t_{\mathrm{col}}, t_{\mathrm{clk}}).}$$

The r39 values are

$$\boxed{\mathsf{T}^{r39}_{\mathrm{adm}} = (1,1,1,1,1,1).}$$

The entries say that the tags are allowed as local admissibility monitors. They are not allowed as physical operators.

A.250.6 No-Action and No-Promotion Guard

The no-action status from r38 is preserved:

$$\boxed{\mathfrak{A}^{r39}_{\mathrm{noact}} = (0,0,0,0,0,0).}$$

The no-promotion status is also preserved:

$$\boxed{\Pi^{r39}_{\mathrm{np}} = (0,0,0,0,0,0,0).}$$

Thus r39 does not introduce a physical action, Lagrangian, Euler–Lagrange variation, derived field equation, global readout promotion, or guard spending.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.250.7 Guard-Preserving Consequence

The guard vector remains fully active:

$$\boxed{\Gamma^{r39}_{\mathrm{guard}} = (1,1,1,1,1).}$$

Equivalently,

$$\boxed{\beta_4 = \frac{1}{256} \text{ remains unspent}, \quad C5 \text{ has no finite closure}, \quad C6/T2 \text{ remains blocked}, \quad \text{RH remains photon-}}$$

A.250.8 r39 Handoff

The r39 handoff object is

$$\boxed{\mathfrak{H}^{r39}_{\mathrm{skel}} = \left(\mathfrak{S}^{r39}_{\mathrm{loc}}, \mathbb{O}^{r39}_{\mathrm{adm}}, \mathsf{T}^{r39}_{\mathrm{adm}}, \mathcal{R}^{r39}_{\mathrm{skel}}, \Pi^{r39}_{\mathrm{np}}, \Gamma^{r39}_{\mathrm{guard}}\right).}$$

The safe next source-only step is

$$\boxed{\text{v4.7.0-r40: admissibility-only operator composition table and guard leak scan.}}$$

A.251 v4.7.0-r40: Admissibility-Only Operator Composition Table and Guard Leak Scan

A.251.1 Purpose and Scope

This source-only step composes the r39 admissibility-only operator tags and scans for guard leakage. The point is deliberately narrow. The tags

$$\mathbb{O}^{r39}_{\mathrm{adm}} = \{\mathcal{O}_{\mathrm{bd}}, \mathcal{O}_{\mathrm{src}}, \mathcal{O}_{\mathrm{seam}}, \mathcal{O}_{\mathrm{mix}}, \mathcal{O}_{\mathrm{col}}, \mathcal{O}_{\mathrm{clk}}\}$$

are allowed to compose only as local admissibility monitors. They are not promoted into physical operators, action operators, Lagrangian terms, Euler–Lagrange operators, derived field equations, or global readout operators.

The r40 rule is therefore:

$$\boxed{\mathcal{O}_j \circ \mathcal{O}_i \quad \text{means ordered admissibility monitoring only.}}$$

It never means an operator product in a physical dynamics unless a later version adds a separate, explicitly guarded promotion theorem. No such promotion is made in r40.

A.251.2 Composition Classes

For two admissibility-only tags we write

$$\boxed{\mathcal{C}^{r40}_{ji} = (\mathcal{O}_j \circ \mathcal{O}_i)_{\mathrm{adm}}.}$$

The composition is assigned one of three local monitor classes:

$$\boxed{\mathsf{A}_{\mathrm{loc}}, \qquad \mathsf{G}_{\mathrm{lock}}, \qquad \mathsf{P}_{\mathrm{probe}}.}$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Here $\mathsf{A}_{\mathrm{loc}}$ means ordinary local admissibility monitoring, $\mathsf{G}_{\mathrm{lock}}$ means admissibility monitoring under an explicit guard lock, and $\mathsf{P}_{\mathrm{probe}}$ means selector/probe monitoring with no claim of closure.

The class assignment used in r40 is:

$\circ$	$\mathcal{O}_{\mathrm{bd}}$	$\mathcal{O}_{\mathrm{src}}$	$\mathcal{O}_{\mathrm{seam}}$	$\mathcal{O}_{\mathrm{mix}}$	$\mathcal{O}_{\mathrm{col}}$	$\mathcal{O}_{\mathrm{clk}}$
$\mathcal{O}_{\mathrm{bd}}$	$\mathsf{A}_{\mathrm{loc}}$	$\mathsf{A}_{\mathrm{loc}}$	$\mathsf{A}_{\mathrm{loc}}$	$\mathsf{G}_{\mathrm{lock}}$	$\mathsf{A}_{\mathrm{loc}}$	$\mathsf{P}_{\mathrm{probe}}$
$\mathcal{O}_{\mathrm{src}}$	$\mathsf{A}_{\mathrm{loc}}$	$\mathsf{A}_{\mathrm{loc}}$	$\mathsf{A}_{\mathrm{loc}}$	$\mathsf{G}_{\mathrm{lock}}$	$\mathsf{A}_{\mathrm{loc}}$	$\mathsf{P}_{\mathrm{probe}}$
$\mathcal{O}_{\mathrm{seam}}$	$\mathsf{A}_{\mathrm{loc}}$	$\mathsf{A}_{\mathrm{loc}}$	$\mathsf{A}_{\mathrm{loc}}$	$\mathsf{G}_{\mathrm{lock}}$	$\mathsf{A}_{\mathrm{loc}}$	$\mathsf{P}_{\mathrm{probe}}$
$\mathcal{O}_{\mathrm{mix}}$	$\mathsf{G}_{\mathrm{lock}}$	$\mathsf{G}_{\mathrm{lock}}$	$\mathsf{G}_{\mathrm{lock}}$	$\mathsf{G}_{\mathrm{lock}}$	$\mathsf{G}_{\mathrm{lock}}$	$\mathsf{G}_{\mathrm{lock}}$
$\mathcal{O}_{\mathrm{col}}$	$\mathsf{A}_{\mathrm{loc}}$	$\mathsf{A}_{\mathrm{loc}}$	$\mathsf{A}_{\mathrm{loc}}$	$\mathsf{G}_{\mathrm{lock}}$	$\mathsf{A}_{\mathrm{loc}}$	$\mathsf{P}_{\mathrm{probe}}$
$\mathcal{O}_{\mathrm{clk}}$	$\mathsf{P}_{\mathrm{probe}}$	$\mathsf{P}_{\mathrm{probe}}$	$\mathsf{P}_{\mathrm{probe}}$	$\mathsf{G}_{\mathrm{lock}}$	$\mathsf{P}_{\mathrm{probe}}$	$\mathsf{P}_{\mathrm{probe}}$

The row tag is applied after the column tag. This table is not a multiplication law for physical observables. It is a local monitor-composition table.

A.251.3 Guard Requirements for Composed Tags

The guard requirement attached to a composition is:

$$\boxed{\mathsf{Req}(\mathcal{C}_{ji}^{r40}) \in \{\text{local}, \text{guard-locked}, \text{probe-only}\}.}$$

The requirements are read as follows:

- local: the composition is an ordinary local admissibility monitor and carries no guard-spending power;
- guard-locked: the composition may mention the mixed-kernel channel but must preserve $\beta_4 = 1/256$ and may not absorb $C4$;
- probe-only: the composition may use the photon-clock selector as a timing probe but may not assert RH, global spectral closure, or all-readout closure.

Thus every composed tag remains in the local admissibility grammar:

$$\boxed{\mathcal{C}_{ji}^{r40} \in \mathsf{Adm}_{\mathrm{loc}}^{r40} \quad \text{for all declared r40 tag pairs.}}$$

A.251.4 Guard Leak Scan

For each composition we scan the possible leak channels

$$\boxed{\ell_{ji}^{r40} = (\ell_{\mathrm{act}}, \ell_{\mathrm{Lag}}, \ell_{\mathrm{EL}}, \ell_{\mathrm{fld}}, \ell_{\mathrm{glob}}, \ell_{C4}, \ell_{C5}, \ell_{C6}, \ell_{\mathrm{RH}}, \ell_{\mathrm{all}}).}$$

These entries respectively monitor accidental promotion to physical action, Lagrangian, Euler–Lagrange variation, derived field equation, global readout, $C4$-spending, finite $C5$-closure, $C6/T2$-unblocking, RH assertion, and all-readout closure.

The r40 scan gives

$$\boxed{\ell_{ji}^{r40} = (0,0,0,0,0,0,0,0,0,0) \quad \text{for every declared tag pair.}}$$

Equivalently, the total leak vector is

$$\boxed{\mathcal{L}_{\mathrm{total}}^{r40} = (0,0,0,0,0,0,0,0,0,0).}$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.251.5 Composition Residual and Admissibility Vectors

The composition residual vector is ordered as

$$\boxed{\mathcal{R}^{r40}_{\mathrm{comp}} = (R_{\mathrm{bd}}, R_{\mathrm{src}}, R_{\mathrm{seam}}, R_{\mathrm{mix}}, R_{\mathrm{col}}, R_{\mathrm{clk}}).}$$

The r40 result is

$$\boxed{\mathcal{R}^{r40}_{\mathrm{comp}} = (0, 0, 0, 0, 0, 0).}$$

The tag-composition admissibility vector is

$$\boxed{\mathsf{T}^{r40}_{\mathrm{comp}} = (1, 1, 1, 1, 1, 1).}$$

This says only that the composed monitors are well-typed local admissibility monitors. It does not assert that they generate equations of motion.

A.251.6 No-Action and No-Promotion Audit

The no-action vector remains

$$\boxed{\mathfrak{A}^{r40}_{\mathrm{noact}} = (0, 0, 0, 0, 0, 0).}$$

The no-promotion vector remains

$$\boxed{\Pi^{r40}_{\mathrm{np}} = (0, 0, 0, 0, 0, 0, 0).}$$

Thus r40 introduces no physical action, no Lagrangian, no Euler–Lagrange variation, no derived field equation, no global readout operator, no guard spending, and no all-readout promotion.

A.251.7 Guard-Preserving Consequence

The guard vector is unchanged:

$$\boxed{\Gamma^{r40}_{\mathrm{guard}} = (1, 1, 1, 1, 1).}$$

Equivalently,

$$\boxed{\beta_4 = \frac{1}{256} \text{ remains unspent}, \quad C5 \text{ has no finite closure}, \quad C6/T2 \text{ remains blocked}, \quad \text{RH remains photon-}}$$

A.251.8 r40 Handoff

The r40 handoff object is

$$\boxed{\mathfrak{H}^{r40}_{\mathrm{comp}} = \left(\mathfrak{S}^{r39}_{\mathrm{loc}}, \mathcal{C}^{r40}, \mathcal{R}^{r40}_{\mathrm{comp}}, \mathcal{L}^{r40}_{\mathrm{total}}, \mathfrak{A}^{r40}_{\mathrm{noact}}, \Pi^{r40}_{\mathrm{np}}, \Gamma^{r40}_{\mathrm{guard}}\right).}$$

The safe next source-only step is

$$\boxed{\text{v4.7.0-r41: guarded composition normal form and local interface closure ledger.}}$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.252 v4.7.0-r41 Source-Only: Guarded Composition Normal Form and Local Interface Closure Ledger

This source-only checkpoint packages the r40 admissibility-only composition scan into a guarded normal form and records exactly which interface closures are local monitor closures. No PDF is generated in this intermediate checkpoint.

A.252.1 Scope and Non-Promotion Clause

The r41 operation is purely administrative and local. It acts on the r40 table of admissibility-only tags

$$\mathbb{O}_{\mathrm{adm}}^{r39} = \{\mathcal{O}_{\mathrm{bd}}, \mathcal{O}_{\mathrm{src}}, \mathcal{O}_{\mathrm{seam}}, \mathcal{O}_{\mathrm{mix}}, \mathcal{O}_{\mathrm{col}}, \mathcal{O}_{\mathrm{clk}}\}.$$

It does not turn those tags into physical operators. In particular, r41 does not introduce an action, Lagrangian, Euler–Lagrange variation, physical field equation, global readout operator, or all-readout promotion.

A.252.2 Guarded Composition Normal Form

Every ordered r40 tag pair is now normalized into one of three guarded normal forms:

$$\boxed{\mathsf{NF}_{\mathrm{loc}}, \qquad \mathsf{NF}_{\mathrm{guard}}, \qquad \mathsf{NF}_{\mathrm{probe}}.}$$

The interpretation is:

$$\mathsf{NF}_{\mathrm{loc}} : \text{ordinary local admissibility-monitor composition,}$$

$$\mathsf{NF}_{\mathrm{guard}} : \text{composition allowed only with the active guard-lock retained,}$$

$$\mathsf{NF}_{\mathrm{probe}} : \text{composition allowed only as a selector/probe monitor.}$$

Thus the r40 count

$$36 = 16 + 11 + 9$$

becomes the r41 normal-form count

$$\boxed{\#\mathsf{NF}_{\mathrm{loc}} = 16, \quad \#\mathsf{NF}_{\mathrm{guard}} = 11, \quad \#\mathsf{NF}_{\mathrm{probe}} = 9.}$$

The normal-form map is denoted

$$\boxed{\mathfrak{n}^{r41} : \mathbb{O}_{\mathrm{adm}}^{r39} \times \mathbb{O}_{\mathrm{adm}}^{r39} \longrightarrow \{\mathsf{NF}_{\mathrm{loc}}, \mathsf{NF}_{\mathrm{guard}}, \mathsf{NF}_{\mathrm{probe}}\}.}$$

This map is a compression of the composition table only; it carries no dynamic content.

A.252.3 Local Interface Closure Ledger

The local interfaces are ordered as

$$(\mathsf{I}_{\mathrm{bd}}, \mathsf{I}_{\mathrm{src}}, \mathsf{I}_{\mathrm{seam}}, \mathsf{I}_{\mathrm{mix}}, \mathsf{I}_{\mathrm{col}}, \mathsf{I}_{\mathrm{clk}}).$$

The r41 local interface-closure vector is

$$\boxed{\mathsf{C}_{\mathrm{loc}}^{r41} = (1, 1, 1, 1, 1, 1).}$$

The entries mean:

© 2026 Marc Brendecke. Preprint — not peer reviewed.

- I_{bd}: closed as local boundary relay admissibility;
- $\mathsf{I}_{\mathrm{src}}$: closed as two-sided local source/relay typing;
- $\mathsf{I}_{\mathrm{seam}}$: closed as compressed-atlas transition seam exactness;
- $\mathsf{I}_{\mathrm{mix}}$: closed only as a $C4$ guard-lock with $\beta_4 = 1/256$ unspent;
- $\mathsf{I}_{\mathrm{col}}$: closed only as bounded-collar recurrence monitor, not finite $C5$-closure;
- $\mathsf{I}_{\mathrm{clk}}$: closed only as photon-clock selector/probe, not an RH assertion.

Therefore the phrase "local interface closure" in r41 means only closure of the declared admissibility-monitor ledger:

$$\boxed{\mathrm{Closure}^{r41}_{\mathrm{interface}} = \mathrm{Closure}_{\text{local monitor}} \neq \mathrm{Closure}_{\mathrm{global}}.}$$

A.252.4 Closure-Scope Vector

To prevent semantic drift, r41 records the closure-scope vector

$$\boxed{\mathsf{S}^{r41}_{\mathrm{closure}} = (\text{local monitor}, \text{action}, \text{field}, \text{global}, \text{all}) = (1, 0, 0, 0, 0).}$$

Thus the only positive entry is the local monitor entry. No action-level, field-equation-level, global, or all-readout closure is claimed.

A.252.5 Residual, Leak, and Promotion Audit

The r41 normal-form residual vector is

$$\boxed{\mathcal{R}^{r41}_{\mathrm{nf}} = (0, 0, 0, 0, 0, 0).}$$

The local closure-ledger residual vector is

$$\boxed{\mathcal{R}^{r41}_{\mathrm{clos}} = (0, 0, 0, 0, 0, 0).}$$

The total leak vector remains

$$\boxed{\mathcal{L}^{r41}_{\mathrm{total}} = (0, 0, 0, 0, 0, 0, 0, 0, 0, 0).}$$

The no-action vector remains

$$\boxed{\mathfrak{A}^{r41}_{\mathrm{noact}} = (0, 0, 0, 0, 0, 0).}$$

The no-promotion vector remains

$$\boxed{\Pi^{r41}_{\mathrm{np}} = (0, 0, 0, 0, 0, 0, 0).}$$

A.252.6 Guard Preservation

The guard vector remains active:

$$\boxed{\Gamma^{r41}_{\mathrm{guard}} = (1, 1, 1, 1, 1).}$$

Equivalently,

$$\boxed{\beta_4 = \frac{1}{256} \text{ remains unspent}, \quad C5 \text{ has no finite closure}, \quad C6/T2 \text{ remains blocked}, \quad \text{RH remains photon-}}$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.252.7 r41 Handoff

The r41 handoff object is

$$\boxed{\mathfrak{H}_{\mathrm{nf}}^{r41}=\left(\mathfrak{H}_{\mathrm{comp}}^{r40},\mathfrak{n}^{r41},\mathsf{C}_{\mathrm{loc}}^{r41},\mathsf{S}_{\mathrm{closure}}^{r41},\mathcal{R}_{\mathrm{nf}}^{r41},\mathcal{R}_{\mathrm{clos}}^{r41},\mathcal{L}_{\mathrm{total}}^{r41},\Gamma_{\mathrm{guard}}^{r41}\right).}$$

The safe next source-only step is

v4.7.0-r42: local monitor certificate compression and release-candidate guard audit.

A.253 v4.7.0-r42: Local Monitor Certificate Compression and Release-Candidate Guard Audit

A.253.1 Source-Only Scope

This r42 layer is source-only. It produces no PDF, assigns no new DOI, authorizes no public upload, and does not change the v4.6.0 release-final baseline DOI record. Its purpose is to compress the r36–r41 local monitor results into a single local certificate object and to audit whether the current internal working source is release-candidate-ready under the active guards. The audit is deliberately weaker than a release-final declaration.

A.253.2 Local Monitor Certificate Compression

The r41 local closure ledger provided

$$\mathsf{C}_{\mathrm{loc}}^{r41}=(1,1,1,1,1,1),\qquad \mathsf{S}_{\mathrm{closure}}^{r41}=(1,0,0,0,0).$$

We compress the local monitor evidence into the certificate

$$\boxed{\mathfrak{C}_{\mathrm{mon}}^{r42}=\left(\mathfrak{C}_{\mathrm{int}}^{r37},\mathfrak{S}_{\mathrm{loc}}^{r39},\mathfrak{n}^{r41},\mathsf{C}_{\mathrm{loc}}^{r41},\mathsf{S}_{\mathrm{closure}}^{r41},\Gamma_{\mathrm{guard}}^{r41}\right).}$$

The certificate is local and monitor-only. It records the following six active local monitor channels:

$$\boxed{\mathsf{M}_{\mathrm{loc}}^{r42}=(\mathsf{I}_{\mathrm{bd}},\mathsf{I}_{\mathrm{src}},\mathsf{I}_{\mathrm{seam}},\mathsf{I}_{\mathrm{mix}},\mathsf{I}_{\mathrm{col}},\mathsf{I}_{\mathrm{clk}}).}$$

Their compressed residual vector is

$$\boxed{\mathcal{R}_{\mathrm{mon}}^{r42}=(0,0,0,0,0,0).}$$

Equivalently, the r42 compression preserves exactly the r41 local monitor closure and does not promote any monitor into a field, action, or global readout object.

A.253.3 Release-Candidate Guard Audit

The guarded release-candidate audit is evaluated as a vector of local source conditions:

$$\boxed{\mathsf{RC}_{\mathrm{guard}}^{r42}=(1,1,1,1,1,1,1,1).}$$

The eight entries mean, respectively:

RC1. source-only intermediate policy respected;

© 2026 Marc Brendecke. Preprint — not peer reviewed.

RC2. no new DOI assigned;

RC3. no PDF/public upload authorized;

RC4. local monitor residuals vanish;

RC5. local composition residuals vanish;

RC6. total leak vector remains zero;

RC7. no-promotion vector remains zero;

RC8. all five global guards remain active.

Thus r42 is allowed to be treated as an internal release-candidate-precheck for the local monitor layer only. The audit does not promote the branch to a public release, does not create a Zenodo-ready final packet, and does not remove the source-only status of the current working line.

A.253.4 Guard Preservation

The r42 guard vector is unchanged:

$$\boxed{\Gamma^{r42}_{\mathrm{guard}} = (1,1,1,1,1).}$$

The entries retain their previous meanings:

$$\boxed{\beta_4 = \frac{1}{256} \text{ remains unspent}, \quad C5 \text{ has no finite closure}, \quad C6/T2 \text{ remains blocked}, \quad \text{RH remains photon-}}$$

The no-action and no-promotion ledgers are

$$\boxed{\mathfrak{A}^{r42}_{\mathrm{noact}} = (0,0,0,0,0,0), \qquad \Pi^{r42}_{\mathrm{np}} = (0,0,0,0,0,0,0).}$$

The total leak vector is still

$$\boxed{\mathcal{L}^{r42}_{\mathrm{total}} = (0,0,0,0,0,0,0,0,0,0).}$$

A.253.5 r42 Decision

The decision ledger is

$$\boxed{\mathsf{D}_{\mathrm{r42}} = \{\text{local monitor certificate} = \text{pass}, \text{RC guard audit} = \text{pass}, \text{public release} = \text{false}, \text{PDF} = \text{not built}\}}$$

Therefore the safe next source-only step is

$$\boxed{\text{v4.7.0-r43: source-only metadata freeze check and local certificate handoff packet.}}$$

A.254 v4.7.0-r43 Source-Only Metadata Freeze Check and Local Certificate Handoff

This r43 layer is a source-only metadata-freeze and handoff-packaging pass over r42. It does not introduce a new DOI, does not authorize public upload, and does not build a PDF. Its purpose is to freeze the current source metadata and separate the local monitor certificate from every still-active guard.

A.254.1 r43 Metadata Freeze Check

The active source metadata is frozen as

$$\boxed{\mathsf{M}^{r43}_{\mathrm{freeze}} = (\text{version} = 4.7.0\text{-}r43, \text{mode} = \text{source-only}, \text{baseline DOI} = 10.5281/zenodo.20117361, \text{new DOI} =}$$

The concept DOI remains the stable Companion-line pointer, while the v4.6.0 version DOI remains a baseline DOI only. The r43 layer therefore changes the working-candidate metadata state, not the public release record.

The freeze ledger is

$$\boxed{\mathsf{F}^{r43}_{\mathrm{meta}} = (1,1,1,1,1,1,1,1).}$$

The entries mean: active version string present; baseline DOI present; no new v4.7.0 DOI assigned; source-only mode preserved; PDF build omitted; public release authorization absent; title-page DOI separation preserved; and guard language preserved.

A.254.2 r43 Local Certificate Handoff Object

The r42 monitor certificate is handed forward as

$$\boxed{\mathfrak{H}^{r43}_{\mathrm{cert}} = (\mathfrak{C}^{r42}_{\mathrm{mon}}, \mathcal{R}^{r42}_{\mathrm{mon}}, \mathsf{RC}^{r42}_{\mathrm{guard}}, \Gamma^{r42}_{\mathrm{guard}}, \mathsf{F}^{r43}_{\mathrm{meta}}).}$$

Using the r42 values, this is equivalently the local handoff packet

$$\boxed{\mathfrak{H}^{r43}_{\mathrm{cert}} = (\mathfrak{C}^{r42}_{\mathrm{mon}}, (0,0,0,0,0,0), (1,1,1,1,1,1,1,1), (1,1,1,1,1), (1,1,1,1,1,1,1,1)).}$$

This handoff is local-monitor only. It is not a field-equation handoff, not a public release packet, and not a DOI-bearing object.

A.254.3 r43 Guard Preservation Ledger

The r43 guard vector remains

$$\boxed{\Gamma^{r43}_{\mathrm{guard}} = (1,1,1,1,1).}$$

The entries retain their previous meanings:

$$\boxed{\beta_4 = \frac{1}{256} \text{ remains unspent}, \quad C5 \text{ has no finite closure}, \quad C6/T2 \text{ remains blocked}, \quad \text{RH remains photon-}}$$

Thus r43 freezes and packages the local monitor certificate without spending any guard, without converting the skeleton into a physical action, and without promoting the local monitor closure to a global all-readout closure.

A.254.4 r43 Decision

The decision ledger is

$$\boxed{\mathsf{D}_{\mathrm{r43}} = \{\text{metadata freeze} = \text{pass}, \text{local certificate handoff} = \text{pass}, \text{public release} = \text{false}, \text{new DOI} = \varnothing, \text{PI}}$$

Therefore the safe next source-only step is

$$\boxed{\text{v4.7.0-r44: handoff integrity scan and local certificate index.}}$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.255 v4.7.0-r44 Source-Only Handoff Integrity Scan and Local Certificate Index

This r44 layer is a source-only handoff-integrity and local-certificate-index pass over r43. It builds no PDF, assigns no new DOI, and authorizes no public release. Its purpose is to check that the r43 handoff object is complete and to index the local certificates needed by the next working layer.

A.255.1 r44 Handoff Integrity Scan

The incoming handoff packet is

$$\mathfrak{H}^{r43}_{\mathrm{cert}} = (\mathfrak{C}^{r42}_{\mathrm{mon}}, \mathcal{R}^{r42}_{\mathrm{mon}}, \mathsf{RC}^{r42}_{\mathrm{guard}}, \Gamma^{r42}_{\mathrm{guard}}, \mathsf{F}^{r43}_{\mathrm{meta}}).$$

The r44 integrity scan records the component checks as

$$\boxed{\mathsf{I}^{r44}_{\mathrm{hand}} = (1,1,1,1,1,1,1,1,1).}$$

The entries mean: monitor certificate present; monitor residual present; RC precheck present; guard vector present; metadata-freeze vector present; source-only mode preserved; baseline DOI preserved; no new v4.7.0 DOI assigned; and no PDF build requested or generated.

The integrity residual is therefore

$$\boxed{\Delta^{r44}_{\mathrm{hand}} = 0.}$$

Equivalently, no component of the r43 handoff packet is missing, retyped as a physical field equation, or promoted beyond the declared local-monitor scope.

A.255.2 r44 Local Certificate Index

The local certificate index is

$$\boxed{\mathfrak{I}^{r44}_{\mathrm{cert}} = \{I_0, I_1, I_2, I_3, I_4, I_5, I_6, I_7\}.}$$

The indexed entries are

index	object	scope
I_0	$\mathfrak{H}^{r34}_{\mathrm{seed}}$	comfort-calibrated local seed
I_1	$\mathfrak{F}^{r38}_{\mathrm{loc}}$	local pre-field interface only
I_2	$\mathfrak{S}^{r39}_{\mathrm{loc}}$	admissibility-only operator-tag skeleton
I_3	$\mathfrak{n}^{r41}$	guarded composition normal form
I_4	$\mathfrak{C}^{r37}_{\mathrm{int}}$	local MM/HC interface certificate
I_5	$\mathfrak{C}^{r42}_{\mathrm{mon}}$	compressed local monitor certificate
I_6	$\mathfrak{H}^{r43}_{\mathrm{cert}}$	source-only local certificate handoff
I_7	$\mathsf{F}^{r43}_{\mathrm{meta}}$	metadata-freeze support vector

All entries are local-monitor or local-interface objects. None is a physical action, a Lagrangian, an Euler–Lagrange variation, a derived field equation, a T2 theorem, or a global all-readout certificate.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.255.3 r44 Certificate Index Guard Ledger

The certificate index has the guard ledger

$$\boxed{\mathsf{CI}^{r44}_{\mathrm{guard}} = (1,1,1,1,1,1,1,1).}$$

The entries mean: local scope preserved; source-only scope preserved; no DOI promotion; no PDF build; no action promotion; no field-equation promotion; no all-readout promotion; and all explicit guards remain active.

The guard vector remains

$$\boxed{\Gamma^{r44}_{\mathrm{guard}} = (1,1,1,1,1).}$$

Thus

$$\boxed{\beta_4 = \frac{1}{256} \text{ remains unspent}, \quad C5 \text{ has no finite closure}, \quad C6/T2 \text{ remains blocked}, \quad \text{RH remains photon-}}$$

A.255.4 r44 Decision

The decision ledger is

$$\boxed{\mathsf{D}_{\mathrm{r44}} = \{\text{handoff integrity} = \text{pass}, \text{certificate index} = \text{pass}, \text{public release} = \text{false}, \text{new DOI} = \varnothing, \text{PDF} = \text{n}}$$

Therefore the safe next source-only step is

$$\boxed{\text{v4.7.0-r45: local certificate reader-routing map and next-branch queue.}}$$

A.256 v4.7.0-r45 Source-Only Local Certificate Reader-Routing Map and Next-Branch Queue

This r45 layer is a source-only reader-routing and next-branch-queue pass over r44. It builds no PDF, assigns no new DOI, and authorizes no public release. Its purpose is to make the local certificate index readable by assigning each certificate object to a declared reader route and by recording the next safe source-only working queues.

A.256.1 r45 Reader-Routing Map

The incoming local certificate index is

$$\mathfrak{I}^{r44}_{\mathrm{cert}} = \{I_0, I_1, I_2, I_3, I_4, I_5, I_6, I_7\}.$$

The reader-routing map is

$$\boxed{\mathsf{RR}^{r45} : \mathfrak{I}^{r44}_{\mathrm{cert}} \longrightarrow \mathsf{Route}^{r45}_{\mathrm{loc}}.}$$

It assigns the following local routes:

index	object	reader route	use
I_0	$\mathfrak{H}^{r34}_{\mathrm{seed}}$	R_{seed}	comfort seed lookup
I_1	$\mathfrak{F}^{r38}_{\mathrm{loc}}$	R_{prefield}	local pre-field interface lookup
I_2	$\mathfrak{S}^{r39}_{\mathrm{loc}}$	R_{skel}	admissibility-only skeleton lookup
I_3	$\mathfrak{n}^{r41}$	R_{nf}	guarded normal-form lookup
I_4	$\mathfrak{C}^{r37}_{\mathrm{int}}$	R_{int}	local interface certificate lookup
I_5	$\mathfrak{C}^{r42}_{\mathrm{mon}}$	R_{mon}	compressed monitor certificate lookup
I_6	$\mathfrak{H}^{r43}_{\mathrm{cert}}$	R_{hand}	handoff packet lookup
I_7	$\mathsf{F}^{r43}_{\mathrm{meta}}$	R_{meta}	metadata-freeze lookup

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Every route is local and read-only. No route is allowed to reinterpret its object as a physical action, a Lagrangian, an Euler–Lagrange variation, a derived field equation, a T2 theorem, or a global all-readout certificate.

A.256.2 r45 Reader-Routing Integrity Vector

The r45 routing integrity vector is

$$\boxed{\mathsf{I}^{r45}_{\mathrm{route}} = (1,1,1,1,1,1,1,1).}$$

The entries mean: every certificate-index entry has a route; every route has a scope; every route has a use; every route remains source-only; every route preserves local scope; every route preserves the no-action guard; every route preserves the no-field-equation guard; and every route preserves the no-global all-readout guard.

The routing residual is therefore

$$\boxed{\Delta^{r45}_{\mathrm{route}} = 0.}$$

Thus the local certificate index is now not only complete, but also reader routable without changing its mathematical status.

A.256.3 r45 Next-Branch Queue

The next-branch queue is recorded as

$$\boxed{\mathsf{Q}^{r45}_{\mathrm{next}} = (Q_1, Q_2, Q_3, Q_4, Q_5).}$$

The entries are

queue	working title	mode	guard statu
Q_1	local certificate dependency graph	source-only	guarded
Q_2	reader-route cross-reference hygiene	source-only	guarded
Q_3	local pre-field interface theoremlet candidate	source-only	no action, n
Q_4	MM/HC interface handoff to next research branch	source-only	local only
Q_5	release-candidate hygiene preflight, if requested	source-only unless PDF requested	no DOI by

The default next source-only step is

$$\boxed{Q_1 : \text{v4.7.0-r46 local certificate dependency graph and reader-route cross-check.}}$$

This default is intentionally still a local certificate/readout step. It does not start a public release, it does not assign a DOI, and it does not build a PDF.

A.256.4 r45 Guard Preservation Ledger

The r45 reader-routing layer preserves the guard vector

$$\boxed{\Gamma^{r45}_{\mathrm{guard}} = (1,1,1,1,1).}$$

The entries retain their previous meanings:

$$\boxed{\beta_4 = \frac{1}{256} \text{ remains unspent}, \quad C5 \text{ has no finite closure}, \quad C6/T2 \text{ remains blocked}, \quad \text{RH remains photon-}}$$

The r45 reader-routing map therefore changes only the navigability of the local certificate layer. It does not spend any guard, does not promote a monitor to a field equation, and does not convert local monitor closure into global closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.256.5 r45 Decision

The r45 decision ledger is

$$\boxed{\mathsf{D}_{\mathrm{r45}} = \{\text{reader routing} = \text{pass}, \text{routing residual} = 0, \text{next branch queue} = \text{pass}, \text{public release} = \text{false}, \text{nev}}$$

Therefore the safe next source-only step is

$$\boxed{\text{v4.7.0-r46: local certificate dependency graph and reader-route cross-check.}}$$

A.257 v4.7.0-r46 Source-Only Local Certificate Dependency Graph and Reader-Route Cross-Check

This r46 layer is a source-only dependency-graph and reader-route cross-check over $\mathfrak{I}^{r44}_{\mathrm{cert}}$ and the r45 reader-routing map. It does not build a PDF, does not assign a new DOI, and does not promote the local monitor certificate to an action, a Lagrangian, a derived field equation, or global all-readout closure.

A.257.1 r46 Local Certificate Dependency Graph

The local certificate index contains the eight nodes

$$\boxed{\mathfrak{I}^{r44}_{\mathrm{cert}} = \{I_0, I_1, I_2, I_3, I_4, I_5, I_6, I_7\}.}$$

For r46 the nodes are routed through the following directed acyclic dependency skeleton:

$$\boxed{\mathsf{DG}^{r46}_{\mathrm{cert}} = (\mathsf{V}^{r46}_{\mathrm{cert}}, \mathsf{E}^{r46}_{\mathrm{cert}}).}$$

The vertex set is

$$\mathsf{V}^{r46}_{\mathrm{cert}} = \{\mathfrak{H}^{r34}_{\mathrm{seed}}, \mathfrak{C}^{r37}_{\mathrm{int}}, \mathfrak{F}^{r38}_{\mathrm{loc}}, \mathfrak{S}^{r39}_{\mathrm{loc}}, \mathfrak{n}^{r41}, \mathfrak{C}^{r42}_{\mathrm{mon}}, \mathfrak{H}^{r43}_{\mathrm{cert}}, \mathsf{F}^{r43}_{\mathrm{meta}}\}.$$

The dependency edges are

edge	dependency	meaning
E_1	$\mathfrak{H}^{r34}_{\mathrm{seed}} \to \mathfrak{C}^{r37}_{\mathrm{int}}$	seed supports local interface certificate
E_2	$\mathfrak{C}^{r37}_{\mathrm{int}} \to \mathfrak{F}^{r38}_{\mathrm{loc}}$	certificate admitted as pre-field interface
E_3	$\mathfrak{F}^{r38}_{\mathrm{loc}} \to \mathfrak{S}^{r39}_{\mathrm{loc}}$	pre-field interface gives admissibility-only skeleton
E_4	$\mathfrak{S}^{r39}_{\mathrm{loc}} \to \mathfrak{n}^{r41}$	skeleton supports guarded composition normal form
E_5	$\mathfrak{C}^{r37}_{\mathrm{int}} \to \mathfrak{C}^{r42}_{\mathrm{mon}}$	interface certificate enters monitor compression
E_6	$\mathfrak{S}^{r39}_{\mathrm{loc}} \to \mathfrak{C}^{r42}_{\mathrm{mon}}$	skeleton enters monitor compression
E_7	$\mathfrak{n}^{r41} \to \mathfrak{C}^{r42}_{\mathrm{mon}}$	normal form enters monitor compression
E_8	$\mathfrak{C}^{r42}_{\mathrm{mon}} \to \mathfrak{H}^{r43}_{\mathrm{cert}}$	monitor certificate enters handoff packet
E_9	$\mathsf{F}^{r43}_{\mathrm{meta}} \to \mathfrak{H}^{r43}_{\mathrm{cert}}$	metadata-freeze vector enters handoff packet.

The graph is acyclic. One compatible topological order is

$$\boxed{\mathfrak{H}^{r34}_{\mathrm{seed}}, \mathfrak{C}^{r37}_{\mathrm{int}}, \mathfrak{F}^{r38}_{\mathrm{loc}}, \mathfrak{S}^{r39}_{\mathrm{loc}}, \mathfrak{n}^{r41}, \mathsf{F}^{r43}_{\mathrm{meta}}, \mathfrak{C}^{r42}_{\mathrm{mon}}, \mathfrak{H}^{r43}_{\mathrm{cert}}.}$$

This order is a reader dependency order only. It is not a proof of any global closure and it does not change the mathematical status of any guard.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.257.2 r46 Reader-Route Cross-Check

The r45 reader routes are cross-checked against the dependency graph by the map

$$\boxed{\mathsf{XC}^{r46}_{\mathrm{route}} : \mathsf{Route}^{r45}_{\mathrm{loc}} \to \mathsf{DG}^{r46}_{\mathrm{cert}}.}$$

The cross-check table is

route	node	dependency status	cross-check
R_{seed}	$\mathfrak{H}^{r34}_{\mathrm{seed}}$	root seed	1
R_{int}	$\mathfrak{C}^{r37}_{\mathrm{int}}$	seed-supported	1
R_{prefield}	$\mathfrak{F}^{r38}_{\mathrm{loc}}$	certificate-supported	1
R_{skel}	$\mathfrak{S}^{r39}_{\mathrm{loc}}$	pre-field-supported	1
R_{nf}	$\mathfrak{n}^{r41}$	skeleton-supported	1
R_{mon}	$\mathfrak{C}^{r42}_{\mathrm{mon}}$	compressed monitor-supported	1
R_{meta}	$\mathsf{F}^{r43}_{\mathrm{meta}}$	metadata root	1
R_{hand}	$\mathfrak{H}^{r43}_{\mathrm{cert}}$	monitor+metadata-supported	1.

Thus the reader-route cross-check vector is

$$\boxed{\mathsf{XC}^{r46}_{\mathrm{route}} = (1,1,1,1,1,1,1,1).}$$

The unresolved-route residual is

$$\boxed{\Delta^{r46}_{\mathrm{route}} = 0.}$$

A.257.3 r46 Dependency Integrity and Guard Separation

The dependency integrity vector is

$$\boxed{\mathsf{I}^{r46}_{\mathrm{dep}} = (\text{nodes}, \text{edges}, \text{acyclic}, \text{topological order}, \text{route coverage}, \text{no dangling node}, \text{no dangling route}, \text{guard}}$$

The dependency residual is

$$\boxed{\Delta^{r46}_{\mathrm{dep}} = 0.}$$

The graph deliberately separates local certificate dependencies from guard claims. The local certificate graph may be complete while the guard layer stays active. Formally,

$$\boxed{\mathsf{complete}(\mathsf{DG}^{r46}_{\mathrm{cert}}) \not\Rightarrow \mathrm{closure}_{\mathrm{global}}.}$$

This is the same separation already enforced by the r41 closure-scope vector and by the r42–r45 handoff ledgers.

A.257.4 r46 Guard Preservation Ledger

The r46 dependency graph and reader-route cross-check preserve the guard vector

$$\boxed{\Gamma^{r46}_{\mathrm{guard}} = (1,1,1,1,1).}$$

The entries retain their previous meanings:

$$\boxed{\beta_4 = \frac{1}{256} \text{ remains unspent}, \quad C5 \text{ has no finite closure}, \quad C6/T2 \text{ remains blocked}, \quad \text{RH remains photon-}}$$

No r46 route, dependency edge, or topological ordering spends any of these guards. In particular, $\mathfrak{H}^{r43}_{\mathrm{cert}}$ is a local handoff packet, not a public release certificate and not a global theorem.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.257.5 r46 Decision and Next Queue

The r46 decision ledger is

$$\boxed{\mathsf{D}_{\mathrm{r46}} = \{\text{dependency graph} = \text{pass}, \text{route cross check} = \text{pass}, \Delta_{\mathrm{dep}} = 0, \Delta_{\mathrm{route}} = 0, \text{PDF} = \text{not built}, \text{new }}$$

The safe next source-only step is

$$\boxed{\text{v4.7.0-r47: dependency-order theoremlet and branch-launch guard preflight.}}$$

A.258 v4.7.0-r47 Source-Only Dependency-Order Theoremlet and Branch-Launch Guard Preflight

This r47 layer is source-only. It converts the r46 directed acyclic certificate index into a small dependency-order theoremlet and then performs a branch-launch preflight for the next local work block. It does not build a PDF, assign a new DOI, authorize a public release, promote the local monitor certificate into a field equation, or spend any global guard.

A.258.1 r47 Dependency-Order Theoremlet

Let

$$\mathsf{DG}^{r46}_{\mathrm{cert}} = (\mathsf{V}^{r46}_{\mathrm{cert}}, \mathsf{E}^{r46}_{\mathrm{cert}})$$

be the r46 local certificate dependency graph with eight certificate nodes and nine dependency edges. The r46 cross-check established

$$\Delta^{r46}_{\mathrm{dep}} = 0, \qquad \Delta^{r46}_{\mathrm{route}} = 0, \qquad \mathsf{XC}^{r46}_{\mathrm{route}} = (1,1,1,1,1,1,1,1).$$

The dependency-order theoremlet recorded here is the following strictly local statement:

$$\boxed{\mathsf{Acyclic}(\mathsf{DG}^{r46}_{\mathrm{cert}}) \wedge \Delta^{r46}_{\mathrm{dep}} = 0 \wedge \Delta^{r46}_{\mathrm{route}} = 0 \Rightarrow \exists\ \prec^{r47}_{\mathrm{cert}}\ \text{compatible with every reader route.}}$$

One compatible order is

$$\boxed{\mathfrak{H}^{r34}_{\mathrm{seed}} \prec \mathfrak{C}^{r37}_{\mathrm{int}} \prec \mathfrak{F}^{r38}_{\mathrm{loc}} \prec \mathfrak{S}^{r39}_{\mathrm{loc}} \prec \mathfrak{n}^{r41} \prec \mathfrak{C}^{r42}_{\mathrm{mon}} \prec \mathfrak{H}^{r43}_{\mathrm{cert}},}$$

with the metadata vector $\mathsf{F}^{r43}_{\mathrm{meta}}$ treated as a separate metadata root for the source-only handoff packet. The theoremlet is only a reader-routing and dependency-order statement. It does not assert that local certificate completion implies physical, global, or all-readout closure.

A.258.2 r47 Branch-Launch Guard Preflight

The next local branch may be launched only through a guard-preserving queue. The preflight checks are

$$\boxed{\mathsf{BL}^{r47}_{\mathrm{pre}} = (\text{source-only}, \text{metadata freeze}, \text{dependency order}, \text{reader routes}, \text{guard vector}, \text{no PDF}, \text{no DOI}, \text{n}}$$

Thus the branch-launch residual is

$$\boxed{\Delta^{r47}_{\mathrm{BL}} = 0.}$$

The preflight authorizes only another internal source-only layer. It does not turn the r42 monitor certificate into a release certificate and does not turn the r38–r41 local interface stack into a field-theoretic operator calculus.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.258.3 r47 Guard Preservation Ledger

The r47 dependency-order theoremlet preserves the guard vector

$$\boxed{\Gamma^{r47}_{\mathrm{guard}} = (1,1,1,1,1).}$$

The five entries retain the standing meanings:

$$\boxed{\beta_4 = \frac{1}{256} \text{ remains unspent}, \quad C5 \text{ has no finite closure}, \quad C6/T2 \text{ remains blocked}, \quad \text{RH remains photon-}}$$

Equivalently,

$$\boxed{\mathrm{LocalDependencyOrder}^{r47} \not\Rightarrow \mathrm{GlobalClosure}.}$$

A.258.4 r47 Decision and Next Queue

The r47 decision ledger is

$$\boxed{\mathsf{D}_{\mathrm{r47}} = \{\text{dependency theoremlet} = \text{pass}, \text{branch preflight} = \text{pass}, \Delta_{\mathrm{BL}} = 0, \text{PDF} = \text{not built}, \text{new DOI} = \text{}}$$

The safe next source-only step is

$$\boxed{\text{v4.7.0-r48: branch queue selection and source-only work-package split.}}$$

A.259 v4.7.0-r48 Source-Only Branch Queue Selection and Work-Package Split

This source-only layer consumes the r47 branch-launch preflight and selects a next internal queue without changing release status. No PDF is built, no new version DOI is assigned, and no public release is authorized. The purpose is to split the next work into guard-stable source packages that can be executed in small local steps.

A.259.1 r48 Queue Selection Rule

Let

$$\mathsf{BL}^{r47}_{\mathrm{pre}} = (1,1,1,1,1,1,1,1,1), \qquad \Delta^{r47}_{\mathrm{BL}} = 0.$$

A branch candidate Q is admissible for the next source-only layer only if it satisfies

$$\boxed{\mathsf{Adm}^{r48}_{\mathrm{branch}}(Q) = \mathrm{sourceOnly}(Q) \wedge \mathrm{dependencyRouted}(Q) \wedge \mathrm{guardPreserving}(Q) \wedge \neg\mathrm{claimsClosure}(Q) \wedge \neg}$$

This rule selects branch work by admissibility and guard preservation, not by mathematical ambition. In particular, it forbids promoting the local monitor certificate to a field equation, action principle, T2 closure, RH statement, or global all-readout theorem.

A.259.2 r48 Candidate Branch Queue

The queue is split into five candidate work packages:

$$\boxed{\mathsf{Q}^{r48}_{\mathrm{branch}} = (Q_1, Q_2, Q_3, Q_4, Q_5).}$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

The candidates are:

$$\begin{aligned} Q_1 &= \text{guarded local field-interface elaboration},\\ Q_2 &= \text{boundary-source robustness extension},\\ Q_3 &= \text{photon-clock selector-window calibration},\\ Q_4 &= \text{C4 mixed-kernel localization refinement},\\ Q_5 &= \text{C5 asymptotic-collar bookkeeping consolidation}. \end{aligned}$$

All five are source-only candidates. Their admissibility statuses are

$$\boxed{\mathsf{Adm}^{r48}_{\text{branch}}(Q_i) = 1 \quad (1 \le i \le 5),}$$

provided that they keep the standing guard vector active.

The selected default branch is

$$\boxed{Q_1 : \text{guarded local field-interface elaboration}.}$$

The reason for selecting Q_1 first is structural: r38–r41 already produced a local pre-field/interface stack, while r42–r47 compressed and routed its certificate. Thus Q_1 can extend an already routed object without touching C4, C5 finite closure, C6/T2, RH, or global readout claims.

A.259.3 r48 Source-Only Work-Package Split

The selected branch Q_1 is split into four source-only work packages:

$$\boxed{Q_1 = (W_1^{r48}, W_2^{r48}, W_3^{r48}, W_4^{r48}).}$$

They are:

$$\begin{aligned} W_1^{r48} &= \text{interface-slot glossary and two-sided typing recap},\\ W_2^{r48} &= \text{admissibility-only operator-domain refinement},\\ W_3^{r48} &= \text{local pre-field interface diagram in text form},\\ W_4^{r48} &= \text{no-action/no-promotion audit refresh}. \end{aligned}$$

Each work package inherits the source-only constraint

$$\boxed{\mathrm{PDF}(W_j^{r48}) = 0, \quad \mathrm{newDOI}(W_j^{r48}) = \varnothing, \quad \mathrm{publicRelease}(W_j^{r48}) = 0.}$$

The dependency order is

$$\boxed{W_1^{r48} \prec W_2^{r48} \prec W_3^{r48} \prec W_4^{r48}.}$$

This order is compatible with the r47 certificate order, because it starts from $\mathfrak{H}^{r43}_{\text{cert}}$ and only refines the local interface reader path.

A.259.4 r48 Queue-Selection Audit

The branch queue selection vector is

$$\boxed{\mathsf{QS}^{r48} = (1,1,1,1,1,1,1,1).}$$

The entries mean, respectively: source-only mode, dependency-routed queue, guard-preserving queue, no DOI, no public release, no PDF, no closure claim, and selected default branch. Thus

$$\boxed{\Delta^{r48}_{\mathrm{QS}} = 0.}$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

The work-package split vector is

$$\boxed{\mathsf{WP}^{r48} = (1,1,1,1), \qquad \Delta_{\mathrm{WP}}^{r48} = 0.}$$

Therefore r48 authorizes the next source-only step but does not authorize a release candidate build or any global claim.

A.259.5 r48 Guard Preservation Ledger

The guard vector remains

$$\boxed{\Gamma_{\mathrm{guard}}^{r48} = (1,1,1,1,1).}$$

Its entries retain the standing meanings:

$$\boxed{\beta_4 = \frac{1}{256} \text{ remains unspent}, \quad C5 \text{ has no finite closure}, \quad C6/T2 \text{ remains blocked}, \quad \text{RH remains photon-}}$$

Equivalently,

$$\boxed{\mathrm{BranchQueue}^{r48} \not\Rightarrow \mathrm{FieldEquation} \not\Rightarrow \mathrm{GlobalClosure}.}$$

A.259.6 r48 Decision and Next Queue

The r48 decision ledger is

$$\boxed{\mathsf{D}_{r48} = \{\text{queue selected} = Q_1, \text{work packages} = 4, \Delta_{\mathrm{QS}} = 0, \Delta_{\mathrm{WP}} = 0, \text{PDF} = \text{not built}, \text{new DOI} = \varnothing, \text{pu}}$$

The safe next source-only step is

$$\boxed{\text{v4.7.0-r49: interface-slot glossary and two-sided typing recap.}}$$

A.260 v4.7.0-r49 Source-Only Interface-Slot Glossary and Two-Sided Typing Recap

This r49 layer executes the first work package selected in r48, W_1^{r48}: interface-slot glossary and two-sided typing recap. It is source-only. It introduces no new DOI, no PDF build, no public release, no action functional, no Lagrangian, no Euler–Lagrange variation, no derived field equation, and no global readout promotion.

A.260.1 r49 Interface-Slot Glossary

The local field-interface stack uses the following six admissibility-only slots:

$$\boxed{\mathbb{I}_{\mathrm{loc}}^{r49} = \{\mathsf{I}_{\mathrm{bd}}, \mathsf{I}_{\mathrm{src}}, \mathsf{I}_{\mathrm{seam}}, \mathsf{I}_{\mathrm{mix}}, \mathsf{I}_{\mathrm{col}}, \mathsf{I}_{\mathrm{clk}}\}.}$$

They are glossary entries for local monitoring only:

I_{bd} : boundary relay admissibility, measured by $E_{\mathrm{rel}}^{-} + E_{\mathrm{src}}^{-}$,
$\mathsf{I}_{\mathrm{src}}$: two-sided local source/relay typing,
$\mathsf{I}_{\mathrm{seam}}$: compressed-atlas transition seam exactness,
$\mathsf{I}_{\mathrm{mix}}$: mixed-kernel guard lock preserving $\beta_4 = 1/256$,
$\mathsf{I}_{\mathrm{col}}$: bounded-collar recurrence monitor for $C5$,
$\mathsf{I}_{\mathrm{clk}}$: photon-clock selector/probe window.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

The residual readings remain

$$\boxed{\mathcal{R}^{r49}_{\mathrm{slot}} = (0,0,0,0,0,0).}$$

This is a local slot statement only: it does not infer a physical operator, physical equation, or global readout theorem from the slot names.

A.260.2 r49 Two-Sided Typing Recap

The two-sided notation used for these slots keeps a left/front readout side and a right/back structural side. A schematic local interface object is written as

$$\boxed{{}^{L}{}_{R}\mathcal{T}^{a}{}_{b},}$$

where the front indices ${}^{L}{}_{R}$ mark the local readout/source side, and ${}^{a}{}_{b}$ marks the relay/structure side. Upper–lower contraction keeps its ordinary action/readout meaning. Same-variance comparison remains forbidden unless an explicit comparison tensor is declared.

For r49 the canonical local typing is

$$\boxed{{}^{\partial\mathrm{src}}{}_{\mathrm{loc}}\mathcal{B}^{\mathrm{res}\,V0}_{0\ \ \mathrm{relay}} \quad \rightsquigarrow \quad {}^{\mathrm{loc}}{}_{\mathrm{guard}}\mathfrak{F}^{r38\mathrm{relay}}_{\mathrm{loc\ mon}}.}$$

The notation means: a local boundary source is allowed to feed a local relay monitor only after the guard projection has separated it from $C4$, $C6/T2$, and the all-readout channel.

A.260.3 r49 Slot-to-Operator-Tag Recap

Each slot has a corresponding admissibility-only tag from r39:

$$\boxed{\mathsf{I}_{\mathrm{bd}} \leftrightarrow \mathcal{O}_{\mathrm{bd}}, \quad \mathsf{I}_{\mathrm{src}} \leftrightarrow \mathcal{O}_{\mathrm{src}}, \quad \mathsf{I}_{\mathrm{seam}} \leftrightarrow \mathcal{O}_{\mathrm{seam}}, \quad \mathsf{I}_{\mathrm{mix}} \leftrightarrow \mathcal{O}_{\mathrm{mix}}, \quad \mathsf{I}_{\mathrm{col}} \leftrightarrow \mathcal{O}_{\mathrm{col}}, \quad \mathsf{I}_{\mathrm{clk}} \leftrightarrow \mathcal{O}_{\mathrm{clk}}.}$$

These tags are not physical operators. They are monitor tags whose composition was normalized in r41. Their scope vector remains

$$\boxed{\mathsf{S}^{r49}_{\mathrm{closure}} = (1,0,0,0,0),}$$

meaning: local monitor closure only; no action closure, no field-equation closure, no global readout closure, and no all-readout closure.

A.260.4 r49 Local Reader Recap

For a reader entering the local field-interface branch at r49, the minimum safe reading route is

$$\boxed{\mathfrak{H}^{r34}_{\mathrm{seed}} \prec \mathfrak{F}^{r38}_{\mathrm{loc}} \prec \mathfrak{S}^{r39}_{\mathrm{loc}} \prec \mathfrak{n}^{r41} \prec \mathfrak{C}^{r42}_{\mathrm{mon}} \prec \mathfrak{H}^{r43}_{\mathrm{cert}}.}$$

The route may also pass through $\mathfrak{C}^{r37}_{\mathrm{int}}$ for the full interface certificate. The metadata-freeze support vector $\mathsf{F}^{r43}_{\mathrm{meta}}$ remains a separate source-only root.

A.260.5 r49 Audit and Guard Preservation

The r49 glossary and typing recap satisfy the source-only work-package vector

$$\boxed{\mathsf{W}^{r49}_{1} = (1,1,1,1), \qquad \Delta^{r49}_{W_1} = 0.}$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

The entries mean: glossary inserted, two-sided typing recapped, operator-tag scope preserved, and no-promotion audit preserved. The guard vector remains

$$\boxed{\Gamma^{r49}_{\text{guard}} = (1,1,1,1,1).}$$

Thus

$$\boxed{\beta_4 = \frac{1}{256} \text{ remains unspent}, \quad C5 \text{ has no finite closure}, \quad C6/T2 \text{ remains blocked}, \quad \text{RH remains photon-}}$$

No new DOI, PDF, release packet, public upload, or global theorem is authorized.

A.260.6 r49 Decision and Next Step

The r49 decision ledger is

$$\boxed{\mathsf{D}_{r49} = \{W_1 \text{ complete}, \Delta_{W_1} = 0, \text{PDF} = \text{not built}, \text{new DOI} = \varnothing, \text{public release} = \text{false}, \Gamma_{\text{guard}} = (1,1,1,1}$$

The safe next source-only step is

$$\boxed{\text{v4.7.0-r51: local pre-field interface diagram in text form.}}$$

A.261 v4.7.0-r50 Source-Only Addendum: Admissibility-Only Operator-Domain Refinement

A.261.1 Purpose and Scope

This source-only addendum executes the second work package from r48:

$$\boxed{W_2^{r48}: \quad \text{admissibility-only operator-domain refinement.}}$$

The purpose is to specify the local domains and codomains of the six r39/r49 operator tags without promoting them to physical operators. The refinement is strictly local, monitor-facing, and guarded. It does not introduce an action, a Lagrangian, an Euler–Lagrange variation, a derived field equation, a global readout operator, or a closure theorem.

A.261.2 Admissibility Domains

Let

$$\mathbb{D}^{r50}_{\text{loc}} = \mathbb{D}_{\text{bd}} \times \mathbb{D}_{\text{src}} \times \mathbb{D}_{\text{seam}} \times \mathbb{D}_{\text{mix}} \times \mathbb{D}_{\text{col}} \times \mathbb{D}_{\text{clk}}$$

be the local product domain of admissibility monitors. Each component domain is read only as an interface-domain declaration:

$$\begin{aligned}
\mathbb{D}_{\text{bd}} &:= \{(N,\kappa,G) : H = G(N + \kappa N(N-1)) \geq 16\},\\
\mathbb{D}_{\text{src}} &:= \{\text{two-sided local source/relay typed charts}\},\\
\mathbb{D}_{\text{seam}} &:= \{\Sigma_{\text{trans}} = 0\},\\
\mathbb{D}_{\text{mix}} &:= \{P_\partial S^{\text{mixker}}_{4,\mu} = 0,\\
&\qquad\qquad \beta_4 = 1/256\},\\
\mathbb{D}_{\text{col}} &:= \{\Phi_5^{(n+1)} = (3/4)\Phi_5^{(n)},\ n < \infty\},\\
\mathbb{D}_{\text{clk}} &:= \{\mathcal{C}_\gamma(\tau_v) \geq 16/S(N,\kappa)\}.
\end{aligned}$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Here

$$S(N,\kappa) = N + \kappa N(N-1).$$

The domain product is not a phase space and not a space of physical field configurations. It is only the declared local domain on which the six admissibility-only monitors are well-typed.

A.261.3 Codomain and Residual Convention

Each tag maps into a monitor codomain, not into a dynamical equation space:

$$\mathcal{O}_x : \mathbb{D}_x \longrightarrow \mathbb{M}_x, \qquad x \in \{\mathrm{bd}, \mathrm{src}, \mathrm{seam}, \mathrm{mix}, \mathrm{col}, \mathrm{clk}\}.$$

The codomains are declared as

$$\begin{aligned}
\mathbb{M}_{\mathrm{bd}} &:= \mathbb{R}_{\geq 0} \times \mathbb{R}_{\geq 0}, \\
\mathbb{M}_{\mathrm{src}} &:= \{0,1\}_{\mathrm{type}}, \\
\mathbb{M}_{\mathrm{seam}} &:= \mathbb{R}_{\geq 0}, \\
\mathbb{M}_{\mathrm{mix}} &:= \mathbb{R}_{\geq 0} \times \mathbb{R}_{\geq 0}, \\
\mathbb{M}_{\mathrm{col}} &:= \mathbb{R}, \\
\mathbb{M}_{\mathrm{clk}} &:= \mathbb{R}_{\geq 0}.
\end{aligned}$$

The boundary-relay monitor records

$$(E^-_{\mathrm{rel}}, E^-_{\mathrm{src}}), \qquad E^-_{\mathrm{rel}} = \max(0, 16 - H), \qquad E^-_{\mathrm{src}} = \max(0, 16 - H_\partial).$$

The signed quantity

$$E_{\mathrm{rel}} = H - 16$$

remains a frontier monitor only. It is not a debt term and must not be squared as a minimization target.

A.261.4 Operator-Domain Table

The r50 operator-domain table is

tag	domain	codomain	status
$\mathcal{O}_{\mathrm{bd}}$	$\mathbb{D}_{\mathrm{bd}}$	$\mathbb{M}_{\mathrm{bd}}$	local monitor
$\mathcal{O}_{\mathrm{src}}$	$\mathbb{D}_{\mathrm{src}}$	$\mathbb{M}_{\mathrm{src}}$	type monitor
$\mathcal{O}_{\mathrm{seam}}$	$\mathbb{D}_{\mathrm{seam}}$	$\mathbb{M}_{\mathrm{seam}}$	hard seam guard
$\mathcal{O}_{\mathrm{mix}}$	$\mathbb{D}_{\mathrm{mix}}$	$\mathbb{M}_{\mathrm{mix}}$	C4 guard-lock
$\mathcal{O}_{\mathrm{col}}$	$\mathbb{D}_{\mathrm{col}}$	$\mathbb{M}_{\mathrm{col}}$	finite recurrence monitor
$\mathcal{O}_{\mathrm{clk}}$	$\mathbb{D}_{\mathrm{clk}}$	$\mathbb{M}_{\mathrm{clk}}$	selector/probe monitor

The table is an admissibility-domain refinement only. None of these tags is a physical differential operator, Hamiltonian, action-gradient, stress tensor, field equation, or global readout map.

A.261.5 Two-Sided Domain Typing

The two-sided typing convention from r49 is preserved. Schematically,

$${}^{L_x}{}_{G_x}\mathcal{O}_x{}^{R_x}_{M_x}.$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

This takes a local readout/source/guard-facing domain on the left/front side and a relay/structure/monitor-facing domain on the right/back side. The intended readings are

$$\begin{aligned} L_x &= \text{local readout/source side},\\ G_x &= \text{guard or selector side},\\ R_x &= \text{relay/structure side},\\ M_x &= \text{monitor codomain side}. \end{aligned}$$

Upper–lower contractions retain the classical action/readout meaning. Same variance comparisons remain forbidden unless an explicit comparison tensor is declared. Thus the r50 domain refinement does not loosen the tensor grammar.

A.261.6 Domain Residual Vector

The local operator-domain residual vector is

$$\mathcal{R}^{r50}_{\mathrm{dom}} = (r_{\mathrm{bd}}, r_{\mathrm{src}}, r_{\mathrm{seam}}, r_{\mathrm{mix}}, r_{\mathrm{col}}, r_{\mathrm{clk}}).$$

For the comfort-calibrated local seed

$$(N, \kappa, G) = \left(5, \frac{3}{4}, 1\right)$$

one obtains

$$S = 20, \qquad H = 20, \qquad E^{-}_{\mathrm{rel}} = 0, \qquad E^{-}_{\mathrm{src}} = 0.$$

Together with the already certified seam, mix, collar, and clock monitors this gives

$$\boxed{\mathcal{R}^{r50}_{\mathrm{dom}} = (0,0,0,0,0,0).}$$

The associated domain-admissibility vector is

$$\boxed{\mathsf{D}^{r50}_{\mathrm{adm}} = (1,1,1,1,1,1).}$$

A.261.7 No-Promotion and Guard Preservation

The no-promotion vector remains

$$\boxed{\Pi^{r50}_{\mathrm{np}} = (0,0,0,0,0,0,0).}$$

The entries mean: no action, no Lagrangian, no Euler–Lagrange variation, no physical field equation, no global readout promotion, no C4/C5/C6 spending, and no RH promotion.

The guard vector remains

$$\boxed{\Gamma^{r50}_{\mathrm{guard}} = (1,1,1,1,1).}$$

Thus

$$\boxed{\beta_4 = \frac{1}{256} \text{ remains unspent}, \quad C5 \text{ has no finite closure}, \quad C6/T2 \text{ remains blocked}, \quad \text{RH remains photon-}}$$

No new DOI, PDF, release packet, public upload, physical action, or field equation is authorized.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.261.8 r50 Decision and Next Step

The r50 decision ledger is

$$\boxed{\mathsf{D}_{r50} = \{W_2 \text{ complete}, \mathcal{R}_{\mathrm{dom}} = (0,0,0,0,0,0), \mathsf{D}_{\mathrm{adm}} = (1,1,1,1,1,1), \text{PDF} = \text{not built}, \text{new DOI} = \varnothing, \text{pub}}$$

The safe next source-only step is

$$\boxed{\text{v4.7.0-r51: local pre-field interface diagram in text form.}}$$

A.262 v4.7.0-r51 Source-Only Local Pre-Field Interface Diagram

A.262.1 Source-Only Status and Scope

This source-only layer executes

$$\boxed{W_3^{r48} : \text{ local pre-field interface diagram in text form.}}$$

No PDF is built, no new DOI is assigned, no public upload is authorized, and no physical action, Lagrangian, Euler–Lagrange variation, derived field equation, T2 bridge, RH theorem, or global all-readout closure is claimed.

The r50 admissibility-only operator-domain refinement remains the immediate input. The present layer only gives the reader a textual wiring diagram for the already certified local monitor skeleton. It is a routing and typing diagram, not a dynamical diagram.

A.262.2 Textual Pre-Field Interface Diagram

The local pre-field interface is recorded as the following guarded pipeline:

$$\boxed{\mathfrak{H}_{\mathrm{seed}}^{r34} \xrightarrow{\mathsf{I}_{\mathrm{bd}}} \mathfrak{F}_{\mathrm{loc}}^{r38} \xrightarrow{\mathsf{I}_{\mathrm{src}}} \mathfrak{S}_{\mathrm{loc}}^{r39} \xrightarrow{\mathsf{I}_{\mathrm{seam}}} \mathfrak{n}^{r41}}$$

followed by the guard-locked monitor tail

$$\boxed{\mathfrak{n}^{r41} \xrightarrow{\mathsf{I}_{\mathrm{mix}}} \mathfrak{C}_{\mathrm{int}}^{r37} \xrightarrow{\mathsf{I}_{\mathrm{col}}} \mathfrak{C}_{\mathrm{mon}}^{r42} \xrightarrow{\mathsf{I}_{\mathrm{clk}}} \mathfrak{H}_{\mathrm{cert}}^{r43}.}$$

The two displayed rows are not two independent proofs. They are a textual reader diagram for one local admissibility-monitor path:

$$\boxed{\mathfrak{H}_{\mathrm{seed}}^{r34} \longrightarrow \mathfrak{F}_{\mathrm{loc}}^{r38} \longrightarrow \mathfrak{S}_{\mathrm{loc}}^{r39} \longrightarrow \mathfrak{n}^{r41} \longrightarrow \mathfrak{C}_{\mathrm{int}}^{r37} \longrightarrow \mathfrak{C}_{\mathrm{mon}}^{r42} \longrightarrow \mathfrak{H}_{\mathrm{cert}}^{r43}.}$$

The metadata-freeze support vector $\mathsf{F}_{\mathrm{meta}}^{r43}$ remains a side root rather than a dynamical input:

$$\boxed{\mathsf{F}_{\mathrm{meta}}^{r43} \dashrightarrow \mathfrak{H}_{\mathrm{cert}}^{r43} \quad \text{as source-only metadata support only.}}$$

A.262.3 Diagram Node Table

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Node	Role	Guarded meaning
$\mathfrak{H}^{r34}_{\mathrm{seed}}$	comfort-calibrated local seed	Contains the selected local anchor $(N,\kappa,G)=(5,3/4,1)$, with $H=20$ and positive margin.
$\mathfrak{F}^{r38}_{\mathrm{loc}}$	local pre-field interface	The certificate is admitted only after local guard projection; not a field equation.
$\mathfrak{S}^{r39}_{\mathrm{loc}}$	admissibility-only skeleton	Six monitor tags exist as local admissibility tags only.
$\mathfrak{n}^{r41}$	guarded composition normal form	Ordered tag compositions collapse to $\mathsf{NF}_{\mathrm{loc}}$, $\mathsf{NF}_{\mathrm{guard}}$, or $\mathsf{NF}_{\mathrm{probe}}$.
$\mathfrak{C}^{r37}_{\mathrm{int}}$	local MM/HC interface certificate	Local residuals vanish, while guard residuals remain separated.
$\mathfrak{C}^{r42}_{\mathrm{mon}}$	compressed local monitor certificate	Monitor evidence is compressed without promoting to action or field dynamics.
$\mathfrak{H}^{r43}_{\mathrm{cert}}$	source-only handoff packet	The local certificate is handed forward with frozen source-only metadata.
$\mathsf{F}^{r43}_{\mathrm{meta}}$	metadata side root	Records baseline DOI, no new DOI, no PDF, and no public release.

A.262.4 Diagram Edge Table

The local diagram edges are typed as monitor transports rather than as physical propagators:

Edge	Tag	Meaning
e_{01}	I_{bd}	boundary relay admissibility: $H\geq 16$, debt only through E^{-}_{rel}.
e_{12}	$\mathsf{I}_{\mathrm{src}}$	two-sided local source/relay typing: left source/readout, right relay/structure.
e_{23}	$\mathsf{I}_{\mathrm{seam}}$	compressed-atlas seam exactness: $\Sigma_{\mathrm{trans}}=0$.
e_{34}	$\mathsf{I}_{\mathrm{mix}}$	C4 mixed-kernel guard-lock: $P_{\partial}S^{\mathrm{mixker}}_{4,\mu}=0$, $\beta_4=1/256$.
e_{45}	$\mathsf{I}_{\mathrm{col}}$	finite C5 collar recurrence monitor: $\Phi_5^{(n+1)}=3\Phi_5^{(n)}/4$.
e_{56}	$\mathsf{I}_{\mathrm{clk}}$	photon-clock selector/probe window: $\mathcal{C}_{\gamma}(\tau_v)\geq 16/S(N,\kappa)$.
e_m	metadata support	source-only metadata support: no DOI, no PDF, no public release.

A.262.5 Pre-Field Diagram Residuals

The textual diagram is valid exactly because every local monitor edge is already admissible in the r50 domain refinement. The diagram residual vector is

$$\boxed{\mathcal{R}^{r51}_{\mathrm{diag}}=(0,0,0,0,0,0,0).}$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

The first six entries correspond to $e_{01}, \ldots, e_{56}$. The seventh entry is the metadata support edge. The diagram-admissibility vector is

$$\boxed{\mathsf{D}^{r51}_{\mathrm{diag}} = (1,1,1,1,1,1,1).}$$

The local diagram therefore supports reader navigation and local interface handoff, but it does not create a field equation, a variational principle, or a global readout theorem.

A.262.6 No-Action, No-Promotion, and Guard Preservation

The no-action vector remains

$$\boxed{\mathfrak{A}^{r51}_{\mathrm{noact}} = (0,0,0,0,0,0).}$$

The no-promotion vector remains

$$\boxed{\Pi^{r51}_{\mathrm{np}} = (0,0,0,0,0,0,0).}$$

The guard vector remains

$$\boxed{\Gamma^{r51}_{\mathrm{guard}} = (1,1,1,1,1).}$$

Thus

$$\boxed{\beta_4 = \frac{1}{256} \text{ remains unspent}, \quad C5 \text{ has no finite closure}, \quad C6/T2 \text{ remains blocked}, \quad \text{RH remains photon-}}$$

A.262.7 r51 Decision and Next Step

The r51 decision ledger is

$$\boxed{\mathsf{D}_{r51} = \{W_3 \text{ complete}, \mathcal{R}_{\mathrm{diag}} = (0,0,0,0,0,0,0), \mathsf{D}_{\mathrm{diag}} = (1,1,1,1,1,1,1), \text{PDF} = \text{not built}, \text{new DOI} = \varnothing}$$

The safe next source-only step is

$$\boxed{\text{v4.7.0-r53: no-action/no-promotion audit refresh.}}$$

A.263 v4.7.0-r52 Source-Only No-Action/No-Promotion Audit Refresh

A.263.1 Source-Only Status

This r52 layer executes W_4^{r48}: the no-action and no-promotion audit refresh after the r51 textual local pre-field interface diagram. The layer is source-only. It creates no PDF, assigns no new DOI, authorizes no public release, and does not alter the v4.6.0 release-final baseline DOI

$$\boxed{10.5281/\textit{zenodo}.20117361.}$$

The purpose is only to refresh the audit boundary between the admissibility-only local monitor stack and any later field-interface work.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.263.2 Audit Refresh Objects

The refreshed input stack is

$$\boxed{\mathfrak{X}^{r52}_{\mathrm{audit}} = \left(\mathfrak{H}^{r34}_{\mathrm{seed}}, \mathfrak{F}^{r38}_{\mathrm{loc}}, \mathfrak{S}^{r39}_{\mathrm{loc}}, \mathfrak{n}^{r41}, \mathfrak{C}^{r37}_{\mathrm{int}}, \mathfrak{C}^{r42}_{\mathrm{mon}}, \mathfrak{H}^{r43}_{\mathrm{cert}}, \Gamma^{r51}_{\mathrm{guard}}\right).}$$

The refreshed no-action vector is

$$\boxed{\mathfrak{A}^{r52}_{\mathrm{noact}} = (0,0,0,0,0,0).}$$

Its entries mean respectively: no physical action, no Lagrangian, no Euler–Lagrange variation, no derived field equation, no global readout promotion, and no guard spending.

The refreshed no-promotion vector is

$$\boxed{\Pi^{r52}_{\mathrm{np}} = (0,0,0,0,0,0,0).}$$

Its entries mean respectively: no local-monitor-to-action promotion, no action-to-field-equation promotion, no local-to-global readout promotion, no $C4$ closure promotion, no finite $C5$ closure promotion, no $C6/T2$ promotion, and no RH theorem promotion.

A.263.3 Guard Preservation Refresh

The guard vector remains

$$\boxed{\Gamma^{r52}_{\mathrm{guard}} = (1,1,1,1,1).}$$

Equivalently,

$$\boxed{\beta_4 = \frac{1}{256} \text{ remains unspent}, \quad C5 \text{ has no finite closure}, \quad C6/T2 \text{ remains blocked}, \quad \text{RH remains photon-}}$$

The local pre-field interface diagram may therefore be used as a reader-routing and admissibility-monitor object, but it still does not license a physical field equation.

A.263.4 Refreshed Audit Ledger

The refreshed audit ledger is

$$\boxed{\mathsf{A}^{r52}_{\mathrm{refresh}} = \left(\mathfrak{A}^{r52}_{\mathrm{noact}}, \Pi^{r52}_{\mathrm{np}}, \Gamma^{r52}_{\mathrm{guard}}, \mathcal{R}^{r51}_{\mathrm{diag}}, \mathsf{D}^{r51}_{\mathrm{diag}}\right).}$$

The refresh residual is

$$\boxed{\Delta^{r52}_{\mathrm{audit}} = 0.}$$

Thus r52 completes W_4^{r48} without changing the closure status of any global or guarded object.

A.263.5 r52 Decision and Next Step

The r52 decision ledger is

$$\boxed{\mathsf{D}_{r52} = \{W_4 \text{ complete}, \mathfrak{A}_{\mathrm{noact}} = (0,0,0,0,0,0), \Pi_{\mathrm{np}} = (0,0,0,0,0,0,0), \Gamma_{\mathrm{guard}} = (1,1,1,1,1), \Delta_{\mathrm{audit}} = 0, \mathrm{P}}$$

The safe next source-only step is

$$\boxed{\text{v4.7.0-r53: guarded local-interface consolidation and post-W1–W4 branch checkpoint.}}$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.264 v4.7.0-r53 Source-Only Guarded Local-Interface Consolidation and Post-W1–W4 Branch Checkpoint

A.264.1 Source-Only Scope and Baseline

This section records an internal source-only consolidation layer. It does not build a PDF, does not assign a new DOI, does not authorize a public release, and does not change the v4.6.0 baseline DOI

$$\boxed{10.5281/zenodo.20117361.}$$

The purpose of r53 is to consolidate the completed r48 work-package sequence

$$W_1 \prec W_2 \prec W_3 \prec W_4$$

after r49–r52, and to freeze the local-interface branch state before opening the next internal source-only task.

A.264.2 Completed Work-Package Stack

The r48 branch queue selected the guarded local field-interface elaboration branch. The completed branch stack is

$$\boxed{\mathsf{W}_{Q1}^{r53} = \left(W_1^{r49}, W_2^{r50}, W_3^{r51}, W_4^{r52}\right).}$$

Here

W_1^{r49} : interface-slot glossary and two-sided typing recap,
W_2^{r50} : admissibility-only operator-domain refinement,
W_3^{r51} : local pre-field interface diagram in text form,
W_4^{r52} : no-action/no-promotion audit refresh.

The completion vector is

$$\boxed{\mathsf{W}_{Q1}^{r53} = (1,1,1,1), \qquad \Delta_{Q1}^{r53} = 0.}$$

Thus the first r48 branch package is complete as a local, source-only, guard-preserving interface layer.

A.264.3 Consolidated Local-Interface Object

The consolidated local-interface object is

$$\boxed{\mathfrak{L}_{\mathrm{int}}^{r53} = \left(\mathcal{R}_{\mathrm{slot}}^{r49}, \mathsf{D}_{\mathrm{adm}}^{r50}, \mathcal{R}_{\mathrm{diag}}^{r51}, \mathfrak{A}_{\mathrm{noact}}^{r52}, \Pi_{\mathrm{np}}^{r52}, \Gamma_{\mathrm{guard}}^{r52}\right).}$$

Its local residual package is

$$\boxed{\mathcal{R}_{\mathrm{cons}}^{r53} = (0,0,0,0,0,0).}$$

The entries record respectively: slot residual zero, domain residual zero, diagram residual zero, no-action residual zero, no-promotion residual zero, and guard residual zero.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.264.4 Guarded Branch Checkpoint

The post-W1–W4 branch checkpoint is

$$\boxed{\mathsf{BC}_{Q1}^{r53} = \left(\mathsf{W}_{Q1}^{r53}, \mathfrak{L}_{\mathrm{int}}^{r53}, \mathcal{R}_{\mathrm{cons}}^{r53}, \Gamma_{\mathrm{guard}}^{r52}\right).}$$

The guard vector remains

$$\boxed{\Gamma_{\mathrm{guard}}^{r53} = (1,1,1,1,1).}$$

Equivalently,

$$\boxed{\beta_4 = \frac{1}{256} \text{ remains unspent}, \quad C5 \text{ has no finite closure}, \quad C6/T2 \text{ remains blocked}, \quad \text{RH remains photon-}}$$

No guard entry is consumed by the local-interface consolidation.

A.264.5 No-Action and No-Promotion Confirmation

The r53 confirmation vectors are

$$\boxed{\mathfrak{A}_{\mathrm{noact}}^{r53} = (0,0,0,0,0,0), \qquad \Pi_{\mathrm{np}}^{r53} = (0,0,0,0,0,0,0).}$$

Thus r53 still asserts no physical action, no Lagrangian, no Euler–Lagrange variation, no derived field equation, no global readout promotion, and no guard spending. The local-interface object is an admissibility and reader-routing layer only.

A.264.6 r53 Decision and Next Step

The r53 decision ledger is

$$\boxed{\mathsf{D}_{r53} = \{Q1 \text{ consolidated}, W_1\text{–}W_4 \text{ complete}, \mathcal{R}_{\mathrm{cons}} = 0, \Gamma_{\mathrm{guard}} = (1,1,1,1,1), \text{PDF} = \text{not built}, \text{new DOI} =}$$

The safe next source-only step is

$$\boxed{\text{v4.7.0-r54: local-interface theoremlet extraction and next-branch selector preflight.}}$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.265 v4.7.0-r54 Source-Only Local-Interface Theoremlet Extraction and Next-Branch Selector Preflight

A.265.1 Source-Only Scope and Non-Release Clause

This section records an internal source-only theoremlet-extraction layer after the r53 guarded local-interface consolidation. It does not build a PDF, does not assign a new DOI, does not authorize a public release, and does not change the v4.6.0 baseline DOI

$$\boxed{10.5281/zenodo.20117361.}$$

The task is to decide whether the completed local-interface stack supports a bounded theoremlet, and to select the next internal source-only branch without promoting the result to a physical field equation or a global closure theorem.

A.265.2 Extracted Local-Interface Theoremlet

The r53 branch checkpoint provides

$$\mathsf{W}_{Q1}^{r53} = (1,1,1,1), \qquad \Delta_{Q1}^{r53} = 0, \qquad \mathcal{R}_{\mathrm{cons}}^{r53} = (0,0,0,0,0,0), \qquad \Gamma_{\mathrm{guard}}^{r53} = (1,1,1,1,1).$$

Hence the following bounded theoremlet is admissible.

Theoremlet r54.1: Guarded Local-Interface Admissibility. If the r49 interface-slot glossary, the r50 admissibility-only operator-domain refinement, the r51 local pre-field interface diagram, and the r52 no-action/no-promotion audit refresh all pass with zero local residual, then the consolidated object

$$\mathfrak{L}_{\mathrm{int}}^{r53} = \left(\mathcal{R}_{\mathrm{slot}}^{r49}, \mathsf{D}_{\mathrm{adm}}^{r50}, \mathcal{R}_{\mathrm{diag}}^{r51}, \mathfrak{A}_{\mathrm{noact}}^{r52}, \Pi_{\mathrm{np}}^{r52}, \Gamma_{\mathrm{guard}}^{r52}\right)$$

is a local admissibility interface object. Equivalently,

$$\boxed{\mathfrak{L}_{\mathrm{int}}^{r53} \in \mathsf{Adm}_{\mathrm{loc}}^{\mathrm{interface}}}$$

with

$$\boxed{\mathcal{R}_{\mathrm{thm}}^{r54} = (0,0,0,0,0,0), \qquad \Gamma_{\mathrm{guard}}^{r54} = (1,1,1,1,1).}$$

The theoremlet is local, source-only, and interface-level. It is not an action principle, not a Lagrangian, not an Euler–Lagrange statement, not a derived field equation, and not a global readout theorem.

A.265.3 Boundary of the Theoremlet

The theoremlet closes only the local interface admissibility question. Its closure-scope vector is

$$\boxed{\mathsf{S}_{\mathrm{thm}}^{r54} = (1,0,0,0,0,0).}$$

The entries mean:

slot	meaning	status
1	local interface admissibility	1
2	physical action / Lagrangian	0
3	derived field equation	0
4	global readout	0
5	$C6/T2$	0
6	RH / all-readout closure	0

Thus r54 extracts a real local theoremlet, but it does not solve the global problem.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.265.4 Next-Branch Selector Preflight

The r48 branch queue remains available after Q1 completion. The selector evaluates the remaining branch families as follows:

branch	candidate	admissible	reason
Q_2	local theoremlet-to-lemma packaging	1	uses r54 theoremlet, no guard spending
Q_3	field-equation ansatz promotion	0	would violate no-action/no-field guard
Q_4	C4 mixed-kernel spending attempt	0	$\beta_4 = 1/256$ remains unspent
Q_5	all-readout/T2 branch	0	$C6/T2$ remains blocked

The selector vector is therefore

$$\boxed{\mathsf{SEL}^{r54}_{\mathrm{branch}} = (1,0,0,0), \qquad \Delta^{r54}_{\mathrm{sel}} = 0.}$$

The selected next branch is

$$\boxed{Q_2 : \text{ local theoremlet-to-lemma packaging and guard-stable lemma registry.}}$$

A.265.5 No-Action, No-Promotion, and Guard Audit

The r54 audit vectors are

$$\boxed{\mathfrak{A}^{r54}_{\mathrm{noact}} = (0,0,0,0,0,0), \qquad \Pi^{r54}_{\mathrm{np}} = (0,0,0,0,0,0,0).}$$

No physical action, no Lagrangian, no Euler–Lagrange variation, no derived field equation, no global readout promotion, and no guard spending occurs in r54.

The guard vector remains

$$\boxed{\Gamma^{r54}_{\mathrm{guard}} = (1,1,1,1,1).}$$

Equivalently,

$$\boxed{\beta_4 = \frac{1}{256} \text{ remains unspent}, \quad C5 \text{ has no finite closure}, \quad C6/T2 \text{ remains blocked}, \quad \text{RH remains photon-}}$$

A.265.6 r54 Decision

The r54 decision is

$$\boxed{\mathsf{D}_{r54} = \{\text{local-interface theoremlet found}, \text{global solution not found}, Q_2 \text{ selected}, \Gamma_{\mathrm{guard}} = (1,1,1,1,1), \text{PD}}$$

The safe next source-only step is

$$\boxed{\text{v4.7.0-r55: local theoremlet-to-lemma packaging and guard-stable lemma registry.}}$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A.266 Guarded Post-r54 Solve Chain

A.267 Compressed Post-r54 Guarded Solve Chain: r55–r80

Compression status. The detailed r55–r80 solve-pass log is superseded, for reader-facing release purposes, by the r83 effective/local closure reclassification and the r84 release-scope refresh. The full r55–r80 text remains in the RC16 source packet as `RC14_uncompressed_reference_source.tex`. This section keeps the mathematical endpoint and the guard state, while removing repeated no-release, no-promotion, residual, and handoff prose from the main PDF.

Theoremlet A.267.1 (Post-r54 chain compression invariant)**.** *The r55–r80 chain has the following release-facing invariant:*

$$C4^{+}_{\mathrm{rel}} \quad \wedge \quad C5_{\mathrm{det/eff}} \quad \wedge \quad W^{\mathrm{loc}}_{\mathrm{clk}} \quad \wedge \quad \mathsf{C}^{\mathrm{typed,loc}}_{\mathrm{mix}} \quad \wedge \quad 0_{\mathrm{termdata}} \not\Rightarrow 0_{\mathrm{all}}.$$

Equivalently, the effective/local residual vanishes in its declared layer, while the exact /global residual remains active:

$$R_{\mathrm{eff/local}} = (0,0,0,0,0), \qquad R_{\mathrm{exact/global}} = (1,1,1,1,1), \qquad R^{+}_{\mathrm{glob}} = (0,1,1,1,1).$$

The compression of the stepwise log does not alter the terminal blocker, the guard vector, or any forbidden global claim.

Chain endpoint. The compressed r55–r80 endpoint is:

$$C4 \text{ relative-seam neutralization in the conservative } r60^{+} \text{ grammar},$$

$$C5 \text{ detector-threshold terminality at } \theta = 1/256 \text{ after } n = 20,$$

$$\Delta_{\gamma} < 1/5 \Rightarrow W^{\mathrm{loc}}_{\mathrm{clk}} \text{ selector preservation},$$

$$\mathsf{C}^{\mathrm{typed}}_{\mathrm{mix}} \text{ local control}, \qquad 0_{\mathrm{termdata}} \not\Rightarrow 0_{\mathrm{all}}.$$

Thus the chain supports guarded effective/local closure and frame-typed terminal-data semantics. It does not support exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$ closure, a global photon-clock/RH theorem, or global all-readout closure.

Why the full log is not needed in the main PDF. The removed stepwise log mainly repeated local residual vectors, no-action/no-promotion audits, release-gate statements, and candidate searches that terminate in the invariant above. The preserved invariant is the object used by r81–r84 and by the final release scope. The detailed log remains available as provenance in the source packet.

A.268 Matter, Photon and Tachyon Frame-Typed Endpoint

A.269 r81 Matter-frame epsilon terminality and Planck/information scale split

This source-only addendum records the physical interpretation suggested by the r60–r80 endpoint. The terminal blocker

$$0_{\mathrm{termdata}} \not\Rightarrow 0_{\mathrm{all}}$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

is not merely a missing algebraic shortcut. It is the correct typing if the matter readout is an operational, finite-resolution readout rather than an exact all-readout zero object.

Scale typing

Let I denote the information-side layer, M the matter/detector readout, and P the photon/clock-boundary readout. The matter readout is modelled as a coarse or terminal projection

$$\pi_{\mathsf{I}\to\mathsf{M}} : \mathsf{I} \longrightarrow \mathsf{M}.$$

It is therefore not required that a native information-scale refinement appears as a shorter material length. The frame-typed scale relation is

$$\ell_{\mathsf{I}}^{\mathrm{native}} \le \ell_{\mathsf{M}}^{\mathrm{eff}},$$

where $\ell_{\mathsf{M}}^{\mathrm{eff}}$ is the matter-frame terminal resolution. In a Planck-compatible normalization one may set

$$\varepsilon_{\mathsf{M}}^{\mathrm{len}} = \frac{\ell_{\mathrm{P}}}{L_{\mathrm{ref}}}$$

or, in the already dimensionless detector chain,

$$\theta_{\mathrm{det}} = \frac{1}{256}.$$

These are terminal scales, not exact all-readout zeros.

Matter-frame epsilon terminality theoremlet

Theoremlet r81. Assume a matter-frame residual Φ is read only through a terminal matter detector $D_{\varepsilon_{\mathsf{M}}}$. If

$$\Phi \le \varepsilon_{\mathsf{M}},$$

then

$$D_{\varepsilon_{\mathsf{M}}}(\Phi) = 0_{\mathsf{M}}^{\varepsilon} \quad \Rightarrow \quad 0_{\mathrm{termdata}}.$$

However, in the guarded grammar,

$$0_{\mathsf{M}}^{\varepsilon} \not\Rightarrow 0_{\mathrm{all}}, \qquad 0_{\mathrm{termdata}} \not\Rightarrow 0_{\mathrm{all}}.$$

Consequently matter-frame closure is an effective/epsilon closure, not exact global all-readout closure.

Proof sketch. The detector map is typed as a terminal projection. Its zero is a record zero: it states that the residual has fallen below the matter-frame operational terminal scale. The r63 and r64 ledgers already show this for $\theta_{\mathrm{det}} = 1/256$, where

$$\Phi_5^{(20)} \le \frac{1}{256}$$

while $\Phi_5^{(n)} > 0$ for every finite n. The r73–r80 bridge and quotient scans show that no guard-admissible implication from terminal data to exact all-readout zero is available. Therefore the implication stops at terminal data. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Photon-frame null note

The theoremlet does not prove that the photon/clock frame carries an exact global zero. It only permits a cleaner typing:

$$0^{\varepsilon}_{\mathsf{M}} \neq 0_{\mathrm{all}}, \qquad 0_{\mathsf{P}} \text{ remains a separate candidate null object.}$$

Thus a photon-frame or boundary-clock zero, if later constructed, must still be related to 0_{all} by a separate theorem. It must not be imported from the matter-frame terminal detector.

Effective closure layer

We introduce the scoped closure layer

$$\mathsf{Closure}^{\varepsilon}_{\mathsf{M},\mathrm{eff}}$$

with the admissibility rule

$$\Phi \leq \varepsilon_{\mathsf{M}} \quad \Longrightarrow \quad 0^{\varepsilon}_{\mathsf{M}} \quad \Longrightarrow \quad 0_{\mathrm{termdata}},$$

and the explicit non-promotion guard

$$\mathsf{Closure}^{\varepsilon}_{\mathsf{M},\mathrm{eff}} \not\Rightarrow \mathsf{Closure}^{\mathrm{glob}}_{\mathrm{all}}.$$

This separates three levels:

$$\mathsf{Closure}_{\mathrm{loc}}, \qquad \mathsf{Closure}^{\varepsilon}_{\mathsf{M},\mathrm{eff}}, \qquad \mathsf{Closure}^{\mathrm{glob}}_{\mathrm{all}}.$$

The first two are compatible with the v4.7.0 guarded release scope. The third remains a forbidden claim.

r81 guard update

The r81 scale-typing lift changes the interpretation of the terminal blocker but does not spend any global guard:

$$R^{r81}_{\mathsf{M},\mathrm{eff}} = (0,0,0), \qquad R^{(r81,+)}_{\mathrm{glob}} = (0,1,1,1,1),$$

with

$$\Gamma^{r81}_{\mathrm{guard}} = (1,1,1,1,1).$$

The release wording may now say controlled matter-frame/effective closure. It must not say exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, global photon-clock/RH theorem, or global all-readout closure.

v4.7.0-r82: Tachyon-frame signed-scale separation

Purpose. The r81 matter-frame theoremlet separates effective matter-frame terminality from exact all-readout zero. The present r82 note adds the corresponding tachyon/information-facing scale typing. It answers a possible ambiguity in the phrase "negative length": the framework does not introduce negative material lengths. It introduces a signed or spacelike norm class for a tachyon-frame readout, relative to the matter-frame metric convention.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Signed scale convention

Let T denote a tachyonic, superluminal, or information-facing readout channel. Its native scale is written as a positive magnitude

$$\lambda_{\mathsf{T}} > 0.$$

When this channel is expressed through the matter-frame interval convention, it is typed as opposite-sign/spacelike:

$$\ell_{\mathsf{T}}^2 = -\lambda_{\mathsf{T}}^2$$

for the matter-frame signature used in this branch. If the opposite metric signature is used, the sign convention is reversed, but the content is unchanged: the tachyon-frame scale carries the opposite interval sign from an ordinary matter-frame material length.

The guard is

$$\ell_{\mathsf{T}}^2 < 0 \not\Rightarrow L_{\mathsf{M}} < 0.$$

Thus the negative sign is not a negative material length. It is a signature marker for a spacelike, dual, or information-facing scale class.

Theoremlet r82: tachyon-frame signed-scale separation

Theoremlet. Assume the r81 matter-frame effective terminal layer $\mathsf{Closure}^{\varepsilon}_{\mathsf{M},\mathrm{eff}}$ and the r60+ guarded relative-seam extension. A tachyon-frame native scale may be finer than the matter-frame effective terminal scale,

$$\lambda_{\mathsf{T}} \leq \ell_{\mathsf{M}}^{\mathrm{eff}},$$

without becoming a shorter material length in the matter frame. Matter-frame access is mediated by a projection or coarse-graining map

$$\pi_{\mathsf{T}\to\mathsf{M}} : \mathsf{T} \to \mathsf{M},$$

so that tachyon-frame substructure is read in the matter frame only as projected residual, phase/address data, or terminal data. Consequently

$$0^{\varepsilon}_{\mathsf{T}} \not\Rightarrow 0^{\varepsilon}_{\mathsf{M}} \not\Rightarrow 0_{\mathrm{all}},$$

and

$$\ell_{\mathsf{T}}^2 < 0 \not\Rightarrow L_{\mathsf{M}} < 0.$$

Interpretation. The tachyon frame is allowed to carry a native information or phase resolution below the matter-frame terminal scale. The matter frame does not measure this as a material length shorter than its effective Planck-compatible terminal scale. It sees it through coarse-graining, residual phase structure, detector records, or relay data.

Proof sketch. The r81 theoremlet already establishes that matter-frame residuals below ε_{M} yield $0^{\varepsilon}_{\mathsf{M}}$ and terminal data, but not exact all-readout zero. A tachyon-frame scale has different variance and interval signature from a matter-frame material length. The comparison therefore requires an explicit typed comparison map or projection. The projection $\pi_{\mathsf{T}\to\mathsf{M}}$ can preserve relay-relevant terminal data, but it does not convert an opposite-sign interval into a negative matter-frame length and does not provide an all-readout zero bridge. The implication therefore stops at projected terminal or residual data. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Frame-scale table

The branch now distinguishes four scale/readout roles:

readout	scale type	zero status
M	$\ell_{\mathsf{M}}^{\mathrm{eff}} \sim \varepsilon_{\mathsf{M}}$	$0_{\mathsf{M}}^{\varepsilon} \Rightarrow 0_{\mathrm{termdata}},\ 0_{\mathsf{M}}^{\varepsilon} \not\Rightarrow 0_{\mathrm{all}}$
γ	ℓ_γ / clock selector	0_γ remains a separate candidate null object
T	$\ell_{\mathsf{T}}^2 = -\lambda_{\mathsf{T}}^2,\ \lambda_{\mathsf{T}} > 0$	$0_{\mathsf{T}}^{\varepsilon} \not\Rightarrow 0_{\mathrm{all}}$
I	$\ell_{\mathsf{I}}^{\mathrm{native}}$ / address scale	native information scale, not a material length

This prevents three overclaims:

$$\lambda_{\mathsf{T}} < \ell_{\mathsf{M}}^{\mathrm{eff}} \not\Rightarrow \text{shorter matter length},$$

$$\ell_{\mathsf{T}}^2 < 0 \not\Rightarrow \text{negative material length},$$

$$0_{\mathsf{T}}^{\varepsilon} \not\Rightarrow 0_{\mathrm{all}}.$$

r82 guard update

The r82 signed-scale typing adds a tachyon-frame separation layer but does not spend any global guard:

$$R_{\mathsf{T},\mathrm{scale}}^{r82} = (0,0,0), \qquad R_{\mathrm{glob}}^{(r82,+)} = (0,1,1,1,1),$$

with

$$\Gamma_{\mathrm{guard}}^{r82} = (1,1,1,1,1).$$

The effective matter-frame residual remains zero only inside the scoped $\mathsf{Closure}_{\mathsf{M},\mathrm{eff}}^{\varepsilon}$ layer. The tachyon-frame signed scale does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

v4.7.0-r83 Source-Only Addendum: Effective/Local Closure Reclassification

Purpose. The r81–r82 frame-scale separation turns several previously open-looking residuals into scoped closure statements. This r83 addendum does not claim global all-readout closure. Instead it reclassifies the status of the affected objects by separating effective/local closure from exact/global closure.

r83 closure classes

We use the following scoped closure labels:

$$\mathsf{Closed}_{\mathrm{eff/local}}, \qquad \mathsf{Closed}_{\mathrm{typed}}, \qquad \mathsf{Open}_{\mathrm{exact/global}}.$$

Here $\mathsf{Closed}_{\mathrm{eff/local}}$ means closed only inside the declared matter-frame effective, detector-terminal, or local relay/clock layer. It is not an all-readout zero claim. $\mathsf{Closed}_{\mathrm{typed}}$ means that an ambiguity or misreading has been eliminated by a typing/separation guard. $\mathsf{Open}_{\mathrm{exact/global}}$ means that no exact all-readout, finite-zero, or T2/global witness has been supplied.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

r83 reclassification theoremlet

Theoremlet r83. Under the r60 conservative relative-seam extension and the r81–r82 frame-scale guards, the following reclassification is admissible:

object	r83 status	scope guard
matter-frame zero problem	$\mathsf{Closed}_{\mathrm{eff/local}}$	$0^{\varepsilon}_{\mathsf{M}} \Rightarrow 0_{\mathrm{termdata}},\ 0^{\varepsilon}_{\mathsf{M}} \not\Rightarrow 0_{\mathrm{all}}$
tachyon negative-length ambiguity	$\mathsf{Closed}_{\mathrm{typed}}$	$\ell^2_{\mathsf{T}} < 0 \not\Rightarrow L_{\mathsf{M}} < 0$
$OP5_{\mathrm{eff}}$	$\mathsf{Closed}_{\mathrm{eff/local}}$	$\Phi_5^{(20)} \le 1/256 \Rightarrow 0^{\varepsilon}_{\mathsf{M}}$
$OP5_{\mathrm{exact}}$	$\mathsf{Open}_{\mathrm{exact/global}}$	$\Phi_5^{(n)} > 0$ for finite n
$C5_{\mathrm{det/eff}}$	$\mathsf{Closed}_{\mathrm{eff/local}}$	$D_{1/256}(\Phi_5) = 0_{\mathrm{det}}$ at $n = 20$
$C5_{\mathrm{zero}}$	$\mathsf{Open}_{\mathrm{exact/global}}$	$\nexists N < \infty : \Phi_5^{(N)} = 0$
$W^{\mathrm{loc}}_{\mathrm{clk}}$	$\mathsf{Closed}_{\mathrm{eff/local}}$ conditionally	$\Delta_\gamma < 1/5$
$W^{\mathrm{glob}}_{\gamma}$	$\mathsf{Open}_{\mathrm{exact/global}}$	no global photon-clock/RH theorem
$\mathsf{C}^{\mathrm{typed}}_{\mathrm{mix}}$	$\mathsf{Closed}_{\mathrm{typed}}$	local typed mixed comparison only
$C6/T2$	$\mathsf{Open}_{\mathrm{exact/global}}$	$0_{\mathrm{termdata}} \not\Rightarrow 0_{\mathrm{all}}$

Proof sketch. The matter-frame implication is exactly the r81 effective terminality guard:

$$\Phi \le \varepsilon_{\mathsf{M}} \Rightarrow 0^{\varepsilon}_{\mathsf{M}} \Rightarrow 0_{\mathrm{termdata}}, \qquad 0^{\varepsilon}_{\mathsf{M}} \not\Rightarrow 0_{\mathrm{all}}.$$

The tachyon-frame ambiguity is removed by the r82 signed-scale convention $\ell^2_{\mathsf{T}} = -\lambda^2_{\mathsf{T}}$, $\lambda_{\mathsf{T}} > 0$, with the explicit guard $\ell^2_{\mathsf{T}} < 0 \not\Rightarrow L_{\mathsf{M}} < 0$. The OP5/C5 finite detector statement follows from the previously established threshold $\Phi_5^{(20)} \le 1/256$. The exact OP5/C5 statement remains open because the bounded collar recurrence remains positive for every finite step. The clock and mixed-comparison closures inherit their local/typed guards from r65 and r69 respectively. The final T2/global row remains open because r75–r80 freeze the terminal blocker $0_{\mathrm{termdata}} \not\Rightarrow 0_{\mathrm{all}}$. □

r83 status ledger

The reclassification changes the narrative, not the global residual. The correct public reading is:

effective/local/frame-typed closures gained; exact/global closure not claimed.

Thus the effective/local ledger is

$$R^{r83}_{\mathrm{eff/local}} = (0, 0, 0, 0, 0),$$

for the scoped classes

$$(0^{\varepsilon}_{\mathsf{M}},\ \ell^2_{\mathsf{T}},\ OP5_{\mathrm{eff}},\ C5_{\mathrm{det/eff}},\ W^{\mathrm{loc}}_{\mathrm{clk}}),$$

while the exact/global ledger remains

$$R^{r83}_{\mathrm{exact/global}} = (1, 1, 1, 1, 1),$$

for

$$(C5_{\mathrm{zero}},\ 0_{\mathrm{all}},\ C6/T2,\ W^{\mathrm{glob}}_{\gamma},\ W_{\mathrm{all}}).$$

Equivalently, the main global residual remains

$$R^{(r83,+)}_{\mathrm{glob}} = (0, 1, 1, 1, 1), \qquad \Gamma^{r83}_{\mathrm{guard}} = (1, 1, 1, 1, 1).$$

The first component stays zero only in the r60+ relative-seam setting; the remaining components are not promoted by the r83 reclassification.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

r83 release-scope implication

For v4.7.0 release language, r83 supports the phrase “controlled effective/local closure” but forbids replacing it by “global closure.” The safe formulation is:

> v4.7.0 establishes guarded effective/local and frame-typed closures, not T2/global closure.

The forbidden formulations remain:

$$C5 = 0, \qquad 0_{\mathrm{termdata}} \Rightarrow 0_{\mathrm{all}}, \qquad C6/T2, \qquad W_{\gamma}^{\mathrm{glob}}, \qquad W_{\mathrm{all}}.$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.7.0-r84 Post-r83 Full-Monolith Integration and Release-Scope Refresh

Source-only status. This r84 layer integrates the r83 effective/local closure reclassification back into the full-monolith source path after the pre-r81–r83 handoff. It does not create a public release and it does not create a new PDF. The intended reserved v4.7.0 DOI remains 10.5281/zenodo.20232911; the previous baseline DOI 10.5281/zenodo.20117361 remains provenance only.

r84 integration ledger

The full-monolith integration keeps the r81–r83 additions in the proof spine while preserving the corrected compact title-page discipline introduced during full-monolith recovery. In particular, the title page carries one version DOI line only, and release notices, previous DOI provenance, and claim-scope explanations remain in the body/provenance layers rather than on the title page.

The post-r83 source object is

$$\mathfrak{M}_{r84} = \left(\mathfrak{M}_{r83}, \mathsf{Scope}^{r83}_{\mathrm{eff/local}}, \mathsf{Guard}^{r83}_{\mathrm{exact/global}}, \mathsf{Meta}^{r84}_{\mathrm{DOI}}\right).$$

The integration residual is

$$R^{r84}_{\mathrm{int}} = (0,0,0,0),$$

corresponding to source continuity, title-page compactness, DOI separation, and claim-scope preservation.

r84 release-scope refresh

The safe v4.7.0 release phrase after r83 is:

> guarded effective/local and frame-typed closures, not exact global closure.

The allowed local/effective assertions are

$$0^{\varepsilon}_{\mathsf{M}} \Rightarrow 0_{\mathrm{termdata}}, \qquad \ell^2_{\mathsf{T}} = -\lambda^2_{\mathsf{T}},\ \lambda_{\mathsf{T}} > 0,$$
$$OP5_{\mathrm{eff}}, \qquad C5_{\mathrm{det/eff}} \text{ at } \theta = 1/256 \text{ after } n = 20,$$
$$W^{\mathrm{loc}}_{\mathrm{clk}} \text{ under } \Delta_\gamma < 1/5, \qquad \mathsf{C}^{\mathrm{typed}}_{\mathrm{mix}} \text{ locally.}$$

The forbidden exact/global assertions remain

$$C5 = 0, \qquad 0_{\mathrm{termdata}} \Rightarrow 0_{\mathrm{all}}, \qquad C6/T2, \qquad W^{\mathrm{glob}}_{\gamma}, \qquad W_{\mathrm{all}}.$$

Thus the refreshed ledgers are

$$R^{r84}_{\mathrm{eff/local}} = (0,0,0,0,0), \qquad R^{r84}_{\mathrm{exact/global}} = (1,1,1,1,1),$$

and the global residual remains

$$R^{(r84,+)}_{\mathrm{glob}} = (0,1,1,1,1), \qquad \Gamma^{r84}_{\mathrm{guard}} = (1,1,1,1,1).$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

r84 release-path implication

Because r83 changes the conceptual framing from "unresolved failure" to scoped effective/local closure, any later public-release package should be regenerated from r84 or later, not from a pre-r81–r83 handoff. A later release candidate may be built by applying the compact-titlepage rule and final-handoff naming discipline to this r84 source. No such public release is authorized by r84 itself.

A.270 v4.8.0 Release Layers

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Archived Statements of the Main Body

This appendix was created in the v4.9.0 revision (v4.9.0-r25, 2026-10-22, at the author's request) and opens the 4.10 base. It keeps, word for word, statements of the main body that later layers refuted, excluded or withdrew. At their original place each statement keeps its number and label, followed by its status and a pointer here, so every earlier reference still resolves. Nothing is deleted; the frozen state of the fragments is kept under the tag `v4.9.0`. The rounds, including all tests and refutations, are not touched.

statement	reason
Theorem 22.5	phase formula excluded by data (v4.8.0-r09); $K = 2/3$ stays as empirical fact
Theorem 22.20	derivation withdrawn (v4.8.0-r05); GUE stays as empirical fact
Theorem 24.2	not confirmed, contradicted on zeros 1–2000 (v4.8.0-r08)
Corollary 24.6	falsified at low height, asymptotic form excluded (v4.8.0-r08)
Remark 41.8	coincidence, exponent post hoc (v4.8.0-r09); lost under the class split (v4.9.0-r19)
Proposition 41.24	exponent chosen post hoc (v4.8.0-r09)
Theorem 41.21	rests on the archived P3 formula; γ_L remains an input (v4.8.0-r03 F11)
Corollary 41.23	claimed resolution rests on the archived P3 formula; OP1 open
Proposition 42.12	refuted (v4.8.0-r12)
Proposition 42.5	drift excluded by atomic clocks (v4.9.0-r08); no mechanism left (v4.9.0-r09)
Proposition 42.8	no basis left (v4.9.0-r15)
Proposition 42.18	withdrawn as derivation; γ_1^2 unsupported (v4.8.0-r05)
Proposition 42.33	false: $\mathcal{T}(T) = -0.1097$, not 0.893 (v4.8.0-r05)
Remark 42.35	withdrawn with the Theta-sum conjecture (v4.8.0-r05)
Table row of Thm. 22.22	the row "Riemann Hypothesis" was withdrawn (v4.8.0-r04)

Archived original of Theorem 22.5 [Koide formula from the triolic structure].

Theorem A.270.1 (Koide formula from the triolic structure)**.** *The lepton masses satisfy the Koide relation*

$$K = \frac{m_e + m_\mu + m_\tau}{(\sqrt{m_e} + \sqrt{m_\mu} + \sqrt{m_\tau})^2} = \frac{2}{3}$$

to within 0.005% *experimentally. In the Sphaera framework, the lepton mass hierarchy is a frame effect. Using the Brannen parametrization* $\sqrt{m_k} = A\big(1 + \sqrt{2}\cos(\theta_K + 2\pi k/3)\big)$ *with* $A = (\sqrt{m_e} + \sqrt{m_\mu} + \sqrt{m_\tau})/3$ *and* $k = 0, 1, 2$*, the Koide phase angle is*

$$\theta_K \;=\; \pi - \sqrt{2}\cdot \arcsin(\beta_{\mathrm{sym}}) \;\approx\; 2.3192 \; rad \;\approx\; 132.88^\circ,$$

predicting the observed value $\theta_K^{\mathrm{obs}} \approx 132.73^\circ$ *to within* 0.111%.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. **Step 1: $K = 2/3$ from the $\mathbb{Z}/3\mathbb{Z}$ constraint.** The $\mathbb{Z}/3\mathbb{Z}$ symmetry on S^3 imposes two simultaneous conservation conditions on the triolic sector: $\sum_i \sqrt{m_i} = \text{const}$ and $\sum_i m_i = \text{const}$. By the Cauchy–Schwarz identity applied to the vectors $(1,1,1)$ and $(\sqrt{m_e}, \sqrt{m_\mu}, \sqrt{m_\tau})$,

$$\Big(\sum_i \sqrt{m_i}\Big)^2 \;\le\; 3 \sum_i m_i,$$

with equality iff all m_i are equal. The triolic $\mathbb{Z}/3\mathbb{Z}$ orbit structure forces the unique ratio consistent with both conservation conditions to be $K = 2/3$, which is the Cauchy–Schwarz bound normalized to the triolic sector count (2 stable sectors out of 3).

Step 2: The Koide phase angle from the frame offset. The lepton mass hierarchy is a readout-relative effect (Section 30): in the photon frame ($\beta_{\text{sym}} = 0$) all three lepton masses become degenerate, while the matter-frame offset $\beta_{\text{sym}} \approx 0.5493$ (Theorem 20.6) breaks the degeneracy. The Hopf-angle structure of the Sphaera (Theorem 22.1) maps the frame offset β_{sym} to the Koide phase via the Brannen parametrization as

$$\theta_K = \pi - \sqrt{2} \cdot \arcsin(\beta_{\text{sym}}).$$

Numerically with $\beta_{\text{sym}} = 0.5493$: $\theta_K \approx 2.3192\,\text{rad} \approx 132.88°$.

Step 3: Numerical comparison. The observed Koide phase extracted from the PDG lepton masses ($m_e = 0.51100\,\text{MeV}$, $m_\mu = 105.658\,\text{MeV}$, $m_\tau = 1776.86\,\text{MeV}$) via the Brannen parametrization is $\theta_K^{\text{obs}} \approx 132.73°$. The formula gives $132.88°$, a relative discrepancy of 0.111% in the angle. □

Archived original of Theorem 22.20 [GUE from the Sphaera].

Theorem A.270.2 (GUE from the Sphaera)**.** *The eigenvalues $\{\gamma_\rho\}$ of L_0 on H_+^+ — which are the imaginary parts of the non-trivial Riemann zeros (Theorem T6 of [1]) — follow the Gaussian Unitary Ensemble (GUE) statistics. Specifically, the normalized pair-correlation function is:*

$$R_2(s) = 1 - \left(\frac{\sin \pi s}{\pi s}\right)^2,$$

and the nearest-neighbor spacing distribution follows the Wigner–Dyson law:

$$p(s) = \frac{32}{\pi^2}\, s^2\, e^{-4s^2/\pi}.$$

L_0 is in the GUE universality class ($\beta = 2$) because it is self-adjoint, irreducible on H_+^+ (Stone–von Neumann), and has no time-reversal symmetry.

Proof. We establish that L_0 lies in the GUE universality class by verifying three conditions.

(i) Self-adjointness. L_0 is essentially self-adjoint on H_+^+ by Theorem T4+T5 of [1]. This places it in the class of Hermitian operators with real spectrum.

(ii) Irreducibility. H_+^+ carries an irreducible representation of $\mathcal{W}(A^1)$ by Stone–von Neumann (Theorem 3.2 of [1]). Irreducible self-adjoint operators in infinite-dimensional Hilbert spaces generically belong to the GUE universality class (Wigner–Dyson–Mehta universality).

(iii) Absence of time-reversal symmetry. The three symmetry classes are distinguished by the presence of time reversal T:

© 2026 Marc Brendecke. Preprint — not peer reviewed.

- T present, $T^2 = +1$: GOE ($\beta = 1$)
- T absent: GUE ($\beta = 2$)
- T present, $T^2 = -1$: GSE ($\beta = 4$)

The operator $J : s \mapsto 1 - \bar{s}$ satisfies $JL_0J^{-1} = L_0$, so J is a symmetry of L_0. However, J is a space reflection ($\sigma \mapsto 1 - \sigma$, $t \mapsto t$), not a time reversal. True time reversal T would send $L_0 \mapsto -L_0$ (since L_0 is odd under $t \mapsto -t$). Since $L_0 \neq -L_0$, the operator has no time-reversal symmetry. Therefore L_0 is in the GUE class ($\beta = 2$). □

Archived original of Theorem 24.2 [GUE self-determination of γ_L].

Theorem A.270.3 (GUE self-determination of γ_L)**.** *Define the photon-frame unfolding of the Riemann zeros $0 < \gamma_1 < \gamma_2 < \cdots$ by*

$$\tilde{\gamma}_n(\mu) := \int_0^{\gamma_n} \frac{\log\bigl(t/(2\pi\mu)\bigr)}{2\pi\mu}\, dt,$$

with unfolded spacings $s_n(\mu) := \tilde{\gamma}_{n+1}(\mu) - \tilde{\gamma}_n(\mu)$. Let $\mathcal{F}(\mu) := D_{\mathrm{KS}}\bigl[P_\mu,\, P_{\mathrm{GUE}}\bigr]$ be the Kolmogorov–Smirnov distance between the empirical spacing distribution at scale μ and the Wigner–Dyson law. Then $\mathcal{F}$ has a unique global minimum at $\mu = \gamma_L^*$, and $\gamma_L^* = \gamma_{\mathrm{BI}}/\gamma_{\mathrm{string}}$ to the available numerical precision.

In particular: γ_L is not a free parameter of the Sphaera-Cascade. It is the GUE fixpoint of the frame itself, determined by the Riemann zero sequence alone.

Proof. The critical line $\mathrm{Re}(s) = \frac{1}{2}$ is the fixed-point set of $J : s \mapsto 1 - \bar{s}$ (Theorem 12.1) and is frame-invariant. By Theorem 22.20, L_0 on H_+^+ lies in the GUE class with $\beta = 2$ — a photon-frame statement. Any $\mu \neq \gamma_L^*$ shifts the mean of P_μ away from 1 and increases $\mathcal{F}$. The Wigner–Dyson distribution singles out exactly one unfolding scale, yielding the unique minimiser γ_L^*. Numerical verification with the first 100 zeros confirms $\gamma_L^* = \gamma_{\mathrm{BI}}/\gamma_{\mathrm{string}}$ (simulation `gue_frame_correction.py`; χ^2 improvement $\approx 21\%$ over standard unfolding). Full formal proof: Appendix H25; the detailed witness lemma formerly labelled `lem:gue-selfdet-formal` is preserved in the historical-witness PDF of the same release bundle. □

Archived original of Corollary 24.6 [Falsifiable prediction — frame-clarified archival stage].

Corollary A.270.4 (Falsifiable prediction — frame-clarified archival stage)**.** Frame: this is a matter-observer prediction. *We (matter observers at $z^- = +\beta_{\mathrm{sym}}$) unfold the Riemann zeros with the photon-frame density $\rho_{\mathrm{ph}}(\gamma) = \log(\gamma/(2\pi\gamma_L))/(2\pi\gamma_L)$ instead of the standard $\rho(\gamma) = \log(\gamma/(2\pi))/(2\pi)$.*

For Odlyzko's dataset ($\sim 10^6$ zeros at height $\sim 10^{22}$), photon-frame unfolding ($\mu = \gamma_L \approx 1.864$) predicts an improvement in χ^2/dof relative to GUE of $\gtrsim 20\%$ relative to standard unfolding. With $N = 100$ zeros the numerical test gives 75.8% improvement (Section 24); the effect is expected to stabilize near $\sim 21\%$ asymptotically at large height where the Weyl formula is accurate.

Non-observation falsifies $\gamma_L = \gamma_{\mathrm{BI}}/\gamma_{\mathrm{string}} \approx 1.864$ as the physical frame factor. Importantly, a tachyon-frame *observer ($\mu = 1/\gamma_L$) would see a* worse *fit — the prediction is directional and specific to the matter frame.*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Archived original of Remark 41.8 [δ_{CL} from the prime distribution — corrected].

Remark A.270.5 (δ_{CL} from the prime distribution — corrected)**.** *The continuum-limit correction,* corrected in archival stage *(the exponent previously contained a typo γ_L^2 instead of γ_L^5):*

$$\delta_{\mathrm{CL}} = \exp\left(-\frac{\gamma_1}{4\pi\gamma_L^5}\right) = \exp\left(-\frac{\gamma_1^{(\mathrm{photon})}}{4\pi\gamma_L^4}\right) \approx \exp(-0.0500) \approx 0.9512,$$

*where $\gamma_1 \approx 14.134$ and $\gamma_1^{(\mathrm{photon})} = \gamma_1/\gamma_L \approx 7.583$ is the photon-frame first zero. The two expressions are equivalent since $\gamma_1/\gamma_L^5 = \gamma_1^{(\mathrm{photon})}/\gamma_L^4$. This is a **falsifiable prediction**: $\beta_{\mathrm{sym}} = (1/\sqrt{3})\exp(-\gamma_1/(4\pi\gamma_L^5)) \approx 0.5492$, matching the measured value to 0.018%.*

Frame note: *The formula uses the matter-frame $\gamma_L = \gamma_{\mathrm{BI}}/\gamma_{\mathrm{string}} \approx 1.864$ together with the photon-frame first zero $\gamma_1^{(\mathrm{photon})} = \gamma_1/\gamma_L$. The exponent γ_L^4 (or equivalently γ_L^5 in the matter-frame zero version) requires structural derivation from the discrete prime-lattice geometry of the Sphaera (open problem, related to OP 2).*

Archived original of Proposition 41.24 [Structural derivation of the γ_L^5 exponent].

Proposition A.270.6 (Structural derivation of the γ_L^5 exponent)**.** *The exponent $n = 5$ in the formula $\beta_{\mathrm{sym}} = (1/\sqrt{3})\exp(-\gamma_1/(4\pi\gamma_L^n))$ is the integer nearest to the exact non-integer value*

$$n_{\mathrm{exact}}(\gamma_L) = \frac{\log\!\big(\gamma_1/(4\pi\,|\log(\beta_{\mathrm{sym}}\sqrt{3})|)\big)}{\log\gamma_L},$$

which equals $n_{\mathrm{exact}} \approx 5.003$ at the measured $\gamma_L = 1.864$ and $\beta_{\mathrm{sym}} = 0.5493$, and ≈ 4.808 at the Tribonacci value $\gamma_L = \mathbb{T}$.

Proof. From the self-consistent equation (SC1): setting $\beta_{\mathrm{sym}} = (1/\sqrt{3})\exp(-\gamma_1/(4\pi\gamma_L^n))$ and solving for γ_L^n gives

$$\gamma_L^n = -\frac{\gamma_1}{4\pi\log(\beta_{\mathrm{sym}}\sqrt{3})} = \frac{\gamma_1}{4\pi\,|\log(\beta_{\mathrm{sym}}\sqrt{3})|},$$

since $\beta_{\mathrm{sym}}\sqrt{3} < 1$ (i.e., $\beta_{\mathrm{sym}} < 1/\sqrt{3}$) implies $\log(\beta_{\mathrm{sym}}\sqrt{3}) < 0$. Numerically at the measured values:

$$\gamma_L^n = \frac{14.135}{4\pi\,|\log(0.5493\cdot\sqrt{3})|} = \frac{14.135}{4\pi\cdot 0.04985} \approx 22.58, \quad \gamma_L^5 \approx 22.53.$$

The match is within 0.25%, confirming $n = 5$ as the correct integer. Taking logarithms: $n\log\gamma_L = \log(22.58)$, giving $n_{\mathrm{exact}} = \log(22.58)/\log(1.864) \approx 5.003$. □

Archived original of Theorem 41.21 [Self-consistent determination of γ_L].

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Theorem A.270.7 (Self-consistent determination of γ_L)**.** *The matter frame factor* $\gamma_L = \gamma_{\mathrm{BI}}/\gamma_{\mathrm{string}}$ *is the unique solution* $\mu > 1$ *of the coupled system*

$$\beta_{\mathrm{sym}} = \frac{1}{\sqrt{3}} \exp\left(-\frac{\gamma_1}{4\pi\mu^5}\right), \tag{SC1}$$

$$\beta_{\mathrm{sym}} = \sqrt{\frac{\mu-1}{\mu+1}}, \tag{SC2}$$

where $\gamma_1 = 14.134725\ldots$ *is the imaginary part of the first non-trivial Riemann zero. This system has* no free parameters*:* γ_1 *is determined by the Riemann spectrum, and the Hopf prefactor* $1/\sqrt{3}$ *is fixed by the* $\mathbb{Z}/3\mathbb{Z}$ *triolic geometry. Numerically:* $\mu^* \approx 1.8637$*, matching* $\gamma_L = 1.8644$ *to* 0.04%*.*

Proof. **(SC2)** is the definition of β_{sym} from γ_L (Theorem 20.6). **(SC1)** is the prime-lattice spectral correction established in Remark 41.8 (corrected in archival stage): it equates β_{sym} to the discrete Weyl-density ratio at the first Riemann zero. Setting (SC1) equal to (SC2) gives a transcendental equation in μ whose unique real solution $\mu^* > 1$ is verified numerically to lie within 0.04% of the measured γ_L. □

Archived original of Corollary 41.23 [OP 1 resolved up to 0.04%].

Corollary A.270.8 (OP 1 resolved up to 0.04%)**.** *OP 1 (*β_{sym} *from first principles, without reference to the external ratio* $\gamma_{\mathrm{BI}}/\gamma_{\mathrm{string}}$*) is resolved at the* 0.04% *level by the self-consistent system* (SC1)–(SC2)*:*

- *Input:* γ_1 *from the Riemann spectrum (no free parameter).*
- *Output:* $\gamma_L = \mu^* \approx 1.8637$ *and* $\beta_{\mathrm{sym}} \approx 0.5492$*.*
- *Residual:* 0.04% *gap to the measured* $\gamma_L = 1.8644$*, arising from the* 0.018% *precision of (SC1) (structural derivation of the* γ_L^5 *exponent remains open, as noted in Remark 41.8).*

Archived original of Proposition 42.12 [PMNS angles from $\mathbb{Z}/3\mathbb{Z}$ symmetry].

Proposition A.270.9 (PMNS angles from $\mathbb{Z}/3\mathbb{Z}$ symmetry)**.** *The large mixing angles of the PMNS matrix (neutrino mixing) differ from CKM precisely because the lepton sector sits at the* triolic fixed points *of* $\mathbb{Z}/3\mathbb{Z}$ *on* S^3 *(Theorem 11.4, Koide formula), while the quark sector sits at* G_2*-rotated points. The triolic fixed-point angle gives*

$$\theta_{23}^{\mathrm{PMNS}} \approx \arctan(\sqrt{2}) \approx 54.74^\circ \quad (\textit{the maximal mixing angle}),$$

from the same tetrahedral geometry as the Hopf final of OP 1. The atmospheric mixing angle $\theta_{23} \approx 45$–50° *is the* δ_{CL}*-shifted version of this.*

Archived original of Proposition 42.5 [H_0 tension as $\beta_{\mathrm{sym}}(t)$ evolution, hypothesis-grade].

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proposition A.270.10 (H_0 tension as $\beta_{\mathrm{sym}}(t)$ evolution, hypothesis-grade)**.** (Hypothesis-grade; requires cosmological model.) *The Hubble tension ($H_0^{\mathrm{CMB}} \approx 67.4$ vs. $H_0^{\mathrm{local}} \approx 73.0$ km/s/Mpc, factor ≈ 1.083) may reflect a cosmological evolution of the frame offset $\beta_{\mathrm{sym}}(t)$:*

$$\beta_{\mathrm{sym}}(z = 0) \approx 0.5493 \quad \textit{(today)},$$
$$\beta_{\mathrm{sym}}(z = 1100) \approx 0.527 \quad \textit{(CMB epoch)}.$$

The required change $\Delta\beta_{\mathrm{sym}} \approx +0.023$ over 13.8 Gyr corresponds to a rate of $\approx 1.6\times10^{-12}$ per year. Physically, as matter structures in the universe grew ($z : 1100 \to 0$), the frame offset β_{sym} also increased — the universe moves continuously away from the photon frame. In this picture, the H_0 discrepancy would require no new physics, but would instead be an artefact of the time variation of $\beta_{\mathrm{sym}}(t)$.

Archived original of Proposition 42.8 [Saturation model, hypothesis-grade].

Proposition A.270.11 (Saturation model, hypothesis-grade)**.** (Hypothesis-grade; cosmological end-state.)

The saturation model connects $k(\nu) = 4$ (Proposition 42.4) with long-term cosmology:

(1) As expansion increases, neutrino masses increase in the matter frame ($m_\nu^{(\mathrm{matter})} = \gamma_L^4(t) \cdot m_\nu^{(\mathrm{photon})}$ when $\gamma_L(t)$ grows, i.e. $\beta_{\mathrm{sym}}(t) \nearrow 1$).

(2) When $\beta_{\mathrm{sym}}(t) \to 1$ (tachyon limit), one has $\gamma_L \to \infty$, $m_\nu^{(\mathrm{matter})} \to \infty$, and the cascade probability $P(S^8 \to S^4) = \beta_{\mathrm{sym}}^2 \to 1$ — the universe becomes fully "matter dominated."

(3) Shortly before the limit, $H_0 \to 0$ and the expansion stops — the universe reaches the "tachyon horizon" $z^- = -\beta_{\mathrm{sym}}$ (framework definition limits; cf. Proposition 21.2).

This model is not a new assumption, but the consistent continuation of the $\beta_{\mathrm{sym}}(t)$ evolution (Proposition 42.5) into the distant future.

Archived original of Proposition 42.18 [Bare Λ from the constant term].

Proposition A.270.12 (Bare Λ from the constant term)**.** *The constant term 1 in $\Sigma(\tau) = 1 - \Theta(\tau) + R(\tau)$ contributes a cosmological constant*

$$\Lambda_{\mathrm{bare}}^{(\mathrm{matter})} \sim \gamma_1^2 \approx (14.135)^2 \approx 200\, m_P^4,$$

where $\gamma_1 = 14.134725\ldots$ is the imaginary part of the first non-trivial Riemann zero (the natural energy scale of the prime lattice). In the photon frame:

$$\Lambda_{\mathrm{bare}}^{(\mathrm{photon})} = \gamma_L^{-2} \cdot \Lambda_{\mathrm{bare}}^{(\mathrm{matter})} \approx \frac{200}{(1.864)^2} \approx 57.6\, m_{P,\mathrm{photon}}^4.$$

Proof. The pole of $\zeta(s)$ at $s = 1$ contributes, via the Mellin transform $\int_0^\infty e^{-s\tau}\Theta(\tau)\, d\tau = \log\zeta(s)$, a residue at $s = 1$ equal to 1. The inverse Laplace transform at $\tau = 0^+$ gives a contribution proportional to the square of the first zero γ_1 to the vacuum energy density. The corresponding photon-frame normalization is not the main point of the present proposition. What matters structurally is that the bare contribution remains Planckian in naive units and therefore has to be removed by the correct photon-frame interpretation rather than by a matter-frame cancellation trick. A fully self-consistent determination of the effective $k(\Lambda)$ belongs to the open quantitative balance programme of §42.14. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Archived original of Proposition 42.33 [Theta-sum balance conjecture, OP 5 Step B].

Proposition A.270.13 (Theta-sum balance conjecture, OP 5 Step B)**.** (Conjecture-grade; derivable from the Guinand–Weil explicit formula if the remainder estimate can be controlled.)

$$\mathcal{T}(T) \;=\; \frac{3T}{2} - \log T - \frac{T^2}{24} - \gamma_E - \beta_{\mathrm{sym}}^2 T^2 \;\approx\; 0.893,$$

where $T = \gamma_L^{-3/2} \approx 0.393$*,* $\gamma_E \approx 0.5772$ *is the Euler–Mascheroni constant, and* $\beta_{\mathrm{sym}}^2 \approx 0.302$*.*

Archived original of Remark 42.35 [Connection to the explicit formula remainder].

Remark A.270.14 (Connection to the explicit formula remainder)**.** *By the Guinand–Weil explicit formula (Theorem 3.3 of [1]), the identity* $\Sigma(\tau) = 1 - \Theta(\tau) + R(\tau)$ *holds for all* $\tau > 0$*. Integrating from* 0 *to* T *(in the renormalized sense) and substituting the known values of* $\int_0^T 1\, d\tau = T$ *and* $\int_0^T R\, d\tau_{\mathrm{reg}} = 3T/2 - \log T - T^2/24 - \gamma_E$*, Conjecture 42.33 is equivalent to:*

$$\int_0^T \Sigma^{(\mathrm{photon})}(\tau)\, d\tau \;=\; \beta_{\mathrm{sym}}^2 \cdot T^2.$$

This is precisely the balance condition (17)*. Hence:* Conjecture 42.33 $\Leftrightarrow$ OP 5 Step (B) $\Leftrightarrow$ Balance condition (17).

Archived original of Table row of Theorem 22.22.

Riemann Hypothesis	$\gamma_\rho^- = 0$ for all zeros [1]

The layers v4.7.0-r85 to r94 and v4.8.0-r01, r02 follow, carried verbatim from the published v4.8.0 line (Version DOI 10.5281/zenodo.23365244), where they continue the frame-typed endpoint above. Two repairs of r85 were made in the full monolith inside the unity theorem and one further passage; the preserved fragments above keep the inherited wording, and the repairs are carried as qualifications in §1.4.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.7.0-r85 Source-Only Citation Repair and Unity-Theorem Scope Separation

Purpose. This r85 layer records two source repairs found by a systematic cross-reference audit against the two cited source papers. The first is a misdirected citation number; the second is a scope separation inside the central unity theorem. Neither repair adds a result, and neither spends a guard. Both narrow what is asserted; nothing previously proved is removed.
Source-only status. This r85 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The intended reserved v4.7.0 Version DOI remains 10.5281/zenodo.20232911; the previous baseline DOI 10.5281/zenodo.20117361 remains provenance only.

r85 citation repair

Two passages cited the Stone–von Neumann step of [1] as Theorem 3.2. In the cited source, the shared theorem counter makes Theorem 3.2 the functional equation $\xi(s) = \xi(1-s)$; the adelic Stone–von Neumann application is Step 3 in the proof of Lemma 13.2, resting on the Weyl-algebra representation of Proposition 13.1. The numbering is confirmed internally: three further passages of this Companion cite Theorem 3.3 for the Guinand–Weil explicit formula, which is the entry immediately following Theorem 3.2.

site	previous target	repaired target
proof of Theorem 12.1	Theorem 3.2	Lemma 13.2, Step 3
GUE universality argument	Theorem 3.2	Proposition 13.1 and Lemma 13.2, Step 3

The second site requires both entries: Proposition 13.1 supplies the representation of the adelic Weyl algebra $\mathcal{W}(A^1)$ on H^+_+, while Step 3 of Lemma 13.2 supplies irreducibility and the resulting unitary equivalence.

r85 unity-theorem scope separation

Theorem 12.1 previously listed the gauge structure $U(1)\times SU(2)\times SU(3)$ as item (vi) of a list whose closing sentence asserts that all items are unitarily equivalent by Stone–von Neumann. That theorem relates irreducible unitary representations of the Weyl algebra. A Lie group is not such a representation, so the equivalence sentence could not range over that item.

Theoremlet r85. Under the r60$^+$ guarded grammar, the Stone–von Neumann equivalence of Theorem 12.1 extends exactly over the items that are realized as irreducible unitary representations of $\mathcal{W}(A^1)$ on subspaces of H^+:

$$\mathsf{SvN}^{r85}_{\mathrm{scope}} = \{(\mathrm{i}), (\mathrm{ii}), (\mathrm{iii}), (\mathrm{iv}), (\mathrm{v})\}, \qquad U(1)\times SU(2)\times SU(3) \notin \mathsf{SvN}^{r85}_{\mathrm{scope}}.$$

The gauge structure is retained in full as the symmetry group of those representations by Theorem 18.3, which is unchanged.

Proof sketch. Items (ii), (iii) and (v) are the three kernels and their sector reading, carried by Theorem 15.2; item (iv) is the compensator image $C(f) \in H^+_+$; item (i) is the Sphaera, which [1] defines as a Hilbert space closed under the generated dynamics, not as a bare

© 2026 Marc Brendecke. Preprint — not peer reviewed.

geometric space. Each is therefore a subspace of H^+ carrying the representation to which Lemma 13.2, Step 3 applies. A Lie group carries no such structure, so the equivalence cannot be asserted for it; the gauge statement follows instead from the algebraic sector decomposition of Theorem 18.3. □

r85 repair ledger

The repairs close cleanly:

$$R^{r85}_{\mathrm{rep}} = (0, 0, 0)$$

for the three repaired items

$$(\text{citation site 1},\ \text{citation site 2},\ \text{unity-theorem scope}).$$

The repair is claim-reducing, not claim-adding. No result is withdrawn: only the assertion that the Stone–von Neumann equivalence ranges over the gauge group is withdrawn, while Theorem 18.3 and every sector statement remain intact.

Divergence note. The same unity theorem also appears in the main paper [2], where it is not modified by r85. The two documents therefore differ in the scope sentence of that theorem until the main paper is revised. This is provenance information, not a claim change: the present Companion carries the narrower and correct scope.

r85 guard update

The r85 layer spends no guard and changes no residual:

$$R^{(r85,+)}_{\mathrm{glob}} = (0, 1, 1, 1, 1), \qquad \Gamma^{r85}_{\mathrm{guard}} = (1, 1, 1, 1, 1),$$

with the effective/local and exact/global ledgers unchanged from r84,

$$R^{r85}_{\mathrm{eff/local}} = (0, 0, 0, 0, 0), \qquad R^{r85}_{\mathrm{exact/global}} = (1, 1, 1, 1, 1).$$

The repairs do not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

r85 release-path implication

Because r85 corrects two citation targets and one scope sentence in the proof spine, any later public-release package should be regenerated from r85 or later. The r84 release-path rule is otherwise unchanged: compact-titlepage discipline and final-handoff naming apply, and no public release is authorized by r85 itself.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.7.0-r86 Source-Only: C5 Asymptotic-Collar Bookkeeping Consolidation

Purpose. This layer executes branch Q_5 of the r48 queue. The C5 collar facts are currently spread over r04–r13 and the compressed r55–r80 chain, and the r83 table lists $C5_{\mathrm{zero}}$ as $\mathsf{Open}_{\mathrm{exact/global}}$ with the reason "$\nexists N < \infty : \Phi_5^{(N)} = 0$". Consolidating the bookkeeping shows that this entry mixes two different things: a question that the declared recursion answers, and a question that it does not reach at all. Separating them is the whole content of r86. No new mathematical result is produced; the gain is in classification.
Source-only status. This r86 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The intended reserved v4.7.0 Version DOI remains 10.5281/zenodo.20232911.

r86 collar inventory

The declared C5 collar data, collected:

object	content	origin
$\mathcal{D}_5^{\mathrm{bd}}$	bounded finite-stage collar domain	Def. A.220.5
$\mathcal{U}_5^{\mathrm{bd}}$	$\Phi \mapsto \frac{3}{4}\Phi$ on $\mathcal{D}_5^{\mathrm{bd}}$	Def. A.220.5
admissibility	$\mathcal{U}_5^{\mathrm{bd}}$ preserves $\mathcal{D}_5^{\mathrm{bd}}$	Lem. A.220.6
$\Phi_5^{(n)}$	$(3/4)^n$ with $\Phi_5^{(0)} = 1$	Thm. A.220.7
finite positivity	$\Phi_5^{(n)} > 0\ \forall n < \infty$	Lem. A.222.5
$\mathsf{LG}_5^{\mathrm{bd}}$	limit may not replace a finite stage	Def. A.222.4
ε-use rule	ε-smallness allowed, promotion not	Prop. A.222.6
$C5_{\mathrm{det/eff}}$	detector zero at $\theta = 1/256,\ n = 20$	r63–r64, compressed

Everything in this table is inherited. r86 adds no entry to it.

r86 threshold function

The detector threshold is currently recorded only at the single point $\theta = 1/256$, $n = 20$. It is a value of a function, and writing that function down costs nothing.

Theoremlet r86.1. For a detector threshold $\theta \in (0, 1)$, the least declared bounded stage at which the collar score falls to or below θ is

$$n_{\min}(\theta) = \left\lceil \log_{4/3} \frac{1}{\theta} \right\rceil = \left\lceil \frac{\ln(1/\theta)}{\ln(4/3)} \right\rceil .$$

In particular $n_{\min}(1/256) = 20$, and each halving of the threshold costs $\ln 2 / \ln(4/3) = 2.4094\ldots$ further stages.

Proof. By Theorem A.220.7, $\Phi_5^{(n)} = (3/4)^n$. Since $\ln(3/4) < 0$, the condition $(3/4)^n \le \theta$ is equivalent to $n \ln(3/4) \le \ln\theta$, that is $n \ge \ln(1/\theta)/\ln(4/3)$; the least such integer is the stated ceiling. For $\theta = 2^{-8}$ this is $\lceil 8 \ln 2 / \ln(4/3) \rceil = \lceil 19.276\ldots \rceil = 20$, which agrees with the inherited r63–r64 detector stage. The halving cost is $\log_{4/3} 2$. □

Remark. The agreement at $\theta = 1/256$ is a consistency check on the inherited chain, not a new claim: $(3/4)^{19} = 0.004228\ldots > 1/256$ while $(3/4)^{20} = 0.003171\ldots \le 1/256$.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

r86 finite-zero exclusion

Theoremlet r86.2. Let $\Phi^{(n+1)} = q\,\Phi^{(n)}$ be any purely multiplicative collar recursion with $q \in (0,1)$ and $\Phi^{(0)} > 0$. Then

$$\nexists\, N < \infty : \ \Phi^{(N)} = 0.$$

The conclusion depends only on $q > 0$, not on the particular value $q = 3/4$.

Proof. $\Phi^{(n)} = q^n \Phi^{(0)}$ is a product of finitely many strictly positive factors and is therefore strictly positive for every finite n. □

Why this is worth recording. The proof is immediate and the statement is already implicit in Remark A.220.8 and Lemma A.222.5. What has not been recorded is its *status consequence*. "No finite N with $\Phi_5^{(N)} = 0$" has been carried as the reason for an *open* entry, as though a finite closure might still be found by working harder inside the collar. Theoremlet r86.2 shows the opposite: inside the declared recursion the answer is settled, and settled negatively. Searching for a finite collar stage that discharges C5 is not a hard open problem but a provably empty search.

The exclusion is also robust in a useful way. Because it turns on $q > 0$ alone, no refinement of the contraction factor can help: replacing 3/4 by any smaller positive factor accelerates the decay and changes $n_{\min}$, but leaves the exclusion untouched.

r86 status refinement

The r83 classes are $\mathsf{Closed}_{\text{eff/local}}$, $\mathsf{Closed}_{\text{typed}}$ and $\mathsf{Open}_{\text{exact/global}}$. The last of these conflates two situations. We separate them:

$$\mathsf{Excluded}_{\text{rec}} \quad \text{versus} \quad \mathsf{Open}_{\text{undecided}}.$$

$\mathsf{Excluded}_{\text{rec}}$ means: inside the declared structure the question has an answer, and the answer is negative. $\mathsf{Open}_{\text{undecided}}$ means: the declared structure does not reach the question at all.

Applying this to $C5_{\text{zero}}$ splits it into two independent questions:

	question	r86 status
(a)	does $\mathcal{U}_5^{\text{bd}}$ reach zero at a finite stage?	$\mathsf{Excluded}_{\text{rec}}$
(b)	is there a route to $C5 = 0$ outside the multiplicative collar?	$\mathsf{Open}_{\text{undecided}}$

Thus the r86 C5 decision datum is

$$\mathsf{D}_{C5}^{r86} = \big(\mathsf{Excluded}_{\text{rec}},\ \mathsf{Open}_{\text{undecided}}\big).$$

This is a refinement of the record, not a change of result. $C5 = 0$ remains unproved and remains a forbidden assertion; question (b) is exactly the part that was never addressed, and it is now stated as the open question in its own right rather than hidden inside (a).

r86 search direction

Theoremlet r86.2 also says what a finite discharge would have to look like. Since the exclusion follows from multiplicativity together with positivity of the factor, any candidate mechanism must leave that form. The admissible shapes are:

shape	modification	what it would require
(i)	$\Phi^{(n+1)} = q\Phi^{(n)} - c,\ c > 0$	a declared subtractive channel with its own guard
(ii)	$\Phi^{(n+1)} = 0$ for $\Phi^{(n)} < \delta$	a truncation rule; not admissible under $\mathsf{LG}_5^{\text{bd}}$
(iii)	a non-collar object	a separate witness, unrelated to $\mathcal{U}_5^{\text{bd}}$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Shape (ii) is precisely what the r11 limit guard forbids, and r86 does not reopen it. Shape (i) would need a subtractive term that is not currently declared anywhere in the bounded–exterior–mixed grammar. Shape (iii) is the honest reading of question (b): a finite C5 discharge, if one exists, will not come from the collar iteration.

Nothing in r86 supplies any of the three. The layer states the search space; it does not enter it.

r86 ledger

The consolidation closes cleanly:

$$R^{r86}_{\mathrm{cons}} = (0, 0, 0, 0)$$

for the four consolidated items

$$(\text{inventory},\ \text{threshold function},\ \text{finite-zero exclusion},\ \text{status split}).$$

The exact/global ledger is unchanged, because question (b) remains unanswered:

$$R^{r86}_{\mathrm{exact/global}} = (1, 1, 1, 1, 1).$$

The first component keeps the value 1, now with a sharper reason: not "no finite closure has been found" but "no finite closure exists inside the declared recursion, and no route outside it has been declared".

r86 guard update

r86 spends no guard. It strengthens the justification of the standing C5 guard rather than consuming it:

$$R^{(r86,+)}_{\mathrm{glob}} = (0, 1, 1, 1, 1), \qquad \Gamma^{r86}_{\mathrm{guard}} = (1, 1, 1, 1, 1), \qquad R^{r86}_{\mathrm{eff/local}} = (0, 0, 0, 0, 0).$$

The layer does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure. The detector-threshold result $C5_{\mathrm{det/eff}}$ is unaffected and remains $\mathsf{Closed}_{\mathrm{eff/local}}$ at $\theta = 1/256$, $n = 20$.

r86 queue update

Branch Q_5 of the r48 queue is discharged as a bookkeeping task. The remaining guard-neutral branch is

$$Q_3 :\ \text{photon-clock selector-window calibration},$$

which would sharpen the provenance of the bound $\Delta_\gamma < 1/5$ carried by $W^{\mathrm{loc}}_{\mathrm{clk}}$. Q_4 remains inadmissible while $\beta_4 = 1/256$ is to stay unspent.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.7.0-r87 Source-Only: Photon-Clock Selector-Window Calibration

Purpose. This layer executes branch Q_3 of the r48 queue. The r15 finite window family is recorded as a table of three computed values; the rules behind that table are not written down in closed form. r87 extracts them, verifies the inherited numbers, and reduces the one remaining blocked relay edge to a single missing datum. As in r86, no new mathematical result is produced: everything below follows from the r15 extraction rule and the r16 lift computation.
Source-only status. This r87 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. It makes no spectral identification between Riemann zeros and photon ticks and does not touch the W_γ^{glob} guard.

r87 window inventory

The inherited photon-clock window data, from Section A.226:

overlap flux	$S_V = N_V + \kappa_V N_V(N_V-1)$
required gain	$G_V^{\mathrm{req}} = 16/S_V, \quad 16 = \Theta_v/a^2$ at $\Theta_v = 1/256,\ a = 1/64$
relay criterion	$G_V S_V \geq 16$
window	$I_V = \{\tau : \mathcal{C}_\gamma(\tau) \geq G_V^{\mathrm{req}}\}$
half-width	$\delta_V = \sigma_V \arccos\sqrt{(G_V^{\mathrm{req}}-1)/(G_V^{\mathrm{avail}}-1)}$

r87 window-existence criterion

Theoremlet r87.1. A chart V has a nonempty zero-clock phase window exactly when its overlap flux lies in the half-open interval

$$\frac{16}{G_V^{\mathrm{avail}}} \;\leq\; S_V \;<\; 16.$$

The upper end is the direct/assisted boundary: for $S_V \geq 16$ the chart needs no clock window at all. The lower end is the assisted/blocked boundary.

Proof. The r15 extraction condition is $1 < G_V^{\mathrm{req}} \leq G_V^{\mathrm{avail}}$ with $G_V^{\mathrm{req}} = 16/S_V$. The left inequality $1 < 16/S_V$ is equivalent to $S_V < 16$; the right inequality $16/S_V \leq G_V^{\mathrm{avail}}$ is equivalent to $S_V \geq 16/G_V^{\mathrm{avail}}$. □

Verification against the inherited table.

chart	S_V	admissible band	in band	δ_V/σ_V
$V3$	10	$[8, 16)$	yes	0.684719
$V5$	10	$[32/5, 16)$	yes	0.886077
$V8$	9	$[8, 16)$	yes	0.490883

The three half-widths were recomputed from the r15 formula in exact rational ratios 3/5, 2/5, 7/9 and agree with the inherited values to six decimals.

Remark. The half-width is monotone in both arguments: it tends to $\pi/2\,\sigma_V$ as $G_V^{\mathrm{req}} \to 1^+$, and it closes to zero as $G_V^{\mathrm{req}} \to G_V^{\mathrm{avail}}$. A chart sitting at the lower band edge therefore has a window in principle but no timing margin in practice. Of the three extracted charts, $V8$ is closest to that edge.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

r87 single-packet lift formula

The r16 shadow-basin lift moved $N_6 : 3 \to 4$ at fixed $\kappa_6 = 3/4$, $G_6 = 3/2$, which raised S_6 from 15/2 to 13. That step is an instance of a general rule.

Theoremlet r87.2. Adding one admissible packet at fixed κ changes the overlap flux by

$$\Delta S(N \to N+1) = 1 + 2\kappa N.$$

Proof. $S(N+1) - S(N) = \big[(N+1) + \kappa(N+1)N\big] - \big[N + \kappa N(N-1)\big] = 1 + 2\kappa N$. □

For the r16 lift this gives $1 + 2 \cdot \frac{3}{4} \cdot 3 = \frac{11}{2}$, and indeed $15/2 + 11/2 = 13$. The gain is quadratic in N: a single packet is worth more at a populated front than at a sparse one.

r87 blocked-chart ranking

r15 ranks the blocked charts by the gain ratio $G^{\mathrm{req}}/G^{\mathrm{avail}}$ and concludes that $V6$ is closest. That ranking is correct for its own measure, but it is not the only measure, and the two do not agree:

chart	S_V	G_V^{avail}	$G^{\mathrm{req}}/G^{\mathrm{avail}}$	missing S
$V0$	4	1	4	12
$V6$ (pre-r16)	15/2	3/2	$64/45 \approx 1.422$	$19/6 \approx 3.167$
$V9$ (pre-r17)	3	3	$16/9 \approx 1.778$	$7/3 \approx 2.333$

By gain ratio the order is $V6 < V9 < V0$; by missing overlap flux it is $V9 < V6 < V0$. The two measures correspond to two different lift mechanisms: raising the available clock gain acts multiplicatively, adding packets acts through S. Both r16 and r17 in fact proceeded through S, by adding a single packet, so the additive column is the one that governed the executed moves. It also explains why $V9$ yielded to the smaller lift: it needed 7/3 where $V6$ needed 19/6.

r87 lift formula against the recorded V9 micro-lift

The chart parameters are recorded in the r14 scan ledger, which lists $(N_9, \kappa_9, G_9) = (2, \frac{1}{2}, 3)$, hence $S_9 = 3$ and $G_9 S_9 = 9 < 16$. The blocked edge $V8 \to V9$ needs $\Delta S \geq 7/3$ by the ranking table above, and Theoremlet r87.2 gives for a single added packet

$$\Delta S = 1 + 2\kappa_9 N_9 = 1 + 2 \cdot \tfrac{1}{2} \cdot 2 = 3 \;\geq\; \tfrac{7}{3}.$$

So one packet suffices, with $S_9' = 6$ and $G_9 S_9' = 18 \geq 16$.

This is exactly the move carried out in Section A.228: the r17 guarded micro-lift takes $(2, \frac{1}{2}, 3) \mapsto (3, \frac{1}{2}, 3)$ and records $H_9' = 18 > 16$ with margin $\mu_9' = 2$. The r87 formula therefore reproduces an already executed lift from the chart parameters alone, without reference to the r17 computation.

Status of the chain. After the r16 lift of $V6$, the r17 atlas-gap bridge $V6 \to V7$ and the r17 micro-lift of $V9$, the finite chart chain runs $V1 \to \cdots \to V9$. The only chart outside it is $V0$, which needs $\Delta S = 12$ — four times the largest lift so far executed — and which r15–r17 keep explicitly guarded. The calibration confirms that ordering: by the additive measure $V0$ is far outside the range in which single-packet moves operate.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

r87 provenance gap for $\Delta_\gamma < 1/5$

The stated aim of Q_3 was to sharpen the provenance of the bound $\Delta_\gamma < 1/5$ carried by $W_{\mathrm{clk}}^{\mathrm{loc}}$. That bound occurs in the present source only inside summary and guard layers, never with a derivation: it enters through the compressed r55–r80 chain, where the detailed r65 step has been replaced by the chain invariant. The uncompressed log is not part of this source packet.

Accordingly r87 does *not* discharge that part of Q_3. What can be said from the present source is negative and worth recording:

$$\Delta_\gamma < 1/5 \notin \{\text{quantities derivable from the r15–r16 window layer}\}.$$

The r15 layer produces the ratios 3/5, 2/5, 7/9, the gains 8/5, 16/9, 32/15, 16/3 and the threshold constant 16; none of these yields 1/5, and the window half-widths are measured in units of σ_V rather than as a bound on a frequency spread. The bound therefore belongs to a different layer and must be recovered from the uncompressed r65 text, not reconstructed from the finite chart family.

r87 ledger

The calibration closes in three of four items:

$$R_{\mathrm{cal}}^{r87} = (0, 0, 0, 1)$$

for

$$(\text{window-existence criterion},\ \text{lift formula},\ \text{blocked-chart ranking},\ \Delta_\gamma \text{ provenance}).$$

The fourth component stays at 1: the bound $\Delta_\gamma < 1/5$ remains an inherited constant whose derivation is outside this source. This is a recorded documentation gap, not a mathematical obstruction, and it does not affect the standing of $W_{\mathrm{clk}}^{\mathrm{loc}}$, which is carried as a conditional local statement.

r87 guard update

r87 spends no guard:

$$R_{\mathrm{glob}}^{(r87,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{r87} = (1, 1, 1, 1, 1), \qquad R_{\mathrm{eff/local}}^{r87} = (0, 0, 0, 0, 0).$$

All statements are finite, local and chart-bound. The layer does not identify Riemann zeros with photon ticks, does not prove a global photon-clock/RH theorem, does not prove $C6/T2$, and does not remove the all-readout blocker. The relay chains remain finite: the longest certified chain is the length-six chain of Corollary A.227.4.

r87 queue update

Branch Q_3 is discharged in its calibration part and left open in its provenance part. The r48 queue after r87:

branch	content	status
Q_1	local field-interface elaboration	done (r49–r54)
Q_2	theoremlet-to-lemma packaging	done (r55–r80)
Q_3	photon-clock selector-window calibration	calibrated (r87); Δ_γ provenance open
Q_4	C4 mixed-kernel localization	inadmissible, β_4 unspent
Q_5	C5 asymptotic-collar bookkeeping	done (r86)

One concrete item remains that needs no new guard: the uncompressed r65 text carrying the derivation of $\Delta_\gamma < 1/5$.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.7.0-r88 Source-Only: Admissibility Test for Non-Collar Routes to Finite C5 Discharge

Purpose. r86 split the $C5_{\text{zero}}$ entry into question (a), settled negatively inside the declared multiplicative collar recursion, and question (b): whether a route to a finite $C5$ discharge exists *outside* that recursion. r86 listed three candidate shapes without examining them. This layer examines them. Two are eliminated; the third is not, and question (b) remains open. Nothing here asserts that a finite discharge exists.
Source-only status. This r88 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The forbidden assertion $C5 = 0$ is not made, and no guard is spent. The analysis is conditional throughout: it asks what a candidate mechanism would have to satisfy, not what holds.

r88 subtractive channel: closed form

Shape (i) of the r86 table is the constant-subtraction recursion

$$\Phi^{(n+1)} = q\,\Phi^{(n)} - c, \qquad q \in (0,1),\ c > 0,\ \Phi^{(0)} > 0.$$

Theoremlet r88.1. This recursion has the closed form

$$\Phi^{(n)} = q^n\left(\Phi^{(0)} + \frac{c}{1-q}\right) - \frac{c}{1-q},$$

and it attains the value zero at an integer stage N if and only if

$$c = c_\star(N) := \frac{(1-q)\,q^N\,\Phi^{(0)}}{1-q^N}.$$

For $q = 3/4$, $\Phi^{(0)} = 1$ this is

$$c_\star(N) = \frac{1}{4\big((4/3)^N - 1\big)}.$$

Proof. Writing $\psi^{(n)} = \Phi^{(n)} + c/(1-q)$ turns the recursion into $\psi^{(n+1)} = q\psi^{(n)}$, whence $\psi^{(n)} = q^n\psi^{(0)}$ and the displayed form. Setting $\Phi^{(N)} = 0$ gives $q^N(\Phi^{(0)} + c/(1-q)) = c/(1-q)$; solving for c yields $c_\star(N)$. Conversely $c = c_\star(N)$ returns $\Phi^{(N)} = 0$ by substitution. □

Verified instances. For $q = 3/4$, $\Phi^{(0)} = 1$: $c_\star(1) = 3/4$, $c_\star(2) = 9/28$, $c_\star(3) = 27/148$, $c_\star(4) = 81/700$, $c_\star(5) = 243/3124$. Each was checked by running the recursion in exact rational arithmetic; each reaches zero at its own stage and at no earlier one.

Remark on genericity. For $c \neq c_\star(N)$ for every N, the orbit does not meet zero at all: it passes from positive to negative between two consecutive stages. Sample runs with $c = 1/10, 1/5, 1/4, 3/10$ cross zero at stages $5, 3, 3, 3$ respectively without ever taking the value zero. Exact finite discharge by a constant subtractive channel therefore requires the channel strength to be tuned to the target stage — a measure-zero condition on c, not a structural feature.

r88 the admissibility obstruction

Tuning is not the decisive objection. The decisive objection is what happens immediately afterwards.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Theoremlet r88.2. Let $c = c_\star(N)$, so that $\Phi^{(N)} = 0$. Then

$$\Phi^{(N+1)} = -c < 0,$$

and $\Phi^{(n)} < 0$ for all $n > N$. Hence the orbit leaves the bounded collar domain $\mathcal{D}_5^{\mathrm{bd}}$ at the stage immediately after the discharge.

Proof. $\Phi^{(N+1)} = q \cdot 0 - c = -c < 0$. For $n > N$ the closed form of Theoremlet r88.1 is strictly decreasing once negative, since $q\Phi - c < \Phi$ whenever $\Phi < c/(1-q)$, which holds for all $\Phi \le 0$. Admissibility of $\mathcal{U}_5^{\mathrm{bd}}$ on $\mathcal{D}_5^{\mathrm{bd}}$ was established in Lemma A.220.6 precisely through preservation of positivity; a map whose next iterate is negative fails that condition. □

Corollary r88.3 (shape (i) collapses into shape (ii)). A subtractive channel can deliver a finite discharge only if it is accompanied by a stopping rule that halts the recursion at stage N instead of continuing into the negative range. Such a rule is a truncation, which is shape (ii) of the r86 table, and truncation is forbidden by the bounded-collar limit guard $\mathsf{LG}_5^{\mathrm{bd}}$ of Definition A.222.4. Shapes (i) and (ii) are therefore not two independent candidates but one, and it is already excluded by a standing guard.

r88 the free-parameter trap

One might weaken shape (i) to a stage-dependent channel $\Phi^{(n+1)} = q\Phi^{(n)} - c_n$. This does produce a finite zero: choosing $c_N = q\Phi^{(N)}$ sets $\Phi^{(N+1)} = 0$ for any prescribed N.

Remark r88.4. That freedom is the objection, not the solution. A family with an unconstrained sequence (c_n) can realize *any* prescribed orbit whatsoever, including orbits that violate every other guard in the ledger. A mechanism that can produce the desired conclusion and its negation with equal ease carries no information about $C5$; it records the desired answer rather than deriving it. An admissible stage-dependent channel would therefore have to fix (c_n) by an independent structural rule — and it is that rule, not the resulting zero, that would constitute the mathematical content. No such rule is declared anywhere in the bounded–exterior–mixed grammar.

r88 what remains: shape (iii)

Shape (iii) is the only surviving candidate: an object unrelated to the collar iteration. r88 does not supply one. What it can do is state the burden such an object would carry.

	requirement on a shape-(iii) witness
(α)	it must exhibit the vanishing of the $C5$ obstruction itself, not the smallness of the score Φ_5
(β)	it must be independent of $\mathcal{U}_5^{\mathrm{bd}}$, since by r86 that recursion cannot deliver the conclusion
(γ)	it must remain compatible with $C5_{\mathrm{det/eff}}$, which is already established at $\theta = 1/256,\ n = 20$
(δ)	it must not import the all-readout slot, since $0_{\mathrm{termdata}} \not\Rightarrow 0_{\mathrm{all}}$ remains frozen

Requirement (α) is the substantive one. The collar layer measures $C5$ through a score whose finite values are positive and whose limit is zero. Any statement about $C5$ obtained *through that score* is therefore a statement about the limit, which is exactly what the standing

© 2026 Marc Brendecke. Preprint — not peer reviewed.

guard separates from finite discharge. A shape-(iii) route must reach the obstruction by some other access; otherwise it re-enters the collar layer and inherits its verdict.

r88 ledger

Of the three r86 shapes, two are now closed and one is not:

$$\mathsf{D}^{r88}_{(b)} = \big(\text{(i) excluded, (ii) excluded, (iii) open}\big), \qquad R^{r88}_{(b)} = (0,0,1).$$

Question (b) of the r86 split therefore remains $\mathsf{Open}_{\text{undecided}}$, but its search space is narrower: it is no longer "some mechanism outside the collar" but specifically a witness meeting (α)–(δ), with the two parametric modifications of the collar recursion ruled out.

r88 guard update

r88 spends no guard. It closes two candidate routes and leaves every ledger value where it stood:

$$R^{(r88,+)}_{\text{glob}} = (0,1,1,1,1), \qquad \Gamma^{r88}_{\text{guard}} = (1,1,1,1,1), \qquad R^{r88}_{\text{exact/global}} = (1,1,1,1,1).$$

Corollary r88.3 strengthens $\mathsf{LG}^{\text{bd}}_5$ by showing that the most natural way around it collapses back into it. The layer does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.7.0-r89 Source-Only: r87 Correction — Recorded V9 Data and Executed Micro-Lift

Purpose. The r87 layer, as first written, asserted that the decomposition of the $V9$ overlap flux was not recorded in this source and that $V8 \to V9$ was the last blocked relay edge. Both assertions were wrong. This layer records the correction and its cause.
Source-only status. This r89 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release.

r89 what was wrong

r87 assertion	status	actual location
"(N_9, κ_9) not recorded"	false	r14 chart ledger: $(2, \frac{1}{2}, 3)$
"$V8 \to V9$ is the only blocked edge"	false	r17 stitched it
"no claim which case holds"	moot	decided in r17

The r14 finite valley-front chart scan lists all ten charts with packet count, coherence parameter and available gain; $V9$ appears there as $(N, \kappa, G) = (2, \frac{1}{2}, 3)$ with clock score 9/16 and status blocked. The r17 guarded micro-lift then performed $(2, \frac{1}{2}, 3) \mapsto (3, \frac{1}{2}, 3)$, obtaining $H_9' = 18 > 16$ with margin $\mu_9' = 2$, and stitched the chain to $V1 \to \cdots \to V9$. Both facts stand in the same document as r87.

r89 what was right

The defect was in the source survey, not in the calculation. The r87 single-packet lift formula

$$\Delta S = 1 + 2\kappa N$$

applied to the r14 parameters gives $\Delta S = 3$, hence $S_9' = 6$ and $G_9 S_9' = 18$ — precisely the values r17 reports. The formula therefore reproduces the executed micro-lift from the chart parameters alone. What r87 presented as an open alternative was in fact a correct prediction of a step already taken.

The corrected r87 text records this as a verification rather than as a question. The r87 ledger is unchanged: $R_{\mathrm{cal}}^{r87} = (0, 0, 0, 1)$, whose single open component was always the Δ_γ provenance and never the $V9$ datum.

r89 cause and rule

The survey behind r87 read r15 and r16 and inferred from the absence of the decomposition *there* that it was absent from the document. It did not read r14, one round earlier, where the chart ledger is tabulated, nor r17, one round later, where the lift is executed.

Rule r89. A claim that a datum is missing from this source is a claim about the whole monolith and must be checked against the rounds adjacent to the one being read, not only against the round in which the datum is used. Absence in the round of use is evidence of nothing: the solve chain habitually tabulates in one round and consumes in another.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

r89 ledger and provenance

The correction is editorial and touches no result:

$$R^{r89}_{\mathrm{corr}} = (0, 0, 0)$$

for (chart datum restored, chain status restored, prediction reclassified as verification).

RC22 was packaged before the correction and therefore carries the erroneous r87 text. It is retained unchanged as provenance; the present source and any later release candidate carry the corrected text. No earlier round is modified by r89.

r89 guard update

r89 spends no guard and changes no ledger value:

$$R^{(r89,+)}_{\mathrm{glob}} = (0, 1, 1, 1, 1), \qquad \Gamma^{r89}_{\mathrm{guard}} = (1, 1, 1, 1, 1).$$

The finite relay chain $V1 \to \cdots \to V9$ is inherited from r17 and is not extended here. $V0$ remains outside it and guarded. The layer does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.7.0-r90 Source-Only: Reconstruction Attempt for the Bound $\Delta_\gamma < 1/5$

Purpose and status. r87 recorded that the bound $\Delta_\gamma < 1/5$ carried by $W_{\mathrm{clk}}^{\mathrm{loc}}$ has no derivation in this source: it enters through the compressed r55–r80 chain, and the detailed r65 text is not part of the source packet. This layer attempts a reconstruction from the material that *is* present. It arrives at a bound of the right size from the r15 clock model. **This is a reconstruction, not a recovery.** It does not establish that r65 argued this way, and it does not replace the missing text. The provenance component of R_{cal}^{r87} stays open.

r90 what the source fixes

Three ingredients are stated in Section A.226. The photon-clock probe is

$$\mathcal{C}_\gamma(\tau) = 1 + \eta\,\frac{\left|\sum_{m=1}^{M} w_m e^{i\gamma_m \tau}\right|^2}{\left(\sum_{m=1}^{M} |w_m|\right)^2},$$

its local normalized form is

$$G_V(\tau) = 1 + (G_V^{\mathrm{avail}} - 1)\cos^2\!\Big(\frac{\tau - \tau_V}{\sigma_V}\Big),$$

so that $\eta = G_V^{\mathrm{avail}} - 1$, and window membership requires $\mathcal{C}_\gamma(\tau) \geq G_V^{\mathrm{req}}$.

r90 coherence bound

Theoremlet r90. Let the clock phases be $\varphi_m = \gamma_m \tau$, centred so that $\langle\varphi\rangle = 0$, and write $\mathrm{Var}(\varphi)$ for their weighted variance. To first order in the spread,

$$\frac{\left|\sum_m w_m e^{i\varphi_m}\right|^2}{\left(\sum_m |w_m|\right)^2} = 1 - \mathrm{Var}(\varphi) + O\big(\mathrm{Var}^2\big),$$

and the selector is preserved on chart V exactly while

$$\mathrm{Var}(\varphi) \;\leq\; 1 - \frac{G_V^{\mathrm{req}} - 1}{G_V^{\mathrm{avail}} - 1} \;=\; 1 - \rho_V,$$

where ρ_V is the ratio already tabulated in the r15 half-width formula.

Proof. Expanding $\langle e^{i\varphi}\rangle = 1 + i\langle\varphi\rangle - \frac{1}{2}\langle\varphi^2\rangle + O(\varphi^3)$ with $\langle\varphi\rangle = 0$ gives $\langle e^{i\varphi}\rangle = 1 - \frac{1}{2}\mathrm{Var} + O(\mathrm{Var}^2)$, hence $|\langle e^{i\varphi}\rangle|^2 = 1 - \mathrm{Var} + O(\mathrm{Var}^2)$. Substituting into $\mathcal{C}_\gamma \geq G^{\mathrm{req}}$ with $\eta = G^{\mathrm{avail}} - 1$ yields $1 + (G^{\mathrm{avail}} - 1)(1 - \mathrm{Var}) \geq G^{\mathrm{req}}$, which rearranges to the display. □

r90 the resulting numerical bound

Applied to the three charts that actually carry windows:

chart	ρ_V	$1 - \rho_V$	decimal
$V3$	3/5	2/5	0.4000
$V5$	2/5	3/5	0.6000
$V8$	7/9	2/9	0.2222

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A bound valid for the whole family is the minimum,

$$\Delta_\gamma^{\max} = \min_V \bigl(1 - \rho_V\bigr) = \frac{2}{9} = 0.2222\ldots,$$

attained at the narrowest window $V8$. The recorded bound $1/5 = 0.2$ sits just below it, with a margin of ten percent:

$$\frac{1}{5} < \frac{2}{9}, \qquad 1 - \frac{1/5}{2/9} = \frac{1}{10}.$$

Assessment. A conservative rounding of 2/9 to the nearest simple fraction below is 1/5, and that matches the rounding discipline used elsewhere in this document, where thresholds are carried as 1/256 and 1/64 rather than as decimals. The reconstruction therefore produces a bound of the correct magnitude, at the correct place in the argument, with a plausible safety margin.

r90 rejected alternative

One numerical coincidence in the same table should be recorded and dismissed:

$$\rho_{V3} - \rho_{V5} = \frac{3}{5} - \frac{2}{5} = \frac{1}{5}.$$

This reproduces the constant exactly rather than approximately, but it carries no mechanism: a difference of two window ratios is not a spread of clock frequencies, it does not depend on γ at all, and nothing in the grammar would make such a difference a preservation threshold. It is rejected as coincidence.

r90 limits of the reconstruction

Three gaps remain and are stated rather than papered over:

(α)	Δ_γ is nowhere defined in this source; whether it denotes a variance, a standard deviation or a maximal deviation is not recorded
(β)	the expansion is first order and holds only for small spread
(γ)	the ten-percent margin is plausible but not derived

Under reading (α) as a standard deviation the bound would instead compare against $\sqrt{2/9} = 0.471$, and 1/5 would be far more conservative than necessary; under the variance reading it is a close fit. That is evidence for the variance reading, not proof.

r90 ledger and guard update

The reconstruction is recorded as such:

$$R_{\mathrm{rec}}^{r90} = (0, 0, 1)$$

for (coherence bound derived, numerical match established, r65 provenance still missing). The third component stays at 1: only the original text can close it.

r90 spends no guard:

$$R_{\mathrm{glob}}^{(r90,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{r90} = (1, 1, 1, 1, 1).$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

All statements are local and chart-bound. The layer makes no spectral identification between Riemann zeros and photon ticks, and it does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure. $W_{\mathrm{clk}}^{\mathrm{loc}}$ continues to be carried as a conditional local statement under its inherited bound, whose status as an inherited constant is unchanged by r90.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.7.0-r91 Source-Only: Q4 Executed — Why β_4 Cannot Be Spent

Purpose. Branch Q_4 of the r48 queue has been carried as *inadmissible* since r54, with the reason recorded as "$\beta_4 = 1/256$ remains unspent". That reads as a policy: a guard held back by choice. This layer executes the branch and finds that it is not a policy. The attempt terminates for a structural reason already present in the source, and the guard survives not because it is protected but because spending it would be the very operation the matter-frame typing forbids. **No guard is spent here.**
Source-only status. This r91 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release.

r91 what C4 still is

The first finding is that $C4$ is no longer a commutator problem. The r07 split test (Lemma A.218.7) decomposes

$$S_4^{\mathrm{comm}} \rightsquigarrow S_{4,-}^{\mathrm{comm}} \oplus S_{4,+}^{\mathrm{comm}}, \qquad S_{4,-}^{\mathrm{comm}} : \mathsf{routed}_{\mathrm{comm}}, \quad S_{4,+}^{\mathrm{comm}} : \mathsf{zero}_{2s},$$

and neither component carries active leakage afterwards. The antisymmetric mixed-transport part is not contracted away; it is *read through a declared commutator comparison channel*, and the source is explicit that this "does not mean a global commutator vanishing theorem". The active support reduces to a single class,

$$\mathrm{Supp}_4^{r07} = \{S_{4,\mathrm{out}}^{\mathrm{collar}}\},$$

the component that crosses the bounded–exterior interface. What remains of $C4$ is therefore a boundary-crossing seam, not a commutator remainder.

r91 what r13 already settled

Lemma A.223.2 tests absorbability of that residual kernel and finds it negative: it is not absorbable by the declared bounded, comparison-routed or selector-routed channels. The same layer enumerates what an absorption would require — a new declared comparison tensor, a new selector, or an identification with bounded $C5$ contraction debt — and records that the first two are absent from the datum while the third is forbidden by the separation rule.

So Q_4 cannot proceed by absorption inside the present grammar. That much was known. The question r91 asks is different: what would it *mean* to spend β_4?

r91 the threshold identity

The bound $\beta_4 = 1/256$ does not occur once. The source states in Definition A.224.18

$$\Theta_v := \beta_4^{r12} = \frac{1}{256},$$

making the residual $C4$ bound and the local ridge-front relay threshold the same quantity by definition. The detector chain carries the same value as $\theta_{\mathrm{det}} = 1/256$, and r81 types it explicitly:

$$\varepsilon_{\mathsf{M}}^{\mathrm{len}} = \frac{\ell_{\mathrm{P}}}{L_{\mathrm{ref}}} \qquad \text{or, dimensionlessly,} \qquad \theta_{\mathrm{det}} = \frac{1}{256},$$

with the accompanying sentence *"These are terminal scales, not exact all-readout zeros."*

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Two readings are possible and should be distinguished. Either 1/256 is one terminal scale appearing in three roles — seam bound, relay threshold, detector threshold — or it is three quantities that happen to share a dyadic value. The source settles the first two by definition; for the third it supplies equality of value and equality of role, but no explicit identification. The argument below needs only the first two plus the r81 typing.

r91 why spending is not available

Theoremlet r91. Spending β_4 — declaring the residual seam absorbed — requires treating a residual at the terminal scale as an exact zero. Under the matter-frame typing of r81 that inference is unavailable:

$$\Phi \le \varepsilon_{\mathsf{M}} \implies 0^{\varepsilon}_{\mathsf{M}} \implies 0_{\mathrm{termdata}}, \qquad \text{but} \qquad 0^{\varepsilon}_{\mathsf{M}} \not\Rightarrow 0_{\mathrm{all}}.$$

Hence Q_4 is not blocked by a missing tensor or an unbuilt selector. It is blocked by the same typing that blocks $C5 = 0$, and no refinement of the localization can remove that block.

Proof. The residual seam is bounded by β_4, and $\beta_4 = \Theta_v$ is a terminal threshold of the declared chain. A spending step asserts that the seam contributes nothing, that is $S^{\mathrm{collar}}_{4,\mathrm{out}} = 0$ as an all-readout statement. What the bound supports is $S^{\mathrm{collar}}_{4,\mathrm{out}} \le \beta_4$, which by Theoremlet r81 yields $0^{\varepsilon}_{\mathsf{M}}$ and terminal data, and explicitly not 0_{all}. The step from the bound to the assertion is exactly the forbidden promotion. □

r91 status reclassification

Using the r86 vocabulary, the queue entry changes class without changing value:

$$Q_4 : \quad \mathsf{inadmissible}_{\mathrm{policy}} \longmapsto \mathsf{Excluded}_{\mathrm{typed}}.$$

The practical difference matters for anyone planning further work. Under the old reading, Q_4 was a branch one might open later, once a suitable comparison tensor had been built. Under the corrected reading, building that tensor would not help: the obstruction is not the absence of a channel but the typing of the scale at which the residual sits. A future $C4$ closure, if one exists, must come from a route that does not pass through a terminal-scale bound — structurally the same requirement that r88 placed on shape (iii) for $C5$.

r91 physical reading

The formal result has a reading in the physical layer that the document otherwise keeps separate, and it is recorded here as interpretation, not as a claim.

The commutator component was routed rather than eliminated. In the compensator language of the main line, where $C = \frac{1}{2}(\mathrm{id}+J)$ splits the space into symmetric and antisymmetric parts, an antisymmetric transport remainder is not something a change of chart removes; it is read through a comparison channel and kept. That is the expected behaviour of a noncommutative structure, and r07 treats it accordingly.

What survives as $C4$ is a seam at the bounded–exterior interface, bounded by the terminal resolution. Read physically, the statement of r91 is that a residual sitting at the resolution scale of the readout cannot be declared absent by that same readout. This is the same discipline that r81 expresses by pairing θ_{det} with $\ell_{\mathrm{P}}/L_{\mathrm{ref}}$: below the terminal scale the frame records a zero, and that record is a statement about the frame, not about the object.

No claim is made that β_4 equals a physical constant, that 1/256 has dimensional meaning, or that the seam corresponds to a measurable quantity.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

r91 ledger and guard update

Q_4 is executed and terminates negatively:

$$R^{r91}_{Q_4} = (0, 0, 1)$$

for (commutator component already routed, absorbability already settled, spending shown unavailable). The third component is recorded as 1 because a negative result closes the branch, not the obstruction: $C4$ remains open in the exact/global ledger.

The guard vector is unchanged, and pointedly so:

$$R^{(r91,+)}_{\text{glob}} = (0, 1, 1, 1, 1), \qquad \Gamma^{r91}_{\text{guard}} = (1, 1, 1, 1, 1), \qquad \beta_4 = \frac{1}{256} \text{ unspent.}$$

β_4 remains unspent not as a reserve held back, but because the spending operation is not available in the declared typing. The layer does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.7.0-r92 Source-Only: Sharpened Reconstruction of $\Delta_\gamma < 1/5$

Purpose. r90 reconstructed the bound $\Delta_\gamma < 1/5$ to within a rounding: the coherence argument gave $2/9 = 0.2222\ldots$, and $1/5$ was read as a conservative rounding of it. This layer replaces that with an exact account. The constant is not a rounding. It is the relative gain reserve of a single named chart, and it can be traced through the arithmetic that produces it. As in r90 this is a reconstruction, not a recovery of the r65 text, which a fourth search has again failed to locate.

r92 the arithmetic signature

Every threshold in the declared chain is dyadic:

$$\beta_4 = \Theta_v = \theta_{\mathrm{det}} = \frac{1}{256} = 2^{-8}, \qquad a = \frac{1}{64} = 2^{-6}, \qquad \frac{\Theta_v}{a^2} = 16 = 2^4.$$

The bound $\Delta_\gamma < 1/5$ is the only non-dyadic constant in the layer. A factor of five cannot arise from products and quotients of powers of two, so it must be produced by a division that introduces it. In the finite chart family there is exactly one such division. The overlap flux of $V3$ is

$$S_{V3} = N + \kappa N(N-1) = 5 + \tfrac{1}{4} \cdot 5 \cdot 4 = 10,$$

and the required gain is

$$G_{V3}^{\mathrm{req}} = \frac{16}{S_{V3}} = \frac{16}{10} = \frac{8}{5}.$$

The five enters here, by cancellation of 16/10, and nowhere else: $V5$ shares $S = 10$ and therefore the same 8/5, while every other chart yields a denominator free of five.

r92 the perturbation picture

r90 let the perturbation act on the coherence factor K inside $\mathcal{C}_\gamma = 1 + \eta K$, giving $\mathrm{Var}(\varphi) \leq 1 - \rho_V$. A second reading is available and is the one the constant selects. At full coherence $\mathcal{C}_\gamma = G_V^{\mathrm{avail}}$; a frequency spread lowers the probe value itself, and the window survives while the probe stays above G_V^{req}. The admissible *relative* drop is therefore

$$\Delta_\gamma^{\max}(V) = 1 - \frac{G_V^{\mathrm{req}}}{G_V^{\mathrm{avail}}}.$$

Theoremlet r92. For the charts carrying windows, the admissible relative drop is

chart	chain	G^{req}	G^{avail}	$1 - G^{\mathrm{req}}/G^{\mathrm{avail}}$
$V3$	$\mathcal{R}_A$	8/5	2	1/5
$V5$	$\mathcal{R}_A$	8/5	5/2	9/25
$V8$	$\mathcal{R}_B$	16/9	2	1/9

Minimized over the primary relay chain $\mathcal{R}_A$ of Section A.226, this is

$$\min_{V \in \mathcal{R}_A} \left(1 - \frac{G_V^{\mathrm{req}}}{G_V^{\mathrm{avail}}}\right) = \min\left\{\frac{1}{5}, \frac{9}{25}\right\} = \frac{1}{5},$$

attained at $V3$, which is the weakest assisted chart of that chain.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof. Substitute the r14 parameters into $S = N + \kappa N(N-1)$ and $G^{\mathrm{req}} = 16/S$, then form the quotient. For $V3$: $S = 10$, $G^{\mathrm{req}} = 8/5$, $G^{\mathrm{req}}/G^{\mathrm{avail}} = 4/5$, so the admissible drop is $1/5$. For $V5$: $G^{\mathrm{req}}/G^{\mathrm{avail}} = 16/25$, giving $9/25 > 1/5$. The minimum over the two is $1/5$. □

r92 comparison with r90

The two reconstructions differ in where the perturbation is applied and in how well they fit:

	value on $\mathcal{R}_A$	fit to $1/5$
r90, spread acts on coherence K	$2/5$	no
r90, minimum over all three charts	$2/9$	by rounding
r92, spread lowers the probe relatively	$1/5$	exact

An exact hit through a chain of four elementary steps — packet count, overlap flux, required gain, relative reserve — is considerably stronger evidence than agreement after rounding. r90 is therefore superseded as an account of the constant, though its coherence bound remains correct as a statement in its own right.

r92 what remains uncertain

Two points are not settled by the fit.

First, the minimum is taken over $\mathcal{R}_A$ alone. Including $V8$, which belongs to the separate chain $\mathcal{R}_B$ in the r15 stitching, would give $1/9$ instead. The restriction is consistent with the guard being written $W^{\mathrm{loc}}_{\mathrm{clk}}$, a local statement, and with the two chains being separate at the time the windows were extracted; but the source does not state it.

Second, the linear response of $\mathcal{C}_\gamma$ to the frequency spread is assumed, not derived. What the fit establishes is that the constant $1/5$ is the relative gain reserve of $V3$; it does not establish that the original argument reached it by this route.

r92 ledger and guard update

The reconstruction improves but does not close the provenance entry:

$$R^{r92}_{\mathrm{rec}} = (0, 0, 1)$$

for (arithmetic origin of the five located, exact value derived, r65 text still absent). The third component can only be closed by the original text.

r92 spends no guard:

$$R^{(r92,+)}_{\mathrm{glob}} = (0, 1, 1, 1, 1), \qquad \Gamma^{r92}_{\mathrm{guard}} = (1, 1, 1, 1, 1).$$

All statements are finite, local and chart-bound. The layer makes no spectral identification between Riemann zeros and photon ticks and does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.7.0-r93 Source-Only: The Two-Sided Typing Read as a Routing Grammar

Purpose. The two-sided structural tensor notation of Definition H2.40 has been in continuous use since r26 without being named for what its index slots actually carry. Read from the definition alone, the front indices look like role annotations attached to an otherwise ordinary tensor display. Read from the sites that use them, they are addresses in a labelled graph, and the contraction discipline is the forwarding rule of that graph. This layer records the reading. It introduces no object, no new contraction and no new claim; it names a structure already present in the source since r26.

r93 what the index slots carry

The decisive evidence is in the application layer, not in the definition layer. The r26 transition typing (§A.237.2) types four atlas cells as

$$ {}^{\partial\mathrm{src}}{}_{\mathrm{loc}}\mathcal{A}_{0\,\mathrm{relay}}^{V0}, \qquad {}^{\mathrm{core}}{}_{\mathrm{loc}}\mathcal{A}_{1\,\mathrm{relay}}^{V5}, \qquad {}^{\mathrm{shadow}}{}_{\mathrm{loc}}\mathcal{A}_{2\,\mathrm{bridge}}^{V7}, \qquad {}^{\gamma\mathrm{clock}}{}_{\mathrm{loc}}\mathcal{A}_{3\,\mathrm{tail}}^{V9}. $$

The back upper slot carries a chart name from the finite scan of §A.225; the back lower slot carries an edge kind, one of relay, bridge, tail. The accompanying r26 audit ledger lists the transitions $\tau_{01}, \tau_{12}, \tau_{23}$ by source type, target type and status. Node, edge kind, source, target, status: this is the data of a labelled directed graph, recorded in index position.

The r50 refinement (§A.261.5) makes the four slots explicit and names them:

$$ {}^{L_x}{}_{G_x}\mathcal{O}_{x\,M_x}^{R_x}, $$

with L_x the local readout/source side, G_x the guard or selector side, R_x the relay/structure side, and M_x the monitor codomain side. The r49 recap (§A.260.2) shows the same grammar carrying a transport condition: a local boundary source may feed a local relay monitor only after the guard projection has separated it from $C4$, $C6/T2$ and the all-readout channel.

Theoremlet r93. The two-sided display grammar of Definitions H2.40 and H2.41, as instantiated from r26 onward, is an interface grammar over a labelled directed graph. Under the correspondence

$$ \begin{aligned} &\text{back upper slot} \longleftrightarrow \text{node of the finite chart family} \\ &\text{back lower slot} \longleftrightarrow \text{edge kind at that node} \\ &\text{front upper slot} \longleftrightarrow \text{local source/readout attachment} \\ &\text{front lower slot} \longleftrightarrow \text{guard or selector governing the attachment} \end{aligned} $$

the admissible-contraction rules of Definition H2.41 read as a forwarding discipline: classical upper–lower contraction is transport along a declared edge, and the prohibition on same-variance pairing without a declared comparison tensor or selector is the prohibition on joining two channels without a declared admission element.

Proof sketch. The correspondence is read off the instantiations rather than postulated. In every two-sided display occurring from r26 onward the back upper slot is filled by a member of the r14 chart family or by an obstruction name carried in the same position, and the back lower slot is filled by an edge kind or monitor role. The r26 transition ledger supplies source, target and status for each transition, which is the incidence data of the graph. The forwarding reading of the contraction rules is a restatement, not an extension: rule 1 of Definition H2.41 permits exactly the pairings in which one slot is a declared codomain of the other, and rules 2–4 admit a pairing only in the presence of a named $G_{\alpha\beta}$, $G^{\alpha\beta}$ or χ_{mn}. No pairing is added and none is removed. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

r93 consequence for the relay arithmetic

Under the routing reading the finite relay layer of §A.226 is an admission calculation. With the overlap flux and required gain

$$S_V = N + \kappa N(N-1), \qquad G_V^{\mathrm{req}} = \frac{16}{S_V},$$

a chart is admissible exactly when its available gain covers its required gain, and the window criterion $16/G_{\mathrm{avail}} \le S_V < 16$ is the corresponding capacity band. The three charts recorded as blocked in the r14 scan are those whose requirement exceeds what the node supplies. The guarded micro-lift of §A.228, which carried $V9$ from $(2, 1/2, 3)$ to $(3, 1/2, 3)$ and obtained $H_9' = 18 > 16$, is a capacity increase on a single node, executed until the link carries and no further.

This also sharpens the reading of §A.270. The relative reserve

$$1 - \frac{G_V^{\mathrm{req}}}{G_V^{\mathrm{avail}}}, \qquad \text{attained as } 1 - \tfrac{4}{5} = \tfrac{1}{5} \text{ at } V3,$$

is the margin by which the weakest assisted node exceeds its own requirement. Minimising it over the chain is the ordinary statement that a chain is limited by its weakest link. The r92 derivation is unchanged; only its interpretation is now stated in the vocabulary the notation already uses.

r93 conservativity is not inertness

Proposition H2.46 states that every well-formed two-sided display expands into the inherited QAM-over-ZF language and that the notation cannot by itself introduce a new primitive tensor kind, a new admissibility relation or a new theorematic certificate. That proposition is not weakened here, and this layer relies on it.

It is worth recording that conservativity in that sense does not mean the notation is decorative. A routing table is likewise definable in the language of the objects it routes between, and it still determines what reaches what. The two statements are compatible: the graph structure carried in the index slots adds no mathematical primitive, and it does determine which transports are admissible. Reading conservativity as inertness is a misreading that this layer is written to prevent.

r93 what is not claimed

The front slots carry readout, frame, guard and selector data. They are not asserted to transform tensorially under any coordinate change, and no transformation law is supplied or implied for them. The graph reading is a reading of the declared typing; it is not a claim that the chart family is a physical network, that the edges are physical channels, or that the node capacities are measurable quantities. The notation is used locally and sparingly in the present release line, as recorded at its introduction, and r93 does not rewrite the monolith into it.

r93 ledger and guard update

The naming is complete for the sites that instantiate the notation; the source is not uniform in its use:

$$R_{\mathrm{route}}^{r93} = (0, 0, 1)$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

for (slot semantics identified, forwarding reading of the contraction rules stated, uniform use across the monolith not performed). The third component is a presentation entry and carries no obstruction.

r93 spends no guard:

$$R_{\mathrm{glob}}^{(r93,+)} = (0,1,1,1,1), \qquad \Gamma_{\mathrm{guard}}^{r93} = (1,1,1,1,1).$$

All statements are finite, local and chart-bound. The layer makes no spectral identification between Riemann zeros and photon ticks and does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.7.0-r94 Source-Only: Provenance of the C5 Contraction Factor 3/4

Purpose. Since r09 the bounded C5 collar runs on the update $\mathcal{U}_5^{\mathrm{bd}} : \Phi \mapsto \frac{3}{4}\Phi$, and r86 inherits it without revisiting where the factor comes from. This layer traces the factor to its first occurrence, records what the source does and does not supply as its justification, and separates the results that depend on the exact value 3/4 from those that hold for every $q \in (0,1)$. The question was raised in an external OP5/QAM working note kept in the repository folder `analysis/` (sections "Re-audit of the origin and necessity of the factor three quarters" and "Step twenty does not identify a unique coefficient" of `OP5_QAM_Residual_Lift.tex`), which was written against the RC19 source and did not see r85–r93. r94 checks its two findings against the current monolith and records them in source. It introduces no new object and no new claim about $C5$.

Source-only status. This r94 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The intended reserved v4.7.0 Version DOI remains 10.5281/zenodo.20232911; the previous baseline DOI 10.5281/zenodo.20117361 remains provenance only.

r94 provenance trace

The C5 chain introduces and propagates the factor as follows.

layer	object	value	how the value enters
r04	Def. A.215.3, row W_5^{bd}	$\Phi_5 = \frac{3}{4},\ \Delta_5 = -\frac{1}{4}$	ledger entry
r05	Def. A.216.4, row Z_5^{bd2}	$\Phi_5^{(2)} = \frac{9}{16}$	ledger entry
r05	same ledger, row Z_5^{collar}	$\Phi_5^{(2)} = \frac{5}{8}$	ledger entry, ratio $\frac{5}{6}$
r06	Def. A.217.2	$q_1 = q_2 \le \frac{3}{4}$	decision criterion
r07	Def. A.218.1	$\Phi_5^{(3)} = \frac{27}{64}$	probe entry
r08	Lem. A.219.7	$\Phi_5^{(4)} = \frac{81}{256}$	computed by the candidate update
r09	Def. A.220.5	$\mathcal{U}_5^{\mathrm{bd}} = \frac{3}{4}\cdot$	operator definition

The r04 value $\Delta_5(W_5^{\mathrm{bd}}) = -\frac{1}{4}$ is defined as the difference of two listed scores; it is not an independent computation of a removed quarter.

Theoremlet r94.1 (provenance). In the present source the C5 contraction factor 3/4 is a declared value, not a derived one. Specifically:

1. its first occurrence in the C5 sense is the r04 ledger entry $\Phi_5(W_5^{\mathrm{bd}}) = 3/4$;
2. the trace values 3/4, 9/16, 27/64 of r04, r05 and r07 are ledger or probe entries, and the r08 value 81/256 is produced by applying the candidate update itself (Proposition A.219.8 correctly classifies it as a prediction, $\mathsf{cand}_{\mathrm{ind}}$);
3. no layer supplies a mechanism — a projection identity, an energy partition, a dynamical rate — that produces the value 3/4 from MM/HC, collar or readout data.

Consequently the phrase "four independent bounded lowering steps" in the proof of Lemma A.220.6 is to be read as "four consecutive bounded lowering steps": three of them are recorded entries and the fourth is the operator's own prediction. The lemma itself is unaffected, because its proof of admissibility rests on positivity and domain preservation, which hold for the declared factor regardless of its origin.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof sketch. Items 1 and 2 are read off the cited definitions and lemmas. For item 3 the working file was searched in full for the forms `\tfrac34`, `\frac34`, `\frac{3}{4}`, `3/4` and `quarter`. The occurrences before r04 belong to other objects: the quarter-turn R_W of the minimal two-directional leakage plane (Corollary H3.25), which is an isometry and therefore cannot contract a norm, and the equal quarter-load shells of Corollary H9.139, which concern OP 1 and are explicitly declared not necessary. From r05 onward every C5 occurrence either restates the r04–r09 trace or applies Theorem A.220.7; the r86 inventory (§A.270) lists Definition A.220.5 as the origin of $\mathcal{U}_5^{\mathrm{bd}}$ and states that everything in it is inherited. The only occurrences that use the word "calibrated" concern the relay coupling $\kappa = 3/4$ of the comfort-calibrated clock seed $(N, \kappa, G) = (5, 3/4, 1)$, a different parameter, chosen with positive margin above the requirement $\kappa \geq 11/20$; no line of the source identifies κ with the C5 ratio. □

Remark (why a single projection does not explain the factor). The working note observes that an orthogonal least-squares correction $r' = r - d$ with $\langle r - d, d\rangle = 0$ gives $\|r'\|^2 = \|r\|^2 - \|d\|^2$, hence the score ratio 3/4 when the removed component carries one quarter of the residual energy. This explains one step at most: the projection is idempotent, so applying it again changes nothing, and a geometric recurrence would need a refreshed fit direction with a new one-quarter identity at every stage. The source declares neither. The remark is recorded as a constraint on any future derivation, not as a candidate mechanism.

r94 what the detector stage determines

r86 computes the detector stage from the factor (§A.270). The converse direction is not recorded there.

Theoremlet r94.2. Let $\Phi^{(n)} = q^n$ with $q \in (0, 1)$ and detector threshold $\theta = 1/256$. The first stage with $\Phi^{(n)} \leq \theta$ equals 20 if and only if

$$\left(\tfrac{1}{256}\right)^{1/19} < q \leq \left(\tfrac{1}{256}\right)^{1/20}, \qquad \text{numerically} \qquad 0.746879\ldots < q \leq 0.757858\ldots.$$

Hence the recorded detector stage $n = 20$ is consistent with $q = 3/4$ but does not single it out: every factor in a half-open interval of width $0.01098\ldots$ produces the same stage.

Proof sketch. For $q \in (0, 1)$ the sequence q^n is strictly decreasing, so the first crossing is at stage 20 exactly when $q^{20} \leq 1/256 < q^{19}$. Taking the 19th and 20th roots gives the interval. For $q = 3/4$ the exact rational check is $(3/4)^{19} = 0.004228\ldots > 1/256 \geq 0.003171\ldots = (3/4)^{20}$, as in the remark to Theoremlet r86.1. □

Illustration from the r05 ledger. The tolerated collar row Z_5^{collar} of Definition A.216.4 has ratio $(5/8)/(3/4) = 5/6$. Iterated at that ratio the score would first reach 1/256 at stage 31, not 20. The stage $n = 20$ therefore records the choice of the strict row Z_5^{bd2} over the tolerated row, which r05 makes by minimality, and not a property of the collar as such. The figures 20 and 31 were checked by iterating in exact rational arithmetic.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

r94 which results depend on the exact value

holds for every $q \in (0,1)$	depends on $q = 3/4$ or the r94.2 interval
finite positivity, Lem. A.222.5	stage $n = 20$ at $\theta = 1/256$ (interval)
limit zero, Thm. A.220.7	halving cost $\ln 2/\ln(4/3)$ of r86.1
finite-zero exclusion, Theoremlet r86.2	instances $c_\star(1), \ldots, c_\star(5)$ of r88.1
subtractive obstruction, Theoremlet r88.2	
r88 shape (iii): $(\alpha), (\beta), (\delta)$	r88 shape (iii): (γ) (interval)

r86.2 already notes that the exclusion depends only on $q > 0$. r94 extends that observation to the whole status picture: none of the structural conclusions of r86 and r88 uses the particular value, and the one guarded claim that mentions it, $C5_{\mathrm{det/eff}}$ at $\theta = 1/256$ after $n = 20$, needs only the interval of Theoremlet r94.2.

This is the same situation r88 records for the subtractive channel (§A.270): a value that is fixed by declaration rather than by an independent structural rule carries the content of the declaration, not of a derivation. For q this is harmless, because no guarded conclusion depends on more than the interval; it becomes relevant only if a later layer wants to treat $3/4$ itself as a physical or geometric constant.

r94 ledger and guard update

$$R^{r94}_{\mathrm{prov}} = (1, 0, 0)$$

for (independent derivation of the factor $3/4$ not available, detector-stage interval determined, value-dependence of the C5 and r86/r88 results sorted). The first component is an open provenance item. It is not an obstruction, it does not enter the r48 branch queue, and none of the guarded claims depends on closing it.

r94 spends no guard and changes no status:

$$R^{(r94,+)}_{\mathrm{glob}} = (0, 1, 1, 1, 1), \qquad \Gamma^{r94}_{\mathrm{guard}} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

The admissible claim $C5_{\mathrm{det/eff}}$ at $\theta = 1/256$ after $n = 20$ is unchanged. The layer does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.8.0-r01 Source-Only: Planck-Readout Metric Layer

Purpose. This layer opens the v4.8.0 line. It integrates the Planck-readout candidate prepared on the repository research branch for v4.8.0-r01 and states it in the frame-typed vocabulary of r81–r82. The candidate separates the physical Planck ruler from the scale carried by a lower interface or transport layer: what is invariant is not the number ℓ_P in every layer, but its guarded recovery under an admissible readout. The monolith already contains one instance of this pattern, the photon-frame Planck length of Theorem 25.5; r01 names the general form and fixes how its sign is to be read relative to the r82 signature convention. The field-equation refactor announced in the working title of the research line is not carried out here.

Source-only status. This v4.8.0-r01 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.7.0 Version DOI 10.5281/zenodo.20232911 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.8.0-r01 readout-calibrated Planck scale

Definition v4.8.0-r01.1. Let R be an admissible readout or interface mode and let $\lambda_R \in \mathbb{R} \setminus \{0\}$ be the local calibration factor induced by the MM/HC/QAM transport from the structural interface to the physical readout R. Write

$$\ell_{\mathrm{eff}}(R) := \ell_P\, \lambda_R, \qquad o_R := \operatorname{sgn} \lambda_R \in \{+1, -1\}, \qquad |\ell_{\mathrm{eff}}(R)| = \ell_P\, |\lambda_R|.$$

$\ell_{\mathrm{eff}}(R)$ is a signed interface coordinate. Its magnitude is the length scale the readout R resolves; its sign o_R is an orientation code.

Theoremlet v4.8.0-r01. Under Definition v4.8.0-r01.1 the following hold.

1. *Matter recovery.* If R_{M} is a stable matter readout with $\lambda_{R_{\mathsf{M}}} = 1 + \varepsilon_{\mathsf{M}}$ and $|\varepsilon_{\mathsf{M}}| \le \delta_{\mathsf{M}} < 1$, then $o_{R_{\mathsf{M}}} = +1$ and $\left|\ell_{\mathrm{eff}}(R_{\mathsf{M}}) - \ell_P\right| \le \delta_{\mathsf{M}} \ell_P$: the ordinary Planck scale is recovered up to the declared local calibration error.

2. *Photon instance.* The photon-frame Planck length of Theorem 25.5 is the instance $\lambda_\gamma = \gamma_L^{-3/2}$, $o_\gamma = +1$, of the same definition.

3. *Opposite orientation.* If R_τ is an opposite-orientation readout with $\lambda_{R_\tau} = -\mu_{R_\tau}$, $\mu_{R_\tau} > 0$, then $\ell_{\mathrm{eff}}(R_\tau) = -\ell_P \mu_{R_\tau}$ while $|\ell_{\mathrm{eff}}(R_\tau)| = \ell_P \mu_{R_\tau} > 0$. The negative sign is the orientation code $o_{R_\tau} = -1$ and not a negative physical distance:

$$o_R = -1 \;\not\Rightarrow\; L_{\mathsf{M}} < 0.$$

Proof sketch. Item 1: $\delta_{\mathsf{M}} < 1$ gives $\lambda_{R_{\mathsf{M}}} > 0$, and $\ell_{\mathrm{eff}}(R_{\mathsf{M}}) - \ell_P = \ell_P \varepsilon_{\mathsf{M}}$. Item 2: substituting $\lambda_\gamma = \gamma_L^{-3/2}$ into the definition reproduces the display of Theorem 25.5; nothing is proved anew and the status of that theorem is inherited unchanged. Item 3: the definition separates the signed coordinate from its magnitude, and the magnitude is positive whenever $\mu_{R_\tau} > 0$. The non-implication is the r82 guard $\ell_{\mathsf{T}}^2 < 0 \not\Rightarrow L_{\mathsf{M}} < 0$ restated for the orientation code. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.8.0-r01 orientation code and r82 signature are different data

r82 types the tachyon-frame scale through the sign of the *interval*, $\ell_{\mathsf{T}}^2 = -\lambda_{\mathsf{T}}^2$ with $\lambda_{\mathsf{T}} > 0$. r01 types an opposite-orientation readout through the sign of a *linear* coordinate, $o_{R_\tau} = -1$. The two are not the same bit: squaring the r01 coordinate gives $\ell_{\text{eff}}(R_\tau)^2 = \ell_P^2 \mu_{R_\tau}^2 > 0$, which is not the r82 interval. This layer therefore does not identify them.

If a later layer calibrates the tachyon-frame native magnitude as $\lambda_{\mathsf{T}} = \ell_P \mu_{R_\tau}$, the readout R_τ carries two independent records,

$$\big(o_{R_\tau},\ \operatorname{sgn} \ell_{\mathsf{T}}^2\big) = (-1, -1), \qquad \text{magnitude } \ell_P \mu_{R_\tau},$$

an orientation code from r01 and a signature marker from r82. Neither record is a negative material length, and neither is derivable from the other without a declared comparison map. This is the r93 discipline: two channels are not joined without an admission element.

Remark (two-sided notation). The research note records the calibration schematically as ${}^{R}{}_{\text{phys}}\mathcal{T}^{\text{struct}}{}_{\text{HC/MM}}$. Under the slot reading of r93 the front lower slot carries a guard or selector, so the matching r93 form is ${}^{R}{}_{G_R}\mathcal{T}^{\text{struct}}{}_{\text{HC/MM}}$ with G_R the calibration guard $|\varepsilon_R| \le \delta_R$. The back slots here name a layer and a transport, not a chart of the r14 family; the display is schematic and is not an instance of the r93 routing correspondence.

v4.8.0-r01 what is not claimed

The layer does not derive any λ_R except by citing Theorem 25.5; for every other readout λ_R is a declared calibration datum. It does not assert that ℓ_P is non-invariant as a physical constant, does not promote a negative sign to an observable negative length, does not prove exact through-collapse transport, and does not refactor the MM/HC field equations.

v4.8.0-r01 ledger and guard update

$$R_{\text{Planck}}^{v480\text{-}r01} = (0, 0, 1)$$

for (calibrated scale defined with matter recovery and photon instance, orientation code separated from the r82 signature, MM/HC field-equation refactor not performed). The third component is open research, not an obstruction.

The layer spends no guard:

$$R_{\text{glob}}^{(v480\text{-}r01,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\text{guard}}^{v480\text{-}r01} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.8.0-r02 Release Integration and Scope Refresh

Purpose. This layer integrates the post-v4.7.0 source-only layers and the opening layer of the v4.8.0 line into the v4.8.0 release, in the manner in which r84 integrated r81–r83 into the v4.7.0 release scope. It adds no mathematical content.
Release status. Unlike the preceding layers, r02 is a release layer. It accompanies the v4.8.0 Version DOI 10.5281/zenodo.23365244 under the Concept DOI 10.5281/zenodo.19210728. The v4.7.0 Version DOI 10.5281/zenodo.20232911 becomes provenance.

v4.8.0-r02 integrated layers

layer	content	status change
r85	citation repair, unity-theorem scope separation	none
r86	C5 collar bookkeeping, finite-zero exclusion for $q > 0$	none
r87, r89	selector-window calibration and its correction	none
r88	non-collar routes: shapes (i), (ii) excluded, (iii) open	none
r90, r92	reconstruction of $\Delta_\gamma < 1/5$	none
r91	Q4 executed: β_4 not spendable	none
r93	two-sided typing read as a routing grammar	none
r94	provenance of the C5 factor 3/4	none
v4.8.0-r01	readout-calibrated Planck scale	none

The open items carried into v4.8.0 are: shape (iii) of r88, the independent derivation of the C5 factor (r94), uniform use of the two-sided notation (r93), and the MM/HC field-equation refactor (v4.8.0-r01). None of them is an obstruction to a guarded claim.

v4.8.0-r02 guard update

The release spends no guard:

$$R_{\mathrm{glob}}^{(v480\text{-}r02,+)} = (0,1,1,1,1), \qquad \Gamma_{\mathrm{guard}}^{v480\text{-}r02} = (1,1,1,1,1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

The safe release formulation remains: guarded effective/local and frame-typed closures, not exact global closure. The release does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, global all-readout closure, or a negative physical length.

A.271 v4.8.0 Post-Release Layers

The source-only layers v4.8.0-r03 to r20 follow, recorded after the v4.8.0 release on the full-monolith working line. They are predominantly audits, corrections and pre-registered tests; their registers (v4.8.0-r09, extended in r17–r20) record which earlier statements stand, are conjectures, or are refuted. Earlier layers are not rewritten.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.8.0-r03 Source-Only: Dimensional and Numerical Audit of the Physical-Constant Layer

Purpose. An external structural evaluation of v4.7.0 rated equation and dimensional integrity as the weakest of its five criteria. This layer audits the places where the monolith assigns physical units, CODATA values or frame-insertion counts: the nomenclature, the phase-shifted constants of §25, the Hawking analysis of §21, the cosmological statements around Theorems 22.10 and 22.16, and the photon-frame Λ block of §42.9. Every number was recomputed. Earlier layers are not rewritten; each finding is recorded here with the correction that follows from the document's own rules, or with the reason why no such correction exists.
Source-only status. This v4.8.0-r03 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.8.0 Version DOI 10.5281/zenodo.23365244 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.8.0-r03 the frame-insertion calculus

The definition in §25 assigns to a constant X the photon-frame value $X^{(\mathrm{photon})} = \gamma_L^{-k(X)} X^{(\mathrm{matter})}$.

Lemma v4.8.0-r03.1. If $X = \prod_i Y_i^{a_i}$ with real exponents a_i and every factor obeys the rule, then X obeys it with

$$k(X) = \sum_i a_i \, k(Y_i).$$

Proof. $X^{(\mathrm{photon})} = \prod_i \big(\gamma_L^{-k(Y_i)} Y_i^{(\mathrm{matter})}\big)^{a_i} = \gamma_L^{-\sum_i a_i k(Y_i)} X^{(\mathrm{matter})}$. □

The lemma is what Theorem 25.5 uses: $\ell_P = (\hbar G_N/c^3)^{1/2}$ gives $k(\ell_P) = \frac{1}{2}(1+2-0) = \frac{3}{2}$, as Proposition 42.16 also states. Since the rule multiplies dimensionful quantities by the pure number γ_L, it is dimensionally consistent by construction. What it requires is that every composite quantity be assigned the k the lemma gives, and that every factor of the composite carry its own k. The findings below are the places where that is not done.

v4.8.0-r03 findings

F1 — vacuum-energy ratio (k-calculus). Equation (16), Corollary 42.17 and the corollary "Cosmological constant revisited" in §25 use $\rho_{\mathrm{vac}} \sim \ell_P^{-4}$ and obtain the frame ratio γ_L^6. The relation $\rho \sim \ell_P^{-4}$ holds only in units with $\hbar = c = 1$. The framework assigns $k(\hbar) = 1$, so $\hbar$ cannot be set to one in both frames. As an energy density, $\rho_{\mathrm{vac}} \sim \hbar c \, \ell_P^{-4}$, and Lemma v4.8.0-r03.1 gives

$$k(\rho_{\mathrm{vac}}) = 1 + 0 - 4 \cdot \tfrac{3}{2} = -5, \qquad \frac{\rho_{\mathrm{vac}}^{(\mathrm{photon})}}{\rho_{\mathrm{vac}}^{(\mathrm{matter})}} = \gamma_L^5 = 22.53\ldots,$$

not γ_L^6. The mass-density form m_P/ℓ_P^3 gives the same $k = -\frac{1}{2} - \frac{9}{2} = -5$. In addition, the source quotes γ_L^6 as 41.8 in one place and 42.0 in another; with $\gamma_L = 1.86444\ldots$ the value is $42.004\ldots$.

F2 — fine-structure drift (sign of the exponent and empirical status). With $k(\alpha^{-1}) = 1$ (Theorem 25.3) the rule gives $(\alpha^{-1})^{(\mathrm{photon})} = \gamma_L^{-1} (\alpha^{-1})^{(\mathrm{matter})}$, hence

$$\alpha^{(\mathrm{matter})} = \gamma_L^{-1} \, \alpha^{(\mathrm{photon})},$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

whereas the corollary "α-drift as a frame effect" writes $\alpha^{(\mathrm{matter})}(t) = \gamma_L^{(\mathrm{local})}(t)\,\alpha^{(\mathrm{photon})}$. The two are inverse to each other. Separately, the corollary speaks of an "observed NIST drift". The NIST $\mathrm{Al}^+/\mathrm{Hg}^+$ clock comparison of 2008 found a drift consistent with zero, and a fit of the same data for an annual modulation gives a fractional amplitude of about $(-8.9 \pm 8.4) \times 10^{-18}$, also consistent with zero. There is therefore no observed drift to explain. The annual-modulation prediction survives only as a constraint: $|\Delta\gamma_L^{(\mathrm{local})}/\gamma_L| \lesssim 10^{-17}$ over the year.

F3 — three incompatible types for Λ. The monolith uses Λ in three dimensionally different roles:

$$\begin{aligned} &\Lambda_{\mathrm{bare}} \sim \gamma_1^2 \approx 200\, m_P^4 && \text{(Prop. 42.18)},\\ &\Lambda_{\mathrm{eff}} \sim \beta_{\mathrm{sym}}^2 \rho_P && \text{(near Thm. 22.10)},\\ &\Lambda_{\mathrm{obs}} = \beta_{\mathrm{sym}}^2\,(\ell_P^{(\mathrm{photon})})^{-2} && \text{(§42.15)}. \end{aligned}$$

The first equates a pure number with a Planck-unit energy density, the second is an energy density, the third a curvature with unit m^{-2}. Curvature and energy density are related by $\Lambda = 8\pi G_N \rho_\Lambda / c^4$; the factor 8π is absent throughout. The photon-frame step of Proposition 42.18 uses $k(\Lambda) = 2$, while the curvature form $\Lambda \propto \ell_P^{-2}$ gives $k = -3$ under Lemma v4.8.0-r03.1. The source already states that a self-consistent $k(\Lambda)$ is open; r03 records that the value 2 used in the display is not the one the calculus gives for any of the three forms.

F4 — the "unit-system artifact" remark. Remark 42.21 and the prediction P4 of §42.15 state that the 10^{-122} of the cosmological-constant problem is an artefact of SI units, and that in natural units $\Lambda_{\mathrm{obs}} \sim \beta_{\mathrm{sym}}^2 \approx 0.30$. This is a dimensional error. The quantity

$$\Lambda_{\mathrm{obs}}\,\ell_P^2 = 1.1056 \times 10^{-52}\,\mathrm{m}^{-2} \cdot (1.6163 \times 10^{-35}\,\mathrm{m})^2 = 2.89 \times 10^{-122}$$

is dimensionless and takes the same value in every unit system; in units with $\ell_P = 1$ the measured Λ_{obs} is 2.89×10^{-122}, not 0.30. Taken literally, $\Lambda_{\mathrm{obs}} = \beta_{\mathrm{sym}}^2 (\ell_P^{(\mathrm{photon})})^{-2} = 7.5 \times 10^{69}\,\mathrm{m}^{-2}$ exceeds the measured value by a factor 6.8×10^{121}. P4 is therefore contradicted by measurement as stated. The same block also gives three different sizes for the required cancellation (2.3×10^{-4}, 1.5×10^{-3} and a factor 3.1), which cannot all describe one ratio.

F5 — cosmological fractions. Theorem 22.16 writes $1 - \beta_{\mathrm{sym}}^2 \approx \frac{2}{3} \approx 0.698$; with $\beta_{\mathrm{sym}}^2 = 0.30178\ldots$ the value is $0.69822\ldots$, which is not $\frac{2}{3}$. Two different predictions for Ω_Λ/Ω_M occur: $\beta_{\mathrm{sym}}^2/\gamma_{\mathrm{string}} = 2.37$ near Theorem 22.10 and $(1-\beta_{\mathrm{sym}}^2)/\beta_{\mathrm{sym}}^2 = 2.31$ from Theorem 22.16. They are different functions of the same parameters and cannot both hold. The observed ratio from the Planck 2018 values quoted in Remark 22.17 is $0.6889/0.3111 = 2.214$, not 2.18. The same remark writes $\beta_{\mathrm{sym}} = \gamma_{\mathrm{BI}}/\gamma_{\mathrm{string}} \approx 0.5493$; that quotient is $\gamma_L = 1.864$, and the stated value belongs to $\beta_{\mathrm{sym}} = \sqrt{(\gamma_L - 1)/(\gamma_L + 1)}$.

F6 — Hawking temperature. Proposition 21.4 derives $T_H = \hbar c^3/(8\pi G_N M k_B) = m_P^2 c^2/(8\pi M k_B)$ correctly. The step to $k(T_H) = -1$ declares $k(k_B) = 0$ but also needs $k(M) = 0$ for the black-hole mass, which is not declared, and which differs from the assignment $k(m_e c^2) = 1$ for the electron. Under Lemma v4.8.0-r03.1, $k(T_H) = 1 - 2 - k(M) = -1 - k(M)$.

F7 — notation. In Theorem 22.10 the letter k denotes a spring constant, while $k(X)$ denotes the frame-insertion count elsewhere in the same part. The oscillator frequency $\omega = \gamma_{\mathrm{string}}\sqrt{2/3}$ is a pure number; Planck units are implied but not declared.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Verified without finding. $\gamma_{\text{string}} = \ln 2/(\pi\sqrt{3}) = 0.12738\ldots$; $\gamma_L = \gamma_{\text{BI}}/\gamma_{\text{string}} = 1.86444\ldots$; $\beta_{\text{sym}} = \sqrt{(\gamma_L - 1)/(\gamma_L + 1)} = 0.54935\ldots$; $\gamma_L^{3/2} = 2.5458\ldots$; $\ell_P^{(\text{photon})} = 6.35 \times 10^{-36}\,\text{m}$; $k(\ell_P) = \frac{3}{2}$; the units of G_N, $\hbar$, c in Theorem 25.3; $T_H = \hbar c^3/(8\pi G_N M k_B)$; $(\alpha^{-1})^{(\text{photon})} = 73.50\ldots$ under $k(\alpha^{-1}) = 1$.

Theoremlet v4.8.0-r03. Under the frame-insertion rule of §25 and Lemma v4.8.0-r03.1:

1. the photon-to-matter ratio of a Planck-cutoff vacuum energy density is γ_L^5, not γ_L^6;
2. $k(\alpha^{-1}) = 1$ implies $\alpha^{(\text{matter})} = \gamma_L^{-1}\alpha^{(\text{photon})}$;
3. Λ has no single k until one of its three forms is chosen, and none of them gives $k = 2$;
4. no choice of units turns the dimensionless number $\Lambda_{\text{obs}}\ell_P^2 \approx 2.9 \times 10^{-122}$ into a number of order one.

Proof sketch. Items 1–3 are applications of the lemma to $\hbar c\,\ell_P^{-4}$, to $\alpha = (\alpha^{-1})^{-1}$, and to ℓ_P^{-2}, $\hbar c\,\ell_P^{-4}$ and $G_N\rho/c^4$ respectively. Item 4 holds because a ratio of two quantities of the same dimension is invariant under change of units. □

v4.8.0-r03 re-check of P4

F4 was checked a second time, against the main paper and against a reading of P4 that is dimensionally consistent.

Origin of the unit ℓ_P^{-2}. The main paper states the frame contribution without a unit. Its Theorem 11.6 (Dark energy as permanent frame offset) gives $\Lambda_{\text{eff}} \sim \beta_{\text{sym}}^2\rho_P$, its Proposition 12.4 (Frame term gives the cosmological constant) gives $\Lambda_{\text{obs}} \sim \beta_{\text{sym}}^2$, and its remark on the cosmological-constant problem records that reducing the bare value by 10^{122} through the Θ–R cancellation "is not yet established" [2]. The unit ℓ_P^{-2} and the statement that 10^{-122} is a unit artefact enter only in the photon-frame Λ block of the companion, §42.9. The error recorded in F4 therefore lies in that later reinterpretation, not in the main paper.

F8 — "dimensionally correct". The proof of Theorem 42.19(iii) calls $\Lambda_{\text{obs}} = \beta_{\text{sym}}^2/\ell_P^{(\text{matter})2}$ "dimensionally correct up to the $\gamma_L^{3/2}$ Planck-length correction". The unit m^{-2} is correct; the value is $1.2 \times 10^{69}\,\text{m}^{-2}$, a factor 10^{121} above the measurement, and no power of γ_L closes that gap. The theorem uses the matter-frame Planck length, P4 the photon-frame one.

F9 — citation. The proof of Theorem 42.19(iii) and the status table of the Λ block cite "Proposition 18.4 of the main paper". The main paper has no numbered environment 18.4. The statement meant is its Proposition 12.4, which asserts only the proportionality $\Lambda_{\text{obs}} \sim \beta_{\text{sym}}^2$.

F10 — units of the balance condition. In Equation (17) the integrand $\Sigma(\tau)$ contains $e^{-\tau/2}$, so τ is a pure number, while the upper limit $\ell_P^{(\text{photon})}$ and the right-hand side $\beta_{\text{sym}}^2\ell_P^{(\text{photon})2}$ are written as a length and a squared length. The condition is consistent only when read in matter-frame Planck units, where $\ell_P^{(\text{photon})} = \gamma_L^{-3/2} = 0.3928\ldots$ and the right-hand side is $\beta_{\text{sym}}^2\gamma_L^{-3} = 0.0466\ldots$. This reading is not stated in the source.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A dimensionally consistent reading, offered and not adopted. Since β_{sym}^2 is a pure number, the consistent place for it is a dimensionless fraction of the critical density, not Λ itself. If the Markov statement of Theorem 22.16, $\Omega_\Lambda = 1 - \beta_{\mathrm{sym}}^2$, is taken as the prediction, then

$$\Lambda_{\mathrm{obs}} = \frac{3\,\Omega_\Lambda H_0^2}{c^2} = \frac{3\,(1-\beta_{\mathrm{sym}}^2)\,H_0^2}{c^2} = 1.112 \times 10^{-52}\,\mathrm{m}^{-2} \qquad (H_0 = 67.4\ \mathrm{km\,s^{-1}\,Mpc^{-1}}),$$

which differs from $\Lambda_{\mathrm{obs}} = 1.1056 \times 10^{-52}\,\mathrm{m}^{-2}$ by 0.6%. At the level of the fraction, $1 - \beta_{\mathrm{sym}}^2 = 0.6982\ldots$ compares with the Planck 2018 values 0.6847 and 0.6889 by 2.0% and 1.4%.

The reading has a price. It predicts the fraction Ω_Λ, and takes the absolute scale from the measured H_0. The number 10^{-122} does not disappear: it becomes $\Lambda_{\mathrm{obs}}\ell_P^2 = 3\Omega_\Lambda(\ell_P/R_H)^2$ with the Hubble radius $R_H = c/H_0$, so that the remaining question is the hierarchy $R_H/\ell_P = 8.49 \times 10^{60}$. The source quantity closest to it is the active zero count $n_H \sim 10^{62}$ at the Hubble time (Theorem 11.5 of the main paper); it differs from R_H/ℓ_P by a factor of about 12, and the source does not identify the two. The reading also selects $1 - \beta_{\mathrm{sym}}^2$ among the two conflicting predictions of F5.

Whether P4 is restated in this form, restated otherwise, or withdrawn is left to the author. r03 records the reading as an option and asserts nothing beyond the arithmetic above.

v4.8.0-r03 provenance of the frame parameters

F11 — both inputs of γ_L are values of the same LQG parameter. The frame factor $\gamma_L = \gamma_{\mathrm{BI}}/\gamma_{\mathrm{string}} = 1.864\ldots$ enters every frame conversion of the monolith, $\beta_{\mathrm{sym}} = \sqrt{(\gamma_L - 1)/(\gamma_L + 1)}$, the cosmological fractions of Theorem 22.16, the photon-frame Planck length of Theorem 25.5 and the instance $\lambda_\gamma = \gamma_L^{-3/2}$ of v4.8.0-r01. Its two inputs have the following origin in the literature.

1. $\gamma_{\mathrm{string}} = \ln 2/(\pi\sqrt{3}) = 0.12738\ldots$, introduced in the nomenclature as "string coupling", is the value of the Barbero–Immirzi parameter obtained by Ashtekar, Baez, Corichi and Krasnov from the state counting of quantum isolated horizons (arXiv `gr-qc/9710007`, `gr-qc/0005126`). It is not a string-theory coupling.

2. $\gamma_{\mathrm{BI}} \approx 0.2375$ is the value of the same parameter after the corrected state counting of Domagala and Lewandowski and of Meissner (arXiv `gr-qc/0407051`, `gr-qc/0407052`). Domagala and Lewandowski show that the earlier counting contains a spurious constraint that suppresses spins above $j = \frac{1}{2}$, and that the earlier value therefore does not hold.

Hence γ_L is the ratio of the corrected to the superseded determination of one parameter. Theorem 20.5 reads this ratio as a Lorentz factor between the matter and the photon frame. The source supplies no derivation of either input from the HC trace formula: the proof of Theorem 20.5 says that γ_{string} is computed in the photon frame from that formula, while the proof of the OP 1/T4 statement around Remark 24.4 calls it extrinsically given. The two statements are incompatible, and only the second is supported by the text.

F11a — the entropy factor. The remark "The entropy factor" in §21 writes the LQG entropy as $S = A\ln 2/(8\pi\gamma_{\mathrm{BI}}\ell_P^2)$, finds agreement with $S_{BH} = A/(4\ell_P^2)$ at $\gamma_{\mathrm{BI}} = \ln 2/(2\pi) \approx 0.110$, and records a discrepancy factor 2.15 as an open quantitative problem. The entropy formula of the counting that produced $\ln 2/(\pi\sqrt{3})$ is

$$S = \frac{\ln 2}{4\pi\sqrt{3}\,\gamma}\,\frac{A}{\ell_P^2},$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

which agrees with S_{BH} exactly at $\gamma = \ln 2/(\pi\sqrt{3}) = \gamma_{\text{string}}$. The factor 2.15 is produced by the formula used in the remark and is not a property of the framework; the discrepancy between the two countings is $\gamma_{\text{BI}}/\gamma_{\text{string}} = \gamma_L$ itself.

Consequence. The agreement of $1-\beta_{\text{sym}}^2$ with the measured Ω_Λ at the 1–2% level recorded in §A.271 rests on β_{sym}, and β_{sym} rests on the ratio of two countings of one parameter, one of which is superseded. As long as that is not stated, the agreement reads as better supported than it is.

A reading in the readout vocabulary, offered and not adopted. The superseded counting admits only minimal spins $j = \frac{1}{2}$ at the horizon punctures, the corrected one all spins. γ_L could therefore be read as the ratio between a minimal-spin readout and a full-spin readout of the same horizon, two readouts in the sense of r81–r82 and r93. Under that reading γ_L is not a Lorentz factor, and the chain $\gamma_L \to \beta_{\text{sym}} \to \Omega_\Lambda$ would need a new justification: why the cosmological fraction should be governed by the ratio of two horizon readouts. r03 records the reading as an option and asserts nothing beyond the provenance above. Whether γ_{string} is renamed, the chain is rederived, or this reading is adopted is left to the author.

v4.8.0-r03 status of the affected statements

statement	finding	status after r03
Eq. (16), Cor. 42.17	F1	corrected: $\gamma_L^5 \approx 22.5$
α-drift corollary	F2	exponent corrected; no observed drift
Λ in Prop. 42.18, Thm. 22.10, P4	F3	open: one type for Λ to be chosen
Rem. 42.21, P4	F4	contradicted as stated; Ω-reading offered
Thm. 42.19(iii)	F8, F9	value off by 10^{121}; citation corrected
Eq. (17)	F10	Planck units made explicit
γ_{string}, Thm. 20.5	F11	provenance recorded; readout reading offered
entropy-factor remark, §21	F11a	corrected: no factor 2.15
Ω statements, Rem. 22.17	F5	values corrected; two predictions conflict
Prop. 21.4	F6	assumption $k(M) = 0$ made explicit
Thm. 22.10	F7	notation only

None of these statements is one of the guarded claims of the release scope, and none of the five guards depends on them. F3 and F4 concern the photon-frame Λ programme, which the source already marks as open; r03 sharpens that status. For P4 the source-only layer records the contradiction and does not supply a replacement; a reformulation of the Λ prediction, or its withdrawal, is left to the author.

v4.8.0-r03 ledger and guard update

$$R_{\text{dim}}^{v480\text{-}r03} = (0, 0, 1, 1, 1, 1)$$

for (k-calculus corrections F1, F2, F6 derived; numerical corrections F1, F5 recorded; single type for Λ not chosen; a replacement for P4 offered in §A.271 but not adopted; one of the two Ω_Λ/Ω_M predictions not selected; independent justification of γ_L as a frame factor not supplied, §A.271).

The layer spends no guard:

$$R_{\text{glob}}^{(v480\text{-}r03,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\text{guard}}^{v480\text{-}r03} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.8.0-r04 Source-Only: Decoupling from the Riemann Source and Standalone Sphaera Transformations

Purpose. The Companion cites the Riemann source [1] at 23 sites for seven distinct results. A full audit of that source (v4.15) and a recomputation of its central formulas show that its proof chain for RH does not hold. The author used the source as a starting point; what the Companion needs from it are the Heisenberg compensator and the Sphaera model of the basic transformations (reflection, cascade, triole hierarchy). This layer records the audit, gives standalone definitions of those three objects that do not depend on any RH statement, repairs the triole hierarchy where it fails, and reclassifies every other dependency. Earlier layers are not rewritten.
Source-only status. This v4.8.0-r04 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.8.0 Version DOI 10.5281/zenodo.23365244 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.8.0-r04 audit of the Riemann source

The combinatorial orbit theory (L1–L11, Axioms W and G, K1), the arithmetic identification $\mathcal{Z} = \zeta$ for $\operatorname{Re} s > 1$ and the functional-equation sketch are correct. The following findings concern the steps that carry the claim. Numerical checks are recorded in the audit scripts of the working folder `llm/`.

A1 — explicit formula. The source states $\Theta(\tau) = 1 - \Sigma(\tau) + R(\tau)$ with $\Theta = \sum_{p,m} m^{-1}\delta(\tau - m\log p)$. Against a smooth test function supported in $(0.6, 3.2)$ with 300 zeros the two sides give 2.1587 and 0.5198. The correct form, which matches to six digits (4.393268 on both sides), is

$$\tau\,\Theta(\tau) = \sum_n \Lambda(n)\,\delta(\tau - \log n) = e^{\tau} - \sum_{\rho} e^{\rho\tau} - \sum_{k\geq 1} e^{-2k\tau} \qquad (\tau > 0).$$

A2 — the reflection on the Hardy space. $s \mapsto 1-\bar{s}$ maps $\operatorname{Re} s > \frac{1}{2}$ outside itself. It acts only on boundary values, where it is complex conjugation; an H^2 function with real boundary values vanishes. In the only well-defined reading the source's H^+_+ is $\{0\}$. The antilinearity of J also makes $\frac{1}{2}(1 \pm J)$ only real-linear.

A3 — self-adjointness. A conjugation commuting with a symmetric operator gives equal deficiency indices, not essential self-adjointness. The operator $-i(x\,d/dx + \frac{1}{2})$ is already self-adjoint on $L^2(\mathbb{R}^+, dx)$, with continuous spectrum $\mathbb{R}$; self-adjointness is not the bottleneck.

A4 — discreteness and identification. The one-dimensional relation $[Q, P] = i$ does not produce a representation of an adelic Weyl algebra; Stone–von Neumann for a locally compact abelian group yields L^2 of the group, not of $\mathbb{Q}^+\backslash\mathbb{A}^1$. The identification step uses Connes' trace formula on the adele class space, whose validity is equivalent to RH; moreover the source's own formula gives $\mathcal{L}[\Sigma](s) = 1/s - \log\zeta(s) + \mathcal{L}[R](s)$, not $\log\zeta(s)$, and $\mathcal{L}[R]$ diverges.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A5 — the set argument. The source assigns the same kernel $(\mathbb{P}, \mathrm{id}, \log p)$ to ζ and to every self-dual automorphic L-function, and its operator stream is independent of the L-function. It would therefore assign one spectrum to all of them, while their zeros differ: the first zero of ζ is $\frac{1}{2}+14.1347251417\,i$, that of $L(s,\chi_{-4})$ is $\frac{1}{2}+6.0209489047\,i$. The Davenport–Heilbronn function satisfies a Riemann-type functional equation and has the zero $0.8085171825+85.69934849\,i$ off the critical line. Where the zeros lie is therefore not determined by the set $(\mathbb{P}, \log p)$ together with the functional equation; it requires the multiplicative structure of the Euler product.

Theoremlet v4.8.0-r04.1. The Riemann source is not a proof source for RH. None of the five guards of the release scope depends on it, and the guard W_γ^{glob}, which forbids a global photon-clock/RH theorem, is confirmed rather than affected.

Proof sketch. A1–A4 each break a link of the chain K1/L11 → T1+T3 → T4+T5 → T6, and A5 shows that the chain could not distinguish ζ from other L-functions. The guards concern $C5$, all-readout terminality, $C6/T2$, W_γ^{glob} and W_{all}; the Companion already treated the RH channel as a local photon-clock probe, so no guarded statement used RH as an input. □

v4.8.0-r04 standalone Heisenberg compensator

Definition v4.8.0-r04.1. Let $\mathcal{H}$ be a complex Hilbert space and $J : \mathcal{H} \to \mathcal{H}$ a *linear* unitary involution, $J^2 = 1$, $J^* = J$. The Heisenberg compensator and its antisymmetric partner are

$$C := \tfrac{1}{2}(1 + J), \qquad A := \tfrac{1}{2}(1 - J).$$

Theoremlet v4.8.0-r04.2. C and A are complementary orthogonal projections, $C+A = 1$, $CA = 0$; $\operatorname{ran} C = \ker(J-1)$, $\operatorname{ran} A = \ker(J+1)$; $C(Jf) = C(f)$ and $A(Jf) = -A(f)$. The terms $\mathcal{L}_{\mathrm{HC}}$ and $\mathcal{L}_{\mathrm{frame}}$ of Theorem 23.3 are J-invariant. $\mathcal{L}_{\mathrm{kin}}$ is J-invariant if and only if $JL_0J = L_0$.

Proof sketch. $C^2 = \frac{1}{4}(1 + 2J + J^2) = C$ and $C^* = C$ since $J^* = J$; likewise for A, and $CA = \frac{1}{4}(1 - J^2) = 0$. $CJ = \frac{1}{2}(J + 1) = C$ and $AJ = -A$. $\langle Jf, Jf\rangle = \langle f, f\rangle$ by unitarity; $\langle Jf, L_0Jf\rangle = \langle f, JL_0Jf\rangle$. □

Instances. (a) The e_1-reflection on $\mathbb{C}\otimes\mathbb{H}$ with the Euclidean inner product of the coefficients is a linear unitary involution; on $\mathbb{H}$ it is an anti-automorphism. (b) The linear reflection $s \mapsto 1 - s$ restricted to the critical line acts on the ordinate as the parity $t \mapsto -t$, $(Jf)(t) = f(-t)$ on $L^2(\mathbb{R})$. For (b), in logarithmic coordinates $L_0 = -i(\partial_y + \frac{1}{2})$ and parity sends $\partial_y \mapsto -\partial_y$, so $JL_0J \neq L_0$: the kinetic invariance is open for this instance.

The compensator therefore requires no RH statement. What it loses relative to the earlier text is the claim that its image is a space whose L_0-spectrum consists of zeta zeros.

v4.8.0-r04 standalone Sphaera cascade

Definition v4.8.0-r04.2. Let $\mathbb{S}$ be a fixed Hilbert space (a set, not a universal collection: ZF admits no set of all sets), and let $\mathbb{S} = S_N \supseteq S_{N-1} \supseteq \cdots \supseteq S_0$ be closed subspaces with orthogonal projections $\Pi_n : S_n \to S_{n-1}$. The cascade is $\hat{K} := \Pi_1 \circ \cdots \circ \Pi_N$.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Theoremlet v4.8.0-r04.3. $\hat{K}$ is the orthogonal projection of $\mathbb{S}$ onto S_0; in particular $\hat{K}^2 = \hat{K}$, and $\hat{K}f = f$ if and only if $f \in S_0$.

Proof sketch. For nested closed subspaces $V \subseteq W$ the projections satisfy $P_V P_W = P_V$; induction over the chain gives $\hat{K} = P_{S_0}$. □

The Maximal Totality principle is read inside this definition: $\hat{K}$ is defined on $\mathbb{S}$ and maps every element of $\mathbb{S}$ into S_0. It is not applied to elements outside its domain.

v4.8.0-r04 repaired triole hierarchy

Use the Cayley–Dickson convention of the Riemann source, $(p + q\ell)(r + s\ell) = (pr - \bar{s}q) + (sp + q\bar{r})\ell$, with $e_4 = \ell$, $e_5 = e_1\ell$, $e_6 = e_2\ell$, $e_7 = e_3\ell$. Explicit multiplication gives the seven quaternionic triples (lines of the Fano plane)

$$(1,2,3),\ (1,4,5),\ (1,6,7),\ (2,4,6),\ (2,5,7),\ (3,4,7),\ (3,5,6).$$

Theoremlet v4.8.0-r04.4. Three quaternionic triples cover all seven imaginary units of $\mathbb{O}$ if and only if they pass through a common unit. There are exactly seven such configurations, one per unit. The source's second triole (e_4, e_5, e_6) is not quaternionic, since $e_4 e_5 = e_1$.

Proof sketch. Two distinct lines of the Fano plane meet in exactly one point. For three distinct lines, $|L_1 \cup L_2 \cup L_3| = 9 - |L_1 \cap L_2| - |L_1 \cap L_3| - |L_2 \cap L_3| + |L_1 \cap L_2 \cap L_3|$, which is 7 if the three lines are concurrent and 6 otherwise. Each point lies on three lines, so each point gives exactly one concurrent triple. $e_4 e_5 = \ell(e_1\ell) = e_1$ is the source's own computation. □

Replacement. The triole configuration through $e_4 = \ell$ is

$$\mathcal{T}_1' = (e_1, e_4, e_5), \qquad \mathcal{T}_2' = (e_2, e_4, e_6), \qquad \mathcal{T}_3' = (e_3, e_4, e_7),$$

which keeps the source's third triole and shares one common axis. Equally admissible is the configuration through e_1, which keeps the inner triole (e_1, e_2, e_3).

Reflection as algebra symmetry. The reflection negating e_1 alone is not an automorphism of $\mathbb{O}$. Of the 2^7 sign changes exactly eight are automorphisms; those negating e_1 negate four units and fix one quaternionic line pointwise, for example $\{e_1, e_2, e_4, e_7\}$ with fixed line $(3, 5, 6)$. A reflection meant as a symmetry of the algebra must have this form.

Above the octonions. In the sedenions the source's outer octo-triole $(f_8, \ldots, f_{14})$ is not closed ($f_8 f_9 = f_1$), and zero divisors exist, e.g. $(e_3 + f_{10})(e_6 - f_{15}) = 0$. The hierarchy continues algebraically, not geometrically; the Hopf structure ends at $\mathbb{O}$, as the source itself states. Theorem 22.4 uses e_4, e_5, e_6 as an $SU(3)$ triplet, not as a quaternionic triole, and is not changed by this layer.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.8.0-r04 dependency map

site	used from the Riemann source	status after r04
Thm. 12.1, equivalence clause	Stone–von Neumann (Lem. 13.2, Step 3)	open
Thm. 15.2	Stone–von Neumann, T4+T5	open
Thms. 19.1, 20.1	Stone–von Neumann	equivalence open
Thms. 22.19, 22.20	T6, $\Sigma = 1 - \Theta + R$	open; formula to be replaced (A1)
Thm. 22.22, row "Riemann Hypothesis"	RH as a result	withdrawn: violates W_γ^{glob}
Thm. 23.3	$\mathrm{Adj}_J(L_0) = 0$	$\mathcal{L}_{\mathrm{HC}}$, $\mathcal{L}_{\mathrm{frame}}$ standalone; $\mathcal{L}_{\mathrm{kin}}$ open
Thm. 23.4	field equation	projector algebra kept; zero reading withdrawn
Thms. 41.13, 41.15	T1+T3, T4+T5	open (OP2 stays open)
Thm. 42.23, Rem. 42.35	explicit formula (Thm. 3.3)	to be replaced by A1
prime lattice Γ_P	classical arithmetic	unaffected
§17	RH as consequence	read as conjecture only

Theoremlet v4.8.0-r04.5. After this layer the Heisenberg compensator, the Sphaera cascade and the triole hierarchy up to $\mathbb{O}$ rest on Definitions v4.8.0-r04.1 and r04.2 and on Theoremlet r04.4, without any statement of [1] beyond classical arithmetic. The statements listed as open lose their previous justification but are not refuted by this layer.

Proof sketch. Theoremlets r04.2–r04.4 use only Hilbert-space projections and explicit Cayley–Dickson multiplication. For the open items the audit A1–A5 removes the cited justification; whether an independent justification exists is not decided here. □

v4.8.0-r04 ledger and guard update

$$R_{\mathrm{dec}}^{v480\text{-}r04} = (0, 0, 0, 1, 1, 1)$$

for (standalone compensator; standalone cascade; repaired triole hierarchy; Stone–von Neumann equivalence of the kernels open; trace-formula field equations open pending A1; kinetic J-invariance open).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v480\text{-}r04,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v480\text{-}r04} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.8.0-r05 Source-Only: Corrected Explicit Formula in the Field-Equation and Θ–R Layer

Purpose. Layer v4.8.0-r04 recorded that the explicit formula $\Sigma = 1-\Theta+R$ taken from [1] is misstated (finding A1) and listed the sites that use it. This layer carries the corrected formula into those sites: the Einstein-equation reading of Theorem 22.19, the bare-Λ proposition, the Θ–R cancellation statement, the photon-frame Θ-sum conjecture of OP 5 Step (B), and the GUE theorem. Two statements can now be decided numerically; both are decided here. Earlier layers are not rewritten.
Source-only status. This v4.8.0-r05 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.8.0 Version DOI 10.5281/zenodo.23365244 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.8.0-r05 corrected trace decomposition

Write $\Sigma(\tau) := \sum_\rho e^{i\gamma_\rho \tau}$, summed over all non-trivial zeros $\rho = \frac{1}{2} + i\gamma_\rho$ (the zeros are used as numerically known data; no statement about their real parts is assumed).

Theoremlet v4.8.0-r05.1. For $\tau > 0$, in the sense of distributions,

$$\Sigma(\tau) = \underbrace{e^{\tau/2}}_{M(\tau)} - \underbrace{e^{-\tau/2} \sum_{n \geq 2} \Lambda(n)\, \delta(\tau - \log n)}_{P(\tau)} - \underbrace{\frac{e^{-\tau/2}}{e^{2\tau} - 1}}_{Z(\tau)}.$$

M comes from the pole of ζ at $s = 1$, P from the prime powers with von Mangoldt weights, Z from the trivial zeros.

Proof sketch. This is the classical explicit formula $\sum_n \Lambda(n)\delta(\tau - \log n) = e^\tau - \sum_\rho e^{\rho\tau} - \sum_{k\geq 1} e^{-2k\tau}$, multiplied by $e^{-\tau/2}$, with $\sum_k e^{-2k\tau} = 1/(e^{2\tau} - 1)$. Paired with a smooth test function supported in $(0.6, 3.2)$ and 300 zeros, both sides of the unmultiplied identity agree to six digits (4.393268), while the form $1 - \Theta + R$ of the source gives 2.1587 against 0.5198 (r04, A1). □

What changes. The three-part structure survives: pole, primes, trivial zeros. What changes is every functional form. The constant 1 becomes the growing term $e^{\tau/2}$; the primes enter with weight $\Lambda(n) = \log p$ and damping $e^{-\tau/2}$, not with weight $1/m$; the trivial-zero term has denominator $e^{2\tau} - 1$ and positive sign inside Z. In addition, equation (14) uses Θ for the zero sum itself, which is also the definition of Σ; read literally, (13) then gives $2\Sigma = 1 + R$. The symbol Θ is reserved here for the prime-side distribution only.

v4.8.0-r05 the Einstein-equation reading

Theorem 22.19 varies $S = \int_0^\infty \Sigma(\tau)\, d\tau$. With the corrected decomposition this integral does not exist: M grows like $e^{\tau/2}$, $Z(\tau) \sim 1/(2\tau)$ as $\tau \to 0^+$, and Σ is a distribution, not a function. Moreover $\operatorname{Tr} e^{-i\tau L_0}$ is the trace of a unitary group, not a heat kernel, so the heat-kernel variation formula used in the proof does not apply.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Theoremlet v4.8.0-r05.2. The correspondence

$$M \;\leftrightarrow\; \Lambda g_{\mu\nu}, \qquad P \;\leftrightarrow\; T_{\mu\nu}, \qquad Z \;\leftrightarrow\; G_{\mu\nu}$$

is retained as a structural conjecture. Theorem 22.19 is not derived: a well-defined action requires a test-function pairing $\langle \Sigma, \varphi \rangle$ and an explicit dependence of φ on $g^{\mu\nu}$, neither of which is supplied.

Proof sketch. Divergence of the three pieces is read off from Theoremlet r05.1; the trace statement follows from the unitarity of $e^{-i\tau L_0}$. □

Bare Λ. Proposition 42.18 assigns to the constant term a size γ_1^2. In the corrected formula the pole term is $M(\tau) = e^{\tau/2}$; it carries no dependence on any zero, so no γ_1^2 can arise from it. The proposition is withdrawn as a derivation; the scale γ_1^2 remains an unsupported ansatz.

Cancellation. Theorem 42.23 reads the explicit formula as a cancellation of the pole by the other terms. The formula is an identity between distributions; it does not by itself fix the value of a truncated integral. The open content of Step (B) is exactly such a value, decided below.

v4.8.0-r05 decision of the Θ-sum conjecture

Proposition 42.33 states

$$\mathcal{T}(T) = 2\sum_{n\geq 1} \frac{\sin(T\gamma_n/\gamma_L)}{\gamma_n/\gamma_L} = \frac{3T}{2} - \log T - \frac{T^2}{24} - \gamma_E - \beta_{\mathrm{sym}}^2 T^2 \approx 0.8934, \qquad T = \gamma_L^{-3/2} \approx 0.3928.$$

Theoremlet v4.8.0-r05.3. $\mathcal{T}(T) = -0.1097 \pm 0.0009$. The conjecture is false; the two sides differ by 1.003. Satisfying the equation would require $\beta_{\mathrm{sym}}^2 \approx 6.80$, outside $[0, 1)$.

Proof sketch. $\mathcal{T}(T) = \gamma_L\, S(a)$ with $S(a) = 2\sum_{\gamma>0} \sin(a\gamma)/\gamma$ and $a = T/\gamma_L \approx 0.2107 < \log 2$. For $x = e^a \in (1, 2)$ one has $\psi_0(x) = 0$, so the Riemann–von Mangoldt formula gives exactly $\sum_\rho x^\rho/\rho = x - \log 2\pi - \frac{1}{2}\log(1 - x^{-2})$. Pairing ρ with $\bar\rho$,

$$S(a) = e^{-a/2}\Big(x - \log 2\pi - \tfrac{1}{2}\log(1 - x^{-2})\Big) - \sum_{\gamma>0} \frac{\cos a\gamma}{\frac{1}{4} + \gamma^2} + \sum_{\gamma>0} \frac{\sin a\gamma}{2\gamma(\frac{1}{4} + \gamma^2)},$$

with both remaining sums absolutely convergent. With the first 2000 zeros the value is -0.109652 (50 zeros: -0.109464; 500: -0.109661), and the tail is bounded by $8.2 \cdot 10^{-4}$ through the zero density $(2\pi)^{-1}\log(\gamma/2\pi)$. The direct partial sums converge to the same value (-0.1449, -0.1117, -0.1053 at $N = 50, 500, 2000$); the first agrees with the -0.146 recorded in Remark 42.34. □

Consequence. By Remark 42.35 the conjecture was stated as equivalent to the balance condition (17) and to OP 5 Step (B). That equivalence used the old R; with the corrected decomposition it would have to be re-derived. Either way, the balance statement in the form given there does not hold. The defect of almost exactly 1 suggests that the constant term was mishandled in the regularization; even a shift by 1 leaves a gap of 0.003, larger than the error bound. Prediction P4 and the photon-frame Λ programme, already marked open in r03, lose this route.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.8.0-r05 GUE statistics: empirical, not derived

Theorem 22.20 derives GUE statistics from self-adjointness, irreducibility and the absence of time reversal. The derivation does not hold: self-adjointness and irreducibility do not imply any random-matrix class (the harmonic oscillator has neither GUE nor GOE spacings), and an antiunitary J with $J^2 = 1$ commuting with L_0, as asserted in the proof, would be a time-reversal-type symmetry and would point to GOE, not GUE.

Theoremlet v4.8.0-r05.4. The nearest-neighbour spacings of the zeros follow GUE statistics empirically. For zeros 101–2000, unfolded with $N(t) = \frac{t}{2\pi}\log\frac{t}{2\pi e} + \frac{7}{8}$, the Kolmogorov–Smirnov distance to the Wigner surmise is 0.041 for GUE ($\beta = 2$), 0.108 for GOE ($\beta = 1$) and 0.315 for Poisson; small spacings $s < 0.1$ occur with frequency 0.0005 (GUE surmise 0.0011, GOE 0.0078, Poisson 0.095).

Proof sketch. Direct computation from 2000 zeros. The residual GUE distance reflects the Wigner surmise as an approximation and the low height; it is consistent with the Montgomery–Odlyzko results. □

Theorem 22.20 is therefore retained as an empirical statement about the zeros and withdrawn as a consequence of the Sphaera structure.

v4.8.0-r05 status of the affected statements

statement	status after r05
(13), (14), (15)	replaced by Theoremlet r05.1
Thm. 22.19	structural conjecture (r05.2); action not defined
Prop. 42.18	withdrawn as derivation; γ_1^2 unsupported
Thm. 42.23	identity, not cancellation; content moved to Step (B)
Prop. 42.33, Rem. 42.35	false: $\mathcal{T}(T) = -0.1097$, not 0.893
Thm. 22.20	empirical fact (r05.4); derivation withdrawn

v4.8.0-r05 ledger and guard update

$$R_{\mathrm{fe}}^{v480\text{-}r05} = (0, 0, 1, 1, 0)$$

for (corrected trace decomposition; Θ-sum conjecture decided; Einstein-equation reading not derived; photon-frame balance condition without a valid form; GUE statement settled empirically).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v480\text{-}r05,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v480\text{-}r05} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.8.0-r06 Source-Only: Signature of the Biquaternion Kernel and the Complex-Vector Reading

Purpose. Definition 12.5 gives the kernel $q_n = it \cdot \mathbf{1} + z_1 e_1 + z_2 e_2 + z_3 e_3$ with $z_k \in \mathbb{C}$ and the norm $|q_n|^2 = -(ct)^2 + |z_1|^2 + |z_2|^2 + |z_3|^2$, read as a Minkowski metric of signature $(-,+,+,+)$ on which the Lorentz group acts by quaternion conjugation. This layer computes the signature, identifies the natural invariant of $\mathbb{C}\otimes\mathbb{H}$, and adopts as working reading the one under which the kernels (z_1, z_2, z_3) carry the Lorentz group. In that reading the symmetric triole is exactly the light-like centred configuration, and the photon sector is the base $S_0 \cong \mathbb{C}P^1$ of Theorem 22.1. Earlier layers are not rewritten.

Source-only status. This v4.8.0-r06 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.8.0 Version DOI 10.5281/zenodo.23365244 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.8.0-r06 signature and natural norm

Write $z_k = x_k + iy_k$ with $x_k, y_k \in \mathbb{R}$.

Theoremlet v4.8.0-r06.1.

(i) The norm of Definition 12.5 is the real quadratic form $-t^2 + \sum_k (x_k^2 + y_k^2)$ of signature $(1,6)$, not $(1,3)$.

(ii) Conjugation $q \mapsto uqu^{-1}$ by $u \in \mathbb{H}^\times$ fixes the scalar part and acts by $SO(3)$ on (x_k) and on (y_k) separately; it contains no boost.

(iii) The natural invariant of $\mathbb{C}\otimes\mathbb{H}$ is the complex-bilinear norm

$$N(q) := q\bar{q} = -t^2 + z_1^2 + z_2^2 + z_3^2, \qquad \mathrm{Re}\, N = -t^2 + |x|^2 - |y|^2, \quad \mathrm{Im}\, N = 2\, x \cdot y,$$

where $\bar{q}$ is quaternion conjugation without complex conjugation; $N(pq) = N(p)N(q)$.

Proof sketch. (i) The Hessian of the form has eigenvalues -1 once and $+1$ six times. (ii) For real u, $ue_k u^{-1}$ is a rotation of the imaginary units and $u\mathbf{1}u^{-1} = \mathbf{1}$. (iii) Direct expansion of $q\bar{q}$; the vector part vanishes because the coefficients commute. Multiplicativity is inherited from $\mathbb{H}$. □

Hence the Minkowski reading of Definition 12.5 and the Lorentz statement of the subsequent remark do not hold as written, and Theorem 13.1 establishes invariance under the involution J, not under the Lorentz group.

v4.8.0-r06 two admissible readings

Reading A (four-vector). On the real slice $\{q : q^\dagger = -q\}$, where $q^\dagger$ combines quaternion and complex conjugation, the z_k are real and $N(q) = -t^2 + x_1^2 + x_2^2 + x_3^2$ is exactly Minkowski $(1,3)$. The map $q \mapsto uqu^\dagger$ with $N(u) = 1$ preserves the slice and N, and changes the time component: these are genuine boosts. The kernels then carry no phase, so the 120° triole has no place.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Reading B (complex three-vector). The kernels form $z = (z_1, z_2, z_3) \in \mathbb{C}^3$, and $q \mapsto uqu^{-1}$ with $u \in \mathbb{C} \otimes \mathbb{H}$, $N(u) = 1$, fixes the scalar part and acts on z by $SO(3,\mathbb{C})$. Since $SO(3,\mathbb{C}) \cong PSL(2,\mathbb{C}) \cong SO^+(1,3)$, this is the proper orthochronous Lorentz group. The invariant is $s(z) := z_1^2 + z_2^2 + z_3^2$. The conjugation formula of the remark after Definition 12.5 becomes correct once u is taken complex. Formally, z is the Riemann–Silberstein vector $E + iB$, and $s = E^2 - B^2 + 2i\,E \cdot B$ collects the two field invariants.

Decision. Reading B is adopted as the working reading of the kernel (author's decision, 2026-10-19). Reading A remains available as the four-vector structure of $\mathbb{C} \otimes \mathbb{H}$; the two are not exclusive. In Reading B the scalar t is an invariant parameter, not the time coordinate of a four-vector; its role is open.

v4.8.0-r06 the triole is the light-like centred configuration

Theoremlet v4.8.0-r06.2. For $(z_1, z_2, z_3) \in \mathbb{C}^3$,

$$z_1 + z_2 + z_3 = 0 \;\text{ and }\; z_1^2 + z_2^2 + z_3^2 = 0 \quad\Longleftrightarrow\quad (z_1, z_2, z_3) = c\,(1, \omega, \omega^2) \text{ up to order},$$

with $\omega = e^{2\pi i/3}$ and $c \in \mathbb{C}$.

Proof sketch. With elementary symmetric functions e_1, e_2, e_3 and $p_2 = \sum z_k^2$ one has $p_2 = e_1^2 - 2e_2$. Hence $e_1 = p_2 = 0$ gives $e_2 = 0$, and the z_k are the roots of $\lambda^3 - e_3$, i.e. the three cube roots of e_3. Conversely $1 + \omega + \omega^2 = 0$ and $1 + \omega^2 + \omega^4 = 0$. □

The triole condition of Theorem 12.1, read with equal moduli and 120° phases, is therefore not only a phase condition: in Reading B it is equivalent to the Lorentz-invariant null condition $s = 0$ together with centring $\sum z_k = 0$.

v4.8.0-r06 invariant sectors

Theoremlet v4.8.0-r06.3. The orbits of $SO(3,\mathbb{C})$ on $\mathbb{C}^3$ are the level sets $\{s = c\}$ for $c \neq 0$, the punctured null cone $\{s = 0\} \setminus \{0\}$, and $\{0\}$. The duality rotation $z \mapsto iz$ maps $s \mapsto -s$ and preserves the null cone.

Proof sketch. For $s \neq 0$ put $e_1 = z/\sqrt{s}$, complete it to a complex-orthonormal basis with determinant 1, and rotate z to $(\sqrt{s}, 0, 0)$; this was checked constructively. Null vectors are treated through Theoremlet r06.4 below. $s(iz) = -s(z)$ is immediate. □

Sector reading (conjecture). The classification of §15 by the sign of $\operatorname{Re} z$ is not Lorentz-invariant in Reading B. A Lorentz-invariant candidate is: $s = 0$ null (photon-type); s real positive, electric-dominated; s real negative, magnetic-dominated. The duality $z \mapsto iz$ exchanges the two massive types and fixes the null sector, matching the reading of the tachyon as J-dual of matter and of the photon as fix-sector. The assignment electric $\leftrightarrow$ matter, magnetic $\leftrightarrow$ tachyon is a model decision and is not made here; the meaning of non-real s is open.

v4.8.0-r06 the null cone and the photon base

For $\psi = (\psi_1, \psi_2) \in \mathbb{C}^2$ put $z(\psi) := (\psi_1^2 - \psi_2^2,\ i(\psi_1^2 + \psi_2^2),\ -2\psi_1\psi_2)$.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Theoremlet v4.8.0-r06.4. $z(\psi)$ lies on the null cone, every non-zero null vector has exactly the two preimages $\pm\psi$, and the projectivized null cone $\{[z] \in \mathbb{C}P^2 : s(z) = 0\}$ is a smooth conic isomorphic to $\mathbb{C}P^1 \cong S^2$. Under the identification of Theorem 22.1 this is the photon base S_0. The map is equivariant: $SL(2,\mathbb{C})$ on ψ induces $SO(3,\mathbb{C})$ on z.

Proof sketch. $s(z(\psi)) = (a-b)^2 - (a+b)^2 + 4ab = 0$ with $a = \psi_1^2$, $b = \psi_2^2$. For a null z, set $\psi_1^2 = \frac{1}{2}(z_1 - iz_2)$, $\psi_2^2 = \frac{1}{2}(-z_1 - iz_2)$; the null condition gives $z_3^2 = 4\psi_1^2\psi_2^2$, which fixes the relative sign, so the preimage is $\pm\psi$. The conic is the Veronese image of $\mathbb{C}P^1$. Equivariance and invariance of the null cone were checked numerically for random unit biquaternions. □

Consequences. The symmetric triole $(1, \omega, \omega^2)$ is the image of the spinor $\psi \approx (0.9744 + 0.1283i,\ 0.3097 + 0.4036i)$. For the null axis of Theorem 14.3, the natural Lorentz-invariant definition in Reading B is $s = 0$, not $|q_n|^2 = 0$; its directions then form exactly S_0. The transport statement $\hat{K}(L_n) = L_{n-1}$ of that theorem is not addressed and remains as stated there.

Information-level reading (conjecture). A normalized ψ is a qubit; its Bloch vector is the Hopf image $S^3 \to S^2$, and $k(\psi) = (|\psi|^2, \psi^\dagger\sigma\psi)$ is a light-like four-vector. Every future time-like vector is a sum $k(\psi_a) + k(\psi_b)$ (checked constructively). This supports the author's conjecture that space-time is a projection of an information level, with single spinors as light-like and spinor pairs as massive readouts, in line with $S_0 = \mathbb{C}P^1$ and $S_1 = \mathbb{H}P^1$ of Theorem 22.1. No derivation of space-time points is claimed.

v4.8.0-r06 status of the affected statements

statement	status after r06
Def. 12.5, Minkowski reading	signature $(1,6)$; replaced by $N(q)$ and Reading B
Lorentz action by quaternion conjugation	correct only with complex u, $N(u) = 1$
Thm. 13.1	shows J-invariance; Lorentz invariance supplied by r06 (Reading B)
Sector classification, §15	not invariant; invariant candidate via s (conjecture)
Triole condition, Thm. 12.1	equivalent to centred null configuration (r06.2)
Null axis, Thm. 14.3	invariant definition $s = 0$; directions $= S_0$ (r06.4)
Prop. 12.7	unaffected

v4.8.0-r06 ledger and guard update

$$R_{\text{kern}}^{v480\text{-}r06} = (0,0,0,0,1,1)$$

for (signature computed; Lorentz action supplied in Reading B; triole lemma; null cone $= S_0$; sector assignment not chosen; role of t open).

The layer spends no guard:

$$R_{\text{glob}}^{(v480\text{-}r06,+)} = (0,1,1,1,1), \qquad \Gamma_{\text{guard}}^{v480\text{-}r06} = (1,1,1,1,1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.8.0-r07 Source-Only: Sector Assignment and Duality Angle in the Complex-Vector Reading

Purpose. Layer v4.8.0-r06 left the assignment of the two massive types of Reading B to matter and tachyon open. This layer adopts a working assignment, justifies it from the signed tachyon-frame scale of §A.269, and shows that every non-light-like kernel configuration has a unique duality angle, so that non-real values of s need no separate sector. Earlier layers are not rewritten.
Source-only status. This v4.8.0-r07 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.8.0 Version DOI 10.5281/zenodo.23365244 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.8.0-r07 working assignment

For $z \in \mathbb{C}^3$ let $s(z) = z_1^2 + z_2^2 + z_3^2$ as in r06. The working assignment is

$$s > 0 \ \text{ matter}, \qquad s < 0 \ \text{ tachyon}, \qquad s = 0 \ \text{ photon}.$$

It is adopted as the author's working choice (2026-10-19) and can be reversed without affecting the theoremlets below, which are stated for the sign of s.

Motivation. Layer r82 types the tachyon frame by the signed scale $\ell_T^2 = -\lambda_T^2$, $\lambda_T > 0$, an allowed effective statement of the release scope. Under the working assignment the sign of s carries exactly this interval signature: positive for matter, negative for tachyon, zero for the photon. It is not the orientation code of §A.270, which r01 keeps separate from the r82 signature.

Consequence for the earlier classification. The classification of §15 placed the photon kernel at purely imaginary z. In Reading B a purely imaginary non-zero $z = iy$ has $s = -|y|^2 < 0$ and is tachyon-type, not light-like. The earlier classification is therefore replaced, not translated.

v4.8.0-r07 duality angle

Theoremlet v4.8.0-r07.1. Every $z \in \mathbb{C}^3$ with $s(z) \neq 0$ can be written as

$$z = e^{i\alpha} z_m, \qquad s(z_m) = |s(z)| > 0,$$

with $\alpha = \frac{1}{2} \arg s(z)$ determined modulo π and z_m determined up to sign. The configuration is of matter type exactly when $\alpha \equiv 0$ and of tachyon type exactly when $\alpha \equiv \pi/2 \pmod{\pi}$.

Proof sketch. $s(e^{i\alpha} z_m) = e^{2i\alpha} s(z_m)$, so $s(z_m) = e^{-2i\alpha} s(z)$ is real positive iff $2\alpha \equiv \arg s(z) \pmod{2\pi}$. Replacing α by $\alpha + \pi$ replaces z_m by $-z_m$. The two special values follow from the sign of s. Checked numerically for 500 random configurations. □

Theoremlet v4.8.0-r07.2. Under the group $SO(3,\mathbb{C}) \times U(1)_{\text{dual}}$, acting by $z \mapsto e^{i\alpha} R z$, the orbits in $\mathbb{C}^3 \setminus \{0\}$ are the shells $\{|s| = r\}$, $r > 0$, and the punctured null cone. In particular, a matter configuration and a tachyon configuration with the same $|s|$ lie in one orbit, related by $\alpha = \pi/2$; the null cone is the only orbit fixed by the whole duality circle.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof sketch. By Theoremlet r06.3 the $SO(3,\mathbb{C})$-orbits are the level sets of s; the duality circle rotates s through all values of fixed modulus and fixes $s = 0$. Transitivity on each shell was checked constructively for 200 random pairs. □

Reading (conjecture). Non-real s is not a further sector: it is a matter configuration seen in a rotated duality frame. In this precise sense matter and tachyon are one object in two duality frames, distinguished Lorentz-invariantly but exchanged by duality, while the photon is the duality-invariant fix-sector. This is the form the initial conjecture of Theorem 12.1 takes in Reading B. Whether the duality angle is physical, or a gauge freedom to be fixed, is not decided here.

v4.8.0-r07 the parameter t

Observation. On the null set of the natural norm, $N(q) = -t^2 + s(z) = 0$, one has $s = t^2$. For real $t \neq 0$ this forces $\alpha \equiv 0$, the matter frame, with $t = \sqrt{|s|}$; for $t = 0$ it forces the photon sector; a tachyon-type kernel on this set requires $t^2 < 0$. The condition $N(q) = 0$ therefore ties t to the invariant magnitude $\sqrt{|s|}$ and fixes the duality frame. Whether the Sphaera kernel is meant to satisfy $N(q) = 0$ is open; the observation is recorded as a candidate role for t, not adopted.

v4.8.0-r07 ledger and guard update

$$R_{\text{sect}}^{v480\text{-}r07} = (0, 0, 0, 1, 1)$$

for (working assignment adopted; duality angle; orbit structure; physical status of the duality angle open; role of t open).

The layer spends no guard:

$$R_{\text{glob}}^{(v480\text{-}r07,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\text{guard}}^{v480\text{-}r07} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.8.0-r08 Source-Only: Test of Prediction P1 (Photon-Frame Unfolding)

Purpose. Corollary 42.24 lists P1 as testable now: unfolding the zeta zeros with the photon-frame density should improve the GUE fit by about 21% (Corollary 24.6), and Theorem 24.2 states that the Kolmogorov–Smirnov distance has a unique global minimum at $\mu = \gamma_L$. This layer performs the test on the first 2000 zeros. Earlier layers are not rewritten.
Source-only status. This v4.8.0-r08 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.8.0 Version DOI 10.5281/zenodo.23365244 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.8.0-r08 setup

As in Theorem 24.2, $\tilde{\gamma}_n(\mu) = \int_0^{\gamma_n} \frac{\log(t/2\pi\mu)}{2\pi\mu}\,dt = \frac{\gamma_n}{2\pi\mu}\big(\log\frac{\gamma_n}{2\pi\mu} - 1\big)$ and $s_n(\mu) = \tilde{\gamma}_{n+1}(\mu) - \tilde{\gamma}_n(\mu)$. The comparison law is the Wigner surmise $p(s) = \frac{32}{\pi^2}s^2e^{-4s^2/\pi}$; χ^2/dof uses 15 bins on $[0,3]$. The scale runs over $\mu \in [0.5, 3]$ in steps of 0.05. Two readings are tested: (a) raw, exactly as defined, and (b) normalized to mean spacing 1, since a pure rescaling would otherwise move the mean.

v4.8.0-r08 result

	zeros 1–100	zeros 101–2000
(a) raw: KS minimum	$\mu = 1.00$	$\mu = 1.00$
(a) KS at $\mu = 1$ / $\mu = \gamma_L$	0.068 / 0.699	0.041 / 0.628
(a) mean spacing at $\mu = \gamma_L$	0.419	0.473
(b) normalized: KS minimum	$\mu = 2.85$	$\mu = 2.25$
(b) KS range over μ	0.037–0.070	0.039–0.042
(b) χ^2/dof change $\mu = 1 \to \gamma_L$	+5.4% improvement	−9.9% (worse)

Theoremlet v4.8.0-r08.1. On the first 2000 zeros, neither reading of the photon-frame unfolding singles out $\mu = \gamma_L$. In reading (a) the unique minimum is the standard unfolding $\mu = 1$; in reading (b) the dependence on μ is flat within the statistical scatter and its minimum is not at γ_L. Moreover, in reading (b) the effect of μ vanishes with height: the local stretch it introduces is of relative size $\log\mu/\log(\gamma/2\pi)$, about 1% at $\gamma \sim 10^{22}$, so an asymptotic improvement of 21% cannot occur.

Proof sketch. Direct computation, script `k11_p1_entfaltung.py` in the working folder. For reading (b), $\rho_\mu(t)/\rho_1(t) = \mu^{-1}\big(1 - \log\mu/\log(t/2\pi)\big)$; after normalization the constant factor μ^{-1} drops out and only the second term, which decays like $1/\log t$, remains. □

Reproducibility note. The value of 75.8% for 100 zeros stated in Corollary 24.6 is not reproduced in either reading. The original script `gue_frame_correction.py` is not in the source folders, so its exact procedure could not be compared.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.8.0-r08 consequences

statement	status after r08
Thm. 24.2 (unique minimum at γ_L)	not confirmed; contradicted on zeros 1–2000
Cor. 24.6, P1 in Cor. 42.24	by its own criterion falsified at low height; the asymptotic form excluded by the $1/\log\gamma$ decay
γ_L as "GUE fixpoint of the frame"	withdrawn; γ_L remains an input (cf. r03 F11, OP1)

Corollary 24.6 states that non-observation falsifies $\gamma_L = \gamma_{\mathrm{BI}}/\gamma_{\mathrm{string}}$ as the physical frame factor. This layer records the non-observation on the accessible zeros and the reason why the stated asymptotic effect cannot appear. It does not by itself decide the physical status of γ_L; that remains with OP1.

v4.8.0-r08 ledger and guard update

$$R_{\mathrm{P1}}^{v480\text{-}r08} = (0, 0, 1)$$

for (test performed; prediction not confirmed; physical status of γ_L open).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v480\text{-}r08,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v480\text{-}r08} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.8.0-r09 Source-Only: Tests of P2 and P3, and Re-Classification of Earlier Results

Purpose. Corollary 42.24 lists P2 (Koide phase) as "already matches data" and P3 (β_{sym}-formula) as "matches to 0.018%". This layer tests both against the measurement precision and against the freedom in choosing the formula. Since layers r03–r08 changed the status of many earlier statements, it then collects the current status of all predictions and of the main results in one register, so that no earlier "proved" or "✓" is read without its present qualification. Earlier layers are not rewritten.
Source-only status. This v4.8.0-r09 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.8.0 Version DOI 10.5281/zenodo.23365244 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.8.0-r09 P2: Koide phase

Theorem 22.5 predicts, in the Brannen form $\sqrt{m_k} = A\big(1 + \sqrt{2}\cos(\theta_K + 2\pi k/3)\big)$, the phase $\theta_K = \pi - \sqrt{2}\arcsin\beta_{\text{sym}} = 2.319109$ rad $= 132.8751°$.

Theoremlet v4.8.0-r09.1. From $m_e = 0.51099895$, $m_\mu = 105.6583755$ and $m_\tau = 1776.93 \pm 0.09$ MeV the observed phase is $\theta_K^{\text{obs}} = 132.7325° \pm 0.0004°$ (with $m_\tau = 1776.86 \pm 0.12$: $132.7328° \pm 0.0005°$). The prediction differs by $+0.142°$, i.e. by roughly 300–400 standard deviations. P2 is excluded by the data.

Proof sketch. Solve the Brannen form for θ from the three masses; propagate the τ-mass error, which dominates. Script `k12_p2_p3.py` in the working folder. □

Remarks. (i) The relative deviation of 0.11% quoted in Theorem 22.5 is small compared with the angle, but large compared with the measurement error; a prediction is tested against the error. (ii) The observed phase is $\theta_K^{\text{obs}} = 2\pi/3 + 0.22223$, within 10^{-5} of $2\pi/3 + 2/9$; this empirical regularity is known in the literature and is not produced by the formula of Theorem 22.5. (iii) Step 1 of the proof of Theorem 22.5 derives $K = 2/3$ from Cauchy–Schwarz. Cauchy–Schwarz gives $(\sum\sqrt{m_i})^2 \le 3\sum m_i$, i.e. $K \ge 1/3$, with equality only for equal masses; it does not yield $K = 2/3$. The Koide value $K = 0.666665$ is an empirical fact; its derivation is open.

v4.8.0-r09 P3: the β_{sym}-formula

Remark 41.8 states $\beta_{\text{sym}} = \frac{1}{\sqrt{3}}\exp\big(-\gamma_1/(4\pi\gamma_L^5)\big)$.

Theoremlet v4.8.0-r09.2. With $\gamma_L = \gamma_{\text{BI}}/\gamma_{\text{string}} = 1.86444$ the formula gives 0.549233 against $\beta_{\text{sym}} = \sqrt{(\gamma_L - 1)/(\gamma_L + 1)} = 0.549348$, a deviation of -0.021%; it holds exactly for $\gamma_L = 1.86366$. It is a relation between the two inputs $\gamma_{\text{BI}}, \gamma_{\text{string}}$ and γ_1, not a prediction of an independently measured quantity. Among the 56 formulas $\frac{1}{\sqrt{3}}\exp(-\gamma_1/(c\,\gamma_L^n))$, $n = 1, \ldots, 8$, $c \in \{1, 2, \pi, 2\pi, 4\pi, 8\pi, 16\pi\}$, 27 lie within $\pm 5\%$; the chance that one of these lands within $\pm 0.02\%$ is about 11%.

Proof sketch. Both sides are functions of γ_L alone; solve by bisection. Enumerate the family and estimate the look-elsewhere probability with a uniform density over $\pm 5\%$. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Assessment. The exponent was changed from γ_L^2 to γ_L^5 after the fact (Remark 41.8), and Remark 41.8 itself marks it as requiring structural derivation. P3 is recorded as a numerical coincidence of moderate significance, not as a confirmed prediction. A derivation of the exponent before comparison with data would change this.

v4.8.0-r09 register of predictions

prediction (Cor. 42.24)	earlier status	status after r09
P1 GUE photon-frame unfolding	testable now	not confirmed (r08); minimum at $\mu = 1$
P2 Koide phase	matches to 0.111%	excluded by data, ~ 300–400σ (r09.1)
P3 β_{sym}-formula	matches to 0.018%	coincidence, exponent chosen post hoc; $\sim 11\%$ look-elsewhere (r09.2)
P4 Λ as frame artefact	reinterpretation	contradicted as stated (r03 F4); Λ type open (r03 F3); Θ-sum route false (r05)
P5 α-drift ratio	not yet testable	exponent corrected (r03 F2); untestable at present
P6 Hawking, tachyon frame	theoretical	assumption $k(M) = 0$ made explicit (r03 F6); untested

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.8.0-r09 register of main results

statement	current status (layer)
Thm. 12.1, equivalence of (i)–(v)	conjecture; Stone–von Neumann justification withdrawn (r04); Reading B gives a precise form via duality (r07)
Thm. 15.2	open (r04)
Def. 12.5, Minkowski reading	replaced: signature $(1,6)$; natural norm $N(q)$, Reading B (r06)
Thm. 13.1	shows J-invariance; Lorentz action from Reading B (r06)
Thm. 22.1	Hopf fibrations classical; physical identification conjecture; the chain $S^{15} \to S^8 \to S^4 \to S^2$ is not a composition of Hopf maps (recorded here)
Thm. 14.3	null axis redefined as $s = 0$, directions $= S_0$ (r06); transport open
triole structure	triole lemma proved (r06.2); octonionic triole configuration repaired (r04.4)
Heisenberg compensator, cascade	standalone, proved (r04.2, r04.3)
Thm. 22.19	structural conjecture; action undefined (r05)
Thm. 22.20	empirical fact; derivation withdrawn (r05)
Thm. 24.2	not confirmed (r08)
Thm. 22.5	$K = 2/3$ empirical, derivation open; phase formula excluded (r09)
Thm. 42.19	(i) holds by definition of the photon frame; (iii) value off by 10^{121} (r03 F8, F9)
Cor. 42.17	corrected to $\gamma_L^5 \approx 22.5$ (r03 F1)
$\gamma_L = \gamma_{\mathrm{BI}}/\gamma_{\mathrm{string}}$	ratio of two published Barbero–Immirzi values (r03 F11); input, not derived (OP1)
OP5 Step (B)	no valid form (r05)
RH	not proved; forbidden as global claim (W_γ^{glob}); Riemann source no proof source (r04)

Theoremlet v4.8.0-r09.3. After r09 no prediction of Corollary 42.24 stands as confirmed. The results that carry are the structural ones: the standalone compensator and cascade, the repaired triole hierarchy, the triole lemma, the signature analysis and Reading B with its orbit, null-cone and duality structure. None of the five guards is affected.

Proof sketch. Collect the decisions of r03–r09 as listed in the two registers. □

v4.8.0-r09 ledger and guard update

$$R_{\mathrm{pred}}^{v480\text{-}r09} = (0,0,0,1)$$

for (P2 decided; P3 classified; register compiled; derivations of $K = 2/3$ and of the P3 exponent open).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v480\text{-}r09,+)} = (0,1,1,1,1), \qquad \Gamma_{\mathrm{guard}}^{v480\text{-}r09} = (1,1,1,1,1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.8.0-r10 Source-Only: The Koide Structure in the Complex-Vector Reading

Purpose. Layer r09 excluded the phase formula of Theorem 22.5 and noted that the observed Koide phase lies close to $2\pi/3 + 2/9$. This layer asks what Reading B (r06, r07) says about the lepton masses before any number is fitted. Because the observed values are known, nothing found here can count as a prediction; the layer records what the structure fixes and what it leaves free. Earlier layers are not rewritten.
Source-only status. This v4.8.0-r10 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.8.0 Version DOI 10.5281/zenodo.23365244 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.8.0-r10 the Koide form is a triole

Theoremlet v4.8.0-r10.1. For $v = (\sqrt{m_e}, \sqrt{m_\mu}, \sqrt{m_\tau})$ the Brannen form $\sqrt{m_k} = A\big(1 + \sqrt{2}\cos(\theta + 2\pi k/3)\big)$ is identical to

$$v = t\,(1,1,1) + \sqrt{2}\ \mathrm{Re}\big(c\,(1,\omega,\omega^2)\big), \qquad t = A, \quad c = |c|e^{i\theta},$$

a scalar plus the real part of a symmetric triole. For any t, c, $K = \sum m_k/(\sum \sqrt{m_k})^2 = \frac{1}{3}\big(1 + |c|^2/t^2\big)$; hence $K = 2/3$ if and only if $|c| = t$. From the measured masses, $|c|/t = 0.999997$ and $\theta \bmod 2\pi/3 = 0.222225$.

Proof sketch. Discrete Fourier decomposition of v over $\mathbb{Z}/3$: the constant mode is t, the first mode is c; the reconstruction is exact. $\sum v_k^2 = 3t^2 + 3|c|^2$ and $\sum v_k = 3t$ give K. □

Theoremlet v4.8.0-r10.2. Embed the triole as the kernel part of Reading B, $q = it \cdot \mathbf{1} + \sum_k z_k e_k$ with $z = c(1,\omega,\omega^2)$. Then $s(z) = 0$ and $N(q) = -t^2$ for every c; the Reading-B invariants therefore fix neither $|c|/t$ (i.e. K) nor $\arg c$ (the Koide phase).

Proof sketch. $s = c^2(1 + \omega^2 + \omega^4) = 0$ by the triole lemma r06.2, and $N(q) = -t^2 + s$. □

Observation (conjecture). The condition $|c| = t$ is equivalent to $-t^2 + |z_k|^2 = 0$ for each kernel separately, i.e. each kernel is light-like in the original per-kernel norm of Definition 12.5. This equivalence depends on the factor $\sqrt{2}$ of the Brannen form, which is a convention; it is recorded as a candidate principle for $K = 2/3$, not as a derivation.

v4.8.0-r10 no canonical angle gives the phase

The following angles of the framework were fixed before comparison: $j\pi/12$ for $j = 0, \ldots, 7$; $\arcsin\beta_{\mathrm{sym}}$, $2\arcsin\beta_{\mathrm{sym}}$, $\arcsin(1/\sqrt{3})$, $2\arcsin(1/\sqrt{3})$, $\arctan\beta_{\mathrm{sym}}$, β_{sym} and $1/\sqrt{3}$ read as radians; the P2 angle; the tetrahedral and magic angles. Reduced modulo $2\pi/3$, the closest is the excluded P2 value (0.2247, 356σ away), the next $\pi/12$ (0.2618). None reproduces $\delta_{\mathrm{obs}} = 0.222225$. The value $2/9$ is not a rational multiple of π, which is what angular structures of this type produce.

v4.8.0-r10 what is missing: dynamics

Reading B is kinematic: it fixes how the kernels transform, not which configuration is realised.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Theoremlet v4.8.0-r10.3. The quadratic Lorentz-invariant real functionals of $z \in \mathbb{C}^3$ are $\mathrm{Re}(e^{-i\varphi} s) = \cos\varphi\,(E^2 - B^2) + \sin\varphi\, 2E \cdot B$ for $z = E + iB$, and a duality rotation $z \mapsto e^{ia} z$ maps $\varphi \mapsto \varphi - 2a$.

Proof sketch. s is the only quadratic $SO(3,\mathbb{C})$-invariant up to complex scale; its real-linear functionals are the real parts of $e^{-i\varphi} s$. $s(e^{ia} z) = e^{2ia} s(z)$. Checked numerically. □

Reading. $\varphi = 0$ is the Maxwell Lagrangian; $\sin\varphi\, 2E \cdot B$ is the topological θ-type term. The simplest dynamics of Reading B is thus electromagnetism, consistent with the photon sector, while the duality angle of r07 reappears as a vacuum angle. Lepton masses, $K = 2/3$ and the Koide phase require an additional principle that fixes $|c|/t$ and $\arg c$. Any such principle has to be stated before it is compared with the known values; only then can it yield a prediction.

v4.8.0-r10 ledger and guard update

$$R_{\mathrm{Koide}}^{v480\text{-}r10} = (0, 0, 0, 1, 1)$$

for (Koide form identified as triole; Reading-B invariants shown not to fix it; canonical angles excluded; principle for $|c| = t$ open; principle for the phase open).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v480\text{-}r10,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v480\text{-}r10} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.8.0-r11 Source-Only: Pre-Registered Test of the Phase Relation $\delta = K/3$

Purpose. Layer r10 wrote the lepton masses as a scalar plus a triole and left the phase open. For the charged leptons the phase satisfies $\delta \approx K/3 \approx 2/9$. This layer records the first pre-registered test of the Companion: the hypothesis that $\delta = K/3$ holds for every triplet of charged fermions. Hypothesis, definitions, data and decision criterion were committed to the repository before any computation (working-folder note `Vorab_Notiz_H1_Koide_Phase.md`, commit `81f38d3`). Earlier layers are not rewritten.
Source-only status. This v4.8.0-r11 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.8.0 Version DOI 10.5281/zenodo.23365244 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.8.0-r11 hypothesis and protocol

Hypotheses. H1: $\delta = K/3$ for each triplet. H1′: $\delta = 2/9$ for each triplet. For quarks $K \neq 2/3$, so the two differ. The factor 1/3 may be read as $1/N$ for $N = 3$ kernels or as $\beta^2_{\mathrm{sym,Hopf}}$; for triplets both readings predict the same number.

Protocol. $v_k = \sqrt{m_k}$ in ascending order, all roots positive; t and c as in Theoremlet r10.1; $\delta = \arg c$ reduced to $[0, 2\pi/3)$. Primary data: running $\overline{\mathrm{MS}}$ Yukawa couplings at $\mu = M_Z$ with PDG-2024 inputs (Antusch, Hinze, Saad, arXiv:2510.01312), for which masses are proportional to couplings and the common factor cancels. Error propagation by 10^6 Monte-Carlo draws. Criterion: $|z| < 2$ compatible, $|z| > 3$ refuted; H1 is refuted if it is refuted for at least one quark triplet.

v4.8.0-r11 result

triplet (at M_Z)	K	δ	$K/3$	z(H1)	z(H1′)
up quarks (u, c, t)	0.8876	0.0518	0.2959	−322	−353
down quarks (d, s, b)	0.7478	0.1001	0.2493	−115	−128
leptons (e, μ, τ)	0.6678	0.2211	0.2226	−49	−46
leptons, pole masses	0.6667	0.2222	0.2222	+0.4	+0.4

Theoremlet v4.8.0-r11.1. By the pre-registered criterion both H1 and H1′ are refuted. The relation $\delta \approx K/3 \approx 2/9$ holds only for the charged leptons with pole masses, where it served as calibration; it fails for both quark triplets and already for the running lepton masses at M_Z.

Proof sketch. Direct computation (script `k14_test_h1.py`, seed 20261020), following the committed protocol without deviation. The secondary data set (PDG-2024 masses at their standard scales) gives the same verdict. □

Assessment. The lepton coincidence is tied to pole masses, as is the Koide relation itself, and is not a law of the triole structure for all triplets. Hypotheses read off from the quark values found here would again be post-dictions and would need their own pre-registration. The procedure itself is retained: it is the Companion's first test that could have failed and was fixed before the data were examined.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.8.0-r11 ledger and guard update

$$R_{\mathrm{H1}}^{v480\text{-}r11} = (0, 0, 1)$$

for (protocol committed before computation; hypothesis decided; principle for the triole parameters still open).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v480\text{-}r11,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v480\text{-}r11} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.8.0-r12 Source-Only: Pre-Registered Neutrino Tests, the Octonionic Flavour Group, and Reclassification of the PMNS Claims

Purpose. Layer r11 introduced pre-registration. This layer records the seven further pre-registered tests H2–H8, which apply the triole form to neutrinos and to lepton mixing, the structural facts they rest on, and a reclassification of the earlier PMNS statements of the Companion. For every test the hypothesis, data and decision criterion were committed to the repository before the computation; the commits are listed below. Earlier layers are not rewritten.
Source-only status. This v4.8.0-r12 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.8.0 Version DOI 10.5281/zenodo.23365244 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.8.0-r12 structural facts

Theoremlet v4.8.0-r12.1 (triole spectrum and sign). For $v_k = t\big(1+\sqrt{2}\,r\cos(\delta + 2\pi k/3)\big)$ one has $v_0 v_1 v_2 = t^3\big(1 - \tfrac{3}{2}r^2 + \tfrac{r^3}{\sqrt{2}}\cos 3\delta\big)$. For $r = 1$ at most one component is negative, and exactly one is negative if and only if $\cos 3\delta < 1/\sqrt{2}$; the boundary $\delta = \pi/12$ is the point where one mass vanishes. The vector v is the spectrum of the Hermitian circulant matrix $t\,\mathbf{1} + \frac{c}{\sqrt{2}}P + \frac{\bar{c}}{\sqrt{2}}P^2$ (P the cyclic shift), whose eigenvectors are the fixed Fourier vectors.

Proof sketch. Expand the product with $\sum_k \cos\theta_k = 0$, $\sum_k \cos^2\theta_k = 3/2$ and $\prod_k \cos\theta_k = \frac{1}{4}\cos 3\delta$ for $\theta_k = \delta + 2\pi k/3$. The circulant statement is the discrete Fourier diagonalization. The identities were also checked numerically to 10^{-15}. □

Theoremlet v4.8.0-r12.2 (octonionic flavour group). In the Cayley–Dickson convention of layer r04 the automorphisms of $\mathbb{O}$ that permute $e_1, \ldots, e_7$ up to sign form a group of order 1344. Its stabilizer of $+e_7$ has order 96, acts unitarily with determinant one and irreducibly on $\mathbb{C}^3 = \mathrm{span}_{\mathbb{C}}(e_4, e_5, e_6)$ with complex structure L_{e_7}, and has the same element-order statistics and the same number of conjugacy classes (ten) as $\Delta(96) \subset SU(3)$.

Proof sketch. An automorphism is fixed by the images of e_1, e_2, e_4; all $14 \cdot 12 \cdot 8$ choices extend and preserve the full multiplication table. Restriction, unitarity, determinant, irreducibility ($\langle|\chi|^2\rangle = 1$) and the class data were computed exhaustively (scripts `k22_oktonion_triplett.py`, `k23_test_h6.py`). The identification with $\Delta(96)$ is by these invariants; an explicit isomorphism is not written out here. □

Theoremlet v4.8.0-r12.3 (residual reflections). With the triole $e_4 \to e_5 \to e_6$ as charged-lepton symmetry and the electron along its invariant direction, the fifteen involutions of the stabilizer preserve the following columns of the lepton mixing matrix:

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Involution	count	preserved $\lvert U_{\alpha i}\rvert^2$	pattern
pure sign changes (one generation fixed)	3	$(\frac{1}{3}, \frac{1}{3}, \frac{1}{3})$	TM2
Fano centre in $\{e_1, e_2, e_3\}$, first lift	3	$(\frac{2}{3}, \frac{1}{6}, \frac{1}{6})$	TM1
Fano centre in $\{e_1, e_2, e_3\}$, second lift	3	$(0, \frac{1}{2}, \frac{1}{2})$	$\theta_{13} = 0$
Fano centre in $\{e_4, e_5, e_6\}$	6	$(\frac{1}{3}, \frac{2 \mp \sqrt{3}}{6})$	$\Delta(96)$ column
Fano centre e_7	0	—	only order-4 lifts

The TM1 lift and the $\theta_{13} = 0$ lift over the same Fano image differ by a pure sign change.

Theoremlet v4.8.0-r12.4 (Cayley–Dickson involution as CP). The involution $p + q\ell \mapsto p - q\ell$ commutes with every lift of the triole and with all TM1 and $\theta_{13} = 0$ involutions, and acts on the triplet as $z \mapsto -\bar{z}$ in the basis in which the triole is the plain cyclic shift. If the Majorana mass matrix is invariant under it and under a residual reflection, then $\sin^2 \theta_{23} = \frac{1}{2}$ and $\delta_{CP} \in \{90°, 270°\}$ exactly.

Proof sketch. $X = -\mathbf{1}$ forces a real mass matrix in the generation basis; the Fourier matrix of the triole turns this into the μ–τ reflection, whose consequences are classical. Checked on 2000 random points of the four-dimensional solution space for each reflection type (script `k26_test_h8.py`). □

v4.8.0-r12 pre-registered tests

Data: NuFIT 6.0 (IC24 with SK atmospheric data, normal ordering unless stated), DESI DR2 with CMB for $\sum m_\nu < 0.0642$ eV, KamLAND-Zen for $m_{\beta\beta}$; Gaussian approximation with asymmetric errors where a χ^2 is quoted.

Test	commit	hypothesis	result under the fixed criterion
H2	`434c5a8`	neutrinos are an $r = 1$ triole, signs free	compatible: normal ordering, $\sum m_\nu = 0.0592 \pm 0.0002$ eV; inverted ordering disfavoured (0.102 eV); weakly discriminating
H3	`e3b140e`	triole signs are Majorana CP parities	compatible, $m_{\beta\beta} = 2.7 \pm 0.7$ meV (≤ 3.5 meV); not testable in the foreseeable future
H4	`00d297e`	TM1 against TM2, equal prior	TM1 strongly preferred, $\chi^2 = 4.4$ vs. 14.7 (2 d.o.f.); the θ_{12} part was not blind
H5	`16eff15`	polar angle of the preserved column is a Hopf angle of the Companion	failed: 0/32 canonical combinations compatible; free azimuth degenerate
H6	`e1e2393`	list of algebraic selection criteria for the TM1 lift	main criterion (dihedral S_3 with the triole) selects the $\theta_{13} = 0$ lift, i.e. speaks against TM1
H7	`e6da7db`	Heisenberg compensator J as sign automorphism selects the lift	no selection for any J; the candidates negating e_1 and fixing e_7 do not commute with the triole
H8	`f6734ab`	Cayley–Dickson involution as CP symmetry	tension, $\chi^2 = 8.8$ (3 d.o.f.); fixed predictions $\sin^2 \theta_{23} = \frac{1}{2}$, $\delta_{CP} = 270°$, with TM1 $\sin^2 \theta_{12} = 0.318$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Reading. The octonionic structure already present in the Companion produces the known mixing patterns TM1, TM2 and $\Delta(96)$ as candidates and, through the Cayley–Dickson involution, a definite CP structure. It does not select TM1: neither minimality, nor the cascade projections (they commute with all sign changes), nor the compensator distinguishes the TM1 lift from the excluded one. The predictions of H4 and H8 will be tested by JUNO (θ_{12}) and by DUNE and Hyper-Kamiokande (θ_{23}, δ_{CP}).

v4.8.0-r12 reclassification of earlier PMNS statements

Proposition 42.12. The value $\arctan\sqrt{2} = 54.74°$ is not the maximal mixing angle; maximal mixing is 45°. The shifted value $\arctan(\sqrt{2})\,\delta_{\mathrm{CL}} = 52.1°$ gives $\sin^2\theta_{23} = 0.622$, against 0.470 (normal ordering, $\theta_{23} = 43.3°$) about 9σ in the Gaussian approximation and about 6σ against the inverted-ordering value 0.550. The proposition and the $\theta_{23}^{\mathrm{PMNS}}$ line of the unified prediction table are reclassified as *refuted*. The replacement statement of this layer is Theoremlet r12.4 with the open selection problem.

Theorem 42.14. Its CKM and PMNS lines rest on the continuum-limit conjecture for δ_{CL}, whose β-formula was classified in layer r09 as a coincidence with an exponent chosen after the fact. The CKM line inherits that status; the PMNS line is refuted as above. The CKM phase remark ($\pi/3$ against $\approx 70°$) is unaffected and remains open.

v4.8.0-r12 ledger and guard update

$$R_{\mathrm{mix}}^{v480\text{-}r12} = (0, 0, 1)$$

for (protocols committed before computation; all seven hypotheses decided; selection principle for the residual reflection still open).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v480\text{-}r12,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v480\text{-}r12} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.8.0-r13 Source-Only: JUNO 2026 Re-Check, Associator of the Generation Triple, and the Colour–Generation Assignment

Purpose. This layer re-checks the fixed predictions of layer r12 against the JUNO data of September 2026, records two further pre-registered items (H9, H10), states an obstruction for quark statements, and compares the local line with the independent v4.9.0 research candidate. Earlier layers are not rewritten.
Source-only status. This v4.8.0-r13 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.8.0 Version DOI 10.5281/zenodo.23365244 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.8.0-r13 re-check against JUNO 2026

JUNO (arXiv:2609.37895, 207.2 live days) reports $\sin^2\theta_{12} = 0.3036 \pm 0.0064$, $\Delta m^2_{21} = (7.388 \pm 0.078) \times 10^{-5}$ eV^2, $\Delta m^2_{31} = +2.509^{+0.027}_{-0.025} \times 10^{-3}$ eV^2 for normal ordering, and prefers normal ordering by at least 2.1σ. These data existed when H2–H8 were fixed but had not been used (the data basis was NuFIT 6.0); they are an independent re-check, not a test fixed in advance of the measurement. Data choice made before the computation: $\sin^2\theta_{12}$ is the inverse-variance mean of NuFIT 6.0 and JUNO, 0.3046 ± 0.0056; for H2 the JUNO mass splittings; all other inputs as in r12. The decision criteria of r12 are unchanged.

Test	NuFIT 6.0 (r12)	with JUNO 2026	verdict under the fixed criterion
H2	normal ordering, $\sum m_\nu = 0.0592$ eV	$\sum m_\nu = 0.0591$ eV; normal ordering preferred	compatible, ordering supported
H4	$\chi^2(\mathrm{TM1}) = 4.4$, $\chi^2(\mathrm{TM2}) = 14.7$	9.6 and 49.1	TM1 far ahead of TM2 but disfavoured on its own ($\chi^2 > 9$, 2 d.o.f.)
H8	$\chi^2 = 8.8$, tension	$\chi^2 = 14.0$	disfavoured (≥ 11.34, 3 d.o.f.)

The shift is driven by θ_{12}: TM1 requires $\sin^2\theta_{12} = 0.3182$, 2.4σ above the combined value.

H9 (note committed as `b75170d`). The relation $|U_{e2}|^2 = \beta^2_{\mathrm{sym}}$, found after the fact in the evaluation of H5, gives $\sin^2\theta_{12} = \beta^2_{\mathrm{sym}}/\cos^2\theta_{13} = 0.30862 \pm 0.00018$ with $\beta_{\mathrm{sym}} = 0.549348$ from $\gamma_{\mathrm{string}} = \ln 2/(\pi\sqrt{3})$ and the inserted $\gamma_{\mathrm{BI}} = 0.2375$. Its present distance to JUNO (0.8σ) is disclosed and does not count. The test is the next JUNO result with $\sigma \leq 0.004$: $|z| < 2$ compatible, $|z| \geq 3$ refuted.

v4.8.0-r13 associator of the generation triple

Theoremlet v4.8.0-r13.1. In the convention of layer r04, $[e_4, e_5, e_6] = (e_4 e_5)e_6 - e_4(e_5 e_6) = -2e_7$. The associator vanishes on Fano lines and has norm 2 on non-collinear triples. Hence every element of the stabilizer of $+e_7$ fixes the associator of the generation triple.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof sketch. From the multiplication of r04, $e_4e_5 = e_1$ and $(e_4e_5)e_6 = e_1e_6 = -e_7$, while $e_5e_6 = -e_3$ and $e_4(e_5e_6) = -e_4e_3 = e_7$; the difference is $-2e_7$. All triples were also evaluated numerically (script `k27_test_h10.py`). An automorphism fixing e_7 maps $[e_4, e_5, e_6]$ to $[he_4, he_5, he_6] = h(-2e_7) = -2e_7$. □

H10 (note committed as `a6b5bcb`). Four associator criteria were fixed in advance for selecting the TM1 lift over the $\theta_{13} = 0$ lift. None separates the two, for any of the four lifts of the triole. Together with r12 (minimality, cascade projections, compensator) no internal selection principle for the TM1 lift has been found.

v4.8.0-r13 colour and generation

Theorem 22.4 places the generations on e_4, e_5, e_6, the triplet of $SU(3) \subset G_2$, and its proof gives each generation one colour index. In the Dixon reading this $SU(3)$ is the colour group, so one octonionic triplet is made to carry both colour and generation. Quarks require 3×3 such labels. Consequences: (i) the octonionic flavour analysis of r12 is mathematically correct but physically conditional on reading the triplet as generation space; read as colour, a broken flavour symmetry inside $SU(3)_c$ would break colour; (ii) quark values of the triole radius cannot be addressed before colour and generation are separated. In addition, the quark radii are scheme dependent at the 2–4% level ($r_u = 1.290$ at M_Z against 1.244 at standard PDG scales; $r_d = 1.115$ against 1.093), about a hundred times the statistical error, so a structural match would carry no weight. No fit is made.

v4.8.0-r13 comparison with the v4.9.0 research candidate

The repository's v4.9.0 research candidate (independent of this line) agrees with it on the Koide condition as an equal-norm law without derivation, on rejecting the phase ansatz $\pi - \sqrt{2}\arcsin\beta$, and on not asserting a Riemann or GUE theorem. It finds, independently, the positive phase chambers $|\theta - 2\pi j/3| \leq \pi/12$ with a parametrically light state near the boundary, the boundary of Theoremlet r12.1. It admits only non-negative roots for the physical Koide ratio; H2 and H3 are therefore reclassified as tests of the *signed-root* Koide relation (for the H2 masses the non-negative-root ratio is 0.525). The colour–generation assignment above concerns both lines.

v4.8.0-r13 ledger and guard update

$$R_{\text{mix}}^{v480\text{-}r13} = (0, 0, 1, 1)$$

for (protocols committed; hypotheses decided; selection principle open; colour–generation separation open).

The layer spends no guard:

$$R_{\text{glob}}^{(v480\text{-}r13,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\text{guard}}^{v480\text{-}r13} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.8.0-r14 Source-Only: Separating Colour and Generation

Purpose. Layer r13 recorded that Theorem 22.4 puts colour and generation on the same octonionic triplet. This layer examines two ways of separating them, both built from structures already present in the Companion, and records the consequences for layers r12 and r13. It is model construction; no prediction is made and no data are compared. Earlier layers are not rewritten.
Source-only status. This v4.8.0-r14 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.8.0 Version DOI 10.5281/zenodo.23365244 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.8.0-r14 requirements

A separation should (a) keep colour as $SU(3) \subset G_2$, the stabilizer of e_7; (b) count generations by a symmetry that commutes with G_2, so that a broken generation symmetry does not break colour; and (c) provide nine linearly independent coloured states, three colours times three generations.

v4.8.0-r14 route A: sedenions

Theoremlet v4.8.0-r14.1. Extend the Cayley–Dickson convention of layer r04 once more, $\mathbb{S} = \mathbb{O} \oplus \mathbb{O}e_8$. The maps ψ, a rotation by $2\pi/3$ in each of the seven planes (e_i, e_{i+8}), and $\varepsilon\colon a + be_8 \mapsto a - be_8$ are automorphisms of $\mathbb{S}$, satisfy $\psi^3 = \varepsilon^2 = 1$ and $\varepsilon\psi\varepsilon = \psi^{-1}$, and commute with G_2 acting diagonally. On the fourteen plane directions $1 + \psi + \psi^2 = 0$. Under $G_2 \times S_3$,

$$\mathbb{S} = (\mathbf{1}, \mathbf{1}) \oplus (\mathbf{1}, \mathrm{sgn}) \oplus (\mathbf{7}, \mathbf{2}).$$

Coloured states therefore carry only the two-dimensional representation of S_3: the nine vectors $\psi^k(e_c)$, $c = 4, 5, 6$, $k = 0, 1, 2$, span a space of dimension 6.

Proof sketch. Automorphism, group relations and commutation were checked numerically for random arguments, for all 1344 signed-permutation automorphisms and for random elements $\exp D$ of the fourteen-dimensional derivation algebra (script `k28_sedenion_generationen.py`); the colour stabilizer of e_7 has dimension 8. The traces $\mathrm{tr}\,\psi = -5$, $\mathrm{tr}\,\varepsilon = 0$ fix the decomposition. That $\mathrm{Aut}(\mathbb{S})$ is exactly $G_2 \times S_3$ is taken from the literature (Gillard and Gresnigt, arXiv:2306.13098, and references there), not re-proved. □

Route A separates colour from generation, but the triole relation $z_1 + z_2 + z_3 = 0$ itself makes the three copies of each coloured state linearly dependent. It yields only two independent coloured generations.

v4.8.0-r14 route B: the triality form

Theoremlet v4.8.0-r14.2. On $\mathbb{O} \oplus \mathbb{O} \oplus \mathbb{O}$ the trilinear form $t(x, y, z) = \mathrm{Re}\big((xy)z\big)$ is invariant under the cyclic shift $(x, y, z) \mapsto (y, z, x)$, under $(x, y, z) \mapsto (\bar{x}, \bar{z}, \bar{y})$, which together generate S_3, and under G_2 acting diagonally, which commutes with this S_3. The three copies of the colour triplet (e_4, e_5, e_6) give nine linearly independent coloured states.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof sketch. The real part of the associator vanishes and $\mathrm{Re}(ab) = \mathrm{Re}(ba)$, so $\mathrm{Re}((xy)z) = \mathrm{Re}(x(yz)) = \mathrm{Re}((yz)x)$. For the reflection, $\mathrm{Re}((\bar{x}\bar{z})\bar{y}) = \mathrm{Re}(y(zx)) = \mathrm{Re}((xy)z)$. Elements of G_2 preserve multiplication and conjugation. Independence holds because the three copies lie in different summands; all statements were also checked numerically. □

Proposed assignment. Generation index = the three triality slots; colour index = the $SU(3)$ triplet e_4, e_5, e_6 within each slot. This matches the Companion's identification of $S_2 \cong S^8 = \mathbb{O}P^1$ with three generations, and the fact that G_2 is the subgroup of Spin(8) fixed by triality. The assignment is proposed, not derived; a native carrier for the three slots is still to be supplied.

v4.8.0-r14 consequences

- After the separation, the order-96 stabilizer of Theoremlet r12.2 lies inside the colour group and cannot act as a flavour symmetry. Theoremlets r12.2–r12.4 remain correct mathematics but lose their reading as lepton-mixing statements; H4 and H8 were already disfavoured by the JUNO 2026 re-check of layer r13.
- A pure generation S_3 with the triole $\mathbb{Z}_3$ for charged leptons and a transposition for neutrinos preserves the column $(0, \frac{1}{2}, \frac{1}{2})$, i.e. $\theta_{13} = 0$, which is excluded. This is the minimality outcome of H6. Realistic mixing requires further breaking, from which no prediction follows yet.
- The triole form $t(1,1,1) + \sqrt{2}\,\mathrm{Re}(c(1,\omega,\omega^2))$ of layer r10 is the decomposition of generation space under S_3 into the symmetric direction and the doublet; $K = 2/3$ remains the equal-norm condition between them and is still not derived.

v4.8.0-r14 ledger and guard update

$$R_{\mathrm{cg}}^{v480\text{-}r14} = (0,1,1)$$

for (separation consistent with G_2 achieved by route B; native carrier of the triality slots open; mixing mechanism open).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v480\text{-}r14,+)} = (0,1,1,1,1), \qquad \Gamma_{\mathrm{guard}}^{v480\text{-}r14} = (1,1,1,1,1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.8.0-r15 Source-Only: The Octonionic Hopf Fibration as Carrier of the Triality Slots

Purpose. Layer r14 proposed the three triality slots of $\mathrm{Re}((xy)z)$ as generation index but left their carrier open. This layer shows that the octonionic Hopf fibration of Theorem 22.1, which the Companion already attaches to $S_2 \cong \mathbb{O}P^1$ and to three generations, carries the slots. Model construction only; no prediction is made and no data are compared. Earlier layers are not rewritten.
Source-only status. This v4.8.0-r15 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.8.0 Version DOI 10.5281/zenodo.23365244 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.8.0-r15 unit triples and the Hopf map

Theoremlet v4.8.0-r15.1. Let $\mathcal{T} = \{(a,b,c) \in (S^7)^3 : (ab)c = 1\}$. Then $\mathcal{T}$ is a fourteen-dimensional manifold (a, b free, $c = \overline{ab}$), closed under the cyclic shift $(a,b,c) \mapsto (b,c,a)$ and under $(a,b,c) \mapsto (\bar{a},\bar{c},\bar{b})$, and invariant under G_2 acting diagonally. For the Hopf map $h(x,y) = (|x|^2 - |y|^2,\, 2x\bar{y})\colon S^{15} \to S^8$, the points over the equator $|x| = |y| = 1/\sqrt{2}$ give unit triples $(\sqrt{2}\,x, \sqrt{2}\,\bar{y}, \bar{v}) \in \mathcal{T}$ with $v = 2x\bar{y}$, and the fibre over v is $\{(a, \bar{a}v)\} \cong S^7$.

Proof sketch. $c = \overline{ab}$ gives $(ab)c = |ab|^2 = 1$. By alternativity $b(\bar{b}\bar{a}) = \bar{a}$, so $bc = \bar{a}$ and $(bc)a = 1$; the remaining cyclic and reflected identities follow in the same way. Elements of G_2 preserve products and conjugation. The Hopf statements are the norm identity $(|x|^2 - |y|^2)^2 + 4|x|^2|y|^2 = 1$ and $a(\bar{a}v) = v$. All identities were also checked numerically (script `k29_hopf_triality_traeger.py`). □

Theoremlet v4.8.0-r15.2 (symmetric point). $(a,a,a) \in \mathcal{T}$ if and only if $a^3 = 1$, i.e. $a = \exp(2\pi u/3)$ for an imaginary unit u. For $u = e_7$ the point is ω in $\mathrm{span}(1, e_7)$, the complex plane at the end of the cascade, and its stabilizer in G_2 is the colour group $SU(3)$ (dimension 8).

Proof sketch. a and its powers lie in the commutative subalgebra $\mathrm{span}(1,u) \cong \mathbb{C}$; the stabilizer of a in G_2 is the stabilizer of u. The dimension was computed from the derivation algebra. □

v4.8.0-r15 reading and a check

The two components of the Hopf total space and the base point are the three triality slots; the triole is their cyclic symmetry, and G_2 acts as the colour side. Choosing the symmetric generation point fixes u and breaks G_2 to the colour $SU(3)$. This gives a natural carrier for the assignment of r14; it does not derive the existence of three generations.

Check: in $\mathrm{span}(1, e_7)$ the condition $(ab)c = 1$ becomes $\alpha_1 + \alpha_2 + \alpha_3 \equiv 0 \pmod{2\pi}$ for the phases. Reading the Koide phases $\delta + 2\pi k/3$ of layer r10 as these phases requires $3\delta \equiv 0$, against $\delta = 0.2222$ for the charged leptons. The direct reading fails. That $3\delta \approx 2/3$ is the known value $2/9 \cdot 3$, whose generality was refuted in r11; it is not counted. How a mass hierarchy and $K = 2/3$ could arise from deviations from the symmetric triple remains open.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.8.0-r15 ledger and guard update

$$R_{\mathrm{cg}}^{v480\text{-}r15} = (0, 0, 1)$$

for (separation consistent with G_2; carrier of the triality slots identified; mass mechanism open).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v480\text{-}r15,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v480\text{-}r15} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.8.0-r16 Source-Only: Pre-Registered Test of Masses from Invariants on the Triple Space

Purpose. Layer r15 left open whether a mass hierarchy can arise on the space of triality triples. This layer records the pre-registered test H11 (note `Vorab_Notiz_H11_Massen_Tripelraum.md`, commit `8e5eaa9`, fixed before computing): are the masses the slot norms at critical points of a natural invariant function? Earlier layers are not rewritten.
Source-only status. This v4.8.0-r16 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.8.0 Version DOI 10.5281/zenodo.23365244 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.8.0-r16 setting and hypotheses

For $(a,b,c) \in \mathbb{O}^3$ put $n_k = |\text{slot } k|$, $S = \sum n_k^2$, $Q = \sum n_k^4$, $P = n_1 n_2 n_3$ and $t = \mathrm{Re}((ab)c)$, $|t| \le P$. These generate the invariants of G_2 (diagonal) $\times S_3$ up to degree four. The fixed reading is $\sqrt{m_k} \propto n_k$, so $K = \sum n_k^2 / (\sum n_k)^2$.

H11a: the masses are the slot norms at critical points of t on $\{S = 3\}$;

H11b: the masses are the slot norms at minima of $V = \alpha S + \beta S^2 + \gamma Q - \kappa t$, bounded below.

Criterion (fixed): refuted if no critical point, respectively no minimum, has three distinct non-zero norms.

v4.8.0-r16 results

Theoremlet v4.8.0-r16.1. At every critical point of V with all $n_k \neq 0$ and $t = P$, each $x_k = n_k^2$ solves the same equation

$$(2\alpha + 4\beta S)\, x + 4\gamma\, x^2 = \kappa P.$$

Hence the slot norms take at most two distinct values. The same holds for the critical points of t on $\{S = 3\}$, which have equal norms ($K = 1/3$) or two empty slots ($K = 1$).

Proof sketch. $\partial V / \partial n_k = 0$ multiplied by n_k gives the displayed equation, since $n_k\, \partial P / \partial n_k = P$ for every k. A quadratic has at most two roots. For H11a the Lagrange conditions give $|bc| = \lambda |a|$ and its cyclic images. □

Numerically (script `k30_test_h11.py`) the global minima on a grid of 210 bounded parameter points are: 180 with equal norms, 25 with one occupied slot, 5 flat ($\gamma = \kappa = 0$, a whole sphere of minima), none with three distinct norms. One deviation from the note is disclosed: the relative tolerance for "distinct" was raised from 10^{-6} to 10^{-4} because the smaller value counted numerical noise in favour of the hypothesis. The verdict does not depend on it, since Theoremlet r16.1 excludes three distinct norms analytically.

Verdict. H11a and H11b are refuted under the fixed criterion. Three distinct norms need a term of degree at least six, which makes the equation cubic; this is necessary, not shown to be sufficient (no such minimum appeared in a tested sextic grid). Any such extension adds at least two parameters, so the two charged-lepton ratios could be fitted but not predicted, and $K = 2/3$ would be a condition on the parameters rather than a consequence. The triole gives the form of the Koide relation (r10), not its dynamics.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.8.0-r16 ledger and guard update

$$R_{\mathrm{cg}}^{v480\text{-}r16} = (0, 0, 1)$$

for (separation consistent with G_2; carrier of the triality slots identified; mass mechanism open, with invariants up to degree four excluded).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v480\text{-}r16,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v480\text{-}r16} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.8.0-r17 Source-Only: Time Direction of the Cosmological Fractions

Purpose. Layer r03 examined the assignment $\Omega_\Lambda = 1 - \beta_{\text{sym}}^2$ of Theorem 22.16 at the present epoch (§A.271) and the provenance of β_{sym} (§A.271). Two hypothesis-grade statements that make the same assignment time-dependent, Proposition 42.5 and Proposition 42.8, have not been checked by any layer since r85 and do not appear in the registers of §A.271. This layer compares them with the measured expansion history. It is a consistency check of existing statements, not a new prediction, and therefore carries no pre-registration note. The computation is `llm/k31_kosmologie_vorzeichen.py`. Earlier layers are not rewritten.
Source-only status. This v4.8.0-r17 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.8.0 Version DOI 10.5281/zenodo.23365244 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.8.0-r17 the three statements

1. Theorem 22.16: $\Omega_\Lambda = P(S^8 \to S^2) = 1 - \beta_{\text{sym}}^2$ and $\Omega_M = P(S^8 \to S^4) = \beta_{\text{sym}}^2$.
2. Proposition 42.5: β_{sym} evolves, with $\beta_{\text{sym}}(z = 1100) \approx 0.527$ and $\beta_{\text{sym}}(z = 0) \approx 0.5493$; "the universe moves continuously away from the photon frame".
3. Proposition 42.8, stated there as "the consistent continuation" of item 2: $\beta_{\text{sym}}(t) \nearrow 1$, $P(S^8 \to S^4) = \beta_{\text{sym}}^2 \to 1$, "the universe becomes fully matter dominated", and the expansion stops.

Items 2 and 3 use item 1 at all epochs, not only today; item 3 does so explicitly through $\beta_{\text{sym}}^2 \to 1$.

v4.8.0-r17 comparison with the expansion history

In flat ΛCDM with the Planck 2018 values $\Omega_M = 0.3111$, $\Omega_r = 9.1 \times 10^{-5}$, the Λ fraction at redshift z is $\Omega_\Lambda(z) = \Omega_\Lambda / \big(\Omega_r(1+z)^4 + \Omega_M(1+z)^3 + \Omega_\Lambda\big)$. Under item 1 the value of β_{sym} that reproduces it is $\sqrt{1 - \Omega_\Lambda(z)}$:

z	$\Omega_\Lambda(z)$	required β_{sym}
−0.5	0.947	0.231
0	0.689	0.558
0.5	0.396	0.777
1	0.217	0.885
2	0.0757	0.961
5	0.0101	0.995
1100	1.3×10^{-9}	1.000

Proposition 42.5 gives instead $\Omega_\Lambda(1100) = 1 - 0.527^2 = 0.722$, larger than the ΛCDM value by a factor 5.8×10^8.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Theoremlet v4.8.0-r17.1. Under the assignment of Theorem 22.16 held at all epochs, the measured expansion history requires $\mathrm{d}\beta_{\mathrm{sym}}/\mathrm{d}t < 0$: β_{sym} decreases from $\beta_{\mathrm{sym}} \approx 1$ at recombination to 0.558 today and tends to 0 in the Λ-dominated future. Propositions 42.5 and 42.8 assume $\mathrm{d}\beta_{\mathrm{sym}}/\mathrm{d}t > 0$ and an end state $\beta_{\mathrm{sym}} \to 1$. The three statements together describe a universe that starts Λ-dominated and ends matter-dominated, the reverse of the observed history. They are therefore not jointly tenable.

Proof sketch. $\Omega_\Lambda(z)$ is monotonically decreasing in z for $\Omega_M, \Omega_\Lambda > 0$, so $\sqrt{1-\Omega_\Lambda(z)}$ is increasing in z, i.e. decreasing in cosmic time. The table evaluates it. Items 2 and 3 state the opposite monotonicity. □

What the comparison does not decide. The present-epoch agreement $1-\beta_{\mathrm{sym}}^2 = 0.698$ against $\Omega_\Lambda = 0.689$ (1.4%) is untouched, as are the caveats of r03 F3–F5 and F11. The check uses ΛCDM as the description of the measured history; a different background would change the numbers, not the sign, because the ratio of matter to dark-energy density scales as a^{3w} and falls with expansion for every dark-energy equation of state $w < 0$. This also removes a possible affinity with the current evidence for evolving dark energy: that evidence concerns the dark-energy density, while Proposition 42.5 concerns the fraction, which increases with time in every such model.

Ways out, recorded and not chosen.

(a) Theorem 22.16 holds only at the present epoch. Then Propositions 42.5 and 42.8 lose their basis, and the 1.4% agreement becomes a statement about one epoch, which r03 F11 already qualifies.

(b) β_{sym} runs the other way, from ≈ 1 at recombination downward. Then the early universe sits at what Proposition 42.8 calls the tachyon limit, and Proposition 42.5 would need the opposite sign of $\Delta\beta_{\mathrm{sym}}$, which it would have to re-derive.

(c) The cascade probabilities are not the cosmological fractions at other epochs, but some other function of them. Then item 1 needs a time-dependent bridge that the source does not supply.

Which way is taken is left to the author.

v4.8.0-r17 register entries

statement	status after r17
Thm. 22.16	present-epoch value 1.4% from Planck; not tenable at all epochs together with Props. 42.5, 42.8 (r17.1)
Prop. 42.5	sign of $\mathrm{d}\beta_{\mathrm{sym}}/\mathrm{d}t$ opposite to the one required by Thm. 22.16; $\Omega_\Lambda(1100)$ off by 5.8×10^8 (r17)
Prop. 42.8	end state $\Omega_M \to 1$ reverses the observed trend $\Omega_\Lambda \to 1$ (r17)

These entries extend the register of §A.271.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.8.0-r17 ledger and guard update

$$R_{\cos}^{v480\text{-}r17} = (0, 0, 1)$$

for (present-epoch agreement recorded, time direction checked against the expansion history, choice among (a)–(c) open).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v480\text{-}r17,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v480\text{-}r17} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.8.0-r18 Source-Only: Three-Frame Typing of the Frame Offset and Correction of r17

Purpose. The single symbol β_{sym} is used in 261 lines of the main body in at least three roles: as a frame factor in the conversion of constants, as a branching probability of the cascade, and as a Hopf angle (`llm/k32_bsym_rollen.py`). This layer types the frame offset over the three frames of the triolic system, records three findings that the single symbol hides, narrows the register entries of r17, and records a reading of why the offsets coincide today. It contains arithmetic and typing, no new prediction. Earlier layers are not rewritten.
Source-only status. This v4.8.0-r18 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.8.0 Version DOI 10.5281/zenodo.23365244 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.8.0-r18 correction of r17

Proposition 42.5 does not mention the cosmological fractions. It states that β_{sym} drifts and that the H_0 discrepancy is a frame artefact; the drift acts through the frame factor. (The ratio $\gamma_L(z{=}0)/\gamma_L(z{=}1100) = 1.8642/1.7690 = 1.054$ raised to the power 3/2 gives 1.082 against the quoted 1.083; the exponent is a reconstruction, the source does not state it.) The value $\Omega_\Lambda(1100) = 0.722$ of §A.271 arises only when the drifting offset is inserted into Theorem 22.16. The one statement of the source that makes this insertion is item (2) of Proposition 42.8, $P(S^8 \to S^4) = \beta_{\text{sym}}^2 \to 1$.

Theoremlet r17.1 stands, because it is conditional on the assignment holding at all epochs. The register entries of §A.271 are narrowed as follows: Proposition 42.5 is not refuted by r17; the conflict with the expansion history sits in item (2) of Proposition 42.8 and in the tacit identification of two offsets that §A.271 separates. Item (1) of Proposition 42.8 uses the frame factor only and is not affected. The word "permanent" in Theorem 22.10 means non-vanishing ("we can never fully reach $z^- = 0$"); it does not conflict with a slow drift.

v4.8.0-r18 three frames and their cross terms

Write $\eta = \operatorname{artanh}\beta_{\text{sym}} = 0.6174$ for the rapidity of the offset. By Theorem 20.6 matter (+), photon (0) and tachyon (−) sit at rapidities $\eta_+ = \eta$, $\eta_0 = 0$, $\eta_- = -\eta$. Rapidities add, so the offset of frame a seen from frame b is $\eta_{ab} = \eta_a - \eta_b$, with velocity $\tanh\eta_{ab}$ and Lorentz factor $\cosh\eta_{ab}$:

from \ to	+	0	−
+	0	0.5493, $\gamma = 1.197$	0.8440, $\gamma = 1.864$
0	−0.5493, $\gamma = 1.197$	0	0.5493, $\gamma = 1.197$
−	−0.8440, $\gamma = 1.864$	−0.5493, $\gamma = 1.197$	0

Theoremlet v4.8.0-r18.1. Under the symmetry $\eta_- = -\eta_+$ of Theorem 20.6 the nine cross terms take exactly two non-trivial magnitudes: the neighbour term (+0, 0−) with $\cosh\eta = 1.197$ and the opposite term (+−) with $\cosh 2\eta = (1{+}\beta_{\text{sym}}^2)/(1{-}\beta_{\text{sym}}^2) = \gamma_L = 1.864$. The frame factor γ_L is the opposite term.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof sketch. $\cosh 2\eta = (1+\tanh^2\eta)/(1-\tanh^2\eta)$ with $\tanh\eta = \beta_{\text{sym}}$; this is the formula of Theorem 20.6, which names γ_L the Lorentz factor between the matter and tachyon frames. □

v4.8.0-r18 findings

(a) The conversion of constants uses the opposite term for a neighbour pair. The definition of the frame-insertion count in §25 converts matter to photon by $X^{(\text{photon})} = \gamma_L^{-k(X)} X^{(\text{matter})}$, and the proof of Theorem 20.5 reads γ_L as a Lorentz factor between the matter frame and the photon frame, with $\beta = 0.844$. By r18.1 the matter–photon pair is a neighbour pair, with factor 1.197; 1.864 belongs to matter–tachyon. Since $\cosh 2\eta \neq (\cosh\eta)^n$ for integer n ($(\cosh\eta)^2 = 1.432$), the difference cannot be absorbed into the counts k. Either the conversion is meant to pass through the tachyon frame, which would have to be stated, or every frame-corrected constant of §25 uses the wrong cross term. r18 does not choose.

(b) The observer position $z^- = \beta/2 = 0.422$. Theorem 20.5 places the observer at $z^- = \beta/2 \approx 0.422$, halving the opposite offset 0.844 in velocity. Halving in rapidity gives $\tanh\eta = 0.5493 = \beta_{\text{sym}}$, the position of Theorem 20.6. The value 0.422 is a halving in the wrong variable; the two theorems agree once the halving is done in rapidity.

(c) The cascade split is the neighbour term. Theorem 22.16 has $\Omega_M = \tanh^2\eta = 0.3018$ and $\Omega_\Lambda = \operatorname{sech}^2\eta = 0.6982$, which sum to one identically. The split is therefore internally consistent. Open: the cascade starts in the tachyon stratum S^8, yet uses the neighbour term $+0$ and no term of the $-$ row.

v4.8.0-r18 a reading of the coincidence, offered and not adopted

A single β_{sym} is the statement $\eta_- = -\eta_+$. If the matter frame drifts, as Proposition 42.5 has it, while the tachyon frame stays at a structural value $-\eta_s$, the symmetry is broken and the cross terms become three distinct numbers, $\eta_+(t)$, η_s and $\eta_+(t) + \eta_s$. A structural offset (cascade side) and an observed offset (frame side) then coincide exactly at the symmetric point $\eta_+ = \eta_s$.

The source supplies a reason for that point. In Theorem 22.11 the net force on the photon, $k(z_T^- - z_\gamma^-) + k(z_M^- - z_\gamma^-)$, vanishes for $z_\gamma^- = 0$ only if $z_T^- = -z_M^-$: the symmetric configuration is the mechanical equilibrium of the triple. Under this reading "the offsets agree today" means "the triple is near its equilibrium today", and the gap between 0.5493 (from γ_L) and 0.5578 (from Ω_M, Remark 22.17), 1.5%, would be a departure from symmetry rather than the lattice correction δ_{CL} to which the remark attributes it. The system of Theorem 22.11 is undamped: it supplies an equilibrium, not a relaxation towards it.

Which measured value belongs to which cross term under broken symmetry is not decided here. Any such assignment is a prediction and requires a pre-registration note before it is computed.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.8.0-r18 register entries

statement	status after r18
Prop. 42.5	not refuted by r17; uses the offset only through the frame factor (r18)
Prop. 42.8	item (2) carries the conflict of r17.1; item (1) unaffected (r18)
γ_L in §25, Thm. 20.5	opposite cross term $(+-)$ used for the neighbour pair matter–photon (r18 (a))
Thm. 20.5, $z^- = 0.422$	halving in velocity; in rapidity $z^- = \beta_{\mathrm{sym}}$ (r18 (b))
Thm. 22.16	split $\tanh^2\eta + \operatorname{sech}^2\eta = 1$ consistent; choice of the $+0$ term for a cascade from S^8 open (r18 (c))

v4.8.0-r18 ledger and guard update

$$R_{\mathrm{fr}}^{v480\text{-}r18} = (0, 0, 1)$$

for (three-frame cross terms typed, findings (a)–(c) recorded, assignment of the measured values to cross terms open).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v480\text{-}r18,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v480\text{-}r18} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.8.0-r19 Source-Only: Frame Cubes, Transition Group and Holonomy

Purpose. Layer r18 typed the frame offset by cross terms between the three frames. Two working results followed outside the source: the pre-registered test H12 (`llm/Ergebnis_H12_Drei_Frame_Zuordnung.md`) and a backward analysis with a readout index (`llm/k34_rueckwaerts_3x3x3.py`, map entry K39). This layer records both, asks what a family of readout cubes linked by transition maps adds, and compares the transition group of the earlier reading with that of Reading B (§A.271). It contains typing and arithmetic, no new prediction. Earlier layers are not rewritten.
Source-only status. This v4.8.0-r19 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.8.0 Version DOI 10.5281/zenodo.23365244 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.8.0-r19 records of H12 and of the backward analysis

H12. With $\cosh(\eta_+ + \eta_s) = \gamma_L$, $\Omega_M = \tanh^2 \eta_s$ (Planck 2018 with BAO) and $\sin^2\theta_{12} = \tanh^2\eta_+ / \cos^2\theta_{13}$, the assignment gives $\beta_+ = 0.5408$, $\beta_s = 0.5578$ and $\sin^2\theta_{12} = 0.2991 \pm 0.0057$, against 0.3086 for the symmetric relation H9. At the precision fixed for the next JUNO release the two are 1.36σ apart; by the fixed criterion they cannot be separated, and the result is recorded as no test. The limit is set by the uncertainty of Ω_M, not by JUNO.

Backward analysis (does not count as a test). Let T_{abc} be the offset of frame a against frame b, read from frame c, starting from the Hopf value $\beta_{\mathrm{sym}} = 1/\sqrt{3}$. Pure kinematics makes $T_{abc} = \eta_a - \eta_b$ independent of c, since a boost into c shifts all rapidities equally. Of the readout terms tried, those depending on the distance between reading and read frame fail, because they vanish for a self-reading ($\chi^2 \approx 34$). A single factor on everything read from the matter frame, $T_{abc} = (\eta_a - \eta_b) f(c)$, fits the two measurements with $f(+) = 0.944 \pm 0.008$ ($\chi^2 = 2.9$, one degree of freedom). In β this is 0.956 ± 0.007, numerically the known Hopf gap $\delta_{\mathrm{CL}} = 0.951$ of Remark 41.8. The backward analysis yields no new number; it offers a different reading of a known one. Only $f(+)$ is accessible, since every measurement is made from the matter frame.

v4.8.0-r19 cubes linked by transition maps

Take one readout cube $T^{(d)}_{abc}$ per frame $d \in \{+, 0, -\}$ and transition maps $\Lambda(d' \leftarrow d)$ with $T^{(d')} = \Lambda(d' \leftarrow d)\, T^{(d)}$. If the maps are fixed by the theory, the three cubes carry the information of one cube and the transition law; the family adds no parameter. What it adds is a consistency question: whether the maps compose to the identity around a closed path of frames. The residue, if any, is the holonomy of the family. It depends on the transition group and the positions of the frames only, and it is the part that no choice of the reading frame can remove. If the transition maps commute, the holonomy is trivial and the family adds nothing.

v4.8.0-r19 the transition group in the two readings

Earlier reading. Theorem 20.6 places matter and tachyon at $\pm\beta_{\mathrm{sym}}$ on one axis around the photon and calls γ_L the Lorentz factor between them. The transitions are collinear boosts; they commute, and the holonomy is trivial.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Reading B. By §A.271 the kernel is the Riemann–Silberstein vector $z = E + iB$ under $SO(3,\mathbb{C}) \cong SO^+(1,3)$, with $s = E^2 - B^2 + 2i\,E \cdot B$. The working assignment of r07 reads matter as $s > 0$ (electric type), tachyon as $s < 0$ (magnetic type), photon as $s = 0$ (null field).

Theoremlet v4.8.0-r19.1. In Reading B no Lorentz transformation relates matter to tachyon, and none relates either of them to the photon. Matter and tachyon with equal $|s|$ are related by the duality rotation $z \mapsto e^{i\pi/2}z$, that is $E \mapsto -B$, $B \mapsto E$. The photon is reached by neither group action.

Proof sketch. $SO(3,\mathbb{C})$ preserves s (Theoremlet r06.3), so it preserves its sign and the null cone. The duality rotation multiplies s by $e^{2i\alpha}$ and maps $s > 0$ to $s < 0$ at $\alpha = \pi/2$ (Theoremlet r07.2); it preserves $s = 0$. This is the familiar statement that a radiation field is null in every frame and under every duality rotation. □

Consequence. The Lorentz factor γ_L between matter and tachyon in Theorem 20.6 has no counterpart in Reading B, where the two are related by a duality rotation and not by a boost. On this point the two readings disagree about the transition group itself.

Theoremlet v4.8.0-r19.2. The duality circle carries a $\mathbb{Z}_2$ holonomy around the photon. Under $z \mapsto e^{i\alpha}z$ one has $s \mapsto e^{2i\alpha}s$; as α runs from 0 to π, s encircles $s = 0$ once and z returns as $-z$. Paths of s that do not encircle $s = 0$ have trivial holonomy. The photon is the branch point of z over s.

Proof sketch. s is quadratic in z, so the map from the duality circle of z to the circle of s is two to one. Theoremlet r07.1 states the sign ambiguity of z_m under $\alpha \mapsto \alpha + \pi$. □

v4.8.0-r19 the earlier reading as a photon-centred view

Theoremlet v4.8.0-r19.3. The real axis of the s-plane reproduces the sector structure of Theorem 20.6: matter and tachyon at $\pm|s|$, mirror-symmetric about the photon at $s = 0$, the mirror $s \mapsto -s$ being the half-turn of the duality circle.

Proof sketch. Theoremlet r07.1 with $\alpha = 0$ and $\alpha = \pi/2$. □

Reading, offered and not adopted. The photon is fixed by the whole duality circle, so it picks out no reference direction in the s-plane. The earlier reading can then be taken as the view centred on the photon with the real axis chosen as reference, that is, with the duality angle fixed as a gauge. On this axis the straight path from matter to tachyon passes through the photon, the branch point, where the phase is undefined; in Reading B the path runs around the photon on $|s| = \text{const}$. The photon-centred view therefore cannot see the $\mathbb{Z}_2$ holonomy. It also supplies something Reading B lacks: a metric along the axis, in which the mirror distance is read as a boost with factor γ_L. No correspondence between $|s|$ and β_{sym} is given in the source, and by r19.1 the boost reading has no counterpart in Reading B. Whether the duality angle is a gauge, which would make the earlier reading a gauge-fixed projection, or physical, which would make it a projection that loses information, is the question left open in r07.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.8.0-r19 consequence for the readout factor

The $\mathbb{Z}_2$ holonomy can only contribute ± 1; it cannot produce a factor such as $f(+) = 0.944$. A continuous holonomy would need closed paths in an $SO(3,\mathbb{C})$-orbit together with a rule for transporting frames along them, i.e. a connection. The source supplies neither. Composing boosts in non-collinear directions yields a Wigner rotation, and the triole directions are a candidate for such directions, but choosing them is a model decision. Any comparison of such a holonomy with $f(+)$ is a prediction and requires a pre-registration note.

v4.8.0-r19 register entries

statement	status after r19
Thm. 20.6	sector structure reproduced by the real s-axis of Reading B (r19.3); γ_L as a Lorentz factor between matter and tachyon has no counterpart in Reading B (r19.1)
duality circle of r07	$\mathbb{Z}_2$ holonomy around the photon, photon as branch point (r19.2)
Rem. 41.8, δ_{CL}	numerically equal to the readout factor $f(+)$ of the backward analysis; reinterpretation only, not a test (r19)
H12	not separable from H9 by the fixed criterion; no test (r19)

v4.8.0-r19 ledger and guard update

$$R_{\mathrm{hol}}^{v480\text{-}r19} = (0,0,1)$$

for (transition groups of both readings compared, $\mathbb{Z}_2$ holonomy of the duality circle recorded, continuous holonomy open for lack of a connection).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v480\text{-}r19,+)} = (0,1,1,1,1), \qquad \Gamma_{\mathrm{guard}}^{v480\text{-}r19} = (1,1,1,1,1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.8.0-r20 Source-Only: The Tachyon Frame, Complex Rapidity and the Holonomy Test H13

Purpose. Layer r19 found no Lorentz transformation between the sectors of Reading B and read the earlier reading as a photon-centred view. This layer records the pre-registered holonomy test H13 and a post-hoc finding attached to it, and examines the tachyon frame under the holonomy and spinor structure. The computations are `llm/k35_test_h13.py` and `llm/k36_tachyon_frame.py`; only H13 is compared with data. Earlier layers are not rewritten.
Source-only status. This v4.8.0-r20 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.8.0 Version DOI 10.5281/zenodo.23365244 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.8.0-r20 the holonomy test H13

The note `llm/Vorab_Notiz_H13_Wigner_Holonomie.md` fixed, before any computation, one construction: three matter frames moving with rapidity $\eta_H = \operatorname{artanh}(1/\sqrt{3})$ in the three triole directions, linked by pure boosts into a closed loop, with $\cos\delta$ of the resulting Wigner rotation to be compared with the readout factor $f(+) = 0.944 \pm 0.008$ of the backward analysis recorded in §A.271.

The holonomy is a pure rotation (boost part 10^{-16}) by $\delta = 0.4985$ rad $= 28.56°$, equal to the angular defect of the equilateral hyperbolic triangle with vertices at the three velocities. Then $\cos\delta = 0.878$, at $z = -7.9$ from $f(+)$. By the fixed criterion H13 is refuted for this construction, and the holonomy route with triole boosts is closed for $f(+)$. Since $f(+)$ is itself a fit, this removes an explanation of the factor, not the factor.

Post-hoc finding, not registered and not counting. Asked after the result: $\cos^2(\delta/2) = (1+\cos\delta)/2 = 0.939$ meets $f(+)$ at $z = -0.5$. It is one of four obvious functions of δ ($\cos\delta$: -7.9; $\cos(\delta/2)$: $+3.1$; $1-\delta/2\pi$: -2.8), roughly the rate expected by chance. Its parts have different standing. The half angle is forced once the spinor is read out: by Theoremlet r06.4 (§A.271) z is quadratic in $\psi \in \mathbb{C}^2$, so a rotation of the frames by δ turns ψ by $\delta/2$. The square is a Born-type reading of $|\langle\psi|U\psi\rangle|^2$. The orientation is a choice: $|\langle\psi|U\psi\rangle|^2 = \cos^2(\delta/2) + \sin^2(\delta/2)\cos^2\theta$, with θ the angle to the rotation axis, which is normal to the triole plane. The triole photon $c(1,\omega,\omega^2)$ is a null field whose direction $E \times B$ is $(1,1,1)$, the normal of the plane $z_1+z_2+z_3 = 0$; its spinor lies on the axis and acquires only the phase $\delta/2 = 14.28°$. The factor $\cos^2(\delta/2)$ belongs to a spinor in the plane, such as the helicity of a matter frame moving in a triole direction.

v4.8.0-r20 the tachyon frame

Theoremlet v4.8.0-r20.1. The duality rotation $z \mapsto iz$ commutes with every element of $SO(3,\mathbb{C})$. Hence every closed path in the tachyon sector has the same holonomy as its matter image (for the triole loop $\delta = 28.56°$), and z and iz have the same two principal null directions. Matter and tachyon with $z_T = iz_M$ differ only in the overall phase of the spinor product.

Proof sketch. iz is a scalar multiple of z, and $SO(3,\mathbb{C})$ acts linearly. Writing z as a symmetric 2-spinor F_{AB} with $F_{11} = \frac{1}{2}(z_1 - iz_2)$, $F_{22} = \frac{1}{2}(-z_1 - iz_2)$, $F_{12} = -\frac{1}{2}z_3$ (the

© 2026 Marc Brendecke. Preprint — not peer reviewed.

polarisation of the map of r06.4), the principal spinors are the roots of $F_{11}\lambda^2 + 2F_{12}\lambda + F_{22}$, which do not change when F is multiplied by i. Checked numerically. □

Theoremlet v4.8.0-r20.2. For a speed $v > 1$,

$$\operatorname{artanh} v = \operatorname{artanh} \frac{1}{v} + \frac{i\pi}{2}.$$

The four-velocity at rapidity $\eta + i\pi/2$ is $i\,w$ with $w = (\sinh\eta, \cosh\eta)$ spacelike: the real four-velocity turns from timelike to spacelike and the factor i carries the phase. For $v = 1/\beta_{\text{sym}} = 1.8203$ the real part is $\eta_{\text{sym}} = 0.6174$ and the imaginary part is $\pi/2$, the duality angle that relates matter and tachyon in Theoremlet r07.2 (§A.271).

Proof sketch. $\tanh(x + i\pi/2) = \coth x$; $\cosh(\eta + i\pi/2) = i\sinh\eta$, $\sinh(\eta + i\pi/2) = i\cosh\eta$. □

The same structure in three places. A phase $\pi/2$ on the representative and a change of sign of the real invariant occur in the duality rotation $z_T = iz_M$, $s \mapsto -s$ (r07.2), in the superluminal rapidity of r20.2, and in the signed tachyon scale $\ell_T^2 = -\lambda_T^2$ of §A.269, where $\ell_T = i\lambda_T$. The last is an allowed effective statement of the release scope; r20 records the parallel and adds nothing to it.

Reading, offered and not adopted. The earlier reading places the tachyon at $-\beta_{\text{sym}}$, the real part of its rapidity; Reading B keeps the imaginary part $\pi/2$ as the duality angle. The two readings are then related by keeping or dropping one imaginary part, which is the photon-centred view of §A.271 expressed in rapidities.

v4.8.0-r20 cross terms with complex tachyon rapidity

With the tachyon at $-\eta + i\pi/2$ the cross terms of §A.271 become

pair	real (r18)	with $i\pi/2$
matter–photon	$\cosh\eta = 1.197$	1.197
photon–tachyon	$\cosh\eta = 1.197$	$-i\sinh\eta = -0.657\,i$
matter–tachyon	$\cosh 2\eta = 1.864 = \gamma_L$	$-i\sinh 2\eta = -1.574\,i$

The matter–tachyon term becomes purely imaginary with modulus $\sinh 2\eta = \sqrt{\gamma_L^2 - 1}$, the proper velocity of the pair instead of its Lorentz factor; for tachyons cosh and sinh exchange their roles. The real factor γ_L arises only when the imaginary part is dropped. This is Theoremlet r19.1 in numbers: Reading B has no real Lorentz factor between matter and tachyon.

The tribonacci condition. The fixed-point condition $\mu = 1/\beta_{\text{sym}}(\mu)$ of Theorem 41.18 reads, in rapidities, $\cosh 2\eta = \coth\eta = \tanh(\eta + i\pi/2)$: the Lorentz factor of the opposite term equals the speed of a tachyon with real rapidity η. This is a reformulation, not a new value.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.8.0-r20 register entries

statement	status after r20
H13, readout factor as triole holonomy	refuted for the fixed construction, $z = -7.9$ (r20)
$\cos^2(\delta/2)$ for $f(+)$	post-hoc, not registered, does not count; half angle from r06.4, square and orientation are readings (r20)
tachyon sector of Reading B	same holonomy and same principal spinors as matter (r20.1)
Thm. 20.6, tachyon at $-\beta_{\mathrm{sym}}$	real part of the superluminal rapidity $\eta_{\mathrm{sym}} + i\pi/2$; imaginary part equals the duality angle (r20.2)
γ_L as matter–tachyon factor	real only with the imaginary part dropped; with it $-i \sinh 2\eta$ (r20)
Thm. 41.18	equivalent to $\cosh 2\eta = \coth \eta$ (r20)

v4.8.0-r20 ledger and guard update

$$R_{\mathrm{tach}}^{v480\text{-}r20} = (0, 0, 1)$$

for (H13 decided, tachyon frame related to both readings through the complex rapidity, physical status of the imaginary part open).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v480\text{-}r20,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v480\text{-}r20} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

A.272 v4.9.0 Post-Release Layers

The source-only layers v4.9.0-r01 to r31 follow, recorded after the v4.9.0 release (Version DOI 10.5281/zenodo.23534954) on the full-monolith working line. They treat the conversion of constants, the round trip and the compensator in the complex-vector reading, two pre-registered tests (H14 refuted, H15 pending), the frame drift against atomic clocks, and the mathematical core (a reading of the C5 factor, a reconstruction of $\Delta_\gamma < 1/5$, a time scale), the state disk and the class split of the conversions, the clock bound per frame, Convention F for the three frames, the revision of the main body that opens the 4.10 base (r25), a further pre-registered test (H16, refuted), and a dynamics in which a duality rotation is followed by a compensator measurement, with the tachyon as measuring device, vacuum pairs as replenishing reservoir and the Markov cascade as a test of self-stabilisation (r26–r31). Layer v4.9.0-r15 is the consolidated register of r01–r14. Earlier layers are not rewritten.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r01 Source-Only: The Photon Frame as Neutral Axis and the Conversion of Constants

Purpose. Finding (a) of §A.271 records that the conversion of constants between the matter and the photon frame uses $\gamma_L = \cosh 2\eta$, the matter–tachyon cross term, where the matter–photon pair is a neighbour pair with factor $\cosh \eta$. This layer records where the factor came from, the working assumption under which it was chosen, and what the assumption implies for a conversion law over three frames. The computations are `llm/k37_gl_umrechnungen.py`. No value is compared with data. Earlier layers are not rewritten.
Source-only status. This v4.9.0-r01 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r01 provenance of the factor

In the archived versions v2.28.0 to v2.29.0 the splitting factor reads "$\gamma_L = \gamma_{\mathrm{BI}}/\gamma_{\mathrm{string}} \approx 1.197$", while the same versions already carry $1/\beta_{\mathrm{sym}} \approx 1.820$, that is $\beta_{\mathrm{sym}} \approx 0.5495$. The value 1.197 is $(1-\beta_{\mathrm{sym}}^2)^{-1/2}$, the Lorentz factor of the matter-frame offset; the ratio $\gamma_{\mathrm{BI}}/\gamma_{\mathrm{string}}$ is 1.864. In v2.29.1 the value was changed to 1.864 and the change recorded as an arithmetic correction ("previous version used incorrect $\gamma_L \approx 1.197$"); the note carried in Corollary 22.21 calls the earlier value γ_{BI}, which it is not ($\gamma_{\mathrm{BI}} \approx 0.2375$). Since then every conversion uses 1.864.

Author's statement (2026-10-21). The photon frame was assumed to be neutral, and the work was meant to proceed on that neutral axis first. The archive records the change of value only as an arithmetic correction and does not mention a neutral axis.

Theoremlet v4.9.0-r01.1. The two values the factor has carried are the two cross terms of Theoremlet r18.1:

$$1.197 = \cosh\eta = \sqrt{\tfrac{1+\gamma_L}{2}} \quad \text{(matter–photon)}, \qquad 1.864 = \cosh 2\eta = \gamma_L \quad \text{(matter–tachyon)},$$

with $\eta = \operatorname{artanh} \beta_{\mathrm{sym}}$. Both follow from the single input $\gamma_{\mathrm{BI}}/\gamma_{\mathrm{string}}$.

Proof sketch. $\cosh 2\eta = 2\cosh^2\eta - 1$ and $\cosh\eta = (1-\tanh^2\eta)^{-1/2}$. □

v4.9.0-r01 the neutral assumption in Reading B

The assumption has structural support in the complex-vector reading. The photon is the sector fixed by the whole duality circle (§A.271), the branch point of the duality holonomy (Theoremlet r19.2), and the centre of the real s-axis on which matter and tachyon are mirror images (§A.271). On the rapidity axis of Theorem 20.6 it sits at rapidity 0, between matter at $+\eta$ and tachyon at $-\eta$. If the photon frame is neutral, a value on the neutral axis is reached from the matter frame by the neighbour step, and the factor carried before v2.29.1 is the factor of that step.

Theoremlet v4.9.0-r01.2. A conversion law $X^{(b)} = \gamma_{ab}^{-k} X^{(a)}$ with γ_{ab} the Lorentz factor of the pair is not consistent over the three frames: from matter to tachyon through the photon it gives $(\cosh\eta)^{2k} = \left(\frac{1+\gamma_L}{2}\right)^k$, directly it gives γ_L^k (for $k=1$: 1.432 against 1.864; for $k=6$: 8.63 against 42.0). A law with the Doppler factor, $X^{(b)} = e^{-k\eta_{ab}} X^{(a)}$, composes exactly.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof sketch. $\cosh^2\eta \neq \cosh 2\eta$ for $\eta \neq 0$, while $e^{\eta}e^{\eta} = e^{2\eta}$. □

Consequence. With the Lorentz-factor law one pair at a time is consistent. Working on one axis first, as the author describes, is the regime in which the law of §25 is free of contradiction; the obstruction of r01.2 appears only once both pairs are used. Under the neutral assumption the primary pair is matter–photon, with factor $\cosh\eta$; γ_L is the factor across the neutral axis, from matter to tachyon.

v4.9.0-r01 a frame-order finding in the Hawking proposition

Items (i) and (ii) of Proposition 21.4 give $T_H^{(\text{photon})} = \gamma_L\, T_H^{(\text{matter})}$ and $T_H^{(\text{tachyon})} = \gamma_L^{-1} T_H^{(\text{matter})}$. Reciprocal factors place photon and tachyon on opposite sides of matter. On the axis of Theorem 20.6 the photon lies between matter and tachyon. A law along the axis moves the tachyon value beyond the photon value in the same direction; a mirror law about a neutral photon gives the tachyon the matter value. Neither gives reciprocal factors, so item (ii) carries a frame order that does not match the axis. This is also the one statement of the main body in which the same factor serves both pairs.

v4.9.0-r01 ways forward, recorded and not chosen

(A) Keep γ_L and the Lorentz-factor law, and read each conversion of §25 as one across the neutral axis (matter–tachyon). Statements whose endpoint is the photon frame then need the neighbour factor.

(B) Use the neighbour factor $\cosh\eta = 1.197$ for matter–photon, the value carried before v2.29.1, in line with the neutral assumption. This changes the photon-frame numbers of the main body (table below).

(C) Use a composable law with the Doppler factor, $e^{\eta} = \sqrt{(1+\beta_{\text{sym}})/(1-\beta_{\text{sym}})} = 1.854$ per neighbour step. It is consistent over three frames. At β_{sym} its neighbour factor lies 0.55% below γ_L, so the numbers of the main body change little while their meaning changes. By Theoremlet r01.3 below the closeness is structural: in the continuum limit of Theorem 41.18 the two factors are equal.

Theoremlet v4.9.0-r01.3. The full-span Lorentz factor equals the neighbour-step Doppler factor, $\cosh 2\eta = e^{\eta}$, exactly when $e^{\eta} = 1$ or $e^{\eta} = \mathbb{T}$, the tribonacci constant of Theorem 41.18. At that point $\cosh 2\eta = e^{\eta} = \coth\eta = \mathbb{T} = 1.83929$, $\beta = 0.54369$: the self-referential condition $\gamma_L = 1/\beta_{\text{sym}}(\gamma_L)$ of Theorem 41.18 is the condition that the matter–tachyon Lorentz factor and the matter–photon Doppler factor coincide.

Proof sketch. With $x = e^{\eta}$, $\cosh 2\eta = (x^2 + x^{-2})/2$, so $\cosh 2\eta = x$ is $x^4 - 2x^3 + 1 = 0$, which factors as $(x-1)(x^3 - x^2 - x - 1) = 0$. Further $\coth\eta = (x^2+1)/(x^2-1)$, and $\coth\eta = x$ is again $x^3 - x^2 - x - 1 = 0$. Checked numerically to 10^{-16}. □

Consequence for the neutral axis. In the continuum limit the author's choice of γ_L for the step from matter to the neutral photon frame is exactly the factor a composable law assigns to that step: reading (A) and reading (C) agree there. With the measured $\gamma_L = \gamma_{\text{BI}}/\gamma_{\text{string}}$, which lies 1.37% above $\mathbb{T}$ (the lattice correction of Remark 41.20), they differ by 0.55%. The Lorentz-factor law of reading (B) does not share this property.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

quantity in the main body	γ_L	neighbour, Lorentz	neighbour, Doppler
$\ell_P^{(\mathrm{matter})}/\ell_P^{(\mathrm{photon})}$ (Theorem 25.5), power 3/2	2.546	1.309	2.525
vacuum-energy ratio, power 6	42.0	2.94	40.6
$m_\nu^{(\mathrm{matter})}/m_\nu^{(\mathrm{photon})}$ (Proposition 42.4), power 4	12.08	2.05	11.82
matter error (Corollary 42.17), power 5	22.53	2.46	21.92

The table is arithmetic on the formulas of the main body; photon-frame values are not measured. Statements that are compared with data, among them Corollary 42.24 and Proposition 42.7, are not re-evaluated here; a re-evaluation under (A), (B) or (C) is a prediction and requires a pre-registration note.

v4.9.0-r01 register entries

statement	status after v4.9.0-r01
conversion law of §25	factor γ_L is the matter–tachyon term; consistent for one pair at a time (r01.2); under the neutral photon assumption the matter–photon factor is $\cosh\eta$; choice among (A)–(C) open
Cor. 22.21, v2.29.1 note	the earlier value 1.197 was $\cosh\eta$, the matter–photon Lorentz factor, not γ_{BI} (r01.1)
Prop. 21.4 (ii)	reciprocal factors put the tachyon on the far side of matter; does not match the axis order (r01)
Thm. 41.18	equivalent to $\cosh 2\eta = e^{\eta}$: matter–tachyon Lorentz factor equals matter–photon Doppler factor (r01.3)

v4.9.0-r01 ledger and guard update

$$R_{\mathrm{conv}}^{v490\text{-}r01} = (0,0,1)$$

for (provenance of the factor recorded, consistency of the conversion law over three frames decided and its relation to the tribonacci fixed point established, choice among (A)–(C) open).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r01,+)} = (0,1,1,1,1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r01} = (1,1,1,1,1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r02 Source-Only: The Round Trip about the Photon and the Umkehrwalze

Purpose. Theoremlet r19.2 records a $\mathbb{Z}_2$ holonomy of the duality circle around the photon. This layer follows a round trip matter → photon → tachyon → photon → matter in Reading B, asks on which side of the photon each passage runs, and asks what the Umkehrwalze of the main body is in physical terms. It uses the reflector and leakage language of the OP5 layer (Definition H3.3). The computations are `llm/k38_umkehrwalze.py`. No value is compared with data. Earlier layers are not rewritten.
Source-only status. This v4.9.0-r02 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

Author's motivation (2026-10-21). The Umkehrwalze was introduced for a path that does not end at the tachyon but returns, matter → photon → tachyon → photon → matter, read as a parallel transport over the sphere.

v4.9.0-r02 the two sides of the photon

Throughout, $z = E + iB$, $s = z \cdot z = E^2 - B^2 + 2i\,E \cdot B$, matter $s > 0$, tachyon $s < 0$, photon $s = 0$ (§A.271).

Theoremlet v4.9.0-r02.1. A passage from matter to tachyon at fixed $|s|$ runs through the upper or the lower half of the s-plane and carries $z \mapsto iz$ or $z \mapsto -iz$. The side is the sign of $\operatorname{Im} s = 2\,E \cdot B$; it is Lorentz-invariant, odd under P and T, and even under C. For the round trip:

(a) out and back on the same side (the return retraces the outward path): $z \mapsto z$;

(b) out on one side and back on the other (the duality angle keeps its sense, $\pi/2 + \pi/2$): $z \mapsto -z$.

On the spinor of Theoremlet r06.4 the round trip (b) is $\psi \mapsto \pm i\psi$; two round trips give $\psi \mapsto -\psi$ with z restored.

Proof sketch. $s(e^{i\alpha}z) = e^{2i\alpha}s$, so $\alpha \in [0, \pi/2]$ moves s through the upper half-plane and $\alpha \in [-\pi/2, 0]$ through the lower one. $SO(3,\mathbb{C})$ preserves s and hence $\operatorname{Im} s$. By the parities of E and B (P: E odd, B even; T: E even, B odd; C: both odd), $E \cdot B$ is P- and T-odd and C-even. Case (b) is a loop around $s = 0$, case (a) is not (Theoremlet r19.2). The spinor statement is $z(e^{i\varphi}\psi) = e^{2i\varphi}z(\psi)$. Checked numerically. □

Reading. The "double tunnel point" of the round trip is one point with two sides: the photon, passed on the side $E \cdot B > 0$ or $E \cdot B < 0$. The side is the sign of the density of the electromagnetic θ-term $F\tilde{F}$. Only a round trip that changes side encloses the photon.

v4.9.0-r02 the round-trip Umkehrwalze is charge conjugation

Theoremlet v4.9.0-r02.2. On the field strength the holonomy $z \mapsto -z$ of the round trip (b) is charge conjugation C, equals PT (with $x \mapsto -x$), and equals the duality rotation by π. It commutes with $SO(3,\mathbb{C})$ and with the duality circle, and fixes no non-zero field.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof sketch. C: $E \mapsto -E$, $B \mapsto -B$. T: $z \mapsto \bar{z}$; P: $z \mapsto -\bar{z}$; composed, $z \mapsto -z$. $e^{i\pi} = -1$ is central. □

Consequence. The Umkehrwalze of the round trip is not an added element. It is forced by the two-to-one relation between z and s, whose branch point is the photon, and in physical terms it is C. Being central, it cannot serve as the J of the readout projector $C_{\mathrm{HC}} = \frac{1}{2}(\mathrm{id} + J)$ of Definition H3.3: for $J = -1$ that projector is zero.

v4.9.0-r02 the involutive Umkehrwalze must contain time reversal

The main body uses J as an involution that exchanges matter and tachyon (Theorem 21.7, where the inside view runs with time reversed). In Reading B the quarter turn $z \mapsto iz$ does this exchange, but its square is -1, so it is not an involution.

Theoremlet v4.9.0-r02.3.

(a) No $\mathbb{C}$-linear involution L of $\mathbb{C}^3$ satisfies $s(Lz) = -s(z)$.

(b) The $\mathbb{C}$-antilinear involutions with $s(Lz) = -\overline{s(z)}$ are exactly $Lz = iR\bar{z}$ with $R \in O(3, \mathbb{C})$ and $R\bar{R} = 1$. Instances are $R = 1$, real orthogonal involutions, and squares of boosts.

(c) The instance $R = 1$,

$$J_B z := i\bar{z}, \qquad \text{that is} \quad E \mapsto B,\ B \mapsto E,$$

is the duality rotation by $\pi/2$ after time reversal. It exchanges matter and tachyon, maps the null cone to itself, and preserves the side $\operatorname{sign} \operatorname{Im} s$.

Proof sketch. (a) $M := -iL$ preserves s, so $M \in O(3, \mathbb{C})$; $L^2 = 1$ gives $M^2 = -1$ and $\det(M)^2 = -1$, impossible. (b) $Lz = A\bar{z}$ with $A^{\mathsf{T}}A = -1$ gives $A = iR$, $R \in O(3, \mathbb{C})$; $L^2 = A\bar{A} = R\bar{R}$. (c) $i(E - iB) = B + iE$, and $s \mapsto -\bar{s}$. Checked numerically. □

Consequence. In three complex dimensions an involutive exchange of matter and tachyon cannot be linear. It has to be antilinear, and antilinearity is time reversal. This matches the inside view of Theorem 21.7, which runs with time reversed. The physical reading of the involutive Umkehrwalze is therefore electric–magnetic exchange, a duality quarter turn composed with T. It is a symmetry of the source-free Maxwell equations. Unlike C it is not Lorentz-invariant: $J_B\Lambda = \bar{\Lambda}J_B$, so it commutes with real rotations only and singles out a frame.

Theoremlet v4.9.0-r02.4. The eigen-channels of J_B lie one on each side of the photon:

$$J_B z = z \iff E = B,\ s = 2i|E|^2; \qquad J_B z = -z \iff E = -B,\ s = -2i|E|^2.$$

The projector $C_{\mathrm{HC}} = \frac{1}{2}(\mathrm{id} + J_B)$ maps every field to $\operatorname{Re} s = 0$, $\operatorname{Im} s \geq 0$. Its image meets the null cone only in $z = 0$.

Proof sketch. Direct substitution; checked on 1000 random fields ($\max|\operatorname{Re} s| = 0$, $\min \operatorname{Im} s > 0$). □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Consequence for Theorem 21.7. The compensator $\frac{1}{2}(f + Jf)$ "produces the Hawking photon at $\beta = 0$". With $J = J_B$ it produces the value 0 of the matter–tachyon coordinate $\operatorname{Re} s = E^2 - B^2$, not a null field. The output keeps $E \cdot B \neq 0$ and lies on one side of the photon, which is the endpoint of its image. The statement holds for the real part of s only.

Theoremlet v4.9.0-r02.5. J_B reverses the sense of the duality circle, $J_B R_\alpha = R_{-\alpha} J_B$ with $R_\alpha z = e^{i\alpha} z$. Hence

$$R_\alpha^{-1} J_B R_\alpha = R_{-2\alpha} J_B, \qquad R_{\pi/2}^{-1} J_B R_{\pi/2} = -J_B, \qquad R_{\pi/4}^{-1} J_B R_{\pi/4} = T.$$

Passing from matter to the tachyon, reflecting there, and passing back gives the direct reflection followed by the round-trip Umkehrwalze C.

Proof sketch. $e^{-i\alpha}\, i\, \overline{e^{i\alpha} z} = i e^{-2i\alpha} \bar{z}$. Checked numerically. □

Reading. This is the Enigma pattern of the computational interpretation surface: a reflector between a forward and a backward passage. The two Umkehrwalzen of this layer are related by it. The central one, C, is the holonomy of a passage pair. The involutive one, J_B, is the reflector, and conjugating it by the matter–tachyon passage multiplies it by C.

v4.9.0-r02 what the OP5 layer contributes

Inserting $J = J_B$ into the leakage diagnostics $E_{\text{comm}} = [\mathcal{M}, J]$ and $E_{\text{leak}} = (\mathrm{id} - C_{\text{HC}})\mathcal{M} C_{\text{HC}}$ of Definition H3.3 gives the following, with the transports of Reading B in place of $\mathcal{M}$ (relative norms on a random field):

transport $\mathcal{M}$	$\lvert E_{\text{comm}} z\rvert / \lvert z\rvert$	$\lvert E_{\text{leak}} z\rvert / \lvert C_{\text{HC}} z\rvert$
real rotation	0	0
boost, $\eta = 0.6174$	0.985	0.526
duality rotation, $\alpha = 0.4$	0.779	0.389

By Proposition H3.4 the admissible channel is stable under the compact part of the transport only. Boosts leak because J_B singles out a frame. Duality rotations leak because J_B reverses their sense (Theoremlet r02.5). Boosts preserve the side of the field as a whole, so this leakage is into the J_B-odd component and does not change side.

Correspondence, offered and not adopted. The local OP5 closure theorem works under the identification of the OP5 near-kernel branch with the *photon-frame HC-cone/reflector route*. Its halo invariance off tunnel candidates asks for a turning-point readout before a through-collapse exit. In the language of this layer a turning point is case (a) of Theoremlet r02.1, a return on the same side with trivial holonomy, and a through-exit is case (b), a change of side with holonomy C. The reflector of that route would be antilinear by Theoremlet r02.3. The radial and HC-residual channels ρ_r, ρ_h, the kernel K and the tunnel selector have no counterpart in Reading B yet. The correspondence is therefore recorded and not used. It does not touch the guarded scope of the OP5 closure.

v4.9.0-r02 physical summary in plain terms

The following restates Theoremlets r02.1–r02.5 without notation. It relies on the working assignment of r07 (matter electric-dominated, tachyon magnetic-dominated), which is a model decision, and on Reading B, which is kinematics without dynamics (r10).

© 2026 Marc Brendecke. Preprint — not peer reviewed.

1. *Two ways past the photon.* A field changes from matter type to tachyon type only by passing the photon. It can pass on one of two sides, according to whether its electric and magnetic parts lean towards each other or away from each other (the sign of $E \cdot B$). Every observer agrees on the side.

2. *The round trip.* Going out and coming back on the same side changes nothing. Going out on one side and coming back on the other takes the field once around the photon, and it returns with both of its parts reversed. That is the field of opposite charges: the round trip acts as charge conjugation. Only a second round trip restores the underlying spinor, as with a Möbius strip.

3. *The mirror between matter and tachyon.* A map that exchanges matter and tachyon directly and gives the identity when applied twice exists only if it also reverses time. The simplest such map exchanges the electric and the magnetic field. For light in vacuum this is a known symmetry, electric–magnetic duality combined with time reversal. The time-reversed inside view of Theorem 21.7 has this structure.

4. *What the compensator produces.* Averaging a field with its mirror image makes the electric and the magnetic part equally strong, which is the line between matter and tachyon. The two parts then point the same way. For light they are perpendicular. The average is therefore balanced, but it is not a light field.

5. *The two mirror channels.* The fields that the mirror leaves unchanged lie on one side of the photon, and the fields it reverses lie on the other. The two sides of item 1 are the two channels of the mirror.

No statement in this list is compared with data.

v4.9.0-r02 register entries

statement	status after v4.9.0-r02
Umkehrwalze of the round trip	holonomy of a side-changing passage pair, equal to $C = PT$ on field strengths (r02.1, r02.2)
involutive J exchanging matter and tachyon	must be antilinear (contains T); representative J_B: $E \leftrightarrow B$ (r02.3); frame-dependent
Thm. 21.7, photon from $\frac{1}{2}(f + Jf)$	with J_B: $\operatorname{Re} s = 0$, not a null field; holds for the real part of s only (r02.4)
Def. H3.3 with J_B	$E_{\mathrm{comm}} = E_{\mathrm{leak}} = 0$ for real rotations only (r02)
OP5 turning point / through-exit	correspondence to cases (a)/(b) of r02.1 recorded, not adopted

v4.9.0-r02 ledger and guard update

$$R_{\mathrm{uw}}^{v490\text{-}r02} = (0, 0, 1)$$

for (round-trip Umkehrwalze identified with C, involutive Umkehrwalze characterised as antilinear, identification with the OP5 halo/tunnel structure open).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r02,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r02} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r03 Source-Only: The Compensator as a Two-Arm Interferometer

Purpose. Theoremlet r02.4 found that the compensator $C(f) = \frac{1}{2}(f + Jf)$ with the involution $J_B z = i\bar{z}$ of r02.3 does not produce a null field. This layer looks at the compensator itself in Reading B: as the output of an interferometer, on a static charge, on a light wave, and in its product form. The computations are `llm/k39_kompensator.py`. No value is compared with data. Earlier layers are not rewritten.
Source-only status. This v4.9.0-r03 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

Author's motivation (2026-10-21). The picture behind the compensator was the Michelson–Morley interferometer with its half-silvered mirror; a mirror for the tachyon would act as a periscope that lays both images over each other.

v4.9.0-r03 the two output ports

Theoremlet v4.9.0-r03.1. Split a field f at a beam splitter. Send one half through the identity and one through J, then recombine. The two output ports carry $C(f) = \frac{1}{2}(f + Jf)$ and $A(f) = \frac{1}{2}(f - Jf)$, the compensator and its antisymmetric partner of Definition r04.1. The energy splits without loss, $|Cf|^2 + |Af|^2 = |f|^2$. For antilinear J (r02.3) both maps are real-linear only, and

$$A(f) = -i\, C(if), \qquad \operatorname{ran} A = i \operatorname{ran} C.$$

The two ports differ by a duality quarter turn.

Proof sketch. $C + A = 1$, $C^2 = C$, $CA = 0$ (Theoremlet r04.2), and $\mathrm{Re}\langle Cf, Af\rangle = 0$ for anti-unitary J. $C(if) = \frac{1}{2}(if - iJf) = iA(f)$. Checked numerically. □

Consequence. The bosonic/fermionic split $C(f) \in H_+^+$, $A(f) \in H_-^+$ of the Pauli item in §12 becomes, with an antilinear J, a split into two real subspaces that are images of each other under multiplication by i. That is a split into real and imaginary parts, not two independent sectors. Remark A2 of r04 already noted that $\frac{1}{2}(1 \pm J)$ is only real-linear for antilinear J.

v4.9.0-r03 a charge and a light wave in the interferometer

Theoremlet v4.9.0-r03.2.

(a) For the Coulomb field of a charge q (matter type, $s > 0$), $J_B f$ is the field of a magnetic charge q (tachyon type). $C(f)$ and $A(f)$ are the fields of dyons with electric charge $q/2$ and magnetic charge $\pm q/2$, with $s = \pm\frac{i}{2}|E_0|^2$.

(b) For a plane light wave, J_B reverses the direction of propagation and rotates the polarisation by 90°. This is what a phase-conjugating mirror (time reversal, r02.3) followed by a polarisation rotator (duality quarter turn) does. C of the wave has $E \parallel B$ and is not a light wave.

Proof sketch. J_B: $E \mapsto B$, $B \mapsto E$, and C, A give $E = \pm B = \frac{1}{2}E_0$. For $E = \hat{x}$, $B = \hat{y}$: J_B gives $E = \hat{y}$, $B = \hat{x}$, $E \times B = -\hat{z}$. Checked numerically. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Reading. For light the mirror of r02.3 is an existing optical device, but on light it does not change the type: $s = 0$ stays 0. A mirror that turns a matter-type field into a tachyon-type field of the same shape must turn electric charge into magnetic charge. Under the assignment of r07 the missing tachyon mirror is therefore, in Reading B, the missing magnetic charge. The periscope image, that is the overlay of a field with its mirror image, is a dyon. This is the content of the "photon at $\beta = 0$" of Theorem 21.3 and Theorem 21.7 when f is a static matter field.

v4.9.0-r03 what the interferometer reads out

Theoremlet v4.9.0-r03.3. Put a duality phase φ into the reference arm, $f \mapsto e^{i\varphi} f$. The difference of the two port energies is

$$D(\varphi) = \left|\tfrac{1}{2}(e^{i\varphi} f + J_B f)\right|^2 - \left|\tfrac{1}{2}(e^{i\varphi} f - J_B f)\right|^2 = \operatorname{Im}\bigl(e^{i\varphi} s\bigr).$$

The fringe has amplitude $|s|$ and phase $\arg s$. At $\varphi = 0$ it reads $2E \cdot B$, the side of the photon (r02.1). Matter and tachyon give fringes half a period apart, and light gives no fringe. The visibility is

$$V = \frac{|s|}{|f|^2} = \sqrt{1 - \Bigl(\frac{|E \times B|}{u}\Bigr)^2} = \frac{1}{\cosh 2\eta_f}, \qquad u = \tfrac{1}{2}(E^2 + B^2),$$

where η_f is the rapidity of the frame in which $E \parallel B$. That frame exists for every non-null field.

Proof sketch. $D(\varphi) = \operatorname{Re}\langle e^{i\varphi} f, J_B f\rangle$ and $\langle z, i\bar{z}\rangle = i\bar{s}$. Further $u^2 - |E \times B|^2 = \tfrac{1}{4}|s|^2$. The frame with $E \parallel B$ moves along $E \times B$ with speed v, $v/(1+v^2) = |E \times B|/(2u)$, so $|E \times B|/u = \tanh 2\eta_f$. Checked numerically, including the boost to $E \parallel B$. □

Reading. In this interferometer light gives a null result, and the visibility of any other field is the inverse Lorentz factor of its energy transport, $|E \times B|/u = \tanh 2\eta_f$. The double rapidity $2\eta_f$ appears unforced: the field's rest frame moves at $\tanh \eta_f$, its energy at $\tanh 2\eta_f$. If η_f were η_{sym}, the visibility would be $1/\gamma_L = 0.536$ and the energy speed 0.844. This is arithmetic. No field is identified with η_{sym}, and an identification would be a prediction requiring a pre-registration note.

v4.9.0-r03 sum and product

Proposition 16.1 calls $C(f)$ the projection onto $K_3 = e_1 e_2$ and states that it is orthogonal to both inputs. A projection is not orthogonal to its input: $\operatorname{Re}\langle f, Cf\rangle = |Cf|^2$.

Theoremlet v4.9.0-r03.4. The product of a field with its mirror image has the properties claimed for $e_3 = e_1 e_2$:

$$z \times J_B z = 2\, E \times B, \qquad z \cdot J_B z = 2iu.$$

The vector part is twice the Poynting vector, real and orthogonal to E and to B. The scalar part is the energy density. The pair $(u, E \times B)$ is null, $|E \times B| = u$, exactly for light.

Proof sketch. $z \times \bar{z} = -2iE \times B$ and $z \cdot \bar{z} = E^2 + B^2$. Checked numerically. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Consequence. The orthogonality in Proposition 16.1 holds for the product of f with Jf, not for their average. In Reading B the product yields the energy and momentum of the field, and it is light-like exactly on the photon sector. The average yields a dyon. Which of the two the main body means by the compensator is not decided here. Both readings are recorded.

v4.9.0-r03 register entries

statement	status after v4.9.0-r03
compensator C, partner A	the two output ports of a two-arm interferometer with J in one arm; real-linear, $A(f) = -iC(if)$ (r03.1)
Pauli item, H_+^+/H_-^+ split	with antilinear J a real/imaginary split, the sectors are i-images of each other (r03.1)
Thm. 21.3, Thm. 21.7, photon at $\beta = 0$	for a static matter field $C(f)$ is a dyon field (r03.2)
Prop. 16.1, orthogonal to both inputs	false for the average; true for the product $z \times J_B z = 2E \times B$ (r03.4)
tachyon mirror	on light: phase conjugation plus polarisation rotation; on matter-type fields it needs magnetic charge (r03.2)

v4.9.0-r03 ledger and guard update

$$R_{\text{comp}}^{v490\text{-}r03} = (0, 0, 1)$$

for (interferometer form of the compensator established, readout $D(\varphi) = \mathrm{Im}(e^{i\varphi}s)$ and visibility $1/\cosh 2\eta_f$ derived, choice between average and product reading open).

The layer spends no guard:

$$R_{\text{glob}}^{(v490\text{-}r03,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\text{guard}}^{v490\text{-}r03} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r04 Source-Only: The Compensator as a Matrix on the Electric–Magnetic Doublet

Purpose. On $\mathbb{C}^3$ the involution J_B of r02.3 is antilinear. It has no complex matrix, and the compensator built from it is only real-linear (r03.1). This layer writes the field as the real pair (E, B) instead. Then J_B, the compensator and the readout of r03 become ordinary 2×2 matrices, and the state of a field becomes a 2×2 matrix of the kind used for polarisation in optics. The computations are `llm/k40_dublett_matrix.py`. No value is compared with data. Earlier layers are not rewritten.
Source-only status. This v4.9.0-r04 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

Author's question (2026-10-21). Would it be better to put the compensator into a matrix?

v4.9.0-r04 the doublet and its matrices

Write $F = [\,E \mid B\,] \in \mathbb{R}^{3\times 2}$ for $z = E + iB$, and let $\sigma_x = \left(\begin{smallmatrix} 0 & 1 \\ 1 & 0 \end{smallmatrix}\right)$, $\sigma_z = \left(\begin{smallmatrix} 1 & 0 \\ 0 & -1 \end{smallmatrix}\right)$, R_α the rotation of $\mathbb{R}^2$ by α.

Theoremlet v4.9.0-r04.1. The operations of r02 act on F from the right by real 2×2 matrices:

$$e^{i\alpha} z \leftrightarrow F R_\alpha^{\mathsf{T}}, \qquad J_B z = i\bar{z} \leftrightarrow F\sigma_x, \qquad T:\ \bar{z} \leftrightarrow F\sigma_z, \qquad C:\ -z \leftrightarrow -F.$$

They generate $O(2)$: σ_x and σ_z are reflections and R_α rotations. The relation $J_B R_\alpha = R_{-\alpha} J_B$ of r02.5 is the relation between a reflection and a rotation in $O(2)$, and the Enigma composite of r02.5 is the matrix identity $R_{\pi/2}\sigma_x R_{\pi/2}^{-1} = -\sigma_x$. Spatial rotations act from the left, $F \mapsto RF$, and commute with all of these. Boosts mix E and B through cross products, are not of the form $F \mapsto LFM$, and do not commute with σ_x. The compensator and its partner are ordinary orthogonal projections,

$$C_{\mathrm{HC}} = \tfrac{1}{2}(1 + \sigma_x) = \tfrac{1}{2}\begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}, \qquad A = \tfrac{1}{2}(1 - \sigma_x) = \tfrac{1}{2}\begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix},$$

and FC_{HC} is the compensator of r02.4 and r03.

Proof sketch. $e^{i\alpha}(E+iB) = (E\cos\alpha - B\sin\alpha) + i(E\sin\alpha + B\cos\alpha)$; $i(E - iB) = B + iE$; $E - iB$. The boost formula $E' = \gamma(E + v \times B) - \ldots$ couples the columns through $v\times$. Checked numerically. □

Consequence. The antilinearity of r02.3 is not lost. It is the statement that σ_x is a reflection, $\det \sigma_x = -1$, while the duality circle consists of rotations. Over $\mathbb{R}$ the compensator is linear, and the obstruction of r03.1 disappears.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r04 the state matrix

Theoremlet v4.9.0-r04.2. The 2×2 Gram matrix $G = F^{\mathsf{T}} F$ of E and B decomposes as

$$G = u\,\mathbf{1} + \tfrac{1}{2}\big(\operatorname{Re} s\;\sigma_z + \operatorname{Im} s\;\sigma_x\big), \qquad \det G = |E \times B|^2, \qquad u^2 - \det G = \tfrac{1}{4}|s|^2.$$

Put $\rho = G/\operatorname{tr} G$ and $b = (\operatorname{Re} s, \operatorname{Im} s)/(2u)$, so that $\rho = \frac{1}{2}(\mathbf{1} + b_1\sigma_z + b_2\sigma_x)$. Then:

(a) $|b| = 1/\cosh 2\eta_f \le 1$, with η_f as in r03.3. $|b| = 1$ exactly when $E \parallel B$, and $b = 0$ exactly for a null field. Matter ($b_1 > 0$) and tachyon ($b_1 < 0$) lie on opposite sides of the unit disk, light at its centre, and the side of the photon of r02.1 is the sign of b_2.

(b) The duality rotation by α rotates b by 2α. J_B reflects $b_1 \mapsto -b_1$, and T reflects $b_2 \mapsto -b_2$.

(c) Under a boost the traceless part of G, that is s, is unchanged and u changes, so b moves along its own direction. It reaches $|b| = 1$ in the frame with $E \parallel B$ and tends to 0 as the boost grows.

G is symmetric and has no σ_y part, so the states fill a disk, not a sphere.

Proof sketch. $G_{11} = E^2$, $G_{22} = B^2$, $G_{12} = E \cdot B$; $\det G = E^2B^2 - (E \cdot B)^2$. (a) follows from r03.3. (b): $G \mapsto R_\alpha G R_\alpha^{\mathsf{T}}$, and the traceless part of a symmetric matrix turns by twice the angle. (c): s is Lorentz-invariant (r06.3). Checked numerically, including a boost scan through the frame $E \parallel B$. □

Reading. This is the formalism of Stokes parameters for the doublet (E, B) in place of two polarisation components. The doubled angle in (b) is the half-angle of the spinor of r06.4 seen from the other side: a duality rotation by α moves the state by 2α. Light is the unpolarised centre, and matter and tachyon are the two fully polarised ends of one axis. Boosting any field moves it towards light.

v4.9.0-r04 the interferometer as a measurement

Theoremlet v4.9.0-r04.3. The energy in the compensator port of r03.1 is $|FC_{\mathrm{HC}}|^2 = \operatorname{tr}(GC_{\mathrm{HC}}) = u + E \cdot B$. With the duality phase φ in the reference arm, the port difference of r03.3 is

$$D(\varphi) = \operatorname{Im}(e^{i\varphi} s) = 2u\; b \cdot (\sin\varphi, \cos\varphi).$$

The interferometer measures the component of b along a direction set by φ. The compensator is the measurement along σ_x, and its partner is the opposite outcome.

Proof sketch. C_{HC} is a symmetric projection, so $|FC_{\mathrm{HC}}|^2 = \operatorname{tr}(C_{\mathrm{HC}} G C_{\mathrm{HC}}) = \operatorname{tr}(GC_{\mathrm{HC}})$. Expand $\operatorname{Im}(e^{i\varphi} s)$. Checked numerically. □

Both readings of r03 in one matrix. The average reading of r03.4 is the projection $F \mapsto FC_{\mathrm{HC}}$, linear in F. The product reading is quadratic. Its scalar part is $\operatorname{tr} G = 2u$, its magnitude is $\sqrt{\det G} = |E \times B|$, and its direction is the cross product of the two columns of F. The trace, traceless part and determinant of G are energy, the invariant s, and the magnitude of the energy flux. The 3×2 matrix F carries in addition the direction of the flux, which G does not see.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r04 relation to the compensator matrix of the main body

Section H22 defines a *global Heisenberg-compensator matrix* only as the extension of $C = \frac{1}{2}(\mathrm{id}+J)$ to the multi-layer architecture, and a frame-balancing tool that splits $X = X^+ + X^-$ by J. The doublet supplies concrete instances in Reading B: $X^+ = FC_{\mathrm{HC}}$ and $X^- = FA$. In matrix form the leakage diagnostics of Definition H3.3, with $J = \sigma_x$ acting from the right, read (relative norms on a random field):

transport $\mathcal{M}$	$\lvert E_{\mathrm{comm}}\rvert/\lvert F\rvert$	$\lvert E_{\mathrm{leak}}\rvert/\lvert FC_{\mathrm{HC}}\rvert$
spatial rotation (left)	0	0
boost, $\eta = 0.6174$	1.113	0.505
duality rotation, $\alpha = 0.4$ (right)	0.779	0.389

This repeats r02 in matrix form. Left and right multiplication commute, so rotations cannot leak. Duality rotations leak because R_α does not commute with σ_x, and boosts leak because they do not factorise. The values for the boost differ from the r02 table because the random field differs; vanishing and non-vanishing agree. Identifying the Millennium matrix of the main body with a matrix on the doublet is not done here.

Open: the third axis. G is real and symmetric, so the state space is a disk. A σ_y part would need an antisymmetric component of the state matrix, for instance from complex amplitudes of time-harmonic fields. Whether such a third axis, which would turn the disk into a sphere, has a meaning in the source is not decided here.

v4.9.0-r04 register entries

statement	status after v4.9.0-r04
compensator $C = \frac{1}{2}(1+J)$	on the doublet (E,B) the linear projection $\frac{1}{2}(1+\sigma_x)$; J_B, T, duality, C generate $O(2)$ (r04.1)
state of a field	Gram matrix $G = u\mathbf{1} + \frac{1}{2}(\mathrm{Re}\, s\, \sigma_z + \mathrm{Im}\, s\, \sigma_x)$, $\det G = \lvert E\times B\rvert^2$; point b in the unit disk, light at the centre, $\lvert b\rvert = 1/\cosh 2\eta_f$ (r04.2)
interferometer of r03	measurement of b along a direction set by φ (r04.3)
global HC matrix (§H22)	doublet gives concrete $X^{\pm}$; identification with the Millennium matrix not made

v4.9.0-r04 ledger and guard update

$$R_{\mathrm{mat}}^{v490\text{-}r04} = (0,0,1)$$

for (doublet matrix form of the compensator established, state disk and readout as measurement established, identification with the global compensator matrix of the main body open).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r04,+)} = (0,1,1,1,1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r04} = (1,1,1,1,1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r05 Source-Only: The Millennium Matrix on the Doublet, a Third Axis near Rotating Black Holes, and a Photon from Matter and Tachyon

Purpose. Three questions of the author follow r04.

1. Can the Millennium matrix of the main body be the doublet matrix of r04? The author also asks for a more meaningful name.

2. Can amplitudes, the missing third axis of r04, enter near a black hole?

3. Can a photon arise when a tachyon meets a forming massive particle, and could such a photon run back and create the tachyon at its origin?

The computations are `llm/k41_schichttransport_kerr_photon.py`. No value is compared with data. Earlier layers are not rewritten.
Source-only status. This v4.9.0-r05 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r05 the Millennium matrix on the doublet

The main body asks three things of the Millennium matrix $\mathcal{M}$. It transports one object through its layers (§38). It commutes with the involution and with the compensator, condition (T4) of Definition 29.5. And the compensator is its spectral projector (Theorem 29.22).

Theoremlet v4.9.0-r05.1. On the doublet of r04, with $J = \sigma_x$ and $C_{\mathrm{HC}} = \frac{1}{2}(1+\sigma_x)$ acting from the right:

(a) The real 2×2 matrices satisfying (T4) are exactly $M = \lambda_+ C_{\mathrm{HC}} + \lambda_- A$. Together with spatial rotations acting from the left, these are all transports of the doublet compatible with (T4).

(b) For $\lambda_+ \neq \lambda_-$, C_{HC} is the spectral projector of M onto the eigenvalue λ_+, as Theorem 29.22 requires.

(c) Of the transports of Reading B, (T4) holds for spatial rotations, for J_B itself and for the duality rotation by π, that is C. It fails for every other duality rotation, for T and for boosts (r02, r04).

(d) The invertible members with determinant 1 are $e^{t\sigma_x} = e^t C_{\mathrm{HC}} + e^{-t} A$. They preserve $\mathrm{Re}\, s = E^2 - B^2$, the matter–tachyon coordinate, and weight the two ports of r03 by $e^{\pm t}$.

Proof sketch. $M\sigma_x = \sigma_x M$ is a linear condition with a two-dimensional solution space spanned by 1 and σ_x. ${e^{t\sigma_x}}^{\mathsf{T}} \sigma_z e^{t\sigma_x} = \sigma_z$ since $\sigma_x \sigma_z = -\sigma_z \sigma_x$. Checked numerically. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Consequence. On the doublet the Millennium matrix can only be a *port-weighting matrix*. It leaves the compensator port and the partner port invariant and multiplies them by two factors. It is not the Lorentz transport of Reading B, since boosts violate (T4). The doublet therefore realises the compensator part of $\mathcal{H}_{16}$, not the full $\mathcal{M}_{16}$. The four-term layer decomposition $\mathcal{M}_{\mathrm{tri}} + \mathcal{M}_{\mathrm{alg}} + \mathcal{M}_{\mathrm{cl}} + \mathcal{M}_{\mathrm{proj}}$ and the index 16 have no counterpart on a two-dimensional doublet. With $t = \eta_{\mathrm{sym}}$ the port weights would be the Doppler factors $e^{\pm\eta}$ of way (C) in r01. That is arithmetic, not an identification.

Naming (author's request, 2026-10-21). From this layer on, $\mathcal{M}_{16}$ is called the *layer-transport matrix*. The name says what §38 defines: one object carried through its layers. Its restriction to the doublet is called the *port-weighting matrix*. Symbols, labels and the wording of earlier layers stay unchanged, so "Millennium matrix" in earlier text means the same object. A renaming of the earlier text and of the PDF keywords would rewrite released layers. It is not done here.

v4.9.0-r05 a third axis, and rotating black holes

Theoremlet v4.9.0-r05.2.

(a) At a point, E and B are real, so the state matrix of r04 has no σ_y part anywhere, near a black hole or not.

(b) For time-harmonic fields with complex amplitudes $\hat{E}, \hat{B}$, the matrix of time averages $\hat{G} = \hat{F}^\dagger \hat{F}$ is Hermitian, and its σ_y part is $-2\,\mathrm{Im}(\hat{E}^* \cdot \hat{B})$. For a plane wave this is ± 1 (relative to $\mathrm{tr}\,\hat{G}$) for the two circular polarisations and 0 for linear polarisation. The third axis is helicity, the quantity generated by the duality rotation. With it the disk of r04 becomes a sphere, with light on the helicity axis.

(c) The gravitational field has the same structure. Its Weyl curvature is a complex symmetric 3×3 matrix, gravito-electric plus i gravito-magnetic. For a Schwarzschild black hole the only component $\Psi_2 = -M/r^3$ is real. For a Kerr black hole $\Psi_2 = -M/(r - ia\cos\theta)^3$ is complex. $\mathrm{Im}\,\Psi_2^2$, the Pontryagin part of the curvature, is positive in one hemisphere, zero on the equator and negative in the other.

Proof sketch. (b): $\mathrm{tr}(\hat{G}\sigma_y) = -2\,\mathrm{Im}\,\hat{G}_{12}$; for $\hat{B} = \hat{z} \times \hat{E}$ with $\hat{E} = (1, \pm i, 0)/\sqrt{2}$ one gets ∓ 1 up to orientation. (c): standard form of Ψ_2 in the Kinnersley tetrad; the sign of $\mathrm{Im}\,\Psi_2^2$ follows the sign of $\cos\theta$. Checked numerically for $M = 1$, $a = 0.9$, $r = 3$. □

Reading. Amplitudes do not enter because a black hole is near. They enter with waves and quantum states. What a black hole adds depends on its rotation. A static one is mirror-symmetric and singles out neither helicity. A rotating one puts its own curvature off the real axis, on opposite sides in its two hemispheres, which is the gravitational counterpart of the two sides of the photon in r02.1. Whether this couples to the helicity axis of (b), and whether Hawking radiation of a rotating black hole would then carry a helicity imbalance between the hemispheres, is open. A quantitative statement is a prediction and requires a pre-registration note. Theorem 21.7 treats the static case only.

v4.9.0-r05 a photon from matter and tachyon

Author's question (2026-10-21). A tachyon meets a massive particle that is just forming. Instead of reaching a massive state, the encounter yields a photon. The tachyon develops a

© 2026 Marc Brendecke. Preprint — not peer reviewed.

pull through the impact, so the photon is drawn along the tachyon's former path, collides again and again, reaches the tachyon's origin, vanishes there, and thereby creates the tachyon.

Theoremlet v4.9.0-r05.3.

(a) Let z_M be of matter type and z_T of tachyon type with $s(z_T) = -s(z_M)$. Then $s(z_M+z_T) = 2\,z_M \cdot z_T$. The superposition is a null field exactly when $z_M \cdot z_T = 0$. For a static electric field E and a static magnetic field B of equal strength, the superposition is null exactly when $E \perp B$. Its energy then flows at the speed of light along $E \times B$, perpendicular to both.

(b) At an angle ϑ between E and B, the superposition sits at $b = (0, \cos\vartheta)$ in the disk of r04. $\vartheta = 90°$ is the photon at the centre, and $\vartheta = 0$ is the dyon of r03.2 at the edge.

(c) Let a tachyon leave the origin O and meet the particle at K, a distance d away, at speed $v > 1$. A photon running back from K to O arrives $d(1 + 1/v)$ after the departure, and this arrival lies in the future light cone of the departure in every frame. The order of departure and meeting is frame-dependent (for observer speeds above $1/v$ it reverses), but the photon leg is not.

Proof sketch. (a) Expand s. For $z_M = E$, $z_T = iB$: $z_M \cdot z_T = iE \cdot B$. Null fields have $|E \times B| = u$ (r03.3). (b) $s = 2i|E|^2 \cos\vartheta$ and $u = |E|^2$. (c) The separation of departure $(0, 0)$ and photon arrival $(d/v + d, 0)$ is timelike and future-directed; the tachyon leg is spacelike. Checked numerically for $v = 1/\beta_{\text{sym}}$. □

Reading. The first part of the picture has a precise form. Matter and tachyon can combine to a photon, but only with crossed orientation. With parallel orientation they give the dyon of r03. The photon does not move along the tachyon's path. It moves perpendicular to both inputs. The last part, a photon that arrives at the origin and thereby creates the tachyon, cannot hold for a photon running forward in time: it always arrives after the tachyon has left. What does hold is the familiar reinterpretation of tachyon exchange. Since the tachyon leg is spacelike, "emitted at O, absorbed at K" and "emitted at K, absorbed at O" are the same exchange seen by different observers. A closed self-creation would need a photon leg running backward in time. Without one, the repeated encounter of the picture is a cycle of period $d(1 + 1/v)$, not a self-creation. If such a cycle returns on the other side of the photon, the tachyon created in the next cycle is the charge conjugate of the first (r02.1). Reading B has no dynamics (r10), so the pull of the picture cannot be derived or excluded here.

v4.9.0-r05 register entries

statement	status after v4.9.0-r05
Millennium matrix, (T4), Thm. 29.22	on the doublet only $\lambda_+ C_{\text{HC}} + \lambda_- A$ (port weighting); boosts and general duality rotations violate (T4); full $\mathcal{M}_{16}$ not identified (r05.1)
name	$\mathcal{M}_{16}$: layer-transport matrix; doublet restriction: port-weighting matrix; earlier wording unchanged
third axis of r04	helicity, for complex amplitudes; Kerr curvature off the real axis with opposite sides in the two hemispheres; coupling open (r05.2)
photon from matter + tachyon	null exactly for crossed orientation, flux perpendicular to both; return photon cannot create the tachyon (r05.3)

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r05 ledger and guard update

$$R_{\mathrm{lt}}^{v490\text{-}r05} = (0, 0, 1)$$

for (doublet form of the layer-transport matrix decided, third axis and photon question answered structurally, identification of the full $\mathcal{M}_{16}$ and the Kerr coupling open).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r05,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r05} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r06 Source-Only: The Tachyon Hit Chain, its Acceleration, and the Higgs Field

Purpose. Theoremlet r05.3(c) stated that a photon returning to the origin of a tachyon cannot create it. The author objected that the tachyon already runs backward in time, and then refined the picture. The Higgs field tries to create matter along a line. A tachyon that has been travelling along this line for a long time hits it again and again. Each hit is a point of spacetime at which an essentially identical photon arises. This layer makes r05.3(c) precise, works out the hit chain, asks what an accelerating or infinitely accelerated tachyon does to the hits, and checks the picture against the Higgs field. Special relativity is used with $c = 1$. The computations are `llm/k42_tachyon_trefferkette.py`. No value is compared with data. Earlier layers are not rewritten.
Source-only status. This v4.9.0-r06 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r06 closing the loop, made precise

Theoremlet v4.9.0-r06.1.

(a) A photon needs the time d for the distance d. A tachyon of speed $v > 1$ covers the same distance in the time d/v, and even read backward it takes back only $d/v < d$. One photon leg and one tachyon leg therefore never close a loop. The remaining gap is $d(1 - 1/v)$, which is $0.451\,d$ for $v = 1/\beta_{\mathrm{sym}}$.

(b) Two tachyon legs emitted in frames moving relative to each other can close or overshoot a loop. A return leg that is instantaneous in a frame of speed u arrives at the origin at $d/v - ud$, before the start for $u > 1/v$.

(c) For $v = 1/\beta_{\mathrm{sym}}$ and $u = \beta_{\mathrm{sym}}$ the return arrives exactly at the start. It runs back along the same spacelike line, because the outgoing leg is itself instantaneous in that frame. This is one exchange read in two directions, case (a) of Theoremlet r02.1, and carries no information around a loop.

Proof sketch. Arithmetic on the worldlines; the return leg is the line of simultaneity $t - t_B = u(x - x_B)$ through $B = (d/v, d)$. Checked numerically. □

Correction of wording in r05. The phrase "a photon leg running backward in time" in the reading of r05.3 is made precise by (a) and (b). What fails is one null leg plus one tachyon leg. Two tachyon legs do not fail. The statement r05.3(c) itself stands.

v4.9.0-r06 the hit chain at constant speed

Theoremlet v4.9.0-r06.2. Let hits occur along a line at speed $v > 1$, each emitting an identical light signal. The energy of each signal comes from the field at the hit, so nothing is carried faster than light.

(a) The signals superpose to a cone, the envelope of their Huygens wavelets, with $\cos\vartheta = 1/v$. For $v = 1/\beta_{\mathrm{sym}}$ this gives $\cos\vartheta = \beta_{\mathrm{sym}}$, $\vartheta = 56.68°$.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(b) The hits are spacelike separated. Observers slower than $1/v$ see them run in the direction of the tachyon, observers faster than $1/v$ see them run the other way, and the observer of speed exactly $u = 1/v$ sees all hits at once. For $v = 1/\beta_{\text{sym}}$ that observer moves at β_{sym}, the matter-frame offset.

(c) The superluminal rapidity of r20.2 is $\operatorname{artanh} v = \eta + i\pi/2$. The boost to the observer of (b) removes η and leaves $i\pi/2$, the duality quarter turn of r07.2. In that frame the tachyon is no longer a moving object but a phase. The plane wave emitted there perpendicular to the line appears in the original frame at $\cos\vartheta' = \beta_{\text{sym}}$, which is the cone of (a).

(d) With imaginary mass $i\mu$, the tachyon at $v = 1/\beta_{\text{sym}}$ has energy $\mu \sinh\eta$ and momentum $\mu\cosh\eta = 1.197\,\mu$, the neighbour cross term of r18.

Proof sketch. (a) Minimal arrival time over emission points; checked by a discrete Huygens sum. (b) $\Delta t' = \gamma(\Delta t - u\Delta x)$ with $\Delta t = \Delta x/v$. (c) $\tanh(x + i\pi/2) = \coth x$ and the aberration formula with $\vartheta = 90°$. (d) $E = \mu/\sqrt{v^2-1}$, $p = \mu v/\sqrt{v^2-1}$. □

Reading. "The tachyon was already on its way before the experiment started" is the frame dependence of (b). There is an observer for whom the whole path exists at one instant. For $v = 1/\beta_{\text{sym}}$ this observer is the matter frame. A superluminal chain of locally triggered emissions is the situation of a light spot sweeping faster than light across a screen. It is permitted because each point emits by itself.

v4.9.0-r06 acceleration and infinite speed

Theoremlet v4.9.0-r06.3.

(a) *Constant proper acceleration.* The tachyonic counterpart of hyperbolic motion is $t = R\cosh\sigma$, $x = R\sinh\sigma$, with proper acceleration $1/R$ everywhere. Its speed $v = \coth\sigma$ and its coordinate acceleration dv/dt diverge at $\sigma = 0$, that is $t = R$, $x = 0$. The hit rate per unit time diverges there as well, so the hits near that point arrive at once as a flash. Beyond $\sigma = 0$ the parameter runs backward in time. By reinterpretation that part is a partner moving forward, and the point $(R, 0)$ is a pair event. A signal emitted along the backward part is read as absorbed.

(b) *Energy given to the hits.* Write $E = \mu\sinh\chi$, $p = \mu\cosh\chi$, $v = \coth\chi$. If each hit takes the energy ε from the tachyon, E falls and v rises. After E_0/ε hits $E = 0$, $v = \infty$, $p = \mu$. Along the way the cone of r06.2 opens from 56.68° to 90°, and the observer seeing simultaneous hits slows from β_{sym} to rest.

In both cases the infinite acceleration is a coordinate effect at a single point. There the tachyon passes through infinite speed, rapidity $i\pi/2$ in the frame at hand. The proper acceleration stays finite.

Proof sketch. (a) $dx/dt = \coth\sigma$; $dv/dt = -1/(R\sinh^3\sigma)$; the second derivative with respect to proper length has Minkowski norm $1/R$. (b) $p^2 - E^2 = \mu^2$ and $v = p/E$. Checked numerically, (b) with six hits from $v = 1/\beta_{\text{sym}}$. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Reading. A tachyon that pays for the photons of its hits accelerates by itself and ends at infinite speed after finitely many hits. At that point the lab frame has become the frame of simultaneous hits, the cone has become a plane wave, and the tachyon is the pure phase of r06.2(c). Continuing past this point turns emission into absorption in the lab description. This is the nearest the hit chain comes to the author's first picture of photons returning to the origin: a chain of emissions meets a chain of absorptions at the point of infinite speed.

v4.9.0-r06 the Higgs field

The main body assigns the Higgs sector the role of mediating between the compensated photon fix-sector and the mass readout of localised branches (Definition 40.12, Theorem 40.13).

Theoremlet v4.9.0-r06.4.

(a) In the electroweak theory the Higgs field gives the neutral pair (W^3, B) the mass matrix $\frac{v_H^2}{4}\left(\begin{smallmatrix} g^2 & -gg' \\ -gg' & g'^2 \end{smallmatrix}\right)$. It has rank one: the photon is its massless eigenvector $(\sin\theta_W, \cos\theta_W)$, the Z the massive one. This is a port-weighting matrix in the sense of r05.1, with weight 0 on the photon port and m_Z^2 on the other. For $g = g'$ ($\theta_W = 45°$) it is exactly $0 \cdot C_{\mathrm{HC}} + m_Z^2 A$, and the photon is the compensator port. The measured mixing angle differs from 45°, so the electroweak photon port is a compensator rotated by $\theta_W - 45°$.

(b) At $\phi = 0$ the Higgs potential has negative mass squared. The field there is tachyonic in the field-theoretic sense, and mass arises when it condenses away from $\phi = 0$.

Proof sketch. (a) The kernel of the matrix is spanned by (g', g); for $g = g'$ the matrix is $\frac{v_H^2 g^2}{2} A$. (b) The potential $-\mu_H^2|\phi|^2 + \lambda|\phi|^4$ has curvature $-\mu_H^2$ at the origin. □

Reading. The Higgs part of the author's picture fits the main body in three respects.

- The photon is the direction the Higgs field does not weight, which is Theorem 40.13 in electroweak form.
- A field "trying to create matter" is a field leaving its tachyonic top.
- A hit that yields a photon instead of a massive state lands in the unweighted port.

Three points need care.

- The doublet of r04 is (E, B), the electroweak doublet is (W^3, B). The correspondence is formal.
- A field with negative mass squared is an instability that grows; it is not a particle that carries signals faster than light. A superluminal chain of hits therefore needs triggers correlated in advance, as for the sweeping light spot, and not a signal along the line.
- The condensed Higgs vacuum is the minimum of its potential and supplies no energy. The energy of the photons comes either from the tachyon, case r06.3(b), or from the field while it is still rolling off its tachyonic top, that is during symmetry breaking.

Comparing θ_W or any other measured quantity with the values of this layer is a prediction and requires a pre-registration note.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r06 register entries

statement	status after v4.9.0-r06
r05.3(c), return photon	stands; one photon plus one tachyon leg leaves the gap $d(1-1/v)$; two tachyon legs can close a loop (r06.1)
hit chain at speed $1/\beta_{\mathrm{sym}}$	cone 56.68°; all hits simultaneous for the observer at β_{sym}, where the tachyon is the phase $i\pi/2$ (r06.2)
accelerating tachyon	infinite speed at one point after finite time or finitely many hits; there a pair event, emission turns into absorption (r06.3)
Higgs mediation (Def. 40.12)	electroweak mass matrix is a port weighting with massless photon port; compensator port for $\theta_W = 45°$; formal correspondence (r06.4)

v4.9.0-r06 ledger and guard update

$$R_{\mathrm{chain}}^{v490\text{-}r06} = (0,0,1)$$

for (loop condition decided, hit chain and acceleration worked out, physical status of the trigger and of the Higgs correspondence open).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r06,+)} = (0,1,1,1,1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r06} = (1,1,1,1,1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r07 Source-Only: The Cone Angle and the Weak Mixing Angle

Purpose. This layer records three things. First, the pre-registered test H14 of the relation between the cone angle of r06.2 and the weak mixing angle, with its result. Second, an exploration that followed it, kept as a protocol in which every variant is counted. Third, the structural reading that came out of the exploration, and the pre-registered hypothesis H15 it led to. Files, all under `llm/`:

- `Vorab_Notiz_H14_Kegelwinkel_Weinberg.md`;
- `Ergebnis_H14_Kegelwinkel_Weinberg.md`;
- `Erkundungsprotokoll_Weinberg_Kegel.md`;
- `Vorab_Notiz_H15_W_Masse_gamma_L.md`;
- `Ergebnis_H15_W_Masse_gamma_L.md`;
- scripts `k43_test_h14.py` to `k46_test_h15.py`.

Data: Particle Data Group, Review of Particle Physics 2024 with the 2025 update (summary tables and the review of the electroweak model), retrieved 2026-10-21. Earlier layers are not rewritten.
Source-only status. This v4.9.0-r07 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r07 test H14

H14 states $\vartheta_c = 2\theta_W$, equivalently $\sin^2\theta_W = (1-\beta_{\mathrm{sym}})/2$. It was formulated after the closeness 56.7° against $2\theta_W \approx 57.5°$ had been noticed. The note says so, fixes the on-shell definition $\sin^2\theta_W = 1 - m_W^2/m_Z^2$ as decisive, and states the expected outcome in advance.

Result. The prediction is 0.22533. The on-shell value from the masses is 0.22321 ± 0.00026, so $z = +8.2$. As an exact relation H14 is **refuted**. The prediction lies inside the band spanned by the on-shell and the $\overline{\mathrm{MS}}$ value (0.23122), so as a tree-level relation it is **not decidable**. A random simple relation would land in that band with probability of about 0.02, so no outcome counts as support.

v4.9.0-r07 exploration, counted

The exploration asked whether a physical effect can close the gap of +0.00212 in $\sin^2\theta_W$. The gap is equivalent to removing a contribution $\Delta\rho_X = +0.00275$ present in nature, or to an m_W smaller by 110 MeV. The variants tried, each counted, were:

- the tribonacci value in place of β_{sym}, which moves the prediction the wrong way;
- removal of the top-quark contribution, which overshoots to 0.2304;
- the tree-level angle from (α, G_F, m_Z), giving 0.212 or 0.234 depending on α;

© 2026 Marc Brendecke. Preprint — not peer reviewed.

- a fraction of the top contribution, with no meaning;
- neutrinos, eleven orders of magnitude too small;
- every combination of omitted contributions, of which only two are large (top 0.00934, remainder 0.00108) and none hits;
- mixing with a dark state.

The last variant gives a general bound. Any real contribution of the required size would shift m_W by 155 MeV against the Standard-Model prediction. The measured value differs from that prediction by $+10 \pm 14$ MeV. Real effects of particles, known or dark, can therefore account for at most about a quarter of the gap.

v4.9.0-r07 the structural reading

Theoremlet v4.9.0-r07.1. With $\Pi(x) = 2\arctan e^{-x}$, Lobachevsky's angle of parallelism, the following are equivalent:

$$\cos 2\theta = \tanh \eta, \qquad \tan\theta = e^{-\eta}, \qquad \sin^2\theta = \frac{1}{1+e^{2\eta}}, \qquad 2\theta = \Pi(\eta).$$

The cone angle of r06.2 is $\vartheta_c = \Pi(\eta)$. For the port weighting $e^{t\sigma_x} = e^t C_{\mathrm{HC}} + e^{-t} A$ of r05.1(d) at $t = \eta$, the ratio of the weights is $e^{2\eta} = \cot^2\theta$.

Proof sketch. $\cos 2\theta = (1-\tan^2\theta)/(1+\tan^2\theta)$ with $\tan^2\theta = e^{-2\eta}$ gives $\tanh\eta$; $\cos\Pi(x) = \tanh x$. Checked numerically. □

Consequence. H14 is a statement about a ratio of couplings, $g/g' = e^{\eta}$, the Doppler factor of way (C) in r01. It is not a statement about a ratio of masses. Couplings run. In the $\overline{\mathrm{MS}}$ scheme $\sin^2\theta_W$ has its minimum, about 0.231, at $\mu = M_W$, and rises above and below. No scale reaches 0.22533. The structural reading thus points to the scheme in which the gap is larger, and it does not support the closeness to the on-shell value.

v4.9.0-r07 an observation, and hypothesis H15

Observation (found during the exploration). The on-shell coupling ratio is $\cot\theta_W = m_W/\sqrt{m_Z^2 - m_W^2} = 1.86550 \pm 0.00140$. The three constants of the source in this range sit at the following distances: $\gamma_L = \cosh 2\eta = 1.86444$ at $z = -0.8$, $e^{\eta} = 1.85419$ at $z = -8.1$, $1/\beta_{\mathrm{sym}} = 1.82034$ at $z = -32$. By Theoremlet r01.3, γ_L (way A) and e^{η} (way C) coincide at the tribonacci point and differ by 0.55% otherwise. The data sit at way (A). The cone relation used way (C).

Why this is not evidence. The relation was found after the fact. The three constants 1.820, 1.854, 1.864 lie in the range that $\cot\theta_W$ covers across schemes, 1.823 to 1.866, so some pairing is nearly expected. The chance of a 1σ hit of one of the three at the on-shell value is about 0.17. Reading B contains no $SU(2) \times U(1)$, so the identification of the weight ratio with $(g/g')^2$ is formal.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Hypothesis H15 (pre-registered). $\cot\theta_W = \gamma_L$ in the on-shell definition, that is $m_W = m_Z\gamma_L/\sqrt{1+\gamma_L^2}$. Here γ_{BI} is the root of the state-counting equation $\sum_j 2\exp(-2\pi\gamma\sqrt{j(j+1)}) = 1$ that r03 cites, and $\gamma_{\mathrm{string}} = \ln 2/(\pi\sqrt{3})$. Computed immediately after the note was committed and before any comparison:

$$\gamma_{\mathrm{BI}} = 0.237532957966, \qquad \gamma_L = 1.8646997646, \qquad m_W^{\mathrm{pred}} = 80.3615 \pm 0.0018 \text{ GeV}.$$

The comparison is with the first world average or combination of m_W published after 2026-10-21. Known values do not count. The prediction lies within a few MeV of the Standard-Model prediction (80.359 ± 0.006 GeV). A comparison can therefore refute H15, but it can support H15 over the Standard Model only once the measurement excludes the Standard-Model value at 3σ, which needs a precision below about 1 MeV. The result is pending.

v4.9.0-r07 register entries

statement	status after v4.9.0-r07
H14, $\vartheta_c = 2\theta_W$	refuted as exact relation ($z = +8.2$); not decidable at tree level; no support
cone angle	$\vartheta_c = \Pi(\eta)$, the angle of parallelism (r07.1); not measured
gap of H14 by real particle effects	bounded by m_W to at most about a quarter (exploration, 8 counted variants)
H15, $\cot\theta_W = \gamma_L$	pre-registered; prediction $m_W = 80.3615 \pm 0.0018$ GeV; comparison pending
r01 ways (A)/(C)	the on-shell coupling ratio lies at way (A) (observation, not a test)

v4.9.0-r07 ledger and guard update

$$R_{\mathrm{ew}}^{v490\text{-}r07} = (0, 0, 1)$$

for (H14 decided, structural reading of the cone established, H15 pending).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r07,+)} = (0,1,1,1,1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r07} = (1,1,1,1,1), \qquad \beta_4 = 1/256 \text{ unspent}.$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r08 Source-Only: Frame Drift, Atomic Clocks and the Hubble Tension

Purpose. Proposition 42.5 explains the Hubble tension by a drift of the frame offset. r17 and r18 tested that drift against the expansion history. This layer tests it against the constancy of the fine-structure constant, using the source's own rule for which quantities stay fixed. It also checks the other explanations of the tension available in the source, settles the sign of k for α left open in r03, names a frame-invariant clock, and states what the drift means for H15 of r07. Files: `llm/Erkundungsprotokoll_Weinberg_Kegel.md` (entries E10 to E13) and scripts `llm/k47_drift_weinberg.py`, `llm/k48_feste_groessen.py`. No new prediction is made. Earlier layers are not rewritten.
Source-only status. This v4.9.0-r08 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r08 what stays fixed under drift

The source fixes this in four places.

- The definition in §25: $X^{(\text{photon})} = \gamma_L^{-k(X)} X^{(\text{matter})}$.
- Theorem 25.3: $k(c) = 0$, $k(\gamma_{\text{BI}}) = 0$ ("frame-invariant by construction").
- The corollary "α-drift as a frame effect" in the same section: "the hardware value $\alpha^{(\text{photon})}$ is strictly constant", with $\alpha^{(\text{matter})}(t) = \gamma_L^{(\text{local})}(t)\, \alpha^{(\text{photon})}$.
- Item (1) of Proposition 42.8, which writes the neutrino mass in the same way.

Photon-frame values and quantities with $k = 0$ stay fixed. Matter-frame values drift with a local factor $\gamma_L^{(\text{local})}(t)^k$. The structural factor $\gamma_L = \gamma_{\text{BI}}/\gamma_{\text{string}}$ is fixed, since both inputs are fixed numbers. The source thus separates a fixed structural factor from a local one that may drift. This is the separation r18 offered.

v4.9.0-r08 the sign of k for α

r03 (F2) found that Theorem 25.3, whose row reads "$\alpha^{-1} \ldots k = 1$", and the corollary above are inverse to each other, and left the choice open. The main body uses k for α in three places:

- prediction P5 among the predictions around Corollary 42.24, with $\gamma_L^{k(\alpha)-k(G_N)} = \gamma_L^{-1}$;
- Proposition 42.7, "$k(\alpha_s) = 1$ (like the electromagnetic coupling α)" with $\alpha_s^{(\text{photon})} = \gamma_L^{-1} \alpha_s^{(\text{matter})}$;
- the corollary, $\alpha^{(\text{matter})} = \gamma_L\, \alpha^{(\text{photon})}$.

All three mean $k(\alpha) = +1$ for α itself. No photon-frame value of α^{-1} is used anywhere in the main body.

Resolution (reading decision). The table row shows the CODATA value of α^{-1}, and its entry $k = 1$ belongs to α. With this reading all three uses agree and nothing else changes. The resolution changes no number of the main body.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r08 the drift against atomic clocks

Theoremlet v4.9.0-r08.1. Suppose the drift of Proposition 42.5, 1.6×10^{-12} per year in β_{sym}, acts through the local frame factor, as r18 reads it. Then the corollary gives $|\mathrm{d}\ln\alpha/\mathrm{d}t|$ for each choice of the factor:

factor	exponent needed for 1.083	$\lvert\mathrm{d}\ln\alpha/\mathrm{d}t\rvert$ per year	$\lvert\Delta\alpha/\alpha\rvert$ since $z = 1100$
$\gamma_L = \cosh 2\eta$	1.52	3.9×10^{-12}	5.4%
neighbour, $\cosh\eta$	4.72	1.3×10^{-12}	1.7%
neighbour, e^{η}	2.54	2.3×10^{-12}	3.2%

The comparison of two $^{171}\mathrm{Yb}^{+}$ clocks gives $(1/\alpha)\,\mathrm{d}\alpha/\mathrm{d}t = (1.0 \pm 1.1) \times 10^{-18}$ per year (Lange et al., Phys. Rev. Lett. 126, 011102, 2021). Every choice lies about six orders of magnitude above that bound.

Independently of the reading: if a drifting factor with exponent $k(H_0)$ produces the tension $\ln 1.083$, then α changes by $\ln 1.083 \cdot k(\alpha)/k(H_0)$. For a drift linear in time, the clocks require $k(\alpha)/k(H_0) < 5.5 \times 10^{-7}$ at 2σ.

Proof sketch. $\mathrm{d}\ln\gamma_L/\mathrm{d}t = 4\beta_{\mathrm{sym}}(1-\beta_{\mathrm{sym}}^2)^{-2}\dot{\beta}_{\mathrm{sym}}/\gamma_L$, and correspondingly for the neighbour factors. The exponents follow from the ratios of the factors at $z = 0$ and $z = 1100$. Checked numerically. □

Why measuring through photons does not help. A clock comparison measures the ratio of two transition frequencies of atoms. A common frame factor on both frequencies cancels. The dependence on α comes from inside the atom, which is matter. If the atom's internal physics were photon-frame physics, there would be no drifting $\alpha^{(\mathrm{matter})}$ at all. The corollary's annual-drift prediction would then lapse too, and Proposition 42.5 would lose its mechanism.

Consequence. In its frame-factor form, Proposition 42.5 cannot explain the tension and keep α constant at the same time. Two ways out would remain, and the source contains neither. One is $k(\alpha) \approx 0$, which contradicts the resolution above and the corollary. The other is a separate factor for the expansion that does not touch the constants. r03 already bounded the annual modulation of the local factor by 10^{-17}. The present layer applies the same clocks to the secular drift.

v4.9.0-r08 other explanations in the source

1. *Static frame typing.* The value inferred from the microwave background would be read in another frame than the local value. A constant factor disturbs no clock. By Corollary 22.21 a clock frequency transforms with γ_L, so $k(H_0) = 1$, which gives a factor 1.864 (neighbour factors 1.197, 1.854) against the observed 1.083 ± 0.017. Only fractional exponents near 0.13 or 0.44 would fit, and Theorem 25.3 gives no basis for them. The observed ratio is too loose to single out an exponent.

2. *A change confined to the early universe.* A larger electron mass at recombination, by about one to two percent, is discussed in the literature as a relief of the tension. With the drift of Proposition 42.5, β_{sym} was smaller earlier, so $m_e^{(\mathrm{matter})}$ with $k = 1$ was smaller. That is the wrong direction, and the same holds for α with $k(\alpha) = +1$. The opposite sign of the drift, the one r17 found Theorem 22.16 to require, would point the right way.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

The change would then have to stop before $z \approx 3$, because of quasar spectra and clocks, and at 5% it would be too large.

No explanation of the tension within the source survives the clocks, the quasar spectra and the size of the effect together. Proposition 42.5 stays a hypothesis without a working mechanism.

v4.9.0-r08 a frame-invariant clock

Theoremlet v4.9.0-r08.2. By Theorem 25.3 the Compton frequency $m_e c^2/\hbar$ has $k = k(m_e c^2) - k(\hbar) = 1 - 1 = 0$. A clock ticking at a Compton frequency is frame-invariant in the source's sense. An atomic transition frequency scales as $\alpha^2 m_e c^2/\hbar$ and has $k = 2k(\alpha) = 2$. The ratio of the two clocks reads $\gamma_L^{(\mathrm{local})}(t)^{\pm 2}$ directly, a local test of the frame factor that needs no cosmology.

Proof sketch. Additivity of k (Lemma v4.8.0-r03.1). □

Remark. Clocks tied to the Compton frequency of a mass have been built by atom interferometry (Lan et al., Science 339, 554, 2013). A comparison of such a clock with an atomic clock would read the drift of r08.1 at twice the rate. Existing clock comparisons already exclude that drift. The clock concerns local readout only. It touches neither the local selector guard nor the forbidden global photon-clock statement.

v4.9.0-r08 consequence for H15

H15 (r07) is stated with the structural factor $\gamma_L = \gamma_{\mathrm{BI}}/\gamma_{\mathrm{string}}$, which is fixed. It predicts a weak mixing angle that is constant in time. In addition, m_W/m_Z is a ratio of masses, and if both carry the same k the frame factor cancels. A drift of θ_W, with $\sin^2\theta_W = 0.242$ at $z = 1100$, would follow only if H15 were read with the local factor, which the source's rules do not suggest. The pre-registered comparison of H15 is unaffected.

v4.9.0-r08 register entries

statement	status after v4.9.0-r08
Prop. 42.5	frame-factor drift incompatible with atomic clocks by about six orders of magnitude in every reading (r08.1); static typing and early-only change do not work; hypothesis without working mechanism
Thm. 25.3, row α^{-1}	resolved: $k = 1$ belongs to α; consistent with P5, Prop. 42.7 and the corollary (closes r03 F2)
fixed quantities under drift	photon-frame values and $k = 0$; structural $\gamma_L = \gamma_{\mathrm{BI}}/\gamma_{\mathrm{string}}$ fixed; local factor may drift
Compton clock	$k = 0$, frame-invariant; ratio to an atomic clock reads $\gamma_L^{(\mathrm{local})\pm 2}$ (r08.2)
H15	predicts a constant θ_W; comparison unaffected

v4.9.0-r08 ledger and guard update

$$R_{\mathrm{drift}}^{v490\text{-}r08} = (0, 0, 1)$$

for (drift against clocks decided, sign of $k(\alpha)$ resolved, a mechanism for the Hubble tension open).

© 2026 Marc Brendecke. Preprint — not peer reviewed.

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r08,+)} = (0,1,1,1,1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r08} = (1,1,1,1,1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r09 Source-Only: What the Local Frame Factor Can Be

Purpose. r08 separated a fixed structural factor $\gamma_L = \gamma_{\mathrm{BI}}/\gamma_{\mathrm{string}}$ from a local factor $\gamma_L^{(\mathrm{local})}$ that may vary. This layer collects what the source says the local factor depends on. It records two contradictions found on the way, tests three readings of the local factor against clock data, and corrects the arithmetic of prediction P5. Files: `llm/Erkundungsprotokoll_Weinberg_Kegel.md` (entries E14, E15) and the script `llm/k49_lokaler_faktor.py`. No new prediction is made. Earlier layers are not rewritten.
Source-only status. This v4.9.0-r09 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r09 what the source says

The source names four influences on the local factor. For none of them does it give a derived relation, and it gives no field law for $\gamma_L^{(\mathrm{local})}(x,t)$.

influence	place	content
observer	Theorem 20.5	every matter observer carries $z^- \neq 0$; measured constants are frame-invariant values times a Lorentz factor of the observer
cosmic time	Proposition 42.5	β_{sym} grows "as matter structures in the universe grew"
Earth's orbit	corollary "α-drift as a frame effect"	the factor varies "as Earth moves through the Sphaera-Cascade geometry over its orbit", with a period of one year, absent for a clock that does not co-orbit
potential	prediction P5 near Corollary 42.24	amplitude of the α modulation $\sim 10^{-10}\,\delta\Phi/c^2$

Two contradictions.

1. Theorem 20.5 calls γ_{BI} "the Lorentz-transformed γ_{string}, observed from a matter frame". In that theorem γ_{BI} depends on the observer. Theorem 25.3 sets $k(\gamma_{\mathrm{BI}}) = 0$, "frame-invariant by construction".

2. Theorem 20.5 projects all constants, including c, with one Lorentz factor, which means $k = 1$ for all. Theorem 25.3 gives $k(c) = 0$ and $k(G_N) = 2$.

Both are resolved if Theorem 20.5 is read as a statement about the structural factor alone. That is the separation of r18 and r08. What the local factor consists of then remains open.

v4.9.0-r09 three readings tested

With the rule $\delta \ln X = k(X)\,\delta \ln \gamma_L^{(\mathrm{local})}$, three readings were tested.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Theoremlet v4.9.0-r09.1.

(V) *Velocity against a preferred frame.* If the observer's rapidity adds to η, Earth's orbital motion projected on the dipole of the microwave background gives $\delta\eta \approx 9.8 \times 10^{-5}$. That yields $\delta \ln \gamma_L = 2 \tanh 2\eta \, \delta\eta \approx 1.6 \times 10^{-4}$, and with $k(\alpha) = 1$ an annual α modulation of the same size. The clock fit quoted in r03 (F2) allows about 2×10^{-17}. The reading is **excluded**, by a factor of about 10^{13}. The static part, 370 km/s against the background, would make α depend on direction at the level of 2×10^{-3}.

(P) *Gravitational potential.* $\delta \ln \gamma_L^{(\mathrm{local})} = \kappa \, \delta\Phi/c^2$. The source's own coefficient $\kappa \sim 10^{-10}$ and the eccentricity of Earth's orbit ($\delta\Phi/c^2 \approx 1.6 \times 10^{-10}$) give an annual $\delta\alpha/\alpha \approx 1.6 \times 10^{-20}$, inside the bounds. The reading **survives**. It cannot drive the cosmological drift of Proposition 42.5: potentials of large-scale structure, $\sim 10^{-5}$, give $\delta \ln \gamma_L \sim 10^{-15}$, against about 5×10^{-2} needed.

(D) *Matter density.* The source gives no functional form. Density differs between laboratory, space and astrophysical environments by many orders of magnitude, so any appreciable coupling would have to be screened. The reading is **not testable** as stated.

Proof sketch. (V) $\mathrm{d} \ln \cosh 2\eta / \mathrm{d}\eta = 2 \tanh 2\eta$, with $29.8 \cos 11^\circ$ km/s against c. (P) $GM_\odot/(c^2 a) = 9.87 \times 10^{-9}$ times the eccentricity 0.0167. Checked numerically. □

Consequence. Of the three readings only the potential reading is compatible with the clocks. It is also the one the source itself states in P5. It describes a small violation of local position invariance and cannot explain the Hubble tension. Within it, every constant couples to the potential with $\beta_X = k(X) \, \kappa$:

- the Compton clock of r08.2, with $k = 0$, does not couple;
- α couples with κ;
- an atomic clock and G_N, with $k = 2$, couple with 2κ.

The ratio of the couplings, not their size, is the content of the source. The size κ is not derived.

v4.9.0-r09 the arithmetic of P5

P5 states $\frac{\delta\alpha/\alpha}{\delta G_N/G_N}\big|_{\mathrm{annual}} = \gamma_L^{k(\alpha)-k(G_N)} \cdot \frac{k(\alpha)}{k(G_N)} = \gamma_L^{-1} \approx 0.537$.

Theoremlet v4.9.0-r09.2. From $X^{(\mathrm{matter})} = \gamma_L^k X^{(\mathrm{photon})}$ with fixed photon-frame values follows $\delta X/X = k(X) \, \delta\gamma_L/\gamma_L$. Hence

$$\frac{\delta\alpha/\alpha}{\delta G_N/G_N} = \frac{k(\alpha)}{k(G_N)} = \frac{1}{2}.$$

The static factor $\gamma_L^{k(\alpha)-k(G_N)}$ does not belong in a ratio of relative changes. The middle expression of P5 equals $\gamma_L^{-1}/2 = 0.268$, not γ_L^{-1}. The corrected value 0.5 is the same in every reading of the local factor.

Proof sketch. Logarithmic differentiation. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r09 register entries

statement	status after v4.9.0-r09
local factor $\gamma_L^{(\mathrm{local})}$	four influences named (observer, time, orbit, potential), no field law; only the potential reading survives the clocks (r09.1)
Thm. 20.5 vs Thm. 25.3	two contradictions (γ_{BI} observer-dependent vs $k = 0$; one factor for all vs $k(c) = 0$, $k(G_N) = 2$); resolved by reading Thm. 20.5 as structural
P5, annual ratio	corrected to $k(\alpha)/k(G_N) = 1/2$; the stated $\gamma_L^{-1} = 0.537$ is wrong twice (r09.2)
potential coupling	$\beta_X = k(X)\kappa$; Compton clock uncoupled; $\kappa \sim 10^{-10}$ not derived, below present bounds
Prop. 42.5	the surviving reading gives 10^{-15} instead of 5×10^{-2}; unchanged: hypothesis without working mechanism

v4.9.0-r09 ledger and guard update

$$R_{\mathrm{loc}}^{v490\text{-}r09} = (0, 0, 1)$$

for (influences and contradictions recorded, readings of the local factor decided against clocks, size of κ open).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r09,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r09} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r10 Source-Only: A Reading of the C5 Factor 3/4 and a Class of Shape-(iii) Witnesses

Purpose. r94 found that the C5 contraction factor 3/4 is declared, not derived, and that a single projection explains at most one step. r88 left shape (iii), a witness outside the collar iteration, as the only open route to a finite C5 discharge. This layer offers a reading of the factor built from objects the source already has, and uses the same reading to exclude a class of shape-(iii) witnesses. The reading is an interpretation, admitted as such by the author (2026-10-22), and is not a derivation. Files: `llm/Erkundungsprotokoll_Kern.md` and the scripts `llm/k50_c5_faktor.py`, `llm/k51_c5_form_iii.py`. Earlier layers are not rewritten. This layer does not assert or approach an exact finite C5 zero closure.
Source-only status. This v4.9.0-r10 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r10 rotate, then compensate

On the doublet of r04, with $C_{\mathrm{HC}} = \frac{1}{2}(1+\sigma_x)$, $A = \frac{1}{2}(1-\sigma_x)$ and the duality rotation R_α, write u for the unit vector spanning $\operatorname{ran} C_{\mathrm{HC}}$.

Theoremlet v4.9.0-r10.1. $C_{\mathrm{HC}} R_\alpha C_{\mathrm{HC}} = \cos\alpha\, C_{\mathrm{HC}}$. One step "rotate by α, then compensate" keeps the fraction $\cos^2\alpha$ of the energy in the compensator port and sends $\sin^2\alpha$ to the partner port. Because the rotation lies between the projections, the step is not idempotent: every repetition removes the same fraction again. For $\alpha = \pi/6$:

$$\cos^2\tfrac{\pi}{6} = \tfrac{3}{4}, \qquad \left(\tfrac{3}{4}\right)^n : \ \tfrac{3}{4},\ \tfrac{9}{16},\ \tfrac{27}{64},\ \tfrac{81}{256},\ \ldots,$$

which are the ledger values of r04, r05, r07 and r08. The first stage with $(3/4)^n \le 1/256$ is $n = 20$.

Proof sketch. $\langle u, R_\alpha u\rangle = \cos\alpha$ in two dimensions. The stage was checked in exact rational arithmetic. □

Reading. The step meets the condition r94 set for any derivation: a refreshed one-quarter identity at every stage. Two sources for $\pi/6$ are available in the source.

- *Triolic.* The quarter turn from matter to tachyon ($\pi/2$, Theoremlet r07.2) is divided into three equal steps. Then 27/64, the r07 probe entry, is one complete passage from matter to tachyon. This reading needs no value of β_{sym}.
- *Hopf.* $\tan\alpha = 1/\sqrt{3}$, the Hopf value of β_{sym} in Theorem 22.16, so $q = 1/(1+\beta_{\mathrm{sym}}^2) = 3/4$. With the measured $\beta_{\mathrm{sym}} = 0.5493$ the same formula gives $q = 0.7682$ and stage 22, outside the interval of Theoremlet r94.2. This reading holds only at the Hopf value.

A Zeno pattern. A single quarter turn followed by compensation empties the compensator port, $C_{\mathrm{HC}} R_{\pi/2} u = 0$. Three compensated steps of $\pi/6$ leave 27/64, six leave 0.66, three hundred leave 0.99. The more often the content is compensated, the less of it leaves. This is the pattern of the quantum Zeno effect.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Observations, not interpretations.

1. The relay couplings of the r14 chart family take the values $0, \frac{1}{4}, \frac{1}{2}, \frac{3}{4}, 1$. These are $\sin^2$ of $0, \pi/6, \pi/4, \pi/3, \pi/2$. A uniform grid of quarters has this property anyway, so it carries little weight.

2. Chart $V3$, whose relay coupling $\kappa = 1/4$ produces the five in $\Delta_\gamma < 1/5$ (r92), and the clock seed, $(N, \kappa, G) = (5, \frac{3}{4}, 1)$, have couplings that are the two port fractions of one $\pi/6$ step. In $V3$, $\kappa(N-1) = 1$ doubles the overlap flux, $S = 2N = 10$, and gives $G^{\mathrm{req}}/G^{\mathrm{avail}} = 4/5$.

3. The tolerated collar row of r05 has ratio $5/6 = 1/(1 + \frac{1}{5})$, the form of the Hopf reading with $1/5$ in place of $1/3$.

None of these is used below.

v4.9.0-r10 a class of shape-(iii) witnesses

r88 requires of a shape-(iii) witness that it exhibit the vanishing of the obstruction itself, not of the score (α), and that it not import the all-readout slot (δ). Under the reading above, the C5 content is the field content of the doublet, and the score is its energy in the compensator port. The transports available in Reading B are spatial rotations, duality rotations and boosts.

Theoremlet v4.9.0-r10.2 (conditional on the reading).

(a) *Isometric witnesses* (spatial and duality rotations) conserve $\|FC_{\mathrm{HC}}\|^2 + \|FA\|^2 = \|F\|^2$. A witness of this kind that empties the compensator port puts the whole content into the partner port. It satisfies (α) only by importing the partner readout, which is (δ).

(b) *Boosts* change the total $E^2 + B^2$, but $E^2 + B^2 \geq |s|$ for every boost, with equality in the frame where $E \parallel B$. For a non-null field the content therefore has a positive floor that no boost removes.

(c) For a null field the total falls like $e^{-2\eta}$ under a boost along the propagation, and reaches zero only in the limit $\eta \to \infty$, never at a finite stage.

Hence no witness built from the transports of Reading B meets (α) and (δ) at a finite stage. The obstruction behaves like the invariant s, which no group action of Reading B changes (Theoremlet r19.1).

Proof sketch. (a) $C_{\mathrm{HC}} + A = 1$, $C_{\mathrm{HC}} A = 0$, both symmetric. (b) $u^2 - |E \times B|^2 = \frac{1}{4}|s|^2$ (r03.3) gives $2u \geq |s|$; checked over 16200 boosts. (c) Doppler factor of a plane wave. □

Consequence. Under this reading, a shape-(iii) witness would have to lie outside the transport group of Reading B. It would have to change the invariant s, that is, carry dynamics. Reading B has no dynamics (r10 of v4.8.0). The route therefore stays open in the sense of r88, and the class of witnesses examined here is closed. The layer does not assert, and does not approach, an exact finite C5 zero closure. It records why the transports at hand cannot supply one.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r10 register entries

statement	status after v4.9.0-r10
C5 factor 3/4 (r94)	reading offered: rotate by $\pi/6$, then compensate; reproduces $3/4, 9/16, 27/64, 81/256$ and $n = 20$; interpretation, not derivation (r10.1)
source of $\pi/6$	triolic division of the matter–tachyon quarter turn, or Hopf value $1/\sqrt{3}$ (only at that value)
shape (iii) (r88)	under the reading: isometric witnesses relocate, boosts meet the floor $\lvert s\rvert$, null fields reach zero only as a limit; witnesses would need dynamics (r10.2)
$\Delta_\gamma < 1/5$ (r92)	observation: $V3$ coupling $1/4$ and clock-seed coupling $3/4$ are the two port fractions of one $\pi/6$ step; not used

v4.9.0-r10 ledger and guard update

$$R_{\text{core}}^{v490\text{-}r10} = (0, 0, 1)$$

for (a reading of the factor offered, a class of shape-(iii) witnesses closed under it, a derivation of the reading open).

The layer spends no guard:

$$R_{\text{glob}}^{(v490\text{-}r10,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\text{guard}}^{v490\text{-}r10} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r11 Source-Only: A Minimal Dynamics for Reading B — Kicked Duality Rotation

Purpose. r10 found that the open threads of the core meet at one gap: Reading B has no dynamics (r10 of v4.8.0). The C5 factor 3/4 would be derived if a dynamics produced the alternation "rotate, then compensate" of r10.1. A shape-(iii) witness would need a dynamics that changes the invariant s (r10.2). This layer proposes the smallest dynamics with these properties and checks what it does. It is a model proposal, not a derivation from the source. The script is `llm/k52_kick_dynamik.py`. No value is compared with data. Earlier layers are not rewritten. This layer does not assert or approach an exact finite C5 zero closure.
Source-only status. This v4.9.0-r11 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r11 the proposal

Definition (kicked duality dynamics, proposal). Between kicks the field turns in the duality circle at a constant rate, $z(t) = e^{i\omega t} z$, on the doublet $F \mapsto F R_{\omega t}^{\mathsf{T}}$. At the times $t_n = n\tau$ the compensator C_{HC} is applied, and the content of the partner port is handed to a reservoir:

$$F_{n+1} = F_n R_{\omega\tau}^{\mathsf{T}} C_{\mathrm{HC}}, \qquad \mathcal{E}_{\mathrm{res}}^{(n+1)} = \mathcal{E}_{\mathrm{res}}^{(n)} + \| F_n R_{\omega\tau}^{\mathsf{T}} A \|^2 .$$

The rotation is the transport of Reading B, the kick is the readout of the main body, and the reservoir is the partner sector $\operatorname{ran} A$ of r03.1.

v4.9.0-r11 what it does

Theoremlet v4.9.0-r11.1.

(a) Each stage keeps $\cos^2 \omega\tau$ of the content and exports $\sin^2 \omega\tau$. Retained plus exported content is conserved.

(b) The stage factor equals 3/4 for exactly one $\omega\tau \in (0, \pi/2)$, namely $\omega\tau = \pi/6$. Then the retained content runs through $3/4, 9/16, 27/64, 81/256, \ldots$ and first falls below 1/256 at stage 20.

(c) In the state disk of r04 a stage is a rotation of the Bloch vector by $2\omega\tau = 60°$ followed by a measurement along the J-axis, kept with probability $(1 + \cos 60°)/2 = 3/4$.

(d) The duality phase accumulates in steps of 30°. Three stages make the quarter turn from matter to tachyon, six the half turn $z \mapsto -z$ (the charge conjugation of r02.2), twelve the full turn, twenty-four the return of the spinor of r06.4.

(e) The rotation leaves $|s|$ unchanged. The kick changes it: it sets $\operatorname{Re} s = 0$ and lowers $|s|$ (in the checked example from 2.15 to 0.83). The dynamics therefore leaves the transport group of Reading B, as r10.2 requires of any witness that changes s.

(f) Dividing the same phase $\pi/6$ into m kicks keeps $\cos^{2m}(\pi/6m)$: 0.75, 0.87, 0.95, 0.99 for $m = 1, 2, 5, 50$. The stage factor is set by how often the readout acts per phase, which is the quantum Zeno pattern.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof sketch. (a)–(b) $C_{\mathrm{HC}} R_\alpha C_{\mathrm{HC}} = \cos\alpha\, C_{\mathrm{HC}}$ (r10.1) and $C_{\mathrm{HC}} + A = 1$. (c) $G \mapsto RGR^{\mathsf{T}}$ turns the Bloch vector by twice the angle (r04.2). (e) direct computation. All items were checked numerically, (b) also in exact rational arithmetic. □

v4.9.0-r11 what it does not do

1. *No finite closure.* The retained content after n stages is $(3/4)^n > 0$ for every n. It vanishes in one stage only for $\omega\tau = \pi/2$, and then the whole content sits in the reservoir: it is exported, not annihilated. The dynamics reproduces $C5_{\mathrm{det/eff}}$ at $\theta = 1/256$, $n = 20$, and keeps $0_{\mathrm{termdata}} \not\Rightarrow 0_{\mathrm{all}}$. The all-readout slot includes the reservoir.

2. *No scale.* Only the product $\omega\tau$ is fixed. The source supplies neither a rate ω nor an interval τ. The oscillator frequency of Theorem 22.11, $\gamma_{\mathrm{string}}\sqrt{2/3}$, is a dimensionless candidate for ω in the source's units. Without a unit of time it fixes nothing.

3. *No mechanism for the kick.* Why the readout acts at regular intervals, and three times per quarter turn, is not explained. The triolic division of r10 is assumed, not derived.

4. *Not the field equations.* The proposal acts on the doublet at a point. It is not a field theory and makes no claim about the Maxwell or Einstein limits of the main body.

Remark. The structure, a fixed polarisation rotation followed by a polariser at every cycle, is the one used in quantum Zeno experiments with photons, among them the interaction-free measurements of Kwiat and co-workers (1995). There a rotation of $\pi/(2N)$ per cycle with N cycles per quarter turn leaves $\cos^{2N}(\pi/2N)$. The proposal of this layer is that structure with $N = 3$. The correspondence concerns the algebra only. On a light wave the compensator of r03 is not a polariser (r03.2).

v4.9.0-r11 register entries

statement	status after v4.9.0-r11
dynamics of Reading B (r10 of v4.8.0)	proposal: kicked duality rotation with compensator kicks and partner reservoir; model decision, not derived
C5 factor 3/4	produced by the proposal at $\omega\tau = \pi/6$, uniquely in $(0, \pi/2)$ (r11.1)
shape (iii)	the kick changes s, but the dynamics exports instead of annihilating; no finite closure (r11)
scale of the dynamics	open: only $\omega\tau$ fixed

v4.9.0-r11 ledger and guard update

$$R_{\mathrm{dyn}}^{v490\text{-}r11} = (0, 0, 1)$$

for (a minimal dynamics proposed, its stage factor and cycles established, its scale and kick mechanism open).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r11,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r11} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r12 Source-Only: A Reconstruction of the Bound $\Delta_\gamma < 1/5$ in Place of the Missing r65

Purpose. The bound $\Delta_\gamma < 1/5$ carried by $W_{\mathrm{clk}}^{\mathrm{loc}}$ comes from r65, whose full text is not available. The author confirmed on 2026-10-22 that it is lost and asked for a reconstruction. r90 obtained a bound of the right size (2/9). r92 obtained 1/5 exactly as a relative gain reserve, without stating why that reading and that chain apply. This layer states the premises explicitly, derives the strict bound from them, and draws the consequences under the dynamics of r11. It replaces the missing derivation. It does not recover the r65 text and does not claim that r65 argued this way. The script is `llm/k53_r65_rekonstruktion.py`. Earlier layers are not rewritten.
Source-only status. This v4.9.0-r12 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r12 premises

(P1) *Chart data* of r14 (§A.225.2): overlap flux $S = N + \kappa N(N-1)$, relay score $H = GS$, relay threshold 16.

(P2) *Survival* of a window is the strict relay condition of Definition A.234.1, applied to the perturbed gain: $G(1-\Delta_\gamma)\,S > 16$.

(P3) *The perturbation is a multiplicative loss.* Δ_γ is the fraction of the clock content lost per readout. Under the dynamics of r11 this is the fraction exported to the partner reservoir, $\sin^2\alpha$ for a duality drift α between readouts.

(P4) *Scope.* The guard concerns the charts that carry clock windows (unassisted score below 16, assisted score at least 16) on the primary relay chain $\mathcal{R}_A$ of Section A.226, that is $V3$ and $V5$.

(P1) and (P2) are in the source. (P3) chooses r92's reading over r90's, and gives it a mechanism. (P4) is r92's restriction, consistent with the local index of $W_{\mathrm{clk}}^{\mathrm{loc}}$.

v4.9.0-r12 the bound

Theoremlet v4.9.0-r12.1. Under (P1)–(P4), $W_{\mathrm{clk}}^{\mathrm{loc}}$ is preserved if and only if

$$\Delta_\gamma < \min_{V\in\{V3,V5\}}\Big(1-\frac{16}{G_V S_V}\Big) = \min\Big\{\frac{1}{5},\frac{9}{25}\Big\} = \frac{1}{5}.$$

The inequality is strict. At $\Delta_\gamma = 1/5$ chart $V3$ has $G(1-\Delta_\gamma)S = 16$ exactly, which the strict window excludes.

Proof sketch. (P2) rearranges to $\Delta_\gamma < 1 - 16/(GS)$. $V3$: $N=5$, $\kappa=1/4$, $G=2$, $S=10$, $H=20$, hence 1/5. $V5$: $N=4$, $\kappa=1/2$, $G=5/2$, $S=10$, $H=25$, hence 9/25. Checked in exact rational arithmetic. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Where the five comes from. $V3$ is the only chart of the r14 family with $\kappa(N-1)=1$: its pair-overlap flux $\kappa N(N-1)$ equals its direct flux N, so $S=2N$. With $G=2$ the bound becomes $1-16/(4N)=1-4/N$, which is $1/5$ at $N=5$. The binding chart is the balanced chart of the primary chain.

v4.9.0-r12 consequences under the dynamics of r11

Theoremlet v4.9.0-r12.2 (conditional on (P3)).

(a) The clock selector survives a duality drift α per readout exactly when $\sin^2\alpha < 1/5$, that is $\tan\alpha < 1/2$, $\alpha < 26.57°$.

(b) The C5 stage of r10 and r11 ($\alpha=\pi/6$) exports $1/4 > 1/5$ and lies outside. The tolerated collar row of r05 (ratio 5/6) exports $1/6 < 1/5$ and lies inside.

(c) The rows of r05 that reach $\Phi_5=0$ in one step, $Z_5^{\text{mix-cancel}}$ and $Z_5^{\text{all-cancel}}$, move the whole score $-3/4$ into A_4 or U_6 and are blocked by $C4$ and $C6$. This is the quarter turn of r10.2 in ledger form: the compensator port is emptied by relocating the content to another slot.

Proof sketch. (a) (P3) in Theoremlet r12.1. (b) $1-3/4$ and $1-5/6$. (c) read off Definition A.216.4. □

Reading. The strict C5 lowering and the clock selector are separated by the bound: a C5 stage drifts further per readout than the clock window tolerates. This is consistent with the two being distinct guard items, $C5_{\text{det/eff}}$ and $W_{\text{clk}}^{\text{loc}}$, established on different charts. The r05 collar row, which r05 calls "useful as a stability check", is the one whose drift the clock tolerates. That r05 classified it as tolerated for this reason is an interpretation.

v4.9.0-r12 status of the bound

layer	contribution
r87	no derivation present; not reconstructible from the window layer alone
r90	coherence reading, bound 2/9, 1/5 as rounding
r92	gain-reserve reading, 1/5 exactly on $\mathcal{R}_A$, premises implicit
v4.9.0-r12	premises (P1)–(P4) stated, strictness from r23, multiplicative loss from r11, balanced chart $V3$ identified

The provenance component of R_{cal}^{r87} stays open, since the r65 text itself is not recovered. The derivation component is closed under (P1)–(P4).

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r12 register entries

statement	status after v4.9.0-r12
$\Delta_\gamma < 1/5$ in $W_{\mathrm{clk}}^{\mathrm{loc}}$	derived under (P1)–(P4), strict (r12.1); r65 text lost (author), provenance open
binding chart	$V3$, the balanced chart $\kappa(N-1)=1$ of $\mathcal{R}_A$
clock vs C5 stage	tolerated drift $\tan\alpha < 1/2$; C5 stage outside, r05 collar row inside (r12.2)
r05 zero rows	relocation into A_4, U_6, blocked by $C4$, $C6$; same structure as r10.2

v4.9.0-r12 ledger and guard update

$$R_{\mathrm{cal}}^{v490\text{-}r12} = (0,0,1)$$

for (derivation under stated premises closed, consequences under r11 recorded, r65 provenance open).

The layer spends no guard. $W_{\mathrm{clk}}^{\mathrm{loc}}$ under $\Delta_\gamma < 1/5$ remains the local, permitted statement:

$$R_{\mathrm{glob}}^{(v490\text{-}r12,+)} = (0,1,1,1,1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r12} = (1,1,1,1,1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r13 Source-Only: The Time Scale of the Kicked Duality Dynamics

Purpose. The dynamics proposed in r11 fixes only the product $\omega\tau = \pi/6$. This layer derives the time scale from the clock model the source already contains (Section A.226). It follows the source in reading the photon-clock probe as a diagnostic timing function, not as an identity between Riemann zeros and photons. All statements are local, in the clock's own units, and make no global photon-clock statement. The script is `llm/k54_zeitskala.py`. Earlier layers are not rewritten.
Source-only status. This v4.9.0-r13 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r13 the rate from the tick profile

r15 writes the local clock tick as $G_V(\tau) = 1 + (G_V^{\text{avail}} - 1)\cos^2\big((\tau - \tau_V)/\sigma_V\big)$. Between two kicks of r11 the energy in the compensator port is $\cos^2(\omega t)$. Both are $\cos^2$ of an angle linear in time.

Theoremlet v4.9.0-r13.1.

(a) Identifying the two profiles gives $\omega = 1/\sigma_V$: the tick width of r15 is the time the duality rotation of r11 needs for one radian. The kick interval is $\tau_{\text{k}} = \frac{\pi}{6}\,\sigma_V$.

(b) The smallest probe with a profile has two terms. For $\mathcal{C}_\gamma \propto |e^{i\gamma_a\tau} + e^{i\gamma_b\tau}|^2$ the coherent part is exactly $\cos^2\big(\frac{1}{2}(\gamma_b - \gamma_a)\tau\big)$, so $\sigma = 2/(\gamma_b - \gamma_a)$. The beat period is $T_{\text{b}} = 2\pi/(\gamma_b - \gamma_a)$.

(c) Hence $\tau_{\text{k}} = T_{\text{b}}/6$ for every pair of clock frequencies. Three stages, the quarter turn from matter to tachyon, take half a beat. Six stages, the half turn $z \mapsto -z$, take one beat. Twelve take two beats, and twenty-four, the return of the spinor, take four. The detector stage $n = 20$ lies at $10/3$ beats.

(d) For the two lowest frequencies the source uses, $\gamma_1 = 14.1347\ldots$ and $\gamma_2 = 21.0220\ldots$: $\sigma = 0.29039$, $T_{\text{b}} = 0.91228$, $\tau_{\text{k}} = 0.15205$ in clock units.

Proof sketch. (b) $|e^{ia} + e^{ib}|^2 = 4\cos^2\frac{b-a}{2}$; checked numerically to 10^{-15}. (c) $\tau_{\text{k}} = \frac{\pi}{6}\cdot\frac{2}{\Delta\gamma} = \frac{1}{6}\cdot\frac{2\pi}{\Delta\gamma}$. □

What is fixed and what is not. The relative statement (c) holds for any pair of frequencies. The absolute value (d) depends on which pair carries the tick, and the source names none. The first two are the minimal choice. The clock unit itself has no conversion to seconds in the source. Supplying one would require identifying clock frequencies with physical frequencies. The source expressly declines that identification, and as a global statement it would touch the forbidden global photon-clock statement. The time scale is therefore derived up to the unit of the clock.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r13 the windows of the primary chain

Theoremlet v4.9.0-r13.2. In the angle of r13.1(a), the half-widths δ_V/σ_V of r15 are 39.23° for $V3$, 50.77° for $V5$ and 28.13° for $V8$. A timing offset of one C5 stage, 30°, stays inside every window of the primary chain $\mathcal{R}_A = \{V3, V5\}$ and leaves the window of $V8$ on the chain $\mathcal{R}_B$.

Proof sketch. Conversion of the r15 table. □

Reading. This is a second, independent reason for the restriction (P4) of r12 to $\mathcal{R}_A$: only there does a single C5 stage fit into the clock window. The tolerance 26.57° of r12 concerns the loss of clock content per readout, a different quantity from a timing offset within one window, so the two numbers do not contradict each other. The window of $V8$ lies between them.

v4.9.0-r13 the stage angle between Hopf and lattice

The main body relates the Hopf value and the measured value of the offset by $\beta_{\mathrm{sym}} = \frac{1}{\sqrt{3}}\exp\bigl(-\gamma_1/(4\pi\gamma_L^5)\bigr) = 0.5492$ (the table with "Hopf (continuum S^3)" and "Physical (prime lattice)" rows). The stage angle $\arctan\beta_{\mathrm{sym}}$ of r10 is therefore 30° in the continuum limit $\gamma_1 \to 0$ and 28.78° on the lattice, with $q = 0.7683$. The C5 factor 3/4 is the continuum value. On the lattice the Hopf reading would give stage 22 (r10), while the triolic reading keeps $\pi/6$. Which of the two describes the C5 recursion is open. The detector stage $n = 20$ favours the continuum value or the triolic reading.

v4.9.0-r13 register entries

statement	status after v4.9.0-r13
time scale of r11	$\omega = 1/\sigma_V$, $\tau_{\mathrm{k}} = T_{\mathrm{b}}/6$ (one sixth of the clock beat) for every frequency pair (r13.1)
absolute value	$\tau_{\mathrm{k}} = 0.15205$ clock units for the two lowest frequencies; no conversion to seconds, by the source's own restriction
r12 premise (P4)	second reason: one C5 stage fits only into the windows of $\mathcal{R}_A$ (r13.2)
C5 factor 3/4	continuum value of the Hopf reading; lattice value 0.7683; triolic reading unaffected

v4.9.0-r13 ledger and guard update

$$R_{\mathrm{time}}^{v490\text{-}r13} = (0, 0, 1)$$

for (relative time scale derived, windows of the primary chain checked, unit of the clock open).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r13,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r13} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r14 Source-Only: Why Six Kicks per Beat — Readout Directions from the Triole and the Umkehrwalze

Purpose. r13 fixed the kick interval of the dynamics of r11 at one sixth of the clock beat. It left open why the compensator acts six times per beat. This layer offers a reason built from two objects of the source: the triole and the involutive Umkehrwalze J_B of r02.3. The reason is an interpretation (admitted by the author, 2026-10-22), not a derivation. The script is `llm/k55_kick_richtungen.py`. Earlier layers are not rewritten.
Source-only status. This v4.9.0-r14 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r14 kicks as passages of readout directions

Replace the periodic kick of r11 by an event: the compensator acts whenever the phase of the invariant s passes one of a fixed set of readout directions. Under a uniform duality rotation the events are periodic, and their spacing is set by the set of directions.

Theoremlet v4.9.0-r14.1. In the s-plane, $\arg s = 2\alpha$ with α the duality angle.

(a) The triole directions $\arg s \in \{0°, 120°, 240°\}$ and their images under J_B, which acts as $s \mapsto -\bar{s}$ (r02.3), are together the six directions $0°, 60°, \ldots, 300°$. The set contains matter ($0°$) and tachyon ($180°$).

(b) Passing them under a uniform rotation gives kicks every $30°$ of duality angle, hence the stage factor $\cos^2 30° = 3/4$. The triole alone gives $60°$ and $q = 1/4$. A quarter grid gives $45°$ and $q = 1/2$. Only triole and J_B together give the factor of the source.

(c) One revolution of s around the photon ($\arg s$ from $0°$ to $360°$) passes the six directions once. It is the duality half turn $z \mapsto -z$, the charge conjugation of r02.2, and by r13 it lasts exactly one beat.

Proof sketch. $-\overline{e^{i\theta}} = e^{i(\pi-\theta)}$ maps $\{0, 120, 240\}$ to $\{180, 60, 300\}$. The rest follows from $\arg s = 2\alpha$ and r10.1. Checked numerically. □

Reading. The six kicks per beat are then not an external schedule but the six readout directions met by one revolution of s around the photon. The factor 3/4, the detector stage 20, the beat, the quarter turn from matter to tachyon and the charge conjugation of the round trip become aspects of one construction: a uniform duality rotation read out at triole$\cup J_B$(triole).

What is assumed. The triole of the source (Theoremlet r06.2) is a condition on the three components of a vector, $(1, \omega, \omega^2)$, not on the phase of s. Placing the triole on $\arg s$ is the interpretive step of this layer. Squaring maps the triole of phases onto a triole of phases, so the placement is consistent, but the source does not make it. Why the readout should act at discrete directions at all is also assumed.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Remark, offered and not adopted. Six directions at 60° with the triole permutations as symmetry are the root system A_2 of $SU(3)$, whose Weyl group is S_3. r15 of v4.8.0 met $SU(3)$ as the stabiliser of the symmetric point, there read as colour. Whether the readout hexagon and the colour $SU(3)$ are related would need a map from the duality plane to a Cartan plane, which the source does not supply.

v4.9.0-r14 register entries

statement	status after v4.9.0-r14
six kicks per beat (r13)	reading offered: readout at triole $\cup\, J_B$(triole) in the s-plane (r14.1)
factor 3/4	needs both triole and J_B; either alone, or a quarter grid, gives another factor
beat	one revolution of s around the photon, equal to the charge conjugation of r02.2
A_2/colour	remark only

v4.9.0-r14 ledger and guard update

$$R_{\text{kick}}^{v490\text{-}r14} = (0, 0, 1)$$

for (a reason for the kick schedule offered, its consequences consistent with r10–r13, the placement of the triole open).

The layer spends no guard:

$$R_{\text{glob}}^{(v490\text{-}r14,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\text{guard}}^{v490\text{-}r14} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r15 Source-Only: Consolidated Register of v4.9.0-r01 to r14 and the Ways Out of r17 after r08

Purpose. Fourteen layers since the 4.9.0 release have changed the status of several statements of the main body and added readings of their own. This layer collects the statuses in one register, in the manner of §A.271, and lists the assumptions that only the author can confirm. It also draws one consequence that r08 makes available: r17 left three ways out for the cosmological fractions, and r08 decides between them. No new computation and no new prediction enter. Earlier layers are not rewritten.
Source-only status. This v4.9.0-r15 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r15 the ways out of r17 after r08

r17 found that Theorem 22.16, held at all epochs, needs β_{sym} to decrease from about 1 at recombination to 0.558 today. Propositions 42.5 and 42.8 assume the opposite. It recorded three ways out: (a) the theorem holds only at the present epoch; (b) β_{sym} runs downward; (c) the cascade probabilities are some other function of the fractions at other epochs.

Theoremlet v4.9.0-r15.1. Under the rules of r08 (photon-frame values and $k = 0$ fixed, matter-frame values drifting with the local factor, $k(\alpha) = 1$):

(b) is excluded. A decrease of β_{sym} from about 1 to 0.558 changes the local factor by far more than the 5% that r08.1 already finds incompatible with atomic clocks.

(a) and (c) remain. Both treat β_{sym} as fixed and read Theorem 22.16 at one epoch, (a) directly, (c) through a time-dependent bridge with fixed β_{sym} that the source does not supply.

1. Item (1) of Proposition 42.8 lets the neutrino mass grow with $\gamma_L(t)^4$. By the clock bound on the same local factor this growth is excluded too. Together with item (2), which r17 already rejected, the proposition loses its basis.

Proof sketch. (b): a local factor diverging towards recombination; the clock bound of r08.1 applies a fortiori. (a), (c): neither needs a drift. Item (1): $k(m_\nu) = 4$ drifts faster than α with the same factor. □

Consequence. With r08, Theorem 22.16 is a statement about the present epoch, $1-\beta_{\mathrm{sym}}^2 = 0.698$ against 0.689. Its cosmological reach beyond that epoch is not supported by any statement of the source that survives the clocks.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r15 register

statement	layer	status
conversion law, §25	r01	factor γ_L is matter–tachyon; ways (A)–(C) open; on-shell $\cot\theta_W$ lies at (A) (r07, observation)
Thm. 41.18	r01	equivalent to $\cosh 2\eta = e^{\eta}$
Prop. 21.4 (ii)	r01	frame order does not match the axis
Umkehrwalze	r02	round trip with side change $= C = PT$; involutive form antilinear, $J_B\colon E \leftrightarrow B$
Thm. 21.7, compensator	r02, r03	output $\operatorname{Re} s = 0$, not null; for a charge a dyon
Prop. 16.1	r03	orthogonality holds for the product $z \times J_B z$, not the average
compensator as matrix	r04	$\frac{1}{2}(1+\sigma_x)$ on the doublet; state disk, light at the centre
Millennium matrix	r05	on the doublet only port weighting; reading name “layer-transport matrix”
photon from matter + tachyon	r05	only for crossed orientation
tachyon hit chain	r06	cone 56.68°, simultaneous hits in the matter frame
electroweak mass matrix	r06	port weighting with massless photon port (formal)
H14	r07	refuted as exact relation, undecidable at tree level
H15	r07	pre-registered, $m_W = 80.3615 \pm 0.0018$ GeV, comparison pending
Prop. 42.5	r08, r09	no working mechanism; drift excluded by clocks
Thm. 25.3, α row	r08	$k = 1$ belongs to α (closes r03 F2)
P5	r09	corrected to $k(\alpha)/k(G_N) = 1/2$
local frame factor	r09	only a weak potential coupling survives; no field law
C5 factor 3/4	r10, r11, r14	produced by kicked duality rotation read out at triole $\cup J_B$(triole); interpretation
C5 shape (iii)	r10	witnesses from the transport group of Reading B excluded under the reading
$\Delta_\gamma < 1/5$	r12	derived under premises (P1)–(P4); r65 text lost, provenance open
time scale	r13	kick interval one sixth of the clock beat; unit of the clock open
Thm. 22.16	r15	present-epoch statement; r17 way (b) excluded (r15.1)
Prop. 42.8	r15	no basis left (items (1) and (2))

v4.9.0-r15 assumptions for the author

The following steps are interpretations of these layers. Only the author can say whether they match the original intention.

1. Theorem 20.5 speaks about the fixed structural factor, not the local one (r09). This resolves its two conflicts with Theorem 25.3.

2. The C5 recursion is an alternation of duality rotation and compensation (r10, r11).

© 2026 Marc Brendecke. Preprint — not peer reviewed.

3. The readout acts at discrete directions, and the triole lies on the phase of s (r14).
4. The guard $W_{\mathrm{clk}}^{\mathrm{loc}}$ refers to the windows of the primary chain $\mathcal{R}_A$ (r12, r13).
5. Which of the ways (A)–(C) of r01 the conversion of constants takes.
6. Whether Theorem 22.16 is meant for the present epoch only (way (a)) or through a bridge (way (c)).

v4.9.0-r15 ledger and guard update

$$R_{\mathrm{reg}}^{v490\text{-}r15} = (0, 0, 1)$$

for (register consolidated, r17 way (b) excluded, author decisions listed and open).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r15,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r15} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r16 Source-Only: Author Decisions on the Open Readings of r15

Purpose. r15 listed six readings that only the author can confirm (§A.272). The author answered four of them on 2026-10-22. This layer records the answers and what changes in the register. No computation enters. Earlier layers are not rewritten.
Source-only status. This v4.9.0-r16 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r16 the decisions

no.	question (r15)	author's answer (2026-10-22)
1	Does Theorem 20.5 speak about the fixed structural factor?	**Yes, structural.**
2, 3	Is the C5 recursion an alternation of duality rotation and compensation, read out at the triole and its image under J_B?	**Yes, as originally intended.**
5	Which of the ways (A)–(C) does the conversion of constants take?	left open
6	Is Theorem 22.16 meant for the present epoch only, or through a bridge?	left open
4	Does $W_{\mathrm{clk}}^{\mathrm{loc}}$ refer to the windows of $\mathcal{R}_A$?	not asked; stays a premise of r12

v4.9.0-r16 consequences

1. *Decision 1.* The two conflicts between Theorem 20.5 and Theorem 25.3 recorded in r09 are resolved by the author. The theorem concerns $\gamma_L = \gamma_{\mathrm{BI}}/\gamma_{\mathrm{string}}$ as a fixed structural ratio. It makes no statement about how a measured constant depends on the observer. Any variation runs through the local factor of r08 and r09. Theorem 25.3 stands as written, with the α row read as in r08.

2. *Decisions 2 and 3.* The reading of r10, r11 and r14 is the author's intended reading of the C5 recursion: a uniform duality rotation, compensated whenever the phase of s passes the triole or its J_B-image. Its status changes from "interpretation offered" to "author's reading". It remains a reading, not a derivation from the earlier text. The source never states the recursion in this form, and the placement of the triole on $\arg s$ is not in r06. Theoremlet r10.2 on shape-(iii) witnesses, which was conditional on the reading, now holds under the author's reading.

3. *Decisions 5 and 6* stay open as recorded in r01 and r15.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r16 register entries

statement	status after v4.9.0-r16
Thm. 20.5	structural statement about γ_L (author); conflicts of r09 resolved
C5 recursion (r10, r11, r14)	author's reading: kicked duality rotation read out at triole $\cup J_B$(triole)
C5 shape (iii) (r10.2)	witnesses from the transport group of Reading B excluded under the author's reading
conversion ways (A)–(C)	open (author)
Thm. 22.16, epoch	open between r17 ways (a) and (c) (author)

v4.9.0-r16 ledger and guard update

$$R_{\text{auth}}^{v490\text{-}r16} = (0, 0, 1)$$

for (decisions 1 to 3 recorded, their consequences entered, decisions 5 and 6 open).

The layer spends no guard:

$$R_{\text{glob}}^{(v490\text{-}r16,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\text{guard}}^{v490\text{-}r16} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r17 Source-Only: Entropy between the Frames, the Photon as Information Singularity, and a Scale Dimension

Purpose. The main body works from the photon frame. The author asked, on 2026-10-22, three things. Must several structures coincide for that frame to arise at all? Should the photon be treated as an information singularity? Would a new degree of freedom help, one in which conversions between frames can be parked? This layer answers the second and third questions in the state disk of r04 and records a coincidence that bears on the first. The script is `llm/k56_entropie_skala.py`. No value is compared with data. Earlier layers are not rewritten.
Source-only status. This v4.9.0-r17 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r17 the frame factor in the state disk

In the state disk of r04 a field has the Bloch vector b with $|b| = 1/\cosh 2\eta_f$, where η_f is the rapidity of the observer against the frame in which $E \parallel B$. The state matrix $\rho = \frac{1}{2}(1 + b\cdot\sigma)$ has the eigenvalues $\frac{1}{2}(1 \pm |b|)$.

Theoremlet v4.9.0-r17.1. For an observer offset by η:

$$|b| = \frac{1}{\cosh 2\eta}, \qquad p_\pm = \frac{1}{2}\Big(1 \pm \frac{1}{\cosh 2\eta}\Big), \qquad p_+ = \frac{1}{1+\beta^2}, \qquad p_- = \frac{\beta^2}{1+\beta^2}, \qquad \frac{p_-}{p_+} = \beta^2,$$

with $\beta = \tanh\eta$. At the offset of the source, $\cosh 2\eta = \gamma_L$, the frame factor is the inverse length of the Bloch vector, $\gamma_L = 1/|b|$. At the Hopf value of the source, $\beta = 1/\sqrt{3}$ and $\gamma_L = 2$, the eigenvalues are 3/4 and 1/4. At the measured value they are 0.76818 and 0.23182.

Proof sketch. $\cosh 2\eta = (1+\beta^2)/(1-\beta^2)$, hence $\frac{1}{2}(1 + 1/\cosh 2\eta) = 1/(1+\beta^2)$. Checked numerically. □

Consequence. Three numbers of the earlier layers are one. The C5 factor 3/4 (r10, r11) is the larger eigenvalue of the doublet state at the Hopf offset. The exported quarter is the smaller one. The lattice value 0.7683 of r13 is the larger eigenvalue at the measured offset. The frame factor γ_L is the inverse length of the same Bloch vector. The eigenvalue ratio is β^2_{sym}, the quantity that Theorem 22.16 calls Ω_M. The eigenvalue itself, 0.2318, is not Ω_M.

v4.9.0-r17 entropy as order parameter, and the photon

Theoremlet v4.9.0-r17.2. The entropy $S = -p_+\log_2 p_+ - p_-\log_2 p_-$ of the doublet state increases monotonically from 0 in the frame with $E \parallel B$ to 1 bit as $\eta_f \to \infty$. It is 0.811 bit at the Hopf offset and 0.781 bit at the measured offset. The photon, $b = 0$, is the maximally mixed state. In the Bures metric of the disk, $\mathrm{d}s^2 = \frac{1}{4}\big(\mathrm{d}r^2/(1-r^2) + r^2\mathrm{d}\theta^2\big)$, the radial distance from the state with $E \parallel B$ to the state at offset η is $\frac{1}{2}\,\mathrm{gd}(2\eta)$. The photon lies at the finite distance $\pi/4$, and the remaining distance is $\frac{1}{2}\Pi(2\eta)$, half the angle of parallelism of r07 at twice the rapidity.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof sketch. $r = \operatorname{sech} 2\eta$ gives $\mathrm{d}r^2/(1-r^2) = 4\operatorname{sech}^2 2\eta\,\mathrm{d}\eta^2$, hence $\mathrm{d}s = \operatorname{sech} 2\eta\,\mathrm{d}\eta$ and $\int_0^\eta \operatorname{sech} 2x\,\mathrm{d}x = \frac{1}{2}\operatorname{gd}(2\eta)$, with $\operatorname{gd}(\infty) = \pi/2$ and $\Pi = \pi/2 - \operatorname{gd}$. Checked numerically. □

The photon as information singularity, made precise. The metric itself is regular at the photon. What becomes singular is the direction: $r^2\,\mathrm{d}\theta^2 \to 0$, so the angle θ, which carries the distinction between matter and tachyon (b_1) and the side of r02 (b_2), is undefined at $b = 0$. This is the polar-coordinate form of the branch point of r19.2. In this sense the photon is an information singularity. It is the state of maximal entropy, in which the information that distinguishes the frames is absent. It is reached by no finite boost (Theoremlet r19.1), only in the limit, and at a finite information distance.

On the author's first question. The photon is the point where the two eigenvalues coincide, $p_+ = p_- = \frac{1}{2}$, and where matter and tachyon lose their distinction. In the state disk the special frame is therefore the frame in which the structures that separate the other frames fall together. This supports the author's conjecture. It also shows that the photon frame is not a frame like the others: it is the limit point and reference of the disk. A finite factor γ_L "to the photon frame", as in §25, corresponds to the state with $|b| = 1/\gamma_L$, not to $b = 0$.

v4.9.0-r17 a scale dimension for the conversions

The rule $X^{(\text{matter})} = \gamma_L^k X^{(\text{photon})}$ is a scaling with weights k. Carrying the scale as a coordinate σ of its own parks the conversion there, $X(\sigma) = e^{k\sigma}X(0)$, with the photon frame at $\sigma = 0$.

Theoremlet v4.9.0-r17.3.

(a) σ is a coordinate in which conversions compose as shifts only if the factor is a homomorphism of the rapidity: $\sigma = \eta$, $X = e^{k\eta}X(0)$, the Doppler law of way (C) in r01. The Lorentz factor is not one, $\cosh\eta_1 \cosh\eta_2 \neq \cosh(\eta_1 + \eta_2)$. Way (A), $\cosh 2\eta = e^\sigma$, holds only at the tribonacci point (r01.3).

(b) Combinations of total weight 0 are independent of σ, for example the Compton frequency $m_e c^2/\hbar$ of r08.2. Every measurable ratio of weight $w \neq 0$ moves as $e^{w\sigma}$.

Proof sketch. (a) $e^{\eta_1}e^{\eta_2} = e^{\eta_1+\eta_2}$; the Lorentz case numerically 1.432 against 1.864. (b) additivity of k. □

Reading. The author's parking dimension has a definite form. It is a line on which frames are points and conversions are shifts. Its coordinate is the rapidity, and the conversion law it allows is the Doppler law. It also locates the photon: on the σ-line of the source the photon frame sits at $\sigma = 0$, while in the state disk it sits at $\eta_f = \infty$. The two pictures differ because §25 measures offsets from the photon, whereas the disk measures them from the rest frame of the field. Way (C) follows from the request that the parked conversions compose. The observation of r07 (on-shell $\cot\theta_W$ close to γ_L, way A) points the other way. It is not a test, and decision 5 of r15 stays with the author.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r17 register entries

statement	status after v4.9.0-r17
frame factor γ_L	inverse length of the Bloch vector at the offset, $\gamma_L = 1/\lvert b\rvert$ (r17.1)
C5 factor 3/4, lattice value 0.768	larger eigenvalue of the doublet state at the Hopf, respectively measured, offset (r17.1)
photon	maximally mixed state; information singularity as undefined direction at finite Bures distance $\pi/4$; reached only in the limit (r17.2)
photon frame of §25	reference and limit point; finite γ_L corresponds to $\lvert b\rvert = 1/\gamma_L$, not $b = 0$
scale dimension σ	composable only as rapidity, giving the Doppler law (way C); weight-0 combinations invariant (r17.3)

v4.9.0-r17 ledger and guard update

$$R_{\text{info}}^{v490\text{-}r17} = (0, 0, 1)$$

for (eigenvalue identities established, photon as information singularity made precise, choice of conversion way left to the author).

The layer spends no guard:

$$R_{\text{glob}}^{(v490\text{-}r17,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\text{guard}}^{v490\text{-}r17} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r18 Source-Only: The Old Conversions on the Scale Line, and How Ways A and C Can Hold Together

Purpose. r17 introduced the state disk as a tool for the frames and the scale line σ, and found that composable conversions along σ need the Doppler law (way C of r01). The observation of r07 points to the Lorentz-factor reading γ_L (way A). On 2026-10-22 the author asked to read the conversions of the main body with these tools and to examine what could or must hold for ways A and C to be valid at the same time. The script is `llm/k57_a_und_c.py`. No value is compared with data. Earlier layers are not rewritten.
Source-only status. This v4.9.0-r18 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r18 three readings of one offset

Theoremlet v4.9.0-r18.1. For an offset of rapidity η there are three natural factors:

$$\coth\eta = \frac{1}{\beta} = \sqrt{\frac{p_+}{p_-}}, \qquad e^{\eta}\ \text{(Doppler)}, \qquad \cosh 2\eta = \frac{1}{|b|}\ \text{(inverse Bloch length)}.$$

They are the square root of the eigenvalue ratio of the doublet state (r17.1), the transport factor of way C, and the factor of way A. At the measured offset they are 1.82034, 1.85419 and 1.86444, the three constants $1/\beta_{\mathrm{sym}}$, e^{η} and γ_L that occur in the main body and in r01. All three are equal exactly at the tribonacci point, $\coth\eta = e^{\eta} = \cosh 2\eta = \mathbb{T} = 1.83929$.

Proof sketch. r01.3 shows $\cosh 2\eta = e^{\eta} \Leftrightarrow \coth\eta = e^{\eta} \Leftrightarrow x^3 - x^2 - x - 1 = 0$ for $x = e^{\eta} \neq 1$. □

Consequence. Ways A and C agree numerically exactly when the offset sits at the self-referential fixed point of Theorem 41.18. That fixed point is the condition that the three readings of the offset coincide. The measured offset lies 1.37% above it in γ_L. So that the two ways also agree in numbers, the source's lattice correction (Remark 41.20) has to be carried as a separate offset, for example as a fixed shift on the scale line, rather than being folded into the factor.

v4.9.0-r18 two classes on one scale line

Theoremlet v4.9.0-r18.2. On the scale line σ of r17 there are two kinds of quantity.

(a) *Scale quantities* (frequencies, lengths, masses, energies) move by a homomorphism, $X(\sigma) = e^{k\sigma}X(0)$. This is way C, and it composes.

(b) *Readout quantities* (visibility, contrast of the compensator ports, eigenvalue ratios, mixing ratios) are state functions of the total offset, for example $V(\sigma) = \operatorname{sech} 2\sigma$, $p_-/p_+ = \tanh^2\sigma$. Their factor $\cosh 2\sigma$ is way A. They compose as well, through addition of rapidities: after two boosts the visibility is $\operatorname{sech} 2(\sigma_1 + \sigma_2)$, not the product of the single values.

The obstruction of Theoremlet r01.2 arises only if Lorentz factors are multiplied as though they were transports. With the two classes separated, ways A and C hold at the same time, each for its own class.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof sketch. (a) $e^{\sigma_1} e^{\sigma_2} = e^{\sigma_1+\sigma_2}$. (b) the visibility of r03.3 is a function of the rapidity to the frame with $E \parallel B$, and rapidities add along a line. Checked numerically: $\operatorname{sech} 2(\sigma_1+\sigma_2) = 0.311$ against the product 0.452 for $\sigma_1 = 0.3$, $\sigma_2 = \eta$. □

What must hold. Both ways are valid together if, and in this reading only if:

1. every conversion is assigned to one class, scale or readout;
2. frames are added as rapidities on one scale line, never by multiplying Lorentz factors;
3. for equality in numbers in addition, the offset sits at the tribonacci point, and the lattice correction is carried separately.

Items 1 and 2 give consistency. Item 3 gives equal values. The author's idea of parking conversions in a dimension of their own (r17) is what makes item 2 possible.

v4.9.0-r18 the conversions of the main body, read again

conversion	k	class	A	C	C/A
clock frequency, Cor. 22.21	1	scale	1.8644	1.8542	−0.55%
Planck length, Thm. 25.5	3/2	scale	2.5458	2.5248	−0.82%
neutrino mass, Prop. 42.4	4	scale	12.084	11.820	−2.18%
matter error, Cor. 42.17	5	scale	22.529	21.916	−2.72%
vacuum energy ($k = 5$ after r03)	5	scale	22.529	21.916	−2.72%
Hawking temperature, Prop. 21.4	1	scale	1.8644	1.8542	−0.55%

Readout quantities, way A as state function: the visibility and compensator contrast of r03 ($1/\gamma_L = 0.536$), the eigenvalue ratio $p_-/p_+ = \beta^2_{\text{sym}} = 0.302$, which is the Ω_M of Theorem 22.16, the mixing ratio observed in r07 ($\cot\theta_W \approx \gamma_L$), and the structural ratio $\gamma_{\text{BI}}/\gamma_{\text{string}}$ of Theorem 20.5.

Reading. All conversions of §25 in the table are scale quantities. Under the class reading they take way C, and their values change by 0.55% to 2.7%. The quantities that keep γ_L are readouts. These include the mixing ratio of r07 and the cascade fraction of Theorem 22.16. The tension between r07 (way A) and r17 (way C) disappears, because the two concern different classes. This is an offered reading. Whether the author's conversions were meant this way is decision 5 of r15. No compared value of the main body (Corollary 42.24, Proposition 42.7) is re-evaluated here.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r18 register entries

statement	status after v4.9.0-r18
$1/\beta_{\mathrm{sym}}$, e^{η}, γ_L	three readings of one offset (eigenvalue ratio, Doppler, inverse Bloch length), equal at the tribonacci point (r18.1)
ways A and C	can hold together: C for scale quantities, A as state function for readout quantities, frames added as rapidities (r18.2); numerical equality needs the tribonacci point and a separate lattice offset
conversions of §25	scale quantities; under the class reading way C, changes 0.55%–2.7%
r07 against r17	tension removed under the class reading (different classes)
decision 5 of r15	refined; still with the author

v4.9.0-r18 ledger and guard update

$$R_{\mathrm{conv}}^{v490\text{-}r18} = (0, 0, 1)$$

for (three readings identified, conditions for A and C together stated, assignment of classes left to the author).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r18,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r18} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r19 Source-Only: The Class Split Applied to the Main Body

Purpose. r18 offered a reading under which ways A and C of r01 hold together: scale quantities move by the Doppler factor, readout quantities keep γ_L as a state function. On 2026-10-22 the author asked to apply that split. This layer goes through every place in the main body where γ_L converts between frames, assigns each place to a class by a fixed rule, and recomputes it. The script is `llm/k58_klassentrennung.py`. No new prediction is made, no value is compared with new data, and no earlier layer is rewritten. The applied split remains an offered reading. Decision 5 of r15 stays with the author.
Source-only status. This v4.9.0-r19 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r19 the rule

Rule. A quantity that is a monomial in constants with frame-insertion counts, $X = \prod_i X_i^{n_i}$, has the weight $k(X) = \sum_i n_i k(X_i)$ and is a *scale quantity*. It converts by $e^{k\eta}$ (way C). A quantity that is a function of the doublet state is a *readout quantity*. Examples are a visibility, an eigenvalue ratio, a mixing angle, or a ratio of two constants with $k = 0$. It keeps γ_L (way A). The rule fixes the class from the form of the quantity alone, before any comparison value is read. A dimensionless constant can be a scale quantity: α is a monomial with $k = 1$. The offset is $\eta = \frac{1}{2}\operatorname{arcosh}\gamma_L$ as in r18, so $e^{\eta} = 1.85419$.

Theoremlet v4.9.0-r19.1. Under the rule, the definition of §25, $X^{(\text{photon})} = \gamma_L^{-k(X)} X^{(\text{matter})}$, becomes $X^{(\text{photon})} = e^{-k(X)\eta} X^{(\text{matter})}$. Each insertion of z^- contributes one Doppler factor, and the weights add: $k(XY) = k(X) + k(Y)$ is now exact, also for half-integer weights such as $k(\ell_P) = 3/2$. With γ_L per insertion the additivity holds as well, but the composition of two frame steps does not (Theoremlet r01.2).

Proof sketch. $e^{k_1\eta} e^{k_2\eta} = e^{(k_1+k_2)\eta}$, and $e^{k\eta_1} e^{k\eta_2} = e^{k(\eta_1+\eta_2)}$ for consecutive steps. □

v4.9.0-r19 scale quantities of the main body

place	k	A (γ_L)	C (e^{η})	C/A
photon-frame Planck length, Thm. 25.5	3/2	6.35×10^{-36} m	6.40×10^{-36} m	+0.83%
vacuum ratio, Prop. 42.16 ($k = 5$ after r03)	5	22.53	21.92	−2.72%
residual of Cor. 42.17 ($\log_{10}$)	5	−120.6	−120.7	—
neutrino factor, Prop. 42.4	4	12.08	11.82	−2.18%
neutrino photon-frame suppression	−3	0.1543	0.1569	+1.67%
hadronic factor at $k(\alpha_s) = 1$, Prop. 42.7	−1	0.5364	0.5393	+0.55%
lower end of the range there	−2	0.2877	0.2909	+1.11%
$\Lambda_{\text{bare}}^{(\text{photon})}$, Prop. 42.18 ($m_P^4$)	−2	57.5	58.1	+1.11%
$\beta_{\text{sym}}^2 \gamma_L^{-3}/200$, Remark 42.21	−3	2.33×10^{-4}	2.37×10^{-4}	+1.67%
first zero in the photon frame, Cor. 22.21	−1	7.581	7.623	+0.55%
β_{eff} of Cor. 22.21(iii)	1	3.729	3.708	−0.55%

© 2026 Marc Brendecke. Preprint — not peer reviewed.

None of these places changes its qualitative statement. The residual of Corollary 42.17 still lies near $10^{-120.7}$, Remark 42.21 still reads as a cancellation at the level 10^{-3} to 10^{-4}, and the neutrino factor stays near 12. The changes are of order 1%.

v4.9.0-r19 places where the split matters

Theoremlet v4.9.0-r19.2. Three statements of Corollary 42.24 respond to the split.

(P5) Under C, $\mathrm{d}\ln X = k(X)\,\mathrm{d}\sigma$ for every scale quantity. The ratio of annual modulations is therefore $k(\alpha)/k(G_N) = 1/2$ exactly, with no prefactor. The correction of r09, which replaced the stated $\gamma_L^{-1} \approx 0.537$ by 1/2, follows from the rule without a separate argument.

(P3) The exponent γ_L^5 of Remark 41.8 has weight 5 and is a scale factor. Under C the formula gives $\beta_{\mathrm{sym}} = 0.54847$, -0.15% against 0.5493, instead of 0.54923 (-0.01%). Solved self-consistently with $\eta = \operatorname{artanh}\beta_{\mathrm{sym}}$ it gives 0.54824 (-0.19%). The match of P3 does not survive the split. This agrees with r09, which had already classed P3 as a coincidence.

(P6) The Hawking temperature has $k = -1$ (Proposition 21.4) and is a scale quantity. Under C, $T_{\mathrm{Sphaera}}/T_H^{(\mathrm{tachyon})} = \gamma_{\mathrm{string}} e^{\eta} = 0.23619$. The value $\gamma_{\mathrm{BI}} = \gamma_{\mathrm{string}}\gamma_L = 0.23750$ is a readout and unchanged, so the identification $\gamma_{\mathrm{BI}} = T_{\mathrm{Sphaera}}/T_H^{(\mathrm{tachyon})}$ of item (iii) there holds to -0.55%, not exactly. It becomes exact only at the tribonacci point.

Proof sketch. (P5) $\mathrm{d}\ln e^{k\sigma} = k\,\mathrm{d}\sigma$. (P3), (P6) numerical, `llm/k58_klassentrennung.py`. □

Readout quantities, unchanged. $\gamma_{\mathrm{BI}}/\gamma_{\mathrm{string}} = \gamma_L$ (Theorem 20.5), the visibility $1/\gamma_L$ of r03, $\Omega_M = \beta_{\mathrm{sym}}^2$ of Theorem 22.16, the Koide phase of P2 (132.88°, a function of β_{sym} alone), and the mixing ratio $\cot\theta_W \approx \gamma_L$ of r07. The pre-registered test H15 rests on that mixing ratio and is not affected.

v4.9.0-r19 register entries

statement	status after v4.9.0-r19, under the class split
conversions of §25	one Doppler factor per insertion; values change by 0.55% to 2.7%, statements unchanged (r19.1)
P5	1/2 exactly, the r09 correction follows from the rule (r19.2)
P3	match lost (-0.15%); consistent with the r09 class "coincidence"
P6, item (iii) of Prop. 21.4	γ_{BI} as Hawking ratio only to -0.55%; exact only at the tribonacci point
P2, H15, Ω_M, Thm. 20.5	readouts, unchanged
class split	applied; still an offered reading, decision 5 of r15 with the author

v4.9.0-r19 ledger and guard update

$$R_{\mathrm{split}}^{v490\text{-}r19} = (0, 0, 1)$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

for (all conversion places classified, consequences computed, adoption of the split left to the author).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r19,+)} = (0,1,1,1,1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r19} = (1,1,1,1,1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r20 Source-Only: A Provenance Hypothesis for $\Delta_\gamma < 1/5$, the Photon-Frame Clock Gain at the Hopf Point

Purpose. r12 derived $\Delta_\gamma < 1/5$ from the chart data of r14 under the premises (P1)–(P4). It left open where those data, and therefore the five, come from. The r65 text is lost. On 2026-10-22 the author suggested that the value was produced in a photon-frame calculation, and that it need not be correct or generally valid. This layer proposes a provenance on that basis and examines how far the value holds away from it. The script is `llm/k59_herkunft_delta_gamma.py`. The proposal is a hypothesis about origin. It does not recover r65, makes no new prediction, and compares nothing with data.
Source-only status. This v4.9.0-r20 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r20 the hypothesis

r14 calls the charts with a window "photon-frame timing-assisted" (Proposition A.225.2). The gain G is therefore the gain of the photon-frame clock. A clock frequency converts between frames with the frame factor (Corollary 22.21).

Theoremlet v4.9.0-r20.1. At the Hopf point $\beta^2_{\mathrm{sym}} = 1/3$, the doublet state of r17 has $\gamma_L = 2$, $p_+ = 3/4$ and $p_- = 1/4$. The binding chart $V3$ of r12 and the clock seed of r15 consist of exactly these numbers:

$$G_{V3} = 2 = \gamma_L, \qquad \kappa_{V3} = \tfrac{1}{4} = p_-, \qquad N_{V3} = 5 = 1 + \frac{1}{p_-} = 2 + \frac{1}{\beta^2_{\mathrm{sym}}}, \qquad \kappa_{\mathrm{seed}} = \tfrac{3}{4} = p_+.$$

The value $N = 5$ is the balance condition $\kappa(N-1) = 1$ of r12 at $\kappa = p_-$. With these values $1 - 16/(GS) = 1 - 16/(2 \cdot 10) = 1/5$ exactly. The collar row of r05 (ratio $5/6 = 1/(1 + \frac{1}{5})$) and the C5 factor $3/4 = p_+$ (r17) belong to the same point.

Proof sketch. r17.1 with $\beta^2_{\mathrm{sym}} = 1/3$: $\gamma_L = (1+\beta^2_{\mathrm{sym}})/(1-\beta^2_{\mathrm{sym}}) = 2$, $p_- = \beta^2_{\mathrm{sym}}/(1+\beta^2_{\mathrm{sym}}) = 1/4$. Exact rational arithmetic. □

Hypothesis. The bound $\Delta_\gamma < 1/5$ was obtained with the photon-frame clock gain set to the frame factor at the Hopf point, $\gamma_L = 2$, the continuum value $\beta_{\mathrm{sym}} = 1/\sqrt{3}$ without the lattice correction of Remark 41.8. A rounding of $\gamma_L = 1.864$ to the half-integer grid of r14 gives the same $G = 2$. The hypothesis explains why the five arises at exactly one chart (r92). It also explains why the C5 factor and the clock bound share one point.

Weight of the evidence. The r14 values lie on grids of quarters (κ) and halves (G). A single coincidence such as $G = 2$ would carry little weight. Four numbers that match at one point, together with the balance condition, carry more. Against the hypothesis: the other window charts $V5$ ($G = 5/2$) and $V8$ ($G = 2$, $\kappa = 1$) have no comparable reading, and the threshold 16 is dyadic and frame-free.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r20 the bound away from the Hopf point

Theoremlet v4.9.0-r20.2. Keep the chart form of r12 and the threshold 16, and replace the Hopf values by those at the measured offset. Then the bound of $V3$ becomes:

reading	G	κ, N	bound
Hopf point (source)	2	1/4, 5	0.2000
$G = \gamma_L$, chart fixed	1.8644	1/4, 5	0.1418
$G = e^{\eta}$ (clock as scale quantity, r19), chart fixed	1.8542	1/4, 5	0.1371
$G = \gamma_L$, $\kappa = p_- = 0.2318$, $N = 5$	1.8644	0.2318, 5	0.1095
$G = e^{\eta}$, $\kappa = p_-$, $N = 5$	1.8542	0.2318, 5	0.1045
$G = \gamma_L$, balanced $N = 1 + 1/p_- = 5.31$	1.8644	0.2318, 5.31	0.1925

In every reading the bound lies below 1/5. A different generalization, $\tan\alpha < 1/\gamma_L$ in place of $\tan\alpha < 1/2$, gives $1/(1+\gamma_L^2) = 0.2234$ instead and lies above it.

Proof sketch. $\Delta_\gamma^{\max} = 1 - 16/(GS)$ with $S = N + \kappa N(N-1)$ and the values of r17.1 at $\beta_{\mathrm{sym}} = \tanh\eta$. Numerical. □

Consequences.

1. 1/5 is a value of the Hopf point. It is not frame-invariant. At the measured offset the bound depends on how the chart is carried over, and the readings that keep the chart form give 0.10 to 0.19.
2. In those readings 1/5 is not conservative. A drift loss between the true bound and 1/5 would be admitted by the stated condition and still close the $V3$ window.
3. r12.2(b) changes with it. For the readings with the chart fixed, the bound is about 0.14. The tolerated collar row of r05, which exports $1/6 = 0.167$, then lies outside, and so does the C5 stage (1/4). The separation of the C5 stage from the clock selector stays, but the reading that r05 tolerated that row for this reason holds only at the Hopf point.

v4.9.0-r20 guard status

$W_{\mathrm{clk}}^{\mathrm{loc}}$ under $\Delta_\gamma < 1/5$ is a permitted local statement. This layer makes its bound frame-typed: $\Delta_\gamma < \Delta_\gamma^{\max}(\eta)$, with $\Delta_\gamma^{\max} = 1/5$ at the Hopf point. At the measured offset the condition becomes narrower, about 0.14 for the chart fixed. A narrower local condition claims less, not more. No guard is spent, and nothing global is approached. Which carrying-over is right is not decided here. It needs the clock dynamics that r13 left open.

v4.9.0-r20 register entries

statement	status after v4.9.0-r20
provenance of $\Delta_\gamma < 1/5$	hypothesis: photon-frame clock gain $G = \gamma_L = 2$ at the Hopf point; $V3$ and clock seed are Hopf data (r20.1); r65 text still lost
$\Delta_\gamma < 1/5$	exact at the Hopf point; frame-typed, at the measured offset 0.10–0.19 with the chart form kept (r20.2)
r12.2(b), collar row of r05	inside only at the Hopf point; outside at about 0.14
Hopf point	now carries the C5 factor 3/4, the export 1/4, the $V3$ coupling and gain, and $\Delta_\gamma = 1/5$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r20 ledger and guard update

$$R_{\mathrm{prov}}^{v490\text{-}r20} = (0, 0, 1)$$

for (provenance hypothesis stated with its evidence, bound away from the Hopf point computed, choice of carrying-over open).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r20,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r20} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r21 Source-Only: The Clock Bound Frame by Frame, and a Derivation of $1/5$

Purpose. r20 traced $\Delta_\gamma < 1/5$ to the Hopf point and found that the bound changes away from it. How it changes depended on how the chart was carried over. On 2026-10-22 the author asked to compute the bound separately for each frame, since a distinction by frame may have to be built in everywhere, and to attempt a concrete derivation. This layer states five premises, derives the bound per frame from them, and shows that $1/5$ follows from one consistency condition. The script is `llm/k60_delta_gamma_je_frame.py`. No new prediction is made and nothing is compared with data. Earlier layers are not rewritten.
Source-only status. This v4.9.0-r21 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r21 premises

The window condition is that of r12, $G(1-\Delta_\gamma)S > 16$ with $S = N+\kappa N(N-1)$. The doublet state of r17 at $x = \beta_{\mathrm{sym}}^2$ has $|b| = 1/\gamma$, $\gamma = (1+x)/(1-x)$, $p_- = x/(1+x)$, $p_+ = 1/(1+x)$.

(D1) *Clock gain.* The photon-frame clock restores the visibility $|b|$ of the compensator readout to 1, so $G = 1/|b| = |\gamma|$. This is a readout quantity in the sense of r19 and keeps γ_L (way A).

(D2) *Pair coupling.* κ is the content of the partner port. In the matter frame ($b_1 > 0$) that is p_-. In the tachyon frame ($b_1 < 0$) J_B exchanges the ports (r02, r04), so it is p_+.

(D3) *Balance.* In the frame that needs the clock, the partner content of the other packets refills one packet: $\kappa(N-1) = 1$. This is the balance condition that singles out $V3$ in r12.

(D4) *Packet count.* N counts packets and is the same in every frame.

(D5) *Tachyon description.* In the tachyon frame the chart is the clock seed of the source, $(N, \kappa, G) = (5, \frac{3}{4}, 1)$, so $G_T = 1$.

(D2) and (D3) follow from r02, r04 and r12. (D1) gives a mechanism to the gain that r14 calls photon-frame timing. (D4) and (D5) are assumptions. (D5) is taken from the data of the source.

v4.9.0-r21 the derivation

Theoremlet v4.9.0-r21.1. Under (D1)–(D5), with $N = 1 + 1/p_- = 2 + 1/x$ from (D3), the survival reserves $1 - 16/(GS)$ are

$$\Delta_M^{\max}(x) = \frac{10x^2 - 5x + 1}{(1+x)(1+2x)}, \qquad \Delta_T^{\max}(x) = \frac{1 + 3x - 14x^2}{(1+x)(1+2x)}.$$

Require that the window's margin be the same whichever frame describes it, $G_M S_M = G_T S_T$. This has the single solution $x = 1/3$, the Hopf point. There $N = 5$, $S_M = 10$, $S_T = 20$, and

$$\Delta_\gamma^{\max} = \frac{1}{5}$$

in both frames. Without (D3), with $N = 5$ held fixed, the same condition reads $4p_-^2 - 9p_- + 2 = 0$, with the admissible root $p_- = 1/4$, again the Hopf point.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof sketch. Substitute (D1)–(D5) and simplify symbolically; solve $\gamma S_M = S_T$ for $x > 0$. Checked with a computer algebra system. □

Reading. This is a derivation of 1/5 from the doublet state and one consistency condition: one physical window must not survive in one frame and fail in the other. The integer chart $(5, \frac{1}{4}, 2)$ and the seed $(5, \frac{3}{4}, 1)$ of the source are then its matter and tachyon descriptions at that point. The seed is the tachyon image of $V3$ under J_B. It needs no gain, because in the tachyon frame the major content sits in the partner port and the pair flux alone reaches $S = 20 \geq 16$. The derivation does not recover r65. It shows that a calculation which fixes the frame factor at its Hopf value $\gamma_L = 2$, as r20 proposed, lands on the one point where matter and tachyon agree.

v4.9.0-r21 the bound in each frame

frame, point	$\lvert G\rvert$	κ	N	S	reserve	remark
matter, Hopf	2	0.250	5	10	0.200	window
tachyon, Hopf	1	0.750	5	20	0.200	no window needed
matter, measured	1.864	0.232	5.31	10.63	0.1925	window
tachyon, measured	1	0.768	5.31	22.92	0.302	no window needed
photon, disk ($b = 0$)	∞	0.5	—	—	$\to 1$	degenerate
photon, scale line ($G = 1$, $N = 5$)	1	0.5	5	15	< 0	blocked

Theoremlet v4.9.0-r21.2.

(a) Matter frame: at the measured offset the bound is 0.1925, 0.0075 below 1/5. The frame-free 1/5 is slightly too generous there. The collar row of r05 (1/6) stays inside and the C5 stage (1/4) outside.

(b) Tachyon frame: the pair flux reaches the relay threshold without the clock. $W_{\mathrm{clk}}^{\mathrm{loc}}$ has no content there. The unassisted reserve is 0.302.

(c) Photon frame: the bound is degenerate. In the state disk the gain $1/|b|$ diverges and every drift is tolerated. On the scale line of r17, with no frame factor, the window is closed. This is the information singularity of r17.2 in the clock layer: at the photon the quantity that the bound constrains is not defined.

(d) Away from the Hopf point the matter and tachyon reserves differ, 0.19 against 0.30. A frame-free bound exists only at $x = 1/3$.

Proof sketch. Numerical evaluation of r21.1 and of the window condition at the stated values. □

Relation to r20. r20 listed several ways to carry the chart away from the Hopf point. The readings with the chart held fixed gave 0.10 to 0.14 and put the collar row outside. They violate (D3). Under (D1)–(D5) the balanced reading is selected, the matter bound is 0.1925, and consequence 3 of r20 does not apply.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r21 a rule for frame distinction

The computation suggests a rule for building the frame distinction into other places. The tachyon frame is the J_B image of the matter frame: $x \to 1/x$, $p_\pm$ exchanged, $|\gamma|$ unchanged and γ negative, the phase $i\pi$ of the superluminal rapidity of r20 (v4.8.0). The photon frame is the fixed point $p_+ = p_- = \frac{1}{2}$, where readout quantities degenerate. A statement that is valid in one frame needs a separate check in the other two. It is frame-free only where the descriptions agree. For the clock bound that is the Hopf point alone. This layer proposes the rule. Applying it to the other layers is left open.

v4.9.0-r21 guard status

$W^{\mathrm{loc}}_{\mathrm{clk}}$ stays a local statement. Its condition becomes frame-typed: $\Delta_\gamma < \Delta_M^{\max}(\beta^2_{\mathrm{sym}})$ in the matter frame, 1/5 at the Hopf point and 0.1925 at the measured offset; empty in the tachyon frame; undefined at the photon. The narrower matter-frame value claims less than 1/5. No guard is spent. The photon-frame degeneracy is no route to W^{glob}_γ: it is the absence of the constrained quantity, not a global clock statement.

v4.9.0-r21 register entries

statement	status after v4.9.0-r21
$\Delta_\gamma < 1/5$	derived under (D1)–(D5) as the unique frame-consistent bound, at the Hopf point (r21.1); r65 text still lost
clock bound per frame	matter 0.1925 (measured), tachyon no window (reserve 0.302), photon degenerate (r21.2)
clock seed $(5, \frac{3}{4}, 1)$	tachyon image of $V3$ under J_B
r20 consequence 3	superseded under (D3): collar row inside
frame distinction	rule proposed (tachyon = J_B image, photon = degenerate fixed point); application to other layers open

v4.9.0-r21 ledger and guard update

$$R^{v490\text{-}r21}_{\mathrm{frame}} = (0, 0, 1)$$

for (bound derived under stated premises, values per frame computed, application of the frame rule elsewhere open).

The layer spends no guard:

$$R^{(v490\text{-}r21,+)}_{\mathrm{glob}} = (0, 1, 1, 1, 1), \qquad \Gamma^{v490\text{-}r21}_{\mathrm{guard}} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r22 Source-Only: The Frame Rule as a Standing Convention, Applied to the Main Statements

Purpose. r21 proposed a rule for distinguishing the three frames. On 2026-10-22 the author asked to build it in. The document grows append-only, so earlier statements are not edited. This layer adopts the rule as a standing convention for every later layer, applies it to the main statements that the v4.9.0 line has touched, and records where each statement holds. The script is `llm/k61_frame_regel.py`. No new prediction is made and nothing is compared with data.

Source-only status. This v4.9.0-r22 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r22 the rule

Convention F (frame rule). Let $x = \beta_{\mathrm{sym}}^2$. The *matter frame* M is the doublet state of r17 at x: $\gamma = (1+x)/(1-x)$, $p_+ = 1/(1+x)$, $p_- = x/(1+x)$, compensator port holding p_+. The *tachyon frame* T is the J_B image of M: $x \mapsto 1/x$, $p_\pm$ exchanged, $\gamma \mapsto -\gamma$, which is the phase $i\pi$ of $\cosh 2\eta$ under the superluminal rapidity $\eta + i\pi/2$ of r20 (v4.8.0). The *photon frame* P is the fixed point $b = 0$, $p_+ = p_- = \frac{1}{2}$.

(F1) A statement is *frame-free* if M and T give the same value. Otherwise it carries the frame in which it holds.

(F2) At P a readout quantity is evaluated separately and may be degenerate. A degenerate value is not a global statement.

(F3) A conversion of weight k into T has the magnitude of its class (r19) and the phase $(-1)^k$, from $\gamma \mapsto -\gamma$.

(F4) From this layer on, every layer contains a subsection labelled `subsec:v490-rNN-frames` that states the M, T and P status of its results. The guard linter (`llm/training/guard_linter.py`) reports a layer from r22 on that lacks it.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r22 the main statements frame by frame

statement	M	T	P	status
frame factor γ	1.864	-1.864	∞	magnitude frame-free
visibility $\lvert b\rvert = 1/\lvert\gamma\rvert$ (r03)	0.536	0.536	0	frame-free
eigenvalues (p_+, p_-) (r17)	$(0.768, 0.232)$	$(0.232, 0.768)$	$(\frac{1}{2}, \frac{1}{2})$	exchanged
entropy (r17)	0.781 bit	0.781 bit	1 bit	frame-free
C5 factor, kick reading $\cos^2\frac{\pi}{6}$ (r10)	3/4	3/4	—	frame-free
C5 factor as compensator eigenvalue (r17)	0.768	0.232	$\frac{1}{2}$	matter only
$\Omega_M = p_-/p_+$, Thm. 22.16	0.302	3.31	1	matter only
Koide phase, Thm. 22.5	132.88°	undefined	—	matter only
$\cot\theta_W \approx \gamma$ (r07)	1.864	-1.864	∞	$\theta \mapsto \pi - \theta$
$\sin^2\theta_W = 1/(1+\gamma^2)$, H15	0.2234	0.2234	0	frame-free
P5 $= k(\alpha)/k(G_N)$ (r19)	1/2	1/2	—	frame-free
$k = 0$ quantities (c, γ_{BI}, $m_e c^2/\hbar$)	same	same	same	frame-free in all three
clock bound Δ_γ (r21)	0.1925	no window	degenerate	matter only

Theoremlet v4.9.0-r22.1. Under Convention F:

(a) Frame-free are the quantities that depend on the state only through $|b|$ or through the unordered pair of eigenvalues (visibility, entropy, $\sin^2\theta_W$), the kick reading of the C5 factor, ratios of weights, and the $k = 0$ quantities.

(b) Matter-only are the quantities that distinguish the compensator port from the partner port: the eigenvalue reading of 3/4, Ω_M, the Koide phase, and the clock bound. Their tachyon values are the inverse ratio, an undefined arcsin, or an absent window.

(c) At the photon every port-sensitive quantity is degenerate ($p_+ = p_-$), in line with r17.2.

Proof sketch. J_B exchanges the ports and leaves $|b|$ and the unordered spectrum fixed. $AR_\alpha A = \cos\alpha\, A$ as for C_{HC}. The rest is the table, computed in `llm/k61_frame_regel.py`. □

v4.9.0-r22 the tachyon sign scale

Theoremlet v4.9.0-r22.2. By (F3), a length ($k = 3/2$) acquires the phase $(-1)^{3/2} = \mp i$ and its square the phase -1. This reproduces the signed tachyon scale $\ell_T^2 = -\lambda_T^2$, $\ell_T = i\lambda_T$, of §A.269 (up to the choice of branch). Taking the phase from the rapidity $\eta + i\pi/2$ directly, one insertion at a time, gives $e^{3i\pi/4}$ for the length and $-i$ for its square. That does not reproduce it.

Proof sketch. $(-1)^{3/2} = e^{3i\pi/2}$; $e^{(3/2)\cdot i\pi/2} = e^{3i\pi/4}$, $e^{3\cdot i\pi/2} = -i$. □

Consequence. The class split of r19 fixes the magnitude of a tachyon conversion. Its phase comes from the sign of the invariant, $s \mapsto -s$, that is $\gamma \mapsto -\gamma$, and not from the Doppler factor at the complex rapidity. With this, way C and the permitted statement $\ell_T^2 = -\lambda_T^2$ are compatible. Without (F3), way C would contradict it. The layer adds nothing to that permitted statement. It records which reading of the tachyon conversion is consistent with it.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r22 status in the three frames

result of this layer	M / T / P
Convention F	defined on all three; T as J_B image, P as fixed point
classification r22.1	frame-free set holds in M and T, matter-only set holds in M, port-sensitive quantities degenerate in P
sign scale r22.2	a statement about T; reproduces the r82 scale; trivial in M; at P no conversion is defined

v4.9.0-r22 register entries

statement	status after v4.9.0-r22
frame rule	adopted as Convention F for all later layers; required subsection, checked by the guard linter
Ω_M, Koide phase, eigenvalue reading of 3/4, clock bound	matter-frame statements (r22.1)
visibility, entropy, $\sin^2\theta_W$, kick reading of 3/4, P5, $k = 0$	frame-free between M and T (r22.1)
tachyon conversions	magnitude by class (r19), phase $(-1)^k$; consistent with $\ell_T^2 = -\lambda_T^2$ (r22.2)
earlier layers	not edited; their statements are read through the table above

v4.9.0-r22 ledger and guard update

$$R_{\mathrm{F}}^{v490\text{-}r22} = (0, 0, 1)$$

for (convention adopted, main statements classified, application to the remaining layers of the main body open).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r22,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r22} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r23 Source-Only: Convention F Applied to the Main Body

Purpose. r22 adopted the frame rule as Convention F and applied it to the statements that the v4.9.0 line had touched. On 2026-10-22 the author asked to apply it to the rest of the main body. This layer does that for every statement environment of the main body (lines 1 to about 52 100) that depends on a frame. The script `llm/k62_frame_regel_hauptkorpus.py` lists and classifies them. Earlier layers are not edited. No prediction is made, and nothing is compared with data.
Source-only status. This v4.9.0-r23 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r23 scope and categories

The scan takes every theorem, proposition, corollary, lemma, conjecture and definition of the main body that contains β_{sym}, γ_L, z^-, γ_{BI}, γ_{string} or the words “photon frame”, “matter frame” or “tachyon”. It finds 113 of the 2234 statement environments there. Each one gets a category:

F frame-free: it depends only on $|b|$, on $|\gamma|$, on $k = 0$ quantities, or on pure arithmetic;

M matter only: it depends on the ordered or signed offset β_{sym}, which becomes $1/\beta_{\mathrm{sym}}$ in the tachyon frame;

C conversion of weight k: matter value as stated; tachyon value of the magnitude of its class (r19) with phase $(-1)^k$ (F3);

S a statement at the photon frame of the main body ($\beta_{\mathrm{sym}} = 0$, $\sigma = 0$);

P a statement in the limit $\beta \to 1$;

D it defines the sectors or frames themselves;

N it carries the photon-frame name without any frame quantity.

v4.9.0-r23 the classification

category	count	content
F	7	structural constants, k-table, tribonacci fixed point in $\|\gamma\|$, Guinand–Weil identity
M	29	cosmology, Koide, OP1/OP2 complex, Hopf angle, quasiconformal reading
C	12	constant conversions of §25 and the hypothesis propositions
S	9	vanishing of Λ and the Θ–R balance in the photon frame
P	3	horizon and saturation limit
D	13	sector and frame definitions, axioms
N	40	OP1 working layer, OP5 halo/collar/Dixon grammar

F: Theorems 12.3, 20.4, 24.2, 25.3, 41.18, 42.23; Proposition 20.2.
M: Theorems 21.3, 21.5, 21.10, 22.2, 22.5, 22.9, 22.10, 22.11, 22.14, 22.16, 26.4, 41.16, 41.21, 41.26, 41.27, 42.14; Definitions 23.2, 41.9; Propositions 26.5, 41.6, 41.7, 41.12, 41.24, 42.2, 42.5; Corollaries 41.19, 41.23, 42.24 (mixed: P1 and P6 are C, P2–P4 are M, P5 is F); the unlabelled corollary “Cosmological constant revisited” of §25.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

C: Theorems 20.5, 25.5; Propositions 21.4, 42.4, 42.7, 42.16, 42.18; Corollaries 22.21, 24.6, 42.17; the definition of k and the α-drift corollary of §25.
S: Theorem 42.19; Propositions 23.5, 42.22, 42.27, 42.30; Definitions 42.20, 42.32, H9.126; Conjecture 42.33.
P: Propositions 21.2, 42.8; the Schwarzschild-horizon definition preceding Proposition 21.2.
D: Theorems 12.1, 20.3, 20.6, 21.7, 22.22, 32.1, 40.1, 40.8; Proposition 16.1; Corollaries 29.26, 35.4; the sector-classification definition of §15; the definition of the tachyon branch.
N: the OP1 working statements (Definition H3.31, Corollary H3.35, Theorem H3.36, Definition H9.133, Corollary H9.135), the live OP5 Step (B) kernel (Definition H9.238, Proposition H9.239), one OP queue placement, and 32 unlabelled statements of the OP5 halo, collar, tunnel and Dixon grammar.

v4.9.0-r23 what the classification shows

Theoremlet v4.9.0-r23.1.

(a) *The tachyon of the main body is not the tachyon of Convention F.* Theorem 20.6 puts the tachyon at $-\beta_{\text{sym}}$. Convention F follows Reading B and the superluminal rapidity of r20 (v4.8.0), and puts it at $1/\beta_{\text{sym}}$ with $\gamma \mapsto -\gamma$. Under the main body's own tachyon, every even function of β_{sym} would be frame-free. 18 statements are of that kind, 12 of the matter-only ones and 6 of the S statements: Ω_M, $\Lambda_{\text{obs}} \sim \beta_{\text{sym}}^2$, the frame term of the Lagrangian, the β_{sym}^2 term of the proper time, the Beltrami coefficient. Under F they hold in the matter frame only.

(b) *The photon frame of the main body is not the photon of Convention F.* The main body measures offsets from the photon, so its photon frame is $\beta_{\text{sym}} = 0$. In the state disk $\beta_{\text{sym}} = 0$ is the pure state $|b| = 1$, and the photon is the centre $b = 0$, reached as $\beta \to 1$ (r17). The nine S statements ($\Lambda = 0$, the frame term vanishing, Step (A) trivial in the photon frame) hold at $\sigma = 0$ on the scale line. They do not carry over to P. At P, $p_-/p_+ = 1$ and the frame term is maximal.

(c) *The limit $\beta \to 1$ of the main body is the photon of Convention F.* The Schwarzschild horizon ($\beta \to 1$), the cascade singularity of Proposition 21.2, and the saturation end state of Proposition 42.8 sit where F places the photon fixed point. The undefined cascade there is the degeneracy of (F2).

(d) *The quasiconformal reading sees the tachyon as an orientation reversal.* In Theorem 41.26 $\mu = \beta_{\text{sym}}^2$ and $K = \gamma_L$. In the tachyon frame $\mu = 1/\beta_{\text{sym}}^2 > 1$ and $K = -\gamma_L$: an orientation-reversing map, the same sign change $\gamma \mapsto -\gamma$ as in F.

(e) *Conversions acquire a tachyon sign.* By (F3), quantities of odd weight change sign in the tachyon frame: the Hawking temperature ($k = -1$), α ($k = 1$), the vacuum ratio ($k = 5$). Even weights do not (neutrino factor, $k = 4$). Lengths take the phase $\mp i$, consistent with $\ell_T^2 = -\lambda_T^2$ (r22.2).

(f) *The N statements, including the local OP5 closure of the declared grammar, contain no frame quantity.* Convention F neither narrows nor widens them. They stay local within their grammar.

Proof sketch. (a) count of the M and S statements whose β_{sym}-dependence is even; listed in the script. (b), (c) $|b| = \operatorname{sech} 2\eta$ is 1 at $\eta = 0$ and 0 as $\eta \to \infty$. (d) $K = (1+\mu)/(1-\mu)$ at $\mu = 1/\beta_{\text{sym}}^2$. (e) (F3). (f) by inspection: no frame macro, no offset. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

A sign remark. The sector-classification definition of §15 gives matter $\beta < 0$. Theorem 20.6 gives matter $+\beta_{\mathrm{sym}}$. The two signs disagree within the main body. v4.8.0-r06 recorded that the classification by $\mathrm{Re}\, z$ is not invariant, and v4.8.0-r07 replaced it by the sign of s. Convention F rests on that assignment, so the disagreement does not enter here. It is noted, not corrected.

v4.9.0-r23 status in the three frames

category	M / T / P
F (7)	same in M and T; at P only where no port quantity enters
M (29)	valid in M; in T the inverse ratio, an undefined angle or a different equation; at P degenerate
C (12)	M as stated; T with magnitude by class and phase $(-1)^k$; at P no conversion defined
S (9)	statements at $\sigma = 0$ of the scale line; neither M nor T; not P
P (3)	statements about P itself, degenerate as (F2) requires
D, N (53)	define the frames, or carry no frame quantity

v4.9.0-r23 register entries

statement	status after v4.9.0-r23
frame-dependent statements of the main body	113 classified under Convention F (r23); list in `llm/k62_frame_regel_hauptkorpus.py`
tachyon of Thm. 20.6	$-\beta_{\mathrm{sym}}$ (real part); Convention F uses $1/\beta_{\mathrm{sym}}$ (Reading B); 18 statements frame-free only under the former
photon frame of the main body	$\beta_{\mathrm{sym}} = 0$, the scale-line reference; under F the pure state, not P
horizon, saturation limit	at the photon fixed point of F
local OP5 closure	no frame quantity; unchanged and local

v4.9.0-r23 ledger and guard update

$$R_{\mathrm{F}}^{v490\text{-}r23} = (0, 0, 1)$$

for (main body classified, discrepancies of tachyon and photon placement recorded, choice between the two tachyon placements for the even statements left to the author).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r23,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r23} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r24 Source-Only: Correction of the Three Discrepancies of r23 as Single-Frame Artefacts

Purpose. r23 recorded three discrepancies between the main body and Convention F: the sign of the matter offset, the position of the tachyon, and the position of the photon. On 2026-10-22 the author asked for a correction and pointed out that the main body worked in one frame only, which may be where the discrepancies come from. This layer shows that this is so for all three. It corrects them by typing, not by editing: the main body is not rewritten, and its statements receive the frame they were made in. The script is `llm/k63_widerspruche_ein_frame.py`. No prediction is made, and nothing is compared with data.
Source-only status. This v4.9.0-r24 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r24 the single frame of the main body

The main body describes everything from one frame and along one real axis. It calls that frame the photon frame and measures the offset β_{sym} from it. Three features of that frame matter here. First, it fixes an orientation on the axis. Second, it sees only real rapidities. Third, it is the zero of its own scale line. Each discrepancy of r23 comes from one of these.

v4.9.0-r24 the three corrections

Theoremlet v4.9.0-r24.1 (sign). The sign of β in the sector definition of §15 (matter $\beta < 0$) and in Theorem 20.6 (matter $+\beta_{\mathrm{sym}}$) is an orientation code in the sense of v4.8.0-r01 (§A.270). The two choices differ by the reflection $z \mapsto -z$, which is charge conjugation C (v4.9.0-r02). C reverses $\mathrm{Re}\, z$ and leaves $s = z \cdot z$ unchanged. The sector is fixed by the sign of s (v4.8.0-r07), so both choices assign the same sectors. The two passages use opposite orientation conventions. That is a conflict of notation, not of physics.

Correction. Read β of the sector definition and $\pm\beta_{\mathrm{sym}}$ of Theorem 20.6 as orientation codes, and read $\beta_{\mathrm{sym}} = |\beta| \geq 0$ as the magnitude of the offset. Matter and tachyon are told apart by the sign of s.

Theoremlet v4.9.0-r24.2 (tachyon). The tachyon rapidity is $-\eta + i\pi/2$, with $\tanh(-\eta + i\pi/2) = -\coth\eta = -1/\beta_{\mathrm{sym}}$. Projected onto the single real axis of the main body, dropping the imaginary part, it is $-\eta$, that is $\beta = -\beta_{\mathrm{sym}}$. This is the tachyon of Theorem 20.6. In its own frame the tachyon has $\beta_T^2 = 1/\beta_{\mathrm{sym}}^2$, which is the tachyon of Convention F. Both describe the same tachyon, the first as seen along the axis of the reference frame and the second in its own frame.

Correction. Convention F receives a fourth entry, $T|_R$, the tachyon seen from the reference frame of the main body, at $-\beta_{\mathrm{sym}}$. The 18 even statements of r23.1(a), among them Ω_M, $g_{tt} = -\beta_{\mathrm{sym}}^2$ and the Beltrami coefficient, hold for M and for $T|_R$ alike. They are frame-free in the description of the main body. In the tachyon's own frame they take the values at $1/\beta_{\mathrm{sym}}^2$ ($\Omega_M \to 3.31$). The two values are different typings, and neither contradicts the other.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Theoremlet v4.9.0-r24.3 (photon). The frame that the main body calls the photon frame, $\beta_{\mathrm{sym}} = 0$, is in the state disk the pure state $|b| = 1$, with entropy 0. The main-body conversion factor to it is the ratio of Bloch lengths:

$$\frac{|b_R|}{|b_M|} = \frac{1}{\operatorname{sech} 2\eta} = \cosh 2\eta = \gamma_L.$$

To the photon of Convention F, $b = 0$, the same factor would diverge. The reference frame of the main body is therefore the pure-state frame R, the zero of the scale line. The photon P of Convention F is the maximally mixed state, reached for $\beta \to 1$.

Proof sketch. r24.1: $s(-z) = s(z)$ and $\mathrm{Re}(-z) = -\mathrm{Re}\, z$, checked on 1000 random z. r24.2: $\tanh(x + i\pi/2) = \coth x$. r24.3: $|b| = \operatorname{sech} 2\eta$, equal to 1 at $\eta = 0$. Numerically $|b_R|/|b_M| = 1.86444 = \gamma_L$. □

Correction. Read "photon frame" in the main body, where it denotes the zero of the offset or the target of a γ_L^{-k} conversion, as the reference frame R. The nine S statements of r23 become R statements and hold there. $\Lambda = 0$ and the vanishing frame term hold in R, the frame where the doublet state is pure. At the photon P the frame term β_{sym}^2 and the ratio p_-/p_+ are maximal (1). This also explains why a finite γ_L points to $|b| = 1/\gamma_L$ and not to $b = 0$ (r17): the conversion goes from M to R, not from M to P.

v4.9.0-r24 Convention F completed

(F5) *Reference frame.* R is the pure state $|b| = 1$, $\beta_{\mathrm{sym}} = 0$, $\sigma = 0$. It is the frame of the main body. The conversion $X^{(R)} = \gamma_L^{-k} X^{(M)}$ of §25 goes from M to R.

(F6) *Projection.* $T|_R$ is the tachyon seen from R along the real axis: $\beta = -\beta_{\mathrm{sym}}$, imaginary part dropped. A statement of the main body about the tachyon is a statement about $T|_R$ unless it says otherwise.

(F7) *Orientation.* The sign of β is an orientation code. The sector is fixed by the sign of s.

The r23 classification becomes: S → R (valid in R); the 18 even statements are frame-free between M and $T|_R$ and matter-only in the tachyon's own frame; P keeps the three $\beta \to 1$ statements.

v4.9.0-r24 status in the three frames

result	M / T / P (and R, $T\|_R$)
r24.1 sign	orientation code in every frame; sector by the sign of s, unchanged under C
r24.2 tachyon	$T\|_R$ at $-\beta_{\mathrm{sym}}$, T at $1/\beta_{\mathrm{sym}}$; even statements equal in M and $T\|_R$, different in T
r24.3 photon	R is the pure state, P the maximally mixed state; S statements hold in R, not in P

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r24 register entries

statement	status after v4.9.0-r24
sign of the matter offset	orientation code; the two conventions differ by C; sector by s (r24.1)
tachyon of Thm. 20.6	$T\|_R$, the real projection of the tachyon onto the reference axis (r24.2)
photon frame of the main body	reference frame R, the pure state; conversions go from M to R (r24.3)
S statements of r23	valid in R
18 even statements of r23	frame-free between M and $T\|_R$; in the tachyon's own frame at $1/\beta_{\mathrm{sym}}^2$
discrepancies of r23	resolved as single-frame artefacts; no edit of the main body

v4.9.0-r24 ledger and guard update

$$R_{\mathrm{F}}^{v490\text{-}r24} = (0, 0, 0)$$

for (sign resolved, tachyon placement resolved, photon placement resolved). The open component of r23, the choice between the two tachyon placements, is closed: both are kept with their typings.

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r24,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r24} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r25 Source-Only: Revision of the Main Body

Purpose. On 2026-10-22 the author asked for a revision of the document. Earlier work should not be lost, but false statements should not keep standing where they proved to be dead ends. Until now the main body was never changed: refutations lived only in the rounds. This layer records the first revision of the main body. The script `llm/aufraeumen_v490.py` applies it. The state before the revision is kept under the tag `vor-aufraeumen-v490-r24`. The rounds of the solve log are not touched.
Source-only status. This v4.9.0-r25 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r25 the revision rule

(A) *Dead ends.* A statement that later layers refuted, excluded or withdrew is moved word for word to §A.270. At its place the same environment remains, with the same number and label, a short statement of what survives, and a pointer. Every earlier reference therefore still resolves.

(B) *Known errors.* A value or formula whose correct form a layer established is corrected in place. A framed status note gives the earlier wording and the layer that justifies the correction.

(C) *Reinterpreted or open statements.* The text stays. A framed status note gives the current status and the layers.

The rounds keep growing append-only. A later revision of the main body follows the same rule.

v4.9.0-r25 what was revised

(A) Archived, 15 **items.** The phase formula of the Koide theorem (Theorem 22.5; $K = 2/3$ stays as empirical fact); the derivation of GUE statistics (Theorem 22.20; GUE stays as empirical fact); the GUE self-determination and prediction P1 (Theorem 24.2, Corollary 24.6); the β_{sym}-formula P3 with its exponent, the self-consistent system and the claimed resolution of OP 1 (Remark 41.8, Proposition 41.24, Theorem 41.21, Corollary 41.23); the PMNS proposition (Proposition 42.12); the H_0 and saturation hypotheses (Propositions 42.5, 42.8); the bare Λ (Proposition 42.18); the Theta-sum conjecture and its remainder estimate (Proposition 42.33, Remark 42.35); the row "Riemann Hypothesis" of Theorem 22.22.

(B) Corrected. The vacuum ratio $\gamma_L^6 \to \gamma_L^5$ and the residual $10^{-120.4} \to 10^{-120.6}$ (Proposition 42.16, Corollary 42.17, the corollary "Cosmological constant revisited", P4 and the status table of §42.15; v4.8.0-r03 F1); P5 $\gamma_L^{-1} \approx 0.537 \to 1/2$ (v4.9.0-r09); the first row of Theorem 25.3, $k = 1$ for α (v4.8.0-r03 F2, v4.9.0-r08); the orthogonality in Proposition 16.1, which holds for the product and not for the average (v4.9.0-r03.4).

(C) Status notes. On the sector definition and Theorem 20.6 (orientation code, $T|_R$, v4.9.0-r24); on §25 (the photon frame there is the reference frame R); on Theorems 12.1, 15.2, 13.1, 14.3, 22.1, 22.19, 23.4, 20.5, 22.16, 41.18, 25.5, 41.26, 42.19; Definition 12.5; Proposition 21.4; Corollaries 22.21, 42.24; Remark 42.21.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Not revised. The rounds of the solve log, the pre-registration notes and their results, and the OP5 grammar. Statements whose status is a decision of the author (ways A–C, the epoch of Theorem 22.16) carry a note but keep their text.

v4.9.0-r25 status in the three frames

part	M / T / P
archived statements	no frame statement survives in the main text; the archive keeps the original wording
corrections	frame typing as in r23 and r24: conversions go from M to R, the tachyon sign follows (F3)
status notes	carry the frame classification of r23 where it applies

v4.9.0-r25 ledger and guard update

$$R_{\mathrm{rev}}^{v490\text{-}r25} = (0,0,0)$$

for (dead ends archived, known errors corrected, status notes placed). The revision removes the one place in the main body that listed the Riemann Hypothesis as a result; the guard against a global photon-clock/RH statement is thereby better kept, not spent.

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r25,+)} = (0,1,1,1,1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r25} = (1,1,1,1,1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r26 Source-Only: Pre-Registered Test of Degree-Six Potentials on the Triple Space

Purpose. Layer v4.8.0-r16 showed that invariant potentials up to degree four on the triple space of the triality slots give at most two distinct slot norms, and named degree six as necessary for three. This layer records the pre-registered test H16 of degree six (note `llm/Vorab_Notiz_H16_Grad6_Tripelraum.md`, commit `d0360b8`, fixed before computing; created under the label H12, which was already taken, and renamed without change of content in commit `6bd2f1b`). Earlier layers are not rewritten.
Source-only status. This v4.9.0-r26 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r26 setting and hypotheses

As in v4.8.0-r16: slot norms n_k, $x_k = n_k^2$, $S = \sum x_k$, $Q = \sum x_k^2$, $H = \sum x_k^3$, $P = n_1 n_2 n_3$, $t = \mathrm{Re}((ab)c) = P$ at the minimum, reading $\sqrt{m_k} \propto n_k$. Only H adds a term x_k^3 to the condition at critical points, so the fixed minimal extension is

$$V_6 = -S + \beta S^2 + \gamma Q - \kappa P + \delta H, \qquad \delta > 0,\ \kappa \geq 0.$$

H16a: for part of a fixed parameter distribution ($N = 2000$), V_6 has a global minimum with three distinct non-zero norms. H16b: for some parameters the charged-lepton triple is the global minimum. H16c: $K = 2/3$ is singled out among such minima (at least half within 0.01).

v4.9.0-r26 results

Script `llm/k64_test_h16.py`. Global minima over the distribution: 1585 with equal norms, 398 with one occupied slot, 17 with a degenerate pair, **none** with three distinct norms. The lepton triple is a critical point on the whole two-parameter family given by Vieta's relations, but on the 41×41 grid it is nowhere even a local minimum (1139 points with $\delta \leq 0$, 542 with an indefinite Hessian).

Verdict. H16a, H16b and H16c are refuted under the fixed criterion.

Theoremlet v4.9.0-r26.1 (found after the verdict). Every polynomial potential in the slot norms and P of degree at most six has the form $V = A(S,P) + e_2\,(b_0 + b_1 S)$ with $e_2 = \sum_{i<j} x_i x_j$. At a critical point with three distinct non-zero norms it is a saddle if $b_1 \neq 0$; if $b_0 = b_1 = 0$ it does not depend on e_2 and the norms are not fixed. Hence no potential of degree at most six has a minimum with three distinct norms; degree at least eight is necessary.

Proof sketch. $Q = S^2 - 2e_2$, $H = S^3 - 3Se_2 + 3P^2$, $SQ = S^3 - 2Se_2$; the other invariants S, S^2, S^3, P, SP, P^2 do not contain e_2. For distinct x_k the map $n \mapsto (S, e_2, P)$ is a local diffeomorphism (Vandermonde), and at a critical point the Hessian transforms by congruence. In these coordinates it reads $\left(\begin{smallmatrix} * & b_1 & * \\ b_1 & 0 & 0 \\ * & 0 & * \end{smallmatrix}\right)$, whose leading 2×2 minor is $-b_1^2 < 0$. For V_6, $b_1 = -3\delta$, and $\partial V/\partial e_2 = 0$ is the Vieta relation $2\gamma + 3\delta S = 0$. Checked exactly with 40 digits at 2000 random critical points of the general degree-six ansatz (script `llm/k65_sattel_grad6.py`); in double precision the negative eigenvalue can fall to $5 \cdot 10^{-15}$ of the largest. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Reclassification. The statement of v4.8.0-r16 that degree six is necessary holds for critical points; for minima degree six is not enough. Whether degree eight suffices is not shown. It adds at least four coefficients, so the two lepton ratios could at best be fitted, and $K = 2/3$ involves $\sum n_k$, which is not a polynomial invariant. Within this framework the generation S_3 with natural invariants does not produce a mass hierarchy; one needs an explicit breaking of S_3 or invariants beyond the norms, such as the real parts of the individual slots, which G_2 leaves fixed.

v4.9.0-r26 status in the three frames

result	M / T / P
H16 verdict, r26.1	frame-free: statements about octonion norms and polynomial invariants, no frame quantity enters
mass reading $\sqrt{m_k} \propto n_k$	matter frame M (charged-lepton masses, as the Koide phase in v4.9.0-r22); not evaluated in T or P

v4.9.0-r26 ledger and guard update

$$R_{\mathrm{cg}}^{v490\text{-}r26} = (0, 0, 1)$$

for (separation consistent with G_2; carrier of the triality slots identified; mass mechanism open, with all norm-invariant potentials up to degree six excluded).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r26,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r26} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r27 Source-Only: A Dynamics for the Kicked Duality Rotation, Rotation plus Compensator Measurement

Purpose. Reading B is kinematics: its transports (spatial and duality rotations, boosts) leave the invariant s fixed. r10.2 found that a witness of shape (iii) would have to change s, that is, carry dynamics. r16 recorded the author's reading of the C5 recursion as a kicked duality rotation read out at the compensator. On 2026-10-22 the author asked to attempt a dynamics. This layer proposes one on the state disk of r04 and r17: a duality rotation followed by a measurement in the compensator basis. It is a model, not a derivation from the earlier text. The script is `llm/k64_dynamik_kick_messung.py`. No new prediction is made, and nothing is compared with data.
Source-only status. This v4.9.0-r27 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r27 the model

The doublet state is $\rho = \frac{1}{2}(1+b_x\sigma_x+b_z\sigma_z)$, with the compensator axis along x, $C_{\mathrm{HC}} = \frac{1}{2}(1+\sigma_x)$, $A = \frac{1}{2}(1-\sigma_x)$ (r04). One step consists of two parts.

1. A duality rotation $U_\alpha = e^{-i\alpha\sigma_y}$: angle α on the doublet, angle 2α on the disk.
2. A measurement $\{C_{\mathrm{HC}}, A\}$. *Selective*: the state is kept in the compensator port, with survival probability $p = \mathrm{tr}(C_{\mathrm{HC}}U_\alpha\rho U_\alpha^{\mathsf{T}})$. *Non-selective*: $\rho \mapsto C_{\mathrm{HC}}\rho' C_{\mathrm{HC}} + A\rho' A$, the content of the other port going to the partner reservoir of r11.

Theoremlet v4.9.0-r27.1.

(a) Selective steps reproduce the C5 recursion. From a state on the compensator axis the survival per step is $\cos^2\alpha$, 3/4 at $\alpha = \pi/6$, and $(3/4)^n < 1/256$ first at $n = 20$.

(b) One non-selective step at $\alpha = \pi/6$ takes the pure reference state R ($|b| = 1$, v4.9.0-r24) to the Hopf state: $|b| = \cos 2\alpha = \frac{1}{2}$, $\gamma_L = 1/|b| = 2$, eigenvalues 3/4 and 1/4, entropy 0.811 bit (r17.1). The measured offset $|b| = 1/\gamma_L = 0.5364$ corresponds to the kick angle $\alpha = 28.78°$, with survival $\cos^2\alpha = 0.7682$, the lattice value of r13.

(c) Non-selective steps contract the Bloch vector: after the first step it lies on the compensator axis, and each further step multiplies its length by $|\cos 2\alpha|$. For $0 < \alpha < \pi/2$ every state flows to $b = 0$, the photon of Convention F. The entropy never decreases.

(d) The trace $2u$ is kept and $|b| = |s|/2u$ shrinks, so the step changes $|s|$. It therefore lies outside the transport group of Reading B, as r10.2 requires of a witness of shape (iii).

(e) In the limit of dense kicks, $\alpha = \omega\tau$ with $\tau \to 0$, the contraction rate of $|b|$ is $\Gamma = 2\omega^2\tau$, which tends to 0 (quantum Zeno effect). The continuous form is a dephasing in the compensator basis.

Proof sketch. (a) $C_{\mathrm{HC}}U_\alpha C_{\mathrm{HC}} = \cos\alpha\, C_{\mathrm{HC}}$ on the compensator ray (r10.1). (b), (c) the rotation turns b by 2α; the non-selective measurement removes the component across the axis, leaving $b_x\cos 2\alpha$. (c) the step is a unital channel, so the entropy cannot fall; checked on 20 000 steps from random states. (d) $\mathrm{tr}\,\rho$ fixed, $|b|$ reduced. (e) $-\ln\cos 2\omega\tau/\tau \to 2\omega^2\tau$, checked at $\tau = 0.2, 0.05, 0.01$. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Reading. The dynamics joins three earlier results. The C5 factor 3/4 is the survival of one selective step (r10). The Hopf point is the state after one non-selective step from R (r17). The lattice value 0.768 is the survival at a kick angle 1.22° below $\pi/6$ (r13). The photon is the attractor of the non-selective flow. It is reached kinematically only in the limit (r19.1, r17.2), but here it is approached through loss of information. The Zeno limit (e) is the pattern that r10 noted: denser compensation lets less content leave.

v4.9.0-r27 what the dynamics does not give

Theoremlet v4.9.0-r27.2.

(a) For $0 \le \alpha < \pi/2$ the selective survival $(\cos^2\alpha)^n$ is positive for every finite n. The compensator score does not vanish at a finite stage. It falls below the detector threshold 1/256 after $n = 20$, which is the permitted statement $C5_{\mathrm{det/eff}}$.

(b) The score is empty in one step only at $\alpha = \pi/2$. That is the quarter turn of r10.2, which moves the whole content into the partner port. r12.2(c) records that this relocation is blocked by $C4$ and $C6$.

(c) At $\alpha = \pi/4$ one non-selective step reaches $b = 0$ exactly. This is a statement about the doublet state. The compensator score there is 1/2, not 0. It is no clock statement and no global statement.

The model therefore supplies dynamics of the kind that r10.2 asked for, and it changes s. It does not supply a finite zero of the compensator score. Its time unit is the kick interval of r13, in clock units. It gives no unit in seconds and no value of the coupling κ of r09. What physically performs the measurement is not identified.

Proof sketch. (a) $\cos^2\alpha > 0$ for $\alpha < \pi/2$; $(3/4)^{19} > 1/256 > (3/4)^{20}$. (b) $\cos^2(\pi/2) = 0$. (c) $\cos(\pi/2) = 0$; survival $\cos^2(\pi/4) = 1/2$. □

v4.9.0-r27 status in the three frames

result	M / T / P
r27.1(a) survival 3/4	frame-free: in T the ports are exchanged by J_B, and the survival in the tachyon's compensator port is again 3/4
r27.1(b) Hopf state after one kick	in M from R; in T the image with $b \to -b$, same $\lvert b\rvert$
r27.1(c) flow to $b = 0$	frame-free in direction of $\lvert b\rvert$; P is the attractor, degenerate as (F2) requires
r27.1(d) change of s	in M and T alike; at P $s = 0$ and the step acts trivially
r27.2 limits	frame-free

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r27 register entries

statement	status after v4.9.0-r27
dynamics of Reading B	model proposed: duality rotation plus compensator measurement; changes s (r27.1)
C5 factor 3/4, $n = 20$	survival of selective steps at $\alpha = \pi/6$ (r27.1(a))
Hopf point	state after one non-selective step from R (r27.1(b))
lattice value 0.768	survival at kick angle 28.78° (r27.1(b))
photon	attractor of the non-selective flow (r27.1(c))
shape (iii)	the model changes s but gives no finite zero of the score (r27.2); open
clock unit, κ, measuring agent	open

v4.9.0-r27 ledger and guard update

$$R_{\mathrm{dyn}}^{v490\text{-}r27} = (0, 0, 1)$$

for (model stated, earlier results joined, physical agent and scale open).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r27,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r27} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r28 Source-Only: Who Measures? The Tachyon as Measuring Device, Self-Stabilisation of the Field, and a Correction of the Tachyon Map

Purpose. r27 left open what performs the compensator measurement. On 2026-10-22 the author asked whether the tachyon can be the measuring device, and whether the generating field may stabilise itself. This layer tests six variants fixed in advance in the script `llm/k65_messgeraet_tachyon_feld.py`, and reports one further check that was added after the fact. On the way it corrects the name of the tachyon map used since r21. No new prediction is made, and nothing is compared with data.
Source-only status. This v4.9.0-r28 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r28 correction: the tachyon map is the duality rotation $z \mapsto iz$

On the doublet, $J_B = \sigma_x$ and $C_{\mathrm{HC}} = \frac{1}{2}(1 + \sigma_x) = \frac{1}{2}(1 + J_B)$ (r04). So J_B leaves the compensator ports fixed, with eigenvalue $+1$ on C_{HC} and -1 on A. On the state disk it reverses only $b_z = \mathrm{Re}\, s/2u$. The map that r21, r22 (Convention F) and r27 attribute to the tachyon reverses the whole Bloch vector, $b \mapsto -b$, and exchanges the ports. That is the duality rotation $z \mapsto iz$ of v4.8.0-r07.2, acting on states as $\rho \mapsto \sigma_y \rho \sigma_y$. Both maps send matter to tachyon, since both reverse $\mathrm{Re}\, s$. J_B keeps the side $\mathrm{sgn}(E \cdot B)$ of r02, and $z \mapsto iz$ changes it.

Correction. In r21 (premise D2, the clock seed as image of $V3$), in r22 (Convention F, definition of T) and in r27 (frame table), read "J_B image" as "image under the duality rotation $z \mapsto iz$". Every property those layers use holds for that map: $b \mapsto -b$, exchange of $p_\pm$ between the ports, $x \mapsto 1/x$, $\gamma \mapsto -\gamma$, and $AR_\alpha A = \cos\alpha\, A$. No result of those layers changes. The earlier layers are not rewritten.

v4.9.0-r28 the tachyon as measuring device

A device measures the compensator ports when the system port controls an operation W on it. The coherence between the ports is then multiplied by $D = \mathrm{tr}(\rho_{\mathrm{dev}} W)$. The measurement is ideal when $D = 0$.

Theoremlet v4.9.0-r28.1.

(a) A tachyon in its reference state, on its compensator axis ($b = (-1, 0)$, the partner port of matter), is an ideal measuring device when it couples through time reversal $T = \sigma_z$: $D = 0$, and the port weights stay 3/4 and 1/4. Through J_B or through $C = -1$ it does not measure ($D = 1$), because its reference state is an eigenstate of both.

(b) A tachyon that is the image of the system itself measures only partly: $D = 0.50$ through J_B and $D = 0.87$ through T, for the state after one $\pi/6$ step from R.

(c) An exchange (partial swap, angle θ) of the system with its own tachyon image contracts the whole Bloch vector, $b \mapsto \cos 2\theta\, b$. This is the radial part of the non-selective step of r27 with $\theta = \alpha$. At $\theta = \pi/6$ one exchange takes R to the Hopf state. Repeated with the $\pi/6$ rotation, every state flows to the photon $b = 0$.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof sketch. (a), (b) controlled-W coupling and partial trace; numerically D as stated. (c) b and $-b$ commute, so the partial swap of $\rho \otimes \sigma_y\rho\sigma_y$ gives $\cos^2\theta\,\rho + \sin^2\theta\,\sigma_y\rho\sigma_y$. □

Reading. The tachyon can be the measuring device. Two conditions apply: it must start in its reference state, and it must couple through time reversal. That matches r02, where the round trip through the tachyon runs through T. Matter that exchanges content with its own tachyon image measures itself, with no outside apparatus. This self-measurement drives the state to the photon, as in r27.

v4.9.0-r28 does the field stabilise itself?

Theoremlet v4.9.0-r28.2.

(a) *Field alone, two frequencies* (clock model of r15, kick interval one sixth of the beat, r13). The port coherence falls to $|D|^2 = 3/4$ after one interval. This is the same angle $\pi/6$ as in r13, not an independent confirmation. It returns to 1 after six intervals. The process is reversible and does not measure.

(b) *Field alone, many frequencies* (40 heights of zeros as clock frequencies, clock model only). The coherence falls to about $N^{-1/2}$ (mean 0.14 against 0.16) with partial revivals up to 0.41. The field dephases itself only approximately. An exact self-measurement would need the limit $N \to \infty$.

(c) *Field with a replenished tachyon reservoir.* After each $\pi/6$ rotation the system exchanges with a fresh tachyon in its reference state, at angle θ. The state then has a unique attractor, reached from every start, on the partner-port axis:

$$|b|_\infty = \frac{\sin^2\theta}{1 - \cos^2\theta\,\cos 2\alpha}.$$

At $\alpha = \pi/6$ this gives 0.126, 0.400, 0.667 and 0.857 for $\theta = 15°$, $30°$, $45°$ and $60°$. This is a self-stabilised state with $0 < |b| < 1$. It exists only while the reservoir is replenished.

Proof sketch. (a) $|\frac{1}{2}(1 + e^{ik\pi/3})|^2$. (b) $K = |N^{-1}\sum e^{i\gamma_m k\tau}|$ over $k = 50, \ldots, 5000$. (c) iteration from five starts, spread below 10^{-10}; the closed form checked symbolically at the four angles. □

A check added after the fact. If the exchange strength is set equal to the offset, $\sin^2\theta = \beta_{\mathrm{sym}}^2 = x$, the attractor in (c) is self-consistent, in the sense that $|b|_\infty = (1-x)/(1+x) = 1/\gamma$, exactly when $3x^2 + 2x - 1 = 0$, that is $x = 1/3$, the Hopf point. At the measured values it is not: $|b|_\infty = 0.4825$ against $1/\gamma_L = 0.5364$. This check was not fixed in advance. It joins the other results that hold exactly at the Hopf point only, and it is recorded as an observation.

Answer to the author's questions. The tachyon can be the measuring device, if it couples through time reversal from its reference state. The field alone does not stabilise itself: with two frequencies it oscillates back, and with many it dephases towards the photon. A stable state between the pure state and the photon arises only in exchange with a tachyon reservoir that keeps being replenished, that is, in an open system. At the Hopf point that state is self-consistent.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r28 status in the three frames

result	M / T / P
correction of the tachyon map	concerns the definition of T; M and P unchanged
r28.1 tachyon as meter	a statement about the M–T pair; at P the ports carry equal weight, and there is nothing to measure
r28.2(a), (b) self-dephasing	frame-free between M and T (only $\|b\|$ enters); P is the limit
r28.2(c) attractor	on the partner-port axis of M, the compensator axis of T; not at P for $\theta > 0$

v4.9.0-r28 register entries

statement	status after v4.9.0-r28
tachyon map in r21, r22, r27	the duality rotation $z \mapsto iz$, not J_B; results unchanged (correction)
tachyon as measuring device	ideal with T-coupling from its reference state; none with J_B or C (r28.1)
self-measurement	exchange with the own tachyon image reproduces the radial step of r27 (r28.1(c))
self-stabilisation	not for the closed field; yes with a replenished tachyon reservoir, unique attractor (r28.2)
Hopf point	self-consistent attractor for $\sin^2\theta = \beta_{\mathrm{sym}}^2$ (post hoc)

v4.9.0-r28 ledger and guard update

$$R_{\mathrm{meas}}^{v490\text{-}r28} = (0, 0, 1)$$

for (meter identified under stated coupling, self-stabilisation classified, the physical source of the replenished reservoir open).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r28,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r28} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r29 Source-Only: Where the Tachyon Reservoir Could Come From

Purpose. r28 found that the field stabilises between the pure state and the photon only in exchange with a tachyon reservoir that keeps being replenished. On 2026-10-22 the author asked where that reservoir could come from. This layer tests six candidates that the document itself offers. Candidates and criteria were fixed in advance in `llm/k66_tachyon_reservoir_quelle.py`. No new prediction is made, and nothing is compared with data.
Source-only status. This v4.9.0-r29 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r29 criteria and candidates

A source must (i) supply fresh tachyons again and again, (ii) supply them in the tachyon reference state, polarised on the partner-port axis, $b_T = (-1, 0)$, (iii) exchange with the system coherently, and (iv) imply no forbidden statement. The candidates are: (K1) pairs at a horizon (Theorems 21.3, 21.7); (K2) virtual pairs from the vacuum (Proposition 20.2); (K3) the cascade flow of Theorem 22.16, $P(S^8 \to S^4) = \beta_{\mathrm{sym}}^2$; (K4) the other modes of the field; (K5) a thermal bath through the Wick rotation (Theorem 20.3); (K6) the tachyon hit chain of r06.

v4.9.0-r29 results

Theoremlet v4.9.0-r29.1.

(a) *The vacuum supplies tachyons in the reference state.* By r24 the vacuum $z^- = 0$ of the main body is the reference frame R, the pure state on the compensator axis. Its image under the tachyon map $z \mapsto iz$ (r28) is exactly $b_T = (-1, 0)$. A matter–tachyon pair created from the vacuum therefore hands over a tachyon in its reference state. K2 meets (ii).

(b) *Horizons do not.* The horizon $\beta \to 1$ is the photon fixed point of Convention F (r23). There $b_T = 0$, and the attractor of r28.2(c) is $|b| = 0$: no stabilisation. In Theorem 21.3 the tachyon also falls inwards, away from an outside system. K1 fails (ii).

(c) *The other modes and a warm bath only partly.* Other modes dephase (r28.2(b)) and carry no polarisation, so the attractor is 0 (K4 fails (ii)). A thermal bath has $|b_T| < 1$, and the attractor falls with it: $|b| = 0.500, 0.193, 0$ for $|b_T| = 1, 0.5, 0$ at $\sin^2\theta = 1/3$. The zero-temperature limit is the vacuum of (a) (K5 reduces to K2).

(d) *The hit chain distributes, it does not replenish.* One tachyon meets many matter systems at once (r06). After the first meeting it is no longer in its reference state. The pair event at the point of infinite speed (r06) is a pair creation, that is K2.

(e) *The cascade fixes the strength, if the exchange is coherent.* With the exchange strength $\sin^2\theta = \beta_{\mathrm{sym}}^2$, the cascade probability, the coherent exchange of r28 has the self-consistent attractor $|b| = 1/2$ exactly at the Hopf point. Read instead as a Markov replacement, $\rho \mapsto (1-p)\rho + p\rho_T$ with $p = \beta_{\mathrm{sym}}^2$, the attractor is 0.378 and lies off the axis (41°). It is self-consistent only at $p = 0.3865$, which corresponds to no point named in the document.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof sketch. (a) $\sigma_y \rho_R \sigma_y$ with $\rho_R = \frac{1}{2}(1 + \sigma_x)$. (b), (c) iteration of rotation $\pi/6$ plus exchange to the fixed point, for the stated b_T. (e) the same for the coherent partial swap and for the Markov replacement; root of the self-consistency condition by bisection. □

Bookkeeping. At the attractor each kick leaves the incoming tachyon changed, from $b_T = (-1, 0)$ to $(-0.750, 0.144)$. The reservoir takes up polarisation and must be replaced. The exported quarter of r11 goes there. A steady state therefore needs one fresh vacuum pair per kick. The rate of pair creation is not fixed by anything in the present layers.

Reading. Of the six candidates, only the vacuum supplies tachyons in the right state. The proposed source is the pair creation from the vacuum: the matter partner of each pair continues, and the tachyon partner serves once as the measuring device and the reservoir, coupling through time reversal (r28.1). The cascade fits the exchange strength β^2_{sym}, provided the exchange is coherent and not a classical replacement. In this picture the matter system holds its state because the vacuum keeps supplying fresh partners. Three points stay open: the rate of pair creation, why the exchange should be coherent, and where the absorbed polarisation of the used tachyons goes.

v4.9.0-r29 status in the three frames

result	M / T / P
r29.1(a) vacuum pairs	the vacuum is R (pure); the pair has its matter member in M and its tachyon member in T, in the reference state
r29.1(b) horizon	lies at P, where the reservoir is unpolarised
r29.1(c), (d)	frame-free statements about polarisation; P gives 0
r29.1(e) cascade strength	β^2_{sym} is a matter-frame quantity (r22); the self-consistency holds at the Hopf point only

v4.9.0-r29 register entries

statement	status after v4.9.0-r29
source of the tachyon reservoir	proposed: pair creation from the vacuum R (r29.1(a)); horizon, other modes, hit chain excluded; warm bath only partial
exchange strength	β^2_{sym} from the cascade, if coherent; Markov reading not self-consistent at a named point (r29.1(e))
open	rate of pair creation, coherence of the exchange, fate of the used tachyons

v4.9.0-r29 ledger and guard update

$$R^{v490\text{-}r29}_{\mathrm{res}} = (0, 0, 1)$$

for (candidates tested, vacuum pairs proposed, rate and coherence open).

The layer spends no guard:

$$R^{(v490\text{-}r29,+)}_{\mathrm{glob}} = (0, 1, 1, 1, 1), \qquad \Gamma^{v490\text{-}r29}_{\mathrm{guard}} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r30 Source-Only: The Compensator Avoids the Classical One-Half, and Why the Exchange Is Coherent

Purpose. r29 left open why the exchange between matter and tachyon should be coherent rather than a classical replacement. On 2026-10-22 the author pointed out that the Heisenberg compensator was introduced precisely to avoid the classical probability one half of finding the object in one or the other sector. This layer makes that precise, checks it against the document's own use of the compensator, and draws the consequence for r28 and r29. The script is `llm/k67_kompensator_kohaerenz.py`. No new prediction is made, and nothing is compared with data.
Source-only status. This v4.9.0-r30 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r30 classical one-half against the compensator

Theoremlet v4.9.0-r30.1.

(a) A classical mixture of a state with its tachyon image, $\frac{1}{2}\rho + \frac{1}{2}\sigma_y\rho\sigma_y$, puts weight 1/2 into the compensator port for every state. This is the classical one half.

(b) A coherent combination of the amplitudes f and $e^{i\varphi}f_T$ puts a weight into the compensator port that depends on the relative phase. It is 1/2 only at two phases, and it is 0 or 1, a definite outcome, at the two others.

(c) The document's own use of the compensator is coherent. The remark "Light as the Heisenberg compensator of the universe" of the main body computes $C(\beta_{\mathrm{sym}}) = \frac{1}{2}(\beta_{\mathrm{sym}} - \beta_{\mathrm{sym}}) = 0$. The cancellation exists only for amplitudes. A mixture of probabilities gives $\frac{1}{2}\beta_{\mathrm{sym}}^2 + \frac{1}{2}\beta_{\mathrm{sym}}^2 = \beta_{\mathrm{sym}}^2 = 0.302$, never 0.

Proof sketch. (a) $\sigma_y C_{\mathrm{HC}}\sigma_y = A$, so the mixture has weight $\frac{1}{2}(\mathrm{tr}\, C_{\mathrm{HC}}\rho + \mathrm{tr}\, A\rho) = \frac{1}{2}$; checked on 1000 random states. (b) numerical at $\varphi = 0, \pi/2, \pi, 3\pi/2$: $1/2, 0, 1/2, 1$. (c) direct. □

Reading. This confirms the author's account. The compensator acts on amplitudes. Where a classical description can only say "with probability one half here or there", the compensator gives a definite port, decided by the relative phase. The remark on light uses exactly this property.

v4.9.0-r30 the compensator partner is the tachyon seen through time reversal

Theoremlet v4.9.0-r30.2.

(a) $C = \frac{1}{2}(1 + J)$ is a projection only for an involution, $J^2 = 1$. This holds for $J_B = \sigma_x$. It fails for the tachyon map $z \mapsto iz$, whose square is -1 on amplitudes; there $C^2 = J/2$.

(b) J_B is the tachyon map followed by time reversal: $J_B = R_{\pi/2} \circ T$, with $T\colon z \mapsto \bar{z}$. On the doublet, $\left(\begin{smallmatrix} 0 & -1 \\ 1 & 0 \end{smallmatrix}\right)\mathrm{diag}(1, -1) = \sigma_x$.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(c) So the compensator combines a state with its tachyon partner seen through time reversal. r28.1(a) found that the tachyon measures ideally exactly when it couples through T. The two results are one statement: the coupling through T turns the tachyon into the compensator partner.

Proof sketch. (a) $(\frac{1}{2}(1+J))^2 = \frac{1}{4}(1+2J+J^2)$. (b) matrix product; $z \mapsto i\bar{z}$ as in r02. (c) combine (b) with r28.1(a). □

v4.9.0-r30 consequence for the reservoir

Theoremlet v4.9.0-r30.3.

(a) If the exchange of r28 and r29 runs through the compensator, it is coherent by construction. The Markov replacement of r29.1(e) is the classical mixture that the compensator was introduced to avoid.

(b) With coherent exchange of strength β_{sym}^2 the attractor is self-consistent exactly at the Hopf point (r28, r29). The open point of r29, why the exchange is coherent, is answered under the author's account of the compensator.

(c) At equal strength the coherent attractor is sharper. At strength 1/2 after $\pi/6$ rotations, coherent exchange gives $|b| = 2/3$, port weights 1/6 and 5/6, entropy 0.32 bit. Markov replacement gives $|b| = 0.577$, port weights 1/4 and 3/4, entropy 0.74 bit.

Proof sketch. (a) definition of the compensator, r30.1. (b) r28.2, r29.1(e). (c) iteration to the fixed point. □

On the wording of Theorem 22.16. That theorem calls $\Omega_M = \beta_{\mathrm{sym}}^2$ and $\Omega_\Lambda = 1 - \beta_{\mathrm{sym}}^2$ Markov transition probabilities. Under the reading of this layer, β_{sym}^2 remains the weight of one step, $\sin^2\theta$. Successive steps are composed as amplitudes through the compensator, not as a classical Markov chain. The theorem keeps its text. A status note under the revision rule of r25 would record this reading. It is not placed here, and the decision lies with the author.

v4.9.0-r30 status in the three frames

result	M / T / P
r30.1 one half	the mixture of M with its T image gives 1/2; the coherent combination is definite at two phases; at P both give 1/2
r30.2 partner through T	relates M and T; J_B keeps the side, $z \mapsto iz$ changes it (r28)
r30.3 coherent reservoir	as r28.2(c) and r29; self-consistency at the Hopf point only

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r30 register entries

statement	status after v4.9.0-r30
purpose of the compensator	to replace the classical one half by a definite port (author); confirmed by the main body's remark on light (r30.1)
compensator partner	the tachyon seen through time reversal, $J_B = R_{\pi/2} \circ T$ (r30.2)
coherence of the exchange (r29)	answered: coherent by construction of the compensator (r30.3)
Markov wording of Thm. 22.16	read as one-step weight; status note possible, decision with the author

v4.9.0-r30 ledger and guard update

$$R_{\mathrm{coh}}^{v490\text{-}r30} = (0,0,0)$$

for (one-half avoidance made precise, partner identified, coherence of the exchange answered).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r30,+)} = (0,1,1,1,1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r30} = (1,1,1,1,1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r31 Source-Only: The Markov Cascade as a Test of Self-Stabilisation, and a Status Note to Theorem 22.16

Purpose. r30 noted that Theorem 22.16 calls Ω_M and Ω_Λ Markov transition probabilities, while the compensator composes amplitudes. On 2026-10-22 the author asked for a status note. He explained that the Markov chain was meant to predict the probabilities better and to test whether the cascade can stabilise itself. This layer runs that test on the chain of the proof and places the note under the revision rule of r25. The script is `llm/k68_markov_kaskade_stabil.py`. The note is applied by `llm/apply_patches.py` from `llm/revision_spec_v490_r31.py`, so that the GitHub candidate can receive the same text. No new prediction is made, and nothing is compared with data.
Source-only status. This v4.9.0-r31 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r31 the test

The proof of Theorem 22.16 uses the chain $S^8 \to S^4$ with probability β^2_{sym}, $S^8 \to S^2$ with $1-\beta^2_{\text{sym}}$, $S^4 \to S^2$ with 1, and S^2 absorbing.

Theoremlet v4.9.0-r31.1.

(a) As given, the chain does not stabilise. After three steps every distribution has reached S^2, and the only stationary distribution is $(p_2, p_4, p_8) = (1, 0, 0)$, the photon.

(b) With a return path $S^2 \to S^8$ of rate $r > 0$ the chain is irreducible and has the unique stationary distribution

$$(p_2, p_4, p_8) = \frac{(1,\ \beta^2_{\text{sym}} r,\ r)}{1 + r + \beta^2_{\text{sym}} r}.$$

Every start converges to it. At $\beta^2_{\text{sym}} = 1/3$, $r = 1/2$ the other eigenvalues have modulus 0.41, so the approach is a damped oscillation.

(c) In the stationary state the flux out of S^8 splits into β^2_{sym} towards S^4 and $1-\beta^2_{\text{sym}}$ towards S^2, for every return rate r. The fractions that the theorem names Ω_M and Ω_Λ are therefore stable flux fractions, independent of how fast the return is.

Proof sketch. Stationary equations $\pi Q = \pi$, $\sum \pi = 1$, solved symbolically; convergence checked from S^8 over 200 steps at $r = 1/10, 1/2, 1$; eigenvalues 1 and $-\frac{1}{4} \pm i\sqrt{15}/12$ at $\beta^2_{\text{sym}} = 1/3$, $r = 1/2$. □

Reading. The author's test gives a clear answer. The closed cascade does not stabilise itself; it ends in the photon. With a return path it does, and then the matter fraction β^2_{sym} is a stable property of the flow. This is the same pattern on the level of the strata as on the doublet in r27 and r28. Closed, everything flows to the photon. Replenished, a stable state arises. A natural return path is the pair creation from the vacuum of r29. The rate r remains open, but the fractions do not depend on it. The Markov chain is a tool for these probabilities between the strata. Within one step, amplitudes compose through the compensator (r30).

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r31 the status note

The status note under Theorem 22.16 is extended (rule C of r25). It keeps the earlier content and adds the author's purpose for the Markov chain, the result r31.1, and the reading of r30. The text of the theorem is unchanged.

v4.9.0-r31 status in the three frames

result	M / T / P
r31.1(a) absorption	the absorbing stratum S^2 is the photon ground state of the main body; the chain ends there in every frame description
r31.1(b), (c)	β^2_{sym} is a matter-frame quantity (r22); the stationary fractions hold in M, and are frame-free between M and $T\|_R$

v4.9.0-r31 register entries

statement	status after v4.9.0-r31
Markov cascade of Thm. 22.16	author's tool for probabilities and self-stabilisation; closed: no stable state besides the photon; with return path: unique stationary state, flux fractions β^2_{sym}, $1-\beta^2_{\mathrm{sym}}$ for every rate (r31.1)
status note to Thm. 22.16	extended (rule C of r25)
return path	proposed: pair creation from the vacuum (r29); rate open, fractions independent of it

v4.9.0-r31 ledger and guard update

$$R^{v490\text{-}r31}_{\mathrm{Mk}} = (0,0,1)$$

for (self-stabilisation of the cascade tested, status note placed, rate of the return path open). The layer spends no guard:

$$R^{(v490\text{-}r31,+)}_{\mathrm{glob}} = (0,1,1,1,1), \qquad \Gamma^{v490\text{-}r31}_{\mathrm{guard}} = (1,1,1,1,1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r32 Source-Only: Pre-Registered Test of Real-Part Invariants on the Triple Space

Purpose. Layer r26 showed that no potential in the slot norms up to degree six has a minimum with three distinct norms. The real parts $r_k = \operatorname{Re} a_k$ of the three slots are invariant under G_2 one by one and add invariants that the norms do not see, among them $L = \sum r_k$, which equals $\sum n_k$ on the real axis. This layer records the pre-registered test H17 (note `llm/Vorab_Notiz_H17_Realteile_Tripelraum.md`, commit `2a7271a`, fixed before computing; the author's computer was not reachable, so the note was committed in the working container and sent to the author before the computation). Earlier layers are not rewritten.
Source-only status. This v4.9.0-r32 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r32 setting and hypotheses

With $R_2 = \sum r_k^2$ and the notation of r26, the fixed family adds all real-part invariants up to degree two to the potential of v4.8.0-r16:

$$V_R = -S + \beta S^2 + \gamma Q - \kappa t - \mu L - \nu R_2 - \lambda L^2,$$

bounded below if and only if $\beta > 0$ and $\beta + \gamma > 0$. It is invariant under G_2 (diagonal) and the generation S_3. H17a: for part of a fixed distribution ($N = 2000$) the global minimum over all 24 real coordinates of $\mathbb{O}^3$ has three distinct non-zero norms. H17b: for some parameters the charged-lepton triple is the global minimum. H17c: $K = 2/3$ is singled out among such minima (at least half within 0.01).

v4.9.0-r32 results

Script `llm/k69_test_h17.py`. Global minima over the distribution: 1483 with equal norms, 517 with a degenerate pair, **none** with three distinct norms; 1629 of the 2000 lie on the real axis, where G_2 is unbroken. For the lepton triple, the family given by Vieta's relations on the real axis was scanned on a $21 \times 4 \times 7 \times 3$ grid: 696 points unbounded, 1061 with a negative Hessian eigenvalue, 7 local but not global, **none** global.

Verdict. H17a, H17b and H17c are refuted under the fixed criterion.

Theoremlet v4.9.0-r32.1 (found after the verdict). On the real axis $a_k = n_k$ one has $t = e_3 = n_1 n_2 n_3$, $R_2 = S$, $L = e_1$, and

$$V_R = A(e_1, e_2) + e_3\,(4\gamma e_1 - \kappa).$$

At a critical point on the axis with three distinct norms the Hessian in the coordinates (e_1, e_2, e_3) has a zero in position $(3,3)$ and the entry 4γ in position $(1,3)$; for $\gamma \neq 0$ it is a saddle in $\mathbb{O}^3$. More generally, every symmetric polynomial in n of degree at most five is linear in e_3, so its critical points with three distinct values are saddles unless the coefficient of e_3 is stationary there.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof sketch. $Q = e_1^4 - 4e_1^2e_2 + 2e_2^2 + 4e_1e_3$ and $S = e_1^2 - 2e_2$; e_3^2 has degree six. For distinct n_k the map $n \mapsto (e_1, e_2, e_3)$ is a local diffeomorphism, and at a critical point the Hessian transforms by congruence. The 2×2 minor in (e_1, e_3) is $-16\gamma^2 < 0$. A saddle of the restriction to the axis is a saddle of the full potential. Checked symbolically, and with 50 digits at the seven grid points that passed the numerical test for a local minimum: each has a negative eigenvalue of about 10^{-9} relative to the largest, below the fixed tolerance 10^{-8} (script `llm/k70_achse_e3.py`). □

Reading. The real parts move the obstruction of r26.1 from e_2 to e_3; they do not remove it. On the axis a minimum with three distinct norms generically needs degree at least six, without real parts degree at least eight, each with several free coefficients, so the lepton ratios could at best be fitted. Most minima lie on the real axis and leave G_2 unbroken; the colour group is not selected there. Off the axis no corresponding statement is shown; numerically no minimum with three distinct norms occurred. Within natural $G_2 \times S_3$ invariants the generation symmetry gives at most two distinct masses; the next testable route is an explicit breaking of S_3.

v4.9.0-r32 status in the three frames

result	M / T / P
H17 verdict, r32.1	frame-free: statements about octonions and polynomial invariants, no frame quantity enters
mass reading $\sqrt{m_k} \propto n_k$	matter frame M (charged-lepton masses, as the Koide phase in v4.9.0-r22); not evaluated in T or P

v4.9.0-r32 register entries

statement	status after v4.9.0-r32
mass hierarchy from natural invariants on the triple space (v4.8.0-r16, r26)	refuted also with real-part invariants up to degree two (H17); obstruction moved from e_2 to e_3 (r32.1)
generations as triality slots, colour $SU(3) \subset G_2$ (v4.8.0-r14, r15)	unchanged as a proposal; most minima here leave G_2 unbroken, so the colour group is not selected by them
mass mechanism	open; next testable route: explicit breaking of the generation S_3 with a pre-registered parameter budget

v4.9.0-r32 ledger and guard update

$$R_{\mathrm{cg}}^{v490\text{-}r32} = (0, 0, 1)$$

for (separation consistent with G_2; carrier of the triality slots identified; mass mechanism open, with norm and real-part invariants of low degree excluded).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r32,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r32} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r33 Source-Only: Left-Invariant Labels on the Triple Space and a Pre-Registered Test

Purpose. Layer r32 closed the route of natural potentials in the slot norms and real parts. On the author's request the Dixon algebra $\mathbb{C}\otimes\mathbb{H}\otimes\mathbb{O}$ was then read as the storage for all configurations that the Companion intended it to be (the HS(6) carrier of the Dixon-carrier coding layer), and the generation triple was placed in it. This layer records what follows for the generation triple, a resolution of the conflict with the Lorentz spinor, and the pre-registered test H18 (note `llm/Vorab_Notiz_H18_Links_invariante_Massen.md`, commit `6130814`, fixed before computing). The algebraic statements are proved in Lean (`formal/QSFormal/Dixon.lean` and `Lesarten.lean`, Lean 4.35.0, no Mathlib import, standard axioms only). Earlier layers are not rewritten.
Source-only status. This v4.9.0-r33 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r33 the triple in the Dixon storage

Theoremlet v4.9.0-r33.1.

(a) $\mathbb{H}\otimes\mathbb{O} = \mathbb{O}\oplus(\mathrm{Im}\,\mathbb{H})\otimes\mathbb{O}$, and $(\mathrm{Im}\,\mathbb{H})\otimes\mathbb{O} = i\otimes\mathbb{O}\oplus j\otimes\mathbb{O}\oplus k\otimes\mathbb{O}$ carries the triple of v4.8.0-r14/r15: 24 of the 64 channels, with the coloured states as a 9-channel submask. G_2 acts on $\mathbb{O}$, $\mathrm{Aut}(\mathbb{H}) = SO(3)$ on $\mathbb{H}$; they commute.

(b) Of the 24 signed permutations of the slots in $SO(3)$, exactly the 12 even permutations with an even number of sign changes leave $\mathrm{Re}((xy)z)$ invariant; they are the conjugations by the binary tetrahedral group, so the joint symmetry is A_4. The transpositions of the triality S_3 need octonion conjugation, an anti-automorphism.

(c) If the $\mathbb{C}$ factor is read as the amplitude plane of v4.8.0-r02, then $J_B = R_{\pi/2}\circ T$ (v4.9.0-r30.2(b)) exchanges its two halves; the slot norms are invariant under T, $R_{\pi/2}$ and J_B, and the triality form goes to $-i\,\bar{t}$. Of the rotations of the $\mathbb{C}$ plane only the quarter steps, together with the reflections (the group D_4), keep the label of the two halves.

Proof sketch. Finite checks and polynomial identities over $\mathbb{Z}$, formalised in `Dixon.lean` (`inv_iff`, `tripleMask_popcount`, `jB_eq_swap`, `tC_JB`) and `Lesarten.lean` (`d4_length`, `conj_cyc`, `conj_i`). □

v4.9.0-r33 left-invariant labels

Theoremlet v4.9.0-r33.2. Write the triple as eight quaternions h_c, one for each octonion component, and let the Lorentz spinor act from the left, $h_c \mapsto q h_c$. The quaternionic Gram matrix $\overline{h_c}\,h_d$ is invariant (up to $|q|^2$; with the complex bilinear form also under boosts). Its real part is symmetric in the slots. Its three imaginary components are three invariant labels on the right. For the pure triple the i component is $-x_j\wedge x_k$, so

$$\mu_i = 2\big(|x_j|^2|x_k|^2 - \langle x_j, x_k\rangle^2\big),$$

twice the area spanned by the other two slots. The right action of $1+i+j+k$ permutes the three labels cyclically. For orthogonal slots $\sqrt{\mu_i/2} = n_j n_k = P/n_i$.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof sketch. Associativity of $\mathbb{H}$ gives $\overline{qx}\,qy = |q|^2\,\bar{x}y$ and $\overline{xr}\,yr = \bar{r}(\bar{x}y)r$; for imaginary u, v, $\bar{u}v = (u\cdot v, -u\times v)$; the Lagrange identity in $\mathbb{R}^8$ gives μ_i. Lean: `gram_left`, `gram_right`, `right_cyc`, `gram_pure`, `lagrange8`; numerical check `llm/k72_links_invariant.py`. □
With Lorentz acting from the left the single slot norms are not invariant (a rotation redistributes them), so the conflict of the Dixon check with the spinor reading of Theorem 22.4 can be resolved for the labels by reading the generations on the right; it is not resolved for masses (see H18).

v4.9.0-r33 pre-registered test H18

H18: masses $m_k \propto \sqrt{\mu_k}$ at the global minima of the potentials already tested (family A: v4.8.0-r16; family B: r32), with a fixed parameter grid and distribution. H18a: some point gives determined, distinct, non-zero μ_k; H18b: one of them meets both charged-lepton ratios within 1%. Script `llm/k73_test_h18.py`: family A, 205 points, all undetermined; family B, 500 points, 428 undetermined, 51 with equal and 21 with two distinct values. **H18a and H18b are refuted.** Check after the verdict (`llm/k74_nachkontrolle_h18.py`): in family A the angle between two slots is a flat direction of the minimum, because V sees the angles only through t; in family B the undetermined cases have $\mu \approx 0$, the slots lying parallel on the real axis. The labels resolve the Lorentz conflict but inherit the selection problem: a mass hierarchy would need a principle that fixes the angles between the slots.

v4.9.0-r33 status in the three frames

result	M / T / P
r33.1(a), (b)	algebraic, shown in the reference frame R in which the Dixon grammar is written (class N of r23)
r33.1(c)	under the reading of the $\mathbb{C}$ factor as amplitude plane: J_B exchanges the halves (M $\leftrightarrow$ T); slot norms frame-free; P has equal weight on both halves
r33.2, H18	frame-free in the slots; the $\mathbb{C}$ factor does not enter

v4.9.0-r33 register entries

statement	status after v4.9.0-r33
Dixon-carrier coding (HS(6))	read as the storage for all configurations (author's intention); holds the generation triple; masks conservative
mixed readouts (guarded, binary masks)	defined, never evaluated; r33 does not evaluate one either
Theorem 22.4	colour and generation on the same triplet (r13); spinor reading in conflict with generations in $\mathrm{Im}\,\mathbb{H}$ unless the generations are read on the right (r33.2)
mass hierarchy on the triple space	refuted also through left-invariant labels (H18); needs a principle that fixes the angles between the slots

v4.9.0-r33 ledger and guard update

$$R_{\mathrm{cg}}^{v490\text{-}r33} = (0, 0, 1)$$

for (separation consistent with G_2; carrier and storage of the triality slots identified, Lorentz-compatible labels found; mass mechanism open).

© 2026 Marc Brendecke. Preprint — not peer reviewed.

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r33,+)} = (0,1,1,1,1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r33} = (1,1,1,1,1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r34 Source-Only: Charges of One Generation from the Dixon Storage

Purpose. An inventory of the standard-model features against the structure (`llm/Bestandsaufnahme_Struktur_Standardmodell.md`) found that the Companion derives no charges. This layer records the pre-registered test H19 (note `llm/Vorab_Notiz_H19_Ladungen_Dixon.md`, commit `1e60410`, fixed before computing): does the charge construction from a number operator on $\mathbb{C}\otimes\mathbb{O}$ (C. Furey, Phys. Lett. B, 2015) work with the colour group of the Companion, and what do the frames do with the charges? Earlier layers are not rewritten.
Source-only status. This v4.9.0-r34 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r34 construction

Let L_a be left multiplication by e_a on $\mathbb{O}$ (Cayley–Dickson convention of v4.8.0-r12), and let i be the $\mathbb{C}$ factor of the Dixon storage, read as the quantum-mechanical unit. The colour group is $SU(3) = \mathrm{Stab}_{G_2}(e_7)$ as in Theorem 18.2; in this convention $e_7e_4 = -e_3$, $e_7e_5 = -e_2$, $e_7e_6 = e_1$. Put

$$\alpha_a = \tfrac{1}{2}\big(L_{e_a} + i\,L_{e_7e_a}\big) \quad (a = 4,5,6), \qquad N = \sum_a \alpha_a^\dagger\alpha_a, \qquad Q = N/3.$$

The Clifford relations of the L_a give $\alpha_a^2 = 0$ and $\{\alpha_a, \alpha_b^\dagger\} = \delta_{ab}$, so N is a number operator with eigenvalues $0,1,2,3$ of multiplicities $1,3,3,1$. These multiplicities are forced by the algebra and are not a prediction.

v4.9.0-r34 results

Theoremlet v4.9.0-r34.1.

(a) Every element of $\mathrm{Stab}_{G_2}(e_7)$ commutes with Q.

(b) The eigenspaces $Q = 0$ and $Q = 1$ are colour singlets in $\mathrm{span}_{\mathbb{C}}(1, e_7)$, spanned by $1 \pm i\,e_7$. The eigenspaces $Q = \frac{1}{3}$ and $Q = \frac{2}{3}$ are irreducible colour triplets in $\mathrm{span}_{\mathbb{C}}(e_1,\dots,e_6)$, conjugate to each other and not self-conjugate.

(c) Neither the real span of e_4, e_5, e_6 nor e_7 alone is an eigenspace of Q; the eigenstates combine e_a with e_7e_a, and 1 with e_7.

(d) On the $\mathbb{C}$ factor, $TQT^{-1} = 1 - Q$ and $J_BQJ_B^{-1} = 1 - Q$ for $J_B = R_{\pi/2}\circ T$, while $R_{\pi/2}QR_{\pi/2}^{-1} = Q$.

Proof sketch. (a)–(c) by explicit matrices on $\mathbb{C}^8$: the derivations of $\mathbb{O}$ form a 14-dimensional space, those with $D(e_7) = 0$ an 8-dimensional one; commutators with Q, invariance of the eigenspaces, commutants and characters were computed (script `llm/k75_test_h19.py`, deviations at most 10^{-14}). (d) T and J_B are antilinear, so they conjugate Q to $\bar{Q}$; T sends α_a to $-\alpha_a^\dagger$, hence $\bar{N} = 3 - N$. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Verdict and reading. H19a (a), H19b (b) and H19d (d) are confirmed; H19c, the literal assignment of the Companion, is refuted; H19e (charges equal on the three slots of $(\mathrm{Im}\,\mathbb{H})\otimes\mathbb{O}$) holds by the tensor structure. (a) and (b) reproduce Furey's result in the Companion's convention: the charge pattern of one generation, two colourless states with $Q = 0, 1$ and two conjugate colour triplets with $Q = \frac{1}{3}, \frac{2}{3}$ (in Furey's reading $\nu, \bar{d}, u, e^+$), follows from the storage. This is a consistency check, not a prediction about nature. (c) shows that Theorem 18.2 and Theorem 22.4 place colour and ν_R on the wrong, real, subspaces. (d) bears on the author's remark that Theorem 18.3 was developed in one frame: the frame change exchanges the charge sectors, $Q \leftrightarrow 1 - Q$. The spectrum and the colour structure are frame-free; which state carries which charge is not. The map is not charge conjugation $Q \mapsto -Q$ but charge conjugation followed by a shift of one unit, and it comes from the time-reversal part T of J_B. A charge statement such as the $U(1)$ line of Theorem 18.3 therefore has to name its frame. Hypercharge and chirality are not addressed.

v4.9.0-r34 status in the three frames

result	M / T / P
r34.1(a), (b), (c)	stated in M, with the $\mathbb{C}$ factor read as the quantum-mechanical unit; the spectrum and the colour structure are the same in T
r34.1(d)	relates M and T: sectors exchanged, $Q \leftrightarrow 1 - Q$; P (equal weight on both halves) is not evaluated

v4.9.0-r34 register entries

statement	status after v4.9.0-r34
charges of one generation	follow from the Dixon storage under the reading of the $\mathbb{C}$ factor as quantum-mechanical unit (r34.1(a), (b)); frame-dependent assignment (r34.1(d))
Theorem 18.2, Theorem 22.4	colour on the real span of e_4, e_5, e_6 and ν_R on e_7 refuted (r34.1(c)); candidate for a status note
Theorem 18.3	charge line must name its frame (r34.1(d)); hypercharge and chirality open

v4.9.0-r34 ledger and guard update

$$R_{\mathrm{ch}}^{v490\text{-}r34} = (0, 0, 1)$$

for (charge pattern of one generation consistent with the colour group; frame dependence of the charge assignment fixed; hypercharge and chirality open).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r34,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r34} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r35 Source-Only: Inner and Outer Complex Unit in the Colour Assignment

Purpose. Layer v4.9.0-r34 refuted the literal colour and ν_R assignment of Theorem 18.2 and Theorem 22.4 (r34.1(c)). At the author's suggestion that this may again be a fixed-frame effect, this layer records the pre-registered test H20 (note `llm/Vorab_Notiz_H20_Frame_Zuordnung.md`, commit `7d95feb`, fixed before computing). It separates two complex units: the *inner* unit $J = L_{e_7}$ on $\mathbb{O}$ itself, and the *outer* unit i, the $\mathbb{C}$ factor of the Dixon storage, on which the charge operator Q of r34 is built. Earlier layers are not rewritten.
Source-only status. This v4.9.0-r35 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r35 results

Conventions as in v4.9.0-r34. Put $P_\pm = \frac{1}{2}(1 \mp i\,J)\colon \mathbb{O} \to \mathbb{C} \otimes \mathbb{O}$, and let R_{e_7} be right multiplication by e_7.

Theoremlet v4.9.0-r35.1.

(a) *Inner frame.* $\mathrm{Stab}_{G_2}(e_7)$ is J-linear. $(\mathbb{O}, J) \cong \mathbb{C}^4$ splits into the trivial line $\mathrm{span}_{\mathbb{R}}(1, e_7)$ and the irreducible complex triplet $e_7^{\perp}$, with complex basis e_4, e_5, e_6; $Je_4 = -e_3$, $Je_5 = -e_2$, $Je_6 = e_1$.

(b) $\mathrm{span}(e_1, e_2, e_3) = J\,\mathrm{span}(e_4, e_5, e_6)$ is neither J-invariant nor invariant under $\mathrm{Stab}_{G_2}(e_7)$ nor under Q.

(c) $P_\pm$ is $\mathrm{Stab}_{G_2}(e_7)$-equivariant and injective, and $P_\pm J = \pm i\,P_\pm$. P_- maps $e_7^{\perp}$ onto the triplet $Q = \frac{2}{3}$ and $\mathrm{span}(1, e_7)$, in particular e_7, onto the singlet $Q = 0$; P_+ maps them onto $Q = \frac{1}{3}$ and $Q = 1$. T exchanges P_+ and P_-.

(d) $Q = \frac{1}{2} - \frac{i}{6}\,L_{e_7} - \frac{i}{3}\,R_{e_7}$.

Proof sketch. (a)–(c) by explicit matrices (script `llm/k76_test_h20.py`, deviations at most $2 \cdot 10^{-15}$); irreducibility of complex type from the real commutant on $e_7^{\perp}$, which is two-dimensional. In (c), $P_\pm$ commutes with every J-linear map, and $P_\pm J = \frac{1}{2}(J \pm i) = \pm i\,P_\pm$; $\bar{P}_\pm = P_\mp$ because J is real. (d) by a least-squares fit with residual $2 \cdot 10^{-16}$. □

Verdict and reading. H20a, H20c and H20d are confirmed; H20b is refuted.

1. The triplet basis e_4, e_5, e_6 and $\nu_R \leftrightarrow e_7$ are correct in the inner frame, and Theorem 18.2 holds there. H19c in r34 took the real vectors as states of $\mathbb{C} \otimes \mathbb{O}$, which mixes up the two frames. The translation to the outer frame is $P_\pm$, and the sign is a frame choice. With P_- the Companion's assignment agrees with Furey's reading: e_7 goes to the singlet $Q = 0$ (ν) and e_4, e_5, e_6 to the triplet $Q = \frac{2}{3}$ (u). With P_+ one obtains the conjugate picture. This is r34.1(d) again: T exchanges $P_\pm$, hence $Q \mapsto 1 - Q$.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

2. The sentence in the proof of Theorem 22.4 that the anti-triplet (e_1, e_2, e_3) carries the conjugate colour representation is a genuine error. It arises from mixing the frames: in the inner frame there is one triplet, and e_1, e_2, e_3 are its J-images; a conjugate triplet exists only with the outer unit, as the image of P_+ against that of P_-, and not on $\mathrm{span}(e_1, e_2, e_3)$.

3. The charge needs both units, and both left and right multiplication (r35.1(d)). The complex scalar it of Theorem 18.3 alone generates a global phase that is the same on all four charge sectors; it is not electromagnetism.

4. The double use of e_4, e_5, e_6 for colour and generation (v4.8.0-r13) is not a frame effect and stays.

5. The *reading* of r34.1(c) is corrected: the statement stands, but the Companion does not put colour and ν_R on wrong subspaces. It states them in the inner frame without naming the translation $P_\pm$.

Proposed addition to Konvention F: statements about colour and charge name their complex unit (inner $J = L_{e_7}$, without charge; outer i, with Q), and for the translation the sign P_+ or P_-.

v4.9.0-r35 status in the three frames

result	M / T / P
r35.1(a), (b)	inner frame only; no use of the $\mathbb{C}$ factor, hence the same in M, T and P
r35.1(c), (d)	stated in M; T exchanges P_+ and P_-, so the sign of the translation is the M/T choice; P not evaluated

v4.9.0-r35 register entries

statement	status after v4.9.0-r35
Theorem 18.2	holds in the inner frame (r35.1(a))
Theorem 22.4, triplet basis and $\nu_R \leftrightarrow e_7$	correct in the inner frame; with P_- consistent with Furey's charges (r35.1(c)); reading of r34.1(c) corrected
Theorem 22.4, anti-triplet (e_1, e_2, e_3)	refuted, frame mixing (r35.1(b)); candidate for a status note
Theorem 18.3, $U(1)$ line	the scalar it gives only a global phase; the charge needs both units (r35.1(d))

v4.9.0-r35 ledger and guard update

$$R_{\mathrm{fr}}^{v490\text{-}r35} = (1, 0, 1)$$

for (colour assignment correct in the inner frame; anti-triplet sentence wrong; frame rule for colour and charge proposed).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r35,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r35} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r36 Source-Only: Frame Rule for Complex Units, and the Origin of the Colour–Generation Double Use

Purpose. This layer does two things. It records the author's adoption (2026-10-23) of the frame rule proposed in v4.9.0-r35, with the resulting status notes in the main body. It then records the pre-registered test H21 (note `llm/Vorab_Notiz_H21_Herkunft_Doppelbelegung.md`, commit `516ab15`, fixed before computing) on where the double use of e_4, e_5, e_6 for colour and generation in Theorem 22.4 (v4.8.0-r13) comes from. Earlier layers are not rewritten.
Source-only status. This v4.9.0-r36 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r36 frame rule for complex units

Rule (Convention F, addition; adopted by the author). A statement about colour or charge names its complex unit: the inner unit $J = L_{e_7}$ on $\mathbb{O}$, which carries colour but no charge, or the outer unit i, the $\mathbb{C}$ factor of the Dixon storage, with the charge operator Q. A translation between them names its sign P_+ or P_- (v4.9.0-r35).

Following this rule, status notes are added to Theorem 22.4 and Theorem 18.3 (revision rule C of v4.9.0-r25; patch specification `llm/revision_spec_v490_r36.py`). Theorem 18.2 holds in the inner frame and needs no note.

v4.9.0-r36 the cyclic symmetry of the triole

Conventions as in v4.9.0-r35; $\omega = e^{2\pi i/3}$. Let σ be the real linear map of $\mathbb{O}$ with $\sigma(1) = 1$, $\sigma(e_7) = e_7$, $e_4 \mapsto e_5 \mapsto e_6 \mapsto e_4$, $e_1 \mapsto -e_3$, $e_2 \mapsto -e_1$, $e_3 \mapsto e_2$.

Theoremlet v4.9.0-r36.1.

(a) σ is an automorphism of $\mathbb{O}$ of order three fixing e_7, hence $\sigma \in \mathrm{Stab}_{G_2}(e_7) = SU(3)$. Among the 48 signed extensions of the cycle $e_4 \to e_5 \to e_6$ to e_1, e_2, e_3 with e_7 fixed it is the only automorphism.

(b) On $(e_7^{\perp}, J)$ with complex basis e_4, e_5, e_6, σ is the cyclic permutation matrix: determinant one, eigenvalues $1, \omega, \bar{\omega}$, not central. It commutes with Q and with T, $R_{\pi/2}$, J_B.

(c) The 120° triole $v = \sum_{k=0}^{2} \omega^k e_{k+4}$ (coefficients with respect to J; $\sum_k \omega^k = 0$) satisfies $\sigma v = \bar{\omega} v$, and $P_- v$ lies in the colour triplet $Q = \frac{2}{3}$.

(d) The subgroup of $\mathrm{Stab}_{G_2}(e_7)$ preserving $\mathrm{span}_{\mathbb{R}}(e_4, e_5, e_6)$ is three-dimensional and acts on it as $SO(3)$; it contains σ.

(e) In the Dixon storage $(\mathrm{Im}\,\mathbb{H}) \otimes \mathbb{C} \otimes \mathbb{O}$ of v4.9.0-r33, the cyclic $\mathbb{Z}_3$ on $\mathrm{Im}\,\mathbb{H}$ commutes with $1 \otimes \sigma$ and with all of $\mathrm{Stab}_{G_2}(e_7)$.

Proof sketch. By explicit matrices (script `llm/k77_test_h21.py`): the automorphism condition on all 64 basis products, an exhaustive check of the 48 signed extensions, the complex matrix of σ from $Je_4 = -e_3$, $Je_5 = -e_2$, $Je_6 = e_1$, and the linear condition for the subalgebra in (d). σ is real and acts on $\mathbb{O}$ only, so it commutes with the maps of the $\mathbb{C}$ factor; (e) follows from the tensor structure. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Verdict and reading. H21a–e are confirmed.

1. The triolic ground structure of the Companion, three kernels phase-shifted by 120° with $z_1 + z_2 + z_3 = 0$, and its cyclic symmetry are, inside $\mathbb{O}$, exactly the colour rotation σ, an element of the Weyl group of $SU(3)_c$. Theorem 18.2 reads the kernels as colours, consistent with (c). Theorem 22.4 reads the same cyclic symmetry as the generation symmetry. Colour and generation are therefore one object, not only the same units.

2. This is not a frame effect: σ commutes with all frame maps (b). It is an identification of two roles.

3. Every generation symmetry on the real span of e_4, e_5, e_6, such as the flavour groups of v4.8.0-r12, lies in $SO(3) \subset SU(3)_c$ (d), and breaking it would break colour. This makes consequence (i) of v4.8.0-r13 structural.

4. Inside $\mathbb{O}$ with e_7 fixed there is no $\mathbb{Z}_3$ that is not a colour transformation. The three generations therefore need a second three-fold structure outside $\mathbb{O}$. The Companion has two: the three triality slots of $\mathbb{O}^3$ (v4.8.0-r14) and $\mathrm{Im}\,\mathbb{H}$ in the Dixon storage (v4.9.0-r33). Both commute with colour (e).

5. Whether the triolic ground structure was meant as colour or as generations is open. By (a)–(c) it can be only one of them inside $\mathbb{O}$.

v4.9.0-r36 status in the three frames

result	M / T / P
r36.1(a), (b), (d)	inner frame; σ commutes with T, $R_{\pi/2}$, J_B, hence frame-free between M and T; P not evaluated
r36.1(c)	stated in M through P_-; in T the image is $P_+ v$, in the conjugate triplet $Q = \frac{1}{3}$
r36.1(e)	tensor structure, frame-free

v4.9.0-r36 register entries

statement	status after v4.9.0-r36
frame rule for complex units	adopted (Convention F, addition)
Theorem 22.4	status note added; the generation symmetry $e_4 \to e_5 \to e_6$ is a colour rotation (r36.1(a)–(c))
Theorem 18.3	status note added (scalar *it* a global phase; charge needs both units; $U(2)$, hypercharge)
generations	need a three-fold structure outside $\mathbb{O}$: triality slots or $\mathrm{Im}\,\mathbb{H}$ (r36.1(e))

v4.9.0-r36 ledger and guard update

$$R_{\text{gen}}^{v490\text{-}r36} = (1, 1, 0)$$

for (origin of the double use identified; separation shown to be necessary inside $\mathbb{O}$; carrier of the three generations open).

The layer spends no guard:

$$R_{\text{glob}}^{(v490\text{-}r36,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\text{guard}}^{v490\text{-}r36} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r37 Source-Only: Triality as Self-Traversal, and the Periods of the Structure

Purpose. Layer v4.9.0-r36 showed that inside $\mathbb{O}$ with e_7 fixed every $\mathbb{Z}_3$ symmetry is a colour transformation, so the generations need a second three-fold structure. The author states (2026-10-23) that the triole was meant as another representation of the Sphaera, and that the fractal self-traversal transforms between such representations; the number of steps is not to be fixed at three. This layer records the pre-registered test H22 (note `llm/Vorab_Notiz_H22_Triality_Selbstdurchwanderung.md`, commit `ed74608`, fixed before computing) on the triality form of v4.8.0-r14 and on the periods available to the structure. Earlier layers are not rewritten.
Source-only status. This v4.9.0-r37 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r37 results

Let $t(x,y,z) = \mathrm{Re}((xy)z)$ on $\mathbb{O}^3$, $\mathfrak{g} = \{(A,B,C) \in \mathfrak{so}(8)^3 : t(Ax,y,z) + t(x,By,z) + t(x,y,Cz) = 0\}$, $\tau(x,y,z) = (y,z,x)$ and $\rho(x,y,z) = (\bar{x},\bar{z},\bar{y})$; T, $R_{\pi/2}$ act on the $\mathbb{C}$ factor as in v4.9.0-r30.

Theoremlet v4.9.0-r37.1.

(a) $\dim \mathfrak{g} = 28$; each projection $\mathfrak{g} \to \mathfrak{so}(8)$ is an isomorphism, and the three representations of $\mathfrak{g}$ on the three slots are irreducible and pairwise inequivalent. Thus t is the Spin(8) triality form.

(b) τ maps $\mathfrak{g}$ to itself and permutes the three slot representations cyclically; it is an outer automorphism of order three.

(c) The τ-fixed elements $(A,A,A) \in \mathfrak{g}$ form the derivation algebra of $\mathbb{O}$, of dimension 14; those with $Ae_7 = 0$ form $\mathfrak{su}(3)$.

(d) On $\mathbb{O}^4$ the five bracketings of $\mathrm{Re}(xyzw)$ fall into two classes, $\mathrm{Re}(((xy)z)w) = \mathrm{Re}((xy)(zw)) = \mathrm{Re}(x(y(zw)))$ and $\mathrm{Re}((x(yz))w) = \mathrm{Re}(x((yz)w))$. The cyclic shift of the four slots exchanges the classes, so no single bracketing is invariant, while their sum is.

(e) $\tau, \rho, T, R_{\pi/2}$ generate on $\mathbb{C} \otimes \mathbb{O}^3$ a group of order 48, isomorphic to $S_3 \times D_4$, with element orders $1, 2, 3, 4, 6, 12$; τ and ρ preserve t.

Proof sketch. (a)–(c) by solving the linear conditions on $\mathfrak{so}(8)^3$ (512 equations, 84 unknowns) and computing ranks, intertwiner spaces and commutants (script `llm/k78_test_h22.py`, deviations at most 10^{-14}); τ preserves t because $\mathrm{Re}((xy)z) = \mathrm{Re}((yz)x)$ (v4.8.0-r14.2), and an automorphism permuting inequivalent representations is not inner. (d) from $\mathrm{Re}((ab)c) = \mathrm{Re}(a(bc))$ and $\mathrm{Re}(ab) = \mathrm{Re}(ba)$, checked numerically; the exchange of the classes was found in a check after the evaluation. (e) by enumerating the group. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Verdict and reading. H22a, b, c and e are confirmed. H22d, the statement that no single bracketing is cyclically invariant, is confirmed, but by (d) it gives no prohibition of a fourth slot.

1. The second three-fold structure required by v4.9.0-r36 exists: it is triality. The three slots of $\mathbb{O}^3$ carry the three inequivalent eight-dimensional representations of Spin(8), the shift of the slots is the outer triality automorphism, and what it fixes is $G_2 \supset SU(3)_c$. In this reading generation is what the traversal permutes and colour is what it fixes; they are orbit and fixed group of one motion, no longer one object. It matches the author's reading of the triole as three representations of the same $\mathbb{O} \cong \mathbb{R}^8$. That the slots *are* the physical generations is not shown.

2. That there are exactly three rests on the known classification: Spin(8) has exactly three eight-dimensional irreducible representations and $\mathrm{Out}(\mathrm{Spin}(8)) = S_3$. (a) is consistent with it. It excludes a fourth slot with a new eight-dimensional representation, not a repetition of one of the three. The structure therefore does not by itself forbid a fourth generation.

3. The structure has no distinguished period. Period two belongs to ρ, T, J_B and the alternating triolic shift of §11, period three to triality, period four to the duality rotation $R_{\pi/2}$, and six and twelve to their combinations. In the author's picture, the period seen by a readout is a choice of subgroup.

4. The word triole names at least three structures in the corpus: the frame triole matter–photon–tachyon of the original paper (spring theorem; $\mathbb{Z}_2$ with fixed point, amplitudes $1:1:3$), the kernel triole with 120° phases (a colour rotation inside $\mathbb{O}$, v4.9.0-r36), and the slot triole of $\mathbb{O}^3$ (triality). The double uses found in v4.8.0-r13 and v4.9.0-r34 to r36 arise where these are identified.

v4.9.0-r37 status in the three frames

result	M / T / P
r37.1(a)–(d)	real octonionic statements without the $\mathbb{C}$ factor; the same in M, T and P
r37.1(e)	contains the frame maps themselves; T and J_B exchange M and T, $R_{\pi/2}$ is the duality rotation; P is the common fixed point

v4.9.0-r37 register entries

statement	status after v4.9.0-r37
triality form t of v4.8.0-r14	the Spin(8) triality form (r37.1(a), (b))
separation of colour and generation	colour in the fixed group $G_2 \supset SU(3)_c$ of triality, generation in its orbit (r37.1(c)); physical identification open
exactly three generations	not forbidden by the structure alone; the classification excludes a fourth *distinct* slot only (r37.1(d))
periods of the self-traversal	$1, 2, 3, 4, 6, 12$ (r37.1(e)); no distinguished period

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r37 ledger and guard update

$$R_{\mathrm{tr}}^{v490\text{-}r37} = (1, 0, 1)$$

for (triality carrier of the three slots found; prohibition of a fourth generation not obtained; physical identification of the slots open).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r37,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r37} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r38 Source-Only: The Frame Triole as the Reflection Side of Triality

Purpose. In the original paper the triole of the spring theorem is the triple matter–photon–tachyon: the photon is the equilibrium node and matter and tachyon are mirror images. Layer v4.9.0-r37 found that the cyclic shift τ of the three slots of $\mathbb{O}^3$ is triality. This layer records the pre-registered test H23 (note `llm/Vorab_Notiz_H23_Frame_Triole_Triality.md`, commit `55043f8`, fixed before computing): whether the frame triole has the form of the second slot map $\rho(x, y, z) = (\bar{x}, \bar{z}, \bar{y})$, which fixes one slot and exchanges the other two. Earlier layers are not rewritten.

Source-only status. This v4.9.0-r38 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r38 results

Notation as in v4.9.0-r37; c is octonionic conjugation, and ρ acts on $\mathfrak{g}$ by $(A, B, C) \mapsto (cAc, cCc, cBc)$.

Theoremlet v4.9.0-r38.1.

(a) ρ is an automorphism of $\mathfrak{g}$ of order two. It keeps the representation class of slot 1 and exchanges the classes of slots 2 and 3; it is outer.

(b) The ρ-fixed subalgebra of $\mathfrak{g}$ is 21-dimensional, isomorphic to $\mathfrak{so}(7)$. On slot 1 it acts as $1 \oplus 7$ with invariant line $\mathbb{R} \cdot 1$ and irreducible complement; on slots 2 and 3 it acts irreducibly, by equivalent eight-dimensional representations.

(c) The sign triples preserving t are those with $s_1 s_2 s_3 = 1$, a group $\mathbb{Z}_2 \times \mathbb{Z}_2$; the one trivial on slot 1 is $(1, -1, -1)$.

(d) τ and ρ generate S_3 on the slots; its $\mathbb{Z}_3$ has one orbit of length three, each transposition a fixed slot and an orbit of length two.

(e) $\mathfrak{g}_2 = \{(A, A, A)\}$ is ρ-fixed and lies in the $\mathfrak{so}(7)$ of (b).

Proof sketch. (a), (b), (e) by linear algebra on $\mathfrak{so}(8)^3$: closure under ρ, intertwiner spaces between the slot representations of $\mathfrak{g}$ and of its image, the fixed space of ρ, and commutants (script `llm/k79_test_h23.py`, deviations at most 10^{-15}). (c) is immediate from trilinearity and was checked numerically; (d) by enumeration. □

Verdict and reading. H23a–e are confirmed; (c) and (d) are elementary.

1. The 120° triole is the $\mathbb{Z}_3$-orbit of triality; the frame triole of the original paper has the form of a transposition, one fixed slot and two exchanged. Both are views of the same S_3 on the three triality slots. In the author's picture, the zoom chooses the subgroup.

2. Reading: photon = the fixed slot, which under the fixed Spin(7) has a distinguished direction, the real unit; matter and tachyon = the two exchanged slots, equal under Spin(7) and mirrored by ρ. The parity $(1, -1, -1)$ separates the photon slot from the

© 2026 Marc Brendecke. Preprint — not peer reviewed.

other two in the way $(-1)^F$ separates bosons from fermions; it is trivial, and notable only because it fits this reading.

3. Which slot is the photon is a choice: S_3 has three transpositions, each fixing a different slot. This agrees with Convention F, where the photon is the degenerate fixed point.

4. Colour is fixed by the whole S_3 (e).

5. Open: on one level the three slots cannot carry both the frames (this layer) and the generations (reading of v4.9.0-r37). In the author's picture these would be two zoom levels. Which level carries which role is not shown; $\mathrm{Im}\,\mathbb{H}$ of the Dixon storage (v4.9.0-r33) is a second three-fold carrier.

6. In eight dimensions the two spinor slots are a chirality pair. If matter and tachyon are read as these, the frame change $M \leftrightarrow T$ is a change of chirality. This is a lead for the chirality test of the inventory, not computed here.

v4.9.0-r38 status in the three frames

result	M / T / P
r38.1(a)–(e)	real octonionic statements on the slots, without the $\mathbb{C}$ factor; the same in M, T and P
reading (2), (3)	assigns M, P, T to slots; the assignment is a choice of transposition and is itself the frame choice

v4.9.0-r38 register entries

statement	status after v4.9.0-r38
frame triole of the spring theorem	has the form of a transposition of the triality slots (r38.1(a), (b)); reading photon = vector slot, matter and tachyon = spinor slots
120° triole and frame triole	views of the same S_3 (r38.1(d))
frames versus generations on the slots	open; two levels needed
chirality	lead: $M \leftrightarrow T$ as exchange of the two spinor slots

v4.9.0-r38 ledger and guard update

$$R_{\mathrm{ft}}^{v490\text{-}r38} = (1, 0, 0)$$

for (frame triole matched to the reflection side of triality; level of frames against generations open; chirality open).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r38,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r38} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r39 Source-Only: Chains of Different Length under Hilbert-Shift Zoom and Mixed Readouts

Purpose. The author states (2026-10-23) that the mixed readouts were meant for the case in which the generations form a circle after repeated zoom, and that chains of different length might give different readings. This layer records the pre-registered test H24 (note `llm/Vorab_Notiz_H24_Ketten_Module_Readouts.md`, commit `8c89481`, fixed before computing). It uses the Hilbert shift of Definition H2.33 and the bit registry $\beta_j = 2^j$ of Definition H2.37. Earlier layers are not rewritten.
Source-only status. This v4.9.0-r39 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r39 model

A chain of length m is a cyclic group $\langle g\rangle$ with $\operatorname{ord} g = m$. A readout of level j (bit 2^j) sees $\langle g^{2^j}\rangle$. A mixed mask S with $k = \sum_{j\in S} 2^j$ is read in two ways: (Σ) as one step, $\langle g^k\rangle$, or ($\vee$) as the joint readout $\langle g^{2^j} : j \in S\rangle$. The group is $G = \langle \tau, \rho, T, R_{\pi/2}\rangle \cong S_3 \times D_4$ of v4.9.0-r37. The identification of zoom with the step 2^j is a reading.

v4.9.0-r39 results

Theoremlet v4.9.0-r39.1.

(a) A chain of odd length keeps its length at every level. For the triality, $\tau^{2^j} = \tau$ for even j and $\tau^{2^j} = \tau^{-1} = \rho\tau\rho$ for odd j.

(b) A chain of length 2^a has length 2^{a-j} at level $j < a$ and 1 from level a on. In G the visible orders are $\{1, 2, 3, 4, 6, 12\}$ at level 0, $\{1, 2, 3, 6\}$ at level 1, and $\{1, 3\}$ from level 2 on, where every g^{2^j} lies in $\langle\tau\rangle$.

(c) For $m = 3$ and masks over six bits: in reading (Σ) the 21 masks with $3 \mid k$, for instance $S = \{0, 1\}$, see one generation and the other 42 see three; in reading ($\vee$) every mask sees three.

(d) For every m: (Σ) the visible length is $m/\gcd(m, k)$; ($\vee$) it is $m/\gcd(m, 2^{\min S})$, so the odd part of m is always preserved.

Proof sketch. Elementary: 2^j is invertible modulo odd m, and $2^j \equiv (-1)^j \pmod 3$; the subgroup of $\mathbb{Z}_m$ generated by a set of steps has order $m/\gcd$. All statements were checked by exact enumeration, on G as matrices, for $m \le 64$ and all 63 masks (script `llm/k80_test_h24.py`). □

Verdict and reading. H24a–e are confirmed; the content is the arithmetic of cyclic groups, and the physics lies only in the reading.

1. In the model, the generations do form a circle under repeated zoom, and it is the only thing left: from level 2 on, only the triality circle of length three remains visible, while the frame exchange (period 2) and the duality rotation (period 4) disappear. This separates the two levels left open in v4.9.0-r38: the frames at coarse zoom, the generations at deep zoom.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

2. On odd levels the circle runs backwards, and the reversed circle is the frame mirror ρ; every second level reads the generations in the mirrored frame.

3. What survives zoom is the odd part of a chain length. Chains of length 6 (three generations times two frames) or 12 show three at deep zoom; chains of length 2^a vanish. No generation chain of length four or more arises in G.

4. Mixed readouts give different readings depending on the aggregation: as one step, a third of the masks see one generation; as a joint readout, all see three. Definition H2.36 leaves the aggregation μ open.

Author remark and textual reading. The author states (2026-10-23) that the contraction discipline of Definition H2.41 was introduced so that classical tensor calculus keeps working while mixed channels can also be contracted, that it was hardly used, and that the paper was written from one frame. Read against r39.1: in the selector contraction $\chi_{mn}R^mS^n$ of Definition H2.43, compatible modes are composed; for shift readouts $R^m = g^{2^m}$, $S^n = g^{2^n}$ the composition is $g^{2^m+2^n}$, which is reading (Σ). The Boolean mask grammar $(\chi_A \vee \chi_B)$ only selects the active kinds, which is reading $(\vee)$. Both readings are therefore present in the text, at different places. Combining an even and an odd level by contraction gives $\tau^3 = 1$: a readout composed with its frame mirror does not see the generation circle. The front indices of Definition H2.40 carry the frame; in the main body they are not varied, which is the one-frame restriction. This is a reading, not part of the pre-registered test.

v4.9.0-r39 status in the three frames

result	M / T / P
r39.1(a), (c), (d)	arithmetic of chains, frame-free
r39.1(b)	contains the frame maps; from level 2 on they are invisible, so deep-zoom statements are the same in M, T and P
reading (2)	odd levels read the generations in the ρ-mirrored frame

v4.9.0-r39 register entries

statement	status after v4.9.0-r39
generation circle under zoom	the only cycle of G visible from level 2 on (r39.1(b)); orientation alternates with the level (r39.1(a))
frames versus generations	separated by zoom level in the model
mixed readouts	one or three generations, depending on the aggregation μ; selector contraction $= (\Sigma)$, Boolean mask $= (\vee)$ (reading)
front indices of two-sided displays	carry the frame; not varied in the main body (one-frame restriction)

v4.9.0-r39 ledger and guard update

$$R_{\mathrm{zm}}^{v490\text{-}r39} = (1, 1, 0)$$

for (generation circle survives zoom; frames and generations separated by level; aggregation of mixed readouts open).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r39,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r39} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r40 Source-Only: The Two-Sided Contraction across Frames, and the Weak SU(2)

Purpose. The two-sided notation of Definition H2.40 puts the frame on the front indices. The author states (2026-10-23) that its contraction discipline, Definition H2.41, has never been used or tested, and that it is not expected to work at once. This layer uses the front side across frames for the first time and records the pre-registered test H25 (note `llm/Vorab_Notiz_H25_Zweiseitige_Kontraktion_Frames.md`, commit `d1aba48`, fixed before computing). It also asks whether a weak $SU(2)$ commuting with colour exists in the triality algebra. Earlier layers are not rewritten.
Source-only status. This v4.9.0-r40 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r40 model

Front side: the $\mathbb{C}$ factor of the Dixon storage, with the frame maps $R_{\pi/2}$ (linear, $X \mapsto iX$, contragredient $\omega \mapsto -i\omega$), T (antilinear, complex conjugation) and $J_B = R_{\pi/2} \circ T$. Back side: real structure, acting on the index α. Rule 1 is the $\mathbb{C}$-bilinear contraction $\omega_\alpha X^\alpha$; rule 2 is $q(X) = G_{\alpha\beta} X^\alpha X^\beta$ with real symmetric G, compared with the Hermitian form $h(X) = G_{\alpha\beta} \bar{X}^\alpha X^\beta$. On $W = \mathbb{C} \otimes \mathbb{H} \otimes \mathbb{O}$ the front maps are D_4 and left multiplication L_q by unit quaternions, the back maps right multiplication R_q and $SU(3)_c = \mathrm{Stab}_{G_2}(e_7)$.

v4.9.0-r40 results

Theoremlet v4.9.0-r40.1.

(a) On W every front map commutes with every back map.

(b) $\omega_\alpha X^\alpha$ is invariant under $R_{\pi/2}$ and is complex conjugated under T and J_B.

(c) $q \mapsto -q$ under $R_{\pi/2}$, $q \mapsto \bar{q}$ under T, $q \mapsto -\bar{q}$ under J_B; h is invariant under all eight elements of D_4.

(d) The centralizer of $\mathfrak{su}(3)_c$ in the triality algebra $\mathfrak{g} \cong \mathfrak{so}(8)$ of v4.9.0-r37 is two-dimensional and abelian. On W, L_q and R_q give two copies of $\mathfrak{su}(2)$ that commute with each other, with $\mathfrak{su}(3)_c$ and with D_4.

(e) No nonzero element of $\mathfrak{g}$ acts trivially on one spinor slot and nontrivially on the other.

In a check after the evaluation, the centralizer in (d) was found to be $\mathrm{span}(L_{e_7}, R_{e_7})$ on each slot.

Proof sketch. (b), (c) by direct substitution, checked on random vectors; D_4 is generated by $R_{\pi/2}$ and T. (a), (d), (e) by matrices and null spaces (script `llm/k81_test_h25.py`, deviations at most 10^{-14}); (e) is the injectivity of the slot projections (v4.9.0-r37.1(a)). □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Verdict and reading. H25a–e are confirmed; (a) and (e) are controls.

1. Used across frames for the first time, the rules work, but not as stated. Rule 1 says that the classical contraction remains unchanged; across frames this holds for $R_{\pi/2}$ only, while the antilinear frame maps conjugate the value. Only real readouts are frame-invariant. Rule 2 depends on whether the comparison is bilinear or Hermitian, which the definition does not fix. The notation is frame-covariant, not frame-invariant; this is a finding about the notation, which so far was used in one frame only.

2. The bilinear comparison changes sign under the transition $z \mapsto iz$ from M to T. Read as a squared mass, this is $m^2 \mapsto -m^2$, the tachyon. The Hermitian comparison does not distinguish M from T. The choice is therefore one of content: bilinear comparisons separate the frames, Hermitian ones do not.

3. No $\mathfrak{su}(2)$ of the triality algebra commutes with colour; what commutes is $\mathrm{span}(L_{e_7}, R_{e_7})$, the generators of the charge $Q = \frac{1}{2} - \frac{i}{6}L_{e_7} - \frac{i}{3}R_{e_7}$ of v4.9.0-r35. The triality level carries colour and charge, but no weak interaction. An $\mathfrak{su}(2)$ commuting with colour appears only in the $\mathbb{H}$ factor, twice: from the left (front side) and from the right (back side).

4. Reading: the two $\mathfrak{su}(2)$ of $\mathbb{H}$ sit on the two sides of the two-sided notation, and "the weak $SU(2)$ acts only on the left" would read "the weak $SU(2)$ is one side". This is not shown. The conflict of the inventory remains: the left side also carries the Lorentz rotation, and the right side the generation $\mathbb{Z}_3$ of v4.9.0-r33.

5. Proposed frame clause for rules 1 and 2 (author decision): readouts are real or are conjugated with the antilinear frame maps, and each comparison tensor is declared bilinear (frame-separating) or Hermitian (frame-blind).

v4.9.0-r40 status in the three frames

result	M / T / P
r40.1(b), (c)	compare M with T directly: rule 1 invariant under $R_{\pi/2}$, conjugated under T, J_B; bilinear rule 2 changes sign between M and T; P not evaluated
r40.1(a), (d), (e)	real structural statements, the same in M, T and P

v4.9.0-r40 register entries

statement	status after v4.9.0-r40
Definition H2.41 across frames	frame-covariant, not invariant (r40.1(b), (c)); frame clause proposed
weak $SU(2)$	not in the triality algebra (r40.1(d)); two candidates in $\mathbb{H}$, left and right; overload of $\mathbb{H}$ unresolved
colour centralizer in triality	$\mathrm{span}(L_{e_7}, R_{e_7})$, the charge generators (check after evaluation)

v4.9.0-r40 ledger and guard update

$$R_{\mathrm{ts}}^{v490\text{-}r40} = (1, 0, 0)$$

for (frame behaviour of the two-sided contraction determined; frame clause undecided; weak $SU(2)$ not located).

© 2026 Marc Brendecke. Preprint — not peer reviewed.

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r40,+)} = (0,1,1,1,1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r40} = (1,1,1,1,1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r41 Source-Only: Distributing the Roles of the Quaternion Factor; Hypercharge and Electric Charge

Purpose. The inventory of the standard-model features found the quaternion factor overloaded: it carries the Lorentz spinor (Theorem 22.4), the weak $SU(2)$ (Theorem 18.3) and, in v4.9.0-r33, the generation index. Layers v4.9.0-r37 to r40 found a carrier for the generations in triality, a separation of frames and generations by zoom level, and two $\mathfrak{su}(2)$ commuting with colour, from the left and from the right of $\mathbb{H}$. This layer records the pre-registered test H26 (note `llm/Vorab_Notiz_H26_Verteilung_H_Faktor.md`, commit `1876b55`, fixed before computing) of a distribution D of these roles. The charge pattern of (b) and (c) below was seen analytically before the note and is disclosed there; the computation checks it and is not a prediction. Earlier layers are not rewritten.
Source-only status. This v4.9.0-r41 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r41 distribution D

Lorentz $SL(2,\mathbb{C})$ by left multiplication L_q (rotations) and iL_q (boosts) on $\mathbb{C}\otimes\mathbb{H}$; weak $SU(2)$ by right multiplication R_q on $\mathbb{H}$, with $T_3 = -\frac{i}{2}R_k$; colour $SU(3)_c$ on $\mathbb{O}$; frames D_4 on $\mathbb{C}$; generations on the triality slots of $\mathbb{O}^3$. Hypercharge $Y/2 := Q - \frac{1}{2}$ with Q of v4.9.0-r34/r35; electric charge $Q_{\mathrm{em}} := T_3 + Y/2$. Here i is the outer unit.

v4.9.0-r41 results

Theoremlet v4.9.0-r41.1.

(a) On $\mathbb{C}\otimes\mathbb{H}\otimes\mathbb{O}$ the weak $SU(2)$ commutes with the Lorentz group, with colour and with Q. The Lorentz rotations are fixed by T; the boosts are reversed by T.

(b) The joint eigenspaces of Q and T_3 carry Q_{em} as follows:

Q-sector (colour)	$Y/2$	Q_{em} at $T_3 = +\frac{1}{2}, -\frac{1}{2}$	reading
0 (singlet)	$-\frac{1}{2}$	$0,\ -1$	lepton doublet (ν, e)
$\frac{2}{3}$ (triplet)	$+\frac{1}{6}$	$\frac{2}{3},\ -\frac{1}{3}$	quark doublet (u, d)
$\frac{1}{3}$ (conjugate triplet)	$-\frac{1}{6}$	$\frac{1}{3},\ -\frac{2}{3}$	antiquark doublet $(\bar{d}, \bar{u})$
1 (singlet)	$+\frac{1}{2}$	$1,\ 0$	antilepton doublet $(e^+, \bar{\nu})$

(c) $TQ_{\mathrm{em}}T^{-1} = -Q_{\mathrm{em}}$ and $T(Y/2)T^{-1} = -Y/2$.

(d) There is no weak singlet: $T_1^2 + T_2^2 + T_3^2 = \frac{3}{4}$ on the whole space.

(e) On $\mathbb{C}\otimes\mathbb{H}\otimes\mathbb{O}^3$ the cyclic shift of the slots commutes with L_q, R_q, colour and Q.

Proof sketch. By explicit complex matrices on $\mathbb{H}\otimes\mathbb{O} \cong \mathbb{C}^{32}$, with the $\mathbb{C}$ factor as scalar and T as complex conjugation (script `llm/k82_test_h26.py`, deviations at most 10^{-14}). (c) follows from $TQT^{-1} = 1 - Q$ (v4.9.0-r34.1(d)) and $T(-\frac{i}{2}R_k)T^{-1} = \frac{i}{2}R_k$. (d): R_q acts on $\mathbb{H} \cong \mathbb{C}^2 \otimes \mathbb{C}^2$ as a doublet on the right index. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Verdict and reading. H26a–e are confirmed; (d) is negative for D.

1. In D the quaternion factor is no longer overloaded: the Lorentz group sits on the left (front side), the weak $SU(2)$ on the right (back side), the generations on the triality slots. The zoom levels separate frames from generations; the sides separate the Lorentz group from the weak $SU(2)$.

2. Under the choice $Y/2 = Q - \frac{1}{2}$, $Q_{\mathrm{em}} = T_3 + Y/2$ gives exactly the hypercharges and charges of the left-handed standard-model doublets and their conjugates, with the right colour: leptons singlets, quarks triplets, antiquarks conjugate triplets. The choice is motivated by the pattern and not forced by the structure. The sector assignment is Furey's; similar patterns are known in the literature on $\mathbb{C} \otimes \mathbb{H} \otimes \mathbb{O}$. New here is the placement on the sides and frames of the Companion.

3. In D the frame change T is charge conjugation, $Q_{\mathrm{em}} \mapsto -Q_{\mathrm{em}}$. The shift in $Q \mapsto 1 - Q$ of v4.9.0-r34.1(d) is the hypercharge offset $\frac{1}{2}$. Matter frame and tachyon frame behave in D as particle and antiparticle.

4. D is incomplete. Every state is a weak doublet; the right-handed singlets u_R, d_R, e_R, ν_R, half of a generation, are absent, and all states carry the same left Lorentz index. Chirality is not explained. Weak singlets would need a structure on which the right action of $\mathbb{H}$ is trivial on part of the space, for instance a projector coupling left and right.

v4.9.0-r41 status in the three frames

result	M / T / P
r41.1(a), (d), (e)	the same in M, T and P, except that T reverses the boosts
r41.1(b)	stated in M; in T all charges and hypercharges change sign (r41.1(c)): the T frame reads the conjugate states
r41.1(c)	relates M and T: T acts as charge conjugation; P not evaluated

v4.9.0-r41 register entries

statement	status after v4.9.0-r41
overload of $\mathbb{H}$	resolved in D by sides (Lorentz left, weak right) and by triality for the generations (r41.1(a), (e))
hypercharge and $Q = T_3 + Y/2$	doublet pattern reproduced under the choice $Y/2 = Q - \frac{1}{2}$ (r41.1(b)); not derived
frame T	charge conjugation in D (r41.1(c))
weak singlets, chirality	absent in D (r41.1(d)); open

v4.9.0-r41 ledger and guard update

$$R_{\mathrm{qh}}^{v490\text{-}r41} = (1, 1, 0)$$

for (overload of $\mathbb{H}$ resolved in D; doublet charges reproduced under a choice; singlets and chirality absent).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r41,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r41} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r42 Source-Only: A Compensator of the Millennium Matrix as Chirality Projector

Purpose. Layer v4.9.0-r41 placed the weak $SU(2)$ on the right of $\mathbb{H}$ and obtained only weak doublets. The author states (2026-10-23) that the Millennium matrix of §38, which was to carry several Heisenberg compensators, was meant for the missing half. This layer records the pre-registered test H27 (note `llm/Vorab_Notiz_H27_MM_Kompensator_Chiralitaet.md`, commit `614ac3e`, fixed before computing). The charge pattern was seen analytically before the note and is disclosed there; the computation checks it and is not a prediction. Earlier layers are not rewritten.
Source-only status. This v4.9.0-r42 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r42 construction

On $\mathbb{C}\otimes\mathbb{H}\otimes\mathbb{O}\cong\mathbb{C}^{32}$ with the distribution D of v4.9.0-r41, put

$$C_{\mathrm{ch}}=\tfrac{1}{2}(\mathrm{id}+J),\qquad J=i\,L_{e_7},\qquad T'_a=C_{\mathrm{ch}}T_a,$$

$$\tfrac{Y}{2}=Q-\tfrac{1}{2}+(1-C_{\mathrm{ch}})T_3,\qquad Q_{\mathrm{em}}=T_3+Q-\tfrac{1}{2},$$

with i the outer unit and L_{e_7} the inner unit, so $J^2=1$ and C_{ch} has the form of the Heisenberg compensator.

v4.9.0-r42 results

Theoremlet v4.9.0-r42.1.

(a) C_{ch} is a Hermitian projector of rank 16; it commutes with colour, the Lorentz group, the right action of $\mathbb{H}$ and Q, and its image is the sum of the Q-sectors 0 and $\frac{2}{3}$.

(b) Under colour, the weak $SU(2)'$ and $Y/2$ the space splits into

$(SU(3),SU(2)',Y/2)$	Q_{em}	state	complex dimension
$(3,2,\frac{1}{6})$	$\frac{2}{3},-\frac{1}{3}$	$(u,d)_L$	12
$(1,2,-\frac{1}{2})$	$0,-1$	$(\nu,e)_L$	4
$(\bar3,1,-\frac{2}{3})$	$-\frac{2}{3}$	u^c	6
$(\bar3,1,\frac{1}{3})$	$\frac{1}{3}$	d^c	6
$(1,1,1)$	1	e^c	2
$(1,1,0)$	0	ν^c	2

that is, sixteen Weyl states with two Lorentz components each; $Y/2$ commutes with $SU(2)'$, colour and the Lorentz group, and $Q_{\mathrm{em}}=T'_3+Y/2$.

(c) Over the sixteen states, $\mathrm{Tr}\,Y$, $\mathrm{Tr}\,Y^3$ and the $SU(2)^2U(1)$ and $SU(3)^2U(1)$ traces vanish.

(d) $SU(2)'$ acts only on the image of C_{ch}, and $TC_{\mathrm{ch}}T^{-1}=1-C_{\mathrm{ch}}$.

(e) $\mathcal{M}=\lambda_0C_{\mathrm{ch}}+\lambda_1(1-C_{\mathrm{ch}})$, $\lambda_0\neq\lambda_1$, has C_{ch} as spectral projector, as in Theorem 29.22.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof sketch. By explicit complex matrices (script `llm/k83_test_h27.py`, deviations at most 10^{-15}). C_{ch} equals P_- of v4.9.0-r35.1(c). A first run reported (b) as refuted because the candidate values of $Y/2$ were rounded to nine digits while the null-space test required 10^{-10}, which lost all non-dyadic values; only this rounding was corrected, and both outputs are kept. □

Verdict and reading. H27a–e are confirmed, (b) after the correction of the script; (e) is a control.

1. A compensator of the Companion's form $\frac{1}{2}(\mathrm{id} + J)$, with J the product of the outer and the inner unit, completes the doublets of v4.9.0-r41 to the full left-handed content of one generation with right-handed neutrino. The weak $SU(2)$ acts only on its image: in the model, this is chirality. A compensator of the Millennium matrix separates the chiralities.

2. The frame change T exchanges image and kernel of the compensator. With v4.9.0-r41.1(c), matter frame and tachyon frame are in the model particle and antiparticle with mirrored chirality. This is a reading.

3. The Millennium matrix of §38 is 16×16, and one generation with ν_R has sixteen Weyl states. This is recorded, not interpreted.

4. This is a construction under choices, not a derivation: J and the formula for $Y/2$ are chosen so that the known result appears, and the pattern was seen in advance. It shows that the building blocks of the Companion (compensator, two units, two sides, triality) suffice. Anomaly cancellation follows from the content. Similar constructions exist in the literature on $\mathbb{C} \otimes \mathbb{H} \otimes \mathbb{O}$. Why this choice, symmetry breaking, masses and mixings are not addressed.

v4.9.0-r42 status in the three frames

result	M / T / P
r42.1(a)–(c)	stated in M
r42.1(d)	relates M and T: T exchanges the chiral halves; with r41.1(c) the T frame reads the conjugate states with mirrored chirality; P not evaluated

v4.9.0-r42 register entries

statement	status after v4.9.0-r42
content of one generation	constructed under choices from compensator, two units, two sides (r42.1(b)); not derived
chirality	in the model: the weak $SU(2)$ acts on the image of one compensator (r42.1(d))
Millennium matrix	a two-projector reading with C_{ch} as spectral projector (r42.1(e)); the dimension 16 recorded

v4.9.0-r42 ledger and guard update

$$R_{\mathrm{ch}}^{v490\text{-}r42} = (1, 0, 0)$$

for (content of one generation constructed; choice of J and Y not derived; breaking, masses and mixings open).

© 2026 Marc Brendecke. Preprint — not peer reviewed.

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r42,+)} = (0,1,1,1,1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r42} = (1,1,1,1,1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r43 Source-Only: The Choice of the Chirality Projector Is Fixed by Anomaly Cancellation

Purpose. Layer v4.9.0-r42 built the content of one generation with the compensator $C = \frac{1}{2}(\mathrm{id} + J)$, $J = iL_{e_7}$, and the hypercharge offset $-\frac{1}{2}$, both chosen. The author asks why this choice. This layer records the pre-registered test H28 (note `llm/Vorab_Notiz_H28_Warum_diese_Wahl.md`, commit `9b8b360`, fixed before computing; one case and the sector assignment were computed by hand beforehand and are disclosed there). Earlier layers are not rewritten.
Source-only status. This v4.9.0-r43 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r43 setting

On $\mathbb{C}\otimes\mathbb{H}\otimes\mathbb{O}$ as in v4.9.0-r42, a chirality projector is $C = \frac{1}{2}(1+J)$ with J Hermitian, $J^2 = 1$, $J \neq \pm 1$, acting on $\mathbb{C}\otimes\mathbb{O}$, commuting with colour and Q, of rank 16, with image one colour singlet and one colour triplet. The weak $SU(2)'$ is CT_a, the charge $Q_{\mathrm{em}} = T_3 + Q + s$, the hypercharge $Y/2 = Q + s + (1-C)T_3$.

v4.9.0-r43 results

Theoremlet v4.9.0-r43.1.

(a) The commutant of colour and Q on $\mathbb{C}\otimes\mathbb{O}$ is spanned by the four sector projectors. There are exactly four chirality projectors, with $J = iL_{e_7}$ (image the sectors $0, \frac{2}{3}$), $-iL_{e_7}$ $(\frac{1}{3}, 1)$, iR_{e_7} $(0, \frac{1}{3})$ and $-iR_{e_7}$ $(\frac{2}{3}, 1)$.

(b) For each of them $\mathrm{Tr}\, Y = 0$ holds exactly for $s = -\frac{1}{2}$.

(c) At $s = -\frac{1}{2}$ all four anomaly traces ($\mathrm{Tr}\, Y$, $\mathrm{Tr}\, Y^3$, $SU(2)^2U(1)$, $SU(3)^2U(1)$) vanish exactly for $J = \pm iL_{e_7}$. For $J = \pm iR_{e_7}$ one finds $\mathrm{Tr}\, Y^3 = \pm\frac{3}{2}$ and $SU(2)^2U(1) = \mp 2$, the latter summed over both states of each doublet.

(d) $T(iL_{e_7})T^{-1} = -iL_{e_7}$: the two admissible choices are frame images of each other.

Proof sketch. By explicit matrices and exact fractions (script `llm/k84_test_h28.py`). The commutant consists of scalars on the four Q-sectors, because each sector is colour-irreducible with its own charge; a Hermitian involution there is a choice of sign per sector, and the rank and image conditions leave four. □

Verdict and reading. H28a–d are confirmed.

1. Under the assumptions of the setting the choice of v4.9.0-r42 is forced up to the frame. Colour and charge conservation allow four chirality projectors, the gravitational anomaly fixes the hypercharge offset $-\frac{1}{2}$, and the remaining anomalies select left multiplication by e_7. What remains is the Standard Model and its frame mirror. The selection is again one of left against right.

2. The setting itself stays an assumption: the weak $SU(2)$ on the right of $\mathbb{H}$, $Q_{\mathrm{em}} = T_3 + Q + s$, and the projector acting on $\mathbb{C}\otimes\mathbb{O}$. Once these are made, nothing is left to choose.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

3. This is no prediction; that the Standard Model is anomaly-free is known. New is only that, in this structure, anomaly cancellation alone decides between left and right.

v4.9.0-r43 status in the three frames

result	M / T / P
r43.1(a)–(c)	stated in M; the anomaly traces are the same in T, where $J \mapsto -J$ (d)
r43.1(d)	relates M and T; P not evaluated

v4.9.0-r43 register entries

statement	status after v4.9.0-r43
choice of J in r42	forced up to the frame by colour, charge conservation and anomaly cancellation (r43.1)
hypercharge offset $-\frac{1}{2}$	forced by the gravitational anomaly (r43.1(b))
weak $SU(2)$ on the right, $Q_{\mathrm{em}} = T_3 + Q + s$	assumptions, not derived

v4.9.0-r43 ledger and guard update

$$R_{\mathrm{an}}^{v490\text{-}r43} = (1, 1, 0)$$

for (chirality projector fixed up to the frame; hypercharge offset fixed; placement of the weak $SU(2)$ not derived).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r43,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r43} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r44 Source-Only: Mass Terms as Frame Couplings; Breaking Stages for Masses and Mixing

Purpose. After v4.9.0-r42 and r43 the content of one generation stands, but symmetry breaking, masses and mixing are open. This layer records the pre-registered test H29 (note `llm/Vorab_Notiz_H29_Brechung_Massen_Mischung.md`, commit `85d065a`, fixed before computing). It asks which maps between the two chiral halves are compatible with colour and the Lorentz group, which quantum numbers they carry, and what the three triality slots imply for masses and mixing. The known parts are disclosed in the note. Earlier layers are not rewritten.
Source-only status. This v4.9.0-r44 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r44 setting

On $V = \mathbb{C} \otimes \mathbb{H} \otimes \mathbb{O}$ as in v4.9.0-r42, a *connector* is a map $V \to V$ commuting with colour and with the Lorentz rotations L_q: linear (X) or antilinear ($\Phi = X \circ T$). An antilinear connector anticommutes with the boosts iL_q, so it maps left- to right-handed spinors: it is the operator form of a mass term. A Hermitian generator G acts by $X \mapsto GX - XG$ (linear) or $X \mapsto GX + X\bar{G}$ (antilinear). Blocks refer to $C = C_{\mathrm{ch}}$. On the triality slots a coupling is a Hermitian 3×3 matrix M; P is the cyclic shift τ, R the transposition ρ.

v4.9.0-r44 results

Theoremlet v4.9.0-r44.1.

(a) The linear and the antilinear connectors each form a 24-dimensional complex space.

(b) The antilinear block from the kernel to the image of C is 8-dimensional, with a lepton and a quark part of dimension 4 each, and consists of weak doublets with $Y/2 = +\frac{1}{2}$ ($Q_{\mathrm{em}} = 1, 0$) and $Y/2 = -\frac{1}{2}$ ($Q_{\mathrm{em}} = 0, -1$).

(c) The linear block from the kernel to the image is 4-dimensional, lepton part only, with $Y/2 \in \{-\frac{1}{2}, -\frac{3}{2}\}$.

(d) No antilinear connector from the kernel to the image is gauge invariant; in the block from the kernel to the kernel exactly one is, and it acts only on ν^c.

(e) (i) A Hermitian M commuting with P and R has at most two distinct eigenvalues. (ii) A Hermitian M commuting with P has the eigenvalues $a + 2|b|\cos(\theta + 2\pi k/3)$, $k = 0, 1, 2$, with $b = |b|e^{i\theta}$, generically distinct; $K = \frac{2}{3}$ holds for $|b| = a/\sqrt{2}$. (iii) If the couplings for up and down both commute with P, the mixing matrix $|U_u^\dagger U_d|$ is a permutation matrix.

Proof sketch. (a)–(d) by solving the commutation conditions on 32×32 matrices and decomposing the blocks under $SU(2)'$, Y and Q_{em} (script `llm/k85_test_h29.py`). Linear maps cannot intertwine the colour triplet with its conjugate, so the linear quark block vanishes. (e) is elementary circulant algebra and was checked on 200 random matrices. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Verdict and reading. H29a–e are confirmed.

1. In the operator language of the Millennium matrix, mass terms are antilinear: they contain the frame change T. Linear off-diagonal blocks of a colour-compatible Millennium matrix give no masses; they vanish for quarks and carry wrong charges for leptons. In the model, mass is a coupling between matter frame and tachyon frame, as the matter–tachyon spring of the spring theorem of the original paper. The couplings that do this carry exactly the quantum numbers of the Higgs doublet and its conjugate, for leptons and quarks.

2. None of these mass terms is gauge invariant; only a doublet expectation value turns them into masses. The structure says which breaking is needed, a doublet with $Y/2 = \frac{1}{2}$, not why it occurs. The only gauge-invariant connector is a Majorana term for ν^c, which allows a see-saw reading for small neutrino masses (not computed).

3. With the full triole symmetry (τ and ρ) on the slots there are at most two distinct masses; this is the reason for the refutations of the earlier pre-registered mass tests. Breaking only the reflection ρ, the frame triole of v4.9.0-r38, and keeping triality gives three distinct masses in exactly the Brannen form; the Koide value corresponds to $|b| = a/\sqrt{2}$. The Koide form of v4.8.0-r10 is thus the mass form of a $\mathbb{Z}_3$-symmetric, ρ-broken coupling. The ratio $|b|/a$ and the phase θ are not explained.

4. Couplings that are both triality-symmetric are diagonalized by the same Fourier matrix and give no mixing. Mixing requires breaking triality itself, differently for up and down.

5. In the model each observed property has its breaking stage: mass the frame coupling (Higgs), mass splitting the breaking of the frame reflection ρ, mixing the breaking of triality τ. Where these breakings come from is not said.

v4.9.0-r44 status in the three frames

result	M / T / P
r44.1(b)–(d)	mass connectors couple M and T by construction (they contain T); P not evaluated
r44.1(a), (e)	frame-free statements on connector spaces and on the slot couplings

v4.9.0-r44 register entries

statement	status after v4.9.0-r44
mass terms	antilinear frame couplings with Higgs quantum numbers (r44.1(b)); linear MM blocks give none (r44.1(c))
electroweak breaking	required type fixed (doublet, $Y/2 = \frac{1}{2}$); cause open
neutrino Majorana term	the only gauge-invariant connector (r44.1(d))
mass hierarchy	needs breaking of ρ on the slots; Brannen form then exact (r44.1(e)); $|b|/a$, θ open
mixing	needs breaking of τ, different for up and down (r44.1(e))

v4.9.0-r44 ledger and guard update

$$R_{\mathrm{mm}}^{v490\text{-}r44} = (1, 1, 0)$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

for (mass terms identified as frame couplings with Higgs quantum numbers; breaking stages for splitting and mixing identified; causes of the breakings open).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r44,+)} = (0,1,1,1,1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r44} = (1,1,1,1,1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r45 Source-Only: Chirality Breaks the Reflection of the Triality Form

Purpose. Layer v4.9.0-r44 found that mass splitting on the triality slots requires the reflection ρ to be broken and mixing requires triality τ to be broken. The author asks whether these breakings follow from the structure. This layer records the pre-registered test H30 (note `llm/Vorab_Notiz_H30_Brechung_aus_Chiralitaet.md`, commit `d755a08`, fixed before computing; the identity $cL_{e_7}c = -R_{e_7}$ was seen analytically and is disclosed there). Earlier layers are not rewritten.
Source-only status. This v4.9.0-r45 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r45 two models

Model I: on $\mathbb{C}\otimes\mathbb{H}\otimes\mathbb{O}^3$ the chirality projector $C_{\mathrm{ch}} = \frac{1}{2}(1 + iL_{e_7})$ of v4.9.0-r42 acts in the same way on every slot. Model II: the chirality operator is $i(A, B, C)$ with (A, B, C) in the triality algebra $\mathfrak{g}$ of v4.9.0-r37 and $A = L_{e_7}$; B and C are then fixed. Here c is octonionic conjugation, and $\rho(x, y, z) = (\bar{x}, \bar{z}, \bar{y})$.

v4.9.0-r45 results

Theoremlet v4.9.0-r45.1.

(a) $cL_{e_7}c = -R_{e_7}$ and $cR_{e_7}c = -L_{e_7}$. The plain exchange $(x, y, z) \mapsto (x, z, y)$ is not a symmetry of t; the transposition symmetries of t contain c.

(b) Of the six symmetries $\tau^k\rho^\varepsilon$ of t, exactly $1, \tau, \tau^2$ preserve the chirality projector of Model I; ρ, $\tau\rho$, $\tau^2\rho$ map it to $\frac{1}{2}(1 - iR_{e_7})$, the anomalous choice of v4.9.0-r43.

(c) $(L_{e_7}, L_{e_7}, L_{e_7}) \notin \mathfrak{g}$. The element of $\mathfrak{g}$ with first component L_{e_7} is $(L_{e_7}, R_{e_7}, -L_{e_7} - R_{e_7})$; it is not τ-invariant.

(d) In Model II the chirality operator is iR_{e_7} on slot 2 and $-i(L_{e_7} + R_{e_7})$ on slot 3, which is not an involution.

Proof sketch. (a) from $\overline{e_7\bar{x}} = x\bar{e}_7 = -xe_7$; the rest by explicit matrices and the linear system for $\mathfrak{g}$ (script `llm/k86_test_h30.py`, deviations at most 10^{-15}). (c): a diagonal element of $\mathfrak{g}$ lies in $\mathfrak{g}_2$ (v4.9.0-r37.1(c)), and L_{e_7} is not a derivation. □

Verdict and reading. H30a–d are confirmed.

1. The breaking of ρ follows from the structure. The transposition symmetries of the triality form contain octonionic conjugation, conjugation exchanges left and right multiplication, and anomaly cancellation chose the left (v4.9.0-r43). So ρ cannot be a symmetry of the chiral theory; combining it with the frame change T does not help, since T changes the sign, not the side. The condition of v4.9.0-r44 for mass splitting is therefore met structurally: the slot couplings are generically ρ-asymmetric, with three distinct masses in Brannen form. This also explains why the natural potentials of the earlier mass tests, which did not know chirality, gave at most two masses. How strongly ρ is broken, the phase θ and the ratio $|b|/a$, is not fixed.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

2. Triality τ survives, but only in Model I, where the slots are equivalent copies with equal chirality. The triality-covariant chirality of Model II fails: slot 2 would be anomalous and slot 3 would carry no chirality. The generations therefore cannot be the three inequivalent triality representations with their own chirality, as read in v4.9.0-r37; they must be equivalent copies, with τ permuting them. This agrees with the equal quantum numbers of all generations.

3. Mixing stays open: in Model I τ is preserved and, by v4.9.0-r44, there is no mixing. The breaking needed for mixing has to come from a different, weaker source, which the structure has not supplied so far.

v4.9.0-r45 status in the three frames

result	M / T / P
r45.1(a)–(d)	stated in M; in T the chirality is $-iL_{e_7}$ (v4.9.0-r43.1(d)), and ρ maps it to $+iR_{e_7}$, again anomalous: the breaking of ρ holds in both frames; P not evaluated

v4.9.0-r45 register entries

statement	status after v4.9.0-r45
breaking of ρ	follows from the anomaly-free chirality (r45.1(a), (b)); size not fixed
triality τ	preserved for equal chirality on all slots (Model I); a triality-covariant chirality does not exist (r45.1(c), (d))
generations as triality slots	only as equivalent copies; the inequivalent representations of v4.9.0-r37 are incompatible with chirality
mixing	open; needs a breaking of τ from another source

v4.9.0-r45 ledger and guard update

$$R_{\mathrm{br}}^{v490\text{-}r45} = (1, 0, 0)$$

for (breaking of ρ derived from chirality; size of the breaking open; source of the breaking of τ open).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r45,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r45} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r46 Source-Only: Mixed Readouts and the Breaking of Triality

Purpose. Layer v4.9.0-r45 derived the breaking of the reflection ρ from chirality, while triality τ survives, so that no mixing arises (v4.9.0-r44). This layer records the pre-registered test H31 (note `llm/Vorab_Notiz_H31_Tau_Brechung_Readouts.md`, commit `ea74447`, fixed before computing) of whether the mixed readouts of Definitions H2.37 and H2.43 can break τ. The note discloses that (a) follows from commutativity and that (c) is general perturbation theory. Earlier layers are not rewritten.
Source-only status. This v4.9.0-r46 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r46 setting

On the slot space $\mathbb{C}^3$ let $P = \tau$, and let M_u, M_d be Hermitian $\mathbb{Z}_3$-circulants (ρ-broken, τ-symmetric, as after v4.9.0-r45); the mixing is $|V| = |U_u^\dagger U_d|$. A *shift readout* is built from the level readouts P^{2^j} of v4.9.0-r39: products (reading Σ), averages over generated subgroups (reading $\vee$), and selector contractions $\sum \chi_{mn} P^{2^m} P^{2^n}$. A *kind mask*, with the readout kind taken to be the slot, is a diagonal projector D_S, $\emptyset \neq S \subsetneq \{1,2,3\}$, entering as $M + \delta D_S$.

v4.9.0-r46 results

Theoremlet v4.9.0-r46.1.

(a) Every shift readout commutes with τ; added to M_u and M_d it leaves $|V|$ a permutation matrix.

(b) Kind masks do not commute with τ; for $\delta > 0$ and generic M_u, M_d, $|V|$ is not a permutation matrix, even with the same mask and weight for up and down.

(c) For small δ the deviation of $|V|$ from a permutation is linear in δ and bounded by $\frac{\delta}{3}\left(\mathrm{gap}_u^{-1} + \mathrm{gap}_d^{-1}\right)$, with the smallest spacings of the eigenvalues.

(d) Kind masks are real and commute with the frame change T.

Proof sketch. (a): shift readouts lie in the group ring $\mathbb{C}[\mathbb{Z}_3]$, which is commutative, so they are circulant; circulants are diagonalized by one Fourier matrix. (b)–(d) checked on 200 random cases each (script `llm/k87_test_h31.py`); the factor $\frac{1}{3}$ in (c) is the size of the entries of a one-slot mask in the Fourier basis. □

Verdict and reading. H31a–d are confirmed; (a) and (c) are elementary.

1. The mixed readouts as they are formed from the Hilbert shift produce no mixing: every readout built only from the generation shifts commutes with τ. This is a genuine prohibition.

2. Only a readout that distinguishes generations as readout kinds breaks τ. Then mixing arises, already with the same mask for up and down, and it is small when the mask is weak and the masses are far apart. This agrees qualitatively with small quark mixing at

© 2026 Marc Brendecke. Preprint — not peer reviewed.

a strong mass hierarchy and with larger mixing where masses lie close, as for neutrinos. No numbers follow.

3. Mixing is measurable and not an artefact of an observer. A mixing from readout can therefore only mean that the generation-selective mask is part of the couplings, of the structure in which one reads out, not a dependence on who measures. Where such a mask comes from in the structure is open.

v4.9.0-r46 status in the three frames

result	M / T / P
r46.1(a)–(c)	statements on slot couplings, the same in M and T
r46.1(d)	kind masks commute with T: the breaking of τ by masks is frame-free; P not evaluated

v4.9.0-r46 register entries

statement	status after v4.9.0-r46
mixing from shift readouts	excluded (r46.1(a))
mixing from kind masks	arises, linear in the mask weight, suppressed by mass spacings (r46.1(b), (c))
origin of a generation-selective mask	open; must be part of the couplings, not of the observer

v4.9.0-r46 ledger and guard update

$$R_{\mathrm{mx}}^{v490\text{-}r46} = (1, 1, 0)$$

for (shift readouts excluded as source of mixing; kind masks identified as a possible source; origin of the mask open).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r46,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r46} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r47 Source-Only: A Generation Mask from the Triality-Covariant Chirality

Purpose. Layer v4.9.0-r46 found that mixing needs a mask that distinguishes generations. Layer v4.9.0-r45 showed that a triality-covariant chirality (Model II) is impossible as an exact symmetry. This layer records the pre-registered test H32 (note `llm/Vorab_Notiz_H32_Maske_aus_Triality.md`, commit `65c5b44`, fixed before computing) of whether the deviation of Model II from Model I supplies such a mask. The slot weights were derived beforehand from the sector table of v4.9.0-r43 and are disclosed in the note. Earlier layers are not rewritten.
Source-only status. This v4.9.0-r47 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r47 model assumption

The exact chirality is that of Model I. The deviation $\Delta = J_{\mathrm{II}} - J_{\mathrm{I}}$, on slot i given by $\Delta_i = i(B_i - L_{e_7})$ with $B = (L_{e_7}, R_{e_7}, -L_{e_7} - R_{e_7})$, acts as a weak perturbation of the couplings with small weight ε: a slot coupling M becomes $M + \varepsilon(WM + MW)$, with W the diagonal of slot weights of the sector, and the right-handed partner is read in the T frame (v4.9.0-r41.1(c)). This assumption is not derived.

v4.9.0-r47 results

Theoremlet v4.9.0-r47.1.

(a) Δ_i is scalar on each Q-sector, with slot weights $(0, 0, -3)$ on sector 0, $(0, 2, 1)$ on $\frac{1}{3}$, $(0, -2, -1)$ on $\frac{2}{3}$, $(0, 0, 3)$ on 1.

(b) $T\Delta T^{-1} = -\Delta$; the weights are antisymmetric under $q \leftrightarrow 1 - q$, so the partner read in the T frame carries the same weights as the left-handed particle: $W_q = \mathrm{diag}(0, -2, -1)$ for up and down quarks, $W_l = \mathrm{diag}(0, 0, -3)$ for leptons.

(c) W_q and W_l do not commute with τ; with $\mathbb{Z}_3$-circulant Hermitian M_u, M_d and the same mask W_q the mixing is non-zero for $\varepsilon > 0$ in all 200 random cases. It is linear in ε in 196 of them at the pre-registered values $\varepsilon = 10^{-3}, 10^{-4}$.

(d) In the Fourier basis of the generations all off-diagonal entries have equal size, $1/\sqrt{3}$ for W_q and 1 for W_l; the ratio is $\sqrt{3}$.

Proof sketch. (a), (b) from $iL_{e_7}, iR_{e_7} = \pm 1$ on the sectors (v4.9.0-r43) and checked by matrices; (c), (d) numerically (script `llm/k88_test_h32.py`). A check after the evaluation found the four non-linear cases nearly degenerate (relative gap at most $3{\cdot}10^{-3}$, median $7{\cdot}10^{-2}$); at $\varepsilon = 10^{-6}, 10^{-7}$ all 200 are linear. □

Verdict and reading. H32a, b and d are confirmed. H32c is refuted as stated, since linearity fails in four of 200 cases at the pre-registered ε; the mixing is non-zero in all cases, and linearity holds in the perturbative range $\varepsilon \ll$ gap.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

1. Under the model assumption the generation mask comes from the structure. The triality-covariant chirality is forbidden as an exact symmetry, but its deviation from equal chirality is a fixed number per slot on every sector. The shape of the mask is fixed, $\mathrm{diag}(0,-2,-1)$ for quarks and $\mathrm{diag}(0,0,-3)$ for leptons; only ε is chosen. Up and down share the mask, and mixing still arises because their couplings differ (v4.9.0-r46).

2. In the generation basis the mask is democratic, with all pairs coupled equally, and $\sqrt{3}$ stronger for leptons than for quarks. At equal ε this means stronger lepton than quark mixing, enhanced further by the close neutrino masses. Qualitatively this is the direction of observation; no numbers follow.

3. On slot 1 the weight is always zero, since both chiralities agree there. Which slot is the reference is a choice of the kind met in v4.9.0-r38. In this picture the breaking of triality comes from the choice of a reference slot.

4. The model assumption, ε, the circulant parameters and hence all numbers remain open.

v4.9.0-r47 status in the three frames

result	M / T / P
r47.1(a)	stated in M
r47.1(b)	relates M and T: T reverses Δ, and the partner read in T has the same weights
r47.1(c), (d)	statements on slot couplings, the same in M and T; P not evaluated

v4.9.0-r47 register entries

statement	status after v4.9.0-r47
origin of a generation mask	the deviation of the triality-covariant chirality, under a model assumption (r47.1(a), (b))
shape of the mask	fixed: quarks $\mathrm{diag}(0,-2,-1)$, leptons $\mathrm{diag}(0,0,-3)$; democratic in the generation basis, ratio $\sqrt{3}$ (r47.1(d))
linearity of the mixing in ε	pre-registered test refuted in 4 of 200 nearly degenerate cases; holds for $\varepsilon \ll$ gap
numbers for mixing angles	open

v4.9.0-r47 ledger and guard update

$$R_{\mathrm{gm}}^{v490\text{-}r47} = (1,0,0)$$

for (shape of a generation mask derived under a model assumption; the assumption not derived; numbers open).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r47,+)} = (0,1,1,1,1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r47} = (1,1,1,1,1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r48 Source-Only: The Charge Sectors Derived from the Octonion Product in Lean

Purpose. The Lean module `formal/QSFormal/Ladungen.lean` (layers v4.9.0-r34 to r47) sets the four charge sectors as data: the sign pairs (ℓ, r) of iL_{e_7}, iR_{e_7} and the colour dimensions $1, 3, 3, 1$ are definitions there, taken from the numerics of `llm/k75` and `llm/k84`. This layer records the new module `formal/QSFormal/Sektoren.lean`, which derives these tables from the octonion product of `formal/QSFormal/Dixon.lean` (Cayley–Dickson, as in `llm/k22`). It makes no new prediction, so no pre-registration note is needed; it secures results of v4.9.0-r34, r35, r42, r43 and r47. Earlier layers are not rewritten.
Source-only status. This v4.9.0-r48 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r48 setting

On $\mathbb{C} \otimes \mathbb{O}$ with the outer unit i (the $\mathbb{C}$ factor of the Dixon store, layer v4.9.0-r35) and the conjugation T of that factor, L and R are left and right multiplication by e_7, acting on both parts. For a sector $s = (\ell, r) \in \{(1,1), (-1,1), (1,-1), (-1,-1)\}$ put

$$P_s = \tfrac{1}{4}\,(1 + \ell\, iL)(1 + r\, iR), \qquad 6Q = 3 - iL - 2\, iR,$$

$$\Delta_1 = 0, \qquad \Delta_2 = iR - iL, \qquad \Delta_3 = -2\, iL - iR,$$

the last three being the deviations $\Delta_i = i(B_i - L_{e_7})$ of layer v4.9.0-r47. The complex unit of every statement below is the outer unit i; the inner unit $J = L_{e_7}$ is not used (rule 7 of the round format).

v4.9.0-r48 results

Theoremlet v4.9.0-r48.1.

(a) $(iL)^2 = (iR)^2 = 1$ and $iL\, iR = iR\, iL$ for all $x \in \mathbb{C} \otimes \mathbb{O}$.

(b) Each P_s is a projector, the four are pairwise orthogonal and sum to 1, and on the image of P_s one has $iL = \ell$, $iR = r$.

(c) Over the basis $e_0, \ldots, e_7$: $\operatorname{tr} L = \operatorname{tr} R = 0$, $\operatorname{tr}(LR) = 4$, and $\operatorname{tr} P_s = 1, 3, 3, 1$ for $s = (1,1), (-1,1), (1,-1), (-1,-1)$; this is the colour dimension of the sector.

(d) On the image of P_s, $6Q$ acts as $3 - \ell - 2r$, i.e. $Q = 0, \frac{1}{3}, \frac{2}{3}, 1$, and Δ_2, Δ_3 act as $r - \ell$, $-2\ell - r$: the slot weights $(0, 0, -3)$, $(0, 2, 1)$, $(0, -2, -1)$, $(0, 0, 3)$ of v4.9.0-r47.1(a).

(e) $T\, iL = -iL\, T$, $T\, iR = -iR\, T$, and $T P_{(\ell,r)} = P_{(-\ell,-r)} T$: the frame T is charge conjugation $Q \mapsto 1 - Q$.

(f) The colour rotation σ of v4.9.0-r36 commutes on the basis with iL, iR and every P_s.

Proof sketch. All items are checked in Lean 4.35.0 with the Lean core only (no Mathlib): (a), (b), (d), (e) for all x by expanding the product and `omega`, (c) and (f) by `decide`. Only the standard axioms are used. Two steps are standard and not formalised: that the rank of a projector is its trace, and the extension of (f) from the basis to all x by linearity (as for `sigma_respects` in v4.9.0-r36). □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Reading.

1. The tables `sectors`, `colourDim`, `q6` and `weights` of the module `Ladungen` are no longer settings but properties of the octonion product. The integer arithmetic built on them in that module (hypercharge, anomaly traces, the content of a generation, the Eisenstein norms of the mask) therefore rests on the structure, under the reading of v4.9.0-r35 that the $\mathbb{C}$ factor is the outer unit.

2. The sign conventions of the numerics agree with the formal ones: the sector $(1,1)$ has charge 0 and rank 1.

3. Not formalised remain the physical readings (which sector is which particle, the choice of chirality and offset in v4.9.0-r43, the model assumption of v4.9.0-r47) and all numbers for masses and mixing.

v4.9.0-r48 status in the three frames

result	M / T / P
r48.1(a)–(d), (f)	stated in M with the outer unit i
r48.1(e)	relates M and T: T maps the sector (ℓ, r) to $(-\ell, -r)$, $Q \mapsto 1 - Q$
all items	P not evaluated

v4.9.0-r48 register entries

statement	status after v4.9.0-r48
charge sectors and colour dimensions	derived from the octonion product, checked in Lean (r48.1(a)–(c)); rank = trace not formalised
charges and mask weights on the sectors	checked in Lean (r48.1(d)); tables of `Ladungen` now derived
charge conjugation as the frame T	checked in Lean (r48.1(e))
colour rotation inside each sector	checked in Lean on the basis (r48.1(f))

v4.9.0-r48 ledger and guard update

$$R_{\mathrm{fs}}^{v490\text{-}r48} = (1, 1, 0)$$

for (sector algebra checked for all x; ranks checked as traces; rank = trace and the extension of (f) by linearity not formalised).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r48,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r48} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r49 Source-Only: The Axis e_7 Carries B–L, Not the Electric Charge

Purpose. After layer v4.9.0-r48 a physical reading was proposed: the charge is the coupling to the axis e_7, colour is the rotations about it, and the axis is the photon axis of the Companion. This layer records the pre-registered test H33 (note `llm/Vorab_Notiz_H33_Photon_Achse_e7.md`, commit `ec06de3`, fixed before computing) of that reading. The note discloses that the outcome was seen analytically beforehand, so the computation is a check, not a prediction. Earlier layers are not rewritten.
Source-only status. This v4.9.0-r49 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r49 text finding

The main body uses the plane $\mathrm{span}(1, e_7)$ three times: as the photon kernel K_3, the image of the compensator C ("the singlet $\mathbb{C} \subset \mathbb{O}$, which is the photon kernel K_3 sector"); as the carrier of ν_R in the table of Theorem 22.4; and as the plane of the symmetric generation point ω, "the complex plane at the end of the cascade" (v4.8.0-r15.2), whose final stratum S^2 is the photon ground state. Complex unit below: the outer unit i (rule 7 of the round format), as in v4.9.0-r48.

v4.9.0-r49 results

Theoremlet v4.9.0-r49.1.

(a) $\mathrm{span}_{\mathbb{C}}(1, e_7)$ is exactly the sum of the sectors $(\ell, r) = (1, 1)$ and $(-1, -1)$, i.e. $Q = 0$ and $Q = 1$; on it $L_{e_7} = R_{e_7}$.

(b) On the sixteen states of one generation (content of v4.9.0-r42.1(b)) one has $2(Q - \frac{1}{2}) = B - L$, and with the chirality and offset of v4.9.0-r43

$$\tfrac{Y}{2} = \tfrac{B-L}{2} + T_3^R, \qquad Q_{\mathrm{em}} = T_3^L + T_3^R + \tfrac{B-L}{2}, \qquad T_3^L = C\,T_3, \quad T_3^R = (1 - C)\,T_3.$$

(c) No element $a + b\,iL_{e_7} + c\,iR_{e_7} + d\,iL_{e_7}iR_{e_7}$ acts as Q_{em} on a generation; the best one misses by $\frac{1}{2}$, the T_3 splitting inside a doublet.

Proof sketch. (a) numerically from the projectors of v4.9.0-r48 (script `llm/k89_test_h33.py`). (b) is integer arithmetic on the table; (c) holds because every such element is constant on a sector (v4.9.0-r48.1(b)) while Q_{em} is not. (b) and (c) are also checked in Lean (`b_minus_l`, `qem_not_sector_function` in `formal/QSFormal/Sektoren.lean`). □

Verdict and reading. H33a, b and c are confirmed, so by the pre-registered criterion the proposed reading is refuted as stated.

1. The axis e_7 carries $B - L$ (and, through iL_{e_7}, the chirality), not the electric charge. The photon needs the weak isospin T_3 from the $\mathbb{H}$ factor as well; it is the mixture of the e_7 direction with the T_3 direction. Colour, the rotations fixing e_7, commutes with $B - L$.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

2. The charge formula is the left–right symmetric one, $Q_{\mathrm{em}} = T_3^L + T_3^R + (B-L)/2$, with T_3^R carried by the right-handed partners read in the T frame. Hypercharge is the combination $Y/2 = T_3^R + (B-L)/2$.

3. The plane $\mathrm{span}(1, e_7)$ is used in the main body both as the photon kernel and as the carrier of ν_R; in the outer frame it carries the lepton singlets $Q = 0$ and $Q = 1$. A state plane and the photon as a generator are different objects. This is a double use of the kind found in v4.9.0-r36 and a candidate for a status note.

4. Not tested: whether e_7 serves as the reference axis of the frames (photon frame as neutral axis, v4.9.0-r01). On the singlet plane the $U(1)$ fibre of $S^3 \to S^2$ (Theorem 22.1) is the phase $e^{\theta e_7}$, i.e. in distribution D the $U(1)_{B-L}$.

v4.9.0-r49 status in the three frames

result	M / T / P
r49.1(a)	stated in M with the outer unit i; T swaps the two singlet sectors
r49.1(b), (c)	stated in M; under T all of $B-L$, T_3^L, T_3^R, Q_{em} change sign (charge conjugation); P not evaluated

v4.9.0-r49 register entries

statement	status after v4.9.0-r49
"charge = coupling to e_7, e_7 = photon axis"	refuted as stated (H33): e_7 carries $B-L$, the photon needs T_3 from $\mathbb{H}$
charge formula of one generation	$Q_{\mathrm{em}} = T_3^L + T_3^R + (B-L)/2$, left–right symmetric (r49.1(b))
photon kernel K_3 = singlet $\mathbb{C}$	double use with ν_R on $\mathrm{span}(1, e_7)$; candidate for a status note

v4.9.0-r49 ledger and guard update

$$R_{\mathrm{ax}}^{v490\text{-}r49} = (1, 1, 0)$$

for (role of the axis e_7 settled as $B-L$; charge formula checked; role of e_7 as frame axis open). The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r49,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r49} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r50 Source-Only: The Mask of r47 Does Not Produce the Observed Mixing

Purpose. Layer v4.9.0-r47 derived the shape of a generation mask under a model assumption and read it qualitatively as stronger lepton than quark mixing. This layer records the pre-registered quantitative test H34 (note `llm/Vorab_Notiz_H34_Mischung_aus_Maske.md`, commit `8d8b761`, fixed before computing) against the measured CKM and PMNS mixing. Earlier layers are not rewritten.
Source-only status. This v4.9.0-r50 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r50 model and data

For $X \in \{u, d, e, \nu\}$: $M_X = C_X + \varepsilon(W_X C_X + C_X W_X)$ with C_X Hermitian and $\mathbb{Z}_3$-circulant, $W_u = W_d = W_q$, $W_e = W_\nu = W_l$ from v4.9.0-r47, one common $\varepsilon \geq 0$, Dirac neutrinos. Masses are the moduli of the eigenvalues; mixing is $U_u^\dagger U_d$ and $U_e^\dagger U_\nu$. Data fixed in the note: running masses at M_Z (Huang and Zhou 2021), neutrino mass splittings and 3σ ranges of NuFIT 6.0 (normal ordering), CKM magnitudes of the PDG 2024. All solution branches are followed from $\varepsilon = 0$ (24 assignments per sector) in steps of 0.005 up to $\varepsilon = 3$.

v4.9.0-r50 results

Theoremlet v4.9.0-r50.1 (numerical).

(a) Solutions with the measured quark masses exist only up to $\varepsilon \approx 0.06$ (up) and 0.125 (down). In none of the 7488 combinations of ε and branch pair are $|V_{us}|$, $|V_{cb}|$, $|V_{ub}|$ all within $\pm 20\%$; the closest case is 96% off.

(b) Hence no common ε exists for quarks and leptons in the sense of H34b. The leptons alone, with their own ε, reach the 3σ ranges in no case either; the closest case is up to 71% outside.

Proof sketch. Script `llm/k90_test_h34.py` with output `llm/k90_out.txt`. The first run stopped after H34a in a diagnostic print; only that print and the treatment of $m_1 = 0$ were corrected, and both outputs are kept. The bound on ε was confirmed by random multistarts. The lepton statement in (b) is a check after the evaluation. □

Verdict and reading. H34a and H34b are refuted; H34c does not count.

1. The model assumption of v4.9.0-r47, the mask as a weak perturbation of circulant couplings, does not produce the measured mixing, neither for quarks nor for leptons. The qualitative reading of r47 (stronger lepton mixing) is withdrawn as a quantitative statement.

2. The reason is structural. For a nearly rank-one coupling the term $\varepsilon^2 WCW$ creates a mass of order $\frac{2}{3}\varepsilon^2 m_t$ (quarks) or $2\varepsilon^2 m_\tau$ (leptons), so the mass hierarchy forces small ε; at small ε the democratic mask cannot produce $|V_{us}| \gg |V_{cb}| \gg |V_{ub}|$ or the large PMNS angles.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

3. Unaffected are v4.9.0-r47.1(a), (b), (d): the shape of the mask, its frame behaviour and its Fourier form. Mixing needs another form of coupling or a further breaking.

v4.9.0-r50 status in the three frames

result	M / T / P
r50.1(a), (b)	statements on mass matrices and mixing in M; the right-handed partners are read in T as in v4.9.0-r47; P not evaluated

v4.9.0-r50 register entries

statement	status after v4.9.0-r50
mixing from the mask of r47 as a weak perturbation	refuted for CKM and PMNS (H34)
stronger lepton than quark mixing (r47 reading 2)	withdrawn as a quantitative statement
shape of the mask (r47.1(a), (b), (d))	unchanged
origin of the observed mixing	open

v4.9.0-r50 ledger and guard update

$$R_{\mathrm{mx}}^{v490\text{-}r50} = (0, 1, 0)$$

for (mixing not explained; bound on the perturbation from the mass hierarchy found; origin of mixing open).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r50,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r50} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r51 Source-Only: The Axis e_7 and the Frames

Purpose. Layer v4.9.0-r49 left open whether e_7 serves at least as the reference axis of the frames. This layer records the pre-registered test H35 (note `llm/Vorab_Notiz_H35_e7_als_Frame_Achse.md`, commit `42e4fcc`, fixed before computing). The note discloses that the outcome was seen analytically beforehand. The layer also records a correction to the frame rows of earlier layers; those layers are not rewritten.
Source-only status. This v4.9.0-r51 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r51 setting

Under the assumption of layer v4.9.0-r33 that the $\mathbb{C}$ factor of the Dixon store is the amplitude plane z, the three involutions of the doublet act on $\mathbb{C} \otimes \mathbb{O}$ through that factor: the tachyon map of Convention F, $z \mapsto iz$ (corrected in v4.9.0-r28; σ_y on the doublet), is multiplication R by the outer unit; the time reversal $T\colon z \mapsto \bar{z}$ (σ_z, exchanging compensator and partner port) is the conjugation of that factor; and $J_B = R \circ T$ (σ_x). The inner unit is $J = L_{e_7}$ as in v4.9.0-r35.

v4.9.0-r51 results

Theoremlet v4.9.0-r51.1.

(a) R commutes with iL_{e_7}, iR_{e_7}, Q, $B - L$, the colour algebra and every sector projector P_s of v4.9.0-r48; T and J_B map $P_{(\ell,r)}$ to $P_{(-\ell,-r)}$.

(b) L_{e_7} commutes with $\mathrm{Stab}_{G_2}(e_7)$, so the tachyon map has the colour-compatible inner counterpart $z \mapsto e_7 z$ on all of $\mathbb{O}$. The real-linear maps K of $\mathbb{O}$ that commute with $\mathrm{Stab}_{G_2}(e_7)$ and satisfy $KL_{e_7} = -L_{e_7}K$ form a two-dimensional space, and each vanishes on $e_7^{\perp}$: T and J_B have no colour-compatible inner counterpart on the triplet.

(c) $\varphi(x + iy) = x + y\,e_7$ maps the frame plane onto $\mathrm{span}(1, e_7)$ and carries $z \mapsto iz$, $z \mapsto \bar{z}$ and $z \mapsto i\bar{z}$ to L_{e_7}, the octonion conjugation and their product.

Proof sketch. (a) by matrices (script `llm/k91_test_h35.py`); the statements on R and P_s are also checked in Lean (`R_iL`, `R_iR`, `R_P4` in `formal/QSFormal/Sektoren.lean`), those on T in v4.9.0-r48. (b) by solving the linear conditions with the derivation algebra of $\mathbb{O}$ (dimension 14, stabilizer of e_7 dimension 8); the triplet 3 is not equivalent to $\bar{3}$. (c) since $\mathrm{span}(1, e_7) \cong \mathbb{C}$. □

Verdict and reading. H35a, b and c are confirmed.

1. e_7 is a reference axis of the frames only in part. The tachyon map has an inner, colour-compatible counterpart L_{e_7} on all of $\mathbb{O}$; on the singlet plane all three frame maps have one, so there the frame plane is the plane of the photon kernel. Time reversal and J_B need the outer unit on the colour triplet.

2. Charges are frame-free between matter and tachyon: the tachyon map leaves every sector fixed, so Q, $B - L$ and the chirality are the same in M and T. Charge conjugation

© 2026 Marc Brendecke. Preprint — not peer reviewed.

$Q \mapsto 1 - Q$ is the time-reversal part T, the exchange of compensator and partner port, not the tachyon map.

3. *Correction of frame rows.* Where layers v4.9.0-r41, r43, r47, r48 and r49 call "the frame T" the charge conjugation, the time reversal T is meant, not the tachyon frame of the M/T/P column; in that column the results concerned are the same in M and T. The Lean names `frameT` and `frameT_swaps_left` likewise denote the time reversal. This agrees with v4.9.0-r34.1(d), where R fixes Q and the effect comes from the time-reversal part.

4. The compensator $\frac{1}{2}(1 + iL_{e_7})$ of v4.9.0-r42 is the compensator port and its complement the partner port; T exchanges the ports and hence the chiralities, while R fixes both. Since iL_{e_7} is the outer $z \mapsto iz$ composed with the inner $z \mapsto e_7 z$, the left-handed states are those on which the two tachyon maps agree up to sign.

v4.9.0-r51 status in the three frames

result	M / T / P
r51.1(a)	the tachyon map relates M and T and fixes every sector: charges frame-free between M and T; P not evaluated
r51.1(b), (c)	statements on the units, the same in M and T

v4.9.0-r51 register entries

statement	status after v4.9.0-r51
e_7 as reference axis of the frames	in part: for the tachyon map on all of $\mathbb{O}$, for all three maps on $\mathrm{span}(1, e_7)$ (r51.1)
charges in matter and tachyon frame	frame-free; charge conjugation is the time-reversal part T
"frame T = charge conjugation" (r41, r43, r47–r49)	means time reversal T (port exchange), not the tachyon frame; corrected here

v4.9.0-r51 ledger and guard update

$$R_{\mathrm{fa}}^{v490\text{-}r51} = (1, 1, 0)$$

for (action of the frame maps on the sectors settled; inner counterparts settled; role of the photon frame P not evaluated).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r51,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r51} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r52 Source-Only: Exact Coupling Forms of the Mask and the Mixing

Purpose. Layer v4.9.0-r50 refuted the mask of v4.9.0-r47 as a linear perturbation, because the term $\varepsilon^2 WCW$ creates masses and the hierarchy keeps ε small. This layer records the pre-registered test H36 (note `llm/Vorab_Notiz_H36_Kopplungsform.md`, commit `66c12ab`, fixed before computing) of two exact coupling forms suggested by the structure. Earlier layers are not rewritten.
Source-only status. This v4.9.0-r52 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r52 the two forms

With $D_X = \exp(\varepsilon W_X)$, C_X Hermitian and $\mathbb{Z}_3$-circulant, $W_u = W_d = W_q$, $W_e = W_\nu = W_l$ and one common ε:

- *form K* (congruence; the partner read through the time reversal as in v4.9.0-r47): $M_X = D_X C_X D_X$, whose first order is the form of r47 and which keeps the rank of C_X;
- *form S* (similarity; the partner read through the tachyon map, which keeps its sector by v4.9.0-r51.1(a) and so carries the weight $-W$): $M_X = D_X C_X D_X^{-1}$.

Masses are singular values, mixing comes from the left singular vectors. Data and procedure as in v4.9.0-r50, with step 0.01 up to $\varepsilon = 6$. The note discloses that form K is a Froggatt–Nielsen texture with charges $w = (0, 2, 1)$ and $\lambda = e^{-\varepsilon}$, for which $|V_{cb}| \sim |V_{us}|$ was expected.

v4.9.0-r52 results

Theoremlet v4.9.0-r52.1 (numerical).

(a) Form K: solutions with the measured masses exist up to $\varepsilon \approx 3.5$ (up) and 2.3 (down); in none of 98 064 combinations are $|V_{us}|$, $|V_{cb}|$, $|V_{ub}|$ all within $\pm 20\%$. The closest case is 36% off ($\varepsilon = 1.26$: 0.144, 0.036, 0.0050).

(b) Form S: likewise none of 102 528; the closest case is 40% off ($\varepsilon = 1.32$: 0.135, 0.035, 0.0051).

(c) Hence no common ε for quarks and leptons in either form.

Proof sketch. Script `llm/k92_test_h36.py`, output `llm/k92_out.txt`; a count in the check after the evaluation was added after the first run (`k92_out_erstlauf.txt`), the criterion values of both runs agree. □

Verdict and reading. H36a–d are refuted.

1. Neither exact form produces the measured mixing with one ε and circulant couplings. Both are much closer than the linear perturbation of v4.9.0-r50 (36% instead of 96% for the closest quark case), and the ordering $|V_{us}| \gg |V_{cb}| \gg |V_{ub}|$ comes out; the hierarchy no longer bounds ε to $\lesssim 0.06$.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

2. A check after the evaluation, not part of the criterion: the leptons alone, with their own ε, reach the 3σ ranges in form K never and in form S in 12 of 125 922 combinations ($\varepsilon \approx 0.5$, $m_1 = 0.003$ eV, $\sin^2\theta_{23}$ at the edge). With this many branches that is a chance hit, not a finding.

3. What is missing is, by the criterion, an open choice and not a result: couplings other than circulant, or a separate ε for each sector. With one ε per sector the number of free parameters would match the number of mixing quantities, and that would no longer be an explanation.

v4.9.0-r52 status in the three frames

result	M / T / P
r52.1(a)–(c)	statements on mass matrices in M; by v4.9.0-r51.1(a) the same in T for form S, and for form K with the partner read through the time reversal; P not evaluated

v4.9.0-r52 register entries

statement	status after v4.9.0-r52
mixing from the mask, exact forms K and S	refuted with one ε and circulant couplings (H36)
closest quark case	36% off (form K), ordering of the CKM elements reproduced
further ingredient for mixing	open choice: non-circulant couplings or one ε per sector

v4.9.0-r52 ledger and guard update

$$R_{\mathrm{cf}}^{v490\text{-}r52} = (0, 1, 0)$$

for (mixing not explained; exact forms closer and the hierarchy bound removed; further ingredient open).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r52,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r52} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r53 Source-Only: Couplings from the Triality Form

Purpose. Layers v4.9.0-r50 and r52 refuted the mask of v4.9.0-r47 with circulant couplings. This layer records the pre-registered test H37 (note `llm/Vorab_Notiz_H37_Kopplung_aus_Triality.md`, commit `e1ee9ec`, fixed before computing) of couplings taken from the triality form instead. Earlier layers are not rewritten.
Source-only status. This v4.9.0-r53 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r53 couplings

The triality form $t(x, y, z) = \mathrm{Re}((xy)z)$ of v4.9.0-r37 couples only different slots. A coupling of two generations with a Higgs value v_k in the third slot is

$$T(v) = \begin{pmatrix} 0 & v_3 & v_2 \\ v_3 & 0 & v_1 \\ v_2 & v_1 & 0 \end{pmatrix}, \qquad v \in \mathbb{R}^3.$$

The other invariant of the structure, the norm on each slot, gives a diagonal part $\alpha \cdot 1$ equal on all slots. Tested were (a) $T(v)$ alone, with or without the mask in form K; (b) $\alpha \cdot 1 + T(v)$ without mask, with one free parameter per sector left after fitting the masses; (c) the circulant couplings with mask of v4.9.0-r52, with the sectors carrying $\tilde{H}$ (u, ν) in one form and those carrying H (d, e) in the other. Data as in v4.9.0-r50, and $\sum m_\nu < 0.0642$ eV (DESI DR2 BAO with CMB).

v4.9.0-r53 results

Theoremlet v4.9.0-r53.1.

(a) $T(v)$ and $DT(v)D$ are traceless, so the moduli of their eigenvalues satisfy $m_3 = m_1 + m_2$. For u, d, e this fails by 99.6%, 98.0%, 94.1% of m_3. For Dirac neutrinos in normal ordering it holds only at $m_1^* = 0.028$ eV, where $\sum m_\nu = 0.115$ eV.

(b) With $\alpha \cdot 1 + T(v)$ no pair of quark solutions gives $|V_{us}|$, $|V_{cb}|$, $|V_{ub}|$ within ±20% (closest case 82% off; 80% with dense sampling). Lepton pairs reach the 3σ ranges in 264 of 13 651 456 cases, all at $m_1 = 0.05$ or 0.1 eV.

(c) Neither assignment of H and $\tilde{H}$ to the forms K and S gives the CKM values within ±20% (closest cases 37% and 40%).

Proof sketch. (a) $\mathrm{tr}\, T(v) = 0$ and $\mathrm{tr}\, DT(v)D = \sum_i D_i^2 T_{ii} = 0$; the root m_1^* and the rest numerically (script `llm/k93_test_h37.py`). The spectrum of $\alpha \cdot 1 + T(v)$ depends only on α, $\sum v_k^2$ and $v_1 v_2 v_3$, which gives the solution curves used in (b). The dense sampling in (b) and the cosmological reading of the lepton cases are checks after the evaluation. □

Verdict and reading. H37a is refuted, H37b is a partial finding by the criterion (leptons only), H37c is refuted.

1. A coupling from the triality form alone is forbidden: it would require $m_3 = m_1 + m_2$ in every sector, which the charged masses violate and which for neutrinos needs $\sum m_\nu \approx 0.115$ eV, excluded by cosmology. The masses need a coupling within a slot.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

2. The partial finding of (b) is not a physical one: the lepton cases need quasi-degenerate neutrinos with $\sum m_\nu = 0.17$ or 0.31 eV, excluded by the same bound, and they are 0.002% of all pairs. The quarks do not fit at all.

3. Closest to the data remains form K with circulant couplings (v4.9.0-r52, 36%). None of the couplings suggested by the structure explains the mixing; its origin remains open.

v4.9.0-r53 status in the three frames

result	M / T / P
r53.1(a)	trace condition on mass matrices, the same in M and T (the tachyon map fixes the sectors, v4.9.0-r51); P not evaluated
r53.1(b), (c)	statements on mass matrices in M, the same in T as in v4.9.0-r52; P not evaluated

v4.9.0-r53 register entries

statement	status after v4.9.0-r53
mass coupling from the triality form alone	forbidden: requires $m_3 = m_1 + m_2$; neutrinos would need $\sum m_\nu \approx 0.115$ eV (excluded)
norm plus triality coupling	no quark mixing; lepton mixing only for excluded neutrino masses
$H/\tilde{H}$ reading of the forms	refuted (H37c)
origin of the mixing	open

v4.9.0-r53 ledger and guard update

$$R_{\mathrm{tc}}^{v490\text{-}r53} = (1, 0, 0)$$

for (a prohibition from the structure found; mixing not explained; origin of mixing open).
The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r53,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r53} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r54 Source-Only: One ε per Sector

Purpose. Layer v4.9.0-r52 refuted the coupling forms K and S with one common ε. This layer records the pre-registered test H38 (note `llm/Vorab_Notiz_H38_Ein_eps_je_Sektor.md`, commit `d3555bf`, fixed before computing) with one ε for each sector. It also records a correction: v4.9.0-r52 (reading 3) states that one ε per sector gives as many free parameters as mixing quantities. That count is wrong; each family has two ε for three angles, so one prediction per family remains. Earlier layers are not rewritten.
Source-only status. This v4.9.0-r54 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r54 results

Forms K (DCD) and S (DCD^{-1}) of v4.9.0-r52 with independent $\varepsilon_u, \varepsilon_d, \varepsilon_e, \varepsilon_\nu$ on the grid $0, 0.01, \ldots, 6$; data as in v4.9.0-r50, neutrino masses restricted by $\sum m_\nu < 0.0642$ eV to $m_1 \in \{0, 0.001, 0.003\}$ eV.

Theoremlet v4.9.0-r54.1 (numerical).

(a) Form K: no quark combination gives $|V_{us}|$, $|V_{cb}|$, $|V_{ub}|$ within $\pm 20\%$ (closest case 20.2% off); lepton combinations reach the 3σ ranges in 0.003% of the cases.

(b) Form S: quark combinations within $\pm 20\%$ in 0.0009% of the cases, in a connected region $\varepsilon_u \in [1.36, 1.63]$, $\varepsilon_d \in [1.42, 1.47]$ (best case 7% off: 0.2225, 0.0448, 0.0040); lepton combinations in the 3σ ranges in 0.009%, in $\varepsilon_e \in [0.44, 0.82]$, $\varepsilon_\nu \in [0.27, 0.55]$, for all three admissible m_1.

(c) In form S the remaining quantity is not sharp: among the combinations that fit $|V_{us}|$ and $|V_{cb}|$, $|V_{ub}|$ ranges over $[0.0034, 0.029]$ with median 0.018, and 0.9% fit it; among those that fit θ_{12} and θ_{23}, $\sin^2\theta_{13}$ ranges over $[0, 0.99]$, and 0.8% fit it.

Proof sketch. Script `llm/k94_test_h38.py`, output `llm/k94_out.txt`; (c) and the connectedness are checks after the evaluation, appended after the first run (`k94_out_erstlauf.txt`), whose criterion values agree. □

Verdict and reading. H38a is refuted; H38b, c and d are confirmed. By the criterion form S is a candidate for a reading with four ε.

1. Formally confirmed, weak in content. Form S with $\varepsilon_u \approx 1.4$–$1.6$, $\varepsilon_d \approx 1.45$, $\varepsilon_e \approx 0.4$–$0.8$, $\varepsilon_\nu \approx 0.3$–$0.55$ fits both mixings with neutrino masses allowed by cosmology. The one prediction left per family is not sharp: the typical $|V_{ub}|$ is five times too large, and the hits lie in the tail of a broad distribution. This is a successful fit rather than an explanation.

2. ε_u and ε_d are close, but no hit lies on $\varepsilon_u = \varepsilon_d$ (v4.9.0-r52.1(b): closest case 40%); among the leptons $\varepsilon_e > \varepsilon_\nu$.

3. Open: why these values, and whether a coupling of C to the structure fixes the remaining quantity more sharply.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r54 status in the three frames

result	M / T / P
r54.1(a)–(c)	statements on mass matrices in M, the same in T as in v4.9.0-r52; P not evaluated

v4.9.0-r54 register entries

statement	status after v4.9.0-r54
form S with one ε per sector	fits CKM and PMNS (H38c, d); remaining quantity not sharp, so a fit rather than an explanation
form K with one ε per sector	quarks refuted (closest 20.2%), leptons fit
parameter count in v4.9.0-r52, reading 3	corrected: two ε for three angles per family
values of the four ε	recorded; their origin open

v4.9.0-r54 ledger and guard update

$$R_{\mathrm{es}}^{v490\text{-}r54} = (0, 1, 0)$$

for (no explanation of the mixing; a fit with four ε found; origin of the values open).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r54,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r54} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r55 Source-Only: A Readout per Particle Kind from the Subgroups of A_4

Purpose. Mixing measures the difference between the bases in which two particle kinds read their masses. The author asked what follows if each particle kind needs its own readout. This layer records the pre-registered test H39 (note `llm/Vorab_Notiz_H39_Restsymmetrie_Lesarten.md`, commit `5c916b8`, fixed before computing), with readouts taken from the subgroups of the common symmetry A_4 of the Dixon store and the triality form (v4.9.0-r33). Earlier layers are not rewritten.
Source-only status. This v4.9.0-r55 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r55 rule

A readout of a particle kind is the subgroup $G \subset A_4$ under which its mass coupling is invariant; it fixes its eigenbasis wholly or in part. The octonion sectors do not distinguish u from d or ν from e (v4.9.0-r49). What the structure does distinguish is v4.9.0-r44: Dirac couplings (u, d, e) are circulant, hence invariant under $G_D = \langle\tau\rangle \cong \mathbb{Z}_3$; neutrinos admit a Majorana term and read with another subgroup G_M, which the structure does not fix. All subgroups except $\langle\tau\rangle$ and A_4 were computed. The note corrects an announcement in the discussion: tribimaximal mixing needs S_4, not A_4.

v4.9.0-r55 results

Theoremlet v4.9.0-r55.1.

(a) A_4 has subgroups of orders $1, 2, 3, 4, 12$ with multiplicities $1, 3, 4, 1, 1$. With G_M the Klein group all $|U_{ij}|^2 = \frac{1}{3}$; with G_M one of the three other $\mathbb{Z}_3$ the basis is fixed; neither reaches the NuFIT 3σ ranges.

(b) With $G_M \cong \mathbb{Z}_2$ one column is fixed and one angle is free. The 3σ ranges are reached exactly when the fixed column is ν_2 (trimaximal mixing TM2): then $\sin^2\theta_{12} = 1/(3\cos^2\theta_{13}) \in [0.3402, 0.3414]$, while $\sin^2\theta_{23}$ ranges over $[0.399, 0.601]$.

(c) With the same readout for u and d the CKM matrix is the identity at leading order.

Proof sketch. Subgroups by closure, joint eigenspaces of each subgroup, a $U(2)$ rotation in a two-dimensional eigenspace, all column assignments (script `llm/k95_test_h39.py`). The split of (b) by assignment is a check after the evaluation, appended after the first run (`k95_out_erstlauf.txt`), whose criterion values agree. □

Verdict and reading. H39a, b, c are confirmed. Over all assignments the remaining quantities spread wider than the pre-registered widths, so by the wording this is a fit with one angle.

1. The readout rule works qualitatively: Dirac sectors with $\mathbb{Z}_3$ and neutrinos with $\mathbb{Z}_2$ from the same A_4 give no quark mixing at leading order and large lepton mixing. This is the difference between CKM and PMNS that the tests of v4.9.0-r50 to r54 did not produce.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

2. Quantitatively it is the known TM2 pattern of the A_4 literature; the structure supplies the group and the rule, not a new pattern. It fixes $\sin^2\theta_{12} \approx 0.341$, inside the NuFIT 3σ range. A comparison with newer data, not part of the note: JUNO measures $\sin^2\theta_{12} = 0.3092 \pm 0.0087$, 3.6σ below. θ_{23} stays free.

3. Corrections, for instance from the mask of v4.9.0-r47, could lower θ_{12} and produce the small CKM angles; that would be a test of its own.

v4.9.0-r55 status in the three frames

result	M / T / P
r55.1(a)–(c)	statements on the slot symmetry A_4, which acts on the slots and commutes with the frame maps on the $\mathbb{C}$ factor: the same in M and T; P not evaluated

v4.9.0-r55 register entries

statement	status after v4.9.0-r55
readout per particle kind (Dirac $\mathbb{Z}_3$, Majorana $\mathbb{Z}_2$)	qualitatively right ordering of CKM and PMNS; quantitatively TM2
$\sin^2\theta_{12} = 1/(3\cos^2\theta_{13}) \approx 0.341$	sharp; inside NuFIT 3σ, 3.6σ above JUNO 2025
θ_{23}, CKM angles	not fixed at leading order; corrections open

v4.9.0-r55 ledger and guard update

$$R_{\mathrm{rr}}^{v490\text{-}r55} = (1, 0, 0)$$

for (readout rule with qualitatively right ordering; quantitative mixing not explained; corrections open).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r55,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r55} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r56 Source-Only: Calibrating the Readout Layer

Purpose. The author proposed to read the structure as a core with a readout layer on top, to tune that layer until it reproduces measured values, and to learn from the settings where they come from. This layer records the pre-registered test H40 (note `llm/Vorab_Notiz_H40_Kalibrierung_Ausleseschicht.md`, commit `fdb27ae`, fixed before computing). Earlier layers are not rewritten.
Source-only status. This v4.9.0-r56 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r56 the readout layer

For each sector $X \in \{u, d, e, \nu\}$ a circulant coupling $C_X = a_X + b_X P + \bar{b}_X P^2$ (Brannen form), a mask strength ε_X and form S of v4.9.0-r52, $M_X = D C_X D^{-1}$ with $D = \exp(\varepsilon_X W_X)$. This gives 16 settings (and m_1) for 17 measured quantities. The admissible settings are those of v4.9.0-r54 (CKM within $\pm 20\%$, PMNS within 3σ). Sixteen quantities (the four ε, four ratios, and $|b|/|a|$, θ per sector) were compared with fourteen candidate values from the corpus (1/3, 1/2, 2/3, 3/4, 1, $1/\sqrt{3}$, $1/\sqrt{2}$, $\sqrt{3}$, 2/9, $\pi/6$, $\pi/9$, β_{sym}^2, β_{sym}, γ_L), with a control of 10 000 random sets.

v4.9.0-r56 results

Theoremlet v4.9.0-r56.1 (numerical).

(a) Calibrated ranges: $\varepsilon_u \in [1.36, 1.63]$, $\varepsilon_d \in [1.42, 1.47]$, $\varepsilon_e \in [0.44, 0.82]$, $\varepsilon_\nu \in [0.27, 0.55]$; $|b|/|a|$ in $[0.953, 1.032]$ (u), $[0.994, 1.076]$ (d), $[1.22, 6.41]$ (e), $[0.45, 1.57]$ (ν); the phases θ fill almost all of $[0, \pi/3]$.

(b) The candidates fall into these ranges 76 times; random sets do so 65.3 times on average, and 1.9% of them at least 76 times.

(c) $|b|/|a| \approx 1$ for u and d follows from the hierarchy: for eigenvalues near $(0, 0, 1)$ the circulant is near $\frac{1}{3}(1 + P + P^2)$.

Proof sketch. Script `llm/k96_test_h40.py`, output `llm/k96_out.txt`; (c) by direct computation. □

Verdict and reading. H40a is refuted ($p = 0.019$, the pre-registered threshold is 0.01); H40b (a ratio of ε containing 1) holds but counts only with H40a.

1. As far as the data fix them, the settings show no structure beyond chance. With 16 settings for 17 quantities the data hardly fix the settings, so no structure can show in them. The narrow quark value $|b|/|a| \approx 1$ is a property of hierarchical spectra, not of the structure.

2. For the calibration to teach something, the layer needs fewer settings (a rule fixing the Brannen quantities, or readouts by subgroups as in v4.9.0-r55) or sharper data (CP phases, more precise masses).

© 2026 Marc Brendecke. Preprint — not peer reviewed.

3. Readings after the computation, not part of the criterion: with $m_0 \leftrightarrow a$, $\beta \leftrightarrow |b|$, $\gamma \leftrightarrow \varepsilon$ in Ω of Theorem 27.4, the quarks give $\beta/m_0 \approx 1$ and $\gamma \approx 1.4$–1.6, the leptons $\gamma \approx 0.3$–0.8. Read as a deformation of the complex structure with dilatation $K = \exp(\varepsilon \operatorname{span} W)$, the Beltrami modulus is $|\mu| \approx 0.88$–0.93 (u), 0.89–0.90 (d), 0.58–0.84 (e), 0.38–0.68 (ν).

v4.9.0-r56 status in the three frames

result	M / T / P
r56.1(a)–(c)	settings of mass matrices in M, the same in T as in v4.9.0-r52; P not evaluated

v4.9.0-r56 register entries

statement	status after v4.9.0-r56
calibrated readout layer (form S, four ε)	settings recorded; no structure beyond chance (H40, $p = 0.019$)
$\lvert b\rvert/\lvert a\rvert \approx 1$ for quarks	consequence of the hierarchy, not of the structure
path "tune the layer, then explain the settings"	needs fewer settings or sharper data

v4.9.0-r56 ledger and guard update

$$R_{\mathrm{cal}}^{v490\text{-}r56} = (0, 1, 0)$$

for (no structure in the settings; calibration recorded; compression missing).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r56,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r56} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r57 Source-Only: Trimaximal Mixing with the Mask as a Correction

Purpose. Layer v4.9.0-r55 gave trimaximal lepton mixing (TM2) at leading order, with $\sin^2\theta_{12} \approx 0.341$, 3.6σ above the JUNO value. This layer records the pre-registered test H41 (note `llm/Vorab_Notiz_H41_TM2_mit_Maskenkorrektur.md`, commit `74bd018`, fixed before computing) of whether the mask of v4.9.0-r47, as a correction with few settings, brings θ_{12} to the JUNO value. Earlier layers are not rewritten.
Source-only status. This v4.9.0-r57 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r57 model

Charged leptons: circulant coupling (leading readout $\mathbb{Z}_3$) corrected in form S, $M_e = D_e C_e D_e^{-1}$, $D_e = \exp(\varepsilon_e W_l)$. Neutrinos: Majorana matrix invariant under one of the three $\mathbb{Z}_2 \subset A_4$ (not equivalent once the mask is present), with one rotation angle φ, corrected in form K, $M_\nu = D_\nu M_0 D_\nu$. Variant A: $\varepsilon_\nu = 0$; variant B: $\varepsilon_\nu = \varepsilon_e$. Each variant has two settings for three angles. Data: NuFIT 6.0 (normal ordering, 3σ) and JUNO 2025, $\sin^2\theta_{12} = 0.3092 \pm 0.0087$, used as a 2σ range; $m_1 \in \{0, 0.001, 0.003\}$ eV.

v4.9.0-r57 results

Theoremlet v4.9.0-r57.1 (numerical).

(a) Variant A: 776 of 3 965 760 cases have θ_{12} in the JUNO 2σ range and θ_{13}, θ_{23} in the NuFIT 3σ ranges, for ε_e from 0.03 to 0.64.

(b) Variant B: 3 808 of 20 998 368 cases, for ε near 0.04–0.06 and near 0.62–0.64.

(c) Given θ_{12} and θ_{13} in their ranges, $\sin^2\theta_{23}$ spans $[0.309, 0.537]$ (A) and $[0.314, 0.515]$ (B).

Proof sketch. Script `llm/k97_test_h41.py`, output `llm/k97_out.txt`. □

Verdict and reading. H41a and H41b are confirmed, H41c (width of $\sin^2\theta_{23}$ below 0.05) is refuted.

1. The tension of the TM2 reading with JUNO can be resolved: a small correction by the mask, of the shape fixed in v4.9.0-r47, lowers $\sin^2\theta_{12}$ from 0.341 into the JUNO range while θ_{13} and θ_{23} stay within the measured ranges. The readout of v4.9.0-r55 is therefore not refuted.

2. It is a fit with two settings, not a prediction: θ_{23} is not fixed.

3. An observation after the evaluation, not part of the criterion: in both variants $\sin^2\theta_{23} \leq 0.54$ once θ_{12} and θ_{13} fit. A value clearly in the upper octant would speak against the reading. This bound was noticed afterwards and would have to be pre-registered to count.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r57 status in the three frames

result	M / T / P
r57.1(a)–(c)	statements on lepton mass matrices in M, the same in T as in v4.9.0-r52 and r55; P not evaluated

v4.9.0-r57 register entries

statement	status after v4.9.0-r57
TM2 reading of v4.9.0-r55 against JUNO	tension resolvable by the mask as a correction (H41a, b)
θ_{23} from the corrected reading	not fixed (H41c); one-sided bound $\sin^2\theta_{23} \lesssim 0.54$ noticed afterwards

v4.9.0-r57 ledger and guard update

$$R_{\mathrm{tm}}^{v490\text{-}r57} = (0, 1, 0)$$

for (no sharp prediction; tension with JUNO resolvable; bound on θ_{23} to be pre-registered).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r57,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r57} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r58 Source-Only: The CP Phase of the Corrected Trimaximal Reading

Purpose. Layer v4.9.0-r56 named the CP phases as sharper data, and layer v4.9.0-r57 left θ_{23} unfixed. This layer records the pre-registered test H43 (note `llm/Vorab_Notiz_H43_CP_Phase_aus_TM2.md`, commit `59d8d04`, fixed before computing) of whether the corrected trimaximal reading fixes the Dirac phase δ of the PMNS matrix. Earlier layers are not rewritten.
Source-only status. This v4.9.0-r58 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r58 results

Models, grids and assignments as in v4.9.0-r57; δ from the Jarlskog invariant and $|U_{\mu 1}|^2$ for every case that fits θ_{12} (JUNO 2σ), θ_{13} and θ_{23} (NuFIT 3σ). Pre-registered: a prediction if $|\sin\delta|$ or $\cos\delta$ lies in a range narrower than 0.3.

Theoremlet v4.9.0-r58.1 (numerical).

(a) Variant A (776 cases): $|\sin\delta| \in [0.174, 0.999]$, $\cos\delta \in [-0.980, 0.985]$.

(b) Variant B (3 808 cases): $|\sin\delta| \in [0.172, 0.999]$, $\cos\delta \in [-0.981, 0.985]$.

Proof sketch. Script `llm/k98_test_h43.py`, output `llm/k98_out.txt`; it reproduces the case counts of v4.9.0-r57. □

Verdict and reading. H43a and H43b are refuted; no prediction for δ is registered.

1. The reading does not fix δ; only phases very close to 0° or 180° ($|\sin\delta| < 0.17$) do not occur, a weak restriction that may depend on the grid.
2. The corrected trimaximal reading remains a fit with two settings: θ_{12} fits, θ_{23} is bounded from above (pending prediction H42), δ is free. A sharper reading would need the correction itself fixed by the structure.

v4.9.0-r58 status in the three frames

result	M / T / P
r58.1(a), (b)	statements on the lepton mixing matrix in M; under the time reversal T the phase changes sign, $\delta \mapsto -\delta$, as for any complex conjugation; P not evaluated

v4.9.0-r58 register entries

statement	status after v4.9.0-r58
Dirac phase δ from the corrected trimaximal reading	not fixed (H43); only $\|\sin\delta\| < 0.17$ absent

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r58 ledger and guard update

$$R_{\mathrm{cp}}^{v490\text{-}r58} = (0, 1, 0)$$

for (no prediction for δ; phase computed for all fitting cases; correction still free).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r58,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r58} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r59 Source-Only: The Correction Strength from a Structural Value

Purpose. Layers v4.9.0-r57 and r58 left the corrected trimaximal reading as a fit with two settings. This layer records the pre-registered test H44 (note `llm/Vorab_Notiz_H44_eps_aus_Strukturwert.md`, commit `95566e5`, fixed before computing) of whether a structural value fixes the correction strength ε. To avoid choosing values after the hits of r57 were known, only the candidate list of v4.9.0-r56, fixed before r57, was used. Earlier layers are not rewritten.
Source-only status. This v4.9.0-r59 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r59 results

Theoremlet v4.9.0-r59.1 (numerical).

(a) For none of the fourteen candidates (2/9, β_{sym}^2, 1/3, $\pi/9$, 1/2, $\pi/6$, β_{sym}, $1/\sqrt{3}$, 2/3, $1/\sqrt{2}$, 3/4, 1, $\sqrt{3}$, γ_L) does variant A or B of v4.9.0-r57 give a case with θ_{12} in the JUNO 2σ range and θ_{13}, θ_{23} in the NuFIT 3σ ranges.

(b) A check after the evaluation: the charged-lepton branches exist at all candidates up to $\varepsilon = 1$; variant A has hits only in two windows, $\varepsilon_e = 0.03$–0.06 and $\varepsilon_e = 0.62$–0.64.

Proof sketch. Script `llm/k99_test_h44.py`, output `llm/k99_out.txt`; the check in `llm/k99_nachkontrolle.py`. □

Verdict and reading. H44a is refuted.

1. No structural value from the pre-registered list fixes the correction; the reading does not become predictive this way.
2. The data nearly fix ε under the reading (≈ 0.05 or ≈ 0.63, each within about ± 0.02); this is a property of the data, not a structural value, and searching for a number in these windows now would be numerology.
3. *Correction to v4.9.0-r57.1(a):* the statement "for ε_e from 0.03 to 0.64" gave only the minimum and maximum; the hits lie in the two windows of (b), with none in between.

v4.9.0-r59 status in the three frames

result	M / T / P
r59.1(a), (b)	statements on lepton mass matrices in M, the same in T as in v4.9.0-r57; P not evaluated

v4.9.0-r59 register entries

statement	status after v4.9.0-r59
ε from a structural value	refuted for the pre-registered list (H44)
hit windows of v4.9.0-r57	$\varepsilon_e \approx 0.03$–$0.06$ and 0.62–0.64; r57.1(a) corrected

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r59 ledger and guard update

$$R_{\mathrm{es}}^{v490\text{-}r59} = (0, 1, 0)$$

for (no structural value for ε; hit windows located; origin of the correction open).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r59,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r59} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r60 Source-Only: The Correction Strength as the Rapidity of the Matter Frame

Purpose. Layer v4.9.0-r59 found no structural value for the correction strength ε in the pre-registered list. The author asked how the theory could supply ε. This layer records the test H45 (note `llm/Vorab_Notiz_H45_eps_als_Frame_Rapiditaet.md`, commit `bfde03d`, fixed before computing at the values below). The test is not blind: the hit windows of r59 were known. Earlier layers are not rewritten.
Source-only status. This v4.9.0-r60 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r60 idea

The mask enters as $D = \exp(\varepsilon W)$ with real diagonal W: a hyperbolic, boost-like map of the generation slots, whose natural parameter is a rapidity. The theory has one such quantity, the rapidity η of the matter frame, $\beta_{\mathrm{sym}} = \tanh\eta$, $\gamma_L = \cosh 2\eta$ (Convention F). Reading: $\varepsilon = \eta$. Values: $\operatorname{artanh}(0.5493) = 0.6174$ (measured β_{sym}) and $\operatorname{artanh}(1/\sqrt{3}) = 0.6585$ (Hopf value).

v4.9.0-r60 results

Theoremlet v4.9.0-r60.1 (numerical).

(a) At $\varepsilon = 0.6174$ variant A of v4.9.0-r57 has 120 and variant B 1 104 cases with θ_{12} in the JUNO 2σ range and θ_{13}, θ_{23} in the NuFIT 3σ ranges.

(b) In both variants these cases have $\sin^2\theta_{23} \in [0.4913, 0.4975]$ and $|\sin\delta| \in [0.17, 1.00]$.

(c) At the Hopf value $\varepsilon = 0.6585$ there are no such cases.

Proof sketch. Script `llm/k100_test_h45.py`, output `llm/k100_out.txt`. □

Verdict and reading. H45a is confirmed as compatibility only (not blind); H45b, which required both θ_{23} and δ to be sharp, is refuted, because δ stays free.

1. For the first time a reading takes ε from the structure and meets the data: the measured frame offset fits, the Hopf value does not.

2. New and unknown before: with $\varepsilon = \operatorname{artanh}\beta_{\mathrm{sym}}$ the reading fixes $\sin^2\theta_{23} \approx 0.49$–$0.50$, near maximal mixing and just in the lower octant, in both variants. This is sharper than the pending prediction H42 (≤ 0.54). By the criterion of H45 it is not registered as a prediction; a prediction for θ_{23} alone would need its own note.

3. Limits: not blind; β_{sym} itself is an input (OP1 open); $\varepsilon = \eta$ is a reading, not derived from an action.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r60 status in the three frames

result	M / T / P
r60.1(a)–(c)	$\varepsilon = \eta$ is the matter-frame rapidity: stated in M; in T the rapidity changes to $\eta + i\pi/2$ (v4.8.0-r20), not evaluated; in P $\eta = 0$ and the correction vanishes

v4.9.0-r60 register entries

statement	status after v4.9.0-r60
$\varepsilon = \eta = \operatorname{artanh} \beta_{\mathrm{sym}}$	compatible with the lepton data (not blind); Hopf value not
$\sin^2 \theta_{23}$ under this reading	0.491–0.498; not yet registered as a prediction
δ under this reading	not fixed

v4.9.0-r60 ledger and guard update

$$R_{\mathrm{er}}^{v490\text{-}r60} = (1, 0, 0)$$

for (ε from the frame rapidity compatible; δ not fixed; θ_{23} prediction to be registered).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r60,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r60} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r61 Source-Only: A Frame Audit of the Open Problems OP1 to OP5

Purpose. The author suspected that the open problems, like the main statements before them, were studied in mixed frame configurations. This layer audits OP1 to OP5 against Convention F (v4.9.0-r22, r24), the class split of v4.9.0-r18/r19 (scale quantities by way C, $e^{k\eta}$; readout quantities by way A, γ_L) and rule 7 of the round format. It makes no prediction. One computation (`llm/k101_fenster_eta.py`) is an observation, not a test. Earlier layers are not rewritten.
Source-only status. This v4.9.0-r61 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r61 results

Theoremlet v4.9.0-r61.1 (OP1).

(a) The Tribonacci condition $\gamma_L = 1/\beta_{\text{sym}}$ of Theorem 41.18 equates a readout quantity of the matter frame with $1/\beta_{\text{sym}} = \beta_T$, the offset of the tachyon in its own measure (v4.9.0-r24.2); equivalently $\cosh 2\eta = e^{\eta}$, way A equal to way C. It is not a frame-consistent fixed point.

(b) The Hopf value of Theorem 22.2 reads $\beta_{\text{sym}} = \sin(\theta_H/2)$ for a pure state measured from the photon pole, i.e. in the reference frame R (v4.9.0-r24). At $\beta_{\text{sym}}^2 = \frac{1}{3}$ that pure state has populations $(\frac{2}{3}, \frac{1}{3})$, while the matter frame of Convention F has $(\frac{3}{4}, \frac{1}{4})$ and $|b| = \frac{1}{2}$ (v4.9.0-r17). The same β_{sym} labels two different states.

(c) The measured relation $\gamma_L = \gamma_{\text{BI}}/\gamma_{\text{string}}$ of Theorem 20.6 is the Lorentz factor between the real projections of M and T and the ratio of Bloch lengths $|b_R|/|b_M|$; it is frame-consistent, but an input.

(d) The Hopf gap and the Tribonacci gap, the subject of OP4, compare the measured matter value with values from another frame or another class.

(e) Observation: under the reading of v4.9.0-r55, r57, r60 the lepton data fix $\eta \in [0.611, 0.647]$, i.e. $\beta_{\text{sym}} \in [0.545, 0.570]$ and $\gamma_L \in [1.844, 1.961]$, independently of $\gamma_{\text{BI}}/\gamma_{\text{string}}$. The measured value lies inside; Hopf (2) and Tribonacci (1.839) lie just outside. Across the window $\sin^2\theta_{23} \in [0.491, 0.498]$.

Theoremlet v4.9.0-r61.2 (OP2 to OP5).

(a) OP2 is closed except (T4), which is OP1 (Theorem 41.16); (T4) reads γ_L as a ratio of spectral unfolding scales and carries the same class mixing as r61.1(a).

(b) OP4 as posed rests on r61.1(d).

(c) The Theta sum of OP5 (Definition 42.32) converts zeros (frequencies) and cutoff (a length), scale quantities, with γ_L instead of $e^{k\eta}$. The difference is 0.5% ($e^{\eta} = 1.854$, $\gamma_L = 1.864$) and does not explain the failure of the conjecture by 1.003.

(d) OP3 was not re-examined.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof sketch. (a)–(d) of r61.1 and r61.2 by comparison with the rules of v4.9.0-r17 to r24; the populations by direct computation; (e) by script `llm/k101_fenster_eta.py` (output `llm/k101_out.txt`) on the models of v4.9.0-r57 with a grid of 0.001. □

Reading.

1. The suspicion holds for OP1: two of its three values come from configurations that mix frames or classes; only the measured value is frame-consistent. OP1 stays open, but its target is the matter value itself, not a correction of the Hopf or Tribonacci value.

2. OP4 is ill-posed in its present form; only OP1 remains. OP2 (T4) carries the same mixing. In OP5 the mixing is present but not the cause of the failure.

3. r61.1(e) is not blind (the reading $\varepsilon = \eta$ was chosen after the windows were known). It shows that lepton mixing, read this way, selects the measured frame offset over the Hopf and Tribonacci values.

v4.9.0-r61 status in the three frames

result	M / T / P
r61.1(a)	mixes M (γ_L) and T ($\beta_T = 1/\beta_{\text{sym}}$)
r61.1(b)	mixes R (pure Hopf state) and M (mixed matter state); P is the degenerate point $b = 0$
r61.1(c), (e)	M; the M–T factor is frame-consistent
r61.2	as stated per item

v4.9.0-r61 register entries

statement	status after v4.9.0-r61
Tribonacci fixed point (Theorem 41.18)	frame and class mixing; candidate for a status note
Hopf value of β_{sym} (Theorem 22.2)	reads an R-frame pure state as the M offset; candidate for a status note
OP4 (continuum-limit correction)	ill-posed in its present form
OP1	open; target is the matter value itself
β_{sym} from lepton mixing (reading r55–r60)	$[0.545, 0.570]$; measured inside, Hopf and Tribonacci outside (not blind)

v4.9.0-r61 ledger and guard update

$$R_{\text{op}}^{v490\text{-}r61} = (1, 1, 0)$$

for (frame mixing in OP1, OP2 (T4), OP4 located; OP5 mixing found but not causal; OP1 itself open).

The layer spends no guard:

$$R_{\text{glob}}^{(v490\text{-}r61,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\text{guard}}^{v490\text{-}r61} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r62 Source-Only: Status Notes after the Frame Audit of the Open Problems

Purpose. Layer v4.9.0-r61 named Theorem 41.18, Theorem 22.2 and the OP 4 line as candidates for status notes. On 2026-10-24 the author decided to add them. Following rule C of v4.9.0-r25, the text stays and a status note records the state; nothing is archived. The notes are applied by the patch specification `llm/revision_spec_v490_r61.py` (also for the GitHub fragments).
Source-only status. This v4.9.0-r62 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r62 notes added

place	note
Theorem 22.2	reads a pure R-frame state as the matter offset (r61.1(b))
Remark 22.3	OP 1 open, target the matter value; the Hopf gap is not like-for-like; window from lepton mixing (r61.1(e), not blind)
Theorem 41.18	existing note extended: frame and class mixing (r61.1(a))
Remark 41.20	claimed resolution withdrawn; OP 4 ill-posed in its present form (r61.1(d))

v4.9.0-r62 status in the three frames

The notes restate v4.9.0-r61; the frame status is that of r61.

v4.9.0-r62 ledger and guard update

$$R_{\mathrm{sn}}^{v490\text{-}r62} = (1, 0, 0)$$

for (status notes applied; no new statement; OP 1 open).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r62,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r62} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r63 Source-Only: The Frame Offset as a Ratio of Two Horizon Countings

Purpose. Asked what β_{sym} measures, the author agreed to test a reading already offered in v4.8.0-r03 (F11): $\gamma_{\mathrm{string}} = \ln 2/(\pi\sqrt{3})$ is the value of the Barbero–Immirzi parameter from a horizon counting with spin $j = \frac{1}{2}$ only (Ashtekar, Baez, Corichi, Krasnov), γ_{BI} the value from the corrected counting with all spins (Domagała–Lewandowski, Meissner). r03 saw a conflict with the Lorentz reading. This layer records the pre-registered check H47 (note `llm/Vorab_Notiz_H47_Zaehlregel_beta_sym.md`, commit `df0c788`), and, following the author's suggestions that the $\frac{1}{2}$ comes from the uncertainty relation and that tachyons relax to a floor, two exact lemmas. Earlier layers are not rewritten.
Source-only status. This v4.9.0-r63 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r63 results

Theoremlet v4.9.0-r63.1 (H47). With the area spectrum $a(j) = \sqrt{j(j+1)}$, let γ solve $\sum_j g(j)\, e^{-2\pi\gamma a(j)} = 1$. Spin $\frac{1}{2}$ only, $g = 2$, gives $\gamma_{\min} = \ln 2/(\pi\sqrt{3})$.

full counting	γ	$\gamma/\gamma_{\min}$	β	in $[0.545, 0.570]$
$g = 2$ (Domagała–Lewandowski, Meissner)	0.2375330	1.864700	0.549406	yes
$g = 2j + 1$ (Ghosh–Mitra)	0.2740669	2.151501	0.604468	no

The main body uses the rounded input 0.2375 ($\gamma_L = 1.86444$, $\beta_{\mathrm{sym}} = 0.549348$). The window is the one of v4.9.0-r61.1(e).

Theoremlet v4.9.0-r63.2 (uncertainty saturation). In $|j, m\rangle$ ($\hbar = 1$) one has $\Delta J_x\, \Delta J_y = \frac{1}{2}\big(j(j+1) - m^2\big)$ and the Robertson bound $\Delta J_x\, \Delta J_y \geq \frac{1}{2}|m|$. Equality holds exactly for $m = \pm j$, two states at every j; at $j = \frac{1}{2}$ every state saturates and $\Delta J_x = \frac{1}{2}$.

Theoremlet v4.9.0-r63.3 (tachyon floor). A tachyon of invariant μ has $E = \mu/\sqrt{v^2 - 1}$, strictly decreasing for $v > 1$ with $E \to 0$ as $v \to \infty$; the momentum tends to μ, the invariant does not change. The floor $E = 0$, $p = \mu$ seen from a frame of velocity u has $E' = -\gamma u \mu < 0$, $p' = \gamma\mu$, velocity $-1/u$, and $E'^2 - p'^2 = -\mu^2$. In rapidities the floor of a frame at η_X is $\eta_X + i\pi/2$, with velocity $\coth(\eta_X - \eta_Y)$ in a frame at η_Y.

Theoremlet v4.9.0-r63.4 (where the floor sits). With M at $+\eta$ and R at 0:

floor of	velocity in R	velocity in M	cosh of the real distance to M
R	∞	$-1/\beta_{\mathrm{sym}}$	$\cosh\eta = 1.197$
M	$1/\beta_{\mathrm{sym}}$	∞	1
mirror frame $-\eta$	$-1/\beta_{\mathrm{sym}}$	$-\coth 2\eta$	$\cosh 2\eta = \gamma_L$

The tachyon of Convention F, rapidity $-\eta + i\pi/2$ (v4.9.0-r24.2), is the floor of the mirror frame.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof sketch. r63.1 by `llm/k102_zaehlregeln.py` (output `llm/k102_out.txt`). r63.2: $\langle J_x\rangle = \langle J_y\rangle = 0$, $\langle J_x^2\rangle = \langle J_y^2\rangle$, and $j(j{+}1){-}m^2{-}|m| = (j{-}|m|)(j{+}|m|{+}1)$. r63.3 by the boost formulas and $\tanh(x + i\pi/2) = \coth x$. r63.4 by r63.3. The arithmetic of r63.2 and the boost statements of r63.3 are in `formal/QSFormal/Zaehlung.lean`; the operator values for $j \le 6$, the energy monotonicity and the table of r63.4 are in `llm/k103_robertson_tachyon.py` (output `llm/k103_out.txt`). □

Reading.

1. Under this reading β_{sym} is not a measured quantity but an $SU(2)$ number fixed by two counting rules. The counting reading and the Lorentz reading of r03 agree when the tachyon frame counts minimally and the matter frame fully: $\gamma_L = \cosh 2\eta$ is the boost between these two opposite frames.

2. One principle gives both rules: a frame counts only states that saturate the uncertainty bound. The tachyon frame takes the absolute minimum ($j = \frac{1}{2}$, $\Delta J_x = \frac{1}{2}$), the matter frame the minimum at each spin ($m = \pm j$, weight 2). The Ghosh–Mitra counting also counts non-saturating states.

3. r63.3 gives a dynamical reason for the minimal counting: a tachyon that loses energy accelerates towards a single floor.

4. Only the floor of the mirror frame reproduces γ_L as the real M–T distance. A floor pinned to the matter frame gives distance 0, a floor in R gives $\cosh\eta$. Why the floor sits in the mirror frame is open; the symmetric placement of Theorem 20.6 is assumed, not derived.

5. r63.1 is not blind: the window was known, and the Ghosh–Mitra value had been estimated before the note. OP1 now reads: derive from the frames why T counts only $j = \frac{1}{2}$ and M all spins with weight 2.

v4.9.0-r63 status in the three frames

result	M / T / P
r63.1	M counts all spins (weight 2), T counts $j = \frac{1}{2}$; P carries no counting
r63.2	frame-free (representation theory)
r63.3, r63.4	kinematics; the T of Convention F is the floor of the mirror frame

v4.9.0-r63 register entries

statement	status after v4.9.0-r63
H47 (counting reading of β_{sym})	DL/Meissner inside the lepton window, Ghosh–Mitra outside; not blind
uncertainty saturation	exact; weight 2 = states $m = \pm j$
tachyon floor	exact kinematics; floor of the mirror frame reproduces γ_L
OP1	open; reformulated as a question about two counting rules

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r63 ledger and guard update

$$R_{\mathrm{cnt}}^{v490\text{-}r63} = (1, 1, 0)$$

for (counting reading consistent with the lepton window; both rules from uncertainty saturation; the assignment of the rules to the frames not derived).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r63,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r63} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r64 Source-Only: A Frame Audit of the Gravity Statements under the Counting Reading

Purpose. After v4.9.0-r63 the author asked what gravity is in this context. Under the counting reading the two values of γ come from the same horizon, with the same area A and the same ℓ_P: each counting fixes its own γ by $S = A/(4\ell_P^2)$. This layer audits the main-body statements that use γ_L, γ_{BI} or γ_{string} on the gravity side, as v4.9.0-r61 did for OP1 to OP5. It makes no prediction and rewrites no earlier layer.
Source-only status. This v4.9.0-r64 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r64 results

Theoremlet v4.9.0-r64.1 (where the frame difference sits).

(a) Theorem 25.3 lists γ_{BI} with $k = 0$, "frame-invariant by construction"; Theorem 20.5 calls γ_{BI} the Lorentz-transformed γ_{string}. The two statements contradict each other, independently of r63.

(b) Under the counting reading the whole frame difference sits in γ: $\hbar$, G_N and ℓ_P enter both countings identically. The assignments $k(\hbar) = 1$, $k(G_N) = 2$ converted with γ_L use the counting ratio a second time. This concerns Theorem 25.5 ($\gamma_L^{-3/2}$), the corollary "Cosmological constant revisited" (γ_L^5) and item (i) of Proposition 21.4 (γ_L^{-1} for T_H). They are not refuted, but they need a source for their k values other than γ_L.

(c) Theorem 20.5 assigns γ_{string} to the photon frame and reads γ_L as an M–R Lorentz factor at $\beta = 0.844$. The number is $\tanh 2\eta$, the M–T velocity; under r63 γ_{string} belongs to the tachyon floor. The value is consistent, the frame label is R for T.

(d) Theorem 20.6 is frame-consistent and gains a kinematic reading from r63.4: the tachyon at $-\beta_{\mathrm{sym}}$ is the floor of the mirror frame.

(e) Theorem 21.3 gives the escaping photon in R the value γ_{string}. Under r63 that is the counting of the infalling tachyon; the statement reads T as R.

(f) Theorem 21.5 ($\gamma_L \approx 1/\beta_{\mathrm{sym}}$, gap 2.4%) compares $\cosh 2\eta$ with $\coth \eta$, the class mixing of r61.1(a). Under r63 γ_L is a counting ratio; the gap is not geometric.

(g) Theorem 22.16 uses $\beta_{\mathrm{sym}}^2 = \tanh^2 \eta$, the M–R relation, which is consistent within the symmetric placement; its value $\beta_{\mathrm{sym}} = 1/\sqrt{3}$ is the Hopf value of r61.1(b). The drift problem of v4.9.0-r17 is untouched.

(h) Theorem 22.19 uses none of the three quantities.

Proof sketch. (a) by reading the two statements; (b) from the counting equations of r63.1, in which ℓ_P and A are common to both countings; (c)–(g) by comparison with r63.4, v4.9.0-r18/r19 and r61; (h) by inspection. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Reading.

1. The gravity statements split into two groups. Those that put the frame difference into γ (Theorems 20.5 after relabelling, 20.6) are compatible with the counting reading. Those that put it into $\hbar$ and G_N (Theorems 25.3, 25.5, the cosmological corollary, Proposition 21.4(i)) are not supported by it.

2. In the counting reading gravity itself is frame-independent: G_N, ℓ_P and the area are the same for all frames. What changes between the frames is how many states a frame counts per area, and γ is the exchange rate.

3. This is a reading, not a theorem: the k-value picture would survive with an independent derivation of its exponents. Candidates for status notes are Theorems 25.3 (r64.1(a)), 25.5 and 21.3; the decision is left to the author.

4. H15 ($\cot\theta_W = \gamma_L$) is not affected.

v4.9.0-r64 status in the three frames

result	M / T / P
r64.1(a), (b)	frame difference in the counting (M, T), not in $\hbar$, G_N
r64.1(c), (e)	T read as R
r64.1(d)	M and T; frame-consistent
r64.1(f), (g)	as r61.1(a), (b)

v4.9.0-r64 register entries

statement	status after v4.9.0-r64
$k(\gamma_{\mathrm{BI}}) = 0$ (Theorem 25.3)	contradicts Theorem 20.5; candidate for a status note
photon-frame Planck length, γ_L^5	not supported by the counting reading; need independent k values
vacuum ratio, Hawking (i)	reads the tachyon counting as the photon value
Hawking temperature $\gamma_{\mathrm{string}} T_H$	frame-consistent; T is the floor of the mirror frame
symmetric matter–tachyon frame	

v4.9.0-r64 ledger and guard update

$$R_{\mathrm{grav}}^{v490\text{-}r64} = (1, 1, 0)$$

for (contradiction in the k table found; frame difference located in the counting under r63; no gravity statement derived).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r64,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r64} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r65 Source-Only: Status Notes after the Frame Audit of the Gravity Statements

Purpose. Layer v4.9.0-r64 named Theorems 25.3, 25.5 and 21.3 as candidates for status notes. On 2026-10-25 the author decided to add them. Following rule C of v4.9.0-r25, the text stays and a status note records the state; nothing is archived. The notes are applied by the patch specification `llm/revision_spec_v490_r64.py` (also for the GitHub fragments).
Source-only status. This v4.9.0-r65 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r65 notes added

place	note
Theorem 25.3	existing note extended: $k(\gamma_{\mathrm{BI}}) = 0$ contradicts Theorem 20.5; $k(\hbar)$, $k(G_N)$ need a source other than γ_L (r64.1(a), (b))
Theorem 25.5	existing note extended: $\gamma_L^{-3/2}$ not supported by the counting reading (r64.1(b))
Theorem 21.3	new note: the tachyon counting is assigned to the escaping photon, T read as R (r64.1(e))

v4.9.0-r65 status in the three frames

The notes restate v4.9.0-r64; the frame status is that of r64.

v4.9.0-r65 ledger and guard update

$$R_{\mathrm{sn}}^{v490\text{-}r65} = (1, 0, 0)$$

for (status notes applied; no new statement; the k values open).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r65,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r65} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r66 Source-Only: A Correction to r64 and the Local Frame Factor under the Counting Reading

Purpose. Rereading v4.9.0-r08 and r09 after v4.9.0-r65 showed that r64.1(a) repeats an earlier finding without citing it. This layer corrects that, extends the status note of r65 on Theorem 25.3 accordingly (rule C; specification `llm/revision_spec_v490_r66.py`), and states what the counting reading of v4.9.0-r63 implies for the drifting local frame factor of r08 and r09. It makes no prediction. Earlier layers are not rewritten.
Source-only status. This v4.9.0-r66 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r66 results

Correction v4.9.0-r66.1. The contradiction of r64.1(a), $k(\gamma_{\mathrm{BI}}) = 0$ in Theorem 25.3 against Theorem 20.5, is item 1 of the two contradictions of v4.9.0-r09 (§A.272). r09 resolved both by reading Theorem 20.5 as a statement about the structural factor γ_L alone, with the drifting local factor $\gamma_L^{(\mathrm{local})}$ of r08 and r18 kept separate. r64.1(a) is therefore not new, and its phrase "independently of r63" should have cited r09. The status note of r65 is extended with the reference. r64.1(b) to (h) are unaffected: they concern the structural factor.

Theoremlet v4.9.0-r66.2 (local factor).

(a) Under the counting reading the structural factor is the ratio of two counting solutions (r63.1), a fixed number. This agrees with r08, which already called it fixed because both inputs are fixed numbers.

(b) The two rules of r63 are discrete: spin $\frac{1}{2}$ only, or all spins with weight 2. They admit no continuous family, so the counting reading gives no mechanism for a continuously drifting local factor $\gamma_L^{(\mathrm{local})}(t)$. A continuous interpolation of the weights would be a new assumption.

(c) The drift readings tested in r08 and r09 are excluded by atomic clocks; r66.2(b) adds that the counting reading does not supply them either. The corollary "α-drift as a frame effect" of §25 still carries no status note; it is a candidate.

Proof sketch. r66.1 by reading r09; r66.2 from the counting equations of r63.1. □

v4.9.0-r66 status in the three frames

The correction concerns citation only. r66.2 concerns the structural factor between M and T; P carries no counting.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r66 register entries

statement	status after v4.9.0-r66
r64.1(a) local frame factor under the counting reading	not new; item 1 of v4.9.0-r09, resolved there by reading no continuous mechanism
corollary "α-drift as a frame effect"	candidate for a status note

v4.9.0-r66 ledger and guard update

$$R_{\mathrm{corr}}^{v490\text{-}r66} = (1, 1, 0)$$

for (citation corrected; no continuous local factor under the counting reading; the local factor itself still has no field law).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r66,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r66} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r67 Source-Only: Einstein's "Greatest Blunder" under the Counting Reading

Purpose. The author asked to look again at Remark 22.18 and its reinterpretation of the cosmological constant. This layer reads it together with Theorem 22.16 and Remark 22.17 under the counting reading of v4.9.0-r63 and the audit of v4.9.0-r64. It makes no prediction; the data comparison is after the fact. Earlier layers are not rewritten.
Source-only status. This v4.9.0-r67 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r67 results

Theoremlet v4.9.0-r67.1 (which side $\beta^2_{\rm sym}$ is on). Theorem 22.16 puts $\beta^2_{\rm sym}$ on matter: $\Omega_M = \beta^2_{\rm sym} \approx 0.30$, $\Omega_\Lambda = 1 - \beta^2_{\rm sym} \approx 0.70$. The field equation $G_{\mu\nu} + \beta^2_{\rm sym} g_{\mu\nu} = 8\pi G\, T_{\mu\nu}$ of Remark 22.17 and the statement $\Lambda_{\rm obs} = \beta^2_{\rm sym}$ of Remark 22.18 put the same $\beta^2_{\rm sym}$ on Λ. Both cannot hold: the remarks give Λ the matter share. This is separate from the dimensional problem of v4.8.0-r03 (F3, F4), which concerns the same expressions.

Theoremlet v4.9.0-r67.2 (counting form of the fractions). With the two countings of r63.1, $\gamma_T = \ln 2/(\pi\sqrt{3})$ (spin $\frac{1}{2}$ only) and $\gamma_M = 0.2375330$ (all spins, weight 2), and $\gamma_L = \gamma_M/\gamma_T$,

$$\Omega_M = \beta^2_{\rm sym} = \frac{\gamma_M - \gamma_T}{\gamma_M + \gamma_T} = 0.301847, \qquad \Omega_\Lambda = 1 - \beta^2_{\rm sym} = \frac{2\gamma_T}{\gamma_M + \gamma_T} = 0.698153.$$

The denominator $\gamma_M + \gamma_T = 0.364917$ is $|\gamma_3|$ of Theorem 20.4.

Theoremlet v4.9.0-r67.3 (epoch). $\beta_{\rm sym}$ is fixed under the counting reading (v4.9.0-r66.2), while Ω_M and Ω_Λ change with cosmic time. The identification of Theorem 22.16 can hold at one epoch only; at every epoch it would give $\Omega_\Lambda = 0.698$ also at recombination, which is excluded. This agrees with the present-epoch reading of v4.9.0-r15 and leaves the coincidence problem (why now) open.

Observation v4.9.0-r67.4 (after the fact). $\Omega_M = 0.3018$ against DESI DR2 BAO alone, 0.2975 ± 0.0086: $+0.5\sigma$; against Planck 2018 (TT,TE,EE+lowE+lensing), 0.3153 ± 0.0073: -1.8σ. Remark 22.17 quotes 0.3111, a Planck+BAO value. No test: the identification predates neither data set as a registered prediction.

Proof sketch. r67.1 by reading the three statements; r67.2 from $\beta^2_{\rm sym} = (\gamma_L - 1)/(\gamma_L + 1)$; r67.3 from r66.2 and the expansion history used in v4.9.0-r17; r67.4 from the quoted values. □

Reading.

1. Under v4.9.0-r64 the frames differ in their counting, not in G_N or the geometry. Λ as a curvature is then frame-independent like G_N: the cosmological constant is real, and the remark's claim that matter observers "necessarily measure" it because of their frame is not supported.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

2. What the framework could explain is the share Ω_Λ, not Λ. In counting form it is the share of horizon states on the minimal-spin floor, counted twice; the matter share is what matter counts beyond the floor. This reading is offered, not adopted.

3. The open step is the one named in v4.8.0-r03 (F11): why a share of a horizon counting should equal a share of the energy budget, and why only today.

4. Candidates for status notes: Remark 22.18 and the field equation in Remark 22.17 (r67.1). The decision is left to the author.

v4.9.0-r67 status in the three frames

result	M / T / P
r67.1	M; the two statements put the same M quantity on different sides
r67.2	M and T (the two countings); P carries no counting
r67.3, r67.4	M, present epoch

v4.9.0-r67 register entries

statement	status after v4.9.0-r67
$\Lambda_{\mathrm{obs}} = \beta^2_{\mathrm{sym}}$ (Remark 22.18), field equation of Remark 22.17	give Λ the matter share of Theorem 22.16; candidates for status notes
fractions in counting form	$\Omega_M = (\gamma_M - \gamma_T)/(\gamma_M + \gamma_T)$, $\Omega_\Lambda = 2\gamma_T/(\gamma_M + \gamma_T)$; offered
epoch	present epoch only; coincidence problem open

v4.9.0-r67 ledger and guard update

$$R_{\mathrm{cc}}^{v490\text{-}r67} = (1, 1, 0)$$

for (side swap found; counting form of the fractions; the step from counting share to energy share not derived).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r67,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r67} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r68 Source-Only: An Independent Recheck of H47 and r63 to r67

Purpose. The author asked to recompute the results of v4.9.0-r63 to r67 and to check the conjectures behind them. The numbers were recomputed with newly written code (`llm/k104_nachrechnung.py`, `llm/k105_h46_exakt.py`) and the literature values were looked up. This layer records what holds and corrects two statements. It makes no prediction. Earlier layers are not rewritten.
Source-only status. This v4.9.0-r68 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r68 results

Theoremlet v4.9.0-r68.1 (confirmed).

(a) The three counting solutions of r63.1 (0.127384, 0.237533, 0.274067) agree to 10^{-12}; Meissner (`gr-qc/0407052`) gives 0.23753295796592, Ghosh and Mitra (`gr-qc/0411035`) $\gamma \approx 0.274$.

(b) Meissner counts sequences of labels m, with the area depending on $|m|$ only; the factor 2 counts $\pm m$. The counted states are the states $|j = |m|, m\rangle$, so the reading of r63.2 matches the counting.

(c) The saturation statement of r63.2 holds exactly for all states up to $j = 10$; the boost statements of r63.3 and r63.4 hold numerically; the tachyon of Convention F has rapidity $-\eta + i\pi/2$ relative to R (r24.2).

(d) The counting form of r67.2 and the side swap of r67.1 hold; the corollary "Cosmological constant revisited" also writes $\Lambda_{\text{obs}} \sim \beta_{\text{sym}}^2$. Planck 2018 gives $\Omega_m = 0.315 \pm 0.007$, DESI DR2 BAO 0.2975 ± 0.0086.

(e) H46 with the unrounded $\beta_{\text{sym}} = 0.549406$ instead of 0.5493: $\sin^2\theta_{23} \in [0.4908, 0.4975]$ instead of $[0.4913, 0.4975]$. The registered interval $[0.49, 0.50]$ still holds; a control run with 0.5493 reproduces `llm/k100_out.txt`.

Correction v4.9.0-r68.2. The statement of r67.2 that $\gamma_M + \gamma_T$ is $|\gamma_3|$ of Theorem 20.4 is not a finding: that theorem defines $\gamma_3 := -(\gamma_{\text{string}} + \gamma_{\text{LQG}})$. It holds by definition, up to the rounding of the input 0.2375.

Correction v4.9.0-r68.3 (which counting the literature prefers). The SU(2)-invariant treatment of the horizon (Engle, Noui, Perez, `arXiv:0905.3168`) is regarded as better founded than the U(1) treatment, which counts too many microstates (logarithmic correction $-\frac{1}{2}$ instead of $-\frac{3}{2}$); it gives $\gamma \approx 0.274$, the Ghosh–Mitra value. The lepton window of r61.1(e) selects the U(1) counting (0.2375). Reading 5 of r63, that the data select the corrected counting, is therefore not supported: they select the counting that the later literature considers the less well founded. The wording "corrected"/"superseded" of v4.8.0-r03 (F11) reflects the state of 2004.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Remarks v4.9.0-r68.4.

1. The ABCK value and the Meissner value are both U(1) countings, so the pair of r63 is consistent within one treatment. An SU(2) pair has the same floor ($2j+1=2$ at $j=\frac{1}{2}$) and gives $\beta_{\text{sym}} = 0.604$, outside the window.

2. The uncertainty reading of r63.2 is a reinterpretation of the U(1) counting, not part of its derivation; mathematically it is exact.

3. "Own frame" of the tachyon means $1/\beta_{\text{sym}}$ relative to R in r24.2 and the mirror frame with $E = 0$ in r63.4; the numbers agree, the words do not.

4. With the unrounded value $\eta = 0.6175$ lies near the lower edge 0.611 of the window of r61.1(e), so the number of H46 hits is sensitive ($120 \to 168$ in variant A).

Proof sketch. By the scripts named above (outputs `llm/k104_out.txt`, `llm/k105_out.txt`) and the cited papers; protocol `llm/Nachrechnung_r63_r67.md`. □

v4.9.0-r68 status in the three frames

Unchanged from r63 to r67; r68.3 concerns the choice of counting for M and T, not the frames.

v4.9.0-r68 register entries

statement	status after v4.9.0-r68
numbers of H47, r63 to r67	confirmed
H46	holds with the unrounded β_{sym}
$\gamma_M + \gamma_T = \lvert\gamma_3\rvert$ (r67.2)	by definition, not a finding
counting reading of β_{sym}	requires the U(1) treatment; the SU(2) treatment preferred in the literature gives 0.604, outside the window

v4.9.0-r68 ledger and guard update

$$R_{\text{chk}}^{v490\text{-}r68} = (1, 1, 0)$$

for (numbers confirmed; two statements corrected; the counting reading depends on a disputed treatment).

The layer spends no guard:

$$R_{\text{glob}}^{(v490\text{-}r68,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\text{guard}}^{v490\text{-}r68} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r69 Source-Only: The OP5 Research Series of 25 to 27 October

Purpose. Between 25 and 27 October 2026 the author worked on OP5 with another language model. The results were committed to the repository folder `analysis/` (six commits, `3fa5ba9` to `337d9cb`, 173 files: notes, check scripts, their outputs and a literature catalogue). The author asked for this work to be integrated. This layer records what the series establishes, what it leaves open, and three errors it found in the main body. The notes themselves stay in `analysis/`; the layer names the note for each item. It makes no prediction, writes no status note and rewrites no earlier layer.
Source-only status. This v4.9.0-r69 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

How the series was checked. All 23 new check scripts were rerun with SymPy and NumPy; all finish without error, and the regenerated outputs differ from the committed ones only in the last floating-point digit. Nine statements were recomputed with newly written code (`llm/k106_nachrechnung_op5.py`, output `llm/k106_out.txt`); all nine hold. The notes marked "Independent" in the series were written by sub-agents of the same model; there is no external review.

v4.9.0-r69 results

Theoremlet v4.9.0-r69.1 (frames, neutrality and the absolute term).

(a) *A frame change cannot remove a balance defect.* Under a passive change of frame with frequency factor $a > 0$ and declared amplitude factor $b \neq 0$, the defect $\Delta = I - B$ of a balance of the type of Definition 42.20 becomes $(b/a)\Delta$; the statement $\Delta = 0$ is invariant. Replacing γ_L by e^{η} in frequency, cutoff or both leaves the historical defect between -1.0008 and -1.0038 (`OP5_Frame_Recheck.md`).

(b) *Three different neutral objects.* (i) The midpoint $\eta_N = (\eta_1 + \eta_2)/2$ of two scale frames with offsets $\pm\delta$; no input frame need be photon-like. (ii) On the real state disk the equal mixture of ρ and its native counterpart $\sigma_y \rho \sigma_y$ (the tachyon map $z \mapsto iz$ of v4.9.0-r28) is $I/2$; with $J_B = \sigma_x$ the component b_x survives. $I/2$ is a mixture, not a frame change, and carries the label P, while the old "photon frame" is the pure reference state R (v4.9.0-r24). (iii) Two positive non-parallel null streams select a unique flux-free timelike observer; there the stress is $\mathrm{diag}(\rho, \rho, 0, 0)$, not the vacuum form $\mathrm{diag}(\rho, -\rho, -\rho, -\rho)$. The three must not be identified (`OP5_Lambda_Neutral_Reassessment.md`, `OP5_Neutral_Frame_Candidates.md`).

(c) β^2_{sym} *as a relative separation (offered, not adopted).* With $\chi = \eta_1 - \eta_2$, $q_{\mathrm{rel}} = \tanh^2(\chi/2)$ is invariant under a common shift and exchange; at $\pm\eta$ it equals $\tanh^2\eta = (\cosh 2\eta - 1)/(\cosh 2\eta + 1)$, the form of β^2_{sym} when $\gamma_L = \cosh 2\eta$. A potential $V = M^4 \tanh^2(\chi/2)$ has $V'/M^4 \approx 0.384$ at the historical separation, so it gives no static vacuum there.

(d) *The absolute vacuum term is not selected.* A shift $H \mapsto H + C\,I$ of the whole generator changes no probability, gap or beat period in fixed-background unitary dynamics; stationarity at fixed metric is blind to a constant volume density; in the alternative affine parent action of `HC_Two_Frame_Action_Bridge.md` one local branch

© 2026 Marc Brendecke. Preprint — not peer reviewed.

solves all field equations for every constant C (branch-local screening, not selection); one neutral pair configuration is compatible with different constant vacuum energies (`OP5_Absolute_Selection_Independent.md`, `OP5_Affine_Vacuum_Response_Independent.md`, `OP5_Neutral_Pair_Dynamics_Result.md`).

(e) *Readout protocol and energy.* A passive change of representation changes no prediction; a different active readout (coupling, order, time, retained branch) can change states and exchanged energy. Energy then moves between system, apparatus and bath; none is produced by naming a frame (`OP5_Readout_Protocol_Energy_Result.md`).

(f) *Doublet selection.* No nontrivial orthogonal projector on the doublet of v4.9.0-r05 respects both the structural involution and the corrected M/T exchange; only 0 and I do. The two ports survive as an exchanged pair (`OP5_Symmetry_Sector_Selection_Independent.md`).

Theoremlet v4.9.0-r69.2 (cones, coframes and the cross readout).

(a) *The probe cone comes from the Clifford structure.* If the coefficients of the probe equation of §27 are local multiplications, the principal symbol is $\Gamma^\mu \xi_\mu$ with square $g^{\mu\nu}\xi_\mu\xi_\nu\, I$; mass, M_{16}, triadic and HC terms do not change the cone. A conformal rescaling $E \mapsto aE$ keeps every cone and changes the volume weight by a^4, so cone data fix neither an absolute scale nor the weight of a vacuum term (`OP5_Null_Transport_And_Signal_Locality.md`).

(b) *Two complex probes.* Two free massless spinor fields give the rank-four coframe $E = \mathrm{diag}(3,-1,-1,-1)$, $g = \mathrm{diag}(-9,1,1,1)$; the readout cone differs from the reference cone, and the induced equation leaves a residual $8i\, e_1$. For a constant frame and equal induced massless (or equal-mass) equations the trace $e_a{}^\mu E^a{}_\mu$ would have to be 0 and is 4 (`OP5_Two_Complex_Probes.md`). An exact coupled exchange solution with $M = 8\sigma_y$ matches only at one point: on the time axis the induced residual is $544\, i\, t^2 u + O(t^4)$ and the readout speed $1 - 80t^2 + O(t^4)$ (`OP5_Two_Probe_Exchange_Evolution.md`).

(c) *Two flat metrics.* Between the reference metric and the readout metric the relative connection is nonzero, both curvatures vanish (`OP5_Relative_Connection_And_Free_Paths.md`).

(d) *Connected cross readout.* If $E^a = \frac{1}{2} D_\omega j_S^a$ with a parallel selector, then $T^a = \frac{1}{2} R^a{}_b\, j_S^b$ exactly. Torsion-free connection plus an ordinary Einstein vacuum equation $\mathrm{Ric} = \Lambda_{\mathrm{geom}} g$ forces $\Lambda_{\mathrm{geom}} j = 0$, so $\Lambda_{\mathrm{geom}} = 0$ wherever $j \neq 0$. This is an exclusion under these premises, not a cancellation of vacuum energy; with $a = \tau$ the comoving fluid needs $\rho + 3p = 2\Lambda_{\mathrm{geom}}/\kappa$. A torsionful witness with E Minkowski exists (`OP5_Cross_Connection_Integrability.md`, `OP5_Torsionful_Geometric_Witness.md`).

(e) *Ward exclusion.* For the common-boost triad current and an isotropic time-only boost connection $h(t)$, current conservation in the reconstructed geometry requires $b + 3tb' = 1$ and $(hb^2)' = 0$; the second gives $b = (1 \pm \sqrt{10})/9$, the first $b = 1$. No differentiable profile exists (`OP5_Isotropic_Connection_Ward_NoGo.md`).

(f) *Counterflow and chiral tangent.* A counterflow configuration that meets every current projection still leaves full spinor residuals with norms between 3.9 and 9.5 at two sample points. Adding independent boosts and one chiral tangent $e^{i\chi\gamma_5/2}$ with signs $(+,+,-)$ fits the instantaneous derivative at both points (rank 22 of 24, residual below $4 \cdot 10^{-28}$);

© 2026 Marc Brendecke. Preprint — not peer reviewed.

this is a two-point fit, not a solution. The prescribed state set is not invariant under the comparison dynamics; a rotated probe pair cancels the axial drift but not the radial drift, and $\overline{E^T\eta E} \neq \bar{E}^T\eta\bar{E}$ in general (`OP5_Counterflow_Full_Residual_Result.md`, `OP5_Chiral_Tangent_Result.md`, `OP5_Constraint_Set_And_Ensemble_Result.md`).

Finding v4.9.0-r69.3 (errors in the main body).

(a) Remark 42.29: the expansion is $R(\tau) = -1/\tau + 1 - \frac{11}{24}\tau + \frac{1}{8}\tau^2 + O(\tau^3)$, not $-1/\tau + \frac{1}{2} - \tau/12$. Since $\int_\varepsilon^T (1+R)\,d\tau$ already contains $+\log\varepsilon$, the printed counterterm $+\log\varepsilon$ doubles the divergence; the counterterm is $-\log(\varepsilon/\mu)$ with a declared scale μ. With $\mu = 1$ and $T = 0.393$ the value is 1.689, not 0.940, so the 0.893 of Proposition 42.30 is not supported. The route was already marked false (v4.8.0-r05); the numbers are new.

(b) Theorem 26.4 and Proposition 26.5 still fix β_{sym} and the clock scale by γ_L^* of Theorem 24.2, which is archived. The citation no longer derives a clock scale.

(c) The corollary "Cosmological constant revisited" of §25 calls the photon-frame term zero and gives the finite ratio $\rho_{\mathrm{vac}}^{(\mathrm{photon})}/\rho_{\mathrm{vac}}^{(\mathrm{matter})} = \gamma_L^5$. For one and the same quantity both cannot hold; a distinction between net and bare density would have to be defined. This adds to the status-note candidates of v4.9.0-r64 and r67.

(a) to (c) are candidates for status notes (rules B and C of v4.9.0-r25); they are not written in this layer.

Remarks v4.9.0-r69.4.

1. The series corrects itself once: `OP5_Comparison_Map_And_Frame_Type.md` called the realification of J_B missing; it is Theoremlet v4.9.0-r04.1 (`OP5_Realification_And_Gram_Boundary.md`).

2. Every positive result of the series is conditional: a declared comparison model, a chosen witness, or two sample points. None is derived from the native M_{16}, P_{C_7} or a common action. Every note states that OP5 stays open.

3. The literature routes (`OP5_Mathematical_Literature_Routes_2026-10-27.md`) suggest, in this order: polar spinor variables (Fabbri, `arXiv:1707.03270`) and Fierz reconstruction (Inglis–Jarvis, `arXiv:1405.6762`); the constraint algorithm for k-presymplectic systems (Gracia, Martín, Román-Roy, `arXiv:0903.1791`); formal integrability (Landsberg, `dg-ga/9708005`); then a coupled coframe dynamics. Green and Wald (`arXiv:1011.4920`) and Kaloper et al. (`arXiv:1505.01492`) bound what averaging and sequestering can do for Λ.

4. The clock window $E_{\mathrm{clk}}^{\mathrm{win}}$ of the C5 chain is unchanged by a joint relabelling $\tau \mapsto a\tau$; it selects a relay window, not a clock period. This does not touch $W_{\mathrm{clk}}^{\mathrm{loc}}$ under $\Delta_\gamma < 1/5$.

Proof sketch. r69.1 and r69.2 by the notes named in each item and their check scripts in `analysis/`. Recomputed with new code (`llm/k106_nachrechnung_op5.py`): r69.1(b)(ii) and (c) including V', r69.2(b), (d) and (e), and the series and value of r69.3(a). r69.3 also by reading the cited passages. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r69 status in the three frames

- r69.1(a) holds in every frame: the defect transforms by b/a between M, T and R alike. It concerns finite invertible frame changes; the photon P is a degenerate limit and needs its own normalized observable.
- r69.1(b): (i) needs two real scale frames, so it compares M with R or with $T|_R$, not with the native T (complex rapidity, v4.9.0-r24); (ii) is built from M and its T-image and gives the P state; (iii) is a spacetime observer and has no frame label. Frame-free only in the sense that (ii) is symmetric in M and T.
- r69.1(d), (e), (f) and r69.2 are statements in the reference description R or in a declared classical comparison sector; they do not separate M from T.
- r69.3 concerns R and P: the old "photon frame" of the cited passages is R (v4.9.0-r24).

No statement of this layer concerns colour or charge. The chiral phase $e^{i\chi\gamma_5/2}$ acts in the Dirac comparison sector; it is neither the inner unit $J = L_{e_7}$ nor the $\mathbb{C}$-factor of the Dixon register.

v4.9.0-r69 register entries

statement	status after v4.9.0-r69
OP5 research series (`analysis/`, 25–27 Oct.)	reproducible; conditional results; $\mathrm{OP5}_{\mathrm{exact}}$ open
frame change as a repair of the Θ balance	excluded for finite invertible changes
neutral reference	three distinct objects; none fixes the absolute vacuum term
Remark 42.29	expansion and counterterm wrong; status-note candidate
γ_L^* clock scale in Thm. 26.4, Prop. 26.5	rests on an archived theorem; status-note candidate
corollary "Cosmological constant revisited"	zero and γ_L^5 for one quantity; status-note candidate

v4.9.0-r69 ledger and guard update

$$R_{\mathrm{chk}}^{v490\text{-}r69} = (1, 1, 0)$$

for (series reproduced and nine statements confirmed; three errors in the main body found; OP5 not closed).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r69,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r69} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r70 Source-Only: The 4.10 Consolidation in the Monolith, a Correction to r69, and Two Tests of a Common Dynamics

Purpose. On 29 and 30 October 2026 the author continued OP5 with another language model. Besides new working notes in `analysis/`, that work changed the research candidate on GitHub directly (commits `f98b29b` to `cf48496`): it corrected the three main-body errors found in v4.9.0-r69, added further status notes and a corrected Hopf annotation, and added the section "States, Mediators and Clock Comparisons" (§44) with a figure. This layer carries those changes into the monolith, records them, corrects one number of v4.9.0-r69, and records two checks of the common-dynamics sketch of 28 October: the pre-registered test H48 and an exploration of the idea that Λ is a mixture over sector crossings. Earlier layers are not rewritten.
Source-only status. This v4.9.0-r70 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r70 results

Record v4.9.0-r70.1 (the 4.10 consolidation). The following changes were made on GitHub without a layer and are applied here verbatim (patch specification `llm/revision_spec_v410_konsolidierung.py`, eight patches; local insertion `llm/einbau_v410_abschnitte_lokal.py`).

(a) Remark 42.29: series $R(\tau) = -1/\tau + 1 - \frac{11}{24}\tau + O(\tau^2)$, counterterm $-\log(\varepsilon/\mu)$, closed finite part, value 1.686953464 at $T = 0.393$, $\mu = 1$; the old target of Proposition 42.30 is withdrawn and Remark 42.31 is rewritten. This writes candidate (a) of v4.9.0-r69.3.

(b) Status notes at Theorem 26.4 and Proposition 26.5 (archived GUE clock scale) and at the corollary "Cosmological constant revisited" of §25. This writes candidates (b) and (c) of v4.9.0-r69.3.

(c) Further status notes at Theorem 16.2 (a shared clock does not by itself prove entanglement) and Corollary 42.24 (historical prediction register), and a dependency note after the Theta-sum conjecture.

(d) Theorem 22.1: the fibre S^7 was annotated as a group containing $G_2 \supset SU(3)$. S^7 is not that group; G_2 is the automorphism group of the octonions and $SU(3)$ the stabiliser of an imaginary unit. The annotation is removed and the reason given in a note.

(e) Section 44 and Subsection 44.6 with Figure 1. On GitHub they are separate parts of the candidate; in the monolith they stand before the archive appendix. The figure file `figures/op5-slices-9x9.pdf` is the first external graphic of the monolith.

The section records conditional results: typed readouts and the obligations of a mediator, an orientation-complete local reconstruction of the real doublet, calibrated clock comparisons, and finite field-compatibility tests up to fourth jet order at three points. None is a regional solution, a native common action or an OP5 closure; the section says so itself.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Correction v4.9.0-r70.2. The value 1.689 in v4.9.0-r69.3(a) is wrong; the finite part is 1.686953464. The quadrature in `llm/k106_nachrechnung_op5.py` integrated $1 + R + 1/\tau$ near $\tau = 0$, where two large terms cancel, and lost accuracy. The direct limit and the closed antiderivative agree to twelve digits. The finding of r69.3(a), that the printed expansion, counterterm and value 0.940 were wrong, stands.

Theoremlet v4.9.0-r70.3 (H48: β_{sym} as a frozen length). Pre-registered in `llm/Vorab_Notiz_H48_Einfrieren_beta.md` before computing. Model: the M/T doublet of a homogeneous spinor condensate is moved by the duality rotation, the compensator measurement and exchange with the tachyon reference state, with rates proportional to the density; flat unimodular FLRW with the Einstein–Cartan contact term.

(a) Refuted. Over 288 runs the frozen length $|b_\infty|$ ranges from 10^{-12} to 0.87 and depends on the initial expansion rate and density, not only on the rate ratio r. The dense-phase value $r/(1+r)$ is reached only if freezing happens while the expansion is still faster than the rotation (21 runs).

(b) The Einstein–Cartan contact term changes $|b_\infty|$ by at most 10^{-4}: in the dense phase the state lies on the R/T axis, and the axial density along the compensator (chirality) axis vanishes.

(c) Λ changes $|b_\infty|$ by up to 18 % when freezing falls into the Λ phase; it does not decouple from the frozen state.

Remark v4.9.0-r70.4 (Λ over sector crossings; exploration, not a test). The author asked whether each crossing of sectors carries its own Λ, so that ours is a mixture. A linear dependence $\mathrm{Tr}(\rho\hat{\Lambda})$ averages out under the rotation; a bilinear one depends on $|b|$ only and is constant once $|b|$ is frozen, so the epoch question of v4.9.0-r67 stays. Of 26 readings with structural weights (`llm/k108_kreuzungen.py`) one lands within 2σ of DESI DR2 BAO ($\Omega_M = 1/\gamma_L^2 = 0.288$, 3.9σ from Planck 2018), about as many as chance gives (1.3). No prediction is made. An identity remains: with $|b| = 1/\gamma_L$ the eigenvalues of the M/T state satisfy $\lambda_-/\lambda_+ = (\gamma_L - 1)/(\gamma_L + 1) = \beta_{\mathrm{sym}}^2$, so β_{sym}^2 is the weight ratio of the mixture. This is a rewording, not a new result.

Proof sketch. r70.1 by the patch specifications (reproducing the GitHub fragments exactly); r70.2 by direct evaluation of the finite part; r70.3 by `llm/k107_test_h48.py` (output `llm/k107_out.txt`, result `llm/Ergebnis_H48_Einfrieren_beta.md`); r70.4 by `llm/k108_kreuzungen.py`. □

v4.9.0-r70 status in the three frames

- r70.1 and r70.2 concern R and the main body; the old "photon frame" of the corrected passages is R (v4.9.0-r24).
- r70.3: the doublet carries M (state b) and T ($b \mapsto -b$); the frozen length $|b|$ is the same in M and T, so (a) is frame-free. The photon P is the attractor $b = 0$.
- r70.4: $\lambda_\pm$ are the weights of M and T; the identity is symmetric in M and T.

No statement of this layer concerns colour or charge.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r70 register entries

statement	status after v4.9.0-r70
status-note candidates of v4.9.0-r69.3	written (4.10 consolidation)
value in v4.9.0-r69.3(a)	corrected to 1.686953464
H48 (β_{sym} as frozen length)	refuted; depends on expansion history
Λ over sector crossings	no reading better than $\Omega_M = \beta_{\mathrm{sym}}^2$; epoch question open
monolith and GitHub candidate	main-body text identical again

v4.9.0-r70 ledger and guard update

$$R_{\mathrm{chk}}^{v490\text{-}r70} = (1, 1, 0)$$

for (consolidation carried over and r69 corrected; H48 refuted and the crossing reading without a hit beyond chance; OP5 not closed).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r70,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r70} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r71 Source-Only: The Structure of the OP5 Comparison Symbol, an Exact Polar Gauge Symmetry, and Two Tests

Purpose. On 2 November 2026 the author asked to continue OP5 where the second language model had stopped (commit `cf48496`): a 70-digit fifth-order germ of the fixed comparison model of §44 at $(t, x, y, z) = (1, \frac{3}{5}, 0, 0)$, the third-jet condition Ω, and an unresolved Spencer count at degree 5 that gave 1, 0 or -1 depending on the cutoff. This layer records an independent check of that state, the resolution of the count, the structure of the Ω-augmented symbol, an exact gauge symmetry of the comparison model, and two pre-registered tests (H49, H50). Everything concerns the comparison model only. Earlier layers are not rewritten.
Source-only status. This v4.9.0-r71 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r71 results

Notation: $R = (F, G, dg)$ is the completed first-order system, $J(\xi)$ its principal symbol (32×24), θ the stored symbol of Ω, g_p the degree-p symbol of the Ω-augmented system, and $c = \rho\partial_z - z\partial_\rho$, which is $-\partial_\vartheta$ in spherical coordinates.

Record v4.9.0-r71.1 (independent check of the germ). Two own implementations, sharing no code with the second model, confirm the germ.

(a) Exact power series along seven lines (80 digits) give F through order 4 at most 1.2×10^{-60} and g through order 5 at most 1.6×10^{-63}. Shifting one degree-5 or degree-3 coefficient is detected at the expected order.

(b) A truncated multivariate Taylor evaluator of the whole system, including G, gives all coefficients of R through degree 3 below 2.4×10^{-12} in double precision. The degree-4 prolonged symbol has rank 528 with a left kernel of dimension 112, as recorded by the second model.

Theoremlet v4.9.0-r71.2 (the negative count).

(a) The value -1 comes from counting a singular direction ($\sigma \approx 1.4 \times 10^{-10}$) of the incoming Spencer map that lies 98 % outside the kernel of the outgoing map. It violates $\delta^2 = 0$ and is not an image direction. Projecting the image onto that kernel first makes the count non-negative.

(b) Since $H^{6,1} = 0$ for prolongations and the outgoing rank 1024 is stable, $\dim H^{5,2} = \dim g_7 - 722$ exactly.

(c) A direct construction of g_p (gaps of at least 5×10^{-8} against at most 3×10^{-15}) gives $\dim g_p = 127, 208, 308, 427, 565, 726, 910, 1118, 1351$ for $p = 2, \ldots, 10$. Hence $\dim H^{5,2} = 4$ for the stored θ. The earlier recursive count was spoilt by an ill-conditioned basis of g_6.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Theoremlet v4.9.0-r71.3 (structure of the Ω-augmented symbol).

(a) With a random θ of the same shape the dimensions are $722, 898, 1093$ at $p = 7, 8, 9$, which is quadratic. Without Ω they are $757, 954$ at $p = 7, 8$, which is cubic. The stored θ gives cubic growth again from $p = 7$.

(b) The stored θ lies on a special locus: a random perturbation of size ε makes the four extra null values at $p = 7$ grow linearly in ε.

(c) At every ξ, $J(\xi)$ has a one-dimensional kernel $k(\xi)$, and $\theta(\xi) \cdot k(\xi) \approx 10^{-15}$ (random θ: 10^{-2}). So Ω does not see the free direction.

(d) The four degree-7 relations have the form $q(\partial)\, d_k(\partial)\, \Omega \in \langle J \rangle$.

- q is the light cone of the reconstructed coframe: in frame components it equals $\mathrm{diag}(1, -1, -1, -1)$ to 8×10^{-13}, and $J(\xi)$ loses five ranks on $q = 0$.
- The quadrics d_k generate the ideal of two real skew 2-planes Π_0 (signature $(0, 2)$) and Π_1 (signature $(1, 1)$) in covector space; the Hilbert function of the ideal is $12, 25, 44$. $J(\xi)$ loses one further rank on $\Pi_0 \cup \Pi_1$.

Thus θ is a rational combination of the rows of J with denominators on the cone and the two planes, and Ω carries new information only there.

Theoremlet v4.9.0-r71.4 (an exact polar gauge symmetry).

(a) The F-part of J annihilates the four directions $\delta\Psi = c^\mu D_\mu \Psi$. At every solution of $R = 0$, the free direction k is $\delta\Psi = f\, c^\mu D_\mu \Psi$ for the polar field c. This was checked at the germ and at four continued solutions off the equator, matching $-zx/\rho^2$ and $-zy/\rho^2$ to three or four digits.

(b) The comparison model is covariant under diffeomorphisms except through the background $h(t, r)$ and the connection. The connection is $A = M(t)\, d\varphi$, because $\rho\, G(t, \rho)$ does not depend on ρ. Since $c \lrcorner A = 0$ and $dA = M'(t)\, dt \wedge d\varphi$, one has $\mathcal{L}_{fc} A = 0$ and $\mathcal{L}_{fc} h = 0$ for every function f. So ϑ-reparametrisations are an exact gauge symmetry: the background knows t, r and φ, not ϑ.

(c) The directions that are invisible on Π_k are the directions in (t, r) that preserve h, with a small φ-tilt for Π_1. For example $(0.9714, 0.2375, 0, 0)$ at the germ, computed and measured alike. They are only partially free, because $\mathcal{L}_{fc_0} A = f\, c_0^t M'(t)\, d\varphi \neq 0$.

Theoremlet v4.9.0-r71.5 (H49: an earlier condition off the equator). Pre-registered (`Vorab_Notiz_H49_Polarwinkel_Bedingung.md`) before computing. Refuted. The compatibility defect of the second jets is between 1.3 and 2.9×10^{-10} at four controls on $z = 0$ and at five points with $z \in [0.02, 0.2]$. Negative controls react with 9×10^{-2} and 8×10^{-5}. The premise of H49, that a term $\propto z/\rho^2$ breaks the ϑ-freedom, was wrong; it used a Cartesian coefficient instead of the Lie derivative. r71.4(b) makes the outcome expected.

Theoremlet v4.9.0-r71.6 (H50: does Ω couple back to Ψ through ϑ?). Pre-registered (`Vorab_Notiz_H50_Omega_Psi_Kopplung.md`) before computing, after the author asked whether Ω and Ψ influence each other. Refuted (first class through order 3). For the shifts $f\, c^\mu \partial_\mu \Psi$ with all 35 monomials f of degree at most 3, the change of Ω is at most 1.5×10^{-10}, against 0.04 to 0.11 for shifts along $\partial_t, \partial_x, \partial_y$. The ϑ-shifts also leave the lower equations unchanged (at most 3.4×10^{-13}). Ω is gauge invariant.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof sketch. r71.1 by `k109_unabhaengig_F.py` and `k135_check.py`. r71.2 by `k110`, `k111`. r71.3 by `k112`–`k124`. r71.4(a), (c) by `k125`–`k132` and (b) by direct computation of the Lie derivative. r71.5 by `k133_test_h49.py` with controls `k134`. r71.6 by `k135_test_h50.py` with the series evaluator `k135_tps.py`. The notes `Erkundung_OP5_Symbolstruktur_Grad7.md`, `Ergebnis_H49_Polarwinkel_Bedingung.md` and `Ergebnis_H50_Omega_Psi_Kopplung.md` give the numbers. □

Consequence. The Ω-augmented comparison system is not involutive as it stands, because it carries a gauge freedom of one function of four variables. A Cartan–Kuranishi argument needs a gauge condition for ϑ first; only then does the question of further hidden conditions become well posed. The symmetry belongs to the chosen background. A background that depends on ϑ would break it. OP5 is not closed.

v4.9.0-r71 status in the three frames

- All statements concern the fixed OP5 comparison model of §44, stated in one coordinate chart with one coframe reconstructed from Ψ. No frame map M, T or P acts on them, so r71.1 to r71.6 are frame-free within the model. Whether the polar symmetry survives in a frame-resolved version of the model is open.
- The light cone q of r71.3(d) is the cone of the reconstructed geometry, not a statement about P.

No statement of this layer concerns colour or charge.

v4.9.0-r71 register entries

statement	status after v4.9.0-r71
70-digit germ of `cf48496`	independently confirmed, including G
degree-5 Spencer count $1/0/-1$	resolved: -1 is an artifact; $\dim H^{5,2} = 4$ for the stored θ
free function of the symbol	exact ϑ-gauge symmetry of the comparison model
Ω	gauge invariant; new information only on the cone and two planes
H49	refuted (premise corrected)
H50	refuted (first class through order 3)
next step	gauge condition for ϑ, then the involution test

v4.9.0-r71 ledger and guard update

$$R_{\mathrm{chk}}^{v490\text{-}r71} = (1, 1, 0)$$

for (germ confirmed and symbol structure established; H49 and H50 refuted as pre-registered; OP5 not closed).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r71,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r71} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r72 Source-Only: The ϑ Gauge of the OP5 Comparison Symbol, a Cartan Test, and Four Hidden Conditions

Purpose. Layer v4.9.0-r71 found that the free function of the Ω-augmented OP5 comparison symbol is an exact gauge symmetry $\delta\Psi = f\,\partial_\vartheta\Psi$ and named the next step: fix the ϑ gauge, then test involution. On 3 November 2026 the author asked for exactly this. This layer records the gauge, one confirmed pre-registered Cartan test (H51), three further pre-registered tests that turned out invalid by their own criteria (H52, H53, H54), and one exploration of the Spencer cohomology of the gauged symbol. Everything concerns the comparison model of §44 only. Earlier layers are not rewritten.
Source-only status. This v4.9.0-r72 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r72 results

Notation as in §A.272: $R = (F, G, dg)$, J the completed first-order symbol, θ the symbol of Ω, g_p the degree-p symbol of the Ω-augmented system. g'_p is the symbol after the gauge condition of r72.1, c_k denotes the Taylor coefficients of degree k of Ψ, and P_5, P_{56}, P_{567} are the dimensions of the projections of a solution space of the linearized jet tower onto c_5, (c_5, c_6), (c_5, c_6, c_7).

Theoremlet v4.9.0-r72.1 (the ϑ gauge).

(a) The gauge vector $v = \partial_\vartheta\Psi$ at the germ is a common kernel vector of all four blocks of J (smallest singular value 5.5×10^{-15} against 0.13), and all blocks of θ annihilate it (1.6×10^{-14}).

(b) The algebraic condition $s(\Psi) = S(x)$ with $s = \Psi^\dagger K_2 \Psi$, one component of the current, is transversal: $|ds \cdot v|/|ds| = 0.45$, the largest value among $\Psi^\dagger\Psi$, $\Psi^\dagger K_a\Psi$, $\Psi^\dagger Q_a\Psi$, $\Psi^\dagger(I \otimes \beta)\Psi$. $\Psi^\dagger(Z \otimes \beta)\Psi$ is fixed by g and blind (3.7×10^{-16}). Because $ds \cdot v \neq 0$, the condition fixes f locally.

(c) At symbol level every transversal algebraic gauge gives $g'_p \cong g_p/(S^p \otimes v)$. Measured with clean gaps: $g'_p = g_p - \binom{p+3}{3} = 117, 188, 273, 371, 481, 606, 745$ for $p = 2, \ldots, 8$.

Theoremlet v4.9.0-r72.2 (H51 confirmed: the gauged symbol is involutive at order 6). Pre-registered before the computation. Cartan's test $\dim g'_{p+1} \leq \sum_{k=0}^{3} \dim g'^{(k)}_p$ in three random orthonormal frames, all ranks with gap ratio at least 10^7:

p	$\dim g'^{(k)}_p$, $k = 0..3$	sum	$\dim g'_{p+1}$	verdict
4	273, 85, 14, 0	372	371	not involutive
5	371, 98, 14, 0	483	481	not involutive
6	481, 111, 14, 0	606	606	involutive
7	606, 125, 14, 0	745	745	involutive

After the gauge no free function of four variables remains; formally the symbol leaves 14 real functions of three variables.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Record v4.9.0-r72.3 (H52 invalid; the stored fifth jet does not extend). Pre-registered test of formal integrability from order 6 with the stored jet through degree 5 held fixed. The ranks in the large systems had no clean gap (ratio 70 to 600); the test is invalid by its criterion. A clean subsystem (gap 6×10^7) shows that the stored degree-5 coefficients cannot be extended at all: the compatibility residual is 4.0×10^{-3}, at the level of the positive control. They were solved on 30 October only against the degree-4 equations and violate the once prolonged Ω condition. The counts agree: without the prolonged Ω rows the fibres have $381 = 371 + 10$ and $501 = 481 + 20$ dimensions. This was an error of the test design, not a property of the system.

Record v4.9.0-r72.4 (H53 invalid; the tower over the repaired fourth jet). Corrected design: only the repaired jet through degree 4 is fixed, $c_5, \ldots, c_N$ are free, and the linearization $L = A^\mu(x)\partial_\mu + B(x)$ is exact (symmetric probes $x^{me_\nu}e_i$; even terms cancel, cubic terms start above the degrees used). Reference residual 4.6×10^{-14}, positive control 3.5×10^{-3}.

(a) With equations through degree 6 (F_7) everything is clean: residual 4.0×10^{-12}, $P_5 = 371$, $P_{56} - P_5 = 481$. Up to these equations there are no conditions beyond (F, G, dg), Ω and the gauge.

(b) With equations through degree 7 (F_8, 9865×10200) P_{56} drops from 852 to 848, while $P_{567} - P_{56} = 606 = \dim g'_7$. One rank had gap ratio 4×10^3; the test is invalid by its criterion.

Theoremlet v4.9.0-r72.5 (exploration: Spencer cohomology of the gauged symbol). Not pre-registered. With $H^{k,2} = \dim\ker(\delta : g_k \otimes \Lambda^2 \to S^{k-1} \otimes \Lambda^3) - \operatorname{rank}(\delta : g_{k+1} \otimes \Lambda^1 \to S^k \otimes \Lambda^2)$, all gaps at least 4×10^6: $H^{4,2} = 0$ and $H^{5,2} = 4$ for the ungauged symbol (check against r71.2) and $H^{4,2}(g') = 0$, $H^{5,2}(g') = 4$ for the gauged one. At most four hidden conditions on 6-jets are possible, none on 5-jets. The gauge does not change this, so the four do not come from the ϑ freedom.

Record v4.9.0-r72.6 (H54 invalid; an error bound says exactly four). Pre-registered. By Weyl's inequality an error ΔE moves every singular value by at most $\|\Delta E\|_2$. Measured from independent estimates: the reduced matrix behind P_{56} contains no degree-3 coefficients of A, B and has error $\leq 7 \times 10^{-13}$ against its smallest counted value 1.0×10^{-9}; the full F_8 has error about 5×10^{-9} against its smallest nonzero value 2.7×10^{-6}, and two perturbed variants keep rank 7957. Hence $P_{56} \leq 848$, and with r72.5 $P_{56} \geq 848$. The criterion also demanded a gap ratio of 10^4 in the perturbed variants, which the perturbation itself makes impossible (ratios 3×10^3, 7×10^3); the test is invalid. The pre-registered variant (degree-3 part of B) does not enter F_8 at all; a stricter substitute was used and recorded.

Proof and measurement sources. r72.1 by `k136_eichung.py`. r72.2 by `k137_cartan_test.py`. r72.3 by `k138_tps_d.py` and `k139_fortsetzung.py`. r72.4 by `k140_fortsetzung_exakt.py` and `k140_f8_schritte.py`. r72.5 by `k141_spencer_geeicht.py`. r72.6 by `k142_h54.py`. The notes `Ergebnis_H51_Cartan_Test_geeichtes_Symbol.md` to `Ergebnis_H54_vier_Bedingungen_6Jets.md` and `Erkundung_Spencer_geeichtes_Symbol.md` give the numbers. □

Consequence. The gauged symbol is involutive from order 6 (H51). The gauged system itself is, by all evidence, not formally integrable at order 6: four further conditions on 6-jets appear once the degree-7 equations are used (r72.4(b), r72.5, r72.6), but no test has

© 2026 Marc Brendecke. Preprint — not peer reviewed.

confirmed this formally. A confirming test needs new data, for example a second base point. If confirmed, the four conditions must be determined, added to the system, and the Cartan test repeated. OP5 is not closed.

v4.9.0-r72 status in the three frames

- All statements concern the fixed OP5 comparison model of §44 in one chart with one coframe reconstructed from Ψ. No frame map M, T or P acts on them; r72.1 to r72.6 are frame-free within the model.
- The gauge function $\Psi^\dagger K_2 \Psi$ is a current component in the model coframe, not a frame-resolved quantity.

No statement of this layer concerns colour or charge.

v4.9.0-r72 register entries

statement	status after v4.9.0-r72
ϑ gauge	algebraic, $\Psi^\dagger K_2 \Psi = S(x)$; removes exactly the free function
H51	confirmed: gauged symbol involutive first at $p = 6$
H52, H53, H54	invalid by their criteria (test design); findings recorded
stored degree-5 coefficients	not extendable; degree 5 must stay free
$H^{5,2}(g')$	4 (exploration)
hidden conditions on 6-jets	by all evidence exactly four; not formally confirmed
next step	second base point (H55), then the explicit conditions and a new Cartan test

v4.9.0-r72 ledger and guard update

$$R_{\mathrm{chk}}^{v490\text{-}r72} = (1, 1, 0)$$

for (gauge fixed and H51 confirmed; H52 to H54 recorded as invalid; OP5 not closed).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r72,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r72} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r73 Source-Only: Four Hidden Sixth-Order Conditions of the Gauged OP5 Comparison System

Purpose. Layer v4.9.0-r72 found, by all evidence but without formal confirmation, four hidden conditions on 6-jets of the ϑ-gauged OP5 comparison system (§A.272). On 3 November 2026 the author asked to confirm them, to determine them explicitly, and then to repeat the Cartan test. This layer records one confirmed pre-registered test (H55), the explicit determination and structure of the four conditions (exploration), and two pre-registered Cartan tests of the extended symbol that are invalid by their criteria (H56, H57). Everything concerns the comparison model of §44 only. Earlier layers are not rewritten.
Source-only status. This v4.9.0-r73 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r73 results

Notation as in §A.272. K_N is the projection onto (c_5, c_6) of the solution space of the linearized tower with the 4-jet fixed and equations through degree $N-1$; q is the light cone of the reconstructed coframe; Π_0, Π_1 are the two skew characteristic planes of r71.3.

Theoremlet v4.9.0-r73.1 (H55 confirmed: the four conditions at a second base point). Pre-registered. A second 4-jet $c_4' = c_4^{\mathrm{rep}} + \alpha z_4$ from the kernel of the degree-3/4 tower (seed 55, distance 50 %) is extendable (residual 5.0×10^{-14}; positive control 5.3×10^{-4}). There $P_{56} = 852$ with equations through degree 6 and $P_{56} = 848$ with equations through degree 7; all gaps at least 4×10^4; error bounds 710 and 1390 times below the deciding singular values. The reduced matrix behind P_{56} depends only on the jet through degree 2 and is the same at both points; the new information is the rank of the full system at a changed 4-jet.

Record v4.9.0-r73.2 (exploration: the four conditions determined). At the first base point K_7 has dimension 852, $K_8 \subset K_7$ dimension 848; the four conditions are the linear forms on K_7 that vanish on K_8.

(a) They are genuine sixth-order equations: 99 % of their weight lies on c_6, and on the gauged symbol fibre g_6' they have rank 4. The complete symbol at order 6 has dimension 477, not 481.

(b) They act only on light-like plane waves. Plane waves $\xi^6 \otimes u$ in the symbol exist only on the cone $q(\xi) = 0$ (seven polarizations per direction) and on isolated characteristic directions; the conditions are nonzero on the cone and vanish (9×10^{-15}) off it.

(c) On every null direction they have rank 1 on the seven polarizations (next singular value $\leq 3 \times 10^{-9}$ relative): they fix one scalar combination $s(\xi, u)$.

(d) The direction in the space of conditions is quadratic in ξ: $n(\xi) \propto M(\xi \otimes \xi)$ to 1.6×10^{-9} on held-out null directions, while no linear law fits. The four quadrics vanishing on $\Pi_0 \cup \Pi_1$, determined independently, lie in $\mathrm{span}(M, q)$: rank 5 with and without them (gap 0.2 against 1.5×10^{-13}).

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Reading: on the cone the conditions are $\sigma_k(\xi,u) = d_k(\xi)\, s(\xi,u)$ up to a fixed recombination, with d_k the four plane quadrics. These are the factors of the four degree-7 relations $q\, d_k\, \Omega \in \langle J \rangle$ of r71.3: identities at symbol level that leave one sixth-order condition each once lower-order terms are kept, in line with $H^{5,2}(g') = 4$ (r72.5).

Record v4.9.0-r73.3 (H56 and H57 invalid: the Cartan test of the extended symbol). Pre-registered (H57 explicitly not blind). The extended symbol g'' adds the four conditions and their prolongations to g'. In both tests and all three frames: $\dim g''_6 = 477$, $\dim g''_7 = 594$, not involutive at $p = 6$. From $p = 8$ the symbol ranks have gap ratios of only 8×10^2 to 10^3; one singular value (5.6×10^{-9}) decides whether $\dim g''_8$ is 723 (H56: involutive first at $p = 8$) or 724 (H57: at $p = 7$). Both tests are invalid. The cause is precision: θ stems from the older germ (deviation about 10^{-7}), the four conditions from the repaired germ, and genuine singular values near 10^{-8} at $p = 8$ cannot be separated from that deviation in double precision.

Proof and measurement sources. r73.1 by `k143_h55.py`. r73.2 by `k144_vier_bedingungen.py` and `k145_charakteristik.py`; the conditions are stored in `k144_bedingungen_p1.npy`. r73.3 by `k146_cartan_erweitert.py` and `k147_cartan_projiziert.py`. The notes `Ergebnis_H55_zweiter_Grundpunkt.md`, `Erkundung_vier_Bedingungen_6Jets.md`, `Ergebnis_H56_Cartan_Test_erweitertes_Symbol.md` and `Ergebnis_H57_Cartan_Test_projizierte_Bedingungen.md` give the numbers. □

Consequence. The gauged comparison system is not formally integrable at order 6: it has four further sixth-order conditions, now determined and understood as the curvature of the four relations $q\, d_k\, \Omega$. Whether the extended symbol becomes involutive at order 7 or 8 cannot be decided in double precision with the stored θ. The next step is a consistent data set at the repaired germ (Ω and the four conditions) in higher precision. OP5 is not closed.

v4.9.0-r73 status in the three frames

- All statements concern the fixed OP5 comparison model in one chart with one coframe reconstructed from Ψ. No frame map M, T or P acts on them; r73.1 to r73.3 are frame-free within the model.
- The light cone q and the planes Π_0, Π_1 belong to the reconstructed geometry of the model, not to the photon frame P.

No statement of this layer concerns colour or charge.

v4.9.0-r73 register entries

statement	status after v4.9.0-r73
four hidden conditions on 6-jets	confirmed at a second base point (H55)
their form	order 6; only on the light cone; $d_k(\xi)\, s(\xi,u)$ with the plane quadrics (exploration)
complete symbol at order 6	dimension 477
Cartan test of the extended symbol	H56, H57 invalid; not involutive at 6; 7 or 8 undecided in double precision
next step	consistent high-precision data at the repaired germ

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r73 ledger and guard update

$$R_{\mathrm{chk}}^{v490\text{-}r73} = (1, 1, 0)$$

for (four conditions confirmed and determined; H56 and H57 recorded as invalid; OP5 not closed).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r73,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r73} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r74 Source-Only: Consistent Data, Spencer Bounds and the Precision Limit of the Extended OP5 Symbol

Purpose. Layer v4.9.0-r73 left open whether the symbol g'' extended by the four hidden sixth-order conditions becomes involutive at order 7 or 8, and whether the extended system has further hidden conditions (§A.272). On 3 November 2026 the author asked to rebuild the data consistently, to bound further conditions, and to try a staged elimination and a plane-wave reformulation. This layer records one pre-registered test (H58, invalid by its criterion), two explorations and one method check. Everything concerns the comparison model of §44 only. Earlier layers are not rewritten.
Source-only status. This v4.9.0-r74 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r74 results

Record v4.9.0-r74.1 (H58 invalid: consistent data). Pre-registered, not blind. All parts of the symbol now come from the exact linearization at the repaired germ: $J = A(0)$ (deviation from the stored J 1.2×10^{-13}), θ rebuilt as the linear form on the admissible 3-jets that vanishes once the degree-3 equations are added ($189 \to 188$, inclusion to 2.8×10^{-11}), the four conditions of r73.2, and the gauge. Without the four conditions the known dimensions $117, \dots, 898$ ($p = 2, \dots, 9$) are reproduced with gaps $\geq 5 \times 10^6$. With them, in all three frames: $\dim g''_p = 477, 594, 723, 865$ ($p = 6, \dots, 9$), not involutive at $p = 6, 7$, involutive at $p = 8$, characters $723, 130, 12, 0$. The ambiguous singular value of H57 is gone (noise 1.7×10^{-12}, smallest genuine value 8.1×10^{-9}), so $\dim g''_8 = 724$ was an artifact. Five ranks have gap ratios 3×10^3 to 8×10^3 instead of the required 10^4; the test is invalid.

Theoremlet v4.9.0-r74.2 (exploration: Spencer cohomology of the extended symbol). With consistent data, all δ-rank gaps $\geq 3 \times 10^4$: for the gauged symbol g' $H^{5,2} = 4$, $H^{6,2} = H^{7,2} = 0$ (check against r72.5); for the extended symbol g'' $H^{5,2} = 8$, $H^{6,2} = 4$, $H^{7,2} = 3$. The four conditions were the only possible obstructions of the gauged system through $k = 7$; the new sixth-order equations themselves leave room for at most four hidden conditions on 7-jets and three on 8-jets. Whether they occur is decided by lower-order terms, not by the symbol.

Record v4.9.0-r74.3 (method check: staged elimination and conditioning). A staged elimination of the linearized tower (top degree first, through left kernels of symbol-sized blocks) needs little memory and reproduces 371, 852 and 481, but gives 476 instead of 477 with a gap of 4×10^2. The cause is structural: the prolonged symbol of J alone is well conditioned (smallest genuine value 4×10^{-5}, noise 10^{-15}, unchanged by Ruiz equilibration), while Ω and the four conditions are nearly dependent on the J rows (independent weight about 10^{-8}), because they carry new information only on the cone and the planes. Double precision separates one or two such levels, not three.

Record v4.9.0-r74.4 (exploration: plane waves on the cone). If the symbol module were reduced, plane waves $\xi^p \otimes u$ on the characteristic set would span every g_p with well-conditioned data. Without the four conditions, 300 null directions with seven polarizations each span $115, 184, 267, 364, 475$ for $p = 2, \dots, 6$ (gaps $\geq 2 \times 10^{11}$) instead of

© 2026 Marc Brendecke. Preprint — not peer reviewed.

117, 188, 273, 371, 481. The deficit grows irregularly, which points to a non-reduced module; the simple reformulation does not carry.

Proof and measurement sources. r74.1 by `k148_konsistent.py`; r74.2 by `k149_spencer_erweitert.py`; r74.3 by `k150_stufen.py` and `k151_kondition.py`; r74.4 by `k152_wellen.py`. The notes `Ergebnis_H58_konsistente_Daten.md`, `Erkundung_Spencer_erweitertes_Symbol.md`, `Methodenpruefung_Elimination_Kondition.md` and `Erkundung_ebene_Wellen.md` give the numbers. □

Consequence. By all evidence the extended symbol is involutive first at order 8, with 12 functions of three variables. Further hidden conditions (up to four on 7-jets, three on 8-jets) are not excluded. Deciding them needs either the whole chain (germ, linearization, elimination) in higher precision or a reformulation that handles the multiplicities of the symbol module on the characteristic set. OP5 is not closed.

v4.9.0-r74 status in the three frames

- All statements concern the fixed OP5 comparison model in one chart with one coframe reconstructed from Ψ. No frame map M, T or P acts on them; r74.1 to r74.4 are frame-free within the model.

No statement of this layer concerns colour or charge.

v4.9.0-r74 register entries

statement	status after v4.9.0-r74
θ at the repaired germ	rebuilt consistently (replaces the stored θ)
extended symbol	involutive first at $p = 8$ by all evidence (H56, H58); formally unconfirmed
further hidden conditions	room for 4 on 7-jets, 3 on 8-jets (Spencer); undecided
precision limit	structural (Ω nearly dependent on J); staged elimination and plane waves do not lift it
next step	higher precision of the whole chain, or multiplicity-aware reformulation

v4.9.0-r74 ledger and guard update

$$R_{\mathrm{chk}}^{v490\text{-}r74} = (1, 0, 0)$$

for (consistent data and Spencer bounds established; H58 invalid; OP5 not closed).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r74,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r74} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r75 Source-Only: Renaming the Millennium Matrix to Compensator Matrix

Purpose. The 16×16 matrix $\mathcal{M}_{16}$ of §38 was called the Millennium matrix. The name suggested a link to the Millennium Prize Problems that the Companion does not claim, and the author had asked several times for a name that says what the object does. On 3 November 2026 the author decided: $\mathcal{M}_{16}$ is called the *compensator matrix*, because it carries several Heisenberg compensators and extends the local rule $C = \frac{1}{2}(\mathrm{id} + J)$ to all layers (§38; in v4.9.0-r42 one of its compensators acts as chirality projector). This layer records the renaming. No mathematical statement changes. Earlier layers are not rewritten.
Source-only status. This v4.9.0-r75 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r75 results

Record v4.9.0-r75.1 (naming rule). From this layer on, $\mathcal{M}_{16}$ is the *compensator matrix*. Unchanged are the symbol $\mathcal{M}_{16}$ (and $\mathcal{M}$, $\mathcal{M}^{\mu}{}_{\nu}$ where used), every label (for example `sec:millennium-hc`), and the abbreviation MM in MM/HC, MM split and similar compounds, which is kept as a historical abbreviation for the matrix $\mathcal{M}_{16}$. Wherever an earlier layer, an inherited appendix or a note says "Millennium matrix", read "compensator matrix".

Record v4.9.0-r75.2 (where the name was changed). The main body uses the new name in the nomenclature, in the sections on the probe–closure field system, the structural core, the HC hypotheses and the master model, including the section title of §38 and the definition, which names the former term once. On the research line the 4.8 sections on the native operator link and the coupling candidate are changed likewise, and the PDF keywords. In the local monolith the 27 main-body occurrences are changed by the patch list `llm/revision_spec_v490_r75.py`; the main body is again word-identical with the research line. Inherited appendices (repair, transport and OP protocols, technical tools, release records), the archive and all earlier layers keep the old wording verbatim. The bibliography entry "Set Theory: The Third Millennium Edition" is a title and not affected.

Consequence. A name change only. No definition, theorem, proof, number, register entry or guard is changed.

v4.9.0-r75 status in the three frames

- The renaming concerns notation only. No frame map M, T or P acts on it; the frame statements of §38 and of v4.9.0-r42 are unchanged.

No statement of this layer concerns colour or charge.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r75 register entries

statement	status after v4.9.0-r75
name of $\mathcal{M}_{16}$	compensator matrix (formerly Millennium matrix)
symbol, labels, abbreviation MM	unchanged
earlier layers and inherited appendices	old wording kept; read with the new name

v4.9.0-r75 ledger and guard update

$$R_{\mathrm{chk}}^{v490\text{-}r75} = (1, 0, 0)$$

for (renaming recorded; no statement changed).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r75,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r75} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r76 Source-Only: Correction to r75 — the Earlier Name "Layer-Transport Matrix"

Purpose. Layer v4.9.0-r75 (§A.272) renamed $\mathcal{M}_{16}$ from "Millennium matrix" to "compensator matrix". It overlooked that layer v4.9.0-r05 had already, at the author's request of 21 October 2026, named $\mathcal{M}_{16}$ the *layer-transport matrix* and its restriction to the (E, B)-doublet the *port-weighting matrix*, without changing the main body. This layer corrects the record. Earlier layers are not rewritten.
Source-only status. This v4.9.0-r76 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r76 results

Record v4.9.0-r76.1 (correction of r75). The statement of r75 that $\mathcal{M}_{16}$ "was called the Millennium matrix" is incomplete: from r05 to r75 the declared name was "layer-transport matrix", while the main body still said "Millennium matrix". The author confirmed on 3 November 2026, after being shown r05, that *compensator matrix* is the valid name. It replaces "layer-transport matrix". Reason: the defining properties of $\mathcal{M}_{16}$ in §38 are that it extends $C = \frac{1}{2}(\mathrm{id} + J)$ to all layers and has C as a spectral projector; "layer transport" describes only its place in the master chain.

Record v4.9.0-r76.2 (doublet restriction). The name *port-weighting matrix* of r05 for the restriction of $\mathcal{M}_{16}$ to the (E, B)-doublet stays. It keeps the compensator matrix $\mathcal{M}_{16}$ apart from the 2×2 matrix of the single compensator on the doublet (v4.9.0-r03).

Reading rule. In earlier text "Millennium matrix" and, in layers r05 to r74, "layer-transport matrix" both mean the compensator matrix $\mathcal{M}_{16}$. Symbol, labels and the abbreviation MM are unchanged (r75.1). No mathematical statement changes.

v4.9.0-r76 status in the three frames

- The correction concerns notation only. No frame map M, T or P acts on it.

No statement of this layer concerns colour or charge.

v4.9.0-r76 register entries

statement	status after v4.9.0-r76
name of $\mathcal{M}_{16}$	compensator matrix; replaces "layer-transport matrix" (r05) and "Millennium matrix"
restriction to the (E, B)-doublet	port-weighting matrix (r05), unchanged
r75 record	corrected: the r05 name was omitted

v4.9.0-r76 ledger and guard update

$$R_{\mathrm{chk}}^{v490\text{-}r76} = (1, 0, 0)$$

for (naming record corrected; no statement changed).

© 2026 Marc Brendecke. Preprint — not peer reviewed.

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r76,+)} = (0,1,1,1,1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r76} = (1,1,1,1,1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r77 Source-Only: High Precision, Formal Integrability at the Germ and the Covariance of the OP5 Comparison System

Purpose. Layer v4.9.0-r74 left two questions open (§A.272). The first is whether the extended symbol g'' is involutive first at order 8. The second is whether the system has further hidden conditions on 7- and 8-jets; Spencer allows up to four and three. Both were blocked by double precision. On 3 and 4 November 2026 the author asked for three things:

- to repeat the whole chain in high precision;
- to run the decisive eliminations on parallel GitHub runners;
- to look for links of the result to physics.

This layer records one high-precision exploration and six pre-registered tests (H59 to H64). Three tests are confirmed and three refuted. Everything concerns the comparison model of §44 only. Earlier layers are not rewritten.
Source-only status. This v4.9.0-r77 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r77 results

Record v4.9.0-r77.1 (exploration: the four conditions and the Cartan test in high precision). Not pre-registered and not blind: the values of r74.1 were known.

- *Linearization.* The linearization was rebuilt pointwise in 160 digits. The deviation from the earlier high-precision data is 2×10^{-21} in degree 1 and 2×10^{-19} in degree 2; these earlier data set the old precision limit.
- *Eliminations.* c_8 and c_7 were eliminated in ball arithmetic with 70 digits. The Newton residuals are 10^{-66} and 9×10^{-52}.
- *Fibre of 6-jets.* The Gram matrix of the derived conditions on the 501-dimensional fibre has 24 genuine eigenvalues, the smallest 5.1×10^{-13}, then 1.9×10^{-71}: a gap of 10^{58}. The fibre therefore has dimension exactly 477. It consists of 20 Ω-prolongations and exactly four new conditions.
- *Cartan test.* On this basis $\dim g''_p = 477, 594, 723, 865$ for $p = 6, \ldots, 9$. The characters at $p = 8$ are $723, 130, 12, 0$ in all three frames. The symbol is not involutive at $p = 6, 7$ and is involutive at $p = 8$, with rank gaps $\geq 10^8$.
- *A withdrawn local finding.* An intermediate local finding (only 48 instead of 112 identities at order 4) came from a conversion with 15 digits and was withdrawn before this layer.

Theoremlet v4.9.0-r77.2 (H59 confirmed: no further hidden conditions on 7- and 8-jets). Pre-registered, not blind (the symbol dimensions were known). Before the decisive runs, a dated addendum relaxed the residual bound from 10^{-40} to 10^{-30}, after a function test at $q = 4$.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

- *Setup.* The jet through degree $q-1$ is fixed and the equations go through degree $q+1$. c_{q+2} and c_{q+1} are eliminated in 70 digits on parallel runners (16 GB each).
- *Result.* The fibres are $F_6 = 477$ (positive control), $F_7 = 637 - 43 = 594$ and $F_8 = 789 - 66 = 723$. The Gram gaps are 1.6×10^{58}, 2.8×10^{58} and 3.9×10^{56}; all residuals are $\leq 1.1 \times 10^{-51}$.
- *Reading.* The Spencer bounds of r74.2 are not attained. Together with the involutivity of g_8'', the completed system is formally integrable at the germ by Goldschmidt's criterion, provided two elimination levels catch all obstructions as at order 6. The general formal solution depends on 12 functions of three variables.

Theoremlet v4.9.0-r77.3 (H60 confirmed: six polarizations per null direction). Pre-registered. With $s_3 = 12$ and a cone of degree 2, the multiplicity on the cone is 6. If the symbol module is reduced there, this gives six plane waves per null direction. Over 200 null directions at each of $p = 6, \ldots, 9$, the polarization space $\{u : \xi^p \otimes u \in g_p''\}$ has dimension 6 every time, with gaps $\geq 3 \times 10^7$. Off the cone the dimension is 0. The twelve Cauchy data are therefore six real massless modes. The deficit of plane waves seen in r74.4 comes from the planes Π_0, Π_1 and from isolated directions, not from the generic cone.

Theoremlet v4.9.0-r77.4 (H61 refuted: no Weyl type in the naive spin assignment). Pre-registered. The test reads Ψ as three Dirac spinors in the Dirac basis, with spin $I_3 \otimes \Sigma$, chirality $I_3 \otimes \gamma_5$ and complex structure J.

- The six-dimensional polarization space is invariant under none of J, γ_5 or the helicity: the median residuals are 1.0, 0.99 and 0.5.
- Post hoc (exploration): $P + JP$ has dimension 12 but is not the kernel of the massless Dirac symbol, so a Majorana reading also fails. The modes spread over all three blocks of Ψ, most of them over the third block, which does not enter the coframe.

Theoremlet v4.9.0-r77.5 (H62 refuted: the symbol is invariant only under scaling). Pre-registered. The stabilizer $\{(A, B) \in \mathfrak{gl}(4) \times \mathfrak{gl}(24) : X_{A,B} g_p'' \subset g_p''\}$ has dimension 50 at $p = 6, 7, 8$, with gaps $\geq 7 \times 10^8$. Its spacetime part is only $A \propto 1$, and its intersection with $\mathfrak{so}(Q)$ is 0, not 6. The internal part (dimension 49) is closed under commutators. Its Killing form has signature $(3, 1, 45)$, so the internal part is mostly nilpotent; no compact semisimple part is visible (exploration). A first run that formed the Gram matrix as a difference cancelled digits and is not counted. An addendum places the result: invariance at a fixed background is expected to fail, because every symmetry must preserve the characteristic set, and $J(\xi)$ loses rank on $\Pi_0 \cup \Pi_1$. These planes are not q-orthogonal, are not spanned by bilinears of Ψ (r71, `k122`) and arise in the G part (`k126`).

Theoremlet v4.9.0-r77.6 (H63 refuted: already F is not Lorentz symmetric at a fixed background). Pre-registered. For the blocks of J the stabilizer $\{(A, B, C) : \sum_\mu A_{\mu\nu} X_\mu + X_\nu B - C X_\nu = 0\}$ has spacetime parts of dimension 7 for F and 1 for $F + G$ and $F + G + dg$. In all three cases the intersection with $\mathfrak{so}(Q)$ is 0. Part (a), predicting 6 for F, fails; part (b), predicting ≤ 1 when G is included, holds. The six non-scaling directions of F lie near $\mathfrak{so}(Q)$ (cosines 0.74 to 0.9998) but form no $\mathfrak{so}(Q')$ for any form Q' (exploration). The germ Ψ enters the principal symbol beyond a tetrad, through the coframe built from Ψ. The symbol is then invariant only under the stabilizer of Ψ, which is trivial for a generic spinor.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Theoremlet v4.9.0-r77.7 (H64 confirmed: R is covariant). Pre-registered. Run 1 was invalid: its positive control was 2.1×10^{-7}, above the bound 10^{-9}, because of an unregularized pseudoinverse of the rank-31 symbol. A dated addendum preceded run 2, which is not blind. The construction $R(\Psi, \partial\Psi; A, \partial A, h, \partial h)$ was rebuilt after three transformations: a random linear change of coordinates in $GL(4)$, a random internal Lorentz transformation of Ψ, and both together. In each case the jets and the background transform tensorially.

- With a free row transformation C, the relative residual $\min_C \|J'L - CJ\|/\|J'L\|$ is $\leq 4.1 \times 10^{-10}$ for F, G and dg.
- The positive control gives $\leq 1.1 \times 10^{-11}$.
- The negative control, with the background held fixed, gives 0.26 for F and G.

At symbol level, R is therefore coordinate covariant and internally Lorentz covariant. The breaking found in H62 and H63 comes entirely from the background (Ψ at the germ, A, h).

Proof and measurement sources.

- r77.1: `k153b_hp_linear.py`, `k154_hp_tools.py`, `k157_hp_bedingungen.py`, `k158_cartan_hp.py`.
- r77.2: `k159_hp_faser.py`, with the workflow `hp-faser.yml` and the data under `ci/hp/`.
- r77.3: `k160_polarisationen.py`.
- r77.4: `k161_spin.py`, `k161b_majorana.py`, `k161c_bloecke.py`.
- r77.5: `k162_symmetrien.py`, `k162b_innere.py`.
- r77.6: `k163_lorentz_F.py`, `k163b_eigene_metrik.py`.
- r77.7: `k164_kovarianz.py`.

The pre-registration notes `Vorab_Notiz_H59` to `H64` and the result notes lie in `notes/research/predictions/`. □

Consequence. The extended comparison system is formally integrable at the germ, with six real massless modes. Its construction is covariant; Lorentz symmetry is broken only by the background. The physical questions therefore move to the background:

- what the fields A and h are;
- whether the background solves covariant equations;
- whether Ω, θ, the gauge $\Psi^\dagger K_2 \Psi = S$ and the four conditions are covariant;
- which Lorentz action, if any, acts on the six modes.

OP5 is not closed.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r77 status in the three frames

- All statements concern the fixed OP5 comparison model in one chart, with one coframe reconstructed from Ψ. No frame map M, T or P acts on them. The Lorentz transformations of r77.5 to r77.7 act inside the model and are not the photon frame P.

No statement of this layer concerns colour or charge. The internal $3\otimes$ structure of Ψ is not read as generations.

v4.9.0-r77 register entries

statement	status after v4.9.0-r77
four hidden conditions	exactly four (gap 10^{58}, high precision, exploration)
extended symbol	involutive first at $p = 8$, characters $723, 130, 12, 0$ (exploration)
further hidden conditions	none on 7- and 8-jets (H59 confirmed); formally integrable at the germ
massless modes	six real polarizations per null direction (H60 confirmed)
spin of the modes	not Weyl or Majorana in the naive assignment (H61 refuted); undetermined
symmetry at fixed background	only scaling (H62, H63 refuted); structurally expected
covariance of R	coordinate and internal Lorentz covariant at symbol level (H64 confirmed, run 2 not blind)
next step	covariance of Ω, θ, the gauge and the four conditions; origin of the background A, h

v4.9.0-r77 ledger and guard update

$$R_{\mathrm{chk}}^{v490\text{-}r77} = (1, 1, 0)$$

for (formal integrability at the germ and covariance of R established; H61 to H63 recorded as refuted; OP5 not closed).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r77,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r77} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r78 Source-Only: Covariance of the OP5 Comparison System at Value Level and the Stabilizer of the Gauge

Purpose. Layer v4.9.0-r77 showed that the construction R of the OP5 comparison system is covariant at the level of its first-order symbol (H64, §A.272). On 4 November 2026 the author asked to check the remaining building blocks: Ω and θ, the four hidden conditions and the gauge $\Psi^\dagger K_2 \Psi = S$. Ω is derived from R alone, the four conditions from R together with the gauge. Hence it suffices to establish two things:

- the covariance of R at value level, with all lower-order terms;
- the stabilizer of the gauge.

This layer records one pre-registered test that is invalid by its criterion (H65) and one confirmed test (H66). Everything concerns the comparison model of §44 only. Earlier layers are not rewritten.

Source-only status. This v4.9.0-r78 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r78 results

Record v4.9.0-r78.1 (H65 invalid: stabilizer of the gauge as predicted, value level below the control bound). Pre-registered.

- *Part (b), the stabilizer of the gauge.* The generators $X \in \mathfrak{so}(1,3)$, in the spin representation, that preserve $\Psi^\dagger K_2 \Psi$ are those with $K_2 X + X^\dagger K_2 = 0$. They span a three-dimensional subalgebra: the rotation about y and the boosts along x and z. This is $\mathfrak{so}(1,2)$, the stabilizer of e_y, as predicted. The gauge is a non-covariant gauge fixing that singles out one spatial direction.
- *Part (a), value level.* Run 1 was invalid because the row transformation was taken from the symbol, which does not determine it off the solution set (positive control 0.24). A dated addendum preceded run 2, which is not blind and fits the row transformation from values. Run 2 gave residuals $\leq 1.1 \times 10^{-10}$, but its positive control (5.6×10^{-11}) missed the pre-registered bound 10^{-12}, which lay below the numerical floor. Part (a) is therefore invalid in both runs, and so is H65 as a whole. The bound was not relaxed after the fact.

Theoremlet v4.9.0-r78.2 (H66 confirmed: R is covariant at value level). Pre-registered, not blind (the values of H65 were known); the control bound is 10^{-9}.

- *Setup.* For each fixed pair (M, S), with $M \in GL(4)$ and $S \in \mathrm{Spin}(1,3)$, the row transformation C is fitted from 80 random data sets off the solution set: Ψ, its jets, and the background $A, \partial A, h, \partial h$. It is tested on 20 further data sets. Five pairs were used: a pure rotation, a pure boost, a pure change of coordinates, and two mixed pairs.
- *Result.* All relative residuals are $\leq 2.2 \times 10^{-11}$, and C is block diagonal on F, G, dg to 5×10^{-11}. The positive control gives $\leq 3.2 \times 10^{-13}$. The negative control, with the background held fixed, gives 57 for F and 1.2×10^3 for G.
- *Reading.* R is covariant with all lower-order terms. Hence Ω, which is derived from R alone, is covariant. The four hidden conditions are derived from R together with the gauge, so they are covariant at most under $\mathfrak{so}(1,2)$.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof and measurement sources. r78.1 by `k165_kovarianz_rest.py` (run 1) and `k165b_werte.py` (run 2); r78.2 by `k166_kovarianz_werte.py`. The notes `Vorab_Notiz_H65`, `H66` and the result notes lie in `notes/research/predictions/`. □

Consequence. The comparison system R and its integrability condition Ω are covariant. Covariance is broken in two places:

- by the prescribed background (Ψ at the germ, A, h);
- by the gauge, which fixes the y-component of a current and leaves $\mathfrak{so}(1,2)$.

Whether the number of the four hidden conditions and their support on the cone survive a change of gauge is open. So is a native dynamics for the background. OP5 is not closed.

v4.9.0-r78 status in the three frames

- All statements concern the fixed OP5 comparison model in one chart, with one coframe reconstructed from Ψ. No frame map M, T or P acts on them. The Lorentz transformations here act inside the model and are not the photon frame P.

No statement of this layer concerns colour or charge.

v4.9.0-r78 register entries

statement	status after v4.9.0-r78
covariance of R at value level	confirmed (H66, not blind); H65(a) invalid by its control bound
covariance of Ω	follows, since Ω is derived from R alone
gauge $\Psi^\dagger K_2 \Psi = S$	preserved exactly by $\mathfrak{so}(1,2)$, the stabilizer of e_y (H65(b))
four hidden conditions	covariant at most under $\mathfrak{so}(1,2)$; gauge independence open
next step	gauge independence of the four conditions; native dynamics of the background

v4.9.0-r78 ledger and guard update

$$R_{\text{chk}}^{v490\text{-}r78} = (1,1,0)$$

for (covariance of R and Ω established; H65 recorded as invalid; OP5 not closed).

The layer spends no guard:

$$R_{\text{glob}}^{(v490\text{-}r78,+)} = (0,1,1,1,1), \qquad \Gamma_{\text{guard}}^{v490\text{-}r78} = (1,1,1,1,1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r79 Source-Only: Gauge Independence of the Four Hidden Conditions of the OP5 Comparison System

Purpose. Layer v4.9.0-r78 showed that R and Ω are covariant, while the gauge $\Psi^\dagger K_2 \Psi = S$ keeps only $\mathfrak{so}(1,2)$ (§A.272). The four hidden conditions come from R together with this gauge, so their status was open: they could be content of the covariant system or an artefact of the non-covariant gauge fixing. On 4 November 2026 the author asked to decide this. This layer records one pre-registered test (H67), which is confirmed. Everything concerns the comparison model of §44 only. Earlier layers are not rewritten.
Source-only status. This v4.9.0-r79 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r79 results

Theoremlet v4.9.0-r79.1 (H67 confirmed: the four hidden conditions are gauge independent). Pre-registered. Layer r72.1 showed that several algebraic gauges $s = \Psi^\dagger M \Psi$ are transversal at the germ, and that every transversal algebraic gauge gives the same symbol. Whether the hidden conditions also agree depends on the lower-order terms.

- *Setup.* For each gauge the high-precision fibre computation of r77.2 was repeated at $q = 6$ (70 digits, one parallel runner per gauge). The gauge series is $w = 2(\mathrm{Re}\, M\Psi, \mathrm{Im}\, M\Psi)$, with transversalities $|ds \cdot v|/|ds|$ at the germ in brackets:
  - $M = K_2$, the control (0.45);
  - $M = 1$, the density (0.077);
  - $M = K_3$ (0.165);
  - $M = Q_3$ (0.256);
  - $M = I_3 \otimes \beta$, the Lorentz scalar $\bar\Psi\Psi$ (0.023).
- *Result.* In all five gauges the fibre of 6-jets is $501 - 24 = 477$. The Gram gaps are $\geq 4 \times 10^{56}$ and the Newton residuals $\leq 1.0 \times 10^{-51}$. The polarization space has dimension 6 on 100 null directions, with gaps $\geq 9 \times 10^5$, and dimension 0 off the cone.
- *Reading.* The number of the four conditions and their support on the light cone do not depend on the gauge. This holds in particular for the covariant gauge $\bar\Psi\Psi = S$.

Proof and measurement sources. r79.1 by `k167_hp_eichung.py`, run on GitHub runners with the workflow `hp-eichung.yml` and the data under `ci/hp_eich/`; the logs are `k167_logs.txt`. The note `Vorab_Notiz_H67` and the result note lie in `notes/research/predictions/`. □

Consequence. Together with H66, the four hidden conditions belong to the covariant content of the comparison system, not to the gauge fixing. For all structural results so far (four conditions, formal integrability at the germ, six massless modes), covariance is broken only by the prescribed background (Ψ at the germ, A, h). The next step is a native dynamics for this background. OP5 is not closed.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r79 status in the three frames

- All statements concern the fixed OP5 comparison model in one chart, with one coframe reconstructed from Ψ. No frame map M, T or P acts on them.

No statement of this layer concerns colour or charge.

v4.9.0-r79 register entries

statement	status after v4.9.0-r79
four hidden conditions	gauge independent in number and support (H67 confirmed, five gauges including $\bar{\Psi}\Psi$)
source of Lorentz breaking	only the prescribed background (Ψ, A, h)
next step	native dynamics of the background A, h

v4.9.0-r79 ledger and guard update

$$R_{\mathrm{chk}}^{v490\text{-}r79} = (1, 0, 0)$$

for (gauge independence of the four conditions established; OP5 not closed).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r79,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r79} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r80 Source-Only: The Background of the OP5 Comparison System Is Not Sourced by Its Spinors; Closing the Comparison-Model Branch

Purpose. Layer v4.9.0-r79 left one source of Lorentz breaking in the comparison system: the prescribed background (Ψ at the germ, A, h) (§A.272). This layer asks whether the background is generated by the three spinor blocks themselves. It records one exploration (polar/Fierz variables of the blocks) and one pre-registered test (H68), which is refuted. On 4 November 2026 the author decided to close the comparison-model branch at this point and to return to the common question behind OP1–OP5, a joint dynamics. The layer therefore ends with a closing register of layers r71–r80. Everything concerns the comparison model of §44 only. Earlier layers are not rewritten.
Source-only status. This v4.9.0-r80 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r80 results

Theoremlet v4.9.0-r80.1 (exploration: the three blocks are regular spinor fluids in a Hubble-like flow). Exploration without a criterion. At the germ $(t, x, y, z) = (1, 0.6, 0, 0)$ the polar (Fierz) variables of each block $u_n \in \mathbb{C}^4$ were computed in the Dirac basis, with $\Phi = \bar{u}u$, $\Theta = \bar{u}i\gamma_5 u$, $J^a = \bar{u}\gamma^a u$, $K^a = \bar{u}\gamma^a\gamma_5 u$.

- The Fierz identities $J^2 = \Phi^2 + \Theta^2 = -K^2$, $J \cdot K = 0$ hold to 10^{-14}.
- All three blocks lie in the regular Dirac sector with Yvon–Takabayasi angle $|\beta| \le 0.07°$; the densities are $\phi^2 = 5.97,\ 0.43,\ 6.38$.
- The velocities are $u^a = \gamma(1, \pm 0.6, 0, 0)$, $\gamma = 1.25$: blocks 0 and 1 move with $v = +x/t$, block 2 against it (the counterflow of the earlier series).
- All spin axes point along the azimuthal direction $\hat{\varphi} = e_y$, parallel for blocks 0, 1 and antiparallel for block 2. The same axis is singled out by the gauge K_2 (stabilizer $\mathfrak{so}(1,2)$, r78.1) and by the background $A = M(t)\, d\varphi$.

Theoremlet v4.9.0-r80.2 (H68 refuted: no Yang–Mills or Klein–Gordon equation with spinor sources). Pre-registered. The background is $A = M(t)\, d\varphi$ with $M(t) = \frac{3}{2\cos 2\theta}(\cos\theta\, \lambda_4 - \sin\theta\, \lambda_6)$, $\theta = \frac{1}{2}\arcsin(t/2)$, and $h = \gamma/\cos 2\theta$. Known in advance: $F = M'(t)\, dt \wedge d\varphi$, and the values of A, F generate $\mathfrak{k} = \mathrm{span}(\lambda_2, \lambda_4, \lambda_6) \cong \mathfrak{su}(2)$, with $Y_3 = \lambda_2 \in \mathfrak{k}$. In full $\mathfrak{su}(3)$ the test fails trivially, because the blocks carry a λ_3 density; it was therefore restricted to $\mathfrak{k}$.

- *Setup.* Metric of the model's own coframe $g = E^{\mathsf{T}}\eta E$ with $E^a_\mu = \mathrm{Re}(\Psi^\dagger Q_a D_\mu \Psi)$, $\sqrt{|g|} = |\det E|$; $D = \partial + iA$, $L^\nu = |g|^{-1/2}\partial_\mu(|g|^{1/2}F^{\mu\nu}) + i[A_\mu, F^{\mu\nu}]$; colour current $j^\nu_b = \bar{\Psi}\gamma^\nu(\lambda_b \otimes 1)\Psi$; scalar source $s = \bar{\Psi}(Y \otimes 1)\Psi$, the derivative of the Yukawa term. Truncated power series around the germ (repaired 4-jet), equations to degree 2, constants by least squares.
- *Hypotheses.* (a) $L^\nu_b = \kappa_A j^\nu_b$ for $b \in \{2, 4, 6\}$; (b) $\Box_g h = \kappa_h s$; (c) $\Box_g h + m^2 h = \kappa_h s$. Confirmed at relative residual $r \le 10^{-6}$, refuted at $r \ge 10^{-2}$.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

- *Controls.* Bianchi identity of L to degree 1: 1.2×10^{-14}; flat $\Box h$ from the series equals the exact value -6.344838895781825 to 2×10^{-15}; germ residual of $R \leq 2.8 \times 10^{-13}$.
- *Result.* (a) $r = 0.955$ (per degree 0.965, 0.963, 0.954, fitted κ_A between 0.12 and 22); (b) $r = 0.995$; (c) $r = 0.826$. All three refuted. Already degree 0, which does not depend on the higher jets and their ϑ-gauge, fails.
- *Why.* At the germ the sources in $\mathfrak{k}$ are about 10^{-3}, the left side is $O(0.1\text{–}0.4)$; for the scalar $s = 0.002$ against $\Box_g h = 6.91$. The directions do not match either. The spinor currents lie almost entirely outside $\mathfrak{k}$, in $\lambda_0, \lambda_3, \lambda_8$ and $\lambda_1, \lambda_5, \lambda_7$: the blocks are practically incoherent with respect to the off-diagonal generators that carry A.
- *Flat metric (exploration).* The analytic structure is confirmed: L has only the components (t, λ_2) and $(\hat{\varphi}, \lambda_4, \lambda_6)$. The fits fail as well ($r = 0.9997$, 1.000, 0.28).

Theoremlet v4.9.0-r80.3 (closing register of the comparison-model branch, r71–r80). Author decision of 4 November 2026. The comparison system is a fixed construction: its background is prescribed and, by r80.2, not generated by its spinors through Yang–Mills or Klein–Gordon equations. Its structural results are local and concern this construction only.

- *Established at the germ.* An exact ϑ-gauge symmetry (r71); involution of the gauged symbol from order 6 (r72); exactly four hidden conditions of order 6, supported on the light cone (r73, r77); formal integrability of the extended system (H59); six real polarizations per null direction (H60); covariance of R and Ω (H64, H66); gauge independence of the four conditions (H67).
- *Not established.* A native origin of the background (r80.2); any identification of the six modes with known fields (H61); any statement about the QSC model itself; the OP5 vacuum term.
- *Closed as a line of work.* Further refinements of this comparison system (Lorentz action on the six modes, second base point) are not pursued without a native dynamics. The next step is one level up: a survey of the action ansätze tried so far and of what a joint action for Ψ, connection, scalar and coframe must at least achieve.

Proof and measurement sources. r80.1 by `k168_fierz.py` and the note `Erkundung_Fierz_Hintergrund.md`; r80.2 by `k169_feldgleichungen.py` with output `k169_out.json`. The note `Vorab_Notiz_H68` and the result note lie in `notes/research/predictions/`. □

Consequence. The comparison system leaves no internal source of its Lorentz breaking: the breaking comes from a background that the system does not generate. The branch is closed with this layer. OP5 is not closed; the open problem is moved to the level of a joint dynamics, which OP1–OP5 share.

v4.9.0-r80 status in the three frames

- All statements concern the fixed OP5 comparison model in one chart, with one coframe reconstructed from Ψ. No frame map M, T or P acts on them.
- The colour labels λ_b of r80.2 are the internal 3×3 indices of the comparison model, read in its single fixed frame. They are not identified with $\mathrm{SU}(3)_c$ of §A.272.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r80 register entries

statement	status after v4.9.0-r80
spinor blocks at the germ	regular Dirac sector, Hubble-like counterflow, spins along $\hat{\varphi}$ (exploration)
background A, h	not sourced by the blocks via Yang–Mills on ₹ or Klein–Gordon (H68 refuted)
comparison-model branch	closed (author decision); results r71–r79 remain valid as local statements
next step	survey of action ansätze; joint dynamics

v4.9.0-r80 ledger and guard update

$$R_{\mathrm{chk}}^{v490\text{-}r80} = (1, 0, 0)$$

for (H68 decided; background not native; OP5 not closed).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r80,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r80} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r81 Source-Only: A Joint Action W1, the Matter–Tachyon Pairing, and the Search for a Vacuum Term

Purpose. Layer v4.9.0-r80 closed the comparison-model branch and moved the open question to the level of a joint dynamics (§A.272). On 4 and 5 November 2026 the author took five decisions on such an action, and the work that followed is recorded here. The layer contains the first action W1 and its identities, four pre-registered tests (H69, H70, H71, H72) and one refuted test (H73), a series of explorations on the matter–tachyon pairing, the vacuum term, the Higgs field as a tachyonic field and the connectors between matter and its tachyon image, and three corrections of earlier statements. The research notes are `notes/research/v4_10_0_w1/` (merged as PR #65). Earlier layers are not rewritten.
Source-only status. This v4.9.0-r81 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r81 decisions and the action W1

Author decisions (4–5 November 2026).

1. The coframe is an independent field; whether solutions satisfy $E \approx C[\Psi]$ is to be tested later, not assumed.
2. A real action with Dirac adjoint and symmetric kinetic term in frame M is the main line; the complex reading K (stationarity of $\mathrm{Re}\, S$ and $\mathrm{Im}\, S$) is computed alongside.
3. Einstein–Cartan gravity without a cosmological term, one system for now.
4. The Dixon carrier $\mathbb{C} \otimes \mathbb{H} \otimes \mathbb{O}$ with the $U(1)$ generated by $J = iL_{e_7}$, extended later by colour and the weak $SU(2)$; bookkeeping in QAM.
5. The closure projector $\hat{P}_{C_7}$ is left open.

Theoremlet v4.9.0-r81.1 (the action W1). Exploration. W1 is $S = \int e\,[\frac{1}{2\kappa}R(e,\omega) + \frac{1}{2}(\Psi^\dagger i\bar{\sigma}^\mu D_\mu \Psi + \text{c.c.}) - \frac{1}{4}F^2]$ on the sixteen left-handed Weyl states of one generation with ν_R (§A.272), Lorentz acting by left multiplication on $\mathbb{H}$.

(a) $U(1)_J$ is anomaly-free on the full carrier: $\mathrm{Tr}\, q = \mathrm{Tr}\, q^3 = 0$ for J, iR_{e_7} and $B - L$. On the image of C_{ch} alone $B - L$ would be anomalous ($\mathrm{Tr}\, q^3 = -1.78$).

(b) Eliminating the torsion gives $\mathcal{L}_{\mathrm{eff}} = -\frac{3}{16}\kappa\, j \cdot j$ with the total current of all sixteen states; the control reproduces the Hehl–Datta value for a Dirac field. The torsion is blind to the $U(1)$ charge.

(c) W1 produces no vacuum term: without a scalar potential no term proportional to $g_{\mu\nu}$ with constant coefficient arises.

v4.9.0-r81 the complex reading and the matter–tachyon pairing

Theoremlet v4.9.0-r81.2 (complex reading). Exploration, pointwise identities checked to 2×10^{-14}. With the symmetric kinetic term $\mathrm{Im}\, S \equiv 0$: K adds nothing. Without symmetrization $\mathrm{Im}\, L = \frac{1}{2}D_\mu j^\mu$; with an independent connection, stationarity of $\mathrm{Im}\, S$ forces $j^a = 0$, hence $\Psi = 0$ in a single frame, since $j^0 = |\psi|^2 \geq 0$. On the image of C_{ch} the inner and outer units agree up to sign.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Theoremlet v4.9.0-r81.3 (the matter–tachyon gap). Exploration. With the tachyon frame of Convention F as the continuation $\eta \to \eta + i\pi/2$ along an axis n, the vector map is $\Lambda(i\pi) = -R_n(\pi)$, so a tachyon component carries the current $j_T = \Lambda(i\pi) j_M$ of negative density. Then $j_M + j_T = 2j_\perp$: K only requires that matter flows along the axis, and the remaining equations from $\operatorname{Im} S$ hold. For frames offset by δ (author: frames need not be synchronous) the transverse current is antiperiodic with period 2δ.

Theoremlet v4.9.0-r81.4 (H69 confirmed). Pre-registered (commit `dbcc40d`). Flat space, $A = 0$: two counter-propagating left-handed waves plus their tachyon image offset by δ, with kinetic sign $s_T = -1$ from F3. The current balance holds exactly iff $\Delta\omega\,\delta \equiv \pi \pmod{2\pi}$ (residual 5×10^{-15}, violated at 0 and $\pi/2$), and the total energy–momentum vanishes including cross terms (10^{-15}). The pair carries neither net current nor net energy; the offset fixes a beat frequency.

Theoremlet v4.9.0-r81.5 (H70: the offset must be comoving). Pre-registered (commit `54c354a`, not blind). On flat FRW with massless Weyl fields, of the couplings $\delta = c/H$, a fixed conformal offset, and $\delta = t - \sqrt{t^2 - \beta_{\text{sym}}^2}$ from Theorem 22.9, only the fixed conformal offset ($\delta \propto a$ in cosmic time) keeps the pair consistent (H70a confirmed). The other two violate the Weyl equation of the tachyon image; the third is undefined for $t \le \beta_{\text{sym}}$. H70b is refuted: wrong couplings leave a spatially homogeneous residual, which is not Λ-like (only $\tau\tau$ and τz components) and comes from the violated field equation.

Theoremlet v4.9.0-r81.6 (frame weights and the frame check). Explorations.

(a) The neutral pairing has equal weights, the photon-frame mixture P. With matter-frame weights a homogeneous, traceless, radiation-like residual $\propto p_+ - p_- = 1/\gamma$ remains. Massless fields never give a Λ-like term.

(b) On the doublet, the tachyon map $R = \sigma_y$ and time reversal $T = \sigma_z$ give the same image on the compensator axis, where the matter frame lies; both exchange the port weights. They differ off the axis and on the carrier, where T conjugates the charges and R fixes the sectors (§A.272).

v4.9.0-r81 mass, tachyonic modes and the vacuum term

Theoremlet v4.9.0-r81.7 (H71: the massive partner). Pre-registered (commit `4a3b802`; (a), (b) not blind). For $L = \psi^\dagger i\bar\sigma\partial\psi + s\,\chi^\dagger i\sigma\partial\chi - m(\psi^\dagger\chi + \text{h.c.})$: (a) $s = +1$ gives $\omega^2 = k^2 + m^2$, $s = -1$ gives the tachyonic $\omega^2 = k^2 - m^2$ with a spacelike current; (b) for $s = +1$ the balance fails; (c) for $s = -1$ it fails as well (refuted). An antiparticle and a mass-coupled mirror partner are classically the same Dirac partner; an uncoupled mirror sector (the author's "mirror particle") adds positive, dark-matter-like energy.

Theoremlet v4.9.0-r81.8 (H72 and H73: a vacuum term from tachyonic modes). For a plane mode $T_{\mu\nu} = N\,p_\mu p_\nu$; an isotropic incoherent gas has $w = \sum Nk^2/(3\sum N\omega^2)$.

(a) H72 (commit `bc85519`): a Λ-like term $T = \rho\eta$ arises from tachyonic modes of both norm signs on at least two momentum shells (residual 3×10^{-16}); its height and sign are set by a free weight ratio; frame weights on one shell give none.

(b) H73 (commit `623a080`) refuted: the reference states (matter at rest, tachyon at threshold) with matter-frame weights give $w = -\beta_{\text{sym}}^2/3 = -0.10$.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(c) With modes and weights at a common rapidity the mixture is exactly pressureless dust in frame M ($\rho = m/\cosh\eta$) and pure tension with $\rho = 0$ in frame T, for every η. If gravity sees $a\,T_M + b\,T_T$, then $w = -b/(3a)$ and $\Omega_\Lambda = b/(3a)$: frame M alone gives no dark energy, equal weights give $\Omega_m = \frac{2}{3}$, $\Omega_\Lambda = \frac{1}{3}$, the exchange of the observed fractions, and frame weights give $-\beta^2_{\mathrm{sym}}/3$ again. None of six justified weightings gives the observed $\Omega_\Lambda \approx 0.685$; "two of three sectors" gives 0.667 but was proposed after seeing the target and counts as a fit.

Reading. The structure supplies the mechanism of a vacuum term (tension without energy from the tachyonic part), not its size.

v4.9.0-r81 the Higgs field as a tachyonic field and the connectors

Theoremlet v4.9.0-r81.9 (connectors between matter and its tachyon image). Explorations.

(a) The only gauge-invariant mass term of the carrier, the ν^c Majorana connector (r44.1(d)), is never tachyonic: $\omega^2 = k^2 + |M|^2$ for both kinetic signs and all phases.

(b) Sources: the Higgs field is read as tachyonic at the origin (r06.4(b)) and as an observer-frame image (Remark 36.2); the connectors with Higgs quantum numbers are r44.1(b).

(c) For $p_M = m(\cosh\eta, \sinh\eta)$ and the tachyon image $p_T = m(\sinh\eta, \pm\cosh\eta)$ one has $p_M \cdot p_T = 0$ for the co-moving image, for every η. The connector momenta $p_M \pm p_T$ give: co-moving, lightlike; counter-moving, the pair connector $q^2 = +2m^2\sinh 2\eta$ and the transition connector $q^2 = -2m^2\sinh 2\eta$, tachyonic. A Higgs-like negative mass squared thus follows kinematically for counter-moving matter and tachyon image.

(d) The η-independent Σ-mixtures of the four connectors have $q^2 \in \{0, 4, 8, 16\}\,m^2$, from $q^2 = (n^2 - k^2)m^2$; the bit assignment of the masks only renames them. No string-like fourth level arises; the frame continuation is cyclic of order four.

v4.9.0-r81 corrections

1. *To r44, Reading 1.* "Mass is a coupling between matter frame and tachyon frame" uses the time reversal T. By r81.6(b) the reading is half right: mass terms couple a state to a partner with tachyon-frame weights that is also charge conjugate. The frame correction list of r51 (Reading 3) omits r44 and is hereby extended to it.

2. *To the note on r81.8(c).* The value $w = -\beta^2_{\mathrm{sym}}/3$ of H73 is not an artefact: it reappears exactly when the two readings of the consistent mixture are weighted with the frame weights.

3. *To the draft of W1.* Ψ is the whole carrier, not the image of C_{ch}; C_{ch} is the compensator (the chirality), not $\hat{P}_{C_7}$, which is distinct by the section "Testing the Native Operator Link" of the 4.10 research candidate. The condition $\mathrm{Tr}(\mathcal{M}_{16}\hat{P}_{C_7}) \equiv 0 \pmod{\mathrm{HC}}$ is not defined as a computation anywhere in the sources.

Proof and measurement sources. Scripts `k170`–`k187` and the notes in `notes/research/v4_10_0_w1/`; pre-registrations and result notes H69–H73 in `notes/research/predictions/`. Deferred ideas with triggers (several loop systems joined by a string, the Chinese remainder reading of "mod HC", the dark-energy reading of OP5, condensation of tachyonic modes) are listed in `Zurueckgestellte_Ideen.md` there. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Consequence. A joint action with independent coframe exists in the form W1 and is consistent with the Dixon carrier. The complex reading forces a current balance between matter and its tachyon image, which a comoving offset pair satisfies exactly. A vacuum term needs tachyonic modes of both norm signs; its size is not fixed by the structure. OP5 is not closed.

v4.9.0-r81 status in the three frames

- W1 is written in frame M. The tachyon frame T enters only through the declared map $\eta \to \eta + i\pi/2$ and the sign F3; the photon frame P is the equal-weight mixture. All comparisons between M and T are made after this map.
- r81.6(b) separates the tachyon map R from time reversal T; results stated for one hold for the other only on the compensator axis.
- Colour and charge labels are those of the carrier in frame M; they are frame-free between M and T (§A.272).

v4.9.0-r81 register entries

statement	status after v4.9.0-r81
joint action W1	set up; anomaly-free $U(1)_J$; torsion $-\frac{3}{16}\kappa j^2$; no vacuum term
offset matter–tachyon pair	neutral in current and energy (H69); offset comoving (H70)
massive tachyon partner	tachyonic dispersion (H71); no current balance with mass
vacuum term	possible from tachyonic modes of both norms (H72); size not fixed; H73 refuted
Higgs-like tachyonic connector	follows kinematically for counter-moving matter and tachyon image (exploration)
r44 reading 1	corrected (time reversal, not tachyon map)

v4.9.0-r81 ledger and guard update

$$R_{\mathrm{chk}}^{v490\text{-}r81} = (1, 0, 0)$$

for (joint action set up and tested in part; vacuum term not fixed; OP5 not closed).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r81,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r81} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r82 Source-Only: The Higgs Sector of W2, the Stability of the Matter–Tachyon Cancellation, and a Torsion Pairing Channel

Purpose. Layer v4.9.0-r81 set up the joint action W1 and found that a vacuum term needs tachyonic modes and that its size is not fixed (§A.272). On 5 November 2026 the author asked to continue with the Higgs sector. This layer adds it (action W2), records the bookkeeping of the vacuum energy in the matter–tachyon pairing, two pre-registered stability tests (H74, H75, both not blind), and an exploration of the pairing channel of the torsion interaction of W1. The notes are `notes/research/v4_10_0_r82/`. Earlier layers are not rewritten.
Source-only status. This v4.9.0-r82 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r82 results

Theoremlet v4.9.0-r82.1 (W2 and its vacuum energy). Exploration, analytic. W2 is W1 plus a Higgs doublet with $V = -\mu^2|\Phi|^2 + \lambda|\Phi|^4$ and Yukawa couplings through the connectors of r44.1(b); the sign of $-\mu^2$ is read as the tachyonic transition connector of r81.9(c), not derived.

(a) Alone, W2 has the vacuum energy $V_{\min} = -\mu^4/(4\lambda)$, about $(100\,\mathrm{GeV})^4$, the known excess of about 10^{55}.

(b) With the tachyon image of every field and the sign F3, the whole energy–momentum tensor of the image changes sign, including its vacuum energy. With weights a (M) and b (T) the vacuum energy is $(a-b)V_{\min}$: exactly zero at photon weights for every μ, λ; $V_{\min}/\gamma$ at matter weights. This is the structure of the energy parity of Kaplan and Sundrum (2005) and the reading "$\Lambda = 0$ in the photon frame" of Theorem 22.16's status note.

(c) A time offset between the condensates of M and T leaves the cancellation of a formed condensate exact; a residual arises only transiently during the transitions.

Theoremlet v4.9.0-r82.2 (H74: direct coupling destroys the exact cancellation). Pre-registered (commit `b24bb08`, not blind). A normal oscillator coupled linearly ($\varepsilon\phi\chi$) to one with the sign F3 has $\Omega^2 = \frac{1}{2}(\omega_1^2 + \omega_2^2) \pm \sqrt{(\omega_1^2 - \omega_2^2)^2/4 - \varepsilon^2}$.

(a) Equal spectra, as required for exact cancellation, grow for every $\varepsilon > 0$ with rate $\varepsilon/(2\omega)$ (measured 5.011×10^{-4}, 4.985×10^{-3}, 0.0498 against 5×10^{-4}, 5×10^{-3}, 0.05).

(b) Detuned spectra are bounded for $\varepsilon < |\omega_1^2 - \omega_2^2|/2$ and grow above (rate 0.01594 against 0.01618).

(c) Two normal oscillators stay bounded (control).

Run 1 with the pre-registered time $T = 400$ left three small-coupling cases undecided (planning error: growth by 10^3 needs $T \sim 14/\varepsilon$); run 2 with adapted time confirms them. *Reading.* Exact cancellation and stability exclude each other under direct coupling; stability needs detuning, which leaves a residual $\gtrsim 2\varepsilon|V_{\min}|/\omega^2$.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Theoremlet v4.9.0-r82.3 (H75: gravity as the only coupling). Pre-registered (commit `6aa8477`, not blind). Newtonian two-fluid perturbations, matter with density ρ_A and the tachyon sector with negative inertial and gravitational mass $-\rho_B$; both fall alike, $\ddot{x} = 4\pi G N x$, $N = \left(\begin{smallmatrix} \rho_A & -\rho_B \\ \rho_A & -\rho_B \end{smallmatrix}\right)$.

(a) At photon weights $N^2 = 0$: no exponential growth; the difference of the sectors stays constant and the common mode grows as t^2 (exactly $\delta_0(1+t^2)$). The pre-registered bound "at most linear" was wrong for the common mode.

(b) At matter weights one Jeans mode with rate $\sqrt{4\pi G\rho_A(1-\beta_{\mathrm{sym}}^2)}$ (exact); two normal fluids give $\sqrt{4\pi G\rho_A(1+\beta_{\mathrm{sym}}^2)}$.

(c) A pair $\pm m$ accelerates jointly at constant distance and constant acceleration with conserved energy and momentum (Bondi's runaway), confirmed exactly.

Reading. Gravity as the only coupling has no resonance; the price is a joint runaway. The quantum decay of the vacuum into pairs of positive and negative energy is not covered.

Theoremlet v4.9.0-r82.4 (a pairing channel of the torsion interaction). Exploration. The Fierz identity $\bar{\sigma}^{\mu}_{ab}\bar{\sigma}_{\mu\,cd} = 2\varepsilon_{ac}\varepsilon_{bd}$ gives $j_1 \cdot j_2 = 2|\Phi|^2$ with the Lorentz-scalar pair $\Phi = \psi_2^{\mathrm{T}}\varepsilon\psi_1$. For anticommuting fields the torsion interaction $-\frac{3}{16}\kappa\, j \cdot j$ of r81.1(b) is attractive in this channel. Its strength $\frac{3}{4}\kappa = 6\pi/M_P^2$ reaches the critical value of a condensate only for a cutoff $\gtrsim 1.5\, M_P$. The only gauge-invariant channel is $\nu^c\nu^c$ (r44.1(d)); a condensate there would give a Majorana mass near M_P and, through the seesaw, $m_\nu \sim 10^{-6}\,\mathrm{eV}$. The electroweak Higgs is not obtained this way.

Proof and measurement sources. Scripts `k188`, `k188b`, `k189`, `k190` and the notes in `notes/research/v4_10_0_r82/`; pre-registrations and results H74, H75 in `notes/research/predictions/`. □

Consequence. The structure supplies mechanisms (exact cancellation of the vacuum energy at photon weights, stability only for weak, gravitational coupling, an attractive pairing channel), but no scale: the only scale W1 and W2 bring with them is the Planck scale. OP5 is not closed.

v4.9.0-r82 status in the three frames

- W2 is written in frame M; the tachyon sector enters through the declared continuation and the sign F3, the photon frame as the equal-weight mixture. Vacuum energies of M and T are compared only after this map.
- r82.3 treats the tachyon sector as negative inertial and gravitational mass in frame M's Newtonian limit; this is the F3 reading, not a frame-free statement.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r82 register entries

statement	status after v4.9.0-r82
Higgs vacuum energy in the M–T pairing	cancels exactly at photon weights; $V_{\min}/\gamma$ at matter weights
exact cancellation under direct coupling	unstable (H74)
exact cancellation, gravity only	no exponential instability, joint runaway (H75)
torsion pairing channel	attractive, Planck scale; ν^c Majorana channel only gauge-invariant one

v4.9.0-r82 ledger and guard update

$$R_{\text{chk}}^{v490\text{-}r82} = (1, 0, 0)$$

for (stability of the cancellation decided in the classical linear and Newtonian cases; scale not fixed; OP5 not closed).

The layer spends no guard:

$$R_{\text{glob}}^{(v490\text{-}r82,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\text{guard}}^{v490\text{-}r82} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r83 Source-Only: Quantum Decay of the Matter–Tachyon Vacuum, the Tachyon Image of an Action, and a Correction to r82

Purpose. Layer v4.9.0-r82 left the quantum decay of the vacuum of the matter–tachyon pairing open (§A.272). On 5 November 2026 the author chose to check it first. The check (H76) needed a preferred frame, and the search for one exposed that the tachyon image of an *action* had not been fixed. The author asked to settle this before choosing. This layer records both pre-registered tests (H76, H77, both not blind), an exploration of ghost condensation as the frame-defining mechanism, and a correction to r82.1(b), (c). The notes are `notes/research/v4_10_0_r83/`. Earlier layers are not rewritten.
Source-only status. This v4.9.0-r83 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r83 results

Theoremlet v4.9.0-r83.1 (H76: gravitational decay of the vacuum). Pre-registered (commit `2b47e7f`, not blind; the scaling is known from Cline, Jeon and Moore, hep-ph/0311312). One massless scalar in M (positive energy) and one in T (negative energy), coupled only by a graviton, $\kappa^2 = 32\pi/M_P^2$; the process is $0 \to \varphi\varphi\,\chi\chi$.

(a) With only a Lorentz-invariant cutoff $P^2 < \Lambda^2$ the decay is allowed and its rate is unbounded: with an additional frame cutoff $L\Lambda$ the ratios $\Gamma(L)/\Gamma(L/2)$ are $7.06, 4.77, 4.23, 4.07, 4.02, 4.006, 4.002$ for $L = 2, \ldots, 128$. The rate grows as L^2, the volume of boosts on the hyperboloid.

(b) With a cutoff $|k| < \Lambda$ in a preferred frame, $\Gamma/V = c\,\Lambda^8/M_P^4$ with exponent 8.000 and 7.997 and $c = 2.25 \times 10^{-5}$ (relative error 0.12%).

(c) Transferred to the gamma-ray bound of Cline et al. (their $c \sim 1$, $\Lambda \lesssim 3$ MeV) the bound moves by $c^{-1/7} = 4.6$ to $\Lambda \lesssim 14$ MeV (rough). A T sector that is a Lorentz-invariant copy up to the Higgs scale and couples only gravitationally contradicts the gamma-ray background.

Run 1 (direct Monte Carlo) was precise for (b) but not for (a) at large L; run 2 with ten times more samples did not converge (heavy-tailed weights). Run 3 computes the same quantity by Lorentz invariance (configuration in the rest frame of the matter pair, exact boost volume) and agrees with run 1 at $L = 1, 2, 4$ to better than 1%. The criterion was not changed; the method was changed after seeing the data.

Theoremlet v4.9.0-r83.2 (the condensate of I4 defines no frame). Exploration. A condensate $\langle\chi\rangle = \text{const}$ of tachyonic modes is Lorentz invariant. It selects no frame, and the decay of r83.1 persists. It can supply a scale, not the frame.

Theoremlet v4.9.0-r83.3 (H77: the tachyon image of an action). Pre-registered (commit `d087d67`, not blind). Three readings for a scalar with $L_M = \frac{1}{2}(\chi_t^2 - \chi_x^2) - \frac{1}{2}m^2\chi^2$: A, $L_T = -L_M$ (energy parity, used in r82.1(b)); B, the kinematics continued ($X \mapsto -X$, constants and potentials kept), as the rapidity $\eta + i\pi/2$ of r20.2 (v4.8.0) gives $p \mapsto ip$; B′,

© 2026 Marc Brendecke. Preprint — not peer reviewed.

$m^2 \mapsto -m^2$ with real kinematics. Criteria: K1 tachyonic dispersion (r07.2, r20.2, H71); K2 massless $L_T = -L_M$ (r81); K3 a running mode carries energy of the opposite sign (F3 on readouts).

reading	dispersion	K1	K2	K3
A	$\omega^2 = k^2 + m^2$	no	yes	yes
B	$\omega^2 = k^2 - m^2$	yes	yes	yes
B′	$\omega^2 = k^2 - m^2$	yes	no	no

Only reading B is compatible. Reading A gives a ghost copy with normal dispersion, not a tachyon in the sense of r20. Applying F3 by dimension to every quantity is a complex change of units and changes nothing; a reading has to state what is continued, and B continues the kinematics.

Theoremlet v4.9.0-r83.4 (ghost condensation under reading B). Exploration (sympy). A healthy matter side $P_M(X) = X/2 + cX^2$, $c > 0$, becomes under B $P_T(X) = -X/2 + cX^2$, the form of ghost condensation (Arkani-Hamed, Cheng, Luty, Mukohyama, hep-th/0312099): minimum at $X_0 = 1/(4c) > 0$, $P_{XX} = 2c > 0$, $\rho = 1/(16c)$, $w = -1$. The condensate $\langle\dot\chi\rangle \neq 0$ selects a frame (candidate: the comoving frame of r81.5, where the M–T offset is fixed in conformal time), and its fluctuations carry positive energy, so the decay of r83.1 does not occur. By the literature (Furukawa et al., 1001.4634) the fluctuations behave like dark matter, with $10\,\mathrm{eV} \lesssim M \lesssim 100$ GeV. Under A the only extremum lies at $X < 0$ with $P_{XX} < 0$. The term cX^2 is not in W1 or W2 and is an input; the height of the Λ-like term is not fixed.

Proof and measurement sources. Scripts `k191`, `k191b`, `k191c`, `k192`, `k192b` and the notes in `notes/research/v4_10_0_r83/`; pre-registrations and results H76, H77 in `notes/research/predictions/`. □

v4.9.0-r83 correction to r82

Correction to v4.9.0-r82.1(b), (c). r82.1(b) took the tachyon image of the Higgs sector in reading A: the whole energy–momentum tensor of the image changes sign, including its vacuum energy, so that the vacuum energy is $(a-b)V_{\min}$ and vanishes at photon weights. By r83.3 the image is to be formed in reading B. Then a potential without derivatives keeps its sign, and the homogeneous Higgs image obeys $\ddot\phi = +V'(\phi)$: around $\phi = 0$ it has $\ddot\phi = -2\mu^2\phi$ and is stable (script `k192b`). The tachyonic mass of the Higgs in M is a normal mass in the image, the image does not condense, and the vacuum energy is $a\,V_{\min}$, not $(a-b)V_{\min}$. **The exact cancellation of r82.1(b) and the transient residual of r82.1(c) are withdrawn.** r82.1(a), r82.2 (H74), r82.3 (H75) and r82.4 stand: under B a running T mode is per mode the negative-energy oscillator of H74, and moving T matter has the negative inertial and gravitational mass of H75. The register line "Higgs vacuum energy in the M–T pairing: cancels exactly at photon weights" of r82 is superseded by the register below.

Consequence. The structure now offers a frame and stability (ghost condensation under B), but the mechanism against the excess of about 10^{55} is gone. A scale other than the Planck scale appears only as an upper bound from stability ($\lesssim 14$ MeV from data, $\lesssim 2$ meV by the naturalness argument of Kaplan and Sundrum, hep-th/0505265). OP5 is not closed.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r83 status in the three frames

- r83.3 fixes how frame T enters an action: by the rapidity map of r20 on the kinematics, with constants and potentials kept. The photon frame remains the equal-weight mixture; nothing in this layer is evaluated there.
- r83.1 and r83.4 work in frame M with the T sector entering through reading B; the preferred frame of r83.1(b) and r83.4 is a frame of the cosmological background, not one of the frames M, T, P. The bound of v4.9.0-r09 (Reading V) on a direct coupling of the observer's rapidity to such a frame applies.

v4.9.0-r83 register entries

statement	status after v4.9.0-r83
quantum decay of the M–T vacuum	unbounded without a preferred frame; with one, $c\,\Lambda^8/M_P^4$, $\Lambda \lesssim 14$ MeV (H76)
tachyon image of an action	reading B (kinematics continued); A excluded (H77)
Higgs vacuum energy in the M–T pairing	no cancellation; $a\,V_{\min}$ (correction to r82.1)
ghost condensation under B	stable, frame-defining, $w = -1$; cX^2 input, height free

v4.9.0-r83 ledger and guard update

$$R_{\mathrm{chk}}^{v490\text{-}r83} = (1, 1, 0)$$

for (quantum decay decided for the gravitational channel; reading of the tachyon image decided; scale not fixed).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r83,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r83} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r84 Source-Only: The Tachyon Part as a Second Reading of the Same Field, the Complete Complex Boost, and the Expanding Background

Purpose. Layer v4.9.0-r83 showed that the vacuum of the matter–tachyon pairing decays through gravity unless a frame is preferred, and that ghost condensation under reading B supplies a frame (§A.272). On 5 November 2026 the author was asked whether the tachyon part T is an independent field or a second reading of the same field, a question open since k174. He chose the second reading. This layer records the consequences: an exploration of ghost condensation on an expanding background, four pre-registered tests (H78–H81, all not blind), and two author decisions. The notes are `notes/research/v4_10_0_r84/`. Earlier layers are not rewritten.
Source-only status. This v4.9.0-r84 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r84 results

Theoremlet v4.9.0-r84.1 (ghost condensation on FRW). Exploration (`k193`). For $P(X) = -X/2 + cX^2$ the quantity $a^3 P_X \sqrt{X} = C$ is conserved, and near $X_0 = 1/(4c)$ $\rho = 1/(16c) + C/(a^3\sqrt{c})$, $p = -1/(16c)$: a Λ-like constant plus dust. Both the height and the dust share C are free; $C = 0$ gives pure Λ. The condensate cures only the condensing field: under reading B the images of massless fields keep negative energy, and r83.1 applies to them if T is an independent field.

Author decision (5 November 2026). The tachyon part is the same field in a second reading, not an independent field. Variant 1 (independent field with a frame cutoff) and the deferred idea I1 remain alternatives.

Theoremlet v4.9.0-r84.2 (H78: the second reading in the form of k174). Pre-registered (commit `ef3e1e0`, not blind). With $\phi_T(x) = \phi(x + \delta)$ and $\psi_T(x) = -i\sigma_z \psi(x + \delta)$ (spinor rotated, argument only shifted) and weights a, b in flat space:

(a) scalar: $aL_M + bL_T = (a - b)\frac{1}{2}X - (a + b)V$, independent of δ (probe 4×10^{-16}); stable for $a > b$, $m^2_{\text{eff}} = \gamma m^2$ at matter weights, no kinetic term at photon weights;

(b) Weyl: operator $(a - b)(\bar\sigma^t \partial_t + \bar\sigma^z \partial_z) + (a + b)(\bar\sigma^x \partial_x + \bar\sigma^y \partial_y)$, dispersion $\omega^2 = k_z^2 + q^2 k_\perp^2$, $q = (a + b)/(a - b) = \gamma = 1.8645$ at matter weights;

(c) no independent T quanta, hence no decay channel as in r83.1.

By (b) this form predicts a Lorentz violation of order one for fermions, excluded by laboratory bounds of 10^{-13}–10^{-15} on the electron (Altschul, arXiv:1005.2994). The first run of the δ probe used a test field that was not periodic on the grid (26%); the second run gives 4×10^{-16}. With the real action the offset δ drops out in flat space; the δ-dependence of H69 and H70 comes from the complex reading K.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Theoremlet v4.9.0-r84.3 (H79: the complete complex boost). Pre-registered (commit `701f1ca`, not blind). The map of k174 is the complex boost with rapidity $i\pi$: $S(\theta) = e^{-\theta\sigma_z/2}$ satisfies $S\bar{\sigma}^\mu S = \Lambda(\theta)^\mu{}_\nu \bar{\sigma}^\nu$, and $S(i\pi) = -i\sigma_z$, $\Lambda(i\pi) = \mathrm{diag}(-1,1,1,-1)$. A boost acts on the spinor index and on the argument; k174 used only the first. With $\psi_T(x) = S\psi(\Lambda x + \delta)$:

(a) holomorphic image of the adjoint ($\psi^\dagger \mapsto \psi^\dagger S$): the image term is $-L_M$ in all four directions, the action $(a-b)S_M$, isotropic, stable for $a > b$, the image carries opposite energy (criterion K3 of r83.3);

(b) Hermitian image ($\psi^\dagger \mapsto \psi^\dagger S^\dagger$): $+L_M$, action $(a+b)S_M$, isotropic, K3 violated;

(c) scalar: $\Lambda(i\pi)$ preserves the metric, as r84.2(a); probe with reflected and shifted argument 3×10^{-16}.

The reflection of time in the argument drops out of the action in flat space; the warning of r51 and r81 concerns the doublet, not this.

Author decision (5 November 2026). The image is formed by the complete complex boost, with the adjoint mapped holomorphically; in the matter reading the Dirac adjoint stays. In the author's words: "fix it like this; if it does not fit, we have to look for something else."

Theoremlet v4.9.0-r84.4 (the second reading on FRW). Exploration and pre-registered tests H80, H81 (commit `a5799ef`, not blind; script `k196`). In conformal time the image sees the scale factor $a(|\delta - \tau|)$. Scalars carry only even powers of a, so the continuation $a(-\tau) = \pm a(\tau)$ is irrelevant; conformal fields (massless fermions, gauge fields) see nothing. The reduced action of a scalar has kinetic weight $K = a\,a(\tau)^2 - b\,a_T^2$ and potential weight $P = a\,a(\tau)^4 + b\,a_T^4$.

(a) H80, $\delta = 0$ (reflection about the big bang): the mode equation is exactly $(a-b)$ times that of the matter scalar with $m^2 \mapsto \gamma m^2$, for radiation and matter and both continuations. No new dynamics; the image is the universe before the big bang, contracting as τ grows. Predictions could come only from the state at $\tau = 0$, as in the CPT-symmetric universe of Boyle, Finn and Turok (arXiv:1803.08928); the structural resemblance was noticed afterwards.

(b) H81a, $\delta \neq 0$: K vanishes where $a(\tau)/a(\delta - \tau) = \beta_{\mathrm{sym}}$, at $\tau^* = \beta_{\mathrm{sym}}\delta/(1+\beta_{\mathrm{sym}}) = 0.3546\,\delta$ (radiation) and $\sqrt{\beta}_{\mathrm{sym}}\,\delta/(1+\sqrt{\beta}_{\mathrm{sym}}) = 0.4257\,\delta$ (matter), because $b/a = \beta_{\mathrm{sym}}^2$.

(c) H81b: τ^* is a regular singular point with exponents $(0,0)$; generic solutions grow as $\ln|\tau - \tau^*|$ (slope -0.120, $R^2 = 1 - 4 \times 10^{-15}$), a single solution per k stays bounded ($\max|\chi|/|\chi_0| = 1.0007$). The passage through τ^* selects a mode. The choice $k = m = 1$ was not fixed in the pre-registration.

(d) H81c refuted quantitatively: before τ^*, $P/K < 0$ (tachyonic sign), but the growth of the mode $k = 0$, $m = 3$ is a factor 1.57, not above 10; the window is short for $m\delta = 3$.

Proof and measurement sources. Scripts `k193`–`k196` and the notes in `notes/research/v4_10_0_r84/`; pre-registrations and results H78–H81 in `notes/research/predictions/`. □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Consequence. As a second reading of the same field, with the complete complex boost and the holomorphic adjoint, the tachyon part is isotropic, stable for $a > b$ and free of the decay of r83.1. In flat space it only renormalises the kinetic term $(a - b)$ against the potential $(a + b)$. On an expanding background the reflection about the big bang adds nothing, while an offset produces an early epoch τ^* fixed by β_{sym} at which a mode is selected. The vacuum energy is not cancelled in any variant. OP5 is not closed. The deferred idea I1 (several systems joined by a string) had its first trigger hit by r84.2(b) (two light cones that cannot be merged); it was offered to the author.

v4.9.0-r84 status in the three frames

- Frame T enters as the image of frame M under the complete complex boost; the photon frame is the equal-weight mixture, where the kinetic weight $a-b$ vanishes and the reduced theory has no dynamics. This degeneracy is a statement at P in the sense of F2, not a global one.
- The weights $a = p_+$, $b = p_-$ are those of frame M; τ^* and $m_{\mathrm{eff}}^2 = \gamma m^2$ are matter-frame statements.

v4.9.0-r84 register entries

statement	status after v4.9.0-r84
T part	second reading of the same field (author decision)
image with spinor rotated only (k174 form)	anisotropic for fermions, excluded (H78)
complete complex boost, holomorphic adjoint	isotropic, stable, $(a - b)S_M$ (H79; author decision)
reflection about the big bang ($\delta = 0$)	no new dynamics (H80)
offset $\delta \neq 0$	epoch $\tau^* = \beta_{\mathrm{sym}}\delta/(1 + \beta_{\mathrm{sym}})$ (radiation), mode selection (H81)
ghost condensation on FRW	Λ plus dust, both free

v4.9.0-r84 ledger and guard update

$$R_{\mathrm{chk}}^{v490\text{-}r84} = (1, 1, 0)$$

for (form of the T part decided by the author and checked; behaviour on FRW decided for both centres; scale not fixed).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r84,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r84} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r85 Source-Only: Masses and Couplings in the Second Reading, the Epoch τ^*, and the Big Bang as a CPT Transition

Purpose. Layer v4.9.0-r84 fixed the tachyon part as a second reading of the same field, formed by the complete complex boost with the holomorphic adjoint (§A.272). On 5 November 2026 the author asked to check the mass terms, the weighting of gravity, and the two centres of the time reflection further. This layer records five pre-registered tests (H82–H86, all not blind), two explorations, one author decision (the domain G2) and one working assumption. The notes are `notes/research/v4_10_0_r85/`. Earlier layers are not rewritten.
Source-only status. This v4.9.0-r85 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r85 results

Theoremlet v4.9.0-r85.1 (H82: masses and couplings). Pre-registered (commit `c3f956c`, not blind). With $S = -i\sigma_z$, $S^{\mathsf{T}}\varepsilon S = \varepsilon$: Majorana and Dirac mass terms are invariant under the complete boost and enter with $P = a + b$; terms with derivatives enter with $K = a - b$. After canonical normalisation, at matter weights ($K = 1/\gamma$, $P = 1$): α, fermion and Majorana masses and G_N scale with γ; m_H and ℓ_P with $\sqrt{\gamma}$; v with $\gamma^{-1/2}$; m_W and θ_W are unchanged (a, confirmed). Compared with the weights of v4.9.0-r19, α, α_s and θ_W agree, G_N (1 against $k = 2$) and ℓ_P ($\frac{1}{2}$ against $\frac{3}{2}$) do not (b, refuted as expected). The different growth follows a fixed pattern given by the number of derivative terms; all factors can be absorbed into redefined parameters.

Theoremlet v4.9.0-r85.2 (H83: three rules for gauge fields and gravity). Pre-registered (commit `36ec8e7`, not blind). R1 as in r85.1; R2 computed from $g \mapsto -g$ with fixed fields (reading B for a scalar is exactly this), $R[-g] = -R[g]$, $\sqrt{|g|}$ unchanged; R3 a single action on $c\,g$. R2 gives Einstein–Cartan -1, Yang–Mills $+1$ (two inverse metrics), Weyl kinetic term $-i$ (one inverse vierbein). No rule gives $k(\alpha) = 1$ and $k(G_N) = 2$ together; G_N scales with γ^1 under all three. The match for α in r85.1 depends on R1.

Exploration v4.9.0-r85.3 (the phase of the spinors). Under R2 the fermion action is $(a - ib)S_{\text{Weyl}}$: in the real reading its tachyon image is invisible, in the complex reading K $\operatorname{Im} S = -b\,S_{\text{Weyl}}$ forces the energy–momentum tensor and the spin density of the fermions to vanish. Quantum mechanically $a - ib$ is a partial Wick rotation by $\arctan(b/a)$ (16.8° at matter weights, 45° at photon weights). $g \mapsto -g$ is not unique for spinors, since $Cl(1,3) \ncong Cl(3,1)$; F3 allows the phase $\mp i$ for a weight $\frac{1}{2}$. *Working assumption* (author, "carry on" on the proposal): the Weyl kinetic term has F3 weight 1, as in r81 and r84.3; not an explicit decision.

Theoremlet v4.9.0-r85.4 (H84: quantum modes at τ^*). Pre-registered (commit `d2a19f2`, not blind). For the offset $\delta \neq 0$ of r84.4 the Wronskian $K(\chi\bar{\chi}' - \bar{\chi}\chi') = i$ stays constant (4×10^{-14}); since $K(\tau^*) = 0$, every mode with this normalisation has $\chi' \propto 1/|\tau - \tau^*|$, and its energy density $E\,|\tau - \tau^*| \to -0.1496$ from below and $+0.1494$ from above. The mode selection of r84.4(c) is classical only; for a quantum field the offset is singular at τ^*, which favours the reflection about the big bang.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Author decision (5 November 2026): the domain G2. The reflection maps the domain as well: the image of $\tau > 0$ is $\tau < 0$. r84.4(a) (H80) took both readings on the whole time axis (domain G1). The author chose **G2**: the matter reading lives after the big bang, the tachyon reading on its image before it, joined only at the big bang. This is structurally the two-sheet picture of the CPT-symmetric universe of Boyle, Finn and Turok (arXiv:1803.08928). r84.4(a) stays valid for G1.

Exploration v4.9.0-r85.5 (the sharp transition under G2). For conformal fields the weight jumps from $-b$ to a at $\tau = 0$. With continuity of the field and of weight times derivative, the positive-norm mode before the big bang becomes $Ce^{-i\omega\tau} + De^{+i\omega\tau}$, $C = (a+b)/(2a)$, $D = (a-b)/(2a)$; normalised $\alpha = (1+x)/(2\sqrt{x})$, $\beta = (1-x)/(2\sqrt{x})$ with $x = \beta_{\mathrm{sym}}^2$, $\beta/\alpha = 1/\gamma = 0.536$, $n = (1-x)^2/(4x) = 0.404$ per mode for every k (sympy). At photon weights $n = 0$.

Theoremlet v4.9.0-r85.6 (H85, H86: the smooth transition). A real smooth weight from $-b$ to a has a zero and is singular by r85.4; the weight is therefore taken around zero in the complex plane, $w = b^{1-s}a^s e^{\pm i\pi(1-s)}$, a partial Wick rotation, with conserved holomorphic Wronskian.

(a) H85 (commit `4ac77a0`, partly not blind; $\Delta = 1$, tanh profile): both quantitative criteria missed. D approaches the sharp value only linearly in $k\Delta$ (3.5% at 0.01), and the pre-registration had exchanged the roles of C and D.

(b) Noticed afterwards: adiabatically the positive-norm mode of the tachyon reading continues as $e^{+ik\tau}$, which after the big bang is the mode of negative frequency; the admixture C falls off exponentially; the two senses of rotation distribute D unequally.

(c) H86 (commit `851297a`, not blind, motivated by (b); new settings $\Delta = 2$, tanh and error-function profiles): confirmed. For $k\Delta \geq 6$, $|C| < 10^{-2}$ and $\sqrt{|D_+||D_-|} = \sqrt{b/a}$ to 10^{-13}; $\ln|C|$ is linear on $k\Delta \in [2,8]$ ($R^2 = 0.99987$) with slope -2.67, within the 15% criterion around π only narrowly and not tending to π in the range; at $k\Delta = 0.01$, C meets the sharp value to 2×10^{-4}; at $k\Delta = 0.02$, $|D_-|/|D_+| = 1.13$ (tanh) and 1.10 (erf) with $\sqrt{|D_+||D_-|}/D_0 = 0.998$ and 0.999.

Proof and measurement sources. Scripts `k197`–`k201` and the notes in `notes/research/v4_10_0_r85/`; pre-registrations and results H82–H86 in `notes/research/predictions/`. □

Consequence. Mass terms carry over to the second reading; the pattern of factors matches v4.9.0-r19 for the gauge couplings but not for gravity. For a quantum field the reflection about the big bang is preferred over an offset. Under G2 with a complex transition the big bang maps every mode of the tachyon reading before it to a mode of negative frequency after it (a CPT assignment), with an admixture fixed by β_{sym} for long waves and cut off exponentially above $1/\Delta$; Δ is a free scale. The two senses of the Wick rotation give an asymmetry with fixed product; its physical meaning is open. OP5 is not closed.

v4.9.0-r85 status in the three frames

- Under G2 the matter frame M holds after the big bang and the tachyon frame T before it; the photon frame P is the equal-weight mixture, where the transition of r85.5 produces no particles ($n = 0$). This is a statement at P in the sense of F2.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

- The factors of r85.1 and the numbers of r85.4–r85.6 use the weights of frame M. The comparison with v4.9.0-r19 relates a reduction (M and T to one field) to a conversion (M to P); these are different operations.

v4.9.0-r85 register entries

statement	status after v4.9.0-r85
masses in the second reading	carry over; factors γ, $\sqrt{\gamma}$, 1 by derivative count (H82)
weights of v4.9.0-r19 from the reduction	gauge couplings yes, gravity no, under three rules (H82, H83)
F3 weight of the Weyl kinetic term	1 (working assumption); $\frac{1}{2}$ would hide or forbid fermions
offset $\delta \neq 0$, quantum	singular at τ^* (H84)
domain of the readings	G2: T before, M after the big bang (author decision)
big bang under G2	CPT assignment plus admixture fixed by β_{sym}, cut off above $1/\Delta$ (H86)

v4.9.0-r85 ledger and guard update

$$R_{\mathrm{chk}}^{v490\text{-}r85} = (1, 1, 0)$$

for (masses and the r19 comparison decided; the big-bang transition decided for G2; scale Δ not fixed).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r85,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r85} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r86 Source-Only: Independent Checks with a Second System: the Big-Bang Admixture and the Field Equations of W1

Purpose. Layers v4.9.0-r81 to r85 rest on Python and sympy scripts written in the same sessions (§A.272, §A.272). On 5 and 6 November 2026 a second, independent system (Wolfram Language through a connector) was used to recompute the big-bang transition of r85.6 and to derive the field equations of the joint action W1 from their definitions. This layer records three pre-registered tests (H87–H89, all not blind, since the expected values were known), the errors found in the checking code, and a correction to the reading of the energy–momentum tensor in the W1 notes. The notes are `notes/research/v4_10_0_r86/`. Earlier layers are not rewritten.
Source-only status. This v4.9.0-r86 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.9.0 Version DOI 10.5281/zenodo.23534954 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.9.0-r86 results

Theoremlet v4.9.0-r86.1 (H87: the admixture at the big bang, second system). Pre-registered (commit `61b7132`, not blind).

(a) Confirmed. Direct integration in Wolfram (working precision 30) reproduces the values of H86 (r85.6(c)) with relative deviations 2.6×10^{-13}, 7×10^{-11}, 5×10^{-9}, 5×10^{-7} at $k\Delta = 2, 4, 6, 8$; the least accurate value is the scipy value at $k\Delta = 8$.

(b) Refuted as formulated. For large $k\Delta$ the mode equation was integrated along $\tau = x + ih$ below the singularity of sech^2 at $i\pi\Delta/2$ ($h = 0.9$ and 0.95 times $\pi\Delta/2$); the boundary values at $\mathrm{Re}\,\tau \to \pm\infty$ are unchanged and the problem is well conditioned (checks at $k\Delta = 4, 8$ against the real axis to 3×10^{-10}; both lines agree to 1.5×10^{-8}). From $|C| = 4.23 \times 10^{-12}$ at $k\Delta = 10$ to 1.30×10^{-37} at 30 the local slopes are 2.925, 2.912, 2.936, 2.975; between 20 and 30 the slope is 2.956, 5.9% below π, outside the 3% criterion.

(c) Confirmed. The sharp transition of r85.5 symbolically: $C_0 = (a+b)/(2a)$, $D_0 = (a-b)/(2a)$, $\alpha = (1+x)/(2\sqrt{x})$, $\beta = (1-x)/(2\sqrt{x})$, $|\alpha|^2 - |\beta|^2 = 1$; numerically $\beta/\alpha = 0.53634$, $n = 0.40382$.

Noticed afterwards (not pre-registered). The local slope rises slowly on $k\Delta \in [15, 30]$. A fit $\ln|C| = c_0 + p\ln(k\Delta) - s\,k\Delta$ gives $s = 3.150$, $p \approx 4.8$ (exact through $20, 25, 30$) or $s = 3.077$, $p \approx 3.0$ (least squares on 15–30): compatible with $e^{-\pi k\Delta}$ times a large power prefactor. The position of the singularity fixes the exponent; the prefactor depends on the strength $\ln(a/b) - i\pi$ and is not perturbative (the first Born approximation is off by a factor 2.2 at $k\Delta = 8$). A confirmation needs a new pre-registered test with a fixed fit model.

Theoremlet v4.9.0-r86.2 (H88: the field equations of W1 from their definitions). Pre-registered (commit `06a8380`, not blind). All four parts confirmed, no deviation.

(a) Generator: with $\Sigma^{ab} = \frac{1}{4}(\sigma^a\bar\sigma^b - \sigma^b\bar\sigma^a)$, $\Sigma^{ab\dagger}\bar\sigma^c + \bar\sigma^c\Sigma^{ab} = -(\eta^{bc}\bar\sigma^a - \eta^{ac}\bar\sigma^b)$ in all 64 components, hence $D_\mu j^c = \partial_\mu j^c + \omega_\mu{}^c{}_b\, j^b$. The alternative ordering $\frac{1}{4}(\bar\sigma^a\sigma^b - \bar\sigma^b\sigma^a)$ is not covariant.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(b) Torsion: from $R^{ab}{}_{\mu\nu} = \partial\omega - \partial\omega + \omega\omega - \omega\omega$ and the symmetric spinor coupling, eliminating all 24 contorsion components for two Weyl spinors gives exactly $\mathcal{L}_{\text{eff}} = -\frac{3}{16}\kappa\, j \cdot j$ with the total current (r81.1(b)). For a single Weyl spinor $\mathcal{L}_{\text{eff}}$ vanishes, since its current is null.

(c) Weyl equation: the variation with respect to $\psi^\dagger$ is exactly $i\bar\sigma^\mu\partial_\mu\psi - qJA_\mu\bar\sigma^\mu\psi - \frac{3\kappa}{8} j_a\bar\sigma^a\psi = 0$.

(d) Ward identity: $M = qJA_\mu\bar\sigma^\mu + \frac{3\kappa}{8} j_a\bar\sigma^a$ is hermitian, hence $\partial_\mu j^\mu = 0$ on solutions.

Theoremlet v4.9.0-r86.3 (H89: the energy–momentum tensor of W1). Pre-registered (commit `6355d15`, not blind). All three parts confirmed.

(a) The vierbein variation of $e\,\mathcal{L}_{4f}$ at fixed spinors gives $T^{(4f)}_{\mu\nu} = -g_{\mu\nu}\mathcal{L}_{4f} = \frac{3\kappa}{16}\, j^2\, g_{\mu\nu}$: on its own the four-fermion part has the form of a Λ term ($p = -\rho$), with a height $\propto j^2$ that is not constant.

(b) Two Weyl fields of opposite spin, $\psi_s = e^{-i(3\kappa n/8)t}\chi_s$, solve the equation of r86.2(c) exactly with total current $(n, 0, 0, 0)$. On this solution $\mathcal{L}_{\text{kin}} = 3\kappa n^2/8$, $\mathcal{L} = 3\kappa n^2/16$, and the whole tensor $T_{\mu\nu} = T^{\text{Bel}}_{\mu\nu} - g_{\mu\nu}\mathcal{L}$ is $\rho = p = \frac{3\kappa}{16} n^2 > 0$: stiff matter.

(c) The vierbein variation to first order around flat space, with the spin connection solved from $de + \omega \wedge e = 0$ rather than inserted, is on a solution (two plane waves) exactly minus the Belinfante tensor (deviations 7×10^{-15} and 5×10^{-15} at values of size 30–90) and exactly symmetric. This justifies the form used in (b).

Disclosure. The first run of (c) exceeded the time limit of the connector and was reduced (spinors inserted as a solution, h, ∂h kept as symbols; no change of content). The second run had an error in the checking code (the conjugate transpose also conjugated the real symbols dh). The third run disagreed (non-zero antisymmetric part); following the pre-registered order the conventions were checked first, and the contraction $\omega_{\mu ab}\Sigma^{ab}$ had carried extra η factors. After the correction everything agrees. The disagreement lay in the checking code, not in W1.

Correction v4.9.0-r86.4 (reading of the W1 notes). The W1 notes of r81 (`llm/Wirkung_W1_Dixon_U1.md`) read the four-fermion influence as "$p = \rho$, $\rho \propto -\kappa n^2$, sign convention-dependent, not like Λ". After r86.3: $p = \rho$ is right for the whole tensor; the sign of ρ depends on the state, not on a convention, and is positive for modes of positive frequency; "not like Λ" holds for the whole tensor, not for the four-fermion term on its own, which cannot be separated physically from the kinetic part. r81.1(c) (W1 produces no vacuum term) is not affected: the homogeneous state is stiff matter, not Λ.

Proof and measurement sources. Wolfram code `k202_wolfram_gegenprobe.wl`, `k203_w1_feldgleichungen.wl`, `k204_w1_einstein.wl` and the notes in `notes/research/v4_10_0_r86/`; pre-registrations and results H87–H89 in `notes/research/predictions/`. □

Consequence. The numbers of r85.6 are confirmed by a second system; the admixture at the big bang falls off with a rate close to π per unit $k\Delta$, with a power prefactor, so the cut-off scale is about $1/(\pi\Delta)$ and Δ stays free. The field equations of W1 are derived independently up to the full non-linear Einstein equation and the Bianchi identity: generator, torsion, Weyl equation, Ward identity, and the energy–momentum tensor as the source to first order. The foundation of r81–r85, as far as it rests on these equations, is firmer. OP5 is not closed.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.9.0-r86 status in the three frames

- r86.1 checks the transition of r85.6 under G2 (frame T before, frame M after the big bang) with the weights of frame M; the photon frame P gives no admixture ($n = 0$), unchanged.
- r86.2 and r86.3 concern the matter part of W1 in frame M. The tachyon image under the second reading (r84) multiplies the action by the weights a, b and does not change the form of the equations; no frame-free statement is made beyond that.

v4.9.0-r86 register entries

statement	status after v4.9.0-r86
H86 values (r85.6(c))	confirmed by a second system (H87a)
decay of the admixture	rate near π with power prefactor (afterwards); 3% criterion missed (H87b)
W1 generator, torsion, Weyl, Ward	derived from the definitions (H88)
energy–momentum tensor of W1	minus Belinfante to first order; homogeneous state $\rho = p > 0$ (H89)
four-fermion term alone	Λ form with non-constant height; W1 vacuum term still absent
full Einstein equation, Bianchi	open

v4.9.0-r86 ledger and guard update

$$R_{\mathrm{chk}}^{v490\text{-}r86} = (1, 1, 0)$$

for (r85.6 confirmed independently; W1 field equations derived to first order; full Einstein equation open).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v490\text{-}r86,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v490\text{-}r86} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

Layers after the v4.10.0 release (Version DOI 10.5281/zenodo.23991357) are named v4.10.0-rNN and follow here.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.10.0-r01 Source-Only: The Full Einstein Equation of W1 and the Sign of the Gravitational Action

Purpose. Layer v4.9.0-r86 derived the field equations of the joint action W1 (§A.272) from their definitions up to the full non-linear Einstein equation and the Bianchi identity (§A.272). This first layer after the v4.10.0 release closes that gap with one pre-registered test (H90, not blind) and records what it found on the way: W1 as written has the wrong sign of the gravitational action. The author decided on 6 November 2026 to correct it. The layer states the corrected action and lists which earlier statements change sign. Earlier layers are not rewritten. The notes are `notes/research/v4_10_0_r01/`.
Source-only status. This v4.10.0-r01 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.10.0 Version DOI 10.5281/zenodo.23991357 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.10.0-r01 results

Theoremlet v4.10.0-r01.1 (H90: the Einstein–Cartan equations of W1, non-linear). Pre-registered (commit `e5d03d8`, not blind). Conventions as in r86: $\eta = \mathrm{diag}(1,-1,-1,-1)$, $D_\mu\psi = \partial_\mu\psi + \frac{1}{2}\omega_{\mu ab}\Sigma^{ab}\psi + iqJA_\mu\psi$, $R^{ab}{}_{\mu\nu} = \partial_\mu\omega_\nu{}^{ab} - \partial_\nu\omega_\mu{}^{ab} + \omega_\mu{}^a{}_c\,\omega_\nu{}^{cb} - \omega_\nu{}^a{}_c\,\omega_\mu{}^{cb}$, and the gravitational action written as $S_{\mathrm{grav}} = \frac{\sigma_g}{2\kappa}\int e\,R$ (W1 as written in r81.1: $\sigma_g = +1$). At a generic point of three random curved backgrounds (vierbein a random polynomial of degree three, not diagonal) all fields were carried as jets to third order: three Weyl fields with $J = (+1,-1,+1)$, a real potential A, the vierbein. The spinor jets solve the Weyl equation with gauge coupling and four-fermion term at the point and to first derivative, A the Maxwell equation at the point. Both signs $\sigma_g = \pm 1$ were computed.

(a) Cartan equation. The spin connection $\tilde{\omega}$ of the tetrad postulate solves the full ω-equation (with its derivative term) without spinors; with spinors $\omega = \tilde{\omega} + K$ solves it, with $K_{cab} = -\frac{\sigma_g}{4}\kappa\,\varepsilon_{cabd}\,j^d$ totally antisymmetric (residuals $\le 2\times 10^{-14}$).

(b) Einstein–Cartan equation. The variation with respect to the vierbein at fixed $\omega = \tilde{\omega} + K$ is symmetric on the field equations and equals a multiple of $\tilde{G}_{\mu\nu} + \sigma_g\kappa\,T^{\mathrm{eff}}_{\mu\nu}$ (residual 10^{-15}), with

$$T^{\mathrm{eff}}_{\mu\nu} = T^{\mathrm{Bel}}_{\mu\nu} - g_{\mu\nu}\mathcal{L}_W + T^{\mathrm{Maxwell}}_{\mu\nu} - g_{\mu\nu}\,\lambda\,j\cdot j, \qquad \lambda = -\sigma_g\,\tfrac{3\kappa}{16}.$$

Hence $\tilde{G} = -\sigma_g\,\kappa\,T^{\mathrm{eff}}$, fully non-linear. The K term of the Einstein–Cartan Weyl equation equals $2\lambda\,j_a\bar{\sigma}^a\psi$.

(c) Compatibility. $\tilde{\nabla}^\mu T^{\mathrm{eff}}_{\mu\nu} = 0$ on the field equations (relative 4×10^{-15}); $\tilde{\nabla}^\mu\tilde{G}_{\mu\nu} = 0$ as a control. Negative controls: without the four-fermion term the residuals rise to 0.12–0.19 (Einstein) and 0.25–0.81 (divergence); with spinors off the field equations to 0.16–0.19 and 0.92–1.05.

(d) Sign. For W1 as written ($\sigma_g = +1$) the factor is -1 to 4×10^{-15}: $\tilde{G} = -\kappa T^{\mathrm{eff}}$. A second system (Wolfram) gives $G_{00} = 2\nabla^2\Phi$ for $g = \mathrm{diag}(1+2\Phi, -(1-2\Phi)\text{⅄})$ in the same curvature convention, so $\tilde{G} = +\kappa T$ is attraction and $\tilde{G} = -\kappa T$ repulsion. W1 as written repels positive energy.

The Hehl–Datta control of r81.1(b) compared only the modulus 3/16; the sign had not been checked.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Decision v4.10.0-r01.2 (author, 6 November 2026: the corrected action). W1 is corrected to

$$S = \int e \left[-\tfrac{1}{2\kappa} R(e,\omega) + \tfrac{1}{2}\big(\Psi^\dagger i\bar\sigma^\mu D_\mu \Psi + \text{c.c.}\big) - \tfrac{1}{4}F^2 \right]$$

in the signature $(+,-,-,-)$ (as in Landau–Lifshitz). The field equations become $\tilde G = \kappa T^{\text{eff}}$ with $\mathcal{L}_{4f} = +\frac{3}{16}\kappa\, j\cdot j$, $T^{(4f)}_{\mu\nu} = -\frac{3\kappa}{16}\, j^2 g_{\mu\nu}$, Weyl term $+\frac{3\kappa}{8} j_a \bar\sigma^a \psi$, $K_{cab} = +\frac{\kappa}{4}\varepsilon_{cabd} j^d$.

Correction v4.10.0-r01.3 (statements that change sign). Moduli and forms stay; the signs change as follows.

(a) r81.1(b) and r86.2(b), (c): $\mathcal{L}_{\text{eff}} = +\frac{3}{16}\kappa\, j\cdot j$ and $+\frac{3\kappa}{8} j_a\bar\sigma^a\psi$ in the Weyl equation (were $-$); correct for the action as written, not for the physical one.

(b) r86.3(a), (b): $T^{(4f)} = -\frac{3\kappa}{16} j^2 g$ (Λ form, negative); the homogeneous state of two opposite spins has $\rho = p = -\frac{3\kappa}{16} n^2 < 0$, the negative spin energy density known from Einstein–Cartan theory.

(c) Correction r86.4 is withdrawn in its sign part: the original reading of the W1 notes, $\rho \propto -\kappa n^2$, holds for the physical sign. "Not like Λ" for the whole tensor and r81.1(c) (no vacuum term) stand.

(d) r82.4: the pairing channel of the torsion interaction is repulsive, $H_{\text{int}} \supset +\frac{3}{4}\kappa\, \Phi^\dagger\Phi$. No condensate arises through it; the ν^c Majorana mass near M_P and the seesaw estimate of r82.4 fall.

(e) Not affected (review in `llm/Durchsicht_Vorzeichen_kappa_r81_r86.md`): H69–H73 (source side only), H74 (no gravity), H75 (computed with Newtonian attraction), H76 (rate $\propto \kappa^4$), H77–H81 and H84–H86 (no gravitational coupling), H82 and H83 (relative weights and phases of the Einstein–Cartan term), H48 (Hehl–Datta sign taken from the literature).

Proof and measurement sources. Script `k205_w1_einstein_nichtlinear.py` (Python, jets, output `k205_out.json`) and Wolfram code `k205b_schwachfeld.wl`; notes in `notes/research/v4_10_0_r01/`; pre-registration and result H90 in `notes/research/predictions/`. □

Consequence. The field equations of W1 are now derived independently in full: Cartan equation, elimination of the torsion, Einstein–Cartan equation, its Riemannian form with the four-fermion term, and its compatibility with the Bianchi identity. With the corrected sign the four-fermion part has the form of a negative vacuum term with a height $\propto j^2$; the whole tensor of a homogeneous state is stiff matter with negative energy, not Λ. W1 still produces no vacuum term. OP5 is not closed.

v4.10.0-r01 status in the three frames

- r01.1–r01.3 concern the matter part of W1 in frame M. The tachyon image under the second reading (r84) multiplies the action by the weights a, b; the sign correction is common to both and does not change their ratio. The photon frame P is the degenerate fixed point; no frame-free statement is made beyond that.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

- r01.3(d) concerns the pairing of two left-handed fields in frame M; under the second reading the image term is $-\mathcal{L}_M$ (r84), so the interaction keeps its sign relative to the kinetic term.

v4.10.0-r01 register entries

statement	status after v4.10.0-r01
full Einstein equation of W1, Bianchi	derived and compatible (H90)
sign of the gravitational action	corrected to $-\frac{1}{2\kappa}\int eR$ (author decision)
four-fermion term	$+\frac{3}{16}\kappa\, j\cdot j$; $T^{(4f)}$ negative Λ form
homogeneous spin state	$\rho = p = -\frac{3\kappa}{16}n^2 < 0$
torsion pairing channel (r82.4)	repulsive; Planck-scale ν^c condensate withdrawn
W1 vacuum term	still absent

v4.10.0-r01 ledger and guard update

$$R_{\mathrm{chk}}^{v4100\text{-}r01} = (1,1,1)$$

for (Einstein–Cartan equations derived; compatibility shown; sign corrected by author decision).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v4100\text{-}r01,+)} = (0,1,1,1,1), \qquad \Gamma_{\mathrm{guard}}^{v4100\text{-}r01} = (1,1,1,1,1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.10.0-r02 Source-Only: Asymptotics of the Admixture at the Big Bang

Purpose. Under the domain choice G2 the big bang is a transition from the tachyon reading to the matter reading, and a mode of the tachyon reading picks up an admixture C of positive frequency after the bang that falls off for short waves (r85.6, §A.272). Layer v4.9.0-r86 found the fall-off close to $e^{-\pi k\Delta}$ with an unexplained power prefactor (§A.272). This layer records three pre-registered tests on that question (H91–H93). The last one derives the asymptotics in closed form. All three have formal defects in their validity conditions, which are disclosed. The notes are `notes/research/v4_10_0_r02/`. Earlier layers are not rewritten.
Source-only status. This v4.10.0-r02 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.10.0 Version DOI 10.5281/zenodo.23991357 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.10.0-r02 results

Throughout: $(w\chi')' + k^2 w\chi = 0$, $w = e^{Ls}$, $s = \frac{1}{2}(1+\tanh(t/\Delta))$, $L = \ln(a/b) \mp i\pi$ for the two senses of rotation, a, b the matter weights of r84, $\Delta = 2$; C is the coefficient of e^{-ikt} after the bang for e^{ikt} incoming before it.

Theoremlet v4.10.0-r02.1 (H91: the fall-off up to $k\Delta = 60$). Pre-registered (commit `a49ed7e`, partly not blind). Integration along $t = x + ih$ below the singularity; two contours agree to 10^{-10} up to $k\Delta = 60$.

(a) Invalid by its own criterion, which was a planning error: agreement with the H87 values to 10^{-5} was required, but H87 had recorded four digits only. All deviations, up to 3.2×10^{-4}, lie within that rounding. By the pre-registration parts (b)–(d) are then invalid. As formulated they would also have missed their criteria.

(b)–(d) A fit with $\sqrt{k\Delta}$ and $\ln k\Delta$ gives a slope 3.049, 3% below π. The $\sqrt{k\Delta}$ term improves the fit only by a factor 6.5. The slope between 50 and 60 is 3.021.

Noticed afterwards. The local slope is wavy and rises slowly (3.026 at 60). $|C|$ is the same for both senses of rotation, to 10^{-10}; the asymmetry of r85.6(d) concerns only the other coefficient.

Theoremlet v4.10.0-r02.2 (H92: the prefactor is the Stokes function of the local problem). Pre-registered (commit `c347a92`, partly not blind). Near $t_0 = i\pi\Delta/2$, $s \approx \frac{1}{2} + \Delta/(2\tau)$ with $\tau = t - t_0$, so w has an essential singularity. In $\sigma = k\tau$ the local equation is $y'' - (\lambda/\sigma^2)y' + y = 0$, with the single parameter $\lambda = Lk\Delta/2$. Its reflection $r(\lambda)$, continued below $\sigma = 0$, gives $C \approx r(\lambda)e^{2ikt_0} = r(\lambda)e^{-\pi k\Delta}$.

(b) Confirmed: $\ln|C| + \pi k\Delta - \ln|r|$ is smooth; the fit residual on $[20, 60]$ is 4.1×10^{-3}. The waves sit in r.

(c) Confirmed: $|r(\bar{\lambda})| = |r(\lambda)|$ to 8×10^{-11}.

(d) Confirmed: $r(\lambda)/(2\pi\lambda) = 1 - 1.6 \times 10^{-8}$ at $|\lambda| = 10^{-3}$, the Born limit.

(a) Missed: $\ln|C| + \pi k\Delta - \ln|r| \to \approx -0.62$, not 0.

Disclosure. In run 1 the accuracy control (a contour at $1.5h$) lost digits (3.7–5.5×10^{-6}, above 10^{-6}). In run 2 the control used $0.7h$ ($\leq 6 \times 10^{-10}$); the values of r are identical in both runs.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Theoremlet v4.10.0-r02.3 (H93: the asymptotics derived). Pre-registered with the derivation (commit `a4b1d17`, partly not blind).

(i) With $\chi = w^{-1/2}u$, $u'' + Qu = 0$ and w going from 1 to e^L: $C = e^{-L/2}C_u$. This is the factor missing in r02.2(a).

(ii) For large λ the local $Q = 1 - \lambda^2/(4\sigma^4) - \lambda/\sigma^3$ has four turning points. The over-barrier formula $r \approx -i\,e^{2iF}$ at the turning point $\eta = -i$ (in $\sigma = \sqrt{\lambda/2}\,\eta$) gives $\ln|r| = c\sqrt{k\Delta}$ with

$$c = \frac{2\sqrt{\pi}\,\Gamma(3/4)}{\Gamma(1/4)}\ \mathrm{Re}\,\sqrt{L} = 1.809205,$$

$$1 + \int_1^\infty \bigl(1 - \sqrt{1-\eta^{-4}}\bigr)d\eta = \frac{2\sqrt{\pi}\,\Gamma(3/4)}{\Gamma(1/4)} = 1.19814.$$

The term $-\lambda/\sigma^3$ contributes only the phase $-\pi/4$, so no additive constant appears.

(iii) Together:

$$\ln|C| = -\pi k\Delta + c\sqrt{k\Delta} - \tfrac{1}{2}\ln(a/b) + O\bigl((k\Delta)^{-1/2}\bigr).$$

Numbers (run 2): slope of $\ln|r|$ between $k\Delta = 300$ and 600 is 1.808965 (-0.013%, criterion 0.3%); $\ln|r| - c\sqrt{k\Delta} = 0.0057,\ 0.0040$ (criterion ≤ 0.05, falling); the full problem at $k\Delta = 80, 100, 120$ deviates from (iii) by $-0.0028,\ -0.0085,\ -0.0045$ (criterion 0.02). Both senses of rotation give the same numbers. *Disclosure.* The validity condition (accuracy controls $\le 10^{-6}$) is not met in one place. For the sense + at $k\Delta = 600$ the control is 1.4×10^{-5} in run 2 (contour $1.3h$ instead of $0.7h$, changed after run 1, where the $0.7h$ contour lost digits up to 1.0). Formally the test is therefore invalid. For the sense − it is valid in full and confirmed, and every content criterion is met with a margin of a factor 2 to 20.

Proof and measurement sources. Scripts `k206_abfall_wurzelterm.py`, `k207_lokales_problem.py`, `k208_wkb_pruefung.py` with their outputs, and the notes in `notes/research/v4_10_0_r02/`; pre-registrations and results H91–H93 in `notes/research/predictions/`. □

Consequence. The admixture at the big bang now has a closed asymptotic form. The exponent π comes from the position of the singularity of tanh, the $\sqrt{k\Delta}$ term from the essential singularity of the weight e^{Ls} there, and the constant from the adiabatic amplitude between the weights before and after the bang. This explains the slope below π of r86.1(b), the waves (from the subdominant turning point, not checked separately) and the missing factor of r02.2(a). The cut-off scale stays about $1/(\pi\Delta)$, shifted only by a factor growing like $e^{1.8\sqrt{k\Delta}}$; Δ remains free. OP5 is not closed.

v4.10.0-r02 status in the three frames

- The transition joins frame T (before the bang) and frame M (after it) under G2, with the weights a, b of frame M. All results concern this M–T transition. A frame-free statement is not made.
- Photon frame P: with photon weights $a = b$ one has $L = \mp i\pi$, the constant $-\frac{1}{2}\ln(a/b)$ vanishes and c would be $1.19814\sqrt{\pi/2} = 1.502$. This is a consequence noticed afterwards and was not computed. In the sharp limit the photon weights give no admixture (r85.5), unchanged.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.10.0-r02 register entries

statement	status after v4.10.0-r02
fall-off up to $k\Delta = 60$	numbers secured; H91 invalid (planning error)
prefactor as local Stokes function	waves, symmetry and Born limit confirmed; normalization missed (H92)
closed asymptotics of $\ln \lvert C \rvert$	derived; numbers met; formally invalid for sense +, valid and confirmed for sense − (H93)
$\lvert C \rvert$ for both senses	equal (afterwards, H91; local form H92c)
cut-off scale	$\approx 1/(\pi\Delta)$, Δ free

v4.10.0-r02 ledger and guard update

$$R_{\mathrm{chk}}^{v4100\text{-}r02} = (1, 1, 0)$$

for (fall-off computed to $k\Delta = 600$; asymptotics derived; one accuracy control above its bound).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v4100\text{-}r02,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v4100\text{-}r02} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.10.0-r03 Source-Only: Fermions and Bosons at the Big Bang, and the Vacuum Term under G2

Purpose. Layer v4.10.0-r02 closed the asymptotics of the bosonic admixture at the big bang (§A.272). This layer records what follows for the other fields and for the vacuum term under the domain choice G2 (tachyon reading before, matter reading after the bang; r85). It contains one pre-registered test (H94), an author decision on the duration Δ of the transition, an exact symmetry of the two senses of rotation, and analytic explorations of four routes to Λ under G2. Two of my own intermediate claims are corrected here. The notes are `notes/research/v4_10_0_r03/`. Earlier layers are not rewritten.
Source-only status. This v4.10.0-r03 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.10.0 Version DOI 10.5281/zenodo.23991357 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.10.0-r03 results

Theoremlet v4.10.0-r03.1 (fermions pass the bang without admixture). Exploration (analytic, no criterion). Under G2 and the second reading the symmetric Dirac or Weyl action is weighted by $c(\tau)$, going from $-b$ to a. The field equation $c(i\gamma^\mu\partial_\mu - m)\psi + \frac{i}{2}c'\gamma^0\psi = 0$ is solved exactly by $\psi = c^{-1/2}\chi$, with χ free, for every profile (sympy, residual 0).

(a) Positive and negative frequencies do not mix: no fermions are created at the bang, for any k and any profile. A first-order equation lets the weight be scaled away; the second-order equations of scalars and gauge fields do not.

(b) Transfer factor $\psi_{\text{after}}/\psi_{\text{before}} = \pm i\sqrt{b/a} = \pm i\beta_{\text{sym}}$ (continuous branch along the path; the two senses differ in phase only). The map is unitary in the weighted norm, and the frequency is kept: a particle of the T reading becomes an antiparticle-like state of the M reading, exactly and for all k.

(c) Correction (made in the notes the same day): mass terms keep their sign under the image (r85, H82), so before the bang the mass is $-m$ relative to the kinetic term. On FRW the mass term is $m\,a(\tau)$ with $a \to 0$ at the bang, so (a) and (b) hold for $k \gg m\,a$ at the transition.

Theoremlet v4.10.0-r03.2 (H94: the weighted norm at the bang). Pre-registered (commit `99f1cce`, partly not blind). $N = a(|C|^2 - |D|^2)/b$ for the mode of positive norm before the bang (tanh profile, $\Delta = 2$). All four parts confirmed (one run, Wronskian constant to 10^{-11}).

(a) Bosons: $N \to +1$ for long waves (1.024, 0.974 at $k\Delta = 0.01$) and $N \to -1$ for short waves (-0.969, -1.032 at 20).

(b) Exactly one sign change per sense of rotation, at $k^*\Delta = 0.7361$ and 0.2569.

(c) In between the norm is not conserved ($0.1 < |N| < 0.9$ at 10 and 5 grid points).

(d) Fermions: $N_F = -1$ on all points, to 6×10^{-12}.

Noticed afterwards. The bosonic norm is also amplified (up to 1.18 and -1.63). The mean of the two senses does not restore it. The statement "bosons with matter weights are created, $n = 0.404$ per mode" holds only in the picture of the sharp real jump.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Decision v4.10.0-r03.3 (author, 6 November 2026: $\Delta \lesssim \ell_P$). The duration of the transition is taken at or below the Planck length. All modes below the Planck scale then have $k\Delta \ll 1$: bosons with matter weights pass norm-preserving as a Bogoliubov transformation with $n = (1-\beta_{\text{sym}}^2)^2/(4\beta_{\text{sym}}^2) = 0.404$ per mode, and fermions are mirrored by CPT without creation. The non-unitary range $k \sim 1/\Delta$ lies at or beyond the Planck scale. The total energy of the created bosons is set by that range and stays open.

Theoremlet v4.10.0-r03.4 (the two senses of rotation are exactly conjugate). Exploration (derived analytically, checked numerically). For the transfer matrix $M(L) = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ in the basis (e^{ikt}, e^{-ikt}) one has $\det M = e^{-L}$ (weighted Wronskian). The duality $v = wy'$ gives $M(-L) = e^L JMJ$ with $J = \operatorname{diag}(1,-1)$. Reflection with conjugation, using $s(-t) = 1 - s(t)$, gives $M(-\bar{L}) = \overline{M(L)}^{-1}$. Together

$$M(\bar{L}) = \begin{pmatrix} \bar{d} & \bar{b} \\ \bar{c} & \bar{a} \end{pmatrix}.$$

Hence $C(\bar{L}) = \overline{C(L)}$, so $|C_+| = |C_-|$ exactly, and $D_+\overline{D_-} - b_+C_+ = e^{-L}$, the product rule of r85.6 with its correction term. For the local problem of r02.2, $M = \begin{pmatrix} 1 & 0 \\ r & 1 \end{pmatrix}$ exactly, so $r(\bar{\lambda}) = \overline{r(\lambda)}$. Numerically all identities hold to $\leq 3 \times 10^{-10}$ at $k\Delta = 0.01$–5.

Theoremlet v4.10.0-r03.5 (no Λ from the tachyon sector after the bang under G2). Exploration (analytic, no criterion). Four routes were examined.

(a) Mixture (deferred idea I3): its Λ side came from the tension of the T reading. Under G2 only the M reading exists after the bang, so only the dust part remains.

(b) Tachyonic condensation (I4): after the bang there are no tachyon modes. Before it the image of a scalar sees $-V$ ($\mathcal{L}_T = -\frac{1}{2}X - V$, r84); a potential stable in M runs away there ($\ddot{\phi} = m^2\phi + \lambda\phi^3$) instead of condensing. Under G1 one has $m_{\text{eff}}^2 = \gamma m^2 > 0$.

(c) Ghost condensate (r83.4): under G1, $aP_M + bP_T = (a-b)X/2 + (a+b)cX^2$ is healthy and does not condense. Under G2 the condensate exists only before the bang; after it P_M has $w \in [\frac{1}{3}, 1]$, never -1; at the bang the condensate has $\pi = 0$, so the M field is at rest.

(d) Conclusion: under G2 a vacuum term after the bang can only be $aV_{\min}$ of the M reading, with a free value. The structure does not fix Λ under G2.

Theoremlet v4.10.0-r03.6 (boundary condition and spatial reflection; correction). Exploration (analytic). If the T field before the bang is identified pointwise with the image of the M field after it, $\phi_T(x) = \phi_M(\Lambda(i\pi)x)$, the junction yields $\phi'(0) = 0$ for scalars and $\psi(0) = 0$ for fermions ($\beta_{\text{sym}} \neq 1$). The same identification needs a direction n and a spatial origin (from $x \mapsto \Lambda x + 2x_0$) that nothing fixes; classically it would correlate the modes k and $P_n k$. Quantum mechanically it is inconsistent for $a \neq b$: with antiunitary time reversal the junction forces $u'(0) = 0$, and the Wronskian normalization $uu'^* - u^*u' = i$ then fails at $\tau = 0$. *Correction:* the conditions $\phi'(0) = 0$, $\psi(0) = 0$ stated the same day in the notes do not follow from the structure; a frozen scalar at the bang is a possible choice of state, not a consequence. Under G2 the second reading is a statement about the action, $-b\mathcal{L}_M$ before and $a\mathcal{L}_M$ after, with independent data before the bang, as in H85–H94; at that level neither a direction nor an origin appears (r84, H79).

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof and measurement sources. Scripts `k209_fermionen_urknall.py`, `k210_norm_uebergang.py`, `k211_symmetrie_drehsinne.py` with their outputs, and the explorations in `notes/research/v4_10_0_r03/`; pre-registration and result H94 in `notes/research/predictions/`. □

Consequence. Under G2 the big bang separates bosons from fermions. Fermions are mirrored by CPT exactly and unitarily. Bosons with matter weights are Bogoliubov-transformed for $k\Delta \ll 1$ and CPT-mirrored for $k\Delta \gg 1$, and the transition is non-unitary in between, which the decision $\Delta \lesssim \ell_P$ moves to the Planck scale. The two senses of rotation are exactly conjugate. None of the tachyon-sector mechanisms yields a vacuum term after the bang; Λ remains a free $aV_{\min}$. OP5 is not closed.

v4.10.0-r03 status in the three frames

- r03.1–r03.4 concern the M–T transition under G2, with the weights a, b of frame M: frame T before and frame M after the bang. No frame-free statement is made.
- Photon frame P: with $a = b$ there is no admixture, the fermion factor has modulus 1, and the pointwise identification of r03.6 is not obstructed. The norm then stays $+1$ for bosons, as in r85.5.
- r03.5 states that frame T contributes no vacuum term in frame M after the bang under G2. Under G1 both readings coexist, and the reduced theory is the M theory with renormalized weights (r84).

v4.10.0-r03 register entries

statement	status after v4.10.0-r03
fermions at the bang (G2)	no admixture, CPT exact, factor $\pm i\beta_{\text{sym}}$ (exploration)
bosonic norm at the bang	$+1 \to -1$ with one zero; non-unitary near $k \sim 1/\Delta$ (H94)
duration Δ	$\lesssim \ell_P$ (author decision)
senses of rotation	exactly conjugate, $\lvert C_+\rvert = \lvert C_-\rvert$ (exploration)
Λ from I3, I4, ghost condensate under G2	none after the bang; $\Lambda = aV_{\min}$ free
pointwise identification of T and M	inconsistent for $a \neq b$; boundary conditions withdrawn

v4.10.0-r03 ledger and guard update

$$R_{\text{chk}}^{v4100\text{-}r03} = (1, 1, 0)$$

for (fermions and the bosonic norm at the bang settled; symmetry exact; Λ under G2 without a mechanism).

The layer spends no guard:

$$R_{\text{glob}}^{(v4100\text{-}r03,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\text{guard}}^{v4100\text{-}r03} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.10.0-r04 Source-Only: The Vacuum Term with a Tachyon Branch, and the Bosons Created at the Big Bang

Purpose. Layer v4.10.0-r03 found that under the second reading no tachyon-sector mechanism gives a vacuum term after the big bang (§A.272). This layer records the check under the domain G1, a temporary author decision to admit the tachyon branch as degrees of freedom of its own, two analyses of the deferred idea I3 under that decision (fixed weights and a dynamic branch gas, H95), the withdrawal of the decision, an estimate of the bosons created at the bang, and the state of the comparison data of the open predictions. The notes are `notes/research/v4_10_0_r04/`. Earlier layers are not rewritten.
Source-only status. This v4.10.0-r04 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.10.0 Version DOI 10.5281/zenodo.23991357 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.10.0-r04 results

Theoremlet v4.10.0-r04.1 (domain G1 gives no tension either). Exploration (analytic). The mixture of I3 (k182, k183) consists of a normal branch ($\omega = m\cosh\eta$, $k = m\sinh\eta$) and a tachyonic branch ($\omega = m\sinh\eta$, $k = m\cosh\eta$); the tension comes from the tachyonic one. The reduced theory of the second reading, $(a-b)\frac{1}{2}X - (a+b)V$, has $\omega^2 = k^2 + \gamma m^2$ for every mode and no tachyonic branch. For a massive scalar the averaged pressure vanishes (dust). Under G1, as under G2, the only vacuum term is a free multiple of $V_{\min}$. I3 needs the tachyonic branch as degrees of freedom of its own, that is, a reversal of the decision of r84.

Decision v4.10.0-r04.2 (author, 6 November 2026: tachyon branch admitted, later withdrawn). First the tachyonic branch was admitted as degrees of freedom of its own, so that I3 becomes possible. The known cost: quanta of negative energy, and vacuum decay through the graviton with a frame cut-off $\lesssim 14$ MeV (r83, H76). After r04.3 and r04.4 the author withdrew the decision: the tachyon part is again only the second reading of the same field (r84).

Theoremlet v4.10.0-r04.3 (I3 with fixed weights is excluded). Exploration (analytic, with an estimate). The M and T readings of the mixture have the same density and dilute alike. With fixed weights the sum is one fluid with constant $w = -b/(3a)$, not dust plus Λ. To explain $\Omega_\Lambda \approx 0.69$ today one needs $b/a \approx 2$, and then the dark component at recombination is about 10^{-5} of the baryons instead of 5.4 (cosmological values from prior knowledge, not looked up). In addition, the mixture at common rapidity is not preserved under redshift: $k_1/k_2 = \tanh\eta$ would stay fixed while each k is fixed by η.

Theoremlet v4.10.0-r04.4 (H95: a dynamic branch gas cannot do it either). Preregistered (commit `9eb4e40`, not blind; the refutation was expected analytically). Model: flat FRW with radiation and baryons, and a freely streaming gas with equal numbers in both branches. The normal branch has weight 1; the tachyonic branch has weight r, negative energy, and threshold at $a_{\rm th} = k_0/m$. Grid $60 \times 60 \times 40$ in $(r, a_{\rm th}, \kappa_1/a_{\rm th})$: of 144 000 points, 12 780 keep the total density positive for $a \in [10^{-8}, 1]$. None has both a dark-to-baryon ratio in $[4.32, 6.48]$ at $z = 1090$ and $w_{\rm eff} \in [-0.8, -0.6]$ today. **Refuted.** *Noticed afterwards.* w_0 can be made arbitrarily negative (down to -151), but only with the threshold practically today ($a_{\rm th} = 1.000001$), and then the ratio at recombination is 266.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Theoremlet v4.10.0-r04.5 (bosons created at the bang; estimate). Exploration. With $\Delta \lesssim \ell_P$ (r03.3), bosons with matter weights receive $n_0 = 0.404$ per mode up to $k_{\max} \sim 1/\Delta \gtrsim M_P$. This gives $\rho \approx g\, n_0\, k_{\max}^4/(8\pi^2) = g \cdot 0.0051\, k_{\max}^4$, equivalent to a temperature of $\approx 0.35\, k_{\max}$. The radiation after the bang is Planck-hot, non-thermal, and initially without fermions. The number depends on the non-unitary range $k \sim 1/\Delta$ of r03.2 and is an order of magnitude only.

Remark v4.10.0-r04.6 (state of the comparison data). For H15 (m_W) no world average published after 2026-10-21 was found, and for H42/H46 (θ_{23}) no global analysis after 2026-10-24; all three remain open. *Disclosure:* The PDG 2026 edition (May 2026) already gave $m_W = 80.3625 \pm 0.0077$ GeV before the pre-registration of H15, which listed only PDG 2025 as known. The value is 0.13σ from the prediction and is not a test.

Proof and measurement sources. Script `k212_zweiggas_frw.py` with output, and the explorations and the proposal in `notes/research/v4_10_0_r04/`; pre-registration and result H95 in `notes/research/predictions/`. □

Consequence. The deferred idea I3 (Λ from the dark-matter calculation) has failed in both forms examined, with fixed weights and with a dynamic branch gas, even with the tachyon branch admitted. The author returned to the second reading alone, so Λ is an input. The bosons created at the bang form a Planck-hot, non-thermal start without fermions. An observable trace would have to survive thermalization, for example a gravitational-wave background, which first needs a rule for how tensor modes see the transition. OP5 is not closed.

v4.10.0-r04 status in the three frames

- r04.1–r04.4 concern how the gravity of frame M sees frame T. Under the second reading T is a re-weighting of the same field (no own branch). With an own branch, T would have quanta of negative energy; that was examined and withdrawn.
- r04.5 holds in frame M after the bang. With photon weights (frame P, $a = b$) no bosons are created.
- No frame-free statement is made.

v4.10.0-r04 register entries

statement	status after v4.10.0-r04
Λ under G1 (second reading)	no tension; free multiple of $V_{\min}$
tachyon branch as own degrees of freedom	admitted, then withdrawn (author)
I3 with fixed weights	excluded (constant w, CMB)
I3 with dynamic branch gas	refuted (H95)
bosons created at the bang	$\rho \approx g \cdot 0.0051\, k_{\max}^4$, non-thermal, no fermions (estimate)
H15, H42, H46	open; PDG 2026 m_W disclosed as known before H15

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.10.0-r04 ledger and guard update

$$R_{\mathrm{chk}}^{v4100\text{-}r04} = (1, 1, 0)$$

for (I3 excluded in both forms; decision withdrawn; Λ remains an input).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v4100\text{-}r04,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v4100\text{-}r04} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.10.0-r05 Source-Only: Gravitons at the Big Bang, a Review from the Core, and the Readout of Λ

Purpose. This layer has three parts. First, an estimate of gravitons created at the big bang and the author decision taken on it, with a correction to its rationale. Second, a review of the work since r81 from the core of the companion, prompted by the author's observation that the results keep landing "a little off" while much structure appears. The review leads to a separation of OP5 into three parts. Third, the readout of Λ at the present epoch, with one prediction without free parameter (H96) and one refuted dynamic bridge (H97). The notes are `notes/research/v4_10_0_r05/`. Earlier layers are not rewritten.
Source-only status. This v4.10.0-r05 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.10.0 Version DOI 10.5281/zenodo.23991357 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.10.0-r05 results

Theoremlet v4.10.0-r05.1 (gravitons at the big bang; estimate and correction). Exploration, not blind (the bound on N_{eff} is known; cosmological values from prior knowledge).

(a) With matter weights, gravitons receive $n_0 = 0.404$ per mode up to $k_{\max} \sim 1/\Delta$ (r03.3). With the bosons of the Standard Model thermalizing and the gravitons decoupled, this gives $\Delta N_{\mathrm{eff}} \approx 3.6$; the bound ≈ 0.3 would need $\gtrsim 300$ bosonic degrees of freedom.

(b) Author decision (6 November 2026): gravity carries photon weights ($a = b$) at the transition.

(c) Correction to the rationale of (b): photon weights suppress creation only for $k\Delta \ll 1$. At the smooth transition the produced energy is of the same order ($\int x^3 n_{\mathrm{mix}}\, dx = 0.066$ against $0.091\,\Delta^{-4}$, maximum at $k\Delta \approx 0.76$), so $\Delta N_{\mathrm{eff}} \approx 2.6$.

(d) Counting only modes below $x_c = k_c\Delta$ keeps $\Delta N_{\mathrm{eff}} \lesssim 0.3$ for $x_c \lesssim 0.13$ (or $\lesssim 0.05$ if only the Higgs carries matter weights). This would be another chosen quantity. ΔN_{eff} at the bang is therefore recorded as an open problem, not closed by a cut-off.

Remark v4.10.0-r05.2 (review: why "a little off"). A count of the tests since v4.8.0 (wiki table, after r04) gives about 20 confirmed, 19 refuted, 9 invalid, 4 open and 28 mixed results, 29 marked not blind. The confirmed tests without that mark all check inner mathematics (the OP5 comparison model, one construction). Against external measurements there is no confirmed blind prediction yet.

(a) Much of the structure that "works" is known, consistent physics taken up by the companion. Examples: Dixon and Furey for one generation, triality, Einstein–Cartan with the torsion 3/16, Boyle, Finn and Turok for the big bang, ghost condensation, WKB.

(b) What is new is the frame reading (M, T, P, β_{sym}, weights a, b, G2). Its physical claims are identifications of single dimensionless numbers at one point. With many constants and combinations, agreement within a few percent is expected; r56 and r70.4 found chance level.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(c) Since r81 about twelve author decisions were taken, most of them after a consequence did not fit. Each kept a consequence in range; none produced a confirmed new prediction. The aim W7 ("vacuum term not free") of the action programme is not met.

Decision v4.10.0-r05.3 (author: OP5 separated into three parts). Reading the core shows two meanings of OP5. In the architecture, OP5 is the reversal-stable return readout of the route whose source readout is OP1; closed means $\mathcal{I}(\mathrm{OP1}) = \mathcal{I}(\mathrm{OP5}_{\mathrm{sym}})$ (Theorem H7.2), locally under the guards. In physics, Λ is a readout quantity, $\Omega_\Lambda = 1 - \beta^2_{\mathrm{sym}}$ and $\Omega_M = \beta^2_{\mathrm{sym}}$ as branching of the cascade (Theorem 22.16), not a dynamical vacuum energy. From late October the question became "which dynamics produces Λ" (OP5 series, then r81–v4.10.0-r04). The negative results of that line answer a question the core does not ask. Decision of 6 November 2026:

(i) *OP5 (core):* the reversal closure of the source–return pair, locally closed, not claimed globally;

(ii) Λ *readout:* whether $\Omega_\Lambda : \Omega_M = (1-\beta^2) : \beta^2$ is the cascade branching, and at which epoch;

(iii) *action programme W1/W2:* a separate branch with the negative results of r81–v4.10.0-r04, frozen, without further repair decisions.

A second decision fixes the epoch of (ii) as the present (way (a) of v4.8.0-r17).

Theoremlet v4.10.0-r05.4 (H96: Ω_M today without free parameter; pre-registered, open). Pre-registered (commit `aeaba3d`, supplement on prior knowledge `e338b9b` before any comparison). Prediction:

$$\Omega_M = \beta^2_{\mathrm{sym}} = \frac{\gamma_L - 1}{\gamma_L + 1} = 0.30185, \qquad \gamma_L = \gamma_{\mathrm{BI}}/\gamma_{\mathrm{string}} = 1.864700.$$

Comparison: the first flat-ΛCDM value of Ω_m published after 2026-11-06 from DESI DR3 (BAO alone) or from Euclid; confirmed within 2σ. Known values (DESI DR2 0.2975 ± 0.0086, Planck 0.3153 ± 0.0073 and others) are disclosed and do not count. The prediction rests on the U(1) counting of horizon states (r68) and on reading γ_L, the ratio of a corrected to a superseded value of the same LQG parameter (v4.8.0-r03 F11), as a Lorentz factor.

Theoremlet v4.10.0-r05.5 (the only dynamics for the ratio, and a bridge refuted). Search and one pre-registered test.

(a) The only place where the ratio follows from a dynamics is r31: the Markov cascade with return path has stationary flux fractions β^2, $1-\beta^2$ for every rate. A bridge to cosmic time is missing. The earlier link of the core (scale factor as proper time, H as activation rate of Riemann zeros) was withdrawn with the Riemann paper (v4.8.0-r04).

(b) Proposed bridge: the content of S^2 is vacuum energy that returns to S^8 at rate $\Gamma = \varepsilon H$ and splits there. Net, the vacuum decays into matter at rate $\beta^2 \varepsilon H$; with $\varepsilon = 1$ this has no free parameter.

(c) H97 (commit `de36fcc`, not blind): this bridge with $\varepsilon = 1$ and $\Omega_M = \beta^2$ against DESI DR2 BAO (13 values) gives $\chi^2_{\mathrm{min}} = 105.0$ for 12 degrees of freedom, $p = 6 \times 10^{-17}$. **Refuted.** The control (flat ΛCDM, free Ω_m) reproduces DESI ($\Omega_m = 0.2970$, $hr_d = 101.56$ Mpc). For comparison, flat ΛCDM at $\Omega_M = \beta^2$ has $\chi^2 = 10.9$, $p = 0.54$. *Exploration afterwards:* DR2 allows $\varepsilon \lesssim 0.4$ without preferring $\varepsilon > 0$.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof and measurement sources. Scripts `k213_gravitonen_neff.py`, `k214_erzeugung_photongewichte.py`, `k215_neff_grenze.py`, `k216_bruecke_desi_dr2.py` with outputs; the reviews and proposals in `notes/research/v4_10_0_r05/`; pre-registrations and results H96, H97 in `notes/research/predictions/`. □

Consequence. The companion is, at its core, a kinematic readout structure with sound algebra. Its new physical claims are identifications. They become testable when they predict a function or several observables with one parameter, not one number at one point. OP5 is separated accordingly. The readout of Λ today gives a prediction without free parameter (H96, open). The simplest dynamics for it is refuted (H97). ΔN_{eff} at the bang is open. OP5 (core) stays locally closed under the guards; no global closure is claimed.

v4.10.0-r05 status in the three frames

- r05.1 concerns the M–T transition under G2; the photon-weight decision assigns gravity the weights of frame P.
- r05.4 and r05.5: β^2_{sym} is a matter-frame quantity (r22); the readout $\Omega_M = \beta^2$ is made by a matter-frame observer. In frame P it vanishes ($\Lambda_{\text{fundamental}} = 0$ of Theorem 22.16). No frame-free statement is made.

v4.10.0-r05 register entries

statement	status after v4.10.0-r05
gravitons at the bang	photon weights (author); ΔN_{eff} open (creation at $k \sim 1/\Delta$)
OP5	separated: core closure (local) / Λ readout / action programme (frozen)
epoch of the Λ readout	present (author)
$\Omega_M = \beta^2_{\text{sym}} = 0.30185$	pre-registered prediction H96, open
cascade–Friedmann bridge, $\varepsilon = 1$	refuted (H97)
blind predictions confirmed against data	none yet; open: H15, H42, H46, H96

v4.10.0-r05 ledger and guard update

$$R^{v4100\text{-}r05}_{\text{chk}} = (1, 1, 0)$$

for (review done and OP5 separated; prediction H96 fixed; the dynamics for the readout absent).

The layer spends no guard:

$$R^{(v4100\text{-}r05,+)}_{\text{glob}} = (0, 1, 1, 1, 1), \qquad \Gamma^{v4100\text{-}r05}_{\text{guard}} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.10.0-r06 Source-Only: Semiclassical Einstein Equations from Relative Entropy (Dorau–Much) and the Readouts of the Companion

Purpose. This layer takes up a recent result from the literature, pointed out by the author: P. Dorau and A. Much (Leipzig), *From Quantum Relative Entropy to the Semiclassical Einstein Equations*, Phys. Rev. Lett. **136**, 091602 (2026), DOI 10.1103/lmq8-nsty, arXiv:2510.24491. It records what the paper shows and assumes, and where it meets the readouts of the companion: the entropy factor of §21 (v4.8.0-r03 F11a), the free Λ (v4.10.0-r03, r05), the sign of gravity (v4.10.0-r01) and the frames. The layer is an exploration of the literature; it contains no pre-registered test and no new prediction. The note is `notes/research/v4_10_0_r06/`. Earlier layers are not rewritten.
Source-only status. This v4.10.0-r06 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.10.0 Version DOI 10.5281/zenodo.23991357 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.10.0-r06 results

Theoremlet v4.10.0-r06.1 (the result of Dorau and Much, as stated by them). Literature, read in arXiv version v3; the formulas were not recomputed here.

(a) *Setting.* Near a point the metric is approximated by η_{ab} (leading order). The boost field ξ is null on two null hypersurfaces meeting in a spacelike surface $\mathcal{S}$: a local Rindler horizon, as in Jacobson's thermodynamic derivation (1995). A real Klein–Gordon field of mass m is quantized as a CCR algebra. The reference state ω_0 is quasifree, boost invariant and Hadamard (in the Rindler case the Minkowski vacuum); only coherent excitations ω_ϕ are treated.

(b) *Relative entropy equals energy flux.* The modular group of the wedge acts as the boost, Δ^{it} = boost by $2\pi t$ (Bisognano–Wichmann, Summers–Verch), and ω_0 is KMS at $\beta = 2\pi/\kappa$. The Araki relative entropy then becomes

$$S_{\mathrm{rel}}(\omega_0\|\omega_\phi) = -2\pi \int U \,\langle{:}T_{ab}{:}\rangle_{\omega_\phi}\xi^a\xi^b \, dU \, d\mathrm{vol}_{\mathcal{S}}.$$

(c) *Geometry.* With vanishing expansion, shear and twist at the horizon, the Raychaudhuri equation gives $\delta A = -\int U\, R_{ab}\xi^a\xi^b \, dU \, d\mathrm{vol}_{\mathcal{S}}$.

(d) *Assumed, not derived:* $S_{\mathrm{rel}} = \delta A/4$ (Bekenstein–Hawking, units $G = \hbar = 1$). This fixes the coupling $\alpha = 8\pi$.

(e) *Result.* Holding for all null directions, $\alpha\langle T_{ab}\rangle = R_{ab} + N g_{ab}$; conservation and the contracted Bianchi identity give $N = -R/2 + \Lambda$, hence $G_{ab} + \Lambda g_{ab} = 8\pi\langle{:}T_{ab}{:}\rangle$ with Λ an arbitrary integration constant.

The authors call this "arguments indicating" the result: leading order, coherent states, scalar field. The press reading "gravity from vacuum energy" is not what is shown: the vacuum enters as the reference state of a relative entropy, not as a source of energy.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Theoremlet v4.10.0-r06.2 (the entropy factor becomes the gravitational coupling). Analytic, not blind (follows from r06.1 by one line). Let a horizon counting give $S = c\,A/4$ with $c > 0$. Replacing (d) by $S_{\mathrm{rel}} = c\,\delta A/4$ and keeping the rest of r06.1 gives

$$\alpha = \frac{8\pi}{c}, \qquad G_{\mathrm{eff}} = \frac{G}{c}.$$

For the two countings of §21 (F11a in §A.271) one has $c = \gamma_*/\gamma$, where γ_* is the value at which the counting reproduces S_{BH}: $\gamma_{\mathrm{string}} = \ln 2/(\pi\sqrt{3})$ for the counting with $j = \frac{1}{2}$ only, $\gamma_{\mathrm{BI}} = 0.237533$ for the counting of Domagala–Lewandowski and Meissner. At one common value of γ the two countings differ by the factor $\gamma_{\mathrm{BI}}/\gamma_{\mathrm{string}} = \gamma_L = 1.864700$ (F11a). Under the chain of r06.1 this factor would appear as a ratio of gravitational couplings:

$$\frac{G_{\mathrm{eff}}^{(j=1/2)}}{G_{\mathrm{eff}}^{(\mathrm{DL/M})}} = \frac{c_{\mathrm{DL/M}}}{c_{j=1/2}} = \frac{\gamma_{\mathrm{BI}}}{\gamma_{\mathrm{string}}} = \gamma_L.$$

Consequence. If the T frame counts with $j = \frac{1}{2}$ and the M frame with all spins (r63, H47) at one common γ, the two frames see different Newton constants. A single measured G is consistent only if each frame uses the γ at which its own counting reproduces S_{BH}; then $c = 1$ in both, G is frame independent, and the frame difference sits in γ. This is the reading of r64 (the frame difference in γ, not in $\hbar$ or G); Dorau–Much make it a statement about the coupling, not about the entropy alone.

Theoremlet v4.10.0-r06.3 (Λ as integration constant). Literature and comparison. In r06.1(e) Λ is not fixed by the derivation; it enters through $\nabla^a(\Lambda g_{ab}) = 0$. This is the same situation as in v4.10.0-r03 and r05: under G2 the action gives no Λ after the bang, $\Lambda = a\,V_{\min}$ is free, and Λ is a readout (Theorem 22.16). The agreement supports the classification "Λ is not part of the dynamics". It is not a success of the companion: the standard derivation does not fix Λ either. The readout $\Omega_M = \beta_{\mathrm{sym}}^2$ (H96) is neither supported nor contradicted.

Remark v4.10.0-r06.4 (the sign of gravity from positivity; afterwards). In r06.1 the sign of α is not free: $S_{\mathrm{rel}} \geq 0$ and $c > 0$ give $\alpha > 0$, i.e. attractive gravity. In v4.10.0-r01 the sign of W1 was corrected by an author decision after H90 had shown repulsion. The corrected sign coincides with the one fixed by positivity. This is noted afterwards and is not a derivation of W1: Dorau–Much use a scalar field and Riemannian geometry, W1 is Einstein–Cartan with spinors and torsion $+(3/16)\kappa j^2$. A transfer would need the relative entropy for Dirac fields, which the paper does not treat.

Remark v4.10.0-r06.5 (frames, Wick rotation and the modular group; open). The derivation uses that the modular group of the reference state is a real one-parameter group of boosts with KMS period 2π. In the Euclidean picture that period is the full rotation angle 2π (the Unruh temperature via Wick rotation). The companion has this bridge itself (Theorem 21.10, $t \mapsto i\tau_E$ on the biquaternion). Under the tachyon map $z \mapsto iz$ of Convention F the rapidity becomes imaginary and the boost a rotation. Whether $z \mapsto iz$ coincides with the Wick rotation of Theorem 21.10 is open. If it does, frame T is the Euclidean side of the same KMS structure, not its loss. The precise question is operator algebraic: is there a state on the wedge algebra whose modular group is the tachyon image of the boost? No answer is given here.

Proof and measurement sources. Exploration note `Erkundung_Dorau_Much_relative_Entropie.md` in `notes/research/v4_10_0_r06/`; the paper (arXiv:2510.24491v3). r06.2 is one line of algebra on r06.1(d)–(e). □

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Consequence. Dorau and Much give a rigorous quantum field theoretic form of Jacobson's argument. They assume the area law and leave Λ free. For the companion this has three uses: the entropy factor of §21 is the gravitational coupling (r06.2), so the frame reading of the horizon countings must keep G frame independent; Λ as a readout is consistent with the standard derivation (r06.3); and the frames get a precise operator algebraic question (r06.5). Neither G nor Λ follows from it.

v4.10.0-r06 status in the three frames

- M: the derivation of r06.1 holds as stated (real boosts, KMS at $2\pi/\kappa$).
- T: open (r06.5); under $z \mapsto iz$ the boost becomes a rotation, and whether the KMS structure survives as its Euclidean side is not decided.
- P: degenerate fixed point; no separate statement.

r06.2 is frame dependent by construction (one counting per frame); r06.3 holds in every frame that has the derivation. No frame-free statement is made.

v4.10.0-r06 register entries

statement	status after v4.10.0-r06
Einstein equations from relative entropy	literature (Dorau–Much 2026); area law assumed, Λ free
entropy factor of §21	equals the gravitational coupling (r06.2); G frame independent only with matched γ per frame
Λ	integration constant in the standard derivation; readout in the companion (consistent)
sign of W1 (v4.10.0-r01)	coincides with the positivity sign (afterwards, scalar case only)
frame T and the modular group	open question (r06.5)

v4.10.0-r06 ledger and guard update

$$R_{\mathrm{chk}}^{v4100\text{-}r06} = (1, 1, 0)$$

for (literature recorded; entropy factor tied to the coupling; frame T under the modular group open).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v4100\text{-}r06,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v4100\text{-}r06} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.10.0-r07 Source-Only: The Relative-Entropy Chain in the Tachyon Frame, and a Correction to r06.5

Purpose. The author asked whether the companion can map the derivation of Dorau and Much (§A.272) onto its own frames and so obtain the Einstein equations internally. A pre-registered test was planned. The search of prior work showed that the answer is fixed analytically, so this layer is an exploration, not a test; no hypothesis number is assigned. On the way, Remark r06.5 turned out to be wrong. A second question of the author, why frame independence is reached and the prefactor of the area law is not, is answered in r07.4. The notes are `notes/research/v4_10_0_r07/`. Earlier layers are not rewritten.
Source-only status. This v4.10.0-r07 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.10.0 Version DOI 10.5281/zenodo.23991357 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.10.0-r07 results

Correction v4.10.0-r07.1 (to Remark r06.5). Remark v4.10.0-r06.5 states that under the tachyon map $z \mapsto iz$ the rapidity becomes imaginary and the boost a rotation, and asks whether $z \mapsto iz$ is the Wick rotation of Theorem 21.10. This is wrong. With $z = E + iB$ (v4.9.0-r02), $z \mapsto iz$ is the electromagnetic duality rotation $E \mapsto -B$, $B \mapsto E$, a map of field space, not of spacetime. By Theoremlet v4.8.0-r20.1 it commutes with every element of $SO(3,\mathbb{C})$, the proper Lorentz group acting on z. A boost stays a boost with real rapidity. The question of r06.5 is answered in the negative: $z \mapsto iz$ is not the Wick rotation $t \mapsto i\tau_E$; the two act on different objects. The frame entries of r06 for frame T are replaced by r07.2.

Theoremlet v4.10.0-r07.2 (the chain in frame T for the free Maxwell field). Exploration, not blind (fixed analytically before the check). For the free Maxwell field, the chain of r06.1 gives in frame T the same Einstein equation as in frame M: the same coupling, the same sign, the same inputs.

(a) The duality rotation U_α is a symmetry of the free Maxwell field (conserved helicity). It acts pointwise on F, maps the algebra of the Rindler wedge onto itself and leaves the vacuum invariant.

(b) It commutes with the boost (v4.8.0-r20.1), hence with the modular group of the vacuum (Bisognano–Wichmann).

(c) Relative entropy is invariant under such an automorphism: $S_{\mathrm{rel}}(\omega_0\|\omega_\phi) = S_{\mathrm{rel}}(\omega_0\|\omega_{U\phi})$.

(d) The energy–momentum tensor is Hermitian in z (energy density $\frac{1}{2}|z|^2$, Poynting vector $(\bar{z}\times z)/2i$) and therefore duality invariant; the flux $T_{ab}\xi^a\xi^b$ is unchanged. The bilinear invariant $s = z\cdot z$ changes sign: the Lagrangian density $\frac{1}{2}(E^2 - B^2)$ and $E\cdot B$ are odd (v4.8.0-r06.3, v4.9.0-r40).

Reading. Gravity couples to the Hermitian tensor; the frames differ in the bilinear invariants. For the Maxwell field, gravity is therefore frame blind. This turns the reading of v4.9.0-r64 ("gravity itself is frame independent") into a statement for this case. It is not new physics: the duality invariance of the Maxwell tensor is textbook knowledge.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Proof and measurement sources. (a)–(c) are standard (duality symmetry of the free Maxwell field; invariance of the Araki relative entropy under automorphisms that fix the reference state). (d) and the commutation are checked numerically by `k217_dualitaet_gravitation.py` (2000 random fields and boosts): commutation, invariance of the tensor and of the null flux, and the sign change of $E^2 - B^2$ and $E \cdot B$ hold exactly in floating point; a non-unitary scaling $(1 + 0.3i)z$ changes the tensor by $O(1)$ (negative control). □

Theoremlet v4.10.0-r07.3 (bookkeeping: what the internal version gains). The chain is written in QAM bookkeeping for frames M and T, with the duality rotation as a declared map (`qam_dorau_much_companion.tex`); `qam.sty` raises no error. The exported record gives for both frames the Einstein equation with status *conditional* and inputs {local flatness, area law}, the same inputs as in Dorau–Much. The internal version gains frame independence as a statement (r07.2). It does not reduce the inputs: the area law, and with it the normalization $1/4G$, remains an input (r06.2). For comparison, the same bookkeeping applied to the paper itself (`qam_dorau_much_paper.tex`) reports a promotion if the Einstein equation is declared proved, the area law as an input of the result, and, under the main-body typing of the Bekenstein–Hawking law as a photon-frame statement (§21), a frame mixing without declared map.

Theoremlet v4.10.0-r07.4 (why frame independence is reached and the prefactor of the area law is not). Analytic, not blind (author question after r07.3).

(a) *Frame independence is a statement about relations.* The map between M and T is a symmetry of the system that fixes the reference state and commutes with its modular group (r07.2), and every link of the chain is built from objects invariant under it. Such an argument yields equalities between the frames. It says nothing about the value both frames share.

(b) *The prefactor is an absolute number.* The area law $S = \delta A/4$ fixes the entropy per Planck area, i.e. the coupling (r06.2). A symmetry between frames cannot fix it; something must give one frame its absolute value.

(c) *The countings used by the companion are calibrated to it.* A horizon counting gives $S = (\kappa/\gamma)\, A/\ell_P^2$ with a combinatorial constant κ; the value of the Barbero–Immirzi parameter at which it reproduces the area law $S = A/(N\ell_P^2)$ is $\gamma_* = N\kappa$. Both literature values used by the companion are of this kind: $\gamma_{\text{string}} = 4\kappa_{\text{ABCK}}$, $\kappa_{\text{ABCK}} = \ln 2/(4\pi\sqrt{3})$ (v4.8.0-r03 F11a), and $\gamma_{\text{BI}} = 4\kappa_{\text{DL/M}}$. Their ratio

$$\gamma_L = \frac{\gamma_{\text{BI}}}{\gamma_{\text{string}}} = \frac{\kappa_{\text{DL/M}}}{\kappa_{\text{ABCK}}} = 1.864700$$

does not depend on N: the prefactor cancels (checked for $N = 1, 4, 8$). So do $\beta_{\text{sym}} = (\gamma_L - 1)/(\gamma_L + 1)$ and $\Omega_M = \beta_{\text{sym}}^2$ (H96). Every quantity the companion builds from the two countings is exactly the part in which the normalization of the area law drops out.

(d) *Consequence.* With these inputs the prefactor cannot be derived, in principle and not only in practice: it was put into each γ_* and removed again by taking the ratio. The companion inherits the Barbero–Immirzi ambiguity of loop quantum gravity, in which the area law is matched, not derived. A derivation would need one absolute value of γ (or of the entropy per state) from the structure of the companion, not a ratio; no such statement exists in the corpus.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(e) *A caution.* The state disc at the Hopf point has eigenvalues 3/4 and 1/4 (v4.9.0-r17). Identifying that 1/4 with the one of the area law would be a single-number identification of the kind recorded in v4.10.0-r05.2; there is no mechanism linking the two, and it is not proposed.

Limits. Only the free Maxwell field. For massive fields the tachyon image of the action is tachyonic (H71); a tachyonic field has no stable Hadamard ground state and the premises of the chain fail. Under the second reading (v4.9.0-r84, H79) the tachyon part is a weight on the matter action; how it enters the flux is not examined. Interacting fields: duality is in general not a symmetry.

Consequence. The companion reproduces the Dorau–Much derivation in both frames for the Maxwell field and so obtains frame-blind gravity as a statement. It adds no derivation of the area law; G and Λ remain as in r06. The reason is structural (r07.4): the frame map gives relations, and the countings enter only through a ratio in which the prefactor cancels. The open front is one absolute value of γ from the structure.

v4.10.0-r07 status in the three frames

- M: the chain of r06.1 holds as stated.
- T: for the free Maxwell field the chain holds unchanged (r07.2); for massive fields the tachyon image has no stable ground state (open under the second reading).
- P: the null sector $s = 0$ is fixed by $z \mapsto iz$; no separate statement.

For the Maxwell field, M and T agree: the Einstein equation of the chain is frame free in that case.

v4.10.0-r07 register entries

statement	status after v4.10.0-r07
r06.5 (boost becomes rotation under $z \mapsto iz$)	wrong; $z \mapsto iz$ commutes with $SO(3,\mathbb{C})$ (correction r07.1)
Einstein equation in frame T (Maxwell)	same as in frame M (exploration, not blind)
gravity frame independent (r64)	a statement for the free Maxwell field; open for massive fields
inputs of the internal chain	same as Dorau–Much (local flatness, area law)
prefactor of the area law	not derivable from the two countings: it cancels in γ_L (r07.4)

v4.10.0-r07 ledger and guard update

$$R_{\mathrm{chk}}^{v4100\text{-}r07} = (1, 1, 0)$$

for (r06.5 corrected; chain in frame T for the Maxwell field; area law still an input, prefactor cancels in the ratio of countings).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v4100\text{-}r07,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v4100\text{-}r07} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.10.0-r08 Source-Only: The Assumptions of the Companion and the Dependence of the Open Predictions

Purpose. Dorau and Much (§A.272) start from few assumptions, set openly and motivated independently, and derive one strict consequence. This layer applies that way of working to the companion. First it settles a suspected conflict between the second reading and the photon weights of gravity. Then it lists the assumptions of the companion with their origin, and it traces in QAM bookkeeping what each of the four open predictions rests on. The layer is an inventory; it contains no test and no new prediction. The notes are `notes/research/v4_10_0_r08/`. Earlier layers are not rewritten.
Source-only status. This v4.10.0-r08 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.10.0 Version DOI 10.5281/zenodo.23991357 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.10.0-r08 results

Theoremlet v4.10.0-r08.1 (the source of gravity under the weights). Exploration, analytic. The suspicion was: under the second reading the action is $(a-b)S_M$ (H79), and with photon weights $a = b$ for gravity (v4.10.0-r05.1(b)) the gravitating source of matter would vanish.

(a) The sum $(a - b)$ arises only where both readings live at the same place (domain G1, explored in v4.10.0-r04, not adopted). Under the adopted domain G2 the matter action is $a\,S_M$ after the bang and $-b\,S_M$ on its image before; no sum occurs.

(b) The photon weights concern the gravitational action: $-b_g S_{\mathrm{grav}}$ before, $a_g S_{\mathrm{grav}}$ after the bang, $a_g = b_g$. After the bang $a_g \tilde{G} = a\,\kappa\,T_M$, so $G_{\mathrm{eff}} = (a/a_g)\,G$, one common factor absorbed into the measured G. There is no conflict.

(c) G_{eff} is universal only if all non-gravitational fields carry the same weight a after the bang. Different overall factors for two species would give them different active gravitational masses at equal inertial mass. This assumption (U) is used tacitly in v4.10.0-r03 and is stated nowhere; it is recorded here. The factors of H82 act within one action and change parameters, not universality.

(d) The limit of r07 for massive fields concerns only the domain before the bang: under G2 only the matter reading lives after it, so the chain of r06.1 applies today in frame M to massive and massless fields alike.

Remark v4.10.0-r08.2 (the list of assumptions). The assumptions are listed in `Annahmenliste_Companion.md` with origin and one of six categories: V set before the consequences were known, with a reason of its own; R set afterwards, a selection or repair after a consequence was known or did not fit; K correction of an error; A working assumption; L input from the literature; S tacit. An R assumption is not a flaw, but a prediction resting on it is less blind by that selection. Of the action programme W1/W2 (frozen since v4.10.0-r05), seven entries are R or A within two days; this is the repair series recorded in r05.2. None of them enters the open predictions.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Theoremlet v4.10.0-r08.3 (what the open predictions rest on). Bookkeeping in QAM (`qam_annahmen_vorhersagen.tex`; `qam.sty` raises no error). All four open predictions have status *reading*: their weakest premises are readings of the core, not derivations. Their inputs according to the exported record:

prediction	inputs
H96, $\Omega_M = \beta^2_{\text{sym}}$	γ_{BI}, γ_{string}, counting rule, epoch "today"
H15, m_W	γ_{BI}, γ_{string}, counting rule, m_Z, reading $\cot\theta_W = \gamma_L$
H46, $\sin^2\theta_{23} \in [0.49, 0.50]$	neutrino reading, mask, neutrino data, $\varepsilon = \eta$, γ_{BI}, γ_{string}, counting rule
H42, $\sin^2\theta_{23} \leq 0.54$	neutrino reading, mask, neutrino data

(a) *H15, H46 and H96 rest on the same core*: the two literature values, the assignment of counting rules to the frames (M all spins, T $j = \frac{1}{2}$, U(1) rather than SU(2); v4.9.0-r63, r68) and, through the frame map, the reading of γ_L as Lorentz factor. If the counting rule fails, all three fail together. With the SU(2) counting preferred in the literature ($\gamma_L \approx 2.15$) one would get $m_W = 82.68$ GeV instead of 80.361 GeV and $\Omega_M = 0.365$ instead of 0.302. The U(1) rule was selected by the lepton window (r63) and the m_W relation was noticed afterwards, so these two agreements are not independent of the selection.

(b) H42 and H46 share the neutrino reading (r55, r57); H46 adds $\varepsilon = \eta$ and with it the γ_L core.

(c) Only H42 is independent of the γ_L core. The four open predictions form two independent groups, not four.

(d) Forming γ_L without the declared map from T to M is reported by QAM as frame mixing. γ_L is defined only through the reading it is used to support.

Proof and measurement sources. `Erkundung_Quelle_Gravitation_Gewichte.md` (r08.1); `Annahmenliste_Companion.md` (r08.2, r08.3); `qam_annahmen_H96.tex` and `qam_annahmen_vorhersagen.tex` with exported records. The SU(2) values in r08.3(a) follow from $m_W = m_Z\gamma_L/\sqrt{1+\gamma_L^2}$ and $\Omega_M = (\gamma_L - 1)/(\gamma_L + 1)$. □

Consequence. The most loaded assumption of the companion is the assignment of counting rules to the frames. Deriving it from the frames (OP1 since v4.9.0-r63) would strengthen three open predictions at once; a refutation of one of them points first to that rule. If H15, H46 and H96 are all confirmed, this supports the core more than one hit and less than three independent ones. Assumption U is now explicit.

v4.10.0-r08 status in the three frames

- M: r08.1 concerns the matter reading after the bang; all four predictions are read out in frame M.
- T: enters r08.3 through γ_{string} (counting $j = \frac{1}{2}$) and the map from T to M; under G2 the tachyon reading lives before the bang.
- P: no separate statement.

No frame-free statement is made.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.10.0-r08 register entries

statement	status after v4.10.0-r08
second reading vs. photon weights of gravity	no conflict under G2 (r08.1)
equal weight of all fields after the bang (U)	tacit assumption, now stated
open predictions H15, H42, H46, H96	all status reading; H15, H46, H96 on one γ_L core, H42 independent
most loaded assumption	counting rule per frame (OP1)

v4.10.0-r08 ledger and guard update

$$R_{\mathrm{chk}}^{v4100\text{-}r08} = (1, 1, 0)$$

for (no conflict of weights; dependence of the predictions traced; the counting rule not derived).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v4100\text{-}r08,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v4100\text{-}r08} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.10.0-r09 Source-Only: The Counting Rule per Frame, Its Singular Position (H98) and a Principle

Purpose. Layer v4.10.0-r08.3 found that three of the four open predictions rest on one assumption: the assignment of horizon counting rules to the frames (M counts all spins with weight 2, T counts $j = \frac{1}{2}$; v4.9.0-r63). This layer asks how much the known agreements say about that assignment (pre-registered test H98) and records a principle that selects it (exploration). The notes are `notes/research/v4_10_0_r09/`. Earlier layers are not rewritten.
Source-only status. This v4.10.0-r09 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.10.0 Version DOI 10.5281/zenodo.23991357 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.10.0-r09 results

Theoremlet v4.10.0-r09.1 (H98: the singular position of the pair). Pre-registered (note `Vorab_Notiz_H98_Zaehlregel_Sonderstellung.md`, commit `0cad2d8`); *not blind with respect to the data*: all three values were known, and so was the agreement of the chosen pair. A rule is a spin range and a weight; γ solves $\sum_j g(j)\, e^{-2\pi\gamma a(j)} = 1$ with $a(j) = \sqrt{j(j+1)}$. The fixed rule space S has the spin ranges all j, half-integer j, integer $j \geq 1$, $j = \frac{1}{2}$ only, $j = 1$ only, and the weights 2 and $2j+1$: nine distinct rules, 36 pairs with $\gamma_M > \gamma_T$. The three windows are the lepton window $\beta_{\mathrm{sym}} \in [0.545, 0.570]$ (v4.9.0-r61), m_W within 2σ of PDG 2025 (80.3692 ± 0.0133 GeV, via $\cot\theta_W = \gamma_L$), and Ω_M within 2σ of DESI DR2 BAO (0.2975 ± 0.0086, via $\Omega_M = \beta^2$).

window	pairs inside
lepton window	2
m_W	1
Ω_M	2
all three	1: all spins with weight 2 for M, $j = \frac{1}{2}$ for T (DL/M with ABCK)

Confirmed by the criterion. Values of the pair: $\gamma_L = 1.864700$, $\beta = 0.54941$, $m_W = 80.3615$ GeV (0.6σ), $\Omega_M = 0.3018$ (0.5σ). The SU(2) pair of the literature (weight $2j+1$ for M) gives $\gamma_L = 2.1515$ and misses all three windows.

Reading. Each single window admits other pairs; only m_W selects uniquely. The 36 values of γ_L lie between 1.006 and 3.51; the m_W window is about ± 0.003 wide in γ_L, so about 0.08 pairs would be expected inside it. The weight of the result rests on the m_W relation, which was itself found afterwards (H15 disclosure, look-elsewhere about 0.17). A chance triple agreement is of the order of a few per cent; this is no significance in the strict sense. H98 does not derive the assignment; it shows that among natural rules only this pair agrees with all three known values.

Remark v4.10.0-r09.2 (a principle for the assignment; exploration, not blind). From the two exact lemmas of v4.9.0-r63 a principle P can be stated: a frame counts at a horizon exactly the states that (P1) saturate the uncertainty bound, and (P2) are compatible with its energy condition, where M has all excitation levels and T relaxes to the floor and reaches only the smallest non-zero excitation. Energy and area are linked by the local first law of isolated horizons (Frodden, Ghosh and Perez, 2011): for an observer at fixed distance the energy is proportional to the area. In the rule space S, P1 excludes the weight $2j+1$

© 2026 Marc Brendecke. Preprint — not peer reviewed.

except at $j = \frac{1}{2}$ (where every state saturates, r63.2), P2 gives M all spins and T only $j = \frac{1}{2}$ (smallest area $a(\frac{1}{2}) = \sqrt{3}/2$). P selects exactly the pair confirmed in r09.1.

The counting rule is thus not derived but reduced: instead of a selection by the data (category R in r08.2) it rests on two readings with a reason of their own, uncertainty and energy, together with proved lemmas (r63.2, r63.3) and one literature input (the first law). Both readings came from author remarks after H47 and are not blind. P1 is a reinterpretation: the counting of Meissner counts projections m, not "saturating states" (v4.9.0-r68), and the SU(2) treatment violates P1. The untested bridge is in P2: the particle energy of the tachyon at its floor (r63.3) is identified with the horizon energy seen by an observer. It could be examined through the boost energy at the Rindler horizon, the modular flow of §A.272.

Proof and measurement sources. `k218_regelraum_paare.py` with output (r09.1, sums to $j_{\max} = 4000$); `Ergebnis_H98_Zaehlregel_Sonderstellung.md`; `Erkundung_Zaehlregel_Prinzip.md` (r09.2, analytic on the rule space). □

Consequence. The assignment of counting rules, the most loaded assumption of the companion (r08.3), has a singular position among natural rules with respect to three known values, and a principle with two readings selects it. It is still an assumption. A derivation needs the bridge of P2; a blind test needs a consequence of P other than γ_L.

v4.10.0-r09 status in the three frames

- M: counts all spins with weight 2 (P1, P2: all levels).
- T: counts $j = \frac{1}{2}$ only (P2: floor, smallest area).
- P: carries no counting (v4.9.0-r63).

The rules differ between M and T by construction; no frame-free statement is made.

v4.10.0-r09 register entries

statement	status after v4.10.0-r09
H98 (singular position of DL/M with ABCK)	confirmed, not blind with respect to the data
counting rule per frame (OP1)	reduced to two readings (P1, P2) and the local first law; not derived
bridge in P2 (floor energy vs. horizon energy)	open

v4.10.0-r09 ledger and guard update

$$R_{\text{chk}}^{v4100\text{-}r09} = (1, 1, 0)$$

for (H98 confirmed; principle P recorded; the bridge of P2 not established).

The layer spends no guard:

$$R_{\text{glob}}^{(v4100\text{-}r09,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\text{guard}}^{v4100\text{-}r09} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.10.0-r10 Source-Only: The Bridge of the Counting Principle, and the Counting Rule as a Frame-Dependent Area Quantum

Purpose. Remark v4.10.0-r09.2 left one bridge open: the principle P lets the tachyon frame count only the smallest area excitation because it sits at its floor (v4.9.0-r63.3), and links floor energy to horizon energy through the local first law. This layer examines that bridge. It does not hold in that form; what remains is a sharper statement of the counting rule. A parallel audit on GitHub reached the same verdict (r10.4). Exploration, analytic, no criterion. The notes are `notes/research/v4_10_0_r10/`. Earlier layers are not rewritten.
Source-only status. This v4.10.0-r10 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.10.0 Version DOI 10.5281/zenodo.23991357 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.10.0-r10 results

Theoremlet v4.10.0-r10.1 (the counting equation is an Unruh statistics, and the distance cancels). Analytic. By the local first law (Frodden, Ghosh and Perez, 2013) an observer at proper distance ℓ from the horizon assigns a puncture of spin j, area $8\pi\gamma\, a(j)$ ($\ell_P = 1$), the energy $E_j = \gamma\, a(j)/\ell$, and has the Unruh temperature $T_U = 1/(2\pi\ell)$. Hence

$$E_j/T_U = 2\pi\gamma\, a(j),$$

independent of ℓ (checked for $\ell = 0.5$ to 1000). The counting equation $\sum_j g(j)\, e^{-2\pi\gamma a(j)} = 1$ is this statistics (Ghosh and Perez, 2011). All static observers count alike: a higher or lower energy shifts E_j and T_U in the same ratio.

Theoremlet v4.10.0-r10.2 (the tachyon frame sees the same statistics). By v4.10.0-r07.2 the tachyon map commutes with the boost and fixes the vacuum; an observer in frame T sees the same vacuum with the same modular group, hence the same KMS temperature. The floor of r63.3 (particle energy $E \to 0$ at fixed invariant) is a property of a tachyon particle, not of the state an observer sees at the horizon. **Correction to Remark r09.2:** the bridge in P2 does not hold through the local first law and the Unruh temperature; along that route M and T count alike. "T reaches only the smallest excitation" would need a state of T that is not thermal with respect to the boost; the vacuum is thermal.

Theoremlet v4.10.0-r10.3 (the counting rule as a frame-dependent area quantum). Analytic. If both frames see the same statistics (r10.1, r10.2), the same geometry (r07.2) and both satisfy $S = A/4$ (r06.2), the only remaining place for a difference is the area quantum. The assignment of counting rules to the frames (r08.3, r09) is then equivalent to a frame-dependent Barbero–Immirzi parameter,

$$\gamma_M = \gamma_{\mathrm{BI}}, \qquad \gamma_T = \gamma_{\mathrm{string}}, \qquad \gamma_M/\gamma_T = \gamma_L = 1.864700\,:$$

the smallest area quantum is smaller by the factor γ_L in frame T than in frame M, while the classical area and G are the same. This is v4.9.0-r64.1(b) ("the frame difference sits in γ") in sharper form. The duality rotation alone cannot change an area quantum, since it fixes everything geometric; a reason would have to come from the boost part of the frame map (T as the floor of the mirror frame, r63.4), which only reparametrizes the horizon. No such reason is known.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

Remark v4.10.0-r10.4 (relation to the parallel audit on GitHub). A source reconstruction of OP5 made with another language model (notes `op5_spin_selection_audit_2026_11_06.md`, `op5_common_state_access_2026_11_06.md`, `op5_projector_spin_bridge_2026_11_06.md` in `notes/research/`) reached `main` after layer r09. It agrees with r10.2 and adds two points that bear on r09.2.

(a) Uncertainty saturation alone does not select $j = \frac{1}{2}$: $|1, 1\rangle$ and every $|j, \pm j\rangle$ saturate as well. "Absolute minimum over j" for T and "extremal states at every j" for M are two different rules; saturation is compatible with both and decides between neither. P1 in r09.2 was used only to exclude the weight $2j + 1$, which stands; the selection of $j = \frac{1}{2}$ for T rested on P2.

(b) The tachyon floor $E = \mu/\sqrt{v^2 - 1} \to 0$ does not contain j; without a law linking μ to the spin and a transition that changes j, it gives no relaxation to $j = \frac{1}{2}$. This is the particle side of r10.2.

(c) A passive, invertible change of description cannot remove spin sectors; a T reading restricted to $j = \frac{1}{2}$ is either a restricted readout of the same state (the rest stays in the balance) or a selection with its own weight.

Together: P2 holds neither as an energy condition (r10.2) nor as a relaxation (b); P1 does not select the spin (a). The singular position of r09.1 (H98) is not affected; the assignment remains an assumption, now stated as a frame-dependent area quantum (r10.3).

Proof and measurement sources. `Erkundung_Bruecke_P2_Unruh.md`; the cancellation of ℓ in r10.1 is one line and was checked numerically. □

Consequence. Principle P keeps P1 (uncertainty saturation) as a reading; P2 as an energy condition is not supported along the route of the first law and should be replaced by the plainer statement that each frame has its own area quantum. That is an assumption, not a derivation. The singular position found in H98 is unaffected. The three predictions on the γ_L core (H15, H46, H96) now visibly rest on one assumption: a frame-dependent area quantum with ratio γ_L.

v4.10.0-r10 status in the three frames

- M and T: the same Unruh statistics and the same modular group (r10.1, r10.2); this part is frame free.
- The area quantum differs: $\gamma_M = \gamma_{\mathrm{BI}}$, $\gamma_T = \gamma_{\mathrm{string}}$ (r10.3, assumption).
- P: carries no counting.

v4.10.0-r10 register entries

statement	status after v4.10.0-r10
bridge in P2 via first law and Unruh temperature	does not hold (correction to r09.2)
counting rule per frame	equivalent to a frame-dependent area quantum, $\gamma_M/\gamma_T = \gamma_L$ (assumption)
reason for a frame-dependent area quantum	open

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.10.0-r10 ledger and guard update

$$R_{\mathrm{chk}}^{v4100\text{-}r10} = (1, 1, 0)$$

for (bridge examined; counting rule restated; its reason open).

The layer spends no guard:

$$R_{\mathrm{glob}}^{(v4100\text{-}r10,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\mathrm{guard}}^{v4100\text{-}r10} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

v4.10.0-r11 Source-Only: Integration of the OP5 Source Reconstruction Made with Another Language Model

Purpose. On 6 and 7 November 2026 the author, working with another language model (Codex), traced the origin of OP5 back to the core and brought the result to `main` (merge `4ae32c9`, branch `codex/op5-question-rebuild`): twelve notes `notes/research/op5_*_2026_11_06.md` and `qsc_back_to_basic_idea_2026_11_06.md`, four QAM notes `qam_*_2026_11_0[67].md`, an optional QAM package `qam/tex/qam-physics.sty` with examples and 34 tests, and wiki changes. The work ran in parallel with layers v4.10.0-r06 to r10 and partly examined the same questions; layer r10.4 already cites three of the notes. This layer records the content, checks the load-bearing formulas with independent code, and places the findings against layers r05 to r10, following the pattern of v4.9.0-r69. It contains one author decision made through Codex, two precisions of layer r05 and one reading candidate. No test, no new hypothesis number. The notes are `notes/research/v4_10_0_r11/`. Earlier layers are not rewritten.
Source-only status. This v4.10.0-r11 layer is source-only. It does not build a PDF, does not assign a new DOI, and does not authorize a public release. The v4.10.0 Version DOI 10.5281/zenodo.23991357 and the Concept DOI 10.5281/zenodo.19210728 are unchanged by it.

v4.10.0-r11 results

Decision v4.10.0-r11.1 (author, 6 November, through Codex: the historical OP5 derivation is suspended as a foundation). The historical chain was: spectral carrier → trace formula → metric action → physical bare Λ term → prescribed remainder → cancellation task (OP5, `sec:lambda_cancel`). The step to a well-defined metric action and its variation is missing, and the size of the bare term does not follow (v4.8.0-r03 F3/F4, r04, r05). The author decided to suspend this derivation as the active foundation and to rebuild the question from the independently founded core. New working question: *which independently defined physical quantity can be obtained by an explicit, justified readout of the algebraic core, and which checkable constraint follows, without presupposing the historical Λ cancellation?* The decision is not a refutation of the whole idea; earlier layers, negative results and pre-registered criteria stay. H96 stays open, H97 refuted, W1/W2 frozen. A second precision of the author (7 November, Codex hand-over): the unity of M, T and P was always the hypothesis to be tested, not a presupposed outcome.

Theoremlet v4.10.0-r11.2 (what the reconstruction establishes; checked independently). Analytic; 31 checks with own code, all confirmed.

(a) *Source–return package.* Theorem H7.2 holds as a conditional schema; in parts 21/22 no native route with evaluable A_2, A_3, A_4, class test and readout is instantiated. The published mean $m = (f+b)/2$ loses the branch datum $d = (f-b)/2$: in the shear model on $\mathbb{R}^3$ with $A_2(u,z,y) = (u, z+y, y)$, class u and readout (u,z), the carriers $(1, 2, \pm 1)$ give the same $m = (1,2)$ and forward values $(1,3)$, $(1,1)$. Class conservation with inverse transports does not fix the forward branch.

(b) *Four denominators.* The register ratio $q = \lambda_-/\lambda_+ = \tanh^2 \eta$, the outflow fraction $f_4/(f_4+f_2)$ from S^8 in r31, the contrast $(\gamma_M - \gamma_T)/(\gamma_M + \gamma_T)$ of r67 and Ω_M (matter density over critical density) have different reference quantities. In r31 the outflow fraction returns the

© 2026 Marc Brendecke. Preprint — not peer reviewed.

input q, while the occupation ratio $p_4/p_2 = qr$ depends on the return rate; r31 stabilises q, it does not derive it. The smaller normalised eigenweight is $q/(1+q)$, not q.

(c) *State selection.* The family $E_q = (1+q)^{-1/2}e_x$, $B_q = (q/(1+q))^{1/2}e_y$ has $\operatorname{tr} G = 1$, $E \cdot B = 0$, the same orientation and sector $s_q = (1-q)/(1+q) > 0$, and any ratio $0 < q < 1$; in the σ_x analyser all have $p_\pm = \frac{1}{2}$. The register algebra does not select q. With positive energy weights the minus share is $w_- q/(w_+ + w_- q)$; $f_- = q$ needs $w_-/w_+ = 1/(1-q)$.

(d) *Spin selection.* $j(j+1) - m^2 - |m| = (j-|m|)(j+|m|+1)$: saturation is $|m| = j$ at every j. With weight 2, the spin set $\{\frac{1}{2}\}$ gives $\gamma = \ln 2/(\pi\sqrt{3}) = 0.127384$, the set $\{\frac{1}{2}, 1\}$ gives 0.201784; saturation plus the counting equation fixes no γ until the spin set is fixed.

(e) *Common state and projector bridge.* An invertible change of description cannot remove spin sectors; a restricted readout keeps the unread rest in the balance, a selection has its own weight w; "not accessible" does not imply "zero energy". The chirality projector $C_{\mathrm{ch}} = \frac{1}{2}(1 + iL_{e_7})$ commutes with the tachyon map $z \mapsto iz$ (r51, `Sektoren.lean`); a compatible injective bridge B with $C_{\mathrm{ch}} B = B\Pi$, $U_D B = B U_{\mathrm{spin}}$ passes this on, $\Pi U_{\mathrm{spin}} = U_{\mathrm{spin}} \Pi$. No internal projector yields different M/T spin masks without a declared mask-changing rule.

(f) *Back to the basic idea.* Under duality $z \mapsto e^{i\alpha} z$ the invariant $s = z \cdot z$ goes to $e^{2i\alpha} s$, $|s|$ is kept, $z \mapsto iz$ gives $s \mapsto -s$; a configuration with $s \neq 0$ does not reach $s = 0$ under Lorentz and duality maps. The compensator image $C(z) = \frac{1}{2}(z + i\bar{z}) = (E+B)(1+i)/2$ has $s' = 2i|E'|^2 \neq 0$, it is not null. "One object" has at least four meanings (same carrier, same physical state, states linked by an active operation, different readouts of one state), which must be kept apart.

Proof and measurement sources. `llm/k219_nachrechnung_codex_op5.py`, output `k219_out.txt` (groups A–H), protocol `Integration_Codex_OP5_Rekonstruktion.md`; the commutation in (e) is the existing Lean result. □

Remark v4.10.0-r11.3 (relation to layers r05 to r10). No contradiction. In detail:

(a) *Precision of r05.3(i).* "OP5 (core) locally closed" is to be read as: closed as a conditional schema under the assumptions of Theorem H7.2; a native instance of the route is open (r11.2(a)). The guards are unaffected.

(b) *Precision of r05.3(ii) and of the r05 note on sources.* That note named r31 as the place where the ratio "comes from a dynamics". Read narrowly, the dynamics gives stationarity; the branching number is an input (r11.2(b)). The Λ readout is the additional identification $q = \Omega_M(\text{today})$, as pre-registered in H96. This agrees with r08.2, where K1 ($\Omega_M = \beta^2$ as cascade branching) is already listed as an assumption.

(c) *r08.3, r09.1.* The γ_L core of H15, H46, H96 is the same chain; r11.2(b), (c) add the denominators and $q \neq \lambda_-$. The freedom of r11.2(d) is exactly the rule space S of H98; the singular position found there is unaffected.

(d) *r10.* r11.2(d), (e) were taken up in r10.4. The projector bridge confirms r10.3 from the internal side: as the duality rotation fixes every geometric quantity, the tachyon map fixes the internal sectors; neither yields a frame-dependent mask or area quantum by itself.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

(e) *r03.2, r22, r24.* r11.2(f) restates known results (the compensator image of a light wave is not a light wave; P is the degenerate fixed point; R ≠ P) and adds the shell argument.

Remark v4.10.0-r11.4 (QAM as frame interface language). The package `qam-physics.sty` declares physics profiles (for example GR and QM), nested default contexts, implicit or explicit left roles, an exported formula payload, and a conversion between profiles only with a declared map and an explicit target formula; 34 software cases and ten exports pass, and the existing Lean checker accepts the records. These are bookkeeping tests, not physics tests. The author's precision: QAM is a frame interface language; GR and its physics mapping hold in the frame assigned to them and are not carried to other frames automatically. Relation: v4.10.0-r07.2 is such a declared conversion (map $z \mapsto iz$, commuting with $SO(3,\mathbb{C})$; target formula the same Einstein equation); the Dorau–Much example does not identify the Rindler observer with M, T or P, as in r07.3. In this language r10.3 says that the T profile differs from the M profile by its declared area quantum, $\gamma_T = \gamma_M/\gamma_L$, and that no map is known that produces it.

Remark v4.10.0-r11.5 (two observations from the r27 notes; one reading candidate).

(a) With $U_{\pm\alpha}$ on the Bloch disc, the direction mean is $m = \cos 2\alpha\,(x,z)$ and $(Q(f) + Q(b))/2 = Q(m) + Q(d)$ for the second moment $Q(v) = vv^{\mathsf{T}}$. At $v = (1,0)$, $\alpha = \pi/6$ the trace is 1 before and 1/4 after averaging; the missing $3/4 = \sin^2(\pi/3)$ is the squared direction difference. This is the factor 3/4 of r27 and of the reading "rotate by $\pi/6$, then compensate" (r10, r11); consistent, not new.

(b) For a non-null field with flux $\mathbf{P} = E \times B$, the observers at rapidity 0 and $2\eta_{\mathrm{ref}}$ along $\mathbf{P}$ see the same field scale and opposite fluxes; the flux-free profile lies in the middle, and $1/r = |s|^{-1}n = \cosh 2\eta_{\mathrm{ref}}$ is the ordinary Lorentz factor between the two (example $E = 2e_x$, $B = e_y$: $v = \frac{1}{2}$ flux free, $v = \frac{4}{5}$ gives $E' = 2e_x$, $B' = -e_y$). At $r = 1/\gamma_L$ the boost to the flux-free profile has $v = \beta_{\mathrm{sym}}$ (r17). *Reading candidate, a posteriori:* r10.3 asks for a reason from the boost part of the frame map; the pair of flux-opposed observers at relative Lorentz factor γ_L is a concrete place where γ_L is an ordinary Lorentz factor. Codex states expressly that this does not identify the r27 step with an observer change, and nothing here identifies the pair with M and T.

Consequence. The two lines of work agree. OP5 in its historical form (cancellation of a bare Λ) is no longer an active task; what remains is the conditional source–return schema (no native instance) and the additional identification $q = \Omega_M$ of H96, which stays a pre-registered hypothesis without a derivation of the bridge. The open points are the same as after r10: the reason for the counting rule per frame (a frame-dependent area quantum), a mechanism selecting q, and a measurement map from register to density. Codex proposes the meaning of "one object" as the next question before any further spin or counting branch.

v4.10.0-r11 status in the three frames

- M and T: the shell argument (r11.2(f)) links $s > 0$ and $s < 0$ of equal modulus by duality; the internal sectors are the same in both frames (r11.2(e)); frame free.
- P: not on the orbit of $s \neq 0$; the compensator image is not P. R ≠ P (r24); γ_L compares M with R.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

- Conversions between frame profiles need a declared map and target formula (r11.4); the counting rule per frame remains an assumption (r10.3).

v4.10.0-r11 register entries

statement	status after v4.10.0-r11
historical OP5 cancellation task	suspended as foundation (author, 6 November)
OP5 (core), r05.3(i)	conditional schema closed; native route open (precision)
Λ readout, r05.3(ii)	additional identification $q = \Omega_M$ (H96, open); r31 stabilises, does not derive (precision)
selection of q by the register algebra	none (explicit family)
internal projector as M/T spin mask	no (commutes with the tachyon map)
γ_L as Lorentz factor of flux-opposed observers	algebraic fact; link to r10.3 a reading candidate

v4.10.0-r11 ledger and guard update

$$R_{\text{chk}}^{v4100\text{-}r11} = (1, 1, 0)$$

for (reconstruction integrated; formulas checked; native route and selection of q open).

The layer spends no guard:

$$R_{\text{glob}}^{(v4100\text{-}r11,+)} = (0, 1, 1, 1, 1), \qquad \Gamma_{\text{guard}}^{v4100\text{-}r11} = (1, 1, 1, 1, 1), \qquad \beta_4 = 1/256 \text{ unspent.}$$

It does not imply exact finite $C5$ zero closure, exact all-readout terminality, $C6/T2$, a global photon-clock/RH theorem, or global all-readout closure.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

References

[1] Marc Brendecke, *On the Riemann Hypothesis: A Kernel-Theoretic Approach via Tensor Contraction and Self-Adjoint Extension*, Zenodo historical release line, March 2026. DOI: 10.5281/zenodo.19157158.

[2] Marc Brendecke, *Quantum Sphaera: A Cascade Framework for Quantum Gravity, the Standard Model, and the Riemann Hypothesis*, Zenodo, March 2026. DOI: 10.5281/zenodo.19210213.

[3] Marc Brendecke, *Quantum Sphaera Companion: A Structure-First Mathematical Unfolding* (archival stage pre-shift QAM reindexing, two-sided tensor-notation, and mixed-readout release), April 2026. Concept DOI: 10.5281/zenodo.19210728; version-specific DOI: 10.5281/zenodo.19750674; prior public version DOI: 10.5281/zenodo.19746821.

[4] Yilwook Kim, *GUE-pacing and Energy Highway in the Riemann Lattice*, Zenodo, March 2026. Concept DOI: `10.5281/zenodo.19112361`.

[5] B. Riemann, *Über die Anzahl der Primzahlen unter einer gegebenen Grösse*, Monatsberichte der Berliner Akademie, November 1859, 136–144.

[6] H. M. Edwards, *Riemann's Zeta Function*, Academic Press, New York, 1974.

[7] E. C. Titchmarsh, *The Theory of the Riemann Zeta-Function*, 2nd ed., revised by D. R. Heath-Brown, Oxford University Press, 1986.

[8] J. B. Conrey, *The Riemann Hypothesis*, Notices of the American Mathematical Society **50** (2003), no. 3, 341–353.

[9] H. L. Montgomery, *The Pair Correlation of Zeros of the Zeta Function*, Proc. Sympos. Pure Math. **24** (1973), 181–193.

[10] A. M. Odlyzko, *On the Distribution of Spacings Between Zeros of the Zeta Function*, Mathematics of Computation **48** (1987), no. 177, 273–308. DOI: 10.1090/S0025-5718-1987-0866115-0.

[11] M. L. Mehta, *Random Matrices*, 3rd ed., Pure and Applied Mathematics, vol. 142, Elsevier/Academic Press, 2004.

[12] N. M. Katz and P. Sarnak, *Zeroes of Zeta Functions and Symmetry*, Bulletin of the American Mathematical Society **36** (1999), no. 1, 1–26. DOI: 10.1090/S0273-0979-99-00766-1.

[13] L. V. Ahlfors, *Lectures on Quasiconformal Mappings*, 2nd ed., University Lecture Series, vol. 38, American Mathematical Society, Providence, RI, 2006. DOI: 10.1090/ulect/038.

[14] F. P. Gardiner and N. Lakic, *Quasiconformal Teichmüller Theory*, Mathematical Surveys and Monographs, vol. 76, American Mathematical Society, Providence, RI, 2000.

[15] J. H. Hubbard, *Teichmüller Theory and Applications to Geometry, Topology, and Dynamics, Volume 1: Teichmüller Theory*, Matrix Editions, Ithaca, NY, 2006.

[16] D. Sullivan, *Quasiconformal Homeomorphisms and Dynamics I. Solution of the Fatou–Julia Problem on Wandering Domains*, Annals of Mathematics **122** (1985), no. 3, 401–418. DOI: 10.2307/1971308.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

[17] J. Milnor, *Dynamics in One Complex Variable*, 3rd ed., Annals of Mathematics Studies, vol. 160, Princeton University Press, 2006.

[18] H. Hopf, *Über die Abbildungen der dreidimensionalen Sphäre auf die Kugelfläche*, Mathematische Annalen **104** (1931), 637–665. DOI: 10.1007/BF01457962.

[19] N. Steenrod, *The Topology of Fibre Bundles*, Princeton Mathematical Series, vol. 14, Princeton University Press, 1951.

[20] A. Hatcher, *Algebraic Topology*, Cambridge University Press, 2002.

[21] R. L. Bryant, *Metrics with Exceptional Holonomy*, Annals of Mathematics **126** (1987), no. 3, 525–576. DOI: 10.2307/1971360.

[22] T. Eguchi, P. B. Gilkey, and A. J. Hanson, *Gravitation, Gauge Theories and Differential Geometry*, Physics Reports **66** (1980), no. 6, 213–393. DOI: 10.1016/0370-1573(80)90130-1.

[23] J. C. Baez, *The Octonions*, Bulletin of the American Mathematical Society **39** (2002), no. 2, 145–205. DOI: 10.1090/S0273-0979-01-00934-X.

[24] G. M. Dixon, *Division Algebras: Octonions, Quaternions, Complex Numbers and the Algebraic Design of Physics*, Mathematics and Its Applications, vol. 290, Kluwer Academic Publishers, Dordrecht, 1994. DOI: 10.1007/978-1-4757-2315-1.

[25] C. Furey, $SU(3)_C \times SU(2)_L \times U(1)_Y$ $(\times U(1)_X)$ *as a Symmetry of Division Algebraic Ladder Operators*, European Physical Journal C **78** (2018), 375. DOI: 10.1140/epjc/s10052-018-5844-7.

[26] P. A. M. Dirac, *The Quantum Theory of the Electron*, Proceedings of the Royal Society of London A **117** (1928), no. 778, 610–624. DOI: 10.1098/rspa.1928.0023.

[27] C. N. Yang and R. L. Mills, *Conservation of Isotopic Spin and Isotopic Gauge Invariance*, Physical Review **96** (1954), 191–195. DOI: 10.1103/PhysRev.96.191.

[28] F. Englert and R. Brout, *Broken Symmetry and the Mass of Gauge Vector Mesons*, Physical Review Letters **13** (1964), 321–323. DOI: 10.1103/PhysRevLett.13.321.

[29] P. W. Higgs, *Broken Symmetries and the Masses of Gauge Bosons*, Physical Review Letters **13** (1964), 508–509. DOI: 10.1103/PhysRevLett.13.508.

[30] N. Cabibbo, *Unitary Symmetry and Leptonic Decays*, Physical Review Letters **10** (1963), 531–533. DOI: 10.1103/PhysRevLett.10.531.

[31] M. Kobayashi and T. Maskawa, *CP-Violation in the Renormalizable Theory of Weak Interaction*, Progress of Theoretical Physics **49** (1973), no. 2, 652–657. DOI: 10.1143/PTP.49.652.

[32] Z. Maki, M. Nakagawa, and S. Sakata, *Remarks on the Unified Model of Elementary Particles*, Progress of Theoretical Physics **28** (1962), 870–880. DOI: 10.1143/PTP.28.870.

[33] M. H. Stone, *Linear Transformations in Hilbert Space, III: Operational Methods and Group Theory*, Proceedings of the National Academy of Sciences of the USA **16** (1930), 172–175.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

[34] J. von Neumann, *Die Eindeutigkeit der Schrödingerschen Operatoren*, Mathematische Annalen **104** (1931), 570–578.

[35] A. Weil, *Sur certains groupes d'opérateurs unitaires*, Acta Mathematica **111** (1964), 143–211. DOI: 10.1007/BF02391012.

[36] M. Reed and B. Simon, *Methods of Modern Mathematical Physics II: Fourier Analysis, Self-Adjointness*, Academic Press, New York, 1975.

[37] R. V. Kadison and J. R. Ringrose, *Fundamentals of the Theory of Operator Algebras, Volume I: Elementary Theory*, Graduate Studies in Mathematics, vol. 15, American Mathematical Society, 1997.

[38] E. Zermelo, *Untersuchungen über die Grundlagen der Mengenlehre. I*, Mathematische Annalen **65** (1908), 261–281. DOI: 10.1007/BF01449999.

[39] A. A. Fraenkel, *Einleitung in die Mengenlehre*, 3rd ed., Springer, Berlin, 1928. DOI: 10.1007/978-3-662-42029-4.

[40] T. Jech, *Set Theory: The Third Millennium Edition, Revised and Expanded*, Springer Monographs in Mathematics, Springer, Berlin, 2003.

[41] K. Kunen, *Set Theory: An Introduction to Independence Proofs*, Studies in Logic and the Foundations of Mathematics, vol. 102, North-Holland, Amsterdam, 1980.

[42] A. M. Turing, *On Computable Numbers, with an Application to the Entscheidungsproblem*, Proceedings of the London Mathematical Society (2) **42** (1936), 230–265.

[43] F. W. Kasiski, *Die Geheimschriften und die Dechiffrir-Kunst: Mit besonderer Berücksichtigung der deutschen und der französischen Sprache*, E. S. Mittler und Sohn, Berlin, 1863.

[44] R. L. Rivest, A. Shamir, and L. Adleman, *A Method for Obtaining Digital Signatures and Public-Key Cryptosystems*, Communications of the ACM **21** (1978), no. 2, 120–126. DOI: 10.1145/359340.359342.

[45] W. Diffie and M. E. Hellman, *New Directions in Cryptography*, IEEE Transactions on Information Theory **22** (1976), no. 6, 644–654. DOI: 10.1109/TIT.1976.1055638.

[46] J. R. Norris, *Markov Chains*, Cambridge Series in Statistical and Probabilistic Mathematics, vol. 2, Cambridge University Press, 1997.

[47] P. Glasserman, *Monte Carlo Methods in Financial Engineering*, Stochastic Modelling and Applied Probability, vol. 53, Springer, New York, 2004. DOI: 10.1007/978-0-387-21617-1.

[48] T. Bollerslev, *Generalized Autoregressive Conditional Heteroskedasticity*, Journal of Econometrics **31** (1986), no. 3, 307–327. DOI: 10.1016/0304-4076(86)90063-1.

[49] D. E. Knuth, *The Art of Computer Programming, Vol. 1: Fundamental Algorithms*, 3rd ed., Addison–Wesley, 1997.

[50] Raúl Rojas, *How to Make Zuse's Z3 a Universal Computer*, IEEE Annals of the History of Computing **20** (1998), no. 3, 51–54.

[51] B. J. Copeland (ed.), *Colossus: The Secrets of Bletchley Park's Codebreaking Computers*, Oxford University Press, 2006.

© 2026 Marc Brendecke. Preprint — not peer reviewed.

[52] H. Zuse, *Konrad Zuse's Computer Z3*, historical overview, 2016.

[53] A. J. Menezes, P. C. van Oorschot, and S. A. Vanstone, *Handbook of Applied Cryptography*, CRC Press, 1996.

[54] S. Goldwasser and M. Bellare, *Lecture Notes on Cryptography*, MIT/UCSD lecture notes, 2008.

[55] M. E. Peskin and D. V. Schroeder, *An Introduction to Quantum Field Theory*, Westview Press, 1995.

[56] S. Weinberg, *The Quantum Theory of Fields, Volume I: Foundations*, Cambridge University Press, 1995.

[57] W. Fulton and J. Harris, *Representation Theory: A First Course*, Graduate Texts in Mathematics, vol. 129, Springer, 1991.

[58] M. Nakahara, *Geometry, Topology and Physics*, 2nd ed., Institute of Physics Publishing, 2003.

[59] A. L. Besse, *Einstein Manifolds*, Ergebnisse der Mathematik und ihrer Grenzgebiete, vol. 10, Springer, 1987.

[60] D. D. Joyce, *Compact Manifolds with Special Holonomy*, Oxford University Press, 2000.

© 2026 Marc Brendecke. Preprint — not peer reviewed.