

“SATELLITE” Documentation

Dr. Stavros Akras

Institute for Astronomy, Astrophysics, Space Applications and Remote Sensing,
National Observatory of Athens
September 2021

1 Introduction

The Spectroscopic Analysis Tool for intEgraL fieLd unIt daTacubEs (SATELLITE) is a newly developed code for the spectroscopic characterization of extended photo-ionised nebulae such as planetary nebulae, H II regions or galaxies in the optical regime observed with any integral field unit (IFU). SATELLITE has been written in PYTHON and was developed to be an automatic and user-friendly code. The user does not need any programming or coding knowledge.

The capabilities and performance of the SATELLITE code (v0.1) were first presented in [1] using the IFU data of the Abell 14 PN obtained with VIMOS@ESO. SATELLITE v1.2 has been applied to three more PNe: Hen 2-108 (VIMOS) (Miranda Marques et al. 2021, submitted), NGC 7009 (MUSE), and NGC 6778 (MUSE) (Akras et al. 2021, submitted).

SATELLITE carries out a spectroscopic analysis of extended ionized nebulae through 1D and 2D approach on a list of 35 emission line (the brightest and more frequently detected in ionized nebulae) via a number of pseudo-slits that simulate slit spectrometry and 2D emission line imaging. The analysis is performed in four different modules:

- (I) rotation analysis,
- (II) radial analysis,
- (III) specific slits analysis and
- (IV) 2D analysis

For each module, SATELLITE computes all the typically used nebular parameters and their uncertainties using as input information the available emission lines maps. For all four modules, the uncertainties of the line intensities as well as those of the nebular parameters are computed following a Monte Carlo approach considering a number of spectra.

- extinction coefficient ($c(\text{H}\beta)$),
- electron temperatures and densities for different diagnostic lines,
- ionic and elemental abundances, abundances ratio relative to oxygen and the ionization correction factors,
- emission line ratios from a pre-defined list

The input parameters necessary to run the code are provided by the user in four ASCII files:

- *input.txt*
- *numerical_input.txt*
- *output.txt* (*the requested outputs from the code!*)
- *diagnostic_diagrams_input.txt*
- *plots_parameters_input.txt*

and they are:

- emission line and error maps (or additional error as percentage of its pixel flux.)
- the pixel scale of the IFU
- the interstellar extinction law
- number of replicate spectra
- the width, length, position angle (PA) and coordinates of the pseudo-slits for the 1D analysis
- the coordinates of the central star or the centre of the nebula
- atomic data
- Te and Ne from different diagnostic lines for the proper calculation of the ionic abundances
- the line ratios that the code will compute
- the emission line diagnostic diagrams that the code will construct
- the module or modules that the code will execute

The philosophy behind the development of the SATELLITE code is, besides the unique 2D imaging spectroscopy that IFU technology provides, to carry out a detailed 1D spectroscopic analysis through a number of pseudo-slits that simulate slit spectrometry and emission line imaging in order to properly compare the results presented in previous studies. The outputs from each of SATELLITE's modules are saved in five different folders:

- `output_angles_plots` ← rotation analysis module
- `output_Diagnostic_Diagrams` ← 2D analysis module
- `output_images` ← 2D analysis module (& slit position testing)
- `output_plots` ← 2D analysis module
- `output_radial_plots` ← radial analysis module

The description of each module as well as the input parameters and outcomes are described in more details in the following sections using as a representative example the analysis of the planetary nebulae NGC 7009 and its MUSE data from the Science Verification phase [5, 6].

2 Set up and Run the SATELLITE Code

SATELLITE has been successfully run/tested in Fedora 23 operation systems and PYTHON version 2.7.11.

To use the SATELLITE code, it is also necessary to install a number of libraries such as MATPLOTLIB, NUMPY, SCIPY, ASTROPY, and SEABORN libraries.

The version of the libraries was: matplotlib v2.2.5, numpy v1.11.1 and scipy v0.18.0, astropy v2.0.12, and seaborn v0.9.1. SATELLITE also make use of the PYNEB package version 1.1.15 [4].

The installation of a library can be done via *pip* and the command: *pip install library*.

To get the version of the installed libraries add the following lines in a PYTHON script and run it.

```
import matplotlib
import numpy
import seaborn
import astropy
import scipy

print(matplotlib.__version__)
print(numpy.__version__)
print(seaborn.__version__)
print(astropy.__version__)
print(scipy.__version__)
```

For the atomic data from the Chianti database, it is necessary to download the data from the website <https://www.chiantidatabase.org/>.

2.1 GitHub repository

1) Download SATELLITE from GitHub (<https://github.com/StavrosAkras/SATELLITE>)

command: “**git clone https://github.com/StavrosAkras/SATELLITE.git** ”

A new folder will be created namely SATELLITE with all the necessary files.

2) run setup.py in the satellite folder

command: “**python setup.py install** ”

Setup.py also installs all the necessary python packages need to use SATELLITE included PYNEB.

3) run the code command: “**./satellite > outputLog.txt.** ”

A number of general comments provided from the different modules of the code are written in the *outputLog.txt* ASCII file.

3 READ INPUT DATA

The number of input data that the user has to provide the code.

3.1 read_input_script

In the current version of SATELLITE, there is a pre-defined list of 35 emission lines from which the user can select the emission line that will be used by the code for the analysis. The list contains the most commonly detected and relatively bright emission line in ionized nebula such as the H and He recombination lines (recombination lines from O or N will be implemented to future versions) and collisionally excited lines from O+, O++, N+, N++, S+, S++, etc. (see Figure 1).

The user can access the list from the *input.txt* file and just add “yes” or “no” in the second column for both flux and error maps. The first column in the *input.txt* lists the name of the emission lines. **The input “FIT files” of the emission line map must have the same name.** Moreover, all the input “FIT files” (maps) must be located in the *image_data* folder. **Hint:** In case error maps are not available, the user can create some fake error maps by multiplying the flux maps by e.g. 0.1 to replicate an uncertainty of 10 percent for each emission line. The option of an extra error is also possible. In the forth column, the user can add an extra error for each line as a percentage of the flux (e.g. 1% of the total flux of [O III] 5007Å line). The even rows for each line (error lines) can take two numerical values “0” and “any integer”. (i) “0” means that the final error for each pixel and each emission line is equal to the percentage of the flux, and (ii) “any number” means that the final error for each pixel and each emission line is equal to the error from the error map + the percentage of the flux.

The third column in the *input.txt* deals with the *radial_analysis* module and the user adds, “radial_yes” or “radial_no”, for the lines(with their error maps) are available and will be used for this module. **Note:** The H α and H β lines must always be selected for the determination of the interstellar extinction coefficient and corrected line intensities.

The READ_INPUT_SCRIPT reads the flux and error maps of each line defined in the *input.txt* file pixel-by-pixel and save the values in 2D arrays.

For the example of NGC 7009, the selected emission lines are presented in Figure. 1

3.2 read_input_lines_parameters_script

Besides the selection of the emission line and error maps, there is also a list of numerical parameters that the code needs to run properly and they are provided by the user in different files. All these parameters are read by the code via the READ_INPUT_LINES_PARAMETERS_SCRIPT. Below, the most important parameters are listed together with the ascii files:

- the pixel scale of the IFU in arcsec ($\times 10^2$) \rightarrow *numerical_input.txt*
- the interstellar extinction law ($R \times 10^1$) \rightarrow *numerical_input.txt*
- atomic data \rightarrow *numerical_input.txt*
- the width, length, position angle (PA) and coordinates of the pseudo-slits for the 1D analysis in the *rotation_analysis*, *radial_analysis*, and *specific_slits_analysis* modules \rightarrow *numerical_input.txt*
- the coordinates of the central star or the centre of the nebula \rightarrow *numerical_input.txt* file

HI_6563s	yes	radial_yes	1
HI_6563e	yes	radial_no	0.01
HI_4861s	yes	radial_yes	1
HI_4861e	yes	radial_no	0.01
HI_4340s	no	radial_no	1
HI_4340e	no	radial_no	0
HI_4101s	no	radial_no	1
HI_4101e	no	radial_no	0
HeI_5876s	yes	radial_yes	1
HeI_5876e	yes	radial_no	0.05
HeI_6678s	yes	radial_yes	1
HeI_6678e	yes	radial_no	0.05
HeII_5412s	yes	radial_yes	1
HeII_5412e	yes	radial_no	0.05
HeII_4686s	no	radial_no	1
HeII_4686e	no	radial_no	0
N2_5755s	yes	radial_yes	1
N2_5755e	yes	radial_no	0.1
N2_6548s	yes	radial_yes	1
N2_6548e	yes	radial_no	0.05
N2_6583s	yes	radial_yes	1
N2_6583e	yes	radial_no	0.05
N1_5199s	yes	radial_yes	1
N1_5199e	yes	radial_no	0.10
O3_4363s	no	radial_no	1
O3_4363e	no	radial_no	0
O3_4959s	yes	radial_yes	1
O3_4959e	yes	radial_no	0.01
O3_5007s	yes	radial_yes	1
O3_5007e	yes	radial_no	0.01
O2_3727s	no	radial_no	1
O2_3727e	no	radial_no	0
O2_3729s	no	radial_no	1
O2_3729e	no	radial_no	0
O2_7320s	yes	radial_yes	1
O2_7320e	yes	radial_no	0.1
O2_7330s	yes	radial_yes	1
O2_7330e	yes	radial_no	0.1
O1_5577s	no	radial_yes	1
O1_5577e	no	radial_no	0
O1_6300s	yes	radial_yes	1
O1_6300e	yes	radial_no	0.1
O1_6363s	no	radial_yes	1
O1_6363e	no	radial_no	0
S2_6716s	yes	radial_yes	1
S2_6716e	yes	radial_no	0.05
S2_6731s	yes	radial_yes	1
S2_6731e	yes	radial_no	0.05
S3_6312s	yes	radial_yes	1
---	---	---	---

Figure 1: Example of the *input.txt* file from which the user can select the emission line that will be used for 1D spectroscopic analysis or radial analysis.

- number of replicate spectra for the determination of the uncertainties → *numerical_input.txt* file
- the minimum radius from which the maximum line flux will be determined and the profiles will be normalized to 1. → *numerical_input.txt* file
- the number of column and row pixels that have to be added to the raw maps in order to put the centre of the nebula at the center of the map → *numerical_input.txt*
- the module or modules that the code will execute → *outputs.txt* file
- the emission line ratios that the code will compute based on the available line maps → *outputs.txt*
- the physical parameters (c , T_e , N_e) that the code will compute based on the available line maps → *outputs.txt*
- the T_e and N_e from specific diagnostic lines that will be used to compute ionic abundances → *outputs.txt*
- the elemental abundances and ICFs that the code will compute based on the available line maps → *outputs.txt*
- the diagnostic diagrams that the code will construct → *diagnostic_diagrams_input.txt*
- the xmin/xmax and ymin/ymax for the diagnostic diagrams → *diagnostic_diagrams_input.txt*
- if the user wants to over-plot the selection criteria from Kewley2001 and Kauffmann2003 → *diagnostic_diagrams_input.txt*
- ymin/ymax for the scatter plots → *plots_parameters_input.txt* file

3.3 Reorder line flux and error maps

The first task the user must perform before run SATELLITE is to compute the number of columns and rows of pixels that need to be added to the raw line flux and error maps in order to put the central star or centre of the nebula at the centre of the new map. This task is very important in order to properly rotate the maps and measure the line fluxes in each pseudo-slit. **Note: The final maps have to have the same number of columns and rows!**

Figure 2 presents a cartoon that illustrates the numbers of rows of pixels need to be added to the top and bottom of the raw map as well as the columns of pixels to the left and right parts of the raw maps. The value these extra pixels have is **zero** and they do not have any impact on the spectroscopic analysis. The extra row and columns added in the flux and error maps are provided by the user in the *numerical_input.txt* file.

CM (15,15) is the centre of the raw map and CS (19,12) is the central star of an ellipsoidal planetary nebula (orange colour). The offset between the CS and CM is 4 pixels in row and 3 pixels is column. It is necessary to move the CS to the left and up by adding some extra columns and rows. So, 8 columns are added to the right side and the CS is now in the middle (19,19) and 6 rows at the bottom so the CS is now at the position (18,18). But the map still does not have the same number of columns and rows (38,36). For this reason, we have to add 1 extra row to the top and 1 extra to the bottom. Finally, the CS is the centre of the new map at position (19,19). These numbers are added in the *numerical_input.txt* file as it is show below:

- add_pixels_above=1




Figure 2: Illustrative example of a line flux map and the rows/columns that have to be added in order to coincide the CS with the CM of the new map.

- `add_pixels_below=7`
- `add_pixels_left=0`
- `add_pixels_right=8`
- `total_num_pixels_verti=30`
- `total_num_pixels_horiz=38`

As for NGC 7009, the number of rows to the top and bottom of the raw maps are 35 and 35, respectively, and the extra columns to the left and right side are 4 and 6, respectively.

3.4 Extinction Coefficient

SATELLITE determines the extinction coefficient ($c(H\beta)$) using the PyNEB package [4] (see: https://github.com/Morisset/PyNeb_devel/blob/master/docs/Notebooks/PyNeb_manual_5.ipynb) and constructs the 2D map and

its error map. All the extinction law available in the the PYNEB can be selected ('No correction', 'CCM89', 'CCM89_Bal07', 'CCM89_oD94', 'S79_H83_CCM89', 'K76', 'SM79_Gal', 'G03_LMC', 'MCC99_FM90_LMC', 'F99', 'F88_F99_LMC']). The R parameter is also given by the user (e.g. R=3.1) in the *numerical_input.txt* file as an integer number (R*10=031). The SAVE_FITSIMAGES_SCRIPT saves the 2D arrays as FITS image.

The code does not take into account all the pixels in the pseudo-slits or estimates the extinction coefficient for all the pixels **BUT** only for those that satisfy the following criteria:

- $F(H\alpha) > 0$,
- $F(H\beta) > 0$
- $F(H\alpha) > F(H\beta) * 2.86$
- **otherwise a value "zero" is applied.**

There is also an option to find the outliers and exclude them from the calculation of the mean/median values. The outliers are defined as those pixels with values that do not satisfy the criteria: $\text{value} > \text{per25} + 1.5 * \text{iqr}$ and $\text{value} < \text{per75} - 1.5 * \text{iqr}$, where per25 and per75 are the 25% and 75% percentiles, respectively, and $\text{iqr} = \text{per75} - \text{per25}$ the inter-quartile-range. These calculations are made in the CALCULATIONS_EXCLUDING_OUTLIERS_SCRIPT. In the current version, the outliers are not excluded from the physical parameters that the code computes.

3.5 Atomic Data

The atomic data can also be changed by the user. All the options available in the PYNEB package can be used by adding in the *numerical_input.txt* the words: IRAF_09_orig, IRAF_09, PYNEB_13_01, PYNEB_14_01, PYNEB_14_02, PYNEB_14_03', 'PYNEB_16_01, , PYNEB_17_01, PYNEB_17_02, PYNEB_18_01, PYNEB_20_01, and PYNEB_21_01. The atomic data from Chianti group can also be used (adding the word "Chianti" in the *numerical_input.txt*) but they have to be downloaded from the website (http://www.chiantidatabase.org/chianti_download.html). For more information the user must refer to the website of PyNeb package (https://github.com/Morisset/PyNeb_devel/blob/master/docs/Notebooks/PyNeb_manual_3.ipynb)

4 SATELLITE's Modules

The modules that the code executes are defined in the second column ("yes" or "no") of the *outputs.txt* file. Each module is described below. **Note:** It is recommended to run the RADIAL_ANALYSIS module separately from the other three modules.

4.1 rotation_analysis module

The *rotation_analysis* module deals with the spectroscopic analysis of a number of radial placed pseudo-slits from the centre to outer parts with position angles (PA) between 0 and 360. Figure 3 illustrates as example the position of these pseudo-slits on the [N II] 6584 emission line map of NGC 7009. The minimum and maximum values of the PA, the step in PA, the width and the length of the pseudo-slits are provided by the user in the file *numerical_input.txt*.

The code first computes the integrated $H\beta$ flux and line fluxes in the ROTATE_LINE_FLUXES_SCRIPT. Then, line intensities (relative to $H\beta$ and corrected for the interstellar excitation) as well as all the nebular parameters defined




Figure 3: An illustrative image of the radial positioned pseudo-slits with PA from 0 to 360 every 20 degrees overlaid the [N II] 6584 image of NGC 7009.

by the user in the *output.txt* file for all the pseudo-slits are computed by the PYNEB package. This analysis allows to explore the variation of line intensities, line ratios and physical parameters (T_e , N_e), chemical abundances as functions of PA.

The outcomes from this module are multiple and are saved in different files:

- an ASCII file with the $c(H\beta)$ and the intensity of each emission line and for each PA \rightarrow *output_linesintensities_per_angles.txt*
- an ASCII file with various line ratios defined by the user for each PA \rightarrow *output_lineratios_per_angles.txt*
- an ASCII file with T_e and N_e defined by the user for each PA \rightarrow *PyNeb_output_Te_and_Ne_per_angles.txt*
- an ASCII file with the ionic abundances for each ion defined by the user for each PA \rightarrow *PyNeb_output_total_abund_ICFs_per_angles.txt*
- an ASCII file with the elemental abundances and ICFs for each element defined by the user for each PA \rightarrow *PyNeb_output_ionic_abund_per_angles.txt*
- plots of $c(H\beta)$, T_e , N_e , ionic, elemental abundances, ICFs and abundances ratios as function of the PA \rightarrow *output_angles_plots* folder

The user can use the ASCII files to carry out any further analysis and/or construct his/her own proper plots.

Figure 4 and 5 present the plots of $c(H\beta)$, T_e , N_e , ionic, elemental abundances and ICFs of N as functions of the position angle of the pseudo-slits for the analysis of NGC 7009. T_e and N_e are shown in the same plot as in Figures 4 and 5 or in two separate plots.




Figure 4: Representative plot of $c(H\beta)$ (upper panel), T_e and N_e for two different diagnostic lines (middle/lower panels) as function of the PA of the pseudo-slits.




Figure 5: Representative plot of the ionic/total abundance of N (upper panel), the ICF(N) (middle panel) and the N/O ratio as functions of the pseudo-slits' PA.

4.1.1 slit_line_flux_script

This script calculates the flux for each emission line along a pseudo-slit with a specific position angle, width and length given by the user in the *numerical_input.txt* file. **Note1:** The width should always be an integer number. The pseudo-slit starts from the centre of the image or of the nebula, and it covers only one half of the nebula. The code sums up the values of all the pixels within the area defined by the width and length of the pseudo-slit, except those pixels which have $F(Ha) < 0$, $F(Hb) < 0$ and/or $F(Ha) < F(Hb) * 2.86$ (negative or unrealistic $c(Hb)$).

For the calculations, the script first rotates the entire image/table by a given angle, then calculates the new size (x,y) of the rotated image as well as the center of the new image. The pseudo-slit is always oriented along the up-down direction of the image. The script computes the total flux for each emission line and the total number of pixels.

Note2: It is necessary the image be large enough to be sure that after the rotation, the entire nebula or galaxy remains inside the image. Sometimes an elongated nebula (or even a galaxy) is observed in a specific PA, so the entire nebula fits the field of view of the instrument (**rotation angle=0 in the SATELLITE code corresponds to observed PA=0**).

Note3: The orientation of the maps/images should always be north up and east to the left. Otherwise the user has to take into account the offset between the sky (North) and image orientation.

In case, the slit width and length are equal to the number of the pixels in the raw map (parameter `total_num_pixels_horiz` in the *numerical_input.txt* file), the code calculates the integrated fluxes of the entire nebula for all the position angles. This specific task was used to verify if the rotation of the images affects the integrated line fluxes.

If the slit width and length are larger than the maximum number the software return the following message: “Sorry, your slit width or/and length are larger than the true size of the image”.

4.1.2 TeNe_angles_script, ionicabundances_angles_script and element_abundances_ICFs_angles_script

T_e , N_e , ionic/elemental abundances, ICFs and abundance ratios are also computed for each pseudo-slit. Various diagnostic lines can be used for T_e/N_e . The user can also choose which T_e/N_e combination will be applied for the abundances of each ion (see Figure 6). All these parameters are defined by the user in the *outputs.txt* file. T_e and N_e parameters are computed in the *TeNe_angles_script*, while the ionic, elemental abundances and ICFs are computed in the *ionicabundances_angles_script* and *element_abundances_ICFs_angles_script*. All the scripts make use of the *PyNeb* package.

4.2 specific_slit_analysis module

In this module, the user can define 10 pseudo-slits for a spectroscopic analysis of specific regions/structures in PN (e.g. knots, blobs, inset or outer regions) or in any extended nebula. All the input information from the 10 specific pseudo-slits are given by the user in the *numerical_input.txt* file:

- PA_for_specific_slit_n (in angle)
- width_for_specific_slit_n (in pixels)
- length_for_specific_slit_n (in pixels)
- x_coord_of_spec_slit_n (in pixels)

Te(NII6548_84)_Ne(SII6716_31)	yes
Ne(SII6716_31)_Te(NII6548_84)	yes
Te(OI6300_63)_Ne(SII6716_31)	no
Ne(SII6716_31)_Te(OI6300_63)	no
Te(OII3727_29_7320_30)_Ne(SII6716_31)	no
Ne(SII6716_31)_Te(OII3727_29_7320_30)	no
Te(OIII4959_5007)_Ne(SII6716_31)	no
Ne(SII6716_31)_Te(OIII4959_5007)	no
Te(SIII6312_9069)_Ne(SII6716_31)	yes
Ne(SII6716_31)_Te(SIII6312_9069)	yes
Te(OII3727_29_7320_30)_Ne(OII3727_29)	no
Ne(OII3727_29)_Te(OII3727_29_7320_30)	no
Te(NII6548_84)_Ne(OII3727_29)	no
Ne(OII3727_29)_Te(NII6548_84)	no
Te(OI6300_63)_Ne(OII3727_29)	no
Ne(OII3727_29)_Te(OI6300_63)	no
Te(OIII4959_5007)_Ne(CIIII5517_38)	no
Ne(CIIII5517_38)_Te(OIII4959_5007)	no
Te(SIII6312_9069)_Ne(CIIII5517_38)	yes
Ne(CIIII5517_38)_Te(SIII6312_9069)	yes
Te(OIII4959_5007)_Ne(ArVI4712_40)	no
Ne(ArVI4712_40)_Te(OIII4959_5007)	no
Te(SIII6312_9069)_Ne(ArVI4712_40)	no
Ne(ArVI4712_40)_Te(SIII6312_9069)	no
Te(NII6548_84)_Ne(CIIII5517_38)	no
Ne(CIIII5517_38)_Te(NII6548_84)	no
He+_Te_Ne	TeNIIINeSII
He++_Te_Ne	TeSIIINeClIII
Oo_Te_Ne	TeNIIINeSII
O+_Te_Ne	TeNIIINeSII
O++_Te_Ne	TeSIIINeClIII
No_Te_Ne	TeNIIINeSII
N+_Te_Ne	TeNIIINeSII
S+_Te_Ne	TeNIIINeSII
S++_Te_Ne	TeSIIINeClIII
Ar++_Te_Ne	TeSIIINeClIII
Ar+++_Te_Ne	TeSIIINeClIII
Ne++_Te_Ne	TeSIIINeClIII
Cl+_Te_Ne	TeNIIINeSII
Cl++_Te_Ne	TeSIIINeClIII
Cl+++_Te_Ne	TeSIIINeClIII
He_adun_KB	yes
He_abund_DIMS	yes
He_ICF_KB	yes
He_ICF_DIMS	yes
O_adun_KB	yes
O_abund_DIMS	yes
O_ICF_KB	yes
O_ICF_DIMS	yes
N_adun_KB	yes
N_abund_DIMS	yes
N_ICF_KB	yes
N_ICF_DIMS	yes
S_adun_KB	yes
S_abund_DIMS	yes
S_ICF_KB	yes
S_ICF_DIMS	yes
Ne_adun_KB	no
Ne_abund_DIMS	no
Ne_ICF_KB	no
Ne_ICF_DIMS	no
Ar_adun_KB	yes
Ar_abund_DIMS	yes
Ar_ICF_KB	yes
Ar_ICF_DIMS	yes
Cl_adun_KB	yes
Cl_abund_DIMS	yes
Cl_ICF_KB	yes
Cl_ICF_DIMS	yes

Figure 6: An example of the *outputs.txt* file for NGC 7009 and the parameters that the user has to define for the calculations of T_e , N_e , ionic/elemental abundances, ICFs and abundance ratios.

```

x_coor_of_CS          157
y_coor_of_CS          125

PA_for_specific_slit_1 79
width_for_specific_slit_1 10
length_for_specific_slit_1 320
x_coor_of_spec_slit_1 153
y_coor_of_spec_slit_1 106

PA_for_specific_slit_2 79
width_for_specific_slit_2 8
length_for_specific_slit_2 25
x_coor_of_spec_slit_2 37
y_coor_of_spec_slit_2 148

PA_for_specific_slit_3 79
width_for_specific_slit_3 8
length_for_specific_slit_3 21
x_coor_of_spec_slit_3 68
y_coor_of_spec_slit_3 142

PA_for_specific_slit_4 79
width_for_specific_slit_4 8
length_for_specific_slit_4 42
x_coor_of_spec_slit_4 104
y_coor_of_spec_slit_4 135

PA_for_specific_slit_5 79
width_for_specific_slit_5 8
length_for_specific_slit_5 18
x_coor_of_spec_slit_5 140
y_coor_of_spec_slit_5 128

PA_for_specific_slit_6 79
width_for_specific_slit_6 8
length_for_specific_slit_6 8
x_coor_of_spec_slit_6 169
y_coor_of_spec_slit_6 123

PA_for_specific_slit_7 79
width_for_specific_slit_7 8
length_for_specific_slit_7 46
x_coor_of_spec_slit_7 202
y_coor_of_spec_slit_7 116

PA_for_specific_slit_8 79
width_for_specific_slit_8 8
length_for_specific_slit_8 25
k_coor_of_spec_slit_8 254
y_coor_of_spec_slit_8 106

```

Figure 7: An example of the *numerical.input.txt* file for NGC 7009 and the parameters that the user has to define for the 10 pseudo-slits in the *specific_slit_analysis* module.




Figure 8: Ten selected regions in NGC 7009 overlaid on the [N II] 6584Å image. The position of the centre (x, y coordinates), position angle, width and length of the slits are free parameters provided by the user. Slits 1 and 10 represent the slits position from [2] and [3], respectively. Numbered regions from 2 to 9 correspond to the sub-structures of knots and jet-like, or sub-regions of rims defined in [3].

- `y_coor_of_spec_slit_n` (in pixels)

where `n` is the number of the pseudo-slit from 1 to 10 (see Figure 7). The `x_coor_of_spec_slit_n`, and `y_coor_of_spec_slit_n` parameters refer to the centre of each slit.

SATELLITE calculates the $H\beta$ flux, line intensities (normalized to $H\beta=100$ and corrected for interstellar extinction), emission line ratios (from the *output.txt* file), nebular parameters (T_e , N_e), ionic/total elemental abundances, ICFs and abundance ratios for all 10 pseudo-slits. This module is executed from the *specificPA_line_fluxes_script*. The scripts *slit_line_flux_script* is employed in this module for each pseudo-slit.

$c(H\beta)$, emission line intensities and line ratios are saved in the *output_linesintensities_per_angles.txt* and *output_lineratios_per_angles.txt* files, respectively. So, the user can also perform any extra analysis he/she wants. Similarly, T_e and N_e parameters are computed in the *TeNe_specific_slits_script* and are saved in the *PyNeb_output_Te_and_Ne_specific_slits* file, the ionic abundances are computed in the *ionicabundances_specific_slits_script* and are saved in the *PyNeb_output_ionic_abund_specific_slits* file, finally and the elemental abundances, ICFs and abundance ratios are computed in the *element.abundances_ICFs_specific_slits_script* and are saved in the *PyNeb_output_total_abund_ICFs_specific_slits* file.

Figure 8 shows the position of the specific areas/regions selected for the study of NGC 7009 overlaid on the [N II] flux map. The selected regions are the same as those defined by [3] for a direct comparison of the results from the *specific_slit_analysis module* with 1D long-slit spectroscopic data. Possible differences between the two studies can be associated with the position of the pseudo-slits.




Figure 9: Representative examples of the output figures produced by the *slit_position_testing* module. The slit from [2] (left panel) and R1 slit from [3] (right panel) are shown overlaid on the $H\alpha$ emission line maps. Scale is in "python counting; usual pixel counting starts in 1.

At this point, it is worth mentioning the *slit_position_testing* module of the SATELLITE code. This module is used to verify the position of the pseudo-slits before use the software. **Hint:** When the *slit_position_testing* module is used first deselect all other modules. Moreover, at least an emission line has to be used and defined in the *input.txt* file in order to properly use this module (e.g., $H\alpha$ and $H\beta$ to avoid multiple maps). The output of this module is 10 figures (in png and pdf formats) with the position of each pseudo-slit overlaid on the emission line map (see Figure 9).

5 2D analysis module

Besides the 1D spectroscopic analysis, SATELLITE also performs a spectroscopic analysis in both spatial dimensions simultaneously using the entire maps. For this module, the ANALYSIS2D_SCRIPT, TeNe_2D_SCRIPT, GENERATE_2D_LINERATIO_MAPS_SCRIPT, IONICABUNDANCES_2D_SCRIPT, ELEMENT_ABUNDANCES_ICFs_2D_SCRIPT and DIAGNOSTIC_DIAGRAMS_SCRIPT are employed.

$c(H\beta)$, line intensities, line ratios, T_e , N_e , ionic, elemental abundances, ICFs and abundances ratios are computed for each individual pixels, if the criteria $F(Ha) > 0$, $F(Hb) > 0$ and $F(Ha) > F(Hb) * 2.86$ are satisfied, otherwise a value equals to "zero" is applied.

The main outcomes from this module are 2D maps for all the aforementioned nebular parameters saved in the *output_images* folder. In Figures 10, 11 and 12, the maps of $c(H\beta)$, T_e and N_e using the [S III] and [S II] diagnostic lines, the line ratios $\log([N II]/[O III])$ and $\log([S II]/[S III])$ are presented as representative examples of the outcomes from this module.

The *2D_spectroscopic_analysis* module also calculates and returns the distribution of each maps (histogram plots), e.g., $c(H\beta)$, N_e and T_e maps (see Figures 13 and 14) as well as emission line diagnostic diagrams (see Figures 15 and 16) using the DIAGNOSTIC_DIAGRAMS_SCRIPT which are selected by the user in the *diagnostic_diagrams_input*




Figure 10: $c(\text{H}\beta)$ map of NGC 7009.

file (see Figure 17).

Note1: At this point, it has to be clarified that when the *specific_slits_analysis* and/or *rotation_analysis* modules are used together with the *2D_analysis* module, the emission lines ratios for the three modules are plotted on the same diagnostic diagrams for a direct comparison between an 1D and 2D analysis.

Last but not least, SATELLITE computes and returns an ASCII file with the mean value, standard deviation and the percentiles of 5%, 25% (Q1), 50% (median), 75% (Q3), 95% for all the nebular parameters and emission line ratios for a thorough statistical analysis of the observed nebula.




Figure 11: N_e and T_e maps obtained from the [S II] and [S III] diagnostic lines of NGC 7009.




Figure 12: $\log([N II]/[O III])$ and $\log([S II]/[S III])$ line ratio maps of NGC 7009.




Figure 13: The histogram of $c(H\beta)$ map.





Figure 14: The histograms of $\text{N}_e[\text{S II}]$ and $\text{T}_e[\text{S III}]$ maps.




Figure 15: A representative example of emission line diagnostic diagram: $[\text{S II}] 6716/6731$ versus $\text{H}\alpha/[\text{N II}] 6548+6584$. Cyan dots correspond to the values of individual pixels, pink circles and yellow diamonds show the values obtained from the simulated long-slits of the rotational analysis task with position angles from 0 to 360 degrees with 10 degrees step and the values from the 10 simulated slits in the specific slits task, respectively. The inset plot illustrate the variation of the line ratios with the position angle of the simulated slits.




Figure 16: A representative example of emission line diagnostic diagram: $[\text{O III}] 5007/\text{H}\beta$ versus $[\text{S II}] 6716+6731/\text{H}\alpha$. Cyan dots correspond to the values of individual pixels, pink circles and yellow diamonds show the values obtained from the simulated long-slits of the rotational analysis task with position angles from 0 to 360 degrees with 10 degrees step and the values from the 10 simulated slits in the specific slits task, respectively. The inset plot illustrate the variation of the line ratios with the position angle of the simulated slits. The regimes of the PNe, H II regions and supernova remnants are also drawn.

Kauffmann2003_BPT_NII	no	0	0	0	0
Kewley2001_BPT_NII	no	0	0	0	0
main_AGN_line_BPT_SII	no	0	0	0	0
LINER/Sy2_line_BPT_SII	no	0	0	0	0
main_AGN_line_BPT_OI	no	0	0	0	0
LINER/Sy2_line_BPT_OI	no	0	0	0	0
Ha/NII+_vs_Ha/SII+	yes	-0.5	3.0	-1.0	2.0
SII6716/6731_vs_Ha/SII+	yes	0.0	3.0	-1.5	2.0
SII6716/6731_vs_Ha/NII+	yes	-1.0	2.5	-1.5	2.0
NII+/SII+_vs_SII+/OI+	yes	-0.5	2.5	-0.4	1.2
NII+/SII+_vs_NII+/OI+	yes	0.0	3.2	-0.4	1.2
OIII5007/Hb_vs_NII6584/Ha	yes	-2.5	1.1	-0.25	1.5
OIII5007/Hb_vs_SII+/Ha	yes	-3.0	0.5	-0.25	1.5
OIII5007/Hb_vs_HeII4686/Hb	no	-1.3	0.0	-2.5	1.4
OIII5007/Hb_vs_HeII5412/Hb	yes	-4.5	-1.0	0.3	1.4
OIII5007/Hb_vs_HeI5876/Ha	yes	-1.5	-1.0	0.3	1.4
OIII5007/Hb_vs_NI5199/Hb	yes	-5.0	0.0	0.3	1.4
OIII5007/Hb_vs_ArIII7136/Ha	yes	-1.8	-0.8	0.3	1.4
OIII5007/Hb_vs_OI6300/Ha	yes	-5.0	0.0	0.3	1.4
OIII5007/Hb_vs_NII5755/Hb	no	-5.0	0.0	0.0	2.0
OIII5007/Hb_vs_OIII5007/OI6300	no	-0.5	3.5	-1.5	2.0
OIII5007/Hb_vs_OII3727+/OIII5007	no	-0.5	3.5	-1.5	2.0
OIII5007/Hb_vs_OII7320+/OIII5007	yes	-4.0	-1.0	0.3	1.4
OIII5007/Hb_vs_OII3727+/Hb	no	-0.5	3.5	-1.5	2.0
OIII5007/Hb_vs_OII7320+/Ha	yes	-3.0	-0.5	0.3	1.4
OIII5007/Hb_vs_NeIII3869/Hb	no	-0.5	3.5	-1.5	2.0
OIII5007/Hb_vs_NII6584/OI6300	yes	-0.0	3.0	0.3	1.4
OIII5007/Hb_vs_NI5199/NII6584	yes	-3.5	0.5	0.3	1.4
OIII5007/Hb_vs_OIII4363/Hg	no	-0.5	3.5	-1.5	2.0
OIII5007/Hb_vs_ArIV4712/4740	no	-0.5	3.5	-1.5	2.0
OIII5007/Hb_vs_CII8727/Ha	no	-0.5	3.5	-1.5	2.0
OIII5007/Hb_vs_CII6461/Ha	no	-0.5	3.5	-1.5	2.0
OI6300/Ha_vs_CII8727/Ha	no	-0.5	3.5	-1.5	2.0
ArIV+/Hb_vs_HeII4686/Hb	no	-0.5	3.5	-1.5	2.0
ArIV+/Hb_vs_HeII5412/Hb	no	-0.5	3.5	-1.5	2.0

Figure 17: An example of the *diagnostic_diagrams.input.txt* file for NGC 7009 and the parameters that the user can select for the diagnostic diagrams in the *2D_analysis.module* module.

5.1 radial analysis module

The final module in the current version of SATELLITE (v1.2) conducts a radial spectroscopic analysis considering a pseudo-slit with specific width, length and position angle (parameter=angle_for_radial_flux) provided by the user in the *numerical_input.txt* file. The user must also select the emission lines that will be used for this analysis. This can be made in the third column of the *input.txt* file: radial_yes, or radial_no.

Note1: It is recommended to disable all other modules when the *radial_analysis* module is executed.

The main outcomes of this module are:

- (I) the radial profiles of all the selected emission lines in the *input.txt* normalized by the peak flux.
- (II) the calculation of all the nebular parameters ($c(\text{H}\beta)$, line intensities, line ratios, T_e , N_e , ionic, elemental abundances, ICFs and abundances ratios) as functions of the distance from the central star or the central point of the nebula or galaxy in general.

The normalization of the radial profiles is made using the peak of the flux of each emission line. However, the user can also select the range from where this peak can be obtained by providing the code with the minimum radius (*limit_radial_in_arcsec* parameter). This option permit to investigate the radial distribution of emission lines for regions/substructures with specific interest.

The radial profile of various emission lines for the example of NGC 7009 are shown in Figure 18 (**Hint:** It is recommended to use maximum 4-6 lines for the construction of more illustrative plots.). All radial profiles are normalized to a peak flux found for distances $r > 20$ arcsec (*_radial_in_arcsec > 20*) focused to the low-ionization structures/knot of NGC 7009. The calculation are made in the *find_maxvlaue_script*. Hence, SATELLITE returns the distance between the peak of each selected line and the central star in arcsec. Table 1 lists the distances for the example of NGC 7009 and it can be seen that there is a spatial offset of 1 arcsec between the high/moderate- and low-ionization lines. The values of each radial step (pixel scale of the IFU) are also saved in an ASCII file, so the user can build his/her own radial profiles.

The radial variation of $c(\text{H}\beta)$, T_e , and N_e parameters of NGC 7009 are shown in Figure 19

One again, it has to be pointed out that the code sums up the values of the pixels which have $F(\text{H}\alpha) > 0$, $F(\text{H}\beta) > 0$ and/or $F(\text{H}\alpha) > F(\text{H}\beta) * 2.86$.

5.1.1 radial distance calculations

At this point, it is necessary to further explain how SATELLITE calculates the fluxes of the emission lines as function of the distance from the central star. Figure 20 shown an example of a pseudo-slit at PA=90 degrees.

The width of the pseudo-slit defines how many pixels will be taken into consideration for the flux at each distance. For instance, the fluxes (and errors) of 5 pixels are summed up for the first column (or radial distance $r = 0.2$ arcsec). Then, the code moves to the second column and computes the flux and the corresponding error from the next 5 pixels at the radial distance $r = 0.4$ arcsec and so on (see Figure 20).

After finishing the computation of the fluxes and errors for all the lines, the code computes the extinction coefficient ($c(\text{H}\beta)$) and corrected line intensities (relative to $\text{H}\beta = 100$) as function of the radial distance from the central star or geometric centre as well as all nebular parameters (T_e , N_e , ionic, elemental abundances, ICFs and abundance ratios) (Figures 18 and 19).




Figure 18: Radial profiles for several emission lines of NGC 7009 at PA=79 degrees. Upper panel shows all the radial profiles, while the lower panel zooms-in to the much fainter emission lines.




Figure 19: Representative examples of the radial distribution of $c(H\beta)$ upper panel and T_e , N_e (lower panel).

Table 1: Distances from the central stars of emission line's peak for a pseudo-slit at 79 degree position angle

Line	distance [†] (arcsec)	Line	distance (arcsec)
H I 4861Å	23.6	[N II] 6548Å	24.8
[O III] 4959Å	23.8	H I 6563Å	23.6
[N I] 5199Å	24.8	[N II] 6584Å	24.8
He II 5412Å	20.2	He I 6678Å	23.8
[Cl III] 5517Å	24.2	[S II] 6716Å	24.8
[Cl III] 5538Å	24.4	[S II] 6731Å	24.8
[N I] 5755Å	24.2	[Ar III] 7136Å	24.8
He I 5876Å	23.6	[O II] 7320Å	24.8
[O II] 6300Å	24.8	[O II] 7330Å	24.8
[S III] 6312Å	23.8	[S III] 9069Å	23.8

[†] The spacial resolution of MUSE maps is 0.2 arcsec.




Figure 20: An illustrative example of how SATELLITE computes the fluxes and radial stances from the central star of the nebula or the geometric centre of the nebula.

6 Uncertainty calculations

The uncertainties of emission lines and all nebular parameters for all four modules are computed following the same Monte Carlo approach. In particular, SATELLITE, first, computes the total error of the flux for each pseudo-slit or pixel, using the provided error maps + an extra error as the percentage of the flux.

$$\Delta F_{tot} = \sum_{i=1}^N (\sigma_{F_i} + \lambda * F_i)^{1/2} \quad (1)$$

where i corresponds to the pseudo-slit or pixel and range from 1 to the total number N , σ_{F_i} is the uncertainties of the flux in the pseudo-slit or pixel i based on the provided error maps, F_i is the flux in the pseudo-slit or pixel i and λ corresponds to the percentage of the flux (from 0 to 1.0). If $\lambda=0$, then the code takes into account only the errors from the maps. The $\lambda=0$ parameter is given to the code by the user in the *input.txt* file (forth column). There is also the option not to use the error maps. This is defined in the *input.txt* file (forth column) even rows (the rows of errors). If a non-zero value is provided, the code uses the formula (1) while for a "0" value, the code uses the formula (2).

$$\Delta F_{tot} = \sum_{i=1}^N (\lambda * F_i)^{1/2} \quad (2)$$

These resultant uncertainties of the line fluxes are then used to replicate the spectrum of a pseudo-slit or pixel, and a number of additional spectra are generated using a Monte Carlo algorithm. The number of the replicate spectra is given by the user in the *numerical_input.txt*. SATELLITE computes all the nebular parameters, e.g. T_e , N_e , ionic, elemental abundances and ICFs for all the replicate spectra and the standard deviation of each parameters is the uncertainty of the parameter that the SATELLITE code provides.

7 General Notes:

- It has to be clarified that the SATELLITE code takes as input a list of emission line fluxes and error maps extracted from the datacubes obtained from any IFU. It does not extract the maps from the datacubes. Therefore, this is a step that has to be done before the use of SATELLITE.
- Moreover, SATELLITE can also be applied to individual emission lines images obtained with narrow band filters (if there are available) or the emission line images obtained from 3D photo-ionization models.

8 Possible error messages

In this section, a number of possible errors that may come out are described.

- In case an emission line is missing, the user has to define that in the *input.txt* file by writing "no" in the second and third columns of the corresponding line. If the user has forgotten to properly change the *output.txt* file or the *diagnostic_diagram_input.txt* file, an error will be return, see Figure 21.
- In case an emission line is missing, but the user has forgotten to properly change the *output.txt* file or the *diagnostic_diagram_input.txt* file and a physical parameters has to be computed such as T_e , N_e , abundances, an error will be return like in Figure 22.

```
[akras@akras SATELLITE_v1.2]$ python main_script.py > specific_slits_output.txt
Traceback (most recent call last):
  File "main_script.py", line 280, in <module>
    flux_spec_slit, flux_spec_slit_error, flux_spec_slit_norm, flux_spec_slit_norm_error, ratio_spec_slit, ratio_spec_slit_err
or=sPALfs.specificPA_line_fluxes(flux2D, flux2D_error, line_names, line_ext_error, lines_available, lines_radial, param_estimate
d, param_required, param_mod_name, param_model_values)
  File "/home/akras/more_projects/SATELLITE/NGC6778_MUSE/SATELLITE_v1.2/specificPA_line_fluxes_script.py", line 1531, in
specificPA_line_fluxes
    ratio_HeIa_HeIIa[slit_number]=np.log10(HeIa[slit_number]/HeIIa[slit_number])
ZeroDivisionError: integer division or modulo by zero
```

Figure 21: An example of error in case the He II line is missing (in the *input.txt* file, it has been set as "no") but the He I/He II ratio is required to be computed (in the *output.txt*, the He I/He II ratio is still "yes").

```
[akras@akras SATELLITE_v1.2]$ python main_script.py > 2D_output.txt
Traceback (most recent call last):
  File "main_script.py", line 311, in <module>
    Te_2D, Ne_2D = TeNe2Ds.TeNe(flux2D, flux2D_error, line_names, line_ext_error, lines_available, param_mod_name, param_model_
values, param_estimated, param_required, hdr, Te_PA, Ne_PA)
  File "/home/akras/more_projects/SATELLITE/NGC6778_MUSE/SATELLITE_v1.2/TeNe_2D_script.py", line 2391, in TeNe
    pxi_num, percent5, percentQ1, medianvalue, percentQ3, percent95, meanvalue, sigmavalue= scs.statistic_numbers(Te.NIIClIII, si
zex, sizey)
  File "/home/akras/more_projects/SATELLITE/NGC6778_MUSE/SATELLITE_v1.2/statistics_calculations_script.py", line 37, in s
tatistic_numbers
    a2=np.percentile(data, 5)
  File "/usr/lib64/python2.7/site-packages/numpy/lib/function_base.py", line 3707, in percentile
    a, q, axis, out, overwrite_input, interpolation, keepdims)
  File "/usr/lib64/python2.7/site-packages/numpy/lib/function_base.py", line 3826, in _quantile_unchecked
    interpolation=interpolation)
  File "/usr/lib64/python2.7/site-packages/numpy/lib/function_base.py", line 3405, in _ureduce
    r = func(a, **kwargs)
  File "/usr/lib64/python2.7/site-packages/numpy/lib/function_base.py", line 3941, in _quantile_ureduce_func
    x1 = take(ap, indices_below, axis=axis) * weights_below
  File "/usr/lib64/python2.7/site-packages/numpy/core/fromnumeric.py", line 189, in take
    return _wrapfunc(a, 'take', indices, axis=axis, out=out, mode=mode)
  File "/usr/lib64/python2.7/site-packages/numpy/core/fromnumeric.py", line 56, in _wrapfunc
    return getattr(obj, method)(*args, **kwds)
IndexError: cannot do a non-empty take from an empty axes.
```

Figure 22: An example of error in case a physical parameter is required to be computed and the corresponding line is missing (in the *input.txt* file, it has been set as "no". In this example, the T_e and N_e from the [N II] and [Cl III] diagnostic lines have to be computed but a diagnostic line is missing.

- In case, the arrays of the emission lines have different sizes an error will be return like in Figure 23. Moreover, the error line may also be related to the parameters *total_num_pixels_verti* and *total_num_pixels_horiz* in the *numerical.input.txt* file.

9 How to Cite SATELLITE in a publication?

There are two papers appropriate as references for SATELLITE in a publication. They are:

1) Akras, Stavros; Monteiro, Hektor; Aleman, Isabel; Farias, Marcos A. F. ; May, Daniel ; Pereira, Claudio B., 2020, MNRAS, 493, 2238A

bibtex code:

@ARTICLE{2020MNRAS.493.2238A,

author = Akras, Stavros and Monteiro, Hektor and Aleman, Isabel and Farias, Marcos A. F. and May, Daniel and Pereira, Claudio B.,

title = "Exploring the differences of integrated and spatially resolved analysis using integral field unit data: the

```
[akras@akras Satellite_project_v1.2]$ python main_script.py > angles_output.txt
Traceback (most recent call last):
  File "main_script.py", line 174, in <module>
    flux2D.Ha_6563,flux2D.Hb_4861,flux2D.Hg_4340,flux2D.Hd_4101,flux2D.HeIa_5876,flux2D.HeIb_6678,flux2D.HeIIa_4686,flux2
D.HeIb_5412,flux2D.NIIa_5755,flux2D.NIIb_6548,flux2D.NIIc_6584,flux2D.NI_5199,flux2D.OIIa_4363,flux2D.OIIb_4959,flux2D
.OIIc_5007,flux2D.OIIa_3727,flux2D.OIIb_3729,flux2D.OIIc_7320,flux2D.OIIb_7330,flux2D.OIa_5577,flux2D.OIb_6300,flux2D.OI
c_6363,flux2D.SIIa_6716,flux2D.SIIb_6731,flux2D.SIIa_6312,flux2D.SIIb_9069,flux2D.CIIa_5517,flux2D.CIIb_5538,flux2D
.ArIII_7136,flux2D.ArIVa_4712,flux2D.ArIVb_4740,flux2D.CI_8727,flux2D.CII_6461,flux2D.NeIIa_3868,flux2D.NeIIb_3967,hdr=
read.read_input_images(line_names,lines_available,param_model_values,"fluxes")
  File "/home/akras/more_projects/SATELLITE/NGC7009_MUSE/Satellite_project_v1.2/read_input_script.py", line 48, in read_i
nput_images
    addupdown1=np.concatenate((z2,dataHa,z1), axis=0)
ValueError: all the input array dimensions except for the concatenation axis must match exactly
```

Figure 23: An example of error in case there is a problem with the dimensions of the arrays that correspond to the emission lines. In this case, the error is because the size of the flux maps is not consistent with the *total_num_pixels_verti* and *total_num_pixels_horiz* parameters in the *numerical_input.txt* file.

case of Abell 14",

journal = \mnras,

keywords = techniques: imaging spectroscopy; techniques: spectroscopic; (stars:) binaries: general, ISM: abundances, (ISM:) planetary nebulae: individual: Abell 14, Astrophysics - Astrophysics of Galaxies, Astrophysics - Solar and Stellar Astrophysics,

year = 2020,

month = apr,

volume = 493,

number = 2,

pages = 2238-2252,

doi = 10.1093/mnras/staa383,

archivePrefix = arXiv,

eprint = 2002.12380,

primaryClass = astro-ph.GA,

adsurl = <https://ui.adsabs.harvard.edu/abs/2020MNRAS.493.2238A>,

adsnote = Provided by the SAO/NASA Astrophysics Data System

2) S. Akras; H. Monteiro; J. R. Walsh; J. García-Rojas; I. Aleman; H. Boffin; P. Boumis; A. Chiotelis; R. M. L. Corradi; D. R. Gonçalves; L. A. Gutiérrez-Soto; D. Jones; C. Morisset, 2021, MNRAS, submitted

10 Licence and Copyright Information

SATELLITE is freely available under the General Public License (GPL).

11 Questions of problems

For questions please write an email to Dr. Stavros Akras (stavrosakras@gmail.com)

References

- [1] S. Akras, H. Monteiro, I. Aleman, M. A. F. Farias, D. May, and C. B. Pereira. Exploring the differences of integrated and spatially resolved analysis using integral field unit data: the case of Abell 14. , 493(2):2238–2252, Apr. 2020.
- [2] X. Fang and X. W. Liu. Very deep spectroscopy of the bright Saturn nebula NGC 7009 - I. Observations and plasma diagnostics. , 415(1):181–198, July 2011.
- [3] D. R. Gonçalves, R. L. M. Corradi, A. Mampaso, and M. Perinotto. The Physical Parameters, Excitation, and Chemistry of the Rim, Jets, and Knots of the Planetary Nebula NGC 7009. , 597(2):975–985, Nov. 2003.
- [4] V. Luridiana, C. Morisset, and R. A. Shaw. PyNeb: a new tool for analyzing emission lines. I. Code description and validation of results. , 573:A42, Jan. 2015.
- [5] J. R. Walsh, A. Monreal-Ibero, M. J. Barlow, T. Ueta, R. Wesson, and A. A. Zijlstra. The extinction and dust-to-gas structure of the planetary nebula NGC 7009 observed with MUSE. , 588:A106, Apr. 2016.
- [6] J. R. Walsh, A. Monreal-Ibero, M. J. Barlow, T. Ueta, R. Wesson, A. A. Zijlstra, S. Kimeswenger, M. L. Leal-Ferreira, and M. Otsuka. An imaging spectroscopic survey of the planetary nebula NGC 7009 with MUSE. , 620:A169, Dec. 2018.