


**Cochrane** Database of Systematic Reviews

# Triage tools for detecting cervical spine injury in pediatric trauma



Triage tools for detecting cervical spine injury in pediatric trauma patients (Review)

Copyright @ 2017 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

www.cochranelibrary.com

i



# TABLE OF CONTENTS

ABSTRACT	1
PLAIN LANGUAGE SUMMARY	2
SUMMARY OF FINDINGS	4
BACKGROUND	5
Figure 1	6
Figure 2	7
Figure 3	ç
OBJECTIVES	ç
METHODS	10
RESULTS	12
Figure 4	13
Figure 5	14
Figure 6	15
Figure 7	16
DISCUSSION	16
AUTHORS' CONCLUSIONS	18
ACKNOWLEDGEMENTS	18
REFERENCES	19
CHARACTERISTICS OF STUDIES	26
DATA	33
Test 1. NEXUS.	34
Test 2. Canadian C-spine Rule.	34
ADDITIONAL TABLES	34
APPENDICES	38
CONTRIBUTIONS OF AUTHORS	47
DECLARATIONS OF INTEREST	48
SOURCES OF SUPPORT	48
DIFFERENCES BETWEEN PROTOCOL AND REVIEW	48
INDEX TERMS	48



[Diagnostic Test Accuracy Review]

# Triage tools for detecting cervical spine injury in pediatric trauma patients

Annelie Slaar<sup>1</sup>, M M Fockens<sup>2</sup>, Junfeng Wang<sup>3</sup>, Mario Maas<sup>4</sup>, David J Wilson<sup>5</sup>, J Carel Goslings<sup>6</sup>, Niels WL Schep<sup>7</sup>, Rick R van Rijn<sup>8</sup>

<sup>1</sup>Department of Radiology, Westfriesgasthuis, Hoorn, Netherlands. <sup>2</sup>Academic Medical Center, University of Amsterdam, Amsterdam, Netherlands. <sup>3</sup>Department of Clinical Epidemiology, Biostatistics and Bioinformatics, Academic Medical Center, Amsterdam, Netherlands. <sup>4</sup>Department of Radiology, Academic Medical Center, Amsterdam, Netherlands. <sup>5</sup>Department of Radiology, St Lukes Hospital, Oxford, UK. <sup>6</sup>Trauma Unit, Department of Surgery, Academic Medical Center, Amsterdam, Netherlands. <sup>7</sup>Department of Surgery/Trauma Unit, Academic Medical Center, Amsterdam, Netherlands. <sup>8</sup>Department of Radiology, Academic Medical Center Amsterdam, Netherlands

**Contact:** Annelie Slaar, Department of Radiology, Westfriesgasthuis, Maelsonstraat 3, Hoorn, Noord Holland, 1624 NP, Netherlands. a.slaar@gmail.com.

Editorial group: Cochrane Back and Neck Group.

Publication status and date: New, published in Issue 12, 2017.

**Citation:** Slaar A, Fockens MM, Wang J, Maas M, Wilson DJ, Goslings JC, Schep NWL, van Rijn RR. Triage tools for detecting cervical spine injury in pediatric trauma patients. *Cochrane Database of Systematic Reviews* 2017, Issue 12. Art. No.: CD011686. DOI: 10.1002/14651858.CD011686.pub2.

Copyright © 2017 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

#### **ABSTRACT**

# Background

Pediatric cervical spine injury (CSI) after blunt trauma is rare. Nonetheless, missing these injuries can have severe consequences. To prevent the overuse of radiographic imaging, two clinical decision tools have been developed: The National Emergency X-Radiography Utilization Study (NEXUS) criteria and the Canadian C-spine Rule (CCR). Both tools are proven to be accurate in deciding whether or not diagnostic imaging is needed in adults presenting for blunt trauma screening at the emergency department. However, little information is known about the accuracy of these triage tools in a pediatric population.

# Objectives

To determine the diagnostic accuracy of the NEXUS criteria and the Canadian C-spine Rule in a pediatric population evaluated for CSI following blunt trauma.

## Search methods

We searched the following databases to 24 February 2015: CENTRAL, MEDLINE, MEDLINE Non-Indexed and In-Process Citations, PubMed, Embase, Science Citation Index, ProQuest Dissertations & Theses Database, OpenGrey, ClinicalTrials.gov, World Health Organization (WHO) International Clinical Trials Registry Platform (ICTRP), Cochrane Database of Systematic Reviews, the Database of Abstracts of Reviews of Effects, the Health Technology Assessment, and the Aggressive Research Intelligence Facility.

## **Selection criteria**

We included all retrospective and prospective studies involving children following blunt trauma that evaluated the accuracy of the NEXUS criteria, the Canadian C-spine Rule, or both. Plain radiography, computed tomography (CT) or magnetic resonance imaging (MRI) of the cervical spine, and follow-up were considered as adequate reference standards.

#### **Data collection and analysis**

Two review authors independently assessed the quality of included studies using the QUADAS-2 checklists. They extracted data on study design, patient characteristics, inclusion and exclusion criteria, clinical parameters, target condition, reference standard, and the



diagnostic two-by-two table. We calculated and plotted sensitivity, specificity and negative predictive value in ROC space, and constructed forest plots for visual examination of variation in test accuracy.

#### Main results

Three cohort studies were eligible for analysis, including 3380 patients; 96 children were diagnosed with CSI. One study evaluated the accuracy of the Canadian C-spine Rule and the NEXUS criteria, and two studies evaluated the accuracy of the NEXUS criteria. The studies were of moderate quality. Due to the small number of included studies and the diverse outcomes of those studies, we could not describe a pooled estimate for the diagnostic test accuracy. The sensitivity of the NEXUS criteria of the individual studies was 0.57 (95% confidence interval (CI) 0.18 to 0.90), 0.98 (95% CI 0.91 to 1.00) and 1.00 (95% CI 0.88 to 1.00). The specificity of the NEXUS criteria was 0.35 (95% CI 0.25 to 0.45), 0.54 (95% CI 0.45 to 0.62) and 0.2 (95% CI 0.18 to 0.21). For the Canadian C-spine Rule the sensitivity was 0.86 (95% CI 0.42 to 1.00) and specificity was 0.15 (95% CI 0.08 to 0.23). Since the quantity of the data was small we were not able to investigate heterogeneity.

#### **Authors' conclusions**

There are currently few studies assessing the diagnostic test accuracy of the NEXUS criteria and CCR in children. At the moment, there is not enough evidence to determine the accuracy of the Canadian C-spine Rule to detect CSI in pediatric trauma patients following blunt trauma. The confidence interval of the sensitivity of the NEXUS criteria between the individual studies showed a wide range, with a lower limit varying from 0.18 to 0.91 with a total of four false negative test results, meaning that if physicians use the NEXUS criteria in children, there is a chance of missing CSI. Since missing CSI could have severe consequences with the risk of significant morbidity, we consider that the NEXUS criteria are at best a guide to clinical assessment, with current evidence not supporting strict or protocolized adoption of the tool into pediatric trauma care. Moreover, we have to keep in mind that the sensitivity differs among several studies, and individual confidence intervals of these studies show a wide range. Our main conclusion is therefore that additional well-designed studies with large sample sizes are required to better evaluate the accuracy of the NEXUS criteria or the Canadian C-spine Rule, or both, in order to determine whether they are appropriate triage tools for the clearance of the cervical spine in children following blunt trauma.

#### PLAIN LANGUAGE SUMMARY

## Clinical tests for detecting cervical spine injury (CSI) in children with injuries

## **Background and Rationale**

The cervical spine is the upper part of the spine between the head and shoulders (the neck). The incidence of traumatic cervical spine injury (CSI) in children is very low. However it is very important not to miss this type of injury. To detect CSI, several types of scan imaging techniques can be used (computed tomography (CT) scan, magnetic resonance imaging (MRI) and plain radiography (x-ray)). CT scan and x-ray use radiation that can lead to an increased risk in the development of cancer, especially in children. We therefore need to use plain radiography or CT scan in children only if really necessary. To avoid unnecessary use of those radiographic imaging techniques, it is important to look for clinical tests that can detect whether children are at risk for cervical spine injury and if radiographic imaging needs to be done.

## **Index tests**

The NEXUS criteria and Canadian C-spine Rules are tools that doctors use to decide whether adults are at risk for cervical spine injury and whether they need to use radiographic imaging. These tools are already being used in children, but little information is known about how accurate they are in children. The goal of this review was therefore to evaluate whether these tools can also be used safely and effectively in children.

## **Study characteristics**

We searched medical literature databases to identify studies which tested how well both decision tools can establish whether children are at risk for CSI after blunt trauma. We performed the search in February 2015.

## Quality of the evidence

We identified three studies of moderate to good quality. All studies tested the accuracy of the NEXUS criteria, and one of them also tested the accuracy of the Canadian C-spine Rules.

# **Key results:**

Since only one study looked at the accuracy of the Canadian C-spine Rules, there is not enough evidence at the moment to determine whether the Canadian C-spine Rules can be used safely in children. The sensitivity and specificity of the NEXUS criteria varied among the three studies, meaning that there is a chance of false-negative test results when using the NEXUS criteria, and as a result there is a chance of missing cervical spine injury if doctors only rely on the NEXUS criteria. We therefore consider that the NEXUS criteria are at best a guide to clinical assessment, with current evidence not supporting strict or protocolized adoption of the tool into pediatric trauma care. The



conclusion of our review is that we need more research to evaluate the accuracy of the NEXUS criteria and the Canadian C-spine rules for routine use in children.



## SUMMARY OF FINDINGS

# Summary of findings 1. New Summary of findings table

Triage tools for detecting cervical spine injury in pediatric trauma patients			
Population	Children evaluated for CSI following blunt trauma		
Setting	Emergency Departments		
Index test	NEXUS criteria, Canadian C-spine Rule (CCR)		
Reference stan- dard	Plain radiography, CT scan, and or MRI of the cervical spine, or follow-up without imaging		
Target Condi- tion	Cervical Spine Injury		

Studies	Accuracy (95%CI)	Number of par- ticipants	CSI prevalence	Quality (QUADAS-2)
Jaffe 1987	NEXUS criteria  Sensitivity: 0.98 (0.91 to 1.00)  Specificity: 0.54 (0.45 to 0.62)  Positive LR*: 2.13  Negative LR: 0.03	206	59 (28.6%)	High risk of bias, low concerns of applica- bility**
Ehrlich 2009	NEXUS criteria Sensitivity: 0.57 (0.18 to 0.9) Specificity: 0.35 (0.25 to 0.45) Positive LR: 0.87 Negative LR: 1.24	108	7 (6.5%)	Low risk of bias. No applicability con- cerns
	Canadian C-spine Rule Sensitivity: 0.86 (0.42 to 1.0) Specificity: 0.15 (0.08 to 0.23) Positive LR: 1.00 Negative LR: 0.97	109	7 (6.4%)	
Viccellio 2001	NEXUS criteria  Sensitivity: 1.00 (0.88 to 1.0)  Specificity: 0.2 (0.18 to 0.21)  Positive LR: 1.25  Negative LR: 0.081	3065	30 (0.98%)	Low risk of bias. No applicability con- cerns

<sup>\*</sup>LR=Likelihood ratio

<sup>\*\*</sup>High risk of bias based on inclusion of not only a consecutive group of children evaluated for the presence of CSI, but also an alternative group of children already diagnosed with CSI.



#### BACKGROUND

## **Target condition being diagnosed**

Pediatric cervical spine injury (CSI) after blunt trauma is rare, accounting for only 1% to 2% in children presenting at the emergency department for blunt trauma evaluation (Patel 2001; Viccellio 2001). Nonetheless, every physician should be concerned about missing these injuries because they can have severe consequences, such as death (up to 40%) or life-changing neurological damage (up to 60%) (Cirak 2004; Givens 1996; Hutchings 2009; Kokoska 2001; Parent 2011; Patel 2001; Platzer 2007). In 5% to 10% of cases where CSI is missed, patients experience worsening of neurological symptoms (Ravichandran 1982; Schuster 2005). These adverse sequelae depend primarily on the anatomical level of injury and the presence of dislocation (Partrick 2000), whereby CSI of the upper cervical spine is associated with higher morbidity and mortality than CSI of the lower cervical spine (Leonard 2014).

Children aged under eight display different patterns of injury than children aged between eight and 18 years (Viccellio 2001). This is because younger children have unique anatomic features, such as a relatively larger head, immature neck musculature, horizontally-articulating facet joints, and ligamentous laxity (Kokoska 2001). Consequently, injury at the cranio-cervical junction and upper cervical spine is seen more frequently in younger children (Cirak 2004; Orenstein 1994; Sun 2000). The same applies to ligamentous injury (Dickman 1989; Dormans 1995; Hadley 1988; Hamilton 1992; Osenbach 1992). CSI in children aged eight and older is comparable to blunt trauma patterns in adults (Hill 1984). In this age group, subaxial injuries are most common (Leonard 2014).

Several blunt trauma mechanisms can cause CSI. In children of all ages, falls and motor vehicle crashes account for a large proportion (Leonard 2011; Patel 2001). In younger children, pedestrian and inflicted injuries are seen more often. Older children sustain more injury due to sports-related and recreational activities (Brown 2001; Garton 2008; Hasler 2012; Leonard 2011).

Symptoms of CSI that have been reported are neck pain, torticollis, altered mental status, sensory loss, motor loss and respiratory arrest (Leonard 2013). In children it may be difficult to discriminate between general discomfort and actual complaints due to blunt trauma.

Multiple imaging modalities are available to the physician to diagnose or rule out CSI:

- Plain radiography is usually performed to detect pediatric CSI, since this technique is portable, readily available, and has a relatively low radiation dosage (Booth 2012). As the sensitivity for detecting bony CSI is greater than 90%, plain radiographs are considered adequate for the screening of alert patients who show no abnormalities on neurological examination (Mower 2001; Nigrovic 2012).
- Computed tomography (CT) is superior to plain radiography in detecting cervical spine fractures, with a sensitivity of nearly 100% for detecting bony CSI (Schenarts 2001). However, it is associated with a 90- to 200-fold increased radiation dose and subsequently a higher risk of developing thyroid cancer (Booth 2012; Jimenez 2008). The routine use of CT of the neck is therefore not recommended (NICE 2014). CT has a sensitivity

- of 23% and a specificity of 100% in detecting soft-tissue injury in children (Henry 2013), but the yield of positive, clinically-significant findings on CT of cervical spine injuries in children under five years old is low (Hernandez 2004).
- Magnetic resonance imaging (MRI) is an imaging modality that
  uses a magnetic field and radio waves instead of ionizing
  radiation. Compared to plain radiographs and CT, it detects a
  larger proportion of soft-tissue abnormalities when performed
  within 48 hours of injury (Benzel 1996; Keiper 1998). The
  properties of MRI might therefore suggest a larger role in blunt
  trauma evaluation of young children (Bagley 2006), but its role in
  screening after blunt trauma has not been widely explored, as it
  comprises multiple limitations (availability, time, costs), and the
  clinical significance of the injuries is sometimes unclear (Booth
  2012).

If plain radiography or CT, or both, are used in every patient, the amount of unnecessary imaging and associated radiation exposure will rise to unacceptable levels. Especially in children, excessive radiation exposure can lead to an elevated lifetime risk of developing cancer, primarily thyroid cancer (Jimenez 2008; Pearce 2012; Ron 2002). Unnecessary imaging will also lead to a rise in healthcare costs. To justify the use of diagnostic imaging, it is necessary to know when - and especially when not - to obtain radiographic imaging in children after blunt trauma, without the risk of missing injuries.

The type of CSI that occurs most often in children aged under two years is atlanto-occipital dislocation. In children aged two to seven years, atlanto-axial rotatory subluxation and atlanto-occipital dislocation occur most frequently. Among children aged eight to 15, subaxial vertebral body fractures are most common (Leonard 2013; Leonard 2014). Spinal cord injury is more frequent in the younger age groups (Leonard 2014).

Spinal cord injury without radiographic abnormalities (SCIWORA) on plain radiography or CT accounts for 4% to 66% of all spinal cord injuries in children (Pang 1982; Pang 2004; Trigylidas 2010; Yucesoy 2008). In younger children, falls and pedestrian motor vehicle accidents are the most common cause of SCIWORA. In older children, sports-related injuries are more common (Mortazavi 2011).

If radiographic imaging detects CSI, the therapeutic options for children depend on the type of injury, but might include fixating the cervical spine with a rigid or soft collar, brace, halo-frame, or by internal fixation (Mortazavi 2011). Corticosteroids are sometimes given, but this practice remains controversial (Bracken 1990; Leonard 2013; Leonard 2014).

Outcomes for patients differ according to the type of injury. Children with axial injury are nearly five times more likely to die than children with injuries in the subaxial region (23% versus 4%, respectively) (Patel 2001).

The long-term prognosis for children who sustain cervical spinal cord injury and survive the first 24 hours is poor; life expectancy is reduced by anywhere from six to 45 years, depending on the anatomical level of injury and the degree of spinal cord involvement (NSCISC 2012).

In the long term, people who suffer CSI can develop two types of progressive spinal deformity; (1) paralytic scoliosis, which



develops in 40% to 97% of children following spinal cord injury; and (2) cervical kyphosis secondary to laminectomy (although laminectomy after trauma is seldom necessary) (Seal 2005).

Due to prolonged hospital stay, the large number of treatments and associated long-term assistance, severe pediatric CSI is associated with very high medical, psychological, and societal costs (Vogel 2002a; Vogel 2002b; Vogel 2002c).

## Index test(s)

The tools under evaluation are the NEXUS criteria and the Canadian C-spine Rule (Viccellio 2001, as reported by Hoffman 2000; Stiell 2001). Both tools are designed to decide whether or not diagnostic imaging is needed to prove or rule out any type of CSI in adult trauma patients. There is notably little information about the accuracy of these two tools in children. Currently the Trauma

Association of Canada (Chung 2011), the guidelines of the Congress of Neurological Surgeons (Rozzelle 2013), and the National Institute for Health and Care Excellence (NICE) criteria recommend the use of a combination of these tools for the clearance of the cervical spine in children (NICE 2014).

Both tools consist of multiple scoring items. The NEXUS criteria observe only clinical parameters, while the Canadian C-spine Rule takes the trauma mechanism into account.

Both tools divide the population into high-risk and low-risk patients, where only the high-risk patients (positive index test) need further evaluation by radiographic imaging (see Figure 1: Viccellio 2001; Stiell 2003; and Figure 2: Stiell 2003; Stiell 2001). A negative index test (low-risk patients) means that the person is at low risk for CSI and therefore subsequent radiographic imaging is not necessary, with only clinical follow-up warranted.

Figure 1. NEXUS Criteria, Viccellio 2001 (from Hoffman 2000 paper); Stiell 2003

## The NEXUS Low-Risk Criteria

Cervical-spine radiography is indicated for patients with trauma unless the meet all of the following criteria:

No posterior midline cervical-spine tenderness,\*

No evidence of intoxication, \*\*

A normal level of alertness, \*\*\*


No focal neurologic deficit, \*\*\*\* and

No painful distracting injuries. \*\*\*\*\*

- \*Midline posterior bony cervical-spine tenderness is present if the patient reports pain on palpation of the posterior midline neck from the nuchal ridge to the prominence of the first thoracic vertebra, or if the patients evinces pain with direct palpation of any cervical spinous process.
- \* \* Patients should be considered intoxicated if they have either of the following: a recent history provided by the patient or an observer of intoxication or intoxicating ingestion, or evidence of intoxication on physical examination such as an odor of alcohol, slurred speech, ataxia, dysmetria, or other cerebellar findings, or any behavior consistent with intoxication. Patients may also be considered to be intoxicated if tests of bodily secretions are positive for alcohol or drugs that affect the level of alertness.
- \* \* \* An altered level of alertness can include any of the following: a Glascow Coma Scale score of 14 or less; disorientation to person, place, time, or event; an inability to remember three objects at five minutes; a delayed or inappropriate response to external stimuli; or other findings.
- \*\* \* \* A focal neurological deficit is any focal neurologic finding on motor or sensory examination.
- \*\*\*\*\* No precise definition of a painful distracting injury is possible. This category includes any condition thought by the clinician to be producing pain sufficient to distract the patient from a second (neck) injury. Such injuries may include, but are not limited to, any long-bone fracture; a visceral injury requiring surgical consultation; a large laceration, degloving injury, or crush injury; large burns; or any other injury causing acute functional impairment. Physicians may also classify any injury as distracting if it is thought to have the potential to impair the patient's ability to appreciated their injuries.



Figure 2. Canadian C-Spine Rules Stiell 2003; Stiell 2001



For patients with trauma who are alert ( as indicated by a score of 15 on the Glascow Coma Scale) and in stable condition and in whom cervical-spine injury is a concern, the determination of risk factors guides the use of cervical spine radiography. A dangerous trauma mechanism is considered to be a fall from an elevation > 3ft or 5 stairs; an axial load to the head (e.g., diving); a motor vehicle collision at high speed (>100km/hr) or with rollover or ejection; a collision involving a motorized recreational vehicle; or a bicycle collision. A simple rear-end motor vehicle collision excludes being pushed into oncoming traffic, being hit by a bus or a large truck, a rollover, and being hit by a high-speeds vehicle.

The low-risk criteria of the NEXUS tool are:

- no posterior midline cervical tenderness
- no evidence of intoxication
- the patient is alert and oriented to person, place, time, and event
- no focal neurological deficit
- no painful distracting injuries (e.g. long-bone fracture)

The high-risk criteria of the Canadian C-spine Rule are:

- age > 65 years
- dangerous trauma mechanism
- · paraesthesia in extremities

The low-risk criteria of the Canadian C-spine Rule are:

- · simple rear-end motor vehicle crash
- · sitting in emergency department
- ambulatory at any time
- delayed onset of neck pain
- absent midline C-spine tenderness
- able to actively rotate neck 45° left and right

The NEXUS criteria were developed in an almost entirely adult population. The study group consisted of 34,069 participants, of which 2.5% were eight years old or younger. In the participants with CSI, 1.3% were eight years old or younger (Ehrlich 2009; Viccellio



2001). The Canadian C-spine Rule included no pediatric patients (Ehrlich 2009; Stiell 2001).

For the development of both decision tools, plain radiography was used as the reference standard, unless a CT scan or MRI of the neck was performed because plain radiography was impractical or impossible (Viccellio 2001; Stiell 2001).

The role of the index test is to estimate whether a child is at high risk for CSI in order to determine whether or not further imaging is necessary. Application of the index test should prevent unnecessary radiographic imaging, thus limiting the amount of radiation exposure and healthcare costs. But most importantly, the main goal of the index tests is not to miss CSI.

## **Clinical pathway**

In general, when children present at the emergency department following blunt trauma, they undergo trauma work-up according to Advanced Trauma Life Support (ATLS 2013) or the Advanced Pediatric Life Support (APLS). One of the goals of the trauma work-up is to identify possible CSI. Clinicians therefore perform a series of anamnestic and physical tests to decide whether or not radiographic imaging of the cervical spine is necessary to rule out or define CSI. In most cases a trauma surgeon, a pediatric surgeon, or an emergency physician makes this decision.

The most commonly-used tools for identifying patients eligible for imaging are the NEXUS criteria and the Canadian C-spine Rule. Both tools divide the population into high-risk and low-risk patients, where only the high-risk patients need further evaluation by radiographic imaging.


If no cervical spine injury is suspected based on clinical findings using one of these decision tools, the neck collar (if worn) can be removed and after clinical follow-up the cervical spine is cleared. However, if cervical spine injury is suspected, radiographic imaging is needed to rule out or define CSI.

The first-choice imaging modality in children is plain radiography of the cervical spine. A lateral view and an anteroposterior view are performed in all age groups; from nine years old an odontoid radiograph (open-mouth view) is also obtained. Under the age of nine, the open-mouth view provides no additional information beyond that found on anteroposterior and lateral views (Buhs 2000).

Several guidelines advise performing a CT scan of the cervical spine if the patient has neurological symptoms, the patient is intubated, or the Glasgow Coma Scale is less than 13 at initial assessment (Chung 2011; NICE 2014; Rozzelle 2013). They also advise a CT scan if plain radiography is technically difficult or inadequate, if bony injury is seen at plain radiography, or if there is a strong clinical suspicion of injury despite normal radiographs. MRI of the cervical spine is recommended in patients in whom plain radiography or CT scans are equivocal, or when neurological or ligamentous injury is suspected (NICE 2014). For a flowchart of the clinical pathway, see Figure 3.



Figure 3. Flow chart clinical practice



## Role of index test(s)

So far, little information is known about the applicability of the NEXUS criteria and the Canadian C-spine Rule in children. However, using one or both tools is already the standard of care in many centres for trauma evaluation in children (Chung 2011; NICE 2014; Rozzelle 2013). If the NEXUS criteria and the Canadian C-spine Rule are accurate in defining the need for radiographic imaging in children after blunt trauma, unnecessary imaging can be prevented and radiation exposure and healthcare costs can be reduced.

## Rationale

Since CSI has severe consequences such as spinal cord injury or death (Cirak 2004; Givens 1996; Hutchings 2009; Kokoska 2001; Parent 2011; Patel 2001; Platzer 2007), identifying these injuries during trauma work-up is of great importance. The main purpose the NEXUS criteria and Canadian C-spine Rule is to help the physician decide whether or not a patient needs radiographic imaging.

Evaluation of the cervical spine in a young patient is challenging. Firstly, young children may not be able to communicate crucial symptoms. The physical examination can be compromised in an anxious, crying, or uncooperative child (Ehrlich 2009). Furthermore,

pediatric cervical anatomy differs from that of adults (Cattell 1965; Ehrlich 2009; Fesmire 1989; Kriss 1997; Swischuk 1977; Swischuk 1984).

In conclusion, pediatric cervical spine assessment is different when compared to adults. It is therefore questionable whether these clinical decision tools developed for clearing the cervical spine in an adult population can be applied directly to a pediatric population.

## OBJECTIVES

To determine the diagnostic accuracy of the NEXUS criteria and the Canadian C-spine Rule in a pediatric population evaluated for CSI following blunt trauma.

## **Secondary objectives**

We expected to observe heterogeneity in the estimates of diagnostic accuracy. Factors that may contribute to heterogeneous results include:

1. Differences in the healthcare setting and study design; previous studies state that CSI is seen less often in general emergency departments than in pediatric trauma hospitals, and that a CT



of the neck is more common in general emergency departments (Adelgais 2014).

- Study quality, as assessed by the QUADAS-2 checklist (Whiting 2011).
- Age-related differences; we expect that both the applicability of the NEXUS criteria and the Canadian C-spine Rule and the type of injury will differ according to age in children younger and older than eight (Leonard 2013; Leonard 2014).

#### **METHODS**

## Criteria for considering studies for this review

## Types of studies

We included studies that compared the accuracy of the NEXUS criteria or the Canadian C-spine Rule, or both, with the reference standard.

We considered diagnostic studies with cross-sectional or cohort designs (retrospective or prospective), and randomized controlled trials. We only included results from full reports.

We preferred to include studies in which plain radiography, CT and MRI of the cervical spine was performed, since the sensitivity and specificity for the detection of ligamentous and skeletal injury differ between those imaging modalities. However, we expected this triple combination to be very rare, due to multiple limitations (radiation consequences, availability, time and costs). We therefore also included studies in which only one of the reference standards was applied. For the same reason we also included studies in which no radiographic imaging was obtained but only clinical follow-up (in case of a negative index test).

We expected that there would be studies in which all patients had undergone the reference standard without the index test being scored positive. We therefore also included studies in which those participants who had a negative result of the index test (no highrisk factors) still underwent one of the reference standards (for example, plain radiography).

We excluded case-control studies because of the bias they might introduce.

We preferred studies that evaluated the accuracy of the NEXUS criteria and the Canadian C-spine Rule in direct comparisons to each other, but we did not exclude studies with indirect comparisons, because we expected the number of studies with direct comparisons to be limited.

## **Participants**

We Included children between the age of 0 and 18 who underwent blunt trauma evaluation in the emergency department. We excluded patients with a history of previous surgery of the cervical spine or congenital cervical spine anomalies, or both.

In case of studies with mixed populations, e.g. that included some participants in the groups and that could not be separated from the eligible participants, we tried to contact the study author to provide the data for the group of interest.

#### **Index tests**

The tests under evaluation are the NEXUS criteria and the Canadian C-spine Rule, or tools in which the clinical variables contained four or more elements that are also part of the NEXUS criteria or the Canadian C-spine Rule.

#### **Target conditions**

The target condition was clinically-important CSI, defined as any fracture, dislocation, or ligamentous instability detectable by diagnostic imaging.

#### **Reference standards**

Radiographic imaging (plain radiography, CT or MRI or both) or clinical follow-up (if the index test was scored negative) were the reference standards. Follow-up was defined as clinical evaluation of the neck after removal of the neck collar (if worn) in the case of low-risk patients. The first clinical follow-up is performed in the emergency room during trauma evaluation. If cervical spine injury is suspected based on the index test, or after the first clinical follow-up, radiographic imaging is needed to rule out or define CSI.

We included patients who underwent an eligible reference standard within 72 hours of presentation at the emergency department following blunt trauma. We also included patients who did not undergo radiographic imaging, but who obtained clinical follow-up directly during trauma evaluation during the first 72 hours.

## Search methods for identification of studies

#### **Electronic searches**

The search strategies for identifying diagnostic test accuracy studies consisted of controlled vocabulary and keyword terms for each of the following concepts: the index or reference test, the target condition, and the patient description.

The Information Specialist of the Cochrane Back and Neck Review Group (CBN) developed the search strategies. Chapter 7 'Searching for Studies' of the *Cochrane Handbook for Systematic Reviews of Diagnostic Test Accuracy* (De Vet 2008), as well as feedback and guidance from Information Specialists of the Cochrane DTA Working Group and Cochrane Netherlands, were used to guide strategy development and the selection of databases to be searched.

We searched the following databases from inception to 24 February 2015:

- Cochrane Central Register of Controlled Trials (CENTRAL, the Cochrane Library, Issue 1)
- MEDLINE (OvidSP, 1946 to February Week 3 2015)
- MEDLINE Non-Indexed and In-Process Citations (OvidSP, 23 February 2015)
- Embase (OvidSP, 1980 to 2015 Week 08)
- Science Citation Index (Web of Science, Core Collection)
- ProQuest Dissertations & Theses Database for relevant conference proceedings, dissertations, and theses
- PubMed
- · OpenGrey for 'grey literature'



We searched PubMed to identify studies not in MEDLINE, using the strategy recommended by Duffy 2014. We also searched ClinicalTrials.gov and the World Health Organization (WHO) International Clinical Trials Registry Platform (ICTRP) for ongoing trials. An Information Specialist from the Cochrane Renal Group searched the Diagnostic Test Accuracy (DTA) Studies Register on 10 March 2015.

We also searched the Cochrane Database of Systematic Reviews (CDSR, Issue 2), the Database of Abstracts of Reviews of Effects (DARE, Issue 1), and the Health Technology Assessment (HTA, Issue 1) databases (through the Cochrane Library); and the Aggressive Research Intelligence Facility (ARIF) database for relevant systematic reviews and health technology assessments on 25 February 2015. We searched The Medion Database October 2013. It was not accessible in February 2015.

The search strategies can be found in Appendix 1. The citations per search engine are shown in Table 1.

#### **Searching other resources**

We searched the reference lists of all primary studies and relevant systematic reviews to identify additional studies. There were no language restrictions. Non-English articles were translated.

## **Data collection and analysis**

#### Selection of studies

Two review authors (AS and MMF) assessed titles and abstracts of the search results provided by the Information Specialist. Two review authors (AS and MMF) independently retrieved all potentially relevant articles in full text and examined them, using an assessment form, and resolving discrepancies by discussion. If there was a difference in coding, we approached a third review author (NWLS) to reach consensus.

## Data extraction and management

Two review authors (AS and JW) extracted data separately to a data extraction form, resolving disagreements by consultation with a third review author (NWLS, MM, or RRvR) if necessary. We contacted study authors to provide additional information.

Data collected included:

- 1. Study ID: year of publication, author
- 2. Participants: number of children, age, sex, country
- 3. Inclusion and exclusion criteria
- 4. Study design: consecutive/random, retrospectively/ prospectively
- 5. Clinical parameters: trauma mechanism, clinical pathway
- 6. Target condition: cervical spine injury
- Reference standard: clinical follow-up, plain radiography, CT scan, MRI
- Results: numerical data from a two-by-two table. If no two-by-two tables were available, we extracted data from the text. If
  there was a difference between the sensitivity or specificity, or
  both, between table and text, we used the numbers provided in
  the text.

## Assessment of methodological quality

Two review authors (AS and MMF) independently assessed methodological quality in duplicate, using QUADAS-2 (Whiting 2003; Whiting 2011; Table 2; Table 3) . We resolved any disagreements in a consensus meeting, involving a third review author when necessary.

#### Statistical analysis and data synthesis

We extracted indices of the diagnostic performance of all clinical tools from data presented in each study. We generated diagnostic two-by-two tables, from which we calculated sensitivities and specificities for each index test with 95% confidence intervals, and presented them in forest plots and also in a receiver operator curve (ROC) space. If data presented in trials were uninterpretable to generate two-by-two tables, we contacted the authors of the study to clarify.

We planned to perform meta-analyses of sensitivity and specificity employing a bivariate logistic normal model using a hierarchical approach (Reitsma 2005). This approach enabled us to calculate summary estimates of sensitivity and specificity while dealing with sources of variation within and between studies and any correlation that might exist between sensitivity and specificity. With the model estimates, we plotted sensitivities and specificities in forest plots and in ROC space.

We also planned to compare the different index tests and tried to find whether these tests have different sensitivities or specificities, employing a bivariate model.

We planned to do test comparison by adding covariates for different types of index tests into the bivariate model and testing the significance (significance level = 0.05) of the parameters of covariates. If almost none of the primary studies directly compared these tools, we would have included all studies that evaluate at least one of the index tests into the test comparison. In other words, test comparison would not be limited to direct comparisons, but would use all the evidence available. We planned to compare them qualitatively if data were insufficient for comparison by statistical test. Verification bias is to be expected, since we enable the use of different types of reference standards in test-positive (plain radiography, CT scan, or MRI) and test-negative (clinical follow-up). All the statistical analyses were performed with Review Manager 5 (Review Manager 2014)

## **Investigations of heterogeneity**

We were unable to formally explore heterogeneity, due to a lack of studies. Should there be sufficient data available we used forest plots and sensitivities and specificities plotted in ROC space for visual examination of heterogeneity between studies. We also added covariates, for example age groups (less than eight years old versus eight and older) and QUADAS-2 items to bivariate model to investigate the heterogeneity between studies in the meta-analysis (Whiting 2011). We could investigate heterogeneity only if there were a sufficient number of studies providing adequate information on the factor of interest.

## Sensitivity analyses

Verification bias is the most important form of bias that we expected to encounter, because we evaluated four different types



of reference standards. We could not conduct sensitivity analyses because we retrieved too few studies.

## **Assessment of reporting bias**

As yet there are no quantitative methods for reporting bias in diagnostic test accuracy studies; we therefore did not assess reporting bias.

## RESULTS

## Results of the search


We identified 14,730 citations. The number of citations by search engine is shown in Table 1. We checked the references of relevant reviews and primary diagnostic studies. After initial title and abstract screening, we retrieved 98 full-text papers, of which we considered five to be eligible for the review. Two papers reported

on the same cohort with pediatric data only extractable from one of the papers, which we used as the primary data source (Viccellio 2001). We excluded Hoffman 1992, an unrelated study, as only a small number of children were included and the authors did not separate out the data for the pediatric population. We therefore include three studies (five papers) in the meta-analysis: one was a cohort study with a prospective design (Viccellio 2001) and two were cohort studies with retrospective designs (Ehrlich 2009; Jaffe 1987). There were no disagreements between the review authors on studies eligible for the review. There were no ongoing studies identified and no studies are awaiting classification.

Figure 4 shows the summary of the search, including the reason for excluding papers. Exclusion mainly concerned studies with a different index test (n = 45) and studies with a different study population (n = 26). The latter was explained by the fact that most of these studies were conducted in adults (n = 18).



Figure 4. Study flow diagram.





## Figure 4. (Continued)


references)
included in critical
appraisal
Direct
comparison: 1
NEXUS criteria
alone: 2

We provide details on the design, setting, population, reference standard and definition of the target condition in the Characteristics of included studies table. Details of the excluded studies are shown in the Characteristics of excluded studies table.

## Methodological quality of included studies

The results of the quality assessment for three included studies are presented in Figure 5.

Figure 5. Risk of bias and applicability concerns summary: review authors' judgements about each domain for each included study



We rated two studies as being at low risk of concern for applicability in all domains (Ehrlich 2009; Viccellio 2001). We judged one study to be at high risk of bias and high concern for applicability in the domain of the index test (Jaffe 1987). This was because this study was a derivation study and did not evaluate the NEXUS criteria or the Canadian C-spine Rule; however, the index test of this study contained many items that are also part of the NEXUS criteria. We also rated Jaffe 1987 at high risk of bias in the domains of 'patient selection' and 'flow and timing'. This was based on them not including a consecutive group of patients, but using a second cohort to increase the number of patients with CSI. We judged the study as being at unclear risk of bias in the domain 'index test and

reference standard', because it was not clear whether the reference standard results were interpreted without knowledge of the results of the index test and vice versa. We also judged it as being at unclear risk of bias, since it was a derivation study and therefore no threshold was described for the index test.

Ehrlich 2009 poorly described whether the index test results were interpreted without knowledge of the results of the reference standard, and we therefore rated the domain of the reference standard and index test as being at unclear risk of bias. Ehrlich 2009 also used different types of reference standards, with patients excluded for no clear reason; we therefore rated the domain 'flow and timing' as being at high risk of bias. However, for this review



we have only focused on the cohort that was used for analysing the accuracy of the NEXUS and Canadian C-spine Rule, and this cohort only used one type of reference standard.

In Viccellio 2001 plain radiography and sometimes a CT scan were obtained as the primary imaging modality, so we judged this study to be at high risk of bias in the domain 'flow and timing'.

None of the studies thoroughly explained the interval between the index test and reference standard, but all three studies took place in the emergency department and confirmed that radiographic imaging was conducted there. The interval was therefore less than one day in all studies.

The review authors (AS and MMF) disagreed on only two items of the QUADAS-2 criteria in three studies (Whiting 2011). All disagreements were resolved during consensus meetings.

## **Findings**

We evaluated three studies (Ehrlich 2009; Jaffe 1987; Viccellio 2001) which included 3380 children, for the presence of CSI by using the NEXUS criteria or the Canadian C-spine Rule. The overall incidence of CSI in these three studies was 2.8%.

The three studies described four patient cohorts. In total 3379 children were evaluated for CSI by applying the NEXUS criteria, and 109 children were evaluated for CSI by applying the Canadian C-spine Rule. Around one-third of the patients included in the studies were younger than eight years old. Ninety-six children were diagnosed with CSI, with 10% to 20% of them aged less than eight. In Ehrlich 2009, all included children where younger than 10 years old, but a further division by age was not provided. Jaffe 1987 made a distinction between children aged up to three and those aged four to 12 years, so we do not know the exact number of children aged under eight.

Jaffe 1987 was a derivation study to develop a clinical decision tool for the clearance of the cervical spine. They included an extra set of patients to increase the amount of those with CSI. The index test criteria differed from the NEXUS criteria, but the items that were analysed were almost the same as the NEXUS criteria, and we therefore included this study in the analysis. Plain radiography was obtained for all children as the primary imaging modality.

Viccellio 2001 evaluated the NEXUS criteria in 3065 children. As well as plain radiography, a CT was also obtained as the primary imaging modality in some of these children. The number of children for whom a CT was obtained was not clear.

Ehrlich 2009 was the only study that tested the accuracy of the Canadian C-spine Rule in a direct comparison to the NEXUS criteria. The other two studies did not test the accuracy of the Canadian C-spine Rule (Jaffe 1987; Viccellio 2001). The sensitivity and specificity presented by Ehrlich 2009 were not the same when we did our own calculations with the numbers provided in the text. We contacted the authors to invite them to explain the differences between the numbers provided in the text and the sensitivity and specificity that they had calculated, but did not receive a response. We therefore produced our own two-by-two table using the numbers provided in the text.

Since the number of eligible studies was fewer than four, and since the outcomes of those studies were too diverse, we did not conduct a meta-analysis and therefore present no summary estimates in this review. Instead, we interpret sensitivity and specificity from each primary study separately. Since there were few data, we were unable to investigate heterogeneity. We summarize the extracted data in the forest plots presented in Figure 6 and in Summary of findings 1.

Figure 6. Forest plot of the NEXUS criteria and Canadian C-spine Rule.

#### NEXUS Study TΡ TN Sensitivity (95% CI) Specificity (95% CI) Sensitivity (95% CI) Specificity (95% CI) Ehrlich 2009 4 66 3 35 0.57 [0.18, 0.90] 0.35 [0.25, 0.45] Jaffe 1987 58 68 1 79 0.98 [0.91, 1.00] 0.54 [0.45, 0.62] Viccellio 2001 30 2432 0 603 1.00 [0.88, 1.00] 0.20 [0.18, 0.21] Canadian C-spine Rule Specificity (95% CI) Sensitivity (95% CI) Study TP FP FN TN Sensitivity (95% CI) Specificity (95% CI) Ehrlich 2009 0.86 [0.42, 1.00] 0.15 [0.08, 0.23]


The sensitivity of the NEXUS criteria in the studies was 0.57 (95% CI 0.18 to 0.90) (Ehrlich 2009), 0.98 (95% CI 0.91 to 1.00) (Jaffe 1987) and 1.00 (95% CI 0.88 to 1.00) (Viccellio 2001). The specificity of the NEXUS criteria varied in the studies and was 0.35 (95% CI 0.25 to 0.45) (Ehrlich 2009), 0.54 (95% CI 0.45 to 0.62) (Jaffe 1987) and 0.2 (95% CI 0.18 to 0.21) (Viccellio 2001). For the Canadian C-spine Rule,

the sensitivity was 0.86 (95% CI 0.42 to 1.00) and specificity was 0.15 (95% CI 0.08 to 0.23) (Ehrlich 2009).

The positive and negative predictive values and the positive and negative likelihood ratios for the individual studies are summarized in Figure 6; and Summary of findings 1. The test accuracy of both tests in individual studies is shown in Figure 7.



Figure 7. Summary ROC Plot of tests: 1 NEXUS, 2 Canadian C-spine Rule.



Verification bias was to be expected, since plain radiography and a CT scan were used as reference standards in Viccellio 2001. We could not perform sensitivity analyses because there were too few studies.

# DISCUSSION

## **Summary of main results**

This review evaluates the accuracy of the NEXUS criteria and Canadian C-spine Rule in children following blunt trauma.

Although the quality of the included studies that assessed the accuracy of the NEXUS criteria or the Canadian C-spine Rule or both was moderate, results are only based on three studies. Due to the limited number of studies and the fact that the heterogeneity between the three studies was significant (as seen in Figure 6 and Figure 7), we could not conduct a meta-analysis.

The incidence of CSI in the included studies was 2.8%. A possible explanation is the non-consecutive group of patients included in Jaffe 1987. Although the incidence of CSI is higher than in previous literature (Patel 2001), it is still very low, meaning that to properly



evaluate the accuracy of the NEXUS criteria and the Canadian C-spine Rule we need a large sample size. The total sample size of the included studies in this review was 3380, of which 3065 patients were from Viccellio 2001.

Ehrlich 2009 was the only study that evaluated the accuracy of the Canadian C-spine Rule in a pediatric population. When we recalculated the sensitivity and specificity reported in the text with the numbers provided, we did not obtain the same results; this undermines the reliability of the sensitivity of the Canadian C-spine Rule reported in this study. Since this study was the only one evaluating the Canadian C-spine Rule, we cannot verify the diagnostic accuracy of the Canadian C-spine Rule in a pediatric population. There is therefore too little information to say that the Canadian C-spine Rule can be used safely for the clearance of cervical spine injury in children following blunt trauma.

The sensitivity of the NEXUS criteria was relatively high in two of the studies (Jaffe 1987; Viccellio 2001). Jaffe 1987 was a derivation study to develop a clinical decision tool, which might have led to a higher sensitivity in this study. They also enriched their cohort with children with CSI, which may have led to a sensitivity that is not representative of the population. The other study was a validation study of the NEXUS criteria (Viccellio 2001). This study included the highest number of children and showed a sensitivity of 1.00 (95% CI 0.88 to 1.00). The third study showed a lower sensitivity of 0.57 (95% CI 0.18 to 0.9) (Ehrlich 2009). In this study patients were referred for radiographic imaging if they met one of the highrisk criteria included in the NEXUS tool. By applying these criteria, seven patients were diagnosed with CSI using plain radiography. However, after re-evaluation using the NEXUS criteria on the same cohort, the authors stated that three fractures would have been missed. This is likely to lead to an underestimation of the accuracy of the NEXUS criteria in this study.

The confidence intervals of the sensitivity in the studies showed a wide range, with a lower limit varying from 0.18 to 0.91. This implies that if physicians use the NEXUS criteria in children, there is a chance of missing CSI. Moreover, they have to keep in mind that the sensitivity differs amongst several studies. Since missing CSI could have severe consequences with the risk of significant morbidity, we consider that the NEXUS criteria are at best a guide to clinical assessment as current evidence does not support strict or protocolized adoption of the tool into pediatric trauma care (Cirak 2004; Givens 1996; Hutchings 2009; Kokoska 2001; Parent 2011; Patel 2001; Platzer 2007).

The confidence interval of the specificity of the studies of both the NEXUS criteria and the Canadian C-spine Rule did not show as wide a range as the confidence interval for sensitivity. The confidence interval of the specificity had a lower limit varying from 0.08 to 0.45. The specificity of all studies varied from 0.2 to 0.54. This means that when the NEXUS criteria or the Canadian C-spine Rule are applied in pediatric trauma patients, one of the effects will be that, in several children, radiographic imaging will be obtained without showing CSI. However the aim of these triage tools is to reduce the chance of missing CSI, so the sensitivity needs to be high, which indirectly leads to a low specificity.

## Strengths and weaknesses of the review

This is the first systematic review on the accuracy of the NEXUS criteria and the Canadian C-spine Rule in children.

One of the strengths of this review is that we performed an extensive search in a large number of databases. We selected articles by using clear inclusion and exclusion criteria. Another strength of this review is that we evaluated the evidence using the QUADAS-2 tool (Whiting 2011). This tool provides important information about potential sources of bias and enables a simple and clear presentation of the assessment.

One of the limitations of our review was that only a small number of studies were eligible for inclusion. We therefore could not conduct sensitivity analyses or formally investigate potential sources of heterogeneity. We excluded Hoffman 1992 after we contacted the authors of the study for pediatric data. They explained that they did not complete a detailed examination of the pediatric population in that cohort, because the derivation study had only a small number of children. For this reason, we do not think that excluding this study has a significant effect on the results of this review.

Only one study tested the accuracy of the Canadian C-spine Rule by direct comparison to the NEXUS criteria (Ehrlich 2009); the other two studies did not evaluate the accuracy of the Canadian C-spine Rule. We therefore cannot state whether the NEXUS or the Canadian C-spine Rule is a better tool to decide if imaging is indicated to detect CSI in children following blunt trauma. Also, the numbers provided in this study for the sensitivity and the specificity of the tools were not correct (Ehrlich 2009), and since this was the only study that evaluated the accuracy of the Canadian C-spine Rule there is not enough evidence to determine its accuracy in children. Another limitation is that one study developed an algorithm with items that are also included in the NEXUS criteria. However, this algorithm did not contain all items of the NEXUS criteria.

Another weakness of this review is that the results are based on only a few children diagnosed with CSI, although the incidence was higher than in previous literature (Patel 2001). A larger sample size would be desirable to better evaluate the accuracy of the two triage tools, which should indirectly lead to a higher number of events (patients with CSI).

In this review we have only focused on two decision tools that determine whether children following blunt trauma need to be referred for radiography. In the last two decades, several institutional decision tools have been developed to evaluate the need for referral for radiography in children after blunt trauma, to rule out or diagnose CSI (Anderson 2006; Anderson 2010; Edwards 2001; Hutchings 2009; Pieretti-Vanmarcke 2011). However, these newly-developed decision tools are mostly used in single institutions and none of them is validated in another cohort, let alone evaluated in an implementation study.

## Applicability of findings to the review question

The aim of this review was to evaluate if the NEXUS criteria and the Canadian C-spine Rule are accurate decision tools for detecting CSI in pediatric trauma patients following blunt trauma. Only one study examined the accuracy of the Canadian C-spine Rule (Ehrlich 2009). The NEXUS criteria were assessed in a pediatric population in three studies, but only one study (Viccellio 2001) evaluated the accuracy of the NEXUS criteria in a manner that corresponded to the research question of this review. Only one-third of the patients included in all three studies were younger than eight years old, and only an estimated 10% to 20% of the children with CSI were younger than eight.



#### **AUTHORS' CONCLUSIONS**

#### Implications for practice

At the moment, there is not enough evidence to determine if the Canadian C-spine Rule is accurate in detecting CSI in pediatric patients following blunt trauma. Therefore the evidence does not support the use of the Canadian C-spine Rule can to detect or rule out CSI in pediatric trauma patients.

The information available on the accuracy of the NEXUS criteria in a pediatric population is sparse and based on a small number of CSI events. Clinicians must keep in mind that the sensitivity of the NEXUS criteria differs among the three studies, with a wide range of individual confidence intervals for their sensitivity. This means that there is a chance of missing CSI when only relying on the NEXUS criteria to evaluate the need for imaging in children following blunt trauma. We therefore consider that the NEXUS criteria are at best a guide to clinical assessment as the evidence does not support strict or protocolized adoption of the tool into pediatric trauma care.

The specificity of the triage tools was low. However, in triage tools the main goal is to reduce the chance of missing CSI, so the sensitivity needs to be high, which indirectly leads to a low specificity. If the specificity were higher, fewer radiographic images would be obtained, but this would lead to a decrease in sensitivity and thus an increase in the chance of missing CSI. Moreover, even a small reduction in the amount of radiographic imaging is better than obtaining radiographic imaging of the cervical spine on a routine basis.

Data on children under the age of eight is especially sparse; there is therefore currently is no evidence to support the use of these clinical decision tools in this age group.

## Implications for research

Since the incidence of CSI in children is low, a large number of patients are needed to test the accuracy of a clinical decision tool.

Hence, future research should focus on large adequately powered multicenter prospective trials to assess the accuracy of the NEXUS criteria and Canadian C-spine Rule in children. Only then can we determine whether the NEXUS criteria or Canadian C-spine Rule are sensitive and specific enough to be applied as a decision tool following blunt trauma. It would be important to include enough patients younger than eight years old to ensure the decision tools could be used in children of all ages. Also, the number of CSI events should be high enough. A study should optimally evaluate both the NEXUS criteria and Canadian C-spine rule in all pediatric trauma patients. The first step would be plain radiographic imaging in pediatric trauma patients after clinical evaluation, according to all the items of the NEXUS criteria and Canadian C-spine Rule, so all items of both triage tools would be taken into consideration to evaluate whether or not further radiographic imaging should be obtained. In this way we can evaluate the sensitivity and specificity of both triage tools for predicting the need for radiographic imaging of the cervical spine after blunt trauma in children. As in other studies, clinical follow-up of patients in whom no radiographic imaging is necessary should be considered according to the NEXUS criteria and the Canadian C-spine Rule, thereby reducing the likelihood of missing CSI. In planning the study, trialists should conduct a power analysis to determine how many children younger and older than eight should be included, and how many events (patients with CSI) would be required.

#### ACKNOWLEDGEMENTS

We thank Rene Spijker, Information Specialist of Cochrane Netherlands and Susan Bayliss, Information Specialist of the Cochrane DTA Working Group for their help with the development of the literature search, and Shireen Harbin of Cochrane Back and Neck for developing the search strategy. We also thank Mariska Leeflang, Cochrane member and epidemiologist at the Academic Medical Centre in Amsterdam (the Netherlands) for her contribution in drafting the protocol. Finally we thank Cochrane Child Health reviewers Dr. Brett Rocos and Dr. Warwick Teague for clinical peer review.



#### REFERENCES

#### References to studies included in this review

#### Ehrlich 2009 (published data only)

Ehrlich PF, Wee C, Drongowski R, Rana AR. Canadian C-spine Rule and the National Emergency X-Radiography Utilization low-risk criteria for C-spine radiography in young trauma patients. *Journal of Pediatric Surgery* 2009;**44**(5):987-91.

## Jaffe 1987 {published data only}

Jaffe DM, Binns H, Radkowski MA, Barthel MJ, Engelhard HH 3rd. Developing a clinical algorithm for early management of cervical spine injury in child trauma victims. *Annals of Emergency Medicine* 1987;**16**(3):270-6.

#### Viccellio 2001 (published data only)

Hoffman JR, Mower WR, Wolfson AB, Todd KH, Zucker MI. Validity of a set of clinical criteria to rule out injury to the cervical spine in patients with blunt trauma. National Emergency X-Radiography Utilization Study Group.. *New England Journal of Medicine* 2000;**343**(2):94-9.

Simon H, Viccellio P, Pressman B, Shah M, Hoffman J, Mower W, et al. A prospective evaluation of cervical spine injuries in children and a decision model for radiography. *Pediatric Research* 2000;**47**(4):119A.

\* Viccellio P, Simon H, Pressman BD, Shah MN, Mower WR, Hoffman JR. A prospective multicenter study of cervical spine injury in children. *Pediatrics* 2001;**108**(2):e20.

#### References to studies excluded from this review

## Adelgais 2014 (published data only)

Adelgais KM, Browne L, Holsti M, Metzger RR, Murphy SC, Dudley N. Cervical spine computed tomography utilization in pediatric trauma patients. *Journal of Pediatric Surgery* 2014;**49**(2):333-7.

#### Anderson 2006 (published data only)

Anderson RC, Kan P, Hansen KW, Brockmeyer DL. Cervical spine clearance after trauma in children. *Neurosurgical Focus* 2006;**20**(2):E3.

#### Anderson 2010 (published data only)

Anderson RC, Kan P, Vanaman M, Rubsam J, Hansen KW, Scaife ER, et al. Utility of a cervical spine clearance protocol after trauma in children between 0 and 3 years of age. *Journal of Neurosurgery* 2010;**Pediatrics**(3):292-6.

## Baker 1999 {published data only}

Baker C, Kadish H, Schunk JE. Evaluation of pediatric cervical spine injuries. *American Journal of Emergency Medicine* 1999;**17**(3):230-4.

## Bandiera 2003 (published data only)

Bandiera G, Stiell IG, Wells GA, Clement C, De V Maio, Vandemheen KL, et al. The Canadian C-spine rule performs better than unstructured physician judgment. *Annals of Emergency Medicine* 2003;**42**(3):395-402.

#### Banit 2000 (published data only)

Banit DM, Grau G, Fisher JR. Evaluation of the acute cervical spine: a management algorithm. *Journal of Trauma-Injury Infection & Critical Care* 2000;**49**(3):450-6.

## **Bayless 1989** {published data only}

Bayless P, Ray VG. Incidence of cervical spine injuries in association with blunt head trauma. *American Journal of Emergency Medicine* 1989;**7**(2):139-42.

#### **Blacksin 1995** {published data only}

Blacksin MF, Lee HJ. Frequency and significance of fractures of the upper cervical spine detected by CT in patients with severe neck trauma. *AJR. American Journal off Roentgenology* 1995;**165**(5):1201-4.

### Borock 1991 (published data only)

Borock EC, Gabram SG, Jacobs LM, Murphy MA. A prospective analysis of a two-year experience using computed tomography as an adjunct for cervical spine clearance. *Journal of Trauma* 1991;**31**(7):1001-6.

#### **Brockmeyer 2012** {published data only}

Brockmeyer DL, Ragel BT, Kestle JR. The pediatric cervical spine instability study. A pilot study assessing the prognostic value of four imaging modalities in clearing the cervical spine for children with severe traumatic injuries. *Child's Nervous System: ChNS* 2012;**28**(5):699-705.

## **Brooks 2001** {published data only}

Brooks RA, Willett KM. Evaluation of the Oxford protocol for total spinal clearance in the unconscious trauma patient. *Journal of Trauma-Injury Infection & Critical Care* 2001;**50**(5):862-7.

## **Brown 2001** {published data only}

Brown RL, Brunn MA, Garcia VF. Cervical spine injuries in children: a review of 103 patients treated consecutively at a level 1 pediatric trauma center. *Journal of Pediatric Surgery* 2001;**36**(8):1107-14.

## Browne 2003 (published data only)

Browne GJ, Lam LT, Barker RA. The usefulness of a modified adult protocol for the clearance of paediatric cervical spine injury in the emergency department. *Emergency Medicine* (Fremantle, W.A.) 2003;**15**(2):133-42.

## Burns 2011 (published data only)

Burns EC, Yanchar NL. Using cervical spine clearance guidelines in a pediatric population: a survey of physician practices and opinions. *CJEM: Canadian Journal of Emergency Medical Care* 2011;**13**(1):1-6.

## Clayton 2012 (published data only)

Clayton JL, Harris MB, Weintraub SL, Marr AB, Timmer J, Stuke LE, et al. Risk factors for cervical spine injury. *Injury* 2012;**43**(4):431-5.



## Coffey 2011 (published data only)

Coffey F, Hewitt S, Stiell I, Howarth N, Miller P, Clement C, et al. Validation of the Canadian c-spine rule in the UK emergency department setting. *Emergency Medicine Journal* 2011;**28**(10):873-6.

#### Como 2011 {published data only}

Como JJ, Leukhardt WH, Anderson JS, Wilczewski PA, Samia H, Claridge JA. Computed tomography alone may clear the cervical spine in obtunded blunt trauma patients: a prospective evaluation of a revised protocol. *Journal of Trauma-Injury Infection & Critical Care* 2011;**70**(2):345-9.

## **Dahlquist 2013** {published data only}

Dahlquist RT, Fischer PE, Desai H, Rogers A, Christmas AB, Gibbs MA, et al. Is cervical spine imaging required in patients with femur fractures?. *Academic Emergency Medicine*. *Conference: 2013 Annual Meeting of the Society for Academic Emergency Medicine*, SAEM 2013 Atlanta, GA United States. 2013 May 14-18 2013;**20**(5 SUPPL 1):S58.

#### **Dickinson 2004** {published data only}

Dickinson G, Stiell IG, Schull M, Brison R, Clement CM, Vandemheen KL, et al. Retrospective application of the NEXUS low-risk criteria for cervical spine radiography in Canadian emergency departments. *Annals of Emergency Medicine* 2004;**43**(4):507-14.

#### **DiGiacomo 2002** {published data only}

DiGiacomo JC, Frankel HL, Rotondo MF. Clearing the cervical spine in victims of blunt trauma. *Military Medicine* 2002;**167**(5):398-401.

## Edwards 2001 {published data only}

Edwards MJ, Frankema SP, Kruit MC, Bode PJ, Breslau PJ, Van Vugt AB. Routine cervical spine radiography for trauma victims: Does everybody need it?. *Journal of Trauma-Injury Infection & Critical Care* 2001;**50**(3):529-34.

## Ersoy 1995 (published data only)

Ersoy G, Karcioglu O, Enginbas Y, Eray O, Ayrik C. Are cervical spine X-rays mandatory in all blunt trauma patients?. *European Journal of Emergency Medicine* 1995;**2**(4):191-5.

## Fischer 1984 {published data only}

Fischer RP. Cervical radiographic evaluation of alert patients following blunt trauma. *Annals of Emergency Medicine* 1984;**13**(10):905-7.

#### Garton 2008 (published data only)

Garton HJ, Hammer MR. Detection of pediatric cervical spine injury. *Neurosurgery* 2008;**62**(3):700-8.

## Gbaanador 1986 {published data only}

Gbaanador GB, Fruin AH, Taylon C. Role of routine emergency cervical radiography in head trauma. *American Journal of Surgery* 1986;**152**(6):643-8.

#### Gonzalez 1999 (published data only)

Gonzalez RP, Fried PO, Bukhalo M, Holevar MR, Falimirski ME. Role of clinical examination in screening for blunt cervical spine injury. *Journal of the American College of Surgeons* 1999:**189**(2):152-7.

#### **Gonzalez 2009** {published data only}

Gonzalez RP, Cummings GR, Phelan HA, Bosarge PL, Rodning CB. Clinical examination in complement with computed tomography scan: an effective method for identification of cervical spine injury. *Journal of Trauma-Injury Infection & Critical Care* 2009;**67**(6):1297-304.

## Gonzalez 2013 (published data only)

Gonzalez RP, Cummings GR, Baker JA, Frotan AM, Simmons JD, Brevard SB, et al. Prehospital clinical clearance of the cervical spine: a prospective study. *American Surgeon* 2013;**79**(11):1213-7.

## **Griffen 2003** {published data only}

Griffen MM, Frykberg ER, Kerwin AJ, Schinco MA, Tepas JJ, Rowe K, et al. Radiographic clearance of blunt cervical spine injury: plain radiograph or computed tomography scan?. *Journal of Trauma-Injury Infection & Critical Care* 2003;**55**(2):222-6.

## **Griffith 2011** {published data only}

Griffith B, Bolton C, Goyal N, Brown ML, Jain R. Screening cervical spine CT in a level I trauma center: overutilization?. *AJR: American Journal of Roentgenology* 2011;**197**(2):463-7.

## Griffith 2013 (published data only)

Griffith B, Kelly M, Vallee P, Slezak M, Nagarwala J, Krupp S, et al. Screening cervical spine CT in the emergency department, Phase 2: A prospective assessment of use. *AJNR: American Journal of Neuroradiology* 2013;**34**(4):899-903.

# **Griffith 2014** {published data only}

Griffith B, Vallee P, Krupp S, Jung M, Slezak M, Nagarwala J, et al. Screening cervical spine CT in the emergency department, phase 3: increasing effectiveness of imaging. *Journal of the American College of Radiology* 2014;**11**(2):139-44.

## Hanson 2000 {published data only}

Hanson JA, Blackmore CC, Mann FA, Wilson AJ. Cervical spine injury: Accuracy of helical CT used as a screening technique. *Emergency Radiology* 2000;**7**(1):31-5.

## Hanson 2000a {published data only}

Hanson JA, Blackmore CC, Mann FA, Wilson AJ. Cervical spine injury: a clinical decision rule to identify high-risk patients for helical CT screening. *AJR* 2000;**American J**(3):713-7.

## Hasler 2011 {published data only}

Hasler RM, Exadaktylos AK, Bouamra O, Benneker LM, Clancy M, Sieber R, et al. Epidemiology and predictors of spinal injury in adult major trauma patients: European cohort study. *European Spine Journal* 2011;**20**(12):2174-80.

# **Heffernan 2005** {published data only}

Heffernan DS, Schermer CR, Lu SW. What defines a distracting injury in cervical spine assessment?. *Journal of Trauma-Injury Infection & Critical Care* 2005;**59**(6):1396-9.



#### Hoffman 1992 (published data only)

Hoffman JR, Schriger DL, Mower W, Luo JS, Zucker M. Low-risk criteria for cervical-spine radiography in blunt trauma: a prospective study. *Annals of Emergency Medicine* 1992;**21**(12):1454-60.

#### Hollingshead 2000 (published data only)

Hollingshead MC, Keenan HT, Chung CJ, Ziglar MK. Use of screening neck/cervical spine CT in pediatric head trauma victims. *Radiology* 2000;**217**:340.

#### **Hutchings 2009** {published data only}

Hutchings L, Atijosan O, Burgess C, Willett K. Developing a spinal clearance protocol for unconscious pediatric trauma patients. *Journal of Trauma-Injury Infection & Critical Care* 2009:**67**(4):681-6.

#### **Keenan 2001** {published data only}

Keenan HT, Hollingshead MC, Chung CJ, Ziglar MK. Using CT of the cervical spine for early evaluation of pediatric patients with head trauma. *AJR: American Journal of Roentgenology* 2001;**177**(6):1405-9.

## **Kerr 2005** {published data only}

Kerr D, Bradshaw L, Kelly AM. Implementation of the Canadian C-spine rule reduces cervical spine x-ray rate for alert patients with potential neck injury. *Journal of Emergency Medicine* 2005;**28**(2):127-31.

## Kokabi 2011 (published data only)

Kokabi N, Raper DM, Xing M, Giuffre BM. Application of imaging guidelines in patients with suspected cervical spine trauma: retrospective analysis and literature review. *Emergency Radiology* 2011;**18**(1):31-8.

## **Lee 2003** {published data only}

Lee SL, Sena M, Greenholz SK, Fledderman M. A multidisciplinary approach to the development of a cervical spine clearance protocol: process, rationale, and initial results. *Journal of Pediatric Surgery* 2003;**38**(3):358-62.

## Malomo 1995 {published data only}

Malomo AO, Shokunbi MT, Adeloye A. Evaluation of the use of plain cervical spine radiography in patients with head injury. *East African Medical Journal* 1995;**72**(3):186-8.

# Mannix 2011 {published data only}

Mannix R, Nigrovic LE, Schutzman SA, Hennelly K, Bourgeois FT, Meehan WP, et al. Factors associated with the use of cervical spine computed tomography imaging in pediatric trauma patients. *Academic Emergency Medicine* 2011;**18**(9):905-11.

## Markuske 1983 (published data only)

Markuske H. [Roentgenologic diagnosis of the cervical spine in childhood]. [German]. *Psychiatrie, Neurologie und Medizinische Psychologie* 1983;**35**(5):257-66.

## Markuske 1988 (published data only)

Markuske H. [The cervical spine of children in the x-ray and clinical picture]. [German]. *Radiologia Diagnostica* 1988;**29**(1):137-41.

#### Martin 2004 (published data only)

Martin BW, Dykes E, Lecky FE. Patterns and risks in spinal trauma. Archives of Disease in Childhood 2004;89(9):860-5.

## Meek 2007 {published data only}

Meek R, McGannon D, Edwards L. The safety of nurse clearance of the cervical spine using the National Emergency X-radiography Utilization Study low-risk criteria. *Emergency Medicine Australasia* 2007;**19**(4):372-6.

#### Meldon 1998 (published data only)

Meldon SW, Brant TA, Cydulka RK, Collins TE, Shade BR. Out-of-hospital cervical spine clearance: agreement between emergency medical technicians and emergency physicians. *Journal of Trauma-Injury Infection & Critical Care* 1998;**45**(6):1058-61.

#### Morrison 2012 (published data only)

Morrison J, Jeanmonod R. Utilization of NEXUS C-spine criteria among clinicians in a community emergency department setting. Annals of Emergency Medicine. Conference: American College of Emergency Physicians, ACEP Research Forum 2012 Denver, CO United States. Conference Start: 2012 October 8-9 2012; Suppl: S115.

## Mower 2001 (published data only)

Mower WR, Hoffman JR, Pollack CV Jr, Zucker MI, Browne BJ, Wolfson AB, et al. Use of plain radiography to screen for cervical spine injuries. *Annals of Emergency Medicine* 2001;**38**(1):1-7.

#### Neifeld 1988 {published data only}

Neifeld GL, Keene JG, Hevesy G, Leikin J, Proust A, Thisted RA. Cervical injury in head trauma. *Journal of Emergency Medicine* 1988;**6**(3):203-7.

## Nguyen 2005 {published data only}

Nguyen GK, Clark R. Adequacy of plain radiography in the diagnosis of cervical spine injuries. *Emergency Radiology* 2005;**11**(3):158-61.

## Omran 2001 {published data only}

Omran HA, Dowd MD, Knapp JF. Removing the pediatric cervical collar: current practice patterns. *Archives of Pediatrics & Adolescent Medicine* 2001;**155**(2):162-6.

## Pieretti-Vanmarcke 2011 {published data only}

Pieretti-Vanmarcke R, Velmahos GC, Nance ML, Islam S, Falcone RA Jr, Wales PW, et al. Clinical clearance of the cervical spine in blunt trauma patients younger than 3 years: a multicenter study of the American Association for the Surgery of Trauma. *Journal of Trauma-Injury Infection & Critical Care* 2011;**67**(3):543-9.

## Platzer 2006a {published data only}

Platzer P, Jaindl M, Thalhammer G, Dittrich S, Wieland T, Vecsei V, et al. Clearing the cervical spine in critically injured patients: a comprehensive C-spine protocol to avoid unnecessary delays in diagnosis. *European Spine Journal* 2006;**15**(12):1801-10.



#### Platzer 2006b (published data only)

Platzer P, Thalhammer G, Jaindl M, Dittrich S, Vecsei V, Gaebler C. Clearing the cervical spine in polytrauma patients: current standards in diagnostic algorithm. *European Journal of Trauma* 2006;**32**(6):570-5.

#### Pulfrey 2002 (published data only)

Pulfrey S, Evans MF. Blunt trauma: Which patients require diagnostic imaging of the cervical spine?. *Canadian Family Physician* 2002;**48**:1061-2.

#### Quigley 2014 (published data only)

Quigley A, Stafrace S. Paediatric trauma: Children are not just small adults-Emergency imaging by the general radiologist. Pediatric Radiology Conference: 37th Postgraduate Course and 51st Annual Meeting of the European Society of Paediatric Radiology Amsterdam Netherlands 2014 June 02-06. 2014; Vol. 44:S326.

#### Raza 2013 (published data only)

Raza M, Elkhodair S, Zaheer A, Yousaf S. Safe cervical spine clearance in adult obtunded blunt trauma patients on the basis of a normal multidetector CT scan: a meta-analysis and cohort study (Provisional abstract). *Injury* 2013;**44**:1589-95.

## Ropele 2009 (published data only)

Ropele D, Blech K, Vander Laan KJ. Cervical spine clearance in the nonalert, noncommunicative, or unreliable pediatric blunt trauma patient. *Journal of Trauma Nursing* 2009;**16**(3):148-59.

#### Rose 2012 (published data only)

Rose MK, Rosal LM, Gonzalez RP, Rostas JW, Baker JA, Simmons JD, et al. Clinical clearance of the cervical spine in patients with distracting injuries: it is time to dispel the myth?. *Journal of Trauma and Acute Care Surgery* 2012;**73**(2):498-502.

#### Ross 1987 (published data only)

Ross SE, Schwab CW, David ET, DeLong WG, Born CT. Clearing the cervical spine: initial radiologic evaluation. *Journal of Trauma-Injury Infection & Critical Care* 1987;**27**(9):1055-60.

## Saddison 1991 {published data only}

Saddison D, Vanek VW, Racanelli JL. Clinical indications for cervical spine radiographs in alert trauma patients. *American Surgeon* 1991;**57**(6):366-9.

# Sanchez 2005 {published data only}

Sanchez B, Waxman K, Jones T, Conner S, Chung R, Becerra S. Cervical spine clearance in blunt trauma: evaluation of a computed tomography-based protocol. *Journal of Trauma-Injury Infection & Critical Care* 2005;**59**(1):179-83.

## Scarrow 1999 (published data only)

Scarrow AM, Levy EI, Resnick DK, Adelson PD, Sclabassi RJ. Cervical spine evaluation in obtunded or comatose pediatric trauma patients: A pilot study. *Pediatric Neurosurgery* 1999;**30**(4):169-75.

#### Schleehauf 1989 (published data only)

Schleehauf K, Ross SE, Civil ID, Schwab CW. Computed tomography in the initial evaluation of the cervical spine. *Annals of Emergency Medicine* 1989;**18**(8):815-7.

## Sheikh 2012 {published data only}

Sheikh K, Belfi LM, Sharma R, Baad M, Sanelli PC. Evaluation of acute cervical spine imaging based on ACR Appropriateness Criteria. *Emergency Radiology* 2012;**19**(1):11-7.

#### Smart 2003 (published data only)

Smart PJ, Hardy PJ, Buckley DM, Somers JM, Broderick NJ, Halliday KE, et al. Cervical spine injuries to children under 11: should we use radiography more selectively in their initial assessment?. *Emergency Medicine Journal* 2003;**20**(3):225-7.

#### Stiell 2003 (published data only)

Stiell IG, Clement CM, McKnight RD, Brison R, SchullM J, Rowe BH, et al. The Canadian C-spine rule versus the NEXUS low-risk criteria in patients with trauma. *New England Journal of Medicine* 2003;**349**(26):2510-8.

#### **Stiell 2009** {published data only}

Stiell IG, Clement CM, Grimshaw J, Brison RJ, Rowe BH, Schull MJ, et al. Implementation of the Canadian C-Spine Rule: prospective 12-centre cluster randomised trial. *BMJ (Clinical research ed.)* 2009;**339**:b4146.

## **Stiell 2010** {published data only}

Stiell IG, Clement CM, O'Connor A, Davies B, Leclair C, Sheehan P, et al. Multicentre prospective validation of use of the Canadian C-Spine Rule by triage nurses in the emergency department. *CMAJ: Canadian Medical Association Journal* 2010;**182**(11):1173-9.

## **Stroh 2001** {published data only}

Stroh G, Braude D. Can an out-of-hospital cervical spine clearance protocol identify all patients with injuries? An argument for selective immobilization. *Annals of Emergency Medicine* 2001;**37**(6):609-15.

## Sun 2013 (published data only)

Sun R, Skeete D, Wetjen K, Lilienthal M, Liao J, Madsen M, et al. A pediatric cervical spine clearance protocol to reduce radiation exposure in children. *Journal of Surgical Research* 2013;**183**(1):341-6.

## Tahvonen 2013 {published data only}

Tahvonen P, Oikarinen H, Pääkkö E, Karttunen A, Blanco Sequeiros R, Tervonen O. Justification of CT examinations in young adults and children can be improved by education, guideline implementation and increased MRI capacity. *British Journal of Radiology* 2013;**86**(1029):20130337.

## Valusek 2010 (published data only)

Valusek PA, Langworthy J, Laituri CA, Garey CL, Ostlie DJ, St Peter SD. Detection of cervical spine injury in pediatric trauma patients. Journal of Surgical Research.Conference: 5th Annual Academic Surgical Congress of the Association for Academic Surgery, AAS and the Society of University Surgeons,



SUS San Antonio, TX United States.2010 February 03-05. 2010:420.

## **Velmahos 1996** {published data only}

Velmahos GC, Theodorou D, Tatevossian R, Belzberg H, Cornwell EE 3rd, Berne TV, et al. Radiographic cervical spine evaluation in the alert asymptomatic blunt trauma victim: much ado about nothing. *Journal of Trauma-Injury Infection & Critical Care* 1996;**40**(5):768-74.

## **Additional references**

#### APLS

Advanced Life Support Group. Advanced Paediatric Life Support: The Practical Approach. 5th Edition. Chichester: John Wiley & Sons, 2011.

#### **ATLS 2013**

ATLS Subcommittee, American College of Surgeons' Committee on Trauma, International ATLS working group. Advanced trauma life support (ATLS®): the ninth edition. *Journal of Trauma and Acute Care Surgery* 2013;**74**(5):1363-6.

#### Bagley 2006

Bagley LJ. Imaging of spinal trauma. *Radiologic Clinics of North America* 2006;**44**(1):1-12, vii.

#### Benzel 1996

Benzel EC, Hart BL, Ball PA, Baldwin NG, Orrison WW, Espinosa MC. Magnetic resonance imaging for the evaluation of patients with occult cervical spine injury. *Journal of Neurosurgery* 1996;**85**(5):824-9.

#### Booth 2012

Booth TN. Cervical spine evaluation in pediatric trauma. American Journal of Roentgenology 2012;**198**(5):W417-25.

#### Bracken 1990

Bracken MB, Shepard MJ, Collins WF, Holford TR, Young W, Baskin DS, et al. A randomized, controlled trial of methylprednisolone or naloxone in the treatment of acute spinal-cord injury. Results of the Second National Acute Spinal Cord Injury Study. *New England Journal of Medicine* 1990;**322**(20):1405-11.

#### **Buhs 2000**

Buhs C, Cullen M, Klein M, Farmer D. The pediatric trauma C-spine: is the 'odontoid' view necessary?. *Journal of Pediatric Surgery* 2000;**35**(6):994-7.

## Cattell 1965

Cattell HS, Filtzer DL. Pseudosubluxation and other normal variations in the cervical spine in children. A study of one hundred and sixty children. *The Journal of Bone and Joint Surgery. American Volume* 1965;**47**(7):1295-309.

## Chung 2011

Chung S, Mikrogianakis A, Wales PW, Dirks P, Shroff M, Singhal A, et al. Trauma association of Canada Pediatric Subcommittee

National Pediatric Cervical Spine Evaluation Pathway: consensus guidelines. *Journal of Trauma* 2011;**70**(4):873-84.

#### Cirak 2004

Cirak B, Ziegfeld S, Knight VM, Chang D, Avellino AM, Paidas CN. Spinal injuries in children. *Journal of Pediatric Surgery* 2004;**39**(4):607-12.

#### De Vet 2008

De Vet HC, Eisinga A, Riphagen II, Aertgeerts B, Pewsner D, editor(s). Chapter 7: Searching for studies. In: Deeks JJ, Bossuyt PM, Gatsonis C, editor(s). Cochrane Handbook for Systematic Reviews of Diagnostic Test Accuracy Version 0.4 (updated September 2008). The Cochrane Collaboration, 2008. Available from srdta.cochrane.org.

#### Dickman 1989

Dickman CA, Rekate HL, Sonntag VK, Zabramski JM. Pediatric spinal trauma: vertebral column and spinal cord injuries in children. *Pediatric Neuroscience* 1989;**15**(5):237-55; discussion 56

#### Dormans 1995

Dormans JP, Criscitiello AA, Drummond DS, Davidson RS. Complications in children managed with immobilization in a halo vest. *Journal of Bone and Joint surgery. American volume* 1995;**77**(9):1370-3.

## **Duffy 2014**

Duffy S, Misso K, Noake C, Ross J, Stirk L. Supplementary searches of PubMed to improve currency of MEDLINE and MEDLINE In-Process searches via OvidSP. medicine.exeter.ac.uk/media/universityofexeter/medicalschool/research/pentag/documents/Steven\_Duffy\_ISSG\_Exeter\_2014\_poster\_1.pdf. Poster presented at the UK InterTASC Information Specialists' Sub-Group (ISSG) Workshop; 9 July 2014; Exeter: UK (accessed 6 August 2014.

## Fesmire 1989

Fesmire FM, Luten RC. The pediatric cervical spine: developmental anatomy and clinical aspects. *Journal of Emergency Medicine* 1989;**7**(2):133-42.

## Givens 1996

Givens TG, Polley KA, Smith GF, Hardin WD. Pediatric cervical spine injury: a three-year experience. *Journal of Trauma* 1996;**41**(2):310-4.

#### Hadley 1988

Hadley MN, Zabramski JM, Browner CM, Rekate H, Sonntag VK. Pediatric spinal trauma. Review of 122 cases of spinal cord and vertebral column injuries. *Journal of Neurosurgery* 1988;**68**(1):18-24.

## Hamilton 1992

Hamilton MG, Myles ST. Pediatric spinal injury: review of 174 hospital admissions. *Journal of Neurosurgery* 1992;**77**(5):700-4.

#### Hasler 2012

Hasler RM, Exadaktylos AK, Bouamra O, Benneker LM, Clancy M, Sieber R, et al. Epidemiology and predictors of cervical



spine injury in adult major trauma patients: a multicenter cohort study. *Journal of Trauma and Acute Care Surgery* 2012;**72**(4):975-81.

#### Henry 2013

Henry M, Riesenburger RI, Kryzanski J, Jea A, Hwang SW. A retrospective comparison of CT and MRI in detecting pediatric cervical spine injury. *Child's Nervous System* 2013;**29**(8):1333-8.

#### Hernandez 2004

Hernandez JA, Chupik C, Swischuk LE. Cervical spine trauma in children under 5 years: productivity of CT. *Emergency Radiology* 2004;**10**(4):176-8.

#### Hill 1984

Hill SA, Miller CA, Kosnik EJ, Hunt WE. Pediatric neck injuries. A clinical study. *Journal of Neurosurgery* 1984;**60**(4):700-6.

#### Hoffman 2000

Hoffman JR, Mower WR, Wolfson AB, Todd KH, Zucker MI. Validity of a set of clinical criteria to rule out injury to the cervical spine in patients with blunt trauma. National Emergency X-Radiography Utilization Study Group.. *New England Journal of Medicine* 2000;**343**(2):94-9.

#### Jimenez 2008

Jimenez RR, Deguzman MA, Shiran S, Karrellas A, Lorenzo RL. CT versus plain radiographs for evaluation of c-spine injury in young children: do benefits outweigh risks?. *Pediatric Radiology* 2008;**38**(6):635-44.

## Keiper 1998

Keiper MD, Zimmerman RA, Bilaniuk LT. MRI in the assessment of the supportive soft tissues of the cervical spine in acute trauma in children. *Neuroradiology* 1998;**40**(6):359-63.

#### Kokoska 2001

Kokoska ER, Keller MS, Rallo MC, Weber TR. Characteristics of pediatric cervical spine injuries. *Journal of Pediatric Surgery* 2001;**36**(1):100-5.

#### Kriss 1997

Kriss VM, Kriss TC. Imaging of the cervical spine in infants. *Pediatric Emergency Care* 1997;**13**(1):44-9.

#### Leonard 2011

Leonard JC, Kuppermann N, Olsen C, Babcock-Cimpello L, Brown K, Mahajan P, et al. Factors associated with cervical spine injury in children after blunt trauma. *Annals of Emergency Medicine* 2011;**58**(2):145-55.

#### Leonard 2013

Leonard JC. Cervical spine injury. *Pediatric Clinics of North America* 2013;**60**(5):1123-37.

## Leonard 2014

Leonard JR, Jaffe DM, Kuppermann N, Olsen CS, Leonard JC. Cervical spine injury patterns in children. *Pediatrics* 2014;**133**(5):e1179-88.

#### Mortazavi 2011

Mortazavi M, Gore PA, Chang S, Tubbs RS, Theodore N. Pediatric cervical spine injuries: a comprehensive review. *Child's Nervous System* 2011;**27**(5):705-17.

#### **NICE 2014**

National Institute for Health and Care Excellence. Triage, assessment, investigation and early management of head injury in children, young people and adults. www.nice.org.uk/Guidance/CG176/Evidence 2014 (accessed 5th November 2017).

#### Nigrovic 2012

Nigrovic LE, Rogers AJ, Adelgais KM, Olsen CS, Leonard JR, Jaffe DM, et al. Utility of plain radiographs in detecting traumatic injuries of the cervical spine in children. *Pediatric Emergency Care* 2012;**28**(5):426-32.

#### **NSCISC 2012**

National Spinal Cord Injury Statistical Center. 2012 Annual Report National Spinal Cord Injury Statistical Center. www.nscisc.uab.edu/PublicDocuments/reports/ pdf/2012%20NSCISC%20Annual%20Statistical%20Report %20Complete%20Public%20Version.pdf November 2012 (accessed 5th November 2017).

#### Orenstein 1994

Orenstein JB, Klein B L, Gotschall CS, Ochsenschlager DW, Klatzko MD, Eichelberger MR. Age and outcome in pediatric cervical spine injury: 11-year experience. *Pediatric Emergency Care* 1994;**10**(3):132-7.

## Osenbach 1992

Osenbach RK, Menezes AH. Pediatric spinal cord and vertebral column injury. *Neurosurgery* 1992;**30**(3):385-90.

## Pang 1982

Pang D, Wilberger JE. Spinal cord injury without radiographic abnormalities in children. *Journal of Neurosurgery* 1982;**57**(1):114-29.

## Pang 2004

Pang D. Spinal cord injury without radiographic abnormality in children, 2 decades later. *Neurosurgery* 2004;**55**(6):1325-42; discussion 1342-3.

## Parent 2011

Parent S, Mac-Thiong JM, Roy-Beaudry M, Sosa JF, Labelle H. Spinal cord injury in the pediatric population: a systematic review of the literature. *Journal of Neurotrauma* 2011;**28**(8):1515-24.

## Partrick 2000

Partrick DA, Bensard DD, Moore EE, Calkins CM, Karrer FM. Cervical spine trauma in the injured child: a tragic injury with potential for salvageable functional outcome. *Journal of Pediatric Surgery* 2000;**35**(11):1571-5.

## **Patel 2001**

Patel JC, Tepas JJ, Mollitt DL, Pieper P. Pediatric cervical spine injuries: defining the disease. *Journal of Pediatric Surgery* 2001;**36**(2):373-6.



#### Pearce 2012

Pearce MS, Salotti JA, Little MP, McHugh K, Lee C, Kim KP, et al. Radiation exposure from CT scans in childhood and subsequent risk of leukaemia and brain tumours: a retrospective cohort study. *Lancet* 2012;**380**(9840):499-505.

#### Platzer 2007

Platzer P, Jaindl M, Thalhammer G, Dittrich S, Kutscha-Lissberg Fl, Vecsei V, et al. Cervical spine injuries in pediatric patients. *Journal of Trauma* 2007;**62**(2):389-96.

#### Ravichandran 1982

Ravichandran G, Silver JR. Missed injuries of the spinal cord. *British Medical Journal (Clinical Research Ed.)* 1982;**284**(6320):953-6.

## Reitsma 2005

Reitsma JB, Glas AS, Rutjes AW, Scholten RJ, Bossuyt PM, Zwinderman AH. Bivariate analysis of sensitivity and specificity produces informative summary measures in diagnostic reviews. *Journal of Clinical Epidemiology* 2005;**58**(10):982-90.

## Review Manager 2014 [Computer program]

Nordic Cochrane Centre, The Cochrane Collaboration. Review Manager 5 (RevMan 5). Version 5.3. Copenhagen: Nordic Cochrane Centre, The Cochrane Collaboration, 2014.

#### Ron 2002

Ron E. Let's not relive the past: a review of cancer risk after diagnostic or therapeutic irradiation. *Pediatric Radiology* 2002;**32**(10):739-44; discussion 751-4.

## Rozzelle 2013

Rozzelle CJ, Aarabi B, Dhall SS, Gelb DE, Hurlbert RJ, Ryken TC, et al. Management of pediatric cervical spine and spinal cord injuries. *Neurosurgery* 2013;**72 Suppl** 2:205-26.

## Schenarts 2001

Schenarts PJ, Diaz J, Kaiser C, Carrillo Y, Eddy V, Morris JA. Prospective comparison of admission computed tomographic scan and plain films of the upper cervical spine in trauma patients with altered mental status. *Journal of Trauma* 2001;**51**(4):663-8; discussion 668-9.

# Schuster 2005

Schuster R, Waxman K, Sanchez B, Becerra S, Chung R, Conner S, et al. Magnetic resonance imaging is not needed to clear cervical spines in blunt trauma patients with normal computed tomographic results and no motor deficits. *Archives of Surgery* 2005;**140**(8):762-6.

# Seal 2005

Seal C, Millbrandt T, Gelb D, Ludwig S. The management of pediatric cervical spine injuries. *Seminars in Spine Surgery* 2005;**17**(2):95-9.

#### Stiell 2001

Stiell IG, Wells GA, Vandemheen KL, Clement CM, Lesiuk H, De Maio VJ, et al. The Canadian C-spine rule for radiography in alert and stable trauma patients. *JAMA* 2001;**286**(15):1841-8.

#### Sun 2000

Sun PP, Poffenbarger GJ, Durham S, Zimmerman RA. Spectrum of occipitoatlantoaxial injury in young children. *Journal of Neurosurgery* 2000;**93**(1 Suppl):28-39.

#### Swischuk 1977

Swischuk LE. Anterior displacement of C2 in children: physiologic or pathologic. *Radiology* 1977;**122**(3):759-63.

#### Swischuk 1984

Swischuk LE. The cervical spine in childhood. *Current Problems in Diagnostic Radiology* 1984;**13**(5):1-26.

## Trigylidas 2010

Trigylidas T, Yuh SJ, Vassilyadi M, Matzinger MA, Mikrogianakis A. Spinal cord injuries without radiographic abnormality at two pediatric trauma centers in Ontario. *Pediatric Neurosurgery* 2010;**46**(4):283-9.

#### Vogel 2002a

Vogel LC, Krajci KA, Anderson CJ. Adults with pediatric-onset spinal cord injury: part 1: prevalence of medical complications. *The Journal of Spinal Cord Medicine* 2002;**25**(2):106-16.

#### Vogel 2002b

Vogel LC, Krajci KA, Anderson CJ. Adults with pediatriconset spinal cord injury: part 2: musculoskeletal and neurological complications. *The Journal of Spinal Cord Medicine* 2002;**25**(2):117-23.

# Vogel 2002c

Vogel LC, Krajci KA, Anderson CJ. Adults with pediatric-onset spinal cord injuries: part 3: impact of medical complications. *Journal of Spinal Cord Medicine* 2002;**25**(4):297-305.

## Whiting 2003

Whiting P, Rutjes AW, Reitsma JB, Bossuyt PM, Kleijnen J. The development of QUADAS: a tool for the quality assessment of studies of diagnostic accuracy included in systematic reviews. BMC Medical Research Methodology 2003;**3**:25.

## Whiting 2011

Whiting PF, Rutjes AS, Westwood ME, Mallett S, Deeks JJ, Reitsma JB, et al. Quadas-2: A revised tool for the quality assessment of diagnostic accuracy studies. *Annals of Internal Medicine* 2011;**155**(8):529-36.

## Yucesoy 2008

Yucesoy K, Yuksel KZ. SCIWORA in MRI era. *Clinical Neurology* and *Neurosurgery* 2008;**110**(5):429-33.

<sup>\*</sup> Indicates the major publication for the study



# CHARACTERISTICS OF STUDIES

# **Characteristics of included studies** [ordered by study ID]

# Ehrlich 2009

Study characteristics				
Patient sampling	Retrospective cohort			
Patient characteristics and setting	Study location: University of Michigan CS Mott Children's Hospital			
	Study period: 2005 - 200	7		
	Patients enrolled: 150 pa ography	atients younger than 1	LO years of age with plain radi	
	Patients included in ana	lysis: 108 NEXUS crite	ria, 109 Canadian C-spine Rul	
	Mean age: 4.3 years (SD	3.1)		
	Sex: 72 boys, 53 girls			
	Patients with CSI: 7			
Index tests	NEXUS criteria and Cana	NEXUS criteria and Canadian C-spine Rule		
Target condition and reference standard(s)	Target condition: clinica	lly-important missed	CSI	
	Reference standard: Pla	in radiography, CT or	both	
Flow and timing	Time between reference	Time between reference standard and index test: 0 days		
Comparative				
Notes	No definition given of clinically-important CSI			
	Referral for radiography	depended on the follo	owing items:	
	- awake or not			
	- presence of motor/sen	sory deficits		
	- presence of neck pain			
	- presence of distracting injuries			
	- presence of intoxicating agents			
	275 patients were included, 150 with the same reference stand diography), and 125 without radiographic imaging. Only 109 po included in their analysis			
Methodological quality				
Item	Authors' judgement	Risk of bias	Applicability con- cerns	
DOMAIN 1: Patient Selection				
Was a consecutive or random sample of patients enrolled?	Yes			



Ehrlich 2009 (Continued)				
Was a case-control design avoided?	Yes			
Did the study avoid inappropriate exclusions?	Yes			
		Low	Low	
DOMAIN 2: Index Test All tests				
Were the index test results interpreted without knowledge of the results of the reference standard?	Unclear			
If a threshold was used, was it pre-specified?	Yes			
		Unclear	Low	
DOMAIN 3: Reference Standard				
Is the reference standards likely to correctly classify the target condition?	Yes			
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear			
		Unclear	Low	
DOMAIN 4: Flow and Timing				
Was there an appropriate interval between index test and reference standard?	Yes			
Did all patients receive the same reference standard?	No			
Were all patients included in the analysis?	No			
		High		

# **Jaffe 1987**

dile 1301	
Study characteristics	
Patient sampling	Retrospective cohort
Patient characteristics and setting	Study location: Children's Memorial Hospital, Chicago.
	Study period: 139 consecutive trauma patients, younger than 16 years old from 1983 - 1984 (1 with CSI) and all children younger than 16 years old, with CSI from 1974 to 1984
	Patients enrolled: 232 patients younger than 16 years of age
	Patients included in analysis: 206
	Age: 0 - 3 years 28%, 4 - 12 years 42%, 13 - 16 years 30%
	Sex: 143 boys, 63 girls



Jaffe 1987 (Continued)	Patients with CSI: 59 (28.6	5%)	
Index tests	Derivative of the NEXUS of	riteria with the followi	ng items:
	-neck pain/tenderness		
	-abnormalities of reflexes	s, strength or sensation	l
	-limitation of neck mobili	ty	
	-abnormal mental status		
Target condition and reference standard(s)	Target condition: predict	ing the presence of CSI	in children
	Reference standard: plair	n radiography	
Flow and timing	Time between reference	standard and index tes	t: 0 days
Comparative			
Notes	The items of the index tes US criteria. The item miss		nme as the items in the NEX- distracting injury
Methodological quality			
Item	Authors' judgement	Risk of bias	Applicability con- cerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	No		
Was a case-control design avoided?	No		
Did the study avoid inappropriate exclusions?	Yes		
		High	High
DOMAIN 2: Index Test All tests			
Were the index test results interpreted without knowledge of the results of the reference standard?	Unclear		
If a threshold was used, was it pre-specified?	No		
		Unclear	High
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear		
		Unclear	Low



## Jaffe 1987 (Continued)

DOMAIN 4: F	low and	l Timing
-------------	---------	----------

Was there an appropriate interval between index test and reference standard?	Yes
Did all patients receive the same reference standard?	Yes
Were all patients included in the analysis?	Yes
	Unclear

# Viccellio 2001

Study characteristics				
Patient sampling	Prospective multicenter cohort study			
Patient characteristics and setting	Study location: UCLA Emergency Medici pating centers. Study sites comprise a racluding academic trauma centers, community EDs	ange of acute care facilities, in-		
	Study period: 1990 - 2000 (estimated)			
	Patients enrolled: 34,069 patients, adult	s and children		
	Patients included in analysis: 3065			
	Age: 0 - 2 years 2.9%, 2 - 8 years 26.7%, 9	- 17 years 70.5%		
	Sex: unknown			
	Patients with CSI: 30 (0.98%)			
Index tests	NEXUS criteria	NEXUS criteria		
Target condition and reference standard(s)	Target condition: evaluating the performance of the NEXUS criter the detection of CSI in children			
	Reference standard: Plain radiography,	CT or both		
Flow and timing	Time between reference standard and index test: 0 days			
Comparative				
Notes Analysis of the pediatric population included in the original 1 2000, the seminal publication of the NEXUS tool				
	Only 4 children with CSI younger than 8	years old		
Methodological quality				
ltem	Authors' judgement Risk of bias	Applicability con- cerns		
DOMAIN 1: Patient Selection				



Viccellio 2001 (Continued)				
Was a consecutive or random sample of patients enrolled?	Yes			
Was a case-control design avoided?	Yes			
Did the study avoid inappropriate exclusions?	Yes			
		Low	Low	
DOMAIN 2: Index Test All tests				
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes			
If a threshold was used, was it pre-specified?	Yes			
		Low	Low	
DOMAIN 3: Reference Standard				
Is the reference standards likely to correctly classify the target condition?	Yes			
Were the reference standard results interpreted without knowledge of the results of the index tests?	Yes			
		Low	Low	
DOMAIN 4: Flow and Timing				
Was there an appropriate interval between index test and reference standard?	Yes			
Did all patients receive the same reference standard?	No			
Were all patients included in the analysis?	Yes			
		High		

# **Characteristics of excluded studies** [ordered by study ID]

Study	Reason for exclusion
Adelgais 2014	Index test not relevant
Anderson 2006	Target condition not relevant
Anderson 2010	Index test not relevant
Baker 1999	Index test not relevant
Bandiera 2003	Different study population (adults)
Banit 2000	Index test not relevant



Study	Reason for exclusion
Bayless 1989	I ndex test not relevant
Blacksin 1995	Index test not relevant
Borock 1991	Different study population (only patients with CSI)
Brockmeyer 2012	Different study population (suspected or proven CSI)
Brooks 2001	Index test not relevant
Brown 2001	Index test not relevant
Browne 2003	T arget condition not relevant
Burns 2011	Different target condition
Clayton 2012	Index test not relevant
Coffey 2011	Different study population (adults)
Como 2011	Index test not relevant
Dahlquist 2013	Index test not relevant
Dickinson 2004	Different study population (adults)
DiGiacomo 2002	Index test not relevant
Edwards 2001	Index test not relevant
Ersoy 1995	Index test not relevant
Fischer 1984	Index test not relevant
Garton 2008	Different study population (only patients with CSI)
Gbaanador 1986	Index test not relevant
Gonzalez 1999	Index test not relevant
Gonzalez 2009	Index test not relevant
Gonzalez 2013	Different study population (pre- hospital evaluation)
Griffen 2003	Index test not relevant
Griffith 2011	different study population (adults)
Griffith 2013	different study population (adults)
Griffith 2014	Different study population (adults)
Hanson 2000	Different study population (adults)
Hanson 2000a	Different study population (adults)



Study	Reason for exclusion								
Hasler 2011	Different study population (adults)								
Heffernan 2005	Different study population (adults)								
Hoffman 1992	Authors did not separate data for the pediatric population (only a small number of children were included)								
Hollingshead 2000	Index test not relevant								
Hutchings 2009	Index test not relevant								
Keenan 2001	Index test not relevant								
Kerr 2005	Different study population (adults)								
Kokabi 2011	Different study population (adults)								
Lee 2003	I ndex test not relevant								
Malomo 1995	I ndex test not relevant								
Mannix 2011	Index test not relevant								
Markuske 1983	Target condition not relevant								
Markuske 1988	Target condition not relevant								
Martin 2004	Index test not relevant								
Meek 2007	Index test not relevant								
Meldon 1998	Different study population (out- of- hospital patients)								
Morrison 2012	Index test not relevant								
Mower 2001	Index test not relevant								
Neifeld 1988	Different study population (adults)								
Nguyen 2005	Index test not relevant								
Omran 2001	Index test not relevant								
Pieretti-Vanmarcke 2011	Index test not relevant								
Platzer 2006a	Index test not relevant								
Platzer 2006b	Index test not relevant								
Pulfrey 2002	Different study population (adults)								
Quigley 2014	Index test not relevant								
Raza 2013	Different study population (adults)								



Reason for exclusion								
Index test not relevant								
Index test not relevant								
Index test not relevant								
Different study population (only patients with CSI)								
Index test not relevant								
Different study population (only patients with CSI)								
Index test not relevant								
Different study population (adults)								
Index test not relevant								
Different study population (adults)								
Different study population (adults)								
Different study population (adults)								
Index test not relevant								
Target condition not relevant								
Index test not relevant								
Index test not relevant								
Index test not relevant								

# DATA

Presented below are all the data for all of the tests entered into the review.


# Table Tests. Data tables by test

Test	No. of studies	No. of participants					
1 NEXUS	3	3379					
2 Canadian C-spine Rule	1	109					



## Test 1. NEXUS.

Review: Triage tools for detecting cervical spine injury in pediatric trauma patients Test: 1 NEXUS



# Test 2. Canadian C-spine Rule.

Review: Triage tools for detecting cervical spine injury in pediatric trauma patients Test: 2 Canadian C-spine Rule

St	udy	TP	FP	FN	TN	Sensitivity	Specificity		5	Sensitivit	ty					Specificit	ty		
	Ehrlich 2009	6	87	1	15	0.86 [ 0.42, 1.00 ]	0.15 [ 0.08, 0.23 ]							-	_				
								0	0.2	0.4	0.6	0.8	1 (	)	0.2	0.4	0.6	0.8	1

## **ADDITIONAL TABLES**

# Table 1. Citations per search engine

Search Engine	Number of citations				
MEDLINE	7985				
MEDLINE In-Process	39				
PubMed (search limited to studies not in MEDLINE)	86				
Embase	9187				
CENTRAL	651				
Science Citation Index	1220				
Proquest Dissertation and Theses	20				
Cochrane Database of Systematic Reviews	250				
DARE	105				
НТА	30				
Medion results	16				
ClinicalTrials.gov	19				
WHO ICTRP	19				
OpenGrey	33				
DTA Trials Register	3				



## **Table 1. Citations per search engine** (Continued)

Results found through other sources	1
-Duplicates	-4934
TOTAL	14,730

Table 2. QUADAS review question and inclusion criteria

Category	Review Question	Inclusion Criteria	
Patients	All children aged 0 to 18 years who underwent trauma evaluation Chi at the emergency room		
Index test	NEXUS criteria, Canadian C-spine Rule	NEXUS criteria, Canadian C-spine Rule	
Target condition	Cervical spine injury defined as any fracture or (sub)luxation of the cervical spine, ligamentous injury, or spinal cord injury	Cervical spine injury	
Reference standard	Radiographic imaging in the form of plain radiography, CT, and/ or MRI, or clinical follow-up (if the index test was scored negative) will be our reference standards	Radiographic imaging or clinical fol- low-up	
Outcome	N/A	Sufficient data to construct a 2x2 table	
Study design	Diagnostic studies with cross-sectional or cohort designs (retrospective or prospective) and randomized controlled trials, most favorable direct comparison studies	Diagnostic studies with cross-sectional or cohort designs (retrospective or prospective) and randomized controlled trials	

N/A: not applicable.

Table 3. Assessment of methodological quality: QUADAS-2 and additional items

	Risk of bias		Applicability		
Quality item	Quality indica- tor	Notes	Quality indica- tor	Notes	
Domain 1 Patient selection	Could the selection of participants have introduced bias? (high/low/unclear)		Are there concerns that the included patients and settings do not match the review question? (high/low/unclear)		
	1. Was a consecutive or random sample of patients enrolled?	Yes: if a consecutive or random sample of patients was enrolled  No: if a selected group of patients was enrolled, or if a case control study was performed  Unclear: if there is insufficient information on enrolled patients	1. Were the patients included in the study children presented at the emergency department for evaluation following blunt trauma?	Yes: The included population were children (0 - 18 years old) presenting at the emergency department following blunt trauma.  No: (a) Only adult patients were included, OR (b) if patients were included without blunt trauma OR (c) if the patients included were	



## Table 3. Assessment of methodological quality: QUADAS-2 and additional items (Continued)

not presenting at the emergency department.

**Unclear:** (a) The study population contained adults and children, OR (b) insufficient information was given on the setting, selection criteria, or selection procedure to make a judgment

2. Did the study avoid inappropriate exclusions?

**Yes:** if there were no inappropriate exclusions

**No:** if there were inappropriate exclusions

**Unclear:** if there is insufficient information on exclusions

2. Were the participants planned for evaluation of the cervical spine?

**Yes:** if the participants were planned for evaluation of the cervical spine

**No:** if participants were not planned for evaluation of the cervical spine

**Unclear:** if there is insufficient information

# Domain 2 Index test

## Could the interpretation of the index test have introduced bias? (high/low/unclear)

Are there concerns that the index test, its conduct, or the interpretation differ from the review question? (high/low/unclear)

1. Were the index test results interpreted without knowledge of the results of the reference standard? This will always be rated as **yes**, because the index test is performed before the reference standard.

1. Were the same clinical data available when test results were interpreted as would be available when the test is used in practice?

**Yes:** if clinical data would normally be available when the test is interpreted in practice and similar data were available when interpreting the index test in the study

**No:** if clinical data were not available when index test(s) was/were interpreted

**Unclear:** if insufficient information was given to explain which clinical data were available at the time of assessment

2. Was the threshold used prespecified?

**Yes:** if a clear description of the threshold is given before start of the study

**No:** if no clear description is given beforehand

**Unclear:** if there is insufficient information to determine whether or not a prespecified threshold was used

2. Did the study provide a clear definition of what was considered to be a "positive" result for the index test? **Yes:** if the study provides a clear definition of a positive test result

**No:** if no definition of a positive test result is given

**Unclear:** if insufficient information was given to permit judgment

3. Could the conduct or interpretation of the index test have introduced bias?

**Yes:** If the items of the triage tool were not the same as the items in the index

**No:** if the items of the index test were the same in all patients

\_



Table 3.	Assessment of methodolo	gical qualit	y: QUADAS-2 a	and additional items	(Continued)
----------	-------------------------	--------------	---------------	----------------------	-------------

**Unclear:** if insufficient information is provided

Domain 3

## Reference standard

Could the interpretation of the reference standard have introduced bias? (high/low/unclear)

Are there concerns that the target condition as defined by the preference standard does not match the review question? (high/low/unclear)

1. Is the reference standard likely to correctly classify the target condition?

**Yes**: if the reference standard was plain radiography, CT, MRI or clinical follow-up

No: if the reference standard was not plain radiography, CT, MRI, or clinical follow-up

**Unclear**: if the reference standard was not clearly reported

1. Did the study provide a clear definition of what was considered to be a "positive" result for the reference

standard?

Yes: if a clear description is given about when the reference standard is positive or negative (CSI yes/no)

No: if there is no clear description of the results of the reference standard

Unclear: if the definition of the reference standard is not clearly reported

2. Were the reference standard results interpreted without knowledge of the results of the index test?

Yes: if the radiologist was blinded for the results of the index tests

No: if the radiologist was aware of the results of the index test

**Unclear:** if insufficient information was given on independent or blind assessment of the reference test

### Domain 4

## Flow & Timing

## Could the patient flow have introduced bias? (low/high/unclear)

1. Is the time period between reference standard and index test short enough to be reasonably sure that the target condition did not change between the two tests?

**Yes:** if the reference standard was performed directly after trauma presentation up to 72 hours after trauma presen-

No: if the average interval between reference standard and index test was longer than 72 hours

**Unclear:** if the interval between the trauma and reference standard was not clearly reported

2. Did all patients receive the same reference standard?

Yes: if all patients underwent the same reference standard

No: if not all patients underwent reference standard; also those with a negative index test did not undergo reference

Unclear: if insufficient information is provided

3. Were all patients included in the analysis? **Yes:** if all patients entered in the study are included in the analysis

No: if not all patients in the study are included in the analysis



## Table 3. Assessment of methodological quality: QUADAS-2 and additional items (Continued)

**Unclear:** if it is not clear whether all patients were accounted for

4. Were withdrawals from the study reported? **Yes**: if the number of and reasons for all withdrawals from the study were explained (ideally by a flow chart) or if no participants were excluded from the analysis

**No:** if the reason for withdrawal from the study was not explained

**Unclear**: if insufficient information was given on the withdrawals

#### **APPENDICES**

## **Appendix 1. Search strategies**

## **MEDLINE**

Search 24 Feb 2015

- 1 (NEXUS or CCR).mp. (5678)
- 2 National Emergency X-Radiography.mp. (48)
- 3 (Canadian c-spine or Canadian cervical spine).mp. (40)
- 4 ((Clinical or critical or treatment) adj3 (pathway\* or protocol\*)).mp. (52575)
- 5 (algorithm\* or guideline\*).mp. (488839)
- 6 (decision adj3 (tree\* or rule\* or tool\*)).mp. (15657)
- 7 (triage or protocol\*).mp. (366206)
- 8 or/1-7 [Triage tool keywords] (856300)
- 9 exp Guideline/ (25808)
- 10 Guideline Adherence/ (21997)
- 11 exp guidelines as topic/ (117788)
- 12 exp algorithms/ (181301)
- 13 exp Clinical Protocols/ [includes antineoplastic protocols] (127941)
- 14 Decision Trees/ (8964)
- 15 exp decision support techniques/ [includes data interpretation, statistical] (61776)
- 16 Critical Pathways/ (4775)
- 17 triage/ (8242)
- 18 or/9-17 [Triage tool MeSH terms] (522506)
- 19 8 or 18 [Triage tools] (907178)
- 20 ((neurolog\* or physical\* or clinical\*) adj3 (exam\* or assess\* or sign\*)).mp. (384485)



- 21 MRI\* .mp. (359278)
- 22 (CT\* or Computed Tomography or CAT scan\*).mp. (384167)
- 23 (X ray\* or x-ray\* or xray\* or radiogra\* or roentgenogra\*).mp. (734670)
- 24 Imaging.mp. (660363)
- 25 or/20-24 [Reference standard keywords] (1748610)
- 26 exp physical examination/ or exp neurologic examination/ (1075565)
- 27 exp trauma severity indices/[includes Glasgow Coma Scale, Injury Severity Score, others] (24531)
- 28 "Severity of Illness Index"/ [not exploded leave out Karnofsky Performance Status cancer ADL measure] (173516)
- 29 X-Rays/ (16413)
- 30 Tomography/ or exp Tomography, Emission-Computed/ or exp Tomography, X-Ray/ [includes tomography, x-ray computed] (386295)
- 31 exp Magnetic Resonance Imaging/ (317050)
- 32 Radiography/ (24883)
- 33 or/26-32 [Reference standard MeSH terms] (1874249)
- 34 25 or 33 [Reference standard] (2858529)
- 35 19 or 34 [Triage tools or reference standard] (3572718)
- 36 ((Cervical spine or c-spine) adj5 clear\*).mp. (241)
- 37 (cervical adj5 (trauma\* or injur\* or fracture\* or sublux\* or dislocat\* or avuls\* or instab\*)).mp. [cervical spine injury, cervical spine trauma] (10352)
- 38 (Spinal cord injury without radiographic abnormality or SCIWORA).mp. (113)
- 39 or/36-38 [cervical trauma keywords] (10494)
- 40 exp Cervical Vertebrae/ [includes axis and atlas] (30752)
- 41 exp Neck Injuries/ [includes whiplash injuries] (6628)
- 42 exp Spinal Injuries/ [includes spinal fractures] (17863)
- 43 exp Spinal Cord Injuries/ [includes spinal cord compression, others] (38273)
- 44 spinal fractures/ (10537)
- 45 or/40-44 [cervial trauma MeSH terms] (81875)
- 46 39 or 45 [cervical trauma terms] (84419)
- 47 (Pediatric\* or paediatric\* or peadiatric\*).mp. (239652)
- 48 (Child\*).mp. (1864223)
- 49 (neonate\* or newborn\* or new-born\*).mp. (639758)
- 50 (infant\* or baby or babies or toddler\*).mp. (1042256)
- 51 (adolescen\* or juvenile\* or youth\* or teen\* or preteen\*).mp. (1709977)
- 52 or/47-51 [pediatric keywords] (3315420)
- 53 exp Pediatrics/ [includes perinataology, neonatology] (44737)
- 54 exp Child/ [includes child, preschool] (1562070)



55 exp Infant/[includes infant, newborn] (946461)

56 Adolescent/ (1632349)

57 or/53-56 [pediatric MeSH terms] (2910260)

58 52 or 57 [Pediatric terms] (3316163)

59 35 and 46 and 58 (7985)

\* .mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept word, rare disease supplementary concept word, unique identifier]

## **MEDLINE In-Process & Other Non-Indexed Citations**

Search 24 Feb 2015

- 1 (NEXUS or CCR).mp. (492)
- 2 National Emergency X-Radiography.mp. (3)
- 3 (Canadian c-spine or Canadian cervical spine).mp. (13)
- 4 ((Clinical or critical or treatment) adj3 (pathway\* or protocol\*)).mp. (3193)
- 5 (algorithm\* or guideline\*).mp. (49484)
- 6 (decision adj3 (tree\* or rule\* or tool\*)).mp. (1156)
- 7 (triage or protocol\*).mp. (29444)
- 8 or/1-7 [Triage tool keywords] (79262)
- 9 exp Guideline/ or / (62)
- 10 Guideline Adherence/ (0)
- 11 exp guidelines as topic//(0)
- 12 exp algorithms/ (0)
- 13 exp Clinical Protocols/ [includes antineoplastic protocols] (0)
- 14 Decision Trees/(0)
- 15 exp decision support techniques/ [includes data interpretation, statistical] (0)
- 16 Critical Pathways/ (0)
- 17 triage/ (0)
- 18 or/9-17 [Triage tool MeSH terms] (62)
- 19 8 or 18 [Triage tools] (79262)
- 20 ((neurolog\* or physical\* or clinical\*) adj3 (exam\* or assess\* or sign\*)).mp. (30580)
- 21 MRI\* .mp. (24133)
- 22 (CT\* or Computed Tomography or CAT scan\*).mp. (39230)
- 23 (X ray\* or x-ray\* or xray\* or radiogra\* or roentgenogra\*).mp. (62975)
- 24 Imaging.mp. (61152)
- 25 or/20-24 [Reference standard keywords] (175595)
- 26 exp physical examination/ or exp neurologic examination/ (0)



- 27 exp trauma severity indices/[includes Glasgow Coma Scale, Injury Severity Score, others] (0)
- 28 "Severity of Illness Index"/ [not exploded leave out Karnofsky Performance Status cancer ADL measure] (0)
- 29 X-Rays/ (0)
- 30 Tomography/ or exp Tomography, Emission-Computed/ or exp Tomography, X-Ray/ [includes tomography, x-ray computed] (0)
- 31 exp Magnetic Resonance Imaging/(0)
- 32 Radiography/ (0)
- 33 or/26-32 [Reference standard MeSH terms] (0)
- 34 25 or 33 [Reference standard] (175595)
- 35 19 or 34 [Triage tools or reference standard] (244188)
- 36 ((Cervical spine or c-spine) adj5 clear\*).mp. (20)
- 37 (cervical adj5 (trauma\* or injur\* or fracture\* or sublux\* or dislocat\* or avuls\* or instab\*)).mp. [cervical spine injury, cervical spine trauma] (833)
- 38 (Spinal cord injury without radiographic abnormality or SCIWORA).mp. (10)
- 39 or/36-38 [cervical trauma keywords] (848)
- 40 exp Cervical Vertebrae/[includes axis and atlas] (0)
- 41 exp Neck Injuries/[includes whiplash injuries] (0)
- 42 exp Spinal Injuries/ [includes spinal fractures] (0)
- 43 exp Spinal Cord Injuries/ [includes spinal cord compression, others] (0)
- 44 spinal fractures/(0)
- 45 or/40-44 [cervial trauma MeSH terms] (0)
- 46 39 or 45 [cervical trauma terms] (848)
- 47 (Pediatric\* or paediatric\* or peadiatric\*).mp. (20264)
- 48 (Child\*).mp. (67273)
- 49 (neonate\* or newborn\* or new-born\*).mp. (9578)
- 50 (infant\* or baby or babies or toddler\*).mp. (19120)
- 51 (adolescen\* or juvenile\* or youth\* or teen\* or preteen\*).mp. (24018)
- 52 or/47-51 [pediatric keywords] (106495)
- 53 exp Pediatrics/ [includes perinataology, neonatology] (0)
- 54 exp Child/ [includes child, preschool] (0)
- 55 exp Infant/[includes infant, newborn] (0)
- 56 Adolescent/(0)
- 57 or/53-56 [pediatric MeSH terms] (0)
- 58 52 or 57 [Pediatric terms] (106495)
- 59 35 and 46 and 58 (39)



#### **EMBASE**

Search 24 Feb 2015

- 1 (NEXUS or CCR).mp. (7884)
- 2 National Emergency X-Radiography.mp. (59)
- 3 (Canadian c-spine or Canadian cervical spine).mp. (74)
- 4 ((Clinical or critical or treatment) adj3 (pathway\* or protocol\*)).mp. (116085)
- 5 (algorithm\* or guideline\*).mp. (665390)
- 6 (decision adj3 (tree\* or rule\* or tool\*)).mp. (16747)
- 7 (triage or protocol\*).mp. (426356)
- 8 or/1-7 (1087993)
- 9 practice guideline/ or clinical pathway/ or clinical protocol/ (315692)
- 10 exp algorithm/ (188334)
- 11 "decision tree"/ (6358)
- 12 exp decision support system/ (14055)
- 13 emergency health service/ [used for triage] (69092)
- 14 or/9-13 (575274)
- 15 8 or 14 (1149485)
- 16 ((neurolog\* or physical\* or clinical\*) adj3 (exam\* or assess\* or sign\*)).mp. (779457)
- 17 MRI\*.mp. (587063)
- 18 (CT\* or Computed Tomography or CAT scan\*).mp. (600538)
- 19 (X ray\* or x-ray\* or xray\* or radiogra\* or roentgenogra\*).mp. (876246)
- 20 Imaging.mp. (1064732)
- 21 or/16-20 (2762608)
- 22 exp physical examination/ (160790)
- 23 exp neurologic examination/ (350398)
- 24 exp injury scale/ [used for trauma severity indices] (29378)
- 25 X ray/ (41737)
- 26 exp tomography/ (728786)
- 27 exp computer assisted tomography/ (636166)
- 28 exp nuclear magnetic resonance imaging/ (576997)
- 29 exp radiography/ (897201)
- 30 or/22-29 (2073134)
- 31 21 or 30 (3328617)
- 32 15 or 31 (4251928)
- 33 ((Cervical spine or c-spine) adj5 clear\*).mp. (344)



- 34 (cervical adj5 (trauma\* or injur\* or fracture\* or sublux\* or dislocat\* or avuls\* or instab\*)).mp. (17235)
- 35 (Spinal cord injury without radiographic abnormality or SCIWORA).mp. (181)
- 36 or/33-35 (17402)
- 37 exp cervical spine/ (27701)
- 38 exp neck injury/ (10805)
- 39 exp spine injury/ (30879)
- 40 exp spinal cord injury/ (54289)
- 41 exp spine fracture/ (15827)
- 42 or/37-41 (112517)
- 43 36 or 42 (115463)
- 44 (Pediatric\* or paediatric\* or peadiatric\*).mp. (389216)
- 45 (Child or children or childhood).mp. (2006835)
- 46 (neonate\* or newborn\* or new-born\*).mp. (567587)
- 47 (infant\* or baby or babies or toddler\*).mp. (778286)
- 48 (adolescen\* or juvenile\* or youth\* or teen\* or preteen\*).mp. (1399956)
- 49 or/44-48 (3309713)
- 50 exp pediatrics/ (77383)
- 51 exp child/ (2059816)
- 52 exp infant/ (857030)
- 53 exp adolescent/ (1253833)
- 54 exp juvenile/ (2715056)
- 55 exp adolescence/ (66747)
- 56 exp childhood/ (50991)
- 57 exp childhood injury/ (7203)
- 58 or/50-57 (2785218)
- 59 49 or 58 (3349155)
- 60 32 and 43 and 59 (9187)

### **CENTRAL**

- Search 24 Feb 2015
- #1 NEXUS or CCR 783
- #2 National Emergency X-Radiography 2
- #3 Canadian c-spine or Canadian cervical spine 59
- #4 ((Clinical or critical or treatment) near/3 (pathway\* or protocol\*)) 8326
- #5 algorithm\* or guideline\* 24279
- #6 (decision near/3 (tree\* or rule\* or tool\*)) 2370



#7 triage or protocol\* 56538

#8 #1 or #2 or #3 or #4 or #5 or #6 or #7 76450

#9 MeSH descriptor: [Guideline] explode all trees 19

#10 MeSH descriptor: [Practice Guideline] 15

#11 MeSH descriptor: [Guideline Adherence] 739

#12 MeSH descriptor: [Guidelines as Topic] explode all trees 2078

#13 MeSH descriptor: [Practice Guidelines as Topic] 1770

#14 MeSH descriptor: [Algorithms] explode all trees 3040

#15 MeSH descriptor: [Clinical Protocols] explode all trees 13095

#16 MeSH descriptor: [Decision Trees] 895

#17 MeSH descriptor: [Decision Support Techniques] explode all trees 3202

#18 MeSH descriptor: [Critical Pathways] 262

#19 MeSH descriptor: [Triage] 258

#20 #9 or #10 or #11 or #12 or #13 or #14 or #15 or #16 or #17 or #18 or #19 22374

#21 #8 or #20 78691

#22 ((neurolog\* or physical\* or clinical\*) near/3 (exam\* or assess\* or sign\*)) 45542

#23 MRI\* 11288

#24 CT\* or Computed Tomography or CAT scan 56407

#25 X ray\* or x-ray\* or xray\* or radiogra\* or roentgenogra\* 23437

#26 Imaging 21622

#27 #22 or #23 or #24 or #25 or #26 123319

#28 MeSH descriptor: [Physical Examination] explode all trees 72073

#29 MeSH descriptor: [Neurologic Examination] explode all trees 16982

#30 MeSH descriptor: [Trauma Severity Indices] explode all trees 993

 $\#31\ MeSH\ descriptor$ : [Severity of Illness Index] this term only 14375

#32 MeSH descriptor: [X-Rays] 44

#33 MeSH descriptor: [Tomography] explode all trees 11885

#34 MeSH descriptor: [Tomography, Emission-Computed] explode all trees 2630

#35 MeSH descriptor: [Tomography, X-Ray] explode all trees 4107

#36 MeSH descriptor: [Magnetic Resonance Imaging] explode all trees 5716

#37 MeSH descriptor: [Radiography] explode all trees 13863

 $\#38\ \#28\ or\ \#29\ or\ \#30\ or\ \#31\ or\ \#32\ or\ \#33\ or\ \#34\ or\ \#35\ or\ \#36\ or\ \#37\ 103001$ 

#39 #27 or #38 199647

#40 #21 or #39 253140

#41 ((Cervical spine or c-spine) near/5 clear\*) 10



#42 (cervical near/5 (trauma\* or injur\* or fracture\* or sublux\* or dislocat\*or avuls\* or instab\*)) 472

#43 Spinal cord injury without radiographic abnormality or SCIWORA 2

#44 #41 or #42 or #43 474

#45 MeSH descriptor: [Cervical Vertebrae] explode all trees 776

#46 MeSH descriptor: [Neck Injuries] explode all trees 205

#47 MeSH descriptor: [Spinal Injuries] explode all trees 720

#48 MeSH descriptor: [Spinal Cord Injuries] explode all trees 906

#49 MeSH descriptor: [Spinal Fractures] explode all trees 636

#50 #45 or #46 or #47 or #48 or #49 2442

#51 #44 or #50 2730

#52 Pediatric\* or paediatric\* or peadiatric\* 41377

#53 Child or children or childhood 94858

#54 neonate\* or newborn\* or new-born\* 19365

#55 infant\* or baby or babies or toddler\* 41412

#56 adolescen\* or juvenile\* or youth\* or teen\* or preteen\* 98728

#57 #52 or #53 or #54 or #55 or #56 176411

#58 MeSH descriptor: [Pediatrics] explode all trees 546

#59 MeSH descriptor: [Child] explode all trees 135

#60 MeSH descriptor: [Infant] explode all trees 13304

#61 MeSH descriptor: [Adolescent] 76925

#62 #58 or #59 or #60 or #61 89391

#63 #57 or #62 176421

#64 40 and 51 and 63 4201

#65 #64 in Trials 651

## **Science Citation Index**

Search 24 Feb 2015

# 4 1,220 #3 AND #2 AND #1

Indexes=SCI-EXPANDED Timespan=All years

#31,508,937

TOPIC: (Pediatric\* OR paediatric\* OR peadiatric\* OR Child OR children OR childhood OR neonate\* OR newborn\* OR new-born\* OR infant\* OR baby OR babies OR toddler\* OR adolescen\* OR juvenile\* OR youth\* OR teen\* OR preteen\*)

Indexes=SCI-EXPANDED Timespan=All years

#219,789

TS=(cervical spine clear\* OR c-spine clear\* OR clearing the c-spine OR clearing the cervical spine OR cervical trauma\* OR cervical injur\* OR cervical fracture\* OR cervical sublux\* OR cervical disloc\* OR cervical avuls\* OR cervical instab\* OR Spinal cord injury without radiographic abnormality OR SCIWORA)



Indexes=SCI-EXPANDED Timespan=All years

#14,736,061

TS=(NEXUS OR CCR OR National Emergency X-Radiography OR Canadian c-spine OR Canadian cervical spine OR Clinical pathway\* OR critical pathway\* OR treatment pathway\* OR clinical protocol\* OR treatment protocol\* OR algorithm\* OR guideline\* OR decision tree\* OR decision tool\* OR decision rule\* OR triage OR protocol\* OR neurolog\* assess\* OR neurolog\* exam\* OR neurolog\* sign\* OR physical\* exam\* OR physical\* assess\* OR physical\* sign\* OR Clinical\* exam\* OR clinical\* assess\* OR clinical\* sign\* OR MRI\* OR CT OR Computed Tomography OR CAT scan\* OR X ray\* OR x-ray\* OR radiogra\* OR roentgenogra\* OR imaging)

Indexes=SCI-EXPANDED Timespan=All years

#### **Proquest Dissertations & Theses database**

Search 24 Feb 2015

Advanced search:

all(((NEXUS OR CCR OR National Emergency X-Radiography OR Canadian c-spine OR Canadian cervical spine OR Clinical pathway\* OR critical pathway\* OR treatment pathway\* OR clinical protocol\* OR treatment protocol\* OR algorithm\* OR guideline\* OR decision tree\* OR decision tool\* OR decision rule\* OR triage OR protocol\* OR neurolog\* assess\* OR neurolog\* exam\* OR neurolog\* sign\* OR physical\* exam\* OR physical\* assess\* OR physical\* sign\* OR Clinical\* exam\* OR clinical\* assess\* OR clinical\* sign\* OR MRI\* OR CT\* OR Computed Tomography OR CAT scan\*OR X ray\* OR x-ray\* OR radiogra\* OR roentgenogra\* OR imaging) AND (cervical spine clear\* OR clearing the c-spine OR clearing the cervical spine OR cervical trauma\* OR cervical injur\* OR cervical fracture\* OR cervical sublux\* OR cervical disloc\* OR cervical avuls\* OR cervical instab\* OR Spinal cord injury without radiographic abnormality OR SCIWORA) AND (Pediatric\* OR paediatric\* OR peadiatric\* OR Child OR children OR childhood OR neonate\* OR newborn\* OR new-born\* OR infant\* OR baby OR babies OR toddler\* OR adolescen\* OR juvenile\* OR youth\* OR teen\* OR preteen\*)))

Additional limits - Source type: Conference Papers & Proceedings, Dissertations & Theses

#### **PubMed**

Search 24 Feb 2015. This search contained population terms.

((NEXUS OR CCR OR National Emergency X-Radiography OR Canadian c-spine OR Canadian cervical spine OR Clinical pathway\* OR critical pathway\* OR treatment pathway\* OR clinical protocol\* OR treatment protocol\* OR algorithm\* OR guideline\* OR decision tree\* OR decision tool\* OR decision rule\* OR triage OR protocol\* OR neurolog\* assess\* OR neurolog\* exam\* OR neurolog\* sign\* OR physical\* exam\* OR physical\* assess\* OR physical\* sign\* OR clinical\* exam\* OR clinical\* assess\* OR clinical\* sign\* OR MRI\* OR CT\* OR Computed Tomography OR CAT scan\*OR X ray\* OR x-ray\* OR xray\* OR radiogra\* OR roentgenogra\* OR imaging) **AND** (cervical spine clear\* OR c-spine clear\* OR clearing the c-spine OR clearing the cervical spine OR cervical trauma\* OR cervical injur\* OR cervical fracture\* OR cervical sublux\* OR cervical disloc\* OR cervical avuls\* OR cervical instab\* OR Spinal cord injury without radiographic abnormality or SCIWORA) **AND** (Pediatric\* OR paediatric\* OR peadiatric\* OR Child OR children OR childhood OR neonate\* OR newborn\* OR new-born\* OR infant\* OR baby OR babies OR toddler\* OR adolescen\* OR juvenile\* OR youth\* OR teen\* OR preteen\*) **AND** (pubstatusaheadofprint OR publisher[sb] or pubmednotmedline[sb]))

Searched 5 March 2015. This search did not contain population terms.

((NEXUS OR CCR OR National Emergency X-Radiography OR Canadian c-spine OR Canadian cervical spine OR Clinical pathway\* OR critical pathway\* OR treatment pathway\* OR clinical protocol\* OR treatment protocol\* OR algorithm\* OR guideline\* OR decision tree\* OR decision tool\* OR decision rule\* OR triage OR protocol\* OR neurolog\* assess\* OR neurolog\* exam\* OR neurolog\* sign\* OR physical\* exam\* OR physical\* assess\* OR physical\* sign\* OR clinical\* exam\* OR clinical\* assess\* OR clinical\* sign\* OR MRI\* OR CT\* OR Computed Tomography OR CAT scan\*OR X ray\* OR x-ray\* OR xray\* OR radiogra\* OR roentgenogra\* OR imaging) AND (cervical spine clear\* OR c-spine clear\* OR clearing the c-spine OR clearing the cervical spine OR cervical trauma\* OR cervical injur\* OR cervical fracture\* OR cervical sublux\* OR cervical disloc\* OR cervical avuls\* OR cervical instab\* OR Spinal cord injury without radiographic abnormality or SCIWORA) AND (pubstatusaheadofprint OR publisher[sb] or pubmednotmedline[sb]))

## OpenGrey

Search 24 Feb 2015

((NEXUS OR CCR OR National Emergency X-Radiography OR Canadian c-spine OR Canadian cervical spine OR Clinical pathway\* OR critical pathway\* OR treatment pathway\* OR clinical protocol\* OR treatment protocol\* OR algorithm\* OR guideline\* OR decision tree\* OR decision tool\* OR decision rule\* OR triage OR protocol\* OR neurolog\* assess\* OR neurolog\* exam\* OR neurolog\* sign\* OR physical\* exam\* OR physical\* assess\* OR physical\* sign\* OR clinical\* exam\* OR clinical\* assess\* OR clinical\* sign\* OR MRI\* OR CT\* OR Computed Tomography OR CAT scan\*OR X ray\* OR x-ray\* OR xray\* OR radiogra\* OR roentgenogra\* OR imaging) AND (cervical spine clear\* OR clearing the c-spine OR clearing the cervical spine OR c



## ClinicalTrialsgov

Search 24 Feb 2015

((cervical spine OR c-spine) AND (fracture OR injury OR trauma OR avulsion OR dislocation OR instability) AND (NEXUS OR "National Emergency X-Radiography" OR "Canadian c-spine" OR clearing OR clearance OR decision OR algorithm OR pathway OR triage))

#### **ICTRP**

Search 24 Feb 2015

(NEXUS OR National Emergency X-Radiography OR Canadian c-spine OR Canadian Cervical Spine) AND (cervical fracture OR cervical injury OR cervical trauma OR cervical dislocation OR cervical instability OR cervical avulsion)

#### CDSR, DARE, HTA

Searched 25 Feb 2015

clearance:ti,ab,kw or cervical spine

#### ARIF

Search 24 Feb 2015

Advanced search, all indexed fields: Clearance or cervical spine

## **DTA Trials Register**

Searched 10 March 2015.

We received the following report from the information specialist of the Renal group:

"There are no studies relating to your review in the DTAS Register. I used keywords from your review title plus other broader target condition words e.g. spinal injur\* spinal trauma, head injur\* etc. I found only 3 studies, all of which were in adults only, and which were using radiological modalities to screen for blunt trauma injuries, including cervical arteries. I also used the test names you mentioned, but did not retrieve anything."

## Medion

Searched October 2013

ICPC code = Musculoskeletal OR Neurological

And

Abstract = clearance or "cervical spine"

## CONTRIBUTIONS OF AUTHORS

- Conceiving the review: Slaar, Fockens
- Designing the review: Slaar
- Co-ordinating the review: Van Rijn, Schep
- Data collection for the review: Information Specialist of the Cochrane Back and Neck Group
- Designing search strategies: Information Specialist of the Cochrane Back and Neck Group
- Undertaking searches: Information Specialist of the Cochrane Back and Neck Group
- Screening search results: Fockens, Slaar, Schep
- Organizing retrieval of papers: Fockens
- Screening retrieved papers against inclusion criteria: Fockens, Slaar, Schep
- · Appraising quality of papers: Slaar, Fockens
- · Extracting data from papers: Slaar, Fockens
- Writing to authors of papers for additional information: Slaar, Van Rijn
- · Providing additional data about papers: Slaar
- Obtaining and screening data on unpublished studies: Slaar, Fockens
- · Data management for the review: Slaar, Wang, Schep
- Entering data into Review Manager 5: Slaar, Wang



- · Analysis of data: Wang
- · Interpretation of data: Schep, Slaar, Wang
- Providing a methodological perspective: Schep, Van Rijn, Wang
- Providing a clinical perspective: Maas, Goslings, Van Rijn, Wilson
- Writing the review: Slaar
- Providing general advice on the review: Goslings, Wilson, Maas

## **DECLARATIONS OF INTEREST**

Annelie Slaar has no conflicts of interest.

M. Matthijs Fockens has no conflicts of interest.

Junfeng Wang has no conflicts of interest.

Mario Maas has no conflicts of interest.

David J Wilson has no conflicts of interest.

J Carel Goslings has no conflicts of interest.

Niels WL Schep has no conflicts of interest.

Rick R van Rijn has no conflicts of interest

#### SOURCES OF SUPPORT

#### Internal sources

• Academic Medical Center, Amsterdam, Netherlands.

#### **External sources**

· No sources of support supplied

## DIFFERENCES BETWEEN PROTOCOL AND REVIEW

We planned to meta-analyse the sensitivity and specificity of the tools with a bivariate model. However, we identified only three studies that met the inclusion criteria, and the outcomes of the studies were too diverse for us to perform meta-analyses in this review. For the same reason an analysis of heterogeneity could not be completed.

We did not anticipate in the protocol that we would encounter studies with mixed populations (adults and children) in which we could not extract the data for both groups ourselves. When this occurred during the review process, we attempted to contact the authors to obtain this data.

## INDEX TERMS

## **Medical Subject Headings (MeSH)**

\*Decision Support Techniques; Cervical Vertebrae [diagnostic imaging] [\*injuries]; Checklist; Cohort Studies; Magnetic Resonance Imaging; Radiography; Reference Standards; Spinal Injuries [\*diagnostic imaging] [etiology]; Tomography, X-Ray Computed; Triage [\*methods]; Wounds, Nonpenetrating [\*complications] [diagnostic imaging]

## MeSH check words

Child; Humans