

No.	Part	Scheme
1	(a)	% abundance $^{24}\text{Mg} = 78.99$ % abundance $^{25}\text{Mg} = x$ % abundance $^{26}\text{Mg} = y$ $78.99 + x + y = 100$ $y = 21.01 - x$ ✓ <i>must have</i> <u>Average atomic mass</u> = $\frac{\sum Q_i M_i}{\sum Q_i}$ @ $24.3 = \frac{(78.99 \times 23.985) + (x)(24.986) + (21.01 - x)(25.983)}{100}$ $x = 10.51$ $y = 21.01 - x$ $= 21.01 - 10.51$ $= 10.50$ ✓ m_3 % abundance $^{25}\text{Mg} = 10.51$ (a) 10.52 % abundance $^{26}\text{Mg} = 10.50$ (a) 10.49
	(b)	$\text{SO}_2 + 2\text{H}_2\text{O} \rightarrow \text{SO}_4^{2-} + 4\text{H}^+ + 2\text{e}^-$ ✓ m_1 $\text{Br}_2 + 2\text{e}^- \rightarrow 2\text{Br}^-$ ✓ m_7 $\text{SO}_2 + \text{Br}_2 + 2\text{H}_2\text{O} \rightarrow \text{SO}_4^{2-} + 2\text{Br}^- + 4\text{H}^+$ ✓ m_9 (c)(i) Assume volume of solution = 1 L Molarity = $\frac{n_{\text{solute}}}{V_{\text{solution}}}$ @ $0.2 \text{ mol L}^{-1} = \frac{n_{\text{solute}}}{1 \text{ L}}$ $n_{\text{solute}} = 0.2 \text{ mol}$ $m_{\text{solute}} = 0.2 \text{ mol} \times 40 \text{ g mol}^{-1} = 8.0 \text{ g}$ ✓ m_{10} In 250 mL, $m_{\text{solute}} = 2.0 \text{ g}$ unit insst ✓ m_9 <u>Alternative:</u> $n_{\text{solute}} = MV = (0.2) \left(\frac{250}{1000} \right) = 0.05 \text{ mol}$ $m_{\text{solute}} = 0.05 \text{ mol} \times 40 \text{ g mol}^{-1} = 2.0 \text{ g}$ unit insst — m_{10} ecf


No.	Paragraf	Scheme
	(c)(ii)	$M_1 V_1 = M_2 V_2$ $(0.2 \text{ M})(250 \text{ mL}) = (0.1 \text{ M})(V_2)$ $V_2 = 500 \text{ mL}$ Volume of $\text{H}_2\text{O} = (500 - 250) \text{ mL}$ $= 250 \text{ mL}$ <i>unt. ingat</i> <i>M₂₂ new addition</i>
	(d)(i)	$\text{O}_3 + \text{NO} \rightarrow \text{O}_2 + \text{NO}_2$ <i>0.015 accepted</i> $n_{\text{O}_3} = \frac{0.740 \text{ g}}{48 \text{ g mol}^{-1}} = 0.0154 \text{ mol}$ <i>0.022</i> $n_{\text{NO}} = \frac{0.670 \text{ g}}{30 \text{ g mol}^{-1}} = 0.0223 \text{ mol}$ From equation; 1 mol of $\text{O}_3 = 1 \text{ mol}$ of NO 0.0154 mol of $\text{O}_3 = 0.0154 \text{ mol}$ of NO (needed) Since n_{NO} given $> n_{\text{NO}}$ needed <i>m.g</i> hence NO is excess reactant while O_3 is limiting reactant. <i>comparison</i> <u>Alternative:</u> $n_{\text{O}_3} = \frac{0.740 \text{ g}}{48 \text{ g mol}^{-1}} = 0.0154 \text{ mol}$ <i>0.022</i> $n_{\text{NO}} = \frac{0.670 \text{ g}}{30 \text{ g mol}^{-1}} = 0.0223 \text{ mol}$ 1 mol O_3 produces 1 mol NO_2 Thus, 0.0154 mol NO_2 is produced and 1 mol NO produces 1 mol NO_2 Thus, 0.0223 mol NO_2 is produced Since O_3 produce less NO_2 hence O_3 is the limiting reagent. <i>only mol</i>
	(d)(ii)	$\text{NO}_2 \text{ produced} = 0.0154 \text{ mol} \times 46 \text{ g mol}^{-1}$ $= 0.7084 \text{ g} @ 0.709 \text{ g}$ <i>0.015</i> <i>0.69</i> <i>m₂₀</i> <i>m₂₁</i>
		TOTAL


must

2 must

2cf

m₂₀ m₂₁m₂₀


NO.	PART	SCHEME
2	(a)	Energy $\textcircled{E}$ ΔE (no accepted) Energy level X  Axis (energy and energy level) Line spacing (energy level getting close as energy increases) Arrow transition $\frac{1}{\lambda} = R_H \left(\frac{1}{n_1^2} - \frac{1}{n_2^2} \right) @$ M_5 $\frac{1}{\lambda} = 1.097 \times 10^7 \left(\frac{1}{1^2} - \frac{1}{6^2} \right)$ if n, n_2 any number accepted for $\lambda = 9.3762 \times 10^{-8} \text{ m} @ 93.762 \text{ nm}$ <u>Alternative:</u> $\Delta E = R_H \left(\frac{1}{n_i^2} - \frac{1}{n_f^2} \right) /$ $\Delta E = 2.18 \times 10^{-18} \left(\frac{1}{6^2} - \frac{1}{1^2} \right)$ $\Delta E = -2.12 \times 10^{-18} \text{ J}$ $\Delta E = \frac{hc}{\lambda} /$ $\lambda = \frac{hc}{\Delta E} = \frac{(6.6256 \times 10^{-34} \text{ J s})(3.0 \times 10^8 \text{ m s}^{-1})}{2.12 \times 10^{-18}}$ $\lambda = 9.375 \times 10^{-8} \text{ m}$ all arrow correct must have both eq.

NO.	JAWAB	SOLUSI
	(b)	${}_{13}\text{X}: 1s^2 2s^2 2p^6 3s^2 3p^1$ Orbital diagram (full arrow) also accepted When $\ell = 0 \rightarrow s$ orbital Thus, number of electrons in s orbital = <u>6</u> Orbital of valence electrons with $\ell = 1$ are 3p orbitals. lorek not insert  3px, must have energy level 3py 3pz Kalan x 6 - 1
		TOTAL

* wrong cancel wright

m_l
 $m_{l,0}$


NO	PART	SCHEME
3	(a)(i)	Total valence electron for $\text{SeO}_2 = (1 \times 6) + (2 \times 6) = 18 / m_1$ $\text{:}\ddot{\text{O}}=\ddot{\text{Se}}=\ddot{\text{O}}\text{:}$ @ $\text{:}\ddot{\text{O}}-\ddot{\text{Se}}=\ddot{\text{O}}\text{:}$ octet
	(a)(ii)	VSEPR theory states that a central atom with M_3 if 4 must (2 bonding pair and 1 non-bonding pair) therefore bent molecular geometry @ or lonepair or drawings also accepted $\text{:}\ddot{\text{O}}=\ddot{\text{Se}}=\ddot{\text{O}}\text{:}$ @ $\text{:}\ddot{\text{O}}-\ddot{\text{Se}}=\ddot{\text{O}}\text{:}$
	(a)(iii)	The orbital hybridization is sp^2
	(a)(iv)	(x da each also accept) Each $\text{Se}=\text{O}$ bond is polar towards the oxygen atom. or $\text{O}=\ddot{\text{Se}}\text{:}$ The bonding dipole moments are not cancelled out. @ $\mu \neq 0$ @ unsym Therefore the molecule is also polar. must have both bond dipole cannot cancel to each other
	(b)	Diagram with label sea of delocalised valence electrons Number of delocalised electrons equal with charge contribute (insist) Ca^{2+} equation Every Ca atom releases two electrons and form Ca^{2+} ion. Ca^{2+} equation Metallic bond forms from the electrostatic attraction between positive ions with the negatively charge sea of valence electrons. or Ca^{2+} attractive between with sea of elec Calcium can conduct electricity due to free electrons / delocalised electrons.



Mg depend on M_7

attractive in
electrostatic
forces not
accepted

No.	Mark	Scheme
	(c)	- AlCl₃ is covalent compound. - It has no free moving <u>electron</u> to conduct electricity. - Al is a metal. <i>insist ionx insist</i> - Each <u>Al atom</u> <i>or donate or its</i> releases/contributes three valence electrons. - It has delocalised electrons to conduct electricity. <u>Alternative:</u> - Al is a metal. - Has overlapping between conduction band and valence band @ <i>@ no forbidden gap</i> No band gap between conduction band and valence band. - Electron can move from the conduction band to the valence band to conduct electricity.
		Minimum
		Maximum

No.	Date	Signature
4	(a)	$P_T = P_{H_2} + P_{H_2O}$ $\textcircled{a}$ $P_{H_2} = (0.980 - 0.0313) \text{ atm}$ $\textcircled{a}$ $= 0.9487 \text{ atm}$ $\textcircled{a}$ m_1 $PV = nRT$ $\textcircled{a}$ $n_{H_2} = \frac{(0.9487 \text{ atm})(7.80 \text{ L})}{(0.08206 \text{ L atm K}^{-1} \text{ mol}^{-1})(298.15 \text{ K})}$ $\textcircled{a}$ m_2 $= 0.302 \text{ mol}$ $\textcircled{0.3}$ From equation: $1 \text{ mol } H_2 \equiv 1 \text{ mol Zn}$ $0.302 \text{ mol } H_2 \equiv 0.302 \text{ mol Zn}$ $\textcircled{a}$ $\text{comparison } m_3$ Mass of zinc $= 0.302 \text{ mol} \times 65.4 \text{ g mol}^{-1}$ M_4 $= 19.78 \text{ g}$ M_5 unit x insist
	(b)(i)	 $P \text{ (atm)}$ 218 1.00 0.006 Solid Liquid Gas 0.01 100 374 273.15 373.15 647.15 K $T \text{ (}^\circ\text{C)}$ Shape and label axis Label phase, triple and critical points Boiling point line (100°C, 1 atm)
	(b)(ii)	Water is denser than ice because melting point of ice decreases with pressure. $\textcircled{a}$ (Density + Volume) $\textcircled{a}$ $\text{Ice has open structure}$ water (also accepted)

$PV = \frac{n}{M_r} RT$ $\textcircled{a}$ m_4 eq only if M_r is correct

volume of ice > volume

NO.	PART	SOLUTION
5	(a)	At $400^{\circ}\text{C} = 400^{\circ}\text{C} + 273.15\text{ K} = 673.15\text{ K}$ <i>n kecil</i> → $K_c = K_p (RT)^{-\Delta n}$ <i>m1</i> $\Delta n = (5 - 4) = 1$ <i>m2</i> $K_c = (8.3 \times 10^{-3})(0.08206 \times 673.15\text{ K})^{-1}$ <i>m3</i> $K_c = 1.5025 \times 10^{-4}$ <i>m4</i>
	(b)	Mass of HCl will increase. <i>m5 m6</i> Since this reaction is <u>endothermic</u>, based on Le Chatelier's principle, <u>increasing temperature</u> of the reaction will <u>shift the equilibrium position to the right</u>. <i>forward</i> ✓ <i>for system to increase</i> <i>equilibrium to the right</i>
	(c)	From (a): $K_c = 1.5025 \times 10^{-4}$ $Q_c = \frac{[\text{HCl}]^4 [\text{O}_2]}{[\text{Cl}_2]^2 [\text{H}_2\text{O}]^2}$ <i>m8</i> $Q_c = \frac{(0.25)^4 (0.05)}{(0.70)^2 (0.35)^2}$ <i>@ m9</i> $= 3.254 \times 10^{-3}$ $Q_c > K_c$ Reaction is NOT at equilibrium. The net reaction proceed to the left. <i>together</i> <i>backward reverse</i> ✓
		TOTAL MAXIMUM

NO	PART	SCHODVI
	(b)(ii)	<u>Addition of $\text{Mg}(\text{NO}_3)_2$</u> $\text{Mg}(\text{OH})_2 (\text{s}) \rightleftharpoons \text{Mg}^{2+} (\text{aq}) + 2\text{OH}^- (\text{aq})$ $\text{Mg}(\text{NO}_3)_2 (\text{aq}) \xrightarrow{\quad} \text{Mg}^{2+} (\text{aq}) + 2\text{NO}_3^- (\text{aq})$ Mg^{2+} is common ion / m_{13} Based on Le Chatelier's Principle: • When concentration of Mg^{2+} ion increases, the equilibrium position is shifted to the left • More $\text{Mg}(\text{OH})_2$ precipitate is formed and solubility of $\text{Mg}(\text{OH})_2$ will decrease / m_{12} arrow in phase x
		TOTAL
		MAXIMUM