Logical Equivalence and Negation

When two propositions say the same thing, and how to say the opposite

Last modified: 2026-10-29 14:19:01 (PDT)

Source

Adapted from Kameron Decker Harris's Fall 2026 lecture notes "Logic 3", which are typed transcriptions of Scott Wehrwein's Spring 2025 handwritten lecture notes. The truth table for the distributive law of "and" over "or" comes from those notes; the exercises are new.

Logical equivalence

Exercise 1 (A rule with a redundant part)**.** A server grants access to a user who is an administrator, or who is an administrator and is on the office network. Let P be "the user is an administrator" and Q be "the user is on the office network".
Make a truth table for $P \vee (P \wedge Q)$. In which rows does it have the same truth value as P?

Solution

Solution 1.

P	Q	$P \wedge Q$	$P \vee (P \wedge Q)$
T	T	T	T
T	F	F	T
F	T	F	F
F	F	F	F

The last column matches the P column in every row, so the rule grants access exactly to the administrators, and the part about the office network changes nothing.

Definition 1 (Logical equivalence)**.** Two propositions A and B, each built from some of the propositions $P_1, \ldots, P_n$, are **logically equivalent**, written $A \equiv B$, if they have the same truth value in every row of a truth table (Definition: Truth table), that is, for every combination of truth values of $P_1, \ldots, P_n$.

Example 1 (An equivalence)**.** By Exercise 1, $P \vee (P \wedge Q) \equiv P$ (Definition 1): the truth table of $P \vee (P \wedge Q)$ has the same column as P, in a table with columns for both P and Q.

Remark 1 (Notation for logical equivalence). $A \equiv B$ (Definition 1) is a claim about A and B, not a new proposition built from them with a logical connective (Definition: Logical connective). Some books write $A \Leftrightarrow B$ or $A \iff B$ for it instead, which is easy to confuse with the biconditional (Definition: If and only if); these notes use $\equiv$.
Logical equivalence can be chained: if $A \equiv B$ and $B \equiv C$, then $A \equiv C$, since in each row A has the value of B, which has the value of C.

Exercise 2 (Not both)**.** A continuous integration server runs two test suites. Let P be "the unit tests pass" and Q be "the integration tests pass".
Is "it is not the case that both suites pass", $\neg(P \wedge Q)$, logically equivalent (Definition 1) to "neither suite passes", $\neg P \wedge \neg Q$?

Solution

Solution 2.

P	Q	$P \wedge Q$	$\neg(P \wedge Q)$	$\neg P$	$\neg Q$	$\neg P \wedge \neg Q$
T	T	T	F	F	F	F
T	F	F	T	F	T	F
F	T	F	T	T	F	F
F	F	F	T	T	T	T

No. The two columns differ in the second and third rows, where exactly one suite passes: then it is true that not both pass, but false that neither passes.

Definition 2 (Tautology)**.** A **tautology** is a proposition that is true in every row of its truth table (Definition: Truth table).

Example 2 (A tautology)**.** $P \vee \neg P$ is a tautology (Definition 2):

P	$\neg P$	$P \vee \neg P$
T	F	T
F	T	T

Definition 3 (Contradiction)**.** A **contradiction** is a proposition that is false in every row of its truth table (Definition: Truth table).

Example 3 (A contradiction)**.** $P \wedge \neg P$ is a contradiction (Definition 3):

P	$\neg P$	$P \wedge \neg P$
T	F	F
F	T	F

Remark 2 (Contradictions in proofs). A proof by contradiction is named for this kind of proposition: it assumes the negation of what it proves, and derives a contradiction (Definition 3), such as $Q \wedge \neg Q$.

Theorem 1 (Equivalence is a tautological biconditional)**.** *For propositions A and B, $A \equiv B$ (Definition 1) exactly when $A \Leftrightarrow B$ is a tautology (Definition 2).*

Proof

Proof. By Definition: If and only if, $A \Leftrightarrow B$ is true in a row of the truth table exactly when A and B have the same truth value in that row. So $A \Leftrightarrow B$ is true in every row exactly when A and B have the same truth value in every row, which is what $A \equiv B$ means. □

Example 4 (An equivalence as a tautology)**.** Since $P \vee (P \wedge Q) \equiv P$ (Example 1), Theorem 1 says that $(P \vee (P \wedge Q)) \Leftrightarrow P$ is a tautology.

Exercise 3 (Does "or" distribute over "and"?)**.** In arithmetic, multiplication distributes over addition: $a(b+c) = ab + ac$. Use a truth table to decide whether "or" distributes over "and": is $A \vee (B \wedge C) \equiv (A \vee B) \wedge (A \vee C)$?

Solution

Solution 3.

A	B	C	$B \wedge C$	$A \vee (B \wedge C)$	$A \vee B$	$A \vee C$	$(A \vee B) \wedge (A \vee C)$
T	T	T	T	T	T	T	T
T	T	F	F	T	T	T	T
T	F	T	F	T	T	T	T
T	F	F	F	T	T	T	T
F	T	T	T	T	T	T	T
F	T	F	F	F	T	F	F
F	F	T	F	F	F	T	F
F	F	F	F	F	F	F	F

Yes: the fifth and last columns match in every row. Addition does not distribute over multiplication in arithmetic: $1 + 2 \cdot 3 = 7$, but $(1 + 2)(1 + 3) = 12$. But "or" distributes over "and".

Theorem 2 (Distributive laws)**.** *For propositions A, B and C:*

1. $A \wedge (B \vee C) \equiv (A \wedge B) \vee (A \wedge C)$;
2. $A \vee (B \wedge C) \equiv (A \vee B) \wedge (A \vee C)$.

Proof

Proof. Law 2 is Solution 3. For law 1, the fourth and fifth columns of this table match in every row:

A	B	C	$A \wedge (B \vee C)$	$(A \wedge B) \vee (A \wedge C)$
T	T	T	T	T
T	T	F	T	T
T	F	T	T	T
T	F	F	F	F
F	T	T	F	F
F	T	F	F	F

F	F	T	F	F
F	F	F	F	F

□

Example 5 (Distributing "and")**.** For an integer x, "x is even, and $x > 0$ or $x < 0$" is logically equivalent, by law 1 of Theorem 2, to "x is even and $x > 0$, or x is even and $x < 0$".

Remark 3 (Checking an equivalence in Racket). A truth table is a loop over every combination of truth values, so a program can check an equivalence. In Racket, `for*/and` runs its body for every combination of `p` and `q`, and returns `#f` as soon as the body returns `#f`, and `#t` if it never does:

```
(for*/and ([p '(#t #f)]
           [q '(#t #f)])
  (equal? (or p (and p q)) p))
```

This returns `#t`, as Exercise 1 found. Replacing the body with `(equal? (not (and p q)) (and (not p) (not q)))` returns `#f`, as Exercise 2 found.

Negation

Exercise 4 (When is the door locked?)**.** A door opens when the badge is valid, the PIN is correct, and it is a weekday. Let B be "the badge is valid", N be "the PIN is correct" and W be "it is a weekday", so the door stays locked when $\neg(B \wedge N \wedge W)$ is true. Describe when the door stays locked with a sentence that puts "not" only in front of B, N or W, never in front of a compound proposition.

Solution

Solution 4. The door stays locked when the badge is not valid, the PIN is not correct, or it is not a weekday: $\neg B \vee \neg N \vee \neg W$. $B \wedge N \wedge W$ is false exactly when at least one of B, N and W is false.

Theorem 3 (Double negation)**.** *For a proposition P, $\neg\neg P \equiv P$ (Definition 1).*

Proof

Proof.

P	$\neg P$	$\neg\neg P$
T	F	T
F	T	F

The first and last columns match in both rows. □

Example 6 (Not unreadable)**.** "It is not the case that the file is not readable" means "the file is readable" (Theorem 3).

Theorem 4 (Replacing a part by an equivalent part)**.** *If $A \equiv B$ (Definition 1), then replacing A by B inside a larger proposition gives a proposition logically equivalent to the larger one. In particular, $\neg A \equiv \neg B$.*

Proof

Proof. The truth value of the larger proposition in a row of its truth table depends on the truth value of A in that row, not on how A is written. A and B have the same truth value in every row, so the two larger propositions do too. □

Example 7 (Replacing a part)**.** Since $P \vee (P \wedge Q) \equiv P$ (Example 1), Theorem 4 gives $R \wedge (P \vee (P \wedge Q)) \equiv R \wedge P$ and $\neg(P \vee (P \wedge Q)) \equiv \neg P$.

Theorem 5 (De Morgan's laws)**.** *For propositions P and Q (Wikipedia contributors 2026):*

1. $\neg(P \wedge Q) \equiv \neg P \vee \neg Q$;
2. $\neg(P \vee Q) \equiv \neg P \wedge \neg Q$.

Proof

Proof. For law 1, the fourth and last columns of this table match in every row:

P	Q	$P \wedge Q$	$\neg(P \wedge Q)$	$\neg P$	$\neg Q$	$\neg P \vee \neg Q$
T	T	T	F	F	F	F

T	F	F	T	F	T	T
F	T	F	T	T	F	T
F	F	F	T	T	T	T

Law 2 follows from law 1. Law 1 holds for any two propositions, so it holds for $\neg P$ and $\neg Q$:

$$\neg(\neg P \wedge \neg Q) \equiv \neg\neg P \vee \neg\neg Q.$$

By double negation (Theorem 3) and Theorem 4, the right side is logically equivalent to $P \vee Q$, so

$$\neg(\neg P \wedge \neg Q) \equiv P \vee Q.$$

Negating both sides (Theorem 4) gives

$$\neg\neg(\neg P \wedge \neg Q) \equiv \neg(P \vee Q),$$

and by double negation the left side is logically equivalent to $\neg P \wedge \neg Q$. So $\neg(P \vee Q) \equiv \neg P \wedge \neg Q$. □

Example 8 (Neither coordinate positive)**.** For real numbers x and y, by law 2 of Theorem 5, "not ($x > 0$ or $y > 0$)" is logically equivalent to "$x \leq 0$ and $y \leq 0$".

Exercise 5 (A broken deployment rule)**.** A team's rule is "if the tests pass, then the build is deployed". Let P be "the tests pass" and Q be "the build is deployed". The rule is broken on the days when $\neg(P \Rightarrow Q)$ is true.
Write $\neg(P \Rightarrow Q)$ without $\Rightarrow$, and in English.

Solution

Solution 5. The rule is broken on a day when the tests pass and the build is not deployed: $P \wedge \neg Q$. On a day when the tests fail, the rule is not broken, whether or not the build is deployed.
"If the tests pass, then the build is not deployed", $P \Rightarrow \neg Q$, is not the negation: on a day when the tests fail, it is true, and so is the rule.

Theorem 6 (Negating a conditional)**.** *For propositions P and Q, $\neg(P \Rightarrow Q) \equiv P \wedge \neg Q$.*

Proof

Proof. By Definition: Implies, $P \Rightarrow Q$ is false exactly when P is true and Q is false, so $\neg(P \Rightarrow Q)$ is true exactly in that row. $P \wedge \neg Q$ is true exactly when P is true and $\neg Q$ is true, which is the same row. □

Example 9 (When a search claim fails)**.** The negation of "if the list is sorted, then binary search finds the key" is "the list is sorted and binary search does not find the key" (Theorem 6).

Corollary 1 (A conditional without the arrow)**.** *For propositions P and Q, $P \Rightarrow Q \equiv \neg P \vee Q$.*

Proof

Proof. By double negation (Theorem 3), $P \Rightarrow Q \equiv \neg\neg(P \Rightarrow Q)$. By Theorem 6 and Theorem 4, $\neg\neg(P \Rightarrow Q) \equiv \neg(P \wedge \neg Q)$. By De Morgan's law 1 (Theorem 5), $\neg(P \wedge \neg Q) \equiv \neg P \vee \neg\neg Q$, and by double negation again, that is logically equivalent to $\neg P \vee Q$. □

Example 10 (A rule as an "or")**.** By Corollary 1, "if the tests pass, then the build is deployed" is logically equivalent to "the tests fail or the build is deployed".

Negating quantified propositions

Exercise 6 (Not every test passes)**.** Let T be the set of tests in a test suite, and let $P(t)$ be "test t passes". The report "every test passes" is $\forall t \in T, P(t)$.
Write the negation of this report in English and in symbols, without putting $\neg$ in front of a quantifier.

Solution

Solution 6. "Some test fails": $\exists t \in T, \neg P(t)$. "Every test passes" is false exactly when at least one test does not pass.
"Every test fails", $\forall t \in T, \neg P(t)$, is not the negation: when some tests pass and some fail, both it and "every test passes" are false.

Theorem 7 (Negating a quantifier)**.** *For a set S and a predicate $P(x)$ (Definition: Predicate) whose domain (Definition: Domain (domain of discourse, universe of discourse)) is S:*

1. *$\neg(\forall x \in S, P(x))$ and $\exists x \in S, \neg P(x)$ have the same truth value;*
2. *$\neg(\exists x \in S, P(x))$ and $\forall x \in S, \neg P(x)$ have the same truth value.*

Proof

Proof.

1. By Definition: Universal quantifier, $\forall x \in S, P(x)$ is false exactly when $P(x)$ is false for at least one x in S, which by Definition: Existential quantifier is when $\exists x \in S, \neg P(x)$ is true.
2. By Definition: Existential quantifier, $\exists x \in S, P(x)$ is false exactly when $P(x)$ is false for every x in S, which by Definition: Universal quantifier is when $\forall x \in S, \neg P(x)$ is true.

□

Example 11 (No natural square root of 2)**.** By part 2 of Theorem 7, $\neg(\exists n \in \mathbb{N}, n^2 = 2)$ has the same truth value as $\forall n \in \mathbb{N}, n^2 \neq 2$.

Remark 4 (De Morgan's laws for quantifiers). Theorem 7 is often written $\neg\forall x, P(x) \equiv \exists x, \neg P(x)$ and $\neg\exists x, P(x) \equiv \forall x, \neg P(x)$. These notes say "has the same truth value" instead, because $P(x)$ is a predicate, not a proposition, so Definition 1, which compares truth tables, does not apply.
On a finite set $S = \{a, b\}$, $\forall x \in S, P(x)$ has the same truth value as $P(a) \wedge P(b)$, and $\exists x \in S, P(x)$ has the same truth value as $P(a) \vee P(b)$, so Theorem 7 is De Morgan's laws (Theorem 5) applied to $P(a)$ and $P(b)$.
Racket shows the same pattern on a list: `(not (andmap even? (list 2 4 5)))` and `(ormap odd? (list 2 4 5))` both return `#t`.

Remark 5 (Pushing a negation inward). To negate a proposition built with $\neg$, $\wedge$, $\vee$, $\Rightarrow$ and quantifiers, move the $\neg$ inward one step at a time:

- across a quantifier, with Theorem 7, which swaps $\forall$ and $\exists$;
- across $\wedge$ or $\vee$, with De Morgan's laws (Theorem 5);
- across $\Rightarrow$, with Theorem 6;
- across another $\neg$, with double negation (Theorem 3).

Stop when every $\neg$ sits in front of a proposition with no connective or quantifier inside it. The result is easier to read and to check than a $\neg$ in front of the whole proposition.

Exercise 7 (Completing a palindrome)**.** Let W be the set of words made of the letters a and b, including the empty word, and write st for word s followed by word t. A palindrome is a word that reads the same forwards and backwards, such as abba.
Write the negation of

$$\forall s \in W,\ \exists t \in W,\ st \text{ is a palindrome}$$

without putting $\neg$ in front of a quantifier.

Solution

Solution 7. Apply Theorem 7 twice, from the outside in. First, the negation has the truth value of

$$\exists s \in W,\ \neg(\exists t \in W,\ st \text{ is a palindrome}).$$

For each word s, Theorem 7 says that $\neg(\exists t \in W,\ st$ is a palindrome) has the truth value of $\forall t \in W,\ st$ is not a palindrome. These have the same truth value for every s, so the $\exists s$ propositions built from them do too (Definition: Existential quantifier), and the negation has the truth value of

$$\exists s \in W,\ \forall t \in W,\ st \text{ is not a palindrome.}$$

In English: some word cannot be made into a palindrome by adding letters at its end.

Exercise 8 (Which one is true?)**.** Which is true: $\forall s \in W,\ \exists t \in W,\ st$ is a palindrome, from Exercise 7, or its negation?

Solution

Solution 8. The original is true, so its negation is false. For any word s, let t be s written backwards; then st reads the same backwards. For example, for $s = \text{aab}$, $t = \text{baa}$ and $st = \text{aabbaa}$.

References

Wikipedia contributors. 2026. *De Morgan's Laws — Wikipedia, the Free Encyclopedia.* https://en.wikipedia.org/wiki/De_Morgan%27s_laws.