# Proving Non-Conditional Statements

**If and only if, existence, and disproof**

Last modified: 2026-10-30 13:24:08 (PDT)

**Source**

Adapted from Kameron Decker Harris's Fall 2026 lecture notes "Proving Non-Conditional Statements, Disproof", which are typed transcriptions of Scott Wehrwein's Spring 2025 handwritten lecture notes. These come from those notes:

- the outline of an "if and only if" proof;
- the outline of an existence proof that names an element;
- proving that several propositions are equivalent with a cycle of conditionals;
- the constructive example of half of a positive number;
- the non-constructive example of an irrational power of an irrational number;
- disproving a proposition by proving its negation.

The exercises are new.

## Proving if and only if

**Exercise 1** (A sum of squares equal to zero)**.** Let $x$ and $y$ be real numbers. Prove: $x^2 + y^2 = 0$ if and only if $x = 0$ and $y = 0$ (Definition: If and only if).
Hint: prove the two directions one at a time.

Solution

*Solution* 1. **If $x = 0$ and $y = 0$, then $x^2 + y^2 = 0$.** Suppose $x = 0$ and $y = 0$. Then $x^2 + y^2 = 0^2 + 0^2 = 0$.
**If $x^2 + y^2 = 0$, then $x = 0$ and $y = 0$.** Suppose $x^2 + y^2 = 0$. A square of a real number is never negative, so $y^2 \geq 0$, and

$$\begin{aligned} x^2 &= -y^2 && \text{(subtract } y^2 \text{ from both sides of } x^2 + y^2 = 0) \\ &\leq 0. && (y^2 \geq 0) \end{aligned}$$

Also $x^2 \geq 0$, so $x^2 = 0$, and so $x = 0$. The same steps with $x$ and $y$ swapped give $y = 0$.

**Theorem 1** (A biconditional is two conditionals)**.** *For propositions $P$ and $Q$,*

$$(P \Leftrightarrow Q) \equiv ((P \Rightarrow Q) \wedge (Q \Rightarrow P)).$$

Proof

*Proof.* The fifth and last columns of this truth table (Definition: Truth table) match in every row (Definition: If and only if, Definition: Implies, Definition: And):

| $P$ | $Q$ | $P \Rightarrow Q$ | $Q \Rightarrow P$ | $(P \Rightarrow Q) \wedge (Q \Rightarrow P)$ | $P \Leftrightarrow Q$ |
|---|---|---|---|---|---|
| T | T | T | T | T | T |
| T | F | F | T | F | F |
| F | T | T | F | F | F |
| F | F | T | T | T | T |

So the two propositions are logically equivalent (Definition: Logical equivalence). □

**Example 1** (Proving both directions)**.** Claim: let $n$ be an integer. Then $n$ is even if and only if $n + 2$ is even.
By Theorem 1, it is enough to prove two conditionals.

- If $n$ is even, then $n = 2k$ for some integer $k$ (Definition: Even integer), so $n + 2 = 2k + 2 = 2(k + 1)$, and $k + 1$ is an integer (Remark: Facts about integers used without proof). So $n + 2$ is even.
- If $n + 2$ is even, then $n + 2 = 2m$ for some integer $m$, so $n = 2m - 2 = 2(m - 1)$, and $m - 1$ is an integer. So $n$ is even.

*Remark* 1 (The shape of an "if and only if" proof). A proof of "$P$ if and only if $Q$" has two parts:

(⇒) Suppose $P$.
⋮
Therefore $Q$.

($\Leftarrow$) Suppose $Q$.
$\vdots$
Therefore $P$.

Each part is a proof of a conditional, so each can be a direct proof (Definition: Direct proof), a proof by contrapositive (Definition: Proof by contrapositive), or a proof by contradiction (Definition: Proof by contradiction), and the two parts need not use the same method.

**Theorem 2** (A cycle of conditionals)**.** *Let $P_1, P_2, \ldots, P_k$ be propositions. If $P_1 \Rightarrow P_2$, $P_2 \Rightarrow P_3$, ..., $P_{k-1} \Rightarrow P_k$ and $P_k \Rightarrow P_1$ are all true, then $P_1, \ldots, P_k$ all have the same truth value.*

Proof

*Proof.* Suppose one of the propositions, $P_i$, is true. By modus ponens (Theorem: Common rules of inference), $P_i \Rightarrow P_{i+1}$ makes $P_{i+1}$ true, then $P_{i+1} \Rightarrow P_{i+2}$ makes $P_{i+2}$ true, and so on around the cycle, with $P_k \Rightarrow P_1$ leading from $P_k$ back to $P_1$, until every proposition in the list is true. So either all of $P_1, \ldots, P_k$ are true, or none of them is true, and then all of them are false. □

**Example 2** (Three ways to say "odd")**.** Let $n$ be an integer, and consider

- $P_1$: $n$ is odd;
- $P_2$: $n+1$ is even;
- $P_3$: $n-1$ is even.

By Theorem 2, three conditionals show that all three have the same truth value.

- $P_1 \Rightarrow P_2$: if $n = 2k+1$ (Definition: Odd integer), then $n+1 = 2k+2 = 2(k+1)$.
- $P_2 \Rightarrow P_3$: if $n+1 = 2m$, then $n-1 = 2m-2 = 2(m-1)$.
- $P_3 \Rightarrow P_1$: if $n-1 = 2j$, then $n = 2j+1$.

In each step the new multiplier, $k+1$, $m-1$ or $j$, is an integer (Remark: Facts about integers used without proof). Proving all six conditionals between pairs would have taken twice as many steps.

## Proving existence

**Exercise 2** (A solution in integers)**.** Show that there are integers $x$ and $y$ with $5x+3y=1$.

Solution

*Solution* 2. Take $x=2$ and $y=-3$. Then $5x+3y=5\cdot 2+3\cdot(-3)=10-9=1$. So integers with $5x+3y=1$ exist (Definition: Existential quantifier). Other pairs work too, such as $x=-1$ and $y=2$.

**Definition 1** (Constructive proof)**.** A **constructive proof** of a proposition $\exists x\in S,\ P(x)$ (Definition: Existential quantifier) names a particular element of $S$ and shows that $P$ is true of it (Wikipedia contributors 2026). A proposition with several existential quantifiers, such as "there are integers $m$ and $n$ with $m+n=1$", gets a constructive proof that names one element for each quantifier.

**Example 3** (Half of a positive number)**.** Claim: for every positive real number $x$, there is a positive real number $y$ with $y<x$.
Let $x$ be a positive real number. We give a constructive proof of "there is a positive real number $y$ with $y<x$" for this $x$: take $y=\frac{x}{2}$. Then $y>0$, because $x>0$, and

$$\begin{aligned} x-y &= x-\frac{x}{2} && (\text{substitute } y=\tfrac{x}{2})\\ &= \frac{x}{2} && (\text{arithmetic})\\ &> 0, && (x>0) \end{aligned}$$

so $y<x$. The element named, $\frac{x}{2}$, depends on $x$, which a "for every $x$, there is a $y$" statement allows.
The solution to Exercise 2 is also a constructive proof: it names the integers 2 and $-3$.

**Definition 2** (Non-constructive proof)**.** A **non-constructive proof** of a proposition $\exists x\in S,\ P(x)$ shows that it is true without naming any element $x$ of $S$ for which $P(x)$ is true (Wikipedia contributors 2026).

**Example 4** (An irrational power of an irrational number)**.** Claim: there are irrational numbers $x$ and $y$ (Definition: Irrational numbers) for which $x^y$ is rational (Definition: Rational numbers).

The number $\sqrt{2}$ is irrational (Theorem: $\sqrt{2}$ is irrational). Consider $\sqrt{2}^{\sqrt{2}}$, and take two

cases (Definition: Proof by cases).

**Case 1: $\sqrt{2}^{\sqrt{2}}$ is rational.** Then $x = \sqrt{2}$ and $y = \sqrt{2}$ work.

**Case 2: $\sqrt{2}^{\sqrt{2}}$ is irrational.** Then take $x = \sqrt{2}^{\sqrt{2}}$ and $y = \sqrt{2}$, both irrational, and

$$\begin{aligned} x^y &= \left(\sqrt{2}^{\sqrt{2}}\right)^{\sqrt{2}} && \text{(substitute } x \text{ and } y\text{)} \\ &= \sqrt{2}^{\sqrt{2}\cdot\sqrt{2}} && \text{(a power of a power multiplies the exponents)} \\ &= \sqrt{2}^{2} && (\sqrt{2}\cdot\sqrt{2} = 2) \\ &= 2, \end{aligned}$$

which is rational, since $2 = \frac{2}{1}$.

Every number is rational or irrational, so one of the cases holds, and the claim is true. The proof does not say which case holds, so it does not name a pair $x, y$ that works.

## Disproof

**Exercise 3** (Prove or disprove)**.** Prove or disprove: for every real number $x$, if $x^2 > 1$, then $x > 1$.

Solution

*Solution* 3. The proposition is false. Take $x = -2$. Then $x^2 = 4 > 1$, but $x > 1$ is false. So $x = -2$ makes "if $x^2 > 1$, then $x > 1$" false (Definition: Implies), and so it is a counterexample (Definition: Counterexample) to the proposition.

**Definition 3** (Disproof)**.** A **disproof** of a proposition $P$ is a proof (Definition: Proof) of its negation, $\neg P$ (Definition: Not).

**Example 5** (No integer squares to two)**.** Claim to disprove: there is an integer $n$ with $n^2 = 2$.

By Theorem: Negating a quantifier, its negation is "for every integer $n$, $n^2 \neq 2$". We prove the negation by cases (Definition: Proof by cases). Let $n$ be an integer.

- If $n$ is $-1$, $0$ or $1$, then $n^2$ is $1$, $0$ or $1$, which is not $2$.
- Otherwise $n \leq -2$ or $n \geq 2$, so $|n| \geq 2$, and $n^2 = |n|^2 \geq 2^2 = 4$, so $n^2 \neq 2$.

**Exercise 4** (An equation with a solution)**.** Prove or disprove: there are integers $m$ and $n$ with $6m + 4n = 2$.

Solution

*Solution* 4. The proposition is true. Take $m = 1$ and $n = -1$. Then $6m + 4n = 6 - 4 = 2$.

**Exercise 5** (An equation without a solution)**.** Prove or disprove: there are integers $m$ and $n$ with $6m + 4n = 3$.

Solution

*Solution* 5. The proposition is false. Applying Theorem: Negating a quantifier to each of its two quantifiers in turn, its negation is "for all integers $m$ and $n$, $6m + 4n \neq 3$", and we prove that. Let $m$ and $n$ be integers. Then $6m + 4n = 2(3m + 2n)$, and $3m + 2n$ is an integer (Remark: Facts about integers used without proof), so $6m + 4n$ is even (Definition: Even integer). But $3 = 2 \cdot 1 + 1$ is odd (Definition: Odd integer), and no integer is both even and odd (Remark: Facts about integers used without proof), so $6m + 4n \neq 3$.

## References

Wikipedia contributors. 2026. *Constructive Proof — Wikipedia, the Free Encyclopedia.* https://en.wikipedia.org/wiki/Constructive_proof.