

# Algebra

Last modified: 2026-10-03 02:55:01 (PDT)

## Equalities

Mastery of [Elementary Algebra](#) (a.k.a. “College Algebra”) is a prerequisite for calculus, which is in turn a prerequisite for most statistics and data science courses. Nevertheless, each year, some students are still uncomfortable with algebraic manipulations of mathematical formulas. Therefore, I include this section as a quick reference.

**Theorem 0.1** (Equalities are transitive). *If  $a = b$  and  $b = c$ , then  $a = c$*

**Theorem 0.2** (Substituting equivalent expressions). *If  $a = b$ , then for any function  $f(x)$ ,  $f(a) = f(b)$*

## Inequalities

**Theorem 0.3** (Adding to both sides of an inequality). *If  $a < b$ , then  $a + c < b + c$*

**Theorem 0.4** (negating both sides of an inequality). *If  $a < b$ , then:  $-a > -b$*

**Theorem 0.5** (Multiplying both sides of an inequality by a positive number). *If  $a < b$  and  $c > 0$ , then  $ca < cb$ .*

**Theorem 0.6** (Negation is multiplication by  $-1$ ).

$$-a = (-1) * a$$

## Minimum, maximum, argmin and argmax

**Exercise 0.1** (Smallest value, and where it occurs).

1. Let  $f(x) = (x - 2)^2 + 1$  for every  $x \in \mathbb{R}$ . What is the smallest value  $f$  takes, and at which  $x$  does it take that value?
2. Now let  $g(x) = (x - 2)^2 + 1$  only for  $x \in \{0, 1, 3, 4\}$ . What is the smallest value  $g$  takes, and at which  $x$  does it take that value?

Solution

*Solution 0.1.*

1. The square of a real number is never negative, so  $(x - 2)^2 \geq 0$  for every  $x$ , and therefore  $f(x) \geq 0 + 1 = 1$ . The square is 0 only when  $x - 2 = 0$ , that is, when  $x = 2$ , and there  $f(2) = 1$ . So the smallest value of  $f$  is 1, and  $x = 2$  is the only input where  $f$  takes it.
2. Evaluate  $g$  at each of its four inputs:

|        |   |   |   |   |
|--------|---|---|---|---|
| $x$    | 0 | 1 | 3 | 4 |
| $g(x)$ | 5 | 2 | 2 | 5 |

The smallest value of  $g$  is 2, and  $g$  takes it at two inputs,  $x = 1$  and  $x = 3$ . The answer to “where?” is the set  $\{1, 3\}$ , not a single number.

**Definition 0.1** (Minimum). Let  $A \subseteq \mathbb{R}$ . A number  $m$  is the **minimum** of  $A$ , written  $\min A$ , if  $m \in A$  and  $m \leq a$  for all  $a \in A$ .

*Remark 0.1* (Not every set has a minimum). The set  $\{3, 1, 4\}$  has minimum 1, and the interval  $[0, 1]$  has minimum 0. The interval  $(0, 1]$  has no minimum: every element  $a$  of it has a smaller element,  $a/2$ , also in it.

**Definition 0.2** (Maximum). Let  $A \subseteq \mathbb{R}$ . A number  $M$  is the **maximum** of  $A$ , written  $\max A$ , if  $M \in A$  and  $a \leq M$  for all  $a \in A$ .

**Definition 0.3** (Argmin). Let  $f : A \rightarrow \mathbb{R}$  be a [function](#). The **argmin** of  $f$  over  $A$  is the set of inputs where  $f$  takes its smallest value:

$$\arg \min_{x \in A} f(x) \stackrel{\text{def}}{=} \{x \in A : \forall a \in A, f(x) \leq f(a)\}$$

When this set has exactly one element  $\hat{x}$ , we write  $\hat{x} = \arg \min_{x \in A} f(x)$ .

*Remark 0.2* (Smallest value versus where it occurs). The smallest value itself is  $\min f(A)$ , the minimum (Definition 0.1) of the [image](#) of  $f$ , and the argmin is where that value is attained. For example, let  $f(x) = x^2$  on  $A = \{-1, 0, 2\}$ . The image is  $f(A) = \{1, 0, 4\}$ , so the smallest value is  $\min f(A) = 0$ , and  $\arg \min_{x \in A} f(x) = \{0\}$ . The argmin is empty when  $f$  has no smallest value, for example  $f(x) = x$  on  $(0, 1]$ .

**Definition 0.4** (Argmax). Let  $f : A \rightarrow \mathbb{R}$  be a [function](#). The **argmax** of  $f$  over  $A$  is the set of inputs where  $f$  takes its largest value:

$$\arg \max_{x \in A} f(x) \stackrel{\text{def}}{=} \{x \in A : \forall a \in A, f(a) \leq f(x)\}$$

When this set has exactly one element  $\hat{x}$ , we write  $\hat{x} = \arg \max_{x \in A} f(x)$ .

## Global and local minimizers

**Exercise 0.2** (A dip that is not the bottom). Let  $f(x) = x^3 - 3x$  for  $x \in \mathbb{R}$ .

1. Show that  $f(x) - f(1) = (x - 1)^2(x + 2)$ .
2. Use part 1 to show that  $f(x) \geq f(1)$  for every  $x$  in the interval  $(0, 2)$ .
3. Compute  $f(-3)$ . Is  $f(1)$  the smallest value  $f$  takes on all of  $\mathbb{R}$ ?
4. Does  $f$  take a smallest value anywhere on  $\mathbb{R}$ ?

### Solution

*Solution 0.2.*

1. Since  $f(1) = 1 - 3 = -2$ ,

$$\begin{aligned} f(x) - f(1) &= x^3 - 3x + 2 \\ &= (x - 1)(x^2 + x - 2) \\ &= (x - 1)(x - 1)(x + 2) \\ &= (x - 1)^2(x + 2). \end{aligned}$$

2. For  $x \in (0, 2)$ ,  $(x - 1)^2 \geq 0$  and  $x + 2 > 0$ , so their product is at least 0. By part 1,  $f(x) - f(1) \geq 0$ , that is,  $f(x) \geq f(1)$ .
3.  $f(-3) = -27 + 9 = -18$ . Since  $-18 < -2 = f(1)$ ,  $f(1)$  is not the smallest value of  $f$  on  $\mathbb{R}$ .
4. No. For each  $n \geq 2$ ,  $f(-n) = -n^3 + 3n = -n(n^2 - 3) \leq -n$ , so no value of  $f$  is smaller than all the others: for any  $x$ , picking  $n \geq 2$  with  $-n < f(x)$  gives  $f(-n) < f(x)$ .

**Definition 0.5** (Global minimizer). Let  $A \subseteq \mathbb{R}^p$  and let  $f : A \rightarrow \mathbb{R}$  be a [function](#). A point  $x^* \in A$  is a **global minimizer** of  $f$  over  $A$  if  $f(x^*) \leq f(x)$  for all  $x \in A$ .

*Remark 0.3* (Global minimizers form the argmin). The global minimizers of  $f$  are exactly the elements of  $\arg \min_{x \in A} f(x)$  (Definition 0.3). A function can have more than one global minimizer. For example,  $f(x) = (x^2 - 1)^2$  on  $\mathbb{R}$  satisfies  $f(x) \geq 0$  for every  $x$ , and  $f(x) = 0$  exactly when  $x = -1$  or  $x = 1$ , so its global minimizers are  $-1$  and  $1$ , and  $\arg \min_{x \in \mathbb{R}} f(x) = \{-1, 1\}$ .

**Definition 0.6** (Local minimizer). Let  $A \subseteq \mathbb{R}^p$  and let  $f : A \rightarrow \mathbb{R}$  be a [function](#). A point  $x^* \in A$  is a **local minimizer** of  $f$  if there is a number  $\delta > 0$  such that  $f(x^*) \leq f(x)$  for all  $x \in A$  with  $\|x - x^*\| < \delta$ , where  $\|\cdot\|$  is the [Euclidean norm](#).

*Remark 0.4* (Local and global minimizers). For  $p = 1$ ,  $\|x - x^*\| = |x - x^*|$ . For example, with  $x^* = 1$  and  $\delta = 1$ , the condition  $|x - 1| < 1$  means  $0 < x < 2$ . Every global minimizer (Definition 0.5) is a local minimizer: take any  $\delta > 0$ . For example,  $x^* = 0$  is a global minimizer of  $f(x) = x^2$  on  $\mathbb{R}$ , so it is also a local minimizer. The converse fails. In Exercise 0.2,  $x^* = 1$  is a local minimizer of  $f(x) = x^3 - 3x$  (take  $\delta = 1$ ), but not a global one.

## Convex functions

**Exercise 0.3** (A chord above a parabola). Let  $f(x) = (x - 2)^2$ , and take the points  $x = 0$  and  $y = 4$  with  $t = \frac{1}{2}$ .

1. Compute  $f(tx + (1 - t)y)$ .
2. Compute  $tf(x) + (1 - t)f(y)$ .

3. Is  $f(tx + (1-t)y) \leq tf(x) + (1-t)f(y)$ ?

**Solution**

*Solution 0.3.*

1. With  $t = \frac{1}{2}$ ,  $tx + (1-t)y = \frac{1}{2} \cdot 0 + \frac{1}{2} \cdot 4 = 2$ , so  $f(2) = (2-2)^2 = 0$ .
2.  $f(0) = (0-2)^2 = 4$  and  $f(4) = (4-2)^2 = 4$ , so  $\frac{1}{2} \cdot 4 + \frac{1}{2} \cdot 4 = 4$ .
3. Yes:  $0 \leq 4$ . The value of  $f$  at the midpoint of 0 and 4 is below the average of its values at the two endpoints.

**Definition 0.7** (Convex function). Let  $f : \mathbb{R}^p \rightarrow \mathbb{R}$  be a [function](#).  $f$  is **convex** if

$$f(tx + (1-t)y) \leq tf(x) + (1-t)f(y)$$

for all  $x, y \in \mathbb{R}^p$  and all  $t \in [0, 1]$ .

*Remark 0.5* (Chords lie on or above the graph). The point  $tx + (1-t)y$  lies on the line segment from  $x$  to  $y$ , and the right-hand side is the height of the chord joining  $(x, f(x))$  and  $(y, f(y))$  above that point. So  $f$  is convex when every chord lies on or above the graph.

For example, take  $f(x) = x^2$ ,  $x = -1$ ,  $y = 3$ , and  $t = \frac{1}{2}$ . The point is  $\frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 3 = 1$ , where the graph has height  $f(1) = 1$  and the chord has height  $\frac{1}{2}f(-1) + \frac{1}{2}f(3) = \frac{1}{2} \cdot 1 + \frac{1}{2} \cdot 9 = 5$ . The chord is above the graph:  $1 \leq 5$ .

**Example 0.1** (A function that is not convex). The function in Exercise 0.2,  $f(x) = x^3 - 3x$ , is not convex. Take  $x = -2$ ,  $y = 0$ ,  $t = \frac{1}{2}$ . The point is  $\frac{1}{2} \cdot (-2) + \frac{1}{2} \cdot 0 = -1$ , where the graph has height  $f(-1) = -1 + 3 = 2$ , but the chord has height  $\frac{1}{2}f(-2) + \frac{1}{2}f(0) = \frac{1}{2}(-8 + 6) + \frac{1}{2} \cdot 0 = -1$ , and  $2 \leq -1$  is false.

**Example 0.2** (A convex and a non-convex function).  $f(x) = (x-2)^2$  is convex (Definition 0.7): by expanding the square,  $f(tx+(1-t)y) - tf(x) - (1-t)f(y) = -t(1-t)(x-y)^2 \leq 0$ . Its local minimizer  $x^* = 2$  is also a global minimizer, since  $f(x) = (x-2)^2 \geq 0 = f(2)$  for every  $x$ .

Without convexity, a local minimizer need not be global:  $f(x) = x^3 - 3x$  is not convex (Example 0.1), and it has a local minimizer at  $x^* = 1$  that is not global (Exercise 0.2).

**Theorem 0.7** (Local minimizers of convex functions are global). *Let  $f : \mathbb{R}^p \rightarrow \mathbb{R}$  be a convex function. Every local minimizer of  $f$  is a global minimizer of  $f$  over  $\mathbb{R}^p$ .*

### **i** Proof

*Proof.* Let  $x^*$  be a local minimizer of  $f$ , so there is a  $\delta > 0$  with  $f(x^*) \leq f(x)$  whenever  $\|x - x^*\| < \delta$  (Definition 0.6). Suppose  $x^*$  is not a global minimizer (Definition 0.5). Then some  $y \in \mathbb{R}^p$  has  $f(y) < f(x^*)$ , and in particular  $y \neq x^*$ .

Let  $t = \min \left\{ \frac{1}{2}, \frac{\delta}{2\|y - x^*\|} \right\}$ , so  $t \in (0, 1)$ , and let  $z = ty + (1-t)x^*$ . Then  $z - x^* = t(y - x^*)$ , so  $\|z - x^*\| = t\|y - x^*\| \leq \delta/2 < \delta$ .

By convexity,

$$f(z) \leq tf(y) + (1-t)f(x^*) < tf(x^*) + (1-t)f(x^*) = f(x^*),$$

where the strict inequality uses  $t > 0$  and  $f(y) < f(x^*)$ . So  $f(z) < f(x^*)$  with  $\|z - x^*\| < \delta$ , which contradicts  $x^*$  being a local minimizer. Hence  $x^*$  is a global minimizer.  $\square$

## Infimum and supremum

**Definition 0.8** (Infimum (greatest lower bound)). Let  $A \subseteq \mathbb{R}$  be nonempty and bounded below, meaning that some  $t \in \mathbb{R}$  satisfies  $t \leq a$  for all  $a \in A$ . The **infimum** of  $A$ , written  $\inf A$ , is the greatest real number  $t$  satisfying  $t \leq a$  for all  $a \in A$ :

$$\inf A \stackrel{\text{def}}{=} \max\{t \in \mathbb{R} : \forall a \in A, t \leq a\}$$

If  $A$  is nonempty but not bounded below, we write  $\inf A = -\infty$  by convention.

*Remark 0.6* (Existence of the infimum, and when it is a minimum). The maximum in Definition 0.8 always exists: that is the completeness (greatest-lower-bound) property of the real numbers (Rudin 1976, Definition 1.8, p. 4, and Theorem 1.19, p. 8). For example, for  $A = (1, 2]$ , the numbers  $t$  with  $t \leq a$  for all  $a \in A$  are those with  $t \leq 1$ , and the largest of them is 1, so  $\inf A = 1$ .

If the infimum belongs to  $A$ , it equals the minimum:  $\inf A = \min A$ . For example,  $\inf[1, 2] = 1 = \min[1, 2]$ . For  $A = (1, 2]$ , the infimum 1 is not in  $A$ , and  $A$  has no minimum.

**Example 0.3** (Numerical examples of infimum).

- $\inf\{1, 2, 3\} = 1$ , since 1 is the smallest element.

- $\inf(0.5, 1] = 0.5 = \min[0.5, 1]$ : for intervals open below, the infimum equals the minimum of the corresponding closed-below interval, even though  $0.5 \notin (0.5, 1]$ . More generally,  $\inf(c, b] = \min[c, b] = c$  for any  $c < b$ .
- $\inf\{t \geq 0 : t > 0.5\} = 0.5$ , even though 0.5 itself is not in the set.
- $\inf\{-1, -2, -3, \dots\} = -\infty$ , because no real number is less than or equal to every element of that set.

**Definition 0.9** (Supremum (least upper bound)). Let  $A \subseteq \mathbb{R}$  be nonempty and bounded above, meaning that some  $t \in \mathbb{R}$  satisfies  $a \leq t$  for all  $a \in A$ . The **supremum** of  $A$ , written  $\sup A$ , is the smallest real number  $t$  satisfying  $a \leq t$  for all  $a \in A$ :

$$\sup A \stackrel{\text{def}}{=} \min\{t \in \mathbb{R} : \forall a \in A, a \leq t\}$$

If  $A$  is nonempty but not bounded above, we write  $\sup A = +\infty$  by convention.

*Remark 0.7* (Existence of the supremum, and when it is a maximum). The minimum in Definition 0.9 always exists: that is the completeness (least-upper-bound) property of the real numbers (Rudin 1976, Definition 1.8, p. 4, and Theorem 1.19, p. 8). For example, for  $A = [1, 2)$ , the numbers  $t$  with  $a \leq t$  for all  $a \in A$  are those with  $t \geq 2$ , and the smallest of them is 2, so  $\sup A = 2$ .

If the supremum belongs to  $A$ , it equals the maximum:  $\sup A = \max A$ . For example,  $\sup[1, 2] = 2 = \max[1, 2]$ . For  $A = [1, 2)$ , the supremum 2 is not in  $A$ , and  $A$  has no maximum.

**Example 0.4** (Numerical examples of supremum).

- $\sup\{1, 2, 3\} = 3$ , since 3 is the largest element.
- $\sup\{t \geq 0 : t < 0.5\} = 0.5$ , even though 0.5 itself is not in the set.
- $\sup\{1, 2, 3, \dots\} = +\infty$ , because no real number is greater than or equal to every element of that set.

## Sums

**Theorem 0.8** (adding zero changes nothing).

$$a + 0 = a$$

**Theorem 0.9** (Sums are symmetric).

$$a + b = b + a$$

**Theorem 0.10** (Sums are associative). *When adding three numbers, it does not matter which pair you add first:*

$$(a + b) + c = a + (b + c)$$

**Example 0.5** (Grouping a sum two ways).  $(2 + 3) + 4 = 5 + 4 = 9$ , and  $2 + (3 + 4) = 2 + 7 = 9$ .

## Products

**Theorem 0.11** (Multiplying by 1 changes nothing).

$$a \times 1 = a$$

**Theorem 0.12** (Products are symmetric).

$$a \times b = b \times a$$

**Theorem 0.13** (Products are associative).

$$(a \times b) \times c = a \times (b \times c)$$

## Division

**Theorem 0.14** (Division can be written as a product). *If  $b \neq 0$ , then*

$$\frac{a}{b} = a \times \frac{1}{b}$$

## Sums and products together

**Theorem 0.15** (Multiplication is distributive).

$$a(b + c) = ab + ac$$

**Exercise 0.4** (Expand a squared sum). Is  $(3 + 4)^2$  equal to  $3^2 + 4^2$ ? Then expand  $(a + b)^2$  for any numbers  $a$  and  $b$ , using only the distributive law and the rules above.

### Solution

*Solution 0.4.* No:  $(3 + 4)^2 = 7^2 = 49$ , while  $3^2 + 4^2 = 9 + 16 = 25$ . The difference,  $49 - 25 = 24$ , is  $2 \cdot 3 \cdot 4$ .

To see where that extra term comes from, write the square as a product and apply the distributive law (Theorem 0.15) twice:

$$\begin{aligned}(a + b)^2 &= (a + b)(a + b) && \text{(definition of a square)} \\ &= (a + b)a + (a + b)b && \text{(distributive law)} \\ &= (a^2 + ba) + (ab + b^2) && \text{(distributive law, twice)} \\ &= a^2 + ab + ab + b^2 && \text{(commutative and associative laws)} \\ &= a^2 + 2ab + b^2 && \text{(collect like terms)}\end{aligned}$$

The step “commutative and associative laws” uses Theorem 0.12 to write  $ba$  as  $ab$ , and Theorem 0.10 to drop the parentheses.

**Theorem 0.16** (Square of a sum). *For any numbers  $a$  and  $b$ ,*

$$(a + b)^2 = a^2 + 2ab + b^2$$

### Proof

*Proof.* By Solution 0.4. □

*Remark 0.8* (Square of a difference). Replacing  $b$  by  $-b$  in Theorem 0.16 gives  $(a - b)^2 = a^2 - 2ab + b^2$ . Squared errors such as  $(y - \hat{y})^2$  are expanded this way. For example, with  $y = 5$  and  $\hat{y} = 3$ ,  $(5 - 3)^2 = 2^2 = 4$ , and  $5^2 - 2 \cdot 5 \cdot 3 + 3^2 = 25 - 30 + 9 = 4$ .

## Summation notation

**Exercise 0.5** (Expand a sum). The expression  $\sum_{i=1}^4 i^2$  is shorthand for a sum of four terms.

1. Guess which four terms, and add them up.
2. Write  $\frac{1}{N} \sum_{i=1}^N (y_i - \hat{y}_i)^2$  for  $N = 3$  without the  $\sum$  symbol.

Solution

*Solution 0.5.*

1. Replace  $i$  by each of 1, 2, 3, 4 in turn, and add the results:

$$\begin{aligned}\sum_{i=1}^4 i^2 &= 1^2 + 2^2 + 3^2 + 4^2 \\ &= 1 + 4 + 9 + 16 \\ &= 30\end{aligned}$$

2. Replace  $i$  by each of 1, 2, 3 in turn inside the parentheses, add the three squares, and multiply the total by  $\frac{1}{3}$ :

$$\frac{1}{3} \sum_{i=1}^3 (y_i - \hat{y}_i)^2 = \frac{1}{3} [(y_1 - \hat{y}_1)^2 + (y_2 - \hat{y}_2)^2 + (y_3 - \hat{y}_3)^2]$$

**Definition 0.10** (Summation notation). Let  $m$  and  $n$  be integers with  $m \leq n$ , and let  $a_m, a_{m+1}, \dots, a_n$  be numbers. The **sum** of  $a_m$  through  $a_n$  is

$$\sum_{i=m}^n a_i \stackrel{\text{def}}{=} a_m + a_{m+1} + \dots + a_n$$

The variable  $i$  is the **index** of the sum;  $m$  and  $n$  are its **lower** and **upper limits**.

*Remark 0.9* (The index is a placeholder). The name of the index does not change the sum:  $\sum_{i=1}^n a_i$  and  $\sum_{j=1}^n a_j$  are the same number. For example,  $\sum_{i=1}^3 i = 1 + 2 + 3 = 6$  and  $\sum_{j=1}^3 j = 1 + 2 + 3 = 6$ .

**Definition 0.11** (Empty sum). When the upper limit is less than the lower limit ( $n < m$ ), the sum  $\sum_{i=m}^n a_i$  has no terms. By convention, such an **empty sum** equals 0. For example,  $\sum_{i=1}^0 a_i = 0$ .

**Exercise 0.6** (Rearrange a sum). Let  $c$  be a number, and let  $a_1, a_2, a_3$  and  $b_1, b_2, b_3$  be numbers. Using Definition 0.10 and the rules of algebra above, show that:

1.  $\sum_{i=1}^3 c a_i = c \sum_{i=1}^3 a_i$ ;
2.  $\sum_{i=1}^3 (a_i + b_i) = \sum_{i=1}^3 a_i + \sum_{i=1}^3 b_i$ .

Does either argument depend on there being exactly three terms?

### Solution

*Solution 0.6.* Each step below applies Definition 0.10, the distributive law (Theorem 0.15), or the commutative and associative laws of addition (Theorem 0.9 and Theorem 0.10).

1. Expand the sum, then factor out  $c$ :

$$\begin{aligned} \sum_{i=1}^3 c a_i &= c a_1 + c a_2 + c a_3 && \text{(expand the sum)} \\ &= c (a_1 + a_2 + a_3) && \text{(distributive law)} \\ &= c \sum_{i=1}^3 a_i && \text{(collect the sum)} \end{aligned}$$

2. Expand the sum, then regroup the terms:

$$\begin{aligned} \sum_{i=1}^3 (a_i + b_i) &= (a_1 + b_1) + (a_2 + b_2) + (a_3 + b_3) && \text{(expand the sum)} \\ &= (a_1 + a_2 + a_3) + (b_1 + b_2 + b_3) && \text{(commutative and associative laws)} \\ &= \sum_{i=1}^3 a_i + \sum_{i=1}^3 b_i && \text{(collect the sum)} \end{aligned}$$

Neither argument uses the number of terms: the same steps work for any lower and upper limits.

**Theorem 0.17** (A constant factor comes out of a sum). *For any number  $c$  and numbers  $a_m, \dots, a_n$ ,*

$$\sum_{i=m}^n c a_i = c \sum_{i=m}^n a_i$$

**i** Proof

*Proof.* By Solution 0.6, part 1. □

**Theorem 0.18** (A sum of sums splits). *For any numbers  $a_m, \dots, a_n$  and  $b_m, \dots, b_n$ ,*

$$\sum_{i=m}^n (a_i + b_i) = \sum_{i=m}^n a_i + \sum_{i=m}^n b_i$$

**i** Proof

*Proof.* By Solution 0.6, part 2. □

**Exercise 0.7** (Sum over a set). Let  $A = \{-1, 2, 5\}$  and let  $f(x) = x^2$ . The expression  $\sum_{x \in A} f(x)$  means: add up  $f(x)$  for each element  $x$  of  $A$ .

1. Write  $\sum_{x \in A} f(x)$  without the  $\sum$  symbol, and compute its value.
2. List the elements of  $A$  in a different order, and add up the same terms in that order. Is the total the same?

Solution

*Solution 0.7.*

1. Replace  $x$  by each element of  $A$  in turn, and add the results:

$$\begin{aligned} \sum_{x \in A} f(x) &= f(-1) + f(2) + f(5) \\ &= (-1)^2 + 2^2 + 5^2 \\ &= 1 + 4 + 25 \\ &= 30 \end{aligned}$$

2. In the order  $5, -1, 2$ , the terms are 25, 1 and 4. Their total is  $25 + 1 + 4 = 30$ . The total is the same, because addition does not depend on the order of the terms.

**Definition 0.12** (Sum over a finite set). Let  $A = \{x_1, x_2, \dots, x_k\}$  be a finite **set** with  $k \geq 1$  different elements. Let  $f$  be a **function** that gives a number  $f(x)$  for each element  $x$  of  $A$ . The **sum of  $f$  over  $A$**  is

$$\sum_{x \in A} f(x) \stackrel{\text{def}}{=} f(x_1) + f(x_2) + \dots + f(x_k)$$

If  $A$  has no elements, the sum has no terms, and by convention it equals 0. For example,  $\sum_{x \in \{\}} x^2 = 0$ .

*Remark 0.10* (The order of the terms does not matter). The order in which we list the elements of  $A$  does not change the sum. The reason is that addition is commutative and associative: we can reorder and regroup the terms of a finite sum without changing the total. For example, if  $A = \{1, 2, 3\}$  and  $f(x) = x^2$ , listing  $A$  as 1, 2, 3 gives  $1 + 4 + 9 = 14$ , and listing  $A$  as 3, 1, 2 gives  $9 + 1 + 4 = 14$ . Exercise 0.7, part 2, shows another example. When  $A = \{m, m+1, \dots, n\}$ , this sum is the same as  $\sum_{i=m}^n f(i)$  from Definition 0.10.

*Remark 0.11* (Leaving the set out). Some authors leave the set out and write  $\sum_x f(x)$ . This shorthand means the sum over every value  $x$  can take. For example, if  $x$  is the outcome of one roll of a six-sided die,  $\sum_x f(x)$  means  $\sum_{x \in \{1, 2, 3, 4, 5, 6\}} f(x)$ . In these notes, we write the set out in full.

## Quotients

**Definition 0.13** (Quotient). For real numbers  $a$  and  $b$  with  $b \neq 0$ , the **quotient** of  $a$  by  $b$  is the result of dividing  $a$  by  $b$ :

$$\frac{a}{b}$$

**Definition 0.14** (Fraction, numerator, and denominator). A quotient  $\frac{a}{b}$  (Definition 0.13) is also called a **fraction**;  $a$  is its **numerator** and  $b$  its **denominator**. For example, the fraction  $\frac{6}{4}$  has numerator 6 and denominator 4.

**Example 0.6** (A quotient). The quotient of 6 by 4 is  $\frac{6}{4} = 1.5$ . The quotient of 6 by 0 is undefined, because Definition 0.13 requires a nonzero denominator.

**Definition 0.15** (Rate). A **rate** is a quotient of two quantities, usually with a denominator that measures time, such as weeks or person-years of follow-up. For example, 12 new cases in 4 weeks is a rate of  $\frac{12}{4} = 3$  new cases per week, and 30 cases over 10,000 person-years of follow-up is a rate of  $\frac{30}{10,000} = 0.003$  cases per person-year.

cf. [https://en.wikipedia.org/wiki/Rate\\_\(mathematics\)](https://en.wikipedia.org/wiki/Rate_(mathematics))

**Definition 0.16** (Ratios). A **ratio** is a quotient in which the numerator and denominator are measured using the same unit scales.

cf. <https://en.wikipedia.org/wiki/Ratio>

**Definition 0.17** (Proportion). In statistics, a **proportion** typically means a ratio where the numerator represents a subset of the denominator.

See [https://en.wikipedia.org/wiki/Population\\_proportion](https://en.wikipedia.org/wiki/Population_proportion).

See also [https://en.wikipedia.org/wiki/Proportion\\_\(mathematics\)](https://en.wikipedia.org/wiki/Proportion_(mathematics)) for other meanings.

**Definition 0.18** (Proportional). Two functions  $f(x)$  and  $g(x)$  are **proportional** if their ratio  $\frac{f(x)}{g(x)}$  does not depend on  $x$ . (cf. [https://en.wikipedia.org/wiki/Proportionality\\_\(mathematics\)](https://en.wikipedia.org/wiki/Proportionality_(mathematics)))

Additional reference for elementary algebra: [https://en.wikipedia.org/wiki/Population\\_proportion#Mathematical\\_definition](https://en.wikipedia.org/wiki/Population_proportion#Mathematical_definition)

## Polynomials

**Exercise 0.8** (Reading a polynomial). Let  $f(x) = 3x^4 - x + 7$ .

1. Write  $f$  in the form  $a_n x^n + a_{n-1} x^{n-1} + \cdots + a_1 x + a_0$ , listing every coefficient  $a_4, a_3, a_2, a_1, a_0$ .
2. What is the degree of  $f$ , and what is its leading coefficient?

Solution

*Solution 0.8.*

1.  $f(x) = 3x^4 + 0 \cdot x^3 + 0 \cdot x^2 + (-1) \cdot x + 7$ , so  $a_4 = 3$ ,  $a_3 = 0$ ,  $a_2 = 0$ ,  $a_1 = -1$ , and  $a_0 = 7$ . Powers of  $x$  that do not appear have coefficient 0.
2. The highest power with a nonzero coefficient is  $x^4$ , so the degree is 4 and the leading

coefficient is  $a_4 = 3$ .

**Definition 0.19** (Polynomial). A **polynomial** (in one variable  $x$ ) is a **function**  $f : \mathbb{R} \rightarrow \mathbb{R}$  that can be written as

$$f(x) = a_n x^n + a_{n-1} x^{n-1} + \cdots + a_1 x + a_0$$

for some integer  $n \geq 0$  and constants  $a_0, a_1, \dots, a_n \in \mathbb{R}$  with  $a_n \neq 0$ .

*Remark 0.12* (Constant and zero polynomials). A constant function  $f(x) = 7$  is a polynomial with  $n = 0$  and  $a_0 = 7$ . The requirement  $a_n \neq 0$  means this definition covers nonzero polynomials only: the zero function  $f(x) = 0$  is excluded here, because it has no nonzero coefficient to serve as  $a_n$ .

**Definition 0.20** (Degree of a polynomial). Let  $f(x) = a_n x^n + \cdots + a_1 x + a_0$  be a **polynomial** with  $a_n \neq 0$ . The integer  $n$  is the **degree** of  $f$ .

**Example 0.7** (Degrees of some polynomials).

- $f(x) = 4x$  has degree 1.
- $f(x) = 5 - 2x^2 + x^3$  has degree 3: the degree is the highest power with a nonzero coefficient, not the power in the first term written.
- A constant polynomial  $f(x) = 7$  has degree 0.

**Definition 0.21** (Leading coefficient). Let  $f(x) = a_n x^n + \cdots + a_1 x + a_0$  be a **polynomial** of **degree**  $n$ . The constant  $a_n$  is the **leading coefficient** of  $f$ .

**Example 0.8** (Leading coefficients).

- $f(x) = 5 - 2x^2 + x^3$  has degree 3, so its leading coefficient is  $a_3 = 1$ , not the 5 written first.
- $f(x) = 3 - x^2$  has leading coefficient  $a_2 = -1$ .

## Exponentials and Logarithms

In these notes,  $\log\{x\}$  is the natural logarithm of  $x > 0$ , the logarithm with base  $e \approx 2.718$ , and  $\exp\{x\} = e^x$  is the exponential function. Some sources write  $\ln x$  for the natural logarithm

and reserve  $\log x$  for base 10.

**Theorem 0.19** ( $\exp\{\}$  and  $\log\{\}$  are mutual inverses).

1. For every  $a > 0$ :  $\exp\{\log\{a\}\} = a$ .
2. For every  $a \in \mathbb{R}$ :  $\log\{\exp\{a\}\} = a$ .

**Theorem 0.20** (Logarithm of a product). If  $a > 0$  and  $b > 0$ , then

$$\log\{a \cdot b\} = \log\{a\} + \log\{b\}$$

**Corollary 0.1** (Logarithm of a quotient). If  $a > 0$  and  $b > 0$ , then

$$\log\left\{\frac{a}{b}\right\} = \log\{a\} - \log\{b\}$$

### Proof

*Proof.* Since  $a > 0$  and  $b > 0$ , the quotient  $\frac{a}{b}$  is positive, so Theorem 0.20 applies to the product  $\frac{a}{b} \cdot b$ :

$$\begin{aligned} \log\{a\} &= \log\left\{\frac{a}{b} \cdot b\right\} && (a = \frac{a}{b} \cdot b) \\ &= \log\left\{\frac{a}{b}\right\} + \log\{b\} && (\text{logarithm of a product}) \end{aligned}$$

The second step applies Theorem 0.20. Subtracting  $\log\{b\}$  from both sides gives  $\log\{a\} - \log\{b\} = \log\{\frac{a}{b}\}$ . □

**Theorem 0.21** (Logarithm of a power). If  $a > 0$  and  $b \in \mathbb{R}$ , then

$$\log\{a^b\} = b \cdot \log\{a\}$$

**Theorem 0.22** (exponential of a sum). The exponential of a sum is equal to the product of the exponentials of the addends:

$$\exp\{a + b\} = \exp\{a\} \cdot \exp\{b\}$$

**Example 0.9** (Exponential of a sum). With  $a = 2$  and  $b = 3$ ,  $\exp\{2 + 3\} = \exp\{5\} \approx 148.41$ , and  $\exp\{2\} \cdot \exp\{3\} \approx 7.3891 \cdot 20.0855 \approx 148.41$ .

**Corollary 0.2** (exponential of a difference). *The exponential of a difference is the exponential of the first term divided by the exponential of the second term:*

$$\exp\{a - b\} = \frac{\exp\{a\}}{\exp\{b\}}$$

**Example 0.10** (Exponential of a difference). With  $a = 5$  and  $b = 2$ ,  $\exp\{5 - 2\} = \exp\{3\} \approx 20.09$ , and  $\frac{\exp\{5\}}{\exp\{2\}} \approx \frac{148.4132}{7.3891} \approx 20.09$ .

**Theorem 0.23** (Power of a power). *If  $a > 0$  and  $b, c \in \mathbb{R}$ , then*

$$a^{bc} = (a^b)^c = (a^c)^b$$

**Example 0.11** (A negative base). With  $a = -1$ ,  $b = 2$ , and  $c = \frac{1}{2}$ :

$$\begin{aligned} a^{bc} &= (-1)^{2 \cdot \frac{1}{2}} \\ &= (-1)^1 \\ &= -1 \end{aligned}$$

but

$$\begin{aligned} (a^b)^c &= ((-1)^2)^{\frac{1}{2}} \\ &= 1^{\frac{1}{2}} \\ &= 1 \end{aligned}$$

So  $a^{bc} \neq (a^b)^c$  here, which is why Theorem 0.23 requires  $a > 0$ . The third expression,  $(a^c)^b = ((-1)^{\frac{1}{2}})^2$ , is not even a real number.

**Corollary 0.3** (natural exponential of a product).

$$\exp\{ab\} = (\exp\{a\})^b = (\exp\{b\})^a$$

**Exercise 0.9.** For  $b, c \in \mathbb{R}$ , when does  $b^c = bc$ ?

Solution

*Solution 0.9.* We only count a pair  $(b, c)$  when  $b^c$  is a real number, so for  $b < 0$  we only consider integer  $c$  ( $\mathbb{R}$  agrees:  $(-8)^{(1/3)}$  is NaN). With that convention,  $bc = b^c$  in each

of the following cases:

1.  $c = 1$ , for every  $b$ .
2.  $b = 0$  and  $c > 0$ , since then  $b^c = 0 = bc$ . ( $b = 0$  and  $c = 0$  fails, since  $0^0 = 1 \neq 0$ .)
3.  $b > 0$ ,  $c > 0$ ,  $c \neq 1$ , and  $b = \exp\left\{\frac{\log\{c\}}{c-1}\right\}$ . For  $b > 0$ , dividing both sides of  $b^c = bc$  by  $b$  gives  $b^{c-1} = c$ , which needs  $c > 0$  because  $b^{c-1} > 0$ ; taking logarithms then gives  $(c-1)\log\{b\} = \log\{c\}$ . For example,  $c = 2$  gives  $b = 2$ , and indeed  $2^2 = 4 = 2 \cdot 2$ .
4.  $b < 0$  and  $c$  is an odd integer with  $c \geq 3$ , with  $b = -c^{1/(c-1)}$ ; for example,  $b = -\sqrt{3}$  and  $c = 3$  give  $b^c = -3\sqrt{3} = bc$ .
5.  $b < 0$  and  $c$  is an even integer with  $c \leq -2$ , with  $b = -(-c)^{1/(c-1)}$ ; for example,  $b = -2^{-1/3}$  and  $c = -2$  give  $b^c = 2^{2/3} = bc$ .

For  $b < 0$ , cases 4 and 5 come from  $b^{c-1} = c$  as well: when  $c-1$  is even,  $b^{c-1} > 0$ , so  $c$  must be positive; when  $c-1$  is odd,  $b^{c-1} < 0$ , so  $c$  must be negative.

See the red contours in Figure 2 for a visualization of the  $b \geq 0$  cases.

*(This 3D surface is interactive; see the HTML or slide version of this page to view it.)*

Figure 1: Graph of  $b * c$  and  $b^c$

*(This 3D surface is interactive; see the HTML or slide version of this page to view it.)*

Figure 2: **Graph of  $b^c - b * c$ .** Red contour lines show where  $b^c = b * c$ .

**Exercise 0.10.** For  $a \geq 0$ ,  $b, c \in \mathbb{R}$ , when does  $(a^b)^c = a^{(b^c)}$ ?

### Solution

*Solution 0.10.* Short answer: rarely (that's all you need to know for this course).

Long answer:

Split on whether  $a > 0$  or  $a = 0$ , because the logarithm we use for  $a > 0$  is undefined at  $a = 0$ .

**Case  $a > 0$ .** By Theorem 0.23,  $(a^b)^c = a^{bc}$ , so the question becomes when  $a^{bc} = a^{(b^c)}$  (for pairs  $(b, c)$  where  $b^c$  is defined). Because  $a > 0$ , both sides are positive, and we can take logarithms (Theorem 0.21):

$$\begin{aligned} a^{bc} &= a^{(b^c)} \\ \log\{a^{bc}\} &= \log\{a^{(b^c)}\} \quad (\text{take logarithms of both sides}) \\ bc \cdot \log\{a\} &= b^c \cdot \log\{a\} \quad (\text{logarithm of a power}) \end{aligned} \tag{1}$$

The last line of Equation 1 holds exactly when

1.  $a = 1$  (so that  $\log\{a\} = 0$ ), or
2.  $bc = b^c$  (see Exercise 0.9).

**Case  $a = 0$ .** Here we cannot take logarithms, so we work from the values of powers of 0:  $0^s = 0$  for  $s > 0$ ,  $0^0 = 1$ , and  $0^s$  is undefined for  $s < 0$ .

- If  $b < 0$ , then  $0^b$  is undefined, so  $(a^b)^c$  is undefined.
- If  $b > 0$ , then  $(0^b)^c = 0^c$  and  $b^c > 0$ , so  $0^{(b^c)} = 0$ ; the two sides agree exactly when  $c > 0$ .
- If  $b = 0$ , then  $(0^0)^c = 1^c = 1$ ; for  $c > 0$ ,  $0^{(0^c)} = 0^0 = 1$ , so the two sides agree, and for  $c \leq 0$  they do not.

So for  $a = 0$ ,  $(a^b)^c = a^{(b^c)}$  exactly when  $b \geq 0$  and  $c > 0$ .

In particular, when  $a = 0$ ,  $b \geq 0$ , and  $c = 0$ , the two sides differ:

$$\begin{aligned}(a^b)^c &= (0^b)^0 \\ &= 1 \\ a^{(b^c)} &= 0^{(b^0)} \\ &= 0^1 \\ &= 0\end{aligned}$$

Rudin, Walter. 1976. *Principles of Mathematical Analysis*. 3rd ed. International Series in Pure and Applied Mathematics. McGraw-Hill.