Direct Sums and Orthogonal Complements

Last modified: 2026-10-16 16:23:18 (PDT)

Sums and direct sums

> **Source**
>
> This section, Section and Section are adapted from Zhou (2024), used under the MIT License (see the license text in Section 2 in Subspaces and Rank). The source proves that a subspace and its orthogonal complement together make up $\mathbb{R}^p$ by extending an orthonormal basis; the proof here uses the rank-nullity theorem (Theorem 8 in Column Space, Null Space and Rank-Nullity) instead.

Definition 0.1 (Sum of subspaces)**.** The **sum** of two subspaces $\mathcal{S}_1$ and $\mathcal{S}_2$ of $\mathbb{R}^p$ (Definition 4 in Subspaces and Rank) is the set of all sums of a vector from $\mathcal{S}_1$ and a vector from $\mathcal{S}_2$:

$$\mathcal{S}_1 + \mathcal{S}_2 \stackrel{\text{def}}{=} \left\{ \underset{\sim 1}{x} + \underset{\sim 2}{x} : \underset{\sim 1}{x} \in \mathcal{S}_1,\ \underset{\sim 2}{x} \in \mathcal{S}_2 \right\}.$$

Example 0.1 (Two lines sum to a plane)**.** Let $\mathcal{S}_1 = \text{span}\{(1,0,0)\}$ and $\mathcal{S}_2 = \text{span}\{(0,1,0)\}$, two lines in $\mathbb{R}^3$. A vector of $\mathcal{S}_1 + \mathcal{S}_2$ has the form $a\,(1,0,0) + b\,(0,1,0) = (a,b,0)$, and a and b can be any numbers, so $\mathcal{S}_1 + \mathcal{S}_2 = \{(a,b,0) : a,b \in \mathbb{R}\}$, the plane $z = 0$.

Example 0.2 (The sum is not the union)**.** For the two lines in Example 0.1, the union $\mathcal{S}_1 \cup \mathcal{S}_2$ contains $(1,0,0)$ and $(0,1,0)$ but not their sum $(1,1,0)$: every multiple of $(1,0,0)$ has second entry 0, and every multiple of $(0,1,0)$ has first entry 0. So the union is not closed under addition and is not a subspace, while the sum, the whole plane $z = 0$, is.

Theorem 0.1 (Sums and intersections of subspaces are subspaces)**.** *If $\mathcal{S}_1$ and $\mathcal{S}_2$ are subspaces of $\mathbb{R}^p$ (Definition 4 in Subspaces and Rank), then $\mathcal{S}_1 + \mathcal{S}_2$ (Definition 0.1) and $\mathcal{S}_1 \cap \mathcal{S}_2$ are subspaces of $\mathbb{R}^p$.*

Proof

Proof. Both $\mathcal{S}_1$ and $\mathcal{S}_2$ contain $\underset{\sim}{0}$ (Theorem 1 in Subspaces and Rank).
The sum. It is not empty: it contains $\underset{\sim}{0} + \underset{\sim}{0} = \underset{\sim}{0}$. Take two of its vectors, $\underset{\sim}{u} = \underset{\sim 1}{u} + \underset{\sim 2}{u}$ and $\underset{\sim}{w} = \underset{\sim 1}{w} + \underset{\sim 2}{w}$, with $\underset{\sim 1}{u}, \underset{\sim 1}{w} \in \mathcal{S}_1$ and $\underset{\sim 2}{u}, \underset{\sim 2}{w} \in \mathcal{S}_2$, and a number c. Then

$$\begin{aligned} \underset{\sim}{u} + \underset{\sim}{w} &= (\underset{\sim 1}{u} + \underset{\sim 2}{u}) + (\underset{\sim 1}{w} + \underset{\sim 2}{w}) && \text{(substitute)} \\ &= (\underset{\sim 1}{u} + \underset{\sim 1}{w}) + (\underset{\sim 2}{u} + \underset{\sim 2}{w}), && \text{(regroup; vector addition is entrywise)} \end{aligned}$$

where $\underset{\sim 1}{u} + \underset{\sim 1}{w} \in \mathcal{S}_1$ and $\underset{\sim 2}{u} + \underset{\sim 2}{w} \in \mathcal{S}_2$, because each subspace is closed under addition. Likewise

$$\begin{aligned} c\,\underset{\sim}{u} &= c\,(\underset{\sim 1}{u} + \underset{\sim 2}{u}) && \text{(substitute)} \\ &= c\,\underset{\sim 1}{u} + c\,\underset{\sim 2}{u}, && \text{(distribute } c \text{ entrywise)} \end{aligned}$$

where $c\,\underset{\sim 1}{u} \in \mathcal{S}_1$ and $c\,\underset{\sim 2}{u} \in \mathcal{S}_2$, because each subspace is closed under scalar multiplication. So $\underset{\sim}{u} + \underset{\sim}{w}$ and $c\,\underset{\sim}{u}$ are in $\mathcal{S}_1 + \mathcal{S}_2$.
The intersection. It is not empty: it contains $\underset{\sim}{0}$. If $\underset{\sim}{u}$ and $\underset{\sim}{w}$ are in both $\mathcal{S}_1$ and $\mathcal{S}_2$, then $\underset{\sim}{u} + \underset{\sim}{w}$ and $c\,\underset{\sim}{u}$ are in $\mathcal{S}_1$, because $\mathcal{S}_1$ is a subspace, and in $\mathcal{S}_2$ for the same reason, so they are in $\mathcal{S}_1 \cap \mathcal{S}_2$. □

Example 0.3 (Two planes in $\mathbb{R}^3$)**.** Let $\mathcal{S}_1 = \{(a, b, 0) : a, b \in \mathbb{R}\}$, the plane $z = 0$, and $\mathcal{S}_2 = \{(0, b, c) : b, c \in \mathbb{R}\}$, the plane $x = 0$.

- **Intersection:** a vector in both has third entry 0 and first entry 0, so $\mathcal{S}_1 \cap \mathcal{S}_2 = \{(0, b, 0) : b \in \mathbb{R}\}$, the y-axis, which is a line through the origin.
- **Sum:** $(a, b, 0) + (0, b', c) = (a, b + b', c)$, and any (x, y, z) arises this way, with $a = x$, $b = y$, $b' = 0$ and $c = z$. So $\mathcal{S}_1 + \mathcal{S}_2 = \mathbb{R}^3$.

Theorem 0.2 (The dimension of a sum of subspaces)**.** *If $\mathcal{S}_1$ and $\mathcal{S}_2$ are subspaces of $\mathbb{R}^p$, then*

$$\dim(\mathcal{S}_1 + \mathcal{S}_2) = \dim(\mathcal{S}_1) + \dim(\mathcal{S}_2) - \dim(\mathcal{S}_1 \cap \mathcal{S}_2).$$

Proof

Proof. All three sets in the formula are subspaces (Theorem 0.1), so each has a dimension (Definition 9 in Subspaces and Rank). Let $k = \dim(\mathcal{S}_1 \cap \mathcal{S}_2)$, $d_1 = \dim(\mathcal{S}_1)$ and $d_2 = \dim(\mathcal{S}_2)$.

Build a list. Choose a basis $\underset{\sim}{z}_1, \ldots, \underset{\sim}{z}_k$ of $\mathcal{S}_1 \cap \mathcal{S}_2$ (Theorem 9 in Subspaces and Rank). These vectors are linearly independent and lie in $\mathcal{S}_1$, so Theorem 9 in Subspaces and Rank extends them to a basis $\underset{\sim}{z}_1, \ldots, \underset{\sim}{z}_k, \underset{\sim}{x}_1, \ldots, \underset{\sim}{x}_{d_1-k}$ of $\mathcal{S}_1$, which has d_1 vectors (Theorem 7 in Subspaces and Rank). In the same way, extend them to a basis $\underset{\sim}{z}_1, \ldots, \underset{\sim}{z}_k, \underset{\sim}{y}_1, \ldots, \underset{\sim}{y}_{d_2-k}$ of $\mathcal{S}_2$. We show that the $\underset{\sim}{z}$'s, $\underset{\sim}{x}$'s and $\underset{\sim}{y}$'s together, $k + (d_1 - k) + (d_2 - k) = d_1 + d_2 - k$ vectors, are a basis of $\mathcal{S}_1 + \mathcal{S}_2$ (Definition 6 in Subspaces and Rank).

They span $\mathcal{S}_1 + \mathcal{S}_2$. Any $\underset{\sim}{v} \in \mathcal{S}_1 + \mathcal{S}_2$ is $\underset{\sim}{v}_1 + \underset{\sim}{v}_2$ with $\underset{\sim}{v}_1 \in \mathcal{S}_1$ and $\underset{\sim}{v}_2 \in \mathcal{S}_2$. $\underset{\sim}{v}_1$ is a linear combination of the $\underset{\sim}{z}$'s and $\underset{\sim}{x}$'s, and $\underset{\sim}{v}_2$ is a linear combination of the $\underset{\sim}{z}$'s and $\underset{\sim}{y}$'s, so their sum is a linear combination of all three groups. Conversely, each vector in the list is in $\mathcal{S}_1 + \mathcal{S}_2$ (for example, $\underset{\sim}{x}_1 = \underset{\sim}{x}_1 + \underset{\sim}{0}$), and $\mathcal{S}_1 + \mathcal{S}_2$ is a subspace, so it contains every linear combination of them.

They are linearly independent. Suppose

$$\sum_{i=1}^{k} a_i \underset{\sim}{z}_i + \sum_{j=1}^{d_1-k} b_j \underset{\sim}{x}_j + \sum_{l=1}^{d_2-k} c_l \underset{\sim}{y}_l = \underset{\sim}{0}.$$

Let $\underset{\sim}{w} \stackrel{\text{def}}{=} \sum_l c_l \underset{\sim}{y}_l$. Then $\underset{\sim}{w} \in \mathcal{S}_2$, because it is a linear combination of vectors of $\mathcal{S}_2$. Subtracting $\sum_i a_i \underset{\sim}{z}_i + \sum_j b_j \underset{\sim}{x}_j$ from both sides of the supposed equation gives

$$\underset{\sim}{w} = \sum_{i=1}^{k} (-a_i) \underset{\sim}{z}_i + \sum_{j=1}^{d_1-k} (-b_j) \underset{\sim}{x}_j,$$

a linear combination of vectors of $\mathcal{S}_1$, so $\underset{\sim}{w} \in \mathcal{S}_1$ too. So $\underset{\sim}{w} \in \mathcal{S}_1 \cap \mathcal{S}_2$, and $\underset{\sim}{w} = \sum_i e_i \underset{\sim}{z}_i$ for some numbers e_i, since the $\underset{\sim}{z}$'s span the intersection. Then

$$\begin{aligned}
\underset{\sim}{0} &= \underset{\sim}{w} - \underset{\sim}{w} && \text{(a vector minus itself)} \\
&= \sum_{i=1}^{k} e_i \underset{\sim}{z}_i - \sum_{l=1}^{d_2-k} c_l \underset{\sim}{y}_l && \text{(substitute the two expressions for } \underset{\sim}{w}\text{)} \\
&= \sum_{i=1}^{k} e_i \underset{\sim}{z}_i + \sum_{l=1}^{d_2-k} (-c_l) \underset{\sim}{y}_l, && \text{(write each subtraction as adding } -1 \text{ times)}
\end{aligned}$$

and because the $\underset{\sim}{z}$'s and $\underset{\sim}{y}$'s are a basis of $\mathcal{S}_2$, every $c_l = 0$. The supposed equation then

reads $\sum_i a_i \underset{\sim}{z}_i + \sum_j b_j \underset{\sim}{x}_j = \underset{\sim}{0}$, and because the $\underset{\sim}{z}$'s and $\underset{\sim}{x}$'s are a basis of $\mathcal{S}_1$, every a_i and b_j is 0.
So $\dim(\mathcal{S}_1 + \mathcal{S}_2) = d_1 + d_2 - k$. When some of the groups are empty (for example $k = 0$, or $\mathcal{S}_1 = \mathcal{S}_1 \cap \mathcal{S}_2$), the sums over them are $\underset{\sim}{0}$ and the argument goes through unchanged. □

Example 0.4 (Two planes in $\mathbb{R}^3$, by dimension)**.** For the planes $z = 0$ and $x = 0$ in Example 0.3, each has dimension 2: the plane $z = 0$ has basis $(1,0,0), (0,1,0)$ (Example 16 in Subspaces and Rank), and the plane $x = 0$ has basis $(0,1,0), (0,0,1)$ by the same argument. Their intersection, the y-axis $\text{span}\{(0,1,0)\}$, has dimension 1, because the single nonzero vector $(0,1,0)$ is linearly independent ($c\,(0,1,0) = (0,c,0)$ is $\underset{\sim}{0}$ only if $c = 0$) and so is a basis of it. Theorem 0.2 gives

$$\begin{aligned} \dim(\mathcal{S}_1 + \mathcal{S}_2) &= 2 + 2 - 1 \\ &= 3, \end{aligned}$$

which agrees with $\mathcal{S}_1 + \mathcal{S}_2 = \mathbb{R}^3$ (Example 0.3) and $\dim(\mathbb{R}^3) = 3$ (Example 16 in Subspaces and Rank).

Corollary 0.1 (The dimension of a sum is at most the sum of the dimensions)**.** *If $\mathcal{S}_1$ and $\mathcal{S}_2$ are subspaces of $\mathbb{R}^p$, then*

$$\dim(\mathcal{S}_1 + \mathcal{S}_2) \leq \dim(\mathcal{S}_1) + \dim(\mathcal{S}_2),$$

with equality exactly when $\mathcal{S}_1 \cap \mathcal{S}_2 = \left\{\underset{\sim}{0}\right\}$.

Proof

Proof. By Theorem 0.2, the two sides differ by $\dim(\mathcal{S}_1 \cap \mathcal{S}_2) \geq 0$. That dimension is 0 exactly when a basis of the intersection is the empty list, that is, when the intersection is the span of the empty list, $\left\{\underset{\sim}{0}\right\}$ (Definition 5 in Subspaces and Rank, Example 16 in Subspaces and Rank). □

Example 0.5 (Equality and strict inequality)**.**

- For the lines $\mathcal{S}_1 = \text{span}\{(1,0,0)\}$ and $\mathcal{S}_2 = \text{span}\{(0,1,0)\}$ in Example 0.1, a common vector satisfies $a\,(1,0,0) = b\,(0,1,0)$, so $(a,-b,0) = \underset{\sim}{0}$ and

$$\begin{aligned} a &= b \\ &= 0. \end{aligned}$$

The intersection is $\left\{\underset{\sim}{0}\right\}$, and indeed

$$\begin{aligned} \dim(\mathcal{S}_1 + \mathcal{S}_2) &= 2 \\ &= 1 + 1 : \end{aligned}$$

the sum is the plane $z = 0$ (Example 0.1), which has dimension 2 (Example 16 in Subspaces and Rank), and each line has dimension 1, with its spanning vector as a basis.

- For

$$\begin{aligned} \mathcal{S}_1 &= \mathcal{S}_2 \\ &= \text{span}\{(1, 0, 0)\}, \end{aligned}$$

the sum is the same line, because $a\,(1,0,0) + b\,(1,0,0) = (a+b)\,(1,0,0)$, so $\dim(\mathcal{S}_1 + \mathcal{S}_2) = 1 < 1 + 1$, a strict inequality.

Theorem 0.3 (Rank is subadditive)**.** *For $m \times n$ matrices $\mathbf{A}$ and $\mathbf{B}$,*

$$\text{rank}(\mathbf{A} + \mathbf{B}) \leq \text{rank}(\mathbf{A}) + \text{rank}(\mathbf{B}).$$

Proof

Proof. $\mathcal{C}(\mathbf{A} + \mathbf{B}) \subseteq \mathcal{C}(\mathbf{A}) + \mathcal{C}(\mathbf{B})$**.** Any vector of $\mathcal{C}(\mathbf{A} + \mathbf{B})$ is $(\mathbf{A} + \mathbf{B})\,\underset{\sim}{x}$ for some $\underset{\sim}{x} \in \mathbb{R}^n$ (Definition 1 in Column Space, Null Space and Rank-Nullity), and $(\mathbf{A} + \mathbf{B})\,\underset{\sim}{x} = \mathbf{A}\underset{\sim}{x} + \mathbf{B}\underset{\sim}{x}$ (Theorem 6 in Matrices, with $\underset{\sim}{x}$ as an $n \times 1$ matrix), a vector of $\mathcal{C}(\mathbf{A})$ plus a vector of $\mathcal{C}(\mathbf{B})$.

Compare dimensions. All the sets involved are subspaces of $\mathbb{R}^m$ (Theorem 1 in Column Space, Null Space and Rank-Nullity, Theorem 0.1), so

$$\begin{aligned} \text{rank}(\mathbf{A} + \mathbf{B}) &= \dim(\mathcal{C}(\mathbf{A} + \mathbf{B})) && \text{(Theorem 7 in Column Space, Null Space and Rank-Nullity)} \\ &\leq \dim(\mathcal{C}(\mathbf{A}) + \mathcal{C}(\mathbf{B})) && \text{(the containment, and part 2 of Theorem 10 in Subspaces and Ra} \\ &\leq \dim(\mathcal{C}(\mathbf{A})) + \dim(\mathcal{C}(\mathbf{B})) && \text{(Corollary 0.1)} \\ &= \text{rank}(\mathbf{A}) + \text{rank}(\mathbf{B}). && \text{(Theorem 7 in Column Space, Null Space and Rank-Nullity)} \end{aligned}$$

□

Example 0.6 (Subadditivity with equality and without)**.** Let $\mathbf{A} = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$ and $\mathbf{B} = \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}$. Each has one nonzero column and one zero column. The nonzero column on its own is linearly independent ($c\underset{\sim}{v} = \underset{\sim_2}{0}$ with $\underset{\sim}{v} \neq \underset{\sim_2}{0}$ forces $c = 0$), and the two columns together are not, since 1 times the zero column is $\underset{\sim_2}{0}$; so each matrix has rank 1 (Definition 2 in Subspaces and Rank). By the same argument, $-\mathbf{A}$ has rank 1.

- $\mathbf{A} + \mathbf{B}$ is the 2×2 identity matrix, which has rank 2 (Example 13 in Column Space, Null Space and Rank-Nullity), so the bound $2 \leq 1 + 1$ holds with equality.
- $\mathbf{A} + (-\mathbf{A})$ is the 2×2 zero matrix, which has rank 0: any nonempty list of its columns contains $\underset{\sim_2}{0}$, and $1 \cdot \underset{\sim_2}{0} = \underset{\sim_2}{0}$. So the bound $0 \leq \text{rank}(\mathbf{A}) + \text{rank}(-\mathbf{A}) = 1 + 1$ is strict.

Definition 0.2 (Complementary subspaces)**.** Let $\mathcal{S}_1$, $\mathcal{S}_2$ and $\mathcal{V}$ be subspaces of $\mathbb{R}^p$. $\mathcal{S}_1$ and $\mathcal{S}_2$ are **complementary** in $\mathcal{V}$ if

$$\mathcal{V} = \mathcal{S}_1 + \mathcal{S}_2 \quad \text{and} \quad \mathcal{S}_1 \cap \mathcal{S}_2 = \left\{\underset{\sim}{0}\right\}$$

(Definition 0.1).

Definition 0.3 (Direct sum)**.** If $\mathcal{S}_1$ and $\mathcal{S}_2$ are complementary in $\mathcal{V}$ (Definition 0.2), then $\mathcal{V}$ is the **direct sum** of $\mathcal{S}_1$ and $\mathcal{S}_2$, written $\mathcal{V} = \mathcal{S}_1 \oplus \mathcal{S}_2$.

Example 0.7 ($\mathbb{R}^3$ is a plane plus a line)**.** Let $\mathcal{S}_1 = \{(a, b, 0) : a, b \in \mathbb{R}\}$, the plane $z = 0$, and $\mathcal{S}_2 = \{(0, 0, c) : c \in \mathbb{R}\}$, the z-axis.

- **Sum:** any $(x, y, z) \in \mathbb{R}^3$ is $(x, y, 0) + (0, 0, z)$, so $\mathcal{S}_1 + \mathcal{S}_2 = \mathbb{R}^3$.
- **Intersection:** a vector in both has third entry 0 and first and second entries 0, so $\mathcal{S}_1 \cap \mathcal{S}_2 = \left\{\underset{\sim}{0}\right\}$.

So $\mathbb{R}^3 = \mathcal{S}_1 \oplus \mathcal{S}_2$.

Example 0.8 (Two planes that sum to $\mathbb{R}^3$ but not directly)**.** The planes $z = 0$ and $x = 0$ sum to $\mathbb{R}^3$ (Example 0.3), but they share the y-axis, so they are not complementary, and $\mathbb{R}^3$ is not their direct sum. The shared line shows up as more than one way to split a vector:

$$
\begin{aligned}
(0,1,0) &= (0,1,0) + \underset{\sim}{0} \\
&= \underset{\sim}{0} + (0,1,0),
\end{aligned}
$$

with the first term in the plane $z = 0$ and the second in the plane $x = 0$ both times.

Theorem 0.4 (Three ways to recognize a direct sum)**.** *Let $\mathcal{S}_1$ and $\mathcal{S}_2$ be subspaces of $\mathbb{R}^p$, and let $\mathcal{V} = \mathcal{S}_1 + \mathcal{S}_2$. The following statements are equivalent:*

1. *$\mathcal{V} = \mathcal{S}_1 \oplus \mathcal{S}_2$ (Definition 0.3).*
2. *$\dim(\mathcal{V}) = \dim(\mathcal{S}_1) + \dim(\mathcal{S}_2)$.*
3. *Every $\underset{\sim}{x} \in \mathcal{V}$ can be written as $\underset{\sim}{x} = \underset{\sim 1}{x} + \underset{\sim 2}{x}$ with $\underset{\sim 1}{x} \in \mathcal{S}_1$ and $\underset{\sim 2}{x} \in \mathcal{S}_2$ in only one way.*

Proof

Proof. Since $\mathcal{V} = \mathcal{S}_1 + \mathcal{S}_2$ is given, statement 1 says exactly that $\mathcal{S}_1 \cap \mathcal{S}_2 = \left\{\underset{\sim}{0}\right\}$.

1 and 2 are equivalent. By Corollary 0.1, $\dim(\mathcal{S}_1 + \mathcal{S}_2) = \dim(\mathcal{S}_1) + \dim(\mathcal{S}_2)$ exactly when $\mathcal{S}_1 \cap \mathcal{S}_2 = \left\{\underset{\sim}{0}\right\}$.

1 implies 3. Every $\underset{\sim}{x} \in \mathcal{V}$ has at least one such expression, because $\mathcal{V} = \mathcal{S}_1 + \mathcal{S}_2$. Suppose

$$
\begin{aligned}
\underset{\sim}{x} &= \underset{\sim 1}{u} + \underset{\sim 2}{u} \\
&= \underset{\sim 1}{v} + \underset{\sim 2}{v},
\end{aligned}
$$

with $\underset{\sim 1}{u}, \underset{\sim 1}{v} \in \mathcal{S}_1$ and $\underset{\sim 2}{u}, \underset{\sim 2}{v} \in \mathcal{S}_2$. Then

$$
\begin{aligned}
\underset{\sim 1}{u} - \underset{\sim 1}{v} &= (\underset{\sim 1}{u} + \underset{\sim 2}{u}) - \underset{\sim 2}{u} - \underset{\sim 1}{v} && \text{(add and subtract } \underset{\sim 2}{u}\text{)} \\
&= (\underset{\sim 1}{v} + \underset{\sim 2}{v}) - \underset{\sim 2}{u} - \underset{\sim 1}{v} && \text{(substitute } \underset{\sim 1}{u} + \underset{\sim 2}{u} = \underset{\sim 1}{v} + \underset{\sim 2}{v}\text{)} \\
&= \underset{\sim 2}{v} - \underset{\sim 2}{u}. && \text{(cancel } \underset{\sim 1}{v}\text{)}
\end{aligned}
$$

The left side is in $\mathcal{S}_1$ and the right side is in $\mathcal{S}_2$, because each subspace is closed under addition and scalar multiplication (the difference is $\underset{\sim 1}{u} + (-1)\underset{\sim 1}{v}$). So both sides are in $\mathcal{S}_1 \cap \mathcal{S}_2 = \left\{\underset{\sim}{0}\right\}$, which gives $\underset{\sim 1}{u} = \underset{\sim 1}{v}$ and $\underset{\sim 2}{u} = \underset{\sim 2}{v}$.

3 implies 1. Take any $\underset{\sim}{x} \in \mathcal{S}_1 \cap \mathcal{S}_2$. Then $\underset{\sim}{x} = \underset{\sim}{x} + \underset{\sim}{0}$, with $\underset{\sim}{x} \in \mathcal{S}_1$ and $\underset{\sim}{0} \in \mathcal{S}_2$, and $\underset{\sim}{x} = \underset{\sim}{0} + \underset{\sim}{x}$, with $\underset{\sim}{0} \in \mathcal{S}_1$ and $\underset{\sim}{x} \in \mathcal{S}_2$ (Theorem 1 in Subspaces and Rank). By statement 3 these two expressions are the same, so $\underset{\sim}{x} = \underset{\sim}{0}$, and $\mathcal{S}_1 \cap \mathcal{S}_2 = \left\{\underset{\sim}{0}\right\}$. □

Example 0.9 (Checking the three statements)**.** For the plane $z = 0$ and the z-axis in Example 0.7:

- the dimensions add up:

$$\begin{aligned} 2 + 1 &= 3 \\ &= \dim(\mathbb{R}^3), \end{aligned}$$

  since the plane has dimension 2 (Example 16 in Subspaces and Rank), the z-axis has the single nonzero vector $(0, 0, 1)$ as a basis, and $\dim(\mathbb{R}^3) = 3$ (Example 16 in Subspaces and Rank);

- the split is unique: if

$$\begin{aligned} (x, y, z) &= (a, b, 0) + (0, 0, c) \\ &= (a, b, c), \end{aligned}$$

  then $a = x$, $b = y$ and $c = z$.

For the planes $z = 0$ and $x = 0$ in Example 0.8, both statements fail:

$$\begin{aligned} 2 + 2 &= 4 \\ &\neq 3 \\ &= \dim(\mathcal{S}_1 + \mathcal{S}_2) \end{aligned}$$

(Example 0.4), and $(0, 1, 0)$ splits in two ways.

Definition 0.4 (Projection onto a subspace along a complement)**.** Let $\mathcal{S}$ and $\mathcal{T}$ be subspaces of $\mathbb{R}^p$ with $\mathbb{R}^p = \mathcal{S} \oplus \mathcal{T}$ (Definition 0.3), and let $\underset{\sim}{y} \in \mathbb{R}^p$. By Theorem 0.4, $\underset{\sim}{y} = \underset{\sim}{u} + \underset{\sim}{v}$ for exactly one $\underset{\sim}{u} \in \mathcal{S}$ and $\underset{\sim}{v} \in \mathcal{T}$. The vector $\underset{\sim}{u}$ is the **projection** of $\underset{\sim}{y}$ onto $\mathcal{S}$ along $\mathcal{T}$ (Banerjee and Roy 2014, Definition 6.3, p. 163).

Example 0.10 (The projection depends on the complement)**.** Let $\mathcal{S} = \text{span}\{(1, 0)\}$, the horizontal axis in $\mathbb{R}^2$, and let $\underset{\sim}{y} = (3, 2)$.

- **Along** $\mathcal{T}_1 = \text{span}\{(1, -1)\}$**:** $(3, 2) = (5, 0) + (-2, 2)$, with $(5, 0) \in \mathcal{S}$ and $(-2, 2) = -2 \cdot (1, -1) \in \mathcal{T}_1$, so the projection of $(3, 2)$ onto $\mathcal{S}$ along $\mathcal{T}_1$ is $(5, 0)$.
- **Along** $\mathcal{T}_2 = \text{span}\{(0, 1)\}$**:** $(3, 2) = (3, 0) + (0, 2)$, with $(3, 0) \in \mathcal{S}$ and $(0, 2) \in \mathcal{T}_2$, so the projection of $(3, 2)$ onto $\mathcal{S}$ along $\mathcal{T}_2$ is $(3, 0)$.

Each of $\mathcal{T}_1$ and $\mathcal{T}_2$ is a line that meets $\mathcal{S}$ only at $\underset{\sim}{0}$, so

$$\begin{aligned}\mathbb{R}^2 &= \mathcal{S} \oplus \mathcal{T}_1 \\ &= \mathcal{S} \oplus \mathcal{T}_2\end{aligned}$$

(Definition 0.3, Theorem 0.4). The same vector has a different projection onto the same subspace when the complement changes.

Orthogonal complements

Theorem 0.5 (The dot product is linear in each slot)**.** *For vectors* $\underset{\sim}{x}, \underset{\sim}{u}, \underset{\sim}{w} \in \mathbb{R}^p$ *and numbers* a, b*,*

$$\underset{\sim}{x} \cdot (a\,\underset{\sim}{u} + b\,\underset{\sim}{w}) = a\,(\underset{\sim}{x} \cdot \underset{\sim}{u}) + b\,(\underset{\sim}{x} \cdot \underset{\sim}{w})$$

and

$$(a\,\underset{\sim}{u} + b\,\underset{\sim}{w}) \cdot \underset{\sim}{x} = a\,(\underset{\sim}{u} \cdot \underset{\sim}{x}) + b\,(\underset{\sim}{w} \cdot \underset{\sim}{x}).$$

Proof

Proof. **Second slot.**

$$\begin{aligned}
\underset{\sim}{x} \cdot (a\,\underset{\sim}{u} + b\,\underset{\sim}{w}) &= \sum_{i=1}^{p} x_i\,(a\,\underset{\sim}{u} + b\,\underset{\sim}{w})_i && \text{(Definition 8 in Vectors)} \\
&= \sum_{i=1}^{p} x_i\,(au_i + bw_i) && \text{(Definition 6 in Matrices, Definition 7 in Vectors)} \\
&= \sum_{i=1}^{p} (x_i a u_i + x_i b w_i) && \text{(distribute each } x_i\text{)} \\
&= \sum_{i=1}^{p} (a\,x_i u_i + b\,x_i w_i) && \text{(commute the factors in each product)} \\
&= \sum_{i=1}^{p} a\,x_i u_i + \sum_{i=1}^{p} b\,x_i w_i && \text{(split the finite sum)} \\
&= a \sum_{i=1}^{p} x_i u_i + b \sum_{i=1}^{p} x_i w_i && \text{(factor } a \text{ and } b \text{ out of the sums)} \\
&= a\,(\underset{\sim}{x} \cdot \underset{\sim}{u}) + b\,(\underset{\sim}{x} \cdot \underset{\sim}{w}). && \text{(Definition 8 in Vectors)}
\end{aligned}$$

First slot.

$$
\begin{aligned}
(a\underset{\sim}{u} + b\underset{\sim}{w}) \cdot \underset{\sim}{x} &= \underset{\sim}{x} \cdot (a\underset{\sim}{u} + b\underset{\sim}{w}) && \text{(Theorem 1 in Vectors)} \\
&= a\,(\underset{\sim}{x} \cdot \underset{\sim}{u}) + b\,(\underset{\sim}{x} \cdot \underset{\sim}{w}) && \text{(second slot)} \\
&= a\,(\underset{\sim}{u} \cdot \underset{\sim}{x}) + b\,(\underset{\sim}{w} \cdot \underset{\sim}{x}). && \text{(Theorem 1 in Vectors, twice)}
\end{aligned}
$$

Two special cases are used often. With

$$
\begin{aligned}
a &= b \\
&= 1,
\end{aligned}
$$

and $1\,\underset{\sim}{u} = \underset{\sim}{u}$, the theorem gives $\underset{\sim}{x} \cdot (\underset{\sim}{u} + \underset{\sim}{w}) = \underset{\sim}{x} \cdot \underset{\sim}{u} + \underset{\sim}{x} \cdot \underset{\sim}{w}$. With $\underset{\sim}{w} = \underset{\sim}{u}$ and $b = 0$, and $0\,\underset{\sim}{u} = \underset{\sim}{0}$, it gives $\underset{\sim}{x} \cdot (a\,\underset{\sim}{u}) = a\,(\underset{\sim}{x} \cdot \underset{\sim}{u})$, and likewise in the first slot. □

Example 0.11 (Splitting a dot product)**.** Let $\underset{\sim}{x} = (1, 2)$, $\underset{\sim}{u} = (3, 0)$, $\underset{\sim}{w} = (0, 1)$, $a = 2$ and $b = -1$. Directly, $a\,\underset{\sim}{u} + b\,\underset{\sim}{w} = (6, -1)$ and

$$
\begin{aligned}
\underset{\sim}{x} \cdot (6, -1) &= 6 - 2 \\
&= 4.
\end{aligned}
$$

By Theorem 0.5,

$$
\begin{aligned}
a\,(\underset{\sim}{x} \cdot \underset{\sim}{u}) + b\,(\underset{\sim}{x} \cdot \underset{\sim}{w}) &= 2 \cdot 3 + (-1) \cdot 2 \\
&= 4,
\end{aligned}
$$

the same number.

Definition 0.5 (Orthogonal complement)**.** The **orthogonal complement** of a set $\mathcal{X}$ of vectors in $\mathbb{R}^p$ is the set of vectors orthogonal (Definition 15 in Vectors) to every vector of $\mathcal{X}$:

$$
\mathcal{X}^{\perp} \stackrel{\text{def}}{=} \left\{ \underset{\sim}{u} \in \mathbb{R}^p : \underset{\sim}{x} \cdot \underset{\sim}{u} = 0 \text{ for all } \underset{\sim}{x} \in \mathcal{X} \right\}.
$$

$\mathcal{X}^{\perp}$ is read "$\mathcal{X}$ perp".

Example 0.12 (Orthogonal complements in $\mathbb{R}^3$)**.**

- **A plane.** Let $\mathcal{X} = \{(a, b, 0) : a, b \in \mathbb{R}\}$, the plane $z = 0$. A vector $\underset{\sim}{u}$ is in $\mathcal{X}^{\perp}$ when

$$
\begin{aligned}
(a, b, 0) \cdot \underset{\sim}{u} &= au_1 + bu_2 \\
&= 0
\end{aligned}
$$

for all a and b. Taking $(a, b) = (1, 0)$ forces $u_1 = 0$, and taking $(a, b) = (0, 1)$ forces $u_2 = 0$; conversely, if

$$\begin{aligned} u_1 &= u_2 \\ &= 0, \end{aligned}$$

then $au_1 + bu_2 = 0$ for all a and b. So $\mathcal{X}^\perp = \{(0, 0, c) : c \in \mathbb{R}\}$, the z-axis.

- **Two vectors.** The two-element set $\{(1, 0, 0), (0, 1, 0)\}$ is not a subspace, and its orthogonal complement is the z-axis too: the two conditions $u_1 = 0$ and $u_2 = 0$ are the same as for the plane.
- **The extremes.** $\left\{\underset{\sim}{0}\right\}^\perp = \mathbb{R}^3$, because $\underset{\sim}{0} \cdot \underset{\sim}{u} = 0$ for every $\underset{\sim}{u}$. $(\mathbb{R}^3)^\perp = \left\{\underset{\sim}{0}\right\}$, because a $\underset{\sim}{u}$ in it is orthogonal to itself, and

$$\begin{aligned} \underset{\sim}{u} \cdot \underset{\sim}{u} &= u_1^2 + u_2^2 + u_3^2 \\ &= 0 \end{aligned}$$

forces $\underset{\sim}{u} = \underset{\sim}{0}$.

i Remark (The orthogonal complement of a set that is not a subspace)

Remark 0.1 (The orthogonal complement of a set that is not a subspace). $\mathcal{X}$ in Definition 0.5 need not be a subspace.

Theorem 0.6 (An orthogonal complement is a subspace)**.** *For any set $\mathcal{X}$ of vectors in $\mathbb{R}^p$, $\mathcal{X}^\perp$ (Definition 0.5) is a subspace of $\mathbb{R}^p$ (Definition 4 in Subspaces and Rank).*

i Proof

Proof. $\mathcal{X}^\perp$ is not empty:

$$\begin{aligned} \underset{\sim}{x} \cdot \underset{\sim}{0} &= \sum_i x_i \cdot 0 \\ &= 0 \end{aligned}$$

for every $\underset{\sim}{x}$, so $\underset{\sim}{0} \in \mathcal{X}^\perp$. Let $\underset{\sim}{u}, \underset{\sim}{w} \in \mathcal{X}^\perp$, let $c \in \mathbb{R}$, and let $\underset{\sim}{x} \in \mathcal{X}$.
Addition.

$$\begin{aligned} \underset{\sim}{x} \cdot (\underset{\sim}{u} + \underset{\sim}{w}) &= \underset{\sim}{x} \cdot \underset{\sim}{u} + \underset{\sim}{x} \cdot \underset{\sim}{w} && \text{(Theorem 0.5, special case } a \text{ and } b \text{ both equal to 1)} \\ &= 0 + 0 && (\underset{\sim}{u}, \underset{\sim}{w} \in \mathcal{X}^\perp) \\ &= 0. && \text{(arithmetic)} \end{aligned}$$

Scalar multiplication.

$$\begin{aligned} \underset{\sim}{x} \cdot (c\,\underset{\sim}{u}) &= c\,(\underset{\sim}{x} \cdot \underset{\sim}{u}) && \text{(Theorem 0.5, special case } b = 0) \\ &= c \cdot 0 && (\underset{\sim}{u} \in \mathcal{X}^\perp) \\ &= 0. && \text{(arithmetic)} \end{aligned}$$

These two equations hold for every $\underset{\sim}{x} \in \mathcal{X}$, so $\underset{\sim}{u} + \underset{\sim}{w}$ and $c\,\underset{\sim}{u}$ are in $\mathcal{X}^\perp$. □

Example 0.13 (The complement of a single vector is a line)**.** The one-element set $\mathcal{X} = \{(1,2)\}$ in $\mathbb{R}^2$ is not a subspace: $2\,(1,2) = (2,4)$ is not in it. Its orthogonal complement is

$$\begin{aligned} \mathcal{X}^\perp &= \left\{ \underset{\sim}{u} \in \mathbb{R}^2 : u_1 + 2u_2 = 0 \right\} \\ &= \{c\,(-2,1) : c \in \mathbb{R}\}, \end{aligned}$$

since $u_1 + 2u_2 = 0$ means

$$\begin{aligned} \underset{\sim}{u} &= (-2u_2, u_2) \\ &= u_2\,(-2,1). \end{aligned}$$

That set is span$\{(-2,1)\}$, a subspace by Theorem 2 in Subspaces and Rank, as Theorem 0.6 says it must be. It contains $(-2,1)$, the vector Remark 8 in Vectors found perpendicular to $(1,2)$.

Theorem 0.7 (The orthogonal complement of a column space is a null space)**.** *For any $m \times n$ matrix* $\mathbf{A}$*,*

$$\mathcal{C}(\mathbf{A})^\perp = \mathcal{N}(\mathbf{A}^\top).$$

Proof

Proof. Let $\underset{\sim}{a}_1, \ldots, \underset{\sim}{a}_n$ be the columns of $\mathbf{A}$, so $\underset{\sim}{a}_j = (a_{1j}, \ldots, a_{mj})$. For $\underset{\sim}{x} \in \mathbb{R}^m$, entry j of $\mathbf{A}^\top \underset{\sim}{x}$ is

$$
\begin{aligned}
(\mathbf{A}^\top \underset{\sim}{x})_j &= \sum_{i=1}^{m} (\mathbf{A}^\top)_{ji}\, x_i && \text{(Definition 13 in Matrices)} \\
&= \sum_{i=1}^{m} a_{ij}\, x_i && \text{(Definition 3 in Matrices)} \\
&= \underset{\sim}{a}_j \cdot \underset{\sim}{x}. && \text{(Definition 8 in Vectors)}
\end{aligned}
$$

So $\underset{\sim}{x} \in \mathcal{N}(\mathbf{A}^\top)$ (Definition 3 in Column Space, Null Space and Rank-Nullity) exactly when $\underset{\sim}{a}_j \cdot \underset{\sim}{x} = 0$ for every column $\underset{\sim}{a}_j$.

$\mathcal{C}(\mathbf{A})^\perp \subseteq \mathcal{N}(\mathbf{A}^\top)$. Each column $\underset{\sim}{a}_j$ is in $\mathcal{C}(\mathbf{A})$ (Theorem 1 in Column Space, Null Space and Rank-Nullity), so a vector orthogonal to all of $\mathcal{C}(\mathbf{A})$ is orthogonal to every column.

$\mathcal{N}(\mathbf{A}^\top) \subseteq \mathcal{C}(\mathbf{A})^\perp$. Suppose $\underset{\sim}{a}_j \cdot \underset{\sim}{x} = 0$ for every j, and take any $\underset{\sim}{y} \in \mathcal{C}(\mathbf{A})$, so $\underset{\sim}{y} = \mathbf{A}\underset{\sim}{z}$ for some $\underset{\sim}{z} \in \mathbb{R}^n$ (Definition 1 in Column Space, Null Space and Rank-Nullity). Then

$$
\begin{aligned}
\underset{\sim}{y} \cdot \underset{\sim}{x} &= \sum_{i=1}^{m} (\mathbf{A}\underset{\sim}{z})_i\, x_i && \text{(Definition 8 in Vectors)} \\
&= \sum_{i=1}^{m} \left(\sum_{j=1}^{n} a_{ij} z_j \right) x_i && \text{(Definition 13 in Matrices)} \\
&= \sum_{i=1}^{m} \sum_{j=1}^{n} a_{ij}\, z_j\, x_i && \text{(distribute each } x_i \text{ over the inner sum)} \\
&= \sum_{j=1}^{n} \sum_{i=1}^{m} a_{ij}\, z_j\, x_i && \text{(swap the order of the finite sums)} \\
&= \sum_{j=1}^{n} z_j \sum_{i=1}^{m} a_{ij}\, x_i && \text{(commute, then factor } z_j \text{ out of the inner sum)} \\
&= \sum_{j=1}^{n} z_j\, (\underset{\sim}{a}_j \cdot \underset{\sim}{x}) && \text{(Definition 8 in Vectors)} \\
&= \sum_{j=1}^{n} z_j \cdot 0 && (\text{each } \underset{\sim}{a}_j \cdot \underset{\sim}{x} = 0) \\
&= 0. && \text{(arithmetic)}
\end{aligned}
$$

So $\underset{\sim}{x}$ is orthogonal to every vector of $\mathcal{C}(\mathbf{A})$. □

Example 0.14 (The complement of the column space in Example 1 in Column Space, Null Space and Rank-Nullity)**.** For $\mathbf{A}$ in Example 1 in Column Space, Null Space and Rank-Nullity, $\mathcal{C}(\mathbf{A}) = \text{span}\{(1, 3)\}$ (1). Example 3 in Column Space, Null Space and Rank-Nullity found $\mathcal{N}(\mathbf{A}^\top) = \text{span}\{(-3, 1)\}$, so by Theorem 0.7 this line is $\mathcal{C}(\mathbf{A})^\perp$.

Directly, for any c,

$$\begin{aligned}(c\,(1,3))\cdot(-3,1) &= c\,((1,3)\cdot(-3,1)) && \text{(Theorem 0.5, special case } b=0\text{, first slot)}\\ &= c\,(-3+3) && \text{(Definition 8 in Vectors)}\\ &= 0. && \text{(arithmetic)}\end{aligned}$$

Theorem 0.8 (A subspace and its orthogonal complement make up the whole space)**.** *Let $\mathcal{S}$ be a subspace of $\mathbb{R}^p$. Then*

1. $\mathcal{S}\cap\mathcal{S}^\perp=\left\{\underset{\sim}{0}\right\}$;
2. $\dim(\mathcal{S})+\dim(\mathcal{S}^\perp)=p$;
3. $\mathbb{R}^p=\mathcal{S}\oplus\mathcal{S}^\perp$ *(Definition 0.3).*

So every $\underset{\sim}{y}\in\mathbb{R}^p$ can be written as $\underset{\sim}{y}=\underset{\sim}{u}+\underset{\sim}{v}$ with $\underset{\sim}{u}\in\mathcal{S}$ and $\underset{\sim}{v}\in\mathcal{S}^\perp$ in only one way (Theorem 0.4).

Proof

Proof. $\mathcal{S}^\perp$ is a subspace (Theorem 0.6), so it has a dimension (Definition 9 in Subspaces and Rank).

Part 1. Both $\mathcal{S}$ and $\mathcal{S}^\perp$ contain $\underset{\sim}{0}$ (Theorem 1 in Subspaces and Rank). If $\underset{\sim}{x}$ is in both, then $\underset{\sim}{x}$ is orthogonal to every vector of $\mathcal{S}$, $\underset{\sim}{x}$ itself included, so

$$\begin{aligned}\underset{\sim}{x}\cdot\underset{\sim}{x} &= x_1^2+\cdots+x_p^2\\ &= 0,\end{aligned}$$

which forces every $x_i=0$.

Part 2. Let $d=\dim(\mathcal{S})$. If $d=0$, then $\mathcal{S}$ has a basis (Theorem 9 in Subspaces and Rank) with 0 vectors (Definition 9 in Subspaces and Rank), the empty list, whose span is $\left\{\underset{\sim}{0}\right\}$ (Definition 5 in Subspaces and Rank); so $\mathcal{S}=\left\{\underset{\sim}{0}\right\}$, $\mathcal{S}^\perp=\mathbb{R}^p$ (every $\underset{\sim}{u}$ satisfies $\underset{\sim}{0}\cdot\underset{\sim}{u}=0$), and $0+p=p$ (Example 16 in Subspaces and Rank). If $d\geq 1$, choose a basis $\underset{\sim}{a}_1,\ldots,\underset{\sim}{a}_d$ of $\mathcal{S}$ (Theorem 9 in Subspaces and Rank), and let $\mathbf{A}$ be the $p\times d$ matrix with these columns. The basis spans $\mathcal{S}$, so $\mathcal{S}=\mathcal{C}(\mathbf{A})$ (Theorem 1 in Column Space, Null Space and Rank-Nullity), and $\mathcal{S}^\perp=\mathcal{N}(\mathbf{A}^\top)$ (Theorem 0.7). $\mathbf{A}^\top$ is $d\times p$, so

$$
\begin{aligned}
\dim(\mathcal{S}^\perp) &= \dim(\mathcal{N}(\mathbf{A}^\top)) && \text{(Theorem 0.7)} \\
&= \text{nullity}(\mathbf{A}^\top) && \text{(Definition 7 in Column Space, Null Space and Rank-Nullity)} \\
&= p - \text{rank}(\mathbf{A}^\top) && \text{(Theorem 8 in Column Space, Null Space and Rank-Nullity, for the } p \text{ columns o} \\
&= p - \text{rank}(\mathbf{A}) && \text{(Theorem 10 in Column Space, Null Space and Rank-Nullity)} \\
&= p - \dim(\mathcal{C}(\mathbf{A})) && \text{(Theorem 7 in Column Space, Null Space and Rank-Nullity)} \\
&= p - d. && (\mathcal{C}(\mathbf{A}) = \mathcal{S})
\end{aligned}
$$

Part 3. $\mathcal{S} + \mathcal{S}^\perp$ is a subspace of $\mathbb{R}^p$ (Theorem 0.1), and

$$
\begin{aligned}
\dim(\mathcal{S} + \mathcal{S}^\perp) &= \dim(\mathcal{S}) + \dim(\mathcal{S}^\perp) - \dim(\mathcal{S} \cap \mathcal{S}^\perp) && \text{(Theorem 0.2)} \\
&= p - \dim(\mathcal{S} \cap \mathcal{S}^\perp) && \text{(part 2)} \\
&= p - 0 && \text{(part 1, and } \dim(\{\underset{\sim}{0}\}) = 0 \text{ by Example 16 in Subspaces} \\
&= p. && \text{(arithmetic)}
\end{aligned}
$$

Since $\mathcal{S} + \mathcal{S}^\perp \subseteq \mathbb{R}^p$, $\mathbb{R}^p$ is a subspace (Example 3 in Subspaces and Rank) and $\dim(\mathbb{R}^p) = p$ (Example 16 in Subspaces and Rank), Theorem 11 in Subspaces and Rank gives $\mathcal{S} + \mathcal{S}^\perp = \mathbb{R}^p$. Together with part 1, $\mathbb{R}^p = \mathcal{S} \oplus \mathcal{S}^\perp$. The uniqueness of the split then follows from Theorem 0.4 (statement 1 implies statement 3). □

Example 0.15 (Splitting a vector of $\mathbb{R}^3$ along a line and its complement)**.** Let $\mathcal{S} = \text{span}\{(1,1,0)\}$ in $\mathbb{R}^3$. A single nonzero vector is linearly independent ($c\,\underset{\sim}{v} = \underset{\sim}{0}$ with $\underset{\sim}{v} \neq \underset{\sim}{0}$ forces $c = 0$), so $(1,1,0)$ is a basis of $\mathcal{S}$ and $\dim(\mathcal{S}) = 1$. A vector $\underset{\sim}{u}$ is in $\mathcal{S}^\perp$ exactly when $u_1 + u_2 = 0$: that condition says $(1,1,0) \cdot \underset{\sim}{u} = 0$, and then

$$
\begin{aligned}
c\,(1,1,0) \cdot \underset{\sim}{u} &= c\,(u_1 + u_2) \\
&= 0
\end{aligned}
$$

for every c. So

$$
\begin{aligned}
\mathcal{S}^\perp &= \{(a, -a, c) : a, c \in \mathbb{R}\} \\
&= \text{span}\{(1,-1,0), (0,0,1)\}.
\end{aligned}
$$

Those two vectors are linearly independent, because $a\,(1,-1,0) + c\,(0,0,1) = (a,-a,c)$ is $\underset{\sim}{0}$ only if

$$
\begin{aligned}
a &= c \\
&= 0,
\end{aligned}
$$

so $\dim(\mathcal{S}^\perp) = 2$, and $1 + 2 = 3$, as part 2 says.
To split $\underset{\sim}{y} = (3, 1, 2)$, look for $\underset{\sim}{u} = t\,(1, 1, 0)$ with

$$\begin{aligned}\underset{\sim}{v} &= \underset{\sim}{y} - \underset{\sim}{u} \\ &= (3 - t, 1 - t, 2)\end{aligned}$$

in $\mathcal{S}^\perp$: that membership needs $(3 - t) + (1 - t) = 0$, so $t = 2$. Then $\underset{\sim}{u} = (2, 2, 0)$ and $\underset{\sim}{v} = (1, -1, 2)$, with $\underset{\sim}{u} + \underset{\sim}{v} = (3, 1, 2)$ and

$$\begin{aligned}(1, 1, 0) \cdot (1, -1, 2) &= 1 - 1 + 0 \\ &= 0.\end{aligned}$$

Theorem 0.9 (The orthogonal complement of the orthogonal complement)**.** *For any subspace $\mathcal{S}$ of $\mathbb{R}^p$,*

$$(\mathcal{S}^\perp)^\perp = \mathcal{S}.$$

Proof

Proof. $\mathcal{S} \subseteq (\mathcal{S}^\perp)^\perp$**.** Take $\underset{\sim}{s} \in \mathcal{S}$ and any $\underset{\sim}{v} \in \mathcal{S}^\perp$. By Definition 0.5, $\underset{\sim}{s} \cdot \underset{\sim}{v} = 0$, so $\underset{\sim}{v} \cdot \underset{\sim}{s} = 0$ too (Theorem 1 in Vectors). So $\underset{\sim}{s}$ is orthogonal to every vector of $\mathcal{S}^\perp$, that is, $\underset{\sim}{s} \in (\mathcal{S}^\perp)^\perp$.
Equal dimensions. $\mathcal{S}^\perp$ is a subspace (Theorem 0.6), so part 2 of Theorem 0.8 applies to it as well as to $\mathcal{S}$:

$$\begin{aligned}\dim((\mathcal{S}^\perp)^\perp) &= p - \dim(\mathcal{S}^\perp) && \text{(Theorem 0.8, applied to } \mathcal{S}^\perp) \\ &= p - (p - \dim(\mathcal{S})) && \text{(Theorem 0.8, applied to } \mathcal{S}) \\ &= \dim(\mathcal{S}). && \text{(arithmetic)}\end{aligned}$$

$(\mathcal{S}^\perp)^\perp$ is a subspace (Theorem 0.6) that contains $\mathcal{S}$ and has the same dimension, so it equals $\mathcal{S}$ (Theorem 11 in Subspaces and Rank). □

Example 0.16 (Back to the line)**.** In Example 0.15, $\mathcal{S} = \text{span}\{(1, 1, 0)\}$ and $\mathcal{S}^\perp = \text{span}\{(1, -1, 0), (0, 0, 1)\}$. Put those two spanning vectors in the columns of the 3×2 matrix $\mathbf{B} = \begin{bmatrix} 1 & 0 \\ -1 & 0 \\ 0 & 1 \end{bmatrix}$, so $\mathcal{S}^\perp = \mathcal{C}(\mathbf{B})$ (Theorem 1 in Column Space, Null Space and Rank-Nullity) and $(\mathcal{S}^\perp)^\perp = \mathcal{N}(\mathbf{B}^\top)$ (Theorem 0.7). Since $\mathbf{B}^\top \underset{\sim}{w} = (w_1 - w_2,\ w_3)$ (Definition 13 in Matrices), $\underset{\sim}{w} \in (\mathcal{S}^\perp)^\perp$ exactly when $w_1 - w_2 = 0$ and $w_3 = 0$. So

$$
\begin{aligned}
(\mathcal{S}^{\perp})^{\perp} &= \{(a, a, 0) : a \in \mathbb{R}\} \\
&= \mathcal{S}.
\end{aligned}
$$

Example 0.17 (A set that is not a subspace does not come back)**.** The one-element set $\mathcal{X} = \{(1, 1, 0)\}$ is not a subspace. Its orthogonal complement is the same as that of $\mathcal{S}$ in Example 0.16, since both are defined by the single condition $u_1 + u_2 = 0$, so $(\mathcal{X}^{\perp})^{\perp} = \text{span}\{(1, 1, 0)\}$, which contains $(2, 2, 0) \notin \mathcal{X}$. Theorem 0.9 needs $\mathcal{S}$ to be a subspace.

The fundamental theorem of linear algebra

Theorem 0.10 (Fundamental theorem of linear algebra)**.** *Let* $\mathbf{A}$ *be an* $m \times n$ *matrix with* $\text{rank}(\mathbf{A}) = r$*. Then*

1. $\mathcal{C}(\mathbf{A})^{\perp} = \mathcal{N}(\mathbf{A}^{\top})$*, and* $\mathbb{R}^m = \mathcal{C}(\mathbf{A}) \oplus \mathcal{N}(\mathbf{A}^{\top})$*;*
2. $\mathcal{C}(\mathbf{A}) = \mathcal{N}(\mathbf{A}^{\top})^{\perp}$*;*
3. $\mathcal{N}(\mathbf{A})^{\perp} = \mathcal{C}(\mathbf{A}^{\top})$*, and* $\mathbb{R}^n = \mathcal{C}(\mathbf{A}^{\top}) \oplus \mathcal{N}(\mathbf{A})$*;*
4. $$
\begin{aligned}
\dim(\mathcal{C}(\mathbf{A})) &= \dim(\mathcal{C}(\mathbf{A}^{\top})) \\
&= r,
\end{aligned}
$$

   $\dim(\mathcal{N}(\mathbf{A})) = n - r$*, and* $\dim(\mathcal{N}(\mathbf{A}^{\top})) = m - r$*.*

Proof

Proof. **Part 1.** The first equation is Theorem 0.7. $\mathcal{C}(\mathbf{A})$ is a subspace of $\mathbb{R}^m$ (Theorem 1 in Column Space, Null Space and Rank-Nullity), so

$$
\begin{aligned}
\mathbb{R}^m &= \mathcal{C}(\mathbf{A}) \oplus \mathcal{C}(\mathbf{A})^{\perp} && \text{(Theorem 0.8)} \\
&= \mathcal{C}(\mathbf{A}) \oplus \mathcal{N}(\mathbf{A}^{\top}). && \text{(first equation)}
\end{aligned}
$$

Part 2.

$$
\begin{aligned}
\mathcal{C}(\mathbf{A}) &= (\mathcal{C}(\mathbf{A})^{\perp})^{\perp} && \text{(Theorem 0.9)} \\
&= \mathcal{N}(\mathbf{A}^{\top})^{\perp}. && \text{(part 1)}
\end{aligned}
$$

Part 3. Apply parts 2 and 1 to the $n \times m$ matrix $\mathbf{A}^{\top}$, using $(\mathbf{A}^{\top})^{\top} = \mathbf{A}$: part 2 gives $\mathcal{C}(\mathbf{A}^{\top}) = \mathcal{N}(\mathbf{A})^{\perp}$, and part 1 gives $\mathbb{R}^n = \mathcal{C}(\mathbf{A}^{\top}) \oplus \mathcal{N}(\mathbf{A})$.

Part 4. $\dim(\mathcal{C}(\mathbf{A})) = r$ by Theorem 7 in Column Space, Null Space and Rank-Nullity. $\dim(\mathcal{C}(\mathbf{A}^\top)) = \operatorname{rank}(\mathbf{A}^\top)$ by Theorem 7 in Column Space, Null Space and Rank-Nullity, and that is r by Theorem 10 in Column Space, Null Space and Rank-Nullity. $\mathbf{A}$ has n columns, so $\dim(\mathcal{N}(\mathbf{A})) = n - r$ (Definition 7 in Column Space, Null Space and Rank-Nullity, Theorem 8 in Column Space, Null Space and Rank-Nullity). $\mathbf{A}^\top$ has m columns and rank r, so $\dim(\mathcal{N}(\mathbf{A}^\top)) = m - r$ in the same way. □

Example 0.18 (The four subspaces of the matrix in Example 1 in Column Space, Null Space and Rank-Nullity)**.** For $\mathbf{A}$ in Example 1 in Column Space, Null Space and Rank-Nullity, $m = 2$, $n = 3$ and $r = 1$ (Example 11 in Column Space, Null Space and Rank-Nullity).

- **In $\mathbb{R}^3$:** the row space $\mathcal{C}(\mathbf{A}^\top) = \operatorname{span}\{(1, -2, -2)\}$

  (1) has dimension $1 = r$, and $\mathcal{N}(\mathbf{A}) = \operatorname{span}\{(2, 1, 0), (2, 0, 1)\}$ has dimension $2 = n - r$ (Example 12 in Column Space, Null Space and Rank-Nullity). The spanning vectors are orthogonal:

$$\begin{aligned}(1, -2, -2) \cdot (2, 1, 0) &= 2 - 2 + 0 \\ &= 0\end{aligned}$$

  and

$$\begin{aligned}(1, -2, -2) \cdot (2, 0, 1) &= 2 + 0 - 2 \\ &= 0.\end{aligned}$$

- **In $\mathbb{R}^2$:** $\mathcal{C}(\mathbf{A}) = \operatorname{span}\{(1, 3)\}$ has dimension $1 = r$, and $\mathcal{N}(\mathbf{A}^\top) = \operatorname{span}\{(-3, 1)\}$ has dimension $1 = m - r$ (Example 0.14).

By part 3, $y = (1, 0, 0)$ splits uniquely into a row-space part $t\,(1, -2, -2)$ and a null-space part. The null-space part $\underset{\sim}{y} - t\,(1, -2, -2)$ lies in $\mathcal{N}(\mathbf{A})$ (part 3), and $\mathcal{N}(\mathbf{A}) = \mathcal{C}(\mathbf{A}^\top)^\perp$ (Theorem 0.7 applied to $\mathbf{A}^\top$, with $(\mathbf{A}^\top)^\top = \mathbf{A}$), so it must be orthogonal to $(1, -2, -2)$:

$$\begin{aligned}
0 &= (1, -2, -2) \cdot ((1, 0, 0) - t\,(1, -2, -2)) && \text{(orthogonality)} \\
&= (1, -2, -2) \cdot (1 \cdot (1, 0, 0) + (-t)\,(1, -2, -2)) && \text{(write the difference as a linear combination)} \\
&= 1 \cdot ((1, -2, -2) \cdot (1, 0, 0)) + (-t)\,((1, -2, -2) \cdot (1, -2, -2)) && \text{(Theorem 0.5)} \\
&= 1 \cdot 1 + (-t) \cdot 9 && \text{(Definition 8 in Vectors)} \\
&= 1 - 9t, && \text{(arithmetic)}
\end{aligned}$$

so $t = \frac{1}{9}$. The row-space part is $\left(\frac{1}{9}, -\frac{2}{9}, -\frac{2}{9}\right)$ and the null-space part is $\left(\frac{8}{9}, \frac{2}{9}, \frac{2}{9}\right)$; as a check, row 1 of $\mathbf{A}$ gives $\frac{8}{9} - 2 \cdot \frac{2}{9} - 2 \cdot \frac{2}{9} = 0$, and row 2 is 3 times row 1, so $\mathbf{A}$ sends the null-space part to $\underset{\sim 2}{0}$.

Banerjee, Sudipto, and Anindya Roy. 2014. *Linear Algebra and Matrix Analysis for Statistics*. Vol. 181. Crc Press Boca Raton. https://www.routledge.com/Linear-Algebra-and-Matrix-Analysis-for-Statistics/Banerjee-Roy/p/book/9781420095388.

Zhou, Hua. 2024. *Orthogonal Projections*. Lecture notes for Biostat 216, Mathematical Methods for Biostatistics, University of California, Los Angeles. https://ucla-biostat-216.github.io/2024fall/slides/06-orthproj/06-orthproj.html.