# Sets

Last modified: 2026-10-17 04:46:25 (PDT)

Remark

*Remark.* Probability and statistics describe events, outcomes, and parameters as sets. This page collects the definitions of sets that the Morrison Lab's probability and statistics notes build on. The Functions page defines functions between sets, and the Expressions and Algorithms page defines the expressions that write functions down and the algorithms that compute them.

## Sets

**Definition 0.1** (Set)**.** A **set** is a collection of distinct objects, called its elements. We write $a \in A$ when $a$ is an element of the set $A$, and $a \notin A$ when it is not. Two sets are equal when they have exactly the same elements.

Remark (Naive and formal set theory)

*Remark* 0.1 (Naive and formal set theory). Definition 0.1 is the informal ("naive") notion of a set, which is all these notes need. Formal set theory states axioms for sets instead (see Wikipedia: Set (mathematics)). For example, the last sentence of Definition 0.1, that two sets are equal when they have exactly the same elements, is one of those axioms, the *axiom of extensionality*.

**Example 0.1** (Sets of die rolls)**.** The possible results of rolling a six-sided die form the set $\{1, 2, 3, 4, 5, 6\}$, with $3 \in \{1, 2, 3, 4, 5, 6\}$ and $7 \notin \{1, 2, 3, 4, 5, 6\}$. The sets $\{1, 2, 3\}$ and $\{3, 2, 1\}$ are equal, because they have the same elements; the order in which the elements are listed does not matter.

**Definition 0.2** (Set-builder notation)**.** For a set $A$ and a statement $P(x)$ about an element $x$, **set-builder notation** $\{x \in A : P(x)\}$ denotes the set of elements $x$ of $A$ for which $P(x)$ is true.

**Example 0.2** (Even die rolls in set-builder notation)**.** $\{x \in \{1,2,3,4,5,6\} : x \text{ is even}\} = \{2,4,6\}$ is the set of even rolls, and $\{x \in \mathbb{R} : x > 0\}$ is the set of positive real numbers.

**Definition 0.3** (Interval)**.** For real numbers $a \leq b$, the **intervals** with **endpoints** $a$ and $b$ are the closed interval (Definition 0.4), the open interval (Definition 0.5), and the half-open intervals (Definition 0.6). The unbounded intervals (Definition 0.7) have at most one endpoint.
A square bracket means the endpoint belongs to the interval, and a parenthesis means it does not. The symbols $\infty$ and $-\infty$ are not real numbers, so they always get a parenthesis.

**Definition 0.4** (Closed interval)**.** For real numbers $a \leq b$, the **closed interval** is $[a,b] \stackrel{\text{def}}{=} \{x \in \mathbb{R} : a \leq x \leq b\}$.

**Definition 0.5** (Open interval)**.** For real numbers $a \leq b$, the **open interval** is $(a,b) \stackrel{\text{def}}{=} \{x \in \mathbb{R} : a < x < b\}$.

**Definition 0.6** (Half-open intervals)**.** For real numbers $a \leq b$, the **half-open intervals** are $[a,b) \stackrel{\text{def}}{=} \{x \in \mathbb{R} : a \leq x < b\}$ and $(a,b] \stackrel{\text{def}}{=} \{x \in \mathbb{R} : a < x \leq b\}$.

**Definition 0.7** (Unbounded intervals)**.** The **unbounded intervals** have at most one endpoint:

- $[a,\infty) \stackrel{\text{def}}{=} \{x \in \mathbb{R} : x \geq a\}$ and $(a,\infty) \stackrel{\text{def}}{=} \{x \in \mathbb{R} : x > a\}$;
- $(-\infty,b] \stackrel{\text{def}}{=} \{x \in \mathbb{R} : x \leq b\}$ and $(-\infty,b) \stackrel{\text{def}}{=} \{x \in \mathbb{R} : x < b\}$;
- $(-\infty,\infty) \stackrel{\text{def}}{=} \mathbb{R}$.

**Example 0.3** (Which numbers are in an interval)**.**

- $1 \in [1,2]$, but $1 \notin (1,2)$.
- $2 \in (1,2]$, but $2 \notin [1,2)$.
- 1.5 is in all four of $[1,2]$, $(1,2)$, $[1,2)$, and $(1,2]$.
- $[0,\infty)$ is the set of non-negative real numbers: $0 \in [0,\infty)$ and $1000 \in [0,\infty)$, but $-0.1 \notin [0,\infty)$.

**Definition 0.8** (Subset)**.** A set $A$ is a **subset** of a set $B$, written $A \subseteq B$, if every element of $A$ is an element of $B$.

**Example 0.4** (Even die rolls)**.** The even die rolls form a subset of the possible rolls: $\{2, 4, 6\} \subseteq \{1, 2, 3, 4, 5, 6\}$, because each of 2, 4, and 6 is a possible roll. The set $\{2, 4, 7\}$ is not a subset of $\{1, 2, 3, 4, 5, 6\}$, because 7 is not a possible roll.

**Theorem 0.1** (Every set is a subset of itself)**.** *For every set $A$, $A \subseteq A$.*

Proof

*Proof.* Every element of $A$ is an element of $A$, which is Definition 0.8 with $B = A$. □

**Definition 0.9** (Strict subset)**.** A set $A$ is a **strict subset** of a set $B$, written $A \subsetneq B$, if $A$ is a subset of $B$ and $A \neq B$.

Remark (Proper subsets and the symbol $\subset$)

*Remark* 0.2 (Proper subsets and the symbol $\subset$). Other sources call a strict subset a **proper subset**. Sources disagree about the symbol $\subset$: some use it for "subset" ($\subseteq$), and others for "strict subset" ($\subsetneq$) (see Wikipedia: Subset). For example, $\{1, 2\} \subset \{1, 2\}$ is true in the first reading, because $\{1, 2\} \subseteq \{1, 2\}$, and false in the second, because the two sets are equal. These notes avoid $\subset$ and write $\subseteq$ or $\subsetneq$.

**Definition 0.10** (Superset)**.** A set $B$ is a **superset** of a set $A$, written $B \supseteq A$, if every element of $A$ is an element of $B$.

Remark (Superset and subset)

*Remark* 0.3 (Superset and subset). $B \supseteq A$ means the same thing as $A \subseteq B$ (subset). For example, $\{1, 2, 3\} \supseteq \{1, 3\}$, and $\{1, 3\} \subseteq \{1, 2, 3\}$. But $\{1, 3\}$ is not a superset of $\{1, 2\}$, because $2 \in \{1, 2\}$ and $2 \notin \{1, 3\}$.

**Definition 0.11** (Strict superset)**.** A set $B$ is a **strict superset** of a set $A$, written $B \supsetneq A$, if $B$ is a superset of $A$ and $B \neq A$.

Remark (Strict superset and strict subset)

*Remark* 0.4 (Strict superset and strict subset). $B \supsetneq A$ means the same thing as $A \subsetneq B$ (strict subset). For example, $\{1, 2, 3\} \supsetneq \{1, 3\}$, because $\{1, 2, 3\}$ contains 2 and $\{1, 3\}$ does not.

**Example 0.5** (Strict subsets and supersets of die rolls)**.** For the even rolls $A = \{2, 4, 6\}$ and the possible rolls $B = \{1, 2, 3, 4, 5, 6\}$:

- $A \subsetneq B$, because $A \subseteq B$ (Example 0.4) and $1 \in B$ but $1 \notin A$, so $A \neq B$;
- $B \supseteq A$ and $B \supsetneq A$, for the same reasons;
- $B \subseteq B$, but $B$ is not a strict subset of itself, because $B = B$; likewise $B \supseteq B$, but $B$ is not a strict superset of itself, because $B = B$.

**Definition 0.12** (Empty set)**.** The **empty set**, denoted $\emptyset$, is the set that has no elements.

Remark (Other notation for the empty set)

*Remark* 0.5 (Other notation for the empty set). Other sources write $\{\}$ for $\emptyset$ (see Wikipedia: Empty set). Some sources call it the **null set**, but in measure theory a "null set" usually means a set of measure zero, which need not be empty. For example, the interval $[0, 0] = \{0\}$ has length $0 - 0 = 0$, so it is a null set in that sense, but it is not empty, because $0 \in \{0\}$.

**Example 0.6** (Impossible die rolls)**.** No roll of a six-sided die is greater than 6, so $\{x \in \{1, 2, 3, 4, 5, 6\} : x > 6\} = \emptyset$.

**Definition 0.13** (Vacuous truth)**.** A statement of the form "every element of $A$ has property $P$" is **vacuously true** when $A$ is the empty set: $A$ has no elements, so no element of $A$ fails to have property $P$.

**Example 0.7** (Every impossible die roll is even)**.** The set of die rolls greater than 6 is $\emptyset$ (Example 0.6). So "every die roll greater than 6 is even" is vacuously true, and so is "every die roll greater than 6 is odd": there is no such roll to fail either statement.

**Theorem 0.2** (There is only one empty set)**.** *If $E_1$ and $E_2$ are sets with no elements, then $E_1 = E_2$.*

Proof

*Proof.* Neither set has any elements, so every element of $E_1$ is an element of $E_2$ and vice versa, vacuously. Sets with the same elements are equal (Definition 0.1). □

**Theorem 0.3** (The empty set is a subset of every set)**.** *For every set $A$, $\emptyset \subseteq A$.*

Proof

*Proof.* By Definition 0.8, $\emptyset \subseteq A$ fails only if some element of $\emptyset$ is not in $A$. The empty set has no elements, so no such element exists. □

**Definition 0.14** (Collection of sets (family of sets))**.** A **collection of sets** (also called a **family of sets**) is a set whose elements are themselves sets.

**Example 0.8** (The collection of all subsets of a two-element set)**.** The subsets of $\{1, 2\}$ are $\emptyset$, $\{1\}$, $\{2\}$, and $\{1, 2\}$ (Theorem 0.3 and Theorem 0.1 give the first and the last), so the collection of all subsets of $\{1, 2\}$ is

$$\{\emptyset, \{1\}, \{2\}, \{1, 2\}\}$$

It has four elements, each of them a set. The collection $\{\emptyset\}$ has one element, the empty set, so it is not equal to $\emptyset$, which has none.

**Definition 0.15** (Label)**.** A **label** is a name or symbol attached to an object, so that we can refer to the object and tell it apart from other objects.

**Example 0.9** (Labels)**.** The numbers $1, 2, 3$ can label three students in a class list. Then "student 2" refers to one particular student. The letters $A$ and $B$ can label two sets.

**Definition 0.16** (Index set)**.** An **index set** is a set whose elements are used as labels for objects. Each element of the index set is called an index.

**Example 0.10** (Index sets)**.** The set $\{1, \ldots, n\}$ labels the entries of a vector with $n$ entries. The set of nonnegative real numbers can label moments in time.

**Definition 0.17** (Indexed family)**.** An **indexed family** is a collection of objects, one for each element of a given index set. The object for the index $i$ is written $a_i$.

**Example 0.11** (Indexed families)**.** The entries $x_1, \dots, x_n$ of a vector are an indexed family of numbers, with index set $\{1, \dots, n\}$. The functions $f_\theta$, one for each parameter value $\theta$ in a parameter set, are an indexed family of functions.

## Combining sets

**Definition 0.18** (Union)**.** The **union** of sets $A$ and $B$, written $A \cup B$, is the set of elements that are in $A$, in $B$, or in both:

$$A \cup B \stackrel{\text{def}}{=} \{x : x \in A \text{ or } x \in B\}$$

**Definition 0.19** (Union of a family of sets)**.** Let $I$ be a set of indices, such as $\{1, \dots, n\}$ or $\{1, 2, 3, ...\}$, and let $A_i$ be a set for each $i \in I$. The **union** of the sets $A_i$, written $\bigcup_{i \in I} A_i$, is the set of elements that are in at least one $A_i$. For $I = \{1, \dots, n\}$, it is also written $\bigcup_{i=1}^{n} A_i$ or $A_1 \cup \cdots \cup A_n$, and for $I = \{1, 2, 3, ...\}$, it is written $\bigcup_{i=1}^{\infty} A_i$.

**Definition 0.20** (Intersection)**.** The **intersection** of sets $A$ and $B$, written $A \cap B$, is the set of elements that are in both $A$ and $B$:

$$A \cap B \stackrel{\text{def}}{=} \{x : x \in A \text{ and } x \in B\}$$

**Definition 0.21** (Intersection of a family of sets)**.** Let $I$ be a nonempty set of indices, such as $\{1, \dots, n\}$ or $\{1, 2, 3, ...\}$, and let $A_i$ be a set for each $i \in I$. The **intersection** of the sets $A_i$, written $\bigcap_{i \in I} A_i$, is the set of elements that are in every $A_i$. For $I = \{1, \dots, n\}$, it is also written $\bigcap_{i=1}^{n} A_i$ or $A_1 \cap \cdots \cap A_n$, and for $I = \{1, 2, 3, ...\}$, it is written $\bigcap_{i=1}^{\infty} A_i$.

**Definition 0.22** (Set difference)**.** The **set difference** of sets $A$ and $B$, written $A \setminus B$, is the set of elements of $A$ that are not in $B$:

$$A \setminus B \stackrel{\text{def}}{=} \{x \in A : x \notin B\}$$

**Definition 0.23** (Complement)**.** Let $A$ be a subset of a set $S$. The **complement** of $A$ in $S$ is the set difference (Definition 0.22) $S \setminus A$: the elements of $S$ that are not in $A$. When the set $S$ is clear from context, the complement of $A$ is written $A^c$:

$$A^c \stackrel{\text{def}}{=} S \setminus A = \{x \in S : x \notin A\}$$

**Example 0.12** (Complements of sets of die rolls)**.** Let $S = \{1, 2, 3, 4, 5, 6\}$ be the possible rolls of a die.

- The complement of the even rolls $A = \{2, 4, 6\}$ is $A^c = \{1, 3, 5\}$, the odd rolls.
- The complement of $S$ itself is $S^c = \emptyset$, and the complement of $\emptyset$ is $\emptyset^c = S$.

The complement depends on $S$: in $S = \{1, 2, 3, 4, 5, 6\}$, the complement of $\{2, 4, 6\}$ is $\{1, 3, 5\}$, but in $S = \{2, 4, 6, 8\}$, it is $\{8\}$.

**Example 0.13** (Combining sets of die rolls)**.** Let $A = \{2, 4, 6\}$ (the even rolls) and $B = \{1, 2, 3\}$ (the rolls of at most 3). Then:

- $A \cup B = \{1, 2, 3, 4, 6\}$, the rolls that are even or at most 3;
- $A \cap B = \{2\}$, the only roll that is both;
- $A \setminus B = \{4, 6\}$, the even rolls greater than 3;
- $\{2, 4, 6\} \cap \{1, 3, 5\} = \emptyset$: no roll is both even and odd.

**Example 0.14** (Unions and intersections of many sets)**.**

- $$\begin{aligned}\bigcup_{i=1}^{3} \{i, i+1\} &= \{1, 2\} \cup \{2, 3\} \cup \{3, 4\} \\ &= \{1, 2, 3, 4\}.\end{aligned}$$
- $\bigcup_{i=1}^{\infty} \{i\} = \mathbb{N}$: each natural number $n$ is in the set $\{n\}$, and each set $\{i\}$ contains only the natural number $i$.
- $\bigcap_{i=1}^{\infty} [0, 1/i] = \{0\}$: 0 is in every interval $[0, 1/i]$; a number $x > 0$ is not in $[0, 1/i]$ once $i > 1/x$; and a number $x < 0$ is in none of them.

**Theorem 0.4** (De Morgan's laws)**.** *Let $S$ be a set, let $I$ be a nonempty set of indices, and let $A_i$ be a subset of $S$ for each $i \in I$. Then the complement of a union is the intersection of the complements, and the complement of an intersection is the union of*

*the complements:*

$$S \setminus \bigcup_{i \in I} A_i = \bigcap_{i \in I} (S \setminus A_i)$$
$$S \setminus \bigcap_{i \in I} A_i = \bigcup_{i \in I} (S \setminus A_i)$$

*With the complement notation of Definition 0.23, these read* $\left(\bigcup_{i \in I} A_i\right)^c = \bigcap_{i \in I} A_i^c$ *and* $\left(\bigcap_{i \in I} A_i\right)^c = \bigcup_{i \in I} A_i^c$.

Proof

*Proof.* Two sets are equal when they have the same elements (Definition 0.1), so it is enough to show that an element $x$ is in the left-hand side if and only if ( $\iff$ ) it is in the right-hand side. For the first law:

$$\begin{aligned}
x \in S \setminus \bigcup_{i \in I} A_i &\iff x \in S \text{ and } x \notin \bigcup_{i \in I} A_i && \text{(Definition 0.22)} \\
&\iff x \in S \text{ and } x \notin A_i \text{ for every } i \in I && \text{(Definition 0.18)} \\
&\iff x \in S \setminus A_i \text{ for every } i \in I && \text{(Definition 0.22)} \\
&\iff x \in \bigcap_{i \in I} (S \setminus A_i) && \text{(Definition 0.20)}
\end{aligned}$$

For the second law:

$$\begin{aligned}
x \in S \setminus \bigcap_{i \in I} A_i &\iff x \in S \text{ and } x \notin \bigcap_{i \in I} A_i && \text{(Definition 0.22)} \\
&\iff x \in S \text{ and } x \notin A_i \text{ for some } i \in I && \text{(Definition 0.20)} \\
&\iff x \in S \setminus A_i \text{ for some } i \in I && \text{(Definition 0.22)} \\
&\iff x \in \bigcup_{i \in I} (S \setminus A_i) && \text{(Definition 0.18)}
\end{aligned}$$

□

**Example 0.15** (De Morgan's laws for die rolls)**.** In $S = \{1, 2, 3, 4, 5, 6\}$, let $A = \{2, 4, 6\}$ and $B = \{1, 2, 3\}$, as in Example 0.13, so $A^c = \{1, 3, 5\}$ and $B^c = \{4, 5, 6\}$.

- $$\begin{aligned}(A \cup B)^c &= \{1, 2, 3, 4, 6\}^c \\ &= \{5\},\end{aligned}$$

  and

$$\begin{aligned} A^c \cap B^c &= \{1,3,5\} \cap \{4,5,6\} \\ &= \{5\}. \end{aligned}$$

- $$\begin{aligned} (A \cap B)^c &= \{2\}^c \\ &= \{1,3,4,5,6\}, \end{aligned}$$

  and

$$\begin{aligned} A^c \cup B^c &= \{1,3,5\} \cup \{4,5,6\} \\ &= \{1,3,4,5,6\}. \end{aligned}$$

**Definition 0.24** (Ordered pair)**.** An **ordered pair** $(a, b)$ is a pair of objects listed in order, with $a$ first and $b$ second; two ordered pairs are equal, $(a, b) = (c, d)$, when $a = c$ and $b = d$.

**Definition 0.25** (Cartesian product)**.** The **Cartesian product** of sets $A$ and $B$, written $A \times B$, is the set of ordered pairs whose first entry is in $A$ and whose second entry is in $B$:

$$A \times B \stackrel{\text{def}}{=} \{(a, b) : a \in A \text{ and } b \in B\}$$

**Example 0.16** (Cartesian products)**.**

- $\{1, 2\} \times \{5, 6\} = \{(1, 5), (1, 6), (2, 5), (2, 6)\}$. The pair $(5, 1)$ is not in it, because its first entry, 5, is not in $\{1, 2\}$; so $(1, 5) \neq (5, 1)$.
- The results of rolling a die twice form $\{1, \ldots, 6\} \times \{1, \ldots, 6\}$, which has $6 \cdot 6 = 36$ elements, such as $(3, 5)$: a 3 on the first roll and a 5 on the second.
- $[0, 2] \times [0, 3]$ is the set of points $(x, y)$ in the plane with $0 \leq x \leq 2$ and $0 \leq y \leq 3$: a rectangle of width 2 and height 3.

### Extended non-negative real numbers

**Definition 0.26** (Extended non-negative real numbers)**.** The **extended non-negative real numbers**, written $[0, \infty]$, are the non-negative real numbers together with an extra element $\infty$ that is greater than every real number:

$$[0, \infty] \stackrel{\text{def}}{=} [0, \infty) \cup \{\infty\}$$

Addition extends to $[0, \infty]$ by setting

$$x + \infty = \infty + x$$
$$= \infty$$

for every $x \in [0, \infty]$.

Remark (Arithmetic with $\infty$)

*Remark* 0.6 (Arithmetic with $\infty$). $\infty$ is not a real number, so some real-number arithmetic does not extend to it: $\infty - \infty$ is left undefined, and $\infty + 1 = \infty + 2$ does not imply $1 = 2$. Measure theory uses $[0, \infty]$ because the size of a set, such as the length of the whole real line, can be infinite (see Wikipedia: Extended real number line).

**Example 0.17** (Adding with $\infty$)**.** In $[0, \infty]$, $2 + 3 = 5$ as usual, while $2 + \infty = \infty$ and $\infty + \infty = \infty$.

## Further reading

These books treat sets and the logic behind them in more depth.

- Devlin (1993) is an undergraduate introduction to axiomatic set theory. It covers the axioms of set theory, ordinals, and cardinals, which go well beyond the countable sets and functions in these notes.
- Halmos (1974) is a short, informal treatment of the same axioms, covering relations, functions, families of sets, and cardinal numbers.
- Enderton (2001) is a standard introduction to mathematical logic. Its first chapter collects the facts about sets, relations, and functions that the later chapters use.
- Barker-Plummer et al. (2011) teaches propositional and first-order logic, the language behind statements such as "for every set $A$ there exists…".

## References


Barker-Plummer, Dave, Jon Barwise, and John Etchemendy. 2011. *Language, Proof and Logic*. 2nd ed. CSLI Publications. https://www.amazon.com/dp/1575866323.

Devlin, Keith. 1993. *The Joy of Sets: Fundamentals of Contemporary Set Theory*. 2nd ed. Undergraduate Texts in Mathematics. Springer. https://doi.org/10.1007/978-1-4612-0903-4.

Enderton, Herbert B. 2001. *A Mathematical Introduction to Logic*. 2nd ed. Academic Press. https://doi.org/10.1016/C2009-0-22107-6.

Halmos, Paul R. 1974. *Naive Set Theory*. Undergraduate Texts in Mathematics. Springer. https://doi.org/10.1007/978-1-4757-1645-0.