Newton's Method

Last modified: 2026-11-10 23:22:44 (PST)

Source

This section is adapted from the Newton-Raphson and gradient descent parts of Zhou [1], used under the MIT License (see the license text in Linear Algebra). The source motivates the Newton step by minimizing a quadratic approximation; here that step is proved to be the exact minimizer of the quadratic model. In the source's zigzag example the starting value of the objective is $\frac{1}{2}(b^2 + b)$, not $\frac{1}{2}(1 + b^2)$ as the source writes; the ratio of successive values is unaffected. In the same example the source's formula for $\underset{\sim}{x}^{(1)}$ has $\underset{\sim}{x}^{(1)}$ on its right side where $\underset{\sim}{x}^{(0)}$ is meant. These parts of the source are not part of this section:

- the quadratic convergence theorem for Newton's method in general, and its remedies for instability (a positive definite substitute for the Hessian, and shortening the Newton step by a search along its direction)
- its remark that Hessians are expensive to compute
- the convergence-rate theorem for strongly convex functions
- gradient descent with momentum, and its optimal parameters
- Nesterov acceleration, and stochastic gradient descent
- its code and animations

Definition 0.1 (Iterative method)**.** An **iterative method** [2] starts from a point $\underset{\sim}{x}^{(0)}$ and computes $\underset{\sim}{x}^{(t+1)}$ from $\underset{\sim}{x}^{(t)}$ by a fixed rule, for $t = 0, 1, 2, \ldots$.

Definition 0.2 (Iterate)**.** In an iterative method (Definition 0.1), the points $\underset{\sim}{x}^{(t)}$ are its **iterates**.

Definition 0.3 (Error of an iterate)**.** For a point $\underset{\sim}{x}^*$ that an iterative method (Definition 0.1) is meant to reach, $\|\underset{\sim}{x}^{(t)} - \underset{\sim}{x}^*\|$ is the **error** of the iterate $\underset{\sim}{x}^{(t)}$ (Definition 0.2).

Definition 0.4 (Convergence of a sequence of vectors)**.** A sequence $\underset{\sim}{x}^{(0)}, \underset{\sim}{x}^{(1)}, \ldots$ in $\mathbb{R}^p$ **converges** to $x^* \in \mathbb{R}^p$ if the sequence of distances $\left\|\underset{\sim}{x}^{(t)} - \underset{\sim}{x}^*\right\|$ converges to $\widetilde{0}$. For $p = 1$, $\left\|x^{(t)} - x^*\right\| = \left|\widetilde{x^{(t)}} - x^*\right|$, and this is the convergence of a sequence of numbers to x^*.

Example 0.1 (Halving a vector)**.** The iterative method $\underset{\sim}{x}^{(t+1)} = \frac{1}{2}\,\underset{\sim}{x}^{(t)}$ from $\underset{\sim}{x}^{(0)} = (4,2)^\top$ has iterates $(4,2)^\top, (2,1)^\top, (1,0.5)^\top, \ldots$, that is, $\underset{\sim}{x}^{(t)} = 2^{-t}\,(4,2)^\top$. With $\underset{\sim}{x}^* = \underset{\sim}{0}$, the error of $\underset{\sim}{x}^{(t)}$ is

$$\begin{aligned}\left\|2^{-t}\,(4,2)^\top - \underset{\sim}{0}\right\| &= 2^{-t}\left\|(4,2)^\top\right\| && \text{(norm of a multiple)}\\ &= 2^{-t}\sqrt{20}, && (4^2 + 2^2 = 20)\end{aligned}$$

which converges to 0, so the iterates converge to $\underset{\sim}{0}$.

Definition 0.5 (Newton's method for minimization)**.** Let $f : \mathbb{R}^p \to \mathbb{R}$ have first and second partial derivatives, and let $\underset{\sim}{x}^{(0)}$ be a starting point. **Newton's method** [3] computes, for $t = 0, 1, 2, \ldots$, as long as $\mathbf{H}_f(\underset{\sim}{x}^{(t)})$ is invertible,

$$\underset{\sim}{x}^{(t+1)} \overset{\text{def}}{=} \underset{\sim}{x}^{(t)} - \left[\mathbf{H}_f(\underset{\sim}{x}^{(t)})\right]^{-1} \frac{\partial}{\partial \underset{\sim}{x}} f(\underset{\sim}{x}^{(t)}).$$

Definition 0.6 (Newton step)**.** In Newton's method (Definition 0.5), the vector $-\left[\mathbf{H}_f(\underset{\sim}{x}^{(t)})\right]^{-1} \frac{\partial}{\partial \underset{\sim}{x}} f(\underset{\sim}{x}^{(t)})$ is the **Newton step** at $\underset{\sim}{x}^{(t)}$.

Example 0.2 (Newton's method for a cubic)**.** Let $f(x) = \frac{1}{3}x^3 - 4x$, so $f'(x) = x^2 - 4$ and $f''(x) = 2x$. For $x^{(t)} \neq 0$, one Newton step gives

$$\begin{aligned}x^{(t+1)} &= x^{(t)} - \frac{(x^{(t)})^2 - 4}{2x^{(t)}} && \text{(Newton's method with } p = 1)\\ &= \frac{2(x^{(t)})^2 - (x^{(t)})^2 + 4}{2x^{(t)}} && \text{(common denominator)}\\ &= \frac{1}{2}\left(x^{(t)} + \frac{4}{x^{(t)}}\right). && \text{(simplify)}\end{aligned}$$

From $x^{(0)} = 2.5$:

- $$\begin{aligned}x^{(1)} &= \tfrac{1}{2}(2.5 + 1.6)\\ &= 2.05;\end{aligned}$$

- $$x^{(2)} = \tfrac{1}{2}(2.05 + 1.951219\ldots) = 2.000610$$

  (to six decimals);

- $x^{(3)} = 2.0000001$ (to seven decimals).

The iterates (Definition 0.2) approach 2, the local minimizer of f (a stationary point, since $f'(2) = 0$, with $f''(2) = 4 > 0$; Theorem 7 in Critical Points and Optimality Conditions).

Definition 0.7 (Quadratic model, second-order Taylor polynomial)**.** Let f be as in Definition 0.5 and let $\underset{\sim}{x}$ be a point, with $g \stackrel{\text{def}}{=} \frac{\partial}{\partial \underset{\sim}{x}} f(\underset{\sim}{x})$ and $\mathbf{H} \stackrel{\text{def}}{=} \mathbf{H}_f(\underset{\sim}{x})$. The **quadratic model** of f at $\underset{\sim}{x}$ is the function of a step $\underset{\sim}{\delta} \in \mathbb{R}^p$

$$q(\underset{\sim}{\delta}) \stackrel{\text{def}}{=} f(\underset{\sim}{x}) + \underset{\sim}{g} \cdot \underset{\sim}{\delta} + \tfrac{1}{2}\, \underset{\sim}{\delta}^{\top} \mathbf{H} \underset{\sim}{\delta},$$

also called the **second-order Taylor polynomial** of f at $\underset{\sim}{x}$.

Example 0.3 (The model at $x = 2.5$)**.** For $f(x) = \frac{1}{3}x^3 - 4x$ at $x = 2.5$ (Example 0.2),

$$g = 2.5^2 - 4 = 2.25$$

and

$$H = [2 \cdot 2.5] = [5],$$

so $q(\delta) = f(2.5) + 2.25\,\delta + 2.5\,\delta^2$. For $f(\underset{\sim}{x}) = \frac{1}{2}(x_1^2 + 3x_2^2)$, whose Hessian is $\begin{bmatrix} 1 & 0 \\ 0 & 3 \end{bmatrix}$ at every point, the model at any $\underset{\sim}{x}$ equals $f(\underset{\sim}{x} + \underset{\sim}{\delta})$ exactly (Example 12 in Critical Points and Optimality Conditions with $\mathbf{S} = \begin{bmatrix} 1/2 & 0 \\ 0 & 3/2 \end{bmatrix}$).

ℹ Remark (The quadratic model and Taylor's theorem)

Remark 0.1 (The quadratic model and Taylor's theorem). When the second partial derivatives of f are continuous, Theorem 6 in Critical Points and Optimality Conditions says that $f(\underset{\sim}{x} + \underset{\sim}{\delta})$ equals the expression for $q(\underset{\sim}{\delta})$ in Definition 0.7 with the Hessian evaluated at a point between $\underset{\sim}{x}$ and $\underset{\sim}{x} + \underset{\sim}{\delta}$.

Theorem 0.1 (The Newton step minimizes the quadratic model)**.** *Let q be the quadratic model of f at $\underset{\sim}{x}$, with $\mathbf{H} = \mathbf{H}_f(\underset{\sim}{x})$ positive definite. Its unique minimizer is the Newton step $\underset{\sim}{\delta}^* = -\mathbf{H}^{-1}\underset{\sim}{g}$.*

Proof

Proof. $\mathbf{H}$ is symmetric, by the definition of positive definite, and invertible, so $\underset{\sim}{\delta}^*$ exists and $\mathbf{H}\underset{\sim}{\delta}^* = -\underset{\sim}{g}$. For any $\underset{\sim}{d} \in \mathbb{R}^p$, first

$$\begin{aligned} \underset{\sim}{d}^\top \mathbf{H}\underset{\sim}{\delta}^* &= -\underset{\sim}{d} \cdot \underset{\sim}{g} \quad (\mathbf{H}\underset{\sim}{\delta}^* = -\underset{\sim}{g}) \\ &= -\underset{\sim}{g} \cdot \underset{\sim}{d}, \quad (\text{a } 1 \times 1 \text{ matrix equals its transpose}) \end{aligned}$$

and $\underset{\sim}{\delta}^{*\top} \mathbf{H}\underset{\sim}{d} = \underset{\sim}{d}^\top \mathbf{H}\underset{\sim}{\delta}^*$ in the same way, using $\mathbf{H}^\top = \mathbf{H}$. Then

$$\begin{aligned} q(\underset{\sim}{\delta}^* + \underset{\sim}{d}) &= f(\underset{\sim}{x}) + \underset{\sim}{g} \cdot (\underset{\sim}{\delta}^* + \underset{\sim}{d}) + \tfrac{1}{2}\,(\underset{\sim}{\delta}^* + \underset{\sim}{d})^\top \mathbf{H}(\underset{\sim}{\delta}^* + \underset{\sim}{d}) && \text{(definition of } q) \\ &= f(\underset{\sim}{x}) + \underset{\sim}{g} \cdot \underset{\sim}{\delta}^* + \underset{\sim}{g} \cdot \underset{\sim}{d} + \tfrac{1}{2}\big(\underset{\sim}{\delta}^{*\top} \mathbf{H}\underset{\sim}{\delta}^* + \underset{\sim}{\delta}^{*\top} \mathbf{H}\underset{\sim}{d} + \underset{\sim}{d}^\top \mathbf{H}\underset{\sim}{\delta}^* + \underset{\sim}{d}^\top \mathbf{H}\underset{\sim}{d}\big) && \text{(distribute)} \\ &= f(\underset{\sim}{x}) + \underset{\sim}{g} \cdot \underset{\sim}{\delta}^* + \underset{\sim}{g} \cdot \underset{\sim}{d} + \tfrac{1}{2}\,\underset{\sim}{\delta}^{*\top} \mathbf{H}\underset{\sim}{\delta}^* - \underset{\sim}{g} \cdot \underset{\sim}{d} + \tfrac{1}{2}\,\underset{\sim}{d}^\top \mathbf{H}\underset{\sim}{d} && \text{(both cross terms equal } -\underset{\sim}{g} \cdot \underset{\sim}{d}) \\ &= f(\underset{\sim}{x}) + \underset{\sim}{g} \cdot \underset{\sim}{\delta}^* + \tfrac{1}{2}\,\underset{\sim}{\delta}^{*\top} \mathbf{H}\underset{\sim}{\delta}^* + \tfrac{1}{2}\,\underset{\sim}{d}^\top \mathbf{H}\underset{\sim}{d} && \text{(cancel } \underset{\sim}{g} \cdot \underset{\sim}{d}) \\ &= q(\underset{\sim}{\delta}^*) + \tfrac{1}{2}\,\underset{\sim}{d}^\top \mathbf{H}\underset{\sim}{d}, && \text{(definition of } q) \end{aligned}$$

which is greater than $q(\underset{\sim}{\delta}^*)$ for every $\underset{\sim}{d} \neq \underset{\sim}{0}_{p\times 1}$, since $\mathbf{H}$ is positive definite. □

Example 0.4 (The Newton step at $x = 2.5$)**.** For the model $q(\delta) = f(2.5) + 2.25\,\delta + 2.5\,\delta^2$ of Example 0.3, $H = [5]$ is positive definite, so by Theorem 0.1 its minimizer is

$$\begin{aligned} \delta^* &= -H^{-1} g \\ &= -2.25/5 \\ &= -0.45. \end{aligned}$$

Setting the derivative $2.25 + 5\delta$ to 0 gives the same δ^*, and


$$\begin{aligned} 2.5 - 0.45 &= 2.05 \\ &= x^{(1)} \end{aligned}$$

(Example 0.2).

Definition 0.8 (Linear approximation of the gradient)**.** Let f be as in Newton's method and let $\underset{\sim}{x}$ be a point. The **linear approximation of the gradient** of f at $\underset{\sim}{x}$ is the

R

```r
nm_x_nm <- seq(0.5, 4, length.out = 301)
nm_f_nm <- nm_x_nm^3 / 3 - 4 * nm_x_nm
nm_f0_nm <- 2.5^3 / 3 - 4 * 2.5
nm_q_nm <- nm_f0_nm + 2.25 * (nm_x_nm - 2.5) + 2.5 * (nm_x_nm - 2.5)^2
nm_q_step_nm <- nm_f0_nm + 2.25 * (2.05 - 2.5) + 2.5 * (2.05 - 2.5)^2
nm_f_min_nm <- 2^3 / 3 - 4 * 2
nm_curve_nm <- tibble::tibble(x = nm_x_nm, y = nm_f_nm)
nm_model_nm <- tibble::tibble(x = nm_x_nm, y = nm_q_nm)
nm_start_nm <- tibble::tibble(x = 2.5, y = nm_f0_nm)
nm_step_nm <- tibble::tibble(x = 2.05, y = nm_q_step_nm)
nm_min_nm <- tibble::tibble(x = 2, y = nm_f_min_nm)
nm_aes_nm <- ggplot2::aes(x, y)
nm_ylim_nm <- c(-8, 0)
ggplot2::ggplot(nm_curve_nm, nm_aes_nm) +
  ggplot2::geom_line(colour = "#0072B2") +
  ggplot2::geom_line(
    data = nm_model_nm, colour = "#D55E00", linetype = "dashed"
  ) +
  ggplot2::geom_point(data = nm_start_nm, colour = "#D55E00", size = 3) +
  ggplot2::geom_point(
    data = nm_step_nm, colour = "#D55E00", size = 3, shape = 17
  ) +
  ggplot2::geom_point(
    data = nm_min_nm, colour = "#0072B2", size = 3, shape = 15
  ) +
  ggplot2::coord_cartesian(ylim = nm_ylim_nm) +
  ggplot2::labs(x = "x", y = "value")
```



Python

$p \times 1$ vector-valued function $\underset{\sim}{g}_q$ of a step $\underset{\sim}{\delta} \in \mathbb{R}^p$ whose jth entry, for $j = 1, \ldots, p$, is the function whose graph is the tangent plane to the graph of the partial derivative $\frac{\partial}{\partial x_j} f$ at $\underset{\sim}{x}$, evaluated at $\underset{\sim}{x} + \underset{\sim}{\delta}$:

$$\left[\underset{\sim}{g}_q(\underset{\sim}{\delta})\right]_j \stackrel{\text{def}}{=} \frac{\partial}{\partial x_j} f(\underset{\sim}{x}) + \left(\frac{\partial}{\partial \underset{\sim}{x}} \left(\frac{\partial}{\partial x_j} f(\underset{\sim}{x})\right)\right)^\top \underset{\sim}{\delta}. \tag{1}$$


Example 0.5 (The linear approximation of the derivative of a cubic at $x = -2.5$)**.** For $f(x) = \frac{1}{3}x^3 - 4x$, with $f'(x) = x^2 - 4$ and $f''(x) = 2x$ (Example 0.2), $p = 1$, so the linear approximation of the gradient at $x = -2.5$ is the tangent line to f' there:

$$\begin{aligned} \ell(\delta) &= f'(-2.5) + f''(-2.5)\,\delta && \text{(Equation 1 with } p = 1) \\ &= ((-2.5)^2 - 4) + 2(-2.5)\,\delta && \text{(substitute } f' \text{ and } f'') \\ &= 2.25 - 5\delta. && \text{(arithmetic)} \end{aligned} \tag{2}$$

Figure 2 shows f' and ℓ.

R

```
gla_x <- seq(-3, -1.5, length.out = 301)
gla_deriv <- gla_x^2 - 4
gla_line <- 2.25 - 5 * (gla_x + 2.5)
gla_curve <- tibble::tibble(x = gla_x, y = gla_deriv)
gla_tangent <- tibble::tibble(x = gla_x, y = gla_line)
gla_start <- tibble::tibble(x = -2.5, y = 2.25)
gla_step <- tibble::tibble(x = -2.05, y = 0)
gla_root <- tibble::tibble(x = -2, y = 0)
gla_ylim <- c(-2, 5)
gla_curve |>
  ggplot2::ggplot() +
  ggplot2::aes(x = x, y = y) +
  ggplot2::geom_hline(yintercept = 0) +
  ggplot2::geom_line(colour = "#0072B2") +
  ggplot2::geom_line(
    data = gla_tangent, colour = "#D55E00", linetype = "dashed"
  ) +
  ggplot2::geom_point(data = gla_start, colour = "#D55E00", size = 3) +
  ggplot2::geom_point(
    data = gla_step, colour = "#D55E00", size = 3, shape = 17
  ) +
  ggplot2::geom_point(
    data = gla_root, colour = "#0072B2", size = 3, shape = 15
  ) +
  ggplot2::coord_cartesian(ylim = gla_ylim) +
  ggplot2::labs(x = "x", y = "derivative")
```



Example 0.6 (The tangent line to f is not the linear approximation of the gradient)**.** For the cubic $f(x) = \frac{1}{3}x^3 - 4x$ at $x = -2.5$ (Example 0.5), the tangent line to f itself is

$$\begin{aligned} f(-2.5) + f'(-2.5)\,\delta &= \left(\tfrac{1}{3}(-2.5)^3 - 4(-2.5)\right) + \left((-2.5)^2 - 4\right)\delta && \text{(substitute } f \text{ and } f'\text{)} \\ &= \tfrac{115}{24} + 2.25\,\delta. && \text{(arithmetic)} \end{aligned} \tag{3}$$

It is not the linear approximation of the gradient: it approximates f, not f'. Its root, $\delta = -\frac{115}{54}$, is a step toward a zero of f, not toward a stationary point of f.

Lemma 0.1 (The linear approximation of the gradient uses the Hessian)**.** *Let f, $\underset{\sim}{x}$, the gradient $\underset{\sim}{g}$ and the Hessian $\mathbf{H}$ be as in the quadratic model, and let the second partial derivatives of f be continuous at every point of an open ball around $\underset{\sim}{x}$. Then the linear approximation of the gradient of f at $\underset{\sim}{x}$ is the gradient plus the Hessian times the step:*

$$\underset{\sim}{g}_q(\underset{\sim}{\delta}) = \underset{\sim}{g} + \mathbf{H}\underset{\sim}{\delta}. \tag{4}$$

Proof

Proof. For each i and j, the ith entry of $\frac{\partial}{\partial \underset{\sim}{x}}\left(\frac{\partial}{\partial x_j} f(\underset{\sim}{x})\right)$ is $\frac{\partial}{\partial x_i}\left(\frac{\partial}{\partial x_j} f(\underset{\sim}{x})\right)$, which is $[\mathbf{H}]_{ij}$ by the definitions of the Hessian and of the derivative of a vector-valued function. So, for each j,

$$\begin{aligned} \left[\underset{\sim}{g}_q(\underset{\sim}{\delta})\right]_j &= \frac{\partial}{\partial x_j} f(\underset{\sim}{x}) + \sum_{i=1}^{p} [\mathbf{H}]_{ij}\,\delta_i && \text{(Equation 1, as a sum)} \\ &= [\underset{\sim}{g}]_j + \sum_{i=1}^{p} [\mathbf{H}]_{ij}\,\delta_i && \text{(definition of } \underset{\sim}{g}\text{)} \\ &= [\underset{\sim}{g}]_j + \sum_{i=1}^{p} [\mathbf{H}]_{ji}\,\delta_i && (\mathbf{H} \text{ is symmetric}) \\ &= [\underset{\sim}{g}]_j + [\mathbf{H}\underset{\sim}{\delta}]_j && \text{(matrix-vector product)} \\ &= [\underset{\sim}{g} + \mathbf{H}\underset{\sim}{\delta}]_j. && \text{(vector addition)} \end{aligned} \tag{5}$$

The third line uses the symmetry of the Hessian, whose hypothesis the lemma assumes. □

Example 0.7 (A gradient that equals its linear approximation)**.** For $f(\underset{\sim}{x}) = \frac{1}{2}(x_1^2 + 3x_2^2)$, the second function of Example 0.3, the gradient is $\underset{\sim}{g} = (x_1,\ 3x_2)^\top$, and the Hessian is

$\begin{bmatrix} 1 & 0 \\ 0 & 3 \end{bmatrix}$ at every point. By Lemma 0.1, the linear approximation of the gradient at $\underset{\sim}{x}$ is

$$\begin{aligned} \underset{\sim}{g}_q(\underset{\sim}{\delta}) &= \begin{bmatrix} x_1 \\ 3x_2 \end{bmatrix} + \begin{bmatrix} 1 & 0 \\ 0 & 3 \end{bmatrix} \begin{bmatrix} \delta_1 \\ \delta_2 \end{bmatrix} && \text{(Equation 4)} \\ &= \begin{bmatrix} x_1 \\ 3x_2 \end{bmatrix} + \begin{bmatrix} \delta_1 \\ 3\delta_2 \end{bmatrix} && \text{(matrix-vector product)} \\ &= \begin{bmatrix} x_1 + \delta_1 \\ 3x_2 + 3\delta_2 \end{bmatrix} && \text{(vector addition)} \\ &= \begin{bmatrix} x_1 + \delta_1 \\ 3(x_2 + \delta_2) \end{bmatrix}, && \text{(factor out 3)} \end{aligned} \tag{6}$$

which is the gradient of f at $\underset{\sim}{x} + \underset{\sim}{\delta}$: the gradient of this f is linear, so its linear approximation is exact.

Theorem 0.2 (The Newton step is the root of the linear approximation of the gradient)**.** *Let f, $\underset{\sim}{x}$, g and the Hessian $\mathbf{H}$ be as in Lemma 0.1, with $\mathbf{H}$ invertible. The only step $\underset{\sim}{\delta}$ at which the linear approximation of the gradient is zero is the Newton step:*

$$\underset{\sim}{g}_q(\underset{\sim}{\delta}) = \underset{\sim}{0}_{p\times 1} \iff \underset{\sim}{\delta} = -\mathbf{H}^{-1}\underset{\sim}{g}. \tag{7}$$

Proof

Proof. By Lemma 0.1, $\underset{\sim}{g}_q(\underset{\sim}{\delta}) = \underset{\sim}{0}_{p\times 1}$ exactly when

$$\begin{aligned} \underset{\sim}{0}_{p\times 1} &= \underset{\sim}{g} + \mathbf{H}\underset{\sim}{\delta} && \text{(Equation 4)} \\ -\underset{\sim}{g} &= \mathbf{H}\underset{\sim}{\delta} && \text{(subtract } \underset{\sim}{g} \text{ from both sides)} \\ -\mathbf{H}^{-1}\underset{\sim}{g} &= \underset{\sim}{\delta}. && \text{(multiply both sides by } \mathbf{H}^{-1} \text{ on the left)} \end{aligned} \tag{8}$$

Each line implies the next and, because $\mathbf{H}$ is invertible, the one before, so $\underset{\sim}{\delta} = -\mathbf{H}^{-1}\underset{\sim}{g}$ is the only solution. □

Example 0.8 (The Newton step at $x = -2.5$ as a root)**.** At $x = -2.5$, the linear approximation of the gradient of the cubic $f(x) = \frac{1}{3}x^3 - 4x$ is $\ell(\delta) = 2.25 - 5\delta$ (Example 0.5). Setting it to 0,

$$\begin{aligned} 0 &= 2.25 - 5\delta && \text{(Equation 2)} \\ 5\delta &= 2.25 && \text{(add } 5\delta \text{ to both sides)} \\ \delta &= 0.45, && \text{(divide both sides by 5)} \end{aligned} \tag{9}$$

so the next iterate is $-2.5 + 0.45 = -2.05$, the point where the tangent line in Figure 2

crosses 0. The Newton's method iteration $x^{(t+1)} = \frac{1}{2}\left(x^{(t)} + 4/x^{(t)}\right)$ of Example 0.2 gives the same iterate. Here $H = [-5]$ is not positive definite, so Theorem 0.1 does not describe this step, but Theorem 0.2 does.

Remark (Two ways to read the Newton step)

Remark 0.2 (Two ways to read the Newton step). Theorem 0.1 reads the Newton step as the minimizer of the quadratic model, which needs the Hessian $\mathbf{H}$ to be positive definite. Theorem 0.2 reads it as the root of the linear approximation of the gradient, which needs only $\mathbf{H}$ invertible. Newton's method solves $\frac{\partial}{\partial \underset{\sim}{x}} f(\underset{\sim}{x}) = \underset{\sim}{0}_{p\times 1}$ by replacing the gradient with its linear approximation at each iterate, so it looks for a stationary point of any kind, not only a minimizer.

Definition 0.9 (Quadratic convergence)**.** A sequence $\underset{\sim}{x}^{(0)}, \underset{\sim}{x}^{(1)}, \ldots$ in $\mathbb{R}^p$ that converges to $\underset{\sim}{x}^*$ **converges quadratically** [4] if there is a constant C with

$$\left\|\underset{\sim}{x}^{(t+1)} - \underset{\sim}{x}^*\right\| \le C\left\|\underset{\sim}{x}^{(t)} - \underset{\sim}{x}^*\right\|^2 \qquad \text{for every } t.$$

Example 0.9 (Halving the error is not quadratic convergence)**.** $x^{(t)} = 2^{-t}$ converges to 0, but it does not converge quadratically: the bound would need $2^{-(t+1)} \le C\,(2^{-t})^2 = C\,2^{-2t}$, that is, $2^{t-1} \le C$, which fails once $t > 1 + \log_2 C$. Each step only halves the error.

Example 0.10 (The error squares at each step)**.** For the iteration $x^{(t+1)} = \frac{1}{2}(x^{(t)} + 4/x^{(t)})$ of Example 0.2,

$$\begin{aligned} x^{(t+1)} - 2 &= \frac{1}{2}\left(x^{(t)} + \frac{4}{x^{(t)}}\right) - 2 && \text{(the iteration)} \\ &= \frac{(x^{(t)})^2 + 4 - 4x^{(t)}}{2x^{(t)}} && \text{(common denominator)} \\ &= \frac{(x^{(t)} - 2)^2}{2x^{(t)}}. && \text{(the numerator is a perfect square)} \end{aligned}$$


So near 2 the error is roughly squared and divided by 4 at each step: the errors are 0.05, then 0.00061, then 0.000000093, and $0.05^2/(2 \cdot 2.05) \approx 0.00061$. The iterates converge quadratically (Definition 0.9): $C = \frac{1}{2}$ works for every $x^{(t)} \ge 1$, since then $\frac{1}{2x^{(t)}} \le \frac{1}{2}$, and every iterate from 2.5 is at least 2, since the identity makes $x^{(t+1)} - 2 \ge 0$ whenever $x^{(t)} > 0$. Zhou [1] states such a bound for Newton's method in general; it holds from starting points close enough to a minimizer where the Hessian is positive definite, when near that minimizer the Hessian changes by at most a constant times the distance moved,

which is not proved here.

Example 0.11 (Newton's method can find a maximum)**.** From $x^{(0)} = -2.5$ the same iteration gives -2.05, -2.000610, -2.0000001, approaching -2. But -2 is not a minimizer of f: it is a stationary point of $-f$, whose second derivative at -2 is $-f''(-2) = 4 > 0$, so it is a strict local minimizer of $-f$ (Theorem 7 in Critical Points and Optimality Conditions), that is, f is strictly larger at -2 than at all nearby points. Newton's method looks only for a stationary point. In this example, the Hessian $H = f''(x) = 2x$ is negative at every iterate. If $H < 0$, then Theorem 0.1 does not apply, because it requires a positive definite Hessian. Instead, the model $q(\delta) = f + g\,\delta + \frac{1}{2}H\delta^2$ tends to $-\infty$ as $|\delta| \to \infty$, so it has no minimizer.
Figure 4 shows the cubic and its model.

R

```r
nc_x_nc <- numeric(4)
nc_x_nc[1] <- 2.5
for (nc_t_nc in 1:3) {
  nc_prev_nc <- nc_x_nc[nc_t_nc]
  nc_x_nc[nc_t_nc + 1] <- 0.5 * (nc_prev_nc + 4 / nc_prev_nc)
}
nc_steps_nc <- tibble::tibble(t = 0:3, error = nc_x_nc - 2)
ggplot2::ggplot(nc_steps_nc, ggplot2::aes(t, error)) +
  ggplot2::geom_line(colour = "#0072B2") +
  ggplot2::geom_point(colour = "#0072B2", size = 3) +
  ggplot2::scale_y_log10() +
  ggplot2::labs(x = "iteration t", y = "error of the iterate")
```



Python

```python
import matplotlib.pyplot as plt

nc_x_ax = [2.5]
for _ in range(3):
    nc_x_ax.append(0.5 * (nc_x_ax[-1] + 4 / nc_x_ax[-1]))
nc_err_ax = [x - 2 for x in nc_x_ax]
fig, ax = plt.subplots(figsize=(5, 3.5), layout="constrained")
ax.semilogy(range(4), nc_err_ax, "o-", color="#0072B2")
#> [<matplotlib.lines.Line2D object at 0x7f86c1515bd0>]
ax.set_xlabel("iteration t")
#> Text(0.5, 0, 'iteration t')
ax.set_ylabel("error of the iterate")
#> Text(0, 0.5, 'error of the iterate')
ax.set_xticks(range(4))
#> [<matplotlib.axis.XTick object at 0x7f86c14ef610>, <matplotlib.axis.XTick object at 0x7f86
plt.show();
```

R

```
nx_x_nx <- seq(-4, -0.5, length.out = 301)
nx_f_nx <- nx_x_nx^3 / 3 - 4 * nx_x_nx
nx_f0_nx <- (-2.5)^3 / 3 - 4 * (-2.5)
nx_dx_nx <- nx_x_nx + 2.5
nx_q_nx <- nx_f0_nx + 2.25 * nx_dx_nx - 2.5 * nx_dx_nx^2
nx_dstep_nx <- -2.05 + 2.5
nx_q_step_nx <- nx_f0_nx + 2.25 * nx_dstep_nx - 2.5 * nx_dstep_nx^2
nx_f_max_nx <- (-2)^3 / 3 - 4 * (-2)
nx_curve_nx <- tibble::tibble(x = nx_x_nx, y = nx_f_nx)
nx_model_nx <- tibble::tibble(x = nx_x_nx, y = nx_q_nx)
nx_start_nx <- tibble::tibble(x = -2.5, y = nx_f0_nx)
nx_step_nx <- tibble::tibble(x = -2.05, y = nx_q_step_nx)
nx_max_nx <- tibble::tibble(x = -2, y = nx_f_max_nx)
nx_aes_nx <- ggplot2::aes(x, y)
nx_ylim_nx <- c(-2, 6.5)
nx_curve_nx |>
  ggplot2::ggplot() +
  nx_aes_nx +
  ggplot2::geom_line(colour = "#0072B2") +
  ggplot2::geom_line(
    data = nx_model_nx, colour = "#D55E00", linetype = "dashed"
  ) +
  ggplot2::geom_point(data = nx_start_nx, colour = "#D55E00", size = 3) +
```

Theorem 0.3 (Newton's method minimizes a quadratic in one step)**.** *Let* $\mathbf{S}$ *be a* $p \times p$ *positive definite matrix,* $\underset{\sim}{c} \in \mathbb{R}^p$*, and* $f(\underset{\sim}{x}) = \frac{1}{2}\underset{\sim}{x}^\top \mathbf{S}\underset{\sim}{x} - \underset{\sim}{c} \cdot \underset{\sim}{x}$*. From any* $\underset{\sim}{x}^{(0)}$*, Newton's method (Definition 0.5) gives* $\underset{\sim}{x}^{(1)} = \mathbf{S}^{-1}\underset{\sim}{c}$*, the unique minimizer of* f*.*

Proof

Proof. Partial derivatives of a sum or of a constant multiple are the sum or multiple of the partial derivatives, so the derivative of a quadratic form applies to $\frac{1}{2}\underset{\sim}{x}^\top \mathbf{S}\underset{\sim}{x}$ and the derivative of a linear combination to $\underset{\sim}{c} \cdot \underset{\sim}{x}$:

$$\begin{aligned}\frac{\partial}{\partial \underset{\sim}{x}} f(\underset{\sim}{x}) &= \tfrac{1}{2}(2\mathbf{S}\underset{\sim}{x}) - \underset{\sim}{c} \\ &= \mathbf{S}\underset{\sim}{x} - \underset{\sim}{c};\end{aligned}$$

by the Hessian of a quadratic form with $\frac{1}{2}\mathbf{S}$ in place of $\mathbf{S}$, and because the gradient $-\underset{\sim}{c}$ of the linear term is constant, $\mathbf{H}_f(\underset{\sim}{x}) = \mathbf{S}$, which is invertible. So

$$\begin{aligned}\underset{\sim}{x}^{(1)} &= \underset{\sim}{x}^{(0)} - \mathbf{S}^{-1}(\mathbf{S}\underset{\sim}{x}^{(0)} - \underset{\sim}{c}) && \text{(definition of Newton's method)} \\ &= \underset{\sim}{x}^{(0)} - \mathbf{S}^{-1}\mathbf{S}\underset{\sim}{x}^{(0)} + \mathbf{S}^{-1}\underset{\sim}{c} && \text{(distribute)} \\ &= \underset{\sim}{x}^{(0)} - \underset{\sim}{x}^{(0)} + \mathbf{S}^{-1}\underset{\sim}{c} && (\mathbf{S}^{-1}\mathbf{S} = \mathbf{I}_p) \\ &= \mathbf{S}^{-1}\underset{\sim}{c}. && \text{(cancel)}\end{aligned}$$

The gradient at $\mathbf{S}^{-1}\underset{\sim}{c}$ is $\mathbf{S}\mathbf{S}^{-1}\underset{\sim}{c} - \underset{\sim}{c} = \underset{\sim}{0}_{p\times 1}$, and f is strictly convex (Theorem 5 in Convexity in Several Variables, part 2), so $\mathbf{S}^{-1}\underset{\sim}{c}$ is a global minimizer (Corollary 1 in Convexity in Several Variables). It is the only one: a strictly convex function has at most one global minimizer, since the midpoint of two distinct global minimizers, which have equal values, would have a strictly smaller value (Definition 3 in Convexity in Several Variables with $t = \frac{1}{2}$). □


Example 0.12 (One step to the minimizer)**.** Let $f(\underset{\sim}{x}) = \frac{1}{2}(x_1^2 + 3x_2^2)$, so $\mathbf{S} = \begin{bmatrix} 1 & 0 \\ 0 & 3 \end{bmatrix}$ and $\underset{\sim}{c} = \underset{\sim}{0}_{2\times 1}$. From $\underset{\sim}{x}^{(0)} = (3,1)^\top$ the gradient is $\nabla f(\underset{\sim}{x}^{(0)}) = \mathbf{S}\underset{\sim}{x}^{(0)} = (3,3)^\top$, and $\mathbf{S}^{-1} = \begin{bmatrix} 1 & 0 \\ 0 & 1/3 \end{bmatrix}$, so the Newton step is $-\mathbf{S}^{-1}(3,3)^\top = -(3,1)^\top$ and

$$\begin{aligned}\underset{\sim}{x}^{(1)} &= (3,1)^\top - (3,1)^\top \\ &= \underset{\sim}{0}_{2\times 1},\end{aligned}$$

the minimizer.
Figure 5 shows the step on the level curves of f.

R

```
nq_x1_nq <- seq(-4, 4, length.out = 101)
nq_x2_nq <- seq(-2, 2, length.out = 101)
nq_grid_nq <- tidyr::expand_grid(x1 = nq_x1_nq, x2 = nq_x2_nq)
nq_grid_nq$z <- (nq_grid_nq$x1^2 + 3 * nq_grid_nq$x2^2) / 2
nq_newton_nq <- tibble::tibble(x = 3, y = 1, xend = 0, yend = 0)
nq_grad_nq <- tibble::tibble(x = 3, y = 1, xend = 0.764, yend = -1.236)
nq_start_nq <- tibble::tibble(x1 = 3, x2 = 1)
nq_head_nq <- grid::unit(0.15, "inches")
nq_arrow_nq <- grid::arrow(length = nq_head_nq)
nq_aes_nq <- ggplot2::aes(x1, x2)
nq_z_aes_nq <- ggplot2::aes(z = z)
nq_seg_aes_nq <- ggplot2::aes(x = x, y = y, xend = xend, yend = yend)
nq_grid_nq |>
  ggplot2::ggplot() +
  nq_aes_nq +
  ggplot2::geom_contour(nq_z_aes_nq, colour = "#0072B2", bins = 10) +
  ggplot2::geom_segment(
    nq_seg_aes_nq,
    data = nq_grad_nq,
    colour = "grey50",
    arrow = nq_arrow_nq
  ) +
  ggplot2::geom_segment(
    nq_seg_aes_nq,
    data = nq_newton_nq,
    colour = "#D55E00",
    arrow = nq_arrow_nq
  ) +
  ggplot2::geom_point(data = nq_start_nq, size = 2) +
  ggplot2::coord_fixed() +
  ggplot2::labs(x = "x1", y = "x2")
```



> Figure 6 shows the same function as a surface.


R

(This 3D surface is interactive; see the HTML or slide version of this page to view it.)

Python

```
import matplotlib.pyplot as plt
import numpy as np

nq_x1_3d, nq_x2_3d = np.meshgrid(np.linspace(-4, 4, 61),
                                 np.linspace(-2, 2, 61))
nq_z_3d = (nq_x1_3d**2 + 3 * nq_x2_3d**2) / 2
fig = plt.figure(figsize=(6, 4.5))
ax = fig.add_subplot(projection="3d")
ax.plot_surface(nq_x1_3d, nq_x2_3d, nq_z_3d, color="#0072B2", alpha=0.6)
#> <mpl_toolkits.mplot3d.art3d.Poly3DCollection object at 0x7f86c15ae900>
ax.scatter([3], [1], [6], color="#D55E00", label="start")
#> <mpl_toolkits.mplot3d.art3d.Path3DCollection object at 0x7f86c15aea50>
ax.scatter([0], [0], [0], color="black", label="minimizer")
#> <mpl_toolkits.mplot3d.art3d.Path3DCollection object at 0x7f86bf01b250>
ax.set_xlabel("x1")
#> Text(0.5, 0, 'x1')
ax.set_ylabel("x2")
#> Text(0.5, 0.5, 'x2')
ax.set_zlabel("f(x)")
#> Text(0.5, 0, 'f(x)')
ax.legend(loc="upper right")
#> <matplotlib.legend.Legend object at 0x7f86c15aeba0>
plt.show();
```



[1] H. Zhou, "Optimization and multivariate calculus." Lecture notes for Biostat 216, Mathematical Methods for Biostatistics, University of California, Los Angeles, 2024. Accessed: Oct. 05, 2026. [Online]. Available: https://ucla-biostat-216.github.io/2024fall/slides/13-optim/13-optim.html

[2] Wikipedia contributors, "Iterative method — Wikipedia, the free encyclopedia." 2026. Available: https://en.wikipedia.org/wiki/Iterative_method

[3] Wikipedia contributors, "Newton's method in optimization — Wikipedia, the free encyclopedia." 2026. Available: https://en.wikipedia.org/wiki/Newton%27s_method_in_optimization

[4] Wikipedia contributors, "Rate of convergence — Wikipedia, the free encyclopedia." 2026. Available: https://en.wikipedia.org/wiki/Rate_of_convergence