

Math prerequisites

Last modified: 2026-09-28 01:20:09 (PDT)

Math is not just a way of calculating numerical answers; it is a way of thinking, using clear definitions for concepts and rigorous logic to organize our thoughts and back up our assertions.

Cheng (2025)

These lecture notes use:

- mathematical notation
- algebra
- precalculus
- univariate calculus
- linear algebra
- vector calculus

Some key results are listed here.

Notation

Mathematical notation is not standardized. This section states the conventions these notes use, and the alternatives you may meet in other sources.

Natural numbers

Exercise 0.1 (Which numbers are natural?). List the elements of the set $\{n \in \mathbb{N} : n < 3\}$. Is your answer the same in every textbook?

Solution

Solution 0.1. The answer depends on whether the source counts 0 as a natural number:

- if $\mathbb{N}$ starts at 0, the set is $\{0, 1, 2\}$;
- if $\mathbb{N}$ starts at 1, the set is $\{1, 2\}$.

Both conventions are in common use, so the answer is not the same in every textbook.

Definition 0.1 (Natural numbers (our convention)). In these notes, the **natural numbers** are the positive integers:

$$\mathbb{N} \stackrel{\text{def}}{=} \{1, 2, 3, \dots\}$$

Definition 0.2 (Non-negative integers). The **non-negative integers** are the natural numbers (Definition 0.1) together with 0:

$$\mathbb{N}_0 \stackrel{\text{def}}{=} \{0, 1, 2, 3, \dots\} = \mathbb{N} \cup \{0\}$$

Example 0.1 (Natural numbers in a data analysis).

- Observation indices start at 1, so we write $i \in \{1, \dots, n\}$ with $n \in \mathbb{N}$.
- A count outcome, such as the number of hospital visits in a year, can be 0, so its support is $\mathbb{N}_0$, not $\mathbb{N}$.

i Other conventions for the natural numbers

Other sources may define $\mathbb{N}$ differently, so check each source's definition before reading its formulas:

- Many sources include 0 in $\mathbb{N}$. The international standard ISO 80000-2 defines $\mathbb{N}$ to include 0, continuing the earlier standard ISO 31-11 (1978).
- Other sources start $\mathbb{N}$ at 1, as these notes do.
- To remove the ambiguity, some sources write $\mathbb{N}_1$ or $\mathbb{Z}^+$ for $\{1, 2, 3, \dots\}$, and $\mathbb{N}_0$ or $\mathbb{Z}^{0+}$ for $\{0, 1, 2, \dots\}$.

- The words vary too. “Positive integers” ($\{1, 2, 3, \dots\}$) and “non-negative integers” ($\{0, 1, 2, \dots\}$) are unambiguous. “Whole numbers” usually includes 0, but can also mean all of the integers, negative ones included. “Counting numbers” usually starts at 1, but some sources include 0.
- Some older texts write J for the natural numbers.

When a formula’s meaning depends on whether 0 is included, we write the set out explicitly, for example $\{0, 1, 2, \dots\}$ or $\{1, \dots, n\}$.

Source: [Wikipedia](#), “[Natural number](#)”, “[Terminology and notation](#)” and “[Zero as natural number](#)”, which cites ISO 80000-2:2019 and the textbooks using each notation.

Algebra

Elementary Algebra

Mastery of [Elementary Algebra](#) (a.k.a. “College Algebra”) is a prerequisite for calculus, which is in turn a prerequisite for most statistics and data science courses. Nevertheless, each year, some students are still uncomfortable with algebraic manipulations of mathematical formulas. Therefore, I include this section as a quick reference.

Equalities

Theorem 0.1 (Equalities are transitive). *If $a = b$ and $b = c$, then $a = c$*

Theorem 0.2 (Substituting equivalent expressions). *If $a = b$, then for any function $f(x)$, $f(a) = f(b)$*

Inequalities

Theorem 0.3 (Adding to both sides of an inequality). *If $a < b$, then $a + c < b + c$*

Theorem 0.4 (negating both sides of an inequality). *If $a < b$, then: $-a > -b$*

Theorem 0.5 (Multiplying both sides of an inequality by a nonnegative number). *If $a < b$ and $c \geq 0$, then $ca < cb$.*

Theorem 0.6 (Negation is multiplication by -1).

$$-a = (-1) * a$$

Infimum and supremum

Definition 0.1 (Infimum (greatest lower bound)). The **infimum** of a nonempty set $A \subseteq \mathbb{R}$, written $\inf A$, is the greatest real number m satisfying $m \leq a$ for all $a \in A$:

$$\inf A \stackrel{\text{def}}{=} \max_{t \in \mathbb{R}} \{t : \forall a \in A, a \geq t\}$$

If the infimum belongs to A , it equals the minimum: $\inf A = \min A$.

Example 0.1 (Numerical examples of infimum).

- $\inf\{1, 2, 3\} = 1$, since 1 is the smallest element.
- $\inf(0.5, 1] = 0.5 = \min[0.5, 1]$: for intervals open below, the infimum equals the minimum of the corresponding closed-below interval, even though $0.5 \notin (0.5, 1]$. More generally, $\inf(c, b] = \min[c, b] = c$ for any $c < b$.
- $\inf\{t \geq 0 : t > 0.5\} = 0.5$, even though 0.5 itself is not in the set.

Definition 0.2 (Supremum (least upper bound)). The **supremum** of a nonempty set $A \subseteq \mathbb{R}$, written $\sup A$, is the smallest real number M satisfying $M \geq a$ for all $a \in A$:

$$\sup A \stackrel{\text{def}}{=} \min_{t \in \mathbb{R}} \{t : \forall a \in A, a \leq t\}$$

If the supremum belongs to A , it equals the maximum: $\sup A = \max A$.

Example 0.2 (Numerical examples of supremum).

- $\sup\{1, 2, 3\} = 3$, since 3 is the largest element.
- $\sup\{t \geq 0 : t < 0.5\} = 0.5$, even though 0.5 itself is not in the set.

Sums

Theorem 0.7 (adding zero changes nothing).

$$a + 0 = a$$

Theorem 0.8 (Sums are symmetric).

$$a + b = b + a$$

Theorem 0.9 (Sums are associative).

When summing three or more terms, the order in which you sum them does not matter:

$$(a + b) + c = a + (b + c)$$

Products

Theorem 0.10 (Multiplying by 1 changes nothing).

$$a \times 1 = a$$

Theorem 0.11 (Products are symmetric).

$$a \times b = b \times a$$

Theorem 0.12 (Products are associative).

$$(a \times b) \times c = a \times (b \times c)$$

Division

Theorem 0.13 (Division can be written as a product).

$$\frac{a}{b} = a \times \frac{1}{b}$$

Sums and products together

Theorem 0.14 (Multiplication is distributive).

$$a(b + c) = ab + ac$$

Quotients

Definition 0.3 (Quotients, fractions, rates).

A **quotient** (also called a *fraction* or *rate*) is a division of one quantity by another:

$$\frac{a}{b}$$

In epidemiology, rates typically have a quantity involving time or population in the denominator.

c.f. [https://en.wikipedia.org/wiki/Rate_\(mathematics\)](https://en.wikipedia.org/wiki/Rate_(mathematics))

Definition 0.4 (Ratios). A **ratio** is a quotient in which the numerator and denominator are measured using the same unit scales.

c.f. <https://en.wikipedia.org/wiki/Ratio>

Definition 0.5 (Proportion). In statistics, a **proportion** typically means a ratio where the numerator represents a subset of the denominator.

See https://en.wikipedia.org/wiki/Population_proportion.

See also [https://en.wikipedia.org/wiki/Proportion_\(mathematics\)](https://en.wikipedia.org/wiki/Proportion_(mathematics)) for other meanings.

Definition 0.6 (Proportional). Two functions $f(x)$ and $g(x)$ are **proportional** if their ratio $\frac{f(x)}{g(x)}$ does not depend on x . (c.f. [https://en.wikipedia.org/wiki/Proportionality_\(mathematics\)](https://en.wikipedia.org/wiki/Proportionality_(mathematics)))

Additional reference for elementary algebra: https://en.wikipedia.org/wiki/Population_proportion#Mathematical_definition

Exponentials and Logarithms

Theorem 0.15 (logarithm of a product is the sum of the logs of the factors).

$$\log a \cdot b = \log a + \log b$$

Corollary 0.1 (logarithm of a quotient).

The logarithm of a quotient is equal to the difference of the logs of the factors:

$$\log \frac{a}{b} = \log a - \log b$$

Theorem 0.16 (logarithm of an exponential function).

$$\log\{a^b\} = b \cdot \log\{a\}$$

Theorem 0.17 (exponential of a sum).

The exponential of a sum is equal to the product of the exponentials of the addends:

$$\exp\{a + b\} = \exp\{a\} \cdot \exp\{b\}$$

Corollary 0.2 (exponential of a difference).

The exponential of a difference is equal to the quotient of the exponentials of the addends:

$$\exp\{a - b\} = \frac{\exp\{a\}}{\exp\{b\}}$$

Theorem 0.18 (exponential of a product).

$$a^{bc} = (a^b)^c = (a^c)^b$$

Corollary 0.3 (natural exponential of a product).

$$\exp\{ab\} = (\exp\{a\})^b = (\exp\{b\})^a$$

Exercise 0.1. For $a \geq 0$, $b, c \in \mathbb{R}$, When does $(a^b)^c = a^{(b^c)}$?

Solution

Solution 0.1. Short answer: rarely (that's all you need to know for this course).

Long answer:

If $(a^b)^c = a^{(b^c)}$, then since $(a^b)^c = a^{bc}$, we have:

$$a^{bc} = a^{(b^c)}$$

$$\log\{a^{bc}\} = \log\{a^{(b^c)}\}$$

$$bc \cdot \log\{a\} = b^c \cdot \log\{a\} \tag{1}$$

Equation 1 holds in each of the following cases:

1. $bc = b^c$ (see Exercise 0.2).
2. $a = 1$ (i.e., $\log\{a\} = 0$).
3. $a = 0$ (i.e., $\log\{a\} = -\infty$) and $\text{sign}\{bc\} = \text{sign}\{b^c\}$.

In particular, when $a = 0$ and $c = 0$, $bc = 0$ and $b^c = 1$ (for any $b \in \mathbb{R}$), so $\text{sign}\{bc\} \neq \text{sign}\{b^c\}$, and $(a^b)^c \neq a^{(b^c)}$:

$$\begin{aligned} (a^b)^c &= (0^b)^0 \\ &= 1 \end{aligned}$$

$$\begin{aligned} a^{(b^c)} &= 0^{(b^0)} \\ &= 0^1 \\ &= 0 \end{aligned}$$

Exercise 0.2. For $b, c \in \mathbb{R}$, when does $b^c = bc$?

Solution

Solution 0.2. $bc = b^c$ in each of the following cases:

1. $c = 1$.
2. $b = 0$ and $c > 0$.
3. $b = \exp\left\{\frac{\log c}{c-1}\right\}$ (for $c \geq 0$).

See the red contours in Figure 2 for a visualization.

(This 3D surface is interactive; see the HTML or slide version of this page to view it.)

Figure 1: Graph of $b * c$ and b^c

(This 3D surface is interactive; see the HTML or slide version of this page to view it.)

Figure 2: **Graph of $b^c - b * c$.** Red contour lines show where $b^c = b * c$.

Theorem 0.19 ($\exp\{\}$ and $\log\{\}$ are mutual inverses).

$$\exp\{\log\{a\}\} = \log\{\exp\{a\}\} = a$$

Derivatives

Theorem 0.1 (Constant rule).

$$\frac{\partial}{\partial x} c = 0$$

Theorem 0.2 (Power rule). *If a is constant with respect to x , then:*

$$\frac{\partial}{\partial x} ay = a \frac{\partial x}{\partial y}$$

Theorem 0.3 (Power rule).

$$\frac{\partial}{\partial x} x^q = qx^{q-1}$$

Theorem 0.4 (Derivative of natural logarithm).

$$\log' \{x\} = \frac{1}{x} = x^{-1}$$

Theorem 0.5 (derivative of exponential).

$$\exp' \{x\} = \exp \{x\}$$

Theorem 0.6 (Product rule).

$$(ab)' = ab' + ba'$$

Theorem 0.7 (Quotient rule).

$$(a/b)' = a'/b - (a/b^2)b'$$

Theorem 0.8 (Chain rule).

$$\begin{aligned} \frac{\partial a}{\partial c} &= \frac{\partial a}{\partial b} \frac{\partial b}{\partial c} \\ &= \frac{\partial b}{\partial c} \frac{\partial a}{\partial b} \end{aligned}$$

or in *Euler/Lagrange notation*:

$$(f(g(x)))' = g'(x)f'(g(x))$$

Corollary 0.1 (Chain rule for logarithms).

$$\frac{\partial}{\partial x} \log f(x) = \frac{f'(x)}{f(x)}$$

i Proof

Proof. Apply Theorem 0.8 and Theorem 0.4. □

Integration

Integration is the inverse operation of differentiation: it recovers a function from its derivative and accumulates quantities such as areas, totals, and probabilities. We begin with antiderivatives, then state basic integration rules, and conclude with the Fundamental Theorem of Calculus and a worked example from probability.

Antiderivatives

Definition 0.1 (Antiderivative). A function F is an **antiderivative** of f on an interval I if:

$$\frac{\partial}{\partial x} F(x) = f(x), \quad \forall x \in I$$

(Larson and Edwards 2018, sec. 4.1, pp. 248–249)

Definition 0.2 (Indefinite integral). The **indefinite integral** of f is the family of all antiderivatives (Definition 0.1) of f :

$$\int f(x) dx = F(x) + C$$

where F is any one antiderivative of f and C is an arbitrary constant of integration.
(Larson and Edwards 2018, sec. 4.1, pp. 248–249)


Example 0.1 (Antiderivative of a power function). For $f(x) = x^2$, an antiderivative is $F(x) = \frac{x^3}{3}$, since $\frac{\partial}{\partial x} \frac{x^3}{3} = x^2 = f(x)$. Adding any constant C gives another antiderivative; for example, with $C = 7$, $F(x) = \frac{x^3}{3} + 7$ also satisfies $F'(x) = x^2$, since adding a constant does not change the derivative. See Figure 3.

```

ggplot2::ggplot() +
  ggplot2::geom_function(fun = \(x) {
    ggplot2::labs(x = "x", y = expression(f(x)))
    ggplot2::theme_minimal()
  })
  x_seq <- seq(x_lim[1], x_lim[2], length.out = 200)
  df <- do.call(rbind, lapply(C_vals, \(C) {
    data.frame(
      x = x_seq,
      y = x_seq^3 / 3 + C,
      C = factor(C)
    )
  }))


ggplot2::ggplot(df, ggplot2::aes(x = x, y = y, color = C)) +
  ggplot2::geom_line(linewidth = 0.8) +
  ggplot2::labs(x = "x", y = expression(F(x)), color = C) +
  ggplot2::theme_minimal()

```



(a)

(b) The function $f(x) = x^2$.



(c)

(d) Family of antiderivatives $F(x) = x^3/3 + C$.

Figure 3: The function $f(x) = x^2$ and five antiderivatives $F(x) = x^3/3 + C$ for $C \in \{-2, -1, 0, 1, 2\}$. Each antiderivative has the same derivative f ; they differ only by a vertical shift.

Theorem 0.1 (Basic integration rules). *Each antiderivative in the table is defined only up to an arbitrary constant C (see Definition 0.1); the table omits $+C$ from every row for brevity.*

Function $f(x)$	Antiderivative $F(x)$	Condition
c	cx	—
x^n	$\frac{x^{n+1}}{n+1}$	$n \neq -1$

$\frac{1}{x}$	$\ln x $	$x \neq 0$
e^x	e^x	(self-antiderivative)
e^{cx}	$\frac{1}{c}e^{cx}$	$c \neq 0$
$c \cdot f(x)$	$c \cdot F(x)$	—
$f(x) + g(x)$	$F(x) + G(x)$	—

The first two rows and the bottom two rows (linearity) are from (Larson and Edwards 2018, sec. 4.1, p. 250 “Basic Integration Rules”); $1/x$ is from (Larson and Edwards 2018, sec. 5.2, Theorem 5.5, p. 324); e^x and e^{cx} are from (Larson and Edwards 2018, sec. 5.4, Theorem 5.12, p. 346).

Example 0.2 (Antiderivative of $3x^2 - 1$). By the power rule ($n = 2$) and linearity from Theorem 0.1:

$$\int (3x^2 - 1) dx = 3 \cdot \frac{x^3}{3} - x + C = x^3 - x + C.$$

Verify by differentiating: $\frac{\partial}{\partial x}(x^3 - x + C) = 3x^2 - 1 = f(x)$, as required.

Regularity Conditions

Definition 0.3 (Differentiable function). A function f is **differentiable at** $x = c$ if the limit

$$f'(c) \stackrel{\text{def}}{=} \lim_{h \rightarrow 0} \frac{f(c+h) - f(c)}{h}$$

exists and is finite.

(Larson and Edwards 2018, sec. 2.1, p. 100)

Definition 0.4 (Differentiable on an interval). A function f is **differentiable on** an interval if it is differentiable (Definition 0.3) at every interior point of the interval; at a closed endpoint, the appropriate one-sided derivative is used.

(Larson and Edwards 2018, sec. 2.1, p. 100)

Definition 0.5 (Continuous function). A function f is **continuous at** $x = c$ if all three conditions hold:

1. $f(c)$ is defined,
2. $\lim_{x \rightarrow c} f(x)$ exists, and
3. $\lim_{x \rightarrow c} f(x) = f(c)$.

(Larson and Edwards 2018, sec. 1.4, p. 73)

Definition 0.6 (Continuous on a closed interval). A function f is **continuous on** a closed interval $[a, b]$ if it is continuous (Definition 0.5) at every point of $[a, b]$.

(Larson and Edwards 2018, sec. 1.4, p. 73)

Definition 0.7 (Riemann integral). For a bounded function f on $[a, b]$, split $[a, b]$ into n equal-width subintervals of width $\Delta x \stackrel{\text{def}}{=} (b - a)/n$, and let x_i^* be any point in the i -th subinterval. The **Riemann integral** of f over $[a, b]$ is the limit

$$\int_a^b f(x) dx \stackrel{\text{def}}{=} \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i^*) \Delta x,$$

when that limit exists.

(Larson and Edwards 2018, sec. 4.3, p. 272)

Definition 0.8 (Riemann integrable). A bounded function f is **Riemann integrable on** $[a, b]$ if its Riemann integral (Definition 0.7) over $[a, b]$ exists and is finite.

(Larson and Edwards 2018, sec. 4.3, p. 272)

Definition 0.9 (General Riemann integrability). More generally, using partitions $\mathcal{P}$ of arbitrary mesh — subintervals of varying widths Δx_i — a bounded function f is **Riemann integrable in the general sense** on $[a, b]$ if

$$\int_a^b f(x) dx \stackrel{\text{def}}{=} \lim_{\|\mathcal{P}\| \rightarrow 0} \sum_{i=1}^n f(x_i^*) \Delta x_i$$

exists and is finite, where $\|\mathcal{P}\| = \max_i \Delta x_i$ is the mesh of the partition.

Theorem 0.2 (Equivalence of Riemann sum formulations). *For continuous f on a closed interval $[a, b]$, the equal-width Riemann sum (Definition 0.8) and the arbitrary-mesh Riemann sum (Definition 0.9) give the same value (Rudin 1976, chap. 6). Applied statistics courses mostly use the equal-width form in Definition 0.8.*

Before stating the Fundamental Theorem of Calculus, we record two prerequisite results. The FTC requires the integrand f to be continuous, and the following two theorems establish


where continuity comes from (differentiability $\Rightarrow$ continuity) and what it buys us (continuity $\Rightarrow$ integrability).

Theorem 0.3 (Differentiability implies continuity). *If f is differentiable at $x = c$, then f is continuous at $x = c$.
(Larson and Edwards 2018, Theorem 2.1, p. 106)*

Example 0.3 (Differentiable, hence continuous: $x^3 - x$). $f(x) = x^3 - x$ is differentiable everywhere (with derivative $f'(x) = 3x^2 - 1$), so by Theorem 0.3 it is continuous everywhere.

Example 0.4 (Continuous but not differentiable: $|x|$). The absolute-value function $f(x) = |x|$ is continuous at $x = 0$ ($\lim_{x \rightarrow 0} |x| = 0 = |0|$), but it is not differentiable at $x = 0$: the left-derivative is -1 and the right-derivative is $+1$. This counterexample shows that the converse of Theorem 0.3 fails: continuity does not imply differentiability. See Figure 4.


```
ggplot2::ggplot() +
  ggplot2::geom_function(fun = abs, xlim = c(-2, 2), linewidth = 1) +
  ggplot2::geom_point(ggplot2::aes(x = 0, y = 0), size = 3) +
  ggplot2::labs(x = "x", y = expression(f(x) == abs(x))) +
  ggplot2::theme_minimal()
```



(a)

Figure 4: $f(x) = |x|$ has a sharp corner at $x = 0$ (not differentiable there) but is continuous everywhere: no gaps or jumps.

Theorem 0.4 (Continuity implies integrability). *If f is continuous on the closed interval $[a, b]$, then f is integrable on $[a, b]$ (i.e., the Riemann integral $\int_a^b f(x) dx$ exists and is finite).*

(Larson and Edwards 2018, Theorem 4.4, p. 272)

Example 0.5 (Continuous, hence integrable: polynomials). Every polynomial is continuous on $\mathbb{R}$, so by Theorem 0.4 every polynomial is integrable on every closed interval $[a, b]$.

Example 0.6 (Integrable but not continuous: a step function). Let $f(x) = 0$ for $x < \frac{1}{2}$ and $f(x) = 1$ for $x \geq \frac{1}{2}$. Then f is discontinuous at $x = \frac{1}{2}$, but it is integrable on $[0, 1]$:

$$\int_0^1 f(x) \, dx = \int_0^{1/2} 0 \, dx + \int_{1/2}^1 1 \, dx = 0 + \frac{1}{2} = \frac{1}{2}.$$


This counterexample shows that the converse of Theorem 0.4 fails: integrability does not imply continuity. See Figure 5.

```

step_df <- data.frame(
  x = c(0, 0.5, 0.5, 1),
  y = c(0, 0, 1, 1),
  segment = c("left", "left", "right", "right")
)

ggplot2::ggplot() +
  ggplot2::geom_rect(
    ggplot2::aes(xmin = 0.5, xmax = 1, ymin = 0, ymax = 1),
    fill = "steelblue", alpha = 0.3
  ) +
  ggplot2::geom_line(
    data = step_df,
    ggplot2::aes(x = x, y = y, group = segment),
    linewidth = 1
  ) +
  ggplot2::geom_point(ggplot2::aes(x = 0.5, y = 0), shape = 1, size = 3) +
  ggplot2::geom_point(ggplot2::aes(x = 0.5, y = 1), shape = 16, size = 3) +
  ggplot2::scale_x_continuous(
    breaks = c(0, 0.5, 1),
    labels = c("0", "1/2", "1")
  ) +
  ggplot2::scale_y_continuous(limits = c(-0.1, 1.2)) +
  ggplot2::labs(x = "x", y = "f(x)") +
  ggplot2::theme_minimal()

```



(19)

Figure 5: Step function: $f(x) = 0$ on $[0, \frac{1}{2})$ (open circle at the jump) and $f(x) = 1$ on $[\frac{1}{2}, 1]$ (filled circle). The shaded rectangle has area $\frac{1}{2}$, matching the integral computed in Example 0.6.

Together, Theorem 0.3 and Theorem 0.4 establish the chain:

$$\text{differentiable} \Rightarrow \text{continuous} \Rightarrow \text{integrable}$$

Example 0.4 and Example 0.6 show that neither implication reverses in general.

Fundamental Theorem of Calculus

Theorem 0.5 (Fundamental Theorem of Calculus). *Let f be a continuous function on a closed interval $[a, b]$.*

Part 1 (Derivative of an integral). *Define $F(x) = \int_a^x f(t) dt$ for $x \in [a, b]$. Then F is differentiable and:*

$$\frac{\partial}{\partial x} \int_a^x f(t) dt = f(x) \quad (2)$$

Note

Continuity on all of $[a, b]$ is a sufficient condition. More generally, Part 1 holds at any individual point x where f is integrable on $[a, b]$ (see Definition 0.8) and continuous at x (see Definition 0.5), even if f has jump discontinuities elsewhere (Rudin 1976, Theorem 6.20, p. 133).

(Larson and Edwards 2018, Theorem 4.11, p. 288)

Part 2 (Evaluation theorem). *The F here may be any antiderivative of f — not just the accumulation function from Part 1. If F is an antiderivative of f on $[a, b]$ (i.e., $\frac{\partial}{\partial x} F(x) = f(x)$ for all $x \in [a, b]$), then:*

$$\int_a^b f(x) dx = F(b) - F(a) \quad (3)$$

Equivalently, with b replaced by a variable upper limit x , integrating the derivative of F recovers the net change in F :

$$\int_a^x F'(t) dt = F(x) - F(a) \quad (4)$$

or equivalently in Leibniz notation:

$$\int_a^x \frac{dF}{dt} dt = F(x) - F(a)$$

(Banner 2007, chap. 18; Larson and Edwards 2018, Theorem 4.9, p. 282)

The two parts of the FTC together express that **differentiation and integration are inverse operations**:

- Part 1: differentiating the integral of f recovers f (Equation 2).
- Part 2: the integral of f over $[a, b]$ equals the difference of any antiderivative's values at the endpoints (Equation 3), which rearranges to “integrating the derivative of F recovers the net change in F ” (Equation 4).

The standard form of the FTC assumes f is continuous on $[a, b]$; continuity is *sufficient* but not strictly necessary (see the callout note inside Theorem 0.5 for the more general statement). Since differentiability implies continuity (Theorem 0.3), the FTC applies in particular whenever f is differentiable — a common situation in applied statistics.

Example 0.7 (FTC Part 1 visualized: accumulation function for $f(t) = 2t$). Take $f(t) = 2t$ on $[0, 2]$. The accumulation function from 0 is

$$F(x) \stackrel{\text{def}}{=} \int_0^x 2t \, dt = [t^2]_{t=0}^{t=x} = x^2 - 0^2 = x^2,$$

so $F(x) = x^2$, and indeed $F'(x) = 2x = f(x)$, as Theorem 0.5 Part 1 predicts. Figure 6 shows the integrand on the left (shaded area equals $F(x)$ at each x) and the accumulation function $F(x) = x^2$ on the right (its slope at x equals $f(x) = 2x$).


```

ggplot2::ggplot() +
  ggplot2::geom_area(
    data = data.frame(t = seq(0, x_f),
      ggplot2::aes(x = t, y = 2 * t),
      fill = "steelblue", alpha = 0.4 )
  ) +
  ggplot2::geom_function(fun = \(t) 2 * t) +
  ggplot2::geom_vline(
    data = data.frame(x = x_marks),
    ggplot2::aes(xintercept = x, color = factor(x),
      linetype = "dashed", linewidth = 0.8)
  ) +
  ggplot2::labs(x = "t", y = "f(t) = 2t") +
  ggplot2::theme_minimal() +
  ggplot2::theme(legend.position = "bottom")

slope_df <- data.frame(
  x = x_marks,
  Fx = x_marks^2,
  slope = 2 * x_marks
)


ggplot2::ggplot() +
  ggplot2::geom_function(fun = \(x) x^2, xlim = c(0, 2.5)) +
  ggplot2::geom_point(
    data = slope_df,
    ggplot2::aes(x = x, y = Fx, color = factor(x),
      size = 3)
  ) +
  ggplot2::geom_segment(
    data = slope_df,
    ggplot2::aes(
      x = x - 0.3, xend = x + 0.3,
      y = Fx - 0.3 * slope, yend = Fx + 0.3 * slope,
      color = factor(x)
    ),
    linewidth = 0.8
  ) +
  ggplot2::labs(x = "x", y = expression(F(x) == x^2)) +
  ggplot2::theme_minimal() +
  ggplot2::theme(legend.position = "bottom")

```



(a)

(b) $f(t) = 2t$; shaded area equals $F(1.5) = 2.25$; vertical lines mark $x \in \{1, 1.5, 2\}$.



(c)

(d) $F(x) = x^2$; tangent slope at each marked x equals $f(x) = 2x$.

Figure 6: Left: $f(t) = 2t$; the shaded area $\int_0^{1.5} 2t \, dt = F(1.5) = 2.25$; vertical lines mark $x \in \{1, 1.5, 2\}$. Right: $F(x) = x^2$; for each marked x , the tangent slope equals $f(x) = 2x$.

Example 0.8 (CDF and PDF of the exponential distribution). In what follows, f denotes the PDF and F the CDF — the same letters as the antiderivative pair in Definition 0.1, because the FTC will show F is exactly an antiderivative of f .

For the exponential distribution with rate parameter $\lambda > 0$, the probability density function (PDF) is (Kleinbaum and Klein 2012, sec. II, p. 295, “Survival and Hazard Functions for Selected Distributions”):

$$f(t) = \lambda e^{-\lambda t}, \quad t \geq 0$$

FTC Part 2 gives the cumulative distribution function (CDF) from the PDF. Apply the e^{cx} rule from Theorem 0.1 with $c = -\lambda$ to antidifferentiate the integrand:

$$\begin{aligned} F(t) &= \int_0^t \lambda e^{-\lambda u} du \\ &= \left[\lambda \cdot \frac{1}{-\lambda} e^{-\lambda u} \right]_{u=0}^{u=t} \\ &= [(-1)e^{-\lambda u}]_{u=0}^{u=t} \\ &= [-e^{-\lambda u}]_{u=0}^{u=t} \\ &= -e^{-\lambda t} - (-e^0) \\ &= -e^{-\lambda t} - (-1) \\ &= 1 - e^{-\lambda t} \end{aligned}$$

FTC Part 1 recovers the PDF from the CDF:


$$\begin{aligned} \frac{\partial}{\partial t} F(t) &= \frac{\partial}{\partial t} (1 - e^{-\lambda t}) \\ &= 0 - (-\lambda) e^{-\lambda t} \\ &= \lambda e^{-\lambda t} \\ &= f(t) \end{aligned}$$

For a concrete instance: with $\lambda = 1$ (standard exponential), the probability that $T \leq 2$ is:

$$F(2) = 1 - e^{-1 \cdot 2} = 1 - e^{-2} \approx 1 - 0.135 = 0.865$$

See Figure 7.


```
ggplot2::ggplot() +
  ggplot2::geom_area(
    data = data.frame(t = seq(0, t_max),
                      y = lambda * exp(-lambda * t)),
    fill = "steelblue", alpha = 0.4
  ) +
  ggplot2::geom_function(
    fun = \(t) lambda * exp(-lambda * t),
    xlim = c(0, t_max), linewidth = 1
  ) +
  ggplot2::labs(x = "t", y = "f(t)")
ggplot2::theme_minimal()
```



(a)

(b) PDF with $\lambda = 1$; shaded area equals $F(2) \approx 0.865$.

```
ggplot2::ggplot() +
  ggplot2::geom_function(
    fun = \(t) 1 - exp(-lambda * t),
    xlim = c(0, t_max), linewidth = 1
  ) +
  ggplot2::geom_point(
    ggplot2::aes(x = t_focus, y = F_at_focus),
    size = 3, color = "steelblue"
  ) +
  ggplot2::geom_segment(
    ggplot2::aes(x = t_focus, xend = t_focus, y = 0, yend = F_at_focus),
    linetype = "dashed", color = "steelblue"
  ) +
  ggplot2::geom_segment(
    ggplot2::aes(x = 0, xend = t_focus, y = F_at_focus, yend = F_at_focus),
    linetype = "dashed", color = "steelblue"
  ) +
  ggplot2::labs(x = "t", y = "F(t)") +
  ggplot2::theme_minimal()
```



(c)

(d) CDF with $\lambda = 1$; point marks $F(2) \approx 0.865$.

Figure 7: Exponential distribution with $\lambda = 1$. Left: the PDF $f(t) = \lambda e^{-\lambda t}$; the shaded area under the curve from 0 to 2 equals $F(2) \approx 0.865$. Right: the CDF $F(t) = 1 - e^{-\lambda t}$; the dashed lines mark the value $F(2)$ computed via FTC Part 2.

Double Integrals

The **Fubini–Tonelli theorem** states conditions under which the order of integration in a double integral can be exchanged. We state two versions: the Riemann version (Theorem 0.1) is what applied courses usually use for double integrals of continuous functions on simple regions; the σ -finite measure-theoretic version (Theorem 0.2) is included to make the [joint-distribution form](#) corollary in the probability chapter of *Regression Models for Epidemiology* follow from a stated theorem rather than from an aside.

Theorem 0.1 (Fubini’s theorem (Riemann version)). *Let f be **continuous** on a plane region $R \subseteq \mathbb{R}^2$.*

1. **Vertically simple region.** *If R is defined by $a \leq x \leq b$ and $g_1(x) \leq y \leq g_2(x)$, where g_1 and g_2 are continuous on $[a, b]$, then*

$$\iint_R f(x, y) dA = \int_a^b \int_{g_1(x)}^{g_2(x)} f(x, y) dy dx.$$

2. **Horizontally simple region.** *If R is defined by $c \leq y \leq d$ and $h_1(y) \leq x \leq h_2(y)$, where h_1 and h_2 are continuous on $[c, d]$, then*

$$\iint_R f(x, y) dA = \int_c^d \int_{h_1(y)}^{h_2(y)} f(x, y) dx dy.$$

When R can be described both ways, the two iterated integrals are equal — so the order of integration can be exchanged.

(Larson and Edwards 2018, Theorem 14.2, p. 982)

Example 0.1 (Changing the order of integration for a non-rectangular region). Adapted from (Larson and Edwards 2018, sec. 14.2, Example 4, pp. 984–985).

Let X and Y be independent Uniform(0, 1) random variables, with joint density $f(x, y) = 1$ on the unit square $[0, 1]^2$. Define the function $g(x, y) = e^{-x^2} \mathbb{1}(y \leq x)$, and compute its expectation $E[g(X, Y)]$.

Because the joint density equals 1 on $[0, 1]^2$, this expectation is the double integral of g over the unit square:

$$E[g(X, Y)] = \iint_{[0, 1]^2} g(x, y) dA.$$

The indicator factor $\mathbb{1}(y \leq x)$ equals 1 on the triangular region where $y \leq x$ and 0 elsewhere, so only that region, namely $D = \{(x, y) : x \in [0, 1], y \in [0, x]\}$ (Figure 8),

contributes, and there $g(x, y) = e^{-x^2}$:

$$\mathbb{E}[g(X, Y)] = \iint_D e^{-x^2} dA.$$

```

region <- data.frame(x = c(0, 1, 1), y = c(0, 0, 1))
ggplot2::ggplot(region, ggplot2::aes(x = x, y = y)) +
  ggplot2::geom_polygon(
    fill = "steelblue", alpha = 0.4, color = "black", linewidth = 0.7
  ) +
  ggplot2::annotate(
    "text", x = 0.65, y = 0.28, label = "D", size = 8, fontface = "italic"
  ) +
  ggplot2::annotate(
    "text", x = 0.36, y = 0.52, label = "y == x",
    parse = TRUE, angle = 45
  ) +
  ggplot2::annotate(
    "text", x = 1.08, y = 0.5, label = "x == 1",
    parse = TRUE, angle = 90, hjust = 0.5
  ) +
  ggplot2::annotate(
    "text", x = 0.5, y = -0.1, label = "y == 0", parse = TRUE
  ) +
  ggplot2::scale_x_continuous(breaks = c(0, 0.5, 1), limits = c(-0.1, 1.3)) +
  ggplot2::scale_y_continuous(breaks = c(0, 0.5, 1), limits = c(-0.2, 1.2)) +
  ggplot2::coord_equal() +
  ggplot2::labs(x = "x", y = "y") +
  ggplot2::theme_minimal()

```




Figure 8: Triangular integration region $D = \{(x, y) : x \in [0, 1], y \in [0, x]\}$, bounded below by $y = 0$, above-left by $y = x$, and on the right by $x = 1$.

Order $dx\,dy$ is intractable. Re-describing D as $D = \{(x, y) : y \in [0, 1], x \in [y, 1]\}$, the inner integral is

$$\int_y^1 e^{-x^2} dx,$$

which has no elementary antiderivative.

Order $dy\,dx$ works. Applying Theorem 0.1 Part 1 (e^{-x^2} is continuous and D is the vertically simple region $x \in [0, 1], y \in [0, x]$):

$$\begin{aligned} \iint_D e^{-x^2} dA &= \int_0^1 \int_0^x e^{-x^2} dy dx \\ &= \int_0^1 e^{-x^2} \left(\int_0^x dy \right) dx \\ &= \int_0^1 x e^{-x^2} dx \\ &= \left[-\frac{1}{2} e^{-x^2} \right]_0^1 \\ &= -\frac{1}{2} (e^{-1} - 1) \\ &= \frac{e - 1}{2e} \\ &\approx 0.316 \end{aligned}$$

The solid whose volume equals this integral is shown in Figure 9.

(This 3D surface is interactive; see the HTML or slide version of this page to view it.)

Figure 9: Surface $z = e^{-x^2}$ over the region $D = \{(x, y) : x \in [0, 1], y \in [0, x]\}$. The surface depends only on x (constant in y), so for each x the inner integral over $y \in [0, x]$ contributes $x \cdot e^{-x^2}$.

Example 0.2 (When conditions fail: a counterexample). The conditions in Theorem 0.1 are not merely technical — when they fail, iterated integrals can exist yet disagree. Let

$$f(x, y) = \frac{x^2 - y^2}{(x^2 + y^2)^2}$$

on the unit square $R = [0, 1] \times [0, 1]$. Strictly, f is defined on $R \setminus \{(0, 0)\}$: the denominator vanishes at the origin, so f is undefined there (we return to this point in the condition check).

Integrating y first, then x :

Using $\frac{\partial}{\partial y} \frac{y}{x^2 + y^2} = \frac{x^2 - y^2}{(x^2 + y^2)^2}$:

$$\begin{aligned} \int_0^1 \int_0^1 f(x, y) dy dx &= \int_0^1 \left[\frac{y}{x^2 + y^2} \right]_{y=0}^{y=1} dx \\ &= \int_0^1 \frac{1}{x^2 + 1} dx \\ &= [\arctan(x)]_0^1 \\ &= \frac{\pi}{4} \end{aligned}$$

Integrating x first, then y :

Using $\frac{\partial}{\partial x} \left(-\frac{x}{x^2 + y^2} \right) = \frac{x^2 - y^2}{(x^2 + y^2)^2}$:

$$\begin{aligned} \int_0^1 \int_0^1 f(x, y) dx dy &= \int_0^1 \left[-\frac{x}{x^2 + y^2} \right]_{x=0}^{x=1} dy \\ &= \int_0^1 \left(-\frac{1}{1 + y^2} \right) dy \\ &= -[\arctan(y)]_0^1 \\ &= -\frac{\pi}{4} \end{aligned}$$

Conclusion: $\frac{\pi}{4} \neq -\frac{\pi}{4}$, so the two iterated integrals are unequal. Theorem 0.1 does not apply here.

Why Theorem 0.1's condition fails: Theorem 0.1 requires f to be **continuous** on R . The denominator $(x^2 + y^2)^2$ vanishes at the origin $(0, 0) \in R$, so f is *not even defined* there — let alone continuous — and the theorem does not apply.

(Wikipedia contributors 2024)

The surface, and the singularity at the origin responsible for the failure, are shown in Figure 10.

(This 3D surface is interactive; see the HTML or slide version of this page to view it.)

Figure 10: Surface $f(x, y) = (x^2 - y^2)/(x^2 + y^2)^2$ on $[0, 1]^2$, sampled away from the origin and clipped to $[-50, 50]$ for display. The function diverges to $+\infty$ along the x -axis (red ridge, $f > 0$ when $|x| > |y|$) and to $-\infty$ along the y -axis (blue ridge, $f < 0$ when $|y| > |x|$). The singularity at $(0, 0)$ is why f is not continuous on R and Theorem 0.1 does not apply.

Corollary 0.1 (Continuous functions on a rectangle (corollary of Theorem 0.1)). *If $f : [a, b] \times [c, d] \rightarrow \mathbb{R}$ is **continuous** on the closed bounded rectangle $[a, b] \times [c, d]$, then:*

$$\begin{aligned}\int_a^b \left(\int_c^d f(x, y) dy \right) dx &= \int_c^d \left(\int_a^b f(x, y) dx \right) dy \\ &= \iint_{[a,b] \times [c,d]} f(x, y) dx dy.\end{aligned}$$

(Larson and Edwards 2018, Theorem 14.2, p. 982)

i Proof

Proof. A closed bounded rectangle $[a, b] \times [c, d]$ is both vertically simple (with $g_1 \equiv c$, $g_2 \equiv d$) and horizontally simple (with $h_1 \equiv a$, $h_2 \equiv b$). Applying both parts of Theorem 0.1 to f on this rectangle gives the two iterated forms shown. $\square$

Example 0.3 (Evaluating a double integral on a rectangle). Structure adapted from (Larson and Edwards 2018, sec. 14.2, Example 2, pp. 982–983); the integrand $x^2 + y^2$ is original, chosen so the integral equals $E[g(X, Y)]$ for $g(x, y) = x^2 + y^2$. Let X and Y be independent Uniform(0, 1) random variables, with joint density $f(x, y) = 1$ on the unit square $R = \{(x, y) : x \in [0, 1], y \in [0, 1]\}$ (Figure 11). Define the function $g(x, y) = x^2 + y^2$, and compute its expectation $E[g(X, Y)]$. Because the joint density equals 1 on R , this expectation is the double integral of g over R :

$$E[g(X, Y)] = \iint_R (x^2 + y^2) dA.$$

```

ggplot2::ggplot() +
  ggplot2::annotate(
    "rect", xmin = 0, xmax = 1, ymin = 0, ymax = 1,
    fill = "steelblue", alpha = 0.4, color = "black", linewidth = 0.7
  ) +
  ggplot2::annotate(
    "text", x = 0.5, y = 0.5, label = "R", size = 8, fontface = "italic"
  ) +
  ggplot2::scale_x_continuous(breaks = c(0, 1), limits = c(-0.15, 1.25)) +
  ggplot2::scale_y_continuous(breaks = c(0, 1), limits = c(-0.15, 1.25)) +
  ggplot2::coord_equal() +
  ggplot2::labs(x = "x", y = "y") +
  ggplot2::theme_minimal()

```




Figure 11: Integration region $R = [0, 1]^2$, the unit square.

The integrand is continuous on R , so Corollary 0.1 applies and either order of integration yields the same value.

Integrating y first, then x :

$$\begin{aligned}
\mathbb{E}[g(X, Y)] &= \int_0^1 \int_0^1 (x^2 + y^2) dy dx \\
&= \int_0^1 \left[x^2 y + \frac{y^3}{3} \right]_0^1 dx \\
&= \int_0^1 \left(x^2 + \frac{1}{3} \right) dx \\
&= \left[\frac{x^3}{3} + \frac{x}{3} \right]_0^1 \\
&= \frac{2}{3}
\end{aligned}$$

Integrating x first, then y (verifying the order can be swapped):

$$\begin{aligned}
\int_0^1 \int_0^1 (x^2 + y^2) dx dy &= \int_0^1 \left[\frac{x^3}{3} + y^2 x \right]_0^1 dy \\
&= \int_0^1 \left(\frac{1}{3} + y^2 \right) dy \\
&= \left[\frac{y}{3} + \frac{y^3}{3} \right]_0^1 \\
&= \frac{2}{3}
\end{aligned}$$

Both orders give $\frac{2}{3}$, as Corollary 0.1 guarantees.

As a cross-check, linearity of expectation gives the same value: since $\mathbb{E}[X^2] = \int_0^1 x^2 dx = \frac{1}{3}$ for $X \sim \text{Uniform}(0, 1)$ (and likewise for Y),

$$\mathbb{E}[g(X, Y)] = \mathbb{E}[X^2 + Y^2] = \mathbb{E}[X^2] + \mathbb{E}[Y^2] = \frac{1}{3} + \frac{1}{3} = \frac{2}{3}.$$

The solid whose volume equals this integral is shown in Figure 12.

(This 3D surface is interactive; see the HTML or slide version of this page to view it.)

Figure 12: Surface $z = x^2 + y^2$ over the unit square $[0, 1]^2$. The double integral $\frac{2}{3}$ is the volume between this surface and the xy -plane, and equals $\mathbb{E}[X^2 + Y^2]$.

Theorem 0.2 (Fubini–Tonelli theorem (measure-theoretic form)). *Let $(\Omega_1, \mathcal{F}_1, \mu_1)$ and $(\Omega_2, \mathcal{F}_2, \mu_2)$ be **-finite** measure spaces, and let $f : \Omega_1 \times \Omega_2 \rightarrow \mathbb{R}$ be measurable with respect to the product σ -algebra $\mathcal{F}_1 \otimes \mathcal{F}_2$. If either*

*(a) $f \geq 0$ almost everywhere with respect to $\mu_1 \otimes \mu_2$ (**Tonelli's theorem**), or*

(b) $\int_{\Omega_1 \times \Omega_2} |f| d(\mu_1 \otimes \mu_2) < \infty$ (**Fubini's theorem**),

then both iterated integrals exist, agree with the double integral, and equal each other:

$$\begin{aligned} \int_{\Omega_1 \times \Omega_2} f d(\mu_1 \otimes \mu_2) &= \int_{\Omega_1} \left(\int_{\Omega_2} f(\omega_1, \omega_2) d\mu_2(\omega_2) \right) d\mu_1(\omega_1) \\ &= \int_{\Omega_2} \left(\int_{\Omega_1} f(\omega_1, \omega_2) d\mu_1(\omega_1) \right) d\mu_2(\omega_2). \end{aligned}$$

(Billingsley 1995, Theorem 18.3; Gut 2013, Theorem 9.1, p. 65; Fubini 1907; Wikipedia contributors 2024)

Applied courses rarely need the measure-theoretic generalization itself, but it is what justifies the [joint-distribution form](#) corollary in the probability chapter of *Regression Models for Epidemiology*: probability measures are finite (hence σ -finite), so the σ -finiteness condition is automatic. The integrability conditions (nonnegativity or absolute integrability) still need to be verified in each application.

Example 0.4 (Positive application of Theorem 0.2). Let X and Y be independent Exponential(1) random variables, with joint density $f(x, y) = e^{-(x+y)}$ for $x, y \geq 0$. Their distributions are probability measures on $[0, \infty)$, and probability measures are finite (hence σ -finite), so the product measure on $[0, \infty)^2$ satisfies the σ -finiteness condition of Theorem 0.2. Since $f(x, y) = e^{-(x+y)} \geq 0$, condition (a) (Tonelli theorem, nonnegativity) is also satisfied.

By Theorem 0.2, both iterated integrals exist and agree:

$$\begin{aligned} P(X \leq 1, Y \leq 1) &= \int_0^1 \int_0^1 e^{-(x+y)} dy dx \\ &= \int_0^1 e^{-x} \left(\int_0^1 e^{-y} dy \right) dx \\ &= \int_0^1 e^{-x} (1 - e^{-1}) dx \\ &= (1 - e^{-1}) [-e^{-x}]_0^1 \\ &= (1 - e^{-1})^2. \end{aligned}$$

The joint probability calculation also equals $\left(\int_0^1 e^{-x} dx \right)^2 = (1 - e^{-1})^2$, since $f(x, y) = e^{-x} \cdot e^{-y}$ factors as a product of independent densities. Both iterated integrals agree, as Theorem 0.2 guarantees when condition (a) holds.

Example 0.5 (When neither Fubini–Tonelli condition is satisfied). The same function $f(x, y) = (x^2 - y^2)/(x^2 + y^2)^2$ from Example 0.2 illustrates a case where neither condition of Theorem 0.2 is satisfied.

Why Theorem 0.2’s conditions fail: $\iint_R |f| dA = \infty$, which violates condition (b). Switching to polar coordinates (r, θ) near the origin, the integrand satisfies $|f(x, y)| = |x^2 - y^2|/(x^2 + y^2)^2 = |\cos 2\theta|/r^2$, so

$$\begin{aligned} \iint_R |f| dA &\geq \int_0^{\pi/2} \int_0^\epsilon \frac{|\cos 2\theta|}{r^2} r dr d\theta \\ &= \left(\int_0^{\pi/2} |\cos 2\theta| d\theta \right) \int_0^\epsilon \frac{dr}{r} \\ &= +\infty, \end{aligned}$$

since $\int_0^\epsilon dr/r$ diverges. Therefore $\iint_R |f| dA = \infty$, and condition (b) of Theorem 0.2 is not satisfied. (Condition (a) also fails: f takes both positive and negative values, so it is not nonnegative a.e.) The unequal iterated integrals from Example 0.2 are thus consistent with Theorem 0.2: the theorem simply does not apply.
(Wikipedia contributors 2024)

Linear Algebra

Vectors

Definition 0.1 (Column vector). A **column vector** of length p is an ordered list of p numbers, written vertically:

$$\tilde{x} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_p \end{bmatrix}$$

Column vectors are the default convention in these notes and in most statistics textbooks. They are also called $p \times 1$ *matrices*.

Definition 0.2 (Transpose). The **transpose** of a column vector $\tilde{x}$ is the row vector with the same sequence of entries, written horizontally:

$$\tilde{x}^\top \equiv \tilde{x}' \equiv [x_1, x_2, \dots, x_p]$$

The transpose operation converts a column vector to a row vector, or more generally, swaps the rows and columns of a matrix (Definition 0.10).

Special vectors

Definition 0.3 (Zero vector). The **zero vector** $\tilde{0}$ of length p has all entries equal to zero:

$$\tilde{0} = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \end{bmatrix}$$

The zero vector is the additive identity for vector addition: $\tilde{x} + \tilde{0} = \tilde{x}$ for any vector $\tilde{x}$ of the same length.

Definition 0.4 (Ones vector). The **ones vector** $\tilde{1}$ of length p has all entries equal to one:

$$\tilde{1} = \begin{bmatrix} 1 \\ 1 \\ \vdots \\ 1 \end{bmatrix}$$

The dot product $\tilde{1}^\top \tilde{x} = \tilde{1} \cdot \tilde{x} = \sum_{i=1}^p x_i$ is the sum of all entries of $\tilde{x}$.

Definition 0.5 (Indicator vector / standard basis vector). The j -th **indicator vector** (or *standard basis vector*) $\tilde{e}_j$ of length p has a 1 in position j and 0s elsewhere:

$$(\tilde{e}_j)_i = \begin{cases} 1 & \text{if } i = j \\ 0 & \text{if } i \neq j \end{cases} \quad \tilde{e}_j = \begin{bmatrix} 0 \\ \vdots \\ 0 \\ 1 \\ 0 \\ \vdots \\ 0 \end{bmatrix} \leftarrow \text{position } j$$

They are also called *unit vectors* or *standard basis vectors*.

Theorem 0.1 (Indicator vectors select entries). *For any vector $\tilde{x}$ of length p and any $j \in \{1, \dots, p\}$:*

$$\tilde{e}_j^\top \tilde{x} = x_j$$

i Proof

Proof. Writing the product componentwise:

$$\begin{aligned} \tilde{e}_j^\top \tilde{x} &= \sum_{i=1}^p (\tilde{e}_j)_i x_i \\ &= \sum_{i=1}^p \begin{cases} 1 \cdot x_i & \text{if } i = j \\ 0 \cdot x_i & \text{if } i \neq j \end{cases} \\ &= x_j \end{aligned}$$

□

Definition 0.6 (Dot product/linear combination/inner product). For any two real-valued vectors $\tilde{x} = (x_1, \dots, x_n)$ and $\tilde{y} = (y_1, \dots, y_n)$, the **dot-product** (also called the *linear combination* or *inner product*) of $\tilde{x}$ and $\tilde{y}$ is:

$$\tilde{x} \cdot \tilde{y} = \tilde{x}^\top \tilde{y} \stackrel{\text{def}}{=} \sum_{i=1}^n x_i y_i$$

i Note

See also the definitions in

- Dobson and Barnett (2018), §1.3 (equation 1.1, page 7)
- Kaplan (2022), chapter on vectors
- [wikipedia](#)

“Linear combination” can also refer to weighted sums of vectors, or in other words matrix-vector multiplication.

The dot-product has a different generalization for two matrices; see [wikipedia](#) for more.

Theorem 0.2 (Dot product is symmetric). *The dot product is symmetric:*

$$\tilde{x} \cdot \tilde{y} = \tilde{y} \cdot \tilde{x}$$

i Proof

Proof. Apply:

- Definition [0.6](#)
- symmetry of scalar multiplication
- Definition [0.6](#) again

□

Example 0.1 (Dot product as matrix multiplication). The dot product of two column vectors $\tilde{x}$ and $\tilde{\beta}$ can be written as a matrix product of the row vector $\tilde{x}^\top$ with the column vector $\tilde{\beta}$:

$$\begin{aligned}
\tilde{x} \cdot \tilde{\beta} &= \tilde{x}^\top \tilde{\beta} \\
&= [x_1, x_2, \dots, x_p] \begin{bmatrix} \beta_1 \\ \beta_2 \\ \vdots \\ \beta_p \end{bmatrix} \\
&= x_1\beta_1 + x_2\beta_2 + \dots + x_p\beta_p
\end{aligned}$$

Orthogonality

Definition 0.7 (Orthogonal vectors). Two vectors $\tilde{x}$ and $\tilde{y}$ of the same length are **orthogonal** (written $\tilde{x} \perp \tilde{y}$) if their dot product is zero:

$$\tilde{x} \perp \tilde{y} \iff \tilde{x}^\top \tilde{y} = 0$$

Orthogonality generalizes the geometric notion of perpendicularity to arbitrary dimensions.

Definition 0.8 (Orthonormal vectors). A set of vectors $\{\tilde{x}_1, \tilde{x}_2, \dots, \tilde{x}_k\}$ is **orthonormal** if the vectors are mutually orthogonal and each has unit length:

$$\tilde{x}_i^\top \tilde{x}_j = \begin{cases} 1 & \text{if } i = j \\ 0 & \text{if } i \neq j \end{cases}$$

The indicator vectors $\tilde{e}_1, \tilde{e}_2, \dots, \tilde{e}_p$ (Definition 0.5) form an orthonormal set.

Matrices

Definition 0.9 (Matrix). A **matrix** of dimensions $m \times n$ is a rectangular array of $m \cdot n$ numbers, arranged in m rows and n columns:

$$\mathbf{A} = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}$$

The entry in row i and column j is denoted a_{ij} or $(\mathbf{A})_{ij}$. A column vector of length p is a special case: a $p \times 1$ matrix. A row vector of length p is a $1 \times p$ matrix.

Matrix transpose

Definition 0.10 (Matrix transpose). The **transpose** of an $m \times n$ matrix $\mathbf{A}$ is the $n \times m$ matrix $\mathbf{A}^\top$ obtained by swapping the rows and columns of $\mathbf{A}$:

$$(\mathbf{A}^\top)_{ij} = a_{ji}$$

Theorem 0.3 (Transpose of a sum).

$$(\mathbf{A} + \mathbf{B})^\top = \mathbf{A}^\top + \mathbf{B}^\top$$

In particular, for column vectors $\tilde{x}$ and $\tilde{y}$:

$$(\tilde{x} + \tilde{y})^\top = \tilde{x}^\top + \tilde{y}^\top$$

Theorem 0.4 (Transpose of a product). *For compatible matrices $\mathbf{A}$ and $\mathbf{B}$:*

$$(\mathbf{AB})^\top = \mathbf{B}^\top \mathbf{A}^\top$$

The order of the factors reverses when transposing a product.

Matrix addition

Definition 0.11 (Zero matrix). The $m \times n$ **zero matrix** $\mathbf{0}_{m \times n}$ (or $\mathbf{0}$ when dimensions are clear from context) has all entries equal to zero:

$$\mathbf{0}_{m \times n} = \begin{bmatrix} 0 & 0 & \cdots & 0 \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 \end{bmatrix}$$

Definition 0.12 (Matrix addition). Two matrices $\mathbf{A}$ and $\mathbf{B}$ of the same dimensions $m \times n$ can be added element-wise; their **matrix sum** is:

$$(\mathbf{A} + \mathbf{B})_{ij} = a_{ij} + b_{ij}$$

Theorem 0.5 (Matrix addition is commutative).

$$\mathbf{A} + \mathbf{B} = \mathbf{B} + \mathbf{A}$$

Theorem 0.6 (Matrix addition is associative).

$$(\mathbf{A} + \mathbf{B}) + \mathbf{C} = \mathbf{A} + (\mathbf{B} + \mathbf{C})$$

Theorem 0.7 (Zero matrix is the additive identity).

$$\mathbf{A} + \mathbf{0} = \mathbf{A}$$

Theorem 0.8 (Additive inverse). *For any matrix $\mathbf{A}$, the matrix $-\mathbf{A}$ (defined by $(-\mathbf{A})_{ij} = -a_{ij}$) satisfies:*

$$\mathbf{A} + (-\mathbf{A}) = \mathbf{0}$$

Scalar multiplication

Definition 0.13 (Scalar multiplication). The **scalar multiple** of a matrix $\mathbf{A}$ by a scalar c is:

$$(c\mathbf{A})_{ij} = c \cdot a_{ij}$$

Matrix multiplication

Definition 0.14 (Matrix multiplication). The **product** of an $m \times k$ matrix $\mathbf{A}$ and a $k \times n$ matrix $\mathbf{B}$ is the $m \times n$ matrix $\mathbf{C} = \mathbf{AB}$ with entries:

$$c_{ij} = \sum_{s=1}^k a_{is} b_{sj}$$

Matrix multiplication is only defined when the number of columns in $\mathbf{A}$ equals the number of rows in $\mathbf{B}$.

Matrix multiplication is **not** commutative in general: $\mathbf{AB} \neq \mathbf{BA}$.

Theorem 0.9 (Matrix multiplication is associative).

$$(\mathbf{AB})\mathbf{C} = \mathbf{A}(\mathbf{BC})$$

Theorem 0.10 (Matrix multiplication is distributive over addition).

$$\mathbf{A}(\mathbf{B} + \mathbf{C}) = \mathbf{AB} + \mathbf{AC}$$

$$(\mathbf{A} + \mathbf{B})\mathbf{C} = \mathbf{AC} + \mathbf{BC}$$

Matrix-vector multiplication

Definition 0.15 (Matrix-vector multiplication). The **matrix-vector product** of an $m \times p$ matrix $\mathbf{A}$ and a $p \times 1$ column vector $\tilde{x}$ is the $m \times 1$ column vector $\mathbf{A}\tilde{x}$ with entries:

$$(\mathbf{A}\tilde{x})_i = \sum_{j=1}^p a_{ij} x_j$$

Matrix-vector multiplication is a generalization of the dot product. Each entry of the result is a dot product of a row of $\mathbf{A}$ with the vector $\tilde{x}$.

Special Matrices

See also Definition 0.11 for the zero matrix.

Definition 0.16 (Square matrix). A matrix is **square** if it has the same number of rows as columns.

Definition 0.17 (Order of a square matrix). The **order** of a square matrix (Definition 0.16) is its number of rows, which equals its number of columns.

Definition 0.18 (Matrix power). For a square matrix $\mathbf{A}$ of order p and a positive integer k , the k -th **power** of $\mathbf{A}$ is:

$$\mathbf{A}^k = \underbrace{\mathbf{A} \mathbf{A} \cdots \mathbf{A}}_{k \text{ copies}}$$

In particular, $\mathbf{A}^2 = \mathbf{A}\mathbf{A}$.

Definition 0.19 (Identity matrix). The $p \times p$ **identity matrix** $\mathbf{I}_p$ (or $\mathbf{I}$ when the size is clear from context) has ones on the main diagonal and zeros elsewhere:

$$(\mathbf{I}_p)_{ij} = \begin{cases} 1 & \text{if } i = j \\ 0 & \text{if } i \neq j \end{cases} \quad \mathbf{I}_p = \begin{bmatrix} 1 & 0 & \cdots & 0 \\ 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 1 \end{bmatrix}$$

Theorem 0.11 (Identity matrix is a multiplicative identity). *For any $m \times p$ matrix $\mathbf{A}$:*

$$\mathbf{A} \mathbf{I}_p = \mathbf{A}$$

$$\mathbf{I}_m \mathbf{A} = \mathbf{A}$$

Definition 0.20 (Symmetric matrix). A square matrix $\mathbf{A}$ is **symmetric** if $\mathbf{A}^\top = \mathbf{A}$, i.e., $a_{ij} = a_{ji}$ for all i and j .

Covariance matrices and information matrices are symmetric.

Definition 0.21 (Diagonal matrix). A square matrix $\mathbf{D}$ is a **diagonal matrix** if all off-diagonal entries are zero: $d_{ij} = 0$ whenever $i \neq j$:

$$\mathbf{D} = \begin{bmatrix} d_1 & 0 & \cdots & 0 \\ 0 & d_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & d_p \end{bmatrix}$$

Diagonal matrices are denoted $\mathbf{D} = \text{diag}(d_1, d_2, \dots, d_p)$, where $d_1, \dots, d_p$ are the diagonal entries.

Definition 0.22 (Matrix inverse). For a square $p \times p$ matrix $\mathbf{A}$, the **inverse** $\mathbf{A}^{-1}$ (if it exists) is the unique matrix satisfying:

$$\mathbf{A} \mathbf{A}^{-1} = \mathbf{A}^{-1} \mathbf{A} = \mathbf{I}_p$$

A matrix that has an inverse is called **invertible** or **non-singular**.

Theorem 0.12 (Inverse of a product). *For invertible matrices $\mathbf{A}$ and $\mathbf{B}$:*

$$(\mathbf{AB})^{-1} = \mathbf{B}^{-1} \mathbf{A}^{-1}$$

Definition 0.23 (Idempotent matrix). A square matrix $\mathbf{A}$ is **idempotent** if

$$\mathbf{A}^2 = \mathbf{A}$$

Definition 0.24 (Projection matrix). A square matrix $\mathbf{P}$ is a **projection matrix** (also called an *orthogonal projector*) if it is both symmetric and idempotent:

$$\mathbf{P}^\top = \mathbf{P} \quad \text{and} \quad \mathbf{P}^2 = \mathbf{P}$$

Theorem 0.13 (Complement of a projection matrix). *If $\mathbf{P}$ is a projection matrix, then $\mathbf{I} - \mathbf{P}$ is also a projection matrix.*

i Proof

Proof. We verify symmetry and idempotency.

Symmetry:

$$(\mathbf{I} - \mathbf{P})^\top = \mathbf{I}^\top - \mathbf{P}^\top = \mathbf{I} - \mathbf{P}$$

Idempotency:

$$\begin{aligned}(\mathbf{I} - \mathbf{P})^2 &= (\mathbf{I} - \mathbf{P})(\mathbf{I} - \mathbf{P}) \\&= \mathbf{I} - \mathbf{P} - \mathbf{P} + \mathbf{P}^2 \\&= \mathbf{I} - \mathbf{P} - \mathbf{P} + \mathbf{P} \\&= \mathbf{I} - \mathbf{P}\end{aligned}$$

□

Theorem 0.14 (Hat matrix is a projection matrix). *In a linear regression model with full-rank design matrix $\mathbf{X}$, the **hat matrix***

$$\mathbf{H} = \mathbf{X}(\mathbf{X}^\top \mathbf{X})^{-1} \mathbf{X}^\top$$

is a projection matrix.

i Proof

Proof. We verify symmetry and idempotency.

Symmetry:

$$\begin{aligned}\mathbf{H}^\top &= (\mathbf{X}(\mathbf{X}^\top \mathbf{X})^{-1} \mathbf{X}^\top)^\top \\&= (\mathbf{X}^\top)^\top \cdot ((\mathbf{X}^\top \mathbf{X})^{-1})^\top \cdot \mathbf{X}^\top \\&= \mathbf{X} \cdot (\mathbf{X}^\top \mathbf{X})^{-1} \cdot \mathbf{X}^\top \\&= \mathbf{H}\end{aligned}$$

where the third line uses $(\mathbf{X}^\top)^\top = \mathbf{X}$ and the fact that $\mathbf{X}^\top \mathbf{X}$ is symmetric, so its inverse is also symmetric $((\mathbf{X}^\top \mathbf{X})^{-1})^\top = (\mathbf{X}^\top \mathbf{X})^{-1}$.

Idempotency:

$$\begin{aligned}\mathbf{H}^2 &= \mathbf{X}(\mathbf{X}^\top \mathbf{X})^{-1} \mathbf{X}^\top \cdot \mathbf{X}(\mathbf{X}^\top \mathbf{X})^{-1} \mathbf{X}^\top \\&= \mathbf{X}(\mathbf{X}^\top \mathbf{X})^{-1} (\mathbf{X}^\top \mathbf{X}) (\mathbf{X}^\top \mathbf{X})^{-1} \mathbf{X}^\top \\&= \mathbf{X}(\mathbf{X}^\top \mathbf{X})^{-1} \mathbf{X}^\top \\&= \mathbf{H}\end{aligned}$$

□

The hat matrix appears in the formula for fitted values in linear regression: $\hat{\tilde{y}} = \mathbf{X}\hat{\tilde{\beta}} = \mathbf{X}(\mathbf{X}^\top \mathbf{X})^{-1} \mathbf{X}^\top \tilde{y} = \mathbf{H}\tilde{y}$. It “puts a hat” on $\tilde{y}$ — hence the name.

Theorem 0.15 (Projection matrices produce orthogonal decompositions). *If $\mathbf{P}$ is a projection matrix and $\tilde{v}$ is any vector of compatible dimension, then the two components of the decomposition*

$$\tilde{v} = \underbrace{\mathbf{P}\tilde{v}}_{\text{projected}} + \underbrace{(\mathbf{I} - \mathbf{P})\tilde{v}}_{\text{residual}}$$

are orthogonal:

$$\mathbf{P}\tilde{v} \perp (\mathbf{I} - \mathbf{P})\tilde{v}$$

i Proof

Proof.

$$\begin{aligned} (\mathbf{P}\tilde{v})^\top (\mathbf{I} - \mathbf{P})\tilde{v} &= \tilde{v}^\top \mathbf{P}^\top (\mathbf{I} - \mathbf{P})\tilde{v} \\ &= \tilde{v}^\top \mathbf{P} (\mathbf{I} - \mathbf{P})\tilde{v} \\ &= \tilde{v}^\top (\mathbf{P} - \mathbf{P}^2)\tilde{v} \\ &= \tilde{v}^\top (\mathbf{P} - \mathbf{P})\tilde{v} \\ &= \tilde{v}^\top \mathbf{0} \tilde{v} \\ &= 0 \end{aligned}$$

where the second line uses symmetry ($\mathbf{P}^\top = \mathbf{P}$) and the fourth line uses idempotency ($\mathbf{P}^2 = \mathbf{P}$). $\square$

Quadratic Forms

Definition 0.25 (Quadratic form). A **quadratic form** is a mathematical expression of the structure

$$\tilde{x}^\top \mathbf{S} \tilde{x}$$

where $\tilde{x}$ is a $p \times 1$ vector and $\mathbf{S}$ is a $p \times p$ matrix.

Quadratic forms are the matrix generalizations of the scalar expression cx^2 . They occur frequently in statistics:

- The residual sum of squares in linear regression (Section) is a quadratic form.
- The variance of a linear combination of estimates (see [Inference about Gaussian Linear Regression Models](#)) is a quadratic form: $\text{Var}\left(\tilde{x}^\top \hat{\beta}\right) = \tilde{x}^\top \text{Var}\left(\hat{\beta}\right) \tilde{x}$.

Theorem 0.16 (Symmetric part of a quadratic form). *If $\mathbf{S}$ is a square matrix, then*

$$\tilde{x}^\top \mathbf{S} \tilde{x} = \tilde{x}^\top \left(\frac{1}{2}(\mathbf{S} + \mathbf{S}^\top) \right) \tilde{x}.$$

So the value of a quadratic form depends only on the symmetric part of $\mathbf{S}$.

Design Matrix

Definition 0.26 (Design matrix). In a regression model with n observations and p predictors, the **design matrix** (or *model matrix*) $\mathbf{X}$ is the $n \times p$ matrix whose i -th row is the covariate vector $\tilde{x}_i^\top$ for observation i :

$$\mathbf{X} = \begin{bmatrix} \tilde{x}_1^\top \\ \tilde{x}_2^\top \\ \vdots \\ \tilde{x}_n^\top \end{bmatrix} = \begin{bmatrix} x_{11} & x_{12} & \cdots & x_{1p} \\ x_{21} & x_{22} & \cdots & x_{2p} \\ \vdots & \vdots & \ddots & \vdots \\ x_{n1} & x_{n2} & \cdots & x_{np} \end{bmatrix}$$

The product $\mathbf{X}\tilde{\beta}$ collects all the linear predictors $\tilde{x}_i^\top \tilde{\beta}$ into a single $n \times 1$ vector:

$$\mathbf{X}\tilde{\beta} = \begin{bmatrix} \tilde{x}_1^\top \tilde{\beta} \\ \vdots \\ \tilde{x}_n^\top \tilde{\beta} \end{bmatrix}$$

The matrix $\mathbf{X}^\top \mathbf{X}$ is a $p \times p$ symmetric matrix that appears in the OLS estimator $\hat{\beta} = (\mathbf{X}^\top \mathbf{X})^{-1} \mathbf{X}^\top \tilde{y}$.

Vector Calculus

(adapted from Fieller (2016), §7.2)

This section covers derivatives of functions of vectors and matrices. Linear algebra prerequisites — including vectors, matrices, transpose, dot product, and quadratic forms — are covered in Section .

Let $\tilde{x}$ and $\tilde{\beta}$ be column vectors of length p (see Definition 0.1 and Definition 0.6).

Definition 0.1 (Vector derivative). If $f(\tilde{\beta})$ is a function that takes a vector $\tilde{\beta}$ as input, such as $f(\tilde{\beta}) = x'\tilde{\beta}$, then its **vector derivative** is:

$$\frac{\partial}{\partial \tilde{\beta}} f(\tilde{\beta}) = \begin{bmatrix} \frac{\partial}{\partial \beta_1} f(\tilde{\beta}) \\ \frac{\partial}{\partial \beta_2} f(\tilde{\beta}) \\ \vdots \\ \frac{\partial}{\partial \beta_p} f(\tilde{\beta}) \end{bmatrix}$$

Definition 0.2 (Row-vector derivative). If $f(\tilde{\beta})$ is a function that takes a vector $\tilde{\beta}$ as input, such as $f(\tilde{\beta}) = x'\tilde{\beta}$, then its **row-vector derivative** is:

$$\frac{\partial}{\partial \tilde{\beta}^\top} f(\tilde{\beta}) = \begin{bmatrix} \frac{\partial}{\partial \beta_1} f(\tilde{\beta}) & \frac{\partial}{\partial \beta_2} f(\tilde{\beta}) & \cdots & \frac{\partial}{\partial \beta_p} f(\tilde{\beta}) \end{bmatrix}$$

Theorem 0.1 (Row and column derivatives are transposes).

$$\frac{\partial}{\partial \tilde{\beta}^\top} f(\tilde{\beta}) = \left(\frac{\partial}{\partial \tilde{\beta}} f(\tilde{\beta}) \right)^\top$$

$$\frac{\partial}{\partial \tilde{\beta}} f(\tilde{\beta}) = \left(\frac{\partial}{\partial \tilde{\beta}^\top} f(\tilde{\beta}) \right)^\top$$

Definition 0.3 (Constant). $\tilde{x}$ is **constant with respect to** $\tilde{\beta}$ if

$$\underbrace{\frac{\partial}{\partial \tilde{\beta}} \tilde{x}^\top}_{p \times p} = \mathbf{0}_{p \times p}$$

Example 0.1 (A constant vector). Let $\tilde{\beta} = (\beta_1, \beta_2)^\top$ and $\tilde{x} = (3, 5)^\top$, so $x_1 = 3$ and $x_2 = 5$ do not depend on $\tilde{\beta}$. Expanding $\frac{\partial}{\partial \tilde{\beta}} \tilde{x}^\top$ into its matrix of scalar partial derivatives (Definition 0.1, applied to each component of the row $\tilde{x}^\top$) and evaluating each entry:

$$\underbrace{\frac{\partial}{\partial \tilde{\beta}} \tilde{x}^\top}_{2 \times 2} = \frac{\partial}{\partial \tilde{\beta}} \begin{bmatrix} x_1 & x_2 \end{bmatrix} = \begin{bmatrix} \frac{\partial}{\partial \beta_1} x_1 & \frac{\partial}{\partial \beta_1} x_2 \\ \frac{\partial}{\partial \beta_2} x_1 & \frac{\partial}{\partial \beta_2} x_2 \end{bmatrix} = \begin{bmatrix} \frac{\partial}{\partial \beta_1} 3 & \frac{\partial}{\partial \beta_1} 5 \\ \frac{\partial}{\partial \beta_2} 3 & \frac{\partial}{\partial \beta_2} 5 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} = \mathbf{0}_{2 \times 2}$$

Every entry is the derivative of a constant, so $\frac{\partial}{\partial \tilde{\beta}} \tilde{x}^\top = \mathbf{0}_{2 \times 2}$ and $\tilde{x}$ is constant with respect to $\tilde{\beta}$ (Definition 0.3).

Theorem 0.2 (Derivative of a dot product). *If $\tilde{x}$ is constant with respect to $\tilde{\beta}$, then:*

$$\underbrace{\frac{\partial}{\partial \tilde{\beta}} (\tilde{x} \cdot \tilde{\beta})}_{p \times 1} = \underbrace{\frac{\partial}{\partial \tilde{\beta}} (\tilde{\beta} \cdot \tilde{x})}_{p \times 1} = \tilde{x}_{p \times 1}$$

i Proof

Proof.

$$\begin{aligned}\frac{\partial}{\partial \tilde{\beta}}(\tilde{x} \cdot \tilde{\beta}) &= \begin{bmatrix} \frac{\partial}{\partial \beta_1}(x_1\beta_1 + x_2\beta_2 + \dots + x_p\beta_p) \\ \frac{\partial}{\partial \beta_2}(x_1\beta_1 + x_2\beta_2 + \dots + x_p\beta_p) \\ \vdots \\ \frac{\partial}{\partial \beta_p}(x_1\beta_1 + x_2\beta_2 + \dots + x_p\beta_p) \end{bmatrix} \\ &= \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_p \end{bmatrix} \\ &= \tilde{x}\end{aligned}$$

□

Example 0.2 (Derivative of a dot product). Let $\tilde{x} = (3, 5)^\top$ (constant with respect to $\tilde{\beta}$; see Example 0.1) and $\tilde{\beta} = (\beta_1, \beta_2)^\top$. Then $\tilde{x} \cdot \tilde{\beta} = 3\beta_1 + 5\beta_2$, and by Theorem 0.2:

$$\underbrace{\frac{\partial}{\partial \tilde{\beta}}(\tilde{x} \cdot \tilde{\beta})}_{2 \times 1} = \underbrace{\tilde{x}}_{2 \times 1} = \begin{pmatrix} 3 \\ 5 \end{pmatrix}$$

Verifying entry-wise:

$$\frac{\partial}{\partial \tilde{\beta}}(3\beta_1 + 5\beta_2) = \begin{pmatrix} \frac{\partial}{\partial \beta_1}(3\beta_1 + 5\beta_2) \\ \frac{\partial}{\partial \beta_2}(3\beta_1 + 5\beta_2) \end{pmatrix} = \begin{pmatrix} 3 \\ 5 \end{pmatrix}$$

Both methods agree.

Theorem 0.3 (Product rule for dot-products). *If $\mathbf{a} = \mathbf{a}(\tilde{x})$ and $\mathbf{b} = \mathbf{b}(\tilde{x})$ are differentiable $p \times 1$ vector functions of $\tilde{x}$, then:*

$$\frac{\partial}{\partial \tilde{x}} \underbrace{\tilde{x}}_{p \times 1} \cdot \underbrace{\mathbf{a}}_{p \times 1} \cdot \underbrace{\mathbf{b}}_{p \times 1} = \left(\frac{\partial}{\partial \tilde{x}} \underbrace{\tilde{x}}_{p \times 1} \underbrace{\mathbf{a}^\top}_{1 \times p} \right) \underbrace{\mathbf{b}}_{p \times 1} + \left(\frac{\partial}{\partial \tilde{x}} \underbrace{\tilde{x}}_{p \times 1} \underbrace{\mathbf{b}^\top}_{1 \times p} \right) \underbrace{\mathbf{a}}_{p \times 1}$$

i Proof

Proof. Entry-wise, for $i = 1, \dots, p$:

$$\begin{aligned}
\left[\frac{\partial}{\partial \tilde{x}} (\mathbf{a} \cdot \mathbf{b}) \right]_i &= \frac{\partial}{\partial x_i} \sum_{k=1}^p a_k b_k \\
&= \sum_{k=1}^p \left(b_k \frac{\partial}{\partial x_i} a_k + a_k \frac{\partial}{\partial x_i} b_k \right) \\
&= \left[\left(\frac{\partial}{\partial \tilde{x}} \mathbf{a}^\top \right) \mathbf{b} \right]_i + \left[\left(\frac{\partial}{\partial \tilde{x}} \mathbf{b}^\top \right) \mathbf{a} \right]_i
\end{aligned}$$

□

Example 0.3 (Example of the dot-product rule). Let $\tilde{x} = (\beta_1, \beta_2)^\top$, $\mathbf{a}(\tilde{x}) = (\beta_1, \beta_1\beta_2)^\top$, and $\mathbf{b}(\tilde{x}) = (\beta_2, \beta_1)^\top$. Then:

$$\mathbf{a} \cdot \mathbf{b} = \beta_1 \cdot \beta_2 + \beta_1\beta_2 \cdot \beta_1 = \beta_1\beta_2 + \beta_1^2\beta_2$$

By direct calculation:

$$\underbrace{\frac{\partial}{\partial \tilde{x}} (\mathbf{a} \cdot \mathbf{b})}_{2 \times 1} = \frac{\partial}{\partial \tilde{x}} (\beta_1\beta_2 + \beta_1^2\beta_2) = \begin{pmatrix} \beta_2 + 2\beta_1\beta_2 \\ \beta_1 + \beta_1^2 \end{pmatrix}$$

By the product rule (Theorem 0.3), using $\underbrace{\frac{\partial}{\partial \tilde{x}} \mathbf{a}^\top}_{2 \times 2} = \begin{pmatrix} 1 & \beta_2 \\ 0 & \beta_1 \end{pmatrix}$ and $\underbrace{\frac{\partial}{\partial \tilde{x}} \mathbf{b}^\top}_{2 \times 2} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$:

$$\begin{aligned}
\underbrace{\frac{\partial}{\partial \tilde{x}} (\mathbf{a} \cdot \mathbf{b})}_{2 \times 1} &= \underbrace{\begin{pmatrix} 1 & \beta_2 \\ 0 & \beta_1 \end{pmatrix}}_{2 \times 2} \underbrace{\begin{pmatrix} \beta_2 \\ \beta_1 \end{pmatrix}}_{2 \times 1} + \underbrace{\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}}_{2 \times 2} \underbrace{\begin{pmatrix} \beta_1 \\ \beta_1\beta_2 \end{pmatrix}}_{2 \times 1} \\
&= \begin{pmatrix} \beta_2 + \beta_1\beta_2 \\ \beta_1^2 \end{pmatrix} + \begin{pmatrix} \beta_1\beta_2 \\ \beta_1 \end{pmatrix} \\
&= \begin{pmatrix} \beta_2 + 2\beta_1\beta_2 \\ \beta_1^2 + \beta_1 \end{pmatrix}
\end{aligned}$$

Both methods agree.

Theorem 0.4 (Derivative of a linear map). *If A is an $m \times p$ matrix that is constant with respect to $\tilde{\beta}$, then:*

$$\underbrace{\frac{\partial}{\partial \tilde{\beta}} (A\tilde{\beta})}_{p \times m} = \underbrace{A^\top}_{p \times m}$$

i Proof

Proof. For entry (i, j) , where row i indexes the denominator $\tilde{\beta}$ (see Definition 0.1) and column j indexes the numerator $A\tilde{\beta}$:

$$\begin{aligned} \left[\frac{\partial}{\partial \tilde{\beta}} (A\tilde{\beta}) \right]_{ij} &= \frac{\partial}{\partial \beta_i} (A\tilde{\beta})_j \\ &= \frac{\partial}{\partial \beta_i} \sum_{k=1}^p a_{jk} \beta_k \\ &= a_{ji} \\ &= [A^\top]_{ij} \end{aligned}$$

□

Example 0.4 (Derivative of a linear map). Let $A = \begin{pmatrix} 2 & 3 \end{pmatrix}$ (1×2) and $\tilde{\beta} = (\beta_1, \beta_2)^\top$. Then $A\tilde{\beta} = 2\beta_1 + 3\beta_2$, and by Theorem 0.4:

$$\underbrace{\frac{\partial}{\partial \tilde{\beta}} (A\tilde{\beta})}_{2 \times 1} = \underbrace{A^\top}_{2 \times 1} = \begin{pmatrix} 2 \\ 3 \end{pmatrix}$$

Theorem 0.5 (Vector-derivative of a matrix-vector product). *If A is an $m \times q$ matrix that is constant with respect to $\tilde{\beta}$, and $\tilde{v} = \tilde{v}(\tilde{\beta})$ is a $q \times 1$ vector that depends on the $p \times 1$ vector $\tilde{\beta}$, then:*

$$\underbrace{\frac{\partial}{\partial \tilde{\beta}} (A\tilde{v})}_{p \times m} = \underbrace{\left(\frac{\partial}{\partial \tilde{\beta}} \tilde{v} \right)}_{p \times q} \underbrace{A^\top}_{q \times m}$$

This result generalizes Theorem 0.4, which is the special case $\tilde{v} = \tilde{\beta}$ (so that $\frac{\partial}{\partial \tilde{\beta}} \tilde{\beta} = \mathbf{I}$ and $\frac{\partial}{\partial \tilde{\beta}} (A\tilde{\beta}) = A^\top$).

i Proof

Proof. For entry (i, j) , where row i indexes the denominator $\tilde{\beta}$ and column j indexes the numerator $A\tilde{v}$:

$$\begin{aligned}
\left[\frac{\partial}{\partial \tilde{\beta}} (A\tilde{v}) \right]_{ij} &= \frac{\partial}{\partial \beta_i} (A\tilde{v})_j \\
&= \frac{\partial}{\partial \beta_i} \sum_{k=1}^q a_{jk} v_k \\
&= \sum_{k=1}^q a_{jk} \frac{\partial}{\partial \beta_i} v_k \\
&= \sum_{k=1}^q \left[\frac{\partial}{\partial \tilde{\beta}} \tilde{v} \right]_{ik} [A^\top]_{kj} \\
&= \left[\left(\frac{\partial}{\partial \tilde{\beta}} \tilde{v} \right) A^\top \right]_{ij}
\end{aligned}$$

□

Example 0.5 (Vector-derivative of a matrix-vector product). Let $A = \begin{pmatrix} 2 & 3 \end{pmatrix}$ (1×2 , constant) and $\tilde{v}(\tilde{\beta}) = (\beta_1^2, \beta_2^2)^\top$. Then $A\tilde{v} = 2\beta_1^2 + 3\beta_2^2$. By Theorem 0.5:

$$\begin{aligned}
\underbrace{\frac{\partial}{\partial \tilde{\beta}} (A\tilde{v})}_{2 \times 1} &= \begin{pmatrix} 2\beta_1 & 0 \\ 0 & 2\beta_2 \end{pmatrix} \begin{pmatrix} 2 \\ 3 \end{pmatrix} \\
&= \begin{pmatrix} 4\beta_1 \\ 6\beta_2 \end{pmatrix}
\end{aligned}$$

Corollary 0.1 (Derivative of a dot product, transpose-product form). *If $\tilde{x}$ is constant with respect to $\tilde{\beta}$, then:*

$$\underbrace{\frac{\partial}{\partial \tilde{\beta}} (\tilde{x}^\top \tilde{\beta})}_{p \times 1} = \tilde{x}$$

This vector derivative formula looks a lot like non-vector calculus, except that you have to transpose the coefficient: in scalar calculus $\frac{\partial}{\partial x}(cx) = c$, but here the coefficient $\tilde{x}^\top$ (a row vector) becomes $\tilde{x}$ (a column vector) in the result.

i Proof

Proof. Using Theorem 0.2:

Since $\tilde{x}^\top \tilde{\beta} = \tilde{x} \cdot \tilde{\beta}$ (Definition 0.6), and $\tilde{x}$ is constant with respect to $\tilde{\beta}$:

$$\frac{\partial}{\partial \tilde{\beta}}(\tilde{x}^\top \tilde{\beta}) = \frac{\partial}{\partial \tilde{\beta}}(\tilde{x} \cdot \tilde{\beta}) = \tilde{x}$$

by Theorem 0.2. □

i Proof

Proof. Using Theorem 0.5:

Since $\tilde{x}$ is constant with respect to $\tilde{\beta}$, $A = \tilde{x}^\top$ is a constant $1 \times p$ matrix. Applying Theorem 0.5 with $\tilde{v} = \tilde{\beta}$ (so $\frac{\partial}{\partial \tilde{\beta}} \tilde{\beta} = \mathbf{I}$):

$$\begin{aligned} \frac{\partial}{\partial \tilde{\beta}}(\tilde{x}^\top \tilde{\beta}) &= \left(\frac{\partial}{\partial \tilde{\beta}} \tilde{\beta} \right) (\tilde{x}^\top)^\top \\ &= \mathbf{I} \cdot \tilde{x} \\ &= \tilde{x} \end{aligned}$$

□

Example 0.6 (Derivative of a transpose product). Let $\tilde{x} = (3, 5)^\top$ and $\tilde{\beta} = (\beta_1, \beta_2)^\top$. Then $\tilde{x}^\top \tilde{\beta} = 3\beta_1 + 5\beta_2$, and by Corollary 0.1:

$$\underbrace{\frac{\partial}{\partial \tilde{\beta}} \begin{pmatrix} \tilde{x}^\top & \tilde{\beta} \\ 1 \times 2 & 2 \times 1 \end{pmatrix}}_{2 \times 1} = \underbrace{\tilde{x}}_{2 \times 1} = \begin{pmatrix} 3 \\ 5 \end{pmatrix}$$

Theorem 0.6 (Derivative of a quadratic form). For a quadratic form (Definition 0.25), if S is a symmetric $p \times p$ matrix that is constant with respect to $\tilde{\beta}$, then:

$$\underbrace{\frac{\partial}{\partial \tilde{\beta}}(\tilde{\beta}^\top S \tilde{\beta})}_{p \times 1} = \underbrace{2S\tilde{\beta}}_{p \times 1}$$

i Proof

Proof. Expanding entry-wise, $\tilde{\beta}^\top S \tilde{\beta} = \sum_{j=1}^p \sum_{k=1}^p s_{jk} \beta_j \beta_k$. Differentiating component-wise with respect to β_i for $i = 1, \dots, p$:

$$\begin{aligned}
\left[\frac{\partial}{\partial \tilde{\beta}} (\tilde{\beta}^\top S \tilde{\beta}) \right]_i &= \frac{\partial}{\partial \beta_i} \sum_{j=1}^p \sum_{k=1}^p s_{jk} \beta_j \beta_k \quad (\text{expand quadratic form}) \\
&= \sum_{k=1}^p s_{ik} \beta_k + \sum_{j=1}^p s_{ji} \beta_j \quad (\text{product rule for } \beta_i \beta_k) \\
&= [S \tilde{\beta}]_i + [S^\top \tilde{\beta}]_i \quad (\text{matrix-vector multiplication definition}) \\
&= [(S + S^\top) \tilde{\beta}]_i \quad (\text{linearity of matrix multiplication})
\end{aligned}$$

When S is symmetric ($S = S^\top$), $S + S^\top = 2S$, so $\frac{\partial}{\partial \tilde{\beta}} (\tilde{\beta}^\top S \tilde{\beta}) = 2S \tilde{\beta}$. $\square$

This operation is like taking the derivative of cx^2 with respect to x in non-vector calculus.

Example 0.7 (Derivative of a quadratic form). Let $S = \begin{pmatrix} 3 & 1 \\ 1 & 2 \end{pmatrix}$ (2×2 , symmetric and constant) and $\tilde{\beta} = (\beta_1, \beta_2)^\top$. Then $\tilde{\beta}^\top S \tilde{\beta} = 3\beta_1^2 + 2\beta_1\beta_2 + 2\beta_2^2$. By Theorem 0.6:

$$\underbrace{\frac{\partial}{\partial \tilde{\beta}} (\tilde{\beta}^\top S \tilde{\beta})}_{2 \times 1} = 2S \tilde{\beta} = 2 \begin{pmatrix} 3 & 1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} \beta_1 \\ \beta_2 \end{pmatrix} = \begin{pmatrix} 6\beta_1 + 2\beta_2 \\ 2\beta_1 + 4\beta_2 \end{pmatrix}$$

Differentiating component-wise directly:

$$\begin{pmatrix} \frac{\partial}{\partial \beta_1} (3\beta_1^2 + 2\beta_1\beta_2 + 2\beta_2^2) \\ \frac{\partial}{\partial \beta_2} (3\beta_1^2 + 2\beta_1\beta_2 + 2\beta_2^2) \end{pmatrix} = \begin{pmatrix} 6\beta_1 + 2\beta_2 \\ 2\beta_1 + 4\beta_2 \end{pmatrix}$$

Both methods agree.

Corollary 0.2 (Derivative of a simple quadratic form).

$$\underbrace{\frac{\partial}{\partial \tilde{\beta}} (\tilde{\beta}^\top \tilde{\beta})}_{p \times 1} = \underbrace{2\tilde{\beta}}_{p \times 1}$$

i Proof

Proof. Applying Theorem 0.6 with $S = \mathbf{I}_{p \times p}$ (which is symmetric and constant with respect to $\tilde{\beta}$):

$$\begin{aligned}
\frac{\partial}{\partial \tilde{\beta}}(\tilde{\beta}^\top \tilde{\beta}) &= \frac{\partial}{\partial \tilde{\beta}}(\tilde{\beta}^\top \mathbf{I}_{p \times p} \tilde{\beta}) \quad (\text{rewrite with identity matrix}) \\
&= 2\mathbf{I}_{p \times p} \tilde{\beta} \quad (\text{apply @thm-quadratic-form with } S = \mathbf{I}_{p \times p}) \\
&= 2\tilde{\beta} \quad (\text{identity matrix property})
\end{aligned}$$

□

This vector derivative is like taking the derivative of x^2 .

Example 0.8 (Derivative of a sum of squares). Let $\tilde{\beta} = (\beta_1, \beta_2)^\top$, so $\tilde{\beta}^\top \tilde{\beta} = \beta_1^2 + \beta_2^2$. By Corollary 0.2:

$$\underbrace{\frac{\partial}{\partial \tilde{\beta}}(\tilde{\beta}^\top \tilde{\beta})}_{2 \times 1} = 2\tilde{\beta} = \begin{pmatrix} 2\beta_1 \\ 2\beta_2 \end{pmatrix}$$

Direct partial differentiation yields the same column vector.

Theorem 0.7 (Vector chain rule).

$$\frac{\partial z}{\partial \tilde{x}} = \frac{\partial y}{\partial \tilde{x}} \frac{\partial z}{\partial y}$$

or in Euler/Lagrange notation:

$$(f(g(\tilde{x})))' = \tilde{g}'(\tilde{x})f'(g(\tilde{x}))$$

See <https://quickfem.com/finite-element-analysis/>, specifically https://quickfem.com/wp-content/uploads/IFEM.AppF_.pdf

See also https://en.wikipedia.org/wiki/Gradient#Relationship_with_Fr%C3%A9chet_derivative

This chain rule is like the univariate chain rule (Theorem 0.8), but the order matters now. The version presented here is for the **gradient** (column vector); the **total derivative** (row vector) would be the **transpose of the gradient**.

Corollary 0.3 (Vector chain rule for quadratic forms).

$$\frac{\partial}{\partial \tilde{\beta}} (\tilde{\varepsilon}(\tilde{\beta}) \cdot \tilde{\varepsilon}(\tilde{\beta})) = \left(\frac{\partial}{\partial \tilde{\beta}} \tilde{\varepsilon}(\tilde{\beta}) \right) (2\tilde{\varepsilon}(\tilde{\beta}))$$

Additional resources

Calculus

- Kaplan (2022)
- Khuri (2003)
- Banner (2007)
- Larson and Edwards (2018)
- Miller (2016)
 - <http://www.youtube.com/watch?v=xYzQL0TUtBA>
 - http://www.youtube.com/watch?v=Ps2SBo_WjoE

Linear Algebra and Vector Calculus

- Fieller (2016)
- Banerjee and Roy (2014)
- Searle and Khuri (2017)

Numerical Analysis

- [Hua Zhou's lecture notes](#) for “UCLA Biostat 216 - Mathematical Methods for Biostatistics” (2023 Fall)

Real Analysis

- Grinberg (2017)

References

- Banerjee, Sudipto, and Anindya Roy. 2014. *Linear Algebra and Matrix Analysis for Statistics*. Vol. 181. Crc Press Boca Raton. <https://www.routledge.com/Linear-Algebra-and-Matrix-Analysis-for-Statistics/Banerjee-Roy/p/book/9781420095388>.
- Banner, Adrian D. 2007. *The Calculus Lifesaver : All the Tools You Need to Excel at Calculus*. A Princeton Lifesaver Study Guide. Princeton University Press. <https://press.princeton.edu/books/paperback/9780691130880/the-calculus-lifesaver>.
- Billingsley, Patrick. 1995. *Probability and Measure*. 3rd ed. Wiley Series in Probability and Mathematical Statistics. Wiley.
- Cheng, Eugenia. 2025. “Opinion | How Math Turned Me from a D.E.I. Skeptic to a Supporter.” *The New York Times*. <https://www.nytimes.com/2025/09/05/opinion/math-dei.html>.
- Dobson, Annette J, and Adrian G Barnett. 2018. *An Introduction to Generalized Linear Models*. 4th ed. CRC press. <https://doi.org/10.1201/9781315182780>.
- Fieller, Nick. 2016. *Basics of Matrix Algebra for Statistics with R*. Chapman; Hall/CRC. <https://doi.org/10.1201/9781315370200>.
- Fubini, Guido. 1907. “Sugli Integrali Multipli.” *Rendiconti Della Reale Accademia Dei Lincei. Classe Di Scienze Fisiche, Matematiche e Naturali* 16: 608–14.
- Grinberg, Raffi. 2017. *The Real Analysis Lifesaver: All the Tools You Need to Understand Proofs*. 1st ed. Princeton Lifesaver Study Guides. Princeton University Press. <https://press.princeton.edu/books/paperback/9780691172934/the-real-analysis-lifesaver>.
- Gut, Allan. 2013. *Probability: A Graduate Course*. 2nd ed. Springer Texts in Statistics. Springer. <https://doi.org/10.1007/978-1-4614-4708-5>.
- Kaplan, Daniel. 2022. *MOSAIC Calculus*. Www.mosaic-web.org. www.mosaic-web.org.
- Khuri, André I. 2003. *Advanced Calculus with Applications in Statistics*. John Wiley & Sons. <https://doi.org/10.1002/0471394882>.
- Kleinbaum, David G, and Mitchel Klein. 2012. *Survival Analysis: A Self-Learning Text*. 3rd ed. Springer. <https://doi.org/10.1007/978-1-4419-6646-9>.
- Larson, Ron, and Bruce H. Edwards. 2018. *Calculus*. 11th ed. Cengage Learning. <https://www.cengage.com/c/calculus-11e-larson/>.

Miller, Steven J. 2016. *The Probability Lifesaver: Calculus Review Problems*. https://web.williams.edu/Mathematics/sjmiller/public_html/probabilitylifesaver/index.htm#:~:text=http%3A//web.williams.edu/Mathematics/sjmiller/public_html/probabilitylifesaver/supplementalchap_calcreview.pdf.

Rudin, Walter. 1976. *Principles of Mathematical Analysis*. 3rd ed. International Series in Pure and Applied Mathematics. McGraw-Hill.

Searle, Shayle R, and Andre I Khuri. 2017. *Matrix Algebra Useful for Statistics*. John Wiley & Sons.

Wikipedia contributors. 2024. *Fubini's Theorem* — *Wikipedia, the Free Encyclopedia*. https://en.wikipedia.org/wiki/Fubini%27s_theorem.