Proof Writing

Last modified: 2026-10-19 13:49:51 (PDT)

This page collects general advice on how to write proofs and derivations. The goal of a proof is not just to convince yourself that a result is true; it is to convince a *reader*, and to show them *why* it is true. Each principle on this page serves that goal. The symbols for logical entailment are listed under Definitions for proofs below.

Definitions for proofs

Definition 0.1 (Proposition (statement))**.** A **proposition** [1], or **statement**, is a declarative sentence that is either true or false.

Example 0.1 (Propositions, and a sentence that is not one)**.**

- "$2 + 3 = 5$" is a proposition, and it is true.
- "7 is even" is a proposition, and it is false.
- "$x > 2$" is not a proposition on its own: it is true when $x = 3$ and false when $x = 1$, so it has no truth value until x is given a value.

Definition 0.2 (Predicate)**.** A **predicate** [2] $P(x)$ is a declarative sentence about a variable x that becomes a proposition (Definition 0.1) whenever a value is substituted for x.

Example 0.2 (A predicate)**.** Let $P(x)$ be the predicate "$x > 2$", for real numbers x.

- $P(3)$ is the proposition "$3 > 2$", which is true.
- $P(1)$ is the proposition "$1 > 2$", which is false.

Definition 0.3 (Logical connective)**.** A **logical connective** [3] builds a new proposition (Definition 0.1) from one or two propositions P and Q, with a truth value that depends only on the truth values of P and Q. These notes use five, defined next: not (Definition 0.4), and (Definition 0.5), or (Definition 0.6), implies (Definition 0.7), and if and only if (Definition 0.8).

Definition 0.4 (Not)**.** The logical connective (Definition 0.3) **not** [4] is written $\neg P$ ("not P"); it is true when P is false, and false when P is true.

Example 0.3 (Negating a true and a false proposition)**.** Let P be "$5 > 3$", which is true, and let Q be "$2 > 3$", which is false.

- $\neg P$, "it is not true that $5 > 3$", is false, because P is true.
- $\neg Q$, "it is not true that $2 > 3$", is true, because Q is false.

Definition 0.5 (And)**.** The logical connective (Definition 0.3) **and** is written [5] $P \wedge Q$ ("P and Q"); it is true when P and Q are both true, and false otherwise.

Example 0.4 (Two propositions joined by "and")**.** Let P be "$3 < 5$", which is true, let Q be "$5 < 4$", which is false, and let R be "$5 < 7$", which is true.

- $P \wedge R$ is true, because P and R are both true.
- $P \wedge Q$ is false, because Q is false.

Definition 0.6 (Or)**.** The logical connective (Definition 0.3) **or** is written [6] $P \vee Q$ ("P or Q"); it is true when at least one of P and Q is true, and false when both are false.

Example 0.5 (Two propositions joined by "or")**.** Let P be "$3 < 2$", which is false, let Q be "$5 < 7$", which is true, and let R be "$5 < 4$", which is false.

- $P \vee Q$ is true, because Q is true.
- $P \vee R$ is false, because P and R are both false.

Definition 0.7 (Implies)**.** The logical connective (Definition 0.3) **implies** is written [7] $P \Rightarrow Q$ ("if P, then Q", or "P implies Q"); it is false when P is true and Q is false, and true otherwise.

Example 0.6 (Implications with a true and a false premise)**.** Take $x = 3$.

- $x > 2 \Rightarrow x > 1$ is true: the premise $3 > 2$ is true and the conclusion $3 > 1$ is true.
- $x > 2 \Rightarrow x > 4$ is false: the premise $3 > 2$ is true, but the conclusion $3 > 4$ is false.
- $x > 5 \Rightarrow x > 4$ is true: the premise $3 > 5$ is false, so the implication is true whatever the conclusion.

Definition 0.8 (If and only if)**.** The logical connective (Definition 0.3) **if and only if** is written [8] $P \Leftrightarrow Q$ ("P if and only if Q"); it is true when P and Q have the same truth value, and false otherwise.

Example 0.7 ("If and only if" with equal and different truth values)**.** Take $x = 4$.

- "x is even $\Leftrightarrow x > 3$" is true: 4 is even and $4 > 3$, so both sides are true.
- "$x > 3 \Leftrightarrow x > 5$" is false: $4 > 3$ is true, but $4 > 5$ is false.

Example 0.8 (Truth values of connectives)**.** Let P be "$2 + 3 = 5$", which is true, and let Q be "7 is even", which is false.

- $\neg Q$, "7 is not even", is true, because Q is false.
- $P \wedge Q$ is false, because Q is false.
- $P \vee Q$ is true, because P is true.
- $P \Rightarrow Q$ is false, because P is true and Q is false.
- $Q \Rightarrow P$ is true, because Q is false.
- $P \Leftrightarrow Q$ is false, because P and Q have different truth values.

Definition 0.9 (Quantifier)**.** A **quantifier** [9] turns a predicate $P(x)$ (Definition 0.2) about the elements x of a set S into a proposition (Definition 0.1). There are two: the universal quantifier (Definition 0.10) and the existential quantifier (Definition 0.11).

Definition 0.10 (Universal quantifier)**.** The **universal quantifier** [10] "for all" is a quantifier (Definition 0.9): $\forall x \in S, P(x)$ ("for every x in S, $P(x)$") is true when $P(x)$ is true for every element x of S, and false otherwise.

Example 0.9 (A universally quantified proposition)**.** Let $P(x)$ be the predicate "x is even".

- "$\forall x \in \{2, 4, 6\}, P(x)$" is true, because 2, 4, and 6 are all even.
- "$\forall x \in \{2, 3, 4\}, P(x)$" is false, because $P(3)$ is false.

Definition 0.11 (Existential quantifier)**.** The **existential quantifier** [11] "there exists" is a quantifier (Definition 0.9): $\exists x \in S, P(x)$ ("there is an x in S such that $P(x)$") is true when $P(x)$ is true for at least one element x of S, and false otherwise.

Example 0.10 (An existentially quantified proposition)**.** Let $P(x)$ be the predicate "x is even".

- "$\exists x \in \{1, 2, 3\}, P(x)$" is true, because $P(2)$ is true.
- "$\exists x \in \{1, 3, 5\}, P(x)$" is false, because none of 1, 3, and 5 is even.

Example 0.11 (Quantified propositions)**.** Let $P(x)$ be the predicate "$x > 2$", for integers x.

- "$\exists x \in \{1, 2, 3\}, P(x)$" is true, because $P(3)$ is true.
- "$\forall x \in \{1, 2, 3\}, P(x)$" is false, because $P(1)$ is false.
- "$\forall x \in \{3, 4, 5\}, P(x)$" is true, because $3 > 2$, $4 > 2$, and $5 > 2$ are all true.

Definition 0.12 (Inequality)**.** An **inequality** is a proposition (Definition 0.1) or a predicate (Definition 0.2) that compares two expressions a and b whose values are real numbers, in one of the forms:

- $a < b$ ("a is less than b");
- $a \leq b$ ("a is less than or equal to b");
- $a > b$ ("a is greater than b");
- $a \geq b$ ("a is greater than or equal to b").

Example 0.12 (Inequalities)**.**

- "$3 < 5$" is an inequality, and it is true.
- "$5 \leq 3$" is an inequality, and it is false.
- "$x \geq 2$" is an inequality and a predicate: it is true when $x = 2$ and false when $x = 1$.
- "$x = 2$" is not an inequality: it is an equality.

Definition 0.13 (Logical entailment (logical consequence, deductive consequence))**.** Propositions or predicates $P_1, \ldots, P_n$ (Definition 0.1, Definition 0.2) **logically entail** [12] a proposition or predicate Q when every choice of values for their variables that makes all of $P_1, \ldots, P_n$ true also makes Q true. Then Q is a **logical consequence**, or **deductive consequence**, of $P_1, \ldots, P_n$.

Example 0.13 (Entailment, and its failure)**.** Let x be an integer.

- "$x = 2$" entails "$x^2 = 4$": the only value that makes "$x = 2$" true is $x = 2$, and $2^2 = 4$.
- "$x > 3$" and "$x < 5$" together entail "$x = 4$": 4 is the only integer greater than 3 and less than 5.
- "$x^2 = 4$" does not entail "$x = 2$": the value $x = -2$ makes "$x^2 = 4$" true, since $(-2)^2 = 4$, but makes "$x = 2$" false.

We can use any of:

- $\therefore$ (`\therefore` in LaTeX),
- $\Rightarrow$ (`\Rightarrow`),
- $\models$ (`\models`)

to denote logical entailment (Definition 0.13). $\Rightarrow$ also writes the connective "implies" (Definition 0.7); P entails Q exactly when $P \Rightarrow Q$ is true for every choice of values for the variables in P and Q.

Let's save $\rightarrow$ (`\rightarrow`) for convergence results.

Definition 0.14 (Proof)**.** A **proof** [13] of a proposition (Definition 0.1) is a finite list of statements that ends with that proposition, in which each statement is

- an assumption,
- a definition,
- a previously proved result, or
- a consequence of earlier statements in the list, by one rule of logic or algebra.

Example 0.14 (The sum of two even integers is even)**.** Let a and b be even integers (definition). Then $a = 2j$ and $b = 2k$ for some integers j and k, and

$$\begin{aligned} a + b &= 2j + 2k && \text{(substitute } a = 2j \text{ and } b = 2k\text{)} \\ &= 2(j + k) && \text{(distributive law)} \end{aligned}$$

The second line uses the distributive law. Since $j+k$ is an integer, $a+b$ is even (definition). For example,

$$\begin{aligned} 4 + 10 &= 2 \cdot 2 + 2 \cdot 5 \\ &= 2 \cdot 7 \\ &= 14. \end{aligned}$$

Definition 0.15 (Derivation)**.** A **derivation** is a proof (Definition 0.14), or a part of one, written as a chain of expressions, one per line, in which each line is joined to the line before it by $=$ or by an inequality sign (Definition 0.12), and carries an annotation that names the reason that step holds: an assumption, a definition, a previously proved result, or a rule of algebra.

Example 0.15 (A derivation with equalities and inequalities)**.** For every real number x:

$$\begin{aligned} (x+1)^2 &= x^2 + 2x + 1 && \text{(square of a sum)} \\ &\geq 2x + 1 && (x^2 \geq 0) \\ &> 2x && (1 > 0) \end{aligned}$$

The first step uses the square of a sum. The steps are $=$, $\geq$, and $>$, so the derivation shows $(x+1)^2 > 2x$. For example, with $x = 3$: $(3+1)^2 = 16$, and $16 > 6 = 2 \cdot 3$.

Theorem 0.1 (What a derivation shows)**.** *Consider a derivation (Definition 0.15) with expressions $E_0, E_1, \ldots, E_n$, where each line E_j is joined to the line E_{j-1} before it by $=$, $\leq$ or $<$.*

1. *If every step is $=$, then $E_0 = E_n$.*
2. *If every step is $=$ or $\leq$, then $E_0 \leq E_n$.*
3. *If every step is $=$, $\leq$, or $<$, and at least one step is $<$, then $E_0 < E_n$.*

The same holds with $\geq$ and $>$ in place of $\leq$ and $<$.

Proof

Proof. First, a fact about two steps in a row. For numbers a, b and c, if $a \; R \; b$ and $b \; S \; c$, where R and S are each one of $=$, $\leq$ and $<$, then $a \; T \; c$, where T is

- $<$ if R or S is $<$;
- otherwise $\leq$ if R or S is $\leq$;
- otherwise $=$.

This holds in each case:

- If R and S are both $=$, then $a = c$ (equalities are transitive).
- If exactly one of R and S is $=$, replace b by the equal expression in the other relation (substitution). This gives the other relation between a and c.
- If neither is $=$, then each of $a \; R \; b$ and $b \; S \; c$ is $\leq$ or $<$, and $\leq$ means $<$ or $=$ (strict and non-strict inequalities). A case with an $=$ is covered by the previous bullet, and

the remaining case is $a < b$ and $b < c$, which gives $a < c$ because $<$ is transitive on the real numbers. So two steps that are each $\leq$ or $<$ give $<$ if either is $<$, and otherwise $\leq$.

Now let T_j be the relation named by the same rule applied to the first j steps of the chain, with T_0 being $=$. We show $E_0 \; T_j \; E_j$ by induction on j. For $j = 0$, $E_0 = E_0$. If $E_0 \; T_j \; E_j$, then the step from E_j to E_{j+1} and the fact above give $E_0 \; T_{j+1} \; E_{j+1}$. At $j = n$, T_n is $=$ when every step is $=$, which is part 1. It is $\leq$ or $=$ when every step is $=$ or $\leq$, and $=$ implies $\leq$, which is part 2. It is $<$ when some step is $<$, which is part 3.
The argument for $\geq$ and $>$ is the same with the inequalities reversed. □

Example 0.16 (Checking a derivation line by line)**.** The derivation in Example 0.15 has the steps $=$, $\geq$ and $>$, so by Theorem 0.1 it shows $(x + 1)^2 > 2x$. The code evaluates the four lines of that derivation at several values of x, and checks each step and the comparison of the first line with the last.

```r
x <- c(-3, -1, 0, 2, 3)
lines <- data.frame(
  x = x,
  line_0 = (x + 1)^2,
  line_1 = x^2 + 2 * x + 1,
  line_2 = 2 * x + 1,
  line_3 = 2 * x
)
steps <- with(
  lines,
  data.frame(
    x = x,
    step_1_equal = line_0 == line_1,
    step_2_geq = line_1 >= line_2,
    step_3_greater = line_2 > line_3,
    first_greater_than_last = line_0 > line_3
  )
)
lines
#>    x line_0 line_1 line_2 line_3
#> 1 -3      4      4     -5     -6
#> 2 -1      0      0     -1     -2
#> 3  0      1      1      1      0
#> 4  2      9      9      5      4
#> 5  3     16     16      7      6
steps
#>    x step_1_equal step_2_geq step_3_greater first_greater_than_last
#> 1 -3         TRUE       TRUE           TRUE                    TRUE
#> 2 -1         TRUE       TRUE           TRUE                    TRUE
#> 3  0         TRUE       TRUE           TRUE                    TRUE
#> 4  2         TRUE       TRUE           TRUE                    TRUE
#> 5  3         TRUE       TRUE           TRUE                    TRUE
```

At all 5 values of x, the three steps hold in 5, 5 and 5 cases, and the first line exceeds the last in 5 cases, as Theorem 0.1 says. At $x = 0$, for example, the first line is 1 and the last is 0.

Definition 0.16 (Result)**.** A **result** is a proposition (Definition 0.1) that has a proof (Definition 0.14). These notes label each result by its role: as a theorem (Definition 0.17), a lemma (Definition 0.18), or a corollary (Definition 0.19).

Example 0.17 (A proposition that is a result, and one that is not)**.** "$2 + 3 = 5$" is a proposition (Definition 0.1) that has a proof, the computation of $2 + 3$, so it is a result. "7 is even" is a proposition that is false, so it has no proof and is not a result.

Remark ("Proposition" as a kind of result)

Remark 0.1 ("Proposition" as a kind of result)*.* Some sources also call a result of moderate importance a "proposition". In these notes, "proposition" always means a sentence that is true or false (Definition 0.1), whether or not it has been proved.

Definition 0.17 (Theorem)**.** A **theorem** [14] is a result (Definition 0.16) of interest in its own right.

Example 0.18 (A theorem)**.** "The sum of two even integers is even" is a result: Example 0.14 proves it. It is of interest in its own right, since it applies whenever two even integers are added, as in $2 + 4 = 6$, so it is a theorem.

Definition 0.18 (Lemma)**.** A **lemma** [15] is a result (Definition 0.16) proved mainly as a step in the proof of another result.

Example 0.19 (A lemma in the proof of a larger result)**.** To show that the sum of four even integers a, b, c, and d is even, write $a + b + c + d = (a + b) + (c + d)$. By the result of Example 0.14, $a + b$ and $c + d$ are even, and applying that result again to these two sums shows that their sum is even. Here that result is proved mainly as a step in the proof of another result, so it serves as a lemma. For example, $2 + 4 + 6 + 8 = (2 + 4) + (6 + 8) = 6 + 14 = 20$, which is even.

Definition 0.19 (Corollary)**.** A **corollary** [16] is a result (Definition 0.16) that follows from a theorem with little extra work.

Example 0.20 (A corollary of a result)**.** Example 0.14 proves the result "the sum of two even integers is even". A corollary follows with little extra work: the sum of three even integers a, b, and c is even, because $a + b + c = (a + b) + c$, $a + b$ is even by that result, and applying the result again to the even integers $a + b$ and c shows that $(a + b) + c$ is even. For example,

$$\begin{aligned} 2+4+6 &= (2+4)+6 \\ &= 6+6 \\ &= 12, \end{aligned}$$

which is even.

Definition 0.20 (Counterexample)**.** A **counterexample** [17] to a proposition of the form "for every x in A, $P(x)$", where $P(x)$ is a predicate (Definition 0.2), is an element x of A for which $P(x)$ is false. A single counterexample shows that the proposition is false.

Example 0.21 (A counterexample)**.** The proposition "for every integer n, $n+n>n$" is false. The integer $n=0$ is a counterexample: $0+0=0$, and $0>0$ is false. The proposition is true for some integers, such as $n=3$, since $3+3=6>3$, but one counterexample is enough to make it false.

Definition 0.21 (Converse)**.** The **converse** [18] of the proposition "if P, then Q" is the proposition "if Q, then P".

Example 0.22 (A true proposition with a false converse)**.** For an integer n, call n a multiple of 4 if $n=4k$ for some integer k.

- "If n is a multiple of 4, then n is even" is true:

  $$\begin{aligned} n &= 4k \\ &= 2\cdot(2k), \end{aligned}$$

  and $2k$ is an integer.

- Its converse, "if n is even, then n is a multiple of 4", is false. The integer $n=2$ is a counterexample (Definition 0.20): 2 is even, but $2=4k$ only for $k=\frac{1}{2}$, which is not an integer.

So a proposition can be true while its converse is false.

Definition 0.22 (Sufficient condition)**.** When the proposition "if P, then Q" is true, P is a **sufficient condition** [19] for Q.

Example 0.23 (A sufficient condition)**.** For a real number x, "if $x > 5$, then $x > 2$" is true, because a number greater than 5 is greater than 2. So $x > 5$ is a sufficient condition for $x > 2$: knowing that $x = 6$ satisfies $x > 5$ is enough to conclude that $6 > 2$.

Definition 0.23 (Necessary condition)**.** When the proposition "if P, then Q" is true, Q is a **necessary condition** [19] for P.

Example 0.24 (A necessary condition)**.** For an integer n, "if n is a multiple of 6, then n is a multiple of 3" is true, because $6k = 3 \cdot (2k)$ for every integer k. So being a multiple of 3 is a necessary condition for being a multiple of 6: $12 = 6 \cdot 2$ is a multiple of 6, and it is also a multiple of 3, since $12 = 3 \cdot 4$.

Definition 0.24 (Necessary and sufficient condition)**.** When both "if P, then Q" and its converse (Definition 0.21) are true, P is a **necessary and sufficient condition** [19] for Q (Definition 0.22, Definition 0.23), written "P if and only if Q".

Example 0.25 (Necessary but not sufficient)**.** In Example 0.22, "if n is a multiple of 4, then n is even" is true, so:

- being a multiple of 4 is a sufficient condition for n to be even;
- being even is a necessary condition for n to be a multiple of 4.

Being even is not a sufficient condition for being a multiple of 4: $n = 2$ is even, but not a multiple of 4.

Definition 0.25 (Proof by contradiction)**.** A **proof by contradiction** [20] of a proposition P assumes that P is false, and derives from that assumption a proposition that is known to be false.

Example 0.26 (No integer is both even and odd)**.** Suppose, for contradiction, that some integer n is both even and odd (definition). Then $n = 2j$ and $n = 2k + 1$ for some integers j and k, and

$$\begin{aligned} 2j &= 2k + 1 && \text{(both equal } n\text{)} \\ 2j - 2k &= 1 && \text{(subtract } 2k \text{ from both sides)} \\ 2(j - k) &= 1 && \text{(distributive law)} \\ j - k &= \tfrac{1}{2} && \text{(divide both sides by 2)} \end{aligned}$$

The third line uses the distributive law. But $j - k$ is an integer, and $\frac{1}{2}$ is not, so the last line is false. So no integer is both even and odd. For example, $6 = 2 \cdot 3$ is even, and $6 = 2k + 1$ would need $k = 2.5$, which is not an integer.

Theorem 0.2 (A proof by contradiction proves its proposition)**.** *Let P and Q be propositions (Definition 0.1). If $\neg P \Rightarrow Q$ is true and Q is false, then P is true.*

Proof

Proof. By Definition 0.7, $\neg P \Rightarrow Q$ is false only when $\neg P$ is true and Q is false. Here Q is false, and $\neg P \Rightarrow Q$ is true, so $\neg P$ is not true. Then $\neg P$ is false. Since $\neg P$ is true whenever P is false, P must be true. □

Example 0.27 (A truth table for a proof by contradiction)**.** There are only four ways to assign truth values to P and Q. The code lists them, with the truth value of $\neg P \Rightarrow Q$, and marks the rows where $\neg P \Rightarrow Q$ is true and Q is false.

```
tt <- expand.grid(Q = c(TRUE, FALSE), P = c(TRUE, FALSE))
tt$not_P_implies_Q <- !(!tt$P) | tt$Q
tt$hypotheses_hold <- tt$not_P_implies_Q & !tt$Q
tt
#>       Q     P not_P_implies_Q hypotheses_hold
#> 1  TRUE  TRUE            TRUE           FALSE
#> 2 FALSE  TRUE            TRUE            TRUE
#> 3  TRUE FALSE            TRUE           FALSE
#> 4 FALSE FALSE           FALSE           FALSE
```

The hypotheses of Theorem 0.2 hold in 1 of the 4 rows, and in 1 of them P is true. So whenever the hypotheses hold, P is true.

Don't skip steps

Show every step of the derivation. Each line should follow from the previous line by a single, identifiable move:

- applying a definition,
- substituting a known result, or
- performing one algebraic operation.

When you skip steps, you force the reader to reconstruct your reasoning, which is exactly the

work the proof was supposed to do *for* them. A step that feels "obvious" while you are writing it is often the step a reader gets stuck on. Phrases like "it can be shown that" or "clearly" usually mark a skipped step; show the step instead.

Annotate each step

For each step, note the definition, theorem, or algebraic rule that justifies it. A brief annotation to the right of the line is usually enough. Annotations make the proof checkable, and they teach the reader which tool to reach for in similar situations.

For example, here is a derivation that the hat matrix $\mathbf{H} = \mathbf{X}(\mathbf{X}^\top\mathbf{X})^{-1}\mathbf{X}^\top$ is idempotent, $\mathbf{H}^2 = \mathbf{H}$, with each step annotated:

$$
\begin{aligned}
\mathbf{H}^2 &= \mathbf{H}\mathbf{H} && \text{(definition of a matrix power)} \\
&= \mathbf{X}(\mathbf{X}^\top\mathbf{X})^{-1}\mathbf{X}^\top\,\mathbf{X}(\mathbf{X}^\top\mathbf{X})^{-1}\mathbf{X}^\top && \text{(definition of } \mathbf{H}\text{)} \\
&= \mathbf{X}(\mathbf{X}^\top\mathbf{X})^{-1}(\mathbf{X}^\top\mathbf{X})(\mathbf{X}^\top\mathbf{X})^{-1}\mathbf{X}^\top && \text{(regroup; matrix multiplication is associative)} \\
&= \mathbf{X}\,\mathbf{I}_p\,(\mathbf{X}^\top\mathbf{X})^{-1}\mathbf{X}^\top && \text{(a matrix inverse times the matrix is } \mathbf{I}_p\text{)} \\
&= \mathbf{X}(\mathbf{X}^\top\mathbf{X})^{-1}\mathbf{X}^\top && \text{(the identity matrix changes nothing)} \\
&= \mathbf{H} && \text{(definition of } \mathbf{H}\text{)}
\end{aligned}
$$

The Linear Algebra page uses this derivation to show that $\mathbf{H}$ is a projection matrix. The probability notes derive the variance identity $\operatorname{Var}(X) = \mathrm{E}[X^2] - (\mathrm{E}[X])^2$ in the same annotated style.

Direct proof

Definition 0.26 (Direct proof)**.** A **direct proof** [21] of a proposition Q starts from the assumptions that Q is stated under, if any, and reaches Q through a chain of steps, each of which applies a definition, a previously proved result, or one rule of logic or algebra to the assumptions or to earlier steps. A direct proof never assumes that Q is false; compare proof by contradiction, which does.

Example 0.28 (The square of an odd integer is odd)**.** Claim: if n is an odd integer, then n^2 is odd.
Since n is odd, there is an integer k with $n = 2k + 1$. Then

$$\begin{aligned} n^2 &= (2k+1)^2 && \text{(substitute } n = 2k+1\text{)} \\ &= (2k)^2 + 2 \cdot 2k \cdot 1 + 1^2 && \text{(square of a sum)} \\ &= 4k^2 + 4k + 1 && \text{(arithmetic)} \\ &= 2(2k^2 + 2k) + 1 && \text{(distributive law, read from right to left)} \end{aligned}$$

and $2k^2 + 2k$ is an integer, because sums and products of integers are integers. So $n^2 = 2 \cdot (\text{an integer}) + 1$, which is the definition of odd. For example,

$$\begin{aligned} n &= 7 \\ &= 2 \cdot 3 + 1 \end{aligned}$$

gives

$$\begin{aligned} n^2 &= 49 \\ &= 2 \cdot 24 + 1, \end{aligned}$$

and $2 \cdot 3^2 + 2 \cdot 3 = 24$.
The proof above that the sum of two even integers is even is also a direct proof.

Follow the golden rule

In general, follow the golden rule: treat your readers the way you want to be treated as a reader.

When you read someone else's proof, you want to be able to follow every step without guessing, to know which result is being used at each line, and to never be left wondering where a quantity came from. Write your own proofs to meet that same standard.

Proof by induction

Many proofs in these notes show that a statement holds for every natural number n, or for every integer n from some starting value on. Checking the statement one n at a time would never finish; induction proves it for all of them with two arguments.

Definition 0.27 (Proof by induction)**.** Let n_0 be an integer, and let $P(n)$ be a predicate about the integers $n \geq n_0$. A **proof by induction** [22] on n shows that $P(n)$ holds for every integer $n \geq n_0$ by showing two things:

1. **Base case:** $P(n_0)$ holds.
2. **Inductive step:** for every integer $n > n_0$, if $P(n-1)$ holds, then $P(n)$ holds.

In the inductive step, the assumption that $P(n-1)$ holds is called the **induction**

hypothesis.

ℹ Remark (Why the two steps suffice)

Remark 0.2 (Why the two steps suffice). The base case gives $P(n_0)$. The inductive step with $n = n_0 + 1$ then gives $P(n_0 + 1)$, with $n = n_0 + 2$ it gives $P(n_0 + 2)$, and so on: each integer $n \geq n_0$ is reached after $n - n_0$ uses of the inductive step.
When $P(n)$ is needed only for $n_0 \leq n \leq r$, it is enough to show the inductive step for $n_0 < n \leq r$; the same chain then stops at $P(r)$.

The following example uses summation notation.

Example 0.29 (Sum of the first n natural numbers)**.** For every $n \in \mathbb{N}$,

$$\sum_{i=1}^{n} i = \frac{n(n+1)}{2}.$$

For example, with $n = 3$, $1 + 2 + 3 = 6$ and $\frac{3 \cdot 4}{2} = 6$. To prove the equation for every n, let $P(n)$ be the equation and use Definition 0.27 with $n_0 = 1$.
Base case, $n = 1$**.** The sum has one term:

$$\sum_{i=1}^{1} i = 1 \quad \text{(definition of a sum)}$$

and the right side is

$$\begin{aligned} \frac{1\,(1+1)}{2} &= \frac{1 \cdot 2}{2} && (1 + 1 = 2) \\ &= \frac{2}{2} && (1 \cdot 2 = 2) \\ &= 1. && (\tfrac{2}{2} = 1) \end{aligned}$$

So $P(1)$ holds.
Inductive step. Let $n \geq 2$, and suppose $P(n-1)$ holds:

$$\sum_{i=1}^{n-1} i = \frac{(n-1)\,((n-1)+1)}{2}.$$

Then

$$
\begin{aligned}
\sum_{i=1}^{n} i &= 1 + 2 + \cdots + (n-1) + n && \text{(definition of a sum)} \\
&= (1 + 2 + \cdots + (n-1)) + n && \text{(associative law)} \\
&= \sum_{i=1}^{n-1} i + n && \text{(definition of a sum)} \\
&= \frac{(n-1)\,((n-1)+1)}{2} + n && \text{(induction hypothesis)} \\
&= \frac{(n-1)\,n}{2} + n && ((n-1)+1 = n) \\
&= \frac{(n-1)\,n}{2} + \frac{2n}{2} && (n = \tfrac{2n}{2}) \\
&= \frac{(n-1)\,n + 2n}{2} && \text{(common denominator)} \\
&= \frac{((n-1)+2)\,n}{2} && \text{(distributive law)} \\
&= \frac{(n+1)\,n}{2} && ((n-1)+2 = n+1) \\
&= \frac{n(n+1)}{2}, && \text{(commutative law)}
\end{aligned}
$$

which is $P(n)$. By Definition 0.27, the equation holds for every $n \in \mathbb{N}$.

Formal logic

Definition 0.28 (Propositional logic)**. Propositional logic** [23] is the study of logical entailment among propositions built from basic propositions $P, Q, R, \ldots$ using only the logical connectives. Whether such propositions entail one another depends only on the truth values of the basic propositions, not on what the basic propositions say.

Example 0.30 (P and $P \Rightarrow Q$ entail Q)**.** Suppose P is true and $P \Rightarrow Q$ is true. If Q were false, then $P \Rightarrow Q$ would have a true P and a false Q, so $P \Rightarrow Q$ would be false, which contradicts the assumption. So Q is true. This holds whatever P and Q say, so P and $P \Rightarrow Q$ logically entail Q.

Definition 0.29 (First-order logic)**. First-order logic** [24] extends propositional logic (Definition 0.28) by also allowing:

- variables $x, y, \ldots$ that stand for elements of a set;

- predicates of one or more of those variables, such as $P(x)$ or $x < y$;
- functions of them, such as $x + y$;
- equality between them;
- quantifiers $\forall$ and $\exists$ over them.

Example 0.31 (From "for all" to one element)**.** Let $P(x)$ be a predicate about the elements of a set S, and let $a \in S$. The propositions $\forall x \in S, P(x)$ and $a \in S$ logically entail $P(a)$: if $P(x)$ is true for every element x of S, then it is true for the element a. For example, $\forall x \in \mathbb{Z}, x^2 \geq 0$ entails $3^2 \geq 0$. Propositional logic alone cannot express this entailment, because it has no way to refer to the elements of S.

Definition 0.30 (Formal proof)**.** A **formal proof** [25] is a proof written in propositional logic (Definition 0.28) or first-order logic (Definition 0.29) in which each step is an assumption or follows from earlier steps by one rule taken from a fixed list of allowed rules, so that checking the proof needs no understanding of what its propositions mean.

Example 0.32 (A three-step formal proof)**.** Suppose the allowed rules include "from P and $P \Rightarrow Q$, conclude Q". Then this is a formal proof of Q from the assumptions P and $P \Rightarrow Q$:

1. P (assumption)
2. $P \Rightarrow Q$ (assumption)
3. Q (from steps 1 and 2, by the rule above)

Further reading

- Velleman [26] is a structured introduction to proof techniques such as direct proof, proof by contradiction, and induction.
- Barker-Plummer et al. [27] treats proofs as formal proofs in propositional logic and first-order logic.

Free course materials on formal logic:

- Thomas Icard's Phil 150/250, Mathematical Logic course notes (Stanford) review sets, relations and functions, then cover propositional, modal and first-order logic.
- John Wilcox's logic handouts supplement the same Stanford course, with a handout per week on the syntax and semantics of propositional and first-order logic, natural deduction and metalogic.
- Eric Pacuit's Phil 151: First Order Logic (Stanford, Winter 2009) posts a syllabus, a modal logic study guide and problem sets.

References

[1] Wikipedia contributors, "Proposition — Wikipedia, the free encyclopedia." 2026. Available: https://en.wikipedia.org/wiki/Proposition

[2] Wikipedia contributors, "Predicate (mathematical logic) — Wikipedia, the free encyclopedia." 2026. Available: https://en.wikipedia.org/wiki/Predicate_(mathematical_logic)

[3] Wikipedia contributors, "Logical connective — Wikipedia, the free encyclopedia." 2026. Available: https://en.wikipedia.org/wiki/Logical_connective

[4] Wikipedia contributors, "Negation — Wikipedia, the free encyclopedia." 2026. Available: https://en.wikipedia.org/wiki/Negation

[5] Wikipedia contributors, "Logical conjunction — Wikipedia, the free encyclopedia." 2026. Available: https://en.wikipedia.org/wiki/Logical_conjunction

[6] Wikipedia contributors, "Logical disjunction — Wikipedia, the free encyclopedia." 2026. Available: https://en.wikipedia.org/wiki/Logical_disjunction

[7] Wikipedia contributors, "Material conditional — Wikipedia, the free encyclopedia." 2026. Available: https://en.wikipedia.org/wiki/Material_conditional

[8] Wikipedia contributors, "Logical biconditional — Wikipedia, the free encyclopedia." 2026. Available: https://en.wikipedia.org/wiki/Logical_biconditional

[9] Wikipedia contributors, "Quantifier (logic) — Wikipedia, the free encyclopedia." 2026. Available: https://en.wikipedia.org/wiki/Quantifier_(logic)

[10] Wikipedia contributors, "Universal quantification — Wikipedia, the free encyclopedia." 2026. Available: https://en.wikipedia.org/wiki/Universal_quantification

[11] Wikipedia contributors, "Existential quantification — Wikipedia, the free encyclopedia." 2026. Available: https://en.wikipedia.org/wiki/Existential_quantification

[12] Wikipedia contributors, "Logical consequence — Wikipedia, the free encyclopedia." 2026. Available: https://en.wikipedia.org/wiki/Logical_consequence

[13] Wikipedia contributors, "Mathematical proof — Wikipedia, the free encyclopedia." 2026. Available: https://en.wikipedia.org/wiki/Mathematical_proof

[14] Wikipedia contributors, "Theorem — Wikipedia, the free encyclopedia." 2026. Available: https://en.wikipedia.org/wiki/Theorem

[15] Wikipedia contributors, "Lemma (mathematics) — Wikipedia, the free encyclopedia." 2026. Available: https://en.wikipedia.org/wiki/Lemma_(mathematics)

[16] Wikipedia contributors, "Corollary — Wikipedia, the free encyclopedia." 2026. Available: https://en.wikipedia.org/wiki/Corollary

[17] Wikipedia contributors, "Counterexample — Wikipedia, the free encyclopedia." 2026. Available: https://en.wikipedia.org/wiki/Counterexample

[18] Wikipedia contributors, "Converse (logic) — Wikipedia, the free encyclopedia." 2026. Available: https://en.wikipedia.org/wiki/Converse_(logic)

[19] Wikipedia contributors, "Necessity and sufficiency — Wikipedia, the free encyclopedia." 2026. Available: https://en.wikipedia.org/wiki/Necessity_and_sufficiency

[20] Wikipedia contributors, "Proof by contradiction — Wikipedia, the free encyclopedia." 2026. Available: https://en.wikipedia.org/wiki/Proof_by_contradiction

[21] Wikipedia contributors, "Direct proof — Wikipedia, the free encyclopedia." 2026. Available: https://en.wikipedia.org/wiki/Direct_proof

[22] Wikipedia contributors, "Mathematical induction — Wikipedia, the free encyclopedia." 2026. Available: https://en.wikipedia.org/wiki/Mathematical_induction

[23] Wikipedia contributors, "Propositional calculus — Wikipedia, the free encyclopedia." 2026. Available: https://en.wikipedia.org/wiki/Propositional_calculus

[24] Wikipedia contributors, "First-order logic — Wikipedia, the free encyclopedia." 2026. Available: https://en.wikipedia.org/wiki/First-order_logic

[25] Wikipedia contributors, "Formal proof — Wikipedia, the free encyclopedia." 2026. Available: https://en.wikipedia.org/wiki/Formal_proof

[26] D. J. Velleman, *How to prove it: A structured approach*, 3rd ed. Cambridge University Press, 2019. doi: 10.1017/9781108539890.

[27] D. Barker-Plummer, J. Barwise, and J. Etchemendy, *Language, proof and logic*, 2nd ed. Stanford, CA: CSLI Publications, 2011. Available: https://www.amazon.com/dp/1575866323