# Double Integrals and Fubini's Theorem

Last modified: 2026-10-14 11:20:47 (PDT)

## Double Integrals

The **Fubini–Tonelli theorem** states conditions under which the order of integration in a double integral can be exchanged. We state two versions: the Riemann version (Theorem 0.1) is what applied courses usually use for double integrals of continuous functions on simple regions; the $\sigma$-finite measure-theoretic version (Theorem 0.2) is included to make the joint-distribution form corollary in *Probability for Data Science* follow from a stated theorem rather than from an aside.

**Definition 0.1** (Double integral)**.** Let $f$ be a bounded function on a closed, bounded plane region $R \subseteq \mathbb{R}^2$. Cover $R$ with a grid of rectangles, keep the $n$ rectangles that lie entirely inside $R$, with areas $\Delta A_1, \dots, \Delta A_n$, and choose a point $(x_i, y_i)$ in the $i$-th rectangle. The **double integral** of $f$ over $R$ is

$$\iint_R f(x,y)\,dA \overset{\text{def}}{=} \lim_{\|\Delta\|\to 0} \sum_{i=1}^{n} f(x_i, y_i)\,\Delta A_i,$$

where $\|\Delta\|$ is the length of the longest diagonal among the $n$ rectangles, when that limit exists and has the same value for every choice of grids and of the points $(x_i, y_i)$. The symbol $dA$ stands for an element of area.
(Larson and Edwards 2018, sec. 14.2)

**Example 0.1** (The double integral of 1 is an area)**.** Let $f(x,y) = 1$ on the rectangle $R = [0,2] \times [0,3]$. Every sum in Definition 0.1 adds up the areas of rectangles inside $R$, and those sums approach the area of $R$ as the grid gets finer, so

$$\begin{aligned}\iint_R 1\,dA &= 2 \cdot 3 \\ &= 6.\end{aligned}$$

**Definition 0.2** (Iterated integral)**.** An **iterated integral** is an integral of an integral:

$$\int_a^b \int_{g_1(x)}^{g_2(x)} f(x,y)\,dy\,dx \stackrel{\text{def}}{=} \int_a^b \left( \int_{g_1(x)}^{g_2(x)} f(x,y)\,dy \right) dx$$

That is, integrate over the inner variable ($y$) first, holding the outer variable ($x$) fixed, and then integrate the result over the outer variable. The inner limits $g_1(x)$ and $g_2(x)$ may depend on the outer variable, or be constants $c$ and $d$, as in $\int_a^b \int_c^d f(x,y)\,dy\,dx$. Iterated integrals in the other order, $\int\int \cdots\, dx\,dy$, are defined the same way with the roles of $x$ and $y$ swapped.

**Example 0.2** (An iterated integral)**.** Integrating over $y$ first, then $x$:

$$\begin{aligned}
\int_0^1 \int_0^2 xy\,dy\,dx &= \int_0^1 \left( \int_0^2 xy\,dy \right) dx && \text{(definition of the iterated integral)} \\
&= \int_0^1 x \left( \int_0^2 y\,dy \right) dx && (x \text{ is constant in } y) \\
&= \int_0^1 x \left[ \frac{y^2}{2} \right]_{y=0}^{y=2} dx && \text{(antiderivative of } y) \\
&= \int_0^1 2x\,dx && \text{(evaluate at the limits)} \\
&= \left[ x^2 \right]_{x=0}^{x=1} && \text{(antiderivative of } 2x) \\
&= 1 && \text{(evaluate at the limits)}
\end{aligned}$$

**Theorem 0.1** (Fubini's theorem (Riemann version))**.** *Let $f$ be **continuous** on a plane region $R \subseteq \mathbb{R}^2$.*

1. ***Vertically simple region.*** *If $R$ is defined by $a \le x \le b$ and $g_1(x) \le y \le g_2(x)$, where $g_1$ and $g_2$ are continuous on $[a,b]$, then*

$$\iint_R f(x,y)\,dA = \int_a^b \int_{g_1(x)}^{g_2(x)} f(x,y)\,dy\,dx.$$

2. ***Horizontally simple region.*** *If $R$ is defined by $c \le y \le d$ and $h_1(y) \le x \le h_2(y)$, where $h_1$ and $h_2$ are continuous on $[c,d]$, then*

$$\iint_R f(x,y)\,dA = \int_c^d \int_{h_1(y)}^{h_2(y)} f(x,y)\,dx\,dy.$$

*When $R$ can be described both ways, the two iterated integrals are equal — so the order of integration can be exchanged.*
*(Larson and Edwards 2018, Theorem 14.2, p. 982)*

**Example 0.3** (Changing the order of integration for a non-rectangular region)**.** Adapted from (Larson and Edwards 2018, sec. 14.2, Example 4, pp. 984–985).
Let $X$ and $Y$ be independent Uniform$(0,1)$ random variables, with joint density $f(x,y) = 1$ on the unit square $[0,1]^2$. Define the function $g(x,y) = \mathrm{e}^{-x^2}\,\mathbb{1}(y \le x)$, where $e$ is Euler's number, and compute its expectation $\mathrm{E}[g(X,Y)]$.
Because the joint density equals 1 on $[0,1]^2$, this expectation is the double integral of $g$ over the unit square:

$$\mathrm{E}[g(X,Y)] = \iint_{[0,1]^2} g(x,y)\,dA.$$

The indicator factor $\mathbb{1}(y \le x)$ equals 1 on the triangular region where $y \le x$ and 0 elsewhere, so only that region, namely $D = \{(x,y) : x \in [0,1],\ y \in [0,x]\}$ (Figure 1), contributes, and there $g(x,y) = \mathrm{e}^{-x^2}$:

$$\mathrm{E}[g(X,Y)] = \iint_{D} \mathrm{e}^{-x^2}\,dA.$$

```r
region <- tibble::tibble(x = c(0, 1, 1), y = c(0, 0, 1))
ggplot2::ggplot(region, ggplot2::aes(x = x, y = y)) +
  ggplot2::geom_polygon(
    fill = "steelblue", alpha = 0.4, color = "black", linewidth = 0.7
  ) +
  ggplot2::annotate(
    "text", x = 0.65, y = 0.28, label = "D", size = 8, fontface = "italic"
  ) +
  ggplot2::annotate(
    "text", x = 0.36, y = 0.52, label = "y == x",
    parse = TRUE, angle = 45
  ) +
  ggplot2::annotate(
    "text", x = 1.08, y = 0.5, label = "x == 1",
    parse = TRUE, angle = 90, hjust = 0.5
  ) +
  ggplot2::annotate(
    "text", x = 0.5, y = -0.1, label = "y == 0", parse = TRUE
  ) +
  ggplot2::scale_x_continuous(breaks = c(0, 0.5, 1), limits = c(-0.1, 1.3)) +
  ggplot2::scale_y_continuous(breaks = c(0, 0.5, 1), limits = c(-0.2, 1.2)) +
  ggplot2::coord_equal() +
  ggplot2::labs(x = "x", y = "y") +
  ggplot2::theme_minimal()
```





Figure 1: Triangular integration region $D = \{(x, y) : x \in [0, 1],\ y \in [0, x]\}$, bounded below by $y = 0$, above-left by $y = x$, and on the right by $x = 1$.

**Order $dx\,dy$ is intractable.** Re-describing $D$ as $D = \{(x, y) : y \in [0, 1],\ x \in [y, 1]\}$, the inner integral is

$$\int_y^1 \mathrm{e}^{-x^2}\,dx,$$

which cannot be computed with FTC Part 2: no antiderivative of $\mathrm{e}^{-x^2}$ can be written as a finite formula built from powers, exponentials, logarithms, and trigonometric functions (an **elementary** antiderivative).

**Order $dy\,dx$ works.** Applying Theorem 0.1 Part 1 ($\mathrm{e}^{-x^2}$ is continuous and $D$ is the vertically simple region $x \in [0, 1]$, $y \in [0, x]$):

$$\begin{aligned}
\iint_D \mathrm{e}^{-x^2}\,dA &= \int_0^1 \int_0^x \mathrm{e}^{-x^2}\,dy\,dx && \text{(Fubini, vertically simple region)} \\
&= \int_0^1 \mathrm{e}^{-x^2}\left(\int_0^x dy\right) dx && (\mathrm{e}^{-x^2} \text{ is constant in } y) \\
&= \int_0^1 x\,\mathrm{e}^{-x^2}\,dx && \left(\textstyle\int_0^x dy = x\right)
\end{aligned}$$

To antidifferentiate $x\,\mathrm{e}^{-x^2}$, let $u = x^2$ be the inner function. By the chain rule (Theorem 10 in Derivatives and Taylor Series), with $\frac{du}{dx} = 2x$,

$$\begin{aligned}
\frac{d}{dx}\left(-\frac{1}{2}\,\mathrm{e}^{-u}\right) &= -\frac{1}{2}\,\mathrm{e}^{-u}\cdot(-1)\cdot\frac{du}{dx} && \text{(chain rule)} \\
&= -\frac{1}{2}\,\mathrm{e}^{-x^2}\cdot(-1)\cdot 2x && (\text{substitute } u = x^2 \text{ and } du/dx = 2x) \\
&= x\,\mathrm{e}^{-x^2} && \text{(multiply)}
\end{aligned}$$

so $-\frac{1}{2}\,\mathrm{e}^{-x^2}$ is an antiderivative of $x\,\mathrm{e}^{-x^2}$, and

$$\begin{aligned}
\int_0^1 x\,\mathrm{e}^{-x^2}\,dx &= \left[-\frac{1}{2}\,\mathrm{e}^{-x^2}\right]_0^1 && \text{(FTC Part 2)} \\
&= -\frac{1}{2}(\mathrm{e}^{-1} - \mathrm{e}^{0}) && \text{(evaluate at the limits)} \\
&= -\frac{1}{2}(\mathrm{e}^{-1} - 1) && (\mathrm{e}^0 = 1) \\
&= \frac{1 - \mathrm{e}^{-1}}{2} && (\text{distribute } -\tfrac{1}{2}) \\
&= \frac{e - 1}{2e} && (\text{multiply numerator and denominator by } e) \\
&\approx 0.316
\end{aligned}$$

The solid whose volume equals this integral is shown in Figure 2.

*(This 3D surface is interactive; see the HTML or slide version of this page to view it.)*

Figure 2: Surface $z = e^{-x^2}$ over the region $D = \{(x, y) : x \in [0, 1],\ y \in [0, x]\}$. The surface depends only on $x$ (constant in $y$), so for each $x$ the inner integral over $y \in [0, x]$ contributes $x \cdot e^{-x^2}$.

**Exercise 0.1** (Double integral over a triangular region)**.** Adapted from Miller (2016), Question 1.1.50.
Evaluate the integral:

$$\int_0^1 \int_0^x xy \, dy \, dx$$

and confirm the result by reversing the order of integration.

Solution

*Solution* 0.1. **1. Given order ($y$ then $x$):**

$$\begin{aligned}\int_0^x xy \, dy &= x\left[\frac{y^2}{2}\right]_0^x \\ &= \frac{x^3}{2} \\ \int_0^1 \frac{x^3}{2} \, dx &= \left[\frac{x^4}{8}\right]_0^1 \\ &= \frac{1}{8}\end{aligned}$$

**2. Reversed order ($x$ then $y$):** The triangular region $T = \{(x, y) : 0 \le x \le 1, 0 \le y \le x\}$ is equivalently described by $T = \{(x, y) : 0 \le y \le 1, y \le x \le 1\}$:

$$\begin{aligned}\int_0^1 \int_y^1 xy \, dx \, dy &= \int_0^1 y\left[\frac{x^2}{2}\right]_{x=y}^1 dy \\ &= \int_0^1 \frac{y(1-y^2)}{2} \, dy \\ \int_0^1 \left(\frac{y}{2} - \frac{y^3}{2}\right) dy &= \left[\frac{y^2}{4} - \frac{y^4}{8}\right]_0^1 \\ &= \frac{1}{4} - \frac{1}{8} \\ &= \frac{1}{8}\end{aligned}$$

Both orders yield 1/8.

**Exercise 0.2** (Switching order of integration on a non-rectangular region)**.** Adapted from Miller (2016), Question 1.1.51.
Express the integral:

$$\int_0^1 \int_0^x ye^{-xy}\,dy\,dx$$

by reversing the order of integration.

Solution

*Solution* 0.2. In the given order, the inner integral $\int_0^x ye^{-xy}\,dy$ requires integration by parts with respect to $y$.
Switching the order over the triangular domain $\{(x, y) : 0 \le y \le 1, y \le x \le 1\}$:

$$\int_0^1 \int_y^1 ye^{-xy}\,dx\,dy$$

Now the inner integral has the factor $y$ in place for direct integration with respect to $x$:

$$\begin{aligned}\int_y^1 ye^{-xy}\,dx &= [-e^{-xy}]_{x=y}^1 \\ &= e^{-y^2} - e^{-y}\end{aligned}$$

The double integral becomes:

$$\begin{aligned}\int_0^1 (e^{-y^2} - e^{-y})\,dy &= \int_0^1 e^{-y^2}\,dy - [-e^{-y}]_0^1 \\ &= \int_0^1 e^{-y^2}\,dy - (1 - e^{-1})\end{aligned}$$

The term $\int_0^1 e^{-y^2}\,dy$ has no elementary antiderivative; it can be written using the error function $\operatorname{erf}(z) = \frac{2}{\sqrt{\pi}} \int_0^z e^{-t^2}\,dt$ as $\frac{\sqrt{\pi}}{2}\operatorname{erf}(1) \approx 0.7468$. The overall value is:

$$\frac{\sqrt{\pi}}{2}\operatorname{erf}(1) + e^{-1} - 1 \approx 0.1147$$

**Example 0.4** (When conditions fail: a counterexample)**.** The conditions in Theorem 0.1 are not merely technical — when they fail, iterated integrals can exist yet disagree.

Let

$$f(x,y) = \frac{x^2 - y^2}{(x^2+y^2)^2}$$

on the unit square $R = [0,1] \times [0,1]$. Strictly, $f$ is defined on $R \setminus \{(0,0)\}$: the denominator vanishes at the origin, so $f$ is undefined there (we return to this point in the condition check).

**Integrating $y$ first, then $x$:**

Holding $x$ fixed and differentiating in $y$ (by the quotient rule, Theorem 9 in Derivatives and Taylor Series),

$$\begin{aligned}\frac{d}{dy}\frac{y}{x^2+y^2} &= \frac{(x^2+y^2) - y \cdot 2y}{(x^2+y^2)^2} \\ &= \frac{x^2 - y^2}{(x^2+y^2)^2}.\end{aligned}$$

(A derivative in one variable with the others held fixed is a partial derivative, defined on the vector calculus page.) The arctangent arctan is the inverse of the tangent function from trigonometry; all this example needs is that $\frac{d}{dx}\arctan(x) = \frac{1}{1+x^2}$, $\arctan(0) = 0$, and $\arctan(1) = \frac{\pi}{4}$.

$$\begin{aligned}\int_0^1 \int_0^1 f(x,y)\,dy\,dx &= \int_0^1 \left[\frac{y}{x^2+y^2}\right]_{y=0}^{y=1} dx \\ &= \int_0^1 \frac{1}{x^2+1}\,dx \\ &= [\arctan(x)]_0^1 \\ &= \frac{\pi}{4}\end{aligned}$$

**Integrating $x$ first, then $y$:**

Holding $y$ fixed and differentiating in $x$,

$$\begin{aligned}\frac{d}{dx}\left(-\frac{x}{x^2+y^2}\right) &= -\frac{(x^2+y^2) - x \cdot 2x}{(x^2+y^2)^2} \\ &= \frac{x^2 - y^2}{(x^2+y^2)^2}\ :\end{aligned}$$

$$\begin{aligned}\int_0^1 \int_0^1 f(x,y)\,dx\,dy &= \int_0^1 \left[-\frac{x}{x^2+y^2}\right]_{x=0}^{x=1} dy \\ &= \int_0^1 \left(-\frac{1}{1+y^2}\right) dy \\ &= -[\arctan(y)]_0^1 \\ &= -\frac{\pi}{4}\end{aligned}$$

**Conclusion:** $\frac{\pi}{4} \neq -\frac{\pi}{4}$, so the two iterated integrals are unequal. Theorem 0.1 does not apply here.

**Why Theorem 0.1's condition fails:** Theorem 0.1 requires $f$ to be **continuous** on $R$. The denominator $(x^2+y^2)^2$ vanishes at the origin $(0,0) \in R$, so $f$ is *not even defined* there — let alone continuous — and the theorem does not apply.

(Wikipedia contributors 2024)

Figure 3 shows the surface. The failure comes from the origin, where $f$ is undefined and unbounded.

*(This 3D surface is interactive; see the HTML or slide version of this page to view it.)*

Figure 3: Surface $f(x,y) = (x^2-y^2)/(x^2+y^2)^2$ on $[0,1]^2$, sampled away from the origin and clipped to $[-50,50]$ for display. Approaching the origin, the function grows without bound along the $x$-axis (red ridge, $f > 0$ when $|x| > |y|$) and falls without bound along the $y$-axis (blue ridge, $f < 0$ when $|y| > |x|$). Because $f$ is undefined at $(0,0)$, $f$ is not continuous on $R$ and Theorem 0.1 does not apply.

**Corollary 0.1** (Continuous functions on a rectangle (corollary of Theorem 0.1))**.** *If $f : [a,b] \times [c,d] \to \mathbb{R}$ is **continuous** on the closed bounded rectangle $[a,b] \times [c,d]$, then:*

$$\begin{aligned}\int_a^b \left(\int_c^d f(x,y)\,dy\right) dx &= \int_c^d \left(\int_a^b f(x,y)\,dx\right) dy \\ &= \iint_{[a,b]\times[c,d]} f(x,y)\,dA.\end{aligned}$$

*(Larson and Edwards 2018, Theorem 14.2, p. 982)*

Proof

*Proof.* A closed bounded rectangle $[a,b] \times [c,d]$ is both vertically simple (with $g_1 \equiv c$, $g_2 \equiv d$) and horizontally simple (with $h_1 \equiv a$, $h_2 \equiv b$). Applying both parts of Theorem 0.1 to $f$ on this rectangle gives the two iterated forms shown. □

**Example 0.5** (Evaluating a double integral on a rectangle)**.** Structure adapted from (Larson and Edwards 2018, sec. 14.2, Example 2, pp. 982–983); the integrand $x^2 + y^2$ is original, chosen so the integral equals $\mathrm{E}[g(X, Y)]$ for $g(x, y) = x^2 + y^2$.

Let $X$ and $Y$ be independent Uniform$(0, 1)$ random variables, with joint density $f(x, y) = 1$ on the unit square $R = \{(x, y) : x \in [0, 1],\ y \in [0, 1]\}$ (Figure 4). Define the function $g(x, y) = x^2 + y^2$, and compute its expectation $\mathrm{E}[g(X, Y)]$.

Because the joint density equals 1 on $R$, this expectation is the double integral of $g$ over $R$:

$$\mathrm{E}[g(X, Y)] = \iint_R (x^2 + y^2)\, dA.$$

```r
ggplot2::ggplot() +
  ggplot2::annotate(
    "rect", xmin = 0, xmax = 1, ymin = 0, ymax = 1,
    fill = "steelblue", alpha = 0.4, color = "black", linewidth = 0.7
  ) +
  ggplot2::annotate(
    "text", x = 0.5, y = 0.5, label = "R", size = 8, fontface = "italic"
  ) +
  ggplot2::scale_x_continuous(breaks = c(0, 1), limits = c(-0.15, 1.25)) +
  ggplot2::scale_y_continuous(breaks = c(0, 1), limits = c(-0.15, 1.25)) +
  ggplot2::coord_equal() +
  ggplot2::labs(x = "x", y = "y") +
  ggplot2::theme_minimal()
```





Figure 4: Integration region $R = [0,1]^2$, the unit square.

The integrand is continuous on $R$, so Corollary 0.1 applies and either order of integration yields the same value.

**Integrating $y$ first, then $x$:**

$$\begin{aligned}
\mathrm{E}[g(X,Y)] &= \int_0^1 \int_0^1 (x^2+y^2)\,dy\,dx \\
&= \int_0^1 \left[x^2 y + \frac{y^3}{3}\right]_0^1 dx \\
&= \int_0^1 \left(x^2 + \frac{1}{3}\right) dx \\
&= \left[\frac{x^3}{3} + \frac{x}{3}\right]_0^1 \\
&= \frac{2}{3}
\end{aligned}$$

**Integrating $x$ first, then $y$** (verifying the order can be swapped):

$$\begin{aligned}
\int_0^1 \int_0^1 (x^2+y^2)\,dx\,dy &= \int_0^1 \left[\frac{x^3}{3} + y^2 x\right]_0^1 dy \\
&= \int_0^1 \left(\frac{1}{3} + y^2\right) dy \\
&= \left[\frac{y}{3} + \frac{y^3}{3}\right]_0^1 \\
&= \frac{2}{3}
\end{aligned}$$

Both orders give $\frac{2}{3}$, as Corollary 0.1 guarantees.
As a cross-check, linearity of expectation gives the same value: since

$$\begin{aligned}
\mathrm{E}[X^2] &= \int_0^1 x^2\,dx \\
&= \frac{1}{3}
\end{aligned}$$

for $X \sim \mathrm{Uniform}(0,1)$ (and likewise for $Y$),

$$\begin{aligned}
\mathrm{E}[g(X,Y)] &= \mathrm{E}[X^2 + Y^2] \\
&= \mathrm{E}[X^2] + \mathrm{E}[Y^2] \\
&= \frac{1}{3} + \frac{1}{3} \\
&= \frac{2}{3}.
\end{aligned}$$

The solid whose volume equals this integral is shown in Figure 5.

*(This 3D surface is interactive; see the HTML or slide version of this page to view it.)*

Figure 5: Surface $z = x^2 + y^2$ over the unit square $[0, 1]^2$. The double integral $\frac{2}{3}$ is the volume between this surface and the $xy$-plane, and equals $\mathrm{E}[X^2 + Y^2]$.

**Exercise 0.3** (Double integral of a polynomial over a rectangle)**.** Adapted from Miller (2016), Question 1.1.47.
Evaluate the double integral:

$$\int_0^2 \int_0^3 5(x^2y + xy^2 + 2)\, dy\, dx$$

Solution

*Solution* 0.3. Integrate with respect to $y$ first:

$$\begin{aligned}\int_0^3 (x^2y + xy^2 + 2)\, dy &= \left[\frac{x^2y^2}{2} + \frac{xy^3}{3} + 2y\right]_{y=0}^{3} \\ &= \frac{9}{2}x^2 + 9x + 6\end{aligned}$$

Now integrate with respect to $x$:

$$\begin{aligned}5\int_0^2 \left(\frac{9}{2}x^2 + 9x + 6\right) dx &= 5\left[\frac{3}{2}x^3 + \frac{9}{2}x^2 + 6x\right]_0^2 \\ &= 5\left[\frac{3}{2}(8) + \frac{9}{2}(4) + 6(2)\right] \\ &= 5[12 + 18 + 12] \\ &= 5(42) \\ &= 210\end{aligned}$$

*(Note: The source text carried an arithmetic slip in the inner coefficient resulting in* 190*; the exact value is* 210*.)*

**Exercise 0.4** (Choosing the order of integration)**.** Adapted from Miller (2016), Question 1.1.48.
Evaluate the double integral:

$$\int_0^6 \int_0^5 xe^{-xy}\, dy\, dx$$

Solution

*Solution* 0.4. Integrating with respect to $x$ first requires integration by parts. Integrating with respect to $y$ first is much simpler because the factor of $x$ is already present:

$$\begin{aligned}\int_0^5 xe^{-xy}\,dy &= [-e^{-xy}]_{y=0}^5 \\ &= 1 - e^{-5x}\end{aligned}$$

Now integrate with respect to $x$:

$$\begin{aligned}\int_0^6 (1 - e^{-5x})\,dx &= \left[x + \frac{1}{5}e^{-5x}\right]_0^6 \\ &= \left(6 + \frac{1}{5}e^{-30}\right) - \left(0 + \frac{1}{5}\right) \\ &= \frac{29}{5} + \frac{e^{-30}}{5}\end{aligned}$$

**Exercise 0.5** (Product of single integrals for separable functions)**.** Adapted from Miller (2016), Question 1.1.49.
Let $m, n > 0$. Evaluate:

$$\int_0^1 \int_0^1 x^m y^n \,dy\,dx$$

Solution

*Solution* 0.5. Because the region of integration is a rectangle $[0,1]\times[0,1]$ and the integrand factors as $g(x)h(y) = x^m \cdot y^n$, the double integral splits into the product of two single-variable integrals:

$$\int_0^1 \int_0^1 x^m y^n \,dy\,dx = \left(\int_0^1 x^m \,dx\right)\left(\int_0^1 y^n \,dy\right)$$

Evaluating each factor:

$$\int_0^1 x^m \,dx = \frac{1}{m+1}, \qquad \int_0^1 y^n \,dy = \frac{1}{n+1}$$

Therefore:

$$\int_0^1 \int_0^1 x^m y^n \,dy\,dx = \frac{1}{(m+1)(n+1)}$$

*Remark:* When joint random variables $X$ and $Y$ are independent, their joint density factors as $f_{X,Y}(x,y) = f_X(x)f_Y(y)$, and joint probabilities over product sets factor in exactly this way.

**Exercise 0.6** (Double integral with polynomial and radical terms)**.** Adapted from Miller (2016), Question 1.1.52.
Evaluate the double integral:

$$\int_0^1 \int_0^1 (x^2 + 2xy + y\sqrt{x})\, dy\, dx$$

Solution

*Solution* 0.6. Integrate with respect to $y$ first:

$$\begin{aligned}\int_0^1 (x^2 + 2xy + y\sqrt{x})\, dy &= \left[x^2 y + xy^2 + \frac{y^2\sqrt{x}}{2}\right]_{y=0}^{1} \\ &= x^2 + x + \frac{1}{2}x^{1/2}\end{aligned}$$

Now integrate with respect to $x$:

$$\begin{aligned}\int_0^1 \left(x^2 + x + \frac{1}{2}x^{1/2}\right) dx &= \left[\frac{x^3}{3} + \frac{x^2}{2} + \frac{1}{2}\cdot\frac{x^{3/2}}{3/2}\right]_0^1 \\ &= \left[\frac{x^3}{3} + \frac{x^2}{2} + \frac{1}{3}x^{3/2}\right]_0^1 \\ &= \frac{1}{3} + \frac{1}{2} + \frac{1}{3} \\ &= \frac{2}{3} + \frac{1}{2} \\ &= \frac{7}{6}\end{aligned}$$

**Exercise 0.7** (Double integral of an affine function)**.** Adapted from Miller (2016), Question 1.1.53.
Let $a, b, c$ be constants. Evaluate:

$$\int_0^1 \int_0^1 (ax + by + c)\, dy\, dx$$

Solution

*Solution* 0.7. By linearity of the integral:

$$\begin{aligned}\int_0^1 \int_0^1 (ax+by+c)\,dy\,dx &= a\int_0^1 x\,dx \int_0^1 1\,dy + b\int_0^1 1\,dx \int_0^1 y\,dy + c\int_0^1 1\,dx \int_0^1 1\,dy \\ &= a\left(\frac{1}{2}\right)(1) + b(1)\left(\frac{1}{2}\right) + c(1)(1) \\ &= \frac{a}{2} + \frac{b}{2} + c\end{aligned}$$

**Exercise 0.8** (Harmonic numbers via double integrals)**.** Adapted from Miller (2016), Question 1.1.54.
Prove that for every positive integer $n$:

$$\begin{aligned}\int_0^1 \int_0^1 n(1-xy)^{n-1}\,dx\,dy &= \sum_{k=1}^{n} \frac{1}{k} \\ &= 1 + \frac{1}{2} + \cdots + \frac{1}{n}\end{aligned}$$

Solution

*Solution* 0.8. **Method 1 (integration then geometric series):** Integrate with respect to $x$ first:

$$\begin{aligned}\int_0^1 n(1-xy)^{n-1}\,dx &= \left[-\frac{(1-xy)^n}{y}\right]_{x=0}^{1} \\ &= \frac{1-(1-y)^n}{y}\end{aligned}$$

Use the finite geometric series identity $\sum_{k=0}^{n-1} r^k = \frac{1-r^n}{1-r}$ with $r = 1-y$:

$$\frac{1-(1-y)^n}{y} = \sum_{k=0}^{n-1}(1-y)^k$$

Now integrate each term with respect to $y$ from 0 to 1:

$$\begin{aligned}\int_0^1 (1-y)^k\,dy &= \left[-\frac{(1-y)^{k+1}}{k+1}\right]_0^1 \\ &= \frac{1}{k+1}\end{aligned}$$

Summing over $k = 0, 1, \ldots, n-1$:

$$\begin{aligned}\sum_{k=0}^{n-1} \frac{1}{k+1} &= \sum_{j=1}^{n} \frac{1}{j} \\ &= 1 + \frac{1}{2} + \cdots + \frac{1}{n}\end{aligned}$$

**Method 2 (binomial expansion):** Expand $(1-xy)^{n-1} = \sum_{k=0}^{n-1} \binom{n-1}{k}(-1)^k (xy)^k$. Integrating over $[0,1]^2$:

$$\int_0^1 \int_0^1 (xy)^k \, dx \, dy = \left(\frac{1}{k+1}\right)^2$$

Multiplying by $n$ and using the identity $\frac{n}{k+1}\binom{n-1}{k} = \binom{n}{k+1}$:

$$\begin{aligned}\int_0^1 \int_0^1 n(1-xy)^{n-1} \, dx \, dy &= \sum_{k=0}^{n-1} \binom{n}{k+1} \frac{(-1)^k}{k+1} \\ &= \sum_{j=1}^{n} \frac{1}{j}\end{aligned}$$

*Remark:* This integral identity connects multivariable integration with harmonic numbers $H_n = \sum_{k=1}^{n} \frac{1}{k} \approx \log n + \gamma$, where $\gamma \approx 0.5772$ is the Euler-Mascheroni constant.

**Theorem 0.2** (Fubini–Tonelli theorem (measure-theoretic form))**.** *Let $(\Omega_1, \mathcal{F}_1, \mu_1)$ and $(\Omega_2, \mathcal{F}_2, \mu_2)$ be measure spaces with $\sigma$-finite measures, and let $f : \Omega_1 \times \Omega_2 \to \mathbb{R}$ be measurable with respect to the product $\sigma$-algebra $\mathcal{F}_1 \otimes \mathcal{F}_2$. If either*

*(a) $f \geq 0$ almost everywhere with respect to the product measure $\mu_1 \otimes \mu_2$ (**Tonelli's theorem**), or*

*(b) $\int_{\Omega_1 \times \Omega_2} |f| \, d(\mu_1 \otimes \mu_2) < \infty$, that is, $f$ is absolutely integrable (**Fubini's theorem**),*

*then both iterated integrals exist, agree with the double integral, and equal each other:*

$$\begin{aligned}\int_{\Omega_1 \times \Omega_2} f \, d(\mu_1 \otimes \mu_2) &= \int_{\Omega_1} \left( \int_{\Omega_2} f(\omega_1, \omega_2) \, d\mu_2(\omega_2) \right) d\mu_1(\omega_1) \\ &= \int_{\Omega_2} \left( \int_{\Omega_1} f(\omega_1, \omega_2) \, d\mu_1(\omega_1) \right) d\mu_2(\omega_2).\end{aligned}$$

*(Billingsley 1995, Theorem 18.3; Gut 2013, Theorem 9.1, p. 65; Fubini 1907; Wikipedia contributors 2024)*

*Remark* 0.1 (Fubini–Tonelli for probability measures). Applied courses rarely need the measure-theoretic generalization itself, but it is what justifies the joint-distribution form corollary in *Probability for Data Science*. A probability measure $P$ on $\Omega$ has $P(\Omega) = 1 < \infty$, so it is finite, and hence $\sigma$-finite (finite and $\sigma$-finite measures); for probability measures, the $\sigma$-finiteness condition is automatic.
The integrability conditions (nonnegativity or absolute integrability) still need to be verified in each application. For example, Lebesgue measure (ordinary length) on $[0, 1]$ is a probability measure, so the $\sigma$-finiteness condition holds for both factors $[0, 1]$, yet the two iterated integrals in Example 0.4 are $\pi/4$ and $-\pi/4$. So $\sigma$-finiteness alone does not make the iterated integrals agree.

**Example 0.6** (Positive application of Theorem 0.2)**.** Let $X$ and $Y$ be independent Exponential(1) random variables, with joint density $f(x, y) = e^{-(x+y)}$ for $x, y \geq 0$.
The probability $P(X \leq 1, Y \leq 1)$ is the integral of $f$ over $[0, 1]^2$ with respect to Lebesgue measure (ordinary length) in each coordinate. Lebesgue measure on $[0, \infty)$ is $\sigma$-finite, because $[0, \infty)$ is the union of the intervals $[0, n]$, $n \in \mathbb{N}$, each of finite length $n$; so the $\sigma$-finiteness condition of Theorem 0.2 holds. Since $f(x, y) = e^{-(x+y)} \geq 0$, condition (a) (Tonelli's theorem, nonnegativity) is also satisfied.
By Theorem 0.2, both iterated integrals exist and agree. Integrating $y$ first, then $x$:

$$\begin{aligned}
P(X \leq 1, Y \leq 1) &= \int_0^1 \int_0^1 e^{-(x+y)}\, dy\, dx \\
&= \int_0^1 \int_0^1 e^{-x} e^{-y}\, dy\, dx && \text{(exponential of a sum)} \\
&= \int_0^1 e^{-x} \left( \int_0^1 e^{-y}\, dy \right) dx && (e^{-x} \text{ is constant in } y) \\
&= \int_0^1 e^{-x} [-e^{-y}]_{y=0}^{y=1}\, dx && \text{(antiderivative of } e^{-y}) \\
&= \int_0^1 e^{-x}(1 - e^{-1})\, dx && \text{(evaluate at the limits)} \\
&= (1 - e^{-1}) \int_0^1 e^{-x}\, dx && (1 - e^{-1} \text{ is constant in } x) \\
&= (1 - e^{-1})[-e^{-x}]_{x=0}^{x=1} && \text{(antiderivative of } e^{-x}) \\
&= (1 - e^{-1})^2 && \text{(evaluate at the limits)}
\end{aligned}$$

Integrating $x$ first, then $y$:

$$
\begin{aligned}
P(X \le 1,\, Y \le 1) &= \int_0^1 \int_0^1 e^{-(x+y)}\, dx\, dy \\
&= \int_0^1 \int_0^1 e^{-y} e^{-x}\, dx\, dy && \text{(exponential of a sum)} \\
&= \int_0^1 e^{-y} \left( \int_0^1 e^{-x}\, dx \right) dy && (e^{-y} \text{ is constant in } x) \\
&= \int_0^1 e^{-y} [-e^{-x}]_{x=0}^{x=1}\, dy && \text{(antiderivative of } e^{-x}) \\
&= \int_0^1 e^{-y}(1 - e^{-1})\, dy && \text{(evaluate at the limits)} \\
&= (1 - e^{-1}) \int_0^1 e^{-y}\, dy && (1 - e^{-1} \text{ is constant in } y) \\
&= (1 - e^{-1})[-e^{-y}]_{y=0}^{y=1} && \text{(antiderivative of } e^{-y}) \\
&= (1 - e^{-1})^2 && \text{(evaluate at the limits)}
\end{aligned}
$$

Both iterated integrals equal $(1-e^{-1})^2 \approx 0.400$, as Theorem 0.2 guarantees when condition (a) holds.

**Example 0.7** (When neither Fubini–Tonelli condition is satisfied)**.** The same function $f(x, y) = (x^2 - y^2)/(x^2 + y^2)^2$ from Example 0.4 illustrates a case where neither condition of Theorem 0.2 is satisfied.

**Why Theorem 0.2's conditions fail:** $\iint_R |f|\, dA = \infty$, which violates condition (b). Switching to polar coordinates $(r, \theta)$ near the origin, the integrand satisfies

$$
\begin{aligned}
|f(x, y)| &= \left|x^2 - y^2\right| / (x^2 + y^2)^2 \\
&= |\cos 2\theta| / r^2,
\end{aligned}
$$

so

$$
\begin{aligned}
\iint_R |f|\, dA &\ge \int_0^{\pi/2} \int_0^{\varepsilon} \frac{|\cos 2\theta|}{r^2}\, r\, dr\, d\theta \\
&= \left( \int_0^{\pi/2} |\cos 2\theta|\, d\theta \right) \int_0^{\varepsilon} \frac{dr}{r} \\
&= +\infty,
\end{aligned}
$$

since $\int_0^{\varepsilon} dr/r$ diverges. Therefore $\iint_R |f|\, dA = \infty$, and condition (b) of Theorem 0.2 is not satisfied. (Condition (a) also fails: $f$ takes both positive and negative values, so it is not nonnegative a.e.) The unequal iterated integrals from Example 0.4 are thus consistent

> with Theorem 0.2: the theorem simply does not apply.
> (Wikipedia contributors 2024)


Billingsley, Patrick. 1995. *Probability and Measure*. 3rd ed. Wiley Series in Probability and Mathematical Statistics. Wiley.

Fubini, Guido. 1907. "Sugli Integrali Multipli." *Rendiconti Della Reale Accademia Dei Lincei. Classe Di Scienze Fisiche, Matematiche e Naturali* 16: 608–14.

Gut, Allan. 2013. *Probability: A Graduate Course*. 2nd ed. Springer Texts in Statistics. Springer. https://doi.org/10.1007/978-1-4614-4708-5.

Larson, Ron, and Bruce H. Edwards. 2018. *Calculus*. 11th ed. Cengage Learning. https://www.cengage.com/c/calculus-11e-larson/.

Miller, Steven J. 2016. *The Probability Lifesaver: Calculus Review Problems*. https://web.williams.edu/Mathematics/sjmiller/public_html/probabilitylifesaver/index.htm#:~:text=http%3A//web.williams.edu/Mathematics/sjmiller/public_html/probabilitylifesaver/supplementalchap_calcreview.pdf.

Wikipedia contributors. 2024. *Fubini's Theorem — Wikipedia, the Free Encyclopedia*. https://en.wikipedia.org/wiki/Fubini%27s_theorem.